Entanglement of four-qubit systems: a geometric atlas with polynomial compass II (the tame world)

Frédéric Holweck $^{a)}$, Jean-Gabriel Luque $^{b)}$ and Jean-Yves Thibon $^{c)}$

We propose a new approach to the geometry of the four-qubit entanglement classes depending on parameters. More precisely, we use invariant theory and algebraic geometry to describe various stratifications of the Hilbert space by SLOCC invariant algebraic varieties. The normal forms of the four-qubit classification of Verstraete *et al.* are interpreted as dense subsets of components of the dual variety of the set of separable states and an algorithm based on the invariants/covariants of the four-qubit quantum states is proposed to identify a state with a SLOCC equivalent normal form (up to qubits permutation).

Keywords: Quantum Information Theory, Entangled states, Dual variety, Classical invariant theory, Symmetric functions.

PACS: 02.40.-k, 03.65.Fd, 03.67.-a, 03.65.Ud

^{a)} frederic.holweck@utbm.fr, IRTES-UTBM, Universit {e} de Bourgogne Franche-Comt {e}, 90010 Belfort Cedex, FR

^{b)} jean-gabriel.luque@univ-rouen.fr, Université de Rouen, Laboratoire d'Informatique, du Traitement de l'Information et des Systèmes (LITIS), Avenue de l'Université - BP 8 6801 Saint-étienne-du-Rouvray Cedex, FR

^{c)} jyt@univ-mlv.fr, Laboratoire d'Informatique Gaspard Monge Université Paris-Est Marne-la-Vallée, 77454 Marne-la-Vallée Cedex 2, FR

I. INTRODUCTION

Entanglement is nowdays considered as a central ressource in quantum information processing. The large amount of work produced since the beginning of the XXIst century to understand the nature of entanglement demonstrates the interest of the community in this subject (see the review papers^{14,27} and the references therein). The question of the classification of entanglement of pure multipartite quantum states under the group SLOCC (Stochastic Local Operations with Classical Communication) is one of the most prominent challenging problems on the road to the understanding of entanglement. The complexity of this question grows up exponentially with the number of parts composing the quantum system. If the classification problem is completely solved for a few cases, it is known to be out of reach for most general situation (see Ref³⁷ for a discussion on the computational complexity of the algebraic invariants of five-qubit systems).

The case of the four-qubit Hilbert space is of special importance. First it has generated a large number of papers by itself^{2,6,7,16,31,33,36,38,43}. Then, as long as qubits are concerned, it is the only case where the number of SLOCC orbits is infinite but for which there still exists a list of 9 normal forms found by Verstraete *et al.*^{7,43} (6 of them depend on parameters) which, up to permutation of the qubits, parametrize all SLOCC orbits (i.e. up to permutation of the qubits, a given state is in one of the orbit of the 9 families). The four-qubit case also showed up in different contexts like in the study of the black-holes-qubits correspondence³, in the study of graph-states²⁰ and in many quantum communication protocols like error correcting codes¹⁹. Thus we consider the four-qubit case as rich enough to motivate more investigation and develop new tools which we hope to be useful to study more difficult cases.

This paper is part of a sequence of articles^{25,26} where we have investigated the structure of entanglement for small multipartite systems by combining two different approches: on one side we consider classical invariant theory and we try to understand what the invariants, covariants of the multipartite systems tell us about the SLOCC-orbits and, on the other side, we look at the geometrical structure of the space by building SLOCC algebraic varieties. Combining the two approaches, we obtain a stratification of the Hilbert space by SLOCC algebraic varieties (closure of classes or of union of classes) and criteria based on invariants/covariants, to distinguish them. In our first paper²⁵ we provided a geometric description of all entanglement classes with invariants/covariants criteria for Hilbert spaces with a finite number of orbits. In Ref²⁶ we started to consider the four-qubit case by first looking at specific subvarieties of this Hilbert space. The first subvariety we considered was the nullcone, i.e. the variety of states which annihilate all SLOCC invariant polynomials. It turns out that the number of orbits contained in this variety is finite and thus the techniques developped in Ref²⁵ could apply, leading to a classification, of the nilpotent four-qubit states, using covariants. In the same paper we also considered the case of the algebraic variety $\sigma_3(X)$, the third secant variety (see below for the definitions), defined by the simultaneous vanishing of L and M, the two degree four generators of the ring of SLOCC invariant polynomials (see below for the definition). This latter case already contains an infinite number of orbits.

In this paper we investigate the geometry of the four-qubit Hilbert space by considering specific SLOCC invariant hypersurfaces: the hypersurfaces defined by the vanishing of the invariants of degree four L, M and N and the so-called hyperdeterminant¹⁸ Δ of format $2 \times 2 \times 2 \times 2$. The idea of studing the geometry of the hypersurface defined by Δ to classify entanglement classes appeared ten years ago in the work of Miyake³⁸. The paper of Miyake is based on the study of the singular locus of the hypersurface described ten years earlier by Weymann and Zelevinsky⁴⁴. In this article we go further in the description of the stratification of the ambient space by singular locus of the hyperdeterminant. We consider singularities leading to deeper stratas and we are able to establish a connection between the statas of the ambient space defined by the singularities of Δ and the 9 families given by Verstraete *et al.*'s classification. Like in our previous papers^{25,26} the geometric approach is combined with a classical invariant theory point of view allowing us to identify the strata to which a given state belongs. It leads to an algorithm based on invariants and covariants which provides, up to a qubit permutation, the family and parameters of a representative SLOCC equivalent to a given state.

The paper is organized as follow. In Section II we illustrate how the adjoint representation of SO(8) on its Lie algebra $\mathfrak{so}(8)$ is connected to the SLOCC (= $SL_2(\mathbb{C})^{\times 4}$) orbits of $\mathcal{H} = \mathbb{C}^2 \times \mathbb{C}^2 \times \mathbb{C}^2 \times \mathbb{C}^2$ by showing that the generators of the ring of invariants of four-qubit can be obtained by restriction of the generators of the SO(8) polynomial invariants on $\mathfrak{so}(8)$. Similarly we show that the discriminant of the adjoint representation of SO(8), also known as the defining equation of the dual of the adjoint variety, leads to the $2 \times 2 \times 2 \times 2$ hyperdeterminant. In the proof, the hyperdeterminant is obtained as the discriminant of a quartic. This quartic depends on the embedding of \mathcal{H} in $\mathfrak{so}(8)$. In Section III we investigate the three quartics corresponding to the three natural embeddings of \mathcal{H} in $\mathfrak{so}(8)$. It leads to a first stratification of the ambient space based on the roots of the quartics. In particular, we show that this stratification provides a diagram of normal forms which is connected to the geometry of a special semiregular polytope: the demitesseract. In Section IV the geometry of SLOCC hypersurfaces corresponding to specific dual varieties is investigated. This section ends with a geometric stratification of the ambient space where the families of the four-qubit classification are in correspondence with geometric stratas. Moreover some of those stratas are given by a simple geometric interpretation in terms of duals of orbits of well known quantum states. Finally, combining Section III and IV we give an algorithm based on invariants and covariants of four-qubit to determine the Verstraete type of a given state. Section VI is dedicated to concluding remarks.

Notations

All along the paper a four-qubit state $|\varphi\rangle$ will be denoted by $|\varphi\rangle = \sum_{i_1,...,i_4 \in \{0,1\}^4} a_{i_1...i_4} |i_1 \cdots |i_4\rangle$ where $|i_1 \cdots |i_4\rangle$ stand for the vectors of the computational basis. When not specified the Hilbert space \mathcal{H} will be the space of pure four-qubit quantum states, i.e. $\mathcal{H} = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes$ $\mathbb{C}^2 \otimes \mathbb{C}^2$ and the corresponding SLOCC group will be SLOCC= $SL_2(\mathbb{C})^{\times 4}$. When we work over the projective space $\mathbb{P}(\mathcal{H})$, an algebraic variety $Z \subset \mathbb{P}(\mathcal{H})$ is defined as the zero locus of a collection of homogeneous polynomials. For any subset $Y \subset \mathbb{P}(V)$, the notation \overline{Y} will refer to the Zariski closure²¹.

II. ADJOINT VARIETY OF SO(8)

The classification of four-qubit pure states under SLOCC is strongly connected to the Lie algebra \mathfrak{so}_8 . In the original paper of Verstraete *et al.*⁴³, the action of the $SO(4) \times SO(4)$ subgroup of SO(8), on the space of 8×8 matrices is used to provide a first classification of SLOCC orbits of $\mathcal{H} = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$. This idea is made more precise, and the classification corrected, in the work of Chterental and Djoković⁷ where the Lie algebra \mathfrak{so}_8 is decomposed under the involution defined by the diagonal matrix $g = \begin{pmatrix} I_4 & 0 \\ 0 & -I_4 \end{pmatrix}$:

$$\mathfrak{so}_8 = \mathfrak{k} \oplus \mathfrak{p}. \tag{1}$$

The $SO(4) \times SO(4)$ invariant subspace **p** is

$$\mathfrak{p} = \left\{ \begin{pmatrix} 0 & R \\ -R^T & 0 \end{pmatrix}, R \in \mathcal{M}_4(\mathbb{C}) \right\}$$
(2)

The action of $SO(4) \times SO(4)$ on \mathfrak{p} corresponds to the action of SLOCC on $\mathcal{H} = \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$. The correspondence is given as follows. Let $|\varphi\rangle = \sum_{i,j,k,l \in \{0,1\}} a_{ijkl} |ijkl\rangle$ be a four-qubit state. One associates with $|\varphi\rangle$ the 4 × 4 matrix

$$M_{\varphi} = \begin{pmatrix} a_{0000} & a_{0010} & a_{0001} & a_{0011} \\ a_{1000} & a_{1010} & a_{1001} & a_{1011} \\ a_{0100} & a_{0110} & a_{0101} & a_{0111} \\ a_{1100} & a_{1110} & a_{1101} & a_{1111} \end{pmatrix}.$$
(3)

Let (A_1, A_2, A_3, A_4) be an element of the group SLOCC= $SL_2(\mathbb{C})^{\times 4}$. The action on M_{φ} is given by $(A_1 \otimes A_2)M_{\varphi}(A_3 \otimes A_4)^t$. This action on the space of 4×4 matrices becomes an action of $SO(4) \times SO(4)$ on \mathfrak{p} . This is described in Ref⁷ via the following unitary matrix (see also Ref⁴³):

$$T = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & i & i & 0 \\ 0 & -1 & 1 & 0 \\ i & 0 & 0 & -i \end{pmatrix}.$$
 (4)

One gets the correspondence: $(A_1, A_2, A_3, A_4) \in \text{SLOCC}$ acts on \mathcal{H} and $\begin{pmatrix} P_1 & 0 \\ 0 & P_2 \end{pmatrix} \in SO(4) \times SO(4)$ acts on \mathfrak{p} with $P_1 = T(A_1 \otimes A_2)T^{\dagger}$ and $P_2 = T(A_3 \otimes A_4)T^{\dagger}$ such that

$$A_1 \otimes A_2 \otimes A_3 \otimes A_4 |\varphi\rangle = \begin{pmatrix} P_1 & 0 \\ 0 & P_2 \end{pmatrix} \begin{pmatrix} 0 & TM_{\varphi}T^{\dagger} \\ -(TM_{\varphi}T^{\dagger})^t & 0 \end{pmatrix} \begin{pmatrix} P_1 & 0 \\ 0 & P_2 \end{pmatrix}^{-1}.$$
 (5)

The action of $SO(4) \times SO(4)$ on \mathfrak{p} is nothing but the trace of the adjoint action of SO(8)on \mathfrak{p} . This embedding of the vector space \mathcal{H} into \mathfrak{so}_8 leads to the following observation.

Proposition II.1. Let $\mathbb{C}[\mathfrak{so}_8]^{SO(8)}$ be the ring of invariant polynomials on \mathfrak{so}_8 for the adjoint action of SO(8) and $\mathbb{C}[\mathcal{H}]^{SLOCC}$ the ring of invariant polynomials for the SLOCC action on the four-qubit Hilbert space. The restriction map

$$\mathbb{C}[\mathfrak{so}_8]^{SO(8)} \longrightarrow \mathbb{C}[\mathfrak{p}]^{SO(4) \times SO(4)} = \mathbb{C}[\mathcal{H}]^{SLOCC}$$
(6)

is an isomorphism. Moreover the restriction of the equation defining the dual variety of the adjoint orbit $X_{SO(8)} \subset \mathbb{P}(\mathfrak{so}_8)$ to \mathfrak{p} is the $2 \times 2 \times 2 \times 2$ hyperdeterminant.

Proof. It is well known⁴¹ that the ring of invariant polynomials $\mathbb{C}[\mathfrak{so}_{2n}]^{SO(2n)}$ is a free algebra generated by homogeneous algebraically independent polynomials of degree 2, 4, ..., 2(n - 1), n. Let us denote by $\mathcal{M}_{2n}^0(\mathbb{C})$ the space of traceless matrices of size $2n \times 2n$, and let us recall that $\mathfrak{so}_{2n} = \{A \in \mathcal{M}_{2n}^0(\mathbb{C}), A = -A^t\}$. The generators of $\mathbb{C}[\mathfrak{so}_{2n}]^{SO(2n)}$ can be obtained as the restriction to \mathfrak{so}_{2n} of the sum of the $i \times i$ principal minors. For skew symmetric matrices only the even dimensionnal principal minors do not vanish. Moreover the determinant of a skew symmetric matrix factorizes as the square of the Pfaffian, Pf. In other words the generators of $\mathbb{C}[\mathfrak{so}_{2n}]^{SO(2n)}$ can be chosen to be $h_2, \ldots, h_{2n-2}, Pf$ where h_{2i} is the sum of the $2i \times 2i$ principal minors. In particular if we consider \mathfrak{so}_8 , the generators are four polynomials of degree 2, 4, 6, 4. The ring of invariant polynomials for four-qubit states under SLOCC³⁶ is also a free algebra generated by homogeneous algebraically independent polynomials of degree 2, 4, 4, 6. Let us denote them respectively B, L, M and D, their descriptions will be given in the next section. The restriction of the sum of principal minors h_2, h_4, h_6 and the Pfaffian Pf to \mathfrak{p} , i.e. to the vector space

$$\begin{pmatrix} 0 & TM_{\varphi}T^{\dagger} \\ -(TM_{\varphi}T^{\dagger})^t & 0 \end{pmatrix}$$
(7)

leads to the following equalities: $h_{2|\mathfrak{p}} = 2B$, $h_{4|\mathfrak{p}} = B^2 + 2L + 4M$, $h_{6|\mathfrak{p}} = 2BL + 4BM - 4D$ and $Pf_{|\mathfrak{p}} = L$. It immediately proves that $\mathbb{C}[\mathfrak{so}_8]^{SO(8)}_{|\mathfrak{p}} = \mathbb{C}[\mathcal{H}]^{\mathrm{SLOCC}}$.

Regarding the dual variety of $X_{SO(8)} \subset \mathbb{P}(\mathfrak{so}_8)$, we recall that an equation of $X^*_{SO(2n)}$ is known in terms for the simple roots of the Lie algebras \mathfrak{so}_{2n} . In fact for all simple Lie algebras the equation defining the dual of the projectivization of the adjoint orbit is given by the vanishing of the discriminant $D_{\mathfrak{g}}$, i.e. the product of the (long) roots of \mathfrak{g} (see Ref⁴² p29).

$$D_{\mathfrak{g}} = \prod_{\alpha \in R_l} \alpha = 0 \tag{8}$$

Let $M \in \mathfrak{so}_8$ and M_s its semi-simple part with eigenvalues $\lambda_1, \ldots, \lambda_n$. The roots¹⁵ of \mathfrak{so}_{2n} are linear forms on \mathfrak{h} , the Cartan subalgebra of \mathfrak{so}_{2n} , of the form $\pm L_i \pm L_j$ with $i \neq j$ and such that $L_i(M_s) = \lambda_i$. Thus if α is a root we have $\alpha(M_s) = \pm \lambda_i \pm \lambda_j$ and the matrix $M \in \mathfrak{so}_{2n}$ belongs to $X^*_{SO_{2n}}$ if and only if

$$D_l(M) = \prod_{1 \le i < j \le n, i \ne j} (\lambda_i \pm \lambda_j)^2 = 0.$$
(9)

The roots of the characteristic polynomial of M, $t^{2n} + h_2(M)t^{2n-2} + \cdots + h_{2n-2}(M)t^2 + Pf^2(M)$ are $\pm \lambda_i$. Using the change of variable $x = t^2$, one gets the polynomial

$$x^{n} + h_{2}(M)x^{n-1} + \dots + h_{2n-2}(M)x + Pf^{2}(M)$$
(10)

whose roots are λ_i^2 . Taking the discriminant one obtains:

$$\Delta(x^{n} + h_{2}(M)x^{n-1} + \dots + h_{2n-2}(M)x + Pf^{2}(M)) = \prod_{1 \le i < j \le n} (\lambda_{i}^{2} - \lambda_{j}^{2})^{2} = \prod_{1 \le i < j \le n, i \ne j} (\lambda_{i} \pm \lambda_{j})^{2}$$
(11)

One concludes that $\Delta(x^n + h_2(M)x^{n-1} + \dots + h_{2n-2}(M)x + Pf^2(M)) = D_{\mathfrak{so}_{2n}}(M).$

In particular $\Delta(x^4 + h_2(M)x^3 + h_4(M)x^2 + h_6(M)x + Pf^2(M)) = 0$ is the defining equation of $X^*_{SO(8)}$. But it can also be checked that $\Delta(x^4 + 2Bx^3 + (B^2 + 2L + 4M)x^2 + (2BL + 4BM - 4D)x + L^2) = 0$ is an equation for the dual variety of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, i.e. $\Delta(x^4 + 2Bx^3 + (B^2 + 2L + 4M)x^2 + (2BL + 4BM - 4D)x + L^2)$ is the hyperdeterminant of format $2 \times 2 \times 2 \times 2 \times 2$. \Box

Remark II.1. The connection between the SLOCC orbit structure of four-qubits and the Lie algebra \mathfrak{so}_8 is also investigated in a different manner in the work of Péter Lévay^{31,32} and more recently in Ref³³ with a construction of the four-qubit invariants from the spin representation of SO(16). From a different perspective a connection between four-qubit states and the Dynkin diagram D_4 (the Dynkin diagram of \mathfrak{so}_8) was also pointed out in Ref²⁴ by constructing simple hypersurface singularities of type D_4 and their deformations from four-qubit states.

Remark II.2. The embedding of \mathcal{H} as a subspace of \mathfrak{so}_8 depends on the embedding of the four-qubit state $|\varphi\rangle$ in $\mathcal{M}_4(\mathbb{C})$ given by Eq (3). There are two more ways of looking at a four-qubit state as a linear map from \mathbb{C}^4 to \mathbb{C}^4 . Those other two representations will give different ways of writing the generators of $\mathbb{C}[\mathcal{H}]^{\text{SLOCC}}$, but also two different types of quartics like the one given by Eq (10) for n = 4. In the next section their role in the classification of the SLOCC orbits is investigated.

III. A FIRST CLASSIFICATION BASED ON INVARIANTS

A. SLOCC-invariant polynomials

In a general setting, a pure k-qudit state is an element of the Hilbert space $\mathcal{H} = V_1 \otimes \cdots \otimes V_k$ with $V_i = \mathbb{C}^{n_i}$ regarded as a multilinear form. Two qudit states are equivalent if they belong to the same orbit for the group SLOCC = $GL_{n_1} \times \cdots \times GL_{n_k}$. In principle, one can determine if two states are equivalent by comparing their evaluations on a sufficiently large system of special polynomials (in the coefficients of the forms and auxiliary variables), called concomitants, which are invariant under the action of SLOCC. In practice, this algorithm can be used only in very few cases, because the number and the size of the polynomials increase exponentially with the number of particles and the dimension of the Hilbert space. Nevertheless, even the knowledge of a little part of the polynomials gives rise to interesting information about the classification.

In the case of a four-qubit system, we deal with the quadrilinear form:

$$f := \sum_{0 \le i,j,k,\ell \le 1} a_{ijk\ell} x_i y_j z_k t_\ell \tag{12}$$

and the algebra of the polynomial invariants (polynomials in the coefficients of the forms with no auxiliary variables) is a free algebra on four generators:

1. One of degree 2:

$$B := a_{0000}a_{1111} - a_{1000}a_{0111} + a_{0100}a_{1011} + a_{1100}a_{0011} - a_{0010}a_{1101} + a_{1010}a_{0101} - a_{0110}a_{1001} + a_{1110}a_{0001},$$

$$(13)$$

2. two of degree 4:

$$L := \begin{vmatrix} a_{0000} & a_{0010} & a_{0001} & a_{0011} \\ a_{1000} & a_{1010} & a_{1001} & a_{1011} \\ a_{0100} & a_{0110} & a_{0101} & a_{0111} \\ a_{1100} & a_{1110} & a_{1101} & a_{1111} \end{vmatrix} \text{ and } M := \begin{vmatrix} a_{0000} & a_{0001} & a_{0100} & a_{0101} \\ a_{1000} & a_{1001} & a_{1000} & a_{1001} \\ a_{0010} & a_{0011} & a_{0110} & a_{0111} \\ a_{0010} & a_{0011} & a_{0110} & a_{0111} \\ a_{1010} & a_{1011} & a_{1110} & a_{1111} \end{vmatrix}$$
(14)

3. and one of degree 6: Set $b_{xy} := \det\left(\frac{\partial^2 f}{\partial z_i \partial t_j}\right)$. By interpreting this quadratic form as a bilinear form on the three dimensional space, one finds a 3×3 matrix B_{xy} satisfying

$$b_{xy} = [x_0^2, x_0 x_1, x_1^2] B_{xy} \begin{bmatrix} y_0^2 \\ y_0 y_1 \\ y_1^2 \end{bmatrix}.$$
 The generator of degree 6 is
$$D_{xy} := -\det(B_{xy}).$$
 (15)

Remark that this set is not unique, for instance one can replace L or M by

$$N := -L - M = \begin{vmatrix} a_{0000} & a_{1000} & a_{0001} & a_{1001} \\ a_{0100} & a_{1100} & a_{0101} & a_{1101} \\ a_{0010} & a_{1010} & a_{0011} & a_{1011} \\ a_{0110} & a_{1110} & a_{0111} & a_{1111} \end{vmatrix}.$$
(16)

We remark that, when evaluated on the Verstraete normal form⁴³

$$G_{abcd} := \frac{1}{2}(a+d)(|0000\rangle + |1111\rangle) + \frac{1}{2}(a-d)(|1100\rangle + |0011\rangle) + \frac{1}{2}(b+c)(|0101\rangle + |1010\rangle) + \frac{1}{2}(b-c)(|0110\rangle + |0011\rangle,$$
(17)

these invariants have nice closed expressions:

$$B(|G_{abcd}\rangle) = \frac{1}{2} \left(a^2 + b^2 + c^2 + d^2\right), \quad (18)$$

$$L(|G_{abcd}\rangle) = abcd, \quad (19)$$

$$M(|G_{abcd}\rangle) = \frac{1}{16}(a+b+c+d)(c+d-a-b)(a-b+c-d)(a-b+d-c), \quad (20)$$

$$N(|G_{abcd}\rangle) = \frac{1}{16}(a+b+c-d)(a-b-c-d)(a-b+c+d)(a+b-c+d), \quad (21)$$

$$D_{xy}(|G_{abcd}\rangle) = \frac{1}{32} \left(b^2 - a^2 + c^2 - d^2 \right) \left(-b^2 + a^2 + c^2 - d^2 \right) \left(b^2 + a^2 - c^2 - d^2 \right).$$
(22)

We consider the space $S := \{|G_{abcd}\rangle : (a, b, c, d) \in \mathbb{C}^4\}$ of the Verstraete normal forms. It is the Chevalley section for the SLOCC action on the Hilbert space with Weyl group D_4 : the closure of each generic orbit intersects S along a D_4 orbit. Indeed, since the polynomials $B(|G_{abcd}\rangle)$, $L(|G_{abcd}\rangle)$, $M(|G_{abcd}\rangle)$, and $D_{xy}(|G_{abcd}\rangle)$ are algebraically independent, the system

$$\begin{cases}
B(|G_{abcd}\rangle) = \alpha \\
L(|G_{abcd}\rangle) = \beta \\
M(|G_{abcd}\rangle) = \gamma \\
D_{xy}(|G_{abcd}\rangle) = \delta
\end{cases}$$
(23)

admits at most 192 solutions (this is the order of D_4). Furthermore, the system is clearly invariant under the permutations of the variables (a, b, c, d) and the transformation $(a, b) \rightarrow$ (-a, -b). For generic values of (a, b, c, d), these transformations generate a group isomorphic to D_4 . Hence, knowing one solution, the other ones are deduced from the action of D_4 . Geometrically, the solutions are symmetric with respect to the reflection group of the demitesseract. See Appendix A.

B. Three quadrics

We consider the three quartics

$$Q_{1}(|\varphi\rangle) := x^{4} - 2Bx^{3}y + (B^{2} + 2L + 4M)x^{2}y^{2} + (4D_{xy} - 4B(M + \frac{1}{2}L))xy^{3} + L^{2}y^{4}, (24)$$
$$Q_{2}(|\varphi\rangle) := x^{4} - 2Bx^{3}y + (B^{2} - 4L - 2M)x^{2}y^{2} + (-2MB + 4D_{xy})xy^{3} + M^{2}y^{4}, (25)$$
$$Q_{3}(|\varphi\rangle) := x^{4} - 2Bx^{3}y + (B^{2} + 2L - 2M)x^{2}y^{2} - (2LB + 2MB - 4D_{xy})xy^{3} + N^{2}y^{4}, (26)$$

Evaluated on $|G_{abcd}\rangle$, the roots of Q_1 are a^2 , b^2 , c^2 and d^2 and the roots of Q_2 (resp. Q_3) are the squares of the four polynomial factors of $M(|G_{abcd}\rangle)$ (resp. $N(|G_{abcd}\rangle)$ which are obtained by applying an invertible linear transformation to (a, b, c, d). Hence, the three quartics have the same invariants.

The invariant polynomials of a quartic $f := \alpha x^4 - 4\beta x^3 y + 6\gamma x^2 y^2 - 4\delta x y^3 + \omega y^4$ are algebraic combinations of $I_2 = \alpha \omega - 4\beta \delta + 3\gamma^2$, an invariant of degree 2, which is the apolar of the form with itself, and $I_3 = \alpha \gamma \omega - \alpha \delta^2 - \beta^2 \omega - \gamma^3 + 2\beta \gamma \omega$, an invariant of degree 3 called the catalecticant³⁹. In particular, the discriminant of the quadric is $\Delta = I_2^3 - 27I_3^2$.

From these definitions, one has

$$I_2(Q_1) = I_2(Q_2) = I_2(Q_3) = \frac{4}{3}L^2 + 2BD_{xy} - \frac{4}{3}B^2M - \frac{2}{3}B^2L + \frac{1}{12}B^4 + \frac{4}{3}LM + \frac{4}{3}M^2, \quad (27)$$

and

$$I_{3}(Q_{1}) = I_{3}(Q_{2}) = I_{3}(Q_{3}) = \frac{4}{2} D_{xy} B M + \frac{2}{3} D_{xy} B L - \frac{5}{9} B^{2} M L + \frac{4}{9} L^{2} M - D_{xy}^{2} - \frac{5}{9} B^{2} M^{2} + \frac{1}{18} B^{4} L + \frac{1}{9} B^{4} M - \frac{4}{9} L M^{2} - \frac{1}{6} B^{3} D_{xy} - \frac{2}{9} B^{2} L^{2} + \frac{8}{27} L^{3} - \frac{1}{216} B^{6} - \frac{8}{27} M^{3}.$$

$$(28)$$

Furthermore, we easily check that $\Delta(Q_i)$ is also the hyperdeterminant of $|\varphi\rangle$ (regarded as a quadrilinar form).

Covariants	Interpretation
$\Delta \neq 0$	Four distinct roots
$\Delta = 0$ and $T \neq 0$	Exactly one double root
$T = 0$ and $I_2 \neq 0$	Two distinct double roots
$I_2 = I_3 = 0$ and $Hess \neq 0$	A triple root
Hess = 0	A quadruple root

Table I. Roots of a quartic

In the aim to describe the roots of the quartics, we will use two other covariant polynomials: the Hessian

$$Hess(f) := \begin{vmatrix} \frac{\partial^2}{\partial x^2} f & \frac{\partial^2}{\partial x \partial y} f \\ \frac{\partial^2}{\partial y \partial x} f & \frac{\partial^2}{\partial y^2} f \end{vmatrix}$$
(29)

and the Jacobian of the Hessian

$$T(f) = \begin{vmatrix} \frac{\partial}{\partial x} f & \frac{\partial}{\partial y} f \\ \frac{\partial}{\partial x} Hess(f) & \frac{\partial}{\partial y} Hess(f) \end{vmatrix}.$$
 (30)

From the values of the covariants one can compute the multiplicity of the roots of a quartic f, according to Table I. Notice that the evaluations of *Hess* and *T* on the forms Q_1 , Q_2 and Q_3 are not equal in the general case.

C. A first classification

In this section, we use Table I to obtain a first classification and we refine it by considering the polynomials L, M and N which allow to decide if a quartic has a null root. The discussion is relegated to Appendix B.

We define the invariants

$$P := D_{xy} - BM, S_1 := B^2 + 4M, S_2 = B^2 - 4L, \text{ and } S_3 = B^2 - 4M.$$
(31)

If S is a set of polynomials, we will denote by \mathcal{V}_S the variety defined by the system $\{E = 0 : E \in S\}$. In a previous paper²⁶, we have investigated the case when L = M = 0. It remains to consider the other cases and, according to Appendix B, one has to refine the diagram of inclusions of Figure 1 which represents a first tentative of classification (see also Figure 2 for

the interpretation of red part in terms of roots of the quartics). The green part corresponds to subvarieties of $\mathcal{V}_{L,M}$ that are already investigated in a previous paper²⁶. Also, notice that the whole diagram of Figure 1 can be deduced from the red part, replacing L = 0 by M = 0(resp. N = 0). Note that $\mathcal{V}_{L,P,S_1,I_2,I_3} = \mathcal{V}_{L,P,S_1}$. Indeed, for any form in $\mathcal{V}_{L,P,S_1,I_2,I_3}$, Q_1 has a zero triple root and this implies automatically that Q_2 has a quadruple root which is equal to one of the parameter of the normal form. The case where one of the quartic has two double roots does not appear explicitly in the diagrams. But, a short calculation shows that it is equivalent to the case where one of the quartics has zero as a double root.



Figure 1. Inclusion diagram of the varieties \mathcal{V}_S .

D. The form problem again

Each of the varieties described in Figure 1 contains orbits whose intersections with the Chevalley section S are finite sets of points with symmetries related to some four dimensional polytopes.

- A generic orbit in V_∅ intersect S in 192 points splitting into 8 subsets of 24 points belonging to the same hyperplane. Each of these subsets is constituted with the permutations of the same vector (a, b, c, d) and centered on one of the vertices of a demitesseract (see Figure 3).
- In \mathcal{V}_L , each generic orbit contains also 192 normal forms which are the permutations of the same vector (0, a, b, c). The set of the normal forms splits into 32 subsets which



Figure 2. Interpretation of the red part of Figure 1 in terms of quartics.



Figure 3. Intersection of a generic orbit of \mathcal{V}_{\emptyset} with the subspace of normal forms.

are constituted of 6 points, belonging to the same space of dimension 2, centered on one of the points $(\alpha, \alpha, \alpha, 0)$, $(\alpha, \alpha, 0, \alpha)$, $(\alpha, 0, \alpha, \alpha)$ or $(0, \alpha, \alpha, \alpha)$ with $\alpha = \frac{\pm a \pm b \pm c}{3}$, that are middles of the edges of a tesseract (see Figure 4).

- A generic orbit in V_∆ is a special orbit in V_∅ with 96 normal forms splitting into 8 subsets of 12 points.
- A generic orbit in $\mathcal{V}_{\Delta,L}$ is a special orbit in \mathcal{V}_L with 96 normal forms splitting into 32 subsets of 3 points centered on the middle of the edges of a tesseract (see Figure 5).



Figure 4. Intersection of a generic orbit of \mathcal{V}_L with the subspace of normal forms and the tesseract.



Figure 5. Intersection of a generic orbit of $\mathcal{V}_{\Delta,L}$ with the subspace of normal forms.

- A generic orbit in \mathcal{V}_{I_2,I_3} is a special orbit in \mathcal{V}_{Δ} with 24 normal forms splitting into 8 subsets of 4 points.
- A generic orbit in V_{L,P} is a special orbit in V_{L,Δ} with 48 normal forms splitting into 24 pairs of points whose the middles are the vertices of a 24-cell polytope (see Figure 6). These vertices are also the centers of the faces of a tesseract. Notice that such an orbit can degenerate and have only 24 normal forms, which are exactly the vertices of the 24-cell. In this case, one has T(Q₁) = 0 (because Q₁ has two double roots). From Appendix B, it follows that L = M = 0 and, since this case has already been investigated in one of our previous papers, we will not examine it here.
- A generic orbit in \mathcal{V}_{L,P,S_1} is a special orbit in $\mathcal{V}_{L,P}$ with 16 normal forms, which are the vertices of a 16-cell polytope (see Figure 7).



Figure 6. Intersection of a generic orbit of $\mathcal{V}_{L,P}$ with the subspace of normal forms and a 24-cell polytope.



Figure 7. Intersection of a generic orbit of \mathcal{V}_{L,P,S_1} with the subspace of normal forms and a 16-cell polytope.

Hence, one can interpret the diagram of Figure 1 in terms of normal forms in Figure 8. Let us end the discussion by an illustration of the fact that the quartics Q_1 , Q_2 and Q_3 have an interchangeable role: we first remark that one can choose a degenerate orbit in \mathcal{V}_{\emptyset} by setting a = b = c = d. In this case, the normal forms are the vertices of the demitesseract and so of a 16-cell. It is similar (up to a rotation) to a generic orbit in $\mathcal{V}_{L,\Delta,P,Q}$. In terms of quartics, this is interpreted by the fact that Q_1 has a quadruple root and Q_2 has a zero triple root (compare to the variety $\mathcal{V}_{L,\Delta,P,Q}$, where Q_1 has a zero triple root and Q_2 has a quadruple root).



Figure 8. Diagram of normal forms.

IV. CLASSIFICATION BASED ON GEOMETRIC STRATIFICATION

We now investigate the SLOCC structure of the four-qubit Hilbert space from a geometrical point of view. Because a quantum state is well-defined up to a phase factor we will consider the projective Hilbert space $\mathbb{P}(\mathcal{H}) = \mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2)$ and we recall that $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, the Segre embedding of four projective lines, is the variety of separable states²⁵. We begin the section by defining the auxiliary varieties already used in our previous work^{25,26} to describe SLOCC invariant varieties built from the knowledge of X. Then we look at various stratifications of the ambient space induced by the singular locus of the four hypersurfaces corresponding to the zero locus of L, M, N and Δ . When combined, we recover the stratification induced by the three quartics Q_1, Q_2 and Q_3 .

A. Secant, tangent and dual varieties

The basic geometric tool introduced in our previous papers^{25,26} was the concept of auxiliary varieties. An auxiliary variety is an algebraic variety built by elementary geometric constructions from a given variety. An example of such is the so called secant variety^{30,45}. If $Y \subset \mathbb{P}(V)$ is a projective variety, the secant variety of Y is the algebraic closure of the union of secant lines:

$$\sigma(Y) = \overline{\bigcup_{x,y \in Y} \mathbb{P}^1_{xy}}.$$
(32)

If Y = X, the set of separable states, then $\sigma(X)$ is the algebraic closure of quantum states which are sums of two separable states. This algebraic variety is SLOCC invariant because so is X and it can be shown, for any multipartite system, that $\sigma(X) = \overline{\text{SLOCC} | GHZ \rangle}$ where $| GHZ \rangle$ is the usual generalization of the GHZ-state and X still denotes the variety of separable states.

In the same spirit, higher dimensional secant varieties can be defined by

$$\sigma_k(Y) = \overline{\bigcup_{x_1,\dots,x_k \in Y} \mathbb{P}^{k-1}_{x_1,\dots,x_k}}.$$
(33)

The secant varieties naturally provides a stratification of the ambient space and states belonging to different secant varieties can not be SLOCC equivalent. The interest of secant varieties in the context of quantum entanglement was first pointed out by Heydari²².

Another auxiliary variety of importance is the so-called tangential variety which corresponds to the union of tangent lines. If Y is a smooth variety and $\tilde{T}_y Y$ denotes the embedded tangent space of Y at y we have.

$$\tau(Y) = \bigcup_{y \in Y} \tilde{T}_y Y. \tag{34}$$

This variety has also a nice quantum information theory interpretation. For any multipartite system we have, if X still denotes the variety of separable states, $\tau(X) = \overline{\text{SLOCC} |W\rangle}$ where $|W\rangle$ is the generalization of the W-states. Those are well-known fact from algebraic geometry^{30,45} and have been restated in the language of quantum information theory in Ref²³.

In our geometric description of the stratification of the ambient space by algebraic varieties we will also use the concept of dual variety. Let us remind what the dual variety of a projective algebraic variety is, and why this concept has been already introduced in the study of entanglement of multipartite systems^{25,38}.

Consider $X \subset \mathbb{P}(V)$ a nondegenerate projective variety (i.e. not contained in a hyperplane). The dual variety of X is the closure of the set of tangent hyperplanes, i.e. is defined by

$$X^* = \overline{\{H \in \mathbb{P}(V^*), \exists x \in X_{\text{smooth}}, \tilde{T}_x X \subset H\}}$$
(35)

where $\tilde{T}_x X$ denote the embedded tangent space of X at x (see Ref²⁸).

If X is a G-invariant variety for a G-action on V and X^* is a hypersurface, then the defining equation of X^* is a G-invariant polynomial. In the case of $X = \mathbb{P}^{k_1} \times \cdots \times \mathbb{P}^{k_r}$ with $k_1 \leq k_2 + \cdots + k_r$ (assuming $k_1 \geq k_i$) the dual variety is always a hypersurface called the hyperdeterminant of format $(k_1 + 1) \times (k_2 + 1) \times \cdots \times (k_r + 1)$ and it is a $G = GL_{k_1+1} \times \cdots \times GL_{k_r+1}$ -invariant polynomial. Hyperdeterminants have been deeply studied by Gelfand Kapranov and Zelevinsky^{17,18}.

In the case where $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, the hyperdeterminant of format $2 \times 2 \times 2$ is the so-called Cayley hyperdeterminant and when $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, the hyperdeterminant is the invariant polynomial Δ introduced in Section III.

Other SLOCC invariant polynomials, or SLOCC invariant algebraic varieties of the Hilbert space of four qubits can be interpreted in terms of dual varieties as we now show.

B. The hypersurfaces $\{L = 0\}$, $\{M = 0\}$ and $\{N = 0\}$

The three hypersurfaces defined by the vanishing of one of the quartic invariant polynomials L, M or N are isomorphic and correspond to the dual varieties of three different types of embeddings of $\mathbb{P}^3 \times \mathbb{P}^3$ in $\mathbb{P}^{15} = \mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2)$.

Indeed, let $A, B \in \mathbb{C}^2 \otimes \mathbb{C}^2$ be two 2×2 matrices which are tensors of rank at most two, i.e. $A = t_1 \otimes u_1 + t_2 \otimes u_2$ and $B = v_1 \otimes w_1 + v_2 \otimes w_2$. Then we have the following three embeddings:

$$\mathbb{P}^{3} \times \mathbb{P}^{3} \hookrightarrow \mathbb{P}^{15}$$

$$\phi_{1} : ([A], [B]) \hookrightarrow [t_{1} \otimes u_{1} \otimes v_{1} \otimes w_{1} + t_{1} \otimes u_{1} \otimes v_{2} \otimes w_{2}$$

$$+ t_{2} \otimes u_{2} \otimes v_{1} \otimes w_{1} + t_{2} \otimes u_{2} \otimes v_{2} \otimes w_{2}]$$

$$\phi_{2} : ([A], [B]) \hookrightarrow [w_{1} \otimes u_{1} \otimes v_{1} \otimes t_{1} + w_{1} \otimes u_{1} \otimes v_{2} \otimes t_{2}$$

$$+ w_{2} \otimes u_{2} \otimes v_{1} \otimes t_{1} + w_{2} \otimes u_{2} \otimes v_{2} \otimes t_{2}]$$

$$\phi_{3} : ([A], [B]) \hookrightarrow [t_{1} \otimes v_{1} \otimes u_{1} \otimes w_{1} + t_{1} \otimes v_{1} \otimes u_{2} \otimes w_{2}$$

$$+ t_{2} \otimes v_{2} \otimes u_{1} \otimes w_{1} + t_{2} \otimes v_{2} \otimes u_{2} \otimes w_{2}]$$

Remark IV.1. There are no other ϕ_i to consider. Indeed, more permutations will not give any new varieties. For instance exchanging the vectors t and u in ϕ_1 to construct an other map $\tilde{\phi}_1$ will not give anything new because $\phi_1([A], [B]) = \tilde{\phi}_1([^tA], [B])$. We will denote by $Seg_i(\mathbb{P}^3 \times \mathbb{P}^3) \subset \mathbb{P}(\mathcal{H})$, for i = 1, 2, 3, the three Segre embeddings of $\mathbb{P}^3 \times \mathbb{P}^3$. From a QIT perspective, it should be pointed out that the Segre embedding of $\mathbb{P}^3 \times \mathbb{P}^3 \subset \mathbb{P}^{15}$ corresponds to the algebraic closure of the product of two $|EPR\rangle$ states, i.e. for ϕ_1 we have

$$\mathbb{P}^3 \times \mathbb{P}^3 = \overline{\text{SLOCC}(|00\rangle + |11\rangle) \otimes (|00\rangle + |11\rangle)} = \overline{\text{SLOCC}(|EPR\rangle \otimes |EPR\rangle)} \subset \mathbb{P}^{15}.$$
 (36)

In other words the Segre embedding of $\mathbb{P}^3 \times \mathbb{P}^3$ by ϕ_1 corresponds to the orbit closure of

$$|\varphi_1\rangle = |0000\rangle + |0011\rangle + |1100\rangle + |1111\rangle.$$
(37)

Similarly, the embeddings provided by ϕ_2 and ϕ_3 are the orbit closures of $|\varphi_2\rangle$ and $|\varphi_3\rangle$, with

$$|\varphi_2\rangle = |0000\rangle + |0101\rangle + |1010\rangle + |1111\rangle \tag{38}$$

and

$$|\varphi_3\rangle = |0000\rangle + |0110\rangle + |1001\rangle + |1111\rangle. \tag{39}$$

Let us denote by $|\varphi^{\sigma}\rangle$ the state obtained from $|\varphi\rangle$ by permuting the qubits by σ . Then, it is clear that $|\varphi_2\rangle = |\varphi^{1324}\rangle$ and $|\varphi_3\rangle = |\varphi^{1432}\rangle$. It is well known that the "usual" Segre product of two projective spaces $\mathbb{P}^m \times \mathbb{P}^n \subset \mathbb{P}(\mathbb{C}^{m+1} \otimes \mathbb{C}^{n+1})$ corresponds to the projectivization of the variety of rank one matrices in the projectivization of the space of $(m + 1) \times (n + 1)$ matrices. Taking the sum of two matrices of rank one, one gets a matrix of rank at most two and therefore the secant variety $\sigma(\mathbb{P}^m \times \mathbb{P}^n) \subset \mathbb{P}(\mathbb{C}^{m+1} \otimes \mathbb{C}^{n+1})$ can be interpreted as the projectivization of the locus of rank at most two matrices. When taking higher secants we get the well known stratification of bipartite systems by their rank^{22,25}

$$\mathbb{P}^m \times \mathbb{P}^n \subset \sigma_2(\mathbb{P}^m \times \mathbb{P}^n) \subset \cdots \subset \sigma_{\min(m,n)-1}(\mathbb{P}^m \times \mathbb{P}^n) \subset \mathbb{P}(\mathbb{C}^{m+1} \otimes \mathbb{C}^{n+1}).$$

In $\mathbb{P}^{15} = \mathbb{P}(\mathcal{H})$ the presence of the three varieties $Seg_i(\mathbb{P}^3 \times \mathbb{P}^3)$ will induce a first stratification of the ambient space. Recall the binary shortened notation $|\varphi\rangle = \sum_{i,j,k,l} a_{ijkl} |ijkl\rangle = \sum_{m=0}^{15} a_m |ijkl\rangle$ where $m = 1^i + 2^j + 4^k + 8^l$. As mentioned in Section II, we can represent a four-qubit state $|\varphi\rangle = (a_0, a_1 \dots, a_{15})$ by a 4 × 4 matrix

$$\mathcal{M}_{1} = \begin{pmatrix} a_{0} & a_{1} & a_{2} & a_{3} \\ a_{4} & a_{5} & a_{6} & a_{7} \\ a_{8} & a_{9} & a_{10} & a_{11} \\ a_{12} & a_{13} & a_{14} & a_{15} \end{pmatrix} \in \mathcal{H}$$

$$(40)$$

Similarly we have two alternative embeddings.

$$\mathcal{M}_{2} = \begin{pmatrix} a_{0} & a_{8} & a_{2} & a_{10} \\ a_{1} & a_{9} & a_{3} & a_{11} \\ a_{4} & a_{12} & a_{6} & a_{14} \\ a_{5} & a_{13} & a_{7} & a_{15} \end{pmatrix}, \\ \mathcal{M}_{3} = \begin{pmatrix} a_{0} & a_{1} & a_{8} & a_{9} \\ a_{2} & a_{3} & a_{10} & a_{11} \\ a_{4} & a_{5} & a_{12} & a_{13} \\ a_{6} & a_{7} & a_{14} & a_{15} \end{pmatrix}$$
(41)

If we think in terms of matrix rank, it is clear that the variety $Seg_i(\mathbb{P}^3 \times \mathbb{P}^3)$ is defined by the zero locus of the two by two minors of the matrix \mathcal{M}_i , the secant variety $\sigma(Seg_i(\mathbb{P}^3 \times \mathbb{P}^3))$ is defined by the zero locus of the 3×3 minors of \mathcal{M}_i and the third secant variety $\sigma_3(Seg_i(\mathbb{P}^3 \times \mathbb{P}^3))$ is defined by the vanishing of det (\mathcal{M}_i) (Figure 9).



Figure 9. Inclusion diagram of X within the stratification defined by ϕ_1

Table III shows how the normal forms, following Verstraete *et al.*'s notation, fit in this stratification.

One can obtain similar tables for the stratification by rank of \mathcal{M}_2 and \mathcal{M}_3 . The corresponding forms are obtained by permuting the qubits by the permutations 1324 and 1432.

We can do a little bit better by using simultaneously the strafication by rank of the three embeddings of $\mathbb{P}^3 \times \mathbb{P}^3$. It leads to a stratification by multirank (Figure 10, Table III).

Let us point out that the variety $\cap_{i \in \{1,2,3\}} \sigma_3(Seg_i(\mathbb{P}^3 \times \mathbb{P}^3))$ is nothing but $\sigma_3(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)$ the third secant variety of the set of separable states X which was described in our previous article²⁶.

The stratification of entanglement classes of four-qubit states by multirank has been

Varieties	Forms		
$\mathbb{P}^{15} \smallsetminus \sigma_3(\mathbb{P}^3 \times \mathbb{P}^3)$	$G_{abcd}, L_{abc_2}, L_{ab_2}$		
	L_{ab_3}, L_{a_4}		
$\sigma_3(\mathbb{P}^3 \times \mathbb{P}^3)$	$L_{a_20_{3\oplus\overline{1}}},L_{0_{5\oplus\overline{3}}},L_{0_{7\oplus\overline{1}}}$	3	
	$G_{0bcd}, G_{a0cd}, G_{ab0d}, G_{abc0}$		
	$L_{0bc_2}, L_{a0c_2}, L_{ab0_2}$		
	L_{0b_2}, L_{a0_2}		
	L_{0b_3}, L_{a0_3}		
	L_{0_4}		
$\sigma_2(\mathbb{P}^3 \times \mathbb{P}^3)$	$L_{0_{3\oplus\overline{1}}0_{3\oplus\overline{1}}}$	2	
	$G_{00cd}, G_{0b0d}, G_{0bc0}, G_{a00d}, G_{a0c0}, G_{ab00}$		
	$L_{00c_2}, L_{a00_2}, L_{0b0_2}$		
	$L_{00_{2}}$		
	$L_{00_{3}}$		
$\mathbb{P}^3 imes \mathbb{P}^3$	$G_{a000}, G_{0b00}, G_{00c0}, G_{000d}$	1	
	L_{000_2}		

Table II. Stratification of the ambient space by rank of S_1

already studied in Ref⁵ but without the geometric interpretation in terms of secants of Segre varieties. Table III is identical to Table 4 of Ref⁵.

Finally let us point out that the stratification by rank of the hypersurface $\{L = 0\}$ (respectively $\{M = 0\}$ and $\{N = 0\}$) corresponds to a stratification by singular locus of the hypersurface. In the next section we will see that the study of the singular locus of the hypersurface $\Delta = 0$ is more challenging.

C. Singularities of the dual variety X^*

In this section we consider the dual variety of X denoted by X^* and given by the zero locus of the hyperdeterminant Δ of format $2 \times 2 \times 2 \times 2$. This invariant polynomial of four qubits is of fundamental importance in the study of four-qubit states. Its importance was first emphasized by Miyake³⁸ and recently Gour and Wallach¹⁶ used $|\Delta|$ as an entanglement



Figure 10. Inclusion diagram of X within the stratification defined by multirank

measure of genuine four-qubit entanglement. It is well-known that the nonvanishing of Δ characterizes semi-simple elements for the SLOCC-action on \mathcal{H} . The SLOCC stratification of $X^* = \{|\varphi\rangle \in \mathbb{P}(\mathcal{H}) : \Delta(|\varphi\rangle) = 0\}$ by invariant subvarieties has been regarded in Ref³⁸ in the context of QIT based on the earlier work of Weyman and Zelevinsky⁴⁴. The purpose of the section is to complete the picture by a finer grained stratification of X^* based on the study of the hyperplane sections of X and the understanding of this stratification in terms of normal forms and vanishing of invariants.

A hyperplane H belongs to X^* if the corresponding H is tangent to X at some smooth point $x \in X$, i.e. $\tilde{T}_x X \subset H$. In other words the hyperplane section $X \cap H$ defines a singular hypersurface of X with a singular point at x. The singularity, i.e. the hyperplane section with a singular point x, will be denoted by $(X \cap H, x)$. When X^* is a hypersurface and Ha smooth point of X^* , it is well-known that the singular hypersurface $X \cap H$ has a unique singular point and the Hessian matrix at this unique singular point is nondegenerate¹⁸. For a hyperplane section $X \cap H$ we will denote by $f_{X \cap H}$ the polynomial defining $X \cap H$ as a hypersurface of X. Therefore H is a smooth point of X^* reads as there exists a unique $x \in X$ such that $f_{X \cap H}(x) = 0$, $\partial_i f_{X \cap H}(x) = 0$ and $Hess(f_{X \cap H}, x)$ is of full rank. Such singular point is called a Morse singularity or a A_1 singularity in Arnold's classification of simple singular germs¹. Thus if H is a smooth point of X^* , there exists a unique $x \in X$ such that

Varieties	Forms	mutli-rank
$\mathbb{P}^{15}\smallsetminus\cup_{i\in\{1,2,3\}}\sigma_3(Seg_i(\mathbb{P}^3\times\mathbb{P}^3))$	G_{abcd}	[4, 4, 4]
	L_{abc_2}	
	L_{ab_3}	
$\sigma_3(Seg_1(\mathbb{P}^3 imes \mathbb{P}^3))$	$L^{4231}_{a_2b_2}, L^{4231}_{a_4}$	[3, 4, 4]
$\sigma_3(Seg_2(\mathbb{P}^3\times\mathbb{P}^3))$	$L_{a_2b_2}, L_{a_4}$	[4, 3, 4]
$\sigma_3(Seg_2(\mathbb{P}^3 \times \mathbb{P}^3))$	$L^{2134}_{a_2b_2}, L^{2134}_{a_4}$	[4, 4, 3]
$\cap_{i \in \{1,2,3\}} \sigma_3(Seg_i(\mathbb{P}^3 \times \mathbb{P}^3))$	$L_{a_20_{3\oplus 1}}, L_{0_{7\oplus \overline{1}}}$	[3,3,3]
$=\sigma_3(X)$		
$\cap_{i \in \{1,2,3\}} \sigma_2(Seg_i(\mathbb{P}^3 \times \mathbb{P}^3))$	$\overline{L_{0_{3\oplus\overline{1}}0_{3\oplus\overline{1}}}}$	[2, 2, 2]

Table III. Stratification of the ambient space by multi-rank (only general values of the parameters are taken into account).

 $(X \cap H, x) \sim A_1$. When X^* is a hypersurface, a singular point of X^* is a hyperplane whose corresponding hyperplane section $X \cap H$ does not have a unique A_1 singularity. Therefore there are two possibilities to not satisfy this condition:

- Either $X \cap H$ has more than one singular point,
- or $X \cap H$ has a unique singular point and $Hess(f_{X \cap H}, x)$ is not of maximal rank.

This leads to the notion of node and cusp components of the singular locus of X^* as defined in Ref⁴⁴.

Definition IV.1. Let $X \subset \mathbb{P}(V)$ a nondegenerate projective variety and X^* its dual variety which is assumed to be a hypersurface. The singular locus $Sing(X^*)$ of X^* is given by

$$Sing(X^*) = X^*_{node} \cup X^*_{cusp} \tag{42}$$

where the node component, X_{node}^* is defined by

$$X_{node}^* = \overline{\{H \in X^* : \exists (x, y) \in X \times X, x \neq y, \tilde{T}_x X \subset H, \tilde{T}_y X \subset H\}}$$
(43)

and the cusp component, X_{cusp}^* is

$$X_{cusp}^* = \overline{\{H \in X^* : \exists x \in X, \tilde{T}_x X \subset H, (X \cap H, x) \not\sim A_1\}}.$$
(44)

As pointed out in Ref^{44} , in the case where X is a Segre product, the node component may be further decomposed. We need to introduce the notion of J-node component. We give the definition in the case of a product of projective spaces but it can easily be extended to a Segre product of algebraic varieties.

Definition IV.2. Let $X = \mathbb{P}^{k_1} \times \cdots \times \mathbb{P}^{k_r} \subset \mathbb{P}^{(k_1+1)\dots(k_r+1)-1}$ be the Segre product of rprojective spaces. Let $J = \{j_1, \dots, j_s\} \subset \{1, \dots, r\}$. We say that $(x, y) \in X \times X$ is a J-pair of points when $x = x_1 \otimes x_2 \otimes \cdots \otimes x_{j_1} \otimes \cdots \otimes x_{j_r} \otimes \cdots \otimes x_r$ and $y = y_1 \otimes y_2 \otimes \cdots \otimes x_{j_1} \otimes \cdots \otimes x_{j_r} \otimes \cdots \otimes x_{j_r}$

$$X_{node}^*(J) = \overline{\{H \in X^* : \exists (x, y) \ a \ J\text{-pair of points of } X \times X, \tilde{T}_x X \subset H, \tilde{T}_y X \subset H\}}.$$
 (45)

In Ref⁴⁴ it is proven that for $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, the irreducible components of $\operatorname{Sing}(X^*)$ are

$$\operatorname{Sing}((\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)^* = X^*_{cusp} \cup X^*_{node}(\emptyset) \cup \bigcup_{1 \le i \le j \le 4} X^*_{node}(\{i, j\})$$
(46)

Cusp and node components have interpretation in terms of duals of auxiliary varieties of X. In the case of four-qubit systems one has the following proposition.

Proposition IV.1. Let $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, then

- 1. $X_{node}^* = \sigma(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)^*$ 2. $X_{node}^*(\{i, j\}) = \sigma_{\{i, j\}}(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)^*$
- 3. $X^*_{cusp} = \tau(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)^*$

where $\sigma_{\{J\}}(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)$ is the secant variety of *J*-pairs of points, i.e. $\sigma_{\{J\}}(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1) = \bigcup_{(x,y)J\text{-pair}} \mathbb{P}_{xy}$.

Proof. The points 1 and 2 are already particular cases of Proposition 4.1 of our previous paper²⁵ where we establish that the node component is always the dual variety of some J-secant variety of the original Segre product. The J-secant variety is the variety of secant lines where the lines are defined by J-pairs of points. The proof of this general statement follows from the application of the Terracini's lemma (see Ref²⁵).

Point 3 is more subtle. First it should be noticed that $\tau(X)^* \subset X^*_{cusp}$. This can be understood from the fact that a hyperplane H tangent to $\tau(X)$ is also tangent to X. Moreover

the fact that H is tangent to $\tau(X)$ at v implies that H is tangent to X along the direction v. Therefore the matrix $Hess(f_{X\cap H}, x)$ is degenerate in the direction v. This implies that $\tau(X)^* \subset X^*_{cusp}$. The equality will follow if we prove equality of dimension. Thus one has to calculate the dimension of $\tau(\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1)^*$. The dimension of a dual variety can be calculated by Katz's formula (see Ref¹⁸) which states that for a projective variety $Y \subset \mathbb{P}(V) = \mathbb{P}^N$, the dimension of Y^* is obtained from

$$\dim(Y^*) = N - \min_{H \in X^*} (\operatorname{corank}(Hess(f_{Y \cap H}, x)) - 1$$
(47)

where x is the point where $f_{Y\cap H}$ is singular. In particular this formula says that in general, the dual variety is a hypersurface, because we expect $\min_{H\in X^*}(\operatorname{corank}(\operatorname{Hess}(f_{Y\cap H}, x)))$ to be zero (when it is not, it means that all tangent hyperplanes are tangent to Y not to a point but to a subspace of positive dimension). This formula can be used to compute the dimension of $\tau(X)^*$. For this purpose, one needs to compute the general form of $\operatorname{Hess}(f_{\tau(X)\cap H}, x)$. Assuming $\dim \sigma(X) = 2n + 1$ (and thus $\dim(\tau(X)) = 2n$ see Ref^{45}), it can be shown using moving frames techniques that

$$Hess(f_{\tau(X)\cap H}, v) = \begin{pmatrix} A & Hess(f_{X\cap H}, x) \\ Hess(f_{X\cap H}, x) & 0 \end{pmatrix}$$
(48)

where v is a general point of $\tau(X)$ (i.e. a general element of $\tilde{T}_x X$) and A is a $n \times n$ full rank block built from the cubic invariants of the Taylor expansion of $f_{X\cap H}$ at x. Because H belongs to $\tau(X)^*$ it is tangent to X along the direction v and necessarily rank $(Hess(f_{X\cap H}, x) \leq n - 1 \text{ confirming the fact that } \tau(X)^*$ is at most of codimension 2 in the ambient space. To prove that $\tau(X)^*$ is, in our case, of codimension 2, one needs to show that we can find a matrix $Hess(f_{X\cap H}, x)$ which is of rank n - 1 for a generic tangent vector v. In our situation $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ and let us assume $x = |0000\rangle$. Let $v = |1000\rangle + |0100\rangle + |0010\rangle + |0001\rangle$ be a generic tangent vector to X at x, and consider a general curve $t \to \gamma(t)$, such that $\gamma(0) = x$ and $\gamma'(0) = v$. Then we have

$$\gamma(t) = |0000\rangle + t(|1000\rangle + |0100\rangle + |0010\rangle + |0001\rangle) + \frac{t^2}{2!}(|1100\rangle + |1010\rangle + |1001\rangle + |0110\rangle + |0101\rangle + |0011\rangle) + O(t^3).$$
(49)

For the hyperplane $a\langle 1100| + b\langle 1010| + c\langle 1001| + d\langle 0110| + e\langle 0101| + f\langle 0011|$, we obtain

$$Hess(f_{X \cap H}, x) = \begin{pmatrix} 0 & a & b & c \\ a & 0 & d & e \\ b & d & 0 & f \\ c & e & f & 0 \end{pmatrix}.$$
 (50)

One can check that for a = f, b = e, d = c = -e - f, then v is a singular direction of $Hess(f_{X\cap H}, x)$) and rank $(Hess(f_{X\cap H}, x)) = 3$. Therefore $\dim(\tau(X)^*) = 13$ and the equality follows by irreducibility of X^*_{cusp} . \Box

Proposition IV.1 gives a description of the singular locus of the hyperdeterminants in terms of the dual varieties of (the orbit closure) of GHZ and W states for the four-qubit systems. Combining the description of $\text{Sing}(X^*)$ established in Ref⁴⁴ and our interpretation of the singular components in terms of tangential and secant varieties we get,

Theorem 1. Let $\mathcal{H} = (\mathbb{C}^2)^{\otimes n}$ the Hilbert space of a n-qubit system with $n \geq 3$. Let $X = \mathbb{P}^1 \times \cdots \times \mathbb{P}^1$ be the set of separable states and X^* its dual variety given by the vanishing of the $2 \times \cdots \times 2$ hyperdeterminant. Then we have

- 1. $Sing(X^*) = \sigma_{\{1\}}(X)^* \cup \sigma_{\{2\}}(X)^* \cup \sigma_{\{2\}}(X)^*$ for n = 3,
- 2. $Sing(X^*) = \tau(X)^* \cup \sigma(X)^* \cup_{1 \le i < j \le 4} \sigma_{\{i,j\}}(X)^*$ for n = 4,
- 3. $Sing(X^*) = \tau(X)^* \cup \sigma(X)^* \text{ for } n \ge 5.$

Proof. Point 1 is already proved in our paper Ref²⁵ where the 3-qubits case is studied in details. Point 2 is Proposition IV.1. Now to prove point 3 we use the result of Weyman and Zelevinsky which states that in this case $\operatorname{Sing} X^* = X^*_{cusp} \cup X^*_{node}(\emptyset)$. Weyman and Zelevinsky also proved that both components are irreducible and of codimension 1 in X^* . The identification $\sigma(X)^* = X^*_{node}(\emptyset)$ follows from Terracini's lemma. The inclusion $\tau(X)^* \subset X^*_{cusp}$ is clear from the description of $\tau(X)$ but to prove the equality one needs to calculate the dimension of $\tau(X)^*$. The same argument as Proposition IV.1 shows that it is equivalent to find a $n \times n$ symmetric matrix Q of rank n - 1, with 0 on the diagonal, such that ker $Q = \langle (1, \ldots, 1) \rangle$. Such a matrix can be constructed as follow:

• If n = 2k we consider the matrix Q defined by $q_{ii} = 0, q_{i,n+1-i} = -2(k-1)$ and $q_{i,j} = 1$ if $(i, j) \notin \{(i, i), (i, n+1-i)\}.$ • If n = 2k + 1 we consider the matrix Q defined by $q_{ii} = 0, q_{i,n+1-i} = -2k + 1$ for $1 \le i \le k$ and $k + 2 \le i \le n, q_{k+1,k} = q_{k+1,k+2} = -k$ and $q_{i,j} = 1$ for $(i,j) \notin \{(i,i), (i,n+1-i), (k+1,k), (k+1,k+2)\}.$

In both cases the matrix Q is symmetric with rank n-1 and ker $Q_n = \langle (1, \ldots, 1) \rangle$. Therefore one can construct a hyperplane section of $\tau(X)$ such that the corresponding Hessian is of rank 2n-1, which implies that $\tau(X)^*$ is of codimension 1 in X^* . \Box .

Remark IV.2. It is interesting to point out that the two main stratas of $Sing(X^*)$ are dual varieties of the orbit closure of GHZ and W-states.

The cusp and node components can be further decomposed by their multiplicity. Let us consider

$$X_n^* = \overline{\{H \in X^* : \text{mult}_H X^* = n\}}, \text{ i.e } H \text{ is a root of multiplicity } n \text{ of } \Delta$$
(51)

Then we have a filtration of X^* by multiplicities: $X^* \supset X_1^* \supset X_2^* \supset \cdots \supset \ldots$

We can define

$$X^*_{node,k}(J) = X^*_{node}(J) \cap X^*_k \tag{52}$$

and similarly

$$X^*_{cusp,k} = X^*_{cusp} \cap X^*_k.$$
(53)

A result of Dimca¹², generalized by Parusinski⁴⁰, shows that the multiplicity of a hyperplane $H \in X^*$ is equal to the Milnor number of the hyperplane section $X \cap H$. The Milnor number of an isolated singularity $(f_{X \cap H}, x)$ is a topological invariant defined by

$$\mu = \dim_{\mathbb{C}} \mathcal{O}_k / (\nabla f) \tag{54}$$

where \mathcal{O}_k is ring of all germs $g: (\mathbb{C}^k, x) \to (\mathbb{C}, 0)$ and $(\nabla f) = (\frac{\partial f}{\partial x_1}(x), \dots, \frac{\partial f}{\partial x_k}(x))$ is the gradient ideal. If $X \cap H$ has only isolated singularities, the Milnor number of the hyperplane section $X \cap H$, denoted by $\mu(X \cap H)$, will be the sum of the Milnor number of each singularity. Under this assuption, the result of Dimca says

$$mult_H(X^*) = \mu(X \cap H). \tag{55}$$

Which, geometrically, leads to the following observation of F. Zak⁴⁶

$$\sigma_n(X)^* \subset X_n^*. \tag{56}$$

In Ref²⁴ we have calculated the isolated singular types of the hyperplane sections of the set of separable states of four qubits, $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$, for all possible linear forms (hyperplane) obtained by Verstraete *et al.*. The construction we employed was the following, let $|\varphi\rangle$ be a state given by Verstraete *et al.* classification and let us consider the linear form $\langle \varphi |$. Then the hyperplane section $H_{\langle \varphi |} \cap X$ defines a hypersurface of $X = \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ which may be smooth or have singularities. When the singularities are isolated, we have tools coming from the classification of simple singularities¹ to discriminate the corresponding hyperplane sections. The type of the hypersurface $X \cap H \subset X$ is SLOCC invariant (because X is a homogeneous SLOCC orbit) and so is the singularity type attached to $X \cap H$.

D. Geometric stratification of the Hilbert space of four-qubit states

The singular type of a hyperplane section $X \cap H$ can be used to discuss the different stratas of the hyperdeterminant Δ as we now explain. For instance it is well known that, for generic a, b, c, d, the states G_{abcd} are such that $\Delta(G_{abcd}) \neq 0$, which means that $H_{G_{abcd}} \cap X$ is a smooth hypersurface. By calculating²⁴ for each Verstraete *et al.* normal forms, the different types of the isolated singularities, we identify which forms belong to which components of the singular locus of Δ . If the form gives a hypersurface with only one isolated singularity of type A_1 then the tested form is a smooth point of the hyperdeterminant. If the form gives several A_1 singularities it is a point of the node locus. If the singularity is not of type A_1 it is a point of the cusp locus. Moreover the Milnor number of the singularity gives information on the multiplicity of the component.

Once the normal forms are interpreted as components of some specific singular locus then we can test the normal forms on the invariant obtainted in Figure 1 to identify geometrically the varieties obtained from the analysis of the three quartics. For instance if we consider the Verstraete form L_{ab_2} , the corresponding hyperplane section $H_{\langle L_{ab_2}|}$ has a unique A_2 singularity. It is the only form to have this property (for a generic choice of parameters) and thus the states $|L_{ab_2}\rangle$ form an open subset of $\tau(X)^*$. We easly check that $|L_{ab_2}\rangle$ corresponds to the vanishing of I_2 and I_3 , i.e. following the notations of Figure 1 we have $\mathcal{V}_{I_2,I_3} = \tau(X)^*$.

The singular type of the hyperplane section $H_{\langle \varphi |} \cap X$ being invariant under permutation of the qubits, the forms are given up to a permutation, i.e. $|L_{ab_2}^{\sigma}\rangle \in \tau(X)^*$. Analysing similarly all forms of the 9 families we obtain: **Theorem 2.** The Hilbert space of four-qubit states can be stratified under SLOCC according to the hyperdeterminant as shown in Figure 11 with the varieties of the stratification being



Figure 11. Stratification of the dual variety of the separable states

described in terms of forms and invariants by Tables IV, V, VI.

The stratification can be completed by the stratification of the varieties $Seg_i(\mathbb{P}^3 \times \mathbb{P}^3)^*$ (Figure 12).



Figure 12. Stratification of the ambient space by SLOCC varieties

Remark IV.3. The study of the singularities of Δ enables us to caracterize the Verstraete normal forms as general points of specific stratas. It was well known that $|G_{abcd}\rangle$, for a

Varieties	Forms	Invariants	Singularities		
$\mathbb{P}^{15}\smallsetminus X^*$	G_{abcd}	$\Delta \neq 0$	smooth hyperplane section		
X*	L_{abc_2}	$\Delta = 0$	hyperplane sections		
			with a unique A_1		

Table IV. Stratification by $\Delta = 0$ (smooth points)

Varieties	Forms	Invariants	Singularities	
$X^*_{cusp} = \tau(X)^*$	L_{ab_3}	$I_2 = I_3 = 0$	hyperplane sections	
			with a unique A_2	
$X^*_{cusp,3}$	L_{a_4}	$L = P = S_1 = 0$	hyperplane sections	
		or $M = D_{xy} = S_2 = 0$ or $N = D_{xy} = S_3$	with a unique A_3	
$X^*_{cusp,4}$	$L_{0_{7\oplus\overline{1}}}$	B = L = M = D = 0	hyperplane sections	
			with a unique D_4	

Table V. Stratification by $\Delta = 0$ (cusp components)

Varieties	Forms	Invariants	Singularities
$X^*_{node}(\emptyset) = \sigma(X)^*$	G_{abcc}	see Remark IV.5	hyperplane sections
			with two A_1 singularities
$\cup_{1 \leq i < j \leq 4} \sigma_{\{i,j\}}(X)^*$	$L_{a_2b_2}$	$M = D_{xy} = 0$	hyperplane sections
		or $N = D_{xy} = 0$	with two A_1
		or $L = P = 0$	
$X^*_{node,3}$	$L_{a_20_{3\oplus\overline{1}}}, L_{aac_2}$	$L = M = D_{xy} = 0$	hyperplane sections
			with three A_1

Table VI. Stratification by $\Delta = 0$ (node components)

generic choice of parameters, will be a state on which Δ does not vanish. Now, one can see that the other forms, for a generic choice of parameters and up to permutation, are general points of specific stratas of the singular locus. The two nilpotent states $L_{0_{5\oplus\overline{3}}}$ and $L_{0_{3\oplus\overline{1}}0_{3\oplus\overline{1}}}$ which do not appear in Theorem 2 belong to the nullcone \mathcal{N} .

Remark IV.4. The varieties, as denoted in the Theorem, are not all irreducible. For instance $X^*_{cusp,3}$ is not irreducible and neither is $X^*_{cusp,4}$ which corresponds to the nullcone

 $\mathcal{N}.$

Remark IV.5. All varieties of Figure 12 correspond to varieties detected in Figure 1 by the invariants of the three quartics, except $\sigma(X)^*$ which does not appear in Figure 1 but which can be detected by computing the hyperplane sections of $|G_{abcc}\rangle$. The defining equations of $\sigma(X)^*$ have been computed by Lin and Strumfels³⁵. This component corresponds to the projective closure of the image of the principal minor map for 4×4 matrices.

Remark IV.6. It is interesting to point out that most of the SLOCC varieties exhibited in Figure 12 have a quantum information theory interpretation in terms of the duals of wellknown quantum states orbit closures. In Figure 13, which is a translation of part of Figure 12, we denote by $\overline{|\varphi\rangle}$ the SLOCC orbit closure of the state $|\varphi\rangle$. We also denote by Sep any separable state and thus $\overline{|Sep\rangle} = X$. We also denote by $|GHZ_3\rangle$ the family of states which can be written as a sum of three separable states (tensor of rank 3).



Figure 13. Stratification of the ambient space by duals of state orbit closures

V. VERSTRAETE TYPE OF A FORM

Let A be a set of variables and $|\varphi(A)\rangle$ be a family of parametrized forms where each variable of A ranges over C. A specialization of A is a system S of algebraic equations in the variables of A. We denote by $|\varphi(A)\rangle|_S$ the subfamilies of $|\varphi(A)\rangle$ such that the values of A satisfy S. For instance, the four-parameter family G_{abcd} together with $S = \{a^2 = d^2, b^2 = c^2\}$ is union of the two dimensional spaces generated by one of the four basis states $\{|0000\rangle + |1111\rangle, |0101\rangle + |1010\rangle\}, \{|0000\rangle + |1111\rangle, |0110\rangle + |1001\rangle\}, \{|1100\rangle + |0011\rangle, |0101\rangle + |1010\rangle\},$ or $\{|1100\rangle + |0011\rangle, |0110\rangle + |1001\rangle\}.$

A form φ has a Verstraete type [F, S], where F is one of the nine Verstraete generic forms and S a specialization of the parameters, if there exists a permutation $\sigma \in \mathfrak{S}_4$ of the qubits such that φ is SLOCC-equivalent to an element of $(F|_S)^{\sigma}$.

In this section, we describe an algorithm allowing to compute the Verstraete type for any given form. First note that if the form is nilpotent it is easy to find a Verstraete equivalent form from our previous paper²⁶ and Table VIII. Now suppose that the form $|\varphi\rangle$ is not nilpotent. We use the 170 covariants computed in our previous paper²⁶ in order to discriminate between the Verstraete forms. In particular, we define:

$$\mathcal{L} = L_{6000} + L_{0600} + L_{0060} + L_{0006}$$

$$\mathcal{K}_{5} = K_{5111} + K_{1511} + K_{1151} + K_{1115}, \ \mathcal{K}_{3} = K_{3311} + K_{3131} + K_{3113} + K_{1331} + K_{1313} + K_{1133},$$
$$\overline{\mathcal{G}} = G_{3111}^{1} G_{1311}^{1} G_{1131}^{1} G_{1113}^{1}, \ \mathcal{G} = G_{3111}^{2} + G_{1311}^{2} + G_{1131}^{2} + G_{1113}^{2},$$
$$\mathcal{D} = D_{4000} + D_{0400} + D_{0040} + D_{0004},$$
$$\mathcal{H} = H_{2220} + H_{2202} + H_{2022} + H_{0222}$$

and $C = C_{1111}^2$. We proceed as follows: first we classify the forms with respect to the roots of the three quartics Q_1 , Q_2 and Q_3 according to the discussion in Appendix B. For each of the cases considered in Appendix B, we determine which Verstraete forms can occur and, when there are several possibilities we use one of the covariants previously defined to discriminate between them. Let V be a vector, we denote by ev(V) the vector such that ev(V)[i] = 0 if V[i] = 0 and ev(V)[i] = 1 if $V[i] \neq 0$.

- 1. If the quartics have only nonzero roots
 - (a) If all the roots are simple then this is the generic case and the Verstaete type is $[G_{abcd}; \emptyset].$
 - (b) If each quartic has double root and two simple roots (equivalently $T_1 = T_2 = T_3 = 0$ and $I_2, I_3 \neq 0$ then two cases can occur. Either the Verstraete type is $[G_{abcd}; c = d]$ or it is $[L_{abcc}; \emptyset]$. We determine the forms remarking that $\mathcal{L}(L_{abc_2}) \neq 0$ and $\mathcal{L}(G_{abcc}) = 0$.
 - (c) If each quartic has a single simple root and a triple root (equivalently $I_2 = I_3 = 0$) then three cases can occur: $[G_{abcd}; b = c = d]$, $[L_{abc_2}; b = c]$ and $[L_{ab_3}; \emptyset]$. In order

to determine the type, we evaluate the vector $V = [\mathcal{K}_5, \mathcal{L}]$ on each forms. We can decide the type of the form according to the values

$$ev(V(G_{abbb})) = [0,0], ev(V(L_{abb_2})) = [1,0], \text{ and } ev(V(L_{ab_3})) = [1,1]$$

- 2. If only one of the quartics Q_i has a zero root then
 - (a) If Q_i has only simple roots then the only possibility is $[G_{abcd}; d = 0]$
 - (b) If Q_i has a double zero root and two simple roots then we have three possibilities $[G_{abcd}; c = d = 0], [L_{abc2}; c = 0]$ or $[L_{a_2b_2}; \emptyset]$. We evaluate the form on the vector $V = [\mathcal{K}_3, \mathcal{L}]$ and compare with

$$ev(V(G_{ab00})) = [0,0], ev(V(L_{ab02})) = [1,0], \text{ and } ev(V(L_{a2b2})) = [1,1]$$

(c) If Q_i has triple zero root and a simple root then we have the five possibilities $[G_{abcd}; c = b = d = 0], [L_{abc2}; b = c = 0], [L_{ab3}; a = 0], [L_{a_2b_2}; a = b]$ and $[L_{a_4}; \emptyset]$. We evaluate the form on the vector $V = [\mathcal{C}, \mathcal{D}, \mathcal{K}_5, \mathcal{L}]$ and compare to the identities

$$ev(V(G_{a000})) = [0, 0, 0, 0], ev(V(L_{a002})) = [1, 0, 0, 0],$$

 $ev(V(L_{0b_3})) = [1, 1, 1, 0], ev(V(L_{a_2a_2})) = [1, 1, 0, 0]$

and $ev(V(L_{a_4})) = [1, 1, 1, 1].$

- (d) If Q_i has a double nonzero root and two simple roots then there are two possibilitites $[G_{abcd}; b = c, d = 0]$ and $[L_{abc_2}; b = 0]$ which can be identified by remarking that $\mathcal{L}(G_{abb0}) = 0$ and $\mathcal{L}(L_{a0c_2}) \neq 0$.
- (e) If Q_i has a triple nonzero root then one has to examine 3 possibilities: $[G_{abcd}; a = b = c, d = 0]$, $[L_{abcc}; b = c, a = 0]$ and $L_{abbb}; b = 0]$. It suffices to consider the vector $V = [\mathcal{D}, \mathcal{L}]$ and remark that

$$ev(V(G_{aaa0})) = [0,0], ev(V(L_{0cc_2})) = [1,0], \text{ and } ev(V(L_{a0_3})) = [1,1].$$

- 3. If each quartic has at least a zero root then
 - (a) If all the roots are simple then the type is $[G_{abcd}; d=0]$.

- (b) If all the zero roots are simple and there is a nonzero double root then we have 2 possibilities $[G_{abcd}; b = a, c = -2a, d = 0], [L_{ab_2}; a = 0, c = \frac{b}{2}]$. We can discriminate between these two cases by remarking $\mathcal{L}(G_{aa(-2a)0}) = 0$ and $\mathcal{L}(L_{0(2b)b_2}) \neq 0$.
- (c) If all the zero roots are double then we have to consider 5 cases: $[G_{abcd}; a = b = 0, c = d], [L_{abc_2}; a = b, c = 0], [L_{abc_2}; a = b = 0], [L_{a_2b_2}; a = 0], and [L_{a_20_{3\oplus\overline{1}}}; \emptyset].$ We consider the vector $V = [\overline{\mathcal{G}}, \mathcal{G}, \mathcal{H}, \mathcal{L}]$. The evaluation of this vector on the different cases gives

$$ev(V(G_{00aa})) = [0, 0, 0, 0], ev(V(L_{aa0_2})) = [0, 1, 1, 0],$$

$$ev(V(L_{00c_2})) = [0, 0, 1, 0], ev(L_{0_2a_2}) = [1, 1, 1, 0], \text{ and } ev(L_{a_20_{3\oplus\overline{1}}}) = [1, 1, 1, 1].$$

Example V.1. Consider the form $|\varphi\rangle = 2|0100\rangle + |1101\rangle + 4|1111\rangle + 3|0010\rangle$. All the quartics are equal $Q_1(|\varphi\rangle) = Q_2(|\varphi\rangle) = Q_3(|\varphi\rangle) = x^2(x+3y)^2$. Hence, we are in the case (3.c) of the algorithm. We compute $ev([\overline{\mathcal{G}}(|\varphi\rangle), \mathcal{G}(|\varphi\rangle), \mathcal{H}(|\varphi\rangle), \mathcal{L}(|\varphi\rangle)]) = [0, 1, 1, 0]$. So the type of $|\varphi\rangle$ is L_{aa0_2} .

Example V.2. Consider the form $|GHZ\rangle = |0000\rangle + |1111\rangle$. All the quartics are equal $Q_1(|\varphi\rangle) = Q_2(|\varphi\rangle) = Q_3(|\varphi\rangle) = x^2(x-y)^2$. Hence, we are also in the case (3.c) of the algorithm. We compute $ev([\overline{\mathcal{G}}(|GHZ\rangle), \mathcal{G}(|GHZ\rangle), \mathcal{H}(|GHZ\rangle), \mathcal{L}(|GHZ\rangle)) = [0, 0, 0, 0]$. So the type of $|\varphi\rangle$ is G_{00aa} . Indeed, from the definition $|GHZ\rangle = G_{1001}$.

In the same way, the product of two EPR states $|\varphi_1\rangle = |0000\rangle + |0011\rangle + |1100\rangle + |1111\rangle$ gives $Q_1 = Q_3 = (x - y)^4$, and $Q_2 = x^3(x - 4y)$. Hence, we are in the case (2.c) and since $ev([\mathcal{C}, \mathcal{D}, \mathcal{K}_{5,\mathcal{L}}]) = [0, 0, 0, 0]$, this implies that φ_1 has the type G_{a000} , as expected. Indeed, it is exactly $2G_{1000}$.

Example V.3. In Table VII, we summarize the type of the forms used to describe the inclusion diagram of the third secant variety²⁶.

Remark V.1. Our algorithm is based on a discussion on the roots of the quartics Q_1, Q_2 and Q_3 . It can also be seen using the geometrical approach of Section IV:

- 1. If $|\varphi\rangle$ do not vanish the hyperdeterminant Δ then we are in cases 1.(a), 2.(a) and 3.(a).
- 2. If $|\varphi\rangle$ is a smooth point of Δ , then we are in cases 1.(b), 2.(d) and 3.(b).

$ \varphi angle$	Type
65257	G_{abc0}
6014	$L_{0(2a)b_2}$
65261, 65513, 65273, 65259	$L_{a_2 0_{3 \oplus \overline{1}}}$
59777	G_{abc0}
59510	$G_{aa(-2a)0}$
65267, 65509, 65507, 65269, 65510, 65231	$L_{0_{2}a_{2}}$

Table VII. Verstraete type of forms in the third secant variety

- If |φ⟩ is a smooth point of the cusp component (τ(X)*) then we are in cases 1.(c) and 2.(e).
- If |φ⟩ is a smooth point of the cusp component of multiplicity 3 (X_{cusp,3}), then we are in case 2.(c).
- 5. If $|\varphi\rangle$ is a smooth point of the node compotents, then we are in the case 2.(b).
- If |φ⟩ is a smooth point of the node component of multiplicity 3 (X_{node,3}), then we are in the case 3.(c).
- 7. Otherwise $|\varphi\rangle$ belongs to the nullcone.

Remark V.2. As shown in Ref^6 two states of the same Verstraete form are SLOCC equivalent (up to a qubit permutation) if they take the same values on the four-qubit invariants. Thus the algorithm can be used to decide whether two given states are equivalent.

VI. CONCLUSION

In our previous papers^{25,26} we have proposed algorithmic methods based on invariants and covariants to identify the entanglement class of a given state when the number of orbits is finite. This approach remains efficient for studying the nilpotent four-qubit states because the nullcone $\mathcal{N} \subset \mathbb{P}(\mathcal{H})$ contains a finite number of SLOCC orbits. However a covariant classification of $\mathbb{P}(\mathcal{H}) \setminus \mathcal{N}$ is hopeless as we know that the parametrization of $\mathbb{P}(\mathcal{H})$ by SLOCC orbits depends on parameters. Nevertheless the three quartics $\mathcal{Q}_1, \mathcal{Q}_2$ and Q_3 obtained by the three natural embeddings of \mathcal{H} in $\mathfrak{so}(8)$ lead to a stratification of the ambient space according to the configuration of their roots. This discussion has a natural geometric counterpart through the concept of dual variety: The existence of a zero root means the state belongs to the dual of one of the three embedding of $\mathbb{P}^3 \times \mathbb{P}^3$ while the existence of a multiple root means that the state belongs to some singular locus of the hyperdeterminant. In the spirit of the earlier work of Miyake³⁸ we pushed forward the investigation of the singular stratas of the hyperdeterminant. We showed the existence of six stratas whose general points correspond (up to a qubit permutation) to the six families of Verstraete *et al.* classification depending on parameters. The three families which do not depend on parameters correspond to stratas of the nullcone and were studied in Ref²⁶. Thus, identifying the Verstraete form of a state is similar to finding to which strata this state belongs. This can be achieved by the use of invariants (of the three quartics) and covariants as explained in Section V. This algorithm, thanks to the result of Chen *et al.*⁶, can be used to determine whether two given four-qubit states are SLOCC equivalent up to a permutation of qubits.

ACKNOWLEDGEMENT

The authors would like to thank Péter Lévay for sharing with them his ideas on the geometry of four-qubit invariants.

Appendix A: Demitesseract

The study of the higher-dimensional regular polytopes was pioneered by the Swiss mathematician Ludwig Schäfli in the middle of the 19th century, introducing Schäfli symbols which describe all tesselations of an *n*-sphere. The list of regular polytopes was extended to complex polytopes by Shephard in 1952. Readers interested in the subject may refer to an impressive series of books and papers written by $Coxeter^{8,11}$. Regular and semiregular polytopes form a family of geometrical objects whose symmetries are generated by mirrors. In dimension 4, the list of the (real) regular polytopes contains 6 Figures: the 5-cell (also called the 4-simplex) which is the 4-dimensional analogue of the tetrahedron, the tesseract (hypercube in dimension 4), the hexadecachoron (also called 16-cell) which is the 4-dimensional analogue of the octahedron, the icositetrachoron (also called 24-cell), the 120-cell and the 600-cell. In the paper, we use only three of them : the tesseract (16 vertices, 32 edges, 24 faces, and 8 cells), the 16-cell (8 vertices, 24 edges, 32 faces, and 16 cells) and the 24-cell (24 vertices, 96 edges, 96 faces, and 24 cells), see Figure 14. To each polytope, one can associate a dual polytope whose vertices are constructed from the center of its cells. The tesseract and the 16-cell are dual to each other (see Figure 16) and the 24-cell is self dual. The middle of the edges of a 16-cell or the center of the faces of a tesseract are the vertices of a 24-cell. The finite reflection groups associated with their respective mirrors are B_4 for the tesseract and the hypercube, and F_4 for the 24-cell. Notice that B_4 is a subgroup of F_4 .



Figure 14. Octogonal⁹ projection of the tesseract, the 16-cell and the 24-cell⁸.

The demitesseract belongs to the family of demihypercubes, also called half measure polytopes and denoted by $h\gamma_n$, which are constructed from hypercubes by deleting half of the vertices and forming new facets in place of the deleted vertices¹⁰ (see Figure 15). The



Figure 15. Projection of a demitesseract (in red) constructed on the vertices of a tesseract. demitesseract is the demihypercube in dimension 4. As a polytope, the demitesseract is

identical to the regular hexadecachoron. Whilst the hexadecachoron and the demitesseract



Figure 16. A hexadecachoron (in red) whose vertices are the centers of the cells of a tesseract.

are identical as polytopes, their associated symmetries are different. Indeed, the reflection group of the hexadecachoron is B_4 that is the same of the tesseract. The reflections of the demitesseract generates a subgroup of index 2 of B_4 whose associated mirrors are

$$x_1 = x_2, x_2 = x_3, x_3 = x_4, \text{ and } x_1 + x_2 = 0.$$
 (A1)

This group is generated by the transpositions $x_1 \leftrightarrow x_2$, $x_2 \leftrightarrow x_3$, and $x_3 \leftrightarrow x_4$ together with an additional generator which transposes x_1 and x_2 while reversing the sign of both. So, clearly these reflections generate a group isomorphic to D_4 . The normalized demitesseract has 8 vertices, whose coordinates are (1, 1, 1, 1), (1, 1, -1, -1), (1, -1, 1, -1), (-1, 1, 1, -1), (1, -1, -1, 1), (-1, 1, -1, 1), (-1, -1, 1, 1), and (-1, -1, -1, -1), 24 edges, 32 triangular faces and 16 tetrahedral cells.

Appendix B: Discussion on the quartics Q_1 , Q_2 and Q_3

First we focus the discussion on the number of zero roots of the quartic. We have to treat several cases.

1. The three quartics have at least a zero root

This implies that L = M = 0 and this has been completely investigated in our previous paper²⁶. The quartics are

$$Q_1 = Q_2 = Q_3 = x^4 - 2Bx^3y + B^2x^2y^2 + 4D_{xy}xy^3.$$
 (B1)

If $D_{xy} = 0$, the quartics have at least two zero roots and we obtain

$$Q_1 = Q_2 = Q_3 = x^2 (By - x)^2.$$
(B2)

So the only special case occurs when B = 0 and corresponds to the nilpotent forms. The classification of nilpotent forms is well-known and is due to Djokovic *et al.*^{7,13} (see also the previous paper of the authors²⁶ for the geometrical interpretation). Note also that the case when $B \neq 0$ has been already algebraically and geometrically described²⁶. If $D_{xy} \neq 0$ then the discriminant is

$$\Delta = D_{xy}(27D_{xy} - B^3). \tag{B3}$$

One has first to examine the case when the roots of each quartic are distinct. This corresponds to $\Delta \neq 0$ and has been investigated in our previous paper²⁶ (no other cases appear in the present discussion).

If the quartics have nonzero multiple roots then the forms belong to the variety defined by

$$27D_{xy} - B^3 = 0. (B4)$$

This variety have been also investigated in our previous $paper^{26}$. Nevertheless the present discussion may refine the classification by regarding the other covariants of the quartics:

$$T = 1152(D_{xy}x^{6} + D_{xy}Bx^{5}y + 10D_{xy}^{2}x^{3}y^{2} - 5D_{xy}^{2}Bx^{2}y^{4} + B^{2}D_{xy}^{2}xy^{5} + 2D_{xy}^{3}y^{6})$$

$$Hess = -12(B^{2}x^{4} - 2(12D_{xy} + B^{3})x^{3}y + B(12D_{xy} + B^{3})x^{2}y^{2} + 4B^{2}D_{xy}xy^{3} + 12D_{xy}^{2}y^{4})$$

$$I_{2} = 2BD_{xy} + \frac{1}{12}H^{4}$$

$$I_{3} = -(D_{xy}^{2} + \frac{1}{216}B^{6} + \frac{1}{6}B^{3}D_{xy}).$$
(B5)

Since $D_{xy} \neq 0$ the covariants T and *Hess* are clearly nonzero. According to Table I the only case to investigate is $I_2 = I_3 = 0$. But this implies $B = D_{xy} = 0$ and this case has already been dealt with. It follows that all the interesting cases have been already investigated in our previous article²⁶.

2. Only one of the quartics has a zero roots

a. The quartic Q_1 has a zero root

In this case, one has L = 0 and M = -N (we will see that the other two cases are symmetrical). The quartics are

$$Q_1 = x^4 - 2Bx^3y + (B^2 + 4M)x^2y^2 + 4(D_{xy} - BM)xy^3,$$
(B6)

$$Q_2 = Q_3 = x^4 - 2Bx^3y + (B^2 - 2M)x^2y^2 - 2(BM - 2D_{xy})xy^3 + M^2y^4.$$
 (B7)

The covariants reads

$$I_2 = 2B(D_{xy} - BM) + \frac{1}{12}(B^2 + 4M)^2,$$
 (B8)

$$I_3 = -(D_{xy} - BM)^2 - \frac{1}{216}(B^2 + 4M)^3 - \frac{1}{6}B(B^2 + 4M)(D_{xy} - BM),$$
(B9)

$$Hess(Q_1) = -12 (B^2 - 8 M)x^4 + 24 (B^3 - 8 B M + 12 D_{xy})x^3y - 12 (12 B D_{xy} - 4 B^2 M + B^4 + 16 M^2)x^2y^2 + 48 (B^2 + 4 M)(-D_{xy} + B M)xy^3 - 144 (-D_{xy} + B M)^2y^4,$$
(B10)

$$Hess(Q_2) = -12 (B^2 + 4M) x^4 + 24 (B^3 - 8BM + 12D_{xy}) x^3 y - 12 (-10B^2M + 12BD_{xy}) + B^4 - 8M^2) x^2 y^2 + 24 (B^3M - 2B^2D_{xy} - 8BM^2 + 4MD_{xy}) x y^3 - 12 (B^2M^2 + 4M^3 - 12D_{xy}BM + 12D_{xy}^2) y^4,$$
(B11)

$$T(Q_1) = -1152(D_{xy}x^6 - (B D_{xy} + 4 M^2)x^5y + 10 M (-D_{xy} + B M)x^4y^2 -10 (-D_{xy} + B M)^2x^3y^3 + 5 B (-D_{xy} + B M)^2x^2y^4 - (B^2 + 4 M)(-D_{xy} + B M)^2xy^5 + 2 (-D_{xy} + B M)^3y^6),$$
(B12)

and

$$T(Q_2) = -1152 (D_{xy} - B M)(x^6 - B x^5 y + 5 M x^4 y^2 - 10 D_{xy} x^3 y^3 + 5(B D_{xy} - M^2) x^2 y^4 - (B^2 D_{xy} - B M^2 - 2 M D_{xy}) x y^5 + (D_{xy} B M - M^3 + 2 D_{xy}^2) y^6).$$
(B13)

According to Table I, we have to investigate several cases.

• The quartic Q_1 has a double zero root

This case is identified by the equation $D_{xy} = BM$ and implies automatically $\Delta = 0$. The quartics are

$$Q_1 = x^4 - 2Bx^3y + (B^2 + 4M)x^2y^2, (B14)$$

$$Q_2 = Q_3 = (-x^2 + Bxy + y^2)^2.$$
(B15)

The covariants simplify as

$$I_2 = \frac{1}{2}(B^2 + 4M)^2,$$
 (B16)

$$I_3 = \frac{1}{216} (B^2 + 4M)^3, \quad (B17)$$

$$T(Q_1) = -1152(BMx^6 - M(B^2 + 4M)x^5y), \quad (B18)$$

$$Hess(Q_1) = -12(B^2 - 8M)x^4 + 24(B^2 + 4M)Bx^3y - 12(B^2 + 4M)^2x^2y^2, \quad (B19)$$

$$Hess(Q_2) = Hess(Q_3) = -12(B^2 + 4M)(Bxy + My^2 - x^2)^2.$$
 (B20)

Since Q_2 has two double roots, one has $T(Q_2) = 0$. It remains to examine the following cases

- 1. If Q_1 has two simple distinct nonzero roots then $B^2 + 4M \neq 0$. It follows that $I_2, I_3 \neq 0$ and Q_2 has no triple root. The only remaining case to consider is the case when Q_2 has a quadruple root. But this implies $B^2 + 4M = 0$ and so it is not possible.
- 2. If Q_1 has a nonzero double root then $T(Q_1) = 0$ and this implies $M = D_{xy} = 0$. This configuration cannot occur since it implies that Q_2 has a zero root.
- 3. If Q_1 has a triple zero root then $B^2 + 4M = 0$ and $Hess(Q_2) = 0$. Hence, Q_2 has a quadruple root.
- 4. If Q_1 has a quadruple zero root then M = 0. So this configuration is not possible.

In conclusion, when L = 0 and $M \neq 0$, we have only to investigate the variety given by $B^2 + 4M = 0$.

• Q_1 has a nonzero double root and two simple roots

Since the zero roots of Q_1 is not double, one has $D_{xy} \neq BM$ and then $T(Q_1), T(Q_2) \neq 0$. This prove that Q_2 has exactly one double root.

• Q_1 has a triple nonzero root

We have $D_{xy} \neq BM$ and $I_2 = I_3 = 0$. The equation $I_2 = I_3 = 0$ admits two solutions: $\{M = -\frac{1}{4}B^2, D_{xy} = -\frac{1}{4}B^3\}$ and $\{M = \frac{1}{12}B^2, D_{xy} = \frac{1}{108}B^3\}$. The first solution implies $D_{xy} = BM$, so it must be excluded. From the second solution, we deduce

$$T(Q_1) = -\frac{32}{2187}B^3(2By - 3x)^6,$$
 (B21)

$$T(Q_2) = \frac{4}{2187} B^3 (By - 6x)^6, Hess(Q_1) = -\frac{4}{81} B^2 (2By - 3x)^4,$$
(B22)

$$Hess(Q_2) = -\frac{1}{81}B^2(By - 6x)^4.$$
 (B23)

The only special case is B = 0 and that the form is nilpotent.

• Q_1 has only simple roots

In this case Q_2 has also only simple roots. This case corresponds to

$$\Delta = (B^3 D_{xy} - B^2 M^2 - 18 D_{xy} BM + 16 M^3 + 27 D_{xy})(BM - D_{xy})^2 \neq 0.$$
(B24)

b. Q_2 has a zero root

In this case, one has M = 0 and L = -N. The quartics are

$$Q_2 = x^4 - 2Bx^3y + (B^2 - 4L)x^2y^2 + 4D_{xy}xy^3,$$
(B25)

$$Q_1 = Q_3 = x^4 - 2Bx^3y + (B^2 + 2L)x^2y^2 - 2(BL - 2D_{xy})xy^3 + L^2y^4.$$
 (B26)

The covariants read

$$I_2 = 3\left(\frac{1}{6}B^2 - \frac{2}{3}L\right)^2 + 2BD_{xy},\tag{B27}$$

$$I_3 = -D_{xy}^2 - \left(\frac{1}{6}B^2 - \frac{2}{3}L\right)^3 - B\left(\frac{1}{6}B^2 - \frac{2}{3}L\right)D_{xy},\tag{B28}$$

$$Hess(Q_2) = -12 (B^2 + 8L) x^4 + 24 (B^3 - 4BL + 12D_{xy}) x^3 y - 12 (-8B^2L + 12BD_{xy}) + B^4 + 16L^2 x^2 y^2 - 48(B^2 - 4L) x y^3 - 144D_{xy}^2 y^4,$$
(B29)

$$Hess(Q_1) = -12 (B^2 - 4L) x^4 + 24 (B^3 - 4BL + 12D_{xy}) x^3 y - 12 (-2B^2M + 12BD_{xy}) + B^4 - 8L^2) x^2 y^2 - 24 (-B^3L + 2B^2D_{xy} + 4BL^2 + 4LD_{xy}) x y^3 - 12 (B^2L^2 - 4L^3 - 12D_{xy}BL + 12D_{xy}^2) y^4,$$
(B30)

$$T(Q_2) = -1152(D_{xy} - BL) x^6 - (B D_{xy} + 4 L^2 - B^2L) x^5y + 10 L D_{xy} x^4 y^2 -10 D_{xy}^2 x^3 y^3 + 5 B D_{xy}^2 x^2 y^4 - (B^2 - 4 L) D_{xy}^2 x y^5 - 2 D_{xy}^3 y^6),$$
(B31)

and

$$T(Q_1) = -1152 D_{xy}(x^6 - B x^5 y - 5 L x^4 y^2 - 10 (D_{xy} - BL) x^3 y^3 + 5(B D_{xy} - B^2 L - L^2) x^2 y^4 - (B^2 D_{xy} - B^3 L + 2L D_{xy} - 3BL^2) x y^5$$
(B32)
$$-(B^2 L^2 - L^3 + 2D_{xy}^2 - 3BL D_{xy}) y^6).$$

Hence, the reasoning is very similar to the case L = 0. Let us summarize it below.

According to table I, we have to investigate several cases.

• The quartic Q_2 has a double zero root

The case is identified by the equation $D_{xy} = 0$ and implies $\Delta = 0$, $Q_1 = Q_3 = (x^2 - Bxy + L^2y^2)^2$, $Q_2 = x^2(x^2 - 2Bxy + (B^2 - 4L)y^2)$, $I_2 = \frac{1}{12}(B^2 - 4L)^2$ and $I_3 = -\frac{1}{216}(B^2 - 4L)^3$. Hence, when $L \neq 0$ and M = 0, we have only to investigate the variety given by $B^2 - 4L = 0$. This corresponds the case when Q_1 has a quadruple root and Q_2 has a triple zero root.

• Q_2 has a nonzero double root and two simple roots

In this case, $D_{xy} \neq 0$ and so $T(Q_1), T(Q_2) \neq 0$. So Q_1 has exactly one double root.

• Q_2 has a triple nonzero root

The only solution of $I_2 = I_3 = M = 0$ satisfying $D_{xy} \neq 0$ is $D_{xy} = -\frac{2}{27}B^3$ and $L = -\frac{1}{12}B^2$. The only special case is B = 0 and implies the nilpotence of the form.

• Q_2 has only simple roots

In this case Q_1 has also four distinct roots. Furthermore, the hyperdeterminant factorizes as

$$\Delta = D_{xy}^2 (36BLD_{xy} + B^4L - B^3D_{xy} - 27D_{xy}^2 + 16L^3 - 8B^2L^2).$$

c. Q_3 has a zero root

In this case, one has N = 0 and L = -M. The quartics are

$$Q_3 = x^4 - 2Bx^3y + (B^2 - 4M)x^2y^2 + 4D_{xy}xy^3, \qquad (B33)$$

$$Q_1 = Q_2 = x^4 - 2Bx^3y + (B^2 + 2M)x^2y^2 - 2(BM - 2D_{xy})xy^3 + M^2y^4.$$
 (B34)

So it is deduced by substituting $L \leftrightarrow M$ and $Q_2 \leftrightarrow Q_3$ in the previous discussion (M = 0).

3. The quartics have no zero root

We have to investigate three cases.

a. Q_1 has four simple roots

This means that $\Delta \neq 0$ and then Q_2 and Q_3 have both four simple roots.

b. $T(Q_1) = 0$

The equation $T(Q_1) = 0$ a has two solutions: $M = D_{xy} = 0$ and $N = D_{xy} = 0$. In this two cases, one of the quartic has a zero root. In the same way, if $T(Q_2) = 0$ or $T(Q_3) = 0$ then one of the quartic has a zero root.

c.
$$T(Q_1), T(Q_2), T(Q_3) \neq 0$$

Since Hess = 0 implies T = 0 (if a form has a quadruple root then it has two double roots which are equal), one has two cases to consider:

1. $I_2, I_3 \neq 0$,

2.
$$I_2 = I_3 = 0.$$

Appendix C: Symmetries of the Verstraete forms

1. Permutations of the qubits

Versraete *et al.*⁴³ gave nine inequivalent normal forms for the four qubic forms. These forms were defined up to a permutation of qubits. One of these forms G_{abcd} has 4 parameters and the set of all the G_{abcd} defines a 4-dimension subspace of the ambient space. The five other forms, $L_{abc_2}, L_{a_2b_2}, L_{ab_3}, L_{a_4}$ and $L_{a_20_{3\oplus\overline{1}}}$ have one to three parameters and define five affine subspaces. The remaining three forms, $L_{0_{5\oplus\overline{3}}}, L_{0_{7\oplus\overline{1}}}$ and $L_{0_{3\oplus\overline{1}}0_{3\oplus\overline{1}}}$ are nilpotent. For a given form φ , we will denote by φ^{σ} the form obtained by applying the permutation σ on the qubits. For the more generic form G_{abcd} , we obtain 5 other non equivalent forms G_{abcd}^{1423} , $G_{abcd}^{1324}, G_{abcd}^{1243}, G_{abcd}^{1324}$ and G_{abcd}^{1432} . Each of them is equivalent to a $G_{a'b'c'd'}$ for one of the following specializations (which are involutions):

- (a, b, -c, d),
- $\left(\frac{a+b-c+d}{2}, \frac{a+b+c-d}{2}, \frac{-a+b+c+d}{2}, \frac{a-b+c+d}{2}\right)$,

Forms	Strata
G_{abcd}	Gr_0
L_{abc_2}	Gr_1
$L_{a_2b_2}$	Gr_2
$L_{a_2 0_{3 \oplus \overline{1}}}$	Gr_3
$L_{0_{3\oplus\overline{1}}0_{3\oplus\overline{1}}}$	Gr_4
L_{ab_3}	Gr_5
L_{a_4}	Gr_6
$L_{0_{5\oplus\overline{3}}}$	Gr_7
$L_{0_{7\oplus\overline{1}}}$	Gr_8

Table VIII. Correspondence between Verstraete forms and nilpotent strata

- (a+b-c+d/2), a+b+c-d/2, a-b-c-d/2, -a+b-c-d/2),
 (a+b+c+d/2), a+b-c-d/2, a-b-c+d/2, -a+b-c+d/2),
- and $\left(\frac{a+b+c+d}{2}, \frac{a+b-c-d}{2}, \frac{a-b+c-d}{2}, \frac{a-b-c+d}{2}\right)$.

The permutations of L_{abc_2} split into 6 families with representatives $L_{abc_2}, L_{abc_2}^{1342}, L_{abc_2}^{2431}, L_{abc_2}^{1432}, L_{abc_2}^{2341}$, and $L_{abc_2}^{2143}$. Furthermore, we have

$$\begin{split} L^{1324}_{\left(\frac{a+b}{2}+c,\frac{a+b}{2}-c,\frac{a-b}{2}\right)} &\sim L^{1243}_{abcc}, \ L^{2431}_{\left(\frac{a+b}{2}+c,\frac{a+b}{2}-c,\frac{a-b}{2}\right)} &\sim L^{2134}_{abcc}, \\ L^{1432}_{\left(\frac{a+b}{2}+c,\frac{a+b}{2}-c,\frac{a-b}{2}\right)} &\sim L^{1423}_{abcc}, \ L^{1234}_{\left(\frac{a+b}{2}+c,\frac{a+b}{2}-c,\frac{a-b}{2}\right)} &\sim L^{1324}_{abcc}, \\ L^{2341}_{\left(\frac{a+b}{2}+c,\frac{a+b}{2}-c,\frac{a-b}{2}\right)} &\sim L^{2314}_{abcc}, \ L^{2143}_{\left(\frac{a+b}{2}+c,\frac{a+b}{2}-c,\frac{a-b}{2}\right)} &\sim L^{2413}_{abcc}. \end{split}$$

There are also 6 different families of permutations of L_{ab_3} whose representatives are $L_{ab_3}^{\sigma}$ for $\sigma \in \{1234, 1423, 3421, 1342, 3412\}$. The permutations of $L_{a_2b_2}$ generate 12 non equivalent families $L_{a_2b_2}^{\sigma}$ for $\sigma \in \{1234, 2314, 2413, 1324, 1243, 2134, 2431, 2143, 3241, 3124, 3214\}$. The permutations $L_{a_4}^{\sigma}$ for $\sigma \in \mathfrak{S}_4$ are pairwise nonequivalent. Finally, the permutations of $L_{a_20_{3\oplus\overline{1}}}$ give 4 nonequivalent families $L_{a_20_{3\oplus\overline{1}}}^{1234}$, $L_{a_20_{3\oplus\overline{1}}}^{2134}$, and $L_{a_20_{3\oplus\overline{1}}}^{2314}$.

If we send each parameter to zero, the permutations of the Verstraete forms specialize to nilpotent orbits. To each form corresponds one of the strata defined in our previous paper²⁶ see Table VIII.

Forms	$\mathcal{Q}(G_{abcd})$	$\mathcal{Q}(L_{abc_2})$	$\mathcal{Q}(L_{ab_3})$	$\mathcal{Q}(L_{a_2b_2})$	$\mathcal{Q}(L_{a_4})$	$\mathcal{Q}(L_{a_2 0_{3 \oplus \overline{1}}})$
G_{abcd}	$\mathcal{Q}(G_{abcd})$					
L_{abc_2}	$\mathcal{Q}(G_{abcc})$					
L_{ab_3}	$\mathcal{Q}(G_{aaab})$	$\mathcal{Q}(L_{aba})$				
$L_{a_2b_2}$	$\mathcal{Q}(G_{aabb})$	$\mathcal{Q}(L_{aab})$				
L_{a_4}	$\mathcal{Q}(G_{aaaa})$	$\mathcal{Q}(L_{aaa})$	$\mathcal{Q}(L_{aa_3})$			
$L_{a_2 0_{3 \oplus \overline{1}}}$	$\mathcal{Q}(G_{aa00})$	$\mathcal{Q}(L_{aa0})$	$\mathcal{Q}(L_{a_20_2})$			
$L_{0_{5\oplus\overline{3}}}$	$\mathcal{Q}(G_{0000})$	$\mathcal{Q}(L_{000})$	$\mathcal{Q}(L_{0_20_2})$	$\mathcal{Q}(L_{00_3})$	$\mathcal{Q}(L_{0_4})$	$\mathcal{Q}(L_{0_20_{3\oplus\overline{1}}})$
$L_{0_{7\oplus\overline{1}}}$	$\mathcal{Q}(G_{0000})$	$\mathcal{Q}(L_{000})$	$\mathcal{Q}(L_{0_20_2})$	$\mathcal{Q}(L_{00_3})$	$\mathcal{Q}(L_{0_4})$	$\mathcal{Q}(L_{0_20_{3\oplus\overline{1}}})$
$L_{0_{3\oplus\overline{1}}0_{3\oplus\overline{1}}}$	$\mathcal{Q}(G_{0000})$	$\mathcal{Q}(L_{000})$	$\mathcal{Q}(L_{0_20_2})$	$\mathcal{Q}(L_{00_3})$	$\mathcal{Q}(L_{0_4})$	$\mathcal{Q}(L_{0_20_{3\oplus\overline{1}}})$

Table IX. Values of \mathcal{Q} for Verstrate forms

2. Quadrics again

In a general setting, permuting the qubits in a form induces a permutations on the quadrics Q_1, Q_2, Q_3 . Let us denote $\mathcal{Q}(\varphi) = \{Q_1(\varphi), Q_2(\varphi), Q_3(\varphi)\}$. We notice also that any of the quadrics in the Verstraete forms can be written as $Q_1(G_{abcd})$ for some specialization of the parameters a, b and c. Indeed we have

$$\mathcal{Q}(G_{abcd}) = \left\{ Q_1\left(G_{abcd}\right), Q_1\left(G_{\frac{a+c-d-b}{2}\frac{a+d-b-c}{2}\frac{a+b-c-d}{2}\frac{a+b+c+d}{2}}\right) Q_1\left(G_{\frac{a+b+c-d}{2}\frac{a+b-c+d}{2}\frac{a-b+c+d}{2}\frac{a-b+c+d}{2}}\right) \right\}.$$

The remaining values are summarized in Table IX.

REFERENCES

- ¹Arnol'd V., "Normal forms for functions near degenerate critical points, the Weyl groups of A k, D k, E k and Lagrangian singularities." Functional Analysis and its applications 6.4 (1972): 254-272.
- ²Borsten L., Dahanayake D., Duff M. J., Marrani A. and Rubens W., "Four-Qubit Entanglement Classification from String Theory", Phys. Rev. Lett. **105**, 100507 (2010).
- ³Borsten L., Duff M. J., and Levay P., "The black-hole/qubit correspondence: an up-to-date review.", arXiv preprint arXiv:1206.3166 (2012).

- ⁴Briand E., Luque J.-G., Thibon J.-Y., "A complete set of covariants of the four-qubit system" Journal of Physics A: mathematical and general. 36.38 (2003): 9915.
- ⁵Cao Y., and Wang. A. M. "Discussion of the entanglement classification of a 4-qubit pure state." The European Physical Journal D 44, no. 1 (2007): 159-166.
- ⁶Chen L. Doković D., Grassl M and Zeng B. "Four-qubit pure states as fermionic states." Physical Review A 88, no. 5 (2013): 052309.
- ⁷Chterental O. and Djokovic D., "Normal forms and tensor ranks of pure states of fourqubit", arXiv preprint quant-ph/0612184 (2006).
- ⁸Coxeter H.S.M., *Regular polytopes*, Dover publications, inc. New York (1973).
- ⁹Coxeter H.S.M., "Two aspects of the regular 24-cell in four dimensions", in *Kaleidoscopes*, selected writing of H.S. Coxeter, Wiley-Interscience publication (1995).
- ¹⁰Coxeter H.S.M., "Regular and semiregular polytopes.III", in *Kaleidoscopes, selected writ*ing of H.S. Coxeter, Wiley-Interscience publication (1995).
- ¹¹Coxeter H.S.M., Complex regular polytopes. Cambridge University Press; 2 edition (April 26, 1991)
- ¹²Dimca, A. (1986). "Milnor numbers and multiplicities of dual varieties". Revue Roumaine de Mathématiques Pures et Appliquées, 31(6), 535-538.
- ¹³Djokovic D., Lemire N. and Sekiguchi J., "The closure ordering of adjoint nilpotent orbits in $\mathfrak{so}(p,q)$ ", Tohoku Mathematical Journal 53.3 (2001): 395-442.
- ¹⁴Eltschka C., and Siewert. "Quantifying entanglement resources." Journal of Physics A: Mathematical and Theoretical 47, no. 42 (2014): 424005.
- ¹⁵W. Fulton, J. Harris, *Representation Theory*, Graduate Text in Mathematics, Springer 1991.
- ¹⁶Gour G. and Wallach N., "On symmetric SL-invariant polynomials in four-qubit." In Symmetry: Representation Theory and Its Applications, pp. 259-267. Springer New York, 2014.
- ¹⁷Gelfand I.M., Kapranov M.M., Zelevinsky A.V., "Hyperdeterminants", Advances in Mathematics 96, vol 2 (1992).
- ¹⁸I.M Gelfand M.M Kapranov A.V. Zelevinsky, Discriminants, Resultants and Multidimensional Determinants, Birkhäuser 1994.
- ¹⁹Grassl, M., Beth, T., and Pellizzari, T. (1997). "Codes for the quantum erasure channel". Physical Review A, 56(1), 33.

- ²⁰Gühne, O., Jungnitsch, B., Moroder, T. and Weinstein, Y. S. (2011). "Multiparticle entanglement in graph-diagonal states: Necessary and sufficient conditions for four qubits". Physical Review A, 84(5), 052319.
- ²¹J. Harris, Algebraic Geometry: a first course, Graduate Texts in Mathematics 133 Springer 1992.
- ²²Heydari H., "Geometrical Structure of Entangled States and the Secant Variety", Quantum Information Processing 7 (1), 3-32 (2008).
- ²³Holweck, F. and Lévay, P., 2016. "Classification of multipartite systems featuring only $|W\rangle$ and $|GHZ\rangle$ genuine entangled states". Journal of Physics A: Mathematical and Theoretical, 49(8), (2016) p.085201.
- ²⁴Holweck F., Luque J.-G., and Planat M. "Singularity of type D4 arising from four-qubit systems." Journal of Physics A: Mathematical and Theoretical 47.13 (2014): 135301.
- ²⁵Holweck F., Luque J.-G., Thibon J.-Y., "Geometric descriptions of entangled states by auxiliary varieties", Journal of Mathematical Physics 53, 102203 (2012).
- ²⁶Holweck F., Luque J.-G., Thibon J.-Y., "Entanglement of four-qubit systems: a geometric atlas with polynomial compass I (the finite world)". a completer.
- ²⁷Horodecki R., Horodecki P., Horodecki M., Horodecki K., "Quantum entanglement", Reviews of Modern Physics, 81(2), 865 (2009).
- ²⁸T. Ivey, J.M. Landsberg, Cartan for beginners: Differential Geometry via Moving Frames and Exterior Differential Systems, Graduate Studies in Mathematics **61** 2003.
- ²⁹Lamata L., León J., Salgado D. and Solano E., "Inductive entanglement classification of four-qubit under stochastic local operations and classical communication", Phys. Rev. A. 72, 022318 (2007).
- ³⁰Landsberg J. M., Tensors: Geometry and applications, Vol. 128. Amer Mathematical Society, 2011.
- ³¹Lévay, P., "On the geometry of four-qubit invariants." Journal of Physics A: Mathematical and General 39.30 (2006): 9533.
- ³²Lévay P., "STU black holes as four-qubit systems", Pys. Rev. D 82, 026003 (2010).
- ³³Lévay P. and Holweck F., "Embedding qubits into fermionic Fock space: Peculiarities of the four-qubit case." Physical Review D 91.12 (2015): 125029.
- ³⁴Li D., Li X., and Huang H., "SLOCC classification for nine families of four-qubits." arXiv preprint arXiv:0712.1876 (2007).

- ³⁵Lin S. and Sturmfels B. "Polynomial relations among principal minors of a 4 × 4-matrix." Journal of Algebra 322, no. 11 (2009): 4121-4131.
- ³⁶Luque J.-G. and Thibon J.-Y, "The polynomial invariants of four-qubit", Phys. Rev. A 67, 042303 (2003).
- ³⁷Luque, J.-G., and Thibon J.-Y, "Algebraic invariants of five qubits" Journal of physics A: mathematical and general 39, no. 2 (2005): 371.
- ³⁸Miyake A., "Classification of multipartite entangled states by multidimensional determinants", Phys. Rev. A 67, 012108 (2003).
- ³⁹Olver P., Classical Invariant Theory, Cambridge University Press, Cambridge UK, 1999.
- ⁴⁰Parusiński, A. "Multiplicity of the dual variety". Bulletin of the London Mathematical Society, 23(5), 429-436. (1991).
- ⁴¹Popov V., Vinberg E., "Invariant theory." Algebraic geometry IV. Springer Berlin Heidelberg, 1994. 123-278.
- ⁴²Tevelev E. A., "Projectively Dual Varieties", Journal of Mathematical Sciences **117** (6), 4585-4732 (2003).
- ⁴³Verstraete F., Dehaene F., De Moor B. and Verschelde H., "Four qubits can be entangled in nine different ways", Phys. Rev.. A 65, 052112 (2002).
- ⁴⁴Weyman J. and Zelevinsky A., "Singularities of Hyperdeterminants", Annales de l'Institut Fourier 46, 591-644 (1996).
- ⁴⁵F. Zak, Tangents and Secants of Algebraic Varieties, AMS Translations of mathematical monographs **127** 1993.
- ⁴⁶F. Zak, Determinants of projective varieties and their degrees, Algebraic Transformation Groups and Algebraic Varieties, Encyclopaedia of Mathematical Sciences, vol. 132, Subseries Invariant Theory and Algebraic Transformation Groups, vol. III, Springer-Verlag, Berlin-Heidelberg-New York, 2004.