

# Solving the time-dependent Schrödinger equation via Laplace Transform

Natascha Riahi \*

University of Vienna, Faculty of Physics, Gravitational Physics  
Boltzmanng. 5, 1090 Vienna, Austria

August 1, 2016

## Abstract

We show how the Laplace transform can be used to give a solution of the time-dependent Schrödinger equation for an arbitrary initial wave packet if the solution of the stationary equation is known. The solution is obtained without summing up eigenstates nor do we need the path integral. We solve the initial value problem for three characteristic piecewise constant potentials. The results give an intuitive picture of the wave packet dynamics, reproducing explicitly all possible reflection and transmission processes. We investigate classical and quantum properties of the evolution and determine the reflection and transmission probabilities.

## 1 Introduction

The most elementary and standard way to solve the time-dependent Schrödinger equation is to express the initial wave function in terms of the eigenfunctions and write the solution as a sum of and/or an integral over the corresponding eigensolutions oscillating with the corresponding frequencies. This method is appropriate to investigate the revival behaviour [1] which is a pure quantum phenomenon that occurs in many quantum systems at late times. But in order to get information about details of the wave packet dynamics including the transition from classical to quantum mechanics taking place much earlier than the revival phenomenon the eigenstates have to be summed up numerically [2],[3]. Moreover there are cases such as the finite square well where the energy eigenvalues  $E(n)$  are only given implicitly [4], which complicates the summation.

An alternative route is the Feynman path integral that contains the classical action and seems to be a good candidate for the study of quasiclassical behaviour. But even attempts to simplify the path integral via semiclassical approximations such as the Gutzwiller trace formula [5] or the method of cellular

---

\*e-mail address: riahi@ap.univie.ac.at

dynamics [6] still lead to involved expressions. Moreover evaluating these semi-classical formulas one can never be sure how long they are a good approximation for the exact behaviour of the wave function.

So the question arises whether there is something in between the brute force method of summing up eigenstates and entering the world of path integrals. For the infinite square well a third possibility is known. The so-called mirror solution [7] consists of a sum of free particle solutions, describing a wave packet (particle) being reflected from one wall to the other. This solution is achieved by just considering the special symmetry of the problem and making a successful guess. So it would be desirable to have a method to derive such intuitive solutions also for more general problems.

The method of Laplace transform that we will introduce gives the exact solution of the time-dependent Schrödinger equation when the solution of the stationary equation is known. For our purpose it is not necessary to identify the eigenvalues. Instead the Laplace transformed Schrödinger equation contains the initial wave packet as an inhomogeneous term which allows to apply the method of variation of constants. The solution is then given in terms of the initial wave packet. The only technical difficulty is to perform the inverse Laplace transform. Here we can use the toolbox of rules and inversion formulas, provided for instance in [8],[9].

Though the technique of Laplace transformation has been applied to several specific problems concerning the time dependent Schrödinger equation (see below), the proposed method has not yet been consequently used and explored as a tool to solve the initial value problem of wave packet evolution.

In [10] and [11] the Laplace transform was used to determine a perturbative solution of the time dependent Schrödinger equation with the help of an eigenfunction expansion of the unperturbed Hamiltonian. In [12] and [13] the Laplace transform was applied to problems with time dependent boundary conditions. [14],[15] explored the late time behaviour of special initial wave functions for tunneling phenomena. Both authors required the identification of complex poles of the Laplace transformed solution which brings back a problem of similar difficulty as the identification of eigenvalues and makes the solutions less explicit.

What comes closer to the proposed method is the derivation of the propagator  $K(x, x', t)$  from the Green function which means to invert the half-sided Fourier transform ([16], [17])

$$G(x, x', E) = \int_0^{\infty} K(x, x', t) e^{iEt} dt.$$

Since this integral can in general not be expected to converge this implies a regularization  $E \rightarrow E + i\epsilon$ . This makes the Laplace transform favorable, where such a procedure is not needed and the existing inversion formulas can be applied.

We start by introducing the method in section 2. In section 3 we show that we can reproduce the mirror solution of the infinite square well in a straightforward way. We proceed with the potential step in section 4. The derivation of the exact solution in this case is significantly easier than the derivation of the

propagator in [18],[17] where the PDX method for path integrals was applied. Apart from being of similar simplicity as the most explicit solutions available in the literature ([19],[17]) the obtained solutions admit a direct insight into the wave packet dynamics. We show directly that sufficiently peaked wave packets with energy higher than the potential step will be slowed down as a corresponding classical particle which was already derived in [16] using the saddle point approximation. And we investigate the wave packet behaviour in the classically forbidden region for wave packets with energy lower than the energy of the step. We are able to derive an exact result for the reflection probability in terms of the transmission coefficient of the time independent solutions (33). This result confirms a conjecture in [20] under a certain condition for the initial wave packet.

In section 5 we investigate the asymmetric square well, a potential that is a combination of the two preceding ones and can be described as a box with exit. As far as we know the initial value problem of this model was not considered before. We find out that the solution is a very intuitive combination of the solutions of the infinite square well and the potential step. We show that for reasonably peaked wave packets and after not too long times the probability of a particle to be found in the box is given by powers of the reflection probability of the potential step, where the power is given by the number of reflections the wave packet has already undergone.

## 2 The method

The Laplace transform [8]  $\varphi(x, s)$  of the wavefunction  $\psi(x, t)$  reads

$$\varphi(x, s) = \mathcal{L}(\psi(x, t)) = \int_0^{\infty} \psi(x, t) e^{-st} dt \quad . \quad (1)$$

Here and from now on we always assume that the Laplace transform of the wave function exists which is ensured if the wavefunction does not grow faster than exponentially in time.

Applying the Laplace transformation to the Schrödinger equation

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} + V(x) \psi(x, t) = i\hbar \frac{\partial \psi(x, t)}{\partial t} \quad (2)$$

we find

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \varphi(x, s)}{\partial x^2} + V(x) \varphi(x, s) = i\hbar s \varphi(x, s) - i\hbar \psi(x, 0), \quad (3)$$

where  $\psi(x, 0)$  is the initial wave function.

Equation (3) is an inhomogeneous linear ordinary second order differential equation for the Laplace-transformed wave function  $\varphi(x, s)$ . When two linearly

independent solutions of the homogeneous equation are known a particular solution of (3) can be achieved by the method of variation of constants. With the two homogeneous solutions  $u_1(x, s)$  and  $u_2(x, s)$  we get the particular solution

$$\varphi_p(x, s) = u_1(x, s) \int_c^x \frac{2mi}{\hbar} \frac{u_2(y, s)}{W[u_1, u_2]} \psi(y, 0) dy - u_2(x, s) \int_c^x \frac{2mi}{\hbar} \frac{u_1(y, s)}{W[u_1, u_2]} \psi(y, 0) dy \quad (4)$$

Here  $W[u_1, u_2]$  is the Wronskian of the two homogeneous solutions:

$$W[u_1, u_2] = \frac{\partial u_1(x, s)}{\partial x} u_2(x, s) - \frac{\partial u_2(x, s)}{\partial x} u_1(x, s) \quad .$$

Therefore the general solution of (3) reads

$$\varphi(x, s) = \varphi_p(x, s) + \alpha(s) u_1(x, s) + \beta(s) u_2(x, s) \quad (5)$$

The functions  $\alpha(s)$  and  $\beta(s)$  are determined by the boundary conditions on  $\varphi(x, s)$ , which stem from the physical boundary conditions on the wave function  $\psi(x, t)$ . What is left is working out the inverse Laplace transformation for the specific problem.

### 3 The infinite square well

The infinite square well is a physical model system which can be interpreted as the one-dimensional description of a particle locked up by infinitely rigid walls. The appropriate potential is

$$\begin{aligned} V(x) &= \infty & \text{for } x \leq 0 \\ V(x) &= 0 & \text{for } 0 < x < d \\ V(x) &= \infty & \text{for } x \geq d \quad . \end{aligned}$$

The dynamics of a wave packet is therefore determined by

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} = i\hbar \frac{\partial \psi(x, t)}{\partial t}, \quad \text{for } x \in [0, d] \quad (6a)$$

with the boundary conditions

$$\psi(0, t) = \psi(d, t) = 0. \quad (6b)$$

The Laplace transformed wave packet fulfills

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \varphi(x, s)}{\partial x^2} = i\hbar s \varphi(x, s) - i\hbar \psi(x, 0) \quad (7a)$$

with the boundary conditions

$$\varphi(0, s) = \varphi(d, s) = 0 \quad . \quad (7b)$$

The solution of the homogeneous equation corresponding to (7a) is a linear combination of the functions

$$u_1(x, s) = e^{i\sqrt{\frac{2msi}{h}}x} \quad u_2(x, s) = e^{-i\sqrt{\frac{2msi}{h}}x} \quad .$$

With the Wronskian  $W[u_1, u_2] = 2i\sqrt{\frac{2msi}{h}}$  the general solution of (7a) reads

$$\begin{aligned} \varphi(x, s) = & \sqrt{\frac{m}{2is\hbar}} \left\{ u_1(x, s) \int_0^x u_2(y, s) \psi(y, 0) dy - u_2(x, s) \int_0^x u_1(y, s) \psi(y, 0) dy \right\} \\ & + \alpha(s) u_1(x, s) + \beta(s) u_2(x, s) \quad . \end{aligned} \quad (8)$$

Solving the equations given by the boundary conditions (7b) we find for  $\alpha(s)$  and  $\beta(s)$

$$\begin{aligned} \alpha(s) = & \sqrt{\frac{m}{2is\hbar}} \left\{ \frac{u_1(d, s)}{u_2(d, s) - u_1(d, s)} \int_0^d u_2(y, s) \psi(y, 0) dy \right. \\ & \left. - \frac{u_2(d, s)}{u_2(d, s) - u_1(d, s)} \int_0^d u_1(y, s) \psi(y, 0) dy \right\} = \\ & - \sqrt{\frac{m}{2is\hbar}} \int_0^d u_1(y, s) \psi(y, 0) dy + \\ & \sqrt{\frac{m}{2is\hbar}} \left\{ \frac{u_1(d, s)}{u_2(d, s) - u_1(d, s)} \left[ \int_0^d u_2(y, s) \psi(y, 0) dy - \int_0^d u_1(y, s) \psi(y, 0) dy \right] \right\} \\ \beta(s) = & -\alpha(s) \quad . \end{aligned}$$

Inserting the result into (8) and rewriting the integrals so that each term goes to zero for  $s \rightarrow \infty$  (which is a necessary condition for the successful application of the inverse Laplace transform), we get

$$\begin{aligned} \varphi(x, s) = & \frac{\kappa}{\sqrt{2is}} \left\{ \int_0^x e^{(i-1)\kappa\sqrt{s}(x-y)} \psi(y, 0) dy + \int_x^d e^{(i-1)\kappa\sqrt{s}(y-x)} \psi(y, 0) dy \right. \\ & \left. - \int_0^d e^{(i-1)\kappa\sqrt{s}(x+y)} \psi(y, 0) dy \right\} + \frac{\kappa}{\sqrt{2i}} s w(s) \int_0^d \left\{ e^{(i-1)\kappa\sqrt{s}(2d+x+y)} \right. \\ & \left. - e^{(i-1)\kappa\sqrt{s}(2d+x-y)} + e^{(i-1)\kappa\sqrt{s}(2d-x-y)} - e^{(i-1)\kappa\sqrt{s}(2d-x+y)} \right\} \psi(y, 0) dy \quad (9) \end{aligned}$$

where we have used the abbreviation

$$\kappa = \sqrt{\frac{m}{h}} \quad (10)$$

and  $w(s)$  stands for

$$w(s) = \frac{1}{s^{\frac{3}{2}}(e^{2(i-1)\kappa d\sqrt{s}} - 1)} .$$

Writing  $w(s)$  as a geometric series

$$w(s) = \frac{1}{s^{\frac{3}{2}}(e^{2(i-1)\kappa d\sqrt{s}} - 1)} = -\frac{1}{s^{\frac{3}{2}}} \sum_{k=0}^{\infty} e^{2(i-1)d\kappa\sqrt{s}k} \quad (11)$$

and using the inverse Laplace transform [9]

$$\mathcal{L}^{-1} \left\{ \frac{1}{\sqrt{s}} e^{-(1-i)\sqrt{as}} \right\} = \frac{e^{\frac{ia}{2t}}}{\sqrt{\pi t}} \quad \text{where } a > 0 \quad (12)$$

we find for the solution of (6a) the so-called mirror solution ([7])

$$\begin{aligned} \psi(x, t) = & \frac{\kappa}{\sqrt{2\pi t}i} \left( \int_0^d \left\{ e^{\frac{i(x-y)^2\kappa^2}{2t}} - e^{\frac{i(x+y)^2\kappa^2}{2t}} \right\} \psi(y, 0) dy \right. \\ & + \sum_{k=1}^{\infty} \int_0^d \left\{ e^{\frac{i\kappa^2(2dk+x-y)^2}{2t}} - e^{\frac{i\kappa^2(2dk+x+y)^2}{2t}} \right. \\ & \left. \left. + e^{\frac{i\kappa^2(2dk-x+y)^2}{2t}} - e^{\frac{i\kappa^2(2dk-x-y)^2}{2t}} \right\} \psi(y, 0) dy \right) . \quad (13) \end{aligned}$$

At times much smaller than the revival time ( $T_{rev} = 4md^2/\hbar$ ) of the infinite square well, a given wave packet is reflected between the walls with the classical period corresponding to its momentum expectation value. At a certain period of motion, a certain term of (13) is dominant. The first term is just the free particle propagator. When the wave packet, initially located between the walls, leaves the box it will no longer yield a relevant contribution to the solution. The second term represents the free dynamics of a wave packet, that has the initial shape  $\psi(-x, 0)$ , since

$$\int_0^d e^{\frac{i(x+y)^2\kappa^2}{2t}} \psi(y, 0) dy = \int_{-d}^0 e^{\frac{i(x-y)^2\kappa^2}{2t}} \psi(-y, 0) dy . \quad (14)$$

Furthermore, if  $\psi(x, 0)$  has the momentum expectation value  $p_0$ , the mirrored wavefunction  $\psi(-x, 0)$  has just the opposite momentum  $-p_0$ . Therefore, if the first term describes a wave packet moving towards the left wall, the second describes a wave packet initially located left outside the box and entering it around the time when the classical reflection takes place. So the first two terms of (13) are the quantum version of the reflection of a classical particle at the left wall. Similarly, each term can be given a certain meaning in terms of classical dynamics. If the initial wave packet moves towards the right wall, the term (14) becomes negligible, and another term of (13) will be responsible for the

description of the first reflection at the right wall. For each of the subsequent reflections there will be two relevant terms describing the reflection process. These terms coincide with the exact solution for a wave packet reflected at a single infinite wall which was already investigated in detail ([22],[23],[24]). As time progresses, of course the spreading of wave packets increases and the semiclassical picture is no longer correct.

## 4 The potential step

Our next model system is the potential step, described by the potential

$$\begin{aligned} V(x) &= 0 & \text{for } x < 0 \\ V(x) &= V & \text{for } x \geq 0 \end{aligned} .$$

The evolution of the wave packet  $\psi(x, 0)$ , is determined by

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} = i\hbar \frac{\partial \psi(x, t)}{\partial t} \quad \text{for } x < 0 \quad (15)$$

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} + V\psi(x, t) = i\hbar \frac{\partial \psi(x, t)}{\partial t} \quad \text{for } x \geq 0, \quad (16)$$

where  $\psi(x, t)$  is supposed to be continuously differentiable. The solution will be square integrable for all times, if the initial function is chosen square integrable. We will from now on only consider initial wave packets that are located left of the potential step

$$\psi(x, 0) = 0 \quad \text{for } x \geq 0.$$

According to our method, we find for the Laplace transformed wave packet  $\varphi(x, s)$

$$\begin{aligned} \varphi(x, s) &= \sqrt{\frac{m}{2is\hbar}} \left\{ u_1(x, s) \int_{-\infty}^x u_2(y, s) \psi(y, 0) dy - u_2(x, s) \int_{-\infty}^x u_1(y, s) \psi(y, 0) dy \right\} \\ &+ \alpha(s) u_1(x, s) + \beta(s) u_2(x, s) \quad \text{for } x < 0 \end{aligned} \quad (17a)$$

$$\varphi(x, s) = \gamma(s) u_3(x, s) + \delta(s) u_4(x, s) \quad \text{for } x > 0, \quad (17b)$$

where

$$u_1(x, s) = e^{i\sqrt{\frac{2msi}{\hbar}}x} \quad u_3(x, s) = e^{i\sqrt{\frac{2msi}{\hbar} - \frac{2mV}{\hbar^2}}x} \quad (18a)$$

$$u_2(x, s) = e^{-i\sqrt{\frac{2msi}{\hbar}}x} \quad u_4(x, s) = e^{-i\sqrt{\frac{2msi}{\hbar} - \frac{2mV}{\hbar^2}}x}. \quad (18b)$$

Since  $\psi(x, t)$  must be square integrable,  $\varphi(x, s)$  must vanish for  $x \rightarrow \pm\infty$ . We get

$$\alpha(s) = \delta(s) = 0.$$

Furthermore  $\varphi(x, s)$  should be continuously differentiable at  $x = 0$  which determines the functions  $\beta(s)$  and  $\gamma(s)$  :

$$\gamma(s) = \sqrt{\frac{m}{2si\hbar}} \int_{-\infty}^0 u_2(y, s) \psi(y, 0) dy \cdot \frac{2}{1 + \sqrt{1 - \frac{V}{\hbar si}}} \quad (19a)$$

$$\begin{aligned} \beta(s) &= \sqrt{\frac{m}{2si\hbar}} \int_{-\infty}^0 u_2(y, s) \psi(y, 0) dy \cdot \left( \frac{2}{1 + \sqrt{1 - \frac{V}{\hbar si}}} - 1 \right) \\ &+ \sqrt{\frac{m}{2si\hbar}} \int_{-\infty}^0 u_1(y, s) \psi(y, 0) dy. \end{aligned} \quad (19b)$$

This yields for the Laplace transformed wave packet  $\varphi(x, s)$  (17)

$$\begin{aligned} \varphi(x, s) &= \sqrt{\frac{m}{2si\hbar}} \left\{ \int_{-\infty}^0 e^{i\sqrt{\frac{2msi}{\hbar}}|x-y|} \psi(y, 0) dy + \int_{-\infty}^0 e^{-i\sqrt{\frac{2msi}{\hbar}}(x+y)} \psi(y, 0) dy \cdot \rho(s) \right\} \\ &\text{for } x < 0 \end{aligned} \quad (20a)$$

$$\begin{aligned} \varphi(x, s) &= \sqrt{\frac{m}{2si\hbar}} \int_{-\infty}^0 e^{i\sqrt{\frac{2msi}{\hbar} - \frac{2mV}{\hbar^2}}x} e^{-i\sqrt{\frac{2msi}{\hbar}}y} \psi(y, 0) dy \cdot (\rho(s) + 1) \\ &\text{for } x > 0, \end{aligned} \quad (20b)$$

where

$$\rho(s) = \frac{2}{1 + \sqrt{1 - \frac{V}{\hbar si}}} - 1. \quad (21)$$

#### 4.1 Solution for $x < 0$

The inverse Laplace transform of (21) is [9]

$$\mathcal{L}^{-1} \{ \rho(s) \} = r(t) = \frac{1}{ti} J_1 \left[ \frac{Vt}{2\hbar} \right] e^{-\frac{iVt}{2\hbar}}, \quad (22)$$

where  $J_1(x)$  denotes the Bessel function. Since the inverse Laplace transform of a product is given by a convolution [8]

$$\mathcal{L}^{-1} \{ f(s) \cdot g(s) \} = \int_0^t F(t - \tau) G(\tau) d\tau, \quad (23)$$

$$\text{where } \mathcal{L}^{-1} \{ f(s) \} = F(t), \mathcal{L}^{-1} \{ g(s) \} = G(t),$$

we find for the inverse Laplace transform of the wave packet (20a) in the region  $x < 0$

$$\psi(x, t) = \frac{\kappa}{\sqrt{2\pi t} i} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \psi(y, 0) dy + \int_0^t \frac{\kappa}{\sqrt{2\pi(t-\tau)} i} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2(t-\tau)}} \psi(y, 0) r(\tau) dy d\tau, \quad (24)$$

where we have applied (12,10). The first term describes the free time evolution of the wave packet  $\psi(x, 0)$ . The second term represents the reflected wave packet and has the form of a wavepacket reflected at an infinite wall (see also the description at the end of the last section), deformed by a convolution.

If we use the momentum representation

$$f(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \psi(x, 0) e^{-\frac{ipx}{\hbar}} dx, \quad (25)$$

we can rewrite the convolution integral in (24) as

$$\int_0^t \frac{\kappa}{\sqrt{2\pi(t-\tau)} i} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2(t-\tau)}} \psi(y, 0) dy r(\tau) d\tau = \quad (26)$$

$$\int_0^t \frac{\kappa}{\sqrt{2\pi(t-\tau)} i} \int_{-\infty}^{\infty} e^{\frac{i(x+y)^2 \kappa^2}{2(t-\tau)}} \psi(y, 0) dy r(\tau) d\tau = \quad (27)$$

$$\int_0^t \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{-\frac{ipx}{\hbar}} e^{-\frac{ip^2(t-\tau)}{2m\hbar}} f(p) dp r(\tau) d\tau = \quad (28)$$

$$\frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{-\frac{ipx}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) R(p, t) dp, \quad (29)$$

where

$$R(p, t) = \int_0^t e^{\frac{ip^2 \tau}{2m\hbar} - \frac{iV\tau}{2\hbar}} J_1 \left[ \frac{V\tau}{2\hbar} \right] \frac{d\tau}{\tau i}.$$

We will approximate this function by

$$R(p, t) \approx R(p) = \int_0^{\infty} e^{\frac{ip^2 \tau}{2m\hbar} - \frac{iV\tau}{2\hbar}} J_1 \left[ \frac{V\tau}{2\hbar} \right] \frac{d\tau}{\tau i}. \quad (30)$$

Since the Bessel function fulfills

$$\left| J_1 \left[ \frac{Vt}{2\hbar} \right] \frac{1}{t} \right| \leq \frac{2\hbar}{V} \frac{1}{t^{3/2}}$$

the additional part of the integral

$$\left| \int_t^{\infty} e^{\frac{ip^2 \tau}{2m\hbar} - \frac{iV\tau}{2\hbar}} J_1 \left[ \frac{V\tau}{2\hbar} \right] \frac{d\tau}{\tau i} \right| \leq \int_t^{\infty} \sqrt{\frac{2\hbar}{V}} \frac{1}{\tau^{3/2}} d\tau = 2\sqrt{\frac{2\hbar}{Vt}}$$

does hardly contribute, if

$$t \gg \frac{\hbar}{V},$$

and can therefore be neglected after a very short time. Moreover, if the initial function is reasonably peaked the second term of (24) does not become relevant until the wave packet approaches the barrier. So the approximation can always be used when

$$t_R \gg \frac{\hbar}{V},$$

where  $t_R$  is the time, a classical particle corresponding to the wave packet would need to reach the barrier.

The solution for the region  $x < 0$ , then reads

$$\begin{aligned} \psi(x, t) \approx & \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \psi(y, 0) dy + \\ & \frac{1}{\sqrt{2\pi \hbar}} \int_{-\infty}^{\infty} e^{-\frac{i p x}{\hbar}} e^{-\frac{i p^2 t}{2m \hbar}} f(p) R(p) dp. \end{aligned} \quad (31)$$

Since the integral (30) is itself a Laplace integral, we can apply (22) and find for  $R(p)$

$$R(p) = \rho \left[ -\frac{i p^2}{2m \hbar} \right] = -1 + 2k - 2\sqrt{k(k-1)} \quad \text{with} \quad k = \frac{p^2}{2mV}. \quad (32)$$

This function coincides with the reflection coefficient of the time independent solution (see for instance [25]).

Defining the reflection probability as

$$\mathcal{R}(t) = \int_{-\infty}^0 |\Psi(x, t)|^2 dx,$$

we are now able to show that

$$\lim_{t \rightarrow \infty} \mathcal{R}(t) \leq \int_{-\infty}^{\infty} |f(p) R(p)|^2 dp, \quad (33)$$

if the momentum expectation value  $p_0$  of the initial wavepacket is positive.

The first term of (31) describes the free time evolution of the wavepacket  $\Psi_F(x, t)$ . For positive momentum expectation values we find

$$\lim_{t \rightarrow \infty} \int_{-\infty}^0 |\Psi_F(x, t)|^2 dx = 0. \quad (34)$$

This result can be derived by determining the correction that has to be applied to a free wave packet solution with  $p_0 = 0$  if the expectation value changes to  $p_0 > 0$ . So only the second term of (31) which describes the reflected wave packet contributes

$$\lim_{t \rightarrow \infty} \mathcal{R}(t) = \lim_{t \rightarrow \text{inf ty}} \int_{-\infty}^0 |\Psi_R(x, t)|^2 dx,$$

where

$$\Psi_R(x, t) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{-\frac{ipx}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) R(p) dp = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{ipx}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(-p) R(p) dp.$$

This is the free time evolution of an initial wave packet  $R(p)f(-p)$  which yields

$$\lim_{t \rightarrow \infty} \mathcal{R}(t) \leq \int_{-\infty}^{\infty} |\Psi_R(x, t)|^2 dx = \int_{-\infty}^{\infty} |f(p)R(p)|^2 dp.$$

If in particular the distortion by  $R(p)$  is small enough so that

$$\int_{-\infty}^{\infty} p |f(-p)R(p)|^2 dp < 0, \quad (35a)$$

we find, arguing as for (34),

$$\lim_{t \rightarrow \infty} \int_{-\infty}^0 |\Psi_R(x, t)|^2 dx = \int_{-\infty}^{\infty} |\Psi_R(x, t)|^2 dx, \quad (35b)$$

which means that the inequality in (33) is saturated. So the expression for the reflection probability that was conjectured in [20] turns out to apply only under the condition (35a) for the initial wave function. Note that (33,35b) are exact results since (31) tends to the exact solution (24) for  $t \rightarrow \infty$ .

## 4.2 Solution for $x > 0$

In order to determine the inverse Laplace transform of (20b) we use the momentum representation of  $\psi(x, 0)$ . This yields for the integral

$$\sqrt{\frac{m}{2si\hbar}} \int_{-\infty}^0 e^{i\sqrt{\frac{2m si}{\hbar} - \frac{2mV}{\hbar^2}} x} e^{-i\sqrt{\frac{2m si}{\hbar}} y} \psi(y, 0) dy = \quad (36)$$

$$\frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{i\sqrt{\frac{2m si}{\hbar} - \frac{2mV}{\hbar^2}} x} \sqrt{\frac{m}{2si\hbar}} \int_{-\infty}^{\infty} e^{\frac{ipy}{\hbar}} e^{i\sqrt{\frac{2m si}{\hbar}} |y|} dy f(p) dp = \quad (37)$$

$$\frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{i\sqrt{\frac{2m si}{\hbar} - \frac{2mV}{\hbar^2}} x} \frac{1}{s + \frac{ip^2}{2m\hbar}} f(p) dp. \quad (38)$$

According to [9]

$$\begin{aligned} & \mathcal{L}^{-1} \left\{ 2(s + b + c)^{-1} e^{-\sqrt{a(s+c)}} \right\} = \\ & e^{-ct - bt} \left\{ e^{-i\sqrt{ab}} \operatorname{Erfc} \left[ \sqrt{\frac{a}{4t}} - i\sqrt{bt} \right] + e^{i\sqrt{ab}} \operatorname{Erfc} \left[ \sqrt{\frac{a}{4t}} + i\sqrt{bt} \right] \right\} \\ & \text{where } \operatorname{Re}[a] \geq 0, \end{aligned} \quad (39)$$

and  $\text{Erfc}(x)$  is the entire function

$$\text{Erfc}(x) = \frac{2}{\sqrt{\pi}} \int_x^\infty e^{-u^2} du.$$

We find with

$$a = -\frac{2mi}{\hbar}x^2, \quad b = \frac{i}{2m\hbar}(p^2 - 2mV), \quad c = \frac{iV}{\hbar}$$

for the inverse Laplace transform of the wave packet (20b) in the region  $x > 0$

$$\psi(x, t) = \int_{-\infty}^{\infty} K(x, p, t) f(p) dp + \int_0^t \int_{-\infty}^{\infty} K(x, p, t - \tau) f(p) dp r(\tau) d\tau, \quad (40a)$$

with

$$K(x, p, t) = \frac{1}{2\sqrt{2\pi\hbar}} e^{-\frac{iVt}{\hbar} - \frac{iZ^2}{2\hbar m}} \cdot \left\{ e^{-\frac{i\pi Z}{\hbar}} \text{Erfc} \left[ -i\sqrt{\frac{2mi}{\hbar t}} \frac{x}{2} - i\sqrt{\frac{it}{2\hbar m}} Z \right] + e^{\frac{i\pi Z}{\hbar}} \text{Erfc} \left[ -i\sqrt{\frac{2mi}{\hbar t}} \frac{x}{2} + i\sqrt{\frac{it}{2\hbar m}} Z \right] \right\}, \quad (40b)$$

where we have again used the convolution theorem (23) and we have introduced the abbreviation

$$Z = \sqrt{p^2 - 2mV}.$$

A further interpretation of (40) is only possible if we distinguish between momentum distributions that are concentrated around  $p_0 > \sqrt{2mV}$ , where a classical transmission of the barrier would be possible and those which are concentrated around  $p_0 < \sqrt{2mV}$ , where the wave packet enters a classically forbidden region.

### 4.3 A wave packet climbing the potential step

We can write any initial wave packet  $\psi(x, 0)$  in momentum space (25) as

$$f(p) = e^{-\frac{ipx_0}{\hbar}} F(p - p_0), \quad (41a)$$

where  $F(p)$  fulfills

$$\int_{-\infty}^{\infty} F^*(p) p F(p) dp = 0, \quad \int_{-\infty}^{\infty} F^*(p) F'(p) dp = 0. \quad (41b)$$

The expectation values of  $\psi(x, 0)$  are then given by

$$\langle \hat{x} \rangle = x_0, \quad \langle \hat{p} \rangle = p_0,$$

and the uncertainties will be denoted by  $\Delta x_0$  and  $\Delta p_0$ .

We will now consider an initial function  $f(p)$  (25) almost exclusively concentrated in a region  $p > p_m > \sqrt{2mV}$ , so that we can assume

$$f(p) \approx 0 \quad \text{for } p < p_m, \quad (42)$$

Moreover we require the wave packet to be sufficiently peaked around the momentum expectation value  $p_0$ , so that

$$\frac{p_0}{mV} F(p - p_0)(p - p_0) \approx 0, \quad (43a)$$

$$e^{-\frac{i\sqrt{q^2+2mV}x_0}{\hbar}} F\left(\sqrt{q^2+2mV} - \sqrt{q_0^2+2mV}\right) \approx \quad (43b)$$

$$F(\lambda(q - q_0)) e^{-\frac{iq_0x_0}{\lambda\hbar} - \frac{ix_0\lambda}{\hbar}(q - q_0) - \frac{ix_0\lambda}{2\hbar q_0\kappa_0}(q - q_0)^2},$$

where

$$p_0 = \sqrt{q_0^2 + 2mV}, \quad \lambda = \sqrt{1 - \frac{2mV}{p_0^2}}$$

Finally the difference  $p_m - \sqrt{2mV}$  should be big enough to ensure

$$\frac{1}{\sqrt{\frac{p^2}{2mV} - 1}} = O(1) \quad \text{for } p \geq p_m \quad (44)$$

which means that this factor does not have relevant influence on the approximations.

If we now insert the wave packet into the solution for  $x < 0$  (31) we will use  $R(p) \approx R(p_0)$  since according to (43a) and the mean value theorem

$$f(p)R(p) = f(p)R(p_0) + f(p)R'(p_1)(p - p_0) \approx f(p)R(p_0), \quad \text{with } p_1 \in (p_0, p)$$

where (44) ensures that the derivative of  $R(p)$  (32) will not spoil the approximation. We find for  $x < 0$

$$\begin{aligned} \psi(x, t) &\approx \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \psi(y, 0) dy + \frac{R(p_0)}{\sqrt{2\pi \hbar}} \int_{-\infty}^{\infty} e^{-\frac{ipx}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) dp = \\ &\frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \psi(y, 0) dy + R(p_0) \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2t}} \psi(y, 0) dy. \end{aligned} \quad (45)$$

The solution consists of an incoming and a reflected wave packet. Within our approximation the reflected wave packet has the same shape as the wave packet reflected by an infinite barrier (compare 14), but its probability density

is reduced by a factor  $R(p_0)^2$ . This factor only depends on  $k_0 = p_0^2/(2mV)$  and goes from 1 to zero when  $k_0$  goes from 1 to infinity (32). If  $p_0^2/2m$ , which corresponds to the energy expectation value within our approximation, is much higher than the potential step, there is hardly any reflection at the step.

For the interpretation of the solution for  $x > 0$  (40) we rewrite the error function according to [21]

$$\operatorname{Erfc}\left[\frac{u}{2z}\right] = \frac{2z}{\sqrt{\pi}} e^{-\frac{u^2}{4z^2}} \int_0^\infty e^{-z^2 y^2 - uy} dy, \quad (46)$$

where we choose

$$z = \sqrt{\frac{-im}{2\hbar t}}, \quad u = -\frac{im}{\hbar t} x \mp \frac{i}{\hbar} Z.$$

This yields

$$K(x, p, t) = e^{-\frac{iVt}{\hbar}} \left\{ \frac{\kappa}{\sqrt{2\pi t i}} \frac{1}{\sqrt{2\pi\hbar}} \int_0^\infty e^{\frac{i(x+y)^2 \kappa^2}{2t}} e^{\frac{iZy}{\hbar}} dy + \frac{\kappa}{\sqrt{2\pi t i}} \frac{1}{\sqrt{2\pi\hbar}} \int_0^\infty e^{\frac{i(x+y)^2 \kappa^2}{2t}} e^{-\frac{iZy}{\hbar}} dy \right\}.$$

Applying (43a,43b,44) we find (see Appendix A)

$$\tilde{\psi}(y, 0) := \frac{1}{\sqrt{2\pi\hbar}} \int_{\sqrt{2mV}}^\infty e^{\frac{iZy}{\hbar}} f(p) dp = \frac{1}{\sqrt{2\pi\hbar}} \int_{\sqrt{2mV}}^\infty e^{\frac{i\sqrt{p^2 - 2mV}y}{\hbar}} f(p) dp \approx \quad (47a)$$

$$\approx \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^\infty e^{\frac{iqy}{\hbar}} \lambda e^{-\frac{iq_0 x_0}{\lambda\hbar} - \frac{ix_0 \lambda}{\hbar}(q - q_0) - \frac{ix_0 \lambda}{2\hbar q_0 k_0}(q - q_0)^2} F(\lambda(q - q_0)) dq. \quad (47b)$$

This function is a deformed version of the original wave packet  $\psi(x, 0)$ , characterized by the following properties:

$$\begin{aligned} \langle \tilde{\psi} | \tilde{\psi} \rangle &= \lambda, & \frac{1}{\lambda} \langle \tilde{\psi} | \hat{p} | \tilde{\psi} \rangle &= q_0, & \frac{1}{\lambda} \langle \tilde{\psi} | \hat{x} | \tilde{\psi} \rangle &= \lambda x_0, \\ \Delta x^2 &= \lambda^2 \Delta x_0^2 + \frac{x_0^2}{q_0^2 k_0^2} \Delta p_0^2, & \Delta p^2 &= \frac{1}{\lambda^2} \Delta p_0^2. \end{aligned}$$

$\psi(x, 0)$  was assumed to be zero for  $x > 0$ . If this function is sufficiently peaked around  $x = x_0 < 0$  and  $p_0$ , then also  $\Delta x$  will be sufficiently small so that the deformed wave function will also vanish for  $x > 0$ , and

$$\int_{-\infty}^\infty \frac{\kappa}{\sqrt{2\pi t i}} \frac{1}{\sqrt{2\pi\hbar}} \int_0^\infty e^{\frac{i(x+y)^2 \kappa^2}{2t}} e^{\frac{iZy}{\hbar}} dy f(p) dp = \quad (48a)$$

$$\frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2t}} \tilde{\psi}(y, 0) dy \approx 0. \quad (48b)$$

Therefore we find

$$\int_{-\infty}^\infty K(x, p, t) f(p) dp \approx \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \tilde{\psi}(y, 0) dy.$$

Inserting this result in (40a), applying (30) for the convolution integral and proceeding further as for  $x < 0$ , we get for  $x > 0$

$$\psi(x, t) \approx (1 + R(p_0)) \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \tilde{\psi}(y, 0) dy. \quad (49)$$

This means that the wave function is deformed after it has passed the potential step. It takes now the form of a free particle wave function that looked like  $\tilde{\psi}(x, 0)$  at  $t = 0$ . Moreover it is slowed down, and has now the momentum expectation value  $\sqrt{p_0^2 - 2mV}$  instead of  $p_0$ , which exactly coincides with the reduced momentum of a corresponding classical particle (see figure 1(c)).

When the reflection/transmission process is finished

$$t \gg \frac{x_0 m}{p_0},$$

only the second term of (45), describing the reflected wave function is relevant and the solution consists of a reflected and a transmitted wave packet.

$$\begin{aligned} \psi(x, t) &\approx \psi_R(x, t) + \psi_T(x, t) = & (50) \\ &R(p_0) \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2t}} \psi(y, 0) dy + (1 + R(p_0)) \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \tilde{\psi}(y, 0) dy & (51) \end{aligned}$$

Here we do not need to restrict  $\psi_R$  and  $\psi_T$  to the regions left and right of  $x = 0$  anymore, since they will be concentrated in the respective regions anyway. The reflected and the transmitted part then fulfill

$$\langle \psi_R | \psi_R \rangle = (R(p_0))^2, \quad \langle \psi_T | \psi_T \rangle = (R(p_0) + 1)^2 \lambda = 1 - \langle \psi_R | \psi_R \rangle$$

The reflection probability  $|R(p_0)|^2$  is then an approximation for (35b) for a sufficiently peaked wave packet.


#### 4.4 A wave packet reflected by the potential step and swapping into the forbidden region

We now consider a wave packet represented by (41), and concentrated within a region  $0 < p < p_m < \sqrt{2mV}$ , so that we can assume


$$f(p) \approx 0 \quad \text{for } p > p_m \quad \text{and } p < 0, \quad (52)$$

and


$$\frac{1}{\sqrt{1 - \frac{p^2}{2mV}}} = O(1) \quad \text{for } 0 \leq p \leq p_m. \quad (53)$$



(a) The relative dimensionless probability density  $|\Psi|^2 \cdot (\pi\alpha)^{1/2}$  of the initial wave packet.



(b) The relative dimensionless probability density  $|\Psi|^2 \cdot (\pi\alpha)^{1/2}$  at  $t_R = |x_0|m/p_0$ .



(c) The relative dimensionless probability density  $|\Psi|^2 \cdot (\pi\alpha)^{1/2}$  at  $t = 2t_R$ . The momentum is reduced by a factor  $\sqrt{\frac{k_0-1}{k_0}} = 0.577$ .

Figure 1: As initial function we choose a Gaussian wave packet with the properties  $\Delta x_0 = \sqrt{\alpha/2}$ ,  $\Delta p_0 = \hbar/\sqrt{2\alpha}$ ,  $x_0 = -10\sqrt{\alpha}$ ,  $p_0 = 100\hbar/\sqrt{\alpha}$ . The contributions of this Gaussian in the region  $x \geq 0$  are so small that we can use (45,49) and we can extend the integrals to the whole real line. We show the wave packet before, during and after the reflection/transmission process that takes place around the classical reflection time  $t_R$ . We have chosen  $k_0 = 1.5$ .

We also require (43a). Proceeding as in the case  $p_0 > \sqrt{2mV}$ , we get the same result (45) for  $x < 0$

$$\psi(x, t) \approx \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \psi(y, 0) dy + R(p_0) \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2t}} \psi(y, 0) dy. \quad (54)$$

According to (32),  $R(p_0)$  is a complex function with absolute value 1 for  $k_0 = \frac{p_0^2}{2m} < 1$ . For  $k_0 \ll 1$ ,  $R(p_0)$  is approximately -1 and  $\psi(x, t)$  is completely reflected at the barrier.

Applying

$$\text{Erfc}[x] = 2 - \text{Erfc}[-x]$$

to the second term in (40b), and using (46) with

$$z = \sqrt{\frac{-im}{2ht}}, \quad u = \mp \frac{im}{ht} x - \frac{i}{h} Z,$$

we get

$$K(x, p, t) = e^{-\frac{iVt}{h}} \left\{ \frac{1}{\sqrt{2\pi h}} e^{-\frac{itZ^2}{2hm} + \frac{ixZ}{h}} + \frac{\kappa}{\sqrt{2\pi t i}} \frac{1}{\sqrt{2\pi h}} \int_0^\infty e^{\frac{i(x+y)^2 \kappa^2}{2t}} e^{\frac{iZy}{h}} dy - \frac{\kappa}{\sqrt{2\pi t i}} \frac{1}{\sqrt{2\pi h}} \int_0^\infty e^{\frac{i(x-y)^2 \kappa^2}{2t}} e^{\frac{iZy}{h}} dy \right\}.$$

With the help of the mean value theorem and (43a),(53) we determine

$$e^{\frac{iZy}{h}} f(p) = e^{-\frac{\sqrt{2mV-p^2}y}{h}} f(p) = e^{-\frac{\sqrt{2mV-p_0^2}y}{h}} f(p) - e^{-\frac{\sqrt{2mV-p_1^2}y}{h}} \frac{p_1 y}{h\sqrt{2mV-p_1^2}} (p-p_0) f(p) \\ \approx e^{-\frac{\sqrt{2mV-p_0^2}y}{h}} f(p) \text{ with } p_1 \in (p_0, p).$$

Since according to (52)

$$\int_0^{\sqrt{2mV}} f(p) dp \approx \int_{-\infty}^\infty f(p) dp = \psi(0, 0) = 0,$$

we conclude

$$\int_{-\infty}^\infty K(x, p, t) f(p) dp \approx e^{-\frac{\sqrt{2mV-p_0^2}x}{h}} \frac{1}{\sqrt{2\pi h}} \int_{-\infty}^\infty e^{\frac{-ip^2 t}{2m h}} f(p) dp.$$

Proceeding as for (49), we finally determine the solution for  $x > 0$

$$\psi(x, t) \approx (1 + R(p_0)) e^{-\frac{\sqrt{2mV-p_0^2}x}{h}} \frac{1}{\sqrt{2\pi h}} \int_{-\infty}^\infty e^{\frac{-ip^2 t}{2m h}} f(p) dp. \quad (55)$$

We can again conclude that after the reflection process, when

$$t \gg \frac{x_0 m}{p_0},$$

the wavefunction only consists of a reflected and a transmitted part.

$$\psi(x, t) \approx \psi_R(x, t) + \psi_T(x, t) = \tag{56}$$

$$R(p_0) \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2t}} \psi(y, 0) dy + (1 + R(p_0)) e^{-\frac{\sqrt{2mV - p_0^2} x}{\hbar}} \frac{1}{\sqrt{2\pi \hbar}} \int_{-\infty}^{\infty} e^{\frac{-ip^2 t}{\hbar}} f(p) dp. \tag{57}$$

But since  $|R(p_0)| = 1$ , we find

$$\langle \psi_R | \psi_R \rangle = \int_{-\infty}^0 \psi^*(x, t) \psi(x, t) dx \approx \int_{-\infty}^{\infty} \psi^*(x, t) \psi(x, t) dx = 1 \quad ,$$

which means that when the reflected wave packet has left the neighborhood of the step the rest of the wave function also leaves the classically forbidden region (see figure 2(c))

$$\psi(x, t) \approx \psi_R(x, t) \quad \text{for} \quad t \gg \frac{x_0 m}{p_0} .$$

## 5 The asymmetric square well

We will consider a potential that is a combination of the infinite square well and the potential step, which can be described as a box with exit.

$$\begin{aligned} V(x) &= \infty & \text{for} & \quad x \leq -d \\ V(x) &= 0 & \text{for} & \quad -d < x < 0 \\ V(x) &= V & \text{for} & \quad x \geq 0 \quad . \end{aligned}$$

The dynamics of the wave packet  $\psi(x, 0)$ , is then governed by

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} = i\hbar \frac{\partial \psi(x, t)}{\partial t} \quad \text{for} \quad -d \leq x < 0 \tag{58}$$

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} + V\psi(x, t) = i\hbar \frac{\partial \psi(x, t)}{\partial t} \quad \text{for} \quad x \geq 0, \tag{59}$$


where  $\psi(x, t)$  is supposed to be continuously differentiable and square integrable and to fulfill the boundary condition

$$\psi(-d, t) = 0. \tag{60}$$


We will further assume that the initial wave packets are located within the box,

$$\psi(x, 0) = 0 \quad \text{for} \quad x \geq 0. \tag{61}$$


We find for the Laplace transformed wave packet  $\varphi(x, s)$



(a) The relative dimensionless probability density  $|\Psi|^2 \cdot (\pi\alpha)^{1/2}$  of the initial wave packet.



(b) The relative dimensionless density  $|\Psi|^2 \cdot (\pi\alpha)^{1/2}$  at  $t_R = |x_0|m/p_0$ . Note that the exponential decay in the classically forbidden region is determined by the factor  $e^{-\sqrt{1/k_0-1}p_0x/\hbar}$



(c) The relative dimensionless probability density  $|\Psi|^2 \cdot (\pi\alpha)^{1/2}$  at  $t = 2t_R$ . The whole wave function is reflected. There is no part of the probability density left in the classically forbidden region.

Figure 2: As initial function we choose a Gaussian wave packet with the properties  $\Delta x_0 = \sqrt{\alpha/2}$ ,  $\Delta p_0 = \hbar/\sqrt{\alpha 2}$ ,  $x_0^{19} = -10\sqrt{\alpha}$ ,  $p_0 = 10\hbar/\sqrt{\alpha}$ . The contributions of this function for  $x \geq 0$  can again be neglected. We see that the wave packet swaps into the region  $x > 0$  around the classical reflection time, but leaves it entirely when the reflection process is over. We have chosen  $k_0 = 1/4$ .

$$\begin{aligned} \varphi(x, s) = \sqrt{\frac{m}{2is\hbar}} \left\{ u_1(x, s) \int_{-d}^x u_2(y, s) \psi(y, 0) dy - u_2(x, s) \int_{-d}^x u_1(y, s) \psi(y, 0) dy \right\} \\ + \alpha(s) u_1(x, s) + \beta(s) u_2(x, s) \end{aligned} \quad \text{for } x < 0 \quad (62a)$$

$$\varphi(x, s) = \gamma(s) u_3(x, s) + \delta(s) u_4(x, s) \quad \text{for } x > 0 \quad , \quad (62b)$$

where the functions  $u_1, u_2, u_3, u_4$  are defined as in the previous section (18). Since  $\varphi(x, s)$  must vanish for  $x \rightarrow \infty$ , we find

$$\delta(s) = 0 .$$

The boundary condition (60) implies

$$\alpha(s) = -\beta(s) e^{(i-1)2d\kappa\sqrt{s}} .$$

Since  $\varphi(x, s)$  should be continuously differentiable at  $x = 0$ , we get

$$\begin{aligned} \gamma(s) &= \frac{\kappa}{\sqrt{2s}i} \cdot \frac{2I_2 - 2I_1 e^{(i-1)2d\kappa\sqrt{s}}}{1 + \sqrt{1 - \frac{V}{s\hbar}} + e^{(i-1)2d\kappa\sqrt{s}} \left(1 - \sqrt{1 - \frac{V}{s\hbar}}\right)} \quad (63) \\ \beta(s) &= \frac{\kappa}{\sqrt{2s}i} \cdot \frac{1}{1 - e^{(i-1)2d\kappa\sqrt{s}}} \left\{ I_1 - I_2 + \frac{2I_2 - 2I_1 e^{(i-1)2d\kappa\sqrt{s}}}{1 + \sqrt{1 - \frac{V}{s\hbar}} + e^{(i-1)2d\kappa\sqrt{s}} \left(1 - \sqrt{1 - \frac{V}{s\hbar}}\right)} \right\} , \quad (64) \end{aligned}$$

where we have introduced

$$I_1 = \int_{-d}^0 e^{(i-1)y\kappa\sqrt{s}} \psi(y, 0) dy, \quad I_2 = \int_{-d}^0 e^{-(i-1)y\kappa\sqrt{s}} \psi(y, 0) dy .$$

Inserting these results in 62, using the abbreviations (21,10) and applying

$$\frac{1 - \sqrt{1 - \frac{V}{s\hbar}}}{1 + \sqrt{1 - \frac{V}{s\hbar}}} = \rho(s)$$

as well as the series expansion

$$\frac{1}{(1 + e^{(i-1)\kappa 2d\sqrt{s}} \rho(s))} = \sum_{k=0}^{\infty} (-1)^k \rho(s)^k e^{(i-1)2d\kappa\sqrt{s}k} , \quad (65)$$

we find for the Laplace transformed wave packet (62),

$$\begin{aligned} \varphi(x, s) = & \left\{ \sum_{k=0}^{\infty} \frac{\kappa}{\sqrt{2si}} (\rho(s) + 1) (-1)^k \rho(s)^k \int_{-d}^0 e^{(i-1)(2dk-y)\kappa\sqrt{s}} \psi(y, 0) dy \right. \\ & \left. - \sum_{k=0}^{\infty} \frac{\kappa}{\sqrt{2si}} (\rho(s) + 1) (-1)^k \rho(s)^k \int_{-d}^0 e^{(i-1)(2d(k+1)+y)\kappa\sqrt{s}} \psi(y, 0) dy \right\} e^{i\sqrt{\frac{2msi}{\hbar} - \frac{2mV}{\hbar^2}} x} \\ & \text{for } x > 0. \end{aligned} \quad (66a)$$

Applying also

$$\begin{aligned} \frac{1}{1 + e^{(i-1)2d\kappa\sqrt{s}} \rho(s)} \cdot \frac{1}{1 - e^{(i-1)2d\kappa\sqrt{s}}} &= \frac{e^{-(i-1)2d\kappa\sqrt{s}}}{1 + \rho(s)} \left\{ \frac{1}{1 - e^{(i-1)2d\kappa\sqrt{s}}} - \frac{1}{1 + e^{(i-1)2d\kappa\sqrt{s}} \rho(s)} \right\} \\ &= \frac{1}{1 + \rho(s)} \sum_{k=0}^{\infty} (1 - (-1)^k \rho(s)^k) e^{(i-1)2d(k-1)\kappa\sqrt{s}}, \end{aligned}$$

we conclude

$$\begin{aligned} \varphi(x, s) = & - \sum_{k=0}^{\infty} \rho(s)^k (-1)^k \frac{\kappa}{\sqrt{2si}} \int_{-d}^0 e^{(i-1)(2d(k+1)+x+y)\kappa\sqrt{s}} \psi(y, 0) dy \\ & + \sum_{k=0}^{\infty} \rho(s)^{k+1} (-1)^{k+1} \frac{\kappa}{\sqrt{2si}} \int_{-d}^0 e^{(i-1)(2d(k+1)+x-y)\kappa\sqrt{s}} \psi(y, 0) dy \\ & + \frac{\kappa}{\sqrt{2si}} \int_{-d}^0 e^{(i-1)|x-y|\kappa\sqrt{s}} \psi(y, 0) dy \\ & + \sum_{k=0}^{\infty} \rho(s)^{k+1} (-1)^{k+1} \frac{\kappa}{\sqrt{2si}} \int_{-d}^0 e^{(i-1)(2d(k+1)-x+y)\kappa\sqrt{s}} \psi(y, 0) dy \\ & - \sum_{k=0}^{\infty} \rho(s)^{k+2} (-1)^{k+2} \frac{\kappa}{\sqrt{2si}} \int_{-d}^0 e^{(i-1)(2d(k+1)-x-y)\kappa\sqrt{s}} \psi(y, 0) dy \\ & + \frac{\kappa}{\sqrt{2si}} \rho(s) \int_{-d}^0 e^{(i-1)|x+y|\kappa\sqrt{s}} \psi(y, 0) dy \\ & \text{for } x < 0. \end{aligned} \quad (66b)$$

In order to determine the inverse Laplace transform of (66a), we will proceed similar as in section (4.2). We will use the shifted momentum representation

$$f(K, p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \psi(x + K, 0) e^{-\frac{ipx}{\hbar}} dx = e^{\frac{ipK}{\hbar}} f(p) \quad (67)$$

and we introduce

$$L(k, t) = \mathcal{L}^{-1} \{ (\rho(s) + 1) \rho(s)^k \}, \quad (68)$$

which can be expressed as a sequence of convolutions of (22).

This yields for the wave function in the region  $x > 0$

$$\begin{aligned}
\psi(x, t) = & \int_{-\infty}^{\infty} K(x, p, t) f(0, p) dp + \int_0^{\infty} \int_{-\infty}^{\infty} K(x, p, t - \tau) f(0, p) dp r(\tau) d\tau + \\
& \sum_{k=1}^{\infty} (-1)^k \int_0^t \int_{-\infty}^{\infty} K(x, p, t - \tau) f(2dk, p) dp L(k, \tau) d\tau \\
& - \int_{-\infty}^{\infty} K(x, p, t) f(-2d, p) dp - \int_0^{\infty} \int_{-\infty}^{\infty} K(x, p, t - \tau) f(-2d, p) dp r(\tau) d\tau \\
& - (-1)^k \sum_{k=1}^{\infty} \int_0^t \int_{-\infty}^{\infty} K(x, p, t - \tau) f(-2dk, p) dp L(k, \tau) d\tau. \tag{69a}
\end{aligned}$$

Note that the first two terms are identical with the solution of the potential step (40). They describe the behavior of a wave packet that starts with positive momentum and undergoes its first transmission process. The following sum describes the transmitted parts of this wave packet after the  $k$ th reflection at the left wall  $x = -d$ . The remaining terms are relevant for the description of an initial wave packet with negative momentum.

The inverse Laplace transform of (66b) yields

$$\begin{aligned}
\psi(x, t) = & - \sum_{k=0}^{\infty} \int_0^t \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i(t-\tau)}} e^{\frac{i\kappa^2(2d(k+1)+x+y)^2}{2(t-\tau)}} \psi(y, 0) dy (-1)^k M(k, \tau) d\tau \\
& + \sum_{k=0}^{\infty} \int_0^t \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i(t-\tau)}} e^{\frac{i\kappa^2(2d(k+1)+x-y)^2}{2(t-\tau)}} \psi(y, 0) dy (-1)^{k+1} M(k+1, \tau) d\tau \\
& + \sum_{k=0}^{\infty} \int_0^t \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i(t-\tau)}} e^{\frac{i\kappa^2(2d(k+1)-x+y)^2}{2(t-\tau)}} \psi(y, 0) dy (-1)^{k+1} M(k+1, \tau) d\tau \\
& - \sum_{k=0}^{\infty} \sum_{l=0}^{\infty} \int_0^t \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i(t-\tau)}} e^{\frac{i\kappa^2(2d(k+1)-x-y)^2}{2(t-\tau)}} \psi(y, 0) dy (-1)^{k+2} M(k+2, \tau) d\tau \\
& + \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i t}} e^{\frac{i\kappa^2(x-y)^2}{2t}} \psi(y, 0) dy + \int_0^t \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i(t-\tau)}} e^{\frac{i\kappa^2(x+y)^2}{2(t-\tau)}} \psi(y, 0) dy r(\tau) d\tau \\
& \text{for } x < 0, \tag{70}
\end{aligned}$$

where we have introduced

$$M(k, t) \equiv \mathcal{L}^{-1}(\rho(s)^k). \tag{71}$$

The last two terms of (70) coincide with the solution of the potential step (24). If we compare (70) with the solution of the infinite square well (13), we find that (70) consists of the terms of (13) and their convolutions with  $M(k, t)$ . If  $t \gg 2\hbar/V$  and as long as only a limited number of terms are involved, the convolution integral can be extended to infinity as for the approximate solutions

of the potential step (see Appendix B). We find for a sufficiently peaked wave packet with momentum expectation value  $p_0$

$$\begin{aligned}
\psi(x, t) \approx & - \sum_{k=0}^{L_1} (-1)^k R(p_0)^k \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i t}} e^{\frac{i\kappa^2(2d(k+1)+x+y)^2}{2t}} \psi(y, 0) dy \\
& + \sum_{k=0}^{L_2} (-1)^{k+1} R(p_0)^{k+1} \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i t}} e^{\frac{i\kappa^2(2d(k+1)-x+y)^2}{2t}} \psi(y, 0) dy \\
& + \sum_{k=0}^{L_3} (-1)^{k+1} R(p_0)^{k+1} \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i t}} e^{\frac{i\kappa^2(2d(k+1)+x-y)^2}{2t}} \psi(y, 0) dy \\
& - \sum_{k=0}^{L_4} (-1)^{k+2} R(p_0)^{k+2} \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi i t}} e^{\frac{i\kappa^2(2d(k+1)-x-y)^2}{2t}} \psi(y, 0) dy \\
& + \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x-y)^2 \kappa^2}{2t}} \psi(y, 0) dy + R(p_0) \frac{\kappa}{\sqrt{2\pi t i}} \int_{-\infty}^0 e^{\frac{i(x+y)^2 \kappa^2}{2t}} \psi(y, 0) dy
\end{aligned}$$

(72)

for  $x < 0$ .

We find that the approximate solution of the asymmetric square well looks like the solution of the infinite well (13) supplemented by the appropriate reflection coefficients that make allowance for the fact that at each time the wave packet is reflected at  $x = 0$ , parts of the wave packet leave the box. If the wave packet is located between the walls (not too close to any of them) it is described by only term of (70). The probability of finding a corresponding particle within the wall is then given by  $|R(p_0)|^{2m}$ , where  $m$  denotes the already undergone number of reflections at the right wall. The transmission probability is then given by  $1 - |R(p_0)|^{2m}$ .

## 6 Conclusions

We applied the method of Laplace transform to reproduce the mirror solution for the infinite square well in a straightforward way and to derive exact and intuitive solutions for the potential step and the asymmetric square well.

We could show that a wave packet with energy higher than the potential step will partly be reflected and partly transmitted and thereby slowed down (1(b)), as it would be expected from a classical particle that has passed the step. We found that the wave function with energy lower than the potential step will swap into the classically forbidden region only during the reflection process. Moreover we derived an inequality for the reflection probability in terms of the reflection coefficient that is saturated for certain initial wave packets (33,35b).

In the case of the asymmetric well the solution inside the well (70) looks like a generalization of the solution of the infinite well (13). Each term of (70) describes the wave packet after a certain number of reflections and the convolution integrals can be approximated by powers of the reflection coefficient

if the wave packet is sufficiently narrow (72) as long as the system still shows semiclassical behaviour. For this case we also determined the probability of finding a particle inside the box at a given time which yields the corresponding transmission probability. Note that this example illustrates the advantages of the method very well since from the point of view of the characterization in terms of eigenstates the box with exit differs fundamentally from the infinite square well since there are bound and unbound eigenstates.

From the technical point of view we needed for the box with exit only the inverse Laplace transformation of functions that were already used for the infinite square well and the potential step (12, 22). We expect that the solutions of all piecewise constant potentials can be characterized by convolutions of these inverse Laplace transformed functions since the asymmetric square well is a generic case with no special symmetry and a finite as well as an infinite wall. The explicit form of the solutions make the method especially useful for the studying of the tunneling time. It would also be interesting to obtain time dependent solutions for other sorts of exactly solvable potentials as the Morse oscillator or the Hydrogen atom. A generalization of the method to more spatial dimensions can be achieved by using the Duhamel principle.

## A The deformed wave function

The substitution  $q = \sqrt{p^2 - 2mV}$ ,  $q_0 = \sqrt{p_0^2 - 2mV}$  yields

$$\begin{aligned}\tilde{\psi}(y, 0) &= \frac{1}{\sqrt{2\pi\hbar}} \int_{\sqrt{2mV}}^{\infty} e^{\frac{i\sqrt{p^2-2mV}y}{\hbar}} f(p) dp = \\ &= \frac{1}{\sqrt{2\pi\hbar}} \int_0^{\infty} e^{\frac{iqy}{\hbar}} f(\sqrt{q^2 + 2mV}) \frac{q}{\sqrt{q^2 + 2mV}} dq = \\ &= \frac{1}{\sqrt{2\pi\hbar}} \int_0^{\infty} e^{\frac{iqy}{\hbar}} e^{-\frac{i\sqrt{q^2+2mV}x_0}{\hbar}} F\left(\sqrt{q^2 + 2mV} - \sqrt{q_0^2 + 2mV}\right) \frac{q}{\sqrt{q^2 + 2mV}} dq.\end{aligned}$$

Applying (43b) and expanding also  $q/\sqrt{q_0^2 + 2mV}$  around  $q_0$  and using (43a), we find

$$\tilde{\psi}(y, 0) \approx \frac{1}{\sqrt{2\pi\hbar}} \int_0^{\infty} e^{\frac{iqy}{\hbar}} \lambda e^{-\frac{iq_0x_0}{\hbar} - \frac{ix_0\lambda}{\hbar}(q-q_0) - \frac{ix_0\lambda}{2\hbar q_0 k_0}(q-q_0)^2} F(\lambda(q - q_0)) dq.$$

Because of (42) we can extend the integral

$$\tilde{\psi}(y, 0) \approx \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{iqy}{\hbar}} \lambda e^{-\frac{iq_0x_0}{\hbar} - \frac{ix_0\lambda}{\hbar}(q-q_0) - \frac{ix_0\lambda}{2\hbar q_0 k_0}(q-q_0)^2} F(\lambda(q - q_0)) dq.$$

Note that in (43b) the argument of  $F$  is approximated by a Taylor expansion around  $q_0$  up to first order, whereas the exponent is expanded to second order due to a factor  $1/\hbar$

## B Approximation for the asymmetric square well

The solution (70) consists of terms of the form

$$U(k, t) = \int_0^t \frac{\kappa}{\sqrt{2\pi(t-\tau)}i} \int_{-d}^0 e^{\frac{i(Q\pm y)^2\kappa^2}{2(t-\tau)}} \psi(y, 0) dy M(k, \tau) d\tau = \quad (73)$$

$$\int_0^t \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{\mp ipQ}{\hbar}} e^{-\frac{ip^2(t-\tau)}{2m\hbar}} f(p) dp M(k, \tau) d\tau \quad \text{with } Q = 2dk \pm x \quad (74)$$

where, according to ([9])

$$M(k, t) = \mathcal{L}^{-1}(\rho(s)^k) = \frac{k}{i^k t} J_k \left[ \frac{Vt}{2h} \right] e^{-\frac{iVt}{2h}}. \quad (75)$$

If we approximate the  $\tau$ -integral by an infinite Laplace integral as for (31), we find

$$\begin{aligned} U(k, t) &\approx \int_0^\infty \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{\mp ipQ}{\hbar}} e^{-\frac{ip^2(t-\tau)}{2m\hbar}} f(p) dp M(k, \tau) d\tau = \\ &\frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{\mp ipQ}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) \left( \rho \left[ \frac{-ip^2}{2m\hbar} \right] \right)^k dp = \\ &\frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{\mp ipQ}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) R(p)^k dp \end{aligned} \quad (76)$$

For the neglected part of the integral

$$u(k, t) = \int_t^\infty e^{\frac{\mp ipQ}{\hbar}} e^{-\frac{ip^2(t-\tau)}{2m\hbar}} e^{-\frac{iV\tau}{2h}} \frac{k}{i^k \tau} J_k \left[ \frac{V\tau}{2h} \right] d\tau, \quad (77)$$

we get the following estimate

$$\left| \int_t^\infty e^{\frac{ip^2\tau}{2m\hbar} - \frac{iV\tau}{2h}} \frac{k}{\tau} J_k \left[ \frac{V\tau}{2h} \right] d\tau \right| \leq \int_{\frac{tV}{2h}}^\infty \left| \frac{k J_k(y)}{y} \right| dy \leq \quad (78a)$$

$$\left( \int_{\frac{tV}{2h}}^\infty \frac{k}{y^{1+2\epsilon}} dy \right)^{\frac{1}{2}} \cdot \left( \int_0^\infty \frac{(J_k(y))^2}{y^{1-2\epsilon}} dy \right)^{\frac{1}{2}} = \quad (78b)$$

$$k \frac{1}{\sqrt{2\epsilon}} \left( \frac{2h}{Vt} \right)^\epsilon \cdot \left( 2^{2\epsilon} \frac{\Gamma[1-2\epsilon]\Gamma[k+\epsilon]}{2(\Gamma[1-\epsilon])^2\Gamma[1+k-\epsilon]} \right)^{\frac{1}{2}} \leq \quad (78c)$$

$$k \frac{1}{\sqrt{2\epsilon}} \left( \frac{2h}{Vt} \right)^\epsilon \cdot \left( 2^{2\epsilon} \frac{\Gamma[1-2\epsilon]}{2(\Gamma[1-\epsilon])^2} \right)^{\frac{1}{2}} \quad \text{with } 0 < \epsilon < \frac{1}{2}, \quad (78d)$$

where we have applied the Schwarz inequality and the integral formula ([21])

$$\int_0^\infty \frac{(J_k(y))^2}{y^{1-2\epsilon}} dy = 2^{2\epsilon} \frac{\Gamma[1-2\epsilon]\Gamma[k+\epsilon]}{2(\Gamma[1-\epsilon])^2\Gamma[1+k-\epsilon]}. \quad (79)$$

Therefore (76) will be a good approximation for  $U(k, t)$  if  $t \gg \frac{1}{\sqrt{V}}$ . Moreover, if the wave packet is sufficiently peaked around the momentum expectation value  $p_0$ , so that  $R(p_0)^k$  can be put before the integral, we can write

$$U(k, t) \approx R(p_0)^k \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{\mp ipQ}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) dp = \quad (80)$$

$$R(p_0)^k \frac{\kappa}{\sqrt{2\pi t i}} \int_{-d}^0 e^{\frac{i(Q \pm y)^2 \kappa^2}{2t}} \psi(y, 0) dy. \quad (81)$$

However, the estimation (78d) gets worse for increasing  $k$  and moreover the solution (70) consists of infinite sums of terms of the form  $U(k, t)$ . Therefore we can use the approximation only if the number of relevant terms is limited.

We will now show how the approximation works for an initial wave function of the form

$$\psi(y, 0) = e^{\frac{ip_0 y}{\hbar}} G(y),$$

where  $G(y)$  is a two times differentiable real function that is zero outside  $[0, d]$  and  $p_0 = mv_0$  denotes the momentum expectation value of  $\psi(y, 0)$ . Without loss of generality we will only discuss the sum

$$S_1 = \sum_{k=1}^{\infty} U_1(k, t), \text{ with} \quad (82a)$$

$$U_1(k, t) = \int_0^t \frac{\kappa}{\sqrt{2\pi(t-\tau)} i} \int_{-d}^0 e^{\frac{i(2dk+x-y)^2 \kappa^2}{2(t-\tau)}} \psi(y, 0) dy M(k, \tau) d\tau. \quad (82b)$$

Considering only terms with

$$X = \kappa \frac{2dk - x + y - v_0(t-\tau)}{2\sqrt{t-\tau}} > 0$$

and applying the estimate [26]

$$\left| -\text{Erfc}[(1-i)X]X + \frac{e^{2iX^2}}{(1-i)\sqrt{\pi}} \right| \leq \frac{1}{4\sqrt{2\pi}X^2}$$

we find after a twofold partial integration for the spatial integral

$$\left| \int_{-d}^0 \frac{\kappa}{\sqrt{2\pi(t-\tau)}i} e^{\frac{i(2dk+x-y)^2\kappa^2}{2(t-\tau)}} e^{\frac{ip_0y}{\hbar}} G(y) dy \right| = \quad (83)$$

$$\left| \int_{-d}^0 \frac{\sqrt{t-\tau}}{\kappa} \left\{ -\text{Erfc}[(1-i)X]X + \frac{e^{2iX^2}}{(1-i)\sqrt{\pi}} \right\} G''(y) dy \right| \leq \quad (84)$$

$$\left( \frac{\sqrt{t-\tau}}{\kappa} \right)^3 \frac{1}{\sqrt{2\pi}} \int_{-d}^0 \frac{1}{(2dk - v_0(t-\tau) - x + y)^2} |G''(y)| dy \leq \quad (85)$$

$$\left( \frac{\sqrt{t-\tau}}{\kappa} \right)^3 \frac{1}{\sqrt{2\pi}} \frac{1}{(2dk - v_0t - d)^2} \int_{-d}^0 |G''(y)| dy. \quad (86)$$

Applying again the Schwarz inequality and (79) for  $\epsilon = -1/2$ , we can write for the convolution integral

$$\left| \int_0^\infty \left( \frac{\sqrt{t-\tau}}{\kappa} \right)^3 \frac{k}{i^k \tau} J_k \left( \frac{\tau V}{2\hbar} \right) d\tau \right| \leq \quad (87)$$

$$\frac{1}{\kappa^3} \{ (t-\tau)^3 d\tau \}^{\frac{1}{2}} \cdot k \left\{ \left( \frac{V}{2\hbar} \right) \int_0^\infty \frac{J_k(y)}{y} dy \right\}^{\frac{1}{2}} = \quad (88)$$

$$\frac{t^2 \hbar}{2} \sqrt{\frac{V}{2m^3}} \cdot \frac{k}{\sqrt{\pi(k^2 - \frac{1}{4})}} \leq t^2 \hbar \sqrt{\frac{V}{6\pi m^3}}. \quad (89)$$

Then we find for the sum of all  $U_1(k, t)$  (82b) with  $k \leq l$ , where  $l$  is the smallest positive integer with  $l2d \geq v_0t + 3d$ ,

$$\left| \sum_{k=l}^\infty U(k, t) \right| \leq \sum_{k=l}^\infty |U(k, t)| \leq \quad (90)$$

$$\frac{1}{\sqrt{2\pi}} \left( \sum_{m=1}^\infty \frac{1}{4d^2 m^2} \right) t^2 \hbar \sqrt{\frac{V}{6\pi m^3}} \int_{-d}^0 |G''(y)| dy \leq \frac{\pi t^2 \hbar}{24d^2} \sqrt{\frac{V}{18\pi m^3}} \int_{-d}^0 |G''(y)| dy, \quad (91)$$

which can be neglected if

$$\frac{t^2 \hbar}{d^3} \sqrt{\frac{V}{m^3}} \ll 1. \quad (92)$$

So we can write for (82a)

$$S_1 \approx \sum_{k=1}^{l-1} U_1(k, t) = \sum_{k=1}^{l-1} \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{\mp ip(2dk+x)}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) R(p)^k dp \quad (93)$$

$$+ \sum_{k=1}^{l-1} u_1(k, t), \quad (94)$$

where  $u_1(k, t)$  is defined as a special case of  $u(k, t)$  (77), namely

$$u(k, t) = \int_t^{\infty} e^{\frac{ip(2dk+x)}{\hbar}} e^{-\frac{ip^2(t-\tau)}{2m\hbar}} e^{-\frac{iV\tau}{2\hbar}} \frac{k}{i^k \tau} J_k \left[ \frac{V\tau}{2\hbar} \right] d\tau. \quad (95)$$

Applying (78d), we find

$$\left| \sum_{k=1}^{l-1} u_1(k, t) \right| \leq \sum_{k=1}^{l-1} |u_1(k, t)| \leq \quad (96)$$

$$\frac{1}{\sqrt{2\epsilon}} \left( \frac{2\hbar}{Vt} \right)^{\epsilon} \cdot \left( 2^{2\epsilon} \frac{\Gamma[1-2\epsilon]}{2(\Gamma[1-\epsilon])^2} \right)^{\frac{1}{2}} \cdot \frac{l(l-1)}{2}. \quad (97)$$

So this sum can be neglected, if

$$\left( \frac{2\hbar}{Vt} \right)^{\epsilon} \cdot \frac{l(l-1)}{2} \ll 1, \quad (98)$$

and we finally find

$$S_1 \approx \sum_{k=1}^{L_1} \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{\frac{\mp ip(2dk+x)}{\hbar}} e^{-\frac{ip^2 t}{2m\hbar}} f(p) R(p)^k dp, \quad (99)$$

if both conditions (92,98) are fulfilled. Here  $L_1 = l - 1$ . Similarly estimates for the other three types of sums in (70) can be found that will limit the number of relevant terms to  $L_2, L_3, L_4$ .

## Acknowledgments

I thank Helmut Rumpf for his careful reading of this article and his valuable advices. I also thank Beatrix Hiesmayr, Helmuth Hüffel and Helmuth Urbantke for their helpful comments.

## References

- [1] R. Robinett(2004): Quantum wave packet revivals, Phys.Rep. 251, 1-119

- [2] R. Robinett(2000): Visualizing the collapse and revival of wave packets in the infinite square well using expectation values, *Am. J. Phys.* , 410-420
- [3] Z .Gaeta, C. Stroud (1990): Classical and quantum dynamics of a quasi-classical state of the hydrogen atom, *Phys.Rev.A* 42/11,6308-6313
- [4] D. Aronstein, C. Stroud (2000): Analytical investigation of revival phenomena in the finite square-well potential, *Phys. Rev. A* 62, 022102
- [5] M. Gutzwiller (1967): Phase integral approximation in momentum space and bound states of an atom, *J. Math. Phys.* 8, 1979
- [6] E. Heller (1991): Cellular Danamics - A new semiclassical approach to time dependent quantum mechanics,*J. Chem. Phys.* 94, 2723-2729
- [7] M. Kleber (1994) : Exact solutions for time-dependent phenomena in quantum mechanics, *Phys. Rep* 236, 331-393
- [8] A. Zayed (1996): *Handbook of Function and Generalized Function Transformations* (Boca Raton: CRC Press)
- [9] A. Erdely (1954): *Tables of Integral Transforms*,Vol. 1 (New York: McGraw-Hill )
- [10] S.H. Lin, H. Eyring (1971): Solution of the Time Dependent Schrödinger equation by the Laplace Transform Method, *Proceedings of the National Academy of Sciences* 68/1, 76-81
- [11] J.Hladik (1972): Resolution de l'equation de Schrödinger dependant du temps par la method de la transformation de Laplace, *C.R. Acad. Sc. Paris Series B* 275, 177-178
- [12] A. Arnold, M.Erhardt, M.Schulte, Numerical simulations of quantum wave guides, in K.Watanabe (2008): *VLSI and Computer Archetecture* (New York: Nova Science Publishers)
- [13] J. Villavicencio, R.Romo, SSY Silva (2002): Quantum evolution in a step potential barriere, *Phys.Rev.A* 66, 042110
- [14] B. U. Felderhof (2008): Time-dependence and line shape of spontaneous quantum tunneling, *Journal of Physics A* 41, 445302
- [15] G. Garcia-Calderon, A. Rubio (1997): Transient effects and delay time in the dynamics od resonant tunneling, *Phys. Rev. A* 55/5, 3361-3370
- [16] P.Moretti (1992): Tunneling and group velocity in the square potential barrier, *Phys.Rev.A* 46/3, 1233-1237
- [17] G.Krylov, M. Belov (2012): Once again on quantum propagator for the step potential, *AIP Conf. Proc.* 1468, 223-232

- [18] T.O.Carvalho (1993): Exakt space time propagator for the step potential, Phys.Rev. A 47/4, 2562-2573
- [19] V.F. Los, A.V. Los (2010): On the quantum mechanical scattering from a potential step, J.Phys.A 43, 055304
- [20] G.Krylov, M. Belov (2012): Quantum wave propagation through a potential step, AIP Conf. Proc. 1468, 38-46
- [21] W. Magnus, F. Oberhettinger, R.P. Soni (1996): Formulas and Theorems for the Special Functions of Mathematical Physics (Berlin: Springer-Verlag)
- [22] V. V. Dodonov, M. A. Andreato (2000): Deflection of quantum particles by impenetrable boundary, Physics Letters A 275,173-81
- [23] M. Belloni, M. A. Doncheski, R.W. Robinett (2005): Exact Results for 'Bouncing' Gaussian Wave Packets, Physica Scripta Vol.71, 136-140
- [24] M. A. Doncheski, R.W. Robinett (1999): Anatomy of a quantum 'bounce', Eur. J. Phys. 20, 29-37
- [25] A. Messiah: Quantum Mechanics (Amsterdam: North-Holland Publ.)
- [26] M.Abramowitz and I. A. Stegun (1964): Handbook of Mathematical functions