

Phase diagram of non-equilibrium phase transition in a strongly-interacting Rydberg atom vapour

Dong-Sheng Ding,^{1,2,*} Hannes Busche,^{3,4} Bao-Sen Shi,^{1,2,†} Guang-Can Guo,^{1,2} and Charles S. Adams^{3,‡}

¹*Key Laboratory of Quantum Information, University of Science and Technology of China, Hefei, Anhui 230026, China.*

²*Synergetic Innovation Center of Quantum Information and Quantum Physics, University of Science and Technology of China, Hefei, Anhui 230026, China.*

³*Department of Physics, Joint Quantum Centre (JQC) Durham-Newcastle, Durham University, South Road, Durham DH1 3LE, United Kingdom*

⁴*Current affiliation: Department of Physics, Chemistry and Pharmacy, Physics@SDU, University of Southern Denmark, 5230 Odense M, Denmark*

(Dated: July 22, 2022)

Far-from equilibrium dynamics that lead to self-organization are highly relevant to complex dynamical systems not only in physics, but also in life-, earth-, and social sciences. It is challenging however to find systems with sufficiently controlled interactions that allow to model their emergent properties quantitatively. Here, we study a non-equilibrium phase transition and observe signatures of self-organized criticality in a dilute thermal vapour of atoms optically excited to strongly interacting Rydberg states. Electromagnetically induced transparency (EIT) provides excellent control over the population dynamics and enables high-resolution probing of the driven-dissipative system, which also exhibits phase bistability. Increased sensitivity compared to previous work allows reconstruct the system's phase diagram including the observation of its critical point. We observe that interaction-induced energy shifts and enhanced decay only occur in one of the phases above a critical Rydberg population. This limits the application of generic mean-field models, however a modified, threshold-dependent approach is in qualitative agreement with experimental data. Near the transition threshold, we observe self-organization dynamics as small fluctuations are sufficient to induce a phase transition.

PACS numbers: 42.65.Pc 32.80.Rm 42.62.Fi 64.60.Ht

I. INTRODUCTION

Self-organization and non-equilibrium dynamics in complex dynamical systems are responsible for a diverse range of phenomena not only in physics, but also other fields such as earth sciences, biology, or economics [1]. For example, many non-equilibrium systems exhibit self-organized criticality [2] as they evolve to an attractor that coincides with a critical point in their phase diagram. This leads to scale-invariance and ‘pink’ $1/f$ -noise. Detecting non-equilibrium dynamics and resulting phase transitions with high sensitivity in simple experiments would thus aid the study of the origin of many physical phenomena [3] such as hysteresis behavior or interaction-mediated self-organization. Finding systems where the underlying interactions are sufficiently controlled to allow comparison of quantitative models with experiments is challenging however. Laser excitation of atoms in dilute gases allows precision measurements of state populations and phase diagrams, but typical interatomic interactions are too weak to affect the excitation dynamics. For example, the observation of optical bistability in non-equilibrium light-matter systems initially required cavity feedback [4, 5], or cooperative interactions between optical dipoles [6] decoupled from the environment in cryogenic setups [7].

Much stronger interactions can be achieved in ensembles of Rydberg atoms [8, 9]. Their high polariz-

ability leads to strong and long-ranged dipolar interactions over many micrometers and renders them sensitive to distant individual charges, e.g. originating from ionized Rydberg atoms. Driven-dissipative ensembles of Rydberg atoms exhibit rich dynamics in ultracold atoms, such as aggregate formation [10, 11] around a seed excitation [12–16] or ions [17], bi- or metastable phases [10, 18], or self-organised criticality [19]. Due to the strong interactions, effects such as non-equilibrium phase transitions [20–24] and aggregate formation [25] can also be observed in the optical response of room-temperature ensembles, significantly simplifying the experimental setup. Non-equilibrium phase transitions lead to frequency-dependent optical bistability, while the formation of aggregates can be detected in the temporal evolution of the light transmission. Both ultracold and room-temperature driven-dissipative Rydberg gases have attracted great theoretical interest to study non-equilibrium dynamics and exotic phases of matter [26–29].

In this work, we investigate the non-equilibrium dynamics and reconstruct the resulting phase diagram of a driven-dissipative ensemble of strongly interacting Rydberg atoms in a room-temperature atomic vapour. For this, we exploit the narrow linewidths achievable with electromagnetically-induced transparency (EIT) [30, 31] while scanning the parameters of the EIT probe and coupling fields prevents the system from reaching equilib-

rium. The system dynamics thus strongly depend on parameter scan rates and directions. The narrow EIT linewidth enhances the frequency resolution by two orders of magnitude compared to previous work [20–22]. As a result, we can map out the full phase diagram including the observation of a critical point and reveal previously unobserved spectral features in the bistable optical response and non-equilibrium dynamics at weak driving. Importantly, we find that interaction-induced lineshifts and broadening occur only in one of the two phases of the system above a critical Rydberg atom density. This implies that mean-field based models of driven-dissipative Rydberg vapours need to be modified in the limit of weak driving and limits the potential interaction mechanism responsible for the phase transition, i.e. whether interaction effects originate from coherent dipolar interactions or ionizing collisions [22]. Recent complementary work [24] has identified avalanche ionization and plasma formation as the dominant mechanism for stronger driving.

We also show that the transition threshold can be shifted by adding a second probe beam that enhances driving to the Rydberg state and demonstrate optical switching between two transmission states. Finally, we investigate the behavior near the phase transition point under quasi-steady-state conditions with constant EIT field parameters. In this regime, the system is highly susceptible to small variations in the Rydberg density such that even short-term fluctuations can induce phase transitions, though the system quickly restores its original phase. This indicates the presence of self-organisation and possibly self-organised criticality as recently observed in a cold Rydberg gas [19].

II. BACKGROUND AND MODEL

Throughout this work, we consider the following non-equilibrium system (figure 1(a)): An optical probe field propagates through a thermal vapor of three-level atoms and drives the transition between a ground state $|g\rangle$ and a low-lying, short-lived excited state $|e\rangle$. A coupling field couples $|e\rangle$ to a highly excited and long-lived Rydberg state $|r\rangle$. The probe (coupling) Rabi frequency, angular frequency, and detuning are denoted by $\Omega_{p(c)}$, $\omega_{p(c)}$, and $\Delta_{p(c)}$, respectively, while $\Gamma_{e(r)}$ are the decay rates of $|e\rangle$ ($|r\rangle$). Since $\Gamma_r \ll \Gamma_e$, this corresponds to a ladder EIT [31] configuration for $\Omega_p \ll \Omega_c$, and the ensemble is rendered transparent for the probe in a narrow frequency window around two-photon resonance $\Delta_p = -\Delta_c$. Altering the strength of the light-matter coupling dynamically, either by scanning the Rabi frequencies or the detunings of the EIT-fields, changes the rates at which the states are populated and introduces non-equilibrium dynamics into the system. For practical reasons, we consider the transmission for a resonant probe ($\Delta_p = 0$) while scanning Δ_c (or $\Omega_{p(c)}$) in the subsequent experiments.

Self-organized criticality relies on an interaction-induced avalanche in the non-equilibrium dynamics [32] to trigger a phase transition. In a thermal Rydberg vapor, such an avalanche can be induced by either dipolar interactions that facilitate the formation of an aggregate excitation around a seed [25], or ionization of Rydberg atoms due to collisions with electrons and ions originating from previously ionized atoms [33–36]. In either case a critical Rydberg density $N_{R,(c)}$ is required for the avalanche-induced phase transition between two distinct phases: a low- N_R , non-interacting (NI) phase and a high- N_R interacting (I) phase in which Δ_c and Γ_r are modified (figure 1b). Near $N_{R,(c)}$, the system cannot equilibrate as small fluctuations can disturb the equilibrium. In practice, the interacting Rydberg atoms can thus act as a seed [2], that induce an avalanche effect on the surrounding atoms (see figure 1(b)) and leads to a macroscopic phase transition [1]. Due to the narrow linewidths possible in EIT, the optical response on two-photon resonance and population dynamics are strongly affected by the level shifts induced by the avalanche. Above $N_{R,(c)}$, they cause an effective, N_R -dependent detuning of the coupling field, $\Delta_c \rightarrow \Delta_c + \Delta'(N_R)$. Besides resonance shifts, the $|nD_J\rangle$ -Rydberg states used in this work are also subject to broadening and enhanced dephasing, $\Gamma_r \rightarrow \Gamma_r + \Gamma'(N_R)$ [37, 38], as a result of motional averaging of the interaction potentials (in case of dipolar interactions) or Stark shifts (for electric fields created by ions) over different m_J -states (see appendix A). Moreover, ionization would represent an additional dissipation channel in itself [36, 39].

In a driven-dissipative system, N_R depends not only on current parameters, but also the residual Rydberg population and thus previous parameter values and their rate of change. Hence, the effective values of Δ_c and Γ_r depend on the scan direction and give rise to hysteresis effects that result in a bistable optical response. Distinct transmission states correspond to the NI-phase in which the optical response is not affected, and the I-phase in which Δ_c and Γ_r are modified (figure 1b).

To include the dynamics of the phase transition in our description of the system, we investigate the time evolution of populations and coherences in the atomic ensembles' density matrix ρ by adopting a threshold-modified mean-field approach where the additional detuning $\Delta' = \eta_1 N_R$ and the enhanced decay $\Gamma' = \eta_2 N_R$ are proportional to the Rydberg density above a critical Rydberg population $\rho_{rr}^{(c)}$ corresponding to $N_{R,(c)}$. This approach is applicable both in the case of dipolar interactions, where N_R relates to the mean spacing between Rydberg atoms and thus interaction strength, and for background fields originating from ionized Rydberg atoms, where N_R relates to the ion density and the resulting Stark shifts.




Figure 1. (a) Overview of experimental setup. A probe and an identical reference beam are propagating in parallel through a heated Rb cell (length 5 cm). The probe beam is overlapped with a counterpropagating EIT coupling beam. Their transmission signals are detected on a differencing photodetector (DD). The probe and reference fields drive the $|g\rangle = |5P_{1/2}, F=3\rangle \rightarrow |e\rangle = |5P_{1/2}, F'=2\rangle$, the coupling light couples the $|e\rangle \rightarrow |r\rangle = |nD_{3/2}\rangle$ transition in ^{85}Rb . A switch beam can be added to enhance the Rydberg population without changing the probe intensity. Labels: PBS - polarising beam splitter, DM - dichroic mirror, $\lambda/2$ - half-wave plate. (b) Excitation and interaction dynamics in the different phases. In the non-interacting (NI)-phase, the probe transmission under EIT conditions remains unaffected. Above a critical Rydberg atom density $N_{R,C}$, the system is found in an interacting (I) phase where strong interactions lead to energy shifts and enhanced decay of $|r\rangle$.

A. Rydberg EIT in a thermal vapour

In absence of interatomic dipole-dipole interactions or frequency shifts due to external fields, the interaction of the EIT light fields with an ensemble of three level atoms is governed by the master equation

$$\frac{d}{dt}\rho = -\frac{i}{\hbar}[H, \rho] + \frac{1}{\hbar}L \quad (1)$$

where ρ is the atomic ensemble's density matrix and $H = \sum_k H[\rho^{(k)}]$ the atom-light interaction Hamiltonian summed over all the single-atom Hamiltonians

$$H[\rho^{(k)}] = -\frac{\hbar}{2} \begin{pmatrix} 0 & \Omega_p & 0 \\ \Omega_p & -2\Delta_p & \Omega_c \\ 0 & \Omega_c & -2(\Delta_p - \Delta_c) \end{pmatrix}. \quad (2)$$

The Lindblad superoperator $L = \sum_k L[\rho^{(k)}]$ comprised of the single-atom superoperators

$$L[\rho^{(k)}] = \frac{\hbar}{2} \begin{pmatrix} 2\Gamma_e \rho_{ee}^{(k)} & -\Gamma_e \rho_{ge}^{(k)} & -\Gamma_r \rho_{gr}^{(k)} \\ -\Gamma_e \rho_{eg}^{(k)} & -2\Gamma_e \rho_{ee}^{(k)} + 2\Gamma_r \rho_{rr}^{(k)} & -(\Gamma_r + \Gamma_e) \rho_{er}^{(k)} \\ -\Gamma_r \rho_{er}^{(k)} & -(\Gamma_r + \Gamma_e) \rho_{re}^{(k)} & -2\Gamma_r \rho_{rr}^{(k)} \end{pmatrix} \quad (3)$$

accounts for the finite lifetime of $|e\rangle$ and $|r\rangle$. The complex susceptibility of the EIT medium including the Doppler effect due to atomic motion is

$$\chi(v)dv = \frac{|\mu_{ge}|^2}{\epsilon_0 \hbar} \rho_{eg}(v)dv \quad (4)$$

with

$$\rho_{eg}(v)dv = \frac{\Gamma_r + 2i(\delta + \Delta_D)}{(2\Delta_p + 2\omega_p v/c - i\Gamma_e)(\Gamma_r + 2i(\delta + \Delta_D) - i\Omega_c^2)} N(v)dv$$

where $\Delta_D = (\omega_p - \omega_c)v/c$ denotes Doppler shift experienced by an atom moving with velocity v and $\delta = \Delta_c + \Delta_p$ the two-photon detuning. The transmission through the EIT medium can be obtained from the susceptibility via

$$T \sim e^{-\text{Im}[\int k L \chi(v)dv]} \quad (5)$$

where L is the medium length and k the wavevector of the probe field.

B. Implementation of threshold model

In the subsequent experiments, we investigate the change in the optical response as either Δ_c or Ω_p are scanned. To model the EIT spectra, we compute $\chi(v)dv$ substituting $\Delta_c \rightarrow \Delta_c + \Delta'(N_R)$ and $\Gamma_r \rightarrow \Gamma_r + \Gamma'(N_R)$ as explained once the critical threshold population $\rho_{rr} > \rho_{rr}^{(c)}$ is reached. We set $\eta_1/2\pi = 1.27 \times 10^{-2} \text{ MHz } \mu\text{m}^3$ and $\eta_2/2\pi = 7.96 \times 10^4 \text{ MHz } \mu\text{m}^3$, to match the calculated EIT spectra best to the experimental results for $|r\rangle = |47D_{3/2}\rangle$, see figure 2. As dissipation is increased in the I-phase, we reduce the threshold population $\kappa \rho_{rr}^{(c)}$ below which the system reverts to the NI-phase. Setting $\kappa = 0.78$ gives the best fit with the experimental results shown in figure 2b.

III. EXPERIMENTAL SETUP

The experimental setup and a scheme of the relevant ^{85}Rb energy levels are depicted in figure 1. The probe and coupling fields counterpropagate through a 5 cm long Rb vapor cell at a temperature of approximately $T \approx 50^\circ\text{C}$. The atomic density is $1.5 \times 10^{11} \text{ cm}^{-3}$ corresponding to a mean inter-atomic spacing of $\approx 1 \mu\text{m}$. The probe beam ($1/e^2$ -waist radius $\approx 500 \mu\text{m}$) couples the ground state $|g\rangle = |5S_{1/2}, F=3\rangle$ to an intermediate state $|e\rangle = |5P_{1/2}, F'=2\rangle$. The coupling beam ($1/e^2$ -waist radius $\approx 200 \mu\text{m}$) is resonant with the transition from $|e\rangle$ to a Rydberg state $|r\rangle = |nD_{3/2}\rangle$. The estimated peak values of $\Omega_c/2\pi$ are $\approx 20 \text{ MHz}$ for $n = 47$ and $\approx 10 \text{ MHz}$ for $n = 70$ unless stated otherwise. A reference beam identical to the probe, but not overlapping with the coupling light, is sent through the cell in parallel and its transmission signal is subtracted from the probe signal following detection on a pair of balanced amplified photodiodes.

The non-equilibrium phase transition is investigated as three parameters are varied: the coupling detuning Δ_c and the Rabi frequencies Ω_p and Ω_c . Scanning Δ_c allows us to obtain EIT transmission spectra while the probe light is kept on resonance, $\Delta_p/2\pi = 0 \text{ MHz}$, such that the Doppler-broadened absorption background can be neglected. Scanning Ω_p alters the Rydberg population and thus mean interaction strength as $N_R \propto \rho_{rr} \propto \Omega_p^2$ (see above).

In addition to the probe, an additional switching beam can be applied to alter the Rydberg density and control the threshold of the phase transition without changing the properties of the probe itself. It also couples $|g\rangle$ and $|e\rangle$ with detuning Δ_p , but an independent Rabi frequency Ω_s . It intersects with probe and coupling beams at the center of the cell at 2° to avoid crosstalk between the switching, probe, and reference beams at the detector.

IV. EXPERIMENTAL RESULTS

A. Optical bistability

In order to observe Rydberg-mediated optical bistability in the optical response of the three-level EIT medium, we initially observe the EIT lineshape as Δ_c is scanned through resonance from negative to positive detuning and vice versa. Fig. 2 shows the results for a range of different Ω_p as well as results of the threshold-model (b, dashed lines indicate where the system is in the I-phase). At $\Omega_p/2\pi = 6.3 \text{ MHz}$ (Fig. 2 a1) identical EIT spectra are observed for either scan direction without bistability. At 6.7 MHz – and consequently only slightly higher Rydberg density N_R – we observe almost symmetric optical bistability above and below resonance (Fig. 2a2). A sudden drop in the transmission occurs on reso-

nance, where the excitation rate is highest, independent of the scan direction indicating the transition from the NI- to the I-phase. In this regime, the energy shift Δ' induced on $|r\rangle$ is small compared to both Δ_c and the EIT linewidth. This behavior at low Ω_p is distinctively different compared to previous experiments on Rydberg-induced optical bistability that used shelving techniques with higher Ω_p and linewidths [20, 22]. These observed bistability at large red detuning as the peak transmission was strongly shifted due to higher ρ_{rr} and transmission in the I- (scanning from positive to negative Δ_c) exceeded the NI-phase. The ability to probe this new regime and detection of the phase-transition with sub-MHz frequency resolution is a consequence of the narrow EIT resonance. The bistability windows observed in Fig. 2(a2), have widths $< 0.5 \text{ MHz}$, two orders of magnitude narrower compared to previous experiments [20, 22].

As Ω_p is increased (Fig. 2a3 to 2a5), the spectra become asymmetric as an increasingly strong spectral shift Δ^* (see definition in Fig. 2a5) of the peak transmission relative to $\Delta_c/2\pi = 0 \text{ MHz}$ is observed. The shift increases with Ω_p as $N_R \propto \rho_{rr} \propto \Omega_p^2$. At $\Omega_p/2\pi = 6.8 \text{ MHz}$ (Fig. 2a3), bistability is still observable within the EIT resonance feature, but the phase transition occurs at lower/higher Δ_c value for positive/negative scan directions respectively. The on-resonance phase transition disappears as the system is in the I-phase for either scan direction.

For Δ' comparable to the EIT linewidth, near-resonant bistability features disappear entirely. Instead, new bistability windows appear on both sides of the EIT window, as in shelving experiments [20, 22]. Unlike above, the transmission in the I- is now exceeding that in the NI-phase. In the bistability window below $\Delta_c = 0$, the system is found in the I-phase for scans from positive to negative Δ_c where the scan has already crossed through resonance and sufficient Rydberg population has been built up to maintain the state. Above $\Delta_c = 0$, the roles are reversed and system is in the I-phase if the scanned from negative to positive detuning, again after crossing through resonance. The bistability window is narrower for $\Delta_c > 0$ as the sign of Δ' implies an overall red-shift of the spectra due to the choice of $|r\rangle$. Bistability for blue detuning was also observed by Weller et al. [20, 22], but not in the original experiment by Carr et al. [20] where the probe was significantly stronger.

Fig. 2(b), shows EIT spectra calculated with Doppler-averaging as given by equation 5 for a critical Rydberg population (the critical density is estimated as $N_{r,(c)} = 2.9 \times 10^{10} \text{ cm}^{-3}$ assuming that each probe photon creates a Rydberg atom) and all other parameters as stated in section II B. The features occurring as $\rho_{rr} \propto \Omega_p^2$ increases are qualitatively consistent with the experimental data. When the system is in the I-phase, dissipation is increased due to many-body dephasing as discussed before. This results in a lower threshold for the tran-




Figure 2. Observation of optical bistability induced by a non-equilibrium phase transition in EIT transmission spectra. (a) Transmission of probe light as Δ_c is scanned from red- to blue-detuning (red) and vice versa (black) for increasing probe Rabi frequencies $\Omega_p/2\pi = 6.3$ (a1), 6.7 (a2), 6.8 (a3), 7.4 (a4), and 7.9 MHz (a5). Panel (a5) shows the definitions of the transmission difference between the states at the point of the phase transition and the spectral shift Δ^* referred to subsequently. (b) Simulated transmission spectra for increasing Rydberg densities with $N_r = 9.5$ (b1), 10.0 (b2), 10.8 (b3), 15.0 (b4), and 18.0 (b5) where N_r is proportional to the number of Rydberg atoms. The dashed line in panel (b) shows where the system is found in the strongly interacting I-phase.

sition from the high- to the low- ρ_{rr} phase compared to the reverse process. We also note an increase in the peak transmission with Ω_p in the experimental data that is not reproduced by the model.

B. Non-equilibrium phase diagram

The enhanced sensitivity of EIT in the detection of Rydberg induced optical bistability and the underlying phases allows to map out the phase diagram of the driven-dissipative system and investigate the character of the non-equilibrium phase transition. The phase diagram in Fig. 3(a) is obtained by scanning $\Delta_c/2\pi$ over 96 MHz around resonance at a frequency of 10 Hz for various Ω_p with $\Omega_c/2\pi = 13.8 \pm 0.5$ MHz and shows the normalized difference in probe transmission between scan directions. As the bistability is a consequence of the scan-direction dependent occupation of the two phases, the corresponding regions represent the boundaries between the NI- and the high- ρ_{rr} I-phase. Outside these areas, there is no transmission difference indicating that the system is in the same phase irrespective of scan direction, i.e. in the I-phase in between the bistable regions, and the low-

ρ_{rr} NI-phase outside. We observe a critical point at a threshold Rabi frequency of $\Omega_p^{(c)}/2\pi = 5.9 \pm 0.2$ MHz with $\Delta_c^{(c)}/2\pi = 0$ MHz. The spectra shown in Fig. 2(a1) and (a2) correspond to regimes slightly below and above $\Omega_p^{(c)}$. The difference in $\Omega_p^{(c)}$ between Fig. 2a and Fig. 3 results from a slight temperature difference between data sets. As above, the bistable branch is wider for negative Δ_c due to the sign of Δ' . In the bistable branch for negative Δ_c , the systems occupies the I-phase for scans from blue to red detuning and vice versa in the branch for positive Δ_c as discussed above. Note that in the direct vicinity of the critical point, the sign of the transmission difference is switched compared to larger Ω_p . This does not correspond to the occupancy of different phases, but the different regimes for the shift Δ' compared the EIT linewidth as discussed above (Fig. 2).

The onset of the phase transition is characterized by a broken symmetry accompanied by a non-zero order parameter [40]. Fig. 3 (b) shows the shift in peak transmission Δ^* (as defined in Fig. 2a5) vs. $\Omega_p^2 \propto \rho_{rr} \propto N_r$. Depending on the origin of the underlying interactions, either the charge density (ion-induced interactions), or the mean spacing between Rydberg atoms (dipolar interactions) would represent the order parameter, both




Figure 3. Characterizing the non-equilibrium phase transition. (a) Color map of the difference in probe transmission (normalized to the respective maximal values) when scanning Δ_c in either direction, effectively representing the phase diagram of the non-equilibrium system. In the white regions, the system is found in the same phase independent of the scan direction. In the shaded regions, the system is bistable (red if transmission is higher for scans from negative to positive Δ_p , blue in the opposite case) and occupies different phases depending on the scan direction. (b) Shift of peak transmission Δ^* vs. $\Omega_p^2 \propto \rho_{rr}$. Inset: Transmission difference at the transition point for different Ω_p (see Fig. 2a5 for definitions). (c) Measurement of transmission spectra vs. Δ_c for $\Omega_c/2\pi$ between 2.7 and 6.54 MHz. (d) Susceptibility of the phase transition to variations in Ω_c measured in terms of the change in transmission $dT/d\Delta_c$. The shaded region indicates where the system is in the I-phase and $dT/d\Delta_c$ is no longer continuous at the transition point.

ultimately related to N_R . To determine the critical point for, we fit

$$\Delta^* \propto \begin{cases} 0 & \text{for } \Omega_p^2 < \Omega_{p,(c)}^2 \\ (\Omega_p^2 - \Omega_{p,(c)}^2)^\beta & \text{for } \Omega_p^2 \geq \Omega_{p,(c)}^2 \end{cases} \quad (6)$$

and find $\Omega_{p,(c)}^2 = 37(2\pi \times \text{MHz})^2$ with $\beta = 1$. The continuity at $\Omega_{p,(c)}^2$ indicates that the system undergoes a continuous phase transition.

The observation of a clear threshold for Δ^* has consequences for possible physical origins of the phase transition and theoretical descriptions of the system. The physical origin of the phase transition is the broadening of the Rydberg excitation above the critical Rydberg population $\rho_{rr}^{(c)}$. It can be observed both in blue and red detuning, thus the direction of bistability appeared is not dependent on the sign of Rydberg polarisability [22]. The threshold behavior implies that the shift cannot originate directly from effects on individual Rydberg atoms, e.g. ionization via collisions with ground state atoms, as one would expect $\Delta^* \propto \rho_{rr}$ for any ρ_{rr} in this case. Hence, the transition must result from pro-

cesses that involve multiple Rydberg atoms, i.e. dipolar interactions or plasma formation. Rydberg-ground state atom collisions have been observed in thermal Rydberg ensembles as a charged mean-field under EIT-conditions [31] and as a dominant ionization mechanism in a beam of thermal Sr atoms, but the absence of significant shifts and broadening below $\rho_{rr}^{(c)}$ rules them out as direct origin of the phase transition. However, even though they have no immediate effect on the spectra, the resulting ions and electrons are crucial for ionization avalanches and plasma formation [24].

For models describing the system, the threshold behavior also implies that a standard mean field model [26], where the interaction is incorporated by introducing $\Delta' \propto \rho_{rr}$, is insufficient to describe the non-equilibrium dynamics near the critical point, but should remain a valid approximation for $\Omega_p \gg \Omega_{p,(c)}$ as in previous experiments [20–22]. However, by introducing a threshold-dependent mean-field shift and broadening, the phenomenological model described in section II B reproduces the characteristic features of transmission spectra qualitatively well (Fig. 2). Data showing analogous behavior

for the linewidth of the transmission feature at $\Omega_{p,(c)}^2$ can be found in appendix B.

The inset in Fig. 3(b) shows the transmission difference between the phases at the transition point. We also measure the transmission difference (definition in Fig. 2a5) at the transition point (for negative detunings) against Ω_p as Δ_c is scanned. While the transmission difference between the phases increases only slowly for larger Ω_p , and the change in sign (see also Fig. 2) near the critical point becomes once again evident as Ω_p becomes large enough for Δ' to exceed the narrow EIT linewidth.

To further characterize the phase transition, we investigate the dynamics near the critical point. This requires to systematically vary further parameters such as the number and excitation rate of Rydberg atoms or the interaction strength, which influence the modified decay rates, energy shift and the transmission level. Here, we change the coupling Rabi frequency Ω_c while keeping $\Omega_p/2\pi = 6.59$ MHz constant, and record the probe transmission vs. Δ_c (Fig. 3c). As Ω_c is increased, the change in transmission $dT/d\Delta_c$ for red detuning becomes steeper up to the point where the phase transition appears. Fig. 3(d) shows $dT/d\Delta_c$ vs. Ω_c and thus the susceptibility of the system to Ω_c . The slope diverges at the transition threshold as expected for a continuous phase transition. In principle, it would be possible to extract an critical exponent from a similar measurement, but we refrain from this as the errors on $dT/d\Delta_c$ are large and unlike before, the condition $\Omega_c > \Omega_p$ for EIT is no longer met, which may significantly affect the excitation dynamics compared to above.

C. Manipulation of the threshold

In the previous experiments the EIT field parameters, i.e. Ω_p , and its current value determine ρ_{rr} and thus N_R and the onset of the phase transition. The bistability illustrates that N_R , and not any specific value of the EIT fields, is the critical quantity. This allows to further explore the threshold behavior and demonstrate control of the threshold as we apply an additional weak switching field with Rabi frequency $\Omega_s/2\pi = 4.1$ MHz (see Fig. 1) to enhance the Rydberg population and observe the effect on the phase diagram. The switching field increases the driving strength of the $|g\rangle \rightarrow |e\rangle$ transition leading to a higher Rydberg excitation rate without increasing Ω_p itself. In the following, $\Delta_c/2\pi$ is scanned over 88 MHz at 20 Hz and $T \approx 55^\circ\text{C}$.

Fig. 4 (a) and (b) compare the phase diagram without (compare Fig. 3a) and with application of the switching field. The general structure of the phase diagram with the switching field applied remains unchanged, however the threshold and thus the critical point of the phase diagram appear at $\Omega_p^{(c)}/2\pi \approx 4$ MHz instead of ≈ 5 MHz

as ρ_{rr} is enhanced by the additional driving. We also measure the spectral shift Δ^* in the I-phase (Fig. 4c, d) and again find the critical point at a lower value of $(\Omega_p^{(c)}/2\pi)^2 = 16.2$ MHz² (Fig. 4d) compared to 24.3 MHz² without switching field (Fig. 4c) as the relation $\rho_{rr} \propto \Omega_p^2$ no longer holds. Without effective many-body interaction, all atoms in the vapor could be described individually and the switching field should not affect the threshold.

The ability to shift the critical value $\Omega_p^{(c)}$ via the switching field can be used to implement an optically controlled switch between two transmission states for the probe light (Fig. 4d-f) similar to experiments on latching detection of THz-fields using Rydberg optical bistability [23]. To implement the switch, we change the Rydberg state to $|r\rangle = |70D_{3/2}\rangle$ thus increasing the sensitivity while all other parameters remain unchanged. Fig. 4(d) shows the transmission vs. Δ_c with and without switching field. When present, the transition to the I-phase is reduced by approx. 2 MHz such that a switch can be implemented by fixing $\Delta_c/2\pi = -20$ MHz in between the two transition points. We record the transmission over time at fixed detuning both with and without the switching field (Fig. 4e). Fig. 4(f) shows corresponding histograms of the probe transmission for the two transmission states that highlight the change in the statistical distribution of the transmission level. Bimodal counting statistics depending on interactions strength have been predicted by theoretical investigations of Rydberg-mediated optical bistability [29]. Multiple switching fields that only intersect with parts of the excitation volume could be used to realize switching between more than two transmission states cascading multiple bistability regions with different thresholds. Analogously, a setup addressing different Rydberg states in different regions could give further insight whether the phase transition originates from dipolar interactions or ions, similar to experiments which used different isotopes excited to different states as field probes [22] and to study energy exchange processes [41, 42].

While the transmission levels are clearly distinguishable, short term fluctuations between them can be observed (Fig. 4e). We attribute these to short term fluctuations in the Rydberg density that result in a phase transition. Remarkably, the system quickly reverts to its initial phase further supporting the presence of self-organization dynamics.

D. Sensitivity to fluctuations

In the following, we further investigate the sensitivity of the phase transition to fluctuations and the resulting self-organizing response. So far, we have investigated the dynamics for probe and coupling fields with parallel linear polarization. By rotating the polarization of the




Figure 4. Manipulating the phase diagram and optical switching. (a,b) Phase diagrams (at $n = 47$) without (a) and with (b) switching field applied (colors as fig. 3). While the structure of the diagram remains unaffected, the transition occurs at a reduced threshold $\Omega_{p,(c)}$ with switching field applied (T2 vs. T1). (c) Interaction-induced shift Δ^* vs. $(\Omega_p/2\pi)^2$ without (black) and with switching field (blue). (d) Transmission as Δ_c is scanned with and without switching field applied (here $n = 70$). (e) Transmission over time at constant detuning $\Delta_c/2\pi = -20$ MHz (line A in panel d) without (top, blue) and with switching field (bottom, red). (f) Histogram of transmitted intensity levels without (top, blue) and with switching field (bottom, red).

two beams with respect to each other, the rate at which different magnetic sub-levels of the Rydberg state are populated, is enhanced. As a result the system becomes more sensitive to fluctuations in the N_R -dependent interaction strength $\Gamma_r + \Gamma'(N_R)$ and $\Gamma_r + \Gamma'(N_R)$ of $|r\rangle$ in the I-phase is further increased as previously uncoupled m_J -states become more populated. Fig. 5(a) shows transmission scans vs. Δ_c for various angles θ of the half-wave plate controlling the polarization ($\theta = 0^\circ$ corresponds to parallel, 45° to orthogonal polarization) at $|r\rangle = |70D_{3/2}\rangle$ with $m_J \in \{\pm 3/2, \pm 1/2\}$. As the parallel polarization component is reduced, the bistability window becomes narrower and is almost closed for orthogonal polarization (Fig. 5b). Interestingly, the polarization dependence of the hysteresis width cannot be observed at lower Rydberg states ($|r\rangle = |47D_{3/2}\rangle$, see appendix D), which further indicates that enhanced dephasing occurs at higher n due to a wider range of polarizabilities ($\propto n^7$) among the m_J -states that lead to a wider spread of level shifts. This is on stark contrast to the fact that Γ_r is reduced an isolated Rydberg atom with higher n due to their longer lifetimes ($\propto n^3$).

For orthogonal polarization, the hysteresis loop is almost closed, the phase of the system is particularly sensitive to fluctuations. Transmission fluctuates between the levels associated with the two phases when scanning Δ_c ($\Delta_c/2\pi$ is scanned over 96 MHz at 10 Hz) over the bista-

bility window (Fig. 5c). Here, even small fluctuations in ρ_{rr} can induce a phase transition due to the strong dynamic nonlinearities in excitation and decay rates, which are even non-continuous at the threshold $\rho_{rr} = \rho_{rr}^{(c)}$ and characteristic for self-organizing systems.

We define the energy E_I and E_{NI} for the high- ρ_{rr} NI- and low- ρ_{rr} I-phases respectively, with the corresponding dissipation rates given by $\Gamma_I = \Gamma_r + \Gamma'(N_R)$ and $\Gamma_{NI} = \Gamma_r$. Since $\Gamma_I > \Gamma_{NI}$ as, the system's phase is unstable if

$$0 < E_I - E_{NI} < \hbar(\Gamma_I - \Gamma_{NI}) = \hbar\Gamma'(N_R) \quad (7)$$

i.e. the energy of the phases is comparable to the change in decay. For $E_I - E_{NI} > \hbar\Gamma'(N_R)$, the system always occupies the I-phase.

To characterize the response time to fluctuations, we fix $\Delta_c/2\pi = 25.5$ MHz and record the probe transmission over time (Fig. 5d). The system is mostly found in the I-phase, but occasionally, fluctuations induce a transition to the NI-phase resulting in a drop in transmission yet quickly reverts to the I-phase again (see also Fig. 4e). We investigate the distribution of response times required to restore the phase. Fig. 5(e) shows the distribution of the time between adjacent phase jumps. We find that most intervals last below 100 μ s corresponding to the brief transitions to the NI-phase apparent in Fig. 5(d). Measuring the width of one of the spikes yields




Figure 5. (a) Transmission spectra vs. Δ_c ($n = 70$) for different angles of the coupling beam polarization with respect to the probe (θ : angle of half-wave plate in coupling beam path, $\theta = 0^\circ$ parallel, 45° orthogonal linear polarization). Red spectra corresponds to scan direction from red- to blue-detuning, black vice versa. (b) Width of the hysteresis window in which bistability is observed vs. θ . (c) For orthogonal polarisation, the bistability window is narrowest and the system’s phase is unstable as Δ_c is scanned. (d) Transmission over time for fixed $\Delta_c/2\pi = 25.5$ MHz near the center of the instability window. (e) Histogram of the time intervals between phase jumps. Inset: Bimodal distribution of transmission levels subtracted from the maximal transmission. (f) Fourier spectrum of the phase fluctuations. The fits in panels (b) and (e) are guides to the eye.

a similar result of $110 \mu\text{s}$ (appendix C). This instability shows a time scale-invariance for a self-organized criticality, which is governed by the random fluctuations. The inset in Fig. 5(e) shows a histogram of the transmission difference between adjacent data points. The bimodal distribution confirms that the observed fluctuations are not only artifacts originating from noise induced in data acquisition, but correspond to fluctuations between two distinct transmission states. Whether the I- or NI-phase is predominantly occupied depends on the value of Δ_c with detuning closer to resonance favoring the I-phase. Here the system restores to the I-phase and the histogram is biased to higher transmission as predicted in [29]. Similar behavior can also be observed in spontaneous recovery processes in dynamical networks [43], where the network switches back and forth between two collective modes characterized by high and low network activity in analogy to the weakly and strongly interacting phases.

Finally, we analyze the Fourier spectrum of the probe transmission (Fig. 5 f). For low frequencies f , the noise spectrum follows a $1/f$ -power law corresponding to pink noise which occurs in systems exhibiting self-organized criticality [2]. If we split the data over time into individual data sets each only including data points corresponding to the same phase, a separate analysis of the Fourier spectra yields white noise instead confirming that

the pink noise is induced at the phase transition.

E. Scale-invariance control

As demonstrated (Fig. 4), the non-equilibrium dynamics are not only susceptible by fluctuations and noise, but may also be guided by an external seed signal. The introduced energy disturbs the boundary between order and chaos in the system, leading to seed-guided self-organization dynamics. This means the self-organized dynamics of the complexity system can be led by the arbitrary external signal, which is somehow essential for studying artificial intelligence [44] and human consciousness [45]. Such self-organized dynamics is particularly important because of its significance in other fields that are underpinned by similar mechanisms. This mechanism not only brings us applications for sensing noise or external field, but also has a prospective for studying exotic phases of many-body physics. Self-organized criticality relies on three basic elements: strong dynamical non-linearity, balance of exploitation and exploration and manybody interactions [32]. The ingredients of self-organized criticality are well-matched with the exaggerated properties of Rydberg atoms.

There are many spikes with different transmission levels when the system is unstable that corresponds to the distinct avalanche energy. In order to capture the non-




Figure 6. Spike distribution versus the time in three regions: low-based distribution (a), Bimodal distribution (c) and high-based distribution (e). The spike numbers versus the energy interval ΔE at situations (a), (c) and (e), here ΔE is the transmission interval. The small block diagrams in (a), (c) and (e) are the statistic distribution. The red line in shaded area for each case are fitted by the linear functions of $y = -2.2x - 0.2$ (figure. 6b), $y = 2x + 2.7$ for $-1.35 < x < -0.7$ and $y = -3x - 0.8$ for $-0.75 < x < -0.35$ and $y = 4x - 1.65$ for $-0.35 < x < -0.11$ and $y = -11x$ for $-0.11 < x < 0$ (figure. 6d) and $y = 2.4(x + 1)$ (figure. 6f), where y and x represent $\log[N_{splashes}]$ and $\log[\Delta E]$ respectively.

equilibrium dynamics in more detail, we need to characterize the avalanches density probability, that quantifies the relative number of avalanches in the interval ΔE . For our experiment, we choose three regions (tuning the detuning $\Delta_c/2\pi$) in which the system are in low-based distribution (Fig. 6a), Bimodal distribution (Fig. 6b) and high-based distribution (Fig. 6e) respectively. We make a histogram for each distribution, and the results are given in the small block diagram in Fig. 6b, d and f. We make a plot for the number of spikes versus the interval energy ΔE to record statistic behaviour, see Fig. 6b, d and f.

Because our system is two non-equilibrium steady states with respective of high- ρ_{rr} and low- ρ_{rr} . Thus, there is obviously a cutoff effect near the position $\log[\Delta E] = 0$, and the avalanches density probability obeys the finite-size scaling ansatz in reference. [46]. The interacted Rydberg atoms would organize itself into a steady state when the atomic density is changed by tuning $\Delta_c/2\pi$. In this sense, the system exhibits a self-organized criticality. This would be very different to the Oslo model where there are no observed cutoff avalanche size but displaying a self-organized criticality [47]. In the specific situations of low- and high-based distribution (for figure. 6b and f), the scaling in the shaded areas are fitted by the power law. This power law can be taken as the evidence of self-organized criticality [48]. The different slopes of the linear functions fitted in figure. 6b (negative slope) and f (positive slope) characterize the different evolution of the system, which can be controlled by the competition between the low- ρ_{rr} and high- ρ_{rr} state. In this case, the system in the shaded areas could be con-

sidered in the low- ρ_{rr} (high- ρ_{rr}) based state which is far away from high- ρ_{rr} (low- ρ_{rr}) based state, this can be in principle understood as the particular case of finite-size model. Furthermore, if we turn the system into Bimodal situation (figure. 6c and the inset of figure. 6d), there are different evolution, exhibiting two peaks in figure. 6d.

V. DISCUSSION AND CONCLUSION

In this work, we have studied an optically driven non-equilibrium phase transition and optical bistability in a thermal vapor of Rydberg atoms. By applying the driving fields in an EIT configuration, the sensitivity in the frequency domain was enhanced by two orders of magnitude compared to previous experiments which allows to map out the phase diagram in the vicinity of the system's critical point where a continuous-order phase transitions can be observed. In particular, the observed threshold behavior for the interaction induced shift Δ' and broadening Γ' limited to the I-phase at low Ω_p suggests that the shift originates not directly from ionizing collisions between Rydberg and ground state atoms but either from avalanche ionization and plasma formation, as it has been confirmed at for stronger excitation fields [24], or dipolar interactions. Further insight on the role of dipolar interactions or avalanche ionization on the phase transition could be made in experiments that combine the sensitivity of a weak EIT-probe demonstrated here with simultaneous charge detection [22, 49, 50] or spectroscopy of a resulting plasma [24].

Independent of the exact mechanism, the many-body

nature of the resulting Rydberg interaction is the key ingredient for the phase transition and self-organization. This could enable the study of “time crystals” in periodically driven open quantum systems [51]. Besides fundamental studies, the susceptibility to noise also plays a crucial role in phase-transition enhanced sensors for electro-magnetic fields such as microwave or THz fields [23].

Our results highlight that a rich range of non-equilibrium phenomena can be studied even in a relatively simple experiment and provide observational data that helps to refine theoretical models as highlighted by the modifications needed to a standard mean-field approach. The underlying physics of self-organization dynamics observed suggest that the system could provide a controllable environment to study analogous dynamical systems from e.g. biology [52], economics [43] many-body physics in condensed matter [53]. Finally, the sensitivity of the phase to small fluctuations could be applied to the sensing of not only optical fields as demonstrated here (section 4) but also THz fields [23], system noise, microwaves, or more generally any weak fluctuations of parameters linked to the EIT laser fields or external perturbations.

ACKNOWLEDGMENTS

We are grateful to Professors Wei-Ping Zhang, Cheng Chin, Min Xiao and Wei Yi for valuable discussions. We thank Jing Qian and Yan Li for help with theory and programming. This work was supported by National Key R&D Program of China (2017YFA0304800), the National Natural Science Foundation of China (Grant Nos. 61525504, 61722510, 61435011, 11174271, 61275115, 11604322), and the Innovation Fund from CAS. CSA acknowledges financial support from EPSRC Grant Ref. No. EP/M014398/1, the EU RIA project ‘RYSQ’ project, EU-H2020-FETPROACT-2014 184 Grant No. 640378 (RYSQ) and DSTL.

APPENDIX A: Energy broadening of Rydberg state

The many-body interaction underlying the non-equilibrium dynamics may either result from pairwise dipolar interactions that effectively introduce a net polarization of the atomic medium, or ionization of multiple Rydberg atoms, i.e. avalanche ionization [36]. The non-equilibrium phase transition between high- ρ_{rr} NI- and low- ρ_{rr} I-phases requires the energy broadening of Rydberg states. We calculate the pair dipole-dipole interaction potentials for $47D_{3/2}$ which is given in Fig. S7, the Rydberg state is broadened when the Rydberg atoms are approaching. In dipole interaction picture, the local




Figure 7. (a) Potential for pairwise dipole-dipole interactions for $47D_{3/2}$ (left). (b) Stark shift for $47D_{3/2}$, $m_J = 3/2$ (right) assuming an electric amplitude $E \sim \rho_{ion}$ proportional to the mean ion density based on the corresponding mean spacing between ions. Calculated using Alkali Rydberg Calculator [54].

arractor occurs when Rydberg atomic dipoles are randomly aligned, resulting in high- ρ_{rr} state. Dipoles align if normalized rate of energy change is less than broadening. Broadening is dipole-dipole interaction at distance of closest approach C_3/r^3 , where $r = 5/9N^{-1/3}$. Scanning the system from high- ρ_{rr} state to low- ρ_{rr} state resulting in bifurcation, and vice versa. We also calculate the stark shift of Rydberg D -state against ions density, and find that there are also a broadening structure in energy diagram. In the ionization picture of the bifurcation of the phase diagram, the shifts and broadening result from ionized Rydberg atoms vanish from the excitation volume at a different rate compared to the Rydberg atoms and are not explicitly included in ρ . If the phase transition is indeed caused by ions, the model I induced in the beginning could be extended by introducing a charge population in ρ that is incoherently populated by including ionization rates in L .

Although the pair dipole-dipole interaction potentials for S state has no obvious splitting structure. This shows that the I-phase of S state is narrow than the D state, which is consistent with the experimental observations. The broadened spectra of S state in I-phase can be caused by the Rydberg atoms with distinct interaction distance and different density and so on.

APPENDIX B: Threshold-dependent modification of the decay rate

In addition to the interaction induced shift in peak transmission (Fig. 3), we also measure the width $\Delta\omega_{EIT}$ of the EIT transmission window as Ω_p and thus N_R is increased (Fig. V). For $\Omega_p^2 \geq \Omega_{p,(c)}^2 = 37 (2\pi \times \text{MHz})^2$, the non-equilibrium phase transition appears. As for $\Delta_c \rightarrow \Delta_c + \Delta'(N_R)$, we observe an increase in the I-phase scaling linearly with $N_R \propto \rho_{rr} \propto \Omega_p^2$ indicating a modified decay rate $\Gamma_r \rightarrow \Gamma_r + \Gamma'(N_R)$ above threshold.




Figure 8. Width of EIT transmission window vs. $\Omega_p^2 \propto N_R$ for the same dataset as in Fig. 3b ($n = 47$). As for the shift in peak transmission Δ^* , a linear increase can be observed for above the phase transition threshold $\Omega_{p,(c)}^2 = 37 (2\pi \times \text{MHz})^2$ as expected for a modified decay rate $\Gamma_r \rightarrow \Gamma_r + \Gamma'(N_R)$.

APPENDIX C: Measurement of response time

The transition between the two non-equilibrium phases has a characteristic response time. By assuming these two phases as two energy levels of system, we use two-level atomic system to model the instability oscillation. The enlarged collective jump given in Fig. 9(a) shows a competing dynamics of decay and recovery. We solve the master equation of a simplified two-level atomic system with decay rate of $\gamma = 9.2 \text{ kHz}$ corresponding to lifetime of $70D_{3/2}$ Rydberg state at 320 K . This response time corresponding lifetime $\tau_0 = 1/\gamma$ characterizes the minimum amount high- ρ_{rr} population to generate phase transition. The duration of interaction is slower than the natural response time, thus it appears optical instability like in Fig. 4e and Fig. 5d.

Fig. 9(b) shows an example of the change in probe transmission as Ω_p is scanned either way away from the from transition point for while $\Delta_c/2\pi \sim -15 \text{ MHz}$ remains fixed. Unlike at the critical point, the difference in transmission is not continuous. It displays a sudden fall in the transmission. A decrease in transmission with increasing intensity—a kind of self-induced opacity that is contrary to the behavior of most media, which normally exhibits self-induced transparency; this behavior is indicative of a certain type of many-body blockade effect from many-body interaction, not analogy to the character of single-body blockade effect. An interesting feature noted in Fig. 9(b) is that the noise in the transmission signal is stronger for high Ω_p , and thus higher Rydberg fractions, indicating a higher degree of instability beyond the transition to the high- ρ_{rr} phase.

Finally, we measure the width of hysteresis associated with the bistability as a function of scan frequency, the results are shown in Fig. 9(c). The reason why there is a hysteresis cycle is that the system undergoes two dissipate processes with distinct decay rates, causing energy loss in the high- ρ_{rr} state. The dependence on sweep speed implies the transition has a long relaxation time on the millisecond scale, which charac-




Figure 9. (a) Details of the instability spike from Fig. 5(d), the fitted curve based on two-level atomic system. (b) The phase transition with sweeping the Rabi frequency of probe laser, red/black line corresponds $\pm$ direction respectively. The detuning is set to be $\Delta_c/2\pi \sim -15 \text{ MHz}$. (c) Recorded hysteresis width of phase transitions dependent on the sweep frequency.

terizes an amount of high- ρ_{rr} population, supporting the structured phase transition. This is attributed to a self-organization dynamics, a non-equilibrium transition from low- ρ_{rr} non-interacted phase to high- ρ_{rr} interacted phase [1]. Alternatively, there is a saturation point occurred around 50 Hz , the corresponding width of hysteresis is $\Delta T \sim 0.48 \text{ ms}$. We thus estimate the maximum amount of attractor for our system $\Delta T/\tau_0 \approx 4.4$, where $\tau_0 = 0.11 \text{ ms}$ represents the measured natural response time of non-equilibrium phase transition from minimum high- ρ_{rr} interacted state, giving four times larger than the threshold of attractor in our case. The natural response time of non-equilibrium phase transition characterizes the amount of minimum high- ρ_{rr} population, the more the high- ρ_{rr} population, the corresponding hysteresis width is larger, thus the system is more robust to the external interference or perturbation. This timescale of τ_0 is comparable to that of avalanche ionization processes in cold Rydberg ensembles [36] and could indicate that the phase transition is induced by ionization processes.

APPENDIX D: Polarization dependence of optical bistability at $n = 47$

In addition to $n = 70$, we also investigate the influence of the relative polarization between the probe and coupling fields at a lower principle quantum number $n = 47$. At lower n , the weaker interaction-induced shifts result in lower relative shifts between the m_J -sublevels of $|r\rangle$ and thus a much weaker modification $\Gamma'(N_R)$ of the de-




Figure 10. (a) Recorded spectra for different polarization of the coupling beam ($n = 47$). Red/black line corresponds $\pm$ scan direction from red- to blue-detunings (red) and vice versa (black). θ is defined as the angle of half-wave plate inserted in the optical path of coupling beam. (b) is angular dependence of hysteresis width for $n = 47$.

cay rate. This is reflected in a much weaker dependence of the width of the bistability window on the probe and coupling polarizations (Fig. 10).

* dds@ustc.edu.cn

† drshi@ustc.edu.cn

‡ c.s.adams@durham.ac.uk

- [1] Hermann Haken, *Information and Self-Organization: A Macroscopic Approach to Complex Systems* (Springer Science & Business Media, 2006).
- [2] TC Per Bak and Kurt Wiesenfeld, “Self-organized criticality: and explanation of $1/f$ noise,” *Phys. Rev. Lett.* **59**, 381–384 (1987).
- [3] Jens Eisert, Mathis Friesdorf, and Christian Gogolin, “Quantum many-body systems out of equilibrium,” *Nature Physics* **11**, 124 (2015).
- [4] HM Gibbs, SL McCall, and TNC Venkatesan, “Differential gain and bistability using a sodium-filled fabry-perot interferometer,” *Phys. Rev. Lett.* **36**, 1135 (1976).
- [5] Hai Wang, DJ Goorskey, and Min Xiao, “Bistability and instability of three-level atoms inside an optical cavity,” *Phys. Rev. A* **65**, 011801 (2001).
- [6] HJ Carmichael and DF Walls, “Hysteresis in the spectrum for cooperative resonance fluorescence,” *Journal of Physics B: Atomic and Molecular Physics* **10**, L685 (1977).
- [7] MP Hehlen, HU Güdel, Q Shu, J Rai, S Rai, and SC Rand, “Cooperative bistability in dense, excited atomic systems,” *Phys. Rev. Lett.* **73**, 1103 (1994).

- [8] Thomas F Gallagher, *Rydberg atoms*, Vol. 3 (Cambridge University Press, 2005).
- [9] Mark Saffman, TG Walker, and Klaus Mølmer, “Quantum information with rydberg atoms,” *Reviews of Modern Physics* **82**, 2313 (2010).
- [10] N Malossi, MM Valado, S Scotto, P Huillery, P Pillet, D Ciampini, E Arimondo, and O Morsch, “Full counting statistics and phase diagram of a dissipative rydberg gas,” *Phys. Rev. Lett.* **113**, 023006 (2014).
- [11] H Schempp, G Günter, M Robert-de Saint-Vincent, CS Hofmann, D Breyel, A Komnik, DW Schönleber, Martin Gärttner, Jörg Evers, S Whitlock, *et al.*, “Full counting statistics of laser excited rydberg aggregates in a one-dimensional geometry,” *Phys. Rev. Lett.* **112**, 013002 (2014).
- [12] Cenap Ates, Thomas Pohl, Thomas Pattard, and Jan M Rost, “Antiblockade in rydberg excitation of an ultracold lattice gas,” *Phys. Rev. Lett.* **98**, 023002 (2007).
- [13] Thomas Amthor, Christian Giese, Christoph S Hofmann, and Matthias Weidemüller, “Evidence of antiblockade in an ultracold rydberg gas,” *Phys. Rev. Lett.* **104**, 013001 (2010).
- [14] Martin Gärttner, Kilian P Heeg, Thomas Gasenzer, and Jörg Evers, “Dynamic formation of rydberg aggregates at off-resonant excitation,” *Phys. Rev. A* **88**, 043410 (2013).
- [15] Igor Lesanovsky and Juan P Garrahan, “Out-of-equilibrium structures in strongly interacting rydberg gases with dissipation,” *Phys. Rev. A* **90**, 011603 (2014).
- [16] R Celistrino Teixeira, C Hermann-Avigliano, Thanh Long Nguyen, T Cantat-Moltrecht, Jean-Michel Raimond, S Haroche, S Gleyzes, and M Brune, “Microwaves probe dipole blockade and van der waals forces in a cold rydberg gas,” *Phys. Rev. Lett.* **115**, 013001 (2015).
- [17] AD Bounds, NC Jackson, RK Hanley, EM Bridge, P Huillery, and MPA Jones, “Coulomb anti-blockade in a rydberg dressed gas,” *arXiv preprint arXiv:1901.05391* (2019).
- [18] Fabian Letscher, Oliver Thomas, Thomas Niederprüm, Michael Fleischhauer, and Herwig Ott, “Bistability versus metastability in driven dissipative rydberg gases,” *Phys. Rev. X* **7**, 021020 (2017).
- [19] S Helmrich, A Arias, G Lochead, M Buchhold, S Diehl, and S Whitlock, “Observation of self-organised criticality in an ultracold atomic gas,” *arXiv preprint arXiv:1806.09931* (2018).
- [20] Christopher Carr, Ralf Ritter, CG Wade, Charles S Adams, and Kevin J Weatherill, “Nonequilibrium phase transition in a dilute rydberg ensemble,” *Phys. Rev. Lett.* **111**, 113901 (2013).
- [21] Natalia R de Melo, Christopher G Wade, Nikola Šibalić, Jorge M Kondo, Charles S Adams, and Kevin J Weatherill, “Intrinsic optical bistability in a strongly driven rydberg ensemble,” *Phys. Rev. A* **93**, 063863 (2016).
- [22] Daniel Weller, Alban Urvoy, Andy Rico, Robert Löw, and Harald Kübler, “Charge-induced optical bistability in thermal rydberg vapor,” *Phys. Rev. A* **94**, 063820 (2016).
- [23] Christopher G Wade, M Marcuzzi, E Levi, Jorge M Kondo, Igor Lesanovsky, Charles S Adams, and Kevin J Weatherill, “A terahertz-driven non-equilibrium phase transition in a room temperature atomic vapour,” *Nature communications* **9**, 3567 (2018).

- [24] Daniel Weller, James P Shaffer, Tilman Pfau, Robert Löw, and Harald Kübler, “The interplay between thermal rydberg gases and plasmas,” *arXiv preprint arXiv:1901.07850* (2019).
- [25] A Urvoy, F Ripka, Igor Lesanovsky, D Booth, JP Shaffer, T Pfau, and R Löw, “Strongly correlated growth of rydberg aggregates in a vapor cell,” *Phys. Rev. Lett.* **114**, 203002 (2015).
- [26] Tony E Lee, Hartmut Haefner, and MC Cross, “Collective quantum jumps of rydberg atoms,” *Phys. Rev. Lett.* **108**, 023602 (2012).
- [27] Matteo Marcuzzi, Emanuele Levi, Sebastian Diehl, Juan P Garrahan, and Igor Lesanovsky, “Universal nonequilibrium properties of dissipative rydberg gases,” *Phys. Rev. Lett.* **113**, 210401 (2014).
- [28] Hendrik Weimer, “Variational principle for steady states of dissipative quantum many-body systems,” *Phys. Rev. Lett.* **114**, 040402 (2015).
- [29] Nikola Šibalić, Christopher G Wade, Charles S Adams, Kevin J Weatherill, and Thomas Pohl, “Driven-dissipative many-body systems with mixed power-law interactions: Bistabilities and temperature-driven nonequilibrium phase transitions,” *Phys. Rev. A* **94**, 011401 (2016).
- [30] Michael Fleischhauer, Atac Imamoglu, and Jonathan P Marangos, “Electromagnetically induced transparency: Optics in coherent media,” *Reviews of modern physics* **77**, 633 (2005).
- [31] AK Mohapatra, TR Jackson, and CS Adams, “Coherent optical detection of highly excited rydberg states using electromagnetically induced transparency,” *Phys. Rev. Lett.* **98**, 113003 (2007).
- [32] Eric Bonabeau, Directeur de Recherches Du Fnrs Marco, Marco Dorigo, Guy Théraulaz, Guy Theraulaz, *et al.*, *Swarm intelligence: from natural to artificial systems*, 1 (Oxford university press, 1999).
- [33] G Vitrant, J M Raimond, M Gross, and S Haroche, “Rydberg to plasma evolution in a dense gas of very excited atoms,” *Journal of Physics B: Atomic and Molecular Physics* **15**, L49–L55 (1982).
- [34] T. C. Killian, S. Kulin, S. D. Bergeson, L. A. Orozco, C. Orzel, and S. L. Rolston, “Creation of an ultracold neutral plasma,” *Phys. Rev. Lett.* **83**, 4776–4779 (1999).
- [35] M. P. Robinson, B. Laburthe Tolra, Michael W. Noel, T. F. Gallagher, and P. Pillet, “Spontaneous evolution of rydberg atoms into an ultracold plasma,” *Phys. Rev. Lett.* **85**, 4466–4469 (2000).
- [36] M Robert-de Saint-Vincent, CS Hofmann, H Schempp, G Günter, S Whitlock, and M Weidemüller, “Spontaneous avalanche ionization of a strongly blockaded rydberg gas,” *Phys. Rev. Lett.* **110**, 045004 (2013).
- [37] Da-wei Wang, Zheng-hong Li, Hang Zheng, and Shi-yao Zhu, “Time evolution, lamb shift, and emission spectra of spontaneous emission of two identical atoms,” *Phys. Rev. A* **81**, 043819 (2010).
- [38] James Keaveney, *Collective Atom–Light Interactions in Dense Atomic Vapours* (Springer, 2014).
- [39] A Reinhard, T Cubel Liebisch, KC Younge, PR Berman, and G Raithel, “Rydberg-rydberg collisions: Resonant enhancement of state mixing and penning ionization,” *Phys. Rev. Lett.* **100**, 123007 (2008).
- [40] V. L. Ginzburg and L. D. Landau, “On the theory of superconductivity,” *J. Exp.Theor. Phys.* **20**, 1064–1082 (1950).
- [41] G Günter, H Schempp, M Robert-de Saint-Vincent, V Gavryusev, S Helmrich, CS Hofmann, S Whitlock, and M Weidemüller, “Observing the dynamics of dipole-mediated energy transport by interaction-enhanced imaging,” *Science* **342**, 954–956 (2013).
- [42] Gregory S Engel, Tessa R Calhoun, Elizabeth L Read, Tae-Kyu Ahn, Tomáš Mančal, Yuan-Chung Cheng, Robert E Blankenship, and Graham R Fleming, “Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems,” *Nature* **446**, 782–786 (2007).
- [43] Antonio Majdandzic, Boris Podobnik, Sergey V Buldyrev, Dror Y Kenett, Shlomo Havlin, and H Eugene Stanley, “Spontaneous recovery in dynamical networks,” *Nature Physics* **10**, 34 (2014).
- [44] Laurie F Thomas and Sheila Harri-Augstein, *Self-organised learning: Foundations of a conversational science for psychology* (BookBaby, 2013).
- [45] JA Scott Kelso, *Dynamic patterns: The self-organization of brain and behavior* (MIT press, 1997).
- [46] Kim Christensen and Nicholas R Moloney, *Complexity and criticality*, Vol. 1 (World Scientific Publishing Company, 2005).
- [47] Vidar Frette, Kim Christensen, Anders Malthes-Sørensen, Jens Feder, Torstein Jøssang, and Paul Meakin, “Avalanche dynamics in a pile of rice,” *Nature* **379**, 49 (1996).
- [48] Per Bak and Kan Chen, “Self-organized criticality,” *Scientific American* **264**, 46–53 (1991).
- [49] Ryan K Hanley, Alistair D Bounds, Paul Huillery, Niamh C Keegan, Riccardo Faoro, Elizabeth M Bridge, Kevin J Weatherill, and Matthew PA Jones, “Probing interactions of thermal sr rydberg atoms using simultaneous optical and ion detection,” *Journal of Physics B: Atomic, Molecular and Optical Physics* **50**, 115002 (2017).
- [50] D. Barredo, H. Kübler, R. Daschner, R. Löw, and T. Pfau, “Electrical readout for coherent phenomena involving rydberg atoms in thermal vapor cells,” *Phys. Rev. Lett.* **110**, 123002 (2013).
- [51] FM Gambetta, F Carollo, M Marcuzzi, JP Garrahan, and I Lesanovsky, “Discrete time crystals in the absence of manifest symmetries or disorder in open quantum systems,” *arXiv preprint arXiv:1807.10161* (2018).
- [52] Tobias Grafke, Michael E Cates, and Eric Vanden-Eijnden, “Spatiotemporal self-organization of fluctuating bacterial colonies,” *Phys. Rev. Lett.* **119**, 188003 (2017).
- [53] George M Whitesides and Bartosz Grzybowski, “Self-assembly at all scales,” *Science* **295**, 2418–2421 (2002).
- [54] Nikola Šibalić, Jonathan D Pritchard, Charles S Adams, and Kevin J Weatherill, “Arc: An open-source library for calculating properties of alkali rydberg atoms,” *Computer Physics Communications* **220**, 319–331 (2017).