Unification in One Dimension

David J. Jackson

February 15, 2016

Abstract

A physical theory of the world is presented under the unifying principle that all of nature is laid out before us and experienced through the passage of time. The onedimensional progression in time is opened out into a multi-dimensional mathematically consistent flow, with the simplicity of the former giving rise to symmetries of the latter. The act of perception identifies an extended spacetime arena of intermediate dimension, incorporating the symmetry of geometric spatial rotations, against which physical objects are formed and observed. The spacetime symmetry is contained as a subgroup of, and provides a natural breaking mechanism for, the higher general symmetry of time. It will be described how the world of gravitation and cosmology, as well as quantum theory and particle physics, arises from these considerations.

Contents

Chapter 1

Introduction

In establishing a conceptual framework for a physical theory of the world one of the most fundamental questions concerns the nature of the ultimate entity out of which the world is made. From the earth, water, air and fire of the ancient Greeks, through various manifestations of elementary extended or point-like particle theories, to the quantum fields of $20th$ century high energy physics, the notion of a fundamental form of matter behaving according to laws of nature, to be identified empirically or through powers of reason, has strongly influenced the development of scientific theories of the world. The general trend has been to dig deeper into the layers of matter such that macroscopic objects are taken to be composed of discrete particle-like entities which in turn are composed of more basic particles which have become themselves to be considered as merely the states detected in physics experiments as a manifestation of yet deeper underlying entities, such as fields or strings. It is a trend which ever poses the question of what may be uncovered at the next layer down, or whether we may reach the ultimate bedrock of the world.

The view taken in the present investigations is that the world can be built out of the one entity within which all our experiments, experiences, perceptions and indeed our thoughts in general are conducted, that is through progression in time. This universal nature of time applies both to inner thought experiences in the mind, as well as outer thoughts of the physical world; for example, perception of a cloud passing by or of a book on the table. With the basic structure of time being identified with, or isomorphic to, that of the real numbers $\mathbb R$ this gives an immediate connection to a purely mathematical world. The mathematical possibility to express an inner one-dimensional sense of time in the form of an outer multi-dimensional space as an intrinsic and elementary property of the real numbers provides a significant motivation for this study. The aim will be to demonstrate how the external world of experience can result directly from the mathematical structure of temporal flow itself without the need to interpose or postulate the notion of an underlying 'material' substratum of any form.

It may be helpful to begin with an analogy using a familiar example in which multi-dimensional structures are enfolded within a lower-dimensional entity, namely a child's 'pop-up' book of cartoon zoo animals, although, of course, it should not be taken too literally to represent the theory to be presented here. We can consider such a book, when closed, to be an essentially 2-dimensional object in space. When opened fully on a given page a figure will 'pop-up', perhaps an elephant, extended in 3-dimensional space; on another page a 3-dimensional crocodile may appear, and so on. It is down to the creativity and origami skills of the bookmakers to form such 3 dimensional structures that can be perfectly folded away into the 2-dimensional plane (when the book closes) within the fixed constraints of the possibilities allowed by the laws of Euclidean geometry.

It is the contention of these investigations that the 1-dimensional flow of time itself naturally opens out, according to necessary mathematical and geometrical forms, into a higher-dimensional space. The mechanism will be somewhat different to that in the above metaphor since time, unlike a book in space, is not experienced 'all at once' and indeed space itself will need to be unfolded out of the temporal flow. However, within the 1-dimensional flow of time we shall find implicitly contained not only the appropriate mathematical structures for 3-dimensional space and 4-dimensional spacetime but also still higher-dimensional possibilities. The intermediate 3 and 4 dimensional cases can be interpreted as subspaces of the higher-dimensional forms, with the properties of physical objects perceived in spacetime being largely determined by the nature of the general higher-dimensional structures. It is claimed that the opening out of the progression of time in this way into a mathematically determined multi-dimensional flow is responsible for not only our perceptions of objects, from books to real elephants themselves, but of the entire physical universe around us.

In a similar way that the laws of geometry constrain the design of pop-up books, so mathematics will constrain the way in which the physical world can open out from the flow of time and hence determine the laws of physics. It is the main aim of this paper to show how far the consequences of this idea resemble the observed laws of nature of the actual world. In traditional theories properties are assigned to underlying particles or fields, out of an enormous range of conceivable choices of such properties, largely for pragmatic reasons to match the empirically observed world. Here, on the contrary, we expect the present theory to make a much more thorough and direct contact with the structure of the physical world.

In aiming for an inclusive and complete theory, as well as accounting for the basic observed scientific phenomena from particle physics to cosmology, the theory might also address the everyday direct manner through which we actually encounter and experience the world. We shall touch upon all these areas, all of which would benefit from further study, in an attempt to gain an overall consistent worldview.

Although this paper is lengthy all of the contents relate to a single unified theory, rather than to a collection of independent ideas, as will be clear from the progression of sections and the mutual cross-referencing within the text. Here we review the contents of the paper to guide the reader towards the sections which may be of most interest. While the overall order of the text has been designed to introduce the various facets of the theory in a reasonably logical sequence there are four main areas in which progress on the theory has been made essentially in parallel. Each area addresses a particular question and related set of issues which might be asked of any candidate for a unified physical theory. The four areas correspond generally to subsets of the subsequent chapters of this paper:

- Chapters 2–5: The main goal here is to describe how an extended spacetime arena may be identified together with an external and internal curvature and the manner in which they are mutually constrained. First, beginning with a onedimensional temporal interval in chapter 2, we make precise the notion of the multi-dimensional flow of time by deriving in elementary terms what is considered to be its general mathematical form and give several relevant examples. We also emphasise the fact that we are dealing here with a general symmetry of time, in contrast to the symmetry of a higher-dimensional spacetime found in a different class of theories. The notion of perception as identified with the interpretation of certain mathematical forms, implicit in the multi-dimensional flow of time, in the shape of a geometrical spacetime. That is, we describe how an extended external spacetime arena for the world can 'pop-out' from the temporal flow. This structure motivates the employment of more sophisticated mathematical tools, and in chapter 3 standard textbook material on differential geometry and general relativity is reviewed. Papers in the literature from around the mid-1970s to mid-1980s regarding non-Abelian Kaluza-Klein theories, which also describe a unified approach relating external and internal curvature within a similar mathematical framework, are then reviewed in chapter 4. In chapter 5 we then pick up the thread from end of chapter 2, in light of the mediating chapters, using the constraints of the present theory to study the relationship between the external and internal curvature in 4-dimensional spacetime and consider further implications such as constraints on the equations of motion.
- Chapters 6–9: Here we consider higher-dimensional forms of temporal flow with the main aim of establishing a connection with the structures of the Standard Model of particle physics in the breaking of the full symmetry of time over the 4-dimensional base manifold. Crucial to this investigation are the references concerning the structure of the Lie group E_6 acting on the space h₃ σ of 3×3 Hermitian octonion matrices which have been published within the past ten years. The relevant details from this literature are followed closely and reviewed in chapter 6 in the context of the present theory, in particular with the determinant preserving action of E_6 on the elements of $h_3\mathbb{O}$ interpreted as a particularly rich symmetry of a 27-dimensional cubic form of temporal flow. In chapter 7 the principle features of the Standard Model, and their relation to mathematical models based on unification groups, are reviewed in order describe the physical structures to be accounted for by the present theory and the kind of theoretical structures which may be relevant. In chapter 8 we investigate the extent to which the E_6 symmetry action on $h_3\mathbb{O}$ in the context of the present theory can account for the properties of the Standard Model. Several successes are noted in terms of a correlation between the transformation properties of components of $h_3\mathbb{O}$ under the external and identified internal symmetries and corresponding properties of Standard Model particle states. The need to incorporate further particle properties and the natural extension to the larger structure of an E⁷ action preserving a quartic from on the space $F(h_3\mathbb{O})$, interpreted as a symmetry of a 56-dimensional form of temporal flow, leads to some further success in chapter 9 and also the suggestion of investigating yet higher-dimensional forms.
- Chapters 10–11: In addition to uncovering Standard Model features here the ambition is to understand how the present theory might accommodate the empirical observations of HEP experiments, in terms of cross-sections and decay rates for example, and incorporate quantum phenomena in general. The essential textbook aspects of quantum field theory are reviewed in chapter 10, with in particular the structure of cross-section calculations analysed into its basic elements in order to establish a correspondence with the present theory. This correspondence is described in chapter 11 in which the conceptual origins of quantum phenomena within the context of the present theory are established. In one sense this involves generalising the relation between the external geometry and a particular internal gauge field, as assessed in relation to Kaluza-Klein theories in the earlier chapters, for an external geometry expressed in terms of a degeneracy of underlying field solutions. These underlying fields include both gauge fields deriving from the symmetry of time and fields deriving from components of the multi-dimensional form of temporal flow itself, mutually related by a set of implicit constraint equations rather than via a dedicated Lagrangian function. Within the scope of these investigations two main and related points concern the conceptual nature of physical particle states, as analysed in laboratory experiments, and the manner in which the phenomena of quantum theory and general relativity coexist as aspects of the same unified theory.
- Chapters 12–14: The theory has been developed with not only laboratory phenomena in mind but also the large scale structure of the universe with the goal of understanding the extent to which observations in cosmology might also be accounted for. To this end in chapter 12 textbook material on both the standard cosmological model and inflationary theory is reviewed. A new feature of the present theory is described in chapter 13 concerning the possibility of non-trivial intrinsic curvature for the spacetime manifold arising from the elementary properties of its projection out of the multi-dimensional form of time. The extent to which this, combined with additional features of the theory identified earlier, might account for both the phenomena of the dark sector in cosmology and the structure of the very early universe is then considered, with the question of uniqueness for the theory in general also discussed. The study of the Big Bang epoch raises broader questions, in addition to the need to describe physical properties, relating to the reason why the universe should exist at all. In the context of the present theory, with everything constructed through a multi-dimensional form of progression in time, this inevitably leads to the question concerning the origin of time itself. In addressing this issue the speculations of chapter 14 include areas which are not necessarily within the traditional bounds of physics but touch upon other scientific fields of study. This detour is however of value in providing an opportunity to elaborate upon the possibility of identifying a firm foundation for the full physical theory.

Following the discussion of the foundations of the theory in chapter 14 in the concluding chapter 15 we look outwards to the prospects for the further development of the theory in the four main areas outline above, which are also depicted as the four fronts in figure 15.1. In section 15.1 the mutual relations between all aspects of this unified theory will also be described.

Broadly, the four subsets of chapters for the four main branches of the theory listed above are each presented with a structure to some extent analogous to a PhD thesis, in terms of combination of the presentation of new ideas and reviews of established material. In particular summaries of standard textbook material and other cited literature are presented mainly in the subsections, sections and chapters: 2.2.2, 3, 4, 6, 7, 9.2 (up to equation 9.41), 10 and 12, although always in the context of the present theory. On the other hand the main novel theoretical developments follow a trail through the sections and chapters: 2, 5, 8, 9, 10.1, 11, 12.1, 13, 14 and 15, with reference to the standard material of the above intervening chapters and section and further citations discussed in the course of the presentation.

Generally speaking the central chapters 6–11 deal more with the microscopic and laboratory scale while the outer sections through to chapter 13 pursue a thread more closely associated with the macroscopic and large scale features typically studied under general relativity. However all aspects of the present theory are relevant for all scales, as further discussed in the concluding chapter. The current point of closest approach between the present theory and empirically established features of the physical world is in terms of a relationship with the Standard Model of particle physics, regarding in particular transformation properties under the external Lorentz symmetry group and the internal $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge group, as alluded to in the synopsis of Chapters 6–9 listed above. The shortest path from introducing the basic ideas of the theory to an elaboration of this connection with the Standard Model is to follow sections: 2.1, the opening of 2.2, 8.1, 8.2, 9.2 (after equation 9.41) and with further discussion in section 9.3.

The purpose of these investigations can be described as an enquiry into the extent to which the form of the physical world can be determined purely from the fact that it is perceived in time; that is, the extent to which the world can be constructed out of the pure mathematical nature of the progression in time itself. To this end we establish in the following section an expression for the potential multi-dimensional flow of time in the appropriate general mathematical form. This will later provide the means to incorporate 4-dimensional spacetime together with the structures of 'extra dimensions' in a naturally unified way. We begin with a particular example of a multidimensional expression implicit in a finite interval of time which exhibits an apparent geometric symmetry.

Chapter 2

The Symmetry of Time

2.1 General Form of Temporal Flow

A finite interval of time represented by the real number $s \in \mathbb{R}$ can be algebraically expressed in terms of other real numbers x^a $(a = 1, 2, 3...)$ in an endless variety of ways. For example $s = x^1 + x^2$ composes time intervals in series, while $s = x^1x^2$ might represent a rescaling of the temporal unit, or more generally we can have $s =$ $x^1(x^2x^3 + x^4)$ and so on simply by employing the basic arithmetic structure of the real line. More specifically, writing the square of the interval in the form, familiar since Pythagoras, $s^2 = (x^1)^2 + (x^2)^2 + (x^3)^2$, the interval s is then invariant under transformations of the set of real numbers $\{x^1, x^2, x^3\} \in \mathbb{R}^3$ by the orthogonal rotation group $O(3)$, as depicted in figure 2.1. In this case the set of possible numbers $\{x^a\}$ consistent with this form for s exhibits the mathematical symmetry of a vector in 3-dimensional Euclidean space that maintains a fixed length under rotations. This is exactly the mathematical structure within which the physical objects of our perceptions appear to us spatially; thus providing a simple example of how the geometric properties of space can be algebraically embedded within structures implicit in the arithmetic properties of time as represented by the real line.

The broad range of possible expressions for a finite interval s in terms of an arbitrary number of variables $\{x^a\}$, $a = 1...n$, will be constrained to a more restrictive structure in the limit of infinitesimally small temporal intervals. We first consider this limit for the trivial case with the flow of time s expressed in terms of a single real variable x^1 only for which we have simply $s = x^1$. This can symbolically be written as $\delta s = \delta x^1$ as we approach the limit of infinitesimal intervals. We then express the rate of change of x with respect to s in this limit as:

$$
v^1 = \frac{dx^1}{ds} \equiv \frac{\delta x^1}{\delta s} \bigg|_{\delta s \to 0} = 1 \tag{2.1}
$$

For the case with multiple real numbers $\{x^a\} \in \mathbb{R}^n$ representing the flow of time s each will be associated with a corresponding rate of change $v^a = dx^a/ds$ with respect

Figure 2.1: With a fixed finite 1-dimensional interval of time s expressed in the form $s = \sqrt{(x^1)^2 + (x^2)^2 + (x^3)^2}$ the *numerical* morphisms of the three numbers ${x^1, x^2, x^3}$ under which s is invariant can be interpreted as mapping out a spherical shell in a 3-dimensional geometrical space.

to pure time. For example, we may consider the propagation of time expressed for an infinitesimal interval as:

$$
(\delta s)^2 = (\delta x^1)^2 + (\delta x^2)^2 + (\delta x^3)^2 \tag{2.2}
$$

$$
= \eta_{ab}\delta x^a \delta x^b \qquad \text{with} \quad \eta_{ab} = \text{diag}(+1, +1, +1) \tag{2.3}
$$

where $\{a, b\} = \{1, 2, 3\}$ (and with the conventional summation over repeated indices implied throughout this paper). Dividing by $(\delta s)^2$ and taking the limit $\delta s \to 0$ this can be written as $\eta_{ab}v^a v^b = 1$ or $(v^1)^2 + (v^2)^2 + (v^3)^2 = 1$, which is invariant under the group, $O(3)$, of orthogonal transformations in three dimensions applied to $v_3 =$ $(v^1, v^2, v^3) \in \mathbb{R}^3$. This is simply the infinitesimal case of the situation depicted in figure 2.1, with $s \to \delta s/\delta s = 1$ and $x^a \to \delta x^a/\delta s = v^a$ (for $\delta s \to 0$), and is again open to a similar Euclidean spatial interpretation, here for the components $\{v^a\} \in \mathbb{R}^3$. The question is then how to express the general case for the composition and symmetries of a multi-dimensional set of velocities $\{v^a\} \in \mathbb{R}^n$.

The infinitesimal elements of time can be written most generally, taking care to balance the order of the vanishing elements in each term, as:

$$
\delta s = \alpha_a \delta x^a + \sqrt{\alpha_{bc} \delta x^b \delta x^c} + \sqrt[3]{\alpha_{def} \delta x^d \delta x^e \delta x^f} + \dots \tag{2.4}
$$

Here the coefficients $\alpha_{abc...}$ are each equal to ± 1 or 0 since we wish to express the δs purely in terms of simple arithmetic relations of the δx^a . In equation 2.4 each term divides δs into a separate portion of time:

$$
\delta s = \delta s_1 + \delta s_2 + \delta s_3 + \dots \qquad (2.5)
$$

where each term δs_p is the p^{th} -root of a homogeneous polynomial of order p in the $\{\delta x^a\}$. Taking each term in turn, dividing by the interval δs_p in each case and taking the limit $\{\delta s_p, \delta x^a\} \rightarrow 0$ we find:

$$
\delta s_p = \sqrt[p]{\alpha_{abc...}\delta x^a \delta x^b \delta x^c \dots} \tag{2.6}
$$

divide by
$$
\delta s_p
$$
:
$$
1 = \sqrt[p]{\alpha_{abc...} v^a v^b v^c \dots}
$$
 (2.7)

that is:
$$
\alpha_{abc...}v^a v^b v^c \dots = 1 \tag{2.8}
$$

which we write:
$$
L(v) = 1
$$
 (2.9)

where L is a homogeneous polynomial of order p in the components v^a ; it can be considered as a map from the elements of a real *n*-dimensional vector space $v \in \mathbb{R}^n$ onto the unit $1 \in \mathbb{R}$.

The pth -root is dropped in stepping from equation 2.7 to equation 2.8 since, trivially, $1^p = 1$. If the equality in equation 2.7 involved a variable quantity on the left-hand side rather than unity, such as in the case of finding a 'path of extremal length' on an extended manifold for a quadratic form, or metric, using a variational method then the root would be needed, as will be described later for equation 3.78. Further, the components of a local 'metric' $\eta_{ab} = \alpha_{ab} \in {\pm 1, 0}$ may be mapped onto a general metric involving components $g_{\mu\nu}(x) \notin {\pm 1, 0}$ under a transformation from 'local coordinate' variables $\{x^a\}$ to a 'general coordinate system' on such an extended manifold, as we shall describe leading up equation 2.16. (In principle this observation could also apply to the other coefficients $\alpha_{abc...}$ of equation 2.4 considered as generalised 'metrics' for the corresponding extended dimensions).

Equation 2.9 is taken to express the general mathematical form of multidimensional temporal flow and it is the central equation of this paper. The symmetries of $L(v) = 1$ will be represented by groups acting on the vector space \mathbb{R}^n such that for all elements g of the group G and all vectors $v \in \mathbb{R}^n$ satisfying $L(v) = 1$ we have $L(\sigma_g(\boldsymbol{v})) = L(\boldsymbol{v}') = 1$ where $\sigma_g(\boldsymbol{v})$ represents the action of the group element $g \in G$ on the vector $v \in \mathbb{R}^n$. As G acts on \mathbb{R}^n over a continuous range of elements beginning at the identity $q = e \in G$ we can think of this as a continuous morphism of the real numbers v^a . (This is equivalent to the symmetry over the 2-sphere in the example with finite intervals $\{x^a\}$ in figure 2.1). This morphism is always consistent with the dissolving of the fundamental temporal flow s into the possible rates of change v^a of the multi-dimensional real quantities x^a conforming to the requirement $L(v) = 1$ and hence may be termed an isochronal symmetry, of which we next describe several examples.

Quadratic forms in general, including the 4-dimensional example of the expression $L(v) = \eta_{ab}v^av^b$, with $v \in \mathbb{R}^4$, Minkowski metric $\eta_{ab} = \text{diag}(+1, -1, -1, -1)$ and $\{a, b\} = \{0, 1, 2, 3\}$, and the norm of an element of a division algebra ($\mathbb{R}, \mathbb{C}, \mathbb{H}$) or O as introduced below), together with their symmetry groups, are expected to be particularly significant forms of $L(v) = 1$. This is due to their close relation to Clifford algebras and Euclidean spatial geometry, describing for example the space within which we perceive objects. Other possible forms of $L(\mathbf{v}) = 1$ include the determinants of matrices, which are homogeneous polynomials in the matrix elements.

The complex numbers $\mathbb C$ had been studied by Hamilton in the 1830s in a manner consistent with his view of algebra as the science of pure time. This program in part led to his discovery of the quaternions in the 1840s, which also however led him to essentially abandon the notion of a close relation between algebra and time owing to the non-commutative property of the quaternion algebra. Subsequently an 8-dimensional algebra, the octonions O, was discovered independently by Graves and Cayley in the mid-1840s and completed the unique series, $\mathbb{R}, \mathbb{C}, \mathbb{H}$ and $\mathbb{O},$ of normed division algebras [1], as will be reviewed in section 6.2. In fact division algebras only exist over vector spaces of dimension 1, 2, 4 or 8. An algebra A is a division algebra if $ab = 0$ implies $a = 0$ or $b = 0$, with $a, b \in A$; it is a normed division algebra if A is also a normed vector space with $|ab| = |a||b|$. This latter property naturally provides a source of structures of form of equation 2.9 together with the corresponding symmetries.

For example, the quaternion algebra H may be used to compose a possible multi-dimensional form of progression in time. On the space of unit norm elements $v \in \mathbb{H}$, with $L(v) = |v| = 1$, the symmetry group G composed of quaternions of unit norm operating on v under left and right algebra multiplication forms the two-to-one cover of SO(4). The 1-dimensional character of temporal flow is represented by the 'norm' function L applied collectively to the components of $v \in \mathbb{H} \equiv \mathbb{R}^4$; with the noncommutative behaviour of the σ_g symmetry operations within $L(\ldots \sigma_{g''}\sigma_{g'}\sigma_g(v))=1$ describing the properties of the multiple apparently 'internal' temporal dimensions. For the case in which G is homomorphic to an orthonormal rotation group (as is the case for H representing three or four dimensional space, with for example the three imaginary units of the quaternions associated with 3-dimensional Euclidean space) the non-commutative algebraic properties correlate directly with the non-commutative property of *spatial* rotations for $n > 2$.

The fourth division algebra, the octonions O, being non-associative, do not themselves form a group in such a direct way as for the complex numbers or the quaternions; they will however play a significant role in the symmetry of time and hence in physics as will be explained in this paper. Here the division algebras will be combined with matrix algebras in considering the 27-dimensional real vector space of 3×3 Hermitian matrices h₃^① over the octonions with the determinant required to be unity: $L(\mathbf{v}_{27}) = \det(\mathcal{X}) = 1$, with $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$. The group G of determinant preserving symmetry transformations on $h_3\mathbb{O}$ is the exceptional Lie group E₆. This group is well known to be of interest for unification models and will be discussed in detail in the context of the present investigations in chapters 6–8.

With various different forms of progression in time to be considered, in general the subscript *n* in the notation $L(v_n) = 1$ indicates collectively the vector space \mathbb{R}^n , the implied form L and the corresponding symmetry group G (respectively $v_{27} \equiv \mathcal{X} \in$ $h_3\mathbb{O} \equiv \mathbb{R}^{27}$, $L(\boldsymbol{v}_{27}) = \det(\mathcal{X}) = 1$ and $G = E_6$ in the above example for $n = 27$), where any case of ambiguity will be clarified in the text.

Given a possible *n*-dimensional form of progression in time, $L(\mathbf{v}_n) = 1$, the vector $v_n \in \mathbb{R}^n$ may be written as the ordered set of velocities:

$$
\mathbf{v}_n = \{v^1, v^2, \dots, v^n\} \tag{2.10}
$$

$$
= \left\{ \frac{dx^1}{ds}, \frac{dx^2}{ds}, \dots, \frac{dx^n}{ds} \right\} \tag{2.11}
$$

the values of which are unchanged by a numerical translation of the real variables,

$$
x^a \to x^a + r^a \tag{2.12}
$$

for any constant set $\{r^a\} = r_n \in \mathbb{R}^n$, or for a subset of \mathbb{R}^n . Above we described a possible symmetry of $L(v) = 1$ with the action of a group G mixing the numerical components v^a , which represent elements of the temporal flow dx^a/ds . Here we have a further symmetry implicit in $L(v) = 1$ with respect to translations of the numerical variables as $x^a \to x^a + r^a$. That is, we also have trivially:

$$
\boldsymbol{v}_n = \left\{ \frac{d(x^1+r^1)}{ds}, \frac{d(x^2+r^2)}{ds}, \dots, \frac{d(x^n+r^n)}{ds} \right\}.
$$
 (2.13)

satisfying $L(\mathbf{v}_n) = 1$. For the 1-dimensional case of equation 2.1 the symmetry $v^1 =$ $\frac{d(x^1+r^1)}{ds}$ can be readily visualised as a flow v^1 present everywhere on the real line parametrised by $r^1 \in \mathbb{R}$ (rather than at a single arbitrary point for example). In the general case since equation 2.13 is equally valid for all possible $r_n \in \mathbb{R}^n$ the temporal flow, under the condition $L(\mathbf{v}_n) = 1$, effectively occupies the entire \mathbb{R}^n manifold as depicted in figure 2.2.

Figure 2.2: Since the real variables $\{x^a\} \in \mathbb{R}^n$ are arbitrary, the flow described within $L(\mathbf{v}_n) = 1$ applies equally for the particular value $\mathbf{x}_0 \in \mathbb{R}^n$ as for $\mathbf{x}' = \mathbf{x}_0 + \mathbf{r}_n$ and over the range $-\infty < r^a > \infty$, for $a = 1 ... n$. This 'translation' symmetry is implied within the form $L(\mathbf{v}_n) = 1$ and is depicted here for $n = 3$.

This *n*-dimensional freedom in \mathbb{R}^n forms a continuous *n*-dimensional parameter space, which may be considered to form an implicit 'base manifold' M_n , upon which the vector v_n naturally resides in the tangent space T_xM_n at every point $x \in M_n$. Hence the internal structure of the form $L(\mathbf{v}_n) = 1$ and its symmetries contain the skeletal form of a mathematical framework for the description of an apparently external and extended spatial structure.

In other theories and models a higher-dimensional *symmetry of spacetime* is considered, extending beyond our familiar 4-dimensional spacetime arena to one with a total of, for example, five or ten spacetime dimensions. Such models, initiated by Kaluza and Klein, will be described in more detail in chapter 4. In these theories it is necessary to explain how our 4-dimensional spacetime world is embedded in the larger arena, and the means by which the 'extra dimensions' are compactified or otherwise evade direct observation.

As described in the introductory chapter we are familiar with the idea that not only all of our scientific experiments but also everything we experience in the world takes place in time. Relative to 4-dimensional spacetime the flow of pure time is an apparently 'lower-dimensional' structure which pervades all observations and events in the universe. This is in contrast to hypothetical extra dimensions, above the four of space and time, which are beyond our domain of experience. Here we begin on a firm footing by treating one-dimensional temporal flow as the fundamental entity of the world.

Hence, in contrast with Kaluza-Klein theory, for the theory presented in this paper we deal instead with a general higher-dimensional symmetry of time, and it is here necessary to explain how the large scale extended 4-dimensional spacetime geometry and physical structures of the universe can arise from a fundamentally 1-dimensional temporal flow. This phenomenon, and the internal mathematical identifications that give rise to it, will intimately involve the nature of perception. It is the means through which time experienced as a purely 1-dimensional progression can be experienced simultaneously as a multi-dimensional flow of physical objects in an extended spacetime. The mathematical basis for obtaining such an extended base manifold will be found in the application of the symmetry described in figure 2.2 to a 4-dimensional spacetime subset of the translational degrees of freedom of a higher-dimensional temporal form.

For the case considered for the real world, in addition to the 27-dimensional space h_3 ^O described above another important example of a form of time involving both a matrix and a division algebra is identified in the determinant of elements of the 4-dimensional real vector space $h_2 \mathbb{C} \equiv \mathbb{R}^4$, that is the 2 × 2 Hermitian matrices over the complex numbers, together with the action of the determinant preserving group $SL(2,\mathbb{C})$. This group is the double cover of the Lorentz group and will also be significant in this paper since $h_2\mathbb{C}$ is naturally embedded as a subspace of $h_3\mathbb{O}$, with the symmetry group $SL(2, \mathbb{C})$ being a subgroup of E_6 .

Applying the translation symmetry of equation 2.13 in four dimensions only, corresponding to the $h_2\mathbb{C}$ components, provides a natural mechanism for breaking the symmetry of the larger group through the necessary identification of a 4-dimensional background manifold M_4 upon which the Lorentz group acts locally, and to a good approximation globally over extended regions of spacetime. Under the overall normalisation $L(\mathbf{v}_{27}) = 1$ the 4-dimensional form will take more general values $L(\mathbf{v}_4) = h^2 \in \mathbb{R}$ for the subcomponent $v_4 \,\subset v_{27}$ local tangent vectors on M_4 (in this paper the relation $v' \subset v$ between two vectors will denote the *projection* of v' out of v). Further consequences of the symmetry breaking are associated with the necessary choice of a particular direction for the vector field $v_4(x) \in h_2\mathbb{C}$, locally a 1-dimensional flow embedded within a 4-dimensional manifold. Comparisons between these symmetry breaking structures and the Standard Model of particle physics will be made in chapter 8.

The relation between the 'translation symmetry' of $L(\mathbf{v}) = 1$ and the 'rotation' symmetry', more generally denoted by the action $\sigma_q(v)$ for $g \in G$, is key to the development of the geometrical structure of the theory and motivates the review of elements of textbook geometry in chapter 3. We begin in the following two sections by describing a simple model universe, based on a small number of dimensions in order to elaborate upon the nature of the geometric structures involved, in particular concerning the identification of the base manifold. The geometric properties of this manifold, which are significant in general relativity, are intrinsically related to the geometry and symmetries of the residual dimensions, which are significant for gauge theories, resulting from the projection of a higher-dimensional form of $L(\mathbf{v}) = 1$ over the base manifold and corresponding symmetry breaking pattern, as will be described in section 2.3. This development of the theory will be continued in chapter 5 where the relation between the external gravitational field and internal gauge fields over a 4 dimensional spacetime manifold in the context of the present theory will be described.

2.2 Perception in Space and Time

The fact that all of our experiences in the world are encompassed within the passage of time motivated the formulation of the general expression for temporal progression, $L(v) = 1$, presented as equation 2.9 of the previous section. However it is also necessary to account for the fact that all of our experiences of such a physical world appear to be distributed through an extended *manifold*, with the immediate and necessary location of observed physical objects in space, as well as in time. While the general mathematical form for the flow of time may be exemplified by a wide range of mathematical structures and symmetry groups it is the identification of relatively simple structures, those which may be most readily suited to the organisation and understanding of experiences in the world with respect to a background arena of space as well as time, that will be designated by the term perception.

The apparent physical form of the world is shaped out of the interplay between these two basic notions: that of the mathematical form of temporal flow and that of a necessary form of perception. It is the act of interpreting *algebraic* structures within the temporal flow v in terms of an extended coherent *geometrical* structure that breaks the symmetry of the general flow of time described by $L(\mathbf{v}) = 1$.

In this and the following section the discussion will be maintained largely at a general level with a simplified model universe, a world with two spatial dimensions only, being used to make the presentation more concrete for a case which is mathematically simpler than our own world and, in particular, one which may be more readily visualised. The notion of a base manifold may be introduced by considering how it would be possible for physical objects in a spatially 2-dimensional world to be perceived propagating in time. This situation brings to mind the image pictured in figure 2.3. (Such illustrations clearly also serve by analogy to represent our own world, with one spatial dimension being suppressed. Indeed, throughout this chapter the model universe described should be considered both as a metaphor for the general case and for our own world in particular).

Objects in such a world are here depicted by figures in a 2-dimensional plane which are animated, presumably according to certain laws of physics in the form of equations of motion, as they propagate through the third dimension on a 3-dimensional base manifold M_3 . The geometrical structure of the 2-dimensional plane may be con-

Figure 2.3: A representation of a model universe with 2-dimensional physical objects propagating through a 3-dimensional base manifold.

sidered to be compatible with the notion of *spatial perception* of objects by beings in this model world if it possesses, at least to a good approximation, an SO(2) rotational symmetry about any point as well as translational symmetry in this plane. Hence the local symmetry group G of the manifold M_3 must:

- i) contain as a subgroup the symmetry of the purely spatial structure of the world; here the group $SO(2)$,
- ii) act on a space of one dimension higher than that of the spatial geometry; in this case 3-dimensional, and
- iii) be a possible symmetry group or subgroup of a form $L(v) = 1$ in order to conform with the present conceptual ideas.

For our model universe we begin with the 3-dimensional form of temporal flow:

$$
L(v_3) = (v^1)^2 + (v^2)^2 + (v^3)^2 = 1
$$
\n(2.14)

that is with $L(v_3) = \eta_{ab}v^av^b = 1$ and the 3-dimensional metric η_{ab} of equation 2.3 as introduced in the previous section. The full 3-dimensional translational symmetry of this form depicted in figure 2.2 provides the framework for an extended 2-dimensional 'spatial' environment, in addition to the temporal one, constituting the background manifold M_3 . Ultimately a metric with a 'spacetime' signature will be required in order to incorporate causal structure on the base manifold, however this feature is neglected for the simple model presented in this chapter. For the case of the model world an unbroken external symmetry SO(3) will be described in this section, before extending to a larger symmetry SO(5) over the same 3-dimensional base manifold in the following section.

2.2.1 The Base Manifold

The metric η_{ab} implies the existence of an orthonormal basis $\{e_a\}$ with respect to which the pure temporal flow v_3 can be expressed in terms of the components $\{v^a\} \in \mathbb{R}^3$ as:

$$
v_3 = v^a e_a = \frac{dx^a}{ds} e_a = \frac{d(x^a + r^a)}{ds} e_a
$$
 (2.15)

With a 3-dimensional translation symmetry $-\infty < r^a > \infty$ as depicted in figure 2.2 the orthonormal basis projects over the base manifold as an orthonormal frame field on M_3 . This smooth differentiable manifold naturally possesses a tangent space T_xM_3 at each point $x \in M_3$, that is the space M_3 has the properties of a 3-dimensional base manifold of a tangent bundle space, as we shall discuss further in section 3.3 for the general and 4-dimensional spacetime cases.

The assignment $v_3(x) = v^a e_a$ is valid for a local orthonormal coordinate basis or a frame field (with index a for such an orthonormal frame, here $a = \{1, 2, 3\}$). General coordinates on the manifold naturally give rise to a coordinate basis for the tangent space $\{\partial_\mu\}$, with $\partial_\mu \equiv \partial/\partial x^\mu$, (with index μ for general coordinates, here $\mu = \{1, 2, 3\}$. Relabelling the parameters $\{r^a\} \in \mathbb{R}^3$ in equation 2.15 as a particular set of 'general coordinates' $x^{\mu} = \delta^{\mu}_{\ \ a}r^{a}$ on M_{3} , there is an implied coordinate frame on the base manifold $\{\partial_{\mu}\}\$ such that $v_3 = v^a e_a$ of equation 2.15 can be expressed as $v_3 = v^{\mu} \partial_{\mu} = v^a e^{\mu}{}_a(x) \partial_{\mu}$, with the 'triad' components $e^{\mu}{}_a(x) = \delta^{\mu}{}_a$.

More generally under a passive reparametrisation to any general coordinates ${x^{\mu}} \in \mathbb{R}^{3}$ in a region of M_3 a frame field consists of a triad of vector fields $e_a =$ $e^{\mu}{}_{a}(x)\partial_{\mu}$ (for $a=1,2,3$) with components with respect to the general coordinate frame given by the matrix function $e^{\mu}_{a}(x)$ which points to the local Euclidean metric structure at any $x \in M_3$. The set of components $e^{\mu}_a(x)$ contains the same information as its matrix inverse $e^a_{\mu}(x)$, and either of these matrices are sometimes referred to as the 'triad' itself. These matrices transform both under general coordinate transformations and local, or gauge, SO(3) transformations.

The kernel symbol v will usually denote a vector or vector field corresponding to the fundamental flow of time in the form $L(v) = 1$, while the kernel symbol u will denote arbitrary tangent vector fields, such as $u(x) = u^{\mu}(x)\partial_{\mu}$, as indicated in figure 2.4. Either type of vector field may be expressed either in a local orthonormal frame or in a general coordinate frame. The components a vector field $u(x)$ belong to the space \mathbb{R}^3 whether presented in a local or a general coordinate basis; these two possibilities are related by the matrix $e^a_{\mu}(x) \in GL(3, \mathbb{R})$ such that $u^a(x) = e^a_{\mu}(x)u^{\mu}(x)$.

Through a frame field $e_a(x)$ on M_3 the flow of time described numerically by $v_3(x)$ is isomorphic to an *external* tangent vector field which may be described in terms of general coordinates on M_3 , and may be considered to be a flow of time on this manifold space itself, even for the case in which the global geometry is not Euclidean. This latter situation will arise when the local tangent space on M_3 is embedded within a higher-dimensional form of temporal flow, as described in the following section. In this case M_3 will necessarily be treated as a differentiable manifold with finite curvature in general for which only the local geometry at any point $x \in M_3$ will be isomorphic to the Euclidean geometry of \mathbb{R}^3 .

Via the triad field $e^a_{\mu}(x)$ the internal space constant metric η_{ab} of equation 2.3 implied in equation 2.14, similarly as for the vector components v^a , may be expressed

Figure 2.4: A local orthonormal basis $e_a \equiv \partial/\partial x^a$ for the vector field $v_3(x) =$ $v^a(x)e_a(x)$ is related to any other such basis of the same orientation at each point $x \in M_3$ by the action of the symmetry group $G = SO(3)$, which can vary arbitrarily over M_3 . In general a tangent vector field $u(x)$ on the manifold may be expressed in terms of an arbitrary frame field, or a particular orthonormal or general coordinate frame.

on the tangent space for a general coordinate basis. This determines the metric tensor:

$$
g_{\mu\nu}(x) = e^a_{\ \mu}(x)e^b_{\ \nu}(x)\eta_{ab}.\tag{2.16}
$$

In the theory of general relativity it is the freedom of the metric field $g_{\mu\nu}(x)$, or equivalently the tetrad field $e^a_{\mu}(x)$, on a 4-dimensional spacetime base manifold as will be described for equation 3.50, with respect to an arbitrary coordinate system that describes gravitation in the world, as we shall review in section 3.4. While in general the components of $g_{\mu\nu}(x)$ differ from those of η_{ab} in a general coordinate system, even for a flat spacetime, in general relativity it is the absence of any global coordinate basis such that $g_{\mu\nu}(x) = \delta^a_{\mu} \delta^b_{\nu} \eta_{ab}$ everywhere that is responsible for gravitational effects.

In order to consider the *curvature* of the base manifold it is necessary to formalise the notion of *parallelism*. The question concerns the way in which the base manifold M_3 originates out of the flow of time as depicted in figure 2.2, specifically with $n = 3$ for the model case here, such that the symmetry $G = SO(3)$, acting upon individual vectors $v_3(x) \in \mathbb{R}^3$ in the equation $L(v_3) = 1$ can act as an approximately global symmetry over scales that are large compared with the objects being perceived. In section 2.1 we began with finite intervals of multi-dimensional time, as depicted in figure 2.1, and then went on to the infinitesimal case $s \to \delta s$ in order to derive the relation $L(v) = 1$ of equation 2.9. We here need to understand how the symmetry of such infinitesimal intervals can apply coherently over finite *distances* on the manifold M_3 . This is required on the manifold in order to frame stable perceptions of 2-dimensional spatial objects propagating through such a world as depicted in figure 2.3.

In terms of the model world the point is that since we are locally free to choose an orthonormal frame within which to specify the numerical values $v^a(x)$ for the components of $v_3(x) \in TM_3$, the values themselves have no absolute meaning. In particular, no conclusion concerning the equality, or parallelism, of two sets of vector components $v^a(x_1)$ and $v^a(x_2)$ at two different points $x_1, x_2 \in M_3$ in figure 2.4 may be drawn since the bases of local frames at x_1 and x_2 may be chosen independently, within the local SO(3) freedom of the relation $L(\sigma_{g(x)}(\boldsymbol{v}_3(x))) = 1$. A triad frame (such as e_a described in equation 2.15 for the coordinate frame $x^{\mu} = \delta^{\mu}{}_a r^a$) could be declared to specify a parallelism on the manifold (that is, $v_3(x_1)$ is parallel to $v_3(x_2)$) if each of the components agree, $v^a(x_1) = v^a(x_2)$, in the specified triad frame $e_a(x)$). However for the case of a curved space or spacetime no global frame field $e_a(x)$ exists in a manner compatible with the parallelism, since the latter now depends on the path taken between x_1 and x_2 .

The underlying notion of parallelism is more generally defined in terms of a connection 1-form on the manifold, which readily extends to the case of non-global parallelism. The connection is a mathematical object, a Lie algebra valued 1-form, that mutually relates the bases on the manifold, with respect to a given path connecting the points $x_1, x_2 \in M_3$, and hence determines whether any two vectors at these locations are parallel with respect to the path. With such a structure the relative values of the components of two vectors at differing locations does acquire meaning. Hence we wish to identify an SO(3) connection form $A(x)$ on the base manifold M_3 . We describe how a *flat* connection arises *canonically* on M_3 through it's relation to the symmetry group G in subsection 2.2.3 (and further in section 3.2 in the context of the principle bundle structure). In the following subsection we first review the standard geometry on a group manifold itself.

2.2.2 The Group Manifold

While the $\{r^a\} \in \mathbb{R}^3$ translational symmetry of $L(\mathbf{v}_3) = 1$ gives rise to the base manifold M3, the algebraic structure of the rotational symmetry constitutes a second differentiable manifold which is identified with the Lie Group $G = SO(3)$ itself. This manifold is also intimately related to the temporal flow v_3 through the expression $L(v_3) = L(\sigma_q(v_3)) = 1$, with the action $g \in G$ realised on the subspace of unit norm vectors in \mathbb{R}^3 .

Elements of a general Lie group $g \in G$ also act as diffeomorphisms on the manifold G itself [2, 3, 4]. An example is the diffeomorphism $L_q: G \to G$, mapping the point $h \to gh$ with $g, h \in G$, called 'left translation' on the manifold. Due to the nature of the algebraic properties of a symmetry group a Lie group manifold G exhibits distinctive canonical geometrical structures. The significance of 'canonical' (in the sense of intrinsic or naturally existing) structures, where relevant, is that they carry the mathematical development of a theory forward in a necessary and non-arbitrary way. As for the base manifold described above, the group G as a manifold also has a tangent space T_qG at each point $g \in G$, through which a tangent vector field $V(g)$ may be described on G. Smooth vector fields $X(g)$ belonging to the subset which satisfy the relation:

$$
L_{g*}X(h) = X(gh) \tag{2.17}
$$

for all $g, h \in G$, where L_{g*} is the 'tangent mapping', or differential, of the left translation L_q (acting upon objects defined on the tangent space of G), are said to be left-invariant. The set of left-invariant vector fields together with their multiplication in terms of the commutator $[X, Y]$ (considering the vector fields X, Y as mappings in the space of scalar functions $f(g)$ on G), which itself describes a left-invariant vector field, defines the Lie algebra $L(G)$ of the Lie group G. As a vector space $L(G)$ is isomorphic to the set of tangent vectors at any location on G, and in particular to the space T_eG , where $e \in G$ is the identity element of the group. Given any point $h \in G$ the orbit of left translation for all $g \in G$ covers the entire group manifold, as a consequence of the transitive property of multiplication within a Lie group, and hence the corresponding tangent mapping L_{q*} carries any vector $V(h)$ into a left-invariant vector field on G.

In general a 1-form $\omega(x)$, or covector field, on a differentiable manifold M maps a vector field $V(x)$ into the space of real functions on M; this map may be denoted by:

$$
\langle \omega(x), V(x) \rangle = f(x) \tag{2.18}
$$

at any point $x \in M$. Over the manifold G a linearly independent set of left-invariant vector fields, $\{X_{\alpha}\}\$ with $\alpha = 1 \ldots n_G = \dim(G)$, forms a global frame field on G. A dual basis of 1-forms $\{\theta^{\alpha}\}\$ with $\alpha = 1...n_G$ such that $\langle \theta^{\alpha}, X_{\beta} \rangle = \delta^{\alpha}{}_{\beta}$, constitutes a coframe field on G. These covector fields are also left-invariant with $L_g^* \theta^\alpha(gh) = \theta^\alpha(h)$ for the 'pull-back' L_g^* of the left translation by $g \in G$.

The 'exterior algebra' of differential forms includes the exterior product ' \wedge ' and exterior derivative 'd' which act on 1-forms such as $\omega(x) = \omega_{\mu} dx^{\mu}$ and $\sigma(x) = \sigma_{\mu} dx^{\mu}$ to produce 2-forms such as:

$$
\omega \wedge \sigma = \omega \otimes \sigma - \sigma \otimes \omega \tag{2.19}
$$

and
$$
d\omega = \frac{\partial \omega_{\mu}}{\partial x^{\nu}} dx^{\nu} \wedge dx^{\mu}
$$
 (2.20)

For any diffeomorphism between manifolds, $f : M \to N$ (where it may be that $M =$ N), the pull-back map f^* is a structure preserving homomorphism of the exterior algebra. Hence, as for the 1-form basis covectors θ^{α} ($\alpha = 1...n_G$), the 2-forms $d\theta^{\alpha}$ and $\theta^{\beta} \wedge \theta^{\gamma}$ are also left-invariant on G and are therefore related via left-invariant, that is constant, scalar coefficients $c^{\alpha}{}_{\beta\gamma}$ as defined in:

$$
d\theta^{\alpha} + \frac{1}{2}c^{\alpha}{}_{\beta\gamma}\theta^{\beta} \wedge \theta^{\gamma} = 0.
$$
 (2.21)

This is the Maurer-Cartan equation which also serves to define the Lie algebra structure constants $c^{\alpha}{}_{\beta\gamma}$, with respect to the basis $\{\theta^{\alpha}\}\$. It is equivalent to the definition of the structure constants in terms of the dual basis of vector fields $\{X_{\alpha}\}\,$ which represents the Lie algebra itself, in the relation:

$$
[X_{\beta}, X_{\gamma}] = c^{\alpha}{}_{\beta\gamma} X_{\alpha}.
$$
\n(2.22)

The Maurer-Cartan 1-form θ is a single, basis independent, canonical object on the manifold G that expresses the properties of the collective set of n_G 1-forms $\{\theta^{\alpha}\}.$ It is a Lie algebra-valued 1-form defined by its action on a general tangent vector field

 $V = V^{\alpha}(g)X_{\alpha}$ on G with $\langle \theta, V(g) \rangle := V^{\alpha}X_{\alpha} \in L(G)$, where V^{α} are the component values of V at g and $V^{\alpha}X_{\alpha}$ is hence the Lie algebra element corresponding to the left-invariant vector field on G with the tangent vector $V(g)$ at the given point $g \in G$.

The Maurer-Cartan form can be written as $\theta = \theta^{\alpha} X_{\alpha}$ in terms of the dual bases. The canonical form θ encapsulates the parallelisable nature of any Lie group manifold by defining a consistent global parallelism on G. That is, θ represents a single reference frame for each of the tangent spaces which resolves each vector $V(g)$ at any $g \in G$ into its components with respect to a left-invariant frame field $\{X_{\alpha}\}\$. For a matrix group such as $SO(3)$, with a matrix basis $\{E_{\alpha}\}\$ for $L(G)$, the Maurer-Cartan form can also be expressed as the left-invariant matrix of 1-forms $\theta = g^{-1} dg$. In this case $\langle \theta, V(g) \rangle$ is the Lie algebra element $V^{\alpha} E_{\alpha}$ represented in matrix form in terms of the components of the corresponding left-invariant vector field at T_eG .

In terms of θ the Maurer-Cartan equation 2.21 can be written as:

$$
d\theta + \frac{1}{2}[\theta, \theta] = 0
$$
\n(2.23)

where the bracket denotes the 'exterior product' for Lie algebra valued 1-forms. Such a product may be defined on vector-valued p-forms in general provided there is a product defined on the vector space of the values. This is the case for Lie algebra valued forms where, with $\theta = \theta^{\alpha} X_{\alpha}$ and $\phi = \phi^{\beta} X_{\beta}$ and with $\{X_{\alpha}\}\$ a basis for $L(G)$, the product is defined as:

$$
[\theta, \phi] := \theta^{\alpha} \wedge \phi^{\beta} \; [X_{\alpha}, X_{\beta}]. \tag{2.24}
$$

For a matrix basis ${E_{\alpha}}$ and the case of a single 1-form as for equation 2.23 the product 1 $\frac{1}{2}[\theta, \theta] = \theta \wedge \theta$ which implicitly incorporates the multiplication of the E_{α} matrices.

On the group manifold G each left-invariant field X generates a one-parameter subgroup described by the flow $\phi_t = \exp(tX_e)$, where $X_e \in T_eG$ denotes the tangent vector belonging to the field X at the identity $e \in G$ and 'exp' is the 'exponential map' from $L(G)$ into the manifold G. The action of this one-parameter group on any point $h \in G$ is by right translation, as indicated in figure 2.5.

Alternatively, a left-invariant field X^A on G can be *induced* by the right action $R_g: h \to hg$ of elements of the one-parameter group $g(t) = \exp(tA)$, with $A \in T_eG$ such that:

$$
X_h^A(f) = \frac{d}{dt} f(h \exp(tA)) |_{t=0}
$$
 (2.25)

where f is a real-valued function on the manifold G . Left-invariant fields are sometimes denoted by a label 'R' since they are generated by right translations; hence X^R denotes a left-invariant field.

Since a right-invariant field Y^L (which can be generated by left translation) is by definition invariant under right translations the Lie derivative of Y^L with respect to the vector field X^R vanishes:

$$
\mathcal{L}_{X} Y^L = \frac{d}{dt} \phi_t^* Y^L |_{t=0} = 0 \tag{2.26}
$$

$$
\text{that is:} \qquad [X^R, Y^L] = 0 \tag{2.27}
$$

For each $g \in G$ a further diffeomorphism on the group manifold called the adjoint map can be defined by $\text{Ad}_g(h) = ghg^{-1}$ for all $h \in G$, that is $\text{Ad}_g = L_g \circ R_{g^{-1}} \equiv$

Figure 2.5: The integral curves of a left-invariant field on G generate right translations.

 $R_{q^{-1}} \circ L_q$ by the associative property of group composition. The adjoint map is an *automorphism* of the group composition. Since $\text{Ad}_{g_2} \circ \text{Ad}_{g_1} = \text{Ad}_{g_2g_1}$ this is a left action of G on itself.

The adjoint map applied to elements near $e \in G$ gives rise to the group representation $\mathrm{Ad}_g = L_{g*} \circ R_{g^{-1}*}|_e \equiv R_{g^{-1}*} \circ L_{g*}|_e$ acting upon the Lie algebra of the group. For a group represented by matrices this takes the form $\text{Ad}_g(Y) = gYg^{-1}$ for any $Y \in L(G)$. The adjoint representation $g \to \text{Ad}_q$ is a group homomorphism of G into $GL(L(G))$. The 'derived homomorphism' of this representation induces the corresponding adjoint representation for the Lie algebra elements with $\text{ad}_XY = [X, Y]$ for $X, Y \in L(G)$, as an automorphism of the Lie bracket algebra, which naturally involves the structure constants of the group through equation 2.22.

Finally we note that a left-invariant field X^R , which also generates right translations, itself transforms under right translation as $R_{g*}X^R = \text{Ad}_{g^{-1}}X^R$ by the definition of left-invariance and the adjoint representation (while $L_{q*}X^L = \text{Ad}_q X^L$ for a right-invariant field X^L). These group properties will be important for the structure of principle bundles described in section 3.1.

2.2.3 Relating the Base and Group Manifolds

As for the basis $\{X_{\alpha}\}\$ on the manifold G, a frame field $\{e_{a}\}\$, with $a = 1...n$, may be introduced on any *n*-dimensional differentiable manifold M_n , forming a linearly independent set of tangent vectors at each point of the manifold. The real quantities $c^a{}_{bc}(x)$ in the relation:

$$
[e_b, e_c] = c^a{}_{bc}(x)e_a \tag{2.28}
$$

or equivalently in:
$$
\mathrm{d}e^{a} = -\frac{1}{2}c^{a}_{bc}(x)e^{b} \wedge e^{c}
$$
 (2.29)

in terms of the dual coframe field $\{e^a\}$, are here variables called structure *coefficients* (or 'coefficients of anholonomy') rather than constants as for equations 2.21 and 2.22.

Given a general coordinate chart $\{x^{\mu}\}\$ on M_n and a holonomic frame $e_{\mu} = \partial_{\mu}$ the corresponding coefficients $c^{\rho}_{\mu\nu}(x)$ are all zero, while $c^{a}_{bc}(x) \neq 0$ implies that a noncoordinate frame is being employed.

On the manifold G frames composed of *left-invariant* vector fields $\{X_{\alpha}(g)\}\$ were identified as being particularly important owing to the group structure. On the base manifold M_3 , possessing the metric of equation 2.16, basis vectors forming *orthonormal* frames $\{e_a(x)\}\$ are particularly significant. As described in subsection 2.2.1 such a triad frames the components of v_3 subject to the pure numerical relation $L(v_3) = 1$ of equation 2.14, which implicitly contains the local 3×3 Euclidean metric η as expressed in equation 2.3. (In this paper indices $a, b, c \dots$ for a basis $\{e_a\}$ will denote an arbitrary smooth frame field, as for M_n above, or an orthonormal frame field, as for M_3 here, or even a coordinate basis depending on the context; while indices μ, ν, ρ ... for a basis ${e_{\mu}} \equiv {\partial_{\mu}}$ will always denote a coordinate frame).

The two manifolds M_3 and $G = SO(3)$, representing the translational and rotational symmetries of the form $L(v_3) = 1$, as described in section 2.1 and in the two subsections above, are linked through the mapping $g(x): M_3 \to G$. An initial orthonormal frame field ${e_a(x)}$ can be transformed to any other orthonormal frame field $\{e'_b(x)\}\$ by the matrix action $e'_b = e_a g^a_{\ b}$ via the group element $g(x) \in SO(3)$ at every $x \in M$. The map $g(x) : M_3 \to G$, as depicted in figure 2.6, expresses the local choice of an orthonormal frame field $\{e_a(x)\}\$, essentially the choice of local $\{x^1, x^2, x^3\}$ axes of figure 2.1 at each point $x \in M_3$, as a basis for tangent vectors $v_3 \in TM_3$. It is this 'gauge' freedom $g(x) \in SO(3)$ in the choice of local orthonormal frames that prevents a particular frame ${e_a(x)}$ from directly representing parallelism on the base manifold, as described towards the end of subsection 2.2.1.

Figure 2.6: The gauge choice of a frame at each $x \in M_3$ described as a map into elements $g \in G$ between the two manifolds.

Since the operations of the exterior algebra of p-forms are preserved under the pull-back of forms through diffeomorphism maps on manifolds the Lie algebra-valued 1-form:

$$
A = g^*\theta \tag{2.30}
$$

on M_3 captures the structural properties of the Maurer-Cartan 1-form θ on G relative to the map g. While on G we have the linear map $\langle \theta, V(g) \rangle \in L(G)$ from $V(g) \in T_gG$ into the Lie algebra of G, on M₃ we have the linear map $\langle A(x),u(x)\rangle \in L(G)$ from $u(x) \in T_xM_3$ into the same Lie algebra. The Lie algebra-valued 1-form $A(x)$ may be written as $A(x) = A^{\alpha}_{\mu}(x) X_{\alpha} dx^{\mu}$ where $\{dx^{\mu}\}\$ is a coordinate basis of 1-forms on M_3 and $\{X_\alpha\}$ is a basis for $L(G)$. In the appropriate 3×3 matrix representation the SO(3) generators can be denoted by $\{E_{\alpha}\}\equiv \{L_{pq}\}\,$, labelled by a single index mnemonic double letter symbol $pq = \{12, 13, 23\}$, with:

$$
(L_{pq})_{ab} = \delta_{pa}\delta_{qb} - \delta_{pb}\delta_{qa}
$$
\n(2.31)

where a and b label the matrix rows and columns respectively, that is:

$$
L_{12} = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \qquad L_{13} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \qquad L_{23} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}
$$
(2.32)

Unlike the canonical 1-form $\theta = \theta^{\alpha} X_{\alpha}$ on G, the 1-form A on M_3 has variable real coefficients $A^{\alpha}_{\mu}(x)$ which, however, are not arbitrary but depend upon the choice of gauge function $g(x)$ as well as upon the choice of coordinates $\{x^{\mu}\}$ on M_3 . Explicitly, for a matrix group G, the 1-form $A = g^*\theta$ on M_3 can be expressed as:

$$
A(x) = g^{-1} \mathrm{d}g = g^{-1} \frac{\partial g}{\partial x^{\mu}} \mathrm{d}x^{\mu}
$$
\n(2.33)

It is this canonical mathematical object that serves as a connection 1-form on the base manifold M_3 , formalising the notion of parallelism in manner which will naturally generalise for the case of finite curvature. Here it is possible to choose a gauge with $A(x) = 0$ everywhere on M_3 , simply by taking $g(x)$ to be constant in equation 2.33, and hence we have a flat connection. Indeed, this connection can always be written in terms of 'pure gauge', as it is in equation 2.33, which is one way of defining a flat connection (to be described in more detail in section 3.2). Given a connection $A(x) \neq 0$ a gauge transformation via any $g(x)$ transforms the connection in the standard way as:

$$
A \to A' = g^{-1} A g + g^{-1} \mathrm{d} g. \tag{2.34}
$$

which can be expressed as pure gauge $A' = g^{-1} \mathrm{d}g'$, that is in the form of equation 2.33, in terms of an appropriate gauge function $g'(x)$.

By the homomorphism of exterior algebra relations across the pull-back map the Lie algebra-valued 1-form $A = g^*\theta$ is also subject to a structure equation corresponding to equation 2.23, that is:

$$
dA + \frac{1}{2}[A, A] = 0 \tag{2.35}
$$

In general the curvature 2-form F on the base manifold can be expressed as:

$$
F = dA + \frac{1}{2}[A, A]
$$
 (2.36)

which transforms under a gauge change $g(x)$ as $F \to F' = g^{-1}Fg$. Equations 2.35 and 2.36 then immediately show that the curvature is equal to zero, with $F = 0$ in any gauge, and further expresses the global parallelism implied by the canonical flat connection of equation 2.30.

While the connection 1-form can be written as $A(x) = A^{\alpha}_{\mu}(x) E_{\alpha} dx^{\mu}$ and curvature 2-form can be written as $F(x) = \frac{1}{2} F^{\alpha}_{\mu\nu}(x) E_{\alpha} dx^{\mu} \wedge dx^{\nu}$ (where the factor of $\frac{1}{2}$ arises from the convention of equation 2.19 and the double counting implicit on the right-hand side since the set of asymmetric $dx^{\mu} \wedge dx^{\nu}$ 2-forms, with nine values of $\{\mu,\nu\} = \{1,2,3\},\$ does not describe a linearly independent basis).

Interest in the group $G = SO(3)$ arose as a symmetry action on the form $L(v_3) = 1$ and hence the Lie algebra values of $A(x)$ and $F(x)$ are composed of elements ${E_{\alpha}} \equiv {L_{pq}}$ of equation 2.32 in a representation of $L(G)$ acting naturally upon the vectors $u \in TM_3$, that is on the tangent space of the base manifold, and in particular on the vector $v_3 \in TM_3$ originating in the form $L(v_3) = 1$. The mathematical objects involved are hence intimately associated with each other, with the base space M_3 and the flat connection $A(x) = g(x)^*\theta$ of equation 2.30 upon it arising out of the translation and rotation symmetry properties of $L(v_3) = 1$, with the vectors v_3 themselves being tangent to M_3 .

As an example of this association the constancy of the scalar function $L(\mathbf{v}_3) = 1$ on M_3 can be expressed as $\partial_{\mu}L(v_3) = 0$, or, consistent with the gauge transformations of $v'_3 = g^{-1}v_3$ and equation 2.34, covariantly as:

$$
D_{\mu}L(v_3) = D_{\mu}(v_3 \cdot v_3) = 2v_3 \cdot D_{\mu}v_3 = 0 \qquad (2.37)
$$

with
$$
D_{\mu}v_3 = \partial_{\mu}v_3 + A_{\mu}v_3 \qquad (2.38)
$$

The 'covariant derivative' D_{μ} relating to a connection A_{μ} will be defined more precisely in the following chapter, leading to equation 3.6. These above two equations show how the connection field $A_\mu(x)$ explicitly acts on the vector field $v_3(x)$ in a constraining relation, and hence there is a 'coupling' between these fields over the base manifold M_3 .

A vector field $u(x)$ which satisfies $D_{\mu}u = 0$ everywhere represents a parallel vector field on the manifold. A frame field $\{e_a\}$ that satisfies $D_\mu e_a = 0$ for each value of a defines a parallel frame field – in which case the frame field itself may be used to define the parallelism on the manifold, which is only possible for a flat connection. With respect to the original global coordinates defined in terms of a parametrisation of the translation symmetry of equation 2.15 on M_3 the triad field with components $e^{\mu}_{a}(x) = \delta^{\mu}_{a}$ was identified. The covariant derivative of the corresponding orthonormal basis vectors e_a is constant with respect to the connection form $A(x) = 0$, with constant $g(x)$ in equation 2.33, and hence $\{e_a\}$ defines the parallelism in this case. The geometric objects $e^{\mu}_{a}(x) = \delta^{\mu}_{a}$ and $A(x) = 0$ may both be associated with the constant gauge function $g(x)$ on M_3 taken as the identity element $e \in G = SO(3)$ of the group. A gauge change by a constant $g(x) \neq e \in G$ changes the frame ${e_a}$ but not the connection $A(x) = 0$. Under a general gauge change $q(x)$ the connection, with $A(x) \neq 0$ in general, can be written as an explicit function of the triad field.

In summary the canonical connection $A = g^*\theta$, constructed as depicted in figure 2.6, defines a global parallelism on M_3 (as θ does on the manifold G), such that the parallel transport of a vector $u(x_1)$ from x_1 to another point x_2 on the base manifold, see for example figure 2.4, results in a definite vector $u(x_2)$ independent of the path taken. This formalises the notion of parallelism on the base manifold in a manner which can be generalised for the case of non-global parallelism.

Since $A(x)$ is SO(3)-valued in acting on the tangent space TM_3 it also describes a 'metric compatible' connection. In being completely determined by the triad field the connection $A(x)$ is also torsion-free, where torsion will be defined in section 3.3. In fact since $A(x)$ is a particular case of a linear connection acting on the tangent space the curvature 2-form F of equation 2.36 may be identified with the Riemann tensor and hence denoted **R**. The so(3)-valued curvature tensor $\mathbf{R} = \frac{1}{2}R^{\alpha}{}_{\mu\nu}E_{\alpha}dx^{\mu} \wedge dx^{\nu}$ on M_3 , with $\alpha = 1 \ldots n_G$, then has components $R_{ab\mu\nu} = R^{\mu q}_{\mu\nu} (L_{pq})_{ab}$. Via the triad field the same tensor can be expressed either fully in a local orthonormal frame or fully in a general coordinate frame – in the latter case with four general coordinate indices as:

$$
R_{\rho\sigma\mu\nu} = e^a_{\ \rho} e^b_{\ \sigma} R_{ab\mu\nu} \tag{2.39}
$$

Hence we have constructed a zero Riemann curvature tensor with all components $R_{\rho\sigma\mu\nu}=0$ as implicit in the identification of a canonical flat connection $A=g^*\theta$.

In section 3.1 we shall review the geometry of a principle fibre bundle, hence incorporating M_3 as the base manifold and G as the structure group together in a single manifold, before reviewing Riemannian geometry itself. A non-zero Riemannian curvature will ultimately be obtained on the original base manifold by expanding the form $L(v) = 1$ into a higher-dimensional temporal flow with a larger symmetry group, as we provisionally describe in the following section.

2.3 Higher Dimensions and Symmetry Breaking

In the previous section the construction of a model world required that we drew attention to the particular form of temporal flow $L(v_3) = 1$, as expressed in equation 2.14, as an example of the general *n*-dimensional case. It was shown how v_3 could be interpreted as a tangent vector field over a base manifold M_3 , represented in figure 2.4, which in turn may be parametrised by a set of real number coordinates $x^{\mu} \in \mathbb{R}^{3}$, and with a choice of a local orthonormal reference frame ${e_a(x)}$ determined within the freedom of the local SO(3) symmetry.

However, in general there are many higher symmetry groups acting upon vector spaces of a larger dimension, with elements conforming to $L(\mathbf{v}) = 1$, which we have no mathematical reason to neglect. Indeed, the reasoning of section 2.1 is consistent with the flow of time being channelled into a space of arbitrarily large dimension. Hence, mathematically, there is nothing to prevent the 3-dimensional space of parameters $v_3 \in \mathbb{R}^3$, representing a 3-dimensional flow of time, from further dividing into a larger multi-dimensional space of parameters described by the vector $v_n \in \mathbb{R}^n$ $(n > 3)$ subject to a new form $L(v_n) = 1$ with a higher symmetry group G. (In later chapters the expression $L(\hat{v}) = 1$ will denote the full form of temporal flow being considered, while the full symmetry group, excluding translations, may be denoted \tilde{G} for clarity, as for the remainder of this section).

The original SO(3) geometric symmetry group may now be identified as a subgroup $\overline{H} \subset \hat{G}$, with the 'overline' denoting an *external* symmetry, acting on the 3-dimensional flow v_3 which is projected onto the tangent space of the base manifold M_3 out of the higher-dimensional temporal flow. In this section we begin to consider the conceptual implications and mathematical possibilities of this generalisation for the necessary existence of such a higher symmetry group \tilde{G} acting upon a higherdimensional form of temporal flow $L(\mathbf{v}_n) = 1$.

Since it will ultimately be required to mathematically support the kind of situation depicted in figure 2.3, in which the smaller symmetry $\overline{H} \subset \hat{G}$ is treated in a distinctive way in giving rise to the global geometrical nature of a perceived universe of physical objects, we shall expect to be dealing with a natural mechanism for breaking the higher symmetry. The full 'rotation' symmetry of the form $L(v_n) = 1$, as a generalisation of that depicted in figure 2.1, is now broken since only the degrees of freedom of a subset of the possible dimensions of translation symmetry, depicted in figure 2.2, is employed to locally construct the base manifold

In subsequent chapters, for the real world, we shall motivate the choice of G as the Lie group E_6 , acting on a 27-dimensional form $L(v_{27}) = 1$, with a Lorentz subgroup acting on the local tangent space of the 4-dimensional spacetime base manifold M_4 . In the meantime here, for the model world, we shall take G to be a symmetry group of a form of $L(v_n) = 1$ large enough to contain SO(3), the orthonormal frame symmetry group, as a subgroup of \hat{G} , while retaining the 3-dimensional base space M_3 .

For the case of the model universe $\overline{H} = SO(3)$, acting upon $v_3 \in \mathbb{R}^3$, could be taken to be embedded within various kinds of larger groups, for example $\hat{G} = SU(3)$ acting upon the 6 real components of v_6 corresponding to a 3-dimensional complex vector $c_3 \in \mathbb{C}^3$ with $L(v_6) = c_3^{\dagger} c_3 = 1$. However, here we consider the vectors $v_3 \in \mathbb{R}^3$ of section 2.2 to be vectors in a subspace of \mathbb{R}^n with $n > 3$ upon which the group $SO(3)$ is a straightforward subgroup of $SO(n)$, the latter being a perfectly acceptable symmetry of $L(\mathbf{v}_n) = 1$, acting upon the vectors $\mathbf{v}_n \in \mathbb{R}^n$. In particular we choose $n = 5$ and consider the Lie group $\hat{G} = SO(5)$ acting on the form $L(\mathbf{v}_5) = 1$:

$$
L(\mathbf{v}_5) = \mathbf{v}_5 \cdot \mathbf{v}_5 = (v^1)^2 + (v^2)^2 + (v^3)^2 + (v^4)^2 + (v^5)^2 = 1 \tag{2.40}
$$

$$
= \eta_{ab}v^av^b + (v^4)^2 + (v^5)^2 = 1 \tag{2.41}
$$

where $a, b = 1...3$ and $\eta_{ab} = \text{diag}(+1, +1, +1)$ represents the 3-dimensional Euclidean metric, which was introduced in equation 2.3 of section 2.1.

The 5-dimensional vector v_5 has components $v^a = dx^a/ds$, with $a = 1...5$. Hence the vectors of $L(v_5) = 1$ implicitly contain the 5-dimensional translational freedom $x^a \to x^a + r^a$, $\{r^a\} \in \mathbb{R}^5$, as a particular example of equation 2.13 and figure 2.2 which generalises the 3-dimensional case of equation 2.15. However, we consider only the 3-dimensional freedom of this parameter space and continue to take M_3 to be the base space as we did in section 2.2 and as depicted for the present case in figure 2.7. This choice will ultimately be justified by the identification of geometrical structures on M_3 which may then be interpreted as the base space for perception of physical events as sketched in figure 2.3. The model described in figure 2.7 provides a convenient picture for a provisional discussion of the symmetry breaking structure which will be picked up again in section 5.1 for the more realistic case over a 4-dimensional spacetime base manifold M_4 .

Figure 2.7: (a) The gauge choice at each $x \in M_3$ depicted in figure 2.6 is extended for $\hat{G} = SO(5)$ with (b) only the *subgroup* $SO(3) \subset SO(5)$ now acting on TM_3 .

A basis for the 10-dimensional Lie algebra so(5) (with the lower case 'so' denoting the Lie algebra corresponding to the $SO(5)$ Lie group), as represented on a 5-dimensional vector space as the generators of a symmetry of $L(v_5) = 1$, is provided by the set of ten 5×5 matrices $(L_{pq})_{ab}$ of the type described in equation 2.31, now with ten distinct labels composed out of $p, q = 1...5, p < q$. The three so(5) Lie algebra elements:

$$
L_{12} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \ -1 & 0 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ \end{pmatrix}, \quad L_{13} = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ -1 & 0 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ \end{pmatrix}, \quad L_{23} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \ 0 & 0 & 1 & 0 & 0 \ 0 & -1 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ 0 & 0 & 0 & 0 & 0 \ \end{pmatrix}
$$
(2.42)

generate an $SO(3) \subset SO(5)$ subgroup, as can be see directly as guided by the horizontal and vertical lines drawn into the matrices in equation 2.42 and by comparison with equation 2.32. This $SO(3)$ subgroup can be taken to act on the tangent space of M_3 and hence upon the subspace of vectors $\overline{v}_3 \subset v_5$ projected onto the base space. Of the other seven SO(5) generators one acts purely on the complementary 2-dimensional subspace $\underline{v}_2 \subset v_5$, namely:

$$
L_{45} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}
$$
(2.43)

With the subspace of vectors $\overline{v}_3 \subset v_5$ with $\overline{v}_3 \in TM_3$ in the external space the SO(2) generator L_{45} can be said to act upon the *internal* 2-dimensional space of vectors $v_2 \in \mathbb{R}^2$, which can be considered to represent *extra dimensions* over those required to describe the extended base space M_3 . In general while an 'overline' denotes an external object an 'underline' will denote an object defined in the internal space. In this representation basis the remaining six so(5) Lie algebra elements are:

$$
L_{14} = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad L_{15}, L_{24}, L_{25}, L_{34}, \quad L_{35} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 \end{pmatrix} (2.44)
$$

This final set of six matrices generate SO(5) group elements that *mix* the external \bar{v}_3 and internal \underline{v}_2 parts of the full 5-dimensional temporal flow $v_5 = \begin{pmatrix} \overline{v}_3 \\ \overline{v}_2 \end{pmatrix}$. The vectors \overline{v}_3 and \underline{v}_2 are physically distinct with respect to the M_3 base space. The full SO(5) symmetry is hence broken down to:

$$
SO(3) \times SO(2) \subset SO(5) \tag{2.45}
$$

as the external $SO(3)$ symmetry, represented by the generators of equation 2.42 'locks on' to the tangent space TM_3 leaving the residual internal symmetry $SO(2)$, represented by the generator in equation 2.43, as depicted in figure 2.7(b). Hence only four of the original ten generators of $SO(5)$ survive the symmetry breaking.

For the original unbroken full symmetry the constancy of $L(v_5) = 1$ can be expressed, in comparison with equations 2.37 and 2.38, as the vanishing of the covariant derivative of $L(v_5) = 1$ on the base manifold:

$$
\hat{D}_{\mu}L(v_5) = 0 \Rightarrow v_5 \cdot \partial_{\mu}v_5 + v_5 \cdot \hat{A}_{\mu}v_5 = 0 \qquad (2.46)
$$

where the 'hat' on \hat{D}_{μ} and $\hat{A}_{\mu}(x)$ signifies that the unbroken 10-component so(5)-valued connection 1-form on M_3 is being considered. With the identification of a Riemannian curvature tensor on M_3 six of the gauge field generator degrees of freedom are lost and the broken, physical, form of equation 2.46 can be written as:

$$
D_{\mu}L(v_5) = 0 \Rightarrow v_5 \cdot \partial_{\mu}v_5 + \overline{v}_3 \cdot A_{\mu}\overline{v}_3 + \underline{v}_2 \cdot Y_{\mu}\underline{v}_2 = 0 \qquad (2.47)
$$

where $A_{\mu}(x)$ represents the external so(3)-valued connection 1-form and the gauge field $Y_\mu(x)$ describes an internal so(2)-valued connection 1-form. The final term in equation 2.47 expresses an 'interaction' between the gauge field $Y_\mu(x)$ and the internal temporal components $\underline{v}_2(x)$ which follows directly from the 'minimal coupling' between them implicit in the covariant derivative (as a generalisation from the purely external field coupling described for equations 2.37 and 2.38). The structure of the pattern of interactions for the breaking of the E_6 symmetry of $L(v_{27}) = 1$ over the base manifold M_4 for the real world will be described in chapter 8.

Here for the model, a 3-dimensional projection of the full temporal flow $\overline{v}_3 \subset v_5$ will form a tangent vector field on the base manifold M_3 . While $|v_5| = \sqrt{L(v_5)}$ is fixed in equation 2.40 the quantity $|\overline{v}_3|$ in principle has a variable magnitude, however here we mainly focus on the breaking of the group symmetry action and in particular the relation between the geometry of the resulting external and internal curvature.

Considering first the case of figure $2.7(a)$ with an unbroken set of ten $SO(5)$ Lie group generators the algebra product is given by the following expression, which is valid in general for the orthogonal $SO(n)$ groups:

$$
[L_{pq}, L_{zs}] = \delta_{qr} L_{ps} - \delta_{qs} L_{pr} - \delta_{pr} L_{qs} + \delta_{ps} L_{qr}
$$
\n(2.48)

From this the so(5) algebra structure constants c^{pq} $_{rs\,tu}$ in this basis can be read off, for example $c^{23}{}_{12\,13} = -1$ since $[L_{12}, L_{13}] = -L_{23}$. Since the group SO(5) is connected and compact any element $g \in SO(5)$ in this matrix representation can be expressed as

$$
g = \exp(\alpha_{pq} L_{pq})
$$
\n(2.49)

with ten real coefficients α_{pq} (summation is implied over repeated index combinations, for the ten labels $\{p, q = 1 \ldots 5, p < q\}$, however the 'upper' or 'lower' location of these indices is of no significance). The 'exponential map' was described in the discussion around figure 2.5 in the context of left-invariant vector fields on the group manifold. Here with the Lie algebra represented by real matrices $L = \alpha_{pq} L_{pq} \in L(\hat{G})$ the exponential map may be explicitly written as $\exp(L) = \sum_{k=0}^{\infty} \frac{1}{k}$ $\frac{1}{k!}L^k$ which converges to a map from $L(\hat{G}) \rightarrow \hat{G}$.

The elements of equation 2.49 satisfy the relation $gg^T = 1_5$ (where $1_5 \in \hat{G}$ is the identity element of the group as represented by the 5×5 unit matrix) and $det(g) = 1$, as required for the special orthogonal group SO(5). As an example, with the notation $c_{pq} = \cos \alpha_{pq}$ and $s_{pq} = \sin \alpha_{pq}$ the element $g_{14} = \exp(\alpha_{14}L_{14})$ has the form:

$$
g_{14} = \begin{pmatrix} c_{14} & 0 & 0 & s_{14} & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ -s_{14} & 0 & 0 & c_{14} & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}
$$
(2.50)

The Maurer-Cartan 1-form θ , now defined on the manifold of the full gauge symmetry group $\hat{G} = SO(5)$, can be pulled-back onto the base manifold M_3 as described in subsection 2.2.3. In this way we canonically identify a flat $so(5)$ -valued connection 1-form $\hat{A}(x) = g^*\theta$ and an so(5)-valued curvature 2-form $\hat{F}(x) = d\hat{A} + \frac{1}{2}[\hat{A}, \hat{A}] =$ 0, following equations 2.35 and 2.36, where the 'hat' on A and F here again denote quantities involving the $full$ $SO(5)$ symmetry group.

Beginning with a choice of constant gauge $g_c(x) : M_3 \to SO(5)$, with fixed $g_c \in SO(5)$, we have $\hat{A}(x) = g_c^* \theta = 0$ from equation 2.33. Under a more general gauge transformation of the form of equation 2.49 but with in particular $g_{14} = \exp(\alpha_{14}L_{14})$ and with small values of the function $\alpha_{14}(x)$, we have from equation 2.34:

$$
\hat{A}'(x) = g_{14}^{-1}\hat{A}(x)g_{14} + g_{14}^{-1} \mathrm{d}g_{14} \tag{2.51}
$$

$$
\simeq \, \mathrm{d}\alpha_{14}(x)L_{14} \qquad \text{for } \hat{A}(x) = 0 \tag{2.52}
$$

This can be written $\hat{A}'(x) = A^{\mathbf{14}}(x) L_{\mathbf{14}}$ with $A^{\mathbf{14}}(x) = d\alpha_{\mathbf{14}}(x) = \partial_{\mu} \alpha_{\mathbf{14}}(x) dx^{\mu}$ as the 1-form coefficient of the Lie algebra element L_{14} . Applying a full sequence of all six 'mixing' actions in the order $g_6 = g_{35} g_{34} g_{25} g_{24} g_{15} g_{14}$ (that is with g_{14} first and g_{35} last, each generated by an element of equation 2.44 and all being functions of $x \in M_3$) a 1-form coefficient for each of the ten Lie algebra basis elements L_{pq} may be found. This is equivalent to taking $\hat{A}(x) = g_6^* \theta = g_6^{-1} dg_6$. To order $O(\alpha_{pq} d\alpha_{zs})$ for small transformations the ten coefficients of this connection $\hat{A}(x) = A^{\mu q}(x) L_{pq}$ are found as:

$$
A^{12} = -\alpha_{24} \, d\alpha_{14} - \alpha_{25} \, d\alpha_{15}, \quad A^{13} = -\alpha_{34} \, d\alpha_{14} - \alpha_{35} \, d\alpha_{15}, \quad A^{23} = -\alpha_{34} \, d\alpha_{24} - \alpha_{35} \, d\alpha_{25},
$$

$$
A^{14} = d\alpha_{14}, \quad A^{15} = d\alpha_{15}, \quad A^{24} = d\alpha_{24}, \quad A^{25} = d\alpha_{25}, \quad A^{34} = d\alpha_{34}, \quad A^{35} = d\alpha_{35},
$$

$$
A^{45} = -\alpha_{15} \, d\alpha_{14} - \alpha_{25} \, d\alpha_{24} - \alpha_{35} \, d\alpha_{34} \tag{2.53}
$$

The full so(5)-valued connection 1-form $\hat{A}(x) = A^{\mu q}(x) L_{\mu q}$, summing over the ten 1form coefficients in equation 2.53, is 'unphysical' in the sense that it is 'pure gauge' with respect to the full $SO(5)$ symmetry, and merely presents the same original flat connection, for which we had all ten $A^{pq}(x) = 0$, in a different choice of gauge, namely $g_6(x) \in SO(5)$.

Of more significance is the structure and interpretation of the ten components of the curvature 2-form \hat{F} in the new gauge. These can be written using the general expression for the curvature 2-from coefficients in terms of the connection 1-form coefficients (consistent with equation 2.36):

$$
\hat{F}^{\alpha} = dA^{\alpha} + \frac{1}{2} \hat{c}^{\alpha}{}_{\beta\gamma} A^{\beta} \wedge A^{\gamma}
$$
\n(2.54)

The Lie algebra basis indices for $L(SO(5))$ are here denoted by $\{\alpha, \beta, \gamma\} = 1 \dots 10$ (with $\alpha = 1, 2...$ corresponding to $pq = 12, 13...$) and the structure constants may be read off from equation 2.48. For the internal curvature 2-form coefficient $\hat{F}^{\ddagger 5}(x)$ in the gauge $g_6(x)$, using the asymmetry in the $\beta\gamma$ indices of the structure constants and of the exterior product of 1-forms, we find:

$$
\hat{F}^{45} = dA^{45} + \frac{1}{2} \hat{c}^{45}{}_{\beta\gamma} A^{\beta} \wedge A^{\gamma}
$$
\n
$$
= dA^{45} + \hat{c}^{45}{}_{44 \text{ L5}} A^{14} \wedge A^{15} + \hat{c}^{45}{}_{24 \text{ L5}} A^{24} \wedge A^{25} + \hat{c}^{45}{}_{34 \text{ L5}} A^{34} \wedge A^{35}
$$
\n
$$
= dA^{45} - A^{14} \wedge A^{15} - A^{24} \wedge A^{25} - A^{34} \wedge A^{35}
$$
\n
$$
= d\alpha_{14} \wedge d\alpha_{15} + d\alpha_{24} \wedge d\alpha_{25} + d\alpha_{34} \wedge d\alpha_{35} - d\alpha_{14} \wedge d\alpha_{15} - d\alpha_{24} \wedge d\alpha_{25} - d\alpha_{34} \wedge d\alpha_{35}
$$
\n
$$
= 0 \qquad (2.55)
$$

where in the penultimate line the connection coefficients from equation 2.53 have been substituted into this expression. This result is as expected and indeed zero curvature,

 $\hat{F}^{\mu q} = 0$ for all ten 2-form coefficients, is associated with the connection 1-form $A^{\mu q}$ coefficients, in any gauge choice such as that for equation 2.53, so long as the full 10-dimensional Lie algebra of SO(5) is retained.

At each point on M_3 the Lie algebra so(5) acts partly on the external space T_xM_3 of the base manifold in figure 2.7(a), via an SO(3) ⊂ SO(5) subgroup, and partly on the internal space through the complementary $SO(2) \subset SO(5)$, while the remaining generators of equation 2.44 straddle the external and internal parts of $v_5 \in \mathbb{R}^5$. By choosing an $SO(5)$ gauge in figure 2.7(a) and then breaking the symmetry through a projection onto the structure in figure 2.7(b) the aim here is to demonstrate how non-zero curvature may be associated with the $SO(3)$ and $SO(2)$ subgroups.

While an so(5)-valued connection $A(x)$ provides a means for the parallel transport of a 5-dimensional vector $v_5(x)$ over M_3 for figure 2.7(a), a single component such as $A^{45}(x)L_{45}$ can be interpreted in the *restricted* sense of a representation in the subgroup SO(2) ⊂ SO(5), that is $\hat{A} \to \underline{A} = A^{45}L_{45}$ with $L_{45} \in L(SO(2))$, acting on the subspace of 2-dimensional vectors $\underline{v}_2 \subset v_5$. In this case $\underline{A}(x)$ is a 1-form connection describing the parallel transport of vectors $\underline{v}_2(x)$ in the internal vector space over M_3 for figure $2.7(b)$.

Hence treating the generator L_{45} in isolation from the other nine generators as a purely $SO(2)$ action on the internal space of vectors $\underline{v}_2 \in \mathbb{R}^2$ a curvature 2form $\underline{F} = \underline{F}^{45}L_{45}$ may be identified for this *restricted* internal SO(2) symmetry. In comparison with equation 2.55, for the restricted $SO(2)$ subgroup a finite curvature may be obtained:

$$
\underline{F}^{45} = dA^{45}
$$
\n
$$
= d\alpha_{14} \wedge d\alpha_{15} + d\alpha_{24} \wedge d\alpha_{25} + d\alpha_{34} \wedge d\alpha_{35}
$$
\n
$$
\neq 0 \quad \text{in general.} \tag{2.56}
$$

That is, under a suitable choice of gauge parameters $\alpha_{pq}(x)$ in the full SO(5) symmetry it is possible to identify non-zero curvature components with respect to subgroups such as $SO(2)$, which may be interpreted as a structure with finite internal physical curvature over the base space manifold.

To obtain equation 2.56 we effectively took a restricted set of structure constants, which is trivial for the Abelian subgroup $SO(2) \subset SO(5)$ with a single generator and hence there are no $\underline{c}^{\alpha}{}_{\beta\gamma}$ terms in place of the $\hat{c}^{\alpha}{}_{\beta\gamma}$ terms of equation 2.54. This results in a non-zero internal curvature $\underline{F}^{45} \neq 0$ for the subgroup \underline{H} under what was purely a change of gauge from the point of view of the full group \hat{G} . In this way a finite SO(2) curvature is essentially carved out of the degrees of freedom implicit within the structure of the unbroken flat SO(5) connection.

Complementary to the subgroup $SO(2) \subset SO(5)$ the $SO(3)$ subgroup is generated by the Lie algebra elements of equation 2.42. Acting on the external vector components $\overline{v}_3(x) \in T_xM_3$ the connection 1-forms $A^{12}L_{12}$, $A^{13}L_{13}$ and $A^{23}L_{23}$, associated with the three generators of SO(3), define parallelism on the external tangent space of the manifold M_3 . However here we temporarily follow the same approach applied to the $SO(2)$ case in leading to equation 2.55, hence treating this $SO(3)$ as an 'internal' symmetry in order to examine any new features that arise for a non-Abelian gauge group such as $SO(3)$. Again, initially substituting the full set of $SO(5)$ connection coefficients from equation 2.53 into equation 2.54, we find $\hat{F}^{12} = \hat{F}^{13} = \hat{F}^{23} = 0$ as for all ten so(5)-valued curvature components as discussed above.

A purely SO(3) Lie algebra-valued curvature 2-form can be obtained, similarly as for the SO(2) case, by using only the restricted Lie algebra values of the connection acting purely on the subspace of vectors $\overline{v}_3 \subset v_5$ to identify the curvature tensor $\overline{F} = \overline{F}^{\hat{\mu}q} L_{pq}$, with $pq = 12$, 13, 23, over M_3 . The three components are obtained from equation 2.54 by curtailing the summations to include only the so(3)-valued parts with a restricted set of structure constants $\{\overline{c}^{\alpha}_{\beta\gamma}\}\subset\{\hat{c}^{\alpha}_{\beta\gamma}\}\$ describing the external SO(3) symmetry only. Following a similar procedure that led to equation 2.56 a set of three so(3)-valued curvature coefficients is found to lowest non-trivial order:

$$
\overline{F}^{12} = d\alpha_{14} \wedge d\alpha_{24} + d\alpha_{15} \wedge d\alpha_{25}
$$
\n
$$
\overline{F}^{13} = d\alpha_{14} \wedge d\alpha_{34} + d\alpha_{15} \wedge d\alpha_{35}
$$
\n
$$
\overline{F}^{23} = d\alpha_{24} \wedge d\alpha_{34} + d\alpha_{25} \wedge d\alpha_{35}
$$
\n(2.57)

Hence the SO(3) curvature is also non-zero in general and clearly correlated with the $SO(2)$ curvature of equation 2.56, with the correlation mediated through the six mixing gauge functions $\alpha_{pq}(x)$, with $pq = 14, 15, 24, 25, 34, 35$ under the full $SO(5)$ symmetry. Either the $SO(3)$ or the $SO(2)$ curvature may be non-zero while the other remains zero for a suitable choice of the $\alpha_{pq}(x)$, while both $\overline{F} = 0$ and $\underline{F} = 0$ are simultaneously attained under any choice of $SO(5)$ gauge with constant $\alpha_{pq}(x)$ for example.

The six full SO(5) curvature components $\hat{F}^{\mu q} = 0$ for $pq = 14, 15, 24, 25, 34,$ 35 are not directly associated with subgroup restrictions giving rise to further finite curvature components. Rather these six mixing degrees of gauge symmetry are lost or *broken* in the projection of the full $SO(5)$ symmetry over M_3 through which finite physical curvature for the four components in equations 2.56 and 2.57 is identified. The four functions $\alpha_{pq}(x)$, with $pq = \{12, 13, 23\}$ and 45, corresponding to four actions $g_{pq} = \exp(\alpha_{pq}L_{pq})$ of the gauge symmetry, survive the symmetry breaking and are retained as the gauge symmetries associated with the non-Abelian SO(3) and Abelian SO(2) subgroups respectively.

However the mechanism of symmetry breaking over M_3 itself implies that there is a more fundamental difference between the $SO(3)$ and $SO(2)$ subgroups in the context of the model world we are considering. The former acts externally on the tangent space of the base manifold, that is on $\overline{v}_3 \in TM_3$ as depicted in figure 2.7(b), and is therefore closely related to the geometry of the background space itself and to the existence of a linear connection on the base manifold. Hence the correlation observed above between the $SO(3)$ and $SO(2)$ curvature, with both effectively treated as *internal* symmetries, merely provides a provisional motivation for seeking a unified framework in which $SO(2)$ remains as an internal gauge symmetry while $SO(3)$ is considered as an external symmetry on TM_3 . A non-zero curvature for the internal symmetry $\underline{F}^{45} \neq 0$ in equation 2.56 was obtained by considering the generator of the subgroup \underline{H} = SO(2) within the full gauge symmetry group \hat{G} = SO(5) over a fixed base space M₃. A non-zero Riemannian curvature with components $R^{\rho}_{\sigma\mu\nu}(x) \neq 0$ on M₃ will transform locally under the complementary external subgroup $SO(3) \subset SO(5)$ acting on the tangent space TM_3 .

In this chapter the base manifold M_3 and group manifold G (where here G may be the full symmetry \tilde{G} of the full form $L(\boldsymbol{v}_n) = 1$ or either the internal H or external \overline{H} subgroups) have been treated as largely independent geometric objects, however their mutual relationship is more precisely defined in terms of a single manifold in the form of a principle bundle with base space M_3 and structure group G. Hence the standard properties of these geometric objects, together with a review of Riemannian geometry and the Lagrangian approach to obtaining equations of motion for the corresponding field entities, with be presented in the following chapter.

A relationship between the internal gauge curvature and external Riemannian curvature might be determined through such a principle fibre bundle $P = (M_3, \hat{G})$ with base space M_3 and structure group \hat{G} , based on the picture in figure 2.7(a), with the canonical zero full curvature $\ddot{F}=0$ providing the constraint that relates the internal and external geometry. However the physical situation is represented by figure 2.7(b) which leads to a consideration of two detached bundle spaces, $\overline{P} = (M_3, SO(3))$ and $P = (M_3, SO(2))$. While \overline{P} directly only contains information about the external symmetry and curvature, the bundle space \underline{P} explicitly contains both the structure of the external geometry on M_3 and that of the internal $H = SO(2)$ curvature in the bundle space. As a unifying framework for combining external and internal symmetries this latter structure is very similar to that employed in non-Abelian Kaluza-Klein theories, which are hence reviewed in chapter 4.

In chapter 5 we consider how the above structures of principle bundles and Kaluza-Klein theory might be adapted for the present theory. There we shall upgrade the model presented in this chapter by considering the real world situation with a 4-dimensional base space M_4 with local Lorentz symmetry. This will be embedded in the 'full' symmetry group taken as $\hat{G} = SO^+(1, 9)$, acting on a 10-dimensional form of temporal flow $L(v_{10}) = 1$, which will be broken to $SO^+(1,3) \times SO(6)$ in the projection onto the spacetime base M_4 . This symmetry breaking structure naturally embeds in the further higher-dimensional extensions considered from chapter 6, which will provide a more realistic framework for the details of the internal structures also.

Chapter 3

Review of Geometry and Equations of Motion

3.1 Principle Bundle Structure

In section 2.2 we introduced two independent differentiable manifolds, the base space M_3 and Lie group $G = SO(3)$ with points labelled by $x \in M_3$ and $g \in G$ respectively, both of which are associated with the 3-dimensional form of temporal flow $L(v_3) = 1$ through the respective 'translational' ${r^a} \in \mathbb{R}³$ and 'rotational' $\sigma_g : g \in G$ symmetries:

$$
L\left(\sigma_g\left\{\frac{d(x^a+r^a)}{ds}\right\}\right) = 1\tag{3.1}
$$

The map between the manifolds $g : M_3 \to G$ described in figure 2.6, mapping $x \to g(x)$ (where $\{x^{\mu}\}\$ may be taken as general coordinates on M_3) represents a local choice of gauge, or orthonormal frame, in which to express the tangent vector $v_3(x)$ on M_3 . This association between M_3 and G may be examined more precisely through the construction of a single differentiable manifold, namely a principle fibre bundle P , which combines the geometric properties of the base space M_3 and Lie group G together with their mutual relation.

In the general case the structure group G of a principle fibre bundle P does not need to be related to a symmetry on the tangent space to the base manifold M , as it is for the $SO(3)$ model as implied in equation 3.1. Indeed in the case of figure 2.7(a) the symmetry group $G = SO(5)$ acts only partially on the tangent space of M_3 , while for figure 2.7(b) the group $G = SO(2)$ does not act on the external tangent space at all. Hence it is the generalisation in which G and M are initially introduced independently that we shall review here for the benefit of the subsequent application to the case of a higher symmetry group such as presented in section 2.3.

A principle bundle P is a G -manifold, that is a differentiable manifold upon which the transformation group G acts, with a particular structure as described, with reference to figure 3.1, by the following properties (see for example [2], [3] chapter Vbis, [4]):

Figure 3.1: The relations between three differentiable manifolds: a principle fibre bundle P , the base space M and the structure group G .

- 1: There is a surjective map $\pi : P \to M$ projecting from the bundle space onto the base manifold. Given a section $\sigma(x) : M \to P$ then $\pi \circ \sigma(x) = x$ is the identity map on points $x \in M$.
- 2: For each $x \in M$ the submanifolds $\pi^{-1}(x) \subset P$, called the fibres of P, are diffeomorphic to each other and to the Lie group G .
- 3: The right action of $g \in G$ on points $p \in P$, that is $R_g: P \to P$ with $R_{gh} = R_h \circ R_g$, preserves the fibres of P, that is $\pi \circ R_q = \pi$, and is free and transitive on each fibre.
- 4: There exist local trivialisations over each open subset $U_r \subset M$, consisting of maps $\psi_r : \pi^{-1}(U_r) \to U_r \times G$ with $\psi_r : p \to (x, h)$, such that $\psi_r : pg \to (x, hg)$ that is the right action on P is compatible with the right action on G .

While the right action of G on itself induces left-invariant fields as described in equation 2.25 of subsection 2.2.2, the right action of G on the manifold P induces 'vertical' vector fields in the tangent space TP as (where f is now a real-valued function on the bundle space):

$$
V_p^A(f) = \frac{d}{dt} f(p \exp(tA)) |_{t=0}
$$
 (3.2)

where $A \in L(G)$ and V_p^A is a tangent vector to the fibre of P at the point p. The map $A \to V_p^A$ described in equation 3.2 represents an *isomorphism* of the Lie algebra $L(G)$ into the space of vector fields residing in the vertical tangent space $VP \subset TP$. That is, the Lie algebra bracket structure $[X^A, X^B] = X^{[A,B]}$ of equation 2.22 for the corresponding left-invariant fields $\{X^A\}$ on the manifold G is respected by the Lie bracket on the P bundle with:

$$
[V^A, V^B] = V^{[A,B]} \tag{3.3}
$$
This structure relates to VP, the space of vectors tangent to the individual fibres of P. Different fibres may be related by an additional structure on P called a *connection* which, conceptually, is smooth assignment of a 'horizontal' subspace H_pP of the full tangent space T_pP at each point $p \in P$ such that:

$$
T_p P = V_p P \oplus H_p P \tag{3.4}
$$

$$
R_{g*}H_pP = H_{pg}P \tag{3.5}
$$

where compatibility of the horizontal subspaces on P with the right action of G is assured by the latter requirement.

At every point $p \in P$ a basis for the tangent space of the principle bundle can be expressed in terms of these complementary subspaces. Such a basis $\{\acute{e}_i\} = \{\acute{e}_{\alpha}, \acute{e}_{\alpha}\}$ consists of the subset $\{\acute{e}_{\alpha}\}\in VP$ (that is vectors of the form V_p^A in equation 3.2, tangent to the fibres G_x over each point $x \in M$) and the subset $\{\hat{e}_a\} \in HP$, where \acute{e}_a is the 'horizontal lift' of the basis vector $e_a \in T_xM$ to the point $p \in P$ such that $\pi_*\acute{e}_a = e_a$. The 'acute' mark above the kernel symbol, such as for \acute{e} , denotes an object defined on a principle bundle space in the horizontal lift basis. In all cases the indices $\{i, j, k \ldots\}$ correspond to basis elements for TP in the total space; $\{\alpha, \beta, \gamma \ldots\}$ in the fibre space on P or on the manifold G; and $\{a, b, c \ldots\}$ in a complementary subspace on P or on the base space M .

It should be noted that \acute{e}_a and e_a are not only different vector fields but are also defined on two different manifolds, P and M respectively, although there is a oneto-one correspondence between them. Similarly, there is a one-to-one correspondence between a vector field \acute{e}_{α} on P and a vector field X_{α} on G, for example as generated by the same element $A \in T_eG$ in equations 3.2 and 2.25 respectively. The relations between these vector fields are indicated in figure 3.2.

Figure 3.2: Vertical and horizontal basis vectors in a local trivialisation $U \times G$ of a principle bundle P , together with their associated basis vectors on the group space G and the base manifold M respectively.

The specification of a connection on the principle bundle allows 'parallel transport' between the fibres to be defined by a path in P for which the tangent vector at any $p \in P$ always lies within the horizontal subspace H_pP . This notion of parallelism over M is used in turn to define a *covariant derivative* for associated fields $\phi(x)$ on the base space that transform under a representation of the structure group G , by tracking a parallel basis for the field $\phi(x)$ over any curve C on the base manifold (technically, $\phi(x)$ is a section in a fibre bundle associated with P).

As depicted in figure 3.3 given a point $p_1 \in P$ with $\pi(p_1) = x_1$ and a curve C on M from x_1 to x_2 a connection on P specifies a unique horizontal lift of the curve C to the curve C' on P , by advancing locally within the horizontal subspace $HP \subset TP$. The path C' then represents the 'parallel transport' of p_1 mapped to the unique point $p_2 \in P$, with $\pi(p_2) = x_2$.

Figure 3.3: Parallel transport C' between fibres on a principle fibre bundle P above the curve C on the base manifold from the point $x_1 \in M$ to $x_2 \in M$.

The geometric structure developed in section 2.2 corresponds to a particular kind of principle bundle, namely a frame bundle with structure group $G = SO(3)$ over the base space M_3 , which may be denoted $P = (M_3, SO(3))$. In this case the mapping from p_1 to p_2 in figure 3.3 provides a unique, path C dependent, transport of an orthonormal basis frame from x_1 to x_2 on the base manifold (such basis frames ${e_a}$ are shown in figure 2.4 for the model on M_3). With respect to such a parallel frame the difference between the values of the vector field $v_3(x)$ (belonging to the vector representation of $SO(3)$ at the two base points of the associated vector bundle can be determined. In particular vectors $\mathbf{v}_3(x_1)$ and $\mathbf{v}_3(x_2)$, as originally depicted in figure 2.4, are defined to be parallel with respect to a given path C and connection HP if each of their components coincide in an orthonormal reference frame transported from x_1 to x_2 along C via the horizontal lift C'. This definition of parallelism is independent of the choice $p_1 \in \pi^{-1}(x_1)$ of initial frame. In addition, any vector $\mathbf{v}_3(x_1)$ may be 'parallel transported' to any point of the curve C by maintaining constant

vector components in the corresponding transported frame of the principle bundle at each point along C.

This construction generalises for an arbitrary structure group G acting via a group representation on the field $\phi(x)$ over the base manifold M. The covariant derivative of the field $\phi(x)$ is defined in a such a way as to quantitatively indicate deviations of the value of the field function from that of the parallel transported field for infinitesimal displacements on the base manifold $-$ i.e. the extent to which the field is not self-parallel along a path on the base manifold. That is, if c is a vector at x_0 tangent to the curve C parametrised by $\lambda \to C(\lambda)$ on M with $C(0) = x_0$, see figure 3.3, then the covariant derivative of the field $\phi(x)$ along C at x_0 is defined as:

$$
D_{\mathbf{c}}\phi|_{x_0} = \lim_{\lambda \to 0} (T_{\lambda,0} \phi(x_\lambda) - \phi(x_0)) / \lambda \tag{3.6}
$$

where $T_{\lambda,0} \phi(x_\lambda)$ is the field value $\phi(x_\lambda) \equiv \phi(C(\lambda))$ parallel transported along C from x_{λ} to x_0 . The covariant derivative for the SO(3) connection applied to vectors $v_3(x)$ in equation 2.37 and 2.38 was denoted D_{μ} corresponding to derivatives with respect to general coordinate parameters $\{x^{\mu}\}\$ on M_3 .

If the connection is such that, for all $p \in P$ and all $X, Y \in H_pP$, the bracket composition on H_pP is closed, that is:

$$
[X,Y] \in H_p P \tag{3.7}
$$

then Frobenius criterion is satisfied and P is 'foliated' into a family of integrable 'leaves'. Each leaf is 'horizontal section' of P, with tangent space HP, that is a smooth *n*-dimensional submanifold of P , where n is the dimension of the base space M . In this case all horizontal lift curves C' of figure 3.3 effectively follow the contours of a single horizontal section submanifold, globally defined over M , and parallelism is independent of the path C taken between any two points on the base manifold. A 'flat' connection is defined by this property, as will be described in more detail in the following section.

Generally a horizontal subpace can be specified by a connection 1-form ω on P. The defining structure for HP of equation 3.4 and 3.5 can be attained via a smooth Lie algebra-valued 1-form $\omega \in L(G) \otimes T^*P$, mapping vectors $X \in TP$ into elements of $L(G)$, with the properties (the first of which is essentially the reverse of equation 3.2):

(i)
$$
\omega(V^A) = A
$$
 with $A \in L(G)$ (3.8)

(ii)
$$
R_g^* \omega = \text{Ad}(g^{-1}) \omega
$$
 i.e. $R_g^* \omega_{pg}(X) = g^{-1} \omega_p(X)g$ with $X \in T_pP$ (3.9)

where
$$
H_p P \equiv \{ X \in T_p P \mid \omega(X) = 0 \}
$$
 (3.10)

is the horizontal subspace. A set of trivialisations $\{U_r, \psi_r\}$ consists of an atlas $\{U_r\}$ covering the base manifold M together with a mapping ψ_r of each $\pi^{-1}(U_r) \subset P$ onto $U_r \times G$ such as depicted figure 3.1 and described in the subsequent 'item 4:'. Each trivialisation ψ_r is canonically associated with a section in $\pi^{-1}(U_r)$ which can be written as $\sigma_r(x) = \psi_{r,x}^{-1} \cdot \iota(x)$, where the map $\iota: U_r \to U_r \times G$ sends $x \to (x, e)_r$, with $e \in G$ being the identity element of the group and $\psi_{r,x}$ is the map ψ_r restricted to the space $\pi^{-1}(x)$. That is, $\sigma_r(x) \in P$ corresponds to the identity element $e \in G$ under the local trivialisation map ψ_r , as depicted in figure 3.4. More generally we have the map $\psi_{r,x}: P \to (U_r \times G)$ mapping between the points $\sigma_r(x)h \to (x,h)_r$, with $h \in G$.

Figure 3.4: Two trivialisations, denoted $\{U_r, \psi_r\}$ and $\{U_s, \psi_s\}$, with an overlap region, on a principle bundle P over a base space M .

In the overlap regions on M, for $x \in U_r \cap U_s$, transition mappings $g_{rs} : U_r \cap U_s \rightarrow$ G between such trivialisations are defined as the functions on M :

$$
x \to g_{rs}(x) = \psi_{r,x} \circ \psi_{s,x}^{-1} \in G \tag{3.11}
$$

The transition functions act on the left on a fibre such that $\psi_{r,x}^{-1}(x, g_{rs}h)_r = \psi_{s,x}^{-1}(x,h)_s$. These relate the corresponding canonical sections $\sigma_r(x)$ via the right action of the structure group on P , which commutes with the left action, in a way that is consistent with both $\psi_s(\sigma_s(x)h) = (x,h)_s$ and $\psi_r(\sigma_r(x)h') = (x,h')_r$ in the respective trivialisations, as:

$$
\sigma_s(x) = \sigma_r(x) g_{rs}(x) \tag{3.12}
$$

Given a general connection 1-form ω on P and a set of trivialisations $\{U_r, \psi_r\}$ a unique family A_r of connection 1-forms may be defined on M. Under a particular section σ_r the connection $A_r(x) = \sigma_r^* \omega(p)$ on M can be expressed as $A_r(x) = A^{\alpha}(x)X_{\alpha}$ $A^{\alpha}{}_{\mu}(x)X_{\alpha}dx^{\mu}$ where $\{X_{\alpha}\}\)$ is a basis for $L(G)$. The field of connection coefficients $A^{\alpha'}_{\mu}(x)$ link the basis $\{X_{\alpha}\}\$ for the Lie algebra, typically expressed in the appropriate representation (such as the set of matrices ${E_{\alpha}}$ of equation 2.32 for the case of the vector representation of $G = SO(3)$, with indices $\alpha = 1 \dots \dim(G)$, to a coordinate basis of 1-forms $\{dx^{\mu}\}\$ with indices $\mu = 0, 1, 2, 3$ in the case of a 4-dimensional spacetime base manifold M_4 . Further, given an $L(G)$ -valued 1-form $A(x)$ on M and any section $\sigma(x)$ then there exists a unique connection 1-form $\omega(p)$ on P such that $A = \sigma^* \omega$.

In a gauge theory, that is a theory which is invariant under transformations of the gauge group G which describes a local internal symmetry, the local Lie algebravalued 1-form $A(x)$ on the base manifold M is also known as a Yang-Mills field or 'gauge potential'. Such fields will be generically denoted $Y(x)$ in this paper, as for example in equation 2.47. The notation $A(x)$ may refer to a general connection 1-form, as described above, the gauge field associated with an internal $U(1)$ gauge symmetry, as for electromagnetism, or a connection associated with an orthonormal frame in the

external space, as was the case in subsection 2.2.3 and as will be the case in relation to general relativity, depending on the context. A gauge theory bases upon an internal symmetry, through the notion of a connection 1-form, involves similar mathematical structures as found in general relativity based upon an external symmetry.

For the present theory it will be assumed that the structure of principle bundles with a trivial global topology will be sufficient. In this case the bundle $P = (M, G)$ can be expressed as $P \equiv U \times G$ where a single 'subset' U of figure 3.4 may be identified with the entire base manifold M . This triviality is implied in deriving the bundle structure through the symmetries of $L(v) = 1$ as described for example in figures 2.2, 2.6 and 2.7. In this case the 'overlap region' for a change of trivialisation, or gauge transformation, may consist of the entire volume of the base space M , rather than a limited patch as depicted in figure 3.4.

In order to study the dynamics of the gauge fields it is helpful to introduce the exterior covariant derivative on the bundle space, which will be important for equations in physics. This derivative essentially combines the properties of the exterior derivative, introduced for equation 2.20, with the structure of the partial derivative ∂_{μ} as augmented to the covariant derivative D_{μ} , as described for equation 3.6.

More explicitly, on a principle bundle P the exterior covariant derivative D maps a V-valued r-form ϕ , which acts upon r vector fields $\{X_1 \ldots X_r\}$ on P, to a V-valued $(r+1)$ -form $D\phi$, where V is the representation space associated with G. The action of D is defined as:

$$
D\phi = (d\phi) \circ hor \tag{3.13}
$$

where 'hor' first maps vectors X on the tangent space of P to their horizontal components (that is, hor : $X \to X_h$, such that $X_h \subset H_pP$ of equation 3.4 and $\omega(X_h) = 0$, with ω the connection 1-form on P), and d is the exterior derivative map acting on the *r*-form ϕ .

In general, for a V-valued r-form ϕ on P which is *horizontal* (that is if any of the r vectors X on which ϕ acts is purely vertical then the map is zero, $\phi(X, \ldots) = 0$ and *equivariant* of type ρ (that is ϕ in the associated bundle transforms as $R_g^*\phi = \rho(g^{-1})\phi$ under the right action by $g(x)$ in the ρ representation) then the exterior covariant derivative of ϕ , equation 3.13, takes the simplified form:

$$
D\phi = d\phi + \rho'(\omega) \wedge \phi. \tag{3.14}
$$

where $\rho'(\omega)$ denotes the appropriate representation of the Lie algebra acting on V.

3.2 Curvature and Flat Connection

The curvature 2-form Ω can be *defined* as the exterior covariant derivative of the connection 1-form, that is $\Omega = D\omega$, on the principle bundle. The connection 1-form ω is equivariant, of type Ad as seen in equation 3.9, but it is clearly not a horizontal form, as seen in equation 3.8. However, for this particular case the exterior covariant derivative of the connection 1-form can also be expressed in a simplified form directly in terms of $\omega = \omega^{\alpha} X_{\alpha}$ itself (with $\{X_{\alpha}\}\$ a basis for $L(G)$) through the Cartan structure equation for the curvature 2-form $\Omega = \Omega^{\alpha} X_{\alpha}$ on P:

$$
\Omega(X,Y) = D\omega(X,Y) = d\omega(X,Y) + [\omega(X), \omega(Y)] \tag{3.15}
$$

$$
= d\omega(X,Y) + \frac{1}{2}[\omega,\omega](X,Y) \tag{3.16}
$$

that is
$$
\Omega^{\alpha}(X,Y) = d\omega^{\alpha}(X,Y) + \frac{1}{2}c^{\alpha}{}_{\beta\gamma}\omega^{\beta} \wedge \omega^{\gamma}(X,Y) \qquad (3.17)
$$

acting upon any pair of tangent vectors $X, Y \in TP$ (the meaning of $[\omega, \omega]$ is explained in the discussion around equation 2.24). The Lie algebra-valued curvature 2-form Ω on P is defined in such a way as to be quantitatively sensitive to deviations of the connection 1-form ω , and hence horizontal subspace HP , from the condition of flatness. This can be seen by substituting any $X, Y \in HP$ as arguments for the curvature 2-form Ω in equation 3.15, or equivalently for any $X, Y \in TP$ in terms of $\Omega = D\omega$ we have:

$$
\Omega(X,Y) = D\omega(X,Y) \tag{3.18}
$$

$$
= d\omega(X_h, Y_h) \tag{3.19}
$$

$$
= X_h \langle \omega, Y_h \rangle - Y_h \langle \omega, X_h \rangle - \langle \omega, [X_h, Y_h] \rangle \tag{3.20}
$$

$$
= -\langle \omega, [X_h, Y_h] \rangle \tag{3.21}
$$

where equation 3.19 follows directly from equation 3.13, equation 3.20 follows from the standard definition of the exterior derivative of a 1-form and the map \langle , \rangle was defined in equation 2.18. Hence it follows that Ω is non-zero only if the local horizontal subspaces on P defined by ω are non-integrable, that is the Frobenius criterion of equation 3.7 is not satisfied, and hence a non-zero curvature Ω indeed indicates a non-flat connection.

The Lie algebra valued curvature 2-form Ω on P is equivariant of type Ad, that is it transforms under the adjoint representation of G as $R_g^*\Omega = \text{Ad}(g^{-1})\Omega$. However, unlike the connection ω , the curvature Ω is also a horizontal form on P. Hence Ω , unlike $ω$, is a tensorial form meaning that, for a given choice of gauge or cross-section $σ$ over a region of the base manifold, it can be mapped via the pull-back σ^* to a geometrical object on the base manifold that transforms homogeneously as a representation of the gauge group. Such quantities may be more naturally equated in the expressions of physics. For the model universe of the previous chapter the curvature form of the $SO(3)$ connection on P is tensorial of type $(Ad, so(3))$, that is it takes values in the SO(3) Lie algebra and transforms under the adjoint representation, as indicated after equation 2.36 for the curvature form F on the base manifold.

In a trivialisation $P \equiv U \times G$ on the principle bundle a direct product basis $\{\ddot{e}_i\} = \{\ddot{e}_\alpha, \ddot{e}_\alpha\}$ for the tangent space consists of the subset $\{\ddot{e}_\alpha\} \in VP$, tangent to the fibres G_x over each point $x \in M$, and the subset $\{\ddot{e}_a\}$ with $\ddot{e}_a = \sigma_* e_a$ for each basis vector $e_a \in T_xM$ such that $\pi_* \ddot{e}_a = e_a$. Each vector \ddot{e}_a defined on the section $\sigma(x)$ is Lie transported via the right action of G on P such that the basis covers the entire principle bundle. The 'double dot' mark above the kernel symbol, such as for \ddot{e} , denotes an object defined on a principle bundle space in the direct product basis.

Since P itself is a differentiable manifold equation 2.28 applies for any frame field $\{e_i\}$ on P and is here expressed as:

$$
[e_j, e_k] = c^i_{jk}(p)e_i \tag{3.22}
$$

with real-valued structure coefficients $c^i_{jk}(p)$. In the direct product basis the bracket relations $[\ddot{e}_j, \ddot{e}_k] = \ddot{c}^i_{jk}(p)\ddot{e}_i$ are simply:

$$
[\ddot{e}_{\alpha}, \ddot{e}_{\beta}] = c^{\gamma}{}_{\alpha\beta} \ddot{e}_{\gamma} \tag{3.23}
$$

$$
[\ddot{e}_{\alpha}, \ddot{e}_{b}] = 0 \tag{3.24}
$$

$$
[\ddot{e}_a, \ddot{e}_b] = 0 \tag{3.25}
$$

where $c^{\gamma}{}_{\alpha\beta}$ are the structure constants of the group G. The zero coefficients for the second equation follow as the vector fields \ddot{e}_{α} generate the right translations which Lie transport the vectors \ddot{e}_b over P, and those in the final equation correspond to the choice of a coordinate basis on M.

By contrast the horizontal lift basis $\acute{e}_i = (\acute{e}_{\alpha}, \acute{e}_{a})$ for the tangent space TP, introduced after equation 3.4, is adapted to a given connection ω such that $\acute{e}_{\alpha} \in V_p$ and $\acute{e}_a \in H_p$, as was depicted in figure 3.2, with $\omega(\acute{e}_\alpha) = X_\alpha$ and $\omega(\acute{e}_a) = 0$, by the definition of the horizontal lift basis. Given a trivialisation the horizontal lift basis $\{\hat{e}_i\}$ can be expressed in terms the direct product basis $\{\hat{e}_i\}$ on P via the coefficients $\omega_{a}^{\alpha}(x, g)$ with:

$$
\dot{e}_{\alpha} = \ddot{e}_{\alpha}, \qquad \dot{e}_{a} = \ddot{e}_{a} - \omega^{\alpha}_{\ a} \ddot{e}_{\alpha} \qquad (3.26)
$$

$$
\acute{e}^{\alpha} = \ddot{e}^{\alpha} + \omega^{\alpha}{}_{a}\ddot{e}^{a}, \qquad \qquad \acute{e}^{a} = \ddot{e}^{a} \qquad (3.27)
$$

where $\{\hat{e}^i\} = \{\hat{e}^\alpha, \hat{e}^a\}$ is the dual basis defined as usual such that $\langle \hat{e}^i, \hat{e}_j \rangle = \delta^i_j$. The relation between the horizontal lift basis $\{\hat{e}_i\}$ on P and a direct product basis $\{\hat{e}_i\}$ on $U \times G$ is indicated in figure 3.5.

Figure 3.5: The adapted tangent space basis $\{\hat{e}_i\}$ on P with respect to a particular local trivialisation $\psi : P \to U \times G$ and the corresponding direct product basis $\{\ddot{e}_i\}.$

Acting on both sides of the second expression in equation 3.26 with the 1-form coefficients ω^{β} of the Lie algebra-valued connection 1-form $\omega = \omega^{\beta} X_{\beta}$ on P determines the connection coefficients $\omega^{\beta}(x, g) = \omega^{\beta}(\ddot{e}_a)$ on $U \times G$, as depicted in figure 3.5. From the transformation property of the connection 1-form ω on P under R_g^* in equation 3.9 and with the vector field \acute{e}_{α} generating right actions on P, it follows that:

$$
\acute{e}_{\alpha}\omega^{\beta}_{\ a} = \ddot{e}_{\alpha}\omega^{\beta}_{\ a} = -c^{\beta}_{\ \alpha\gamma}\omega^{\gamma}_{\ a} \tag{3.28}
$$

as the infinitesimal form of the adjoint transformation under the right action of the group.

Covariant differentiation on the base space is intimately related to the directional derivative \acute{e}_a on the principle bundle. Using equation 3.26 the bracket $[\acute{e}_a, \acute{e}_b]$ may be expressed in a direct product basis as:

$$
[\acute{e}_{a},\acute{e}_{b}] = [(\ddot{e}_{a} - \omega^{\alpha}{}_{a}\ddot{e}_{\alpha}),(\ddot{e}_{b} - \omega^{\beta}{}_{b}\ddot{e}_{\beta})]
$$
\n
$$
= [\ddot{e}_{a},\ddot{e}_{b}] - [\omega^{\alpha}{}_{a}\ddot{e}_{\alpha},\ddot{e}_{b}] - [\ddot{e}_{a},\omega^{\beta}{}_{b}\ddot{e}_{\beta}] + [\omega^{\alpha}{}_{a}\ddot{e}_{\alpha},\omega^{\beta}{}_{b}\ddot{e}_{\beta}]
$$
\n
$$
= 0 + \ddot{e}_{b}(\omega^{\alpha}{}_{a})\ddot{e}_{\alpha} - \ddot{e}_{a}(\omega^{\beta}{}_{b})\ddot{e}_{\beta} + \omega^{\alpha}{}_{a}\omega^{\beta}{}_{b}\dot{c}^{\gamma}{}_{\alpha\beta}\ddot{e}_{\gamma} + \omega^{\alpha}{}_{a}(\ddot{e}_{\alpha}\omega^{\beta}{}_{b})\ddot{e}_{\beta} - \omega^{\beta}{}_{b}(\ddot{e}_{\beta}\omega^{\alpha}{}_{a})\ddot{e}_{\alpha}
$$
\n
$$
= \ddot{e}_{b}(\omega^{\gamma}{}_{a})\ddot{e}_{\gamma} - \ddot{e}_{a}(\omega^{\gamma}{}_{b})\ddot{e}_{\gamma} + (\omega^{\alpha}{}_{a}\omega^{\beta}{}_{b}\dot{c}^{\gamma}{}_{\alpha\beta}\ddot{e}_{\gamma} - \omega^{\alpha}{}_{a}\dot{c}^{\gamma}{}_{\alpha\beta}\omega^{\beta}{}_{b}\ddot{e}_{\gamma} + \omega^{\beta}{}_{b}\dot{c}^{\gamma}{}_{\beta\alpha}\omega^{\alpha} - \ddot{e}_{\gamma})
$$
\n
$$
= (\ddot{e}_{b}\omega^{\gamma}{}_{a} - \ddot{e}_{a}\omega^{\gamma}{}_{b} - \omega^{\alpha}{}_{a}\omega^{\beta}{}_{b}\dot{c}^{\gamma}{}_{\alpha\beta})\ddot{e}_{\gamma}
$$
\n
$$
= -\Omega^{\gamma}{}_{a}b\acute{e}_{\gamma}
$$
\n(3.29)

using the first of equations 3.26 and where

$$
\Omega^{\gamma}{}_{ab}(x,g) = \ddot{e}_a \omega^{\gamma}{}_{b} - \ddot{e}_b \omega^{\gamma}{}_{a} + c^{\gamma}{}_{\alpha\beta} \omega^{\alpha}{}_{a} \omega^{\beta}{}_{b} \tag{3.30}
$$

are the curvature components on the principle bundle expressed in a particular trivialisation, as can be shown explicitly by substituting (\ddot{e}_a, \ddot{e}_b) for (X, Y) in equation 3.17. At any point $p \in P$ the components of $\Omega(p)$ are numerically the same in the horizontal lift basis as for a direct product basis, that is $\Omega^{\gamma}_{ab} = \Omega^{\gamma}(\dot{e}_a, \dot{e}_b) = \Omega^{\gamma}(\ddot{e}_a, \ddot{e}_b)$, since Ω is a horizontal form and \acute{e}_a and \ddot{e}_a differ only by a vertical vector, as seen in the second of equations 3.26 and figure 3.5. From equation 3.30, using equation 3.28, it can be shown that:

$$
\acute{e}_{\alpha}\Omega^{\beta}_{ab}(x,g) = \ddot{e}_{\alpha}\Omega^{\beta}_{ab}(x,g) = -c^{\beta}_{\alpha\gamma}\Omega^{\gamma}_{ab}(x,g) \tag{3.31}
$$

again transforming infinitesimally under the adjoint representation, as for the gauge field $\omega^{\alpha}{}_{a}(x,g)$, on the principle bundle.

In summary in the horizontal lift basis the full set of structure coefficients on P are considered with:

$$
[\acute{e}_{\alpha}, \acute{e}_{\beta}] = c^{\gamma}{}_{\alpha\beta} \acute{e}_{\gamma} \tag{3.32}
$$

$$
[\acute{e}_{\alpha}, \acute{e}_{b}] = 0 \tag{3.33}
$$

$$
[\acute{e}_a,\acute{e}_b] \hspace{2mm} = \hspace{2mm} \acute{c}^\alpha_{\hspace{2mm} ab}\acute{e}_\alpha = -\Omega^\alpha_{\hspace{2mm} ab}\acute{e}_\alpha \hspace{2cm} (3.34)
$$

Equation 3.33 follows directly from equations 3.24 and 3.26. Since right translations induce the basis vectors of the subspace VP , via equation 3.2, equation 3.33 expresses the right-invariance of the fields $\acute{e}_b \in HP$, consistent with equation 3.5, and may be compared with equation 2.27 in which Y^L is right-invariant. For the third equation the structure coefficients c_{ab}^d are set to zero since here a coordinate basis is taken for $\{e_a\}$ on the base manifold M in order to simplify the expressions. The fibre dependence of the structure coefficients \hat{c}^{α}_{ab} may be deduced by application of the Jacobi identity with:

$$
[\acute{e}_{\alpha}, [\acute{e}_{a}, \acute{e}_{b}]] + [\acute{e}_{a}, [\acute{e}_{b}, \acute{e}_{\alpha}]] + [\acute{e}_{b}, [\acute{e}_{\alpha}, \acute{e}_{a}]] = 0
$$

\n
$$
= [\acute{e}_{\alpha}, \acute{c}^{\beta}_{ab} \acute{e}_{\beta}] + 0 + 0 = 0
$$

\n
$$
\Rightarrow (\acute{e}_{\alpha} \acute{c}^{\beta}_{ab}) \acute{e}_{\beta} + \acute{c}^{\gamma}_{ab} \acute{c}^{\beta}_{\alpha\gamma} \acute{e}_{\beta} = 0
$$

\n
$$
\Rightarrow \acute{e}_{\alpha} \acute{c}^{\beta}_{ab} = -c^{\beta}_{\alpha\gamma} \acute{c}^{\gamma}_{ab}
$$
(3.35)

The final expression describes the directional derivative of the coefficients \hat{c}^{β}_{ab} with respect to the vector field \acute{e}_{α} , and hence expresses the transformation of \acute{c}^{β}_{ab} under the action of right translation, that is the gauge transformation generated by \acute{e}_{α} . This is consistent with the transformation property in equation 3.31, for the components the curvature 2-form under infinitesimal gauge transformations, as expected since by equations 3.29 and 3.34 we have simply:

$$
\acute{c}^{\alpha}{}_{ab} = -\Omega^{\alpha}{}_{ab} \tag{3.36}
$$

Given a curvature 2-form $\Omega(p)$ on a principle bundle P and a local section $\sigma(x)$ on P, for $x \in U \subset M$, the local representative of Ω on the base space is defined by the pull-back map as the 2-form $F(x) = \sigma^* \Omega(p)$, which also takes values in the Lie algebra, that is $F(x) = F^{\alpha}(x)X_{\alpha}$.

Another significant property of the curvature on the principle bundle P is that the exterior covariant derivative of Ω itself vanishes as a consequence of the definitions used to construct it, that is $D\Omega = 0$, which is called the *Bianchi* identity. The object $D\Omega = 0$ is also a tensorial form on P, like Ω itself, and since the exterior algebra structure pulls back through a section map $\sigma(x)$ we have a similar property for the corresponding object on M, that is on the base space we have $DF = 0$, which is also referred to as the Bianchi identity.

Through the section map σ the structure equation for the curvature 2-form Ω on P , for example in equation 3.16, pulls back to the base space M as:

$$
F = dA + \frac{1}{2}[A, A]
$$
 (3.37)

which was introduced in equation 2.36. In a particular trivialisation the components of the 'Yang-Mills field strength' on the base manifold M are $F^{\alpha}_{ab}(x) = \Omega^{\alpha}_{ab}(x, e)$, while the 'gauge potentials' are $A^{\alpha}_{a}(x) = \omega^{\alpha}_{a}(x, e)$. Consistent with equation 3.30 the above expression for F can be written in components, in a coordinate basis on M , as:

$$
F^{\alpha}_{\ \mu\nu}(x) = \partial_{\mu}A^{\alpha}_{\ \nu} - \partial_{\nu}A^{\alpha}_{\ \mu} + c^{\alpha}_{\ \beta\gamma}A^{\beta}_{\ \mu}A^{\gamma}_{\ \nu} \tag{3.38}
$$

while the 2-forms F^{α} are related to the 1-forms A^{α} according to equation 2.54.

For a connection on a principle bundle for which the structure group G as a subgroup of $GL(m,\mathbb{R})$ exhibits a matrix representation acting upon objects $v(x) \in V$ of an m-dimensional vector space (where m is not necessarily equal to the dimension n of the base manifold) the vector and curvature fields transform under a change of gauge $g(x) \in G$ on the base space M as:

$$
\mathbf{v} \to \mathbf{v}' = g^{-1} \mathbf{v} \tag{3.39}
$$

$$
F \to F' = g^{-1} F g \tag{3.40}
$$

This form of transformation follows from the choice of a right action of G on P , as featuring for example in equation 3.12, and in turn ultimately on the choice for $L(G)$ to be represented by left-invariant vector fields on G as described in subsection 2.2.2.

Connection 1-forms $A_r(x) = \sigma_r(x)^* \omega$ on the base manifold with respect to different trivialisations are related under the local gauge transformations by $g_{rs}(x)$ between the sections of equation 3.12 as:

$$
A_s(x) = \mathrm{Ad}(g_{rs}^{-1}(x))A_r(x) + (g_{rs}^*\theta)_x
$$

where Ad is the transformation of the adjoint representation on the Lie algebra values of $A_r(x)$ and θ is the Maurer-Cartan 1-form on the group manifold G, which here is pulled back onto M via the transition function map $g_{rs}(x): M \to G$. For a matrix representation, dropping the subscript labels, this transformation can be written as:

$$
A \to A' = g^{-1} A g + g^{-1} \mathrm{d} g \tag{3.41}
$$

where the second term is needed to take into account general gauge changes $g(x)$ between sections over M since ω is not a horizontal form on P. Under a change of section $\sigma'(x) = \sigma(x)g(x)$ via the local gauge function $g(x)$, the transformations of equations 3.39–3.41 are considered a passive symmetry from a physical point of view.

The connection 1-form ω on the principle bundle, which is a Lie algebra valued map on the tangent space T_pP of equation 3.4, may be restricted to a mapping on elements of V_pP tangent to the fibres of the bundle space, as it is in equation 3.8 for example. Under this restriction the properties of ω are equivalent to the Maurer-Cartan 1-form θ , described in subsection 2.2.2, which maps left-invariant vector fields on the manifold G as $\theta(X^A) = A$ and which transforms under right translation as $R_g^* \theta = \text{Ad}(g^{-1})\theta$, to be compared with equations 3.8 and 3.9.

Indeed, for a trivial bundle we have $P = M \times G$ and through the natural projection $\pi_2 : M \times G \to G$, the canonical Maurer-Cartan 1-form θ on G can be pulled back to $\omega = \pi_2^* \theta$ on P. Since the pull-back map captures the structure of the exterior algebra as seen through the map itself the Maurer-Cartan equation, that is equation 2.23, pulls back to:

$$
d\omega + \frac{1}{2}[\omega, \omega] = 0 \tag{3.42}
$$

By comparison with equation 3.16 it can be seen that for this connection the curvature vanishes, $\Omega = 0$, that is ω is the canonical flat connection on P.

In general for a continuous map between two differentiable manifolds $f : M \rightarrow$ N, with a vector field u on M and a 1-form ξ on N, the pull-back of the 1-form ξ onto M can be defined as $\langle f^*\xi, u \rangle_x = \langle \xi, f_*u \rangle_{f(x)}$. For the present case the canonical flat connection on the base manifold M, expressed as $A(x) = \sigma^* \omega = A^{\alpha}{}_{\mu}(x) X_{\alpha} dx^{\mu}$ is a Lie algebra-valued map on tangent vectors $u \in T_xM$ and we have:

$$
\langle A, \mathbf{u} \rangle_x = \langle \sigma^* \circ \pi_2^* \theta, \mathbf{u} \rangle_x = \langle \theta, \pi_{2*} \circ \sigma_* \mathbf{u} \rangle_{g = \pi_2 \circ \sigma(x)} \tag{3.43}
$$

where in the latter expression the vector $u \in T_xM$ has been 'pushed forward' through the two maps to a vector in the tangent space of the group manifold. In general $\langle A, u \rangle \neq 0$, even for a flat connection, since an arbitrary trivialisation can be used to define the section map $\sigma_r(x) \equiv \psi_r^{-1}(x, e)_r$. However, for the canonical flat connection on P the horizontal subspace is everywhere tangent to a submanifold $M \times \{g\}$ for some $g \in G$ and the Frobenius criterion of equation 3.7 is satisfied. Hence in this case the section map from M to P may be chosen to coincide with the horizontal section of the canonical flat connection and we have:

$$
\langle A, \mathbf{u} \rangle_x = \langle \sigma^* \omega, \mathbf{u} \rangle_x = \langle \omega, \sigma_* \mathbf{u} \rangle_{p = \sigma(x)} = 0 \tag{3.44}
$$

since for all $u \in T_xM$ we have $\sigma_*u \in H_pP$ in this case, and hence we have $A(x) = 0$ in this choice of gauge section. In general the cross-section σ and horizontal subspace H_pP are distinct objects on P , as indicated for example in figure 3.5, relating to the gauge choice $g(x)$ and connection ω respectively. As can be seen from equations 3.26 and 3.27 if it is possible to choose a direct product basis to coincide with the horizontal lift basis on P then $\omega^{\alpha}_{a}(x, g) = 0$, that is all connection coefficients vanish for this choice of section.

Here we have described the flat connection that was introduced in equations 2.30 and 2.35 directly on the base manifold without constructing the principle fibre bundle. The use of the principle bundle will be more significant for the case of an enlarged symmetry group of $L(v) = 1$ as introduced in section 2.3 and studied further in section 5.1.

3.3 Riemannian Geometry

Any n-dimensional differentiable manifold M is canonically associated with the principle fibre bundle of frames FM, with structure group $GL^+(n,\mathbb{R})$, which preserves the orientation of the frames, over M as the base manifold. A linear connection $\tilde{\omega}$ can be defined on a frame bundle as a $gl(n, \mathbb{R})$ -valued 1-form on FM which may be written $\widetilde{\omega} = \widetilde{\omega}_b^a E_{a}^b$. The quantities $\widetilde{\omega}_b^a = \widetilde{\omega}_{b i}^a e^i$ (with $\{e^i\}$ a basis of 1-forms on the frame bundle) are a set of n^2 1-forms on FM. Each 1-form $\tilde{\omega}_b^a$ is associated with a basis element of $gl(n, \mathbb{R})$ represented by the $n \times n$ matrix E_{a}^{b} for which the only non-zero entry is a '1' in the a^{th} -row and b^{th} -column, that is $(E^b{}_a)^d{}_c = \delta^b_c \delta^d_a$ (where $\{a, b\}$ label the matrices and $\{c, d\}$ label the matrix elements. By comparison the generators of $SO(n)$, as described in equation 2.31, form a subalgebra of $gl(n, \mathbb{R})$ with matrices of the form $L_{pq} = E^q{}_p - E^p{}_q$.

The frame field $\{e_a\}$ on the base space M is a general basis which in some situations may be taken to be an orthonormal or coordinate basis. A section σ on FM corresponds to a choice of frame, that is a basis $\{e_a\}$, at each point of the base space M, with the pull-back $\Gamma = \sigma^* \tilde{\omega}$ being the representative of $\tilde{\omega}$ under this section. This linear connection 1-form Γ on M has components $\Gamma^a_{\ b} = \Gamma^a_{\ bc} e^c$, where $\{e^a\}$ is a coframe basis for T^*M .

In general for a gauge symmetry group with generators represented by $m \times m$ matrices $E_{\alpha} \in L(G)$ the connection components, for an arbitrary coframe $\{e^{a}\}\$ on the base manifold, may be written $A^r{}_s = A^{\alpha}{}_a (E_{\alpha})^r{}_s e^a = A^r{}_{sa} e^a$, with $\{r, s\} = 1 \dots m$, composing a matrix of 1-forms. In the case of a linear connection on M, with $u(x)$ as any tangent vector field, $\Gamma^a_{\ b}(\boldsymbol{u}) = \Gamma^a_{\ bc} u^c$ is a matrix element with $\Gamma^a_{\ \ bc}(x)$ being the components of the linear connection.

The covariant derivative D_a for the case of a linear connection on the external tangent space will be denoted by the kernel symbol ∇. With respect to a general frame field $\{e_a\}$, the components of the corresponding linear connection Γ^a_{bc} satisfy the relation $\nabla e_b = \Gamma^a{}_{bc} e^c \otimes e_a$, that is:

$$
\nabla_c e_b = \Gamma^a{}_{bc} e_a
$$
\nand hence,
$$
\Gamma^a{}_{bc} = \langle e^a, \nabla_c e_b \rangle
$$
\n(3.45)

where in the final term the angular brackets, defined in equation 2.18, denote the

1-form e^a mapping the vector field $\nabla_c e_b$ into the space of real numbers, that is the coefficients Γ^a_{bc} .

The linear connection coefficients Γ^a_{bc} transform under a general change of basis to $e_{b'} = e_a e^a{}_{b'}(x)$, with primed indices denoting the new frame and the matrix $e^{a}_{b'}(x) \in \mathrm{GL}^{+}(n,\mathbb{R}),$ as:

$$
\Gamma^{a'}_{\ b'c'} = (e^{-1})^{a'}_{\ d}\, e^{e}_{\ b'}\, e^{f}_{\ c'} \Gamma^{d}_{\ e f} + (e^{-1})^{a'}_{\ d}\, e_{c'}\, e^{d}_{\ b'} \tag{3.46}
$$

Compared with the gauge transformation of equation 3.41 an extra e^f $_{c'}$ factor appears here for the 3-index affine connection to reflect the tensor-like transformation law of the 1-form part of the connection under a local change of frame on the manifold M.

A subset of frames is provided by a general coordinate chart on the patch $U \subset M$ for which a section of the general frame bundle $\sigma(x): U \to FM$ is given by the coordinate basis $x \to {\partial_\mu}_x$. This defines a holonomic frame ${\partial_\mu}$, with ${\partial_\mu}, {\partial_\nu} = 0$, through which a local representative of the linear connection $\Gamma = \sigma^* \tilde{\omega}$ may be obtained. A second general coordinate chart with coordinate frame section $\{\partial_{\mu'}\}$ defines a further representative of the linear connection $\Gamma' = \sigma'^* \widetilde{\omega}$. The transition function $j(x) : M \to$
 $\widetilde{\Omega}^+(A, \mathbb{R})$ is a model in the set of $GL^+(4,\mathbb{R})$ for all $x \in M$ relates coordinate frames as:

$$
\partial_{\mu'}(x) = \partial_{\nu}(x) j^{\nu}{}_{\mu'}(x) \tag{3.47}
$$

where $j^{\nu}_{\mu'}(x) = \partial x^{\nu}/\partial x^{\mu'}$ is the Jacobian matrix of the general coordinate transformation. These transformations form a special case for equation 3.46 corresponding to a change of coordinate system $\{x^{\mu}\}\rightarrow \{x^{\mu'}\}$ on M.

If M is an *n*-dimensional Riemannian or pseudo-Riemannian manifold (M, g) , that is given a metric field with components $g_{\mu\nu}(x)$ on the manifold, a subset of distinguished frames may be identified which are orthonormal with respect to the metric. This subset of frames over M reduces the total space of FM to a submanifold $OM \subset FM$ which is itself a principle fibre bundle with structure group $SO^+(p,q)$ (or more generally $O(p,q)$) with $p + q = n$. There is a one-to-one correspondence between metric fields $g_{\mu\nu}(x)$ on M and reductions of the structure group $GL^+(n,\mathbb{R})$ to $SO^+(p,q)$ on FM, with each choice of field $g_{\mu\nu}(x)$ isolating one out of the many possible isomorphic copies of principle $SO^+(p,q)$ -bundles.

From the above general case we next consider specifically the spacetime symmetry of a 4-dimensional manifold M_4 . Matrices $l^a_{\ b}(x) \in \text{SO}^+(1,3)$ of the Lorentz group describe spacetime orientation preserving gauge transformations between sections of the principle bundle of orthonormal frames. With the set of vector fields ${e_a(x)}$ for each $x \in M$ now representing such an orthonormal frame, any other orthonormal frame can be expressed as:

$$
e_{b'}(x) = e_a(x) l^a_{b'}(x)
$$
\n(3.48)

while the dual coframe transforms as $e^{b'}(x) = (l^{-1})^{b'}$ $a'_a(x)e^a(x)$. Equation 3.48 expresses the right action of elements of the Lorentz group on the frame field. Since the set of orthonormal frames on the tangent space at any one point $x \in M_4$ is isomorphic to the Lorentz group, through equation 3.48, a principle fibre bundle over M_4 is obtained, with both the fibre space and structure group being $SO^+(1,3)$ itself. It is a reduction of the principle bundle of general linear frames $FM₄$, the latter having fibres isomorphic to the larger group $GL^+(4,\mathbb{R})$.

We can consider a tetrad field $e^a_{\mu}(x)$ as describing an element of a restricted set of the gauge group $GL^+(4,\mathbb{R})$ of all possible orientation-preserving frame transformations over M_4 or, in bridging local orthonormal frames with general coordinate frames, as a mapping between the principle bundle of Lorentz frames and the principle bundle of coordinate frames. That is, $e^a_{\mu}(x)$ relates a section of orthonormal frames ${e_a}_x$ with a coordinate frame basis ${\partial_\mu}_x$ via the right action:

$$
\partial_{\mu}(x) = e_a(x) e^a_{\mu}(x) \tag{3.49}
$$

with $e^a_{\mu}(x) \in GL^+(4,\mathbb{R})$, which can be directly compared to equation 3.48 with the transformation $l^a{}_b(x) \in \text{SO}^+(1,3)$.

For the spacetime metric $g(x)$ on M_4 any local orthonormal frame $\{e_a\}$ is associated with the Minkowski metric $\eta_{ab} = g(e_a, e_b) = \text{diag}(+1, -1, -1, -1)$, while in a general coordinate system the components of the metric are determined by the tetrad field $e^a_{\mu}(x)$ (similarly as we had in equation 2.16 for the 3-dimensional model):

$$
g_{\mu\nu}(x) = e^a{}_{\mu}(x)e^b{}_{\nu}(x)\eta_{ab} \tag{3.50}
$$

The $SO^+(1,3)$ bundle OM_4 may be *extended* to the frame bundle FM_4 with an $SO^+(1,3)$ -valued Lorentz connection $A(x)$ uniquely inducing a linear connection $\Gamma(x)$ for the extended bundle space. Such a $\text{GL}^+(4,\mathbb{R})$ -valued linear connection Γ is compatible with the metric, that is $\nabla g = 0$, while Γ and g need not be related in the general case. The principle bundle of orthonormal frames $OM₄$, equipped with a Lorentz connection, as a subbundle of the principle bundle of general linear frames $FM₄$ over the base manifold hence induces a unique metric connection on the latter space.

Expressing the Lorentz connection in a coordinate basis on M_4 as $A(x) =$ $A_{\mu}(x)dx^{\mu}$ the tetrad components may be considered as a local gauge transformation – that is as a change from a choice of local orthonormal Lorentz frames to the general coordinate frames over the base manifold, within the $GL^+(4,\mathbb{R})$ freedom of the principle bundle $FM₄$. In this way, and by comparison with equation 3.41 for example, the metric preserving linear connection Γ for a general coordinate system may be defined by:

$$
\Gamma^{\lambda}_{\ \mu\nu} = e^{\lambda}_{\ a} A^{a}_{\ b\nu} e^{b}_{\ \mu} + e^{\lambda}_{\ a} \partial_{\nu} e^{a}_{\ \mu} \tag{3.51}
$$

The identification of the linear connection Γ in this form implies that the covariant derivative of the tetrad field vanishes identically:

$$
\nabla_{\mu} e^{a}{}_{\nu} = \partial_{\mu} e^{a}{}_{\nu} + A^{a}{}_{b\mu} e^{b}{}_{\nu} - \Gamma^{\lambda}{}_{\nu\mu} e^{a}{}_{\lambda} = 0 \tag{3.52}
$$

This condition itself implies that A and Γ are *compatible* connections, regardless of the value of the torsion (defined below). In this case the tetrad field $e^a_{\mu}(x)$ 'commutes' with the operation ∇ of covariant differentiation. This means that the operation of interchanging between local field components, such as $u^a(x)$, and general coordinate tangent space field components, such as $u^{\mu}(x)$, via the tetrad field $e^a_{\mu}(x)$, applies in a straightforward manner even for equations involving covariant derivatives.

In particular, since $g_{\mu\nu}(x)$ has the form of equation 3.50 and the Minkowski metric is a constant, the metric field $g(x)$ is preserved by covariant differentiation defined in terms of the linear connection $\Gamma(x)$, which in turn is defined in terms of the Lorentz connection through equation 3.51, that is $\nabla g = 0$ as cited above. If $A^a_{b\mu}(x)$ is chosen to be the unique torsion-free Lorentz connection for a given tetrad field $e^a_{\mu}(x)$, then the corresponding linear connection Γ is the unique torsion-free metric connection expressed in a general coordinate system. This is the Levi-Civita connection, significant for general relativity, which can be written uniquely as a function of the metric tensor components $g_{\mu\nu}(x)$ as:

$$
\Gamma^{\sigma}{}_{\mu\nu} = \frac{1}{2} g^{\sigma\rho} (\partial_{\mu} g_{\rho\nu} + \partial_{\nu} g_{\mu\rho} - \partial_{\rho} g_{\mu\nu}) \tag{3.53}
$$

On the space of the frame bundle over any n -dimensional differentiable manifold M, even without a metric, a canonical \mathbb{R}^n -valued 1-form $\theta_{\rm C} = \theta^a E_a$ can be identified, with each θ^a being a 1-form on FM and $\{E_a\}$ a basis for \mathbb{R}^n , such that at any point $f \in FM$ and for any vector $X \in T_fFM$ we have:

$$
\langle \theta^a, X \rangle := \langle e^a, \pi_* X \rangle = (\pi_* X)^a \tag{3.54}
$$

which is just the components of the projection of X onto the base space M in the frame $f = \{e_a\}$ itself. Given a section $\sigma(x) = f$ on FM the pull-back $e^a = \sigma^* \theta^a$ describes the dual basis vectors of the general $GL(n, \mathbb{R})$ frame f.

The canonical 1-form $\theta_{\rm C}$ is therefore horizontal and equivariant and hence a tensorial form on FM. Given a linear connection $\tilde{\omega}$ on FM the exterior covariant derivative $\Theta = D\theta_C$ is called the *torsion* 2-form on FM. With $\Theta = \Theta^a E_a$, and following equation 3.14, the torsion can be expressed as:

$$
\Theta^a = d\theta^a + \tilde{\omega}^a{}_b \wedge \theta^b \tag{3.55}
$$

This object in turn pulls back to the torsion 2-form $T = \sigma^* \Theta$ on the base manifold M with coefficients T^a_{bc} defined in $T^a = \frac{1}{2}$ $\frac{1}{2}T^a_{\ bc}e^b\wedge e^c$, with:

$$
T^a = de^a + \Gamma^a{}_b \wedge e^b \tag{3.56}
$$

$$
= -\frac{1}{2}c^a{}_{bc}e^b \wedge e^c + \Gamma^a{}_{bc}e^c \wedge e^b \tag{3.57}
$$

$$
= (-\frac{1}{2}c^{a}_{bc} - \frac{1}{2}(\Gamma^{a}_{bc} - \Gamma^{a}_{cb}))e^{b} \wedge e^{c}
$$
 (3.58)

where each term above is a 2-form. Hence for a general linear connection on the manifold M the torsion components can be written as:

$$
T^{a}_{\;bc} = -2\Gamma^{a}_{\;[bc]} - c^{a}_{\;bc} \tag{3.59}
$$

with [...] denoting $\frac{1}{n!}$ times the antisymmetrised sum of the n! terms obtained through permuting the *n* enclosed indices. Via the vielbein field $e^a_{\mu}(x)$ this may be written in a general coordinate frame as:

$$
T^{\rho}_{\ \mu\nu} = -\Gamma^{\rho}_{\ \mu\nu} + \Gamma^{\rho}_{\ \nu\mu} \tag{3.60}
$$

since $[\partial_{\mu}, \partial_{\nu}] = 0$ for such a holonomic frame.

The curvature of the linear connection may also be defined on the frame bundle FM as $\Omega = D\tilde{\omega}$, that is as the exterior covariant derivative of the connection in the usual way, to obtain the tensorial form Ω of type $(Ad, gl(n, \mathbb{R}))$. However, here we deal directly with objects on the base manifold M for an arbitrary frame field $\{e_a\}$ and study the Riemannian curvature $\mathbf{R} = \sigma^* \tilde{\Omega} = R^a{}_b E^b{}_a$, where the matrices $E^b{}_a$ were defined in the opening of this section. From the definition of the curvature 2 form in equations 3.15–3.17 and the gl(n, R) commutators $[E^b_{a}, E^d_{c}] = \delta^b_c E^d_{a} - \delta^d_a E^b_{c}$ (which can be compared with the commutators for the L_{pq} matrices describing the so(n) subalgebra in equation 2.48) the components of curvature $R^a_{\ b}$ may be written for any linear connection Γ in any choice of frame field as:

$$
R^{a}_{b} = d\Gamma^{a}_{b} + \Gamma^{a}_{d} \wedge \Gamma^{d}_{b}
$$

\n
$$
= (d\Gamma^{a}_{bc})e^{c} + \Gamma^{a}_{bd}de^{d} + \Gamma^{a}_{dc}e^{c} \wedge \Gamma^{d}_{be}e^{e}
$$

\n
$$
= (e_{e}\Gamma^{a}_{bc})e^{e} \wedge e^{c} - \frac{1}{2}\Gamma^{a}_{bd}c^{d}_{ce}e^{c} \wedge e^{e} + \Gamma^{a}_{dc}\Gamma^{d}_{be}e^{c} \wedge e^{e}
$$

\n
$$
= \frac{1}{2}(e_{c}\Gamma^{a}_{be} - e_{e}\Gamma^{a}_{bc} + \Gamma^{a}_{dc}\Gamma^{d}_{be} - \Gamma^{a}_{de}\Gamma^{d}_{bc} - \Gamma^{a}_{bd}c^{d}_{ce})e^{c} \wedge e^{e}
$$
(3.61)

In terms of the components of the rank-4 Riemann tensor the curvature can be expressed as $\bm{R} = \frac{1}{2}R^a{}_{bcd}e^c \wedge e^d E^b{}_{a}$. Hence the curvature components on the base manifold M can be written in terms of the linear connection and structure coefficients as:

$$
R^{a}_{\;\; bcd} = e_{c} \Gamma^{a}_{\;\; bd} - e_{d} \Gamma^{a}_{\;\; bc} + \Gamma^{a}_{\;\; ec} \Gamma^{e}_{\;\; bd} - \Gamma^{a}_{\;\; ed} \Gamma^{e}_{\;\; bc} - c^{e}_{\;\; cd} \Gamma^{a}_{\;\; be} \tag{3.62}
$$

If a metric g is also defined on M then $\{e_a\}$ may represent a local orthonormal frame field. In the dual covector basis ${e^a}$ the Riemann tensor may be written as:

$$
\mathbf{R} = \frac{1}{2} R^{\mu q}_{cd} L_{pq} e^c \wedge e^d
$$

$$
= R^{\mu q}_{cd} L_{pq} e^c \otimes e^d
$$
(3.63)

where the latter follows due to the asymmetric arrangement of the ${c, e}$ indices for the coefficients in the final line of equation 3.61. Under the group $SO^+(p,q)$ this object transforms as a rank-4 tensor which can be expressed in components in several equivalent ways, including:

$$
R^{a}_{bcd} = R^{pq}_{cd} (L_{pq})^{a}_{b}
$$

and
$$
R_{abcd} = \eta_{ae} R^{e}_{bcd}
$$
 (3.64)

This latter object is asymmetric in the indices $\{a, b\}$ as well as in $\{c, d\}$. The Riemann tensor in a general coordinate system, as described towards the end of section 2.2 in the context of the $SO(3)$ model on M_3 , may be obtained through the vielbein field $e^a_{\mu}(x)$, with the resulting components:

$$
R_{\rho\sigma\mu\nu} = e^a_{\ \rho} e^b_{\ \sigma} e^c_{\ \mu} e^d_{\ \nu} R_{abcd} \tag{3.65}
$$

Both the curvature and torsion may be considered properties of a linear connection Γ in general. Although they are related through the Ricci and Bianchi identities, respectively:

$$
R^{\rho}_{\ [\sigma\mu\nu]} = -T^{\rho}_{\ [\sigma\mu;\nu]} - T^{\rho}_{\ \kappa[\sigma}T^{\kappa}_{\ \mu\nu]} \tag{3.66}
$$

$$
R^{\rho}_{\sigma[\mu\nu;\tau]} = -R^{\rho}_{\sigma\kappa[\tau}T^{\kappa}_{\mu\nu]} \tag{3.67}
$$

(where ; τ denotes the covariant derivative ∇_{τ} with respect to the x^{τ} coordinate) the curvature and torsion are independent geometric concepts where either one may be non-zero while the other is zero. For example for the complete parallelism exhibited on a Lie group manifold G in terms of the self-parallel frame composed of left-invariant vector fields X_{α} on G, with each $\Gamma^{\alpha}_{\ \beta\gamma} = 0$, the curvature vanishes, as can be seen trivially from equation 3.62, while the torsion is finite, with $T^{\alpha}_{\ \beta\gamma} = -c^{\alpha}_{\ \beta\gamma}$, as determined directly by equation 3.59. On the other hand for the linear connection $\Gamma^{\alpha}_{\ \beta\gamma} = -\frac{1}{2}c^{\alpha}_{\ \beta\gamma}$, in the same basis on G , the curvature is finite while the torsion vanishes, as can also be seen from equations 3.62 and 3.59. This latter case is the unique Levi-Civita connection on a group manifold defined in terms of the Killing metric on G. In general the identities of equations 3.66 and 3.67 clearly simplify for the torsion-free case.

Returning to the case of 4-dimensional spacetime M_4 the quantities $R_{\rho\sigma\mu\nu}$ of equation 3.65 are the components of a general coordinate frame rank-4 tensor with transformations j^{μ} $V_{\nu'} \in GL^+(4,\mathbb{R})$, introduced in equation 3.47, acting on all indices under a change of coordinates. The most general rank-4 tensor on a 4-dimensional manifold has $4^4 = 256$ independent components. However the geometric origin and structure of the Riemann tensor results in considerably less freedom. In components $R_{\rho\sigma\mu\nu}$ is asymmetric in the first two indices $\{\rho,\sigma\}$ since it derives from a Lorentzvalued metric connection and also asymmetric in the final two indices $\{\mu, \nu\}$ since the curvature originates as a 2-form object. This reduces the number of free components down to $(6 \times 6) = 36$. For the torsion-free case considered here the Ricci identity in a general coordinate system of equation 3.66 reduces to simply:

$$
R_{\rho[\sigma\mu\nu]} = 0 \qquad (3.68)
$$

or
$$
R_{\rho\sigma\mu\nu} + R_{\rho\nu\sigma\mu} + R_{\rho\mu\nu\sigma} = 0
$$

where the second equation follows from the asymmetry of $R_{\rho\sigma\mu\nu}$ in the final two indices. This further constraint results in a final total of 20 independent components for the Riemann curvature tensor for the metric and torsion-free case.

The Ricci tensor may be defined as the 'trace' of the Riemann tensor $R_{\sigma\mu} =$ $R^{\rho}{}_{\sigma\mu\rho}$. This is also termed a 'contraction' of upper and lower indices in $R^{\rho}{}_{\sigma\mu\nu}$, which transform in a dual manner to each other under the action of $GL^+(4,\mathbb{R})$. Also for the Lorentz curvature tensor components $R^a{}_{b\mu\nu}$ transformations in the $\{a, b\}$ indices via the group $SO^+(1,3)$ are closely related to those in the $\{\mu,\nu\}$ indices via the holonomic subgroup of $GL^+(4,\mathbb{R})$ through the components of the tetrad field $e^a_{\mu}(x)$, and it is through the latter field that tensor contractions are again possible. In both cases the Lie algebra valued part of the curvature form possesses a transformation symmetry closely related to that of the r-form part in the tangent space of the base manifold. This, of course, is not the case for curvature forms derived for general principle bundles with the symmetry group composing the fibres unrelated to the local symmetry of the base space manifold, and hence an equivalent contraction does not exist for a gauge theory based on such an internal symmetry.

The Ricci tensor is symmetric and hence possesses 10 independent degrees of freedom, including the scalar curvature $R = g^{\mu\nu} R_{\mu\nu}$ (as distinct from the Riemann tensor denoted by a bold \boldsymbol{R} , as on the left-hand side equation 3.63). The utility of such expressions follows from the fact that the operation of contraction maps a tensor object onto another tensor, that is the contracted tensor also transforms as a representation of $GL^+(4,\mathbb{R})$. This tensor preserving property is shared by the operations of the covariant derivative and exterior algebra as we described earlier, and hence all of these operations are useful for identifying the equations of physics.

The remaining 10 components of the Riemann tensor, the non-Ricci part, are described by the Weyl tensor $C_{\rho\sigma\mu\nu}$, it is the trace-free part of $R_{\rho\sigma\mu\nu}$ (all contractions are zero) with which it shares the same symmetries. The trace-free property implies ten relations $C_{\sigma\mu} = C^{\rho}{}_{\sigma\mu\rho} = 0$ between the components of the Weyl tensor $C_{\rho\sigma\mu\nu}$ and hence only ten of them are independent. The Weyl tensor is also the conformally invariant part of the Riemann tensor, that is it is unchanged under a conformal transformation of the metric $g_{\mu\nu}(x) \to f(x)g_{\mu\nu}(x)$ where $f(x)$ is any smooth real function on M_4 . The twenty components of the Riemann tensor can be decomposed explicitly in terms of those of the Weyl tensor and Ricci tensor as:

$$
R_{\rho\sigma\mu\nu} = C_{\rho\sigma\mu\nu} + \frac{1}{2} (g_{\rho\mu} R_{\sigma\nu} - g_{\rho\nu} R_{\sigma\mu} - g_{\sigma\mu} R_{\rho\nu} + g_{\sigma\nu} R_{\rho\mu}) + \frac{1}{6} (g_{\rho\nu} g_{\sigma\mu} - g_{\rho\mu} g_{\sigma\nu}) R
$$
 (3.69)
that is:
$$
R^{\rho\sigma}{}_{\mu\nu} = C^{\rho\sigma}{}_{\mu\nu} + 2R^{[\rho}{}_{[\mu} g^{\sigma]}{}_{\nu]} - \frac{1}{3} R g^{[\rho}{}_{\mu} g^{\sigma]}{}_{\nu}
$$

The Bianchi identity of equation 3.67 for the curvature tensor in the torsion-free case is simply:

$$
R^{\rho}_{\sigma[\mu\nu;\tau]} = 0 \tag{3.70}
$$

$$
\Rightarrow \qquad (R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu})_{;\mu} = 0 \qquad (3.71)
$$

where the latter expression follows from the double contraction of the former. The Einstein tensor is defined as $G^{\mu\nu} := R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu}$. Hence the Einstein tensor $G^{\mu\nu}$, unlike its 'dual' geometric object the Ricci tensor $R^{\mu\nu}$, represents an identitically conserved quantity, that is $G^{\mu\nu}{}_{;\mu} = 0$, which is the origin of its central importance in the field equation of general relativity.

For general relativity in regions of 'empty space' with $T^{\mu\nu} = 0$ by the Einstein equations 3.75 we also have $G^{\mu\nu} = 0$ and hence $R^{\mu\nu} = 0$ and the manifold is said to be 'Ricci flat'. In this Ricci vacuum the Riemann tensor is simply $R^{\rho}{}_{\sigma\mu\nu} = C^{\rho}{}_{\sigma\mu\nu}$, as can be seen explicitly from equation 3.69. The spacetime curvature is then described in terms of the Weyl tensor $C_{\rho\sigma\mu\nu}$, yet in a way dependent upon the matter content in other spacetime regions as will be reviewed alongside equation 5.44 in section 5.2.

We note here that the various possible sign conventions for the expressions of general relativity can be distilled down to the \pm sign used for the right-hand side of just three expressions in the Riemannian geometry:

1) The metric tensor:

$$
\eta_{ab} = \text{diag}(+1, -1, -1, -1) \tag{3.72}
$$

With '+1' for the time component this is a natural convention for the present theory based on forms of temporal flow.

2) The Riemann tensor:

$$
R^{\rho}_{\sigma\mu\nu} = \partial_{\mu}\Gamma^{\rho}_{\sigma\nu} - \partial_{\nu}\Gamma^{\rho}_{\sigma\mu} + \Gamma^{\rho}_{\lambda\mu}\Gamma^{\lambda}_{\sigma\nu} - \Gamma^{\rho}_{\lambda\nu}\Gamma^{\lambda}_{\sigma\mu} \tag{3.73}
$$

Where the final term of equation 3.62 is zero when expressed in a coordinate frame as is the case here.

3) The Ricci tensor:

$$
R_{\mu\nu} = R^{\rho}_{\ \mu\nu\rho} \quad (= -R^{\rho}_{\ \mu\rho\nu}) \tag{3.74}
$$

This is equivalent to choosing the sign convention for the Einstein field equation as $G^{\mu\nu} = -\kappa T^{\mu\nu}$ with positive normalisation constant κ (as will be justified after equation 5.35).

The convention for these three signs chosen here is the same as used for example in ([5] p.24) that is with signs '(- + -)' relative to the original discussion of these conventions in [6]. The Einstein equation, and general relativity itself, will be reviewed in the following section.

3.4 General Relativity

In his 1854 work 'On the Hypotheses which lie at the Foundation of Geometry' Riemann, building upon the study of the intrinsic curvature of 2-dimensional surfaces by Gauss, considered more generally spaces of *n*-dimensions and introduced tensor analysis, in particular incorporating the metric tensor and the Riemann curvature tensor. At the same time Riemann also speculated on the possible curvature for the space of our own world, both on small and large scales, and its possible physical implications.

At around the same time (1861,1865) Maxwell, building upon the 'field' concept introduced earlier by Faraday based on empirical observations, formulated the equations of motion for the electromagnetic field, providing a unified description of electric fields, magnetic fields and also the properties of light.

The mathematical structure of general relativity was developed leading up to 1915 as an application of Riemann's work in geometry, with the dimension of time now included along with space in a 4-dimensional spacetime manifold. Influenced by the work of Maxwell on electromagnetism objects such as the metric and Riemann curvature tensor, as mathematical functions describing the phenomena of gravitation, were now considered as *fields* in spacetime.

In search of a relativistic gravitational field equation consistent with the 'equivalence principle', defined below, and under the empirical guidance that the Newton-Poisson equation $\nabla^2 \Phi = 4\pi G_N \rho$ (a second order differential equation, with Laplacian operator $\nabla^2 = \partial_x^2 + \partial_y^2 + \partial_z^2$, relating the gravitational scalar potential Φ , via Newton's constant G_N , to the scalar mass density distribution ρ) should emerge in the nonrelativistic limiting case for small distortions from a flat spacetime, Einstein converged in 1915 upon the field equation:

$$
G^{\mu\nu} = -\kappa T^{\mu\nu} \tag{3.75}
$$

with κ a constant and $T^{\mu\nu}$ the energy-momentum tensor for the distribution of matter in 4-dimensional spacetime. From the limit of Newtonian gravity the normalisation constant is found to be $\kappa = \frac{8\pi G_N}{c^4}$ $\frac{\sigma G_N}{c^4}$.

In general relativity, it is considered always possible to have a local *inertial* coordinate system on M_4 that is valid within a sufficiently small region of curved 4-dimensional spacetime – strictly an infinitesimal neighbourhood about any point $x \in M_4$, with local metric $\eta = \text{diag}(+1, -1, -1, -1)$.

The strong equivalence principle states that within such a local coordinate system, within a sufficiently small region about the point $x \in M_4$, all laws of physics, other than gravity, take the same form that applies for special relativity in an unaccelerated Cartesian coordinate system in the absence of gravity. These assumptions augment the weak form of the equivalence principle for which the 'laws of physics' are limited to 'the laws of motion of freely falling particles' corresponding to the equivalence of gravitational and inertial mass, and the observation of the apparent lack of gravitational effects within a freely falling lift.

The motion of a freely falling particle in such a local inertial coordinate system ${x^a}$ satisfies the equation $d^2x^a/d\tau^2 = 0$, in choosing the proper time τ to parametrise the trajectory. Transforming to a general coordinate system $\{x^{\mu}\}$ this becomes:

$$
\frac{d^2x^{\lambda}}{d\tau^2} + \Gamma^{\lambda}{}_{\mu\nu}\frac{dx^{\mu}}{d\tau}\frac{dx^{\nu}}{d\tau} = 0
$$
\n(3.76)

which is called the *geodesic* equation of motion and which is valid also in an extended curved spacetime. The quantities $\Gamma^{\lambda}{}_{\mu\nu}$ are the coefficients of the linear connection and the proper time τ itself can be defined in terms of an integral of the invariant intervals $d\tau = (g_{\mu\nu}dx^{\mu}dx^{\nu})^{1/2}$ along the trajectory. In terms of the 4-velocity $u^{\mu} = dx^{\mu}/d\tau$ the above geodesic equation can be written as simply:

$$
u^{\mu}\nabla_{\mu}u^{\nu} = 0 \tag{3.77}
$$

The equivalence principle states that all gravitational effects can be locally transformed away and can be interpreted to mean that we may always choose a local inertial coordinate frame at any $x \in M_4$ such that all the coefficients $\Gamma^{\lambda}{}_{\mu\nu} = 0$. Hence, although the coefficients of the non-tensor object Γ will be frame dependent the torsion tensor T vanishes in all reference frames, by equation 3.60. This torsionfree assumption for Einstein's theory of general relativity has the benefit of simplifying some of the mathematics of the theory, as for example in equations 3.68 and 3.70 of the previous section.

Given a metric $g_{\mu\nu}(x)$ on M_4 the Levi-Civita connection is the unique metric $(\nabla g = 0)$, torsion-free $(T = 0)$ linear connection. The corresponding connection coefficients may be written in a general coordinate frame uniquely in terms of those of the metric tensor as described in equation 3.53. For such a connection equations 3.76 and 3.77 describe the trajectory which extremises the path length between any given end points:

$$
L = \int (g_{\mu\nu}u^{\mu}u^{\nu})^{1/2}d\tau
$$
\n(3.78)

and hence earns the name 'geodesic'. Further, for this connection with $\Gamma(x)$ determined uniquely by $g(x)$, as implied by the equivalence principle, the metric alone determines all gravitational effects and hence can be considered to be the gravitational field for Einstein's general relativity. Since the tetrad field $e^a_{\mu}(x)$ may be considered to be the 'square-root' of the metric, with $g_{\mu\nu} = e^a{}_{\mu}e^b{}_{\nu}\eta_{ab}$ in equation 3.50, the tetrad field itself, which everywhere exhibits the presence of the local inertial frames, may also be considered to represent the gravitational field.

As well as being able to express the metric as $g_{\mu\nu} = \text{diag}(+1, -1, -1, -1)$ at any spacetime location there is sufficient freedom under coordinate transformations such that at any $x \in M_4$ all 40 components of the metric derivatives can be set to zero, that is $g_{\mu\nu,\rho}(x) = 0$, corresponding to coordinate frames with $\Gamma = 0$ as can be seen from equation 3.53. However there is insufficient freedom under general coordinate transformations to set all 100 second derivative quantities $g_{\mu\nu,\rho\sigma}(x)$ to zero and there remain 20 irreducible degrees of freedom which are described by the Riemann curvature tensor, as deduced earlier after equation 3.68.

The components of the metric tensor field $g_{\mu\nu}(x)$ may be determined by solving the second order differential field equation $G^{\mu\nu} = -\kappa T^{\mu\nu}$ for a distribution of matter described by the energy-momentum tensor $T^{\mu\nu}$, in practice by introducing 'boundary conditions' as described in the following section. For a particular physical state for the geometry of the world there will be a range of possible solutions for $g_{\mu\nu}(x)$ and $e^a_{\mu}(x)$ in spacetime (over and above the local Lorentz freedom for the latter field) all with equivalent physical content.

Essentially there is only *one* 'coordinate system' \mathbb{R}^4 through which any region of spacetime may be described, as depicted in figure $3.6(a)$, as a simple space of 4 independent real parameters upon which a solution for the field $g_{\mu\nu}(x)$ may be inscribed.

Figure 3.6: (a) Alternative metric solutions on \mathbb{R}^4 for the same physical state and (b) as apparently represented through an 'alternative' coordinate system overlaid upon the 'original' coordinates.

An alternative expression of the same physical solution then corresponds to a different metric function $g'_{\mu\nu}(x)$ inscribed upon the same \mathbb{R}^4 space. For example in the Schwarzschild solution for the metric field associated with a single massive body located at one point in space, to be presented in equation 5.49, the physical point where the curvature scalar R is largest, and perhaps even singular, will in general have different coordinate values $x \in \mathbb{R}^4$ under a 'coordinate transformation', as indicated by the two small circles in figure $3.6(a)$. However the transformed solution could be conceived of as a new set of 'curvilinear' coordinates overlaying the same physical configuration (explicitly represented by the *same* metric field) as shown in figure $3.6(b)$.

In general it is less useful to think of any coordinates as curvilinear, indeed it is always the case that $[\partial_\mu, \partial_\nu] = 0$ with all structure coefficients $c^{\rho}_{\mu\nu} = 0$. In this sense all coordinate systems can be pictured as a 'flat' purely mathematical parameter space, which for the case of \mathbb{R}^4 can be visualised as the set of 'Euclidean' real number parameters as represented in figure 3.6(a). Physical curvature is a property of the fields on M_4 itself with the Riemann curvature tensor describing the geometrical structure and warping of the corresponding physical spacetime. The set of components $R^{\rho}_{\sigma\mu\nu}(x)$ are given at points on the manifold labelled $x \in M_4$ under the coordinate chart map $\phi: M \to \mathbb{R}^4$, or on a $U \subset M$ subset. A general coordinate transformation is then a mapping between solutions represented on different choices of the *map* $M \to \mathbb{R}^4$ onto a unique \mathbb{R}^4 (assuming here a non-degenerate Jacobian matrix $j^{\mu}_{\ \nu}(x)$, that is neglecting the artificial difference of a 'coordinate singularity' for example for polar coordinates at the corresponding Cartesian coordinate origin).

In general relativity a general coordinate system $\{x^{\mu}\}$ is of no physical significance; all the physics is in the 'fields' on the manifold (see for example [7] chapter 2), with the gravitational field $e^a_{\mu}(x)$ giving rise to the spacetime geometry of the manifold. It is the possibility of relating field quantities on M_4 , such as the coincidence of physical events or the equating of the Einstein tensor with the energy-momentum tensor, that determines the physical content of the theory.

While the coordinate system plays a passive unphysical role, in particular circumstances it may be associated with physical structure. This is true in the case of the Schwarzschild solution in which the origin of a polar coordinate system is associated with the central massive object. This is an example with non-zero Riemann curvature in which the exact spherical symmetry of the physical state is assumed to be exhibited by the metric for which a solution may be found in a greatly simplified form in a naturally preferred system of spherical polar coordinates. For similar reasons, but with finite 4-dimensional curvature considered on a much larger scale, cosmological models also employ a preferred system of coordinates to study solutions of Einstein's field equation, as we shall describe in section 12.2.

In general, however, there will be no preferred solutions and hence no privileged coordinate systems on the base manifold. In this sense all coordinate systems are 'equally bad', or at least on a equal footing, and this expresses the relevance of general covariance for general relativity. Other theories may also be 'generally covariant', but if there is always a particular kind of distinguished coordinate reference frame then the general covariance may be of no relevance. This is the case for special relativity and also for Newtonian mechanics formulated against a flat absolute background of an independent space and time.

Even for general relativity, if the curvature is very small, as it is in practice in a laboratory on the surface of the Earth or even locally within the solar system with respect to the 'fixed stars' of the galaxy, then there will be 'preferred' solutions with everywhere $e^a_{\mu}(x) \simeq \delta^a_{\mu}$ and $g_{\mu\nu}(x) \simeq diag(1, -1, -1, -1)$ found for a coordinate system which is then implicitly pseudo-Euclidean to a very good approximation. In the limit of flat Minkowski spacetime there is a preferred coordinate systems with $g_{\mu\nu}(x) = \text{diag}(1, -1, -1, -1)$ exactly. The corresponding tetrad field is $e^a_{\mu}(x) = \delta^a_{\mu}$, within a global Lorentz transformation (which leaves the metric invariant). In this case a coordinate transformation such that in general $g'_{\mu\nu}(x) \neq \text{diag}(1, -1, -1, -1)$, while the Riemann tensor necessarily remains zero, may be considered as an introduction of a new 'curvilinear' coordinate system, as pictured for example by the transformation in figure $3.6(b)$.

Newtonian mechanics in Euclidean spacetime takes its simplest form when expressed using Cartesian coordinates; however even for a flat space the description of parallel transport and the form of the covariant derivative is non-trivial when expressed in a curvilinear coordinate system. It is only for the choice of a Cartesian coordinate system that a trivial linear connection may be adopted. In general it is the fact that we can not assume a 'flat' geometry over macroscopic distances that necessitates the introduction of the more general notion of parallelism as described by a connection form, as is also the case for a gauge theory based on an internal symmetry as described in section 3.1. For an externally curved geometry the lack of a preferred coordinate system, with a preferred description of parallelism, highlights the significance of general covariance for the theory of general relativity.

As described above a choice of coordinates may be useful in order to express some metric solutions in a simple mathematical form but they are a non-physical, and in this sense a 'gauge', artifact that drop out of all expressions for observable quantities. Working with a general coordinate system and the corresponding use of holonomic reference frames $\{\partial_\mu\}$ does not allow for arbitrary frame transformations as elements of $GL^+(4,\mathbb{R})$. Rather the transition functions $j^{\mu}_{\ \nu}(x)$ of equation 3.47 are restricted to a 'holonomic subgroup' of all possible $GL^+(4,\mathbb{R})$ transformations over the manifold, sometimes called the 'Einstein gauge', and this to some extent disguises underlying gauge structure of general relativity.

Although the 'coordinate invariance' symmetry of the kind implied by general covariance is mathematically rather different from the usual concept of a 'gauge invariance' symmetry, there is a close analogy between them. In both cases there is a loosening of a global symmetry or absolute structure that would otherwise be arbitrarily imposed. In both cases also the equations of motion, together with their solutions, are mapped on to equally valid equations and solutions under the coordinate or gauge transformations. Further, while a particular choice of coordinates greatly assists with calculations for some solutions in general relativity a particular choice of gauge is frequently employed to assist with calculations in a gauge theory.

For general relativity to be considered in terms of a $GL^+(4,\mathbb{R})$ gauge theory of gravity, within the framework of general covariance, the equivalence principle is needed to distinguish the local Minkowski metric as being physically significant in that it marks the transition to special relativity in local inertial coordinate frames. That is, the metric or tetrad field needs to be introduced everywhere on M_4 (there is no equivalent of such fields for an internal symmetry gauge theory). This implies the possibility to contract the $GL^+(4,\mathbb{R})$ structure group down to the Lorentz subgroup (which is then the holonomy group of the general frame bundle). The local Lorentz symmetry itself has mathematical properties very closely related to those of the local symmetry of a gauge theory.

Indeed, while gravitation in Einstein's original theory of 1915 is described through the freedom of the metric field $g_{\mu\nu}(x)$ field, together with its relation to the Levi-Civita connection $\Gamma(x)$, an equivalent formulation of general relativity can be given in terms of the tetrad field $e^a_{\mu}(x)$ together with a Lorentz connection $A(x)$. This latter approach was introduced in 1956 by Utiyama [8] in which general relativity is considered as a type of gauge theory invariant under local Lorentz transformations. Such local transformations are displayed in equation 3.48 and map one local inertial coordinate frame onto another. As well as tensor representations the Lorentz group also has spinor representations. Hence spinor fields can be introduced on a spacetime

manifold with an arbitrary metric $g_{\mu\nu}(x)$ via the tetrad field $e^a_{\mu}(x)$. This also permits gravitation to be considered in terms of an $SL(2,\mathbb{C})$ gauge theory, where $SL(2,\mathbb{C})$ is the double cover of the Lorentz group, as will be described in section 7.3.

The fundamental structures on the base manifold are the local Minkowski spaces, together with their mutual relations through the Lorentz connection on M_4 . With respect to a given coordinate system either the tetrad $e^a_{\mu}(x)$ or metric $g_{\mu\nu}(x)$ field identifies the local inertial frames. In 1920 Einstein postulated that the metric field should be considered to be the fundamental entity of general relativity, referring to it as the 'new ether'. However, whichever fields are considered as fundamental, field equations are still required in order to determine the nature of the field dynamics. At the same time that Einstein arrived at equation 3.75 via the heuristic arguments outlined in the opening of this section Hilbert was in the process of deriving the same equation via a Lagrangian approach. This latter argument, and the employment of Lagrangian methods more generally, will be reviewed in the following section.

3.5 Lagrangian Formalism

In this section we review the standard use of the Lagrangian formalism to derive physical equations of motion, including those for general relativity and gauge theories. In the 4-dimensional spacetime of general relativity the scalar curvature R is adopted as the principle geometric contribution to the total scalar Lagrangian function, with the field equations determined from the Einstein-Hilbert action integral ([9] p.75):

$$
I = \int (\alpha (R - 2\Lambda) + \mathcal{L}) \sqrt{|g|} \, d^4x \tag{3.79}
$$

Here Λ is the cosmological constant, $\mathcal L$ is the Lagrangian function for matter fields and α is a normalisation constant. The magnitude of the metric determinant |g| is employed in the 4-dimensional invariant volume element $\sqrt{|g|} d^4x$. The vacuum equations for general relativity, that is with $\mathcal{L} = 0$ and $\Lambda = 0$, are obtained by requiring that $\delta I = 0$ in equation 3.79 under variation of the metric $\delta g_{\mu\nu}$. With $\delta g^{\mu\nu} = -\delta g_{\mu\nu}$ to first order, $\delta\sqrt{|g|}=\frac{1}{2}$ $\frac{1}{2}\sqrt{|g|} g^{\mu\nu}\delta g_{\mu\nu}$ and with $R = R_{\mu\nu}g^{\mu\nu}$ we have:

$$
\delta I = \int \alpha (R \,\delta \sqrt{|g|} + R_{\mu\nu} \,\delta g^{\mu\nu} \sqrt{|g|} + \delta R_{\mu\nu} \, g^{\mu\nu} \sqrt{|g|}) \, d^4x \tag{3.80}
$$

$$
= \int \alpha \left(\frac{1}{2} R g^{\mu\nu} - R^{\mu\nu}\right) \delta g_{\mu\nu} \sqrt{|g|} \, d^4 x \tag{3.81}
$$

where the final term in equation 3.80 contributes zero to the integral since $g^{\mu\nu}\delta R_{\mu\nu} =$ $(g^{\mu\nu}\delta\Gamma^{\rho}{}_{\mu\nu} - g^{\mu\rho}\delta\Gamma^{\nu}{}_{\mu\nu})_{;\rho}$ and $\delta\Gamma^{\rho}{}_{\mu\nu}$ vanishes on the boundary of integration ([9] p.75). Requiring the action to be stationary, $\delta I = 0$, for any variation of the metric, $\delta g_{\mu\nu}$, leads directly from equation 3.81 to the Einstein vacuum equation:

$$
G^{\mu\nu} := R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu} = 0 \tag{3.82}
$$

For the non-vacuum case the energy momentum tensor $T^{\mu\nu}$ for a general matter Lagrangian $\mathcal{L} \neq 0$ can be defined under variations of the metric $\delta g_{\mu\nu}$ through:

$$
\delta I = \delta \int \mathcal{L} \sqrt{|g|} d^4 x = \int \frac{1}{2} T^{\mu\nu} \delta g_{\mu\nu} \sqrt{|g|} d^4 x \qquad (3.83)
$$

Hence for the full action integral of equation 3.79 stationarity $\delta I = 0$ under the metric variation gives Einstein's field equation for the general case, with $\kappa \equiv \frac{-1}{2\alpha}$ adopted as the normalisation constant:

$$
G^{\mu\nu} + \Lambda g^{\mu\nu} = -\kappa T^{\mu\nu} \tag{3.84}
$$

Assuming that the matter Lagrangian may be a function of $g_{\mu\nu}(x)$, but not of the metric derivatives, the energy-momentum tensor itself, consistent with these equations, can be expressed directly in terms of the matter Lagrangian as:

$$
T^{\mu\nu} = \frac{2}{\sqrt{|g|}} \frac{\partial (\mathcal{L}\sqrt{|g|})}{\partial g_{\mu\nu}} = 2 \frac{\partial \mathcal{L}}{\partial g_{\mu\nu}} - \mathcal{L}g^{\mu\nu}
$$
(3.85)

A simpler application of the principle of least action in the context of general relativity was described earlier for equation 3.78 regarding the derivation of the geodesic equation of motion for a body moving in a gravitational field. Generalising from equation 3.78 for a body with mass m and charge q moving in a curved spacetime through an electromagnetic field with 4-potential $A_\mu(x)$ an action S may be constructed including both the kinematic and an interaction Lagrangian term respectively in:

$$
S = \int \left(m \left(g_{\mu\nu} u^{\mu} u^{\nu} \right)^{1/2} + q u^{\mu} A_{\mu} \right) d\tau \tag{3.86}
$$

Requiring $\delta S = 0$ under variation of the trajectory of the charged body leads to the equation of motion:

$$
m\left(\frac{du^{\lambda}}{d\tau} + \Gamma^{\lambda}{}_{\mu\nu}u^{\mu}u^{\nu}\right) = +F^{\lambda}{}_{\sigma}J^{\sigma}
$$
\n(3.87)

where F^{λ}_{σ} are components of the electromagnetic field tensor and $J^{\mu} = q u^{\mu}$ is the 4-current of the charged body having 4-velocity u^{μ} with respect to the proper time τ . The above equation hence describes a correction to the purely geodesic trajectory of equation 3.76. In the limit of a flat Minkowski spacetime, and will respect to a Cartesian coordinate frame, equation 3.87 simplifies to:

$$
\frac{\partial p^b}{\partial \tau} = +F^b_{\ c} J^c \tag{3.88}
$$

which is the relativistic Lorentz force law, for the charged body with 4-momentum $p^{b} =$ mu^b . Further, in the non-relativistic limit equation 3.88 becomes $m\mathbf{a} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$, the original form of the Lorentz force law, where v and a are the 3-velocity and 3acceleration of the body respectively.

These examples, for the trajectory of a body in a gravitational and/or electromagnetic field, demonstrate the flexibility and generality of the Lagrangian approach. As well as applying to macroscopic physical bodies the use of Lagrangian functions is a standard tool in classical field theory. In general the form of the Lagrangian \mathcal{L} , a function of the fields such as $\phi(x)$, guided by considerations of symmetry, is constructed in order that the requirement for the action integral $S = \int \mathcal{L}(\phi, \partial_{\mu}\phi)\omega$ (where ω is the volume 4-form) to be stationary, $\delta S = 0$, under variations of the fields, such as $\delta \phi$, yields the required equations of motion for the fields via the Euler-Lagrange equation:

$$
\partial_{\mu} \frac{\partial \mathcal{L}}{\partial(\partial_{\mu} \phi)} - \frac{\partial \mathcal{L}}{\partial \phi} = 0.
$$
 (3.89)

In a flat spacetime, in terms of the electromagnetic curvature tensor $F_{\mu\nu}$, Maxwell's equations are:

$$
F_{\left[\mu\nu,\rho\right]} = 0 \tag{3.90}
$$

$$
F^{\mu\nu}{}_{,\mu} = +J^{\nu} \tag{3.91}
$$

which can also be written as $dF = 0$ and $d^*F = *J$ respectively (where '*' denotes the 'Hodge dual' as employed in equation 5.24). These equations are equally valid in a curved spacetime on replacing the partial derivatives ', ρ ' by the covariant derivatives \cdot ; ρ , as an application of the strong principle of equivalence. The first of these equations is simply the Bianchi identity, introduced in section 3.2, for the curvature tensor of a U(1) gauge theory. Here working in the Lorenz gauge with $\partial_{\mu}A^{\mu} = 0$ the inhomogeneous Maxwell equation 3.91 can be written as:

$$
\Box A^{\mu} = +J^{\mu} \tag{3.92}
$$

The Maxwell Lagrangian for the electromagnetic field is constructed as:

$$
\mathcal{L}_{em} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \tag{3.93}
$$

Under variation of the electromagnetic gauge field $A_\mu(x)$ the Euler-Lagrange equation for \mathcal{L}_{em} yields Maxwell's equation for the source-free $J^{\nu} = 0$ case, that is $F^{\mu\nu}{}_{,\mu} = 0$. In combining the Lagrangian of equation 3.93 with the final term of that in equation 3.86, hence including a term coupling the electromagnetic field to a classical charged body, the corresponding Euler-Lagrange equation for $\delta A_\mu(x)$ yields equation 3.91 with the source term on the right-hand side.

The form of the Lagrangian for non-Abelian gauge theory is guided by the Abelian case of electromagnetism, motivating the Lorentz and gauge invariant Yang-Mills Lagrangian:

$$
\mathcal{L}_{\text{YM}} = -\frac{1}{4} F_{\alpha\,\mu\nu} F^{\alpha\,\mu\nu} \tag{3.94}
$$

as a direct generalisation of equation 3.93. For the non-Abelian case there is a further contraction over the index $\alpha = 1 \dots n_G$, for the group generators, between the adjoint and coadjoint representations, which are related by the Killing metric $K_{\alpha\beta}$ (which in a suitable basis is simply $-\delta_{\alpha\beta}$ for the compact simple Lie groups relevant for the internal gauge symmetries in particle physics). In this case the Euler-Lagrange equation 3.89 for \mathcal{L}_{YM} under variation of the gauge field components $Y^{\alpha}_{\mu}(x)$ yields the non-linear second order differential equation:

$$
D_{\mu}F^{\alpha\,\mu\nu} = \partial_{\mu}F^{\alpha\,\mu\nu} + c^{\alpha}_{\beta\gamma}Y^{\beta}_{\mu}F^{\gamma\,\mu\nu} = 0 \tag{3.95}
$$

where D_{μ} is the gauge covariant derivative, which also appears in the Bianchi identity $D_{\lbrack\rho}F^{\alpha}_{\mu\nu]}=0$ as the non-Abelian generalisation of equation 3.90. The immediate distinctive feature of equation 3.95, in comparison with the Maxwell equation 3.91, is the additional non-linear term of the form $[Y, F]$ appearing for the non-Abelian case. Such terms are interpreted as self-interactions of the gauge fields $Y_\mu(x)$, which do not occur for the Maxwell theory. This self-interaction is intrinsically geometric in origin and is implied in the Lagrangian of equation 3.94 itself given that the curvature for a non-Abelian gauge field has the form of equation 3.38 with non-trivial structure constants.

Additional terms in the Lagrangian, either for the Maxwell or Yang-Mills case, may lead to further sources of interactions. In the Standard Model of particle physics interactions between fermion and gauge fields in the corresponding equations of motion are introduced through the 'minimal coupling' in the covariant derivative terms included in a Lagrangian. For example by including \mathcal{L}_{YM} alongside the Dirac Lagrangian for a massless spinor field $\psi(x)$, which transforms as a multiplet under the internal symmetry, together with the conjugate field $\overline{\psi} = \psi^{\dagger} \gamma^0$ (where the γ -matrices will be defined in section 7.1), we have combined:

$$
\mathcal{L}_{\text{YMD}} = -\frac{1}{4} F_{\alpha\,\mu\nu} F^{\alpha\,\mu\nu} - \overline{\psi} \gamma^{\mu} D_{\mu} \psi \tag{3.96}
$$

where
\n
$$
D_{\mu} = \partial_{\mu} + Y^{\alpha}_{\ \mu} E_{\alpha}
$$
\nand
\n
$$
\overline{\psi} \gamma^{\mu} D_{\mu} \psi = \overline{\psi} \gamma^{\mu} \partial_{\mu} \psi + Y^{\alpha}_{\ \mu} j^{\mu}_{\ \alpha}
$$
\nwith
\n
$$
j^{\mu}_{\ \alpha} = \overline{\psi} \gamma^{\mu} E_{\alpha} \psi
$$
\n(3.97)

where in the appropriate representation the E_{α} are $n \times n$ matrices acting on the *n*dimensional field ψ in the internal space. Here a lower case 'j' will generally denote a current such as the Lorentz vector in equation 3.97 composed of elementary fields, as opposed to the upper case analogue $J^{\mu} = qu^{\mu}$ for the macroscopic current featuring in equation 3.87 for example. Under variation of the gauge field $Y^{\alpha}_{\mu}(x)$ the extra term $Y^{\alpha}_{\mu}j^{\mu}_{\alpha}$ in this Lagrangian leads to a modification of equation 3.95 with the source $j^{\mu}(\alpha)$ now appearing in the right-hand side to give:

$$
D_{\mu}F^{\alpha\,\mu\nu} = \partial_{\mu}F^{\alpha\,\mu\nu} + c^{\alpha}{}_{\beta\gamma}Y^{\beta}{}_{\mu}F^{\gamma\,\mu\nu} = j^{\nu\,\alpha} \tag{3.98}
$$

In practice factors of $i = \sqrt{-1}$ and differing \pm signs in the above equations will depend upon the conventions adopted, with coupling constants such as q also appearing in expressions for specific applications in the Standard Model as will be reviewed in section 7.2. In addition to the requirements of symmetry the form of the scalar Lagrangian function is typically heavily guided by the need to obtain the desired equations of motion. As a further example the above Lagrangian of equation 3.96, augmented with a fermion mass term $+m\psi\psi$, under variation of the field $\psi(x)$ yields the Euler-Lagrange equation:

$$
(\gamma^{\mu}D_{\mu} - m)\psi = 0 \tag{3.99}
$$

which is the Dirac equation for the spinor field ψ (within conventional factors of i). The interaction between the fermion field $\psi(x)$ and the gauge field $Y_{\mu}(x) = Y^{\alpha}_{\mu}(x)E_{\alpha}$ is here found in the 'minimal coupling' in the action of the covariant derivative $D_{\mu}\psi =$ $\partial_{\mu}\psi + Y_{\mu}\psi$ in the kinetic term of the Lagrangian in equation 3.96.

As for the case of the charged macroscopic body in equation 3.86 here also the mass m for the field ψ in equation 3.99 has been introduced through a Lagrangian mass term, in this case with $+m\psi\psi$ appended to equation 3.96. Mass terms are generally added to the Lagrangian by hand in this way, although this may not be straightforward to achieve. For example, a corresponding Lagrangian term such as $m^2 Y_\mu Y^\mu$ for a gauge field mass is prohibited by the requirement of gauge invariance, and even the fermion mass term $m\overline{\psi}\psi$ is prohibited in the Standard Model Lagrangian due to the left-right asymmetry of the $SU(2)_L$ -valued gauge field relating to electroweak interactions. In both cases mass terms are incorporated into the Lagrangian through interactions with the Higgs field and spontaneous symmetry breaking, involving the addition of further, apparently ad hoc, terms to the Lagrangian, as will be described in section 7.2.

As described above interactions may be introduced into the Lagrangian by the requirement of invariance under a local gauge symmetry. Such a local symmetry incorporates a corresponding global symmetry of the equations of motion and hence Noether's theorem applies. The theorem states that each *global* continuous symmetry is associated with a conserved current, written in terms of the field $\phi(x)$ as:

$$
j^{\nu}_{\ \alpha} := \left(\frac{\partial \mathcal{L}}{\partial(\partial_{\nu}\phi^{a})}\right) (E_{\alpha})^{a}{}_{b}\phi^{b} \tag{3.100}
$$

for each generator E_{α} of the global symmetry. For the Dirac Lagrangian with a U(1) gauge symmetry, that is the final term of equation 3.96 for the Abelian case, the global $U(1)$ symmetry with a single generator is associated with the Dirac current:

$$
j^{\mu} = \overline{\psi}\gamma^{\mu}\psi \tag{3.101}
$$

and the conservation law is simply $\partial_{\mu}j^{\mu} = 0$.

In contrast to the case of an *internal* global symmetry of the Lagrangian applying Noether's theorem for the external symmetry of global translational invariance of $\mathcal{L}(\phi)$ in a flat Minkowski spacetime leads to the quantity ([10] p.27):

$$
t^{\mu\nu} = \left(\frac{\partial \mathcal{L}}{\partial(\partial_{\mu}\phi)}\right)\partial^{\nu}\phi - \mathcal{L}\eta^{\mu\nu} \tag{3.102}
$$

which satisfies the conservation law $\partial_{\mu}t^{\mu\nu} = 0$, again owing to the Euler-Lagrange field equation. In field theory equation 3.102 can be taken as a definition of the energymomentum tensor. There are four 'conserved charges' associated with $t^{\mu\nu}$, namely the 4-momentum $P^{\mu} = \int d^3x \, t^{\mu 0}$. These include the Hamiltonian $H = P^0$ and the 3-vector P which is interpreted as the physical 3-momentum carried by the field.

However in general the form of $t^{\mu\nu}$ defined in equation 3.102 is neither symmetric nor gauge invariant. For example, with the Lagrangian for the electromagnetic field $\mathcal{L}_{em} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu}$ of equation 3.93, as a function of $A_{\mu}(x)$, equation 3.102 yields:

$$
t^{\mu\nu} = -F^{\mu\rho}\partial^{\nu}A_{\rho} + \frac{1}{4}\eta^{\mu\nu}F_{\rho\sigma}F^{\rho\sigma}
$$
\n(3.103)

for which the lack of symmetry is clear in the $\mu\nu$ indices in the first term and the lack of gauge invariance is clear from the form of the explicit A_{ρ} in this term. The standard interpretation of this observation is that in Lagrangian field theory the energymomentum $t^{\mu\nu}$ is not a directly measurable quantity and the corresponding ambiguity allows for the addition of a extra terms, leading for example to the quantity ([10] p.101):

$$
T^{\mu\nu} = t^{\mu\nu} + \partial_{\rho} (F^{\mu\rho} A^{\nu}) \tag{3.104}
$$

For the source-free case considered here with $\partial_{\mu}F^{\mu\nu} = 0$ this produces a symmetric gauge invariant form of the Maxwell energy-momentum tensor, in fact in the form

of equation 3.105 below with $\eta^{\mu\nu}$ in place of $g^{\mu\nu}$. However, in addition to the ad hoc nature this procedure is clearly flawed in that it is incompatible with general relativity. That is, for any $T^{\mu\nu} \neq 0$ the spacetime geometry, described by the Einstein equation 3.84, is not flat and hence the assumption of spacetime translation symmetry which led to equation 3.102 itself is invalid.

On the other hand, the electromagnetic energy-momentum tensor $T^{\mu\nu}$ can be derived directly by a different standard procedure, in general relativity, from the stationarity of the matter Lagrangian $\delta \int \mathcal{L} = 0$ with respect to variation in the metric tensor $g_{\mu\nu}(x)$, as described towards the opening of this section. Substituting the Maxwell Lagrangian of equation 3.93 into equation 3.85 gives directly:

$$
T^{\mu\nu} = +F^{\mu\rho}F_{\rho}{}^{\nu} + \frac{1}{4}g^{\mu\nu}F^{\rho\sigma}F_{\rho\sigma}
$$
 (3.105)

This general relativistic method yields an energy-momentum tensor $T^{\mu\nu}$ which is symmetric, gauge invariant and complies necessarily with the Einstein equation 3.84 since it derives from the Einstein-Hilbert action of equation 3.79.

For general relativity the four relations $G^{\mu\nu}_{;\mu} = 0$ of the contracted Bianchi identity of equation 3.71, together with the identity $(\Lambda g^{\mu\nu})_{;\mu} = 0$, places four constraints $T^{\mu\nu}_{\ \ ;\mu} = 0$ on the energy-momentum tensor for the general case via the Einstein equation 3.84, which in turn implies that only six of the ten field equations are independent. Hence the metric $g_{\mu\nu}(x)$ is not determined uniquely by $G^{\mu\nu} + \Lambda g^{\mu\nu} = -\kappa T^{\mu\nu}$, but rather four degrees of freedom remain for arbitrary coordinate transformations. Indeed, the field equation is only required to define $g_{\mu\nu}(x)$ up to an equivalence class (M_4, g) of geometries on the manifold M_4 related by coordinate transformations θ such that (M_4, g) and (M_4, θ^*g) are physically equivalent, as described in the discussion of figure 3.6 in the previous section.

Within the Lagrangian framework it is also possible to derive the contracted Bianchi identity $G^{\mu\nu}_{\mu\nu} = 0$ itself. Taking $\mathcal L$ to be the Ricci scalar R the Einstein-Hilbert action $I = \int R \sqrt{|g|} d^4x$ (equation 3.79 for the vacuum case and setting $\alpha = 1$) is a scalar quantity and hence invariant under coordinate transformations. Indeed while the variational method can be employed, via the Einstein equation, to determine the 4-dimensional spacetime geometry it is unable to deduce a specific choice of metric function and coordinates, by the principle of general covariance. However, the fact that $\delta I = 0$ for coordinate transformations can be shown ([6] p.503) to imply the identity $G^{\mu\nu}_{;\mu} = 0$ of equation 3.71.

The examples of this section have shown some of the great variety of circumstances in which the Lagrangian method may be employed. These include cases in Newtonian mechanics, special relativity and general relativity as well as for electromagnetism and non-Abelian gauge theories. However all of these examples also rest on the assumption of the validity of the Lagrangian approach. One of the aims of the present theory is to derive all equations of motion without employing a Lagrangian

Already it has been described for equation 3.71 how the relation $G^{\mu\nu}_{;\mu} = 0$ is a geometric identity which stands alone as a 'conserved' geometric quantity without the need for a Lagrangian formulation. In the present theory it stands at the head as central to the derivation of physical equations of motion, as we shall investigate in section 5.2. This is universally true both for equations of motion at the effective macroscopic level, relating to classical phenomena such as the Lorentz force law, and also at the microscopic level of the fundamental underlying fields, relating to quantum phenomena, where the constraint of the full form of temporal flow $L(\hat{\mathbf{v}}) = 1$ will also prove central to the physics.

In contrast to the Lagrangian approaches in general relativity via equation 3.85 and in field theory via equation 3.102 in the present theory the Einstein equation will essentially be interpreted as the definition of energy-momentum, that is $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$, where a possible $\frac{\Lambda}{\kappa} g^{\mu\nu}$ term may be implicitly included in the left-hand side. (In subsequent chapters this relation may be written simply as $T^{\mu\nu} := G^{\mu\nu}$ to emphasise the equivalence of the two tensors, with the implied normalisation factor of $-\kappa$ explicitly introduced for practical applications). Since the geometric content of $G^{\mu\nu}$ is measurable in general relativity, in principle at least as the gravitational influence on test bodies, defining the energy-momentum tensor this way does have an unambiguous meaning. In principle the structure of the energy-momentum tensor in such a theory may be uniquely specified, distinguishing between equations 3.103 and 3.105 in the example of the electromagnetic field.

This is the case for Kaluza-Klein theory in which equation 3.105, generalised for non-Abelian internal symmetry, is derived from the structure of a higher-dimensional geometry as will be reviewed in the following chapter leading to equation 4.17. With the Yang-Mills equation 3.95 also being derived in equation 4.18 within this framework the Kaluza-Klein approach achieves a degree of unification with less dependence upon the introduction of Lagrangian terms, such as equation 3.94. In section 5.1 we describe how the techniques of Kaluza-Klein theory might be adopted within the present theory before continuing in section 5.2 to explore some of the consequences of these structures in terms of avoiding the need to postulate Lagrangian functions.

Chapter 4

Kaluza-Klein Theory

4.1 General Relativity with Extra Dimensions

Theories with an extra spatial dimension were initially proposed [11, 12] within a few years of the publication of the general theory of relativity, with the aim of accounting for non-gravitational forces of nature through the higher-dimensional geometry, at a time when only two fundamental forces were known, namely gravitation and electromagnetism. A generalisation of the original Kaluza-Klein theory for the case of a non-Abelian internal symmetry, incorporating further dimensions, was elaborated in detail around half a century later ([13], see also [14], [15] sections I–V and [16]).

This unifying framework for gravitation and gauge theories, reviewed here, is constructed in the mathematical setting of a principle fibre bundle. Keeping within the spirit of Einstein's original 4-dimensional spacetime theory of gravitation and the extension to a 5-dimensional arena by Kaluza and Klein, the geometric unification with non-Abelian gauge theory is founded upon a metric tensor \check{g} , now defined upon the manifold of the principle bundle $P = (M_4, G)$ itself (with the 'check' on \check{g} denoting an object on the bundle space).

We note that conventions vary in the literature – in particular with respect to the assignment of index labels such as $\{a, b, \ldots\}$, $\{\alpha, \beta, \ldots\}$, and $\{i, j, \ldots\}$ which in this paper are associated with objects on the manifolds M_4 , G and P respectively, in the manner described shortly before and in figure 3.2. The conventional order of the indices for the linear connection coefficients Γ^a_{bc} also varies, with the convention of equation 3.45 adopted here, while the sign of the Ricci tensor $R_{\mu\nu} = R^{\rho}{}_{\mu\nu\rho}$ of equation 3.74 also differs in some of the references. Hence in turn a number of derived expressions here will have signs differing to those in the literature.

In addition to the metric g_{ab} on the base manifold M_4 a natural metric for the group manifold G is provided by the Killing form K , which as a matrix of components is invertible provided G is a semi-simple Lie group and negative definite if G is compact. In the latter case a basis for the Lie algebra can be chosen such that the Killing form has components $K_{\alpha\beta} = -\delta_{\alpha\beta}$, is described after equation 3.94. Here we choose metric components $g_{\alpha\beta} = +K_{\alpha\beta}$ in order to match the signature convention of equation 3.72,

with spacelike components having a negative norm.

The $\text{Ad}(G)$ -invariant Killing form defines a bi-invariant metric on the manifold G; that is with both the left L_a and right R_a group actions, for any $a \in G$, being isometries on G, with for example $(R_a^* g)_b(X,Y) = g_b(X,Y)$ for all $X,Y \in T_bG$ for the Killing metric g at any point $b \in G$ (subsequently the Killing metric will often be denoted by $g_{\alpha\beta}$, rather than simply the kernel letter g, as the notation used for the indices helps identify the space to which the object belongs). In particular, in terms of the group structure constants $c^{\alpha}{}_{\beta\gamma}$ in a left-invariant basis $\{X_{\alpha}\}$ on the group manifold, the components of the Killing metric are:

$$
g_{\alpha\beta} = K_{\alpha\beta} = c^{\rho}{}_{\alpha\sigma} c^{\sigma}{}_{\beta\rho} \tag{4.1}
$$

A gauge connection 1-form ω on a principle bundle P specifies a right-invariant horizontal subspace H_pP for all points $p \in P$, as described in section 3.1. A unique metric \check{g} may be defined on such a principle bundle space, aligned with the gauge connection structure with:

$$
\check{g}(X,Y) = g(\pi_* X, \pi_* Y) + K(\omega(X), \omega(Y))
$$
\n(4.2)

where $X, Y \in TP$, while here g and K are the metrics on the base space M_4 and group space G respectively. This construction yields an intuitively natural metric on the bundle space in the sense that the vertical VP and horizontal HP subspaces of the tangent space of P, as depicted in figure 3.2, are then *orthogonal* with respect to \check{g} , with $\check{g}(X, Y) = 0$ if $X \in VP$ and $Y \in HP$ for example.

Alternatively, and perhaps more in the spirit of the original Kaluza-Klein theory, the metric \check{q} rather than the connection ω can be taken as the fundamental entity on P. That is, the bundle is initially endowed with a pseudo-Riemannian metric \check{g} with *certain restrictions* – namely compatibility with a metric g_{ab} on M_4 and metric $g_{\alpha\beta}$ on the fibres G_x and the requirement of invariance under the right action of G on P:

$$
R_a^* \check{g}_{pa}(X, Y) = \check{g}_p(X, Y) = \check{g}_{pa}(R_*X, R_*Y) \tag{4.3}
$$

for any $p \in P$, $a \in G$ and $X, Y \in TP$. This latter property then implies the existence of a subspace HP , orthogonal to VP , which is right-invariant and hence is equivalent to the existence of a connection 1-form ω on the bundle P, which is related to \check{g} as described in equation 4.2.

From either perspective from the relation of \check{g} to ω in equation 4.2 in the horizontal lift basis $\acute{e}_i = {\acute{e}_{\alpha}, \acute{e}_{a}}$, with $\acute{e}_{\alpha} \in VP$ and $\acute{e}_{a} \in HP$, for the tangent space on P the metric \acute{g} , and its inverse, take respectively the simple forms:

$$
\acute{g}_{ij} = \left(\begin{array}{c|c} g_{ab} & \\ \hline & g_{\alpha\beta} \end{array}\right) \qquad \text{and} \qquad \acute{g}^{ij} = \left(\begin{array}{c|c} g^{ab} & \\ \hline & g^{\alpha\beta} \end{array}\right) \tag{4.4}
$$

That is with the components of the metric on the base space M_4 being $g_{ab} = \acute{g}(\acute{e}_a, \acute{e}_b)$ and those of the Killing metric on the group space being $g_{\alpha\beta} = \acute{g}(\acute{e}_{\alpha}, \acute{e}_{\beta})$. The offdiagonal components in equation 4.4 are all zero, with for example $\acute{g}(\acute{e}_a, \acute{e}_\beta) = 0$ describing the orthogonality of any $X \in H_pP$ to any $Y \in V_pP$ with respect to this right-invariant metric \acute{g} .

Under a change of frame to a direct product basis $\{\hat{e}_i\} \rightarrow \{\hat{e}_i\}$, that is the reverse of equation 3.26 with $\ddot{e}_{\alpha} = \acute{e}_{\alpha}$ and $\ddot{e}_{a} = \acute{e}_{a} + \omega^{\alpha}{}_{a}\acute{e}_{\alpha}$ for a choice of trivialisation $\psi: P \to U \times G$, see figure 3.5, we have:

$$
\ddot{g}_{ij} = \left(\frac{g_{ab} + g_{\alpha\beta}\omega^{\alpha}{}_{a}\omega^{\beta}{}_{b}}{g_{\alpha\beta}\omega^{\beta}{}_{b}}\right) \quad \text{and} \quad \ddot{g}^{ij} = \left(\frac{g^{ab}}{-\omega^{\alpha}{}_{a}g^{ab}}\left|\frac{-g^{ab}\omega^{\beta}{}_{b}}{g^{\alpha\beta} + g^{ab}\omega^{\alpha}{}_{a}\omega^{\beta}{}_{b}}\right|\right)
$$
\n(4.5)

In this latter basis the non-Abelian gauge fields $\omega^{\alpha}_{a}(p)$ on P for the internal symmetry are found alongside the external spacetime metric elements g_{ab} framed within the components of the full metric \ddot{g}_{ij} on the bundle space. This is a generalisation of the original 5-dimensional Kaluza-Klein theory in which the electromagnetic 4-vector potential A_a appears alongside the components of the spacetime metric g_{ab} within the extended 5×5 metric tensor.

Any differentiable manifold M_n is canonically associated with a principle bundle of linear frames FM_n with structure group $GL(n, \mathbb{R})$, where n is the dimension of the base manifold M_n , as described in the opening of section 3.3. This includes the case in which the base manifold is actually the space of a given principle fibre bundle P itself.

While the metrics g and K on the manifolds M_4 and G can be naturally extended to the metric \check{g} of equation 4.2 on the principle bundle $P = (M_4, G)$ with a connection ω , linear connections on the manifolds M_4 and G may also be generalised to the domain of the larger manifold P. As described for equation 3.45 such a linear connection $\check{\Gamma}$ will define covariant differentiation with $\check{\nabla} \check{e}_i = \check{\Gamma}^j_{\;i} \check{e}_j \equiv \check{\Gamma}^j_{\;ik} \check{e}^k \otimes \check{e}_j$ in a general tangent space basis $\{\check{e}_i\}$ for TP with dual basis $\{\check{e}^i\}$ for T^*P . The identification of the smooth symmetric gauge covariant rank-2 tensor \check{q} everywhere on P endows the principle bundle itself with the structure of a pseudo-Riemannian manifold. In turn a connection Γ compatible with the metric \check{g} , and hence with the geometric structure of the underlying manifold P, may be extended from the notion of a metric connection on M_4 .

Indeed, and further guided by Einstein's general theory of relativity in 4 dimensional spacetime, the unique linear connection which is torsion-free, $\check{T}^i_{jk} = 0$, and compatible with the metric, $\tilde{\nabla}_k \tilde{g}_{ij} = 0$, that is the Levi-Civita connection, may be defined on the bundle space P . The corresponding connection coefficients can be expressed, with $\Gamma_{ijk} = g_{il} \Gamma^{l}_{jk}$ and $c_{ijk} = g_{il} c^{l}_{jk}$, as:

$$
\check{\Gamma}_{ijk} = \frac{1}{2} (\check{e}_j(\check{g}_{ik}) + \check{e}_k(\check{g}_{ij}) - \check{e}_i(\check{g}_{jk})) - \frac{1}{2} (\check{c}_{ijk} + \check{c}_{kji} + \check{c}_{jki})
$$
(4.6)

which expresses equation 3.53 in a general frame. These coefficients take a relatively simple form in the horizontal lift basis, as employed for the metric in equation 4.4, while a coordinate basis will also be adopted on the base space M_4 . In this basis the connection coefficients Γ^a_{bc} on the base space M_4 contribute to the set in equation 4.6 with:

$$
\acute{\Gamma}^{a}_{bc} = \Gamma^{a}_{bc} = \frac{1}{2}g^{ad}(e_b(g_{cd}) + e_c(g_{bd}) - e_d(g_{bc}))
$$
\n(4.7)

which is simply equation 3.53, since the structure coefficients on P are related to the structure coefficients on the base manifold with $\acute{c}_{abc} = c_{abc} = 0$ in this basis (and with

the corresponding term hence absent in equation 3.34). The connection coefficients $\check{\Gamma}_{ijk}$ are also related to the internal curvature through equation 4.6 since in the horizontal lift basis, by equation 3.36, we have $\hat{c}^{\alpha}_{ab} = -\Omega^{\alpha}_{ab}$. Here we adopt the convention of denoting the components of curvature Ω^{α}_{ab} on the principle bundle by F^{α}_{ab} , which may then represent the curvature components on P or M_4 depending on the context, in order to match the notation in many of the references. Ultimately the curvature $F^{\alpha}_{\;\;ab}$ will feature in gauge invariant expressions on the base manifold. From equation 4.6 we find in the horizontal lift basis on the bundle P terms such as (see [13] equation (22)):

$$
\acute{\Gamma}^{\alpha}{}_{ab} = +\frac{1}{2} F^{\alpha}{}_{ab} \qquad \text{and} \qquad \acute{\Gamma}^{a}{}_{b\alpha} = \acute{\Gamma}^{a}{}_{\alpha b} = +\frac{1}{2} g^{ac} g_{\alpha\beta} F^{\beta}{}_{bc} \tag{4.8}
$$

The complete set of coefficients for the Levi-Civita connection on P are listed under 'Cho [13]' as the first case in table 4.1 in the following section.

Hence the Levi-Civita connection of equation 4.6 on the total bundle space P is intimately related to the external curvature on the base space as well as the internal curvature of the gauge group. In turn the components of the Riemann curvature tensor \check{R}^i_{jkl} calculated for this Levi-Civita connection on P according to equation 3.62 is intimately related to *both* the *external* curvature on M_4 via equation 4.7 and the *internal* curvature, associated with gauge group G , which is drawn into the Riemannian geometry through equation 4.8.

It is important to clarify the relation between the linear connection $\Gamma(p)$ and gauge connection $\omega(p)$ on the manifold P. In fact from the point of view of frame bundles and principle fibre bundles in general a *linear* connection $\tilde{\omega}$ (see the opening of section 3.3) on FP would be the same kind of object as the *gauge* connection ω on P. Here we are dealing with Riemannian geometry of the manifold P itself, which is hence the base space upon which the gl (m, \mathbb{R}) -valued 1-form $\Gamma(p) = \Sigma^* \tilde{\omega}$ is defined, where $m = \dim(P) = \dim(M_4) + \dim(G)$ and $\Sigma(p)$ is a section map $P \to FP$ over P. The same manifold P is also the *principle bundle* upon which the gauge connection $ω$ is defined, with $A(x) = σ^*ω(p)$ being the gauge field on M_4 , for a section map $\sigma(x): M_4 \to P$ over the space M_4 .

Having the metric \check{g}_{ij} on P the Ricci tensor $\check{R}_{jk} = \check{g}^{il} \check{R}_{ijkl}$ (equation 3.74) and scalar curvature $\check{R} = \check{g}^{ij} \check{R}_{ij}$ may also be computed, where the latter is found to be (with differing sign convention to [13]):

$$
\check{R} = R_M + R_G + \frac{1}{4}F^2
$$
\n(4.9)

Here R_M is the usual scalar curvature on the base manifold (which varies with the point $x = \pi(p) \in M_4$ under $p \in P$) and R_G is the scalar curvature on the group manifold G (which can be interpreted as a, problematically very large, cosmological constant in this version of Kaluza-Klein theory). The term $F^2 = F^{\alpha}_{ab} F_{\alpha}{}^{ab}$, constructed from the non-Abelian gauge fields, is also gauge invariant and the curvature components $F^{\alpha}_{ab}(p)$ on P can be interpreted as the corresponding gauge covariant curvature components $F^{\alpha}_{ab}(x)$ on the base space M_4 , for example in table 4.1.

As a scalar \hat{R} in equation 4.9 is a quantity which is independent of the basis $\{\check{e}_i\}$ in which it is determined (for example in the direct product or horizontal lift basis respectively for equations (17) and (24) of reference $[13]$). The equations of motion for the theory are then derived by adopting the Lagrangian function $\check{R}\sqrt{|\check{g}|}$, where $|\check{g}|$ is the magnitude of the determinant of the metric \check{g}_{ij} on P, in the Einstein-Hilbert action integral:

$$
A_m = \int \check{R}\sqrt{|\check{g}|} \, d^4x \, d^{n_G}G \tag{4.10}
$$

with $m = 4 + n_G$. The integration over the group manifold G, with volume V_G , is trivial and the above expression reduces to the 4-dimensional action integral:

$$
A_4 = V_G \int \check{R}\sqrt{|g|} \; d^4x \tag{4.11}
$$

where |g| is here the determinant of the metric g_{ac} on M_4 . The variational principle is then applied under the constraint $\delta A_m = 0$, and hence $\delta A_4 = 0$, with respect to restricted variations of the metric $\delta \check{g}$ on the bundle space, consistent with equation 4.3, as explained before equation 4.16 in the following section. Within this restriction this again follows the prescription for the original theory of general relativity on a 4-dimensional spacetime manifold M_4 with scalar curvature R for which the field equations can be determined from the Einstein-Hilbert action integral of equation 3.79.

By comparison of equations 4.9 and 4.11 with 3.79 the constant R_G of the Kaluza-Klein theory indeed appears as a cosmological constant term (albeit too large by a factor of $\sim 10^{120}$ if a natural normalisation is used with the length scale of the group space G taken to be of order the Planck length [13]), while the term F^2 effectively contributes the content for the matter Lagrangian \mathcal{L} . Hence, as a particularly elegant feature of Kaluza-Klein theory, the geometry of the 4-dimensional spacetime manifold along with a *matter* contribution is identified within a single *geometrical* object in the form of R on the principle bundle space.

4.2 Theories with Torsion on the Bundle

One way to remove the problematic cosmological term R_G in equation 4.9 would be to redefine the Lagrangian for the Kaluza-Klein theory by simply adding by hand a counter-cosmological constant term Λ_c to R^{\dot{R}} in equation 4.10 to cancel R_G . However this would be an ad hoc measure, similar in spirit to the inclusion of the original cosmological constant term Λ in equation 3.79, contrived largely to match empirical observation.

However there is flexibility within the Kaluza-Klein approach on a principle fibre bundle if the metric \check{g}_{ij} is not treated as the fundamental object of the theory (see for example [17, 18, 19, 20]). While the same natural metric \check{g}_{ij} of equation 4.4 is employed the linear connection $\check{\Gamma}^i_{jk}$ on P may be defined with some independence from \check{g}_{ij} , unlike for the Levi-Civita connection of equation 4.6. In this case it is possible to derive a curvature scalar R on P such that the cosmological term R_G vanishes and equation 4.9 reduces to simply:

$$
\check{R} = R_M + \frac{1}{4}F^2
$$
\n(4.12)

One way to achieve this is to require the linear connection $\check{\Gamma}^i_{jk}$ to incorporate a description of absolute parallelism on the bundle fibres G_x . As reviewed in subsection 2.2.2 on the manifold G itself the list of canonical geometric objects include a

basis of left-invariant vector fields $\{X_{\alpha}\}\$ and the Maurer-Cartan 1-form $\theta = \theta^{\alpha}X_{\alpha}$ as well as the structure constants $c^{\alpha}{}_{\beta\gamma}$ and the Killing form metric $g_{\alpha\beta}$ of equation 4.1. As described below equation 3.67 in the basis $\{X_{\alpha}\}\$ the choice of linear connection coefficients $\Gamma^{\alpha}{}_{\beta\gamma} = 0$ is equivalent to inducing parallel transport on the group manifold via the left action L_a of G on itself, for any $a \in G$, that is with parallelism defined by the left-invariant vector fields $\{X_\alpha\}$ on G, while $\Gamma^\alpha_{\ \beta\gamma} = -c^\alpha_{\ \beta\gamma}$ corresponds to the parallelism described by a right-invariant frame field under R_a . In either case the resulting Riemann curvature is zero with $R^{\alpha}_{\ \beta\gamma\delta} = 0$, as can be shown using equation 3.62 together with the Jacobi identity expressed in terms of the structure constants.

More generally, employing the derivative action of the left-invariant basis vectors $\{X_{\alpha}\}\$, the right-invariance of the Killing metric implies that the covariant derivative of the metric on G vanishes:

$$
\nabla_{\alpha} g_{\beta \gamma} = X_{\alpha} g_{\beta \gamma} - \Gamma^{\delta}{}_{\beta \alpha} g_{\gamma \delta} - \Gamma^{\delta}{}_{\gamma \alpha} g_{\beta \delta} = 0 \tag{4.13}
$$

provided
$$
\Gamma^{\alpha}{}_{\beta\gamma} = -\rho c^{\alpha}{}_{\beta\gamma}
$$
 for any $\rho \in \mathbb{R}$ (4.14)

since $X_{\alpha}g_{\beta\gamma}=0$ and by the antisymmetry of the $c_{\alpha\beta\gamma}$ indices. Hence for any value of $ρ$ this linear connection is metric compatible, with $\nabla g = 0$ on G. The torsion is zero only for $\rho = \frac{1}{2}$ which hence represents the unique Levi-Civita connection on G. On the other hand the Riemannian curvature is zero on G only for the cases of $\rho = 0$ and $\rho = 1$, which with finite torsion are not Levi-Civita connections. However these latter two cases in describing an absolute parallelism on G can be considered as geometrically natural metric connections on G.

For the linear connection $\Gamma^{\alpha}{}_{\beta\gamma} = 0$ or $\Gamma^{\alpha}{}_{\beta\gamma} = -c^{\alpha}{}_{\beta\gamma}$ employed on the bundle fibres G_x a subset of the torsion components on P are also necessarily non-zero, with $\check{T}^{\alpha}_{\ \beta\gamma}\neq 0$. Hence with the torsion allowed to be non-zero on the bundle space P this version of Kaluza-Klein theory resembles the Einstein-Cartan theory on 4-dimensional spacetime for which Γ and q are treated as independent geometric objects. Here we review four such approaches in the literature.

In Kopczyński [17] a G-invariant linear connection Γ is constructed in terms of the structure on the principle bundle with a gauge connection ω without reference to a metric and with non-zero torsion. This generalises from the Levi-Civita connection described in the previous section (as employed by [13] and others) with the 'gravitational field' on P now being described by the combination of both \check{q} of equation 4.2 and the components of Γ as listed in the corresponding column under 'Kop [17]' in table 4.1. With these components the scalar curvature on P is found to be $\check{R} = R_M + \alpha(\alpha - 1)K^2$, with $K^2 = K^{\alpha\beta}K_{\alpha\beta}$. For the case $\beta = 0$ the connection is metric compatible, resembling Einstein-Cartan general relativity in 4-dimensional spacetime. While this reference shows that the connection coefficients can be greatly simplified compared with the Levi-Civita case, listed under 'Cho [13]' in the first column of table 4.1, in order to achieve the correct dynamics a more complicated Lagrangian function is postulated with $\mathcal{L} = \check{R} + \frac{\mu}{2}$ $\frac{\mu}{2} \check{T}^i_{jk} \check{T}^{\;jk}_i$ including a quadratic torsion term. The cosmological constant Λ obtained in this approach is arbitrary, and may be set to be zero or very small by a suitable choice of the parameters α and μ .

In Orzalesi and Pauri [18] the main motivation is to describe a linear connection Γ on the principle bundle which is gauge covariant. In particular requiring the Ricci curvature on the fibre space to be gauge invariant implies the adoption of zero curvature on the group manifold, that is the case $\rho = 0$ or $\rho = 1$ as described above for equation 4.14. This form differs in a relatively minimal way from the Levi-Civita connection, as can be seen by comparing the entries of column [18] with column [13] in table 4.1. Here the simple scalar Lagrangian $\mathcal{L} = \mathcal{R}$ on the bundle space is again adopted, with the resulting vanishing of $\Lambda \equiv R_G$ on the base space M_4 interpreted as a consequence of the underlying gauge G-symmetry of the Riemannian geometry on P. Without an R_G term the vacuum solution corresponds to zero external curvature $R_M = 0$ together with zero internal curvature $F = 0$.

In Kalinowski [19] the linear connection 1-forms $\check{\Gamma}^i_{\;j} = \check{\Gamma}^i_{\;jk} \check{e}^k$ on P are defined as the horizontal part of the Levi-Civita connection 1-forms Γ^i_j of equation 4.6, that is $\check{\Gamma}^i_{\;j} = \text{hor}(\Gamma^i_{\;j})$ (with 'hor' introduced in equation 3.13) which maps the vertical component of tangent vectors on TP to zero. The components of this linear connection $\check{\Gamma}^i_{jk}$ in the horizontal lift basis are listed in column [19] of table 4.1. The factors of λ arise as here the metric on G is taken to be $g_{\alpha\beta} = \lambda^2 K_{\alpha\beta}$. This linear connection $\check{\Gamma}^i_{jk}$ is metrical, invariant under the G-action, again with non-zero torsion and, while motivated in the context of gauge derivatives of spinor fields, again leads to a vanishing cosmological constant.

In Katanaev [20] an initially completely general $\check{\Gamma}^i_{jk}$ on the principle bundle is considered. Four conditions are postulated for $\check{\Gamma}$ in a geometrically meaningful way related to the structure group G over P and, as for the previous reference, with emphasis on horizontal propagation. In particular for column [20] of table 4.1 on taking $c = 1$ for entry '5)' $\hat{\Gamma}^{\alpha}{}_{ab} = cF^{\alpha}{}_{ab}$ the change in a tangent vector to P under parallel transport using these linear connection coefficients equals the change in the vector due to the basis transformation under parallel transport of the fibres using the gauge connection, with the latter depicted in figure 3.3. The entry '4)' in this column is included for metric compatibility. The coefficients listed represent the case presented in [20] with finite torsion and the absence of a cosmological constant term, although a different choice of Γ consistent with the postulates is possible. A further possibility within this framework would be to set the first two entries, '1)' and '2)', equal to zero in column [20]. This reference is of significance for the present paper in that it highlights the possibility of a geometric origin of Γ on P without any appeal to the Levi-Civita connection.

The complete set of linear connection coefficients for reference [13], augmenting equation 4.8, are collected in the first column of table 4.1. These are listed alongside the linear connection coefficients $\check{\Gamma}^i_{jk}$ on the bundle space P for the above four cases with non-zero torsion. Where necessary signs have been aligned to the conventions used here, with for example linear connection 1-forms $\check{\Gamma}^i_{\;j} = \check{\Gamma}^i_{\;jk}\check{e}^k$. The motivation for the final column headed 'minimal' will be explained in section 5.1.

Only the first case in table 4.1 describes a torsion-free linear connection, yet each of the six cases is a Kaluza-Klein theory providing a unifying framework for general relativity together with gauge field theory. The purpose of collecting together the range of linear connection coefficients is to demonstrate that a significant degree of flexibility is possible within Kaluza-Klein theory while still maintaining this unified framework.

The derivation of Einstein's equations on 4-dimensional spacetime from the Einstein-Hilbert action of equation 3.79 was described in the opening of section 3.5.
$\acute{\Gamma}^i_{jk}$	Cho $[13]$	Kop [17]	$O+P$ [18]	Kal $[19]$	Kat $[20]$	minimal
1) $\hat{\Gamma}^{\alpha}{}_{\beta\gamma}$	$-\frac{1}{2}c^{\alpha}{}_{\beta\gamma}$	$-csc^{\alpha}{}_{\beta\gamma}$	$-c^{\alpha}{}_{\beta\gamma}$ or 0	θ	$-c^{\alpha}{}_{\beta\gamma}$	Ω
2) $\hat{\Gamma}^{\alpha}_{\gamma a}$	0	Ω	θ	$\overline{0}$	$-\omega^{\beta}_{\ \ a} \acute{\Gamma}^{\alpha}_{\ \ \beta\gamma}$	$\overline{0}$
3) $\hat{\Gamma}^a_{\ b\gamma}$	$\frac{1}{2}g^{ac}g_{\gamma\beta}F^{\beta}_{bc}$	$\overline{0}$	$\frac{1}{2}g^{ac}g_{\gamma\beta}F^{\beta}_{\ \ bc}$	$\overline{0}$	$\overline{0}$	$\eta g^{ac}g_{\gamma\beta}F^{\beta}_{bc}$
4) $\dot{\Gamma}^a_{\gamma b}$	$\frac{1}{2}g^{ac}g_{\gamma\beta}F^{\beta}_{bc}$	$\overline{0}$		$\frac{1}{2}g^{ac}g_{\gamma\beta}F^{\beta}_{bc}\left \right.\frac{\lambda}{2}g^{ac}g_{\gamma\beta}F^{\beta}_{bc}\left \right.cg^{ac}g_{\gamma\beta}F^{\beta}_{bc}\right $		θ
5) $\acute{\Gamma}^{\alpha}{}_{ab}$	$\frac{1}{2}F^{\alpha}_{ab}$	βF^{α}_{ab}	$rac{1}{2}F^{\alpha}_{ab}$	$rac{\lambda}{2}F^{\alpha}_{ab}$	cF^{α}_{ab}	θ
6) $\hat{\Gamma}^a_{bc}$	Γ^a_{bc}	Γ^a_{bc}	Γ^a_{bc}	Γ^a_{bc}	Γ^a_{bc}	$\Gamma^a_{\;\;bc}$

Table 4.1: Linear connection components $\tilde{\Gamma}^i_{jk}$ on a principle bundle extracted from [13] equation (22), [17] p.367, [18] equation (19), [19] equation (29), the case in [20] with non-zero torsion on G and for a 'minimal' model. All components are expressed in the horizontal lift basis and $\dot{\Gamma}^a_{\beta\gamma} = \dot{\Gamma}^{\alpha}_{a\gamma} = 0$ in all six cases. Each of $\lambda > 0$, α , β , c and γ , where used as coefficients, are real constant parameters.

In the vacuum case with $\mathcal{L} = 0$ and $\Lambda = 0$ variation of the metric $\delta g_{\mu\nu}$ on M_4 leads to the equation of motion $G^{\mu\nu} = 0$ of equation 3.82. For the Kaluza-Klein extension to the scalar curvature \tilde{R} for the Levi-Civita connection on a principle bundle space the same steps lead to the requirement of the stationarity of the action integral over the full bundle space in equation 4.10, that is $\delta A_m = 0$, under variation of the extended metric \check{g}_{ij} on P, which results in the expression:

$$
\check{G}^{ij} = \check{R}^{ij} - \frac{1}{2}\check{R}\,\check{g}^{ij} = 0\tag{4.15}
$$

In some versions of Kaluza-Klein theory, in particular for the 5-dimensional case, equation 4.15, which implies $\tilde{R}^{ij} = 0$, is quoted as an ansatz at the outset in order to derive equations of motion for the 4-dimensional world by imposing this higher-dimensional 'vacuum equation' (see for example [21], in which the 5-dimensional metric $\check{g}_{ij}(x)$ may depend on the $5th$ coordinate).

However for the extended Kaluza-Klein theories described here the variations in the metric \check{g}_{ij} are not arbitrary since the structure of the symmetries of \check{g}_{ij} on the bundle space P need to be preserved under the variations $\delta \check{g}_{ij}$. That is, the rightinvariance of \check{g}_{ij} of equation 4.3 and the general form of the metric in equation 4.5 should be preserved. This limits the metric variations to the components δg_{ac} and $\delta\omega^{\alpha}_{a}$ on P and leads to two equations of motion on the base manifold M_4 . Applying the variation δg_{ac} under $\delta A_m = 0$ for the action in equation 4.10, with the curvature R of equation 4.12, in a general coordinate basis on M_4 leads to ([19] equation 38):

$$
G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu} = -\kappa T^{\mu\nu}
$$
 (4.16)

with
$$
-2\kappa T^{\mu\nu} = -F^{\alpha\mu}{}_{\rho}F_{\alpha}{}^{\rho\nu} - \frac{1}{4}g^{\mu\nu}F^{\alpha}{}_{\rho\sigma}F_{\alpha}{}^{\rho\sigma} \qquad (4.17)
$$

The left-hand side of the top line would read $G^{\mu\nu} + R_G g^{\mu\nu}$ if the scalar curvature of equation 4.9 based on a Levi-Civita connection is used instead. On the other hand the variation $\delta \omega_{a}^{\alpha}$ leads to

$$
D_{\mu}F^{\alpha\,\mu\nu} = 0 \tag{4.18}
$$

Equation 4.16 is the Einstein field equation with the energy-momentum tensor $T^{\mu\nu}$ composed as equation 4.17 purely from the gauge fields, with the latter being subject to equation 4.18 which is the Yang-Mills field equation (or Maxwell's equation $F^{\mu\nu}_{\ \ ;\mu} = 0$ in the case of the Abelian internal symmetry group $G = U(1)$, see also the discussion after equation 3.91). Hence the source-free Yang-Mills field equation 3.95 has been derived without the explicit introduction of the Yang-Mills Lagrangian of equation 3.94. Rather such a 'Lagrangian term' $F^2 = F_{\alpha\mu\nu} F^{\alpha\mu\nu}$ has been incorporated within the Einstein-Hilbert action based purely on the geometry of the bundle space. In this way the non-Abelian Kaluza-Klein theory provides a unified framework for the combined Einstein-Yang-Mills field equations.

4.3 Theories with Homogeneous Fibres

A further generalisation of Kaluza-Klein theory is also of relevance for the framework presented in this paper. In the present theory the symmetry group G rather than being motivated independently is introduced in terms of the set of symmetry actions on a form $L(v) = 1$ of multi-dimensional temporal flow. This structure is reminiscent of Kaluza-Klein theories with homogeneous fibres in which G acts on a k-dimensional manifold S_k . A bundle space E is constructed with each fibre being a copy of S_k over the base space M_4 . Based on the references [22, 23, 24] this approach will be collectively summarised in this section.

In these models the bundle $E = (M_4, S_k)$ is constructed over the base space M_4 while the fibres S_k may be considered to represent k 'extra dimensions'. For our purposes it is sufficient to consider the trivial bundle with $E \equiv M_4 \times S_k$. While either a left or right action may be considered here we take the gauge group G to act on the space S_k on the left (as for reference [22] for example, and as will be the case for the E_6 action on the space $h_3\mathbb{O}$ as discussed alongside equation 6.55) such that each Lie algebra element $X_{\alpha} \in L(G)$ generates a vector field K_{α} on S_k with the bracket composition exhibiting the negative of the structure constants $c^{\alpha}{}_{\beta\gamma}$ of $L(G)$, that is:

$$
[K_{\beta}, K_{\gamma}] = -c^{\alpha}{}_{\beta\gamma} K_{\alpha} \tag{4.19}
$$

The group actions may also be considered to be one-to-one with the isometry transformations for an inner product defined on the tangent space TS_k . That is, a G-invariant metric may be defined on S_k with Killing vector fields:

$$
K_{\alpha} = K^{\dot{\alpha}}_{\ \alpha} e_{\dot{\alpha}} \tag{4.20}
$$

where $K^{\dot{\alpha}}_{\alpha}$ are the components of K_{α} in a linearly independent tangent space basis ${e_{\alpha}}$ on S_k , with indices $\alpha = 1...k$ and $\alpha = 1...$ dim(G). Such a G-invariant metric $g_{\alpha\beta}$ on S_k may be induced from the Killing metric $K_{\alpha\beta}$ on G itself.

If G acts upon S_k transitively then S_k is a homogeneous space. Given any point $y_0 \in S_k$ the elements $h \in G$ for which $h \cdot y_0 = y_0$ under the left action of the group form the isotropy subgroup H, with $h \in H \subset G$. The homogeneous space S_k is then diffeomorphic to the space of left cosets gH as identified for varying $g \in G$, that is $S_k \equiv G/H$ where H is the isotropy subgroup of the isometry group G. As a vector space the Lie algebra of G may be decomposed as:

$$
L(G) = L(H) + B \tag{4.21}
$$

with $[L(H), L(H)] \subset L(H)$ and $[L(H), B] \subset B$, where $B \equiv T(G/H)$ forms a basis for the tangent space at $y_0 \in S_k$.

Such a linearly independent basis ${e_{\alpha}}$ for TS_k forms a basis for the vertical subspace of the tangent space on the fibre bundle E . A complete 'horizontal basis' on E, written $\acute{e}_i(x, y) = {\acute{e}_\alpha, \acute{e}_a}$, in place of the horizontal lift basis for the principle bundle P of figure 3.2, can be expressed as:

$$
\acute{e}_{\dot{\alpha}} = \ddot{e}_{\dot{\alpha}}, \qquad \acute{e}_{a} = \ddot{e}_{a} - A^{\alpha}_{a}(x) K^{\dot{\alpha}}_{\alpha}(y) \ddot{e}_{\dot{\alpha}}(x, y) \equiv \ddot{e}_{a} - A^{\alpha}_{a}(x) \ddot{e}_{\alpha} \tag{4.22}
$$

in terms of a direct product basis $\ddot{e}_i(x, y) = \{\ddot{e}_{\dot{\alpha}}, \ddot{e}_a\}$ on E, by comparison with equation 3.26 and figure 3.5, using the Killing vector components $K^{\dot{\alpha}}_{\alpha}$ defined in equation 4.20. As implied in equation 4.22 the construction of such a horizontal basis on E corresponds to the introduction of a connection form ω on the associated principle bundle $P \equiv M_4 \times G$. This connection form is written in terms of the coefficients $A^{\alpha}_{a}(x)$ rather than $\omega_{a}^{\alpha}(x, g)$ since the vertical basis is here defined through the *left* action of G (see the discussion in [22] after equation (7.2) for example).

Consistent with the horizontal basis of equation 4.22 a natural metric on the bundle space E may be defined, for which horizontal and vertical vectors are mutually orthogonal, and expressed in a direct product basis as:

$$
\ddot{g}_{ij} = \left(\frac{g_{ab}(x) - g_{\hat{\alpha}\hat{\beta}}(y)K^{\hat{\alpha}}_{\alpha}(y)A^{\alpha}_{\alpha}(x)K^{\hat{\beta}}_{\beta}(y)A^{\beta}_{\ b}(x)}{K^{\hat{\beta}}_{\alpha}(y)A^{\alpha}_{\ b}(x)g_{\hat{\alpha}\hat{\beta}}}\right) (4.23)
$$

which may be compared with equation 4.5 for the case of a principle fibre bundle. Changes in the vertical coordinates $\{y\}$ on E described by the infinitesimal isometries $\varepsilon^{\alpha}(x)$ induce changes in the metric components with respectively:

$$
y^{\hat{\alpha}} \rightarrow y^{\hat{\alpha}} + \varepsilon^{\alpha}(x) K^{\hat{\alpha}}_{\alpha}(y)
$$

\n
$$
A^{\alpha}_{a} \rightarrow A^{\alpha}_{a} + \partial_{a} \varepsilon^{\alpha}(x) + c^{\alpha}_{\beta \gamma} \varepsilon^{\beta}(x) A^{\gamma}_{a}
$$
\n(4.24)

Hence such isometries effectively simulate non-Abelian gauge transformations with $A^\alpha{}_a(x)$ identified as the Yang-Mills gauge field on the base space.

Following the Kaluza-Klein prescription described in section 4.1 the Levi-Civita connection, that is the unique torsion-free linear connection compatible with the metric, and curvature can be constructed on the manifold E based on the metric $\ddot{g}_{ij}(x, y)$ of equation 4.23. In turn an action principle may be employed on this $(4+k)$ -dimensional space with action $A_{4+k} = \int \tilde{R}\sqrt{|\tilde{g}|} d^4x d^ky$ in comparison to equation 4.10 where now the curvature scalar \hat{R} on the bundle $E \equiv M_4 \times S_k$ is found to take the form:

$$
\check{R} = R_M + R_{S_k} + \frac{1}{4} g_{\mathring{\alpha}\mathring{\beta}} K^{\mathring{\alpha}}{}_{\alpha} K^{\mathring{\beta}}{}_{\beta} F^{\alpha}{}_{ab} F^{\beta ab} \tag{4.25}
$$

where R_{S_k} is the scalar curvature of the homogeneous space S_k . With F^{α}_{ab} being the gauge curvature components for the internal symmetry group G the above equation again demonstrates a relation between the external Riemann curvature with scalar R_M and a quadratic term in the internal curvature. This relationship derived for G acting on homogeneous fibres is hence in turn similar to that obtained in equations 4.9 and 4.12 with G itself composing the fibres of a principle bundle. A linear connection on E differing from the Levi-Civita connection may be employed to remove the cosmological term S_k by analogy with the examples cited in the previous section. The Einstein-Yang-Mills equations also follow from a prescription analogous to that described for equations 4.16–4.18.

For models with homogeneous fibres in which the metric $g_{\hat{\alpha}\hat{\beta}}(y)$ is replaced by the more general field components $g^{\hat{\alpha}\hat{\beta}}(x,y) = K^{\hat{\alpha}}_{\ \ \alpha}(y)K^{\hat{\beta}}_{\ \ \beta}$ $\mathcal{L}_{\beta}(y) \Phi^{\alpha\beta}(x)$, which depend on $x \in M_4$ and where Φ is a non-Killing metric on G, a set of scalar fields are introduced into the theory with a number of further terms featuring the derivatives $\partial_a g^{\dot{\alpha}\dot{\beta}}(x,y)$ appearing in the corresponding generalisation of equation 4.25 (see for example [22] equation (8.6)).

On the other hand on constructing \ddot{g}_{ij} in equation 4.23 for the case of trivial isotropy group $H = \{e\}$, where $e \in G$ is the identity element, then by equation 4.21 we have $B = L(G)$ and the α, \ldots indices can be replaced by α, \ldots indices, with $K^{\dot{\alpha}}{}_{\alpha} = \delta^{\dot{\alpha}}{}_{\alpha}$. In this case the theory simplifies to that described in section 4.1 based on the metric of equation 4.5 with the set of vector fields $\{\hat{e}_{\alpha}\}\$ spanning the vertical subspace of the tangent space on $P \equiv M_4 \times G$ (with care for the convention choice of a left or right group action, see for example [13] equations (8′) and (12)).

Even for the case with $H \neq \{e\}$ the full G-symmetry Yang-Mills dynamics is obtained so long as G acts *effectively* on the fibres S_k . This is also the case for G acting on $L(v) = 1$ for the present theory in this paper, and in particular for the E_6 action to be described in chapter 6, and for the broken internal subgroups of G.

The action of G on the set of elements v underlying the form $L(v) = 1$ is also transitive, and hence this set forms a homogeneous space, motivating the review of this section. However with the observation that the Kaluza-Klein unification achieved with homogeneous fibres, given an effective group action, is closely related to that achieved on the associated principle bundle, in the following section we apply some of the observations of the previous section to the present theory. This in particular picks up from the development of section 2.3 with the goal of relating the external curvature to the internal curvature in the context of the new theory.

Chapter 5

Geometry Unified through Temporal Flow

5.1 Relating External and Internal Curvature

In this section, ultimately guided by the framework of Kaluza-Klein theories described in the previous chapter, the aim is to determine a relation between the external and internal geometry over the base manifold arising out of the symmetries of a form of temporal flow $L(v) = 1$, building upon the structures described in chapter 2. In place of the base space M_3 with local symmetry $SO(3)$, introduced for the model universe in section 2.2 with the 3-dimensional form $L(v_3) = 1$ of equation 2.14, here we consider the form:

$$
L(\mathbf{v}_4) = (v^0)^2 - (v^1)^2 - (v^2)^2 - (v^3)^2 = 1
$$
\n(5.1)

that is $L(\mathbf{v}_4) = \eta_{ab}v^a v^b = 1$ with Minkowski metric $\eta = \text{diag}(+1, -1, -1, -1)$, with Lorentz $SO^+(1,3)$ symmetry, projected into the 4-dimensional base space M_4 . Over the spacetime manifold M_4 a globally defined orthonormal basis arises in the manner of equation 2.15 with the natural parallelism on M_4 described by a linear connection with components $\Gamma^a{}_{bc}(x) = 0$ in this basis. With the local symmetry group $SO^+(1,3)$ over the base manifold M_4 the principle bundle $P = (M_4, SO^+(1, 3))$ is implicitly identified in this structure, which in fact can be expressed as the manifold product $P \equiv M_4 \times SO^+(1,3)$ owing to the triviality of the bundle as described towards the end of section 3.1.

However, following section 2.3, here we study initially the geometry of the principle fibre bundle $P \equiv M_4 \times G$, where $G = SO^+(1, 9)$ is provisionally taken as the full symmetry group for the form $L(\hat{v}) = L(v_{10}) = 1$, which in turn is the 10dimensional extension of equation 5.1. The extended base manifold M_4 now arises out of four of the ten translational degrees of freedom of $L(\mathbf{v}_{10}) = 1$, in the manner described in equation 2.13. In place of figure 2.7 for the $SO(5)$ model over M_3 described earlier, for this more realistic model we now have the structures described in figure 5.1.

Figure 5.1: (a) The full symmetry group $\hat{G} = SO^+(1, 9)$ over the base space M_4 (b) broken to the internal symmetry SO(6) with external subgroup $SO^+(1,3) \subset SO^+(1,9)$ acting on the tangent space TM_4 .

The structure of figure 5.1 is associated with a canonical flat connection on M_4 , as described by $A(x) = g^*\theta$ of equation 2.30 where here θ is the Maurer-Cartan 1-form on the group manifold $\hat{G} = SO^+(1, 9)$. This canonical flat connection defines a horizontal lift basis { ϵ_i } on the corresponding principle bundle structure $P \equiv M_4 \times$ $SO⁺(1, 9)$, as a particular case of figure 3.2. In turn a collection of affine connection coefficients $\hat{\Gamma}^i_{jk}$ can be defined in this basis on P.

While $\Gamma^a{}_{bc} = 0$ represents the initial parallelism on M_4 the set $\Gamma^{\alpha}{}_{\beta\gamma} = 0$ describes an absolute parallelism on the manifold \tilde{G} , as described in section 4.2. Extending to the full bundle space P here we provisionally consider the $\acute{\Gamma}^{i}_{~jk}$ set of reference [18] listed in the third column of table 4.1. This set of linear connection coefficients are gauge covariant on P and compatible with the metric of equation 4.2 deriving from the gauge connection ω on P, that is with $\nabla \check{g} = 0$. On adopting such a linear connection on P, based on compatibility with the structures of the form $L(\mathbf{v}_{10}) = 1$ here, we then consider the implications of incorporating this element of Kaluza-Klein theory into the present framework.

The components of the Riemann curvature on the manifold $P \equiv M_4 \times SO^+(1,9)$ can be written in terms of the linear connection and structure coefficients, such as the set $\tilde{\Gamma}^i_{jk}$ described above, directly from equation 3.62 as:

$$
\acute{R}^{i}_{jkl} = \acute{e}_{k}\acute{\Gamma}^{i}_{jl} - \acute{e}_{l}\acute{\Gamma}^{i}_{jk} + \acute{\Gamma}^{i}_{mk}\acute{\Gamma}^{m}_{jl} - \acute{\Gamma}^{i}_{ml}\acute{\Gamma}^{m}_{jk} - \acute{c}^{m}_{kl}\acute{\Gamma}^{i}_{jm} \tag{5.2}
$$

In the present theory we begin with the translational symmetry of the form $L(\mathbf{v}_{10}) = 1$ over the manifold M_4 with a flat Minkowski metric $g_{ac}(x) = \eta_{ac}$ and the canonical flat $SO^+(1, 9)$ -valued connection ω on P. As described in chapter 2, initially in equations 2.35 and 2.36, this latter property means that the full curvature is zero $F=0$, or in components $\hat{F}^{\alpha}_{ab} = 0$. Hence, given that $\hat{\Gamma}^{\alpha}_{\ \beta\gamma} = \Gamma^{\alpha}_{\ \beta\gamma} = 0$ and $\hat{\Gamma}^{a}_{\ bc} = \Gamma^{a}_{\ bc} = 0$, all the linear connection coefficients in column 'O+P [18]' of table 4.1 are zero, $\acute{\Gamma}^{i}_{jk} = 0$, and in turn all components of the Riemann curvature tensor of equation 5.2 vanish on the principle bundle manifold P.

Here the natural absolute parallelism on M_4 and G has been generalised to a natural parallelism on $P \equiv M_4 \times G$ with $\dot{\Gamma}^i_{jk} = 0$ for all coefficients of the linear connection in the horizontal lift basis. In fact for the canonical zero full curvature $\hat{F}^{\alpha}_{ab} = 0$ on the principle bundle all five non-Levi-Civita choices for $\hat{\Gamma}^{i}_{jk}$ in table 4.1 lead via equation 5.2 to $\hat{R}^{i}_{jkl} = 0$, which as the components of a tensor vanish for any frame field on P, expressed generally as:

$$
\check{R}^i_{jkl} = 0 \tag{5.3}
$$

On the other hand there are torsion components with $\check{T}^i_{jk} \neq 0$ and hence the torsion is finite, as it is for the case of a self-parallel frame composed out of left-invariant vector fields on a Lie group manifold G , a copy of which here forms part of the total parallel frame field on P, as described in section 4.2.

In this way the full zero gauge curvature $\hat{F} = 0$ for \hat{G} over M_4 has been translated into zero Riemannian curvature $\check{R}^i_{jkl} = 0$ on the bundle space P. The question then remains regarding how this structure might provide the link through which the external gravitational field will relate to the internal gauge fields over the base space M_4 when the full symmetry is broken.

On the principle bundle $P \equiv M_4 \times SO^+(1,9)$ a trivialisation may be chosen such that the corresponding direct product basis for the tangent space TP is identical to the horizontal lift basis associated with the canonical flat connection ω , which in turn is derived from the full symmetry group. Such a trivialisation represents a gauge choice for which the section σ_0 , depicted in figure 5.2, on the principle bundle P coincides with the submanifolds of the integrable horizontal subspaces of ω on P, and hence with gauge connection components $\omega_{a}^{\alpha}(x, g) = 0$ on the bundle space. The linear connection components are identical $\ddot{\Gamma}^i_{jk} = \dot{\Gamma}^i_{jk} = 0$ in the respective direct product and horizontal lift bases for this choice of gauge, describing the absolute parallelism defined in the frame field adapted to this section on P.

While the canonical flat connection $\omega = \pi_2^* \theta$ on P describes a unique horizontal subspace and the corresponding horizontal lift basis, a direct product basis may be defined in terms of any section on the bundle. Indeed, more generally geometric objects over the base space M_4 may be described with respect to a choice of gauge on the bundle P, as for example determined by the section $\sigma' = \sigma_0 g$, with $g(x) \in \hat{G} = SO^+(1, 9)$, as also represented in figure 5.2. The gauge connection components $\omega^{\alpha}{}_{a}(x, g) \neq 0$ in the new trivialisation are such that the vectors of the horizontal lift basis $\{\hat{e}_i\}$ are expressed as for equation 3.26 with:

$$
\dot{e}_a = \ddot{e}_a - \omega^{\alpha}{}_{a}\ddot{e}_{\alpha} \qquad \text{with} \qquad \omega^{\alpha}{}_{a} \neq 0 \tag{5.4}
$$

while
$$
[\dot{e}_a, \dot{e}_b] = -\hat{F}^{\alpha}_{ab}\dot{e}_{\alpha}
$$
 with $\hat{F}^{\alpha}_{ab} = 0$ (5.5)

since the full $SO^+(1,9)$ zero curvature is a gauge independent structure. However the full group $SO^+(1,9)$ does not act purely as an internal symmetry but is broken by the action of the subgroup $SO^+(1,3) \subset SO^+(1,9)$ on the external tangent space TM_4 . While the choice of gauge $g(x) \in SO^+(1, 9)$ remains arbitrary with respect to the full unbroken symmetry it will affect the physics of the broken symmetry over M_4 . For the

Figure 5.2: Geometric objects Γ and ω on the principle bundle P in relation to the linear connection Γ on the base space M_4 and Maurer-Cartan 1-form θ on the group manifold \ddot{G} .

restricted set of internal SO(6) generators the horizontal lift vectors extracted from equations 5.4 and 5.5 have the properties:

$$
\underline{\dot{e}}_a = \underline{\ddot{e}}_a - \underline{\omega}^\alpha{}_a \underline{\ddot{e}}_\alpha \qquad \text{with} \qquad \underline{\omega}^\alpha{}_a \neq 0 \tag{5.6}
$$

while
$$
[\underline{\acute{e}}_a, \underline{\acute{e}}_b] = -\underline{F}^{\alpha}{}_{ab}\underline{\acute{e}}_{\alpha}
$$
 with $\underline{F}^{\alpha}{}_{ab} \neq 0$ (5.7)

Here the $\underline{\omega}^{\alpha}{}_{a}$ are the components of an so(6)-valued connection 1-form, with the sums over α restricted to the SO(6) generators, resulting in a generally non-zero internal curvature $\underline{F}^{\alpha}{}_{ab}$, as was demonstrated in equation 2.56 for the finite internal SO(2) curvature achieved for small $SO(5)$ gauge transformations over M_3 for the model world of section 2.3. Here we are reproducing the symmetry breaking approach of section 2.3 in the light of the principle bundle structure and Kaluza-Klein theories described in the previous two chapters.

As well as the transformation of the gauge connection ω for a different choice of basis on P the linear connection $\check{\Gamma}$ also transforms. For any change of frame $e_{i'} =$ $e_i e_{i'}^i$ with $e_{i'}^i \in GL(m, \mathbb{R})$ the transformation of a linear connection, displayed in equation 3.46, can be written as:

$$
\Gamma^{i'}_{j'k'} = e^{i'}_{i} e^{j}_{j'} e^{k}_{k'} \Gamma^{i}_{jk} + e^{i'}_{l} e_{k'} e^{l}_{j'} \tag{5.8}
$$

The gauge choice associated with the section $\sigma' = \sigma_0 g(x)$ on P corresponds to a transformation from the horizontal lift basis to an arbitrary direct product basis $\ddot{e}_{i'} =$ $\acute{e}_{i}e_{i'}^{i}$ on a principle bundle, that is the reverse of equation 3.26 or 5.4, and we have:

$$
\left(\begin{array}{cc}\n\ddot{e}_{a'} & \ddot{e}_{\alpha'}\n\end{array}\right) = \left(\begin{array}{cc}\n\epsilon_a & \epsilon_\alpha\n\end{array}\right) \left(\begin{array}{cc}\n\delta^a_{\ a'} & 0 \\
\omega^\alpha_{\ a'} & \delta^\alpha_{\ \alpha'}\n\end{array}\right) \tag{5.9}
$$

and hence
$$
e_{i'}^i = \begin{pmatrix} \delta^a_{a'} & 0 \\ \omega^{\alpha}_{a'} & \delta^{\alpha}_{a'} \end{pmatrix}
$$
 with inverse $(e^{-1})_{i}^{i'} = \begin{pmatrix} \delta^{a'}_{a} & 0 \\ -\omega^{\alpha'}_{a} & \delta^{\alpha'}_{\alpha} \end{pmatrix}$ (5.10)

As a consistency check the same transformation is applied to the full set of Levi-Civita connection coefficients in the horizontal lift basis $\acute{\Gamma}^i_{jk}$ as listed in the 'Cho [13]' column of table 4.1 (as extracted from [13] equation (22)). The expressions for $\overline{\text{the }\Gamma^{i'}}$ $j'_{j'k'}$ components obtained in the direct product basis using equations 5.8 and 5.10 is found to agree with the original reference (the $\bar{\Gamma}$ components in the notation of [13] equation (15)).

The general aim of this approach is to use equations 5.2 and 5.3, with $\tilde{R}^{i}_{jkl} = 0$ deriving from the full zero curvature $\hat{F} = 0$, on the principle bundle $P \equiv M_4 \times SO^+(1, 9)$ as a mutual constraint on the form of the external and internal curvature that results from the symmetry breaking. Once the full symmetry is broken non-zero internal gauge curvature components $\underline{F}^{\alpha}{}_{ab} \neq 0$ from equation 5.7 will be introduced quadratically into the terms of equation 5.2 via relations to the linear connection Γ of the kind listed in table 4.1 (by adopting for example the coefficients of column [18] as provisionally suggested above) and hence into correlation with the external curvature $R^a_{\;\; bcd} \neq 0$ on the base manifold as identified within the appropriate components of $\tilde{R}^{i}_{jkl} = 0$ in a suitable basis.

These structures emerge in the symmetry breaking as represented by the transition from figure 5.1(a) to (b). The structure of figure 5.1(a) implies that the total symmetry SO⁺(1,9) of $L(\mathbf{v}_{10}) = 1$ is associated with a canonical flat gauge field $A = g^*\theta$ with full curvature $\ddot{F}=0$, under which a correlation between the external curvature \overline{R} and internal curvature F is implied in the symmetry breaking to the structure of figure 5.1(b), in particular with the case of both $\overline{R} = 0$ and $F = 0$ simultaneously possible.

In this picture a non-zero external curvature $\overline{R} \neq 0$ on M_4 is absorbed under $\check{R}^i_{jkl} = 0$ on the extended bundle space as the 'buckling' of the geometry of the base manifold is countered by a corresponding finite internal curvature $\underline{F} \neq 0$. The external and internal curvature is hence generated in a necessarily mutually consistent way under the symmetry of $L(\mathbf{v}_{10}) = 1$ in a choice of $SO^+(1, 9)$ gauge over the base space M_4 . The invariance of the zero Riemann tensor of equation 5.3 under a change of frame adapted to choice of section is analogous to the invariance of the action integral of equation 4.10, defined in terms of a scalar curvature \tilde{R} , under variations of the metric \check{g}^{ij} of the kind described in section 4.2. This motivates the *conjecture* that this framework leads to a similar unification of the Einstein-Yang-Mills equations of motion, that is equations 4.16–4.18, as found for non-Abelian Kaluza-Klein theory but ultimately without the need to postulate a Lagrangian function, coupled with the variational principle for the corresponding action integral, to obtain these equations.

Resulting from the projection of the structure of figure $5.1(a)$ over that of figure 5.1(b), with a choice of an $SO^+(1,9)$ gauge section over M_4 for the former, two further bundle structures, associated with the latter figure, may be identified and considered separately. The subgroup $SO^+(1,3)$ is distinguished in that it acts on tangent space vectors $\overline{v}_4 \in TM_4$ of the base manifold, as depicted in figure 5.1(b), and therefore is designated as an *external* symmetry, with the residual $SO(6)$ acting

on the remaining components of $v_6 \text{ } \subset v_{10}$ of the form $L(v_{10}) = 1$ and constituting an internal symmetry. This results in consideration of the complementary subbundles $\overline{P} \equiv M_4 \times SO^+(1,3)$ and $\underline{P} \equiv M_4 \times SO(6)$ which effectively *decouple* from each other as mathematical structures, although related through the correlated geometrical structures they support, as they are mutually *carved out* of the initial unbroken bundle $P = M_4 \times SO^+(1,9).$

Indeed it is the extraction of the subgroup $SO^+(1,3) \subset SO^+(1,9)$, with the action of $SO^+(1,3)$ identified as the external symmetry and absorbed into the local tangent space geometry on M_4 , that *breaks* the full $SO^+(1,9)$ symmetry. The base space M_4 is naturally associated with the frame bundle FM_4 , which is itself a particular type of principle fibre bundle as described in section 3.3. The bundle space $\overline{P} \equiv$ $M_4 \times SO^+(1,3)$, obtained as a *restriction* of the $P \equiv M_4 \times SO^+(1,9)$ bundle, can also be interpreted as a *reduction* of the $FM₄$ frame bundle. In turn an so⁺(1,3)-connection on \overline{P} may be *extended* to a gl $(4,\mathbb{R})$ -connection on the frame bundle, together with the associated tetrad $e^{\mu}_{a}(x)$ and metric $g_{\mu\nu}(x)$ fields on M_4 , as familiar in the theory of general relativity and also described in section 3.3.

As described towards the end of section 3.4 the $SO^+(1,3)$ symmetry can be treated by analogy with an 'internal' Yang-Mills gauge structure. Indeed, as described above for the full $SO^+(1, 9)$ symmetry, quadratic terms in the external $SO^+(1, 3)$ 'gauge curvature' $\overline{F}^{\alpha}{}_{ab}$ will appear in the third and fourth terms of equation 5.2 (essentially as described in [25], which adopts the Levi-Civita connection on the bundle space, leading to equations (3.14) and (3.15) there). However this same external geometry, from the action of SO⁺(1,3) on TM₄, is represented by the Riemannian curvature $R^a{}_{bcd}(x)$, which is also contained within the corresponding \check{R}^i_{jkl} components in a suitable basis (as also described in [25]). Hence the bundle \overline{P} appears to incorporate a redundant description of the external geometry while lacking an explicit reference to the internal curvature.

On the other hand the subbundle $\underline{P} \equiv M_4 \times SO(6)$ is closely related to both the frame bundle FM_4 , upon which the external $SO^+(1,3)$ geometry is expressed in terms of fields such as $g_{\mu\nu}(x)$, as well as the structures of the internal SO(6) geometry with the associated gauge field $Y^{\alpha}_{\mu}(x)$ and curvature $\underline{F}^{\alpha}_{\mu\nu}(x)$ components constructed on $M₄$. Hence in principle all the necessary geometric structures for relating the external and internal curvature can be identified on the bundle P .

Rather than dealing with a connection form ω over M_4 for the full $\ddot{G} =$ $SO⁺(1, 9)$ symmetry it is precisely through the symmetry breaking action, with the degrees of freedom of the $SO^+(1,3) \subset SO^+(1,9)$ subgroup part of the *gauge* connection being converted into the freedom of a *linear* connection on M_4 , that the bundle space $\underline{P} \equiv M_4 \times SO(6)$ emerges. This in turn implies that the structure of the zero curvature $\hat{F} = 0$ for the full canonical flat connection does not explicitly survive the symmetry breaking transition from figure $5.1(a)$ to (b) .

This motivates the study of a unified framework on the space $\underline{P} \equiv M_4 \times SO(6)$ considered from now as a principle bundle standing independently by itself, and not as subbundle 'carved out' of a larger bundle space such as $P \equiv M_4 \times SO^+(1,9)$. It remains then to explicitly define the mathematical nature of the constraint between the internal $SO(6)$ curvature and external $SO⁺(1, 3)$ geometry in terms of the bundle P.

Earlier in this section an absolute parallelism on the bundle $P \equiv M_4 \times SO^+(1,9)$ was constructed in the horizontal lift basis with all $\acute{\Gamma}^{i}_{jk} = 0$, taken from the set of reference [18] listed in the third column of table 4.1 for the canonical zero full curvature $F=0$, implying the identity of equation 5.3. Now, beginning directly on the bundle $\underline{P} = M_4 \times SO(6)$ in itself, the question arises concerning the possible definition of a linear connection on this space. Since there is a gauge connection (which now derives from the internal $SO(6)$ symmetry and in general is not flat) on \underline{P} the horizontal lift basis may be employed, and in turn the natural metric structure with components \dot{g}_{ij} of equation 4.4 introduced.

Hence it is possible to define the unique Levi-Civita connection on this bundle, as described in section 4.1, with the components of equation 4.6 as listed for the horizontal lift basis in the first column of table 4.1 under 'Cho [13]'. However in the present theory at no stage is \underline{P} considered to be a *physical* space or spacetime structure, hence neither the metric \acute{g}_{ij} nor a linear connection $\acute{\Gamma}^{i}_{jk}$ on \underline{P} have a physical geometric meaning, as they do on the base space M_4 . Hence the unique metric-compatible torsion-free Levi-Civita connection is not here considered to be a natural structure on the bundle space as it is for the base manifold, and an alternative argument for the form of Γ on P is sought.

In particular the linear connection on \underline{P} is expected to be closely associated with the linear connection Γ on the base space M_4 , which does describe a physical geometry. Since this is a gl $(4,\mathbb{R})$ -valued 1-form $\Gamma(x) = \Gamma^a{}_{bc} E^b{}_a e^c$ on M_4 , with respect to the distinguished horizontal lift basis on \underline{P} the components $\acute{\Gamma}^a{}_{bc}$ and $\acute{\Gamma}^a{}_{b\gamma}$ alone may be favoured for a linear connection Γ on M_4 in some sense lifted onto \overline{P} , and hence the only non-trivial coefficients of $\hat{\Gamma}^i_{~jk}$ on \underline{P} might be taken to be:

$$
\acute{\Gamma}^{a}{}_{bc} = \Gamma^{a}{}_{bc} \quad \text{and} \quad \acute{\Gamma}^{a}{}_{b\gamma} = \gamma g^{ac} g_{\beta\gamma} F^{\beta}{}_{bc} \tag{5.11}
$$

as listed as the 'minimal' set in the final column of table 4.1. The form of $\hat{\Gamma}^a_{\ \, b\gamma}$ in the equation above and the third row of the table as adopted from the other models in the table, consistent with the requirement that $\hat{\Gamma}$ should transform in a gauge covariant manner on \underline{P} as appropriate for any object relating to a physical entity on the base manifold M_4 . As will be described below this proposal will amount to a minimal structure on \underline{P} linking the present theory with Kaluza-Klein theory with a manifest correlation between the external Riemannian geometry and internal gauge curvature.

While the Levi-Civita connection, $\Gamma = f(g)$ of equation 3.53, on M_4 provides a unique description of the geometry on the base manifold in terms of the metric $g_{\mu\nu}(x)$, the linear connection Γ of equation 5.11 represents an attempt to extend this structure onto P while maintaining the character of the connection Γ on M_4 , concerning in particular the gl $(4, \mathbb{R})$ -valued property. However any linear connection on \underline{P} is intrinsically a gl (m, \mathbb{R}) -valued 1-form (where $m = 4 + 15$ for the internal SO(6) gauge group). For example under the transformation to a direct product basis, as described in equations 5.8–5.10, the components of equation 5.11 in general give rise to linear connection coefficients $\ddot{\Gamma}^{\alpha}_{bc} \neq 0$ and $\ddot{\Gamma}^{\alpha}_{b\gamma} \neq 0$ in addition to $\ddot{\Gamma}^{a}_{bc} \neq 0$ and $\ddot{\Gamma}^{a}_{b\gamma} \neq 0$. Since the character of being $gl(4, \mathbb{R})$ -valued cannot be upheld for a linear connection on \underline{P} an alternative proposal, and one for which parallel transport in the horizontal and vertical directions on \underline{P} more directly reflects the geometry of the base manifold M_4 , will be considered.

A direct way to obtain a linear connection $\Gamma(p)$ on P closely related to $\Gamma(x)$ on M_4 would be to define $\check{\Gamma} = \pi^* \Gamma$ as the pull-back of the gl(4, R)-valued 1-form Γ through the bundle projection $\pi : \underline{P} \to M_4$, by analogy with the identification of the canonical Lie algebra-valued 1-form $\omega = \pi_2^*\theta$ as the pull-back of the Maurer-Cartan 1-form θ through the projection map π_2 : $P \equiv M_4 \times \hat{G} \rightarrow \hat{G}$ for the full bundle P as described in figure 5.2. Indeed, the gl(4, R)-valued linear connection Γ on M_4 , associated with the external symmetry, and $L(G)$ -valued 1-form θ on \tilde{G} , associated with the full symmetry, each describe the parallelism on their respective manifolds.

The canonical flat connection $\omega = \pi_2^* \theta$ on $P \equiv M_4 \times SO^+(1, 9)$ itself is an unambiguous geometric object, completely independent of any particular choice of section or gauge over the base manifold. It derives purely from the properties of θ on the gauge group $\hat{G} = SO^+(1, 9)$. Similarly, a linear connection $\check{\Gamma} = \pi^* \Gamma$ on P or <u>P</u> has no physical significance in itself other than that derived from its relation to a linear connection Γ, and the related Riemannian geometry, on the base manifold M_4 . For the case $\check{\Gamma}(p) = \pi^* \Gamma(x)$ on \underline{P} and for any vector field $X(p) \in T\underline{P}$ we have:

$$
\langle \check{\Gamma}, X \rangle_p = \langle \pi^* \Gamma, X \rangle_p = \langle \Gamma, \pi_* X \rangle_x \tag{5.12}
$$

with $X(p)$ projected in the final line onto the vector $\pi_* X \in TM_4$ and with $\pi : p \in$ $\underline{P} \to x \in M_4$. Hence for any vector in the vertical subspace $X(p) \in V\underline{P}$ we have $\langle \Gamma, X \rangle_p = 0$ since $\pi_* X = 0$. This structure is related to the linear connection on the bundle described for the case of Kalinowski [19] in section 4.2 for which all tangent vectors are mapped onto their horizontal parts, again with the property $\langle \Gamma, X \rangle_p = 0$ for any vertical vector X , and hence again with emphasis on the horizontal structure, which in turn is closely associated with the geometry of the base space M_4 . In fact consideration of all cases collected in table 4.1 leads to the following proposal for the properties of $\check{\Gamma}$ on \underline{P} appropriate for the present theory:

- a) parallel propagation via $\dot{\Gamma}$ in the vertical directions is taken to be trivial in the manner of [19] in the fourth column of table 4.1.
- b) parallel propagation via $\check{\Gamma}$ in the horizontal directions on \underline{P} is taken to relate to the contours of the gauge curvature over the base space M_4 , following [20] in the fifth column of table 4.1 for the case $c = 1$.
- c) with a view to deriving physical equations on the base space M_4 compatibility with gauge covariance should be observed, as emphasised in [18].
- d) consistent with c) the linear connection may be compatible with the natural, but non-physical, metric \check{g}_{ij} of equation 4.4, although the torsion may be arbitrary as initially emphasised in [17].
- e) the bundle \underline{P} serves as an arena to relate the external and internal symmetry structures compatible with the simultaneous possibility of $\overline{R} = 0$ and $\underline{F} = 0$, as derived from consideration of figure 5.1 for the present theory.

Based on these observations and the broader discussion of Kaluza-Klein theory in chapter 4 the conjectured linear connection components on \underline{P} , as extracted from table 4.1, can be summarised as:

4)
$$
\dot{\Gamma}^{a}_{\gamma b} = g^{ac} g_{\gamma \beta} F^{\beta}_{\;bc}, \qquad 5) \dot{\Gamma}^{a}_{\;ab} = F^{\alpha}_{\;ab}, \qquad 6) \dot{\Gamma}^{a}_{\;bc} = \Gamma^{a}_{\;bc} \tag{5.13}
$$

with all other $\acute{\Gamma}^{i}_{jk} = 0$. Hence these are essentially the set of [19] in the fourth column of table 4.1 with $\lambda = 2$, with the motivation for employing this latter value derived from the geometrical argument in [20]. This latter argument also has the benefit of fixing the geometry of Γ without any reference to the Levi-Civita connection on P.

The whole purpose of constructing a linear connection Γ on \underline{P} , as described above, is to provide a means through which a correlation between the external and internal curvature may be explicitly described. On the spacetime manifold M_4 any relationship between the external geometry, expressed in terms of the Einstein tensor with components $G_{\mu\nu}(x)$, and the internal geometry, expressed in terms of the gauge curvature with components $\underline{F}^{\alpha}{}_{\mu\nu}(x)$, must transform covariantly both under general coordinate transformations and under gauge transformations, as described in particular in section 3.4. One technique for obtaining such a relation is to first identify a scalar 'Lagrangian' function which has these invariance properties, as described in section 3.5. This approach, again following the Kaluza-Klein theories, will be adopted provisionally here, although a more direct geometric argument leading to equation 4.16–4.17, which itself has the desired symmetry properties, would ultimately be preferred. (Since in the following components such as $F^{\alpha}_{\ \mu\nu}(x)$ will always refer to the purely internal gauge curvature we now omit the underscore for these objects).

While earlier in this section the Riemannian curvature \acute{R}^{i}_{jkl} was constructed on the full bundle $P \equiv M_4 \times SO^+(1,9)$ we are now focusing on the bundle $P \equiv M_4 \times SO(6)$, upon which the gauge curvature is generally finite. For any linear connection on the bundle space \underline{P} , such as defined by any of the six sets of connection coefficients $\acute{\Gamma}^i_{jk}$ listed in table 4.1, the Riemann curvature tensor can be determined according to equation 5.2, which is specified in the horizontal lift basis. The corresponding Ricci curvature components $\acute{R}_{\alpha\beta}$ and \acute{R}_{ac} are listed here in the first and fourth rows of table 5.1 for the six familiar examples. In all cases the entries in this table calculated here agree with the corresponding equations of the given references – within the sign conventions such as that of equation 3.74 and as alluded to near the opening of chapter 4.

The scalar curvature constructed in the horizontal lift basis on the principle bundle space can be written as:

$$
\acute{R} = \acute{g}^{ij}\acute{R}_{ij} = g^{ac}\acute{R}_{ac} + K^{\alpha\beta}\acute{R}_{\alpha\beta} \tag{5.14}
$$

owing to the simple form of the metric \acute{g} in this basis as expressed in equation 4.4. Hence the Ricci curvature components $\hat{R}_{a\alpha}$ and $\hat{R}_{\alpha a}$ are not required in order to determine the scalar curvature on the bundle.

If each of the four factors of $\frac{1}{2}$ in the 'Cho [13]' column in table 4.1, for the case of Levi-Civita connection coefficients $\tilde{\Gamma}^i_{jk}$ on the bundle, listed in rows 1), 3), 4) and 5) are replaced by the real factors f_1 , \dot{f}_3 , f_4 and f_5 respectively then the scalar curvature in the horizontal lift basis is found to be:

$$
\hat{R} = R_M + R_G + \chi F^2 \tag{5.15}
$$

with
$$
\chi = f_3 - f_3 f_4 + f_4 f_5 - f_3 f_5
$$
 (5.16)

This expression agrees with the scalar curvature for the Levi-Civita case, with each $f_i = \frac{1}{2}$ $\frac{1}{2}$, as quoted originally in equation 4.9, and with each subsequent case of table 4.1 as quoted in the final row of table 5.1. Equations 5.15 and 5.16 show that f_3 is the only

	Cho $[13]$	Kop [17]	$O+P$ [18]	Kal $[19]$	Kat $[20]/\text{min}$
$\hat{R}_{\alpha\beta} = R_{(G)\alpha\beta} + \Vert$	$-\frac{1}{4}F_{\alpha bd}F_{\beta}^{bd}$		$-\frac{1}{4}F_{\alpha bd}F_{\beta}^{\ \ bd}$		
$R_{(G)\alpha\beta} =$	$\frac{1}{4}K_{\alpha\beta}$	$(\alpha - \alpha^2)K_{\alpha\beta}$	θ		$\left(\right)$
$K^{\alpha\beta}\acute{R}_{\alpha\beta} =$	$R_G - \frac{1}{4}F^2$	R_G	$-\frac{1}{4}F^2$		Ω
$\hat{R}_{ac} = R_{(M)ac} +$	$\frac{1}{2}F^{\beta}_{ad}F_{\beta c}^{\ \ d}$	$\overline{0}$	$\frac{1}{2}F^{\beta}_{ad}F_{\beta c}^{\ \ d}$	$\frac{\lambda^2}{4}F^{\beta}_{ad}F_{\beta c}^{\ \ d}$	$c^2F^{\beta}_{ad}F_{\beta c}^{\ \ d}$
$g^{ac}\dot{R}_{ac}=R_M+\Vert$	$\frac{1}{2}F^2$		$\frac{1}{2}F^2$	$\frac{\lambda^2}{4}F^2$	c^2F^2
	$\acute{R}=\acute{g}^{ij}\acute{R}_{ij}=\left\ \right. R_{\!M}+R_{\!G}+\frac{1}{4}F^2$	$R_M + R_G$	$R_M + \frac{1}{4}F^2$	$R_M + \frac{\lambda^2}{4}F^2$	$R_M + c^2 F^2$

Table 5.1: Composition of the scalar curvature \hat{R} on the bundle space for the six cases of table 4.1. Contributions to the components of the Ricci curvature on the bundle include $R_{(G)\alpha\beta}$ and $R_{(M)ac}$ from the group manifold and base space respectively, with $R_G = K^{\alpha\beta} R_{(G)\alpha\beta}$ and $R_M = g^{ac} R_{(M)ac}$ being the respective scalar curvatures. The results for the sixth, 'minimal', case in table 4.1 are identical to those listed for [20] in the final column above with $c^2 = \gamma$.

coefficient which is sufficient in itself to introduce a non-trivial F^2 term, alongside R_M , into the scalar curvature \hat{R} , and this observation in part motivated the consideration of this simplest set of $\tilde{\Gamma}^i_{jk}$ coefficients, as listed in the 'minimal' column of table 4.1 and described in equation 5.11 above. While perhaps not developed as a serious physical proposal this minimal model further demonstrates the flexibility within the Kaluza-Klein framework, obtaining the appropriate link between the external geometry and internal curvature with a seemingly much simpler linear connection on the bundle compared with the Levi-Civita case. More generally, equations 5.15 and 5.16 display the mutual consequences of the non-zero $\tilde{\Gamma}^i_{jk}$ terms for the models listed in table 4.1.

Since $\hat{R}(p)$ is a scalar field on the bundle at any given point p it takes the same value in any local frame. Hence for example in a direct product basis, corresponding to a section σ on P, the scalar value is simply $\ddot{R}(p) = \dot{R}(p)$. Further, since each of the scalar terms in the bottom line of table 5.1 is gauge invariant, a corresponding scalar function on the base space M_4 may be deduced as:

$$
\tilde{R}(x) = \sigma^* \ddot{R}(p) = R_M + R_G + \chi F^2 \tag{5.17}
$$

which is equivalent to $\hat{R}(p)$ for any $p \in \underline{P}$ such that $\pi(p) = x \in M_4$. Hence $\tilde{R}(x)$ is a real scalar function on M_4 which contains information about both the external and internal geometry, is invariant both under coordinate and gauge transformations on the base space, and therefore makes a suitable 'Lagrangian' candidate on M_4 . Whether or not R_G vanishes and the real value χ in equation 5.17 depend upon the particular model, as can be seen for the examples of table 5.1 and via equation 5.16 respectively. For the case of most interest for the present theory, with non-zero linear connection coefficients listed in equation 5.13, corresponding to setting $\lambda = 2$ in the 'Kal [19]' columns of tables 4.1 and 5.1, we have simply $\tilde{R}(x) = R_M + F^2$.

The starting point for the Kaluza-Klein theories reviewed in sections 4.1 and 4.2 is the mathematical structure of a principle fibre bundle $P = (M_4, G)$, such as described in section 3.1 and pictured in figure 3.1. This structure features an extended base space M_4 over which a gauge connection may be introduced on the bundle space P transforming under the internal symmetry gauge group G. In these theories the bundle space is typically interpreted as a higher-dimensional physical spacetime. For example in reference ([18] p.190) the authors write: 'Our general attitude is to regard the n_G vertical dimensions as physically real, and hence the vertical Einstein equations as true dynamical equations of the $(n + n_G)$ -theory.'

A similar perspective is generally adopted for the theories with homogeneous fibres, described in section 4.3, in this case for the bundle space $E = (M_4, S_k)$. In the introduction of reference [24] the authors write: 'Kaluza-Klein theories are theories in which the gravitational potential together with the gauge potentials of various interactions are interpreted as manifestations of (pseudo-) Riemannian structure of the Universe which is $4 + k$ dimensional.' The analogy between coordinate transformations in general relativity and gauge transformations in gauge theory, discussed in section 3.4, is more explicitly realised in these theories as demonstrated for example in equations 4.24.

In Kaluza-Klein theories restrictions on the form of the metric \check{g}_{ij} on the higherdimensional space, in particular a necessary conformity with equation 4.3, induce a 'dimensional reduction' or 'spontaneous compactification' of the larger space. The latter is then interpreted as a bundle structure with fibres, corresponding to the n_{G} dimensional gauge group G or an associated k-dimensional homogeneous space S_k , over the smaller $n = 4$ -dimensional spacetime M_4 .

The origin of the bundle structure in Kaluza-Klein theories hence contrasts sharply with that for the present theory. Here the geometric structure $M \times G$ arises out of the symmetries of a general form of temporal flow $L(v) = 1$ as described in chapter 2. In particular for the 10-dimensional form $L(\mathbf{v}_{10}) = 1$, considered in this section and employed in figure 5.1, the base space M_4 arises out of a parametrisation of a 4-dimensional subset of the 'translational' degrees of freedom of the components v_{10} under $L(\mathbf{v}_{10}) = 1$, with gauge fields drawn over the base space out of the 'rotational' degrees of freedom of the same temporal form.

Here the only *physical* space is the manifold M_4 , providing the arena for general relativity in a 4-dimensional spacetime, with no 'compactification' from a higherdimensional extended spacetime required. The spacetime geometry on M_4 derives from the local Minkowski metric η_{ab} implicit in the 4-dimensional temporal form of equation 5.1, now written $L(\mathbf{v}_4) = h^2$ in the projection out of the higher-dimensional form $L(v_{10}) = 1$. On the other hand the Killing metric $g_{\alpha\beta} = K_{\alpha\beta}$ does not describe the geometry of a physical space, either on the group manifold G or bundle space P. It relates the Lie algebra adjoint and coadjoint representations as usual, with for example $F_{\alpha ab} = g_{\alpha\beta} F^{\beta}_{ab}$, and it may be employed as a mathematical structure on \underline{P} in the derivation of scalar quantities, as for example in equation 5.14.

The roles of the metric $g_{\mu\nu}(x)$ and gauge field $Y^{\alpha}_{\mu}(x)$ in the laws of physics on M_4 are well defined. When lifted to the principle bundle \underline{P} these objects can be augmented by the Killing metric $g_{\alpha\beta}$ on G to define a metric \check{g}_{ij} in the form of equation 4.5 on the bundle space. This latter metric could be employed on \underline{P} , for example to construct a curvature scalar \check{R} from the Riemann tensor \check{R}^i_{jkl} based on a Levi-Civita connection $\check{\Gamma}^i_{jk}$, but no *physical* significance should be attached to the

geometric connotations of the metric \check{g}^{ij} introduced in this way.

Indeed \check{g}^{ij} , as described in equation 4.4, consists of an unnatural marriage with the external local metric η_{ab} originating within the form of $L(\mathbf{v}_{10}) = 1$ upon which the group G, with Killing metric $g_{\alpha\beta}$, acts. This is the case whether the group describes full symmetry $\tilde{G} = SO^+(1,9)$, as considered earlier in this section, or the internal symmetry $G = SO(6)$ as considered here. Such a hybrid metric \check{g}_{ii} , composed of parts of quite different character, hence seems an unnatural object to endow with a physical geometric meaning. Hence here the construction of a Levi-Civita connection on the bundle space as described in subsection 4.1 is not well motivated, with the bundle not considered as representing an extension of general relativity to a higher-dimensional space. On the other hand with this unifying framework taking the shape of a principle fibre bundle over the base space the present theory is naturally related to Kaluza-Klein theories, in particular those of the kind reviewed in section 4.2.

While the structure of these Kaluza-Klein theories rests on a deliberate *exten*sion of the formalism of general relativity into a space with extra dimensions, in the present theory the construction of a linear connection on the bundle space is motivated rather as a mathematical means to relate the physical Riemannian curvature on the base space M_4 to that of the internal gauge fields. Indeed it is still possible to define a linear connection Γ on the bundle which is closely associated with both the linear connection $\Gamma^a{}_{bc}$ on the base space M_4 and the internal gauge curvature $F^{\alpha}{}_{ab}$, however only the linear connection Γ on M_4 has a significance in terms of describing a physical space.

As for other branches of this theory, including its connections with the Standard Model, quantum theory and cosmology to be presented subsequently in this paper, the aim is to develop the theory naturally out of the basic conceptual ideas. Here it is the basic geometric structures relating to the symmetries of $L(v) = 1$, in particular in the symmetry breaking over the M_4 base manifold pictured in figure 5.1, that provides the unified framework for the external and internal curvature. The resulting geometric structure, exemplified here by the principle bundle $\underline{P} \equiv M_4 \times SO(6)$, while not forming a physical spacetime itself, provides the mathematical arena for a unification of the external and internal geometry arising out of the breaking of the full $L(v_{10}) = 1$ symmetry over the base space M_4 .

The general form of the relation between the external Riemannian geometry \boldsymbol{R} and internal gauge curvature F is conjectured to arise naturally in this framework, in a generally and gauge covariant manner, essentially taking the form of equation 4.16– 4.17. This relation is provisionally derived here via the scalar function $R(x)$ of equation 5.17, interpreted as a geometric perturbation to the Einstein-Hilbert action on the base space M_4 arising from the higher-dimensional form of temporal flow $L(\mathbf{v}_{10}) = 1$. In particular, from the range of models studied, with linear connection coefficients $\acute{\Gamma}^{i}_{jk}$ on the bundle listed in table 4.1 and the corresponding scalar curvature \hat{R} determined in table 5.1, the argument outlined in points $(a) - e$)' earlier in this section leads to the proposed set of equation 5.13. This argument focuses on the horizontal transport in \underline{P} skirting over the base manifold M_4 , and in appealing in particular to references [19] and [20] meets half-way with Kaluza-Klein theory. Further progress might be made for example by placing more complete emphasis on point 'b)' with a full set of $\acute{\Gamma}^{i}_{jk}$ coefficients defined in terms of the parallel transport associated with the internal gauge

curvature as described for figure 3.3.

In standard Kaluza-Klein theory the action A_m for the scalar curvature \check{R} defined on the bundle space P in equation 4.10 reduces to the 4-dimensional action integral A_4 of equation 4.11 owing to the trivial integration over the fibre degrees of freedom. The point of view adopted here is that the scalar field $R(x) = R_M + \chi F²$ of equation 5.17 (with $R_G = 0$ and $\chi = 1$ for the model of equation 5.13 constructed here) is defined *directly* on the base space M_4 itself. In turn the action integral is defined directly on the base space as:

$$
\tilde{I} = \int (R_M + \chi F^2) \sqrt{|g|} \, d^4x \tag{5.18}
$$

as a coordinate and gauge invariant expression with all fields defined on M_4 . As denoted by the 'tilde' on \tilde{I} this function is considered as a perturbation of the Einstein-Hilbert action for the vacuum case, equation 3.79 with $\alpha = 1$, $\Lambda = 0$ and $\mathcal{L} = 0$, which was described in the opening of section 3.5. That is, equation 5.18 incorporates the perturbation to the scalar curvature $R_M(x) \to R(x)$ on the base space M_4 . The full Einstein-Hilbert action of equation 3.79 can be written:

$$
I = \int (\alpha R_M + \mathcal{L}) \sqrt{|g|} \, d^4 x \tag{5.19}
$$

where the cosmological constant Λ has been dropped in correspondence with the lack of a finite R_G term in equation 5.18. Further comparison between the above two equations shows that equation 5.18 describes a perturbation to general relativity equivalent to the introduction of a Lagrangian term $\mathcal{L} = +\alpha \chi F^2$ in the original Einstein-Hilbert action. While the mathematical conclusion is identical to Kaluza-Klein theory, here the interpretation involves a more minimal impact on the arena of general relativity in 4-dimensional spacetime, namely without a physical augmentation into a higherdimensional extended spacetime.

The choice of $\chi = \frac{1}{4}$ $\frac{1}{4}$ and $\alpha = \frac{-1}{16\pi G_N}$ respectively in the two equations above represents the standard normalisation for the incorporation of gauge fields into the Einstein-Hilbert action, as described in section 3.5. This standard action if also discussed in ([26] section 20.6) where the shortcomings of the Lagrangian approach are highlighted. The intention of the present theory is ultimately to avoid any direct reference to the Lagrangian formalism entirely. For the present case the form of $\tilde{R} = R_M + \chi F^2$ in equation 5.18, in deriving from equation 5.17, arises from the geometry on the bundle $\underline{P} = M_4 \times SO(6)$ in a physically meaningful way in terms of entities on the base space M_4 . This structure can be considered as a perturbation to general relativity deriving from the need to take into account the internal space of the form $L(\mathbf{v}_{10}) = 1$ and the geometric structures entailed.

If the 4-dimensional form $L(v_4) = 1$ of equation 5.1 alone is considered no symmetry breaking is involved in the identification of the bundle $P \equiv M_4 \times SO^+(1,3)$ out of the symmetries of this form. As described in section 2.2, in the context of the $SO(3)$ model, this structure incorporates a *canonical* flat connection with zero curvature, that is $R^{\rho}_{\sigma\mu\nu}(x) = 0$, without any reference to a Lagrangian. This result is however identical to that achieved in equation 3.82 for the vacuum case using the stationarity of the Einstein-Hilbert action under variation of the metric field $g_{\mu\nu}(x)$

on M_4 ; since $R^{\rho}_{\sigma\mu\nu}(x) = 0$ if $G^{\mu\nu}(x)$ vanishes everywhere in spacetime. Hence the conjecture here is that a perturbation to this Einstein-Hilbert action, in the form of equation 5.18, carries with it the consequences for the Riemannian geometry on M_4 that follow from an embedding in the structures of a larger form of temporal flow such as $L(v_{10}) = 1$.

Here the provisional adoption of a 'Lagrangian function' has a direct conceptual motivation. This is unlike for example the case of the Standard Model Lagrangian for particle physics, elements of which will be reviewed in section 7.2, for which both the fields and Lagrangian terms are generally introduced and contrived by hand with the aim of achieving the desired equations of motion and particle interactions for the known phenomena of high energy physics. The means of bypassing the Standard Model Lagrangian for the present theory will then be described in subsequent chapters, while the avoidance of a necessary Lagrangian to derive classical equations of motion will be considered further here in the following section.

Within this caveat for the employment of a Lagrangian approach, the equation of motion obtained by requiring $\delta \tilde{I} = 0$ for equation 5.18, under variations $\delta g_{\mu\nu}(x)$ of the metric on M_4 , follows the derivation of equation 4.16–4.17 and can be written here as:

$$
G^{\mu\nu} = 2\chi(-F^{\alpha\mu}_{\quad\rho}F_{\alpha}^{\;\;\rho\nu} - \frac{1}{4}g^{\mu\nu}F^{\alpha}_{\;\rho\sigma}F_{\alpha}^{\;\;\rho\sigma}) =: -\kappa T^{\mu\nu} \tag{5.20}
$$

At the purely theoretical level the factor of χ in this equation arises directly in equations 5.15 and 5.16, which in turn derive from the relation of linear connection $\dot{\Gamma}^i_{\;ik}$ on the bundle to the gauge curvature F^{α}_{ab} as listed in the columns of table 4.1. For the present theory the correlation between the external and internal geometry in the breaking of the full form $L(\hat{v}) = 1$ over the base space M_4 has been considered provisionally in terms of the set of linear connection coefficients of equation 5.13, and hence with $\chi = 1$.

With gravitational and gauge field phenomena historically studied independently in practice the normalisation factor connecting the left-hand side and central expressions of equation 5.20 is a matter for empirical convention, as for the factor of $\kappa = \frac{8\pi G_N}{c^4}$ $\frac{t_{GN}}{c^4}$ on the right-hand side of this equation. Here for normalisation in practice we shall set $\chi = \frac{\kappa}{2}$ $\frac{\kappa}{2}$ implying a choice of physical units such that the energy-momentum tensor can be expressed directly in terms of the gauge curvature, as will be the case for the electromagnetic field tensor $F_{\mu\nu}$ in the following section (see for example equation 5.28).

Equation 5.20 reduces to the vacuum solution $G^{\mu\nu}(x) = 0$ for the case in which curvature of the internal gauge field vanishes $F^{\alpha}_{\mu\nu} = 0$. More generally, with the Einstein tensor $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2} R_M g^{\mu\nu}$, contracting the equation 5.20 with $g_{\mu\nu}$ leads to the conclusion $R_M = 0$, the standard vanishing of the scalar curvature associated with a classical gauge field, while the Ricci curvature is generally finite with $R^{\mu\nu} = G^{\mu\nu} \neq 0$.

Hence while for a general solution we have $\tilde{R} = \chi F^2 \neq 0$, the full expression $\tilde{R} = R_M + \chi F^2 \neq 0$ is needed in equation 5.18 in order to derive the field equation 5.20 through the method of variation. A similar observation applies for the vacuum equations of general relativity, namely the derivation of equation 3.82, and further suggests that the Lagrangian approach may not be entirely satisfactory. Ideally the aim here would be to derive equation 5.20 purely by geometrical means and without reference to a Lagrangian. In the meantime, by further considering $\delta\tilde{I}=0$ for the action in

equation 5.18, now with respect to variation in the gauge fields $\delta Y^{\alpha}_{\mu}(x)$, leads, as described earlier for equation 4.18, to the Yang-Mills vacuum equation:

$$
D_{\mu}F^{\alpha\,\mu\nu} = 0 \tag{5.21}
$$

For the case of an Abelian internal U(1) symmetry this relation expresses Maxwell's equation for a source-free electromagnetic field.

While the unification has been described here in terms of the principle bundle space $P \equiv M_4 \times SO(6)$, for the broken group symmetry action, a bundle of homogeneous fibres $E \equiv M_4 \times S_k$ might also be constructed, with fibres composed of the purely internal \underline{v}_6 components of $L(v_{10}) = 1$, complementary to the projection onto the external spacetime with $\overline{v}_4 \in TM_4$ as pictured in figure 5.1(b). A transitive action of $SO^+(1, 9)$ on the space underlying $L(v_{10}) = 1$ can be identified, as for the action of $SO(6)$ on the internal space which hence forms the homogeneous space S_k employed for the fibres. Since these actions are also effective the complete internal gauge symmetry dynamics will be represented for the theory formulated in terms of a bundle with homogeneous fibres, rather than the principle fibre bundle, as was reviewed in section 4.3.

In the models of section 4.3 the internal group G can be considered as a global *isometry*, that is a symmetry preserving a metric $g_{\alpha\dot{\beta}}$ on S_k , with $H \subset G$ as the *isotropy* subgroup leaving any point $y_0 \in S_k$ fixed. By contrast for the present theory $G = SO⁺(1, 9)$ can be considered as an *isochronal* symmetry preserving the temporal form $L(v_{10}) = 1$ with $\overline{H} = SO^+(1,3) \subset \hat{G}$ as the local *isometry* subgroup preserving the metric on TM₄, while the complementary $\underline{H} = SO(6) \subset SO^+(1,9)$ leaves any vector $\overline{v}_4 \in TM_4$ fixed. The bundle structures on $E \equiv M_4 \times S_k$ may ultimately shed further light on the derivation of equation 5.20 together with the theoretical value of χ.

While a consistent and rigorous *mathematical* framework needs to be established a full understanding of the appropriate conceptual picture for the extraction of the geometry on the base manifold derived from, and breaking, the symmetries of the full form $L(\hat{\boldsymbol{v}}) = 1$ is also required. It is out of the marriage of these mathematical and conceptual ideas that an ultimate form for the relationship between the external Riemannian curvature \boldsymbol{R} and internal gauge curvature F on the base space M_4 might be arrived at. This section has described the evolution of ideas arising out of the symmetries of $L(v) = 1$ described in chapter 2, steered by the structures of differential geometry and Kaluza-Klein theory as described in chapters 3 and 4, aiming towards such a unification. Attempting to justify all the steps along the way, via the linear connection on the bundle of equation 5.13, scalar function on the base space of equation 5.17 (with $R_G = 0$) and action integral of equation 5.18, the aim has been to arrive provisionally at the relation of equation 5.20 with minimal assumptions. This equation shows how a relation between the external and internal curvature might be achieved in the present theory with non-zero values for $\mathbf{R} \neq 0$ and $F \neq 0$ closely correlated. The possibility of deriving equation 5.20 via purely geometric means without any reference to a Lagrangian formulation remains as a conjecture of the theory.

It should be further noted that only classical fields have been considered so far and it may be that, given the symmetry of the classical picture described originally in figure 2.2, a quantum field description of the theory will be required to provide the mechanism through which non-flat structures ultimately arises on the base manifold in general. This in turn relates to the concept of 'many solutions' for the geometry $G^{\mu\nu}(x)$ on the base space as will be described in chapter 11. In the meantime, given the Kaluza-Klein relation of equation 5.20 itself, a number of further equations of motion may be deduced without the need for a Lagrangian formalism. Hence these consequences are conjectured also to apply in the present theory, as we review in the following section.

5.2 Equations of Motion for Fields and Matter

In standard field theory the Lagrangian, being a scalar, provides a means to introduce arbitrary, although generally empirically motivated, symmetries into the theory with such symmetries generally preserved in the resulting equations of motion, as reviewed in section 3.5. In the Lagrangian approach the compatibility of the equations of motion with energy-momentum conservation $\partial_{\mu}T^{\mu\nu} = 0$ is ensured through the Euler-Lagrange equation if the energy-momentum tensor is defined according to equation 3.102, as an application of Noether's theorem.

In the present theory equation 5.20 emerges out of the constraint of the simple form $L(\hat{v}) = 1$ projected over the base space M_4 , in principle without the need for a Lagrangian formalism, as described in the previous section for a model based on the form $L(v_{10}) = 1$. The new theory avoids the ambiguity inherent in the choice of a scalar Lagrangian function and replaces the need to impose the principle of extremal action with a firm conceptual grounding in the physical manifestation of the full form of temporal flow $L(\hat{v}) = 1$ and its symmetries. Hence in contrast to the Lagrangian approach here we begin with $T^{\mu\nu}_{;\mu} = 0$ as a direct consequence of the definition of energy-momentum as $T^{\mu\nu} := G^{\mu\nu}$, within a conventional normalisation factor in relations such as equation 5.20, together with the contracted Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$. In the limit of vanishingly small spacetime curvature, with a linear connection $\Gamma \to 0$ in a suitable choice of coordinates, this constraint can be written as $\nabla_{\mu}T^{\mu\nu} = 0 \rightarrow \partial_{\mu}T^{\mu\nu} = 0$ and interpreted as energy-momentum conservation. The question then regards the extent to which this constraint determines the equations of motion, both in a curved spacetime and in the limit of flat Minkowski spacetime, for the entities which apparently compose $T^{\mu\nu}$, without appealing to a Lagrangian structure.

This also contrasts with a more standard approach to general relativity, reviewed in section 3.4, in which the Einstein tensor $G^{\mu\nu}$ is first equated with a generic energy-momentum tensor, $G^{\mu\nu} = -\kappa T^{\mu\nu}$ in equation 3.75, via a normalisation constant κ . In the meantime various examples of possible forms $T^{\mu\nu}$ may be postulated, or deduced from a Lagrangian method, for example for the energy-momentum of a perfect fluid or an electromagnetic field, again with appropriate normalisation factors. Only then are the Einstein tensor and the chosen form for $T^{\mu\nu}$ linked together via equation 3.75. This standard approach distances the relation between the external curvature $G^{\mu\nu}$ and internal curvature $F^{\alpha\mu\nu}$ by the insertion of the apparently mediating object $T^{\mu\nu}$, which may be considered to act as a 'source' for the gravitational field. It is this structure that motivates the form of equations 4.16 and 4.17. One of the

main reasons for considering $T^{\mu\nu}$ to be the source term in the Einstein equation is that material phenomena (such as the properties of everyday tables and chairs) are generally more readily observable than their counterparts in the warping of the spacetime geometry, particularly within the local laboratory environment.

In the present theory the more intimate relation of equation 5.20 arises *directly* from the basic conceptual ideas of the theory, as described in the previous section, with the symmetry groups of both the external and internal geometry mutually related through the unifying symmetry of the full form $L(\hat{v}) = 1$. The motivation for the righthand side of equation 5.20 to subsequently be interpreted as an energy-momentum tensor corresponding to $G^{\mu\nu}$ will be found in the empirical usefulness of such a concept. This will be more apparent when 'quantum effects' are introduced and augment the possible forms of $T^{\mu\nu}$ beyond that of continuous classical fields, as we alluded to at the end of the previous section.

Here, beginning from the unified point of view for classical fields, the external and internal curvatures appear on a similar footing in equation 5.20, with the contracted Riemann curvature on the left-hand side equated identically with terms quadratic in the internal curvature in the central expression. The great difference in the relative strengths of the respective physical forces encountered empirically in nature will later need to be accounted for through the respective interactions and couplings of the fields to be identified in the theory. These will give rise to a variety of laboratory phenomena and will lead to normalisation factors replacing χ in relations such as equation 5.20 once practical units are employed for measured quantities. While the bare mathematical relations are needed to understand the theoretical basis of the unification, for a discussion of the empirical consequences here we set $\chi = \frac{\kappa}{2}$ $rac{\kappa}{2}$ as suggested following equation 5.20 in the previous section.

The tensor $T^{\mu\nu}$ is composed of effective macroscopic quantities or as a function of fundamental fields, to be determined in the theory, which in turn mutually constrains the form of $G^{\mu\nu}$. Here the initial aim will be to demonstrate the extent to which the equations of motion for both external gravitational and internal gauge fields are implied within the unifying form of equation 5.20.

First we consider the classical field for the particular case of $U(1)$ as the internal symmetry, that is the case of electromagnetism. In terms of the components $F_{\mu\nu} =$ $\partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$ of the electromagnetic field tensor the components of the Einstein tensor $G^{\mu\nu}$ of equation 5.20, with a single generator for the internal group, can be written as:

$$
-\frac{1}{\kappa}G^{\mu\nu} = F^{\mu}_{\ \rho}F^{\rho\nu} + \frac{1}{4}g^{\mu\nu}F_{\rho\sigma}F^{\rho\sigma} \tag{5.22}
$$

Hence through this equation direct contact is made between gravitation in the form of the geometric curvature of spacetime and the familiar laboratory phenomena of the electromagnetic field. The fact that powerful electromagnetic effects may be observed for which the associated gravitational field is immeasurably small is an indication of the need to explain the origin of laboratory normalisation units, as mentioned above.

Given the tetrad field components $e^{\mu}_a(x)$ of a local orthonormal frame field ${e_a(x)}$ the components of the electromagnetic curvature tensor in a local Lorentz frame $F_{ab} = e^{\mu}{}_{a}e^{\nu}{}_{b}F_{\mu\nu}$ may be written out as the 4×4 asymmetric matrix:

$$
[F]_{ab} = \begin{pmatrix} 0 & E_1 & E_2 & E_3 \ -E_1 & 0 & -B_3 & B_2 \ -E_2 & B_3 & 0 & -B_1 \ -E_3 & -B_2 & B_1 & 0 \end{pmatrix}.
$$
 (5.23)

This is also the conventional form for the electromagnetic field tensor defined globally for the flat Minkowski spacetime of special relativity. The special symbols E_i and B_i $(i = 1, 2, 3$ with $E_i = F_{0i} = F(e_0, e_i)$ and $-\varepsilon_{ijk}B_k = F_{ij} = F(e_i, e_j)$ for the six independent components of the electromagnetic curvature 2-form F in a particular Lorentz frame $\{e_a\}$ represent the electric and magnetic fields respectively. These six components transform non-trivially under external Lorentz transformations but are trivially unchanged under an internal $g(x) \in U(1)$ gauge transformation, equation 3.40, since $g^{-1}Fg = F$ for an Abelian group.

Historically it was realised that Maxwell's equations 3.90 and 3.91 exhibit a U(1) symmetry before an understanding of gauge theories had been developed, although it was not considered to be a fundamental physical symmetry of nature since it is not a spacetime symmetry. However in the present theory fundamental symmetries are not of spacetime (in any dimension) but of multi-dimensional forms of temporal flow expressed as $L(v) = 1$. These include *both* the familiar 4-dimensional spacetime symmetry associated with perception on an extended manifold M_4 and equally the gauge symmetry groups, including the $U(1)$ of electromagnetism that arises here as will be described in section 8.2. Here both external and internal symmetries, together with their respective physical phenomena, originate naturally from the fundamental concepts of the theory.

In an approximately Minkowskian spacetime the electromagnetic field F_{ab} may be defined and measured operationally by observing the motion of a body of mass m and charge q in the field and using the Lorentz force law of equation 3.88. In that equation $F_c^b = \eta^{ba} F_{ac}$ is a mixed index form of the electromagnetic curvature tensor. The metric is needed to define this tensor, as it is for $F^{cd} = \eta^{ca} \eta^{db} F_{ab}$ and hence in turn to define the 'Hodge dual' of the electromagnetic curvature tensor:

$$
{}^{*}F_{ab} = \frac{1}{2} \varepsilon_{abcd} F^{cd} \quad \text{with} \quad [{}^{*}F]_{ab} = \begin{pmatrix} 0 & -B_{1} & -B_{2} & -B_{3} \\ B_{1} & 0 & -E_{3} & E_{2} \\ B_{2} & E_{3} & 0 & -E_{1} \\ B_{3} & -E_{2} & E_{1} & 0 \end{pmatrix} . \tag{5.24}
$$

In Minkowski spacetime $\varepsilon_{abcd} = \varepsilon_{[abcd]}$ are the components of the completely antisymmetric rank-4 tensor $\varepsilon \equiv e^0 \wedge e^1 \wedge e^2 \wedge e^3$, with $\varepsilon_{0123} = \varepsilon(e_0, e_1, e_2, e_3) = +1$ implying the choice of right-handed orientation for the orthonormal basis ${e_a}$, while the cotensor components are simply $\varepsilon^{abcd} = -\varepsilon_{abcd}$. In a general coordinate system, including the case of a curved spacetime, the metric volume form ω with components:

$$
\omega_{abcd} = \sqrt{|g|} \varepsilon_{abcd} \tag{5.25}
$$

$$
\omega^{abcd} = (1/\sqrt{|g|}) \varepsilon^{abcd} \tag{5.26}
$$

where $g(x)$ is the determinant of the metric $g_{\mu\nu}(x)$, is employed for the Hodge dual operator of equation 5.24 since ε , unlike ω , does not transform as a tensor under general coordinate transformations. The Levi-Civita symbol ε_{abcd} is equivalent to the components of the volume form ω in Minkowski spacetime with global coordinates employed such that the metric $g_{\mu\nu}(x) = \delta^a_{\mu} \delta^b_{\nu} \eta_{ab}$ everywhere.

In general on an *n*-dimensional manifold the space of p -forms has the same number of degrees of freedom as the space of $(n-p)$ -forms with a canonical isomorphism between the two sets given by the metric volume form ω . The isomorphism map is the Hodge dual of a form which contains precisely the same information reorganised into the components of the dual form. For example the map from F in equation 5.23 to $*F$ in equation 5.24 corresponds to a rearrangement of matrix components with $(E_i, B_j) \to (-B_i, E_j).$

The Einstein tensor $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu}$ is the 'trace-reversed' Ricci tensor, it can also be defined as the contraction ([6] p.325):

$$
G^{\beta}_{\gamma} := G^{\tau \beta}_{\gamma \tau}
$$

with
$$
G^{\alpha \beta}_{\gamma \delta} := \frac{1}{2} \omega^{\alpha \beta \rho \sigma} R_{\rho \sigma}^{\mu \nu} \frac{1}{2} \omega_{\mu \nu \gamma \delta}
$$

and in this sense is 'dual' to the Ricci tensor $R^{\mu\nu}$. The tensor G carries exactly the same information, and possesses the same rank-4 tensor symmetries, as the Riemann tensor \bf{R} and hence also has 20 independent components. It is analogous to the dual tensor $*F$ for the electromagnetic curvature tensor F .

The electromagnetic energy-momentum tensor identified with $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$ for equation 5.22, as guided by the Kaluza-Klein framework, is identical to that obtained in equation 3.105 in the Lagrangian formalism since effectively the same matter Lagrangian $\mathcal{L} \sim F^2$ is introduced in both cases, via equations 5.18 and 3.93 respectively. This expression can also be written in an equivalent but more symmetric form ([26] p.456):

$$
T^{\mu\nu} = \frac{1}{2} (F^{\mu}_{\ \rho} F^{\rho\nu} + {}^{*}F^{\mu}_{\ \rho} {}^{*}F^{\rho\nu}) \tag{5.27}
$$

$$
= F^{\mu}_{\ \rho} F^{\rho\nu} + \frac{1}{4} g^{\mu\nu} F_{\rho\sigma} F^{\rho\sigma} \tag{5.28}
$$

From either of these equations the energy density of the electromagnetic field is found to be $T^{00} = \frac{1}{2}$ $\frac{1}{2} (E^2 + B^2)$, as originally expressed by Maxwell. There are two Lorentz invariants of the electromagnetic field, the scalar norm $\frac{1}{2}F_{\mu\nu}F^{\mu\nu} = -(E^2 - B^2)$ and the pseudo-scalar $\frac{1}{4}F_{\mu\nu} * F^{\mu\nu} = \vec{E} \cdot \vec{B}$, although expressions of the latter kind (composing $F_{\mu\nu}$ with its dual) do not feature in $T^{\mu\nu}$. Both of these quantities are functions on the spacetime manifold which locally take the same value in any Lorentz frame and are also invariant under (orientation preserving) general coordinate transformations.

The energy-momentum tensor for the electromagnetic field is also traceless, $T^{\mu}_{\ \mu}=0$, from which the trace of the Einstein equation implies that the scalar curvature vanishes, $R = 0$, and hence in this case the Einstein equation can be written $G^{\mu\nu} =$ $R^{\mu\nu} = -\kappa T^{\mu\nu}$, as described shortly after equation 5.20 in the previous section. Hence in the Einstein-Maxwell theory while the Maxwell tensor $F_{\mu\nu}$ and its dual $*F_{\mu\nu}$ appear in a symmetric way in equation 5.27 the Einstein tensor $G^{\mu\nu}$ is *identical* to its 'dual' $R^{\mu\nu}$.

From this underlying theoretical point of view electromagnetism arises as a U(1) gauge theory with the electromagnetic field tensor being the exact 2-form $F = dA$ as defined in terms of the $U(1)$ connection 1-form $A(x)$. Hence by the exterior algebra property $d^2 = 0$ the curvature 2-form is in turn necessarily *closed* $dF = 0$ as an identity that gives immediately the homogeneous Maxwell equations summarised in equation 3.90.

With the electric current 1-form *defined* as $J := {}^*d*F$ (that is ${}^*J := d*F$ consistent with the inhomogeneous Maxwell equation 3.91) from the property $d^2 = 0$ it also follows immediately that dd[∗] $F = 0$ and we also find the identity d[∗] $J = 0$. In Minkowski spacetime this in turn implies that $\partial_a J^a = 0$ corresponding to the conservation of electric charge expressed in terms of the components of the conserved current J associated with the internal $U(1)$ symmetry. This is very closely analogous to the fact that *defining* the energy-momentum tensor to be $T^{\mu\nu} := G^{\mu\nu}$ leads immediately to the local conservation of energy-momentum $T^{\mu\nu}_{;\mu} = 0$ via the contracted Bianchi identity for the Einstein tensor $G^{\mu\nu}$. Hence Noether's theorem, based on a Lagrangian approach as described in section 3.5, is not needed to identify either of these conserved quantities, which are both purely geometric in origin.

It can be shown ([6] p.472) that for the case $J = 0$ the Einstein equation, in the form of equation 5.22, mutually constrains the evolution of both the gravitational and electromagnetic field, with the latter usually expressed by the source-free Maxwell equation $d*F = 0$, that is equation 3.91 for $J = 0$, as we review here. Applying the identity $G^{\mu\nu}_{;\mu} = 0$ to both sides of equation 5.22 gives:

$$
0 = F^{\mu\tau}{}_{;\mu}F_{\tau}{}^{\nu} + F^{\mu\tau}F_{\tau}{}^{\nu}{}_{;\mu} + \frac{1}{2}g^{\mu\nu}F_{\rho\sigma;\mu}F^{\rho\sigma}
$$

\n
$$
= F_{\tau}{}^{\nu}F^{\mu\tau}{}_{;\mu} + g^{\mu\nu}F^{\rho\sigma}F_{\sigma\mu;\rho} + \frac{1}{2}g^{\mu\nu}F_{\rho\sigma;\mu}F^{\rho\sigma}
$$

\n
$$
= F_{\tau}{}^{\nu}F^{\mu\tau}{}_{;\mu} + \frac{1}{2}g^{\mu\nu}F^{\rho\sigma}(F_{\sigma\mu;\rho} + F_{\mu\rho;\sigma} + F_{\rho\sigma;\mu})
$$

\n
$$
\Rightarrow F_{\tau}{}^{\nu}F^{\mu\tau}{}_{;\mu} = 0
$$

\n(5.29)

The final term in the penultimate equation vanishes by the identity $dF = 0$, that is the homogeneous Maxwell equation 3.90, or $F_{[\sigma\mu;\rho]} = 0$ in components (again here '; μ ' is the covariant derivative with respect to the linear connection Γ in a general curved spacetime). The remaining expression in the bottom line involves a linear combination of the four quantities $F^{\mu\tau}{}_{;\mu}$. The determinant of the coefficients F_{τ}^{ν} is the Lorentz pseudo-scalar $|F_\tau^{\ \nu}| = -(\mathbf{E} \cdot \mathbf{B})^2$ ([6] p.472). For a general electromagnetic field this quantity is non-zero, except that it may vanish on hypersurfaces, and hence in general the source-free form of the Maxwell equation 3.91 does not need to be imposed, rather it may instead be deduced from the Einstein equation for the electromagnetic field that:

$$
F^{\mu\tau}{}_{;\mu} = 0 \tag{5.30}
$$

On defining $J^{\tau} = F^{\mu\tau}{}_{;\mu}$ this result shows that vanishing current $J = 0$ is implied for the relation of equation 5.22 under the Bianchi identity $G^{\mu\nu}_{;\mu} = 0$. For this vacuum case $J = 0$ both the curvature F and its dual ^{*}F satisfy a similar equation, $dF = 0$ and $d *F = 0$ respectively, while for the external curvature there is a greater symmetry with $G^{\mu\nu}$ equal to its 'dual' $R^{\mu\nu}$, as described above.

A similar argument may be followed for the non-Abelian case, beginning with equation 5.20 and following the sequence of expressions leading to equation 5.29 except with $F_{\mu\nu} \rightarrow F^{\alpha}_{\mu\nu}$ and an extra contraction over the index α , representing the group generators, for each quadratic term in the internal curvature. Sandwiched between the two complementary constraining identities for the external and internal curvature, that is the Bianchi identities $G^{\mu\nu}_{\ \ ;\mu} = 0$ and $DF = 0$ respectively, this leads to the Yang-Mills equation $D_{\mu}F^{\alpha\mu\nu}=0$, which was derived from a Lagrangian in equation 3.95, and includes self-interaction terms for the non-Abelian gauge field $Y^{\alpha}_{\mu}(x)$. The same equation was also derived as a consequence of Kaluza-Klein theory in equation 4.18 from the stationarity of the action integral of equation 4.10 on a principle bundle. Generally for the non-Abelian case, as for the Abelian case of Maxwell's equations, a conserved current can be obtained in terms of a geometric identity.

For the present theory the Maxwell and Yang-Mills equations are also proposed to arise through a purely geometric argument, similar to that described for equation 5.29, directly from the identity $G^{\mu\nu}_{;\mu} = 0$ as applied to equation 5.20. This relation itself arose in equation 4.16-4.17 under the stationarity of an action integral in Kaluza-Klein theory, although in the previous section we described how equation 5.20 might be obtained ultimately in the present theory without any appeal to the Lagrangian formalism. Here equation 5.20 is considered to arise as a perturbation to the Einstein vacuum equations, derived for equation 3.82 in terms of the stationarity of the Einstein-Hilbert action under variations of the metric $\delta g_{\mu\nu}(x)$. Consistent with this approach the above discussion suggests that the variation of the gauge field $\delta Y^{\alpha}_{\mu}(x)$ is not needed in order to derive the vacuum Yang-Mills equation 5.21; rather, as for general relativity, only the $\delta g_{\mu\nu}(x)$ variation is needed in order to derive equation 5.20, which in turn itself implies the relation of equation 5.21 as a consequence of the geometric structure. With equation 5.20 itself conjectured to arise inevitably out of the geometric constraints implied in the breaking of the full $L(\hat{\mathbf{v}}) = 1$ symmetry over M_4 any explicit reference to the Lagrangian formalism might be avoided entirely.

In the present framework non-Abelian symmetries arise, as for the case of $U(1)$ above, within the internal symmetry action on the full form $L(\hat{\mathbf{v}}) = 1$. The symmetry breaking is pictured in figure 5.1 for the $L(v_{10}) = 1$ model, for which the internal symmetry is identified simply as $SO(6)$. Internal symmetries deriving from yet higherdimensional forms of $L(\hat{v}) = 1$ will be considered in chapters 8 and 9.

Returning to the Abelian case of electromagnetism, more generally for $J \neq 0$, in applying to the Maxwell tensor F and not to the dual tensor $*F$ the Bianchi identity $dF = 0$ introduces a clear break in the mathematical symmetry between these two tensors. This in turn is directly associated with the empirical asymmetry between the observed roles of the electric and magnetic fields. The field components (E_i, B_j) are oriented within the Maxwell tensor in equation 5.23 such that they are distinguished by the particular properties that $\nabla \cdot \mathbf{B} = 0$ while $\nabla \cdot \mathbf{E} = \sigma$, where σ is the charge density for the case of static fields . (From the historical empirical point of view the asymmetry between the expressions for dF in equation 3.90 and $d*F$ in equation 3.91 is a physical observation in the sense it 'might have been' observed that $dF = *J_M$ with a 'magnetic monopole current' J_M , however empirically such a current has never been seen.)

Here we next consider how equations of motion describing the broad macroscopic properties of matter arise. The microscopic details of fields and quantum physics which underlie these properties need not be considered in any detail here. Rather the general freedom inherent in the Einstein equation, beyond a specific form such as equation 5.22, will be opened up to a more general structure $G^{\mu\nu} = -\kappa T_{\epsilon}^{\mu\nu}$, where ϵ here denotes an effective energy-momentum tensor describing coarse macroscopic phenomena. This macroscopic form of $T_{\epsilon}^{\mu\nu}$ will include terms for the effective flow of physical matter, either charged or uncharged, as well as for the original electromagnetic field, all combinations of which will be collectively subject to $T_{\epsilon}^{\mu\nu}{}_{;\mu} = 0$ through the Einstein equation.

Under the symmetry transformations of a higher-dimensional form of temporal flow $L(\hat{\mathbf{v}}) = 1$ the projection over the base manifold M_4 , as described in the previous section, leads to a relation between classical external and internal fields culminating in a relation of the form of equation 5.20, which may be written:

$$
G^{\mu\nu} = f(Y) \tag{5.31}
$$

The identity $G^{\mu\nu}{}_{;\mu}=0$ then leads to constraints on the equations of motion for the internal gauge fields $Y(x)$, that is the Yang-Mills-Maxwell equations, as described above. A particular form for the energy-momentum tensor is identified as $T^{\mu\nu}$:= $-\frac{1}{\kappa}G^{\mu\nu}$, that is via the Einstein equation.

So far we have considered only the case in which $G^{\mu\nu}(x)$ is equated with a function of the curvature $F^{\alpha}_{\mu\nu}(x)$, in turn derived from a classical continuous gauge field $Y^{\alpha}_{\mu}(x)$, in the form of equation 5.31, which exhibits a relatively even significance for the external gravitational field on the left-hand side and the internal gauge field on the right-hand side. This structure was motivated to obtain $G^{\mu\nu}$ on the left-hand side of equation 5.20 corresponding to a global continuous external linear connection field $\Gamma(x)$ as required to define a geometric perceptual arena on the base manifold as described in section 2.2.

More generally a continuous *internal* gauge field $Y(x)$ is only a local requirement so long as the central expression of equation 5.20 can be modified in a manner compatible with the identity $G^{\mu\nu}{}_{;\mu} = 0$. With the components of the internal symmetry gauge fields $Y^{\alpha}_{\mu}(x)$ coupled with the internal temporal components, through a relation of the form of equation 2.47, only the combined effect is required to be compatible with the necessary smooth geometric structure on the left-hand side of equation 5.31 and we can write:

$$
G^{\mu\nu} = f(Y, \hat{\mathbf{v}}) \tag{5.32}
$$

implying in turn a more flexible expression for the energy-momentum tensor $T^{\mu\nu}$:= $-\frac{1}{\kappa}G^{\mu\nu}$. This extra freedom, not tied to the constraint of a continuous internal gauge field on M_4 , allows for field exchanges between the internal gauge connection $Y(x)$ and components of temporal flow $\hat{v}(x)$, which will be of the kind described in chapters 8 and 9 for more realistic forms $L(\hat{v}) = 1$ in comparison with the observations of high energy physics experiments. The possibility of multiple solutions for $G^{\mu\nu}$ involving exchanges between the field values of Y and \hat{v} will be interpreted as quantum and particle phenomena via the local indistinguishability of the field components, as will be described chapter 11.

While equation 5.31 might be expressed as $G^{\mu\nu} = -\kappa T^{\mu\nu}(Y)$ the more general non-classical extension to equation 5.32 can also be written as $G^{\mu\nu} = -\kappa T^{\mu\nu}(Y,\hat{\boldsymbol{v}})$ with the identification of the rank-2 tensor fields on either side of this expression remaining valid since both sides transform the same way and the contracted Bianchi identity will still apply to both. While $G^{\mu\nu}$ and $T^{\mu\nu}$ are identical in form they denote and possess a differing internal compositions; while the right-hand side can be interpreted as a *source* in terms of the fragmented temporal flow composed of apparent '*matter* fields', $Y(x)$ and $\hat{v}(x)$, the left-hand side represents the same mathematical object interpreted as the Einstein tensor for a linear connection describing the external geometry, as required for perception.

Equation 5.32 expresses the relation between the gravitational field described by the metric $g_{\mu\nu}(x)$ underlying $G^{\mu\nu}$ and the matter fields $Y_{\mu}(x)$ and $\hat{v}(x)$, together with the implicit interaction between these latter 'microscopic' fields themselves. Alternatively the term 'matter field' can refer to an effective macroscopic form for the energy-momentum tensor such as $T_{\epsilon}^{\mu\nu}$ averaging over the microscopic field interaction effects. We begin by looking more generally at properties of the symmetric Einstein tensor $G^{\mu\nu}$ in terms of $T_{\epsilon}^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$. A timelike eigenvector u may be defined for the energy-momentum tensor such that $([27] \text{ p.174})$:

$$
T_{\epsilon}^{\mu\nu}u_{\nu} = \rho u^{\mu} \tag{5.33}
$$

with the vector field $u(x)$ normalised as $|u| = g_{\mu\nu}u^{\mu}u^{\nu} = u^{\mu}u_{\mu} = +1$ such that $\rho = T_{\epsilon}^{\mu\nu} u_{\mu} u_{\nu}$ (= $\rho u^{\mu} u_{\mu}$) which will be identified as the effective 'proper energy density' or mass density, effectively averaging over underlying microscopic field interactions. In the general case:

$$
T_{\epsilon}^{\mu\nu} = \rho u^{\mu} u^{\nu} - S^{\mu\nu} \tag{5.34}
$$

defines the stress tensor $S^{\mu\nu}$ ([27] p.175). This is a symmetric tensor with four constraints $S^{\mu\nu}u_{\nu}=0$ (as can be seen by contracting equation 5.34 with u_{ν}) and hence with six degrees of freedom. The simplest example is that in which the effective energymomentum tensor represents a pressureless perfect fluid (such as a dust cloud) with:

$$
T_{\epsilon}^{\mu\nu} = \rho u^{\mu} u^{\nu}.
$$
\n(5.35)

In this case $G^{\mu\nu} = -\kappa \rho u^{\mu} u^{\nu}$ and we have $g_{\mu\nu}G^{\mu\nu} = -\kappa g_{\mu\nu}\rho u^{\mu}u^{\nu} = -\kappa \rho$. With $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu}$ this in turn implies $R = +\kappa\rho$ with the matter density ρ therefore directly associated with the spacetime scalar curvature R and hence with gravitational effects. The sign convention of equation 3.74, with $G^{\mu\nu} = -\kappa T^{\mu\nu}$ and positive constant κ determined in the Newtonian limit, is motivated in part by the resulting sign in the relation $R = +\kappa \rho$, that is such that *positive* scalar curvature is associated with *positive* matter density.

Applying the contracted Bianchi identity $G^{\mu\nu}{}_{;\mu}=0$ to the right-hand side of

equation 5.35 we then have $([27] \text{ p.175})$:

$$
T_{\epsilon}^{\mu\nu}{}_{;\mu} = 0 \Rightarrow (\rho u^{\mu})_{;\mu} u^{\nu} + \rho u^{\mu} (u^{\nu})_{;\mu} = 0
$$

\n
$$
\Sigma_{\nu}(\times u_{\nu}) \Rightarrow (\rho u^{\mu})_{;\mu} = 0 \text{ since } u_{\nu} (u^{\nu})_{;\mu} = 0
$$

\nhence $\rho u^{\mu} (u^{\nu})_{;\mu} = 0.$ (5.36)

Here the continuity equation $(\rho u^{\mu})_{;\mu} = 0$, describing the conservation of mass-energy, in the second line is substituted back into the first line to deduce the expression in the final line. From this we see that the form of equation 5.35, with $\rho \neq 0$, implies that $u^{\mu}(u^{\nu})_{;\mu} = 0$, that is the flow lines of the fluid are geodesics. Such a result could be derived from the simple Lagrangian of equation 3.78, with the requirement $\delta L = 0$ under variation of the path implying equation 3.77. However here in the case of a perfect fluid the geodesic law for the motion of bodies in general relativity is an inescapable consequence of the Einstein field equation and the Bianchi identity, which is a well-known result.

More generally the effective energy-momentum tensor $T_{\epsilon}^{\mu\nu}$ can describe a perfect fluid with non-zero effective pressure p in the form:

$$
-\frac{1}{\kappa}G^{\mu\nu} =: T_{\epsilon}^{\mu\nu} = (\rho + p)u^{\mu}u^{\nu} - pg^{\mu\nu}.
$$
 (5.37)

with, by comparison with equation 5.34, $S^{\mu\nu} = p(g^{\mu\nu} - u^{\mu}u^{\nu})$ which satisfies $S^{\mu\nu}u_{\nu} = 0$. The material flow u is again subject to $|u|=1$ with ρ and also now p as effective macroscopic terms irrespective of the classical or quantum fields underlying this structure. Again here the structure of matter perceived in spacetime is constrained by the geometrical properties of $G^{\mu\nu}$. Applying the Bianchi identity $G^{\mu\nu}{}_{;\mu}=0$ to the right-hand side of equation 5.37, similarly as above for equation 5.35 leading to equation 5.36, we now find that in general $u^{\mu}(u^{\nu})_{;\mu}$ is non-zero and proportional to the pressure gradient ([27] p.176), as a deviation from pure geodesic flow of the fluid due to the pressure term.

Alternatively we may consider a pressureless fluid carrying charge, that is a fluid with energy density ρ and also a charge density σ . Here we are dealing with continuous classical fields and bodies corresponding to the motions of macroscopic entities, where $T_{\epsilon}^{\mu\nu}$ may represent charged metal plates, wires and so on and $T_{\epsilon m}^{\mu\nu}$ describes a classical electromagnetic field, for example in a laboratory setting. For the original case with the classical electromagnetic gauge field only and $T_{\text{em}}^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$ from equation 5.22 consistency with $G^{\mu\nu}{}_{;\mu} = 0$ required that $J^{\nu} := F^{\mu\nu}{}_{;\mu} = 0$, as described for equation 5.30. It is then through the introduction of effective matter terms that the equations for the electromagnetic field allow for a charged current $J \neq 0$ in combination with energy-momentum in the form $T_{\epsilon}^{\mu\nu} = \rho u^{\mu} u^{\nu}$, both of which are composed in terms of the effective matter content.

We have *defined* $T^{\mu\nu} := G^{\mu\nu}$ and argued, following the previous section, that for an internal U(1) symmetry identified within the full symmetry of $L(\hat{\boldsymbol{v}}) = 1$ this naturally leads to $T_{\text{em}}^{\mu\nu}$ in the form of equation 5.22. Similarly here with $J^{\nu} := F^{\mu\nu}{}_{;\mu}$ we would like to understand the form of J that results as microscopic field transitions over M_4 are considered such that equation 5.22 breaks down giving:

$$
-\frac{1}{\kappa}G^{\mu\nu} = T^{\mu\nu}(Y,\hat{\mathbf{v}}) \neq F^{\mu}_{\ \rho}F^{\rho\nu} + \frac{1}{4}g^{\mu\nu}F_{\rho\sigma}F^{\rho\sigma} \quad \text{and} \quad J^{\nu} = F^{\mu\nu}_{\ \ ;\mu} \neq 0 \tag{5.38}
$$

with a specific form for the first equation relating to a specific form for the latter. In the phenomenological macroscopic limit the effective energy-momentum tensor $T_{\epsilon}^{\mu\nu}$ = $\rho u^{\mu} u^{\nu}$ arose as a possible form for a non-trivial $G^{\mu\nu}$ field for the external spacetime geometry. With charge density defined by $\sigma := \nabla \cdot \boldsymbol{E}$ in the electrostatic limit, under a Lorentz transformation we may associate the 4-vector $J^{\nu} = \sigma u^{\nu}$ with a charged body, such that $\sigma = u_{\nu}J^{\nu}$ is closely analogous to $\rho = u_{\mu}u_{\nu}T_{\epsilon}^{\mu\nu}$ for the matter density of a pressureless fluid. Hence in addition to the 4-momentum density ρu^{μ} the fluid carries an effective charge 4-current $J^{\nu} = \sigma u^{\nu}$, which is identified as a possible form of $F^{\mu\nu}_{;\mu}$ and with the identity $J_{\mu}^{\nu} = 0$ implying the conservation of charge. That is we consider the flow of matter to be simultaneously associated with:

$$
-\frac{1}{\kappa}G^{\mu\nu} =: T_{\epsilon}^{\mu\nu} = \rho u^{\mu}u^{\nu}
$$
\n(5.39)

$$
+F^{\mu\nu}_{\quad;\mu} =: J^{\nu} = \sigma u^{\nu} \tag{5.40}
$$

as the respective definitions of matter density ρ and charge density σ . Here the 4velocity $u(x)$ with $|u| = 1$ represents a fluid carrying both the mass and the charge. The fluid body is interpreted to be immersed in and passing through the electromagnetic field $F_{\mu\nu}$ such that the Einstein equation reads:

$$
-\frac{1}{\kappa}G^{\mu\nu} = T_{\epsilon}^{\mu\nu} = \rho u^{\mu}u^{\nu} + F^{\mu\tau}F_{\tau}{}^{\nu} + \frac{1}{4}g^{\mu\nu}F_{\rho\sigma}F^{\rho\sigma}
$$
(5.41)

That is the form of the energy-momentum tensor for the electromagnetic field from equation 5.22 has been combined with the pressureless perfect fluid term. Here the 4 velocity u of the fluid differs from the 4-velocity eigenvector U defined in $T_{\epsilon}^{\mu\nu}U_{\nu} = \rho' U^{\mu}$ by equation 5.33. With $T_{\epsilon}^{\mu\nu} = \rho' U^{\mu} U^{\nu} - S^{\mu\nu}$ from equation 5.34, the 4-velocity U represents a synthesis of the charged fluid and the electromagnetic field ([27] p.357).

Applying $T_{\epsilon}^{\mu\nu}{}_{;\mu} = 0$ the effect on the terms on the right-hand side of equation 5.41 has already been worked out separately in equations 5.36 and 5.29 respectively. Combined together we find that under the Bianchi identity equation 5.41 becomes (based on [27] p.358):

$$
T_{\epsilon}^{\mu\nu}{}_{;\mu} = 0 \Rightarrow (\rho u^{\mu})_{;\mu} u^{\nu} + \rho u^{\mu} (u^{\nu})_{;\mu} + F_{\tau}^{\nu} F^{\mu\tau}{}_{;\mu} = 0
$$

\n
$$
\Sigma_{\nu}(\times u_{\nu}) \Rightarrow (\rho u^{\mu})_{;\mu} + 0 + F_{\tau}^{\nu} F^{\mu\tau}{}_{;\mu} u_{\nu} = 0
$$

\n
$$
\Rightarrow (\rho u^{\mu})_{;\mu} + g_{\lambda\tau} F^{\lambda\nu} J^{\tau} u_{\nu} = 0
$$

\n
$$
\Rightarrow (\rho u^{\mu})_{;\mu} + F^{\lambda\nu} \sigma u_{\lambda} u_{\nu} = 0
$$

The final term in the fourth line above is asymmetric in the indices of $F^{\lambda\nu}$ while symmetric in the indices of $u_{\lambda}u_{\nu}$ and is therefore equal to zero. The same line then implies that $(\rho u^{\mu})_{;\mu} = 0$ (as for the second line of equation 5.36) which can be substituted into the first line giving:

$$
\rho u^{\mu} (u^{\nu})_{;\mu} + F_{\tau}^{\ \nu} J^{\tau} = 0. \qquad (5.42)
$$

Each term in equation 5.42 was found to be zero for the individual cases of a perfect pressureless fluid alone or an electromagnetic field alone, giving equation 5.36 for geodesic motion and Maxwell's vacuum equation 5.30 respectively. However for the combined case only the total vanishes and hence $G^{\mu\nu}{}_{;\mu} = 0$ implies that:

$$
\rho u^{\mu} (u^{\nu})_{;\mu} = +F^{\nu}_{\ \tau} J^{\tau} \tag{5.43}
$$

This is the relativistic Lorentz force law for a charged fluid in a curved spacetime, which is equivalent to the corresponding law of equation 3.87 for discrete bodies in the appropriate limit (27 p.359) as is similarly the case for the geodesic motion of equation 5.36 considered above. Again Lagrangian terms, such as those in equation 3.86, are not required.

As we described earlier for the effective energy-momentum tensor of equation 5.37 the geodesic flow of an uncharged fluid is modified by the pressure gradient. Similarly for the energy-momentum tensor of equation 5.41 for charged matter the geodesic law is modified by the presence of an electromagnetic field to a form, equation 5.43, which precisely gives the Lorentz force law of equation 3.87. This law, typically in the flat spacetime limit of equation 3.88 or the further non-relativistic limit, can be used to determine the strength of charges and electromagnetic fields in the laboratory and establish appropriate empirical normalisation factors.

The possibility of incorporating electromagnetism and the Lorentz force law within a higher-dimensional approach to general relativity is well known and dates back to Kaluza in 1921 ([11] equation 12). There it was shown that the *five*-dimensional geodesic equation automatically incorporates the Lorentz force law in 4-dimensional spacetime, in the approximation of low 5-velocity. In the present theory the internal gauge fields, such as that for electromagnetism, arise as a higher-dimensional form $L(\hat{v}) = 1$ is projected onto the base space M_4 , with charged matter arising through the interaction properties of the internal fields underlying the smooth spacetime geometry.

For the case in which there is no electromagnetic field $F_{\mu\nu} = 0$ or in which the material flow is uncharged $J^{\mu} = 0$ the geodesic flow is recovered from equation 5.43. On the other hand $J^{\nu} := F^{\mu\nu}_{;\mu} \neq 0$ represents the case for which $T^{\mu\nu}$ as a function of $F^{\mu\nu}$ only, equation 5.28, itself is not conserved, as can be seen from the inconsistency with equation 5.29, while the total $T_{\epsilon}^{\mu\nu}$ of equation 5.41, augmented to include the flow of macroscopic charged matter, is conserved. The Lorentz force law results from the consistency of this total energy-momentum tensor bound together under the requirement of $T^{\mu\nu}_{\epsilon;\mu} = 0$, which itself is a direct consequence of the definition $T^{\mu\nu} := G^{\mu\nu}$ and the Bianchi identity.

While the Bianchi identity implies $T^{\mu\nu}_{;\mu} = 0$ further conservation laws follow from further geometric identities, principally of the form $d^2 = 0$ which for example given $J^{\nu} := F^{\mu\nu}{}_{;\mu}$ implies that $J^{\mu}{}_{;\mu} = 0$, as described in the discussion following equation 5.28. This leads to conserved charges associated with the internal symmetries both for Maxwell and Yang-Mills theories. However, while the Maxwell equations with source $J \neq 0$ imply the conservation of charge, this conservation law is limited to physical entities that carry charge. This marks a fundamental difference with the consequences of the Einstein equation, which can be interpreted as $T^{\mu\nu} := G^{\mu\nu}$, in that, assuming that all fields are associated with energy-momentum defined this way, all fields are covered under the identity $G^{\mu\nu}{}_{;\mu}=0$ and in principle 'no physical entity escapes this surveillance' ([6] p.475).

In the above only the contracted Bianchi identity for the Riemann curvature tensor has been employed. Further, the Einstein equation $G^{\mu\nu} = -\kappa T^{\mu\nu}$ only directly yields certain linear combinations of the Riemann curvature tensor components. However, although the Weyl tensor, introduced before equation 3.69, is that part of the Riemann tensor which is not directly equated with matter $T^{\mu\nu}$ in the Einstein equation it is not arbitrary. Applying the full Bianchi identity $R_{\rho\sigma[\mu\nu;\tau]} = 0$ of equation 3.70 to equation 3.69, rearranging the terms and contracting once leads to ([9] p.85):

$$
C^{\rho\sigma\mu\nu}{}_{;\nu} = R^{\mu[\rho;\sigma]} + \frac{1}{6}g^{\mu[\sigma}R^{;\rho]} =: K^{\rho\sigma\mu} \tag{5.44}
$$

Hence the full Bianchi identity, which contains more information than the contracted form, can be regarded as a field equation for the Weyl tensor in which the source $K^{\rho\sigma\mu}$ is defined as a function of the Ricci tensor. This is analogous to the Maxwell equation 3.91 for the electromagnetic field, which can be written in a curved spacetime as $F^{\mu\nu}_{\ \ ;\mu} = J^{\nu}$, with the electromagnetic current J^{ν} as the source. For equation 5.44 the source $K^{\rho\sigma\mu}$ depends on $R^{\mu\nu}$ which in turn is intimately related to the matter content $T^{\mu\nu}$ through the Einstein equation, which can be written $R^{\mu\nu} = -\kappa (T^{\mu\nu} - \frac{1}{2})$ $\frac{1}{2}Tg^{\mu\nu}$) where $T = g_{\mu\nu}T^{\mu\nu}$. Hence, by substituting $T^{\mu\nu}$ into equation 5.44, the Weyl curvature at any given location on M_4 depends on the matter content elsewhere in spacetime, in a similar way that the electric and magnetic fields depend on the charges elsewhere. The Weyl tensor represents the non-flat part of the Riemann tensor in the matter vacuum, this includes the phenomena of gravity waves (in analogy with electromagnetic waves) as well as gravitational tidal forces and lensing effects. Further, since gravitational waves carry energy even in regions of spacetime where $G^{\mu\nu} = 0$ the association of $T^{\mu\nu} := G^{\mu\nu}$ with 'energy-momentum' itself has a degree of ambiguity, while being of great value for many practical applications.

In this section we have reviewed how a number of equations of motion arise out of the geometry of the Bianchi identities for the external and internal symmetries, given the relation of equation 5.20 obtained by comparison with Kaluza-Klein theory. However the equations of motion are derived we note that in order to empirically test a theory solutions of the field equations need to be determined and compared with actual observations in the world. This in turn requires the specification of initial conditions, or more general boundary conditions, in order to obtain such solutions. With care for the role of the implicit degrees of freedom of gauge and general coordinate transformations the 'initial value problem' is well posed for both classical electromagnetism and general relativity respectively. The evolution of the spacetime geometry is in principle fully obtainable from Einstein's equation and the equations of motion for the matter fields together with suitable boundary conditions.

It is generally not possible to begin with a given source term on the righthand side of the Einstein field equation $G^{\mu\nu} = -\kappa T^{\mu\nu}$ since a coordinate system is required in order to specify the components of $T^{\mu\nu}(x)$, and further the distribution of matter itself is dynamically intertwined with the spacetime geometry through which it propagates. One procedure would be to begin with arbitrary metric functions $g_{\mu\nu}(x)$ and catalogue $(g_{\mu\nu}(x), T^{\mu\nu}(x))$ pairs via equations 3.53, 3.73 and the field equation with $-\kappa T^{\mu\nu} := G^{\mu\nu} = f(g_{\mu\nu})$, in an attempt to converge upon a particular physical system.

In practice exact solutions for the metric $g_{\mu\nu}(x)$ have been found for the cases in which $T^{\mu\nu}$ represents the vacuum $(T^{\mu\nu}=0)$, a perfect fluid or the electromagnetic field (or a combination of the latter two, as described for a pressureless fluid in equation 5.41) and then only for spaces with a high degree of symmetry with a simple form of matter content. All solutions in general relativity consist of a metric description for a complete spacetime geometry, which will be relevant for the study of cosmology, while only a limited region of the manifold may be of physical interest in other cases such as the study of planetary orbits using the Schwarzschild solution, described in the following section, for example.

In summary, many of the equations of motion derived from a Lagrangian in section 3.5 have been shown to arise directly as a consequence of the identity $G^{\mu\nu}_{;\mu} = 0$ given a solution for $G^{\mu\nu}(x)$ for example in the form of equation 5.20. This latter relation itself arose as guided by Kaluza-Klein theory and equation 4.16–4.17 through the employment of a single 'Lagrangian function' on a principle bundle space. As described in the previous section in the present theory it is conjectured that the Lagrangian approach might be ultimately side-stepped entirely and that this one remaining pivotal Lagrangian, in the action of equation 5.18, may also be discarded. In principle it may always be possible to work backwards from the present theory to obtain apparent Lagrangian functions for the theory, but from the present point of view the Lagrangian method is ultimately effective due to its conformity with $G^{\mu\nu}_{\mu\nu} = 0$ through the compatibility of the Euler-Lagrange equation with the requirement $T^{\mu\nu}_{;\mu} = 0$, as described in the opening of this section.

In this section the form of the 4-current $J^{\nu} := +F^{\mu\nu}{}_{;\mu}$, in equations 3.91 and 5.40, has been taken to emerge macroscopically and does not necessarily apply for 'elementary particles'. The origin and role of 4-currents for microscopic fields of the form $j^{\mu}_{\;\alpha} = \overline{\psi}\gamma^{\mu}E_{\alpha}\psi$ in equation 3.97, as well as the Dirac equation 3.99, in the present theory will be addressed in section 11.1, in particular as exemplified by the Abelian $U(1)$ case of electromagnetism. In order to consider the properties of microscopic elementary particles (electrons, photons etc.) it will first be necessary to address the more fundamental questions concerning the quantisation of the theory and the concept of an elementary particle itself. In addition to the identity $G^{\mu\nu}_{;\mu} = 0$ the full form of temporal flow $L(\hat{\mathbf{v}}) = 1$, projected over the M_4 base space, will provide constraints on possible field interactions which are closely analogous to those provided by the Lagrangian for the Standard Model of particle physics, as will be described in chapters 8 and 9.

5.3 Spacetime Manifold and Time Dilation

Here we consider some of the geometric properties on the 4-dimensional spacetime manifold M_4 as arising in the present theory and in relation to general relativity. Here the 4-dimensional base manifold M_4 carries the four coordinate degrees of freedom of our spacetime experience of physical objects in the universe. The symmetry of the Lorentz group fits naturally on such a manifold since it acts on a 4-dimensional vector space which corresponds to the tangent space of M_4 . Selecting a 4-dimensional base space in this way is a provisional empirical input. It is empirical for the obvious reason and provisional since at this point the choice of four dimensions seems theoretically arbitrary and there remains the question of whether a base space of a different dimension could in principle be considered as a background for experience in another possible world. We shall return to this issue, and the question of the uniqueness of the theory in general, in section 13.3.

Hence the study of the Lorentz symmetry is motivated by the fact that it

contains SO(3), the rotational symmetry of the background space within which we perceive physical objects, together with its respect for temporal causality, as well as its central importance in established physical theories of the world. In conformity with the present theory $SO^+(1,3)$ is also the symmetry of a possible form of progression in time, denoted $L(v_4) = 1$ and presented explicitly in equation 5.1, over a 4-dimensional vector space.

Here we are considering the proper orthochronous Lorentz group $SO^+(1,3)$, sometimes denoted L^{\uparrow}_{+} , which is the part of the full Lorentz group that is continuously connected to the identity element. It is hence a continuous symmetry group acting on vectors in the 4-dimensional vector space $\mathbb{R}^{1,3}$, denoting the space \mathbb{R}^{4} with Minkowski metric $\eta_{ab} = \text{diag}(+1, -1, -1, -1)$, as will be reviewed in more detail in section 7.1. Elements of the Lorentz group $l \in SO^+(1,3)$ generate the symmetry transformations $\sigma_l: \nu_4 \to \nu'_4$ such that $L(\nu_4) = L(\nu'_4)$ as an invariant form of temporal flow in four dimensions. At any $x \in M_4$ on the spacetime manifold $v_4(x) \in TM_4$ is a vector in the local tangent space.

The base space M_4 itself originates out of the four dimensions of the translation symmetry of the form $L(v_4) = 1$ which is trivially invariant under $x^a \to x^a + r^a$ for the four components $v_4^a = dx^a/ds$ with $a = \{0, 1, 2, 3\}$, as described more generally in equations 2.10–2.13 of section 2.1 and in section 2.2 for the model world. Here the set of four numbers $r^a \in \mathbb{R}^4$ can be identified with an initial set of four coordinates $x^{\mu} \in \mathbb{R}^4$, with $x^{\mu} = \delta^{\mu}_{\ \ a} r^a$.

The Lorentzian structure of the vector space to which $v_4(x)$ belongs is transferred onto the tangent space of the parameter space M_4 and hence the latter acquires the properties of a 4-dimensional pseudo-Riemannian manifold. That is, since the flow $v_4(x)$ necessarily exists on the manifold, with components $v^a = dx^a/ds$, on M_4 the metric η_{ab} derives locally from the form $L(\mathbf{v}_4) = \eta_{ab}v^a v^b$, and it is described by the metric $g_{\mu\nu}$ in a general coordinate system via a tetrad field $e^a_{\mu}(x)$ as:

$$
g_{\mu\nu} = e^a{}_\mu e^b{}_\nu \eta_{ab} \tag{5.45}
$$

Hence the manifold M_4 inherits its pseudo-Riemannian structure from the Lorentz symmetry of $L(\mathbf{v}_4)$; with the SO(3) subgroup implying the possibility of a suitable 3-dimensional background space which appears to us to be of a more fundamental *a priori* existence than the objects we perceive moving through it.

For such a manifold in which there exist global coordinates such that $g_{\mu\nu}(x) =$ diag(+1, -1, -1, -1) for all $x \in M_4$, that is the constant Minkowski metric, we have the 4-dimensional spacetime of special relativity. In this case the local η_{ab} metric has been drawn out and made global through the existence of large scale coordinates with respect to which the tetrad field can be simply be expressed as $e^a_{\mu}(x) = \delta^a_{\mu}$. For such a Minkowski spacetime manifold the $SO(3)$ subgroup of $SO⁺(1, 3)$, now acting globally, provides the symmetry of the 3-dimensional space through which a physical world of objects might be perceived.

The point of view taken in this paper is that it is the nature of perception itself that implicitly requires an approximately flat background manifold, at least for the extended neighbourhood of the observer, and hence essentially inflates the local Minkowski metric into the extended spacetime arena and thus draws the Lorentz structure of the form $L(\mathbf{v}_4)$ out onto an approximately uniform background spacetime within which objects are perceived. The mathematical expression for such a spacetime structure, to be utilised by perception, arises spontaneously out of the translational symmetry of the form $L(\mathbf{v}_4)$.

While the identification of the 4-dimensional spacetime manifold through the 4-dimensional form of temporal flow $L(v_4) = 1$ will result in a flat spacetime geometry, as described for the model world in subsection 2.2.3, ultimately the base manifold M_4 will be obtained through a subset of four translational degrees of freedom breaking the symmetry of a higher-dimensional form of temporal flow $L(\mathbf{v}_n) = 1$ with $n > 4$, representing the full form $L(\hat{v}) = 1$. A specific expression of for $L(\hat{v}) = 1$ will be introduced in the following chapter, extending beyond the case of $L(v_{10}) = 1$ described in section 5.1. This results in general in a non-zero external Riemannian curvature, complemented by a non-zero internal gauge curvature, as described in sections 2.3 and 5.1. On the M_4 manifold the Lorentz form of equation 5.1, now embedded within the full form $L(\hat{v}) = 1$, locally expresses the relation between the components $v^a = dx^a/ds$ of tangent vectors in an ordered orthonormal basis of the tangent space. Such a local basis, or frame field, $\{e_a\}$ satisfies $g(e_a, e_b) = \eta_{ab}$, and with $\mathbf{v}_4(x) = v^a(x)e_a(x)$ the local relation of equation 5.1 is replaced by the looser constraint on the four components $v^a(x)$ projected onto TM_4 with:

$$
L(\mathbf{v}_4) = (v^0)^2 - (v^1)^2 - (v^2)^2 - (v^3)^2 = \eta_{ab} v^a v^b = \eta(\mathbf{v}_4, \mathbf{v}_4) = h^2.
$$
 (5.46)

with $h \in \mathbb{R}$. While local coordinates $\{x^a\}$ necessarily exist to express the form $L(\mathbf{v}_4)$ = h^2 we may also introduce an arbitrary global coordinate system $\{x^{\mu}\}$ over M_4 which naturally gives rise to a coordinate frame basis, denoted $\{\partial_{\mu}\}\$, for the tangent space at any $x \in M_4$. The coordinate frame is related to the orthonormal frame with $\partial_{\mu} = e_a e_{\mu}^a(x)$ as described in equation 3.49 and section 3.3.

In addition to the observation that in general $h^2 \neq 1$ in equation 5.46 the further consequence of the embedding in the larger form $L(\hat{v}) = 1$ is the possibility of finite Riemannian curvature $\mathbf{R} \neq 0$ as alluded to above. This implies a warping of the geometry such that global coordinates no longer exist such that $e^a_{\mu}(x) = \delta^a_{\mu}$ in general. The tetrad field $e^a_{\mu}(x)$ now describes the necessarily non-trivial relation between global and local coordinates. As described in section 3.4 the unphysical nature of general coordinates is implied under general covariance, while a tetrad field with respect to a set of coordinates \mathbb{R}^4 , as depicted in figure 3.6(a), indicates physically distinguished local orthonormal frames as utilised by the equivalence principle.

In the present theory 'general covariance' is significant since in general the Lorentz symmetry of the form $L(\mathbf{v}_4) = h^2$ cannot be expressed globally with respect to a single coordinate chart on the manifold. Without such a preferred global reference frame all arbitrary coordinate systems are equally valid for the description of the equations of physics on the manifold. In the context of this theory the metric $g_{\mu\nu}(x)$ has particular physical significance for the nature of perception and describes the geometric form through which we literally see the world, motivating its prominent role as the gravitational field; described as the 'new ether' by Einstein as discussed at the end of section 3.4.

The general *global* coordinates do not correspond to an underlying Euclidean or any other geometric structure on the manifold. However, the manifold exists as a space for the flow $v_4(x)$ of $L(v_4) = h^2$ itself and we naturally have a frame field

 ${e_a(x)}$ of local orthonormal basis vectors and local coordinates ${x^a}$ with respect to which this flow can be written with the components $v^a = dx^a/ds$, corresponding to the tangent vector components $v^{\mu} = dx^{\mu}/ds$ in a general coordinate system, and hence we necessarily have a *local* Lorentzian structure on M_4 . While in principle the torsion on such a manifold may be finite the geometry described above is compatible with the 'equivalence principle' which may hence be adopted, together with the implication of vanishing torsion, as a provisional simplifying assumption which will be discussed further in section 13.3.

With respect to a set of general coordinates $\{x^{\mu}\}$ on the M_4 manifold arbitrary vector fields $u(x)$, that is cross-sections of the tangent bundle TM_4 , can be expressed as $u^{\mu}(x)\partial_{\mu}$ with the numbers $u^{\mu} \in \mathbb{R}^{4}$ regarded as the components of a tangent vector on the 4-dimensional manifold M_4 . The situation is similar to that depicted in figure 2.4, except now for a 4-dimensional manifold. For any vector field $u(x)$ on M_4 the quantity $g(\boldsymbol{u},\boldsymbol{u})=g_{\mu\nu}(x)u^{\mu}(x)u^{\nu}(x)$ may be determined at any point $x\in M_4$ and the vector $u(x)$ described as 'timelike', 'null' or 'spacelike' according to whether this quantity is positive, zero or negative respectively. This range of possibilities is also the origin of the name 'space-time' manifold. The 'time' in 'spacetime' refers to the existence of timelike vectors and coordinates rather than explicitly to the actual pure temporal flow s which underlies the particular field $v_4(x)$ as constrained by the equation $L(\hat{v}) = 1$.

Since the Lorentzian manifold structure arises out of the flow of time the light cone geometry of the tangent space is time-orientable over the 4-dimensional volume of the spacetime manifold M_4 . That is, the time-orientation of the light cones is necessarily continuous on M_4 as determined by the directed line element field $v_4(x)$ of temporal flow itself as an extension of the original 1-dimensional progression in time. This time-directed vector field is locally $SO(3)$ invariant and provides a local $(1 + 3)$ dimensional decomposition of spacetime for all $x \in M_4$ with temporal and spatial parts identified in the local reference frames.

Choosing the local coordinate x^0 to be aligned with $v_4(x)$, with components $v^a = \left(\frac{dx^0}{ds}, 0, 0, 0\right)$, then x^0 effectively acts as a parameter for the pure values of time, that is $ds = dx^0/h$ for $L(\mathbf{v}_4) = h^2$, which is a particular case of the more general local expression described in equation 5.47 below. Three spacelike local coordinates x^1, x^2 and x^3 can also be constructed orthogonal to each other and to x^0 with respect to η_{ab} , with local spatial frames related via the SO(3) subgroup.

Whereas embedding the perceptual background of an effective 3-dimensional space and 1-dimensional time within the symmetry structures of the mathematical form $L(\mathbf{v}_4) = 1$ led to their incorporation into the 4-dimensional Minkowski spacetime of special relativity, that is with zero Riemannian curvature, extracting the same base manifold out of a higher-dimensional form of temporal flow $L(\hat{v}) = 1$ results in a more flexible and dynamic 4-dimensional spacetime structure as employed in general relativity. With M_4 itself still originating out of a 4-dimensional translational symmetry of $L(\hat{\boldsymbol{v}}) = 1$, even for the generalisation in which the external geometry is expressed in terms of underlying interacting fields as implied equation 5.32, the Minkowski metric η_{ab} implicit in the form $L(\mathbf{v}_4) = h^2$ is sewn into the local tangent space structure everywhere on the base manifold. This defines a possible metric structure $g_{\mu\nu}(x)$ on M_4 associated in a one-to-one manner with the existence of an $SO^+(1,3)$ orthonormal frame bundle OM_4 within the canonical $GL(4,\mathbb{R})$ general frame bundle FM_4 over the base manifold, as described in section 3.3.

With the external geometry related to the internal geometry via equation 5.20, or more generally with equation 5.31 augmented to equation 5.32, in principle the metric itself might be obtained by adopting the Levi-Civita linear connection on M_4 . The connection is metric compatible, since it derives from the local $SO^+(1,3)$ symmetry of the form $L(\mathbf{v}_4) = h^2$, and assumed to be torsion-free as described above. Hence as for general relativity the metric itself may be extracted by solving the second order differential equation $G^{\mu\nu} = -\kappa T^{\mu\nu}$ given a form for the energy-momentum tensor $T^{\mu\nu}$ under appropriate boundary conditions, as described towards the end of the previous section. An example is given in equation 5.49 below.

The tetrad field $e^a_{\mu}(x)$ with 16 independent components carries two kinds of information. The 10 degrees of freedom of the symmetric metric field $g_{\mu\nu} = e^a_{\mu} e^b_{\nu} \eta_{ab}$ correspond to the gravitational field for the torsion-free metric connection in general relativity, and hence the tetrad field itself can be considered to represent the gravitational field. The remaining 6 degrees of freedom correspond to the local choice of Lorentz frames implicit in $e^a_{\mu}(x)$. This local symmetry provides a link with the framework of local gauge theories as well as with the application of the spinor representations of the Lorentz group, as also alluded to towards the end of section 3.4, which are important in particle physics as will be described in chapter 7.

Here we consider the physical significance of a non-flat Riemannian geometry, described by the metric field $g_{\mu\nu}(x)$, in particular on the *relative* passage of time itself. We also consider the relation of the original pure temporal flow s with the proper time τ which may be recorded by physical objects such as clocks in the material flow of the world.

The underlying pure temporal flow s, subject to the full form $L(\hat{v}) = 1$, exists everywhere on the base manifold M_4 . The $v_4 \subset \hat{v}$ projection onto the tangent space TM_4 to the base manifold is a timelike vector, as is the tangent to any world line on M_4 , with components $v_4^a = dx^a/ds$ restricted under $L(\hat{\boldsymbol{v}}) = 1$ such that:

$$
|\mathbf{v}_4|^2 = \eta_{ab} v^a v^b = h^2 \quad \text{implying} \quad ds^2 = \frac{\eta_{ab}}{h^2} dx^a dx^b \tag{5.47}
$$

However gravitational time dilation will not be directly observed from the perspective of the microscopic flow v_4 . Indeed the underlying pure temporal flow s is not *measured* directly by physical instruments. Rather it is through the structure and symmetries of the form $L(\hat{v}) = 1$ that the physical world emerges on M_4 through relations such as equation 5.20, and with more general expressions for the apparent energy-momentum tensor as implied in equation 5.32. This more general apparent material world may be described empirically in part by the effective energy-momentum tensor $T_{\epsilon}^{\mu\nu} = \rho u^{\mu} u^{\nu}$, as introduced in equation 5.35 of the previous section and leading to the geodesic equation 5.36, where $\rho(x)$ is the matter density and the 4-velocity $u^{\mu}(x) = dx^{\mu}/d\tau$ is defined as the tangent vector at $x \in M_4$ to the world line of the physical body, which may be an element of a pressureless fluid. It is through the motion of physical bodies, such as the hands of a mechanical clock, that time dilation effects may be observed. With the *proper time* τ parametrising the motion of the body for a general coordinate system $\{x^{\mu}\}$ on M_4 we have:

$$
g_{\mu\nu}u^{\mu}u^{\nu} = 1 \quad \text{and with} \quad d\tau^2 = g_{\mu\nu} dx^{\mu} dx^{\nu} \tag{5.48}
$$
identifying an *interval* of proper time $d\tau$. These expressions are invariant under general coordinate transformations. The normalisation for the components of the metric $g_{\mu\nu}(x)$ will depend on the choice of *empirical* units adopted, for example seconds and metres for temporal and spatial dimensions, in recording the motions of the parts of a physical 'clock'.

The local orthonormal coordinates $\{x^a\}$ constructed empirically for the macroscopic proper time interval with $d\tau^2 = \eta_{ab} dx^a dx^b$ will in general not be identical to those of equation 5.47 arising directly out of the mathematical properties of the pure form of temporal flow $L(\hat{v}) = 1$. However with the physical world unfolding through the progression of the fundamental time parameter, and with s and τ represented by the 4-vectors v_4 and u in TM_4 respectively, both temporal parameters are subject to time dilation effects in the same way. The proper time τ , in 4-dimensional spacetime, is implicitly linearly proportional to the pure underlying temporal flow s , which may be expressed in any number of dimensions. This proportionality is expressed through the fixed parameter γ in equation 13.3 in section 13.1 where the relationship between τ and s is further explored.

Hence along a shared world line the fundamental time interval ds is related to the proper time interval $d\tau$ by a constant scaling and the two temporal parameters are equivalent in this sense – that is, within a fixed normalisation factor physical clocks do measure the progression of pure time s. As described in the introductory chapter, and to be expanded in chapter 14, the fundamental underlying mathematical time s is ultimately identified with 'experienced' time, while proper time τ is associated with measurable empirical phenomena, which include for example 'physical brain processes'. Hence these subjective and objective temporal phenomena, which might be exemplified by an observer located within the same inertial frame as a physical clock, are intimately connected. We next consider a particular example of time dilation effects.

The physical manifestation of the metric $g_{\mu\nu}(x)$ in a general coordinate system on M_4 resides in observable *relative* temporal and spatial distortion effects at different locations on the manifold itself. For example the Schwarzschild solution for the metric of a spatially spherically symmetric geometry around a single massive body of mass M is given by the line element:

$$
d\tau^2 = \left(1 - \frac{2G_N M}{r}\right)dt^2 - \left(1 - \frac{2G_N M}{r}\right)^{-1}dr^2 - r^2d\theta^2 - r^2\sin^2\theta d\phi^2 \tag{5.49}
$$

in the 4-dimensional, spatially spherical polar, coordinates $\{t, r, \theta, \phi\}$, where G_N is Newton's gravitational constant. In addition to the assumption of a spatially spherically symmetric metric this solution is obtained by imposing the boundary condition that $g_{\mu\nu}(x)$ approaches the flat Minkowski limit as $r \to \infty$ spatially. This limit can be seen explicitly on taking $r \to \infty$ in equation 5.49 (this example is closely analogous to the case of the Coulomb field for a central electric charge).

The coordinate r parametrises, but does not determine, radial distances. This is consistent with the arbitrary nature of coordinates and all coordinate systems in general, as described in section 3.4 and figure 3.6. The actual radial distance, for given parameters $\{t, \theta, \phi\}$, is measured by the integral of intervals $dR = (1 - 2G_N M/r)^{-1/2} dr$. Similarly a clock at a fixed coordinate location in space records the proper time τ elapsed along its world line through the intervals:

$$
d\tau = \left(1 - \frac{2G_N M}{r}\right)^{\frac{1}{2}} dt \tag{5.50}
$$

relative to the time $\tau_{r\to\infty} = t$ measured by a clock in the flat spacetime limit at $r \to \infty$, and is a function of radial distance from the central mass, as parametrised by the coordinate r. While at any location $x \in M_4$ it is possible to choose *local* inertial coordinates $\{x^a\}$, for which $g_{\mu\nu}(x) = \text{diag}(1, -1, -1, -1)$, the absence of such a global frame for non-zero mass $M > 0$ leads to a relative time dilation effect recorded by clocks at differing radial distances from the central massive object.

As described above this dilation effect applies for the fundamental temporal flow s in exactly the same way as for the proper time τ . Hence with the interval $ds \equiv d\tau$ the same metric $g_{\mu\nu}(x)$ represents the relative temporal dilation on M_4 for the fundamental flow of time s . An observer, named 'twin A ', accompanied by a clock measuring the physical temporal flow τ_A carries an equivalent universal time parameter s_A through which the entire universe unfolds through the realisation and symmetry breaking of the full form of temporal flow $L(\hat{v}_A) = 1$, deriving from s_A as described for equation 2.9. A second observer, 'twin B ', at a separate spacetime location carries a second personal temporal parameter s_B through which B perceives the same universe to unfold through the form $L(\hat{v}_B) = 1$ in a mutually consistent way. This 'dovetailing' of the 'temporalisation' experienced by twins A and B as manifested in the same physical world will be described further in section 14.2 in the discussion of figure 14.7.

The same metric solution $g_{\mu\nu}(x)$ for the single consistent universe, expressed in a particular coordinate system (or equivalently a particular metric expression of a given geometry in terms of a unique set of coordinates \mathbb{R}^4 , adopting the perspective of figure 3.6(a)), provides the relation between the intervals ds_A and ds_B and the equivalent gravitational temporal dilation effect observed between $d\tau_A$ and $d\tau_B$ measured by the clocks of twin A and twin B respectively. The time dilation effect is determined by the empirically constructed metric $g_{\mu\nu}(x)$ in the coordinate system $\{x^{\mu}\}$ since it implicitly determines local inertial coordinates $\{x^a\}$ which are related to those of equation 5.47 by a constant scale factor (again, as will be discussed further near the opening of section 13.1 and alongside equation 13.3). In turn the local coordinates of equation 5.47 directly parametrise the fundamental temporal flow s, within a factor of h^2 , via the projection of the form $L(\mathbf{v}_4)$ onto the tangent space of M_4 .

So far we have implicitly considered only the case of constant $h(x)$ in equation 5.47. In this case all geometric time dilation effects can be considered as having a 'source' in the right-hand side of the Einstein equation 3.75 in terms of the apparent energy-momentum $T^{\mu\nu}(x)$ of ordinary matter. This is the case for the Schwarzschild solution of equation 5.49 for a central massive body. On the other hand possible variations in the magnitude of $h(x)$ in equation 5.47 will act as conformal transformations of the geometry the possible consequences of which will be considered in section 13.1, initially alongside figure 13.1.

5.4 Beyond Kaluza-Klein Theory

For Kaluza-Klein theory, originating as a pure higher-dimensional spacetime extension of general relativity, to be interpreted as a unified theory of gravitation and gauge fields in a 4-dimensional spacetime the symmetry group of general coordinate transformations in the extended spacetime has to be broken down to 4-dimensional general covariance together with the local gauge symmetry. This is equivalent to placing restrictions on the metric of the extended space which then possesses a set of isometries described by Killing vector fields which have a one-to-one relationship with the leftinvariant vector fields on the manifold of an apparent gauge group G. In this way a principle fibre bundle structure emerges on the extended space, exhibiting symmetries such that the freedom in variation of the metric \ddot{g}_{ij} , as expressed in a direct product basis in equation 4.5, is effectively reduced to the components g_{ac} and ω^{α}_{a} . The construction of an action integral on the bundle space then leads to corresponding equations of motion such as those of equations 4.16–4.18. A dynamical mechanism for this process in which an extended 4-dimensional base manifold M_4 of general relativity survives while the extra dimensions lose any sense of external spatial significance, sometimes called 'spontaneous compactification', then remains to be specified, as alluded to in section 5.1. That is, the origin of the above restrictions on the metric for the full space remains to be accounted for.

The Kaluza-Klein models, reviewed in chapter 4, contrast with the idea presented in this paper since here the 'extra dimensions', beyond four, are not required to satisfy an explicitly geometric, or spacetime, symmetry. In turn for the present theory there is no need to explain such a 'compactification', rather the base manifold M_4 is the *only* physically extended manifold to consider as it emerges as a background arena for perception through the translational symmetry of the full form of temporal flow $L(\hat{\mathbf{v}}) = 1$. In section 5.1 we presented these ideas as a *mathematical* possibility taking as an example the SO⁺(1,9) symmetry of $L(\mathbf{v}_{10}) = 1$ projected over M_4 , but the significant conceptual question concerning why this situation should be found in nature also needs to be addressed. We review here the conceptual motivation that led to this framework in the context of this provisional $SO^+(1,9)$ model world.

Out of the purely *algebraic* symmetries of $L(v_{10}) = 1$ the possibility of a local so⁺(1,3)-valued connection 1-form gives *geometric* meaning to M_4 as being not just a numerical parameter space for translational degress of freedom but rather implicitly possessing a Riemannian structure with local metric $g_{ac}(x)$ as an arena for the perception of physical objects in *time* and *space*. The identification of an extended base space is possible since there is a 'spacetime' symmetry as a subgroup of the full symmetry of $L(\hat{v}) = 1$ which acts on the local tangent space of M_4 . This innate possibility of such an interpretation is sufficient for such structures to 'freeze out' from the full symmetry of $L(\hat{v}) = 1$ as a kind of 'gestalt' through which by necessity the physical world is created and perceived.

Given this geometrical realisation of the perceptual 'external' symmetry on the base manifold, out of the full symmetry there remain 'internal' residual gauge fields and surplus temporal components which will collectively contribute to the apparent 'matter' content of the world through which the properties of physical entities will be perceived and identified on the base space. The symmetry of $L(\mathbf{v}_{10}) = 1$ is broken in the identification of the extended M_4 parameter space, with a $\overline{v}_4 \subset v_{10}$ component of the temporal flow projected onto the tangent space TM_4 as depicted in figure 5.1(b). Since the \bar{v}_4 components are distinguished in this way from the residual internal part $v_6 \subset v_{10}$ the full symmetry of the original $SO^+(1, 9)$ action on v_{10} is lost. The surviving symmetry, as gauge freedom over M_4 , is resolved into two pieces with corresponding connection 1-forms identified for both the external and internal spaces.

The combination of the general flow of time, expressed as $L(v_{10}) = 1$, with the implied symmetry properties and canonical mathematical structures existing for these objects, together with the conceptual need for a perceptual base for observation in a world, all taken collectively, has resulted in the identification of a background manifold. The full symmetry of the temporal flow $L(\mathbf{v}_{10}) = 1$ has been 'sacrificed' in the creation of the non-trivial external and internal geometrical entities, but remains as a 'ghostly' presence through which these entities are related. This correlation between the external and internal curvature tensors $\overline{R} \neq 0$ and $F \neq 0$ (while both can be zero together) was described originally for the $SO(5)$ model over M_3 in section 2.3 and for the $SO^+(1,9)$ model over M_4 , in the light of Kaluza-Klein theory, in section 5.1. This latter structure will also apply to the full symmetry action considered for the real world from the following chapter.

The use of geometrical pictures, such as those of figure 5.1, as a visual aid to understanding mathematical structures comes very naturally when the space pictured represents the way we actually perceive those structures in the world. However, the underlying properties of a purely *mathematical* space, such as those demanded here by the concept of the symmetry of time, need to be worked out within the appropriate algebraic rules, which are not necessarily visualisable even by analogy with lowerdimensional structures. Hence while possibly serving as a guide a reliance on such geometric pictures is ultimately likely to prove misleading. This in particular will be the case in the following chapter in which the internal dimensions will no longer have a spatial interpretation (*unlike* the case for the 6-dimensional space of vectors $\underline{v}_6 \in \mathbb{R}^6$ with an internal SO(6) rotational symmetry for the $SO^+(1,9)$ model described above).

On the other hand it can be asked what the perceived part of the mathematics actually looks like, and geometric pictures only really make sense in terms of a literal interpretation in this context. Perception is our window into the world of mathematical forms. It is a window which is both opened up and limited through the possibility of the internal mathematical relations which frame our experiences in a 4-dimensional spacetime. It is also part of the difficulty in theorising beyond the 4-dimensional world of general relativity, for which visualisation is a key tool.

In conclusion then, here a spacetime geometric symmetry is only required to exist on the base manifold, hence in four dimensions for our world. It is also required to be an approximately global symmetry, such that the base manifold may be identified as a suitable arena for perception in the world, at least for extended regions on the scale of everyday observations although not necessarily on the larger scales considered in cosmology.

In Kaluza-Klein theory, as described in chapter 4, while a unified framework is provided for gravity and gauge boson fields, equations 4.16–4.18, there is no energymomentum tensor for fermion fields – that is the matter fields for the leptons and quarks of our world are absent. These fields may be added by hand as sections of fibre bundles over M_4 , associated to the principle bundle P , transforming as spinors

under the external $SO^+(1,3)$ symmetry and in representation multiplets of the internal gauge symmetry group. Coupling between the gauge fields and fermions may then be introduced through interaction terms, also added by hand for example via 'minimal coupling' involving covariant derivatives, in the Lagrangian constructed for the theory.

A more mathematically self-contained approach is through a supersymmetric extension of the Kaluza-Klein framework (see for example [15] section VI, [28], [29] and [30] sections 1 and 2). Fermions may be included for example through generalising the gauge group G of the principle bundle to a 'supergroup' by augmenting the Lie algebra $L(G)$ into a 'graded' Lie algebra. Here the rule for multiplication in the Lie algebra by commutation of elements, as exemplified in equation 2.22, is extended algebraically to include anticommutation which can be used to accommodate the properties of fermion fields. The Einstein-Yang-Mills theory may be extracted as the purely bosonic sector of such extended supergravity theories.

Of the many formulations of supergravity the most attractive model involves a single supersymmetry generator, $N = 1$, so that each Standard Model particle has a single superparticle partner forming a supersymmetric doublet, and is constructed in an 11-dimensional spacetime, that is ' $d = 11$ '. The pairing of bosons with fermions through supersymmetry also tends to naturally lead to the attainment of finite calculations in the corresponding quantum field theory. However, even for the most favourable version in 11-dimensional spacetime a fully renormalisable version of supergravity has not been realised ([26] p.880). Further generalisation of supergravity to a superstring theory, obtaining a finite theory of quantum gravity by modifying QFT at the Planck scale, addresses some of the technical difficulties.

Through this geometrisation of matter in the spirit of Kaluza-Klein models based on higher dimensions of spacetime, extended to the supersymmetric theories of 11-dimensional supergravity and 10-dimensional superstrings, the aim is to incorporate the degrees of freedom of the full set of Standard Model gauge interactions within the geometry of the 7 or 6 extra spatial dimensions. In some cases the extra dimensions are considered to be small and topologically compactified while in other models our own universe may be conceived as a 4-dimensional brane-world embedded as a 4 dimensional hypersurface within the higher-dimensional spacetime bulk (see [31] for a simpler case with a 5-dimensional bulk).

Einstein's theory of gravitation based on a metric tensor in 4-dimensional spacetime hence stimulated a chain of extensions and generalisations that we have briefly reviewed above and summarise below in table 5.2.

One of the attractions of using a symmetry of extra spatial dimensions, as well as its intuitive appeal as an extension of 4-dimensional spacetime geometry, is that it limits the set of possible higher symmetries and mathematical structures to consider. In this paper instead of considering arbitrary symmetries, general geometric symmetries or specifically the symmetry of a spacetime in higher dimensions we consider general symmetries of pure time alone, as expressed through the relation $L(\mathbf{v}) = 1$ and described in chapter 2. This also greatly limits the choice of symmetry groups and their representations. As well as naturally extending to general higher-dimensional mathematical forms of the progression of time $L(\mathbf{v}) = 1$, at the same time we retain the significance of the $(1 + 3)$ -dimensional metrical manifold as a form of observation in the world as having a necessary and a priori nature.

Theoretical Framework	Physical Scope
General Relativity	Gravitation
Kaluza-Klein in 5-dimensions	Electromagnetism
Non-Abelian Kaluza-Klein Theory	Non-Abelian Gauge Fields
(with non-Levi-Civita Γ on P)	(avoid large Cosmological term)
$(\text{with } G \text{ acting on homogeneous fibres})$	(keep full $L(G)$ -valued theory)
Supergravity	Fermions as well as Bosons
Superstrings	Finite Quantum Gravity

Table 5.2: A series of increasingly general frameworks is listed in the first column with their *cumulative* extent of application listed in the second column for the nonparenthetical entries. The means of including fermions and quantum theory within the present framework will be described in section 8.1 and chapter 11 respectively.

The important point here is that the symmetry of the space part of spacetime, such as that of the SO(3) subgroup of the Lorentz symmetry $SO^+(1,3)$ central to general relativity, can be experienced in a different, geometrical, way compared with other higher symmetries of $L(v) = 1$. It may be that higher symmetries, such as $SO^+(1, 9)$, could be interpreted in a geometrical way, but this feature is relatively incidental in comparison with the fundamental requirement that it must describe a symmetry of time.

However, through investigating possible symmetries of time a significant example is identified for the symmetry group $SL(2, \mathbb{O})$ acting on the 10-dimensional space $h_2\mathbb{O}$, constructed in terms of the octonion algebra as described in the following chapter and in particular section 6.3. With $SL(2,\mathbb{O})$ being the covering group of the 10-dimensional Lorentzian symmetry $SO⁺(1, 9)$ this structure will naturally correlate with some of the properties of models based on extra spatial dimensions for which 10-dimensional spacetime is significant. Further, the 16-dimensional Majorana-Weyl spinor representation of the 10-dimensional Lorentz group, highlighted in table 7.1 of section 7.3 and here represented by the θ components appearing in the extension to the space h_3 ^O introduced in equation 6.26 of section 6.4 and described near the opening of section 8.1, is significant in various branches of string theory.

In the present theory by exploring the physical interpretation of the higherdimensional forms of $L(v) = 1$, together with the associated isochronal symmetry groups, expressed over a base space M_4 , contact is made with the series of generalisations mid-way down table 5.2, with the items listed parenthetically, with a framework very similar to non-Abelian Kaluza-Klein theories. The use of a non-Levi-Civita Ginvariant linear connection Γ such as described for equation 5.13 defined on a principle fibre bundle P , or on a bundle of homogeneous fibres E , makes a significant area of contact with the corresponding literature (including $[13, 14, 15, 16]$, $[17, 18, 19, 20]$, [22, 23, 24], [25]). From this point we then immediately diverge away from the progression towards supersymmetry and string theory in table 5.2 and in this context we shall need to explain how mathematical structures identified in the present theory correspond to the inclusion of fermion states as well as the physical concepts of quantum and particle phenomena in general.

Here we describe how field interactions arise in the context of the $SO^+(1,9)$ model. Returning to the bundle space $\underline{P} \equiv M_4 \times SO(6)$ under the breaking of the full $SO^+(1,9)$ symmetry of the model world, the local $SO^+(1,3) \subset SO^+(1,9)$ symmetry acting on the tangent space TM₄ is associated with the linear connection 1-form $\Gamma(x)$ on M_4 , which is central to the theory of general relativity and is subject to the Bianchi identity $D\overline{R} = 0$, while the so(6)-valued connection 1-form on \underline{P} is interpreted as the gauge field $Y(x)$ on M_4 , central to the gauge theory arising from the internal symmetry, and is subject to the Bianchi identity $D\underline{F} = 0$. The structures of the external and internal geometry are correlated and the corresponding equations of motion constrained as described in sections 5.1 and 5.2, with self-interactions arising for the gauge fields for the non-Abelian internal symmetry.

Further dynamical equations of motion will arise out of the full 10-dimensional temporal flow in the broken form of $D_{\mu}L(v_{10}) = 0$, by a direct generalisation of equations 2.46 and 2.47 from the $SO(5)$ model. For the $SO^+(1, 9)$ model the symmetry breaking leads to interactions between the gauge field $Y(x)$ and the internal degrees of freedom deriving from the components of $\underline{v}_6 \subset v_{10}$. That is, in comparison with equation 2.47, we have:

$$
D_{\mu}L(\mathbf{v}_{10}) = 0 \Rightarrow \mathbf{v}_{10} \cdot \partial_{\mu}\mathbf{v}_{10} + \overline{\mathbf{v}}_4 \cdot A_{\mu}\overline{\mathbf{v}}_4 + \underline{\mathbf{v}}_6 \cdot Y_{\mu}\underline{\mathbf{v}}_6 = 0 \tag{5.51}
$$

where $A_\mu(x)$ is the external Lorentz connection on M_4 . Through the interactions between the internal fields $Y_{\mu}(x)$ and $\underline{v}_6(x)$ the apparent matter content of the world on the base manifold arises, together with its quantum properties, as outlined for equation 5.32 and alluded to near the opening of this section.

As described in section 5.1 the principle bundle $\underline{P} \equiv M_4 \times SO(6)$ is not considered here to represent a physical space or spacetime, and neither is the associated bundle with homogeneous fibres. In the absence of a structure of extra spatial dimensions in general the full form of purely *temporal* flow $L(\hat{v}) = 1$ is not required to be associated with a metric geometry. The question then concerns the mathematical structure of the higher-dimensional forms of $L(\mathbf{v}) = 1$ of relevance for the physical world. In the following chapter a particular 27-dimensional form $L(\mathbf{v}_{27}) = 1$ together with its full symmetry group $G = \mathbb{E}_6$ will be introduced.

Given the extra dimensions of the full vector object $v_{27} \in h_3 \mathbb{O}$ the need to identify a Riemannian curvature parametrised over a locally approximately flat 4 dimensional base manifold M_4 breaks the full E_6 symmetry. The geometry on M_4 drawn out of the underlying structures and symmetries implied in the form $L(v_{27}) = 1$ can be described generically by the 4-dimensional relation $-\kappa T^{\mu\nu} := G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ of equation 5.32, with $G^{\mu\nu}_{;\mu} = 0$, shaping the perceptual background of our observable world. The external symmetry, acting on the extended manifold M_4 itself, is a priori essential for perception in the world as geometrically described by the linear connection and Riemannian curvature which are smoothly dependent upon $x \in M_4$. For the present theory this natural and necessary mechanism of symmetry breaking over the manifold M_4 forms a significant part of the conceptual framework through which the mathematical structures are realised in the physical world.

For the full theory based on the action of E_6 on $h_3\mathbb{O}$ the internal coupling in the final term of equation 5.51 will be replaced by an interaction between internal gauge fields and fermion fields, where the latter are identified in the internal components of $v_{27} \in h_3 \mathbb{O}$ under the action of the external symmetry on $L(v_{27}) = 1$ as will be explained in section 8.1. Hence a particular form of extra dimensions can be identified for the present theory which ultimately provides the source for the interacting gauge $Y(x)$ and fermion $\psi(x)$ fields, each of which transforms in the appropriate way under the Lorentz symmetry on 4-dimensional spacetime, underlying the matter and particle effects observed in the real world. A 'supersymmetry' is not required in order to introduce gauge fields alongside fermions fields, together with their mutual interactions, in the unified theory presented here.

Having identified fermion states the question remains concerning the origin of more specific structures of the Standard Model of particle physics, as implemented through Lagrangian terms in the form of equation 3.96 for example and as reviewed more generally in chapter 7. The origin of a series of Standard Model properties in the context of the present theory through the breaking of the full form $L(\hat{\mathbf{v}}) = 1$ will be presented in chapters 8 and 9. The constraints implied in the full form $L(\hat{\mathbf{v}}) = 1$ augment the surveillance of the external geometry with $G^{\mu\nu}_{;\mu} = 0$, described at the end of section 5.2, and the need to postulate any form of Lagrangian approach will recede further, implying ultimately that it may be avoided entirely. The full collection of constraints will also be utilised in order to address the origin of quantum phenomena for the present theory in chapter 11.

In the meantime, before considering the empirical implications for observed laboratory phenomena, in the following chapter we leave the model worlds behind and motivate consideration of E_6 as the symmetry group acting upon $L(\mathbf{v}_{27}) = 1$ as a natural higher-dimensional form of temporal flow.

Chapter 6

E_6 Symmetry on h_3 ^O

6.1 Early Formulations

In order to determine the physical effects, observable on the base manifold, of more general morphisms of the flow of time through a higher-dimensional form we shall need to consider a suitable larger symmetry group acting on an appropriate higher-dimensional vector space. The motivation leading to the identity $L(v) = 1$ of equation 2.9 as the general mathematical form acting on the n real number components of temporal flow v in an *n*-dimensional vector space was described in chapter 2. We are particularly interested here in finding such an expression with n somewhat larger than four (since the case of the external Lorentz symmetry of the form $L(v_4)$ in equation 5.46 corresponds to $n = 4$) and with a significant degree of symmetry. The vector space $h_3\mathbb{O}$ is the set of 3×3 Hermitian matrices over the octonions [1] with elements:

$$
\mathcal{X} = \begin{pmatrix} p & \bar{a} & c \\ a & m & \bar{b} \\ \bar{c} & b & n \end{pmatrix} \in \mathbf{h}_3 \mathbb{O}
$$
 (6.1)

with $p, m, n \in \mathbb{R}$ (here the component labels are chosen to conform with the notation in the relevant references, and n here is of course not the dimension of any space), $a, b, c \in \mathbb{O}$ and \bar{a} denotes the octonion conjugate of a reversing the sign of the 7dimensional imaginary part (the octonion algebra is described in the following section). Hence the vector space $h_3\mathbb{O}$ is 27-dimensional over the real numbers. It is a space with particularly rich symmetry properties largely owing to the nature of the 8-dimensional octonion subspaces.

The dimensions of the vector and spinor representations of the rotation group $SO(n)$ converge in the case of $n = 8$. That is, as well as the 8-dimensional vector representation of SO(8) the 16-dimensional spinor representation reduces to two distinct 8-dimensional spinor spaces, dual to each other. The three 8-dimensional spaces undergo different $SO(8)$ transformations, however mappings may be defined which interchange the transformation behaviour between the three spaces, with a two-to-one map from a spinor to the vector representation. The existence of such maps is due to a property known as the 'principle of triality' [32, 33] and it is unique to spaces of eight dimensions.

Three such 8-dimensional spaces can be represented by three copies of the octonions, in particular under an appropriate $SO(8)$ symmetry operation on the space $h_3\mathbb{O}$ in equation 6.1, as will be described later around equation 6.50. While the actions on the vector and two spinor representations differ for particular SO(8) transformations, collectively as three sets of transformation actions they are isomorphic by triality and it is a matter of convention which octonion space is assigned as the vector or spinor of either kind. Further, a 14-dimensional subgroup of the rotation group $SO(8)$ (itself 28-dimensional) acts on the three octonion spaces in exactly the same way. This is G₂, the automorphism group of the octonion algebra. In fact $G_2 \subset SO(7)$ as the automorphisms only act upon the seven imaginary units of the octonions. (In general an algebra automorphism Φ acts on any two elements a, b of the algebra such that $\Phi(a+b) = \Phi(a) + \Phi(b)$ and $\Phi(ab) = \Phi(a)\Phi(b)$, with the order of the latter product being reversed in the case of an algebra anti-automorphism such as the map $a \to \bar{a}$ of equation 6.7 described in the following section).

The elements of the vector space $h_3\mathbb{O}$ belong to a Jordan algebra for which the algebra product is given by:

$$
\mathcal{X} \circ \mathcal{Y} = \frac{1}{2} (\mathcal{X}\mathcal{Y} + \mathcal{Y}\mathcal{X}) \tag{6.2}
$$

with $\mathcal{X}, \mathcal{Y} \in \mathrm{h}_3\mathbb{O}$ and where $\mathcal{X}\mathcal{Y}$ is the ordinary multiplication of the 3×3 matrices, with the order of matrix components in products matching the order of the matrices in the product, since in general the components may not commute. The Jordan product can also be defined in terms of the triality mappings ([1] $p.30$). This h₃^O algebra is known as the exceptional Jordan algebra since it cannot be expressed in terms of matrices with associative elements (such as real or complex numbers). The algebra itself is commutative but non-associative (as is generally the case for all Jordan algebras) with the exceptional Lie group F_4 being the automorphism symmetry of the algebra.

However, there is a larger symmetry group involving another structure which can be defined on the space $h_3\mathbb{O}$ which is of particular interest here. This is a cubic norm, or determinant, det(\mathcal{X}) \equiv ($\mathcal{X}, \mathcal{X}, \mathcal{X}$) for $\mathcal{X} \in h_3\mathbb{O}$ which will be presented explicitly in section 6.4. A subspace of the vectors $v \in \mathbb{R}^{27}$ map onto elements $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ that satisfy the homogeneous cubic polynomial equation $\det(\mathcal{X}) = 1$ which expresses a form of the principle relation of equation 2.9 denoted $L(v_{27}) = 1$. This subspace is locally 26-dimensional and hence may be denoted S_{26} , as a homogeneous space, following the convention in the opening of section 4.3, although here S_{26} represents the full space of temporal flow rather than a purely internal fibre space.

The symmetry of this 27-dimensional form $L(v_{27}) = 1$ corresponds to a group of morphisms of the elements of $h_3\mathbb{O}$ which preserve the unit cubic norm; that is the set of actions such as $A_\lambda(\mathcal{X})$, parametrised by $\lambda \in \mathbb{R}$, with $(A_\lambda(\mathcal{X}), A_\lambda(\mathcal{X}), A_\lambda(\mathcal{X})) =$ $(\mathcal{X}, \mathcal{X}, \mathcal{X})$. With the identity transformation labelled by $\lambda = 0$ elements of the corresponding Lie algebra may be represented by the objects $D = \partial A_{\lambda}/\partial \lambda|_{\lambda=0}$. More generally the Lie algebra can be defined directly in terms of the set of operators D that annihilate the norm, that is with:

$$
(D\mathcal{X}, \mathcal{X}, \mathcal{X}) + (\mathcal{X}, D\mathcal{X}, \mathcal{X}) + (\mathcal{X}, \mathcal{X}, D\mathcal{X}) = 0
$$
\n(6.3)

These elements are found to comprise a 78-dimensional Lie algebra of rank 6 (the Cartan subalgebra consists of 6 mutually commuting generators); and these properties, together with the fact that it has a 27-dimensional representation, lead to the identification of the Lie algebra E_6 associated with the exceptional Lie group E_6 . In fact it is one of four real non-compact forms of this Lie algebra denoted $E_{6(-26)}$ (since the Killing form signature is -26 (= 26 – 52). In this paper the same upper case kernel letter, such as 'E', is used to denote either the group or the algebra, depending on the context, for the exceptional Lie groups, although notation such as $L(E_6)$ may be used to emphasise the Lie algebra. Lower case kernel letters are used to denote a classical Lie algebra, such as $\mathfrak{so}(n)$, corresponding to a Lie group, such as $\mathfrak{SO}(n)$, although again whether a statement refers to the Lie group, its algebra or both should generally be clear from the context).

The first construction of the E_6 Lie algebra in terms of action on the space $h_3\mathbb{O}$ dates from 1950 [34] and combined the 52-dimensional algebra of derivations D^R of the Jordan algebra $h_3\mathbb{O}$ (that is, the generators of the automorphism group F_4) with the 26-dimensional set D^B composed of operations of right action on h₃ \mathbb{O} by traceless elements of h₃ \mathbb{O} itself. The total set of elements in this $(52 + 26) = 78$ -dimensional space may be written:

$$
D^{R,B} = D^R : \{ \mathbf{F}_4 \text{ automorphism group of } \mathbf{h}_3 \mathbb{O} \text{ Jordan algebra} \} +
$$

$$
D^B : \{ \text{maps } \mathcal{X} \in \mathbf{h}_3 \mathbb{O} \to \mathcal{X} \circ x, \text{ with } x \in \mathbf{h}_3 \mathbb{O}, \text{ tr}(x) = 0 \} \qquad (6.4)
$$

It can also be shown that the commutator $[D_x^B, D_y^B] = D_x^B D_y^B - D_y^B D_x^B \in D^R$. (The D^R and D^B are analogous to the rotations and boosts, respectively, for the Lorentz group, as we shall see later in this chapter). All elements of the set $D^{R,B}$ have the property exhibited by D in equation 6.3 and are therefore associated with an E_6 group action that preserves $\det(\mathcal{X})$ for any $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ ([1] pp.44–46).

Alternatively the Lie algebra $E_{6(-26)}$ can be expressed in terms of mappings induced on h₃ Ω by the 14 generators of G₂ acting on Ω , supplementing the actions of a basis of 64 independent tracefree 3×3 octonion matrices. This construction of a basis for the E_6 algebra, dating from the 1960s ([35] pp.162–164), in terms of a $(14 + 64) = 78$ -dimensional decomposition can be denoted by $D^{G,S}$ and is composed of two sets:

$$
D^{G,S} = D^G : \{G_2 \text{ automorphism group of } \mathbb{O} \text{ algebra}\} +
$$

$$
D^S : \{\text{maps } \mathcal{X} \in \text{h}_3\mathbb{O} \to x_0\mathcal{X} + \mathcal{X}x_0^{\dagger}, \text{ with } x_0 \in \text{sl}_0(3,\mathbb{O})\} \qquad (6.5)
$$

where $sl_0(3, \mathbb{O})$ is the 64-dimensional set of traceless 3×3 matrices over the octonions and D^G is isomorphic to the 14-dimensional Lie algebra G_2 . All elements of this combined set satisfy equation 6.3 and hence $L(v_{27}) = 1$, in the form of $det(\mathcal{X}) = 1$, is preserved by the associated E_6 group action.

More generally ([1] p.28, [36]) denoting by $\text{sl}_0(n,\mathbb{K})$ the set of traceless $n \times n$ matrices with entries in the division algebra $\mathbb{K} = \mathbb{R}, \mathbb{C}, \mathbb{H}$ or \mathbb{O} for $n > 1$ the commutator of the set $sl_0(n, K)$ is closed only if K is commutative and associative (i.e. for R and C only). However, in all cases $\text{sl}(n,\mathbb{K})$ may be defined to be the Lie algebra of operators on \mathbb{K}^n generated by the elements of $\text{sl}_0(n, \mathbb{K})$ under an appropriate matrix commutation rule (the Lie algebra $L(E_6)$ is identified with $sl(3, \mathbb{O})$ in [36] p.950). The Lie group $SL(n, K)$ of operators on $Kⁿ$, with an associative multiplication even for $K = \mathbb{O}$, may be generated by the elements of this $\text{sl}(n, \mathbb{K})$ Lie algebra. For the case $n = 2$ the group $SL(2, \mathbb{K})$ also has a representation on $h_2\mathbb{K}$ that preserves the determinant.

However, this approach of first defining the Lie algebra purely in itself is not followed here. Rather finite group transformations will be constructed first [37, 38, 39, 40, 41. Here $SL(n, \mathbb{K})$ will be defined principally in terms of a set of group transformations that preserve a particular norm on a vector space, essentially by generalisation from $SL(2,\mathbb{C})$ as a determinant preserving action on $h_2\mathbb{C}$. The need for such a realvalued 'norm' is here motivated by the form $L(v) = 1$. In obtaining the full set of symmetry actions on the form $L(v) = 1$ it is partly a matter of convention whether this group of transformations is given a name of the type $SL(n, K)$. A representation of the corresponding Lie algebra $sl(n, K)$ will be defined and derived subsequently through the group action on the representation space. In particular the $L(E_6) \equiv sl(3, \mathbb{O})$ Lie algebra will be described in terms of a basis of vector fields on the tangent space to the hypersurface S_{26} embedded within the space $h_3\mathbb{O}$, which itself can be considered as a 27-dimensional manifold.

Associativity is required of any group of operations in general and it is the case for the elements of E_6 which will be described explicitly in section 6.4. While in general multiplication between octonions is non-associative it is possible to use them in the construction of algebraic elements such that the multiplication defined between these latter elements is in fact associative. Indeed it will be possible to conceive of the elements of E_6 acting on $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ of equation 6.1 in a manifestly associative way represented as a subgroup of $GL(27, \mathbb{R})$ acting on $v_{27} \in \mathbb{R}^{27}$, with $h_3 \mathbb{O} \equiv \mathbb{R}^{27}$ as vector spaces, such that $L(v_{27}) = det(\mathcal{X}) = 1$ is invariant, similarly as for all forms of $L(v) = 1$ under symmetry operations. Since the construction of the Lie group $E_6 \equiv SL(3, \mathbb{O})$ here relies on the composition properties of the octonions in the following section we first turn to the octonion algebra itself.

6.2 Octonion Algebra and Geometric Symmetries

Having introduced the division algebras in section 2.1 in relation to possible multidimensional forms of temporal flow for equation 2.9 here we focus on the largest such algebra. For the remainder of this chapter we follow references [37, 38, 39, 40, 41] in leading from the properties of octonions through their relation with Lorentz transformations to the construction of the symmetry group E_6 and the corresponding Lie algebra. The main reference for these latter structures in particular is ([38] chapters 3 and 4). The above references are extensively reviewed in this chapter, owing to their importance for the present work, and they also provide the source for much of the notation adopted here.

We begin then with a general octonion which, as an element of an eightdimensional vector space, has eight real parameters $\{a_1 \ldots a_8\}$ and can be written:

$$
a = a_1 + a_2 i + a_3 j + a_4 k + a_5 k l + a_6 j l + a_7 i l + a_8 l \tag{6.6}
$$

The first term could be written as a_1e with $e \equiv 1$ representing the real unit through which real numbers such as $a_1 \in \mathbb{R}$ are embedded in the octonions as $a_1 e \in \mathbb{O}$. The seven imaginary units in this basis $\{i, j, k, k\}, \, j\mathcal{l}, \, k\mathcal{l}, \, j\mathcal{l}, \, k\math$ $l^2 = -1$, are mutually anticommuting, with $il j = -j ul$ etc., with their full algebraic composition described in figure 6.1.

Figure 6.1: The multiplication of any two octonion units is given by \pm the third octonion on the same directed line, with $a + (or -)$ sign for composition aligned with (or against) the arrow on the line, for example $k\ell \,il = -j$.

Hamilton's quaternions are contained as a subalgebra of the octonions with imaginary units $\{i, j, k\}$ composed as $i, j = k$ with cyclic permutations as represented by the arrowed circle in figure 6.1. The six other arrowed lines represent six further equivalent H subalgebras embedded in \mathbb{O} . As can be seen from examples such as $kl = kl$ the notation for the imaginary units is chosen as a mnemonic for these relations, where it should be understood that $k\ell$ is a single imaginary base unit on equal footing with any of the other six, with $k \neq k = l$ and so on.

While multiplication within any of the seven quaternion subalgebras is associative, for example $(kl k) = kl(kl) = -1$, multiplication between any three imaginary base units not situated on the same line in figure 6.1 is anti-associative, with for example $(i j)l = -i(j l) = +kl$. Care needs to be taken due to the possible ambiguity in expressions involving products of octonions due to this lack of general associativity. However, the algebra does satisfy the weaker condition that products involving only two distinct octonions $a, b \in \mathbb{O}$ are associative, for example $(aa)b = a(ab)$, and hence the octonions form an alternative algebra.

Octonion conjugation is defined as a real linear map $a \to \bar{a}$ on \mathbb{O} such that for the real unit $a_1 \rightarrow a_1$ while for the seven imaginary units $a_h \rightarrow -a_h$ ($h = 2 \dots 8$). The octonion conjugate of $a \in \mathbb{O}$ in equation 6.6 is therefore:

$$
\bar{a} = a_1 - a_2 i - a_3 j - a_4 k - a_5 k l - a_6 j l - a_7 i l - a_8 l \tag{6.7}
$$

which applies to any product as $\overline{ab} = \overline{ba}$ and is hence an algebra anti-automorphism. For a given octonion a the norm $|a|$ is a real number defined by:

$$
|a|^2 = a\bar{a} = \sum_{h=1}^{8} a_h^2 \tag{6.8}
$$

which applies to any product as $|ab| = |a||b|$ since the algebra is alternative. Hence the norm is compatible with octonion multiplication and it is these properties, which also imply the existence of a unique inverse:

$$
a^{-1} = \frac{\bar{a}}{|a|^2} \tag{6.9}
$$

for any element $a \neq 0$, which make the octonions a 'normed division algebra' ([1] p.9). By a theorem of Hurwitz from 1898 only four such algebras, of real dimension 1, 2, 4 and 8, exist; these are $\mathbb{R}, \mathbb{C}, \mathbb{H}$ and \mathbb{O} which hence form a unique set of algebras, as listed in section 2.1. With the octonions being the largest normed division algebra and possessing a rich symmetry structure they naturally find use in the present context for identifying possible forms $L(v) = 1$ for temporal flow together with the associated symmetries. In the simplest case for $a, b \in \mathbb{O}$ with $|a| = 1$ and $b \equiv v$ the composition ab with $|ab| = |a||b|$ provides a set of symmetry transformations leaving the form $L(\boldsymbol{v}) := |b|^2 = 1$ invariant.

Octonion conjugation can be used to extract the real and imaginary parts of $a \in \mathbb{O}$ as $\text{Re}(a) = \frac{1}{2}(a + \bar{a})$ and $\text{Im}(a) = \frac{1}{2}(a - \bar{a})$ respectively. While for a complex number $z = x + yi \in \mathbb{C}$ the imaginary part is usually defined such that $\text{Im}(z) = y$ is itself a *real* number, as for example in equation 10.96, for the quaternion and octonion cases the imaginary part is defined as an imaginary number, with for example Im(a) = $a_2i + a_3j + \dots$ for equation 6.6, since such an object in general involves several distinct imaginary units. An inner product for any two octonions $a, b \in \mathbb{O}$ may be defined by:

$$
\langle a, b \rangle = \frac{1}{2} (a\overline{b} + b\overline{a}) = \text{Re}(a\overline{b}) = \sum_{h=1}^{8} a_h b_h.
$$
 (6.10)

For a single octonion $|a|^2 = \langle a, a \rangle$, while for any two octonions $\langle a, b \rangle = \langle b, a \rangle$ and *geometric* orthogonality can be defined by the *algebraic* property $\langle a, b \rangle = 0$. From equation 6.10 it can be seen that any real element of \mathbb{O} is orthogonal to any imaginary element and also any pair of anticommuting octonions (such as $\{i, j\}$ with $ij = -ji$ etc.) are orthogonal to each other. In general a unit imaginary s is an element $s \in \text{Im}(\mathbb{O})$ with unit norm $|s| = 1$ which is not necessarily one of the basis units $\{i, j, k \dots\}$. The real unit 1 together with any two orthogonal imaginary units s, s' define a quaternion subalgebra with basis $\{1, s, s', ss'\} \in \mathbb{H}$, which includes any of the seven $\mathbb H$ subalgebras as described by the seven lines in figure 6.1. More generally any two non-parallel imaginary units in O generate a basis for a quaternion algebra.

For any octonion a with $\text{Im}(a) \neq 0$ the unit imaginary s as the point on the 6sphere of unit imaginary octonions in the direction of $\text{Im}(a)$ can be identified. Any such $s \in S^6$, together with the real unit 1, generate a complex subalgebra of \mathbb{O} with basis $\{1, s\} \in \mathbb{C}$. In particular any $a \in \mathbb{O}$ of equation 6.6 may be written as $a = |a|e^{s\alpha}$, with $\alpha \in \mathbb{R}$ and the Euler identity $e^{s\alpha} = \cos \alpha + s \sin \alpha$ applying in the complex subalgebra. Since any two octonions involve at most two complex subspaces, with bases $\{1, s\}$ and $\{1, s'\}$, it follows from the previous paragraph that any calculation involving only two octonions reduces to the case of the quaternion algebra, which being associate hence accounts for the alternative property of the octonion algebra.

As distinct from the 'octonion conjugation' $a \to \bar{a}$, for each $q \in \mathbb{O}$ with $q \neq 0$ a conjugation map on $a \in \mathbb{O}$ is a linear transformation expressed by the following algebraic composition (which is well defined since O is an alternative algebra):

$$
\phi_q: a \rightarrow qaq^{-1}
$$

that is $\phi_q: a \rightarrow qa\overline{q}$ for $|q| = 1$ (6.11)

where the second expression follows using equation 6.9 and in fact describes the complete set of possible transformations since the first expression is insensitive to $|q|$. Selecting $|q| = 1$ implies not only $q^{-1} = \bar{q}$ but also $q = e^{s\frac{\alpha}{2}}$ where $s \in \mathbb{Q}$ is a unit imaginary and $\alpha \in \mathbb{R}$. Since

$$
|\phi_q(a)| = |q||a||\bar{q}| = |a| \tag{6.12}
$$

and
$$
\langle a, b \rangle = \frac{1}{2}(|a+b|^2 - |a|^2 - |b|^2)
$$
 (6.13)

(the latter by equations 6.8 and 6.10) the map in equation 6.11 represents an *isometry* for the elements of O, since geometric relations are preserved. This isometry, leaving $Re(a)$ invariant and being continuously connected to the identity transformation, represents an action of SO(7) upon the seven-dimensional space of imaginary octonions.

For the quaternion subalgebra, with $a, q \in \mathbb{H}$ in the basis $\{1, i, j, k\}$, the map $\phi_q: a\to qa\bar{q}$, in $\mathbb{H}\to \mathbb{H}$, with $q=e^{i\frac{\alpha}{2}}$ rotates a vector $(a_1, a_2, a_3, a_4)\in \mathbb{R}^4$ by the angle α radians in the (j-k) plane (with $\alpha = 2\pi n$, $n \in \mathbb{Z}$, being the identity transformation). That is, applying equation 6.11 and anticommutation for the imaginary units:

$$
\phi_{e^{i\frac{\alpha}{2}}} : a_1 + a_2i + a_3j + a_4k \rightarrow a_1 + a_2i + e^{i\alpha}(a_3j + a_4k)
$$

and hence
$$
\begin{pmatrix} a_3j \\ a_4k \end{pmatrix} \rightarrow \begin{pmatrix} (a_3 \cos \alpha - a_4 \sin \alpha)j \\ (a_3 \sin \alpha + a_4 \cos \alpha)k \end{pmatrix}
$$
 (6.14)

when considered as an *active* transformation relative to a set of constant basis elements $\{1, i, j, k\}$, which is the point of view adopted for such transformations here (in contrast to passive transformations such as exemplified in equations 3.39–3.41 and described in sections 3.1 and 3.2 for gauge transformations on a principle bundle space). On employing left and right multiplication by independent quaternions of unit norm on the full space \mathbb{H} , with 4 real dimensions, the two-to-one cover of $SO(4)$ is obtained, as alluded to in section 2.1.

Transformations in the space $\text{Im}(\mathbb{H})$, in the form of equation 6.11, may be constructed about any unit imaginary element (taken as i in the example of equation 6.14) as the axis of rotation. In this case for quaternions the map ϕ_q is a universal

two-to-one covering map of S^3 into the group of rotations SO(3) (elements $q \in \mathbb{H}$ with $|q| = 1$ describe the 3-sphere, with q and $-q$ mapping to the same rotation). This is a group homomorphism from the algebraic composition of equation 6.11 for quaternions with group structure $\phi_r \circ \phi_s = \phi_{rs}$ into the geometric transformations SO(3) in a 3-dimensional space, which is also the automorphism group of the algebra H. This group may also be generated by composing several actions of the form in equation 6.11 requiring $|q| = 1$ and $q \in \text{Im}(\mathbb{H})$, that is from the 2-sphere $S^2 \subset S^3$ alone. Similar compositions of actions will actually be required for the octonion case in order to construct the full symmetry, as described in the following.

In the case of the octonions the map in equation 6.11 for $\phi_{e^{i\frac{\alpha}{2}}}$ again fixes the $(1-i)$ -plane but now rotates all *three* mutually orthogonal planes, corresponding to the three quaternion subalgebras containing i identified in figure 6.1, simultaneously by α radians. Here the full set of maps ϕ_q , with $q \in \mathbb{O}$ and $|q| = 1$, does not form a group homomorphism of the 7-sphere S^7 into $SO(7)$ since in general there may be no value of $q \in \mathbb{O}$ for which the map $\phi_q(a)$ is equivalent to the composition $\phi_r(\phi_s(a))$, with $r, s \in \mathbb{O}$, due to the non-associativity of the octonions. However it is precisely through this property of octonion composition that the set of maps ϕ_q of equation 6.11 can generate the full Lie group $SO(7)$ by including ordered, or *nested*, combinations such as $\phi_r(\phi_s(a))$ on $a \in \mathbb{O}$ as elementary symmetry operations.

In general a representation R on a vector space V is a structure preserving homomorphism from group elements $\{g_1, g_2, g_3, e\} \in G$ with $g_1g_2 = g_3$ into representation matrices with $R(g_1)R(g_2) = R(g_3)$ and $R(e) = 1$, where the unit matrix 1 represents the identity transformation on V . Since group structure is associative the above actions ϕ_r, ϕ_s , owing to the octonion non-associativity, do not technically represent the Lie group $SO(7)$.

The group structure of these transformations can however be seen when these actions are instead represented by matrices $R(\phi) \in GL(7, \mathbb{R})$ acting on the vector space \mathbb{R}^7 . Consider the example of $\phi_r(\phi_s(a)) = r(s(a)\overline{s})\overline{r}$ with $r = i \in \mathbb{O}$ and $s = l \in \mathbb{O}$. The map $\phi_l : a \to la\bar{l}$ is a linear transformation of the components of Im(a), that is $\{a_2, \ldots, a_8\}$ of equation 6.6, which can be represented on \mathbb{R}^7 by the action of the diagonal 7×7 matrix $R(\phi_l) = \text{diag}(-1, -1, -1, -1, -1, -1, +1) \in GL(7, \mathbb{R})$. Similarly the map $\phi_i: a \to ia\overline{i}$ is represented by $R(\phi_i) = \text{diag}(+1, -1, -1, -1, -1, -1, -1)$. These combine together as the map $R(\phi_i)R(\phi_i) = \text{diag}(-1, +1, +1, +1, +1, +1, -1),$ by matrix multiplication, which does not correspond to any single conjugation action of equation 6.11 but precisely represents the combined action $a \to i(l(a)\overline{l})\overline{i}$. Hence while $i(l(a)\overline{l})\overline{i} \neq (i\overline{l})a(\overline{l}\overline{i})$, due to the octonion non-associativity, the nested action can be representated in $GL(7,\mathbb{R})$ with $R(\phi_i)R(\phi_i) = R(\phi_i \circ \phi_i)$, and with matrix compositions in general, which fully represents the $SO(7)$ Lie group structure. (A similar situation is found for the spinor representation of SO(7) obtained from the one-sided composition action $a \to r(s(a))$, and also for the dual spinor using right actions alone).

The octonion algebra provides a way to express these symmetry transformations in a compact algebraic form, which uses the non-associativity in order to describe the full symmetry, and which may be unfolded into a more explicit group representation in terms of matrices in $GL(n,\mathbb{R})$. In fact by using these octonion properties the full SO(7) rotation group can be generated with the elements $|r| = 1$ and $r \in Im(\mathbb{O})$, that is on the 6 sphere $S^6 \subset S^7$ alone, as described below.

Firstly, setting $\alpha = \pm \pi$ for a single action the conjugation map $\phi_{e^{r\frac{\pm \pi}{2}}}$, with r an imaginary octonion unit, corresponds to rotating the three planes orthogonal to the (1-r)-plane by $\pm 180^\circ$, hence reflecting, or 'flipping' these three planes. This can be readily seen since $e^{r\frac{\pm\pi}{2}} = \cos\frac{\pi}{2} + r\sin\frac{\pm\pi}{2} = \pm r$ and hence equation 6.11 is simply a conjugation map by a unit imaginary element $r \in S^6$. For example with $r = i$ the map $\phi_{e^{i\frac{\pi}{2}}}(j) = i j \overline{i} = -j$ acts on j, as well as each of the other five imaginary units ${k, kl, il, il, l},$ as a sign flip.

Performing a second reflection based on the same (1-r)-plane naturally cancels the first and leaves no total effect. However in performing the second flip with respect to a *different* plane, namely the $(1-(r \cos \frac{\beta}{2} + s \sin \frac{\beta}{2}))$ -plane with s a unit imaginary orthogonal to r, while the combined reflections still cancel for most components a residual rotation by β radians in the (r-s)-plane remains as the net effect on \mathbb{R}^8 . That is the two reflections applied to any $a \in \mathbb{O}$ as:

$$
\phi_{r,s,\frac{\beta}{2}}(a) = (r \cos \frac{\beta}{2} + s \sin \frac{\beta}{2})(-r \ a \ (-\bar{r}))(\overline{r \cos \frac{\beta}{2} + s \sin \frac{\beta}{2}}) \tag{6.15}
$$

rotates the components of a in the $(r-s)$ -plane by β radians, corresponding to two reflections in two mirror lines in this plane, while giving the identity map on the remaining components. Although two discrete flips are involved in this equation the total effect on vectors in \mathbb{O} is of a rotation in the $(r-s)$ -plane varying *continuously* with the parameter $\beta \in \mathbb{R}$, with the identity transformation for $\beta = 0$. In the sevendimensional space of $\text{Im}(\mathbb{O})$ the 21 possible choices of rotation planes from the 21 sets of imaginary base unit pairs for $\{r, s\}$ describes the full Lie group SO(7). The 14parameter automorphism group of the octonions, that is the exceptional Lie group G_2 , is contained as a subgroup of this SO(7) as will be discussed in section 6.4.

The first rotation in equation 6.15 is taken as $-\pi$, that is with $e^{r\frac{-\pi}{2}} = -r$, followed by the second rotation by $+\pi$. The corresponding minus signs for the $-\pi$ rotation in the middle brackets on the right-hand side trivially cancel here but the minus sign is needed for the one-sided spinor actions $a \to (r \cos \frac{\beta}{2} + s \sin \frac{\beta}{2})(-r(a))$ in order for $\beta = 0$ to correspond to the identity transformation in the spinor representation. This latter expression is also compatible with the identify transformation for the spinor case corresponding to $\beta = 4\pi n$ with $n \in \mathbb{Z}$, rather than any multiple of 2π as for the vector representation of equation 6.15.

6.3 Lorentz Transformations on Spacetime Forms

As well as rotations in spaces with a Euclidean metric, such as the case of $SO(7)$ above, composition of division algebra elements can also be used to describe transformations in spaces with a Lorentzian metric, such as on the tangent space of a spacetime manifold. The content of this section is largely based on reference [37]. We begin here with an Hermitian 2×2 octonion matrix X which may be written as:

$$
X = \begin{pmatrix} t+z & \bar{a} \\ a & t-z \end{pmatrix} \in \mathbf{h}_2 \mathbb{O}
$$
 (6.16)

with $a \in \mathbb{O}$ in the general form of equation 6.6 and $\{t, z\} \in \mathbb{R}$, and hence X is 10-dimensional over the real numbers. Since the components of X , involving only a single octonion a, can be taken to lie within a single complex subalgebra of $\mathbb O$ there are no problems with commutativity or associativity in unambiguously defining the determinant of the matrix X in the usual way as:

$$
\det(X) = (t+z)(t-z) - a\bar{a} = t^2 - a_1^2 - a_2^2 \dots - a_8^2 - z^2 \tag{6.17}
$$

This expression has the same form as the square of an invariant interval represented by a Lorentz 10-vector x (or interval of 'proper time' τ), with 10-dimensional spacetime metric $\eta = \text{diag}(+1, -1, \ldots, -1)$, which can be written as:

$$
|\mathbf{x}|^2 = \mathbf{x}^T \eta \mathbf{x} = x_0^2 - x_1^2 - x_2^2 \dots - x_8^2 - x_9^2 \tag{6.18}
$$

Closely analogous structures are obtained for all four normed division algebras, $\mathbb{K} =$ $\mathbb{R}, \mathbb{C}, \mathbb{H}$ or $\mathbb{O},$ with $h_2\mathbb{K}$ representing Lorentz vectors in $(k+2)$ -dimensional spacetime where $k = \dim_{\mathbb{R}}(\mathbb{K})$. For example in the familiar case of 4-dimensional spacetime a Lorentz 4-vector (t, x, y, z) can be represented by:

$$
X = \begin{pmatrix} t+z & x-yi \\ x+yi & t-z \end{pmatrix} = t\sigma^0 + x\sigma^1 + y\sigma^2 + z\sigma^3 \in h_2\mathbb{C}
$$
 (6.19)

This case will be considered in more detail in the section 7.1 where the σ -matrices are presented in equation 7.14. The above expression can be generalised by replacing $\sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ in equation 7.14 with $\sigma^q = \begin{pmatrix} 0 & -q \\ q & 0 \end{pmatrix}$ for $q = i, j$ and k for the quaternion case or $q = i, j, k, kl, il$, il and l for the octonion case of equation 6.16.

A Weyl spinor can be expressed as the 2-component object $\theta = \begin{pmatrix} a \\ b \end{pmatrix}$ $\binom{a}{b} \in \mathbb{K} \oplus \mathbb{K}$, with the Hermitian conjugate $\theta^{\dagger} = (\bar{a} \bar{b})$, and hence each spinor has 16 real components for the octonion case. As an element of $h_2\mathbb{K}$ the square of a spinor:

$$
\theta \theta^{\dagger} = \begin{pmatrix} |a|^2 & a\bar{b} \\ b\bar{a} & |b|^2 \end{pmatrix}
$$
 (6.20)

has
$$
\det(\theta \theta^{\dagger}) = 0 \tag{6.21}
$$

and hence corresponds to a null-vector in $(k + 2)$ -dimensional spacetime. The 'time' component of this null-vector can be expressed in a scalar spinor product $t = \frac{1}{2}$ $\frac{1}{2}\theta^{\dagger}\theta=$ 1 $\frac{1}{2}(|a|^2+|b|^2)$, while the time component of a general element of $X \in h_2\mathbb{K}$ is given by $\overline{t}=\frac{1}{2}$ $\frac{1}{2}$ tr(X), as can be seen in the examples of equations 6.16 and 6.19.

Lorentz transformations in $(k+2)$ -dimensional spacetime are defined as actions Λ which preserve proper time intervals, that is with $|\Lambda(x)| = |x|$. The subset of actions continuously connected to the identity transformation may be composed together to form the Lorentz group $SO^+(1, k+1)$. Since $|\mathbf{x}|^2 \equiv \det(X)$ the rotations and boosts of these geometric spacetime symmetries can be associated with algebraic compositions in the relevant division algebra which preserve the determinant, and Hermitian property, of $h_2\mathbb{K}$. To represent a 10-dimensional Lorentz transformation the Hermitian requirement can be achieved by a conjugation map on $X \in h_2\mathbb{O}$ with the 2×2 matrix M which is well defined if there is no associativity ambiguity:

$$
R: X \to M X M^{\dagger} := (MX) M^{\dagger} = M (X M^{\dagger}) \tag{6.22}
$$

This in turn is achieved if the components of M all belong to a single complex subspace of \mathbb{O} (an alternative possibility is for the columns of $\text{Im}(M)$ to be real multiples of each other [37] p.21). In this case $\det(M)$ is well defined and the further requirement that $\det(MM^{\dagger}) = 1$ is sufficient to ensure that the conjugation map $X \to MXM^{\dagger}$ leaves $\det(X)$ invariant. These two-sided transformations on the vector X are required to be *compatible* with the one-sided actions on the spinor θ and its Hermitian conjugate θ^{\dagger} , meaning that there should also be no associativity problems in relating these representations as:

$$
M(\theta \theta^{\dagger})M^{\dagger} = (M\theta)(\theta^{\dagger}M^{\dagger}) = (M\theta)(M\theta)^{\dagger}
$$
\n(6.23)

where on the left-hand side an octonionic vector is composed as $\theta\theta^{\dagger}$, which is not a general element of $h_2\mathbb{O}$ due to equation 6.21. This compatibility, which will be needed in the following section for the 3×3 matrix case, is satisfied, along with equation 6.22, if the components of each individual M all belong to the same complex subalgebra of \mathbb{O} and also $\det(M) \in \mathbb{R}$. Together with the requirement that the vector transformation preserves $\det(X)$ this implies that $\det(M) = \pm 1$. A complete set of such transformation matrices is listed in table 6.1.

Category 1: Boosts
$$
B_{tz}(\alpha)
$$
, $B_{tx}(\alpha)$ and $B_{tq}(\alpha)$ with:
\n
$$
M_{tz}(\alpha) = \begin{pmatrix} e^{+\frac{\alpha}{2}} & 0 \\ 0 & e^{-\frac{\alpha}{2}} \end{pmatrix}, \ M_{tx}(\alpha) = \begin{pmatrix} \cosh \frac{\alpha}{2} & \sinh \frac{\alpha}{2} \\ \sinh \frac{\alpha}{2} & \cosh \frac{\alpha}{2} \end{pmatrix}, \ M_{tq}(\alpha) = \begin{pmatrix} \cosh \frac{\alpha}{2} & q \sinh \frac{\alpha}{2} \\ -q \sinh \frac{\alpha}{2} & \cosh \frac{\alpha}{2} \end{pmatrix}
$$

Category 2: Rotations $R_{xq}(\alpha)$, $R_{xz}(\alpha)$ and $R_{zq}(\alpha)$ with:

$$
M_{xq}(\alpha) = \begin{pmatrix} e^{-q\frac{\alpha}{2}} & 0\\ 0 & e^{+q\frac{\alpha}{2}} \end{pmatrix}, \ M_{xz}(\alpha) = \begin{pmatrix} \cos\frac{\alpha}{2} & \sin\frac{\alpha}{2} \\ -\sin\frac{\alpha}{2} & \cos\frac{\alpha}{2} \end{pmatrix}, \ M_{zq}(\alpha) = \begin{pmatrix} \cos\frac{\alpha}{2} & -q\sin\frac{\alpha}{2} \\ -q\sin\frac{\alpha}{2} & \cos\frac{\alpha}{2} \end{pmatrix}
$$

Category 3: Transverse Rotations $R_{r,s}(\alpha)$ with:

$$
M_{r,s2}(\alpha) = \begin{pmatrix} r\cos\frac{\alpha}{2} + s\sin\frac{\alpha}{2} & 0\\ 0 & r\cos\frac{\alpha}{2} + s\sin\frac{\alpha}{2} \end{pmatrix} \text{ nested with } M_{r,s1} = \begin{pmatrix} -r & 0\\ 0 & -r \end{pmatrix}
$$

Table 6.1: Three categories of matrices [37, 38] for conjugation action on $X \in h_2\mathbb{O}$ preserving $\det(X)$; with $q, r, s \in \{i, j, k, k\}, \mathcal{Y}, \mathcal{Y}, \mathcal{Y}, \mathcal{Y}\}$ there are $(1 + 1 + 7) = 9$ boosts, $(7 + 1 + 7) = 15$ rotations and 21 $R_{r,s}(\alpha)$ transverse rotations (where the subscript r, s denotes the ordered pair $\{r, s\}$ of imaginary units) representing the 45-dimensional group of Lorentz transformations on a 10-dimensional spacetime.

In the first two categories $\det(M) = +1$ for each of the 24 actions. The third category is a simple 2×2 diagonal matrix form of equation 6.15, with the parameter β replaced by α . The action of the category 3 matrices is ordered by nesting the conjugation as:

$$
R_{r,s}(\alpha)X = M_{r,s2} \left(M_{r,s1} \left(X \right) M_{r,s1}^{\dagger} \right) M_{r,s2}^{\dagger} \tag{6.24}
$$

with $\det(M_{r,s1}) = \det(M_{r,s2}) = -1$. Since the latter matrices are always combined in pairs, and hence are analogous to the action of a single matrix with a determinant of $+1$, the full group of transformations is denoted $SL(2, \mathbb{O})$. It is composed of the 45 actions in table 6.1, each of which describes a one-parameter subgroup with $R(\alpha)R(\beta) = R(\alpha+\beta)$ and each of which represents transformations in a single 2-dimensional plane in 10 dimensional spacetime.

For each of the 45 transformations with $\alpha = 0$ and $\alpha = 2\pi$ it can be seen that $M = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $M = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ $0 -1$ respectively (this is effectively true for the category 3 case since these can be expressed by conjugation with a single such matrix M for these α values). Hence $SL(2,\mathbb{O})$ is the double cover of the 10-dimensional Lorentz group $SO^+(1,9)$, that is $SL(2,\mathbb{O}) \rightarrow SO^+(1,9)$ is a two-to-one homomorphism with kernel ${M = \pm 1_2}$, where 1_2 is the 2×2 identity matrix, since both cases for M give the identity transformation on X due to the two-sided action in equation 6.22.

A number of subgroups may also be identified. The subgroup leaving $tr(X)$ invariant, composed of the 36 category 2 and 3 transformations, defines $SU(2,\mathbb{O})$ which is the two-to-one cover of the purely rotational Lorentz subgroup $SO(9)$, leaving the t-component in equation 6.16 invariant. In turn the 21 category 3 transformations alone form the $Spin(7)$ subgroup as the double cover of $SO(7)$. The structure of these subgroups, including the $SO(8)$ obtained by augmenting the $SO(7)$ with an additional 7 $R_{xx}(\alpha)$ actions from category 2, will also be important for enlarging beyond SL(2, 0) for the 3×3 matrix case in the following section.

Finally in this section we note that for the quaternion case, obtained by restricting all transformations in table 6.1 for $q, r, s \in \{i, j, k\}$, there remain 15 transformations $(5, 7 \text{ and } 3 \text{ for category } 1, 2 \text{ and } 3 \text{ respectively})$ acting on $h_2 \mathbb{H}$ forming $SL(2, \mathbb{H})$ as the double cover of the Lorentz group $SO^+(1, 5)$ on 6-dimensional spacetime. Here, loosening the restriction $\det(M) = \pm 1$ for the transformation matrices, each of the three transverse rotations can be achieved by a single unnested conjugation map such as:

$$
R_i(\alpha)(X) = M_i X M_i^{\dagger} \quad \text{with} \quad M_i(\alpha) = \begin{pmatrix} e^{i\frac{\alpha}{2}} & 0\\ 0 & e^{i\frac{\alpha}{2}} \end{pmatrix}
$$
 (6.25)

which acts on the quaternion component $a \in \mathbb{H}$ of $X = \binom{p}{a} \overline{a}$ \in h₂ \mathbb{H} by fixing the (1-*i*)plane while performing a rotation in the $(j-k)$ -plane of α radians as was described in equation 6.14. Taking a similar form to equation 6.25 the two actions $R_i(\alpha)$ and $R_k(\alpha)$ rotate the $(k-i)$ -plane and $(i-j)$ -plane respectively. This is possible for the quaternions since there is only one imaginary plane orthogonal to each imaginary base unit. This is unlike the case for the octonions in which the nested transverse rotations are needed to describe all 21 such single plane rotations (as explained towards the end of the previous section) and hence account for the complete subgroup $SO(7) \subset SO^+(1, 9)$. In addition the restriction $\det(M) = \pm 1$ is imposed for the octonion case in order to

meet the compatibility requirement of equation 6.23 as will be needed for extension to the 3×3 case as noted after that equation.

For the case of 4-dimensional spacetime the six Lorentz transformations are represented on h₂C by the six category 1 and 2 matrices M in table 6.1 with q taking a single value such as i. This set of actions with $\det(M) = +1$ forms the group $SL(2,\mathbb{C})$ as the double cover of the Lorentz group $SO^+(1,3)$, as will be studied in more detail in section 7.1. In this case there are no imaginary orthogonal planes for the transformation of equation 6.25 to rotate and the action $R_i(\alpha)$, which may be considered as a residue from the 2×2 matrix cases for \mathbb{H} and \mathbb{O} , not only preserves $\det(X)$ but also leaves each *component* of any $X \in h_2\mathbb{C}$ unchanged. In this sense the action in equation 6.25 may be interpreted as an *internal* $U(1)$ symmetry, relative to the external Lorentz symmetry of 4-dimensional spacetime, as will be relevant for the case of the embedding $h_2 \mathbb{C} \subset h_3 \mathbb{O}$ in section 8.2.

6.4 E_6 Transformations on a Form of Time

In this paper the emphasis is on symmetries of forms of multi-dimensional temporal flow $L(\mathbf{v}) = 1$, that is *isochronal* symmetries as introduced in section 2.1, rather than on isometries of a higher-dimensional space or spacetime, as described for equations $6.11-$ 6.13 and in the previous section for example. The $SL(2, \mathbb{O})$ action preserving $det(X)$ with $X \in h_2\mathbb{O}$, described in the previous section, can be interpreted in either way, but there is no reason to restrict multi-dimensional forms of $L(\mathbf{v})$ to have such a spacetime interpretation, as it does in taking the quadratic form of $\det(X)$ for example. Further, with O being the largest division algebra, there is no clear extension of this construction based on $h_2\mathbb{K}$ to a higher-dimensional spacetime symmetry. This leads to the consideration of the extension of h₂ σ to the 27-dimensional space of 3×3 Hermitian octonion matrices $h_3\mathbb{O}$ which has richer symmetry properties while still possessing an underlying structure appropriate for a form of temporal flow.

An element $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ of equation 6.1 may be written as (again closely following [38] chapters 3 and 4 together with [39, 40, 41] and generally adopting the notation therein):

$$
\mathcal{X} = \begin{pmatrix} p & \bar{a} & c \\ a & m & \bar{b} \\ \bar{c} & b & n \end{pmatrix} = \begin{pmatrix} X & \theta \\ \theta & \theta \end{pmatrix} \in \mathbf{h}_3 \mathbb{O} \tag{6.26}
$$

with $p, m, n \in \mathbb{R}$ and $a, b, c \in \mathbb{O}$, while X and θ have the structure of octonionic 2×2 vectors (equation 6.16) and 1×2 spinors respectively, familiar from the previous section.

Under the Jordan product of equation 6.2 elements $\mathcal{X}, \mathcal{Y} \in \mathrm{h}_3\mathbb{O}$ form the exceptional Jordan algebra. However it is the structure of a cubic norm, or determinant, which may be defined on $h_3\mathbb{O}$, without any ambiguity due to the non-associativity of the octonions, that is of interest here. The cubic norm is a homogeneous polynomial form in the components of h₃ \mathbb{O} as a mapping $\mathcal{X} \to \det(\mathcal{X}) \in \mathbb{R}$ into the real numbers,

and hence has the correct structure for a form of $L(v) = 1$. This determinant may be expressed in several equivalent ways including:

$$
\det(\mathcal{X}) = \det(X)n + 2X \cdot (\theta \theta^{\dagger}) \tag{6.27}
$$

$$
= \ pmn - p|b|^2 - m|c|^2 - n|a|^2 + 2\text{Re}(\bar{a}\bar{b}\bar{c}) \tag{6.28}
$$

where the 10-dimensional Lorentz inner product $X \cdot Y = \frac{1}{2}(\text{tr}(X \circ Y) - \text{tr}(X)\text{tr}(Y)),$ with $X, Y \in h_2\mathbb{O}$, in the first line together with equation 6.26 can be used to derive the second line in which the cubic composition of components, consistent with the homogeneous form of equation 2.9, is explicitly seen.

The 2×2 matrices M of $SL(2, \mathbb{O})$ actions listed in table 6.1 can be embedded in the upper-left corner of 3×3 matrices M to obtain the conjugation action for the 3×3 case $R : \mathcal{X} \to \mathcal{M} \mathcal{X} \mathcal{M}^{\dagger}$ with:

$$
\mathcal{M}\mathcal{X}\mathcal{M}^{\dagger} = \left(\begin{array}{c|c} M & 0 \\ \hline 0 & 1 \end{array}\right) \left(\begin{array}{c|c} X & \theta \\ \hline \theta^{\dagger} & n \end{array}\right) \left(\begin{array}{c|c} M & 0 \\ \hline 0 & 1 \end{array}\right)^{\dagger} = \left(\begin{array}{c|c} M\mathcal{X}M^{\dagger} & M\theta \\ \hline \theta^{\dagger}M^{\dagger} & n \end{array}\right) \tag{6.29}
$$

This expression contains the vector $X \to R(X) = M X M^{\dagger}$, spinor $\theta \to R(\theta) = M \theta$ and scalar $n \to 1n$ representations of $SL(2, \mathbb{O})$, each transforming in the appropriate way with the form of the action R determined correspondingly. These transformations respect the 3×3 block structure, as do nested compositions in augmenting the 2×2 matrix actions such as equation 6.24 to expressions of the form:

$$
\mathcal{X} \to R(\mathcal{X}) = \mathcal{M}_n(\dots(\mathcal{M}_1(\mathcal{X})\mathcal{M}_1^{\dagger})\dots)\mathcal{M}_n^{\dagger}
$$
(6.30)

which acts, for example, on the spinor as $\theta \to R(\theta) = M_n(\dots(M_1(\theta)))$. As well as preserving $\det(X)$ with $X \in h_2\mathbb{O}$ the 45 actions of 2×2 matrices M from table 6.1 when embedded in the 3×3 matrices M for equation 6.29 also preserve det(X) for $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ since, from equation 6.27:

$$
\det(R(\mathcal{X})) = \det(R(X))n + 2R(X) \cdot (R(\theta)R(\theta^{\dagger}))
$$

=
$$
\det(R(X))n + 2R(X) \cdot R(\theta \theta^{\dagger})
$$

=
$$
\det(X)n + 2X \cdot \theta \theta^{\dagger}
$$

=
$$
\det(\mathcal{X})
$$
 (6.31)

where the second equality is a result of 'compatibility', and motivates the introduction of this requirement in equation 6.23, and the third equality follows from the Lorentz symmetry of the $SL(2, \mathbb{O})$ action. It is also by compatibility that the 45 $SL(2, \mathbb{O})$ transformations act as one-parameter subgroups on the spinor θ (given the minus signs for the M_1 components for the transverse rotations, originating in equation 6.15, as for $M_{r,s1}$ in table 6.1) as well as on the vector X.

These $SL(2,\mathbb{O})$ actions, called Lorentz transformations when acting on $X \in$ h2O representing 10-dimensional spacetime, also identify 45 one-parameter subgroups acting on $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$, with $R(\alpha)R(\beta)\mathcal{X} = R(\alpha+\beta)\mathcal{X}$, preserving det (\mathcal{X}) (where $R(\alpha)\mathcal{X}$ denotes a particular action $R(\mathcal{X})$, for example from table 6.1, for a particular transformation parameter α). Hence these 45 actions are one-parameter subgroups of $E_6 := SL(3, \mathbb{O})$ which is defined as the group of symmetry transformations under which the determinant on $h_3\mathbb{O}$ is invariant. Again we emphasise that the key here is the structure of a higher-dimensional form of temporal flow which, while necessarily containing a 4-dimensional form perceived as spacetime, does not itself need to possess a higher-dimensional spacetime interpretation.

The exceptional Lie group E_6 is 78-dimensional, as described in section 6.1 and hence the 45 actions adopted from $SL(2, \mathbb{O})$ represented on $h_3\mathbb{O}$ is only part of the full symmetry picture. However the scope of the $SL(2, \mathbb{O})$ action can be enlarged by noting that there are three similar and natural ways to embed the vector X, spinor θ and scalar n representations of $SL(2, \mathbb{O})$ in the 3×3 matrix $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$. The original 'type 1' action described in equation 6.29 for the embedding depicted in equation 6.26 can be written more explicitly in terms of the matrix components:

$$
\mathcal{M}^{(1)} = \begin{pmatrix} M_{11} & M_{12} & 0 \\ M_{21} & M_{22} & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{acting on} \quad \begin{pmatrix} X_{11} & X_{12} & \theta_1 \\ X_{21} & X_{22} & \theta_2 \\ \overline{\theta}_1 & \overline{\theta}_2 & n \end{pmatrix} \tag{6.32}
$$

Maintaining the variables $p, m, n \in \mathbb{R}$ and $a, b, c \in \mathbb{O}$ in the same component locations of the 3×3 matrix $\mathcal{X} \in \mathrm{h}_{3}\mathbb{O}$ in equation 6.26 their placement within the 2×2 vector $X = \begin{pmatrix} X_{11} & X_{12} \\ X_{21} & X_{22} \end{pmatrix}$ and 1×2 spinor $\theta = \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$ $\begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$ under SL(2, 0) may be reassigned by permuting the components of the matrices $\mathcal M$ as follows:

$$
\mathcal{M}^{(a)} = \mathcal{TM}^{(b)}\mathcal{T}^{\dagger} \text{ for } (a, b) = (2, 1), (3, 2), (1, 3) \text{ with } \mathcal{T} = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}
$$
(6.33)

With $\mathcal{T}^{\dagger} = \mathcal{T}^{-1} (= \mathcal{T}^2)$ it can be seen that $\det(\mathcal{T}\mathcal{X}\mathcal{T}^{\dagger}) = \det(\mathcal{X})$ and since $\det(\mathcal{T}) = 1$ the action $\mathcal{X} \to \mathcal{T}\mathcal{X}\mathcal{T}^{\dagger}$ can itself be considered as a transformation of the SL(3, 0) symmetry. The matrices $\mathcal{M}^{(2)}$ and $\mathcal{M}^{(3)}$ then correspond to 'type 2' and 'type 3' transformations respectively with:

$$
\mathcal{M}^{(2)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & M_{11} & M_{12} \\ 0 & M_{21} & M_{22} \end{pmatrix} \quad \text{acting on} \quad \begin{pmatrix} p & \bar{\theta}_1 & \bar{\theta}_2 \\ \theta_1 & X_{11} & X_{12} \\ \theta_2 & X_{21} & X_{22} \end{pmatrix} \tag{6.34}
$$

and

$$
\mathcal{M}^{(3)} = \left(\begin{array}{c|c|c}\nM_{22} & 0 & M_{21} \\
\hline\n0 & 1 & 0 \\
\hline\nM_{12} & 0 & M_{11}\n\end{array}\right) \quad \text{acting on} \quad \left(\begin{array}{c|c}\nX_{22} & \theta_2 & X_{21} \\
\hline\n\bar{\theta}_2 & m & \bar{\theta}_1 \\
\hline\nX_{12} & \theta_1 & X_{11}\n\end{array}\right) \tag{6.35}
$$

The three $\mathcal{M}^{(a)}$ represent three embeddings of the 2×2 matrix actions of table 6.1 into a 3×3 matrix form acting on the same h₃^① components of equation 6.1. Each type 1, 2 or 3 action, even for the nested case of equation 6.30 with $\mathcal{M} \to \mathcal{M}^{(a)}$ for $a = 1, 2$ or 3, respects the corresponding block structure in equation 6.32, 6.34 or 6.35 respectively. Indeed the type 2 and 3 cases are effectively obtained by a simple symmetric permutation of the three octonion and three real entries in $h_3\mathbb{O}$ under the original type 1 action of equation 6.29, and hence for all three types of transformation $\det(\mathcal{X})$ is invariant, as was shown for the type 1 case in equation 6.31. (In addition to the discrete actions of equation 6.33 continuous type transformations may also be defined as described in [38] section 4.4).

With three possible embeddings of the 45-dimensional $SL(2, \mathbb{O})$ transformations there are now a total of $3 \times 45 = 135 \text{ det}(\mathcal{X})$ -preserving one-parameter subgroup actions for $E_6 := SL(3, \mathbb{O})$, which cannot be independent since E_6 is known to be a 78-dimensional group. A basis for the E_6 actions on $h_3\mathbb{O}$ may be obtained by requiring linear independence at the Lie algebra level. However a $G = \mathrm{E}_6$ manifold is not welldefined in terms of the space of 3×3 matrices M upon which to identify tangent vectors with Lie algebra elements. This is in contrast to a case such as $G = SO(3)$ represented by real 3×3 matrices $R \in SO(3)$ acting on vectors $v_3 \in \mathbb{R}^3$. In this case the space of matrices R, with $RR^T = 1_3$ and $\det(R) = 1$ describing topologically the 3-sphere $S³$ with antipodal points identified, defines the group space G upon which tangent vector fields may represent the Lie algebra, as was depicted for the general case in figure 2.5. The Lie algebra may be described in terms of left-invariant vector fields on the group manifold or in terms of the tangent vectors at the identity $e \in G$ through the isomorphism $L(G) \equiv T_e G$. (An example for the latter case was listed in the set of Lie algebra elements $\{L_{pq}\}\$ of equation 2.32 for $G = SO(3)$.

This situation can be understood by considering how a Lie group manifold, on the tangent space of which the Lie algebra may be defined, might also be identified for the Lorentz groups in $(k + 2)$ -dimensional spacetime, with $k = \dim_{\mathbb{R}}(\mathbb{K})$ and $\mathbb{K} =$ $\mathbb{C}, \mathbb{H}, \mathbb{O}$, represented by the action of $SL(2, \mathbb{K})$ on $h_2\mathbb{K}$ matrices. In the first case for $\mathbb{K} = \mathbb{C}$ since complex 2×2 matrices have $(4 \times 2) = 8$ real parameters and $\det(M) =$ $1 \in \mathbb{C}$ represents two constraints on the matrices $M \in SL(2,\mathbb{C})$ these actions are described by $(8 - 2) = 6$ real parameters, which equals the dimension of the Lorentz group $SO^+(1,3)$. This set of 2×2 matrices can take the form of the first six matrices in table 6.1 for $q = i$ as described at the end of section 6.3. Hence these six degrees of freedom of the matrices $SL(2, \mathbb{C})$ fully describe the corresponding group manifold $G \equiv SL(2,\mathbb{C})$ (as the double cover of $SO^+(1,3)$), upon which the Lie algebra of tangent vector fields may be constructed (having the same Lie algebra structure as $\mathrm{so}^+(1,3)$). This Lie algebra, in terms of tangent vectors at the identity $e \in G$, will be explicitly listed as the set of 2×2 matrices $\{M\}$ in equations 8.7 and 8.8 of section 8.1.

For 2×2 quaternion matrices under the constraint $\det(M) = 1 \in \mathbb{H}$ there are $(4 \times 4) - 4 = 12$ free parameters for $M \in SL(2, \mathbb{H})$, insufficient alone to describe the 15-dimensional Lorentz group $SO^+(1,5)$. However, including the 3 transverse rotations via the 2×2 matrix actions of equation 6.25 (one for each imaginary unit of \mathbb{H}) by loosening the constraint on the matrix determinant to $|\det(M)| = 1 \in \mathbb{R}$ results in a total of $(4 \times 4) - 1 = 15$ free parameters. Hence, as for the complex case, a subset of 2×2 quaternion matrices can be identified with a group manifold structure as the double cover of the Lorentz group, here for a 6-dimensional spacetime, and the tangent space to this manifold hence used to describe the Lie algebra so^{$+(1, 5)$}.

However for the 45-dimensional Lorentz group in 10-dimensional spacetime the maximum of $(4 \times 8) = 32$ parameters available in a 2×2 octonion matrix are clearly insufficient to parametrise the full group, and hence the matrices of $SL(2, \mathbb{O})$ in table 6.1 cannot immediately be related to a Lie group manifold as they could for the complex $(SL(2, \mathbb{C})$ with $\det(M) = 1$ and quaternion $(SL(2, \mathbb{H})$ with $|\det(M)| = 1$ cases. Indeed this is why nested $SL(2, \mathbb{O})$ actions are required in the octonion case to make up the extra transformations. Similarly for $SL(3, \mathbb{O})$, with a maximum $(9 \times 8) = 72$ real parameters available in the 3×3 octonion matrices, such objects are insufficient to represent the full 78-dimensional group manifold for E_6 .

The nested action $X \to \mathcal{M}_2(\mathcal{M}_1 \mathcal{X} \mathcal{M}_1^{\mathsf{T}}) \mathcal{M}_2^{\mathsf{T}}$, in the form of equation 6.30, for the case in which the elements of the matrices \mathcal{M}_1 and \mathcal{M}_2 belong to the same $\mathbb{C} \subset \mathbb{O}$ subspace is an associative composition, that is it is equal to $(\mathcal{M}_2\mathcal{M}_1)\mathcal{X}(\mathcal{M}_1^{\dagger}\mathcal{M}_2^{\dagger})$. This is because each matrix element of $\mathcal X$ involves at most only one further complex subspace, and hence each multiplicative action on these elements in the linear transformation on $\mathcal X$ takes place in an associative quaternion subalgebra. Hence these particular cases of nested transformations do behave like a group representation. More generally however, and as for the case of SO(7) generated by composition of the ϕ_q maps with $q \in \text{Im}(\mathbb{O})$ in equation 6.11 as described in section 6.2, here there does not exist a group homomorphism of the full set of E_6 transformations into the set of 3×3 octonionic matrices $\mathcal{M}^{(a)}$.

However, associative group matrices could be constructed here by representing the linear transformations of the E_6 symmetry by 27×27 matrices in $GL(27, \mathbb{R})$ acting on the space $\mathbb{R}^{27} \equiv h_3 \mathbb{O}$, as was the case for SO(7) represented by matrices in GL(7, \mathbb{R}) acting on $\mathbb{R}^7 \equiv \text{Im}(\mathbb{O})$ in section 6.2. Indeed with such large matrices there is plenty of freedom in which to express the full symmetry with elements $R(g) \in GL(27, \mathbb{R})$ which naturally form an associate algebra and with $R(q_1)R(q_2) = R(q_1q_2)$ composing as a true representation of E_6 .

Given the 135 one-parameter subgroup actions on $h_3\mathbb{O}$, collectively implied in equations 6.32, 6.34 and 6.35, it would be straightforward, although laborious, to construct 135 matrices in $GL(27, \mathbb{R})$ acting upon \mathbb{R}^{27} , with the latter containing the 27 parameters of an element of $h_3\mathbb{O}$ in equation 6.1 drawn out into the real column vector $(p, m, n, a_1, \ldots, a_8)^T$. All such actions would preserve the cubic norm of equation 6.28 considered as a map $\mathbb{R}^{27} \to \mathbb{R}$. The multiplication of such elements of E₆ represented as matrices in $GL(27, \mathbb{R})$ is clearly associative, as only \mathbb{R} -valued matrices are involved. Together with the identity element given by the unit matrix 1_{27} and an inverse obtained for any matrix by reversing the transformation with real parameter $\alpha \to -\alpha$, the Lie group structure is evident. Combinations of the 135 one-parameter subgroup actions would carve out a $G \equiv E_6$ submanifold (for this non-compact real form of E_6) embedded within the (27×27) -dimensional space of GL $(27, \mathbb{R})$.

In principle left-invariant tangent vector fields, generated by right translations on G and associated with the one-parameter subgroups, could be constructed upon this E_6 group manifold, as depicted generically in figure 2.5 on a group manifold, and linear dependency used to reduce these to a basis set of 78 vector fields to describe the E_6 Lie algebra. Hence in this representation the Lie algebra may also be identified in terms of the transformation matrices themselves, in the form of elements $D = \partial A_{\lambda}/\partial \lambda|_{\lambda=0}$ as described before equation 6.3, here with $A \in GL(27, \mathbb{R})$. Alternatively the *left* translations of these symmetry transformations on \mathbb{R}^{27} may be associated with vector fields in the tangent space $T\mathbb{R}^{27}$ which also represent the Lie algebra generators of the symmetry. This construction applies generally (see also the discussion in the opening of section 4.3) – for example in the case of SO(3) acting on \mathbb{R}^3 the Lie algebra $L(SO(3))$ may be represented by vector fields in the space $T\mathbb{R}^3$ tangent to the 2-sphere S^2 .

This latter possibility of employing the left or right action of G on the representation space itself to construct the Lie algebra can be employed for the representation of E_6 of relevance here, that is on the space $h_3\mathbb{O}$. Indeed the theoretical motivation for studying E_6 here is precisely owing to its representation on $h_3\mathbb{O}$, together with the subgroup representations on subspaces of $h_3\mathbb{O}$ obtained under symmetry breaking, rather than the pure E_6 group structure in itself. It is the fact that the space h₃ σ with unit determinant has the appropriate structure for a form of temporal flow $L(\mathbf{v}_{27}) = 1$ that provides the primary motivation, with E_6 identified in turn as the corresponding symmetry group. This symmetry is expressed in a very compact cubic form as the determinant preserving actions on $h_3\mathbb{O}$, such as described in equation 6.29, and indeed the origin of the very high degree of symmetry, involving the triality relation for the largest division algebra O, is evident explicitly in this form. These structures would be far from manifest in a 27×27 real matrix representation. The non-associativity of the octonion algebra is employed in folding the full set of E_6 actions into this highly compact 3×3 matrix form.

For subgroups it will be possible to 'straighten-out' or unfold this action into familiar group representation form. This will be the case for the broken symmetry components, involving the external Lorentz group (in the form of the $SL(2,\mathbb{C})$ subgroup already described above) and internal symmetry groups, as we shall study in chapter 8 in comparison with the Standard Model of particle physics as reviewed in chapter 7. Since the representations of these subgroups are to be identified in the components of h₃O, which is ultimately motivated as the space underlying $L(v_{27}) = 1$, the tangent space $Th_3\mathbb{O}$ provides an apt arena for describing the E_6 Lie algebra.

The homomorphism of the Lie algebra $L(E_6) \equiv sl(3, \mathbb{O})$ into the space of vector fields in Th₃O, the tangent space to the 27-dimensional manifold of h₃O, is in fact an isomorphism since the group action of $SL(3, \mathbb{O})$ on $h_3\mathbb{O}$ is effective. This isomorphism is used both to identify individual E_6 generators and also, as described in the following section, the Lie algebra structure itself in terms of the commutators of the algebra elements. Clearly the broken subgroups also act effectively on the components of h_3 ^O and hence, following the discussion toward the end of section 4.3, the gauge field dynamics for the full internal symmetry group will be obtained. More generally the breaking of the isochronal symmetry of $L(v) = 1$ over the base space M_4 will ultimately need to be incorporated into the unification scheme described in section 5.1 with a structure in principle resembling Kaluza-Klein theory based on homogeneous fibres as reviewed in section 4.3.

With the space h₃^O considered as a manifold the map $R(\alpha)X_0$, for any point X_0 on h₃O, is a left action on the manifold with $R(0)\mathcal{X}_0 = \mathcal{X}_0$. This action also describes a curve as a mapping from $\alpha \in \mathbb{R}$ into h₃^① which sends the real number $\alpha = 0$ to the point \mathcal{X}_0 . Acting on all values of $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ this one-parameter group $R(\alpha)$ is associated with the tangent vector field:

$$
\dot{R} = \frac{\partial (R(\alpha)\mathcal{X})}{\partial \alpha}\Big|_{\alpha=0} \in T\mathbf{h}_3 \mathbb{O}
$$
 (6.36)

where a 'dot' over the kernel symbol such as for ' \dot{R} ' will generally denote a tangent vector field on the space $h_3\mathbb{O}$. The local tangent space on $h_3\mathbb{O}$ under the one constraint $\det(\mathcal{X}) = 1$ is 26-dimensional, however the space of vector fields over the 26-dimensional manifold S_{26} is infinite. The task is then to identify the E_6 Lie algebra through a one-to-one isomorphic correspondence with a subset of 78 linearly independent vector fields in Th₃ \mathbb{O} , of the form of equation 6.36, within this ∞ -dimensional space. The search is narrowed down by adopting a starting point based on the 135 one-parameter subgroup actions on h₃ \mathbb{O} obtained through the three types of $SL(2, \mathbb{O})$ conjugation described in equations 6.29 and 6.32–6.35.

The first stage, at the level of these one-parameter subgroups, is to find a convenient new basis for the 21 category 3 transverse rotations $R_{r,s}(\alpha)$ described in table 6.1. For each imaginary base unit q in figure 6.1 the three pairs of imaginary units each describing a quaternion subalgebra with q also form a right-handed 3-dimensional 'coordinate frame' with q. That is, for example with $q = i$, we have $jk = +i$, $k!j! = +i$ and $l\ddot{u} = +i$, matching the pairs listed in the top row of table 6.2.

$q \in \text{Im}(\mathbb{O})$	1 st pair	$2^{\rm nd}$ pair	$3^{\rm rd}$ pair
$\dot{\imath}$	j, k	kl, 1j	l, il
\dot{j}	k, i	il, kl	l, 2l
\boldsymbol{k}	i, j	$\ddot{\mathbf{u}}, \ddot{\mathbf{u}}$	l, k
kl	$j\!l, i$	j, \underline{i}	k, l
$\ddot{\mu}$	i, k	il, k	j, l
\dot{u}	kl, j	$k, \underline{j}l$	i, l
l	il, i	$j\!l, j$	k , k

Table 6.2: (Adopted directly from [38] p.107, table 4.2). The 3 right-handed quaternion subalgebras for each imaginary octonion base unit q ordered as $1st$, $2nd$ and $3rd$ (associated with the rotations R_{q1} , R_{q2} and R_{q3} respectively) as appropriate for the new basis for transverse rotations listed in equations 6.37–6.39.

For each choice of q the associated $1st$, $2nd$ and $3rd$ planes, from the same row of the table, are mutually orthogonal and rotated independently by $R_{q1}(\alpha)$, $R_{q2}(\alpha)$ and $R_{q3}(\alpha)$ respectively, where for example $R_{i1}(\alpha) = R_{j,k}(\alpha)$ by taking the appropriate pair, here $\{r, s\} = \{j, k\}$ from table 6.2, to construct the corresponding category 3 transverse rotation $R_{r,s}(\alpha)$ from table 6.1. Adopting the point of view of *active* transformations these individual plane rotations are in a clockwise sense about the q-axis for positive α and counterclockwise for negative α . They are then composed

together in the following combinations:

$$
A_q(\alpha) = R_{q1}(\alpha) \circ R_{q2}(-\alpha) \tag{6.37}
$$

$$
G_q(\alpha) = R_{q1}(\alpha) \circ R_{q2}(\alpha) \circ R_{q3}(-2\alpha) \tag{6.38}
$$

$$
S_q(\alpha) = R_{q1}(\alpha) \circ R_{q2}(\alpha) \circ R_{q3}(\alpha) \tag{6.39}
$$

Since in all cases each of the two or three plane rotations are independent of each other their order may be interchanged. (The three actions A_q , G_q and S_q may also be recombined to recover the original single plane rotations, for example $R_{q1}(\alpha)$ = $A_q(\alpha/2) \circ G_q(\alpha/6) \circ S_q(\alpha/3)$.

The $(3 \times 7) = 21$ actions defined in equations 6.37–6.39 hence provide a new basis for the Spin(7) transverse rotations applied in table 6.1 on the space $h_2\mathbb{O}$. Since each of these actions is represented by diagonal 2×2 matrices they also apply to the Spin(7) action on the space $\mathbb O$ itself, as the double cover of SO(7) acting on Im($\mathbb O$). However the mathematical motivation for introducing the new basis is seen when applied to the 3×3 matrix case, implicitly due to the triality relation between the three octonion components of $h_3\mathbb{O}$. Indeed when embedded in the type 1, 2 and 3 actions of equations 6.32, 6.34 and 6.35 respectively and determining the tangent vectors of the new transverse rotations in $Th_3\mathbb{O}$ using equation 6.36 it can be shown by direct comparison that:

$$
\dot{A}_q^1 = \dot{A}_q^2 = \dot{A}_q^3 \tag{6.40}
$$

$$
\dot{G}_q^1 = \dot{G}_q^2 = \dot{G}_q^3 \tag{6.41}
$$

$$
\dot{S}_q^1 + \dot{S}_q^2 + \dot{S}_q^3 = 0 \tag{6.42}
$$

for each of the seven cases of $q \in \{i, j, k, k\}, \mathcal{Y}, \mathcal{Y}, \mathcal{Y}, \mathcal{Y}\}$. The superscript a on tangent vectors, as for A^a here, will always denote the type and since raising such a vector to a power has no meaning the a is not placed inside brackets (in cases of ambiguity brackets will be used for the type index '(a)' as for $\mathcal M$ in equations 6.32–6.35). The new choice of equations 6.37–6.39 for the category 3 actions on $h_3\mathbb{O}$ is hence justified by the manifest clarity of the linear dependencies seen in this basis.

Each of the 14 independent generators $\{\dot{A}_q \equiv \dot{A}_q^a, \dot{G}_q \equiv \dot{G}_q^a\}$ (for any $a = 1, 2, 3$) acts on the three octonion elements $a, b, c \in \mathbb{O}$ in equation 6.1 in exactly the same way (while vanishing on the $p, m, n \in \mathbb{R}$ elements as for all transverse rotations). The 14 corresponding group actions of equations 6.37 and 6.38 preserve the multiplication table for $\mathbb O$ continuously as a function of the parameter α , forming the proper continuous automorphism group of the octonions (this group is SO(3) for the quaternion case). Hence, taken together A_q and G_q compose the exceptional group G_2 , which justifies the notation ' G_q ' introduced in equation 6.38.

The notation ' A_q ' in equation 6.37 is introduced owing to the similarity of the kernel symbol to ' λ ' which denotes the Gell-Mann matrices, as listed in table 8.5, which generate the Lie group $SU(3)$, a basis for the Lie algebra of which can also be composed of the 8 generators $\{\dot{A}_q, \dot{G}_l\}$, as will be described in section 8.2. In fact the automorphism group of the octonions may be reduced to the subgroup $SU(3) \subset G_2$ by fixing an imaginary unit such as $l \in \mathbb{O}$. The identification of this SU(3) subgroup also provides a significant motivation for adopting the basis of equations 6.37–6.39 from the potential physical perspective (see also the discussion following equation 6.58).

The notation ' S_q ' in equation 6.39 originates from the symmetric action of three synchronised rotations of α radians in three different planes. Applied to the 2×2 vector $X \in h_2\mathbb{O}$ this synchronised action is identical to the original single action of equation 6.25 (with i generalised to any $q \in \{i, j, k, k, \mathcal{J}, \mathcal{U}, \mathcal{U}\}\)$ in rotating three planes of the $a \in \mathbb{O}$ component of X, although due to the transformation of the spinor $\theta = \begin{pmatrix} c \\ b \end{pmatrix}$, as described below, the action of $S_i^{(1)}$ $i^{(1)}$ for example, that is with $q = i$ in equation 6.39, on h₃ \circ in the 3 \times 3 case is *not* equivalent to the action of equation 6.25 embedded in equation 6.29 or 6.32.

On the other hand equation 6.25 can be augmented to a single unnested 3×3 matrix action $R(\alpha)X = \mathcal{M}_{S_q}^{(a)}\mathcal{X}\mathcal{M}_{S_q}^{(a)\dagger}$ with $\mathcal{M}_{S_q}^{(a)}$, for the type $a = 1$ case, expressed as:

$$
\mathcal{M}_{S_q}^{(1)}(\alpha) = \begin{pmatrix} e^{q\frac{\alpha}{2}} & 0 & 0 \\ 0 & e^{q\frac{\alpha}{2}} & 0 \\ 0 & 0 & e^{-q\alpha} \end{pmatrix}
$$
 (6.43)

with a corresponding permutation of the diagonal entries for the type 2 and type 3 cases. These actions are denoted by kernel symbol S with the '\' as a mnemonic for the diagonal form of equation 6.43. The action $\mathcal{M}_{S_q}^{(1)}(\alpha) \mathcal{X} \mathcal{M}_{S_q}^{(1)\dagger}(\alpha)$ may be considered as a 'phase transformation' in rotating three orthogonal imaginary planes of the a component of X in equation 6.26 by the same angle α . These actions are also related to a demonstration of triality in h₃ Ω involving SO(8) transformations on the three octonion subspaces of $h_3\mathbb{O}$ (see the discussion below alongside equations 6.49 and 6.50 and in [39] around equation 43).

Each of these three 3×3 matrices, including the type 1 case in equation 6.43 for a given q, contains entries in a single complex subalgebra, satisfies $\det(\mathcal{M}_{S_q}^{(a)}) = 1$ and preserves the form $\det(\mathcal{X})$ of equation 6.27 or 6.28, consistent with the requirements for an $SL(3,\mathbb{O})$ action. However, these actions will not lead to elements of the preferred E_6 algebra basis under construction here since they are not of the form of equations 6.32– 6.35 with $\det(M) = \pm 1$ as required for the 'compatible' 2×2 matrix actions described in the previous section. Given the form of M_{xq} in table 6.1 and the type 1, 2 and 3 embeddings of equations 6.32, 6.34 and 6.35 the diagonal matrix of equation 6.43 can be expressed by the matrix product:

$$
\mathcal{M}_{S_q}^{(1)}(\alpha) = \mathcal{M}_{R_{xq}}^{(1)}(-\alpha) \times \mathcal{M}_{R_{xq}}^{(2)}(-2\alpha) \tag{6.44}
$$

and hence
$$
\dot{S}_q^1 = -\dot{R}_{xq}^1 - 2\dot{R}_{xq}^2
$$
 (6.45)

The group action $S_q^{(1)}(\alpha)$ of equation 6.39 on the 10-dimensional subspace $h_2\mathbb{O} \subset h_3\mathbb{O}$, consisting of three independent rotations of imaginary planes of the octonion a, is precisely the same as the $S_q^{(1)}(\alpha)$ action using equation 6.43. On the spinor components $\theta = \begin{pmatrix} \frac{c}{b} \end{pmatrix} \in \mathbb{Q}^2 \subset h_3\mathbb{O}$ these actions are only equivalent for small transformations to order α and diverge at $O(\alpha^2)$ and higher powers. However since all 27 components of h₃ \mathbb{O} transform the same way to $\mathcal{O}(\alpha)$ we have $\dot{S}_q^1 = \dot{S}_q^1$ (and similarly for the type 2 and 3 cases) as vector fields in the space $Th_3\mathbb{O}$, and hence these two objects are interchangeable in expressions of linear dependence.

Since at the group level $S_q^{(a)}(\alpha)$ and $S_q^{(a)}(\alpha)$ differ at $O(\alpha^2)$ the Lie bracket, to be described in the following section, of the corresponding generators with the same Lie algebra element \dot{R} will also differ with $[\dot{S}, \dot{R}] \neq [\dot{S}, \dot{R}]$ in general even though $\dot{S} = \dot{S}$ as elements of a vector space. The Lie bracket in these two cases agrees for the $h_2\mathbb{O} \subset h_3\mathbb{O}$ subspace but differs for the spinor components. The transformations on the components of the spinor θ are expected to be important for the internal symmetries in comparison with the Standard Model and hence care will need to be taken in choosing an appropriate Lie algebra basis. The E_6 Lie algebra table in [38] uses the actions of $S_q^{(1)}(\alpha)$ from equation 6.39 rather than $S_q^{(1)}(\alpha)$ based in equation 6.43, which is hence significant for the Lie algebra structure, and in turn it is the former transformations which are also used in this paper.

Adding the 7 actions $S_q^{(a)}$ for $a = 1, 2$ or 3 to the set of 14 actions $\{A_q, G_q\} \equiv \text{G}_2$ completes a $Spin(7)$ double cover of $SO(7)$ for the type 1, 2 or 3 transverse rotations respectively. These three $SO(7)$ s are mutually related by equation 6.42. In addition to equations 6.40–6.42 further linear dependencies amongst the generators expressed on $T_{\rm h_3}$ O are found (the second of which is equivalent to equation 6.45):

$$
\dot{R}_{xq}^1 + \dot{R}_{xq}^2 + \dot{R}_{xq}^3 = 0 \tag{6.46}
$$

$$
\dot{R}_{xq}^2 = -\frac{1}{2}\dot{R}_{xq}^1 - \frac{1}{2}\dot{S}_q^1 \tag{6.47}
$$

$$
\dot{S}_q^2 = +\frac{3}{2}\dot{R}_{\text{sq}}^1 - \frac{1}{2}\dot{S}_q^1\tag{6.48}
$$

Appending the set of 7 actions $R_{\underline{xq}}^{(a)}$ to the SO(7) of type a (for $a = 1, 2$ or 3) completes a set of 28 actions forming the group SO(8). The three SO(8)s formed this way are actually the *same* $SO(8)$, that is they are composed of the same subset of E_6 transformations on h₃ \mathbb{O} , due to the triality relation between the h₃ \mathbb{O} components. The triality symmetry is described explicitly in [38, 39, 40]. The transformations of the $SO(8)$ subgroup of the type 1 $SL(2, \mathbb{O})$ action in equation 6.32 can be obtained by a nested composition with 3×3 matrices of the form:

$$
\mathcal{M} = \begin{pmatrix} q & 0 & 0 \\ 0 & \overline{q} & 0 \\ 0 & 0 & 1 \end{pmatrix}
$$
 (6.49)

with $q \in \mathbb{O}$ and $|q| = 1$. The action of such type 1 transformations on an element $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ of equation 6.1 leaves the diagonal elements $\{p,m,n\}$ invariant while the three off-diagonal octonion elements transform non-trivially as (see [40] equation 46 and discussion):

$$
a \rightarrow \overline{q}a\overline{q}
$$

\n
$$
b \rightarrow bq
$$

\n
$$
c \rightarrow qc
$$

\n(6.50)

These generate and correspond to the three SO(8) 8-dimensional representations of vector, dual spinor and spinor exhibited via symmetric, right and left and octonion

multiplication respectively, with an implicit triality mapping between the above three octonion actions identified by simply employing the same q for each of the three actions, as alluded to in the opening paragraphs of section 6.1. Corresponding to the triality isomorphism the three actions of SO(8) are permuted into each other via the action of the matrices $\mathcal T$ in equation 6.33, such that we effectively have the *same* copy of $SO(8)$ in common within each of the three types of $SL(2,\mathbb{O})^a$ actions on h₃ \mathbb{O} .

This subgroup $SO(8) \subset E_6$ is in fact precisely the subgroup of E_6 transformations on $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ that leaves invariant the diagonal entries, that is $\{p,m,n\}$ of equation 6.1. This unique $SO(8)$ then contains three different $SO(7)$ s, each built in turn on a unique G_2 . Only this subset of 14 G_2 transformations needs to be described in the form of nested actions while the remaining $SO(8)$ transformations may be composed of seven unnested actions from $S_q^{(a)}$ (as for example from equation 6.43, and replacing the nested $S_q^{(a)}$ actions to obtain Spin(7) from G_2) together with seven $R_{\mathbf{x}\mathbf{q}}^{(a)}$ actions for type $a = 1, 2$ or 3 [39].

Here in this paper the initial importance of triality lies in the fact that it explains in part the rich symmetry of E_6 on h₃^O as an expression of $L(\mathbf{v}) = 1$. Indeed the triality symmetry is responsible for the large degree of redundancy in the set of $(3\times 45)=135$ generators for three types of $\mathrm{SL}(2,\mathbb O)^a$ transformation described above. The relations in equations 6.46–6.48 show that given the type 1 actions it is possible to exclude \dot{S}_q^2 (and hence, from equation 6.42, also \dot{S}_q^3) as well as \dot{R}_{xq}^2 and \dot{R}_{xq}^3 from a linearly independent basis for the E_6 Lie algebra.

Building on the 28 generators of $SO(8)$ (taking a type 1 basis) with any one of the three sets of 8 generators $\{\dot{R}_{xz}^a, \dot{R}_{zq}^a\}$, for type $a = 1, 2$ or 3, leads to one of three copies of SO(9). Each of these $28 + (3 \times 8) = 52$ rotations preserves $tr(\mathcal{X})$, with $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$, and they collectively define the group $F_4 := SU(3, \mathbb{O})$. The trace on $h_3\mathbb{O}$ is analogous to the time component of the Lorentz vector represented by $h_2\mathbb{K}$ (described after equation 6.21 for equations 6.16 and 6.19), but does not itself have a simple temporal interpretation here for the 3×3 case. In the present theory the cubic norm $\det(\mathcal{X}) \equiv L(v_{27}) = 1$ itself expresses a multi-dimensional form of temporal flow, having the form of equation 2.9 as introduced in section 2.1.

Extending further to reproduce the type $a = 1, 2$ and 3 Lorentz transformations $SL(2,\mathbb{O})^a$ by including the 9 boost generators $\{\dot{B}^a_{tz}, \dot{B}^a_{tz}, \dot{B}^a_{tq}\}\$ for each case, and taking into account the further linear dependence:

$$
\dot{B}_{tz}^1 + \dot{B}_{tz}^2 + \dot{B}_{tz}^3 = 0 \tag{6.51}
$$

a total of $52 + (3 \times 9) - 1 = 78$ actions are obtained, accounting for a complete basis of determinant preserving $E_6 := SL(3, \mathbb{O})$ transformations of h₃ \mathbb{O} . The entire group is then described in terms of the actions of complex matrices $\mathcal M$ on the space h₃ $\mathbb O$, with the preferred basis for the Lie algebra represented on $Th_3\mathbb{O}$ reproduced below in table 6.3.

The generators, as described above, of the subalgebras corresponding to the various stages of the subgroup chain:

$$
E_6 \supset SO^+(1,9)_{45} \supset SO(9)_{36} \supset SO(8)_{28} \supset SO(7)_{21} \supset (G_2)_{14} \supset SU(3)_8 \qquad (6.52)
$$

(here, other than for the 78-dimensional E_6 , the subscripts give the dimension of the algebra) can be identified within the three type 1 lines of table 6.3. These can be built

Category 1: Boosts			#
$\dot{B}^1_{\underline{t}z}$	$\dot B^1_{tx}$	$\dot{B}^1_{{\bf \textit{\textbf{L}}q}}$	9
\dot{B}_{tz}^2	\dot{B}_{tx}^2	\dot{B}_{tq}^2	9
	\dot{B}_{tx}^3	\dot{B}^3_{ta}	8
	Category 2: Rotations		
\dot{R}_{xq}^1	\dot{R}^1_{xz}	\dot{R}^1_{zq}	15
	\dot{R}^2_{xz}	\dot{R}_{zq}^2	8
	\dot{R}^3_{xz}	\dot{R}_{zq}^3	8
Category 3: Transverse Rotations			
\AA_q	\dot{G}_q	\dot{S}^1_a	21
		Total Generators	78

Table 6.3: The complete basis for the Lie algebra of E_6 , in terms of tangent vector fields on $Th_3\mathbb{O}$, reproduced from ([38] p.177, table A.1). The actual tangent vector fields are determined and listed in tables 6.6 and 6.7 in the present paper at the end of the following section.

up from $su(3)_8 \equiv \{\dot{A}_q, \dot{G}_l\}$ to $so^+(1, 9)_{45}$ which includes $\{\dot{A}_q, \dot{G}_q\}$ together with all of the type 1 generators in table 6.3.

The rotation subgroup of E_6 , as the compact real form of $F_4 := SU(3, \mathbb{O})$, is generated by the 52 category 2 and 3 transformations in table 6.3. The generator composition of a subalgebra chain leading down from $E_6 \supset F_4$ is presented in ([38] p.119, table 4.4). However, although both preserving $tr(\mathcal{X})$ (for any $\mathcal{X} \in h_3(\mathbb{D})$) and being the automorphism group of the exceptional Jordan algebra (equation 6.2), the group F_4 is not of great significance in the present paper.

At the group level in equation 6.52 each 'SO' might more strictly be replaced by the corresponding double cover 'Spin' group. As the group $SL(3,\mathbb{Q})$ necessarily includes the one-sided spinor actions $\theta \to M\theta$ in equation 6.32 (as well as in equations 6.34 and 6.35) the action for $M = -1₂$ (obtained for any of the category 2 rotations in table 6.1 with $\alpha = 2\pi$) on h₃^O does not give the identity transformation. However SL(3, 0) is not a double cover, rather it is a real simply connected form of $E_6 := SL(3, \mathbb{O})$ itself ([40] section 2, with the same situation applying for $F_4 := SU(3, \mathbb{O})$ acting on h₃ \mathbb{O}). On the other hand the action of $SL(2, \mathbb{O})$ is a double cover of the rotation group $SO^+(1,9) \equiv SL(2,0)/\mathbb{Z}^2$, and similarly for the further rotation subgroups. With an awareness of these issues of group manifold topology groups such as $SO^+(1,9)$ and $SO^+(1,3)$ can be considered to be embedded within the full group E_6 .

6.5 Lie Algebra of E_6

At the group level the E_6 action on $h_3\mathbb{O}$ is composed of 52 rotations, that is the unitary 3×3 matrix actions with $\mathcal{MM}^{\dagger} = \mathbf{1}_3$, and 26 boosts, that is the Hermitian actions with $M = M^{\dagger}$, as can be deduced from the embedded 2×2 matrices M listed in table 6.1 for the category 2 and 3 rotations and category 1 boosts respectively. At the Lie algebra level, in a normalised basis for which the Killing metric K is diagonal with entries in $\{-1, +1\}$ (or more generally negative or positive entries for a diagonal but unnormalised Killing form, explicit values for which will be determined in subsection 8.3.1), base vectors X for which $K(X, X) = -1$ (or $\lt 0$) are called compact generators, corresponding to 'rotations' of the Lie group, while those with $K(X, X) = +1$ (or > 0) are called non-compact generators, corresponding to 'boosts'.

With Killing form signature of -26 , also denoted (52, 26) for 52 rotations and 26 boosts, the non-compact real form of E_6 constructed in the previous section may be denoted as $E_{6(-26)}$, and describes the generator composition $D^{R,B}$ introduced in section 6.1 and displayed in equation 6.4. The Killing form employed in equations 4.1 and 4.2 of section 4.1 for Kaluza-Klein theory was chosen with components $K_{\alpha\beta} = -\delta_{\alpha\beta}$ corresponding to the choice of a compact gauge group. In the symmetry breaking of $E_{6(-26)}$ over the base space M_4 such compact internal symmetry groups will be identified.

An alternative description of E_6 in terms of 14 G_2 actions together with 64 non-G₂ transformations, composed from the actions of the 64 tracefree octonion 3×3 matrices, was also introduced in section 6.1 where $D^{G,S}$ denoted the generator composition as displayed in equation 6.5. In the previous section the G_2 subgroup was identified explicitly as the set of 14 $\{A_q, G_q\}$ transverse rotations. As described shortly after equation 6.50 the remaining 64 actions may be expressed with unnested compositions consisting for example of the 57 group actions corresponding to the category 1 and 2 generators of table 6.3 together with seven S_q^1 actions from equation 6.43 (in place of S_q^1). Hence both the (52+26) and (14+64) decompositions can be clearly seen in table 6.3 in terms of the respective subsets of generators.

Here all 78 generators are explicitly presented in tables 6.6 and 6.7, for the category $\{1, 2\}$ and 3 transformations respectively, as vector fields $R \in Th_3\mathbb{O}$ which, from equation 6.1, are of the form:

$$
\dot{R} = \begin{pmatrix} \dot{p} & \dot{\bar{a}} & \dot{c} \\ \dot{a} & \dot{m} & \dot{\bar{b}} \\ \dot{\bar{c}} & \dot{b} & \dot{n} \end{pmatrix} \in T\mathbf{h}_3\mathbb{O}
$$
 (6.53)

These 78 matrices are themselves Hermitian and hence also belong to the space h₃O. While there is no constraint on the determinant of any $\hat{R} \in Th_3\mathbb{O}$ the matrices are tracefree for all of the category 2 rotations and category 3 transverse rotations. The type 1 transformations act on the $\{p, m, n; a, b, c\}$ components on $h_3\mathbb{O}$ in the same way that the type 2 transformations act on the $\{m, n, p; b, c, a\}$ components and type 3 transformations act on the $\{n, p, m; c, a, b\}$ components as can be seen in equations 6.32, 6.34 and 6.35, for example by following the explicit invariant components n, p and m respectively in these three equations. This same cyclic permutation, consistent with the action of $\mathcal T$ in equation 6.33, is reflected in the tangent vectors in table 6.6 and for \dot{S}_q^a in table 6.7.

These tables describe in intimate detail the anatomy of the E_6 action as expressed on the tangent space Th₃^O. With $p = t + z$ and $m = t - z$, embedding equation 6.16 into h₃ \mathbb{O} , each type 1 tangent vector can be seen to 'point' in the appropriate direction in the relevant $Th_2\mathbb{O}$ components for the subspace $h_2\mathbb{O}$ plane transformations resulting from the action of the matrices in table 6.1, with a similar correspondence identifiable for the type 2 and 3 cases. For example the non-zero components of the category 1 boost and category 2 rotation generators for the type 1 actions on the 10-dimensional subspace $h_2 \mathbb{O}$ are simply:

$$
\dot{B}_{1z}^1 : \dot{t} = +z, \ \dot{z} = +t, \qquad \dot{B}_{1x}^1 : \ \dot{t} = +x, \ \dot{x} = +t, \qquad \dot{B}_{1q}^1 : \ \dot{t} = -a_q, \ \dot{a}_q = -t, \n\dot{R}_{2q}^1 : \ \dot{x} = -a_q, \ \dot{a}_q = +x, \qquad \dot{R}_{2z}^1 : \ \dot{x} = -z, \ \dot{z} = +x, \qquad \dot{R}_{2q}^1 : \ \dot{z} = +a_q, \ \dot{a}_q = -z
$$
\n(6.54)

where, here and in tables 6.6 and 6.7, $x \equiv a_x \equiv a_1$ and similarly a_q refers to the real coefficient in equation 6.6 corresponding to the imaginary unit q, (that is $a_l \equiv a_8$ etc.). The category 3 transverse rotations of equations 6.37, 6.38 and 6.39 each act on several planes in h₂ \mathbb{O} . The transformations of the spinor components of h₃ \mathbb{O} induced by the 3×3 matrix action are also included in tables 6.6 and 6.7.

In subsection 2.2.2 the Lie algebra of a group G was defined in terms of the set of left-invariant vector fields on the group manifold G , as also recalled in the paragraphs leading to equation 6.36 in the previous section. Through any point $h \in G$ each such vector field X generates a one-parameter group of right translations $\phi_t(h) = h \exp(tA)$ where $A = X_e \in T_e G$ is the vector of the field X at the identity $e \in G$, as depicted in figure 2.5 . If G acts by *right* translation on another manifold M this realisation of G induces vector fields $V^A \in TM$ such that $V_x^A(f) := \frac{d}{dt} f(x \exp(tA))|_{t=0}$, at $x \in M$ where $f(x)$ is a real function on M, represents a homomorphism of the Lie algebra with $[V^A, V^B] = V^{[A,B]}$. If the action of G on M is effective there is a one-to-one isomorphism between the Lie algebra $L(G)$ and the set of such vector fields ${V^A}$ in TM (as is the case for the action of G on a principle fibre bundle P as described in section 3.1, see equations 3.2 and 3.3).

If G acts on the manifold M by *left* translations then this relationship is an anti-homomorphism. This is the case for right-invariant vector fields on G itself, which are generated by left translations. The structure constants $c^{\alpha}{}_{\beta\gamma}$ for the Lie bracket of such right-invariant fields $\{Y_{\alpha}^{L}\}\$ on G are precisely the negative values of the Lie algebra structure constants defined in terms of the corresponding left-invariant fields ${X^R_\alpha}$ (which match the right-invariant fields as elements of the tangent vector space at the identity $e \in G$, that is each $Y_\alpha^L(e) = X_\alpha^R(e)$. This anti-homomorphism was also noted for left translations applied to the space of homogeneous fibres for equation 4.19 in the opening of section 4.3.

In the present case the group manifold for $G = \mathrm{E}_6$ is not constructed itself but rather the group acts *transitively* on $h_3\mathbb{O}$, which is hence a homogeneous space, such that det(\mathcal{X}) = 1 is preserved for $\mathcal{X} \in \mathrm{h}_{3}\mathbb{O}$. The action of E₆ on the underlying space h₃ \mathbb{O} is also *effective* and hence the Lie algebra $L(E_6)$ may be constructed in terms of vector fields on the tangent space Th₃O. The E₆ transformations composed as $\mathcal{X} \rightarrow$ $M\mathcal{X}M^{\dagger}$ are left translations as opposed to *right* translations, as has been described

in the previous section, and as will be seen explicitly for subgroups such as $SL(2,\mathbb{C})$ in section 8.1. The Lie algebra commutator, which determines the structure constants of the E_6 Lie algebra, for any two elements \dot{R}_1, \dot{R}_2 is defined through the action of the respective one-parameter subgroups $R_1(\alpha)$ and $R_2(\alpha)$ at any point $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$:

$$
[\dot{R}_2, \dot{R}_1] = \left. \frac{\partial}{\partial(\alpha^2)} [R_2(-\alpha) \circ R_1(-\alpha) \circ R_2(\alpha) \circ R_1(\alpha) \mathcal{X}] \right|_{\alpha=0} \tag{6.55}
$$

Here the four $\pm \alpha$ signs inside the square brackets are chosen so that this Lie algebra structure deriving from left translations is isomorphic to the standard definition of $L(G)$ of equation 2.22 described in subsection 2.2.2. In the general case for a Lie group G equation 6.55 holds with the *opposite* signs for α in the square brackets for the *right* translation mapping of one-parameter subgroup curves $\mathbb{R} \to G$ to the manifold of the Lie group space itself. These curves passing through the identity point $e \in G$ allow a bracket to be constructed on the vector space T_eG isomorphic to the Lie algebra of the group, leading for example to the basis of equation 2.32 for the case $G = SO(3)$.

In acting upon a representation space with a lower dimension than G , as is the case here for the group E_6 acting in the space $h_3\mathbb{O}$, the Lie bracket is constructed necessarily in terms of vector *fields* on the representation space. The choice of signs in equation 6.55 means that the various subalgebras will be defined in the usual way, equivalent to left-invariant fields on the broken subgroup manifolds. Indeed in principle the same E_6 Lie algebra could be constructed in terms of left-invariant fields on the submanifold of GL $(27,\mathbb{R})$ identified as an E₆ group representation acting on \mathbb{R}^{27} as described in the previous section.

Here the term in square brackets on the right-hand side of equation 6.55 represents a curve that passes through any chosen point $\mathcal{X}_0 \in \mathrm{h}_3\mathbb{O}$ for $\alpha = 0$. While the first derivative $\frac{\partial}{\partial \alpha}$ of this same term vanishes identically at $\alpha = 0$ the second derivative $\left(\frac{1}{2}\right)$ 2 $\frac{\partial^2}{\partial \alpha^2}$, or equivalently $\frac{\partial}{\partial (\alpha^2)}$ is non-zero and yields a tangent vector field as $\mathcal X$ varies over $h_3\mathbb O$ corresponding to the Lie bracket of the two vector fields \dot{R}_1 and \dot{R}_2 . For example, by direct calculation taking the type 1, 2 or 3 embeddings of the appropriate matrix actions from table 6.1, applying equation 6.55 and by comparison with tables 6.6 and 6.7 the twelve brackets listed in table 6.4 are determined explicitly.

1) $[\dot{R}_{x\dot{x}}^1, \dot{R}_{xz}^1] = \dot{R}_{z\dot{x}}^1$ (5) $[\dot{R}_{x\dot{x}}^1, \dot{B}_{tx}^1] = -\dot{B}_{t\dot{x}}^1$	9) $[\dot{R}_{xi}^1, \dot{R}_{xl}^1] = -\frac{1}{3}\dot{G}_{\underline{y}} + \frac{1}{3}\dot{S}_{\underline{y}}^1$
2) $[\dot{R}_{\dot{x}i}^1, \dot{R}_{zi}^1] = -\dot{R}_{xz}^1$ 6) $[\dot{R}_{\dot{x}i}^1, \dot{R}_{xz}^2] = -\frac{1}{2}\dot{R}_{zi}^2$	10) $[\dot{R}_{xi}^1, \dot{R}_{xi}^1] = \frac{1}{2}\dot{A}_k + \frac{1}{6}\dot{G}_k + \frac{1}{3}\dot{S}_k^1$
3) $[\dot{B}_{tz}^1, \dot{B}_{tx}^1] = \dot{R}_{xz}^1$ 7) $[\dot{R}_{zi}^1, \dot{B}_{ti}^3] = -\frac{1}{2}\dot{B}_{tx}^2$	$ 11 \rangle [\dot{S}_i^1, \dot{R}_{xj}^1] = \dot{R}_{xk}^1$
4) $[\dot{B}_{tz}^1, \dot{B}_{\dot{u}}^1] = -\dot{R}_{z\dot{u}}^1 \mid 8] [\dot{R}_{xz}^2, \dot{R}_{z\dot{d}}^2] = \dot{R}_{x\dot{d}}^2$	12) $[\dot{R}_{xz}^3, \dot{R}_{z\dot{z}}^3] = \dot{R}_{x\dot{z}}^3$
	$=-\frac{1}{2}\dot{R}_{\text{nl}}^1-\frac{1}{2}\dot{S}_l^1$ $=-\frac{1}{2}\dot{R}_{\text{nl}}^1+\frac{1}{2}\dot{S}_i^1$

Table 6.4: The Lie algebra bracket composition determined for twelve cases by applying equation 6.55 to the left-hand sides and tracking terms to $O(\alpha^2)$ for the sequence of $R(\pm\alpha)\chi$ compositions and matching the right-hand sides with elements in tables 6.6 and 6.7.

All cases in table 6.4 were calculated in full with two exceptions: in 'case 10)' the b component only on the right-hand side was determined and for 'case 11)' the a

component only was determined and the action $S_i^{(1)}$ $i^{(1)}(a)$ was used in place of $S_i^{(1)}$ $\binom{1}{i}(\alpha)$ in the calculation since these actions are identical on the $h_2\mathbb{O} \subset h_3\mathbb{O}$ subspace, as described in the discussion following equation 6.43. The purpose of these calculations is to cross-check the notation and conventions used here. This is useful since there are several sign differences between quantities in this paper and the corresponding expressions in reference [38] as listed for example in table 6.5. (We also note that the conventions used in the present paper differ in the sign of $\pm \alpha$ for $R_{zq}(\alpha)$ and $B_{tq}(\alpha)$ with respect to $([41]$ table 1)).

Action:	$R_{\mathbf{z}\mathbf{q}}(\alpha),$	$R_{xz}(\alpha),$	$R_{zq}(\alpha)$
Sign:	$e^{\pm q\frac{\alpha}{2}}$	$\pm \sin \frac{\alpha}{2}$,	$\pm q \sin \frac{\alpha}{2}$

Table 6.5: Sign differences between table 6.1 in the present paper and ([38] p.90, table 3.1).

Of the 78 basis tangent vectors listed here in tables 6.6 and 6.7 one is explicitly presented in reference [38]. The calculation of this tangent vector, namely for \dot{A}_{l} in table 6.7 here, differs by an overall \pm sign from that presented in ([38] p.112, equation 4.1), but agrees with the sign convention for the same components quoted on ([38] p.121). There is also a factor of two difference between the expression for $[\dot{R}_2, \dot{R}_1]$ displayed here in equation 6.55 and that described in the equations of ([38] p.109).

Whether each of these discrepancies is due to a typographical error or the conventions used in [38] the choice of signs and factors adopted in this paper is necessary in order that the calculations here in table 6.4 are both self-consistent and agree with the corresponding twelve entries in the full E_6 Lie algebra commutation table available in [38] for which the full set of $(78 \times 78 - 78)/2 = 3003$ independent entries were found by computer program, and which is used for this paper particularly in chapter 8.

The references [37, 38, 39, 40, 41] are essential here for describing the anatomy of the E_6 action on $h_3\mathbb{O}$ in a tractable form which may be dissected for the analysis of symmetry breaking patterns. The few inconsistencies in the notation as described above may be accounted for and will not affect the conclusions for physics. In this paper these conventions have been tuned for internal consistency and to be able to consistently read off entries from the full $L(E_6)$ table [38] as the principal point of reference. This in turn means that the correspondence between the generators of subgroups of E_6 , such as an external Lorentz group or an internal $SU(3)$ gauge symmetry group, may not neatly match the conventions generally employed in physical theories, as will be seen in chapter 8, for example in equation 8.9. Hence ultimately a new basis for $L(E_6)$ may be desired as tuned through a foreknowledge of the details of the physical application in the context of the present theory.

In general applying equation 6.55 for any two basis vectors on $Th_3\mathbb{O}$ will itself result in a basis vector field as listed in table 6.3 (or tables 6.6 and 6.7) or a linear combination of such elements as is the case for the brackets numbered 8), 9), 10) and 12) in table 6.4 above. While elements such as $\dot{R}^1_{z_i}$ on the right-hand side of 'case 1)' in this table may be 'integrated up' to the group action $R_{zi}^{(1)}$ $\chi_{\mathbf{z}i}^{(1)}(\alpha)$ on $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ in general it is not straightforward to associate an element, or linear combination of elements, of the E_6 Lie algebra with a one-parameter action of the Lie group describing curves on $h_3\mathbb{O}$. This is
due to the non-associativity of the octonions and the necessary employment of a nested structure to describe the transverse rotations. This is hence unlike the case in general for Lie algebra elements defined on the tangent space of a group manifold such as $G = SO(3)$ which may be associated with Lie group elements by an 'exponential map', as described alongside figure 2.5 and exemplified in equation 2.49 for $SO(5)$. However, of interest here will be broken subgroups, such as the Lorentz group for 4-dimensional spacetime and the SU(3) colour symmetry which may be expressed without the above difficulties. (Again, alternatively, the full E_6 action could in principle be expressed in terms of $G \subset GL(27, \mathbb{R})$ actions on \mathbb{R}^{27} and the consequences of non-associativity and nested actions sidestepped completely).

In the full Lie algebra table [38] with the basis vectors listed in table 6.3 a total of six mutually commuting elements, that is with $[\dot{R}_2, \dot{R}_1] = 0$ for any pair of these six elements, can be identified as the set:

$$
\{\dot{B}_{tz}^1, \dot{B}_{tz}^2, \dot{R}_{xl}^1, \dot{A}_l, \dot{G}_l, \dot{S}_l^1\}
$$
\n(6.56)

which hence forms the Cartan subalgebra for the rank-6 Lie algebra E_6 . There is some flexibility in this choice, due for example to equation 6.51, with $(\dot{B}_{tz}^2 - \dot{B}_{tz}^3)$ replacing the second element \dot{B}_{tz}^2 in ([41] equation 3.8(15)).

In [39, 40] 'symmetry breaking' is considered in terms of making a choice of a preferred h₂ $\mathbb{O} \subset h_3 \mathbb{O}$ together with a preferred imaginary unit for the octonion element in h₂ \mathbb{O} . Here we take a subspace h₂ $\mathbb{C} \subset h_3 \mathbb{O}$, using the isomorphism of the 4-dimensional space $h_2\mathbb{C}$ to the space of the Lorentz vectors which in turn we have identified with tangent vectors on M_4 , the base space for our perception of objects in the world; as described in the previous chapters. The Lorentz group, $SO^+(1,3)$, is seen correspondingly as a non-compact subgroup of E_6 . With $SO^+(1,3)$ being an *external* symmetry on M_4 , which is also a global symmetry to a very good approximation in a laboratory setting, this will then provide the mechanism for the breaking of the E_6 symmetry down to local gauge symmetries which may be compared with the $SU(3)_c \times$ $SU(2)_L \times U(1)_Y$ gauge group and representations in the Standard Model of particle physics.

The Lorentz subgroup for 4-dimensional spacetime can be taken to be generated by the subset of Lie algebra elements in $L(E_6)$:

$$
\{\dot{B}_{1z}^1, \dot{R}_{z\perp}^1, \dot{B}_{1x}^1, \dot{B}_{1y}^1, \dot{R}_{zz}^1, \dot{R}_{z\perp}^1\} \tag{6.57}
$$

acting on
$$
\mathbf{h}_2 = \begin{pmatrix} t+z & x-yl \\ x+yl & t-z \end{pmatrix} \in h_2 \mathbb{C} \subset h_3 \mathbb{O}
$$
 (6.58)

where here the first two generators for this rank-2 subgroup are taken from the Cartan subalgebra for E_6 in equation 6.56. The octonion unit l of the component a in equation 6.1 (rather than i as for equation 6.19 and as discussed at the end of section 6.3) is chosen to represent an external spatial component since then the internal symmetry is more readily identified using the preferred basis of table 6.3, which in turn derived from the conventions of equations 6.37-6.39 and table 6.2 in which l only appears in the $3rd$ pair column. The use of the unit l, rather than i, in this way also serves as a reminder that the Lorentz transformations here are embedded within expressions based on the octonion algebra. This external Lorentz symmetry will be studied in detail in section 8.1.

In section 8.2 an internal symmetry will be provisionally defined here as any operation that fixes the external spacetime components (t, x, y, z) of equation 6.58 for any Lorentz 4-vector. This will include in particular the subgroup $SU(3) \equiv \{A_q, G_l\}$, which from table 6.2 and equations 6.37 and 6.38 leaves the l component invariant, highlighting the significance of this basis choice for physics. The full E_6 Lie algebra commutation table in [38] can be used to identify further internal symmetry groups, as we shall explore in chapter 8.

In the following chapter we first review the Standard Model, and in particular the relationship between the external and internal symmetries found there, before turning to the group E_6 in general in section 7.3 as a candidate for unification of these symmetries as employed in particle physics. Then in chapter 8 the detailed structure of the action of E_6 on h₃ \mathbb{O} , as reviewed in this chapter, will be applied to deduce the properties of the external Lorentz symmetry in relation to the complementary internal symmetry for the present theory.

Through the historical development from the real numbers to the complex numbers, continuing on through the quaternions to the octonions, composed then in 2×2 and further in 3×3 matrix form, the construction of E₆ as a determinant preserving action on $h_3\mathbb{O}$ has been presented as an expression of the symmetry of temporal flow in the form of $L(v_{27}) = 1$. It is of course possible that there may be other, higher-dimensional, forms for $L(v) = 1$, with yet higher symmetry groups that will have consequences for the physics of the world. Further generalisation should be, however, a well defined mathematical problem.

In chapter 9 higher-dimensional forms of temporal flow and the possible role of the largest exceptional Lie groups E_7 and E_8 will be considered. For such cases the E_6 symmetry will be an intermediary on the way up to, or operate in some way parallel to, the larger symmetries for the higher-dimensional forms of $L(v) = 1$. Even in this case, given that the richness of h₃ \mathbb{O} and its symmetries, as 3×3 matrices expressing the triality relation between three elements of the largest normed division algebra, the octonions, is much greater than that of Lorentz 4-vectors and the symmetry of 4-dimensional spacetime, we might still hope to uncover elements of empirically established physical structure in a study based on this E_6 symmetry, assuming that the overall conceptual framework that we are considering here broadly corresponds to the real physical world. This, in the very least, would provide a proof of principle for the conceptual scheme being developed in this paper.

\dot{B}_{Lz}^1	$\dot B_{tx}^1$	\dot{B}^1_{tq}
		$\begin{pmatrix} +p & 0 & +\frac{1}{2}c \\[0.4em] 0 & -m & -\frac{1}{2}\bar{b} \\[0.4em] +\frac{1}{2}\bar{c} & -\frac{1}{2}b & 0 \end{pmatrix} \qquad \begin{pmatrix} +a_x & \frac{1}{2}(p+m) & +\frac{1}{2}\bar{b} \\[0.4em] \frac{1}{2}(p+m) & +a_x & +\frac{1}{2}c \\[0.4em] +\frac{1}{2}b & +\frac{1}{2}\bar{c} & 0 \end{pmatrix} \; \begin{pmatrix} -a_q & \frac{1}{2}(p+m)q & +\frac{1}{2$
\dot{B}^2_{tz}	\dot{B}_{tx}^2	
		$\left(\begin{matrix} 0&+\frac{1}{2}\bar{a}&-\frac{1}{2}c\\ +\frac{1}{2}a&+m&0\\ -\frac{1}{2}\bar{c}&0&-n \end{matrix} \right) \qquad \left(\begin{matrix} 0&+\frac{1}{2}c&+\frac{1}{2}\bar{a}\\ +\frac{1}{2}\bar{c}&+b_{x}&\frac{1}{2}(m+n)\\ +\frac{1}{2}a&\frac{1}{2}(m+n)&+b_{x} \end{matrix} \right) \; \left(\begin{matrix} 0&-\frac{1}{2}cq&+\frac{1}{2}\bar{a}q\\ +\frac{1}{2}q\bar{c}&-b_{q}&\$
\dot{B}_{tz}^3	\dot{B}^3_{tx}	\dot{B}^3_{tq}
		$\begin{array}{ c c c c c c } \hline \begin{array}{cccc} -p & -\frac{1}{2}\bar{a} & 0 \\ -\frac{1}{2}a & 0 & +\frac{1}{2}\bar{b} \\ 0 & +\frac{1}{2}b & +n \end{array} & \hline \begin{array}{cccc} +c_x & +\frac{1}{2}b & \frac{1}{2}(n+p) \\ +\frac{1}{2}\bar{b} & 0 & +\frac{1}{2}a \\ \frac{1}{2}(n+p) & +\frac{1}{2}\bar{a} & +c_x \end{array} & \begin{array}{cccc} -c_q & -\frac{1}{2}qb & -\frac{1}{2}(n+p$
\overline{R}^1_{xq}		$\overline{R^1_{zq}}$
		$\begin{pmatrix} 0 & -a_q - a_x q & -\frac{1}{2}qc \\ -a_q + a_x q & 0 & +\frac{1}{2}q\overline{b} \\ +\frac{1}{2}\overline{c}q & -\frac{1}{2}bq & 0 \end{pmatrix} \begin{pmatrix} +a_x & -\frac{1}{2}(p-m) & +\frac{1}{2}\overline{b} \\ -\frac{1}{2}(p-m) & -a_x & -\frac{1}{2}c \\ +\frac{1}{2}b & -\frac{1}{2}\overline{c} & 0 \end{pmatrix} \begin{pmatrix} +a_q & \frac{1}{2}(p-m)q & -\frac{1}{2}q\overline{b} \\$
	\dot{R}_{xz}^2	
\dot{R}_{xq}^2		$\begin{pmatrix} 0 & +\frac{1}{2}\bar{a}q & -\frac{1}{2}cq \\ -\frac{1}{2}qa & 0 & -b_q-b_xq \\ +\frac{1}{2}q\bar{c} & -b_q+b_xq & 0 \end{pmatrix} \begin{pmatrix} 0 & +\frac{1}{2}c & -\frac{1}{2}\bar{a} \\ +\frac{1}{2}\bar{c} & +b_x & -\frac{1}{2}(m-n) \\ -\frac{1}{2}a & -\frac{1}{2}(m-n) & -b_x \end{pmatrix} \begin{pmatrix} 0 & +\frac{1}{2}cq & +\frac{1}{2}\bar{a}q \\ -\frac{1}{2}q\bar{$
		$\begin{pmatrix} \dot{R}_{xq}^3 & \dot{R}_{xq}^3 & \dot{R}_{zq}^3 \\ 0 & +\frac{1}{2}q\bar{a} & -c_q + c_xq \\ -\frac{1}{2}aq & 0 & +\frac{1}{2}\bar{b}q \\ -c_q - c_xq & -\frac{1}{2}qb & 0 \end{pmatrix} \begin{pmatrix} -c_x & -\frac{1}{2}b & -\frac{1}{2}(n-p) \\ -\frac{1}{2}\bar{b} & 0 & +\frac{1}{2}a \\ -\frac{1}{2}(n-p) & +\frac{1}{2}\bar{a} & +c_x \end{pmatrix} \begin{pmatrix} -c_q &$

Table 6.6: Vector fields on $Th_3\mathbb{O}$ generated by the 26 Category 1 Boosts and 31 Category 2 Rotations from table 6.3 in the form of equation 6.53 $(\dot{B}_{tz}^3, \dot{R}_{xq}^2, \dot{R}_{xq}^3$ are non-basis elements).

 $\dot{A}_i: \quad \dot{a} = -a_4j \quad +a_3k \quad +a_6k l \quad -a_5j l$ $\dot{A}_j: \quad \dot{a} = +a_4i \qquad -a_2k \qquad -a_7k$ +a₅il $\dot{A}_k: \quad \dot{a} = -a_3i + a_2j \qquad \qquad +a_7jl -a_6i l$ $\dot{A}_{\underline{M}}: \quad \dot{a} = -a_6 i + a_7 j \qquad -a_2 j l \qquad -a_3 i l$ $\dot{A}_{jl}: \quad \dot{a} = -a_5i \qquad -a_7k \quad +a_2k l \qquad +a_4i l$ $\dot{A}_{\underline{y}}: \quad \dot{a} = +a_5j +a_6k -a_3k^2 -a_4j^2$ $\dot{A}_l: \quad \dot{a} = -a_7i \quad -a_6j \quad +a_3jl \quad -a_2il$ $\dot{G}_i: \quad \dot{a} = -a_4j \quad +a_3k \quad -a_6k\ell \quad +a_5j\ell \quad -2a_8i\ell \quad +2a_7l$ $\dot{G}_j: \quad \dot{a} = +a_4i \qquad -a_2k \quad +a_7k\! \quad -2a_8j\! \quad -a_5i\! \quad +2a_6l$ \dot{G}_k : $\dot{a} = -a_3i + a_2j -2a_8k^2 -a_7j^2 +a_6i^2 +2a_5k^2$ \dot{G}_{kl} : $\dot{a} = +a_6i \quad -a_7j \quad +2a_8k \quad -a_2jl \quad +a_3il \quad -2a_4l$ $\dot{G}_{\underline{j}\underline{l}}: \quad \dot{a} = -a_5i + 2a_8j + a_7k + a_2k\underline{l}$ $-a_4i\underline{l} - 2a_3l$ $\dot{G}_{\underline{d}}: \quad \dot{a} = +2a_8i \quad +a_5j \quad -a_6k \quad -a_3k\! \quad +a_4j\! \quad -2a_2l$ $\dot{G}_l: \quad \dot{a} = +a_7i +a_6j -2a_5k +2a_4k^2 -a_3j^2 -a_2i^2$ $\dot{S}_{q}^{1}:$ $\sqrt{ }$ $\begin{matrix} \end{matrix}$ $\overline{}$ $\dot{a} = q \sum$ $r \neq 1, q$ $a_r r$ $\dot{b} = +\frac{3}{2}b_q - \frac{3}{2}b_1q - \frac{1}{2}q\sum$ $r \neq 1, q$ $b_r r$ $\dot{c} = -\frac{3}{2}$ $\frac{3}{2}c_q + \frac{3}{2}$ $\frac{3}{2}c_1q-\frac{1}{2}$ $\frac{1}{2}q\sum$ $r \neq 1, q$ $c_r r$ $\dot{S}_q^2 : \dot{a} = -\frac{3}{2}a_q + \frac{3}{2}a_1q - \frac{1}{2}q\sum a_r r$, $\dot{b} = q\sum b_r r$, $\dot{c} = +\frac{3}{2}c_q - \frac{3}{2}c_1q - \frac{1}{2}q\sum c_r r$ \dot{S}_q^3 : $\dot{a} = +\frac{3}{2}a_q - \frac{3}{2}a_1q - \frac{1}{2}q\sum a_r r$, $\dot{b} = -\frac{3}{2}b_q + \frac{3}{2}b_1q - \frac{1}{2}q\sum b_r r$, $\dot{c} = q\sum c_r r$

Table 6.7: Vector fields on $Th_3\mathbb{O}$ generated by the 21 Category 3 Transverse Rotations from the lower section of table 6.3. In the case of \dot{A}_q and \dot{G}_q the form of $\dot{b} = f(b)$ and $\dot{c} = f(c)$ is identical to $\dot{a} = f(a)$. With reference to equation 6.53, in all cases $\dot{p} = \dot{m} = \dot{n} = 0$ with $\{\dot{a}, \dot{b}, \dot{c}\}$ implied from $\{\dot{a}, \dot{b}, \dot{c}\}$. $(\dot{S}_q^2$ and \dot{S}_q^3 are non-basis elements, with $\sum := \sum_{r \neq 1,q}$ here).

Chapter 7

Review of the Standard Model

7.1 Lorentz Symmetry and Spinors

Having introduced the higher 27-dimensional form for the flow of time with symmetry group E_6 acting on $h_3\mathbb{O}$ in the previous chapter we shall address the embedding of the Lorentz symmetry $SO^+(1,3)$, acting on a 4-dimensional spacetime associated with the components of the subspace $h_2\mathbb{C} \subset h_3\mathbb{O}$, within the larger structure in the opening section of the following chapter. Here we first consider the properties of the group $SO⁺(1, 3)$ itself together with its representations.

In general symmetries implicit in the form $L(v) = 1$ may include rotation groups, such as $SO(3) \subset SO^+(1,3)$, which are significant due to their geometrical interpretation as employed in the construction of the background manifold for perception. These rotation groups may also be embedded within a wider set of elements belonging to the Clifford algebra associated with the (pseudo-) Euclidean space to which the rotations apply. These algebras also have spinor representations, which are as mathematically natural as the vector representations. For example, as alluded to in the opening paragraphs of chapter 6 and again explicitly in equation 6.50, the vector and spinor representations of SO(8) are equally significant for the structure and symmetry of $L(\mathbf{v}_{27}) = 1$.

Although most of the discussion below applies to Clifford algebras in general here we focus on the case of the 4-dimensional vector space $\mathbb{R}^{1,3}$, with real Clifford algebra $C(1,3)$ represented by 4×4 γ -matrices satisfying the relations:

$$
\gamma^a \gamma^b + \gamma^b \gamma^a = 2\eta^{ab} \mathbf{1}_4 \tag{7.1}
$$

with indices $\{a, b\} = 0 \dots 3$, Minkowski metric η^{ab} and where $\mathbf{1}_4$ denotes the 4×4 identity matrix. For any vectors $v, w \in \mathbb{R}^{1,3}$ the associated algebra product with $v = v_a \gamma^a$ and $w = w_b \gamma^b$ as elements of $C(1, 3)$ satisfies $vw + wv = 2\eta(v, w)\mathbf{1}_4$, implying for example the relation $v^2 = |v|^2 \mathbf{1}_4$ for all $v \in \mathbb{R}^{1,3}$ which is also sufficient to generate the full algebra. A general element u of the Clifford algebra $C(1,3)$ has the form:

$$
u = u_0 + u_a \gamma^a + u_{ab} \gamma^a \gamma^b + u_{abc} \gamma^a \gamma^b \gamma^c + \cdots
$$

$$
\in C(1,3) = C^0 \oplus C^1 \oplus C^2 \oplus C^3 \oplus \cdots
$$
 (7.2)

with $u_0, u_a, u_{ab} \dots \in \mathbb{R}$, with index values ordered as $a < b < c \dots$, and where C^i denotes the subspace of $C(1,3)$ formed by the product of i basis elements $\{\gamma^a\}$ in this representation. Owing to equation 7.1 the Clifford algebra itself has dimension 2^n , with $n = 4$ here, that is the elements of the $C(1,3)$ algebra describe a vector space with 16 linearly independent elements.

The Clifford algebra itself does not form a group since in general an inverse element may not exist for any given $u \in C(1,3)$. However, the elements belonging to the subset of $C(1,3)$ generated by elements $v \in C^1$ with $\eta(v,v) = \pm 1$ do have an inverse and upon composition generate a group denoted $Pin(1, 3)$. Further, given such elements $v \in C^1 \cap \text{Pin}(1, 3)$ the map ϕ_v from $w \in C^1$ into C^1 :

$$
\phi_v: w \rightarrow v w v^{-1} \tag{7.3}
$$

$$
= 2\eta(v, w)v^{-1} - wvv^{-1}
$$

$$
= \frac{2\eta(v, w)v}{\eta(v, v)} - w
$$
 (7.4)

is a reflection of w through the line containing the origin and v in the (psuedo-) Euclidean space $\mathbb{R}^{1,3}$. These reflections may be combined to describe a representation of $Pin(1,3)$ as orthogonal transformations on the space $\mathbb{R}^{1,3}$ (which is equivalent to $C^1(1,3)$ as a vector space). The application of Clifford algebra composition to induce representations of the rotation groups via equation 7.3 is similar to the use of the conjugation action for elements of a division algebra such as the octonions, described by equation 6.11 in section 6.2, also to represent rotations.

In fact the Lie group $\text{Pin}(1,3)$ is the two-to-one cover of the full Lorentz group $O(1, 3)$, which has four disconnected pieces. Restricting the elements of $Pin(1, 3)$ to those in the even subalgebra $C^{e}(1,3) := \{C^{i}(1,3); i \text{ even}\}\$ of equation 7.2 identifies the subgroup Spin(1,3), which has a representation on the space $\mathbb{R}^{1,3}$ as the group of special orthogonal transformations $SO(1, 3)$. In both cases these actions are two-to-one surjective homomorphisms π with:

$$
\pi: \quad \text{Pin}(1,3) \to \text{O}(1,3) \tag{7.5}
$$

$$
\pi: \quad \text{Spin}(1,3) \to \text{SO}(1,3) \tag{7.6}
$$

Hence the respective Lie algebras are isomorphic, for example $spin(1,3) = so(1,3)$. The part of the group $Spin(1,3)$ as a manifold connected to the identity is in fact 'simply connected' and is denoted $Spin^+(1,3)$, the two-to-one covering group of $SO^+(1,3)$ – which in turn is the part of the full Lorentz group (described above equation 6.22 for the $k = 2$ case) which preserves both the time and the space orientations, as well as the metric relations, of Lorentz 4-vectors.

The set of matrices:

$$
\sigma^{\mathbf{a}\mathbf{b}} = \frac{1}{4} (\gamma^a \gamma^b - \gamma^b \gamma^a) = \frac{1}{4} [\gamma^a, \gamma^b]
$$
\n(7.7)

with $a < b$ and $\gamma^a \gamma^b \in C^2(1,3)$, is isomorphic to the Lie algebra spin⁺(1,3) = so⁺(1,3) under matrix commutation of the six independent $\sigma^{\boldsymbol{\omega}}$ elements. This algebra generates group elements $R(\omega_{cd}) = \exp(\omega_{cd}\sigma^{cd})$ with $\omega_{cd} \in \mathbb{R}$ (summing over the set of six index pairs with $c < d$, in a similar way to the group actions described in equation 2.49). These describe $SO^+(1,3)$ vector transformations on the γ^a matrices themselves:

$$
\phi_R: \ \gamma^a \to R(\omega_{cd})\gamma^a R^{-1}(\omega_{cd}) \equiv (A^{-1})^a{}_b \gamma^b \tag{7.8}
$$

with $A \in SO^+(1,3)$, as well as the spinor representation of $SO^+(1,3)$ on 4-component Dirac spinors $\psi \in \mathbb{C}^4$:

$$
\psi \rightarrow \psi' = e^{\omega_{cd}\sigma^{cd}}\psi = R(\omega_{cd})\psi \tag{7.9}
$$

For a *complex* Clifford algebra in any dimension n this Dirac representation is irreducible. However for the real forms of these algebras with even $n = p+q$ the space of the Dirac representation for the group $Spin^+(p,q)$ decomposes into two halves, known as chiral (left and right) spinors, upon which inequivalent representations act. This may be shown by defining the matrix $\Gamma^5 := \gamma^1 \gamma^2 \dots \gamma^n \in C^n(p,q)$ which anticommutes with each γ^a and hence (by equation 7.7 for the general case) commutes with all elements of $Spin^+(p,q)$, and hence in turn by Schur's lemma the Dirac representation is reducible (unless Γ^5 is a proportional to the unit $2^{\frac{n}{2}} \times 2^{\frac{n}{2}}$ matrix, which is generally not the case).

In the case of (1, 3) spacetime, with 4×4 matrices γ^a acting on the elements $\psi \in \mathbb{C}^4$ of the spinor space, Γ^5 is denoted γ^5 and the usual convention is to take:

$$
\gamma^5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3 \qquad \text{for which} \qquad (\gamma^5)^2 = +\mathbf{1}_4 \tag{7.10}
$$

Due to the factor of i this object does not belong to the *real* Clifford algebra. However as a 4×4 matrix γ^5 does commute with each element of $Spin^+(1,3)$ and can be used to extract the chiral spinors ψ_L and ψ_R via the projection operators P_L and P_R :

$$
\psi_L = P_L \psi \qquad \text{with} \qquad P_L = \frac{1}{2} (1 - \gamma^5) \tag{7.11}
$$

$$
\psi_R = P_R \psi \qquad \text{with} \qquad P_R = \frac{1}{2} (1 + \gamma^5) \tag{7.12}
$$

By Schur's lemma this decomposition into left and right-handed spinors $\psi = \psi_L + \psi_R \in$ \mathbb{C}^4 is maintained under the 4×4 matrix actions of the group Spin⁺(1,3). Hence the Dirac representation is reduced into two $Spin⁺(1, 3)$ invariant and irreducible pieces called Weyl spinors. A suitable explicit representation for the γ -matrices is the Weyl basis with:

$$
\gamma^0 = \begin{pmatrix} 0 & +\mathbf{1}_2 \\ +\mathbf{1}_2 & 0 \end{pmatrix}, \qquad \gamma^a = \begin{pmatrix} 0 & +\sigma^a \\ -\sigma^a & 0 \end{pmatrix}, \qquad \gamma^5 = \begin{pmatrix} -\mathbf{1}_2 & 0 \\ 0 & +\mathbf{1}_2 \end{pmatrix}
$$
(7.13)

where each entry is a 2×2 matrix and the three Pauli matrices σ^a for $a = 1, 2, 3$ are included in the following set:

$$
\sigma^0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}
$$
(7.14)

In the γ -matrix basis of equation 7.13 the Spin⁺(1, 3) action of equation 7.9 can be expressed on the Weyl spinors $\psi_L, \psi_R \in \mathbb{C}^2$ simply as:

$$
\begin{pmatrix} \psi \\ \psi \end{pmatrix} = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix} \rightarrow \begin{pmatrix} R_L & 0 \\ 0 & R_R \end{pmatrix} \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix}
$$
(7.15)

For particle states chirality itself is an observable only for massless fermions, that is $m_f = 0$, in which case it is equivalent to the particle helicity.

The 'spin' group for the Clifford algebra of the real pseudo-Euclidean vector space $\mathbb{R}^{1,3}$ may also be approached directly via the group $SL(2,\mathbb{C})$, which is closely related to the representations R_L and R_R in equation 7.15. The 6-dimensional Lorentz Lie algebra $\text{so}^+(1,3)$ can be expressed in a conventional basis of anti-Hermitian rotation generators $\{J^1, J^2, J^3\}$ and Hermitian boost generators $\{K^1, K^2, K^3\}$ in terms of a 2×2 matrix basis for $sl(2, \mathbb{C})$ in the form:

$$
J^a = -\frac{i}{2}\sigma^a \qquad \text{and} \qquad K^a = -\frac{1}{2}\sigma^a \tag{7.16}
$$

for $a = 1, 2, 3$. The signs are chosen such that the following algebra commutators hold:

$$
[J^a, J^b] = \varepsilon^{abc} J^c \tag{7.17}
$$

$$
[K^a, K^b] = -\varepsilon^{abc} J^c \tag{7.18}
$$

$$
[J^a, K^b] = \varepsilon^{abc} K^c \tag{7.19}
$$

with $\varepsilon^{123} = +1$. In other conventions the signs may vary, and factors of $i = \sqrt{-1}$ may appear if J is defined to be Hermitian, as is the case in quantum mechanics in order to identify real observable quantities for angular momentum. In the standard treatment a general element of the group $SL(2, \mathbb{C})$ is represented by the 2×2 matrix:

$$
S = e^{r_a J^a + b_a K^a} = e^{\frac{1}{2}(-ir_a - b_a)\sigma^a}
$$
\n(7.20)

with the rotations parametrised by $r_a \in \mathbb{R}$, $a = 1, 2, 3$, and the boosts parametrised by $b_a \in \mathbb{R}$, $a = 1, 2, 3$. For the complex linear combinations $A^a = \frac{1}{2}(J^a + iK^a)$ and $B^a=\frac{1}{2}$ $\frac{1}{2}(J^a - iK^a)$ the Lie bracket reads:

$$
[A^a, A^b] = \varepsilon^{abc} A^c \tag{7.21}
$$

$$
[B^a, B^b] = \varepsilon^{abc} B^c \tag{7.22}
$$

$$
[A^a, B^b] = 0 \tag{7.23}
$$

demonstrating that the complexified Lie algebra of sl(2, \mathbb{C}) is isomorphic to su(2) \oplus su(2) (as will be represented in figure 7.2(d) and described in the accompanying text) which is used to label the representations of the Lorentz group by the half-integer values (j_A, j_B) . After the trivial $(0,0)$ scalar case the two lowest-dimensional possibilities are the representations of $SL(2, \mathbb{C})$ denoted $R_L(S)$ and $R_R(S)$ with:

$$
(j_A, j_B) = (\frac{1}{2}, 0) \Rightarrow R_L(S) = e^{\frac{1}{2}(-ir_a - b_a)\sigma^a}
$$
(7.24)

$$
(A^a = -\frac{1}{2}\sigma^a, B^a = 0; \qquad J^a = -\frac{i}{2}\sigma^a, K^a = -\frac{1}{2}\sigma^a)
$$

and
$$
(j_A, j_B) = (0, \frac{1}{2})
$$
 \Rightarrow $R_R(S) = e^{\frac{1}{2}(-ir_a + b_a)\sigma^a}$ (7.25)
\n $(A^a = 0, B^a = -\frac{1}{2}\sigma^a; \qquad J^a = -\frac{i}{2}\sigma^a, K^a = +\frac{1}{2}\sigma^a)$

The first of these representations $R_L(S)$ can be identified with the original set of 2×2 matrices $S \in SL(2, \mathbb{C})$, that is $\{S \in \mathbb{C}(2) : \det(S) = 1\}$, as parametrised in the form of equation 7.20. The representation $R_R(S)$ in equation 7.25 is a *different* map from the *same* complete set of $SL(2, \mathbb{C})$ elements, considered as an abstract group, into 2×2 matrix transformations on a 2-dimensional complex vector space \mathbb{C}^2 . The two representation spaces are given different subscript labels L and R to denote that they belong to different $SL(2, \mathbb{C})$ representations with the left-handed Weyl spinor transforming as $\psi_L \to R_L(S)\psi_L$ and the right-handed Weyl spinor transforming as $\psi_R \to R_R(S)\psi_R.$

Under a discrete parity transformation the sign of a Lorentz boost is reversed while the sign of a rotation is invariant. The naming convention of 'left' and 'right' representations originates since $R_L(S)$ and $R_R(S)$ are related by the sign of the boost generator contributions in equations 7.24 and 7.25 and are hence interchanged under a parity transformation. Indeed in general the parity operation switches between the two Lorentz representations (j_1, j_2) and (j_2, j_1) .

Since there is no 2 × 2 matrix D such that $R_L(S) = DR_R(S)D^{-1}$ for all $S \in$ $SL(2,\mathbb{C})$ the representations $R_L(S)$ and $R_R(S)$ are inequivalent. However the following relationships between equations 7.24 and 7.25 hold (with σ^2 defined in equation 7.14):

$$
R_L^*(S) = \sigma^2 R_R(S) (\sigma^2)^{-1}
$$
\n(7.26)

$$
R_L^{\dagger^{-1}}(S) = R_R(S) \tag{7.27}
$$

$$
R_L^{T}(S) = \sigma^2 R_L^{-1}(S) (\sigma^2)^{-1}
$$
\n(7.28)

showing respectively that the complex conjugate of $R_L(S)$ is equivalent to $R_R(S)$, the contragredient of $R_L(S)$ is equal to $R_R(S)$ and the transpose of $R_L(S)$ is equivalent to its inverse.

The Dirac representation $R_D(S)$ of $SL(2,\mathbb{C})$ has the reducible form $(\frac{1}{2},0) \oplus$ $(0, \frac{1}{2})$ $\frac{1}{2}$), acting on spinors in the space \mathbb{C}^4 , and via equation 7.27 it can be written as:

$$
R_D(S) = \begin{pmatrix} R_L(S) & 0 \\ 0 & R_R(S) \end{pmatrix} = \begin{pmatrix} S & 0 \\ 0 & S^{\dagger^{-1}} \end{pmatrix}
$$
(7.29)

which is the same action as described in equation 7.15, there derived from the Clifford algebra structure, with $R_L = R_L(S)$ and $R_R = R_R(S)$. Hence $S \in SL(2, \mathbb{C})$ acts on the left-handed components of the \mathbb{C}^4 Dirac spinor and $S^{\dagger^{-1}}$ acts on the right-handed components as an inequivalent representation of $SL(2,\mathbb{C})$. Equation 7.29 describes how the $Spin^+(1,3)$ Dirac representation can be constructed by combining left and right spinors as the $(\frac{1}{2},0) \oplus (0,\frac{1}{2})$ $\frac{1}{2}$) representation of SL(2, C). In fact Spin⁺(1,3) is isomorphic to the group $SL(2, \mathbb{C})$ (such isomorphisms for the spin groups only exist in low dimensions and for a handful of cases), each expressing the two-to-one cover of $SO^+(1,3)$.

Equations 7.15 and 7.29 and the comparison of the Dirac representation constructed as a reducible representation of $Spin⁺(1, 3)$ via the Clifford algebra or as a combination of two representations of $SL(2,\mathbb{C})$ expresses the relation between the 4component and 2-component spinor formalism. The 2-component Weyl spinors are more fundamental in the sense that ψ_L and ψ_R are treated differently in important features of the Standard Model, as we shall describe in the following section.

The two-to-one relationship between $SL(2,\mathbb{C})$ and $SO^+(1,3)$ may be exhibited by mapping a Lorentz vector $v_4 \in \mathbb{R}^{1,3}$ into the space of 2×2 complex Hermitian matrices as:

$$
\boldsymbol{v}_4 = (v^0, v^1, v^2, v^3) \rightarrow \boldsymbol{h}_2 = \boldsymbol{v}_4 \cdot \boldsymbol{\sigma} = \begin{pmatrix} v^0 + v^3 & v^1 - v^2 i \\ v^1 + v^2 i & v^0 - v^3 \end{pmatrix} \subset \mathbf{h}_2 \mathbb{C} \tag{7.30}
$$

where σ denotes the 2 \times 2 identity matrix σ ⁰ together with the three Pauli matrices σ^a of equation 7.14. This is the same object introduced in equation 6.19 of section 6.3 and also in equation 6.58 of section 6.5, based on the imaginary unit l in the latter case. We see from this equation, and in comparison with section 6.3, that $det(h_2)$ = $(v^0)^2 - (v^1)^2 - (v^2)^2 - (v^3)^2 = h^2$, with $h \in \mathbb{R}$, which may be expressed as the form $L(\mathbf{v}_4) = h^2$ (as employed in equation 5.46). While the fundamental representation of SL(2, \mathbb{C}) acts on the space \mathbb{C}^2 , the group action for elements $S \in SL(2, \mathbb{C})$ on the space $h_2\mathbb{C}$ provides another representation given by:

$$
h_2 \to h'_2 = S h_2 S^{\dagger} \tag{7.31}
$$

This maps $h_2 \to h'_2$ onto a new 2×2 complex Hermitian matrix with the same determinant; hence mapping the components $v_4^a \rightarrow v_4^{\prime a}$ according to a Lorentz transformation of the real 4-vector $v_4 \in \mathbb{R}^{1,3}$. With $S \in \{1_2, -1_2\}$ giving the identity transformation, $h'_2 = h_2$, the group $SL(2, \mathbb{C})$, isomorphic to $Spin^+(1,3)$ as described above, is the two-to-one covering spin group for $SO^+(1,3)$; that is $SO^+(1,3) = SL(2,\mathbb{C})/\mathbb{Z}^2$.

In fact the components of v_4 transform under the 4-dimensional vector $(\frac{1}{2}, \frac{1}{2})$ $(\frac{1}{2})$ representation of $SL(2,\mathbb{C})$. The matrix h_2 , and hence the vector v_4 , can be considered to be constructed out of two 2-component left-handed Weyl spinors χ and ϕ such that:

$$
h_2 = \chi \chi^\dagger + \phi \phi^\dagger \tag{7.32}
$$

as implied in equations 6.20 and 6.21 of section 6.3, with the elements of the group $SL(2, \mathbb{C})$ acting on the spinor components in the appropriate way.

This spinor substructure of vectors v_4 has some similarity to the situation discussed in section 5.2 for the relation $-\frac{1}{\kappa}G^{\mu\nu} = \rho u^{\mu}u^{\nu} - S^{\mu\nu}$, as implied in equation 5.34 via the Einstein equation, which describes the possibility of composing the rank–2 Einstein tensor in terms of a substructure involving the apparent 4-dimensional macroscopic vector flow $u(x)$ on the base manifold. The natural algebraic substructure of the field $v_4(x)$ in terms of the spinor decomposition of equation 7.32 may in turn be intimately related to the possible field interactions implied within the higher-dimensional form of time $L(\mathbf{v}_{27}) = 1$ at the *microscopic* level, underlying the composition of the Einstein tensor $G^{\mu\nu} = f(Y, \hat{\mathbf{v}})$ as expressed in equation 5.32.

Equation 7.31 describes the determinant preserving action $h_2 \to S h_2 S^{\dagger}$ of the elements $S \in SL(2,\mathbb{C})$ upon elements of the vector space of matrices $h_2 \in h_2\mathbb{C}$ that was extended in equations 6.16 and 6.22, by augmenting the complex numbers to the octonions, to identify an $SL(2, \mathbb{O})$ action on $h_2\mathbb{O}$ as the covering group of the 10-dimensional Lorentz group, as an intermediary for the E_6 action on $h_3\mathbb{O}$. For infinitesimal transformations we write $S = \exp(a) \simeq 1 + a$, where $a \in sl(2, \mathbb{C})$ is an infinitesimal element of the Lie algebra of $SL(2,\mathbb{C})$, and we have:

$$
\mathbf{h}_2 \rightarrow (1+a)\mathbf{h}_2(1+a^{\dagger}) \simeq \mathbf{h}_2 + \delta \mathbf{h}_2 \tag{7.33}
$$

with
$$
\delta h_2 = a h_2 + h_2 a^{\dagger}
$$
 (7.34)

where δh_2 has the same form as the 64 D^S actions on h₃ $\mathbb O$ of equation 6.5 in section 6.1, and may here be considered to represent the Lie algebra $sl(2,\mathbb{C})$ on the tangent space Th₂C. This corresponds to a possible substructure embedding of h₂C \subset h₂O \subset h₃O with respective group actions $SL(2,\mathbb{C}) \subset SL(2,\mathbb{O}) \subset SL(3,\mathbb{O})$. Before moving to the action of $SL(2,\mathbb{C})$ on full space $h_3\mathbb{O}$ in the following chapter (see equation 8.4), we first here consider the action of $SL(2, \mathbb{C})$ on the space h₃C. The 2×2 matrices $S \in SL(2, \mathbb{C})$ can be embedded in 3×3 matrices acting on $\mathcal{X} \in \mathrm{h}_3\mathbb{C}$ as:

$$
\mathcal{X} \rightarrow \left(\begin{array}{c|c} S & 0 \\ \hline 0 & 1 \end{array}\right) \left(\begin{array}{c|c} h_2 & \psi_L \\ \hline \psi_L^{\dagger} & n \end{array}\right) \left(\begin{array}{c|c} S^{\dagger} & 0 \\ \hline 0 & 1 \end{array}\right) \tag{7.35}
$$

This combines the vector representation of $SL(2, \mathbb{C})$ on $h_2 \in h_2\mathbb{C}$ and the spinor representation on $\psi_L \in \mathbb{C}^2$, together with the scalar $n \in \mathbb{R}$, in a single symmetry transformation which preserves det(\mathcal{X}) $\in \mathbb{R}$. In section 8.1 the spinor ψ_L will be identified with $\theta_l \in \mathbb{C}^2$ in a complex subspace of $\theta \in \mathbb{O}^2$ under the full E_6 action on h₃ \mathbb{O} , compatible with the embedding of the SL(2, \mathbb{C}) action of equation 7.35 within the $SL(2, \mathbb{O}) \subset E_6$ action of equation 6.29.

7.2 Internal Symmetry and Electroweak Theory

Together with the external Lorentz symmetry internal gauge symmetries are key to the properties of particle states observed in the laboratory. In this section we review the internal symmetries of the Standard Model with a particular emphasis on electroweak theory and the phenomenon of symmetry breaking (see for example [42]).

The quarks and leptons of one generation of Standard Model fermions transform under an internal symmetry described by the group product $SU(3)_c \times SU(2)_L \times$ $U(1)_Y$ (with the subscripts 'c', 'L' and 'Y' denoting colour, left-handed and hypercharge respectively). The corresponding representation of $SU(3)_c \times SU(2)_L \times U(1)_Y$ is composed as a sum of five irreducible pieces each labelled according to their transformation properties by $(n_3, n_2, n_1)_{L,R}$ with the subscript L or R denoting left or right chiral Weyl spinors, represented as four-component Dirac spinors, under the external Lorentz group. The five pieces are of dimension 6, 3, 3, 2 and 1 respectively (without an extra piece $(1,1,0)_R$ for a right-handed neutrino ν_R):

$$
(3,2,\frac{1}{6})_L + (3,1,\frac{2}{3})_R + (3,1,-\frac{1}{3})_R + (1,2,-\frac{1}{2})_L + (1,1,-1)_R
$$

\n
$$
q_L = \binom{u_L(\frac{2}{3})}{d_L(-\frac{1}{3})} \qquad u_R(\frac{2}{3}) \qquad d_R(-\frac{1}{3}) \qquad l_L = \binom{\nu_L(0)}{e_L(-1)} \qquad e_R(-1)
$$
\n
$$
(7.36)
$$

with the corresponding set of 15 particle states named on the second line alongside their electromagnetic charges. The components of particle multiplets transforming as triplets under $SU(3)_c$, $(n₃ = 3)$, couple to the strong interaction and consist of u-type and d-type quarks, while the $SU(3)_c$ singlet components consist of the neutrino ν and electron e leptonic states.

In the Standard Model electroweak theory weak eigenstates, that is fields transforming according to definite $SU(2)_L$ representations, are composed as left-handed doublets $(n_2 = 2)$ and right-handed singlets $(n_2 = 1)$, transforming for example in the case of leptons as $l_L \to l'_L = e^{-i\omega^\alpha \tau^\alpha} l_L$, with $\omega^\alpha \in \mathbb{R}$ and

$$
\tau^{\alpha} = \frac{1}{2}\sigma^{\alpha} \tag{7.37}
$$

for $\alpha = 1, 2, 3$ (see equation 7.14, with Greek indices used here for the generators of a gauge group), and $e_R \to e'_R = e_R$. With left and right-handed fermions hence undergoing different interactions with the $SU(2)_L$ gauge field this construction describes the empirical observation of parity violation in weak interactions. When additional generations of fermions are considered the weak eigenstates generally consist of a linear combination of physical mass eigenstates leading to the phenomena of mixing between the generations, as will be described towards the end of this section.

The electromagnetic charge of each particle in a multiplet is given by:

$$
Q = T^3 + \frac{Y}{2}
$$
 (7.38)

where T^3 is the eigenvalue under the third, diagonal, $SU(2)_L$ generator and the hypercharge Y labels the U(1)_Y representations, $(n_1 = Y/2)$ in equation 7.36, which are all one-dimensional for this Abelian group. For the right-handed states $T^3 = 0$ and the hypercharge is simply the electric charge of the fermion $Q(\psi_R) = \frac{Y}{2}(\psi_R)$. All fields transform as $\psi \to \psi' = e^{-i\omega \frac{Y}{2}(\psi)}\psi$, with $\omega \in \mathbb{R}$, under the hypercharge gauge symmetry $U(1)_Y$. The hypercharge $\frac{Y}{2}(\psi)$ itself is ultimately defined to give the correct electromagnetic charge Q, via the relation in equation 7.38, which is the same for the L and R parts of each fermion type with $Q(e_L) = Q(e_R) = -1$ for example. The charge Q determines the coupling to the electromagnetic field corresponding to the $U(1)$ _O gauge symmetry that survives electroweak symmetry breaking. Equation 7.38 may be considered as a relation either between the eigenvalues or the operators Q, T^3 and $\frac{Y}{2}$, depending on the context.

The representations of equation 7.36 can be expressed purely in terms of lefthanded fields by applying 'charge conjugation' to the right-handed cases, under which $(3, 1, \frac{2}{3})$ $\frac{2}{3}$) $_R \rightarrow (\bar{3}, 1, -\frac{2}{3})$ $\frac{2}{3}$)_L for example. Having all fields expressed in terms of the same Lorentz representation in this way is useful for unification models, in which individual pieces of equation 7.36 are combined in a larger representation of a single unifying gauge group. Since gauge transformations commute with Lorentz transformations, without interchanging L and R states, such a unifying gauge group then respects Lorentz invariance in the theory. While the states in equation 7.36 are all considered as 'particles' the action of charge conjugation also introduces 'antiparticle' states. Hence both particle and antiparticle states may be combined in unified multiplets, as for the case of the SU(5) model [43] cited regarding figure 7.3 in the following section.

The dynamics of the Standard Model fields is heavily based on a Lagrangian formalism. The Standard Model Lagrangian includes kinetic terms for the fermions in the form of the final term of equation 3.96, which for the lepton doublet l_L , with a conventional factor of i and covariant derivative D_{μ} , can be expressed as:

$$
\mathcal{L}_{\text{kin}} = i\bar{l}_L \gamma^\mu D_\mu l_L \tag{7.39}
$$

with
$$
D_{\mu} = \partial_{\mu} + ig W_{\mu}^{\alpha}(x) \tau^{\alpha} + ig' B_{\mu}(x) \frac{Y}{2}(l_{L}) \qquad (7.40)
$$

where τ^{α} is defined in equation 7.37 (and with an additional $D_{\mu} = \ldots + ig_s G_{\mu}^{\beta}(x) \lambda_{\beta}$ term, with $\beta = 1 \dots 8$ and the λ_{β} matrices listed in table 8.5, for SU(3)_c gauge interactions in the case of quarks). Hence the interaction between the gauge fields $W^{\alpha}_{\mu}(x)$, $B_{\mu}(x)$, with respective couplings g, g', and left-handed leptons has the Lagrangian form:

$$
\mathcal{L}_{int} = -\frac{g}{2} (\bar{\nu}_L \ \bar{e}_L) \gamma^{\mu} \left(\begin{pmatrix} W_{\mu}^3 & W_{\mu}^1 - iW_{\mu}^2 \\ W_{\mu}^1 + iW_{\mu}^2 & -W_{\mu}^3 \end{pmatrix} - \frac{g'}{g} \begin{pmatrix} B_{\mu} & 0 \\ 0 & B_{\mu} \end{pmatrix} \right) \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}
$$
 (7.41)

$$
\mathcal{L}_{\nu} = -\frac{g}{2} \bar{\nu}_L \gamma^{\mu} (W_{\mu}^3 - \frac{g'}{g} B_{\mu}) \nu_L
$$
 (7.42)

where the part \mathcal{L}_{ν} describes the gauge coupling to the neutrino alone, as implied in equation 7.39. Physical gauge boson fields $A_{\mu}(x)$ and $Z_{\mu}(x)$ are defined as a linear combination of $B_{\mu}(x)$ and $W_{\mu}^{3}(x)$ via the orthogonal transformation:

$$
A_{\mu} = \cos \theta_W B_{\mu} + \sin \theta_W W_{\mu}^3
$$

\n
$$
Z_{\mu} = -\sin \theta_W B_{\mu} + \cos \theta_W W_{\mu}^3
$$
\n(7.43)

that is with:

$$
B_{\mu} = \cos \theta_W A_{\mu} - \sin \theta_W Z_{\mu}
$$

\n
$$
W_{\mu}^3 = \sin \theta_W A_{\mu} + \cos \theta_W Z_{\mu}
$$
\n(7.44)

where θ_W is the weak mixing angle. Hence from equation 7.42 the coupling of the neutrino to the physical gauge field A_μ is:

$$
\mathcal{L}_{\nu A} = -\frac{g}{2} \bar{\nu}_L \gamma^\mu (\sin \theta_W A_\mu - \frac{g'}{g} \cos \theta_W A_\mu) \nu_L \tag{7.45}
$$

which is zero for:

$$
\tan \theta_W = \frac{g'}{g} \tag{7.46}
$$

This value of the weak mixing angle θ_W hence describes the electric charge neutrality of the neutrino with $A_\mu(x)$ interpreted as the electromagnetic field, the quanta of which are photons. More generally the coupling terms for the photon can be extracted from the relevant part of the covariant derivative D_{μ} , of the form in equation 7.40, acting on any field $\psi(x)$ as (with T^3 representing the third component of $su(2)_L$ and the hypercharge $\frac{Y}{2}$ as operators acting on the field ψ):

$$
D_{\mu} \sim ig W_{\mu}^{3} T^{3}(\psi) + ig' B_{\mu} \frac{Y}{2}(\psi)
$$
 retaining only W_{μ}^{3}, B_{μ} field parts
\n
$$
= ig \sin \theta_{W} A_{\mu} T^{3} + ig' \cos \theta_{W} A_{\mu} \frac{Y}{2}
$$
 by equation 7.44, dropping Z_{μ} parts
\n
$$
= ig \sin \theta_{W} A_{\mu} T^{3} + ig \sin \theta_{W} A_{\mu} \frac{Y}{2}
$$
 using equation 7.46
\n
$$
= ig \sin \theta_{W} A_{\mu} (T^{3} + \frac{Y}{2})
$$

$$
= ie A_{\mu} Q
$$
 (7.47)

Hence the electromagnetic coupling of any particle state to the photon is always proportional to eQ where the particle charge Q is defined in equation 7.38 and the electromagnetic coupling e is given by:

$$
e = g \sin \theta_W \tag{7.48}
$$

As described after equation 7.38 the different values of $\frac{Y}{2}$ compensate for the different $T³$ values for the L and R states of a given particle such that the respective coupling of each chiral component to the gauge field $A_u(x)$ is the same, as can be seen for each particle type in equation 7.36. Following the same lines of reasoning in equation 7.47 except instead retaining the gauge field $Z_{\mu}(x)$ and dropping the $A_{\mu}(x)$ field parts in the second line leads to:

$$
D_{\mu} \sim ig \cos \theta_{W} Z_{\mu} T^{3} - ig' \sin \theta_{W} Z_{\mu} \frac{Y}{2}
$$

\n
$$
= ig \cos \theta_{W} Z_{\mu} T^{3} - ig \frac{\sin^{2} \theta_{W}}{\cos \theta_{W}} Z_{\mu} \frac{Y}{2}
$$

\n
$$
= ig Z_{\mu} \left(\left(\cos \theta_{W} + \frac{\sin^{2} \theta_{W}}{\cos \theta_{W}} \right) T^{3} - \frac{\sin^{2} \theta_{W}}{\cos \theta_{W}} \left(T^{3} + \frac{Y}{2} \right) \right)
$$

\n
$$
= \frac{ig}{\cos \theta_{W}} Z_{\mu} (T^{3} - Q \sin^{2} \theta_{W}) \qquad (7.49)
$$

Hence there are two terms for the weak neutral interactions. The second term is proportional to the electromagnetic charge Q and is hence the same for L and R particle states. However since the eigenvalues of T^3 are only non-zero for the lefthanded states the first term only couples to the ψ_L components. The combination of the two terms in equation 7.49 implies that parity violation is only partial for neutral weak interactions. On the other hand for the charged weak interactions mediated via the $W^{\pm}_{\mu}(x)$ gauge fields, introduced in equation 7.57 below and involving only $\mathrm{SU}(2)_L$ components, parity violation is maximal. In the Standard Model Lagrangian the lefthanded chiral states ψ_L are projected out of the Dirac spinor states for the fermions using the P_L operator of equation 7.11, as seen for example in equations 7.77 and 7.78 at the end of this section.

In addition to the spin- $\frac{1}{2}$ fermions and spin-1 gauge bosons the Standard Model also introduces a spin-0 Higgs field, which is massive itself and closely associated with the origin of mass for the W^{\pm} and Z^{0} gauge bosons as well as the fermion states. Indeed electroweak theory is inextricably linked to the Higgs sector with the breaking of the electroweak symmetry $SU(2)_L \times U(1)_Y$ to the $U(1)_Q$ of electromagnetism mediated through the action of the gauge group on the Higgs field:

$$
\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{pmatrix}
$$
(7.50)

Transforming as a scalar under the external Lorentz symmetry the Higgs field is also invariant under the internal $SU(3)_c$ symmetry. On the other hand the above complex doublet of scalar fields ϕ transforms as a doublet under $SU(2)_L$ while also possessing hypercharge with $\frac{Y}{2} = +\frac{1}{2}$, which also accounts for the notation ϕ^+ and ϕ^0 in equation 7.50 by reference to equation 7.38. This collection of properties may be denoted

 $(1, 2, \frac{1}{2})$ by comparison with the list of Standard Model fermions in equation 7.36. The Lagrangian for the Higgs sector is:

$$
\mathcal{L}_H = (D_\mu \phi)^\dagger D^\mu \phi - V(\phi) \tag{7.51}
$$

where
$$
D_{\mu}\phi = \left(\partial_{\mu} + i\frac{g}{2}W^{\alpha}_{\mu}\sigma^{\alpha} + i\frac{g'}{2}B_{\mu}\sigma^{0}\right)\phi
$$
 (7.52)

is the gauge covariant derivative which is similar in form to equation 7.40 except with $\frac{Y}{2}$ = $+\frac{1}{2}$ here, and also σ^0 and $\sigma^{\alpha} = {\sigma^1, \sigma^2, \sigma^3}$ have been adopted directly from equation 7.14 rather than via equation 7.37. The fields $W_\mu^\alpha(x)$ and $B_\mu(x)$ are the $SU(2)_L$ and $U(1)_Y$ gauge fields, with couplings g and g' respectively, as introduced in equation 7.40. The breaking of the electroweak symmetry relies on the 'Mexican hat' potential term in the Lagrangian of equation 7.51 with:

$$
V(\phi) = -\mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2 \tag{7.53}
$$

with real coefficients $\mu^2 > 0$ and $\lambda > 0$. From equation 7.50 it can be seen that the potential $V(\phi)$ is a function of $\phi^{\dagger} \phi = \frac{1}{2} \sum_{i=1}^{4} \phi_i^2$ only. The vacuum expectation value for this field $\langle \phi \rangle$, that is the minimum in the potential, can be taken without loss of generality (in the 'unitarity gauge') to be:

$$
\langle \phi \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} \quad \text{with} \quad v = \frac{\mu}{\sqrt{\lambda}} \tag{7.54}
$$

This charge neutral component of the Higgs field $\phi^0 = \frac{v}{\sqrt{2}}$ is invariant under the action of the charge generator $Q = T^3 + \frac{Y}{2} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}$ $\frac{0}{2}$ $+\binom{\frac{1}{2}+0}{0}$ $0\frac{1}{2}$ $=$ $\binom{10}{00}$, from equation 7.38 applied for the Higgs field, which remains unbroken. Hence the gauge symmetry is broken from $SU(2)_L \times U(1)_Y$ down to $U(1)_Q$, identified in a linear combination of the third component of $su(2)_L$ and the hypercharge generator $u(1)_Y$, as the symmetry which leaves the vacuum value $\langle \phi \rangle$ in equation 7.54 invariant.

Masses arise for the gauge fields corresponding to the broken $SU(2)_L \times U(1)_Y$ generators from the kinetic term in the Higgs Lagrangian of equation 7.51. Acting on the vacuum state the covariant derivative of equation 7.52 can be written as:

$$
D_{\mu}\phi = \begin{pmatrix} \partial_{\mu} + \frac{i}{2}gW_{\mu}^{3} + \frac{i}{2}g'B_{\mu} & \frac{i}{2}g(W_{\mu}^{1} - iW_{\mu}^{2}) \\ \frac{i}{2}g(W_{\mu}^{1} + iW_{\mu}^{2}) & \partial_{\mu} - \frac{i}{2}gW_{\mu}^{3} + \frac{i}{2}g'B_{\mu} \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix}
$$
(7.55)

Hence \mathcal{L}_H in equation 7.51 contains the expression (for now neglecting fluctuations about the vacuum value v :

$$
(D_{\mu}\phi)^{\dagger}D^{\mu}\phi = \frac{g^2}{4}(W_{\mu}^1 + iW_{\mu}^2)(W^{1\mu} - iW^{2\mu})\frac{v^2}{2} + \frac{1}{4}\left(W_{\mu}^3 - B_{\mu}\right)\left(\frac{g^2}{-gg'} - gg'\right)\left(\frac{W^{3\mu}}{B^{\mu}}\right)\frac{v^2}{2}
$$
(7.56)

With physical gauge fields $W^{\pm}_{\mu}(x)$ associated respectively with 2×2 matrices σ^{\pm} in the complexified $SU(2)_L$ Lie algebra defined in turn as:

$$
W^{\pm}_{\mu} = \frac{1}{\sqrt{2}} (W_{\mu}^{1} \mp iW_{\mu}^{2})
$$
\n(7.57)

$$
\sigma^{\pm} = \frac{1}{2} (\sigma^1 \pm i \sigma^2) \tag{7.58}
$$

the relation:

$$
\frac{1}{\sqrt{2}}(W^{+}_{\mu}\sigma^{+} + W^{-}_{\mu}\sigma^{-}) = \frac{1}{2}(W^{1}_{\mu}\sigma^{1} + W^{2}_{\mu}\sigma^{2})
$$
\n(7.59)

may be substituted in for the W^1_μ, W^2_μ piece of the covariant derivative in equation 7.52. In turn the first term in equation 7.56 explicitly takes the form of a mass term for the $W^{\pm}_{\mu}(x)$ fields in the Lagrangian:

$$
\mathcal{L}_H = \frac{g^2 v^2}{8} (W_\mu^+ W^{\mu} + W_\mu^- W^{-\mu}) + \dots
$$

hence with $M_W = \frac{1}{2}gv$ (7.60)

being the W^{\pm} mass.

The second term in equation 7.56 contains a 2×2 mass matrix composed of quadratic terms in the couplings g, g' . Applying the same orthogonal transformation of equations 7.43 and 7.44 to the fields W^3_μ and B_μ with the weak mixing angle θ_W as specified in equation 7.46 diagonalises the mass matrix with respect to the fields Z_{μ} and A_{μ} such that:

$$
\mathcal{L}_H = \dots + \frac{1}{2} \left(Z_\mu \ A_\mu \right) \begin{pmatrix} M_Z^2 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} Z^\mu \\ A^\mu \end{pmatrix} \tag{7.61}
$$

with
$$
M_Z = \frac{1}{2}\sqrt{g^2 + g'^2} v = \frac{M_W}{\cos \theta_W}
$$
 (7.62)

Hence the same weak mixing angle θ_W that accounts for the electromagnetic charge neutrality of the neutrino ν through the covariant derivative D_{μ} acting on the lepton field l_L in equation 7.45, deriving from the kinetic term in the Lagrangian for the lepton field in equations 7.39 and 7.40, also diagonalises the above mass matrix and leaves the photon field A_μ massless through D_μ acting on the Higgs field ϕ , deriving from the kinetic term in the Lagrangian for the Higgs field in equations 7.51 and 7.52.

Considering fluctuations about the vacuum value with $v \to v + H(x)$ in equation 7.54 (as neglected in writing down equation 7.56) in the quantum theory the real field $H(x)$ is associated with a massive scalar particle known as the Higgs boson. In terms of the parameters of the theory the Higgs mass is determined to be $M_H = \sqrt{2} \mu = \sqrt{2\lambda} v$. While the vacuum value is empirically constrained to the order of the weak scale with, via equation 7.60, $v = 2\frac{M_W}{g} \sim (\sqrt{2}G_F)^{-\frac{1}{2}} \sim 246$ GeV, where G_F is the Fermi constant, this does not determine the two parameters of the potential in equation 7.53. These latter parameters can now be deduced given the discovery of the Higgs at the LHC and the empirical measurement of $M_H \simeq 125$ GeV [44].

At tree level the relations in the quantum field theory described in equations 7.60 and 7.62 lead to the definition of the parameter:

$$
\rho = \frac{M_W^2}{M_Z^2 \cos^2 \theta_W} = 1\tag{7.63}
$$

The fact that this expression holds approximately for the corresponding empirically measured values can be explained in terms of a further symmetry associated with the Higgs sector. Expressing the Higgs field components in the form of a bi-doublet, that is the 2×2 complex matrix:

$$
\Phi = \frac{1}{\sqrt{2}} \left(\epsilon \phi^*, \phi \right) = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi^{0^*} & \phi^+ \\ -\phi^{+^*} & \phi^0 \end{pmatrix}
$$
 (7.64)

with $\epsilon = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, the Higgs potential term of equation 7.53 may be rewritten as:

$$
V(\Phi) = -\mu^2 \operatorname{tr} \Phi^\dagger \Phi + \lambda (\operatorname{tr} \Phi^\dagger \Phi)^2 \tag{7.65}
$$

This is invariant under the $L \in SU(2)_L$ action $\Phi \to L\Phi$ and $U(1)_Y$ action $\Phi \to \Phi e^{-\frac{i}{2}\theta \sigma^3}$ with $\theta(x) \in \mathbb{R}$ as local gauge transformations. While ϕ and $\epsilon \phi^*$ transform in the same way under $SU(2)_L$, they have opposite hypercharge, with $\frac{Y}{2}(\phi) = +\frac{1}{2}$ and $\frac{Y}{2}(\epsilon \phi^*) = -\frac{1}{2}$, and hence the generator for $U(1)_Y$ transformations here is σ^3 rather than σ^0 (see for example [45] section 3). The Higgs Lagrangian of equation 7.51, which is also invariant under these gauge transformations, can be written in the form:

$$
\mathcal{L}_H = \text{tr}(D_\mu \Phi^\dagger D^\mu \Phi) - V(\Phi) \tag{7.66}
$$

where
$$
D_{\mu}\Phi = \partial_{\mu}\Phi + i\frac{g}{2}W^{\alpha}_{\mu}\sigma^{\alpha}\Phi - i\frac{g'}{2}B_{\mu}\Phi\sigma^3
$$
 (7.67)

is the gauge covariant derivative for the bi-doublet. In the limit $g' \to 0$ this Lagrangian also has an additional, global, symmetry denoted $SU(2)_R$ with action $\Phi \to \Phi R^{\dagger}$ for any $R \in SU(2)_R$, as can be seen by cyclic permutation of the arguments under the trace, with $tr(R\Phi^{\dagger}\Phi R^{\dagger}) = tr(R^{\dagger}R\Phi^{\dagger}\Phi) = tr(\Phi^{\dagger}\Phi)$ for example. This symmetry in the Standard Model is considered to be 'accidental' in the sense that it was not explicitly introduced in constructing the Higgs field to break the electroweak symmetry. It enlarges the complete global symmetry of the Higgs field to the action of $SU(2)_L \times$ $SU(2)_R$, as $\Phi \to L\Phi R^{\dagger}$ (where L here represents a global action of the local $SU(2)_L$ symmetry), which is simply the SO(4) symmetry of the quantity $\sum_{i=1}^{4} \phi_i^2$ described below equation 7.53. The vacuum expectation value of equation 7.54 can be written in the form:

$$
\langle \Phi \rangle = \frac{1}{2} \begin{pmatrix} v & 0 \\ 0 & v \end{pmatrix}
$$
 (7.68)

This vacuum value breaks the global $SU(2)_L \times SU(2)_R$ down to a single $SU(2)$ symmetry denoted $SU(2)_{L+R}$, with the action $\langle \Phi \rangle \to L \langle \Phi \rangle L^{\dagger}$ for $L \in SU(2)_{L+R}$ leaving equation 7.68 invariant. This is equivalent to the $SO(3) \subset SO(4)$ symmetry acting on the four components ϕ_i when taking the values of an arbitrary fixed Euclidean 4-vector, such as $(\phi_1, \phi_2, \phi_3, \phi_4) = (0, 0, v, 0)$ in equation 7.68. The global $SU(2)_{L+R}$ symmetry itself is broken for hypercharge coupling $g' \neq 0$, which involves gauging the U(1) $_Y \subset SU(2)_R$ subgroup via the σ^3 action of equation 7.67, which is both the hypercharge generator itself and also the third component of the $SU(2)_R$ action.

For the Standard Model in the limit $g' \to 0$ the three W^{α}_{μ} gauge fields transform as a triplet under the unbroken $SU(2)_{L+R}$ global symmetry, and hence the masses gained from electroweak symmetry breaking are identical, with $M_{W^{\pm}} = M_{Z^0}$ (as can be seen from equations 7.60 and 7.62 for $g' \to 0$, in this limit. For small g' the unbroken $U(1)_Q$ symmetry corresponding to the massless photon determines a weak mixing angle θ_W with $\cos^2 \theta_W = \frac{g^2}{g^2 + g^2}$ $\frac{g}{g^2+g'^2}$ which also determines the mass ratio of the heavy gauge bosons at tree level according to equation 7.63. This relation $\rho = 1$ is protected from radiative corrections by the approximate $SU(2)_{L+R}$ symmetry, which is hence named 'custodial symmetry' [46, 45].

Masses for all three generations of fermions are included in the Standard Model Lagrangian by appending gauge invariant terms with Yukawa couplings to the Higgs field:

$$
\mathcal{L}_Y = -\Gamma_{u}^{ij} \,\bar{q}_L^i \,\epsilon \phi^* u_R^j - \Gamma_{d}^{ij} \,\bar{q}_L^i \,\phi \,d_R^j - \Gamma_{e}^{ij} \,\bar{l}_L^i \,\phi \,e_R^j + \text{h.c.} \tag{7.69}
$$

(where 'h.c.' is the Hermitian conjugate of all the preceding terms). Here the Yukawa couplings Γ_u , Γ_d and Γ_e are 3×3 complex matrices in generation space with fermion flavour indices $i, j = \{1, 2, 3\}$ and hence, for example, $u_R^i \equiv \{u_R, c_R, t_R\}$ denotes the three generations of u-type right-handed quarks. When the Higgs field acquires the vacuum value $\langle \phi \rangle$ as expressed with the gauge choice of equation 7.54 the fermion states acquire Dirac mass terms via the Yukawa couplings:

$$
\mathcal{L}_M = -M_u^{ij} \bar{u}_L^i u_R^j - M_d^{ij} \bar{d}_L^i d_R^j - M_e^{ij} \bar{e}_L^i e_R^j + \text{h.c.}
$$
\nwhere
$$
M_{u,d,e}^{ij} = \Gamma_{u,d,e}^{ij} \frac{v}{\sqrt{2}}
$$
\n(7.70)

are the three fermion mass matrices. Physical particle states may be identified by diagonalising each M^{ij} matrix using independent unitary transformations applied to each left and right-handed fermion set via 3×3 unitary matrices A^{ij} , such as for:

$$
\boldsymbol{u}_L \to \boldsymbol{u}'_L = A_{u_L}^\dagger \boldsymbol{u}_L \tag{7.71}
$$

$$
\boldsymbol{u}_R \to \boldsymbol{u}_R' = A_{u_R}^\dagger \boldsymbol{u}_R \tag{7.72}
$$

Hence $u'_L \equiv \{u'_L, c'_L, t'_L\}$ and u'_R are mass eigenstate fields with the masses of the three u-type quarks read off from the diagonal elements of:

$$
M'_u = A_{u_L}^{\dagger} M_u A_{u_R} = \begin{pmatrix} m_u & 0 & 0 \\ 0 & m_c & 0 \\ 0 & 0 & m_t \end{pmatrix}
$$
 (7.73)

with
$$
\mathcal{L}_M = -m_u \bar{u}'_L u'_R - m_c \bar{c}'_L c'_R - m_t \bar{t}'_L t'_R + \text{h.c.}
$$
 (7.74)

as the Lagrangian Dirac mass terms for the *u*-type quarks (with u', c' and t' here denoting the individual first, second and third generation u -type quarks). The u quark itself hence has mass $m_u = Y_u \frac{v}{\sqrt{2}}$ with the Yukawa coupling $Y_u = \Gamma_u^{\prime 11}$ extracted from the diagonalised basis. From equation 7.60 the u-quark mass can be related to the W^{\pm} gauge boson mass as:

$$
m_u = \frac{\sqrt{2}}{g} Y_u M_W
$$

with
$$
Y_f = \frac{g m_f}{\sqrt{2} M_W}
$$
 (7.75)

where Y_f is the Yukawa coupling for each fermion f to the Higgs field ϕ , including the similar cases for the d-type quarks and charged leptons as following also from equation 7.70. (The neutrino mass may be treated differently and may not involve a Yukawa coupling, [42] chapter 7). The couplings Y_f are typically small since $m_f \ll M_W$ except for the case of the top quark – with the mass m_t observed to be approximately the sum of M_W and M_Z . All of the Yukawa couplings are added by hand in order to match the empirically determined fermion masses.

In the physical mass eigenstate basis there is no Yukawa mixing between generations, as can be seen in equation 7.74 in comparison to equation 7.70 where in the latter expression the quark states coupling to the weak $SU(2)_L$ gauge fields are generally composed of a linear combination of the physical quark states. The weak $SU(2)_L$ doublets in the quark sector may be written as $\binom{u}{d}_L$, $\binom{c}{s}$ $\left(\frac{c}{\tilde{b}}\right)_L$, with the inter-generation mixing expressed purely in terms of the \overline{d} -type quark states:

$$
\begin{pmatrix} \tilde{d} \\ \tilde{s} \\ \tilde{b} \end{pmatrix} = V_{\text{CKM}} \begin{pmatrix} d' \\ s' \\ b' \end{pmatrix}
$$
 (7.76)

Here the weak states $\tilde{d}, \tilde{s}, \tilde{b}$ are related to the physical states d', s', b' via the 3 \times 3 unitary Cabibbo-Kobayashi-Maskawa mixing matrix $V_{\text{CKM}} = A_{u_L}^{\dagger} A_{d_L}$. With five relative global phase transformations between the six quarks (u, d, c, s, t, b) only four of the nine parameters of the unitary matrix V_{CKM} are physical. These four parameters describe three real mixing angles between the three generations and one complex phase which gives rise to CP violating phenomena. Together with the six quark masses a total of ten physical parameters (contributing just over half of the 18 Standard Model parameters listed in table 15.2) may hence be deduced from the Lagrangian for the quark sector after the above field redefinitions. (Again, the description of neutrino mixing in the leptonic sector is a little different, [42] chapter 7).

The weak interaction terms for the quarks with the charged gauge bosons W^{\pm} may be described by the Lagrangian:

$$
\mathcal{L}_{qW} = -\frac{g}{2\sqrt{2}} \bar{u}^{i} \gamma^{\mu} (1 - \gamma^{5}) \tilde{d}^{i} W_{\mu}^{-} + \text{h.c}
$$
 (7.77)

with the implied sum for $i = 1, 2, 3$ over the weak states (and where the Hermitian conjugate contains the W^+_μ term). Expressing the d-type quarks as a linear combination of the mass states the above Lagrangian can be written in terms of the six physical quarks as:

$$
\mathcal{L}_{qW} = -\frac{g}{2\sqrt{2}} \left(\bar{u} \quad \bar{c} \quad \bar{t} \right) \gamma^{\mu} (1 - \gamma^5) V_{\text{CKM}} \begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} W_{\mu}^- + \text{h.c} \qquad (7.78)
$$

In these equations the operator $P_L = \frac{1}{2}$ $\frac{1}{2}(1-\gamma^5)$ of equation 7.11 has been put in by hand to project out the left-handed components of the Dirac spinors, describing maximal parity violation for the charged weak current. This CKM mixing originates from the mismatch between the Yukawa and weak interactions in the Standard Model Lagrangian, with the corresponding mass and weak quark eigenstates for the u and d type quarks related via unitary transformations such as equations 7.71 and 7.72. On the other hand the neutral currents are flavour-diagonal and such terms are unchanged by the unitary transformations relating the mass and weak states, that is with $A_{u_L}^{\dagger} A_{u_L} =$ $1₃$ and so on. Hence there are no flavour changing neutral currents coupled to the Z_{μ} or A_{μ} fields, and only the W^{\pm} fields mediate mixing between the generations.

7.3 Unification Models and Dynkin Analysis

While the action of E_6 on $h_3\mathbb{O}$ studied in chapter 6 describes a symmetry of time it is also of course desirable that the mathematical structures arising in the present theory should bear a close resemblance to the symmetries and structures experimentally identified in particle physics. This data is summarised in the Standard Model, as reviewed in the previous two sections, which describes the non-gravitational interactions between fundamental particles in terms of the gauge symmetry group $SU(3)_c \times SU(2)_L \times U(1)_Y$. Hence in this section we make a preliminary assessment of the suitability of the Lie group E_6 , both generally and as constructed in chapter 6, as a unifying symmetry.

It is well known that the three subgroup components of the Standard Model gauge symmetry are related to the series of normed division algebras, as introduced here in section 2.1 in the context of forms of temporal flow and discussed further in section 6.2. Indeed, $U(1)$ is isomorphic to the complex numbers $\mathbb C$ of unit magnitude under multiplication, while $SU(2)$ is similarly isomorphic to the quaternions $\mathbb H$ of unit magnitude, and $SU(3)$ is the subgroup of G_2 , the automorphism group of the octonions O, that leaves invariant a given imaginary octonion element. The aesthetic appeal and elegance of such observations have led a number of authors to speculate on a direct connection between the existence of these unique mathematical objects and the nature of the physical structure of the world (see for example [47, 48, 49, 50]). However, while identifying a relationship between the mathematical properties of the division algebras and features of the Standard Model of particle physics much of this work is lacking in any underlying conceptual motivation for the importance of such mathematical objects in nature.

Since the octonion algebra features significantly in the present paper, in the action of the group $E_6 \equiv SL(3, \mathbb{O})$ on the space $h_3 \mathbb{O}$, the references cited above suggest a reasonable likelihood of identifying some relation between the structures of the present theory and those of the Standard Model. Such a correspondence will be described in the following chapter. In the present theory we have both a clear conceptual understanding of the source of these algebras through the symmetry of the flow of time and in turn a well defined constraint on the introduction of these algebraic structures into the equations of physics through the relation $L(\mathbf{v}) = 1$ and its symmetries.

Also in the present theory, as well as aiming to account for the internal gauge interactions of the Standard Model through the higher-dimensional structures, gravitation is included on the base manifold M_4 with a subspace $h_2 \mathbb{C} \subset h_3 \mathbb{O}$ locally identified with the tangent space TM_4 and with the subgroup $SL(2,\mathbb{C}) \subset E_6$ being the covering group of the external Lorentz group. As described towards the end of section 3.4 general relativity can be presented in the form of a gauge theory with a local Lorentz symmetry constructed in terms of the components of both a Lorentz Lie algebra-valued connection $A^a_{\ b\mu}(x)$ and a tetrad field $e^a_{\ \mu}(x)$. In terms of the covering group it can in turn be considered to be an $SL(2,\mathbb{C})$ gauge theory with an $sl(2,\mathbb{C})$ -valued connection, which can accommodate a description of both vector and spinor objects in spacetime.

While the dynamics of such an $SL(2,\mathbb{C})$ 'gauge theory' of gravitation [51, 52] are different to those of a standard Yang-Mills gauge theory, as also described in section 3.4, an extension for internal gauge symmetries might be more readily achieved with such a theory of gravity. (Considering gravity as a gauge theory contrasts with the Kaluza-Klein approach reviewed in chapter 4 for which an internal gauge theory derives from general relativity with extra spatial dimensions.) Indeed an $SL(2,\mathbb{C})\times U(1)$ theory of gravitation and electromagnetism can be obtained by introducing an additional $e^{i\alpha(x)}$ phase factor element for the group $U(1)$. This can be achieved by augmenting the set of symmetry actions $S \in SL(2,\mathbb{C})$ with $\det(S) = 1$ to include also the actions $U = e^{i\alpha(x)}\mathbf{1}_2 \in U(1)$. The mapping of equation 7.31, now incorporating also the U(1) action $h_2 \to Uh_2U^{\dagger}$, then remains one that preserves the value of $\det(h_2)$ and leaves the metric $g_{\mu\nu}(x)$ on M_4 invariant, as for the original $SL(2,\mathbb{C})$ action. Further, the U(1) action in fact leaves each of the four components of h_2 invariant and hence effectively acts as an 'internal symmetry', as described also at the end of section 6.3. Within the set of E₆ symmetry actions on h₃^① the action of S_q^1 in equation 6.43, particularly on a type 1 h₂ $\mathbb{C} \subset h_3 \mathbb{O}$ subspace, is most reminiscent of the above U(1) symmetry action on $h_2\mathbb{C}$ and this property is suggestive for the choice of the $U(1)_Q$ action for the electromagnetic gauge symmetry in the present theory.

By further augmenting the internal degrees of freedom such unification schemes which begin with an $SL(2,\mathbb{C})$ theory of gravity can be extended to an $SL(2,\mathbb{C})\times G$ theory where G may be the gauge symmetry group for the internal forces as identified experimentally in the Standard Model, that is $SU(3)_c \times SU(2)_L \times U(1)_Y$. For such a model there remains the task of introducing states which transform as fermions under the external $SL(2,\mathbb{C})$ symmetry and under the appropriate representations of the internal symmetry as summarised in equation 7.36. However such an approach, with the appropriate interpretation of the gauge groups and their empirically motivated representations, only serves to describe gravity together with internal field interactions in a more unified framework.

In the present theory, however, the unification group $\hat{G} = \mathbb{E}_6$ includes the external spacetime symmetry central to general relativity in the form of the subgroup $SL(2,\mathbb{C}) \subset E_6$. It is then through the distinctive role of this subgroup, in the identification of the necessary perceptual background for the world, that the larger symmetry is broken down to local gauge groups with representations on the broken fragments of the space $h_3\mathbb{O}$. The local gauge groups themselves will be initially identified as the 'stability' group leaving the space of vectors $v_4 \in TM_4$, via equation 7.30 equivalent to $h_2 \in h_2 \mathbb{C} \subset h_2 \mathbb{O}$, invariant, generalising from the above case of the $e^{i\alpha(x)}\mathbf{1}_2 \in U(1)$ action on $h_2\mathbb{C}$.

Having at hand the *real* form of E_6 acting on $h_3\mathbb{O}$, as described in the previous chapter, a detailed study of this symmetry breaking over TM_4 is possible. Initially, however, in this section the symmetry breaking patterns for E_6 and the question of whether this group is large enough to actually contain both $SL(2,\mathbb{C})$ and $SU(3)$ × $SU(2) \times U(1)$ will be addressed at the level of the *complex* Lie algebras, in order to gain an overview, before returning to the specific real forms of these algebras in the following chapter.

One of the main motivations for studying the complexified forms of real Lie algebras in general is the existence of a concise classification scheme. Indeed, every complex simple Lie algebra belongs to one of just four sets of classical algebra types, which include the complex forms of the rotation algebras $\mathfrak{so}(p,q)$, or is otherwise identified with one of the five exceptional cases, which include $L(E_6)$. A further motivation is that each complex simple Lie algebra has a one-to-one correspondence with a 'Dynkin diagram', with semi-simple Lie algebras likewise corresponding to disconnected Dynkin diagrams. The analysis of such diagrams gives a good deal of guidance towards the possible symmetry breaking patterns for a complex Lie algebra and its real forms as encountered in the context of a theoretical model for physical phenomena.

Firstly, we briefly review the relationship between Lie algebras and their representations. In general, each complex simple Lie algebra, as exemplified by the Dynkin diagrams shown later in this section, and taking its place amongst the systematic classification of such algebras, may be associated with several real forms, with each real algebra in turn associated with one of more Lie group, and finally each Lie group possesses an unlimited number of representations. This situation is depicted in figure 7.1.

Figure 7.1: Any given complex Lie algebra $L_{\mathbb{C}}$, which has a unique Dynkin diagram, is in general associated with a multiplicity of real algebra forms $L_{\mathbb{R}}$, groups G and representations R.

While Dynkin analysis at the level of $L_{\mathbb{C}}$ is described in this section, in this paper we generally deal with the structures of $L_{\mathbb{R}}$ and G , with notation such as so (p, q) used for a real Lie algebra and $SO(p,q)$ for the related Lie group, with the distinction being otherwise understood from the context. As an example of the chain of relations in figure 7.1 the case for $L_{\mathbb{C}} =$ so(10) with links through to the $R = 16$ representation, of particular interest here and featuring for example in equation 8.3 in the opening of the following chapter, is described in table 7.1.

$L_{\mathbb{C}}$	$L_{\mathbb{R}}$	G	$_{\it R}$
		O(1,9)	
		SO(1,9)	1
	so(10)	$SO^+(1,9)$	10
so(10)	\Leftrightarrow \rightarrow so(1,9)	\rightarrow Spin ⁺ (1,9) $\overset{\nearrow}{\Leftrightarrow}$	$\rightarrow 16$
	so(2,8)	Spin(1,9)	$\overline{16}$
		Pin(1,9)	٠

Table 7.1: The complex Lie algebra so(10), called D_5 in Cartan's notation, with corresponding real algebra forms $L_{\mathbb{R}}$, groups G and representations R (labelled by their dimension), as an example of the general case depicted in figure 7.1, with a particular chain of forms discussed in the text highlighted by the horizontal arrows.

In developing a theoretical model the initial motivation often begins from the left-hand side of figure 7.1, by identifying a complex Lie algebra which exhibits an appropriate symmetry breaking pattern to account for the gauge groups of the Standard Model as described in previous section; and then the task remains to identify the appropriate representations for particle states such as those of equation 7.36. In this paper such an approach also serves as a useful guide, as we describe in this section. However, here our starting point is rather more anchored in the right-hand side of figure 7.1 since the mathematical form $L(v) = 1$ strongly motivates the possible representations, with the set of real numbers composing the vector \boldsymbol{v} already belonging to a representation space transforming under the relevant symmetries of $L(v) = 1$.

As a preliminary observation we note that given our use of the $R = 27$ representation of the particular group $G = E_{6(-26)}$, this uniquely leads back via the real Lie algebra $L_{\mathbb{R}} = L(E_{6(-26)})$ to the complex Lie algebra $L_{\mathbb{C}} = L(E_6)$ as we step from right to left through figure 7.1. The structure of symmetry breaking feeding down from the complex Lie algebra is largely preserved in terms of semi-simplicity of the algebra and group and in terms of the reducibility of the algebra and group representations. Hence we here consider the Dynkin diagrams for the relevant complex Lie algebras and the significant Lie subalgebras involved.

The 'rank' of a Lie algebra is the dimension of the Cartan subalgebra, composed of a maximal subset of mutually commuting generators, which is unique up to automorphisms of the Lie algebra. For a rank-n Lie algebra there are n 'simple roots' in the dual root space which is constructed out of the eigenvalues in the adjoint representation of the algebra in the Cartan-Weyl basis. The properties of a rank-n Lie algebra can be described in terms of geometric relations between these simple roots in the Euclidean \mathbb{R}^n root-space and encoded in the topology relating the n nodes of the corresponding Dynkin diagram, such as those depicted in figure 7.2 for the rank-6 $L(E_6)$, rank-5 so(10), rank-4 su(5) and rank-2 Lorentz Lie algebras. For example, the Dynkin diagram for the Lorentz algebra consists of two disconnected nodes, meaning that the corresponding two simple roots are at 90^0 in root space, whereas nodes

Figure 7.2: The four Dynkin diagrams for the (a) $L(E_6)$, (b) so(10), (c) su(5) and (d) Lorentz or $sl(2,\mathbb{C})$ Lie algebras.

connected by a single line denote an angle of 120^0 . At the level of the complexified Lie algebra $L_{\mathbb{C}}$ the Lorentz algebra has the semi-simple composition su(2) \oplus su(2), as described earlier in equations 7.21–7.23, which in this case is not respected by the corresponding real form $L_{\mathbb{R}} = \text{so}^+(1, 3)$ of the Lorentz Lie algebra which is simple. An explicit basis for the Cartan subalgebra for the real form of $L(E_6)$ of importance in this paper was given in equation 6.56 as represented by vector fields on the space $Th_3\mathbb{O}$.

Regular subalgebras, that is those respecting the Cartan-Weyl decomposition of the complex Lie algebra, may be readily obtained from the Dynkin diagrams. A maximal subgroup $G' \subset G$ is one for which there is no intermediate G'' such that $G' \subset G$ $G'' \subset G$ as a series of proper subgroups, with a similar definition for the corresponding maximal subalgebra. A regular maximal subalgebra can be obtained from a Dynkin diagram by the prescription of removing one node and including an extra $U(1)$ factor, which also means that the algebra obtained is not 'semi-simple'. For example figure 7.3 shows a possible symmetry breaking pattern for the $su(5)$ algebra for the well-known case [43], as alluded to in the previous section, in which the Standard Model local gauge group is obtained.

Figure 7.3: Removing a node from the Dynkin diagram for the Lie algebra of the group $SU(5)$ reveals a breaking to $SU(3) \times SU(2) \times U(1)$, which motivates the use of $SU(5)$ in unified theories.

Similarly from figure 7.2 it can be seen that $SO(10)$ contains $SU(5)$ as a subgroup, by removing either of two appropriate end nodes. Hence the full Standard Model gauge group can be obtained by first breaking $SO(10)$ to $SU(5)$ and then breaking SU(5) as described in figure 7.3. Hence the 45-dimensional group $SL(2,\mathbb{O}) \equiv Spin^+(1,9)$ constructed in section 6.3 as the double cover of $SO^+(1,9)$, which is generated by a real form of the complex Lie algebra $\mathfrak{so}(10)$, is also potentially of great interest for internal gauge group unification in particle physics.

In the context of the discussion of section 6.4 following equation 6.42, the G_2 automorphism group of $\mathbb O$ is reduced to the subgroup $SU(3) \subset G_2$ if a complex subspace, for example with the imaginary unit $l \in \mathbb{O}$, is fixed, as also alluded to near the opening of this section. Similarly the subgroup $SU(3) \subset G_2 \subset SO^+(1,9)$ may be obtained through the selection of a preferred subspace $h_2\mathbb{C} \subset h_2\mathbb{O}$, since this choice also fixes an imaginary unit of $h_2\mathbb{O}$. Here the mechanism for such a selection is provided by the nature of perception on the base manifold M_4 with the vector space $TM_4 \equiv h_2 \mathbb{C}$ and h₂C ⊂ h₂^O through the identification of an external SO⁺(1,3) ⊂ SO⁺(1,9) symmetry.

However breaking the rank-2 Lorentz group out of the rank-5 so(10) clearly does not leave sufficient symmetry to describe the full rank-4 Standard Model gauge group.

It was also shown in section 6.4 how 3 copies of $SL(2,\mathbb{O})$, described with a total of $(3 \times 45) = 135$ generator actions, lock tightly together as an independent basis set of 78 generators, summarised in table 6.3, for the E_6 action on $\mathcal{X} \in h_3\mathbb{O}$ preserving $\det(\mathcal{X})$. This space hence describes a highly symmetric form of temporal flow $L(\mathbf{v}_{27}) = 1$ motivating the study of this exceptional Lie group.

As well as composing a rich symmetry of a multi-dimensional form of $L(\mathbf{v}) = 1$, additional motivation for the use of E_6 indeed comes from the fact that this Lie group is well known as a good candidate for the unifying symmetry group in models describing a unification of the non-gravitational fundamental forces of nature. Further, unlike the two larger exceptional Lie groups, E_7 and E_8 , the group E_6 has complex representations and these are needed to describe the observed multiplets of states in particle physics of equation 7.36 which are not left-right symmetric. From figure 7.2 it can be seen that E_6 contains $SO(10)$ and hence in turn $SU(5)$ and finally also the Standard Model gauge symmetry, with the chain of subgroups: $E_6 \supset SO(10) \supset SU(5) \supset SU(3) \times SU(2) \times U(1)$. The potential of E_6 as a unifying group has been known since the early history of the Standard Model of particle physics even as it was still taking shape in the 1970s (see for example [53]) and continues today (see also, for example [54] pp.302–308).

In this case the higher rank of E_6 over that of $SU(5)$, with 2 additional Dynkin nodes, suggests that in principle the physical phenomena of the rank-2 Lorentz transformations might be described alongside the rank-4 Standard Model gauge group within the full the rank-6 symmetry group E_6 . However it is not possible to break E_6 into the combined Lorentz and Standard Model algebras by the Dynkin analysis prescribed above. While it can be shown that E_6 contains subgroups such as $SU(3) \times SU(2) \times U(1) \times SU(3)$, for example by removing the central node in figure 7.2(a), a similar decomposition but with a rank-2 $SU(3)$ replaced by the rank-2 Lorentz group is not possible. An alternative prescription for obtaining semi-simple regular maximal subalgebras via an intermediate 'extended' Dynkin diagram does not help this situation. However, to some extent this Dynkin analysis oriented within the Cartan-Weyl basis for complex forms of the Lie algebras represents a ball-park picture and is not tailored to fit the fine details for a real form of E_6 represented within the context of a specific theory.

To study these details not only is the real form of the group action needed but also an understanding of how the dynamics arises, and the means by which a symmetry subgroup of $L(v_{27}) = 1$ might be associated with gauge field interactions, in order to account for the phenomena observed in the laboratory. In particular the structure of the symmetry breaking itself, involving the extended spacetime manifold M_4 , will need to be considered more explicitly. In the meantime, the observation that the Lorentz group and Standard Model gauge groups almost fit together at the level of this static Dynkin diagram analysis is an encouraging feature.

In principle then, the possibility of identifying features of the full gauge symmetry group for the strong and electroweak particle interactions for the theory presented here based on the E_6 symmetry of $L(\boldsymbol{v}_{27}) = 1$ is worth pursuing, as we explore in the following chapter. It is further noted that the 16 representation in table 7.1, that is the Majorana-Weyl spinor introduced in section 5.4 and described in the following section, as exemplified by the $Spin^+(1, 9)$ spinor θ of equations 6.26 and 8.2, possesses a branching pattern under the $SU(3) \times SU(2) \times U(1) \subset SO(10)$ subgroup into representation multiplets corresponding to the 15 particle types of a complete generation of Standard Model fermions of equation 7.36 (plus a right-handed neutrino). However a different approach will be followed here, involving both the incorporation of the external Lorentz symmetry within $Spin^+(1, 9)$ as well as the extension to the E_6 symmetry. Indeed we begin in the opening section of the following chapter by identifying objects which transform as fermions under the external symmetry.

We also note here the possible significance of the three possible embeddings of an h₂ Ω subspace, as represented by the components $X \in h_2 \Omega$ in equations 6.32, 6.34 and 6.35, within the space $h_3\mathbb{O}$, with equivalent symmetry transformation properties, and in particular three copies of the Spin⁺(1, 9) spinor θ representation. These three embeddings are related by the matrix $\mathcal T$ of equation 6.33, as described in section 6.4, and in terms the octonion triality isomorphism as discussed alongside equations 6.49 and 6.50, relating to the rich symmetry of this form of $L(v_{27}) = 1$. This is suggestive since we shall have to account for three generations of fermion families, related through the CKM matrix of equation 7.76 in the case of the quarks, which might here be related through the full set of E_6 symmetry transformations. On the other hand only one embedding of h₂C \subset h₃O will be associated with the local tangent space TM₄ in the symmetry breaking, potentially lifting the degeneracy between the three generations of fermions in the present theory.

Again, while the connection between some of these algebraic structures and the Standard Model is well known, here there is an underlying motivation for the origin of these mathematical forms in a physical theory based on the symmetries of $L(v_{27}) = 1$ representing a multi-dimensional form of temporal flow.

Considering then the demands from both ends of figure 7.1 at the same time, with the choice of $L_{\mathbb{C}}$ guided by general features of the Standard Model and the space R identified under a highly symmetric form of $L(\mathbf{v}) = 1$, we naturally converge upon the group E_6 acting on the representation space $h_3\mathbb{O}$, such that the matrix determinant is invariant, as being of particular interest. Indeed this motivated the detailed study in chapter 6 based on references [37, 38, 39, 40, 41]. Further, the identification of the Lorentz subgroup of E_6 acting upon the subspace $h_2\mathbb{C}$ representing 4-dimensional spacetime explicitly provides the symmetry breaking mechanism through which the broken internal subgroups of the larger symmetry may be realised as the local gauge groups. The symmetry breaking was pictured in figure 5.1 for the provisional model with an $SO^+(1,9)$ symmetry acting on the form $L(\mathbf{v}_{10}) = 1$. That case for a 10dimensional spacetime symmetry, now described by $Spin^+(1, 9)$ acting on $h_2\mathbb{O}$, constitutes a significant intermediate stage between the full 27-dimensional form of temporal flow and the external 4-dimensional spacetime structure.

In order to analyse the physical content of this theory it will be necessary to dissect the anatomy of the explicit real form of E_6 constructed in chapter 6 in the context of symmetry breaking over the extended M_4 manifold. In the following chapter we first study the action of the external Lorentz symmetry on the full set of h3O components, building on the analysis of equation 7.35 presented at the end of section 7.1, and then assess how the properties of the internal symmetry, surviving the symmetry breaking, compare with the Standard Model.

Chapter 8

E_6 Symmetry Breaking

8.1 External Symmetry on h_3 ^O

Having at hand a complete mathematical description of the E_6 symmetry action from chapter 6, preserving the determinant on the space h₃ \mathbb{O} as a form of $L(\mathbf{v}) = 1$, the physical significance of various subgroup actions can be considered locally with respect to the spacetime manifold M_4 . In particular a distinguished set of symmetry transformations will act on the components of $v_{27} \in h_3 \mathbb{O}$ lying in the local spacetime tangent space TM_4 . These transformations form the subgroup $SL(2,\mathbb{C})$, the double cover of the Lorentz group, which is identified then as the *external* symmetry group. This spacetime symmetry is central to general relativity, while in the flat spacetime limit these Lorentz transformations form a global symmetry on M_4 as for the theory of special relativity. With the flow of time expanded into the 27-dimensional space of 3×3 Hermitian octonion matrices h₃ \circ there are 23 *extra dimensions* beyond those needed to locate events taking place in our 4-dimensional spacetime world. The explicit action of the external Lorentz symmetry on all components of the space h3O will be described this section, based on the real form of E_6 as constructed in chapter 6.

The form of h₃ $\mathbb O$ matrices transforming under the type 1 $SL(2,\mathbb C)$ and $SL(2,\mathbb O)$ subgroups of E_6 , with the structure described in equation 6.29, is compatible with the isomorphism of vector spaces ([1] p.30):

$$
h_3 \mathbb{O} \quad \cong \quad \mathbb{R} \oplus h_2 \mathbb{O} \oplus \mathbb{O}^2 \tag{8.1}
$$

$$
\left(\begin{pmatrix} & & \\ & X & \\ & & \end{pmatrix}\begin{pmatrix} \theta \\ & \theta \end{pmatrix}\right) \rightarrow (n, X, \theta) \tag{8.2}
$$

 $27_{\text{E}_6} \rightarrow (1+10+16)_{\text{Spin}^+(1,9)}$ (8.3)

The three parts of this decomposition are respectively the scalar, vector and spinor

representations of the 10-dimensional spacetime symmetry group $SO^+(1, 9)$, for which the covering group is $Spin^+(1,9) \equiv SL(2, \mathbb{O})$. A spinor representation with both Majorana and Weyl properties is only possible for $d = (2, \text{ mod } 8)$ spacetime dimensions, as is the case for $SO^+(1, 9)$. The object θ corresponds to the Majorana-Weyl spinor representation, denoted 16, which can be described by 16 real numbers owing to the reality condition for Majorana spinors (in general a Majorana spinor ψ is one which is equal to its 'charge conjugate' ψ^c , this reality condition is also possible in 4-dimensional spacetime).

As described in [1] the decomposition of equations 8.1–8.3 gives a representation of Spin⁺(1,9) as linear transformations of h₃ Ω which do not preserve the Jordan algebra but do, importantly for the present considerations, preserve the determinant of $h_3\mathbb{O}$, as presented explicitly in equation 6.31 of section 6.4. The relationship between the complex Lie algebra $L_{\mathbb{C}} =$ so(10), its real forms, the group Spin⁺(1,9) and its representations was presented explicitly in table 7.1. Similarly as for so(10) in the Dynkin analysis of section 7.3 we can consider the above decomposition as a mathematically intermediary stage in studying the Lorentz subgroup, $Spin^+(1,3) \equiv SL(2,\mathbb{C})$, in E₆.

While the 27-dimensional irreducible representation of E_6 decomposes as a reducible representation of $Spin^+(1, 9)$, as shown in equation 8.3, further decomposition is to be expected under smaller subgroups such as the external Lorentz transformations of 4-dimensional spacetime considered in this section, and also for the internal symmetry groups to be identified in the following section.

We can identify the Lorentz 4-vector $v_4 = (v^0, v^1, v^2, v^3) \equiv h_2$ in the upper left-hand 2×2 matrix embedded within the larger 3×3 matrices in h₃ \mathbb{O} , as was the case for $h_2\mathbb{C} \subset h_3\mathbb{C}$ in equation 7.35. The relation $\det(\mathcal{X}) = 1$ with $\mathcal{X} \in h_3\mathbb{O}$ is preserved under operations of $SL(2, \mathbb{C})$ representing the Lorentz group upon this space as:

$$
\mathcal{X} \rightarrow \left(\begin{array}{c|c} S & 0 \\ \hline 0 & 1 \end{array}\right) \left(\begin{array}{cc|c} h^{00} & h^{01} + \overline{a}(6) & c \\ \hline h^{10} + a(6) & h^{11} & \overline{b} \\ \hline \overline{c} & b & n \end{array}\right) \left(\begin{array}{c|c} S^{\dagger} & 0 \\ \hline 0 & 1 \end{array}\right) \tag{8.4}
$$

with $h^{00} = v^0 + v^3$, $h^{01} = v^1 - v^2l$ $h^{10} = v^1 + v^2 l$, $h^{11} = v^0 - v^3$ (8.5)

with $S \in SL(2,\mathbb{C})$, and with '1' describing the identity transformation in the trivial 1-dimensional representation of this group, acting upon the components of $\mathcal X$ of equation 6.26. This action preserves the value of $\det(\mathbf{h}_2) = h^2$, as it is simply the transformation of equation 7.31, as well as leaving $\det(\mathcal{X}) = 1$ invariant. In equation 8.4 $a(6)$ denotes the 6-dimensional imaginary part of $a \in h_3 \mathbb{O}$ of equation 6.26, that is excluding the real $a_1 = v^1$ and imaginary $a_8 l = v^2 l$ components of $a \in \mathbb{O}$ which are associated with the external 4-vector $v_4 \in TM_4$.

The four components of the projected $v_4(x) \subset v_{27}(x)$, forming a tangent vector in TM_4 locally on the spacetime manifold M_4 , transform as the components of a Lorentz 4-vector. These components are embedded within the space $h_3\mathbb{O}$ via the 2×2 matrices $h_2 \in h_2\mathbb{C}$. While in section 7.1 $\{1, i\}$ denoted the base units for the space \mathbb{C} , for

example for σ^2 in equation 7.14 as used in equation 7.30 (and also in section 6.3, for example equation 6.19), here the preferred subspace $\mathbb{C} \subset \mathbb{O}$ basis is taken to be $\{1, l\}$ for v_4 , as indicated in equation 8.5, in conformity with the conventions of sections 6.4 and 6.5, and in particular equation 6.58, and as employed in the following section. Since the $SL(2,\mathbb{C})$ actions, based on this $\{1, l\}$ complex subspace are embedded in the 'type 1' location of equation 6.32 this group will be denoted $SL(2,\mathbb{C})^1$.

The full set of actions of the real form of E_6 on the space $h_3\mathbb{O}$ was constructed in section 6.4. With the group action of $SL(2,\mathbb{C})^1$ on $h_3\mathbb{O}$ in equation 8.4 embedded within the type 1 group action of $SL(2,\mathbb{O})^1$ on the same space as displayed in equation 6.29 we can write:

$$
SO^+(1,3) \equiv SL(2,\mathbb{C})^1 \subset SL(2,\mathbb{O})^1 \subset SL(3,\mathbb{O}) \equiv E_6 \tag{8.6}
$$

where the first '≡' strictly applies at the Lie algebra level. This shows explicitly how the action of the Lorentz group may be embedded within the higher symmetry group E_6 acting on the space h₃ \mathbb{O} . The direct physical interpretation of the former symmetry in the shape of the perceptual background of the spacetime manifold M_4 provides a direct source for the breakdown of the latter symmetry.

The six Lorentz group generators as a subset of the 78 E_6 generators were listed in equation 6.57 of section 6.5. They can be read off from the full E_6 Lie algebra table [38] and seen to satisfy the $SO⁺(1, 3)$ algebra which is reproduced here in table 8.1.

	$\left[\bullet,\bullet\right] \left[\begin{array}{cccc} \dot{R}_{z l} & \dot{R}_{x z} & \dot{R}_{x l} & \dot{B}_{t x} & \dot{B}_{t l} & \dot{B}_{t z} \end{array}\right]$			
	$\dot{R}_{z l}$ 0 $-\dot{R}_{x l}$ $\dot{R}_{x z}$ 0 $-\dot{B}_{t z}$ $\dot{B}_{t l}$			
\dot{R}_{xz}		$\dot{R}_{x\perp} \qquad 0 \qquad -\dot{R}_{z\perp} \qquad \dot{B}_{tz} \qquad 0 \qquad -\dot{B}_{tx}$		
	$\begin{array}{ccc} \dot{R}_{x l} & -\dot{R}_{x z} & \dot{R}_{z l} & 0 & -\dot{B}_{t l} & \dot{B}_{t x} & 0 \\ 0 & -\dot{B}_{t z} & \dot{B}_{t l} & 0 & \dot{R}_{x l} & -\dot{R}_{x z} \end{array}$			
		\dot{B}_{tz} 0 $-\dot{B}_{tx}$ $-\dot{R}_{xl}$ 0 \dot{R}_{zl}		
	$\begin{vmatrix} -\dot{B}_{\rm H} & \dot{B}_{\rm fx} & 0 & \dot{R}_{\rm zz} & -\dot{R}_{\rm zd} & 0 \end{vmatrix}$			

Table 8.1: (Extracted from the E_6 Lie algebra table in [38]). The Lie algebra structure for the set of Lorentz generators of equation 6.57, with bracket composition $[\dot{R}_{z}^1, \dot{R}_{xz}^1] = -\dot{R}_{x}^1$ etc. The type superscripts '1' are omitted in the table entries, which are all generators of $SL(2,\mathbb{C})^1$. (Each entry is equivalent to that for the corresponding 6×6 table for the generators $\{J^1, -J^2, J^3, -K^1, K^2, -K^3\}$ with the Lie bracket of equations 7.17–7.19, via the correspondence of equation 8.9).

The corresponding 2×2 matrix actions for the category 1 boosts and category 2 rotations can be read off for the case $q = l$ in table 6.1 of section 6.3. Since each of these actions involves the composition of matrix elements from a single complex subspace, with base units $\{1, l\}$, and with each of $a, b, c \in \mathbb{O}$ (or $p, m, n \in \mathbb{R}$) as elements of $h_3\mathbb{O}$ appearing in separate product terms, the symmetry transformations are equivalent to those based on H subalgebras and are hence associative. Consistent with the discussion in the paragraphs following equation 6.35 this means that the symmetry group and corresponding Lie algebra can be represented in terms of the transformation matrices themselves. A matrix representation for the Lorentz Lie algebra is therefore provided by defining $\dot{M} = \frac{\partial}{\partial \alpha} M \Big|_{\alpha=0}$ for the corresponding six matrix actions in table 6.1 (here presented in a different order), that is:

$$
\dot{M}_{2l} = \begin{pmatrix} 0 & -\frac{l}{2} \\ -\frac{l}{2} & 0 \end{pmatrix}, \qquad \dot{M}_{22} = \begin{pmatrix} 0 & +\frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix}, \qquad \dot{M}_{2l} = \begin{pmatrix} -\frac{l}{2} & 0 \\ 0 & +\frac{l}{2} \end{pmatrix}, \qquad (8.7)
$$

$$
\dot{M}_{Lx} = \begin{pmatrix} 0 & +\frac{1}{2} \\ +\frac{1}{2} & 0 \end{pmatrix}, \qquad \dot{M}_{Lx} = \begin{pmatrix} 0 & +\frac{l}{2} \\ -\frac{l}{2} & 0 \end{pmatrix}, \qquad \dot{M}_{Lz} = \begin{pmatrix} +\frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}, \quad (8.8)
$$

where the latter three are the boost generators as can be identified by the time component 't' label in the subscript. Expressing the three Pauli matrices as $\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma^2 = \binom{0 - l}{l \; 0}, \, \sigma^3 = \binom{1 \; 0}{0 \; -1}$ $0 - 1$), that is equation 7.14 with i replaced by the imaginary unit l , the six elements of this Lorentz Lie algebra can be written as:

$$
\begin{pmatrix}\n\dot{M}_{\mathbf{z}} \\
\dot{M}_{\mathbf{z}\mathbf{z}} \\
\dot{M}_{\mathbf{z}\mathbf{z}}\n\end{pmatrix} = \begin{pmatrix}\n-\frac{l}{2}\sigma^1 \\
+\frac{l}{2}\sigma^2 \\
-\frac{l}{2}\sigma^3\n\end{pmatrix} \sim \begin{pmatrix}\n+J^1 \\
-J^2 \\
+J^3\n\end{pmatrix}, \qquad \begin{pmatrix}\n\dot{M}_{\mathbf{z}\mathbf{z}} \\
\dot{M}_{\mathbf{z}\mathbf{z}}\n\end{pmatrix} = \begin{pmatrix}\n+\frac{1}{2}\sigma^1 \\
-\frac{1}{2}\sigma^2 \\
+\frac{1}{2}\sigma^3\n\end{pmatrix} \sim \begin{pmatrix}\n-K^1 \\
+K^2 \\
-K^3\n\end{pmatrix}
$$
\n(8.9)

The associations with the Lorentz rotation ${J^a}$ and boost ${K^a}$ generators of equation 7.16 are such that with $\{J^2, K^1, K^3\} \rightarrow \{-J^2, -K^1, -K^3\}$ the Lie algebra of equations 7.17–7.19 matches that of the commutators in table 8.1. Hence in the context of the SL(2, \mathbb{C})¹ action in equation 8.4 these sign conventions, $\{\dot{R}, \dot{B}\} \sim \pm \{J, K\},$ in equation 8.9 should be noted, which at the group level simply corresponds to a sign flip for a subset of the six real parameters $\{r_a, b_a\}$ in equation 7.20, and hence in turn will relate to the definition of left and right-handed spinors. As described in the discussion following table 6.5 in section 6.5 here the key orientation for such conventions is provided by the E_6 Lie algebra table of reference [38] from which table 8.1 is extracted. Ultimately a different set of $L(E_6)$ sign conventions may be preferred, in alignment with the physical application.

We next address the action of the external Lorentz symmetry on a general element $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$, including the full set of 16 real components of $\theta = \begin{pmatrix} \varepsilon \\ b \end{pmatrix} \in \mathbb{O}^2$, that is the 16-dimensional Majorana-Weyl spinor under $SL(2,\mathbb{O})^1$, composed of the octonion entries c and \bar{b} , as introduced in equation 6.26 and described after equations 8.1–8.3.

The two-sided $SL(2,\mathbb{C})^1$ action on $h_2\mathbb{O}$ in equation 8.4 only transforms the real diagonal entries h^{00} and h^{11} together with the $h^{10} = a_1 + a_8l$ and $h^{01} = a_1 - a_8l$ components of $a \in \mathbb{O}$. The six components of $a(6) \in \text{Im}(a)$ remain invariant as may be deduced from the form of the six $SL(2,\mathbb{C})^1$ generators in table 6.6 or from equation 6.54 for the case $q = l$. (This is equivalent to the invariance of \mathbf{v}_6 under $SO^+(1,3)$ for the model of figure 5.1). Of the additional 17 components in h₃ \circ the real diagonal *n* entry is also invariant, as is clear from equation 8.4, while all 16 components of $b, c \in \mathbb{O}$ transform non-trivially under the one-sided $SL(2,\mathbb{C})^1$ action.

The spinor $\theta_l = \begin{pmatrix} c \\ b \end{pmatrix}_l \in \mathbb{C}^2$ will denote the $\{1, l\}$ components of c and \bar{b} in θ , that is $\binom{c}{b} \in \mathbb{O}^2$ restricted to the $\{1, l\}$ complex subspace. By comparison with equation 7.35 this object transforms as a left-handed Weyl spinor $\psi_L = \theta_l$ under the $SL(2,\mathbb{C})^1$ action in equation 8.4. Consistent with the above comments on the sign conventions for the Lorentz generators here we take this $S \in SL(2,\mathbb{C})^1$ action on θ_l to define the left-handed spinor representation, guided but not constrained by the standard definitions of section 7.1.

Due to the anticommuting property, for example in equation 7.1, Clifford algebras are also related to the division algebras ([1] section 2.3), with analogous rotational properties as alluded to following equation 7.4. Indeed it can be shown, for example, that $C(0, 2) = \mathbb{H}$ for the Clifford algebra associated with the 2-dimensional vector space $\mathbb{R}^{0,2}$, while $C(1,3) = \mathbb{H}(2)$, that is the Clifford algebra for 4-dimensional spacetime is isomorphic to the algebra of 2×2 quaternionic matrices under multiplication. (However, since $C(p, q)$ is in all cases an *associative* algebra there are no such isomorphisms involving the octonion algebra).

As a representation of $C(1,3)$ the algebra $\mathbb{H}(2)$ acts, by matrix multiplication, on the spinor space \mathbb{H}^2 rather than the usual Dirac spinor space \mathbb{C}^4 . We consider first the quaternionic spinor as a subspace of the octonionic spinor $\theta = \begin{pmatrix} c \\ b \end{pmatrix}$ with base units $\{1, l, i, u\}$:

$$
\theta_{\mathbb{H}} = \begin{pmatrix} c \\ \overline{b} \end{pmatrix}_{\mathbb{H}} = \begin{pmatrix} c_1 + c_8l + c_2i + c_7il \\ b_1 - b_8l - b_2i - b_7il \end{pmatrix} \in \mathbb{H}^2 \subset \mathbb{O}^2 \tag{8.10}
$$

Upon restriction to the subset of 2×2 matrix actions of $SL(2,\mathbb{C})^1 \subset SL(2,\mathbb{H})^1 \subset \mathbb{H}(2)$ (which is isomorphic to the group $Spin^+(1,3)$ as identified within the Clifford algebra $C(1,3) = \mathbb{H}(2)$, with base units $\{1, l\}$, the spinor space $\theta_{\mathbb{H}}$ decomposes into two parts:

$$
\theta_l = \begin{pmatrix} c_1 + c_8 l \\ b_1 - b_8 l \end{pmatrix} \text{ and } \theta_i = \begin{pmatrix} c_7 \mathbf{i} l + c_2 i \\ -b_7 \mathbf{i} l - b_2 i \end{pmatrix}
$$
 (8.11)

which transform independently. As was described for equation 6.14 the group actions are considered as active transformations. Further, under the left action by the imaginary unit l the components of c transform as:

$$
(c_1 + c_8l) \rightarrow l(c_1 + c_8l) = (-c_8 + c_1l)
$$

\n
$$
(c_7il + c_2i) \rightarrow l(c_7il + c_2i) = (-c_2il + c_7i)
$$
\n(8.12)

and hence the (c_7, c_2) components of θ_i transform under left multiplication by l in an identical manner to the respective components (c_1, c_8) of θ_l , which is also trivially true for multiplication by the real unit 1. This observation applies also to the \bar{b} components of θ_l and θ_i , while the structure of the 2 × 2 matrix action of $SL(2,\mathbb{C})^1$ applies in the same way on both of these objects. In fact the transformations of θ_l and θ_i in equation 8.11 are identical both for the generators of $SL(2,\mathbb{C})^1$ and for the finite group actions, as can be readily seen by explicit calculation. For example applying the Lorentz symmetry rotation matrix $M_{z}(\alpha)$ from table 6.1 to θ_l and θ_i results in the respective transformations:

$$
R_{zI}(\alpha) \theta_{l} = \begin{pmatrix} \cos\frac{\alpha}{2} & -l\sin\frac{\alpha}{2} \\ -l\sin\frac{\alpha}{2} & \cos\frac{\alpha}{2} \end{pmatrix} \begin{pmatrix} c_{1} + c_{8}l \\ b_{1} - b_{8}l \end{pmatrix}
$$

\n
$$
= \begin{pmatrix} (\cos\frac{\alpha}{2}c_{1} - \sin\frac{\alpha}{2}b_{8}) & + & (-\sin\frac{\alpha}{2}b_{1} + \cos\frac{\alpha}{2}c_{8})l \\ (\cos\frac{\alpha}{2}b_{1} + \sin\frac{\alpha}{2}c_{8}) & + & (-\sin\frac{\alpha}{2}c_{1} - \cos\frac{\alpha}{2}b_{8})l \end{pmatrix}
$$

\n
$$
R_{zI}(\alpha) \theta_{i} = \begin{pmatrix} \cos\frac{\alpha}{2} & -l\sin\frac{\alpha}{2} \\ -l\sin\frac{\alpha}{2} & \cos\frac{\alpha}{2} \end{pmatrix} \begin{pmatrix} c_{7}il + c_{2}i \\ -b_{7}il - b_{2}i \end{pmatrix}
$$

\n
$$
= \begin{pmatrix} (\cos\frac{\alpha}{2}c_{7} - \sin\frac{\alpha}{2}b_{2})il + (\sin\frac{\alpha}{2}b_{7} + \cos\frac{\alpha}{2}c_{2})i \\ (-\cos\frac{\alpha}{2}b_{7} + \sin\frac{\alpha}{2}c_{2})il + (-\sin\frac{\alpha}{2}c_{7} - \cos\frac{\alpha}{2}b_{2})i \end{pmatrix}
$$

Here it can be seen that the four real coefficients $\{c_1, c_8, b_1, -b_8\}$ of the spinor θ_l map onto the components of $R_{\mathcal{A}}(\alpha)\theta_l$ in precisely the same way that the coefficients $\{c_7, c_2, -b_7, -b_2\}$ of θ_i map onto the components of $R_{\alpha}(a)\theta_i$. A similar observation applies for θ_l and θ_i under the remaining five Lorentz symmetry actions. Hence as well as the original left-handed Weyl spinor θ_l the components of θ_i also transform exactly as a left-handed spinor of $SL(2,\mathbb{C})^1$. This representation of $SL(2,\mathbb{C})$ on the two lefthanded spinors θ_l and θ_i of equation 8.11 in \mathbb{H}^2 contrasts with the representation constructed in equations 7.15 and 7.29 on the left and right-handed spinors ψ_L and ψ_R in \mathbb{C}^4 .

Considering the further two quaternionic subspaces with base units $\{1, l, j, j\}$ and $\{1, l, k, k\}$ it can be seen that the original full octonionic spinor $\theta = \left(\frac{c}{b}\right)$, with 16 real components, reduces to a total of four left-handed Weyl spinors under the action of $SL(2,\mathbb{C})^1$, augmenting the set in equation 8.11 to:

$$
\theta_l = \begin{pmatrix} c_1 + c_8l \\ b_1 - b_8l \end{pmatrix}, \ \theta_i = \begin{pmatrix} c_7il + c_2i \\ -b_7il - b_2i \end{pmatrix}, \ \theta_j = \begin{pmatrix} c_6jl + c_3j \\ -b_6jl - b_3j \end{pmatrix}, \ \theta_k = \begin{pmatrix} c_5kl + c_4k \\ -b_5kl - b_4k \end{pmatrix}
$$
\n(8.13)

There is an equivalent decomposition for a corresponding set of conjugate spinors in $\theta^{\dagger} = (\bar{c} \; b)$ under the right action of S^{\dagger} on $h_3 \mathbb{O}$ as implied in equation 8.4. This set of four Weyl spinors in equation 8.13 will be important for interpreting further symmetries, *internal* to the action of $SL(2,\mathbb{C})^1$ on $h_3\mathbb{O}$, in the following section.

In this section we have described how the decomposition of the 27 representation of E_6 under the subgroup $Spin^+(1, 9)$ of equation 8.3 further reduces under the subgroup $SL(2, \mathbb{C})^1$ as summarised in table 8.2.

This may be compared with the $SL(2,\mathbb{C})^1$ action on the subspace $h_3\mathbb{C} \subset h_3\mathbb{O}$ as described in equation 7.35 for which the nine real components of h_3C transform as one 4-vector h_2 , one Weyl spinor ψ_L and one scalar n (closely relating to v_4 , θ_l and n respectively in table 8.2). The six extra scalars and three extra spinors in table 8.2 result from the additional $27 - 9 = 18$ real components in h₃ \mathbb{O} . In both cases each Weyl spinor, as for the space \mathbb{C}^2 , has four real parameters.

	$Spin^+(1,9)$	$SL(2,\mathbb{C})^1$	Components	
	1 scalar	$(0,0)$ scalar	\boldsymbol{n}	
	10 vector	$\left(\frac{1}{2},\frac{1}{2}\right)$ vector	\boldsymbol{v}_4	
		$6 \times (0,0)$ scalars	a(6)	
16	spinor	$4 \times (\frac{1}{2}, 0)$ spinors	$\theta_{l,i,j,k}$	

Table 8.2: The further decomposition of the $(1 + 10 + 16)$ representation of $Spin^+(1,9) \subset E_6$ of equation 8.3 under the subgroup of external 4-dimensional spacetime symmetry $SL(2, \mathbb{C})^1 \subset Spin^+(1, 9)$ actions of equation 8.4, and the corresponding components of $h_3\mathbb{O}$ transformed.

The spinor components of $\theta_{l,i,j,k}$ represent 'internal' dimensions of the space h₃ \mathbb{O} in the sense that, unlike $v_4 \in TM_4$, they are not tangent to the external spacetime M_4 , but they *do* transform in a non-trivial manner, as spinors, under the external $SL(2,\mathbb{C})$ ¹ symmetry, and in this sense they are not purely internal objects. This feature for the cubic form of temporal flow $L(v_{27}) = 1$ is hence distinct from that seen for a quadratic form with a spacetime symmetry. For example for the 10-dimensional spacetime form considered in section 5.1 the external $SO^+(1,3)$ symmetry acts on the external $\overline{v}_4 \subset v_{10}$ components *only*, as pictured in figure 5.1(b) and applies also for the corresponding gauge field in equation 5.51, as for the external symmetry of any higher-dimensional spacetime structure. For the present theory based on temporal progression, here taking a cubic form, of particular interest in the following section will be the nature of the internal symmetry transformations on the four $SL(2,\mathbb{C})^1$ spinors from the final line of table 8.2.

8.2 Internal $SU(3)_c \times U(1)_Q$ Symmetry

Physically the $SL(2,\mathbb{C})$ ¹ symmetry studied in section 8.1 is considered 'external' as it is the two-to-one cover of the Lorentz group which in the full theory acts on the tangent space TM_4 of the extended 4-dimensional spacetime manifold. This structure is central to the theory of general relativity and gravitation, as described in sections 3.3, 3.4 and 5.3. On the other hand the 'internal' symmetry will consist of further subgroups of E_6 , which will be central to the structure of local gauge theories and the Standard Model of particle physics as reviewed in the previous chapter.

At the end of the previous section the branching of the 16 representation of $Spin⁺(1, 9)$ into a set of four Weyl spinors under the external Lorentz subgroup $SL(2,\mathbb{C})^1$ was described, as listed in table 8.2. Independently it is also known that the same $Spin⁺(1, 9)$ Majorana-Weyl 16-dimensional representation branches into a set of multiplets describing the 15 states of one generation of Standard Model quarks and leptons, as listed in equation 7.36, together with a right-handed neutrino, all expressed uniformly in terms of left-handed fields, under the internal subgroup $SU(3)_c \times SU(2)_L \times$ $U(1)_Y$, as noted in section 7.3, but it is not the approach we follow here. In this section we consider the internal symmetry derived from the subgroup Stab(TM_4) ⊂ E₆, defined below, and its relation to the set of four Weyl spinors derived from the external symmetry $SL(2, \mathbb{C})^1 \subset Spin^+(1, 9)$.

In contrast to the 6 generators of the external $SL(2,\mathbb{C})^1$ symmetry, of the remaining $(78 - 6) = 72$ generators of E₆ those which leave all tangent space vectors $v_4 \in TM_4$ untouched may literally be considered to constitute an *internal* symmetry, surviving the symmetry breaking, and are expected to be significant for the physics of local gauge theories. While leaving TM_4 invariant these internal symmetries will in general have non-trivial actions on the remaining, 'extra dimensions' within v_{27} , through which we may seek to identify a relation with the phenomena of physical particle interactions as observed in the laboratory and described by the Standard Model.

Here then, as a preliminary definition, and in contrast to the external symmetry, the internal symmetry will be obtained from the set of all E_6 actions on $h_3\mathbb{O}$ which leave the four components for any $v_4 = (v^0, v^1, v^2, v^3)$ in equation 8.5 (that is, $h_2 \in h_2 \mathbb{C}$ of equation 8.4) invariant. These components, including $v^2 \equiv a_8$ associated with the imaginary unit l of $a \in \mathbb{O}$, can also be expressed in the combination (p, m, a_1, a_8) with respect to the parametrisation of equation 6.1. The corresponding symmetry group is complementary to the actions of $SL(2,\mathbb{C})^1$ and will be denoted $Stab(TM_4)$ as the stability group of all vectors $v_4 \in TM_4$. By inspection from tables 6.6 and 6.7, for the 78 elements in the preferred basis for the Lie algebra of E_6 defined on the space $T_{\text{h}_3\text{O}}$, the group Stab (TM_4) is generated by the 31 elements listed in table 8.3. In particular we shall be looking to identify closed subgroups within $Stab(TM₄)$ for which each generator is independent of $SL(2,\mathbb{C})^1$ in terms of Lie bracket composition.

Category 1 and 2: Boosts and Rotations	#	
$(\dot{R}_{xz}^2 - \dot{B}_{tx}^2), \qquad (\dot{R}_{xz}^3 + \dot{B}_{tx}^3)$	$\overline{2}$	
$(\dot{R}_{zq}^2 + \dot{B}_{tq}^2), \qquad (\dot{R}_{zq}^3 - \dot{B}_{tq}^3)$	14	
Category 3: Transverse Rotations		
\dot{A}_q , \dot{G}_l , \dot{S}_l^1	9	
$(\dot{G}_q + 2\dot{S}_q^1)$ $q = \{i, j, k, kl, \dot{\jmath}, d\}$	6	
Total	31	

Table 8.3: The Lie algebra generators of the 31 dimensional group Stab $(TM₄)$. The subscript q denotes any of the seven imaginary octonion units $\{i, j, k, k, \mu, \mu, \nu\}$ unless stated otherwise.

The 16 vector fields on T_{13} ^O generating the Category 1 and 2 elements of Stab(TM_4) are written out explicitly in equations 8.14 and 8.15 in which the invariant action on the 4-dimensional subspace $h_2\mathbb{C} \subset h_3\mathbb{O}$ is clear. (In fact they leave all 10

components of $h_2 \mathbb{O} \subset h_3 \mathbb{O}$ invariant, see also [41] equations 4.12(27) and 4.13(28)).

$$
\dot{R}_{xz}^2 - \dot{B}_{tx}^2 = \begin{pmatrix} 0 & 0 & -\bar{a} \\ 0 & 0 & -m \\ -a & -m & -2b_x \end{pmatrix}, \quad \dot{R}_{xz}^3 + \dot{B}_{tx}^3 = \begin{pmatrix} 0 & 0 & p \\ 0 & 0 & a \\ p & \bar{a} & 2c_x \end{pmatrix}
$$
\n
$$
\dot{R}_{zq}^2 + \dot{B}_{tq}^2 = \begin{pmatrix} 0 & 0 & \bar{a}q \\ 0 & 0 & mq \\ -qa & -mq & -2b_q \end{pmatrix}, \quad \dot{R}_{zq}^3 - \dot{B}_{tq}^3 = \begin{pmatrix} 0 & 0 & pq \\ 0 & 0 & aq \\ 0 & 0 & aq \\ -pq & -q\bar{a} & 2c_q \end{pmatrix}
$$
\n(8.15)

Of the 15 transverse rotations in table 8.3 the first 9 are basis vectors which explicitly leave the components of h_2 invariant, while for each of the remaining six $(\dot{G}_q + 2\dot{S}_q^1)$ actions the non-zero $\dot{a}_8 l$ contributions in table 6.7 cancel.

Although the category 1 and 2 transformations of type 1 as originally composed on the 10-dimensional space $h_2\mathbb{O}$ each act as a simple rotation or boost in a 2-dimensional plane the effect on the components of the spinor θ is less straightforward in the full $h_3\mathbb{O}$ action, as was seen for the case of the external symmetry $SL(2,\mathbb{C})^1$ in the previous section. This is also seen for the type 2 and 3 internal transformations of equations 8.14 and 8.15. Hence these actions, together with the 15 internal transverse rotations, stir up the θ components in non-trivial ways.

Of particular interest is the SU(3) subgroup introduced below equation 6.42 and discussed shortly after figure 7.3 (as described in [38] pp.115 and 136, following [55]). This SU(3) is defined in terms of the transverse rotations acting on the octonion space \mathbb{O} alone as the subgroup SU(3) \subset G₂ of the octonion automorphism group that leaves one imaginary unit, here l , invariant. The corresponding Lie algebra $su(3)$ is described by the set of 8 generators $\{\dot{A}_q, \dot{G}_l\}$ which, as transformations of E_6 on the full space h₃O, act on each of the octonion elements $a, b, c \in \mathbb{O}$ in the same way leaving invariant the complex $\{1, l\}$ subspaces, and as elements of table 8.3 identified within stab(TM_4) may be provisionally associated with the colour $su(3)_c$ of the Standard Model. This algebra is also independent of $SL(2,\mathbb{C})^1$ in terms of the Lie bracket composition, that is $[X, Y] = 0$ for all $X \in sl(2, \mathbb{C})^1$ and $Y \in sl(3)_{c}$, and hence we have the semi-simple subgroup:

$$
SL(2,\mathbb{C})^1 \times SU(3)_c \subset E_6 \tag{8.16}
$$

The Lie algebra composition of the $\{\dot{A}_q, \dot{G}_l\} \in \text{su}(3)_{c}$ elements from the E₆ commutation table in [38] is reproduced here in table 8.4. The Lie algebra in table 8.4 is isomorphic to the su(3) Lie algebra represented by the eight 3×3 Gell-Mann matrices listed in table 8.5.

The two algebras in tables 8.4 and 8.5 are identical within the choice of sign conventions, numerical coefficients and the fact that the Gell-Mann matrices are taken to be Hermitian. The latter property results in an extra factors of i accompanying the λ_{α} matrices in the algebra isomorphism listed in table 8.6. The factors of i belong to the same complex algebra C used in the components of the Gell-Mann matrices themselves, but are independent of the octonion algebra elements on the left-hand side. (That is the isomorphism is between the basis $\{\dot{A}_q, \dot{G}_l\}$ and the anti-Hermitian matrices $\sim i\lambda_\alpha$

$\left[\bullet,\bullet\; \right]$ $\left.\rule{0pt}{12pt}\right]$			\dot{A}_i \dot{A}_j \dot{A}_k \dot{A}_{kl} \dot{A}_{jl} \dot{A}_{il} \dot{A}_l \dot{G}_l			
	\dot{A}_i 0 \dot{A}_k $-\dot{A}_j$ $-\dot{A}_{jl}$ \dot{A}_{kl} $\dot{A}_l - \dot{G}_l$ $-\dot{A}_{il}$ $3\dot{A}_{il}$					
\dot{A}_i	$-\dot{A}_k$		$0 \qquad \qquad \dot{A}_i \qquad -\dot{A}_{i l} \qquad -\dot{A}_l - \dot{G}_l \qquad \dot{A}_{kl} \qquad \qquad \dot{A}_{j l} \qquad 3 \dot{A}_{j l}$			
	\dot{A}_k \dot{A}_j $-\dot{A}_i$ 0 $-2\dot{A}_l$ $-\dot{A}_{il}$ \dot{A}_{il} $2\dot{A}_{kl}$ 0					
	\dot{A}_{kl} $\begin{vmatrix} \dot{A}_{jl} & \dot{A}_{il} & 2\dot{A}_{l} & 0 & -\dot{A}_{i} & -\dot{A}_{j} & -2\dot{A}_{k} & 0 \end{vmatrix}$					
\dot{A}_{il}	$\left \begin{array}{cccc} -\dot{A}_{kl} & \dot{A}_l + \dot{G}_l & \dot{A}_{il} & \dot{A}_i & 0 \end{array} \right $				$-\dot{A}_k$ $-\dot{A}_j$ $-3\dot{A}_j$	
\dot{A}_{il}	$-\dot{A}_l + \dot{G}_l$ $-\dot{A}_{kl}$ $-\dot{A}_{jl}$ \dot{A}_j			\dot{A}_k 0 \dot{A}_i -3 \dot{A}_i		
	\dot{A}_l $\dot{A}_{\underline{d}}$ $-\dot{A}_{\underline{d}}$ $-2\dot{A}_{\underline{d}}$ $2\dot{A}_k$		\dot{A}_i		$-\dot{A}_i \hspace{1.6cm} 0 \hspace{1.4cm} 0$	
	$\dot G_l \quad \Big \quad - 3 \dot A_{\not\perp} \quad \quad - 3 \dot A_{\not\perp} \quad \quad \ \ 0 \qquad \quad \ \ \, 0$		$3\dot{A}_i$		$3\dot{A}_i \hspace{1.6cm} 0 \hspace{1.6cm} 0$	

Table 8.4: (Extracted from the E_6 Lie algebra table in [38]). The Lie algebra structure for the SU(3)_c generators $\{\dot{A}_q, \dot{G}_l\}$, with bracket composition $[\dot{A}_i, \dot{A}_j] = \dot{A}_k$ etc.

$$
\lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}
$$

$$
\lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \quad \lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}
$$

$$
\lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \quad \lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}
$$

$$
\text{with } [\lambda_\alpha, \lambda_\beta] = i f_{\alpha\beta\gamma} \lambda_\gamma \quad \text{and} \quad f_{123} = 2, f_{458} = f_{678} = \sqrt{3},
$$

$$
f_{147} = -f_{156} = f_{246} = f_{257} = f_{345} = -f_{367} = 1
$$

Table 8.5: The set of eight complex Hermitian Gell-Mann matrices of su(3), with representatives of the completely antisymmetric structure constants $f_{\alpha\beta\gamma}$ which are non-zero.
rather than directly with the Hermitian Gell-Mann matrices. This is analogous to the relation between the external $SL(2, \mathbb{C})^1$ generators and the conventional Lorentz algebra in equation 8.9, where factors of i would also appear if the J^a were defined as Hermitian rather than anti-Hermitian in equation 7.16).

$$
\begin{array}{|c|c|} \hline \dot{A}_k \sim -i\lambda_1 & \dot{A}_\mu \sim -i\lambda_2 & \dot{A}_l \sim i\lambda_3 \\ \\ \dot{A}_i \sim -i\lambda_4 & \dot{A}_\mu \sim i\lambda_5 \\ \\ \dot{A}_\mu \sim -i\lambda_6 & \dot{A}_j \sim -i\lambda_7 & \dot{G}_l \sim -i\sqrt{3}\lambda_8 \\ \hline \end{array}
$$

Table 8.6: The isomorphism between the su(3)_c \subset E₆ Lie algebra basis $\{\dot{A}_q, \dot{G}_l\}$ and the eight Gell-Mann matrices λ_{α} ([38] p.137, table 4.5).

The 3×3 Gell-Mann matrices transform the components of complex vectors $u \in \mathbb{C}^3$ corresponding, in the context of an $\mathrm{SU}(3)_c$ gauge theory, to the interactions between 'red', 'blue' and 'green' quark states encountered in quantum chromodynamics. Similarly the $\{\dot{A}_q, \dot{G}_l\}$ algebra elements, as transformations on the space h₃ \mathbb{O} mix the components of the Spin⁺(1,9) spinor $\theta = \begin{pmatrix} \frac{c}{b} \end{pmatrix} \in \mathbb{O}^2$. For example the tangent vector field \dot{A}_i on the $\binom{c}{b}$ components of h₃ \mathbb{O} , obtained from table 6.7, are:

$$
\dot{A}_i : \begin{pmatrix} \dot{c} \\ \dot{b} \end{pmatrix} = \begin{pmatrix} \dot{c}_1 + \dot{c}_8 l, & +\dot{c}_7 \dot{\mathbf{u}} + \dot{c}_2 i, & +\dot{c}_6 \dot{\mathbf{u}} + \dot{c}_3 j, & +\dot{c}_5 \dot{\mathbf{u}} + \dot{c}_4 k \\ \dot{b}_1 - \dot{b}_8 l, & -\dot{b}_7 \dot{\mathbf{u}} - \dot{b}_2 i, & -\dot{b}_6 \dot{\mathbf{u}} - \dot{b}_3 j, & -\dot{b}_5 \dot{\mathbf{u}} - \dot{b}_4 k \end{pmatrix}
$$

$$
= \begin{pmatrix} 0 + 0l, & +0\dot{\mathbf{u}} + 0i, & -c_5 \dot{\mathbf{u}} - c_4 j, & +c_6 \dot{\mathbf{u}} + c_3 k \\ 0 - 0l, & -0\dot{\mathbf{u}} - 0i, & +b_5 \dot{\mathbf{u}} + b_4 j, & -b_6 \dot{\mathbf{u}} - b_3 k \end{pmatrix} (8.17)
$$

The components here have been ordered to match those of the four left-handed Weyl spinors $(\theta_l, \theta_i, \theta_j, \theta_k)$ of equation 8.13. The fact that each real component of c transforms in the same way as the corresponding component of b is expected since $SU(3)_c$ acts on each of $a, b, c \in \mathbb{O}$ in precisely the same way. However, it is also noted that the action \dot{A}_i in equation 8.17 respects the 4-way spinor decomposition, with for example \dot{c}_6 and \dot{c}_3 of $\dot{\theta}_j$ taking the respective values of $-c_5$ and $-c_4$ from the spinor θ_k . This apparently non-trivial observation applies to all eight $SU(3)_c$ generators, which hence represent a mixing of the four Weyl spinors, as a structure maintained within the mixing of the eight real components of the octonion elements.

The extraction of the components of a spinor θ into a matrix of real numbers will be denoted by $[\theta]$. For example, from equation 8.13 the spinor θ_i can be mapped to the 2×2 matrix of real numbers $[\theta_i] = \begin{pmatrix} c_7 & c_2 \\ -b_7 & -b_8 \end{pmatrix}$ $-b_7-b_2$ (with components ordered to match those of the spinor θ_l under $SL(2,\mathbb{C})$ ¹ transformations, as described for equations 8.11– 8.13). With this notation and the Lorentz spinor definitions in equation 8.13 the above equation 8.17 can be expressed as:

$$
\dot{A}_i : [\dot{\theta}] = ([\dot{\theta}_l], [\dot{\theta}_i], [\dot{\theta}_j], [\dot{\theta}_k]) \n= (0_2, 0_2, -[\theta_k], [\theta_j])
$$
\n(8.18)

where 0_2 represents the 2×2 zero matrix. This expression shows explicitly how the internal SU(3)_c generator \dot{A}_i mixes the external SL(2, \mathbb{C})¹ spinors θ_j and θ_k identified in the previous section. The tangent vectors of all eight generators $\{\dot{A}_q, \dot{G}_l\}$ of $SU(3)_{c}$ on the spinor space $\theta \in \mathbb{O}^2$ are listed in table 8.7 alongside the actions of the Gell-Mann matrices, using the correspondence in table 8.6, on the vectors $u \in \mathbb{C}^3$. On the left-hand side the elements $\{\dot{A}_q, \dot{G}_l\}$ are already expressed as tangent vectors, while on the right-hand side the tangents are obtained by matrix multiplication of the λ_{α} into $u \in \mathbb{C}^3$.

		$([\dot{\theta}_l], \qquad [\dot{\theta}_i], \qquad [\dot{\theta}_j], \qquad [\dot{\theta}_k])$		$(\dot{u}_1, \dot{u}_3,$		\dot{u}_3)
	$A_i = (0_2, 0_2, -[\theta_k],$ $A_{il} = (0_2, 0_2, [l\theta_k],$	$[\theta_i]$ $[l\theta_j]$	\sim \sim	$\lambda_4 \Rightarrow (u_3, 0,$ $\lambda_5 \Rightarrow (-iu_3, 0,$		u_1) iu_1)
		$A_j = (0_2, \quad [\theta_k], \quad 0_2, \quad -[\theta_i])$ $A_{\underline{i}\underline{l}} = (0_2, -[l\theta_k], 0_2, -[l\theta_i])$	\sim \sim	$\lambda_7 \Rightarrow (0,$ $\lambda_6 \Rightarrow (0, u_3,$	$-iu_3,$	iu_2) u_2)
	$A_k = (0_2, -[\theta_j], [\theta_i],$ $A_{kl} = (0_2, \quad [l\theta_j], \quad [l\theta_i],$	$0_2)$ $0_2)$	\sim \sim	$\lambda_1 \Rightarrow (u_2, u_1, 0)$ $\lambda_2 \Rightarrow (-iu_2, iu_1, 0)$		
	$\dot{A}_l = (0_2, \quad [l\theta_i], \quad -[l\theta_j],$	0_2) $G_l = (0_2, \quad [l\theta_i], \quad [l\theta_j], \quad -2[l\theta_k])$	\sim \sim	$\lambda_3 \Rightarrow (u_1,$ $\lambda_8 \Rightarrow \frac{1}{\sqrt{3}}(\ u_1, \quad u_2, \quad -2u_3)$		$-u_2, 0)$

Table 8.7: The tangent vector generators for the SU(3) representations on \mathbb{O}^2 and \mathbb{C}^3 . The column vectors of \mathbb{C}^3 are displayed as a row vectors for convenience in the table.

In table 8.7 a term such as $[l\theta_i]$ denotes multiplying the spinor θ_i on the left by l before extracting the coefficients of $l\theta_i$ with the Im(\mathbb{O}) units ordered as in equation 8.13. This notation is used to isolate the mixing effect on the real number coefficients, with care for the joint effects of the division algebra composition as well as matrix algebra composition. For the case of $u \in \mathbb{C}^3$, the two real degrees of freedom for each of the u_1, u_2, u_3 belong to the *same* complex space $\mathbb C$ (with base units $\{1, i\}$) but occupy different components of the 1×3 column matrix vector \mathbb{C}^3 . For the case of $\theta \in \mathbb{O}^2$ the four real degrees of freedom for each of the $\theta_l, \theta_i, \theta_j, \theta_k$ belong to a *different* 2dimensional subspace of \mathbb{O} (with base units $\{1, l\}$, $\{\underline{i}, i\}$, $\{\underline{j}, j\}$, $\{k, k\}$ respectively) but occupy the *same* components of the 1×2 column matrix vector \mathbb{O}^2 .

Hence, as seen in table 8.7, the six transformations \dot{A}_q ($q \neq l$) mix the components of the three Weyl spinors $\theta_i, \theta_j, \theta_k$ in a similar manner that the Gell-Mann

matrices λ_{α} ($\alpha \neq 3, 8$) mix the three \mathbb{C}^3 components u_1, u_2, u_3 , with the correspondence between the objects of each representation space depending on the form of the isomorphism in table 8.6, which is arbitrary up to the automorphism group of $su(3)$. In both cases there are two remaining diagonal generators as listed at the bottom of table 8.7. (The physics here is determined by the $\{\dot{A}_q, \dot{G}_l\}$ transformations as generators of $SU(3)_c$ rather than the particular choice of correspondence with the λ_α matrices, as was similarly the case for the external $SL(2,\mathbb{C})^1$ action of the previous section as described after equation 8.9). In the case of the full set of $\{\dot{A}_q, \dot{G}_l\}$ acting on the components of $h_3\mathbb{O}$ there is a copy of the same set of mixing transformations within the components of the Hermitian conjugate spinor $\theta^{\dagger} = (\bar{c} \, b)$ of equation 6.26 which also transforms under the internal $SU(3)_c$ symmetry (similarly as described for the $SL(2,\mathbb{C})^1$ spinors below equation 8.13).

In conclusion the internal $SU(3)_c$ symmetry action in the left-hand column of table 8.7 dovetails neatly with the external $SL(2,\mathbb{C})^1$ spinor structure of equation 8.13. The mixing action of $SU(3)_c$ in table 8.7 takes a form summarised as:

$$
\theta = (\theta_l, \quad \underbrace{\theta_i, \quad \theta_j, \quad \theta_k}_{\text{SU(3)}_c \text{ action}})
$$
\n(8.19)

which implies that as a gauge theory the $SU(3)_c$ internal symmetry will mediate interactions between the Weyl spinors $\theta_i, \theta_j, \theta_k$, transforming under the fundamental representation, which in turn will hence be identified with the three colour degrees of freedom of the quark states. On the other hand the invariance of θ_l , transforming under the trivial representation of $SU(3)_c$, suggests that these components should be associated with the leptonic sector of the Standard Model (with the subscript l originating from the $\{1, l\}$ base units for θ_l also then serving as a mnemonic for its leptonic character). Further aspects of the Standard Model might then be expected to be uncovered by exploring further aspects of the internal symmetry group within E_6 , which will occupy the remainder of this chapter.

In particular the Standard Model Abelian gauge group $U(1)_Q$, underlying Maxwell's equations and the phenomena of electromagnetism, might also be sought as an internal symmetry within $Stab(TM_4)$. Of the 31 generators for the internal symmetry group Stab(TM_4) listed in table 8.3 there is a $(31-8) = 23$ -dimensional set which as a vector space is independent of the internal $SU(3)_c$ generators. Of these 23 there are 3 sets each of 6 elements:

$$
(\dot{R}_{zq}^2 + \dot{B}_{tq}^2), \qquad (\dot{R}_{zq}^3 - \dot{B}_{tq}^3), \qquad (\dot{G}_q + 2\dot{S}_q^1) \tag{8.20}
$$

with $q \neq l$, totalling 18 elements each of which fails to commute with some of the internal $SU(3)_c$ generators in the set $\{\dot{A}_q, \dot{G}_l\}$. As a cross-check this observation appears to hold for any linear combination of elements selected from the 18 in equation 8.20, by further inspection of the E_6 Lie algebra table [38]. Hence none of the 18 elements in equation 8.20 can belong to a group which may be appended to the subgroup decomposition $SL(2,\mathbb{C})^1 \times SU(3)_{c}$ in equation 8.16 (in fact the first 12 elements in equation 8.20 also fail to commute with $SL(2,\mathbb{C})^1$). This then leaves a set of only $(31 - 8 - 18) = 5$ internal generators which in terms of Lie algebra composition, and not only as a vector space, is independent of $su(3)_c$. These are the elements:

$$
(\dot{R}_{xz}^2 - \dot{B}_{tx}^2), \quad (\dot{R}_{xz}^3 + \dot{B}_{tx}^3), \quad (\dot{R}_{z\dot{z}}^2 + \dot{B}_{t\dot{z}}^2), \quad (\dot{R}_{z\dot{z}}^3 - \dot{B}_{t\dot{z}}^3), \quad \dot{S}_l^1 \tag{8.21}
$$

Indeed, each of the nine individual component parts listed within equation 8.21 commute with all eight elements of the internal $su(3)_c$ basis set. However the first 4 elements in equation 8.21 each fail to commute with the external $SL(2,\mathbb{C})^1$ generators. This leaves \dot{S}_l^1 as the *only* E₆ Lie algebra generator of Stab(*TM*₄) which is independent of both $SL(2,\mathbb{C})^1$ and $SU(3)_c$. Hence of the many possible $U(1) \subset E_6$ subgroups the one generated by \dot{S}^1_l is identified as the most suitable candidate for the internal $\mathrm{U}(1)_Q$ gauge symmetry of electromagnetism.

Moreover the generator \dot{S}_l^1 is also closely associated with the diagonal symmetry action S_l^1 , described by equation 6.43, and leaves the 4-dimensional spacetime components in $h_2\mathbb{C}$ invariant as a residual of the $SL(2,\mathbb{O})$ action on $h_2\mathbb{O}$ as described at the end of section 6.3. A similar internal $U(1)$ symmetry associated with electromagnetism has been considered for the $SL(2,\mathbb{C}) \times U(1)$ gauge theories as discussed in the opening paragraphs of section 7.3. While as elements of the vector space $Th_3\mathbb{O}$ we have $\dot{S}_l^1 = \dot{S}_l^1$, as discussed following equation 6.45, the group actions $S_l^{(1)}$ $l_l^{(1)}(\alpha)$ and $\mathcal{S}^{(1)}_l$ $\ell_l^{(1)}(\alpha)$ diverge at $O(\alpha^2)$ and in any case, although suggestive, this argument alone is insufficient in itself to associate the $U(1)_Q$ symmetry with S_l^1 out of many possible $U(1) \subset E_6$ subgroups. Here the main case for this association is the observation that the U(1) subgroup S^1_l uniquely both belongs to $\text{Stab}(TM_4)$ and at the Lie algebra level is independent of the $SL(2,\mathbb{C})^1 \times SU(3)_{c}$ subgroup of equation 8.16.

Hence here the internal $U(1)$ generated by \dot{S}_l^1 is a natural candidate to consider for the $U(1)_Q$ component of the Standard Model gauge symmetry group. From table 6.7 it can be seen that the generator \dot{S}_l^1 impacts on all 8 real components of both c and \overline{b} of $\theta \in \mathbb{O}^2$. In fact, and in comparison with equation 8.17, the tangent vector \dot{S}_l^1 on the spinor components $\theta = \begin{pmatrix} c \\ b \end{pmatrix}$ is given explicitly by:

$$
\dot{S}_{l}^{1}:\begin{pmatrix}\n\dot{c} \\
\dot{b}\n\end{pmatrix} = \begin{pmatrix}\n\dot{c}_{1} + \dot{c}_{8}l, & +\dot{c}_{7}\dot{d} + \dot{c}_{2}i, & +\dot{c}_{6}\dot{d} + \dot{c}_{3}j, & +\dot{c}_{5}kl + \dot{c}_{4}k \\
\dot{b}_{1} - \dot{b}_{8}l, & -\dot{b}_{7}\dot{d} - \dot{b}_{2}i, & -\dot{b}_{6}\dot{d} - \dot{b}_{3}j, & -\dot{b}_{5}kl - \dot{b}_{4}k\n\end{pmatrix}
$$
\n
$$
= \begin{pmatrix}\n-\frac{3}{2}c_{8} + \frac{3}{2}c_{1}l, & +\frac{1}{2}c_{2}\dot{d} - \frac{1}{2}c_{7}i, & +\frac{1}{2}c_{3}\dot{d} - \frac{1}{2}c_{6}j, & +\frac{1}{2}c_{4}kl - \frac{1}{2}c_{5}k \\
\frac{3}{2}b_{8} + \frac{3}{2}b_{1}l, & -\frac{1}{2}b_{2}\dot{d} + \frac{1}{2}b_{7}i, & -\frac{1}{2}b_{3}\dot{d} + \frac{1}{2}b_{6}j, & -\frac{1}{2}b_{4}kl + \frac{1}{2}b_{5}k\n\end{pmatrix}
$$
\n
$$
(8.22)
$$
\nwith $[\dot{\theta}] = \begin{pmatrix}\n+\frac{3}{2} \left[l\theta_{l} \right], & -\frac{1}{2} \left[l\theta_{i} \right], & -\frac{1}{2} \left[l\theta_{j} \right], & -\frac{1}{2} \left[l\theta_{k} \right] \n\end{pmatrix}$

which may be compared with the $\mathrm{su}(3)_{c}$ action on θ in equation 8.18 and table 8.7. Here the two components of $c \in \mathbb{Q}$ within each of the four Weyl spinors are mixed, and similarly for the corresponding pair of $\bar{b} \in \mathbb{O}$ components, with no mixing of components between different spinors. This is consistent with the nature of the electromagnetic interaction which does not transform between different fermion types.

A further observation from equation 8.23 regards the factor of $\frac{3}{2}$ found for the θ_l spinor in contrast to the factors of $\frac{1}{2}$ aligned with the three remaining spinors θ_i , θ_j , θ_k . Hence, with \dot{S}^1_l provisionally associated with electromagnetism and by comparison with equation 8.19, the apparent 'electromagnetic charge' assigned to the leptonic sector is three times larger than that assigned to the quark sector. Associating θ_i , θ_j , θ_k with the three colour states of a d-quark this observation in principle accounts for the 'fractional charge' of magnitude $\frac{1}{3}$ as theoretically ascribed and empirically confirmed for d-quark states relative to the electron charge. Based on this observation we introduce the notation:

$$
\dot{S}_l^a = \frac{2}{3}\dot{S}_l^a\tag{8.24}
$$

(for $a = 1, 2, 3$) such that the above charge values $\frac{3}{2}$ and $\frac{1}{2}$ are normalised to 1 and 1 $\frac{1}{3}$ under \dot{S}_l^1 , representing the generator of $U(1)_Q$, for ease of comparison with the Standard Model convention for which the electron charge is −1. The 'bar' through \dot{S}_l^1 is a mnemonic symbol for this normalisation of fractional charges relative to the e^- charge. (The corresponding normalisation for components of the *group* action S_l^1 , which is not needed here, would need to take into account the *nested* composition of equation 6.39. This group normalisation would hence be different for the single action of S_l^1 of equation 6.43).

Hence the subgroup in equation 8.16 may be augmented to:

$$
SL(2,\mathbb{C})^1 \times SU(3)_c \times U(1)_Q \subset E_6 \tag{8.25}
$$

with the internal group $SU(3)_c \times U(1)_Q$ generated by $\{\dot{A}_q, \dot{G}_l, \dot{S}_l^1\} \in \text{stab}(TM_4)$. The action of this larger internal symmetry on the four $SL(2,\mathbb{C})^1$ spinors also augments equation 8.19 as:

$$
\theta = (\theta_l, \quad \underbrace{\theta_i, \quad \theta_j, \quad \theta_k}_{U(1)_Q})
$$

\nSU(3)_c : 1 3 (8.26)
\nU(1)_Q : +1 - $\frac{1}{3}$ - $\frac{1}{3}$ - $\frac{1}{3}$

With the generator \dot{S}_l^1 hence associated with electromagnetic charge it is instructive to consider this action on the full set of $h_3\mathbb{O}$ components. From table 6.7 the diagonal components of \dot{S}_l^1 are trivial, with $\dot{p} = \dot{m} = \dot{n} = 0$, while action on the remaining components $a, b, c \in \mathbb{O}$, via equation 8.24, may be summarised as:

$$
\dot{S}_l^1 = \begin{pmatrix} \dot{a} \\ \dot{b} \\ \dot{c} \end{pmatrix} = \begin{pmatrix} 0 \, l \, a_{1,l} + \frac{2}{3} \, l \, a(6) \\ -1 \, l \, b_{1,l} - \frac{1}{3} \, l \, b(6) \\ +1 \, l \, c_{1,l} - \frac{1}{3} \, l \, c(6) \end{pmatrix} \tag{8.27}
$$

where $a_{1,l} \equiv (a_1 + a_8l)$ and $a(6) \equiv (a_7i\ell + a_2i + a_6j\ell + a_3j + a_5k\ell + a_4k)$, with similar expressions for b and c, following the component order of the spinors in equation 8.13. The same definition of $a(6)$ is implied in equation 8.4. By comparison with the above discussion leading to equation 8.26 the expression for \dot{S}^1_l in equation 8.27 incorporates 'charges' of 0 and $\frac{2}{3}$ for the \dot{a} components, that is we have:

$$
a = (a_{1,l}, \underbrace{a_{\underline{d},i}, a_{\underline{d},j}, a_{\underline{d},k}}_{\text{SU(3)}_c})
$$

\nSU(3)_c: 1 3
\nU(1)_Q: 0 + $\frac{2}{3}$ + $\frac{2}{3}$ + $\frac{2}{3}$ (8.28)

where the $SU(3)_c$ action on $a \in \mathbb{O}$ is identical to that on the octonion components of $\theta = \begin{pmatrix} c \\ b \end{pmatrix}$ in equation 8.26. While physical lepton states are invariant under SU(3)_c and

are hence associated with the Weyl spinor θ_l in equation 8.26, the neutrino states are also invariant under the $U(1)_{\mathcal{Q}}$ of electromagnetism, that is with zero charge, and are provisionally associated with the $a_{1,l}$ components in equations 8.27 and 8.28; while a set of u-quarks with $\frac{2}{3}$ fractional charges is similarly associated with the $a(6)$ components.

However, unlike $\theta = \begin{pmatrix} c \\ b \end{pmatrix}$ the $a \in h_3\mathbb{O}$ component does not correspond to a set of $SL(2,\mathbb{C})$ ¹ Weyl spinors, as can be seen from table 8.2. Further, the 'neutrino' components $a_{1,l} = a_1 + a_8l = v^1 + v^2l$ have already apparently been accounted for as part of the external vector $v_4 \in TM_4$ on the base manifold, as described in equations 8.4 and 8.5. These features clearly require further investigation.

While in the Standard Model the e^- lepton charge is -1 and the d-quark charge is $-\frac{1}{3}$, with positive charges for their antimatter counterparts, the convention and interpretation of the \pm -signs of equations 8.23 and 8.26 will depend upon the conventions used and the identification of particle and antiparticle states as relating to the spacetime dynamics of the theory. As for GUT theories in which particle and antiparticle states may coexist within the same $SU(5)$ multiplet [43], see sections 7.2 and 7.3, the apparently opposite charges in equation 8.23 may relate, for example, to a combination of 'antimatter' electrons and 'matter' d-quarks in the components of $h_3\mathbb{O}$ (which may in turn ultimately relate to the nature of the asymmetry between matter and antimatter in the universe).

Within the above caveats, aligned with the charges of 1 and $\frac{1}{3}$ for the electron and d-quark Weyl spinors of equation 8.26 the respective $U(1)_Q$ charges of 0 and $\frac{2}{3}$ in equation 8.28 correlate with charges of $\binom{0}{-1}$ −1) for the $\binom{\nu}{e}$ e^{ν} lepton doublet and $\binom{+2/3}{-1/3}$) for the $\binom{u}{d}$ $_d^u$) quark doublet of the Standard Model. In addition the states associated with each *left-handed* doublet of charges interact via the exchange of W^{\pm} gauge bosons in the Standard Model. Hence it remains to be understood how interactions within each of these doublets may be mediated via an $SU(2)_L$ symmetry, and how such ν -lepton and u-quark components of $a \in \mathbb{O} \subset h_3\mathbb{O}$ gain a Weyl spinor structure under the external $SL(2,\mathbb{C})^1$ action.

While the empirical charge structure of the Standard Model fermions is in principle accounted for by a $U(1)_Q$ symmetry associated with the generator \dot{S}_l^1 of equation 8.27, further elaboration of this theory is required in order to further reconstruct the pattern of particle multiplets listed in equation 7.36. Guided by the Standard Model it will be necessary to understand the origin of weak interactions in order to address these details. Hence in the following section we investigate the possible identification of an $SU(2)_L$ gauge symmetry within the structure of the broken E_6 action on $h_3 \mathbb{O}$ in the present theory.

8.3 Elements of Electroweak Theory

8.3.1 SU(2) Transformations and $SU(3)$, Symmetry

Within the set of 31 internal basis elements in table 8.3 it is possible to identify a number of SU(2) subgroups, for example generated by the three elements $\dot{G}_q + 2\dot{S}_q^1$ with $q = \{i, j, k\}$ or a different triplet of imaginary units (excluding l) belonging to a common line in figure 6.1 and hence generating a quaternion subalgebra. While

independent of $\{A_q, G_l\}$ as a vector space none of these su(2) generator sets is independent of the su(3)_c algebra in terms of the Lie bracket (that is with $[X, Y] = 0$ for all $X \in \text{su}(3)_c, Y \in \text{su}(2)$, as discussed after equation 8.20.

It is an open question whether all possible internal symmetry subgroups should have physical significance. In the above case the generator $\dot{G}_q + 2\dot{S}_q^1$ for $q = i, j$ and k mixes the components of θ_l with those of θ_i , θ_j and θ_k respectively, hence mixing between 'leptons' and 'quarks', and would apparently correspond to 'new physics' with respect to the Standard Model. However this particular SU(2) action does not describe a 'fundamental representation' on the set four Weyl spinors, as was the case for $SU(3)_c$ on the left-hand side of table 8.7 or for $U(1)_Q$ in equation 8.23.

In any case here we attempt to identify an $SU(2)$ symmetry which, as for the case of the \dot{S}^1_l generator identified for equation 8.25 for an internal $\mathrm{U}(1)_Q$ symmetry, is independent of the internal $SU(3)_c$. The other four internal generators in equation 8.21 form a trivial algebra with zero Lie bracket for all products – although non-zero commutators are obtained if \dot{S}_l^1 is included (with $[(\dot{R}_{xz}^2 - \dot{B}_{zx}^2), \dot{S}_l^1] = \frac{3}{2}(\dot{R}_{zt}^2 + \dot{B}_{tl}^2)$ for example) but this is still insufficient structure to form an su(2) algebra. It is also the case that none of these four elements commute with $SL(2,\mathbb{C})^1$ and in fact none of the 31 − 9 = 22 remaining elements of Stab(TM_4) in table 8.3 commute with the subgroup $SL(2,\mathbb{C})^1 \times SU(3)_c$ of equation 8.16, as implied in the discussion following equation 8.21. Further, it is to be expected from the Dynkin analysis described in section 7.3 that in fact there is no possibility of identifying an $SU(2)$ subgroup of E_6 which is independent of *both* an external $SL(2, \mathbb{C})$ and an internal $SU(3) \times U(1)$ symmetry group.

However, although the full internal gauge symmetry group of the Standard Model reads $SU(3)_c \times SU(2)_L \times U(1)_Y$ there are a number of features of weak interactions associated with $SU(2)_L$, as observed in high energy physics experiments and written into the Standard Model, which qualitatively differ from the strong and electromagnetic interactions associated with $SU(3)_c$ and $U(1)_Q$ respectively. If an internal SU(2) were to be found at this stage, at the level of symmetry groups and their representations, in a similar manner as for the internal $SU(3)_c \times U(1)_Q$ in the previous section, it seems unlikely that the kind of distinctive properties observed for the weak interactions could arise purely in the dynamics of the full theory. The differences in empirical properties between the strong and electromagnetic interactions themselves originate largely out of the differences between the non-Abelian SU(3) and Abelian U(1) symmetries at the group and representation level, with many more interactions possible in the quantum theory for the former case. However while the non-Abelian group $SU(2)$ is mathematically intermediate in size between $SU(3)$ and $U(1)$ the physically observed features associated with the gauge group SU(2) are of a quite different nature, as described in section 7.2 and summarised in the following paragraph.

Firstly the weak interactions violate parity symmetry, prompting the subscript 'L' for the left-handed character of this chiral $SU(2)_L$ gauge theory. Secondly, the Standard Model $SU(2)_L$ is closely association with a $U(1)_Y$ gauge symmetry, with $U(1)_Q$ surviving the electroweak symmetry breaking, which is in turn associated with the Lorentz scalar Higgs field transforming as an $SU(2)_L$ doublet and providing the mechanism by which three gauge bosons, the W^{\pm} and Z^{0} , gain a non-zero mass. Thirdly, the weak interactions mix particle states from the three distinct generations of fermions, as described by the CKM matrix.

Here we initially focus upon the simple fact that weak $SU(2)_L$ transformations act on fermion doublets of the form $\binom{\nu}{e}$ $\binom{u}{e}$ and $\binom{u}{d}$ $\binom{u}{d}$, which have been associated with the \int_{0}^{a} $\binom{a}{\theta}$ components of h₃ $\mathbb O$ for the present theory. This was described at the end of the previous section where it was noted that the $U(1)_Q$ electromagnetic charges associated with the \dot{S}_l^1 action on $\theta = \begin{pmatrix} c \\ b \end{pmatrix}$ and the a component are respectively aligned with the charges of the (e -lepton, d -quark) and (ν -lepton, u -quarks) particle states.

The type 1 $SL(2,\mathbb{C})^1$ action on the four Weyl spinors of equation 8.13 is complemented by $SL(2,\mathbb{C})^2$ and $SL(2,\mathbb{C})^3$ transformations of type 2 and 3, all involving quaternion algebra composition with $l \in \mathbb{O}$ being the only imaginary octonion unit appearing in the transformation matrices. Two $SU(2)$ s are immediately identifiable in terms of the rotation subgroups of the type 2 and type 3 Lorentz groups, as denoted by SU(2)², generated by the set $\{\dot{R}_{z_l}^2, \dot{R}_{xz}^2, \dot{R}_{x_l}^2\}$, and SU(2)³, as generated by $\{R_{\underline{z}\underline{l}}^3, R_{\underline{x}\underline{z}}^3, R_{\underline{x}\underline{l}}^3\}$. Neither SU(2)² ⊂ SL(2, C)² nor SU(2)³ ⊂ SL(2, C)³ is independent of $\overline{\text{SL}(2,\mathbb{C})^1}$ within the E₆ Lie algebra, with for example $[\dot{R}_{xz}^2, \dot{R}_{xz}^1] = \frac{1}{2}\dot{R}_{xz}^3 \neq 0$, and neither of them forms a subgroup of $Stab(TM_4)$, and hence they do not appear to form an internal symmetry by the original definition which led to table 8.3. However owing to the properties described below in exploring further the structure of these transformations the groups $SU(2)^{2,3}$ are found to be of some interest in relation to the structure of electroweak theory.

By reference to equations 6.32, 6.34 and 6.35 of section 6.4, and with the spinor components $\theta^a = \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$ $\begin{bmatrix} \theta_1 \\ \theta_2 \end{bmatrix} \in \mathbb{Q}^2$ represented by $\left(\frac{c}{b}\right), \left(\frac{a}{c}\right)$ $\binom{a}{\bar{c}}$ and $\binom{b}{\bar{a}}$ $\binom{b}{\bar{a}}$ for the type $a = 1, 2$ and 3 transformations respectively, the three types of $M^{(a)} \in SL(2,\mathbb{C})^a$ action, with each set generated by equations 8.7 and 8.8, are of the form:

$$
\left(M^{(1)}\right)\left(\begin{array}{c}c\\ \bar{b}\end{array}\right), \qquad \left(M^{(2)}\right)\left(\begin{array}{c}a\\ \bar{c}\end{array}\right), \qquad \left(M^{(3)}\right)\left(\begin{array}{c}b\\ \bar{a}\end{array}\right) \qquad (8.29)
$$

with an equivalent right composition $\theta^{\dagger}M^{\dagger} = (M\theta)^{\dagger}$ associated with each action above, as seen in the example of the full type 1 embedding of equation 6.29. In all cases however the group action is by *left* translation, that is with group representations $R(g_1)R(g_2) = R(g_1g_2)$ as discussed in section 6.2 after equation 6.14, and involves elements of the non-commutative quaternion algebra.

The type 1 action of $SL(2,\mathbb{C})^1$ decomposes the space $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix} \in \mathbb{O}^2$ into the four Weyl spinors of equation 8.13. The transformations $SL(2,\mathbb{C})^{2,3}$ of type 2 and 3, with complementary transformation matrices also based on the units $\{1, l\}$, similarly respect the octonion decomposition aligned to the four base unit sets:

$$
\{1, l\}, \{i\}, \{j\}, \{j\}, \{k\}, \} \tag{8.30}
$$

based on the same quarternion subalgebras, now for all three of $a, b, c \in \mathbb{O}$. Hence the subgroups $SU(2)^{2,3} \subset E_6$ describe transformations between the components of equation 8.26 and those of equation 8.28 respecting the alignment of the four component pieces, and hence acting independently on the corresponding doublets of leptonic and quark states as appropriate for weak interactions. With respect to the embedding of $a, b, c \in \mathbb{O}$ as components of h₃ \mathbb{O} in equation 6.1, the spinor representation mixing actions of $SL(2,\mathbb{C})^{1,2,3}$ can also be displayed graphically as:

This again shows how the $\binom{c}{b}$ spinor components under $SL(2,\mathbb{C})^1$ are replaced by $\binom{a}{b}$ and $\binom{b}{a}$ spinors under $SL(2,\mathbb{C})^2$ and $SL(2,\mathbb{C})^3$ respectively, depending on the \bar{c}) and \bar{a} alignment of the $\theta^a = \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix}$ $\begin{bmatrix} \theta_1 \\ \theta_2 \end{bmatrix}$ components in equations 6.32–6.35. It is the observation that the SL(2, \mathbb{C})^{2,3} actions relate the $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix} \in \mathbb{O}^2$ components with the $a \in \mathbb{O}$ component in equations 8.29 and 8.31, while respecting the four-way octonion decomposition of equation 8.30, that suggests that these transformations might be closely related to the weak interactions.

In section 8.1 the Weyl spinors $\theta_i, \theta_j, \theta_k$ were identified alongside θ_l in equation 8.13 originating from the *one-sided* action of $SL(2, \mathbb{C})^1 \subset SL(2, \mathbb{H})^1 \subset \mathbb{H}(2)$ on $\left(\frac{c}{b}\right)_{\mathbb{H}} \in \mathbb{H}^2$. The quark spinors $\theta_i, \theta_j, \theta_k$ are formed out of subspaces of \mathbb{H}^2 with quarternion base units $\{\underline{\mathbf{i}}, i\}, \{\underline{\mathbf{i}}, j\}, \{\underline{\mathbf{k}}, k\} \in \mathbb{O}$ respectively, with the actions on these objects by matrices composed of the base units $\{1, l\}$, completing the 3 sets of H subalgebras of the octonions, involving in particular the quaternionic left multiplication by l as demonstrated in equation 8.12. Although the full set of $\mathbb{H}(2)$ matrix actions are not involved this asymmetric one-sided action is apparently incomplete in terms of the set of possible actions of the non-commutative quaternion algebra on these components.

This observation might in principle relate to a possible mechanism for the origin of chirality in SU(2) interactions in the Standard Model. This situation can be contrasted with left-right symmetric gauge theories with the internal symmetry group $\text{SU}(2)_L \times \text{SU}(2)_R \times \text{U}(1)$ formulated in terms of fields defined over the quaternion algebra (see for example [56] and the references therein). A mechanism is then required through which the symmetry in these parity conserving models is broken to $SU(2)_L \times U(1)$ to match the observed parity violating phenomena of weak interactions.

Here since the set of *external* Lorentz transformations of $SL(2,\mathbb{C})^1$ act asymmetrically on the *left* on $\theta^1 \in \mathbb{O}^2$ and on the subspaces of Weyl spinors $\theta_i, \theta_j, \theta_k$ we may expect to identify a set of actions on these spinor components algebraically composed from the right, which have a complementary effect owing to the non-commuting property of the H algebra, potentially forming a distinct internal symmetry, at least with regards to the quark states represented by these three Weyl spinors. Similarly, in the present context, for actions involving multiplications by elements belonging to $SL(2, \mathbb{H})^a \subset \mathbb{H}(2)$ some *chiral* behaviour might be expected to arise in this theory as the type $a = 2, 3$ actions complement the symmetry breaking action of the external Lorentz transformations $SL(2,\mathbb{C})^1 \subset SL(2,\mathbb{H})^1$. Further, although only one fermion generation has been considered explicitly, the structure of equation 8.31 is suggestive in terms of the need to ultimately account for the CKM mixing between three generations of fermions.

However while these possibilities provided some of the initial motivation for studying the actions of the groups $SU(2)^{2,3}$ the mechanism for the above physical phenomena will require further developments. The source of parity violation in the present theory will be described in section 9.2, having explicitly constructed both left and right-handed Weyl spinors by extending the form of temporal flow beyond the action of E_6 on h₃ \mathbb{O} . As will be described in section 9.3 a further expansion to a yet higher-dimensional flow of time may be required in order to account for three generations of fermions and the phenomena of CKM mixing.

Here the main motivation for studying the SU(2)^{2,3} ⊂ E₆ subgroups is the structure of the action on the doublet components of $h_3\mathbb{O}$ as described for equations 8.29–8.31 above in relation to the weak interaction transformations for doublets of fermions in the Standard Model. In this subsection we hence further explore this group structure before focusing on a pattern of symmetry breaking that closely parallels the properties of electroweak symmetry breaking in the remainder of this section. In particular the subgroup $SU(3)_c \times SU(2)^2 \times U(1)^2 \subset E_6$, provisionally considered as an 'internal symmetry' (where $U(1)^2$ is the type 2 equivalent of $U(1)^1 = U(1)_Q$ identified in the previous section), is analogous to the Standard Model gauge symmetry $SU(3)_c \times SU(2)_L \times U(1)_Y$; with the impingement of the action $SU(2)^2 \times U(1)^2$ on the external spacetime components of h₂C ⊂ h₃^O breaking this symmetry down to U(1)_{*Q*}. This will be described in the following subsection and constitutes a 'mock electroweak theory'. We will then ultimately need to address how to combine these structures with the external $SL(2,\mathbb{C})^1$ symmetry, which within the E_6 structure is *not* independent of the $SU(2)^{2,3}$ actions.

In fact with $SU(2)^a \subset SL(2,\mathbb{C})^a$ for $a = 1,2,3$ these structures are found together with three types of $U(1)^a$ action described by \dot{S}^a_l for $a=1,2,3$, as introduced in equation 8.24, within the full E_6 action on the space h₃ \mathbb{O} . The SU(3)_c action of table 8.7, corresponding to the set of eight generators $\{\dot{A}_q, \dot{G}_l\}$, not only transforms $a, b, c \in \mathbb{O} \subset h_3\mathbb{O}$ in precisely the same way (table 6.7), acting on the components of $a(6)$ as a triplet and $a_{1,l}$ as a singlet (in the notation of equation 8.27), but is also independent of both the $SL(2,\mathbb{C})^{1,2,3}$ and $S_l^{1,2,3}$ ^{1,2,3} actions in terms of the E_6 algebra Lie bracket. This means that the $SL(2,\mathbb{C})^a$ and S_l^a actions may effectively be stripped out and considered independently of the $SU(3)_c$ action. This may aid the identification of an internal $SU(2)_L \times U(1)_Y$ symmetry, and its relation with the external Lorentz symmetry $SL(2,\mathbb{C})^1$, bearing in mind that the former is expected to be 'broken' to the $U(1)_Q$ symmetry associated with \dot{S}_l^1 .

The nine generators of the combined type $a = 1, 2$ and 3 rotations $SU(2)^a$ form a closed subalgebra of E_6 , which is eight dimensional due to the linear dependence of the \dot{R}_{nl}^a generators as displayed in equation 6.46. This subalgebra is in fact an su(3), a linearly independent basis for which can be described by the eight rotation generators ([38] p.128):

$$
su(3)_s \equiv \{\dot{R}_{\text{nl}}^1, \dot{R}_{\text{nl}}^2, \dot{R}_{\text{nz}}^1, \dot{R}_{\text{nz}}^2, \dot{R}_{\text{nz}}^3, \dot{R}_{\text{nl}}^3, \dot{R}_{\text{nl}}^1, \dot{R}_{\text{nl}}^2, \dot{R}_{\text{nl}}^3\} \tag{8.32}
$$

These generate a group denoted $SU(3)_s$ (where 's' denotes the 'standard' representation or embedding of this group in E₆ [38]). As implied above within E₆ the subgroup $SU(3)_s$ is independent of the colour subgroup $SU(3)_c$, as generated by the eight elements of table 8.4, with the Lie bracket composition of any element of equation 8.32 with any element of $\{\dot{A}_q, \dot{G}_l\}$ being zero. The generators of $SU(3)_c$ are explicitly 'type

independent', in that there are no type labels on any of the eight generators $\{\dot{A}_q, \dot{G}_l\}$ ([38] p.128), none of which distinguish between the three types. The subgroup $SU(3)_s$ is also 'type independent', in that all three types play an equivalent role, however the individual generators do carry type labels as for example in equation 8.32.

The group product $SU(3)_s \times SU(3)_c \subset E_6$ is a rank-4 subgroup of the complete rank-6 symmetry group E_6 . In fact $SU(3)_s \subset SL(3,\mathbb{C})_s$ where $SL(3,\mathbb{C})_s$ is the 16dimensional rank-4 group generated by the type 1, 2 and 3 rotations of equation 8.32 together with the a linearly independent set of the type 1, 2 and 3 boosts also based on the $\{1, l\}$ complex subspace. Taking into account equation 6.51 we have ([38] p.128):

$$
sl(3,\mathbb{C})_s \equiv su(3)_s \cup \{\dot{B}_{1z}^1, \dot{B}_{1z}^2, \dot{B}_{1x}^1, \dot{B}_{1x}^2, \dot{B}_{1x}^3, \dot{B}_{1z}^1, \dot{B}_{2z}^2, \dot{B}_{2z}^3\}
$$
(8.33)

In fact $sl(3,\mathbb{C})_s$ is the closed subalgebra formed collectively out of the three types of Lorentz generators $sl(2,\mathbb{C})^a$ for $a=1,2,3$, with group actions as pictured in equation 8.31, which also act on the complex subspace $h_3C \subset h_3O$ formed with base units $\{1, l\}$ with for example the type 1 action of equation 7.35.

At the level of complex Lie algebras $L_{\mathbb{C}}$ we have the semi-simple decomposition $sl(3,\mathbb{C}) \equiv su(3) \times su(3)$, and hence the rank-6 subgroup obtained for this real form of E_6 :

$$
SL(3,\mathbb{C})_s \times SU(3)_c \subset E_6 \tag{8.34}
$$

is closely related to an $SU(3) \times SU(3) \times SU(3) \subset E_6$ decomposition, which may be readily obtained by the analysis described in section 7.3 via the extension of the Dynkin diagram for the complex E_6 Lie algebra of figure 7.2(a). In the present theory it is the $SL(2,\mathbb{C})^1 \subset SL(3,\mathbb{C})_s \subset E_6$ Lorentz symmetry of external spacetime that breaks the full E_6 symmetry.

In fact E_6 also contains the following rank-6 subgroup (listed as one of a number of possible decompositions from a mathematical point of view in [38] p.187) which augments equation 8.25:

$$
SL(2,\mathbb{C})^1 \times U(1)_Q \times D(1)_B \times SU(3)_c \subset E_6 \tag{8.35}
$$

with $SL(3,\mathbb{C})_s$ broken to $SL(2,\mathbb{C})^1 \times U(1)_Q \times D(1)_B$, and where $U(1)_Q$ is generated by $\dot{S}_{l}^{1}=\frac{2}{3}% {\textstyle\sum\nolimits_{i}} \left[\frac{\dot{a}_{i}^{2}+1}% {a_{i}^{2}+1}+2\left(\frac{\dot{a}_{i}^{2}+1}{2}\right) ^{2} \right] ,$ $\frac{2}{3}\dot{S}_l^1$ with $\dot{S}_l^1 = (-\dot{R}_{\alpha l}^1 - 2\dot{R}_{\alpha l}^2)$, via equations 8.24 and 6.47, and $D(1)_B$ is generated by $(\dot{B}_{tz}^1 + 2\dot{B}_{tz}^2)$. This latter generator is presented explicitly in equation 13.5 together with a possible physical interpretation of the $D(1)_B$ subgroup in the context of the present theory as described in section 13.2. Further contained within this symmetry breaking pattern is the choice of $SU(2)^{1} \times U(1)_{Q} \subset SU(3)_{s} \subset SL(3,\mathbb{C})_{s}$ with the identification of $U(1)_Q = U(1)^1$, which, in relation to the three possible type $a =$ 1, 2, 3 embeddings $SU(2)^a \times U(1)^a \subset SU(3)_s$ will be seen to be closely related to the phenomena of electroweak symmetry breaking in the Standard Model.

Before describing this connection we note that within the context of the present theory in principle it may be possible to mutually constrain the values of the gauge field couplings associated with a range of internal subgroups in terms of the normalisation of the underlying simple E_6 Lie algebra as expressed by the Killing form. The Killing metric $K_{\alpha\beta} = c^{\rho}{}_{\alpha\sigma}c^{\sigma}{}_{\beta\rho}$ in terms of the algebra structure constants $c^{\alpha}{}_{\beta\gamma}$ was introduced in the discussion leading to equation 4.1. Using this expression the components $K_{\alpha\beta}$ of the complete Killing form for the 78 generators of E_6 in the preferred basis of table 6.3 can in principle be determined directly from the rows of the E_6 Lie algebra table in [38]. For example for the su(3)_c $\equiv \{\dot{A}_i, \dot{G}_l\}$ generators of the colour symmetry described in the previous section we find:

$$
K(\dot{A}_i, \dot{A}_i) = -48,
$$
 $K(\dot{A}_l, \dot{A}_l) = -48,$ $K(\dot{G}_l, \dot{G}_l) = -144$

The Killing metric elements for $SU(3)_c$ can be compared with those for the $SU(2)^2 \times$ $U(1)²$ generators identified in $SU(3)_s$, as adopted in a 'mock electroweak theory', and in principle used to mutually normalise all coupling constants, including $\alpha_s = \frac{g_s^2}{4\pi}$ for the strong interactions, under the unifying simple group E_6 , for comparison with the relative couplings adopted for Standard Model gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y$. With a view towards studying such a mock electroweak theory here we analyse the Killing form for the generators relevant to $su(3)_s$, and calculate from the rows of the E_6 Lie algebra table in [38]:

$$
K(\dot{R}_{\text{nl}}^1, \dot{R}_{\text{nl}}^1) = -24, \qquad K(\dot{R}_{\text{nl}}^1, \dot{S}_l^1) = 0, \qquad K(\dot{S}_l^1, \dot{S}_l^1) = -72
$$

$$
K(\dot{R}_{\text{nl}}^1, \dot{R}_{\text{nl}}^1) = -24, \qquad K(\dot{R}_{\text{nl}}^2, \dot{R}_{\text{nl}}^2) = -24, \qquad K(\dot{R}_{\text{nl}}^2, \dot{R}_{\text{nl}}^2) = -24
$$

The negative values are consistent with the nature of the corresponding group actions as 'rotations', as described in the opening of section 6.5. The bilinear property of the Killing form can be used to deduce further elements as appropriate for a change of basis within the linearly dependent set of elements \dot{R}_{x}^a and \dot{S}_l^b $(a, b \in \{1, 2, 3\})$ with:

$$
\dot{R}_{\text{nl}}^2 = -\frac{1}{2}\dot{R}_{\text{nl}}^1 - \frac{1}{2}\dot{S}_l^1 \quad \Rightarrow \quad K(\dot{R}_{\text{nl}}^2, \dot{R}_{\text{nl}}^2) = -24
$$
\n
$$
\dot{S}_l^2 = +\frac{3}{2}\dot{R}_{\text{nl}}^1 - \frac{1}{2}\dot{S}_l^1 \quad \Rightarrow \quad K(\dot{S}_l^2, \dot{S}_l^2) = -72
$$

via equations 6.47 and 6.48, while:

$$
\dot{R}_{\text{nl}}^3 = -\frac{1}{2}\dot{R}_{\text{nl}}^1 + \frac{1}{2}\dot{S}_l^1 \qquad \Rightarrow \qquad K(\dot{R}_{\text{nl}}^3, \dot{R}_{\text{nl}}^3) = -24
$$
\n
$$
\dot{S}_l^3 = -\frac{3}{2}\dot{R}_{\text{nl}}^1 - \frac{1}{2}\dot{S}_l^1 \qquad \Rightarrow \qquad K(\dot{S}_l^3, \dot{S}_l^3) = -72
$$

Hence the three sets of basis elements $\{\dot{R}_{\underline{x}\underline{l}}^{\underline{a}},\frac{1}{\sqrt{\underline{x}}}\}$ $\frac{1}{3}\dot{S}_l^a$, for either $a = 1, 2$ or 3 have a suitably normalised Killing form. Further, from the bilinearity of the Killing form it is also found for example that:

$$
K(\dot{R}_{\text{nl}}^2, \dot{S}_l^2) = 0
$$
 while $K(\dot{R}_{\text{nl}}^2, \dot{S}_l^1) = +36$

indicating that the Killing form is not diagonal in the latter basis.

Alternatively, restricting the computation of $K_{\alpha\beta} = c^{\rho}{}_{\alpha\sigma}c^{\sigma}{}_{\beta\rho}$ to the SU(3)_s subalgebra all elements of the corresponding 8×8 Killing metric K_8 for the subalgebra basis of equation 8.32 are determined, with for example:

$$
K_8(\dot{R}_{\text{nl}}^1, \dot{R}_{\text{nl}}^1) = -3,
$$
 $K_8(\dot{R}_{\text{nl}}^2, \dot{R}_{\text{nl}}^2) = -3,$ $K_8(\dot{R}_{\text{nl}}^1, \dot{R}_{\text{nl}}^2) = +\frac{3}{2}$

where the latter element is the only non-zero off-diagonal entry of the symmetric Killing form. Hence we replace the basis element \dot{R}_{at}^2 in equation 8.32 with $\frac{1}{\sqrt{2}}$ $\frac{1}{3}\dot{S}_l^1$ such that the $SU(3)_s$ basis:

$$
su(3)_s \equiv \{\dot{R}_{\rm 2L}^1, \frac{1}{\sqrt{3}}\dot{S}_l^1, \dot{R}_{\rm 2Z}^1, \dot{R}_{\rm 2Z}^2, \dot{R}_{\rm 2Z}^3, \dot{R}_{\rm 2L}^1, \dot{R}_{\rm 2L}^2, \dot{R}_{\rm 2L}^3\} \tag{8.36}
$$

has normalised Killing metric $K_8 = -3(1_8)$, where 1_8 is the 8×8 unit matrix. As for the generators of $SU(3)_c$ in table 8.6 a correspondence may be found between the $SU(3)$ _s generators of equation 8.36 and the representation of $SU(3)$ in terms of Gell-Mann λ matrices, as described here in table 8.8.

$$
\begin{array}{|c|c|} \hline \dot R^1_{zI} \sim \frac{1}{2} i \lambda_1 & \dot R^1_{xz} \sim \frac{1}{2} i \lambda_2 & \dot R^1_{xI} \sim \frac{1}{2} i \lambda_3 \\ \\ \hline \dot R^2_{zI} \sim -\frac{1}{2} i \lambda_4 & \dot R^2_{xz} \sim \frac{1}{2} i \lambda_5 \\ \\ \hline \dot R^3_{zI} \sim \frac{1}{2} i \lambda_6 & \dot R^3_{xz} \sim \frac{1}{2} i \lambda_7 & \frac{1}{\sqrt{3}} \dot S^1_{l} \sim \frac{1}{2} i \lambda_8 \\ \\ \hline \end{array}
$$

Table 8.8: The isomorphism between the su(3)_s \subset E₆ Lie algebra basis of equation 8.36 and the eight Gell-Mann matrices of table 8.5.

The choices of basis elements $\{\dot{R}_{x}^{a}, \frac{1}{\sqrt{x}}\}$ $\frac{1}{3}\dot{S}_l^a$ for type $a = 2$ or 3 in place of $a = 1$ correspond to two further possible correlates of the basis matrices $\{\lambda_3, \lambda_8\}$ in the Gell-Mann representation of $su(3)$. These three possibilities correspond to three closely related ways to embed the subgroup $SU(2) \times U(1)$ in $SU(3)$. Here we first study $SU(2)^2 \subset SU(3)$, and the corresponding set of generators $\{\dot{R}_{z_l}^2, \dot{R}_{xz}^2, \dot{R}_{x1}^2\}$. These are the type 2 versions of the three actions of equation 8.7 which, as described in equation 8.9, are respectively associated with the three Pauli matrices $\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $\sigma^2 = \begin{pmatrix} 0 & -l \\ l & 0 \end{pmatrix}, \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ $0 -1$), within factors of $\pm \frac{l}{2}$ $rac{l}{2}$.

In the Standard Model electroweak theory the $su(2)_L$ Lie algebra-valued connection 1-form $W(x) = W^{\alpha}(x)\tau^{\alpha}$, with $W^{\alpha}(x) = W^{\alpha}_{\mu}(x)dx^{\mu}$, $\tau^{\alpha} = \frac{1}{2}$ $\frac{1}{2}\sigma^{\alpha}$ from equation 7.37 and $\alpha = 1, 2, 3$, is parametrised by the three gauge fields $W_\mu^{\alpha}(x)$. The charged gauge boson fields $W^{\pm}_{\mu}(x)$ are associated with complex linear combinations of the SU(2)_L generators $\sigma^{\pm} = \frac{1}{2}$ $\frac{1}{2}(\sigma^1 \pm i\sigma^2)$ as was described in equations 7.57 and 7.58. Guided by this construction based on $SU(2)_L$ generators, here in the complex algebra for $SU(2)^2 \subset E_6$ we define:

$$
\dot{\Sigma}^{(2)\pm} := \dot{R}_{z\perp}^2 \pm i\dot{R}_{xz}^2 \tag{8.37}
$$

Here the imaginary unit $i \in \mathbb{C}$ in the complexification of the E₆ Lie algebra commutes with the elements of $Th_3\mathbb{O}$, which are based on an independent octonion algebra \mathbb{O} . This is the standard notion of a complexified Lie algebra $L_{\mathbb{C}} \equiv L_{\mathbb{R}} + iL_{\mathbb{R}}$, as for example described for figure 7.1, applied here to $L_{\mathbb{R}}$ as the real E_6 Lie algebra represented in the space of vector fields in $Th_3\mathbb{O}$.

The generator \dot{S}_l^1 was associated with the internal symmetry $U(1)_Q$ and electromagnetic charge in the previous section. As a rotation the corresponding group transformation S_l^1 takes the form of unitary 3×3 matrix actions, as described in the opening of section 6.5. Hence considering $i\dot{S}_l^1$ to be an Hermitian generator in the complexified E_6 algebra *real* eigenvalues may be obtained under the adjoint representation. In particular, reading off the corresponding entries in the Lie algebra table in [38] for the complex element of equation 8.37 it is found that:

$$
[\dot{S}_l^1, (\dot{R}_{z1}^2 + i\dot{R}_{xz}^2)] = \frac{3}{2}\dot{R}_{xz}^2 - i\frac{3}{2}\dot{R}_{z1}^2 = -i\frac{3}{2}(\dot{R}_{z1}^2 + i\dot{R}_{xz}^2)
$$
 (8.38)
ace
$$
[i\dot{S}_l^1, (\dot{R}_{z1}^2 + i\dot{R}_{xz}^2)] = +(\dot{R}_{z1}^2 + i\dot{R}_{xz}^2)
$$

hence
$$
[i\dot{S}_l^1, (\dot{R}_{zl}^2 + i\dot{R}_{xz}^2)] = +(R_{zl}^2 + i\dot{R}_{xz}^2)
$$

and $[i\dot{S}_l^1, \dot{\Sigma}^{(2)\pm}] = \pm \dot{\Sigma}^{(2)\pm}$ (8.39)

with real charge eigenvalues ± 1 . Hence the generators $\dot{\Sigma}^{(2)\pm}$ of equation 8.37 are associated with the *same* magnitude of $U(1)_Q$ charge under \dot{S}_l^1 as was found for the electron in the leptonic components $\theta_l \subset h_3 \mathbb{O}$ as described in equations 8.22–8.26. Since such factors of $\frac{3}{2}$ as seen in equation 8.38 are relatively sparse in the E₆ Lie algebra table [38], with none appearing for example here in table 6.4, this seems to be a non-trivial correspondence of \dot{S}_l^1 charges. In the su(3)_s basis of equation 8.36 the generators $\dot{\Sigma}^{(2)\pm}$ are in fact two of the eigenvectors of elements of the Cartan subalgebra, which in turn has a basis $\{\dot{R}_{\underline{x}l}^1, \frac{1}{\sqrt{x}}\}$ $\frac{1}{3}\dot{S}_l^1$, under the adjoint representation in the complex $su(3)$ _s algebra. Indeed we find also:

$$
[i\dot{R}_{x}^{1}, \dot{\Sigma}^{(2)\pm}] = \pm \frac{1}{2}\dot{\Sigma}^{(2)\pm}
$$
\n(8.40)

More generally for a Lie algebra of rank-n the elements of the Cartan subalgebra ${H_i}, i = 1...n$, are mutually commuting and any element, or linear combination of elements, in ${H_i}$ generates a U(1) symmetry. In any representation of the Lie algebra the eigenvalues, or 'weights', of such a $U(1)$ generator can be considered as 'charges'. In the present case the $U(1)_Q$ generator \dot{S}_l^1 , which also belongs to the E_6 Cartan subalgebra as can be seen from equation 6.56, is associated with electromagnetic charge.

In the Cartan-Weyl basis of a complex Lie algebra the eigenvectors E_{α} of elements of the Cartan subalgebra ${H_i}$ in the adjoint representation have real eigenvalues α_i :

$$
[H_i, E_\alpha] = \alpha_i E_\alpha \tag{8.41}
$$

with
$$
[E_{\alpha}, E_{-\alpha}] = (K^{ij} \alpha_j) H_i
$$
 (8.42)

(where K^{ij} are components of the Killing metric restricted to the Cartan subalgebra). The eigenvalues, or 'weights', α_i of the adjoint representation are also called 'roots', the full set of which under $\{H_i\}$ is central to the classification of complex Lie algebras, as alluded to in section 7.3 alongside figure 7.2. Since the elements of the Lie algebra form the vector space $\text{span}(H_i, E_\alpha)$ upon which the adjoint representation acts, the dimension of this representation is equal to the dimension of the Lie algebra itself. Generally in a given representation r of a Lie algebra on a vector space V with eigenvectors $|v\rangle$ and weights λ_i , that is with:

$$
H_i^{(r)}|v\rangle = \lambda_i|v\rangle
$$

then:
$$
H_i^{(r)}(E_\alpha^{(r)}|v\rangle) = E_\alpha^{(r)}H_i^{(r)}|v\rangle + [H_i^{(r)}, E_\alpha^{(r)}]|v\rangle = (\lambda_i + \alpha_i)(E_\alpha^{(r)}|v\rangle) \quad (8.43)
$$

using equation 8.41. That is, the $E_{\alpha}^{(r)}$ act as 'raising' operators (while the $E_{-\alpha}^{(r)}$ $\frac{d^{(r)}}{d-\alpha}$ act as 'lowering' operators) on the eigenstates in the representation.

Hence by comparison of equation 8.39 with equation 8.41 above the complex linear combinations $\dot{\Sigma}^{(2)\pm}$ of equation 8.37 as eigenvectors \dot{S}_l^1 under the 78-dimensional adjoint representation of E_6 indeed have charges of ± 1 under the same generator of U(1)_Q which acts on the e-lepton and d-quark states identified in the $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ components of the 27-dimensional representation of the E_6 symmetry on the space h₃O. Further, according to equation 8.43, the $\dot{\Sigma}^{(2)\pm}$ actions are expected to transform states in the h₃^O representation with a change of ± 1 units of the electron charge. Based on the type 2 subgroup $SU(2)^2 \subset SL(2,\mathbb{C})^2$ these raising and lowering operations are associated with the $\theta^2 = \left(\frac{a}{c}\right)$ $\frac{a}{c}$ components of h₃ \mathbb{O} as shown explicitly in equations 8.29 and 8.31. In this subsection we have focussed precisely upon this doublet action of the $SU(2)^2$ symmetry which appears to be closely related to transformations within the lepton $\binom{\nu}{e}$ $\binom{u}{e}$ and quark $\binom{u}{d}$ $\binom{u}{d}$ doublets as mediated by the W^{\pm} gauge bosons in the Standard Model.

In the Cartan-Weyl basis generally the Lie bracket $[E_{\alpha}, E_{-\alpha}]$ describes an element of the Cartan subalgebra, as can be seen from equation 8.42. From the E_6 Lie algebra table in [38] we find:

$$
[\dot{\Sigma}^{(2)+}, \dot{\Sigma}^{(2)-}] = [(\dot{R}_{\rm z1}^2 + i\dot{R}_{\rm zz}^2), (\dot{R}_{\rm z1}^2 - i\dot{R}_{\rm zz}^2)] = -i\dot{S}_l^1 - i\dot{R}_{\rm zz}^1 \tag{8.44}
$$

which is indeed in the Cartan subalgebra of equation 6.56, for the complexified E_6 Lie algebra, and also for the complex $su(3)_s$ subalgebra. This is consistent with the identification of the $\dot{R}_{\nu l}^1$ 'charges' for $\dot{\Sigma}^{(2)\pm}$ in equation 8.40. However it the S_l^1 action that has been associated with the internal symmetry $U(1)_Q$ in the previous section and in turn the eigenvalues of \dot{S}_l^1 associated with physical electromagnetic charges. It is the latter charges of ± 1 for the states $\dot{\Sigma}^{(2)\pm}$ which will be provisionally associated with the $W^{\pm}_{\mu}(x)$ charged gauge fields in the mock electroweak theory.

In quantum field theory the creation and annihilation operators associated with real fields do not describe charged particles, rather conserved charges are associated with complex fields, or complex linear combinations of real fields, as will be described in section 10.3. A complex scalar field $\mathcal{Y}(x)$ has charge q under a U(1) symmetry if it transforms as $\mathcal{Y} \to e^{iq\alpha} \mathcal{Y}$, with $\alpha \in \mathbb{R}$ and $e^{iq\alpha} \in U(1)$, with q also labelling the irreducible representation of U(1). The derivative of this transformation at $\alpha = 0$ can be written as $\partial \mathcal{Y}/\partial \alpha = \mathcal{Y} = +q(i\mathcal{Y})$. This has the same form as equation 8.23, which via equation 8.24 implies the U(1)_Q action \dot{S}_l^1 on the field components $\theta_l(x)$ reads $[\dot{\theta}_l] = +1[l\theta_l],$ with the complex imaginary unit l and charge represented by the real eigenvalue $q = +1$.

As for the electron and d-quarks charges identified in the components of $h_3\mathbb{O}$ it remains to be seen how the charges for gauge bosons derived from generators such as $\Sigma^{(2)\pm}$ relate to the likelihood of physical processes such as observed in high energy physics experiments for the present theory. This will be discussed in section 11.2 in comparison with standard QFT for which the charges are placed by hand into Lagrangian terms, leading to calculations of transition amplitudes and cross-sections. The phenomenon of running coupling, as described in section 11.3, will ultimately also need to be considered for any comparison between theoretical couplings derived from a normalised Killing form for a simple Lie algebra and the couplings measured empirically in the laboratory. As well as accounting for quantisation a full dynamical theory will also be required, incorporating for example self-interactions for non-Abelian gauge fields, as explored in relation to Kaluza-Klein theories here in chapters 4 and 5.

8.3.2 $SU(2)^2 \times U(1)^2$ Mixing Angle

The four type 1 actions $\{\dot{R}_{z_l}^1, \dot{R}_{xz_l}^1, \dot{R}_{xl}^1\}, \dot{S}_l^1$ generate the group $SU(2)^1 \times U(1)_Q$. Here $SU(2)^1$, generated by $\{\dot{R}_{z_l}^1, \dot{R}_{xz}^1, \dot{R}_{xl}^1\}$, is the rotation subgroup of the external Lorentz transformations, as studied in section 8.1, which commutes with the internal symmetry $U(1)_Q$, underlying Maxwell's electromagnetic field, generated by \dot{S}_l^1 as identified in section 8.2. For the case of the corresponding set of four type 2 actions $\{\dot{R}_{\rm zl}^2, \dot{R}_{\rm zz}^2, \dot{R}_{\rm zz}^2, \dot{R}_{\rm zl}^2\}, \dot{S}_l^2$, a similar structure can be identified for $SU(2)^2 \times U(1)^2$. In a similar way that \ddot{S}_l^1 commutes with su(2)¹, and indeed with the Lorentz group $sl(2,\mathbb{C})^1$, it is also the case that \dot{S}_l^2 commutes with su(2)² and hence with $\dot{\Sigma}^{(2)\pm}$ of equation 8.37:

$$
[\dot{S}_l^2, \dot{\Sigma}^{(2)\pm}] = 0 \tag{8.45}
$$

This commutator is consistent with those in equations 8.39 and 8.40 given the linear dependence obtained from equations 6.48 and 8.24:

$$
\dot{S}_l^2 = \dot{R}_{\text{nl}}^1 - \frac{1}{2} \dot{S}_l^1 \tag{8.46}
$$

While the generator \dot{S}_l^1 is associated with electric charge Q the generator \dot{R}_{nl}^2 is associated with T^3 , the third component of $SU(2)^2$. The linear dependencies in the E₆ Lie algebra of equations 6.47 and 6.48 also imply the relation:

$$
-\dot{S}_l^1 = \dot{R}_{2l}^2 + \frac{1}{2}\dot{S}_l^2\tag{8.47}
$$

which is closely reminiscent of the relation:

$$
Q = T^3 + \frac{1}{2}Y \tag{8.48}
$$

of equation 7.38, within the choice of sign conventions. This suggests associating $\frac{1}{2}\dot{S}_l^2$ with $\frac{1}{2}Y$ as a candidate for the generator of the *hypercharge* symmetry U(1)_Y ~ U(1)² which commutes with $SU(2)^2$, as generated by $\{\dot{R}_{z_l}^2, \dot{R}_{xz}^2, \dot{R}_{xt}^2\}$, and as provisionally associated with $SU(2)_L$ for a mock electroweak theory in the previous subsection. The generator \dot{S}_l^2 may also be expressed as the linear combination of type 1 elements in equation 8.46, which lies in the Cartan subalgebra of the E_6 Lie algebra. Hence the 'weights' of \dot{S}_l^2 may indeed be considered as 'charges', which are termed hypercharges for the corresponding $U(1)_Y$ symmetry.

More generally opening up consideration of the three $SL(2,\mathbb{C})^a$ actions in the previous subsection also motivates an examination of the $U(1)$ charge structure associated with \dot{S}^a_l for all three types $a = 1, 2, 3$. To understand the relationships between these charges all three generators \dot{S}_l^a , for $a = 1, 2, 3$, from table 6.7 with a factor of $\times \frac{2}{3}$ 3 from equation 8.24, are explicitly written out in terms of $T_{\text{B}}\mathbb{O}$ components in equation 8.49. Each entry of the form (x, y) represents the factors of l which multiply the components of h₃ Ω algebraically from the left side – where x is the 'leptonic part', that is on the real and l components, while η is the 'quark part', that is on the remaining six imaginary units of each $a, b, c \in \mathbb{O}$. The components for \dot{S}_l^1 in equation 8.49 contain the same information as equation 8.27 rearranged into the 3×3 matrix of Th₃^O. It can be seen here that $\dot{S}_l^1 + \dot{S}_l^2 + \dot{S}_l^3 = 0$, for each component a, b and c, consistent with equation 6.42. Also shown are the corresponding components of \dot{R}_{nl}^2 as obtained from table 6.6, which can be seen to be consistent with equation 8.47.

$$
\begin{aligned}\n\dot{S}_{l}^{1} & \dot{S}_{l}^{2} & \dot{S}_{l}^{3} \\
(0, +\frac{2}{3}) & (1, -\frac{1}{3}) \\
(-1, -\frac{1}{3}) & (1, -\frac{1}{3})\n\end{aligned}\n\begin{aligned}\n&\text{(1, } -\frac{1}{3}) & \text{(1, } -\frac{1}{3}) & (0, +\frac{2}{3}) \\
&\text{(1, } -\frac{1}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) \\
&\text{(1, } -\frac{1}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) \\
&\text{(1, } -\frac{1}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) \\
&\text{(1, } -\frac{1}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) & (1, -\frac{1}{3}) & (0, +\frac{2}{3}) \\
&\text{(2, } -\frac{1}{3}) & \text{(2, } -\frac{1}{2}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) \\
&\text{(3, 4)} & \text{(4, } -\frac{1}{2}, -\frac{1}{2}) & (0, +\frac{1}{2}) & (0, +\frac{1}{2}) & (0, +\frac{2}{3}) \\
&\text{(4, } -\frac{1}{2}, -\frac{1}{2}) & (0, +\frac{1}{2}, -\frac{1}{2}) & (0, +\frac{1}{2}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) \\
&\text{(5, } -\frac{1}{2}, -\frac{1}{2}) & (0, +\frac{1}{2}, -\frac{1}{2}) & (0, +\frac{1}{2}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) \\
&\text{(4, } -\frac{1}{2}, -\frac{1}{2}) & (0, +\frac{1}{2}, -\frac{1}{2}) & (0, +\frac{1}{2}) & (0, +\frac{2}{3}) & (0, +\frac{2}{3}) & (0, +\frac{2}{
$$

Hence the $\frac{1}{2}\dot{S}_l^2$ 'hypercharge values' of $\left(-\frac{1}{2}, -\frac{1}{6}\right)$ on the \bar{a} and c components in equation 8.49 match the Standard Model hypercharge values of $\frac{Y}{2}(l_L) = -\frac{1}{2}$ and Y $\frac{Y}{2}(q_L) = +\frac{1}{6}$ for the left-handed doublets of leptons and quarks respectively of equation 7.36 (up to a sign convention, which again will ultimately depend on the definition of particle and antiparticle states in spacetime). As can be seen in equation 8.31 and described in the previous subsection these components (\bar{a} c) are also linked by the $SU(2)^2 \subset SL(2,\mathbb{C})^2$ actions and corresponding $\dot{\Sigma}^{(2)\pm}$ operators provisionally associated with the $W^{\pm}_{\mu}(x)$ charged gauge fields. Although some of these observations are naturally mutually correlated, the $(1, 2, -\frac{1}{2})$ $(\frac{1}{2})_L$ and $(3, 2, \frac{1}{6})$ $(\frac{1}{6})_L$ pieces of equation 7.36 are hence closely associated respectively with the $\binom{a_1, l}{\theta_l}$ and $\binom{a(6)}{\theta_{i,j,k}}$ $\binom{a(6)}{\theta_{i,j,k}}$ components of h₃ $\mathbb O$ in equations 8.26 and 8.28.

While right-handed fermion states remain to be identified, the hypercharges of the right-handed fermion singlets in equation 7.36 are also closely correlated with the \dot{S}_l^a charges in equation 8.49. This is expected since $Q = \frac{Y}{2}$ $\frac{Y}{2}$ for these cases and the electric charge Q is well described by \dot{S}_l^1 . Also in the top row of \dot{S}_l^3 the values $(1, -\frac{1}{3})$, $(0, +\frac{2}{3})$ have the same magnitude as the $\frac{Y}{2}$ values for the right-handed singlets e_R , d_R , ν_R and u_R respectively, although these components do not correspond to the correct electromagnetic charges Q under \dot{S}^1_l for those respective fermion states. However these observations do suggest opening up consideration of the (correlated) charges for all three \dot{S}_l^a generators. Indeed as well as \dot{S}_l^2 the generator \dot{S}_l^3 should also relate to hypercharge as $SU(2)^3 \times U(1)^3$ also forms a possible mock $SU(2)_L \times U(1)_Y$ action with the following linear dependence also found within the E_6 algebra:

$$
\dot{S}_l^1 = \dot{R}_{\text{z}l}^3 - \frac{1}{2}\dot{S}_l^3 \tag{8.50}
$$

This equation is the type 3 version of equation 8.47. Further linear relations include $\dot{S}_l^1 = -2\dot{R}_{nl}^2 + \dot{S}_l^3$ and $\dot{S}_l^1 = 2\dot{R}_{nl}^3 + \dot{S}_l^2$ which combine non-commuting type 2 and 3 actions on the right-hand side. Such equations of linear dependence, relating the generators \dot{S}_l^a and $\dot{R}_{\underline{\tau}\underline{l}}^b$ for $a, b = \{1, 2, 3\}$, are fixed by the structure of the E₆ Lie algebra and closely resemble equation 7.38 which is constructed to relate the electric charge Q, third component of weak isospin T^3 and hypercharge $\frac{Y}{2}$ in the Standard Model. However a fuller understanding of this structure in the present theory will require the identification of right-handed fermion states, and in particular such states with $T^3 = 0$. The origin of both left and right-handed states, together with their mutual relation will be considered explicitly in section 9.2, while in the meantime we further consider the structure of the mock electroweak theory within the E_6 framework.

In particular, moving away from the a *static* analysis of the E_6 symmetry breaking pattern to a more dynamic perspective, we next study the structure of an $SU(2)^2 \times U(1)^2$ gauge theory based on the symmetry generators $\{\dot{R}_{z_l}^2, \dot{R}_{xz}^2, \dot{R}_{x_l}^2, \dot{S}_l^2\}.$ These act on the doublet components of the type 2 spinor $\theta^2 = \left(\frac{a}{c}\right)^2$ (\vec{c}) in h₃^O. Restricted to the complex subspace $\mathbb{C} \subset \mathbb{O}$ with $\{1, l\}$ basis units the components $\theta_l^2 = \left(\frac{a}{c}\right)^2$ $\binom{a}{\bar{c}}_l$ provisionally represents the lepton doublet $\binom{\nu}{e}$ e^{ν} . Since these components do not correspond to complete $SL(2,\mathbb{C})$ ¹ Weyl spinors for either the neutrino *or* the electron part this $SU(2)^2 \times U(1)^2$ symmetry is clearly not directly equivalent to the $SU(2)_L \times U(1)_Y$ symmetry of electroweak theory. However the components of θ_l^2 do transform under the internal symmetry $SU(3)_c \times U(1)_Q$ appropriately to represent such a lepton doublet, as described in section 8.2, and hence the $SU(2)^2 \times U(1)^2$ symmetry serves as a useful intermediate model, considered as a mock electroweak theory. The equations of motion for the corresponding field $\theta_l^2(x)$ in spacetime M_4 will then involve the gauge covariant derivative (essentially as described in section 3.1):

$$
D_{\mu}\theta_l^2(x) = \partial_{\mu}\theta_l^2(x) + \tilde{g}\,\tilde{W}_{\mu}^{\alpha}(x)\,\dot{R}^{(2)\alpha}(\theta_l^2) + \tilde{g}'\,\tilde{B}_{\mu}(x)\,\frac{1}{2}\dot{S}_l^2(\theta_l^2) \tag{8.51}
$$

where $\alpha = 1, 2, 3$ and $\dot{R}^{(2)\alpha} \equiv {\{\dot{R}_{zl}^2, \dot{R}_{xz}^2, \dot{R}_{xl}^2\}},$ and for example $\dot{R}_{2l}^2(\theta_l^2)$ denotes the θ_l^2 components of \dot{R}_{2l}^2 . The couplings \tilde{g}, \tilde{g}' and the gauge fields $\tilde{W}_{\mu}^{\alpha}(x$ with the $SU(2)^2 \times U(1)^2$ gauge symmetry are introduced by analogy with the Standard Model case in equation 7.40 and hence similar notation is adopted. However it is important to contrast the corresponding gauge coupling terms implied in equations 7.40 and 8.51 with for example respectively:

$$
D_{\mu}l_{L} \sim i g' B_{\mu}(x) \frac{Y}{2}(l_{L}) l_{L} \quad \text{and} \quad D_{\mu} \theta_{l}^{2} \sim \tilde{g}' \tilde{B}_{\mu}(x) \frac{\dot{S}_{l}^{2}}{2}(\theta_{l}^{2}) \quad (8.52)
$$

In the former case, apart from the conventional factor of i there are four factors: the coupling g', the gauge field $B_{\mu}(x)$, the hypercharge generator $\frac{Y}{2}(l_L) = -\frac{1}{2}$ $\frac{1}{2}$ $\binom{1}{0}$ and the lepton doublet $l_L = \binom{\nu}{e}$ e_{L}^{ν} . In the latter case there are only three factors: with the coupling \tilde{g}' and gauge field $\tilde{B}_{\mu}(x)$ having a similar role as for the first case, while the third part $\frac{\dot{S}_l^2}{2}(\theta_l^2)$ corresponds to the action of the hypercharge generator represented directly on the $\theta_l^2 = \begin{pmatrix} a \\ \bar{c} \end{pmatrix}$ $\epsilon_{\bar{c}}^{a}$ components of h₃ \mathbb{O} , which is equivalent to the combination $\frac{Y}{2}(l_L) l_L$ for the standard case.

In principle in the second case the coupling \tilde{g}' may be absorbed into the gauge field $\tilde{B}_{\mu}(x)$ since we are here dealing with the pure covariant derivatives, as originally expressed in equation 2.38 of subsection 2.2.3 for the gauge field $A_\mu(x)$ without any coupling constant. Adopting couplings such as $\tilde{g}' = 1$ is also compatible with the construction of a direct relationship between the curvature for the external linear connection and that for the internal gauge connection as described in section 5.1 and equation 5.20 (with a factor such as $\chi = 1$ in principle determined by the geometric structure), in comparison with Kaluza-Klein theory. A similar observation applies for

the coupling \tilde{g} associated with the gauge field $\tilde{W}_{\mu}^{\alpha}(x)$ in equation 8.51. Ultimately both \tilde{g} and \tilde{g}' will be absorbed into the relevant gauge fields and effectively set equal to one.

In turn the 'charges' of individual states will depend upon the representation which is already determined directly by the values of $\frac{\dot{S}_l^2}{2}(\theta_l^2)$ in the second expression of equation 8.52, which are closely analogous to the case for the electromagnetic charges obtained from $\dot{S}_l^1(\theta^1)$ in equations 8.23 and 8.24 and further discussed towards the end of the previous subsection. More generally this will require a suitable mutual normalisation of the generators $\dot{R}^{(2)\alpha}$ and $\frac{1}{2}\dot{S}_l^2$ based on the Killing form of the full E_6 Lie algebra, as also described in the previous subsection, to relate the charges for the various subgroups of the internal gauge symmetry. With the gauge groups represented directly on the space $Th_3\mathbb{O}$ this structure parallels that employed for Kaluza-Klein theory based on homogeneous fibres as described in section 4.3.

For now considering \tilde{g} and \tilde{g}' as free parameters in equation 8.51 allows a closer comparison with the structure of the electroweak theory in the Standard Model for which the couplings g and g' are independent. However here neither the $SU(2)^2$ generated by $\{\dot{R}_{z_l}^2, \dot{R}_{xz}^2, \dot{R}_{xl}^2\}$ nor the U(1)² generated by \dot{S}_l^2 are internal symmetries in the sense of table 8.3, that is within $Stab(TM_4)$, with each of these four generators impacting upon the components of the type 1 subspace $h_2\mathbb{C} \subset h_3\mathbb{O}$, which represent components of the external spacetime TM_4 , as can be seen explicitly from the form of these four generators in tables 6.6 and 6.7. The breaking of the full E_6 symmetry action on h₃ \mathbb{O} in this identification of the type 1 subspace h₂ \mathbb{C} with the local tangent space of the external spacetime hence includes the breaking of the $SU(2)^2 \times U(1)^2 \subset E_6$ subgroup.

The covariant derivative applied to the θ_l^2 components in equation 8.51 can be applied to the components of $h_3\mathbb{O}$ more generally and written out explicitly using tables 6.6 and 6.7. In particular we find that applied to the type 1 embedding of the 2×2 matrix of components $X \in h_2 \mathbb{O} \subset h_3 \mathbb{O}$ and $h_2 \in h_2 \mathbb{C} \subset h_3 \mathbb{O}$ this covariant derivative reads respectively:

$$
D_{\mu}X = \partial_{\mu}X + \tilde{g}\tilde{W}_{\mu}^{1}(\frac{1}{2}cl) + \tilde{g}\tilde{W}_{\mu}^{2}(\frac{1}{2}cl) + \tilde{g}\tilde{W}_{\mu}^{2}(\frac{1}{2}cl) + \tilde{g}\tilde{W}_{\mu}^{3}(\frac{1}{2}\bar{a}l) + \tilde{g}'\tilde{B}_{\mu}\frac{1}{2}\dot{S}_{l}^{2}(\bar{a})
$$

\n
$$
\tilde{g}\tilde{W}_{\mu}^{1}(-\frac{1}{2}l\bar{c}) + \tilde{g}\tilde{W}_{\mu}^{2}(\frac{1}{2}\bar{c}) + \tilde{g}\tilde{W}_{\mu}^{3}(-\frac{1}{2}la) + \tilde{g}'\tilde{B}_{\mu}\frac{1}{2}\dot{S}_{l}^{2}(a)
$$
\n
$$
\tilde{g}\tilde{W}_{\mu}^{1}(b_{l}) + \tilde{g}\tilde{W}_{\mu}^{2}(b_{x}) + 0 + 0
$$
\n(8.53)

$$
D_{\mu}\mathbf{h}_{2} = \partial_{\mu}\mathbf{h}_{2} + \begin{pmatrix} \frac{\tilde{g}}{2} \left(\tilde{W}_{\mu}^{1}(c_{1}l - c_{8}) + \tilde{W}_{\mu}^{2}(c_{1} + c_{8}l) + \tilde{W}_{\mu}^{3}(a_{1}l + a_{8}) \right) + \frac{\tilde{g}'}{2} \tilde{B}_{\mu}(-a_{1}l - a_{8}) \\ \frac{\tilde{g}}{2} \left(\tilde{W}_{\mu}^{1}(-c_{1}l - c_{8}) + \tilde{W}_{\mu}^{2}(c_{1} - c_{8}l) + \tilde{W}_{\mu}^{3}(-a_{1}l + a_{8}) \right) + \frac{\tilde{g}'}{2} \tilde{B}_{\mu}(a_{1}l - a_{8}) \qquad \tilde{g} \tilde{W}_{\mu}^{1}(b_{8}) + \tilde{g} \tilde{W}_{\mu}^{2}(b_{1}) \end{pmatrix}
$$
\n(8.54)

where the second equation shows that indeed each of the four gauge fields $\tilde{W}^{\alpha}_{\mu}(x)$, $\tilde{B}_{\mu}(x)$ has non-zero impact on the $\{1, l\}$ components of $X \in h_2\mathbb{O}$, that is on the 4dimensional vector $h_2 \in h_2 \mathbb{C}$ of equations 8.4 and 8.5, unlike the case of equations 8.14 and 8.15 for example, and hence are not associated with a purely internal symmetry in the sense of $\text{Stab}(TM_4)$. However an orthogonal linear combination of gauge fields may be taken with:

$$
\tilde{B}_{\mu} = \cos \theta_{M^2} \tilde{A}_{\mu} - \sin \theta_{M^2} \tilde{Z}_{\mu}
$$
\n
$$
\tilde{W}_{\mu}^3 = \sin \theta_{M^2} \tilde{A}_{\mu} + \cos \theta_{M^2} \tilde{Z}_{\mu}
$$
\n(8.55)

by analogy with equations 7.43 and 7.44, where θ_{M^2} (with subscript M^2 denoting 'mock mixing angle' of type 2) plays a similar role to the weak mixing angle θ_W . The corresponding contribution from the $\tilde{B}_{\mu}(x)$ and $\tilde{W}^{3}_{\mu}(x)$ fields to the $\bar{a}_{1,l}$ components in the top-right element of equation 8.54 is then:

$$
D_{\mu}\bar{a}_{1,l} = ... +
$$

\n
$$
\frac{\tilde{g}}{2}\sin\theta_{M^2}\tilde{A}_{\mu}(a_1l + a_8) + \frac{\tilde{g}'}{2}\cos\theta_{M^2}\tilde{A}_{\mu}(-a_1l - a_8) + \frac{\tilde{g}}{2}\cos\theta_{M^2}\tilde{Z}_{\mu}(a_1l + a_8) - \frac{\tilde{g}'}{2}\sin\theta_{M^2}\tilde{Z}_{\mu}(-a_1l - a_8)
$$
\n(8.56)

Hence the gauge field $\tilde{A}_{\mu}(x)$ represents a purely internal field, with no action on the external $h_2 \in h_2 \mathbb{C}$ components, provided:

$$
\tilde{g}\sin\theta_{M^2} = \tilde{g}'\cos\theta_{M^2}
$$
\nthat is:

\n
$$
\tan\theta_{M^2} = \frac{\tilde{g}'}{\tilde{g}} \tag{8.57}
$$

This relation is closely analogous to equation 7.46 for electroweak theory in the Standard Model. However here in the case of equation 8.57 neither a Lagrangian formalism, using for example equation 7.41, nor a Higgs field is required to break the $SU(2)^2 \times U(1)^2$ symmetry down to a $U(1)$ symmetry associated with the gauge field $\tilde{A}_{\mu}(x)$. Considering more generally the $\tilde{A}_{\mu}(x)$ field part of the covariant derivative D_{μ} of equation 8.51, via equation 8.55, on all of the components of $X \in h_3\mathbb{O}$, with for example $\dot{R}_{\text{nl}}^2 \equiv \dot{R}_{\text{nl}}^2(\mathcal{X})$, we have:

$$
D_{\mu}\mathcal{X}(x) = \partial_{\mu}\mathcal{X}(x) + \tilde{g}\sin\theta_{M^2}\tilde{A}_{\mu}(x)\dot{R}_{\mu}^2 + \tilde{g}'\cos\theta_{M^2}\tilde{A}_{\mu}(x)\frac{1}{2}\dot{S}_l^2
$$

\n
$$
= \partial_{\mu}\mathcal{X}(x) + \tilde{g}\sin\theta_{M^2}\tilde{A}_{\mu}(x)\dot{R}_{\mu}^2 + \tilde{g}\tan\theta_{M^2}\cos\theta_{M^2}\tilde{A}_{\mu}(x)\frac{1}{2}\dot{S}_l^2
$$

\n
$$
= \partial_{\mu}\mathcal{X}(x) + \tilde{g}\sin\theta_{M^2}\tilde{A}_{\mu}(x)(\dot{R}_{\mu}^2 + \frac{1}{2}\dot{S}_l^2)
$$

\n
$$
= \partial_{\mu}\mathcal{X}(x) + \tilde{g}\sin\theta_{M^2}\tilde{A}_{\mu}(x)(-\dot{S}_l^1)
$$
(8.58)

where the final line is fixed by the *linear dependence* of equation 8.47 for the generators of the E₆ Lie algebra. The gauge field $\tilde{A}_{\mu}(x)$ is hence associated with \dot{S}^1_l which as an element of stab(TM_4) has been identified as the generator of the internal gauge symmetry $U(1)_{\mathcal{O}}$ of electromagnetism in the previous section (see the discussion following equation 8.21). The zero charge of the ν -lepton, associated with the $a_{1,l}$ components in equation 8.28, is here taken to be entirely equivalent to the fact that the action \dot{S}_l^1 l does not impinge on the $\{1, l\}$ components of $a \in \mathbb{O} \subset h_3\mathbb{O}$. The apparently ambiguous nature of these $a_{1,l}$ components, which have been associated *both* with the neutrino state and with part of the vector space $h_2C \equiv TM_4$ on the external spacetime, will be resolved in section 9.2.

The lines of equation 8.58 are closely analogous to those of equation 7.47 from electroweak theory, with $\tilde{A}_{\mu}(x) \sim A_{\mu}(x)$ and $-\dot{S}_{l}^{1} \sim Q$. The apparent electromagnetic coupling \tilde{e} may be identified directly in equation 8.58 as:

$$
\tilde{e} = \tilde{g}\sin\theta_{M^2} \tag{8.59}
$$

which is also analogous to equation 7.48 in the Standard Model.

We next employ a basis for the E_6 algebra with a normalised Killing form with components proportional to the unit 78×78 matrix. In this case it will be possible to determine the value of the mixing angle θ_{M^2} in the breaking of the $SU(2)^2 \times U(1)^2$ symmetry to $U(1)_Q$. For the su(3)_s subalgebra such a normalised basis is provided by equation 8.36 with three possible choices of $\{\dot{R}_{\text{nl}}^a, \frac{1}{\sqrt{n}}\}$ $\frac{\dot{S}^a}{3}\dot{S}^a_l$, for type $a = 1, 2, 3$, for the first two elements. The covariant derivative of equation 8.51 may be rewritten in the normalised Killing form basis, with $\frac{1}{\sqrt{2}}$ $\frac{1}{3}\dot{S}_l^a = \frac{\sqrt{3}}{2}$ $\frac{\sqrt{3}}{2}\dot{S}_l^a$ (via equation 8.24) and with the couplings \tilde{g} and \tilde{g}' absorbed into the gauge fields as:

$$
D_{\mu}\theta_l^2(x) = \partial_{\mu}\theta_l^2(x) + \tilde{W}_{\mu}^{\alpha}(x)\dot{R}^{(2)\alpha}(\theta_l^2) + \tilde{B}_{\mu}(x)\frac{\sqrt{3}}{2}\dot{S}_l^2(\theta_l^2)
$$
(8.60)

where again $\alpha = 1, 2, 3$ and $\dot{R}^{(2)\alpha} \equiv \{\dot{R}_{\alpha}^2, \dot{R}_{\alpha}^2, \dot{R}_{\alpha}^2\}$. The gauge fields $\{\tilde{W}_{\mu}^3(x), \tilde{B}_{\mu}(x)\}$ aligned with the generators $\{\dot{R}_{xl}^2, \frac{\sqrt{3}}{2}$ $\frac{\sqrt{3}}{2}\dot{S}_l^2$ may be expressed in a new basis with gauge fields $\{\tilde{Z}_{\mu}(x), \tilde{A}_{\mu}(x)\}\$ aligned with the generators $\{\dot{R}_{x}^{1}, \frac{\sqrt{3}}{2}\}\$ $\frac{\sqrt{3}}{2}\dot{S}_l^1$. In this basis the 'internal' gauge field $\tilde{Z}_{\mu}(x)$ is associated with \dot{R}_{μ}^{1} which as a generator of $SL(2,\mathbb{C})^1$, as originally listed in equation 6.57, is in fact a purely external action! However here we are dealing with a mock electroweak theory for which some inappropriate features may be observed, as was the case for the ambiguity of the $a_{1,l}$ components noted above. In any case the electromagnetic gauge field $\tilde{A}_{\mu}(x)$ associated with $\frac{\sqrt{3}}{2}$ $\frac{\sqrt{3}}{2}\dot{S}_l^1$ in the new basis does represent a purely internal action. Transferring to the new basis we have:

$$
\tilde{W}_{\mu}^{3} \dot{R}_{\nu}^{2} + \tilde{B}_{\mu} \frac{\sqrt{3}}{2} \dot{S}_{l}^{2} \Rightarrow \tilde{Z}_{\mu} \dot{R}_{\nu}^{1} + \tilde{A}_{\mu} \frac{\sqrt{3}}{2} \dot{S}_{l}^{1}
$$
\nhence:
\n
$$
\sin \theta_{M^{2}} \tilde{A}_{\mu} \dot{R}_{\nu}^{2} + \cos \theta_{M^{2}} \tilde{A}_{\mu} \frac{\sqrt{3}}{2} \dot{S}_{l}^{2} = \tilde{A}_{\mu} \frac{\sqrt{3}}{2} \dot{S}_{l}^{1}
$$
\n
$$
\sin \theta_{M^{2}} \tilde{A}_{\mu} \left(-\frac{1}{2} \dot{R}_{\nu}^{1} - \frac{3}{4} \dot{S}_{l}^{1}\right) + \cos \theta_{M^{2}} \tilde{A}_{\mu} \frac{\sqrt{3}}{2} (\dot{R}_{\nu}^{1} - \frac{1}{2} \dot{S}_{l}^{1}) = \tilde{A}_{\mu} \frac{\sqrt{3}}{2} \dot{S}_{l}^{1} \qquad (8.61)
$$

where in the second line the orthogonal transformation of equation 8.55 has been applied to the left-hand side and only the \tilde{A}_{μ} field part has been retained on both sides. Equations 6.47 and 6.48, together with equation 8.24, have been used for the bottom line. By equating the basis vector $\dot{R}_{\nu l}^1$ and \dot{S}_l^1 parts separately in this final line above it can be deduced that:

$$
\sin \theta_{M^2} = -\frac{\sqrt{3}}{2}
$$
 and $\cos \theta_{M^2} = -\frac{1}{2}$ (8.62)

and hence:

$$
\sin^2 \theta_{M^2} = \frac{3}{4}
$$
 with $\theta_{M^2} = 240^0$ (8.63)

as the mixing angle. Performing a similar analysis for the type 3 case of $SU(2)^3 \times U(1)^3$ breaking to $U(1)_Q$ leads to a similar result, except with $\sin \theta_{M^3} = +\frac{\sqrt{3}}{2}$ $\frac{\sqrt{3}}{2}$ and $\theta_{M^3} = 120^0$. Setting $\tilde{g} = 1$ the magnitudes of the coupling constants to substitute into equations 8.51 and the bottom line of equation 8.58 in order to match the normalised expressions of equations 8.60 and the right-hand side of equation 8.61 are, relative to \tilde{g} :

$$
\tilde{g}
$$
 : $\tilde{g}' = \sqrt{3}\tilde{g}$: $\tilde{e} = \frac{\sqrt{3}}{2}\tilde{g}$ (8.64)

These values are consistently obtained from equations 8.57 and 8.59 by substituting in the value of θ_{M^2} from equations 8.62 and 8.63, with a similar observation applying for the type 3 case.

This analysis is useful for comparison with the Standard Model for which the gauge groups $SU(2)_L$ and $U(1)_Y$ are not obtained from a single unifying group and hence the respective gauge couplings g and g' of equation 7.40 are independent. Indeed equation 8.60 above may be compared with the form of the covariant derivative of a left-handed doublet of leptons in the Standard Model, which from equations 7.37 and 7.40 can be written as:

$$
D_{\mu} = \partial_{\mu} + igW_{\mu}^{\alpha}(x)\frac{1}{2}\sigma^{\alpha} - ig'B_{\mu}(x)\frac{1}{2}\sigma^{0}
$$
\n(8.65)

In this case the third component of weak isospin $T^3 = \frac{1}{2}$ $\frac{1}{2}\sigma^3$ and hypercharge $\frac{Y}{2} = -\frac{1}{2}$ $rac{1}{2}\sigma^0$ combine to form the charge operator $Q = \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}$ $\frac{0}{7}$ -1) via equation 7.38 for the lepton doublet. In this particular case for equation 7.47 we have the weak mixing combination:

$$
igW^3_\mu \frac{1}{2}\sigma^3 \quad - \quad ig'B_\mu \frac{1}{2}\sigma^0 \quad \Rightarrow \quad ieA_\mu Q
$$

which may be directly compared with:

$$
\tilde{W}_{\mu}^{3} \dot{R}_{\mu}^{(2)} + \tilde{B}_{\mu} \frac{\sqrt{3}}{2} \dot{S}_{l}^{2} \Rightarrow \tilde{A}_{\mu} \frac{\sqrt{3}}{2} \dot{S}_{l}^{1}
$$

from the top line of equation 8.61. In the former case the set of 2×2 matrix actions $\{\frac{1}{2}\sigma^1, \frac{1}{2}\sigma^2, \frac{1}{2}\sigma^3, \frac{1}{2}\sigma^0\}$, as well as forming a basis for elements of the vector space h₂C ⊂ C(2), forms a basis for the Lie algebra $SU(2)_L \times U(1)_Y$ with the normalisation convention $tr(\tau^{\alpha} \tau^{\beta}) = \frac{1}{2} \delta^{\alpha \beta}$, here including $\tau^0 = \frac{1}{2}$ $\frac{1}{2}\sigma^0$ with $\alpha, \beta = 0 \dots 3$. The couplings g and g' are introduced in this basis. For the empirically measured case the electroweak mixing angle is determined to be $\sin^2 \theta_W \simeq 0.23$ at the energy scale of M_Z [44], with corresponding electroweak couplings from equations 7.46 and 7.48 approximately in the proportions:

$$
g \qquad : \qquad g' \simeq 0.55 \, g \qquad : \qquad e \simeq 0.48 \, g \tag{8.66}
$$

Given the unit electron charge for the leptonic component θ_l^1 in equation 8.26 the action of the generator \dot{S}_l^1 is analogous to that of the unit 2×2 matrix σ^0 of equation 7.14. Similarly the normalisation of the type 1 actions $\{\dot{R}_{z_l}^1, \dot{R}_{xz}^1, \dot{R}_{xz}^1\}$, as seen in equations 8.7 and 8.9, parallels the set of 2×2 matrices $\{\tau^1, \tau^2, \tau^3\} = \{\frac{1}{2}\sigma^1, \frac{1}{2}\sigma^2, \frac{1}{2}\sigma^3\}$. Transferring this analysis to the type 2 case, the set of generators $\{\dot{R}_{z_l}^2, \dot{R}_{xz}^2, \dot{R}_{z_l}^2, \frac{1}{2}\dot{S}_l^2\}$ $l \}$ also parallels the set of matrix actions $\{\frac{1}{2}\sigma^1, \frac{1}{2}\sigma^2, \frac{1}{2}\sigma^3, \frac{1}{2}\sigma^0\}.$

Since this is the generator normalisation used initially in equation 8.51 it may naively be expected that the couplings obtained for the $SU(2)^2 \times U(1)^2$ symmetry breaking via the constraint of the E_6 algebra Killing form in equation 8.64 may be directly compared with the corresponding values for $SU(2)_L \times U(1)_Y$ electroweak theory obtained empirically as displayed in equation 8.66. The significant differences in these values hinges on the differing values for the calculated $\sin^2 \theta_{M2} = \frac{3}{4}$ of equation 8.63 and the empirical $\sin^2 \theta_W \simeq 0.23$. However, as emphasised earlier in this subsection the $SU(2)^2$ symmetry does not act on $SL(2,\mathbb{C})$ ¹ Weyl spinors in the appropriate way to describe weak interactions, and here we are dealing with a provisional 'mock electroweak theory', which nevertheless exhibits some of the features associated with corresponding structures of the Standard Model such as the identification of a mixing angle itself.

It is also noted that in the mock theory the calculated value of $\sin^2 \theta_{M^2} = \frac{3}{4}$ 4 effectively corresponds to a 'unification scale' whereas the empirical value of $\sin^2 \theta_W \simeq$ 0.23 is determined at the practical energy scale of $M_Z \sim 10^2 \text{ GeV}$. In standard quantum field theory the phenomena of 'running coupling' for an Abelian compared with a non-Abelian gauge theory implies that the ratio g' : g increases with the energy scale as will be described in section 11.3 and depicted in figure 11.10. Hence the need of a quantisation scheme for the present theory, as alluded to at the end of the previous subsection and as proposed in chapter 11, with the consequence of running coupling, may be one factor leading to the large calculated mixing angle for the present theory. This observation would apply even if the gauge group $SU(2)_L \times U(1)_Y$ were to be correctly identified in the theory.

In any case in this subsection it has been demonstrated how the relative couplings of the internal gauge groups may in principle be related through unification within the simple Lie group E_6 . Finally here we consider how a more realistic electroweak theory might be constructed within this framework. In the above we have assumed a symmetry breaking pattern of $SU(2)^2 \times U(1)^2 \rightarrow U(1)_Q$, whereas these subgroups are actually embedded in a larger symmetry breaking structure with $SU(3)_s \rightarrow U(1)_Q$. That is, instead of equation 8.60 we might rather begin with the gauge covariant derivative:

$$
D_{\mu}\theta_l^2(x) = \partial_{\mu}\theta_l^2(x) + W^{\alpha}_{\mu}(x)\dot{R}^{\alpha}(\theta_l^2)
$$

where now $\alpha = 1...8$ summing over the full basis of eight $SU(3)_s$ generators in equation 8.36. Here all three embeddings of $SU(2)^a \times U(1)^a \subset SU(3)_s$, for type $a = 1, 2, 3$, must come into play with the choice of $SU(2)^1 \subset SL(2,\mathbb{C})^1$ as the rotation subgroup of the Lorentz group acting on external spacetime breaking the symmetry.

While this symmetry breaking structure requires further study the fact that the mock electroweak symmetry $SU(2)^a \times U(1)^a$ may be embedded in $SU(3)_s$ in two ways, of type $a = 2$ or $a = 3$, while the symmetry $SU(2)^{1} \times U(1)^{1}$ has only *one* embedding, of type $a = 1$, may be of some significance. Within $\mathrm{su}(3)_s$ the three $\mathrm{U}(1)^a$ generators are linearly dependent by equation 6.42, while by equation 6.46 only the \dot{R}_{nl}^a part of the three $SU(2)$ ^a generators are linearly dependent. These observations offer a hint that for an internal $SU(2)^a$ combining types $a = 2$ and 3 the ratio of the effective coupling \tilde{g} to that for the effective U(1)_Q coupling \tilde{e} maybe somewhat larger than that for the type $a = 2$ case alone which led the final expression of equation 8.64, once the linear dependencies of the generators are taken into account, which may result in a closer correspondence with the Standard Model case in equation 8.66.

Type 2 gauge fields:

$$
\tilde{W}_{\mu}^{(2)\pm}(x) = \tilde{W}_{\mu}^{(2)1}(x) \mp i\tilde{W}_{\mu}^{(2)2}(x) \tag{8.67}
$$

may be associated with the type 2 generators $\dot{\Sigma}^{(2)\pm}$ of equation 8.37 in the complex $SU(2)^2$ subalgebra (by comparison with equations 7.57 and 7.58 for the Standard Model, although neglecting possible factors of $\frac{1}{\sqrt{2}}$ $\frac{1}{2}$ or $\frac{1}{2}$ here). As described above and as can be seen from table 6.6 the relevant generators \dot{R}_{z}^2 and \dot{R}_{xz}^2 mix the a component of $h_3\mathbb{O}$ with the c component only. The fact that in the mock theory the $SU(2)^2 \times U(1)^2$ symmetry acts on the $\theta^2 = \left(\frac{a}{c}\right)^2$ $\binom{a}{c} \subset h_3 \mathbb{O}$ components, and *not* physical fermion doublets, is one reason not to expect the calculated mixing angle to match the empirical case. That is, while the type 2 symmetry $SU(2)^2 \times U(1)^2$ has some of the properties associated with the electroweak symmetry $SU(2)_L \times U(1)_Y$, the $SU(2)^2$ transformations do not relate the $a \in h_3 \mathbb{O}$ component to *both* components of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$.

On the other hand type 3 gauge fields $\tilde{W}_{\mu}^{(3)\pm}(x) = \tilde{W}_{\mu}^{(3)1}(x) \pm i \tilde{W}_{\mu}^{(3)2}(x)$ may be associated with similar generators in the complex $SU(2)^3$ subalgebra:

$$
\dot{\Sigma}^{(3)\pm} = \dot{R}_{\rm 2d}^3 \mp i \dot{R}_{\rm 2z}^3 \tag{8.68}
$$

The \pm signs are chosen such that the generators $\dot{\Sigma}^{(3)\pm}$, as for $\dot{\Sigma}^{(2)\pm}$ in equation 8.39, carry charges of ± 1 , that is:

$$
[i\dot{S}_l^1, \dot{\Sigma}^{(3)\pm}] = \pm \dot{\Sigma}^{(3)\pm} \tag{8.69}
$$

Adding to the discussion towards the end of the previous subsection, together $\dot{\Sigma}^{(2)\pm}$ and $\dot{\Sigma}^{(3)\pm}$ describe four of the six eigenvectors of the Cartan subalgebra in the Cartan-Weyl basis for the adjoint representation of the complexified $su(3)_s$ algebra. The full set of six eigenvectors are sometimes denoted U^{\pm} , V^{\pm} and T^{\pm} in the su(3) root space diagram (as for example in the context of the $SU(3)$ flavour symmetry between u, d and s-type quarks).

For the case of equation 8.68, as can also be seen from table 6.6, the generators $\dot{R}_{\rm z1}^3$ and $\dot{R}_{\rm x2}^3$ mix the a component of h₃ \odot with the b component only, that is within the $\theta^3 = \begin{pmatrix} b \\ \bar{a} \end{pmatrix}$ $\binom{b}{\bar{a}} \subset h_3\mathbb{O}$ components as shown in equation 8.31. Hence it appears that physical charged gauge boson fields $W^{\pm}_{\mu}(x)$ must indeed be related to both type $2 \dot{\Sigma}^{(2)\pm}$ and type $3 \Sigma^{(3)\pm}$ operators to act on fermion doublets. Ultimately the interactions of the physical W^{\pm} particle states will need to be appropriately oriented with respect to physical fermion states. The latter will in turn require a possible $SL(2,\mathbb{C})^1$ Weyl spinor interpretation of the $a \in h_3\mathbb{O}$ components, which have been provisionally associated with neutrino and u-quark states according the internal $SU(3)_c \times U(1)_Q$ transformations of equation 8.28.

The possible means of identifying Weyl spinor states for the ν -lepton and ν quarks within the $a \in \mathbb{O} \in \mathrm{h}_3\mathbb{O}$ components will be addressed in section 9.1. The identification of both left and right-handed Weyl spinors together with the Dirac representation of the external $SL(2,\mathbb{C})^1$ symmetry will then be described in section 9.2. Finally the possibility of identifying three generations of fermions and the phenomena of CKM mixing will be outlined in section 9.3. All of the above features may need to come together in order to fully identify the physical $SU(2)_L$ symmetry together with standard phenomena of electroweak theory within the context of the present theory. In the meantime in the following subsection we study further suggestive features of the $SU(2)^2 \times U(1)^2$ mock electroweak theory based within the E₆ framework, and in particular concerning the source of finite mass for the both the gauge bosons and the fermion states. Then we shall briefly consider further possible $SU(2) \subset E_6$ subgroups as candidate components of an electroweak symmetry, before extending beyond E_6 in the following chapter.

8.3.3 Origin of Mass and Higgs Phenomena

The empirical weakness of the weak interaction relative to electromagnetic phenomena owes not to the value of the coupling q , in equation 7.40 for example, which is around twice the value of e, equation 8.66, but to the large values for the masses of the W^{\pm} and $Z⁰$ gauge bosons. Although in this chapter we are dealing primarily at the level of the Lie algebra structure, together with the simple dynamic expressions introduced in the previous subsection, it will be considered here how mass terms for particle states may originate in the symmetry breaking structure, not only for the massive gauge bosons but also for leptons and quarks in the full theory. Here \tilde{W}^{\pm} and \tilde{Z}^{0} gauge bosons will be provisionally associated with the appropriate fields of the $SU(2)^2 \times U(1)^2$ mock electroweak theory, and hence we first look in more detail at the field $\tilde{Z}_{\mu}(x)$.

The gauge field $\tilde{Z}_{\mu}(x)$ appearing in the top line of equation 8.61 was identified along with $\tilde{A}_{\mu}(x)$ as aligned to the choice of basis elements $\{\dot{R}_{x}^1, \frac{\sqrt{3}}{2}\}$ $\frac{\sqrt{3}}{2}\dot{S}_l^1$. As described earlier the apparent association of the 'internal' field $\tilde{Z}_{\mu}(x)$ with the 'external' generator \dot{R}_{x}^{1} is one of a number of significant caveats associated with the mock electroweak theory. Through the orthogonal transformation of equation 8.55; that is with $\tilde{Z}_{\mu} = \cos \theta_{M^2} \tilde{W}_{\mu}^3 - \sin \theta_{M^2} \tilde{B}_{\mu}$, in analogy with electroweak theory and equation 7.43, as for the 'photon' field $\tilde{A}_{\mu}(x)$, the field $\tilde{Z}_{\mu}(x)$ is associated with a linear combination of the generators \dot{R}_{nl}^2 and \dot{S}_l^2 . Since in the E₆ Lie algebra $[\dot{S}_l^1, \dot{R}_{\text{nl}}^2] = 0$ and $[\dot{S}_l^1, \dot{S}_l^2] = 0$ any such linear combination of \dot{R}_{nl}^2 and \dot{S}_l^2 has zero electromagnetic charge. Hence the \tilde{Z}^0 gauge boson and $\tilde{\gamma}$ photon, associated with the fields $\tilde{Z}_{\mu}(x)$ and $\tilde{A}_{\mu}(x)$ respectively, are neutral, unlike the case of the charged \tilde{W}^{\pm} gauge bosons associated with fields $\tilde{W}^{(2)\pm}_{\mu}(x)$ of equation 8.67 corresponding to the generators $\dot{\Sigma}^{(2)\pm}$ of equations 8.37 and 8.39. This is also the case when such linear combinations are extended to include the type 3 form of these generators since also $[\dot{S}_l^1, \dot{R}_{ul}^3] = 0$ and $[\dot{S}_l^1, \dot{S}_l^3] = 0.$

From the type 2 \tilde{W}_{μ}^{3} and \tilde{B}_{μ} terms in equation 8.53 it can be seen that the transformations associated with the fields $\tilde{Z}_{\mu}(x)$ and $\tilde{A}_{\mu}(x)$ mix the components of $a \in \mathbb{O}$ within $h_2\mathbb{O}$. This is unlike the more involved transformations associated with the fields $\tilde{W}^{(2)1}_{\mu}(x)$ and $\tilde{W}^{(2)2}_{\mu}(x)$, as can also can be seen in table 6.6 for the corresponding generators \dot{R}_{z}^2 and \dot{R}_{xz}^2 which mix components of $h_2 \mathbb{O} \subset h_3 \mathbb{O}$ with those not in $h_2 \mathbb{O}$. Isolating the interaction of the \tilde{Z}_{μ} field with the a and $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ components separately may allow a determination of the coupling of the \tilde{Z}^0 to the lepton pairs as well as quark pairs, which might be directly compared with the electromagnetic coupling of the photon to the same components as summarised in equations 8.23, 8.26 and 8.28.

This may be more straightforward than for interactions involving the W^{\pm} gauge

bosons as here not only is the interaction restricted to single components but also one generation of fermion states may suffice since there are no flavour changing neutral currents in the Standard Model, as described at the end of section 7.2. Hence a more detailed study of interactions for the field $\tilde{Z}_{\mu}(x)$ in comparison with the field $\tilde{A}_{\mu}(x)$ may prove enlightening in comparison with the relevant properties of the Standard Model described in section 7.2 and in particular with respect to the determination of the relative couplings.

This may involve linear combinations of type 2 and type 3 actions on $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ with generators of the weak neutral field $\tilde{Z}_{\mu}(x)$ being complementary to the generator \dot{S}_l^1 of the electromagnetic field $\tilde{A}_{\mu}(x)$ with respect to the full SU(3)_s $\subset E_6$ symmetry, as considered towards the end of the previous subsection, with the $\tilde{Z}_{\mu}(x)$ field associated with a different linear combination of charge neutral $SU(3)$ _s generators. Ultimately however for comparison with weak neutral interactions described in the Standard Model via equation 7.49 both left-handed and right-handed fermions will need to be identified, such that $T^3 = 0$ for right-handed states, and this itself will require an extension beyond the study of E_6 on the space h₃ \mathbb{O} . Such an extension will also be required to identify the physical $SU(2)_L \times U(1)_Y$ symmetry, independent of the external $SL(2,\mathbb{C})^1$ generators, and fully account for both W^{\pm} and Z^0 interactions.

In the meantime here we consider broader features of the mock electroweak theory as described in the previous two subsections, and in particular how masses may arise for gauge bosons through the impingement of the $SU(2)^2 \times U(1)^2 \subset E_6$ symmetry on the 4-dimensional subspace $h_2\mathbb{C} \subset h_3\mathbb{O}$ associated with the tangent space TM_4 of the external spacetime. Having in mind comparisons with the Standard Model we return to the convention of equation 8.51 with coupling parameters \tilde{g} and \tilde{g}' in place of employing generators normalised according to the E_6 Killing form.

In equation 8.56 the coupling of the $\tilde{A}_{\mu}(x)$ and $\tilde{Z}_{\mu}(x)$ fields to the $\bar{a}_{1,l} = a_1 - a_8l$ components of h₂C \subset h₃^O was extracted. The constraint tan $\theta_{M^2} = \tilde{g}'/\tilde{g}$ was derived in equation 8.57 in order for the $\tilde{A}_{\mu}(x)$ contribution to vanish. With this constraint the impingement of the field $\tilde{Z}_{\mu}(x)$ on the $\bar{a}_{1,l}$ subcomponent part of $\mathrm{h}_2\mathbb{C}$ from equation 8.56 can be written:

$$
D_{\mu}\bar{a}_{1,l} = \dots + \frac{\tilde{g}}{2}\cos\theta_{M^2}\tilde{Z}_{\mu}(a_1l + a_8) - \frac{\tilde{g}'}{2}\sin\theta_{M^2}\tilde{Z}_{\mu}(-a_1l - a_8)
$$

\n
$$
= \frac{1}{2}(\tilde{g}\cos\theta_{M^2} + \tilde{g}'\sin\theta_{M^2})\tilde{Z}_{\mu}(a_1l + a_8)
$$

\n
$$
= (\tilde{g}\cos\theta_{M^2} + \tilde{g}\tan\theta_{M^2}\sin\theta_{M^2})\tilde{Z}_{\mu}\frac{1}{2}(a_1l + a_8)
$$

\n
$$
= \frac{\tilde{g}}{\cos\theta_{M^2}}\tilde{Z}_{\mu}\frac{1}{2}(a_1l + a_8)
$$
(8.70)

This compares with the impingement of the type 2 fields $\tilde{W}^{(2)\pm}_{\mu}(x)$ of equation 8.67, as composed of $\tilde{W}^{(2)1}_{\mu}(x)$ and $\tilde{W}^{(2)2}_{\mu}(x)$, on the same off-diagonal elements of $h_2\mathbb{C}$ in equation 8.54, which is proportional to $\tilde{g}/2$. Hence the coupling of the corresponding gauge fields to the subcomponent $\bar{a}_{1,l}$ of $h_2 \mathbb{C} \subset h_3 \mathbb{O}$ is in the following ratio:

$$
\begin{array}{rcl}\n\tilde{Z}_{\mu} & : & \tilde{W}_{\mu}^{\pm} & : & \tilde{A}_{\mu} \\
\frac{\tilde{g}}{\cos \theta_{M^2}} & : & \tilde{g} & : & 0\n\end{array} \tag{8.71}
$$

This suggests, given the Standard Model expression for M_W in equation 7.60 and its relation to M_Z in equation 7.62, that the interaction with components of h₂C originating here in equation 8.54 is closely related to the masses of the gauge bosons in the present theory (within the caveats that the impingement on all four components of h₂C may need to be addressed and factors of 2 or $\sqrt{2}$ may appear for some terms in a more thorough analysis, but here we are merely noting certain general features of the mock electroweak theory). This structure arises here without the need to introduce a Lagrangian or a custom-built scalar Higgs field ϕ .

Mass terms such as for equations 7.60 and 7.62, arising in the Standard Model Lagrangian, are quadratic in the gauge boson fields due to the quadratic composition $(D_{\mu}\phi)^{\dagger}D^{\mu}\phi$ constructed for the Lorentz invariant initial Lagrangian \mathcal{L}_H in equation 7.51. In the present theory the composition of the gauge fields with the components of h₃^O in expressions such as $D_{\mu}L(v_{27}) = 0$ has a different structure, linear in the gauge fields. Here the concept and nature of particle 'mass' is yet to be identified, and will require an understanding of quantisation and physical particle states as will be described in chapter 11. However the impingement of the 'internal' $SU(2)^{2,3}$ symmetry upon the components of the external spacetime tangent space TM_4 is expected to correlate closely with the phenomenology of the massive W^{\pm} and Z^{0} , involving the kinematic properties of these gauge boson states in spacetime, and hence accounting for the short-range nature of the weak interaction. If gauge boson masses may be obtained through these interactions this raises the question of how further elements of the present theory might correspond to the Higgs sector of the Standard Model.

In the present theory the Lorentz SO⁺(1, 3) symmetry acts on the form $L(\mathbf{v}_4)$ = $(v^{1})^{2} - (v^{2})^{2} - (v^{3})^{2} - (v^{4})^{2} = h^{2}$ of equation 5.46 with the components of $v_{4} \in h_{2} \mathbb{C}$ embedded in h₃^O under the $SL(2,\mathbb{C})^1 \subset E_6$ subgroup action. In a suitable choice of frame a Lorentz 4-vector can be expressed as $v_4 = (v^0, 0, 0, 0)$ which in turn can be written as

$$
\boldsymbol{h}_2 = \left(\begin{array}{cc} v^0 & 0\\ 0 & v^0 \end{array}\right) \tag{8.72}
$$

that is with the three components $v^1 = v^2 = v^3 = 0$ in equations 8.4 and 8.5. This 4vector is invariant under the $SU(2)^1 \subset SL(2,\mathbb{C})^1$ transformations $h_2 \to Sh_2S^{\dagger}$, which preserve the form of equation 8.72, with the type 1 rotations $S \in SU(2)^1 \subset SL(2,\mathbb{C})^1$ generated by $\{\dot{R}_{\underline{z}}^1, \dot{R}_{\underline{z}z}^1, \dot{R}_{\underline{z}l}^1\}$; that is the subset of Lorentz generators in equation 6.57 leaving the $t \equiv v^0$ component in equation 6.58 fixed.

As described in section 7.2 for the Higgs sector an SU(2) custodial symmetry originates as a subgroup of the SO(4) symmetry of the form of the potential $V(\phi)$ = $f(\phi_1^2 + \phi_2^2 + \phi_3^2 + \phi_4^2)$ implicit in equation 7.53. The vacuum value of the Standard Model Higgs field can be expressed in terms of the bi-doublet Φ of equation 7.64 in the form of equation 7.68, that is with $\langle \Phi \rangle = \frac{1}{2}$ $rac{1}{2}$ $\binom{v}{0}$ $\begin{pmatrix} v & 0 \\ 0 & v \end{pmatrix}$, which is invariant under the transformations $\langle \Phi \rangle \to L \langle \Phi \rangle L^{\dagger}$ with $L \in SU(2)_{L+R}$, where $SU(2)_{L+R}$ is the custodial symmetry, highlighting the close similarity to the symmetry of a given Lorentz 4-vector $h_2 \in h_2 \mathbb{C}$ such as that in equation 8.72 in the present theory.

Rather than a Higgs complex doublet field ϕ and Lagrangian \mathcal{L}_H with 'accidental' global SO(4) symmetry (for $g' \to 0$), here we have the Lorentz 4-vector v_4 with an external $SO^+(1,3)$ symmetry for the form $L(\mathbf{v}_4) = h^2$ which is 'spontaneously broken'

by the non-zero particular projected value of $v_4 \in TM_4$. Here the value $v^0 \neq 0$ in equation 8.72 is simply the magnitude of the Lorentz 4-vector v_4 , projected out of the components of $v_{27} \in h_3\mathbb{O}$, onto the tangent space TM_4 ; without the need for a 'Mexican hat potential' such as equation 7.53 to provide the mechanism for 'spontaneous symmetry breaking' and induce a non-zero 'vacuum value' for the field. This choice of vacuum value for $v_4 \in TM_4$ is in addition to the E_6 symmetry breaking through the necessary choice of a Lorentz subgroup $SL(2,\mathbb{C})^1 \subset E_6$ associated with the external spacetime and as explored in the earlier sections of this chapter.

Here the Lorentz symmetry itself, expressed with $h_2 = v_4 \cdot \sigma_4$ as $h_2 \to Sh_2S^{\dagger}$ in equation 7.30 and 7.31, is reduced to the choice of $S \in SU(2)^1 \subset SL(2,\mathbb{C})^1$ for a particular 4-vector $v_4 \in TM_4$, as a close analogy to the custodial symmetry $\text{SU}(2)_{L+R} \subset \text{SU}(2)_L \times \text{SU}(2)_R$ for the Higgs case as described following equation 7.68. However the present theory may need to be developed beyond the model based on the $SU(2)^2 \times U(1)^2$ symmetry towards a more standard $SU(2)_L \times U(1)_Y$ electroweak theory before a more precise correlate of the 'custodial symmetry' might be identified.

In the Standard Model electroweak theory three of the four $SU(2)_L \times U(1)_Y$ generators are spontaneously broken since they change the vacuum expectation value of the Higgs field $\langle \phi \rangle$ in equation 7.54 or $\langle \Phi \rangle$ in equation 7.68), while maintaining the minimum of the Higgs potential $(V(\phi))$ in equation 7.53 or $V(\Phi)$ in equation 7.65). The three degrees of freedom of the Higgs field associated with the broken generators give rise to the mass terms for the W^{\pm} and Z^{0} gauge bosons in the Lagrangian. Fluctuations around the vacuum $v+H(x)$ in the fourth degree of freedom are associated with a mass term for the Higgs scalar particle, which is also proportional to the vacuum value of $v \simeq 246$ GeV, as described shortly after equation 7.62. The unbroken U(1)_Q generator Q leaves both $\langle \phi \rangle$ and $V(\phi)$ invariant, with the vacuum carrying zero electric charge, as described just after equation 7.54.

In the present theory, while the $v_4 \n\in h_2\mathbb{C}$ vacuum value of equation 8.72 is also invariant under the U(1)_Q symmetry, in fact with the S_l^1 action leaving all components of $h_2 \mathbb{C} \subset h_3 \mathbb{O}$ unchanged, the $SU(2)^2 \times U(1)^2$ generators associated with the \tilde{W}^{\pm}_{μ} and \tilde{Z}_{μ} fields mix the v_4 components in h₃^O such that $|v_4|$ is not invariant, unlike the case for $V(\phi)$ in the standard electroweak theory as described above. While here we are dealing with a mock theory the possibility of associating three of the four degrees of freedom for $\delta v_4(x)$ (in particular for the spatial components $\{v^1(x), v^2(x), v^3(x)\}\$) with longitudinal components for the gauge bosons and hence masses for the \tilde{W}^{\pm} and \tilde{Z}^{0} particles may assist in the identification of the physical $SU(2)_L \times U(1)_Y$ electroweak theory within the present framework. In any case here fluctuations in the Lorentz scalar magnitude $|v_4|$ (closely related to variation in the remaining temporal component $v^0 + \tilde{H}(x)$ will be associated with the Higgs field and corresponding massive boson particle state. That is, $H(x) \sim \delta |\mathbf{v}_4(x)|$ in the present theory is provisionally correlated with the scalar field $H(x)$ in the Standard Model.

In the full dynamical quantum theory it will of course be necessary to explain how the phenomenology of the Standard Model Lorentz scalar Higgs field and particle state, as observed in the laboratory, may be derived in detail from the components of the fundamental 4-vector field v_4 , which will be referred to in this context as a 'vector-Higgs'. This structure brings to mind other models for which there is no fundamental Higgs scalar field, with the latter for example composed out of fermion states. Hence here we briefly review some of the properties of technicolor models ([57, 58], see also [46]) for comparison and contrast with the present theory.

For QCD with two flavours $Q_{L,R} = \begin{pmatrix} u & u \\ d & v \end{pmatrix}$ $\binom{u}{d}_{L,R}$ in the massless fermion limit the manifold of vacuum states, that is the 2×2 matrix of scalars $\langle Q_L \overline Q_R \rangle \neq 0$, breaks the global symmetry of the Lagrangian resulting in three Goldstone bosons corresponding to the three pion states π^{\pm} and π^{0} . Coupling the quarks to $SU(2)_{L} \times U(1)_{Y}$ this gauge symmetry is broken by the vacuum since the SU(2) only couples to the lefthanded fermions, while a $U(1)_O$ gauge symmetry is preserved. The symmetry breaking generates masses for the corresponding W^{\pm} and Z^{0} gauge bosons which are, however, too small compared to the empirical values since the pion decay constant f_{π} is only around 93 MeV.

Motivated by these observations and difficulties associated with a fundamental scalar Higgs in the Standard Model, a new strongly interacting sector of fermions called 'techniquarks' is postulated which couple to a new 'technicolor' gauge symmetry $\text{SU}(N)_{tc}$. The techniquarks $T_{L,R} = {U \choose D}_{L,R}$ also transform under $\text{SU}(2)_L \times \text{U}(1)_Y$ but are singlets under the standard colour symmetry $SU(3)_c$. Scalar combinations of T and \overline{T} condense in the vacuum owing to the new strong technicolor interaction. As for the QCD case above the vacuum state is termed a 'condensate' by analogy with phenomena in condensed matter physics, and in particular the formation of BCS pairs of electrons in superconductivity.

For such a model the technipion decay constant may be taken to be $F_{\Pi} \simeq$ 246 GeV to replicate the masses of the W^{\pm} and Z^{0} as previously obtained with a scalar Higgs sector. The Standard Model relation $M_W/M_Z = \cos \theta_W$ of equation 7.62 is also reproduced. The techniquark Lagrangian includes kinetic terms of the form:

$$
\mathcal{L}_{tc} \sim \overline{T}_{L,R} \gamma^{\mu} (\partial_{\mu} + i g_N G_{tc\mu} + i g W_{\mu} + i g' B_{\mu}) T_{L,R}
$$
(8.73)

with technicolor gauge field $G_{tc\mu}$ and coupling g_N as well as the $SU(2)_L \times U(1)_Y$ gauge fields and couplings. In the quantum field theory masses for the W^{\pm} and Z^{0} gauge bosons are generated by the corrections introduced into the corresponding gauge boson propagators through the interaction terms in equation 8.73, with massless technipions effectively appearing as the longitudinal components of massive W^{\pm} and Z^{0} bosons. The low energy behaviour can be described by an effective phenomenological Lagrangian for the vacuum expectation value $\Psi(x) = \langle T_L \overline{T}_R \rangle \neq 0$ with:

$$
\mathcal{L}_{\Psi} \sim F_{\Pi}^{2} \operatorname{tr}(D_{\mu} \Psi^{\dagger} D^{\mu} \Psi) \tag{8.74}
$$

For the two techniquark model the scalar $\Psi(x)$ is a 2×2 matrix which plays the role of the scalar doublet of Higgs fields. The form of equation 8.74 is analogous to the kinetic term in the Higgs Lagrangian in the form of equation 7.66 for the bi-doublet field $\Phi(x)$.

Masses for ordinary quarks and leptons are introduced by replacing the scalar Higgs in the Standard Model Yukawa terms of equation 7.69 by techniquark bilinears resulting in 4-fermion interactions with quartic terms such as the scalar:

$$
\mathcal{L} \sim \overline{Q}_L \left(\mathbf{1}_2 T_L \overline{T}_R \right) Q_R \tag{8.75}
$$

Here $Q_{L,R}$ are ordinary quarks which gain mass when the techniquarks form a condensate $\langle T_L \overline{T}_R \rangle \neq 0$. A suitable variety of quartic interactions and coupling parameters are needed to reproduce the empirical values for the standard quarks and leptons. Higher-order Lagrangian terms such as 6-fermion interactions may also be considered.

As a theory of electroweak symmetry breaking without a fundamental scalar Higgs the above technicolor model has some resemblance with the present theory. The structure of the 2×2 scalar condensate $\langle T\overline{T}\rangle$ as composed out of fermions indeed bares some resemblence to the spinor decomposition of the vector $h_2 = \chi \chi^{\dagger} + \phi \phi^{\dagger}$ of equation 7.32. However the latter expression merely represents the algebraic substructure within the components of h_2 without the need of a new *technicolor interaction* with gauge group $SU(N)_{tc}$ to condense fermions into a single object. Hence for the present theory the association of the scalar Higgs with the scalar magnitude $|v_4|$ of the 'vector-Higgs' $v_4 \equiv h_2$ is analogous to technicolor models with a scalar condensate composed of a new set of fermions, in that in both cases the need to postulate a fundamental scalar Higgs field is avoided. The absence of any observation of states belonging to a technihadron spectrum rules out a number of technicolor models.

Unlike the case of the Standard Model for the present theory the mass for the W^{\pm} and Z^{0} states is expected to arise from terms in $D_{\mu}L(v_{27})=0$ which are linear in the gauge fields, as suggested in part by the relative couplings listed in equation 8.71 and described earlier in this subsection. These gauge field interaction terms are in fact similar in structure to those of the Lagrangian of equation 8.73 and hence the origin of the gauge boson masses in the present theory also resembles the corresponding structure of the technicolor model. At a suitably low energy scale the present theory might also be compatible with an effective Lagrangian term quadratic in the gauge fields similar to equation 8.74 for the technicolor case.

An origin for the masses of the fermion states in the present theory is also required, as a correlate of the 'Yukawa interactions' introduced in the Standard Model Lagrangian. As described above for the W^{\pm} and Z^{0} gauge bosons, mass terms for the fermions might also be expected to involve a form of coupling with the external components $v_4 \n\t\in h_2 \mathbb{C} \subset h_3 \mathbb{O}$, which have been shown to exhibit properties analogous to those of the Standard Model Higgs field. The expression of the full form of $L(\mathbf{v}_{27}) =$ 1 as the determinant of $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ matrices, as written out in equation 6.28, includes the terms $p|b|^2$, $m|c|^2$ and $n|a|^2$. With the projected components on TM_4 related under $L(v_4) = h^2$ in equation 5.46 and adopting the $v_4 \equiv h_2$ components of equation 8.72 with $v^0 = h$ embedded within h₃ \mathbb{O} as for equations 8.4 and 8.5 the determinant can be written as:

$$
\det(\mathcal{X}) = h^2 n - h|b|^2 - h|c|^2 - n|a|^2 + 2\text{Re}(\bar{a}\bar{b}\bar{c}) = 1 \tag{8.76}
$$

The b, c internal components of $h_3\mathbb{O}$ hence have a multiplicative 'coupling' with the vacuum value h from the components of v_4 in the form $h(b\bar{b}+c\bar{c})$. These are analogous to the Yukawa coupling terms between the Higgs field and fermion fields in Standard Model Lagrangian of equation 7.69. This suggests that the fermion masses may be proportional to h , in a similar way that they are proportional to the Higgs field vacuum value v, equation 7.70, in the Standard Model.

Since $L(v_{27}) = 1$ is invariant under the transformations of E_6 , and hence also under the external and internal subgroups, in the dynamics of the theory the actual mass terms may correspond to gauge invariant expressions as for the case of the Lagrangian approach. Possible quartic or higher-order terms within a higherdimensional form of $L(\mathbf{v}) = 1$ as a source of mass for the standard fermions, considered

towards the end of section 9.2, are also analogous to the technicolor Lagrangian terms of the form of equation 8.75, at least with regard to their non-standard quartic nature.

As for the case of the massive gauge bosons, for which quadratic mass terms do not arise in the basic elements of the present theory as discussed above, ultimately comparison between this theory and the Standard Model should be made at the level of empirical phenomena rather than a Lagrangian, which in any case is absent in the present theory. In addition to the field dynamics the role of mass in the calculations of quantum field theory and its relation to 'renormalisation' and physical particle states as studied in high energy physics experiments will need to be understood, as will be discussed in chapter 11.

Although only one generation of fermions has so far been considered in relation to the components of h₃ \mathbb{O} , in the discussion following equation 8.31 in subsection 8.3.1 it was hinted that for the full theory the existence of three generations of physical fermion states might ultimately be correlated with the existence of three types of $SL(2,\mathbb{O}) \subset E_6$ subgroup action, as introduced in equations 6.32–6.35. From this perspective, given the asymmetric structure of the three terms hbb, hcc and na \bar{a} with respect to h in equation 8.76, and the need for renormalisation in the full theory, it is possible that the physical mass eigenstates of empirically studied particles will not be aligned neatly with the type $a = 1, 2$ and 3 spinor θ^a components of h₃ \mathbb{O} . Instead the choice of the external $v_4 \in h_2 \mathbb{C} \subset h_3 \mathbb{O}$ may be skewed relative to the three generations of physical fermions, which may each then be related to v_4 via a *continuous* (defined in [38] p.127, as alluded to here after equation 6.35) rather than *discrete* type transformation, leading to the spectrum of masses observed for the leptons and quarks.

In the Standard Model the phenomena of CKM mixing in the quark sector relates to a mismatch between weak interaction and mass eigenstates as was reviewed in section 7.2. In this section we have established a correlation between the weak interaction and the subgroups $SU(2)^2$ and $SU(2)^3$ of E_6 in the context of the present theory. If the mass states of three generations of quarks are skewed into the components of $h_3\mathbb{O}$ via a continuous type transformations as described above this contrasts with the charged weak interaction of the \tilde{W}^{\pm} gauge bosons associated with the SU(2)^{2,3} actions which constitute a discrete type complement to the external $SL(2,\mathbb{C})^1$ symmetry. This structure hence provides a possible basis for the mismatch between weak and mass eigenstates responsible for the CKM mixing between three generations of quarks, with a similar structure accounting for neutrino oscillations in the leptonic sector.

As described earlier the transformations for the symmetry group $SU(3)_c \subset E_6$, generated by $\{\dot{A}_q, \dot{G}_l\}$, act on each of the three $a, b, c \in \mathbb{O}$ components of $h_3\mathbb{O}$ in exactly the same way, in manner that is independent of both discrete and continuous type transformations (as contrasted with the $SU(3)_s$ actions after equation 8.32). In the present context this symmetry of the $SU(3)_c$ action on the three octonion components in h₃ \mathbb{O} , together with its independence from the SL $(2,\mathbb{C})^{1,2,3}$ and $S_l^{1,2,3}$ $\iota^{1,2,3}$ transformations, is likely to be physically relevant for the observation of three generations of fermions, at least for the quark content and the corresponding phenomena of CKM mixing between generations with each of the three generations of quarks subject to an identical coupling to the $SU(3)_c$ strong interaction gauge bosons.

However while the existence of three generations of fermions may ultimately be correlated with the three types of embedding of the $\theta^{1,2,3}$ components in h₃ \mathbb{O} , as presumed for the discussion above, a somewhat larger space will be required to explicitly house all of the degrees of freedom, as we shall explore in the following chapter. The expansion of the form of temporal flow $L(\mathbf{v}) = 1$ will be accompanied by a corresponding expansion of the group of symmetry transformations, opening up the possibility of identifying an internal $SU(2)_L \times U(1)_Y$ symmetry matching all the properties of the Standard Model.

Finally in this section we consider further possible candidates for the Standard Model $SU(2)_L$ gauge symmetry in terms of generators confined to the E_6 Lie algebra in the present theory. We return to $\{\dot{R}_{z_l}^a, \dot{R}_{xz}^a, \dot{R}_{xt}^a, \dot{B}_{tx}^a, \dot{B}_{tt}^a, \dot{B}_{tz}^a\}$ as the three sets of six generators for $SL(2,\mathbb{C})^a$ for each of $a=1,2,3$ (the $a=1$ set was listed in equation 6.57) with the Lorentz Lie algebra of table 8.1 satisfied in all three cases, as considered in subsection 8.3.1. In particular we look more generally to construct explicit $SU(2)$ subgroups out of the collection of 12 generators of this form with $a = 2$ or 3. These form a subset of the 16 generators for the $sl(3,\mathbb{C})_s$ subalgebra described in equation 8.33 and presented explicitly within table 6.6, taking $q = l$, including the elements $\dot{R}_{rl}^{2,3}$ $x^{2,3}$ and \dot{B}_{tz}^3 which do not belong to the preferred 78-dimensional basis for E_6 .

As for the case of the six generators of the Lorentz algebra, listed equation 6.57 and table 8.1, in the *complexified* Lie algebra the $SL(2,\mathbb{C})^2$ subalgebra of type 2 is also isomorphic to $SU(2) \times SU(2)$. The generators of these two $SU(2)$ s may be denoted A^a and B^b , in correspondence with equations 7.21–7.23 and 8.7–8.9 (within the choice of sign conventions as noted for the latter equations), with:

$$
\{A^{1}, A^{2}, A^{3}\} = \{\frac{1}{2}(\dot{R}_{\mathcal{A}}^{2} + i\dot{B}_{\mathcal{L}}^{2}), \frac{1}{2}(\dot{R}_{\mathcal{Z}\mathcal{Z}}^{2} + i\dot{B}_{\mathcal{L}}^{2}), \frac{1}{2}(\dot{R}_{\mathcal{Z}\mathcal{L}}^{2} + i\dot{B}_{\mathcal{L}}^{2})\}
$$

and
$$
\{B^{1}, B^{2}, B^{3}\} = \{\frac{1}{2}(\dot{R}_{\mathcal{Z}\mathcal{I}}^{2} - i\dot{B}_{\mathcal{L}}^{2}), \frac{1}{2}(\dot{R}_{\mathcal{Z}\mathcal{Z}}^{2} - i\dot{B}_{\mathcal{L}}^{2}), \frac{1}{2}(\dot{R}_{\mathcal{Z}\mathcal{L}}^{2} - i\dot{B}_{\mathcal{L}}^{2})\}
$$

such that:
$$
[i\dot{S}_{l}^{1}, (A^{1} \pm iA^{2})] = \pm(A^{1} \pm iA^{2})
$$

and
$$
[i\dot{S}_{l}^{1}, (B^{1} \pm iB^{2})] = \pm(B^{1} \pm iB^{2})
$$

with the latter two expressions hence describing charge eigenstates. Such eigenstates might in principle be correlated with charged gauge bosons \tilde{W}^{\pm} as described in the previous two subsections. A similar analysis follows for the $SL(2,\mathbb{C})^3$ subalgebra of type 3. In addition to this by using the full set of 12 generators for both $SL(2,\mathbb{C})^2$ and $SL(2,\mathbb{C})^3$ two further $SU(2)$ s can be identified in the complexified algebra in this case with A^a and B^b composed as:

$$
A^{1} = \frac{1}{\sqrt{2}} (\dot{R}_{z}^{2} + \dot{R}_{z}^{3} + i\dot{B}_{tz}^{2} + i\dot{B}_{tx}^{3})
$$
\n
$$
B^{1} = \frac{1}{\sqrt{2}} (\dot{R}_{z}^{2} + \dot{R}_{z}^{3} - i\dot{B}_{tz}^{2} - i\dot{B}_{tx}^{3})
$$
\n
$$
A^{2} = \frac{1}{\sqrt{2}} (\dot{R}_{xz}^{2} + \dot{R}_{xz}^{3} + i\dot{B}_{tt}^{2} + i\dot{B}_{tt}^{3})
$$
\n
$$
B^{2} = \frac{1}{\sqrt{2}} (\dot{R}_{xz}^{2} + \dot{R}_{xz}^{3} - i\dot{B}_{tt}^{2} - i\dot{B}_{tt}^{3})
$$
\n
$$
B^{3} = (\dot{R}_{x1}^{2} + \dot{R}_{x1}^{3} - i\dot{B}_{tz}^{3} - i\dot{B}_{tz}^{3})
$$
\n
$$
B^{3} = (\dot{R}_{x1}^{2} + \dot{R}_{x1}^{3} - i\dot{B}_{tz}^{2} - i\dot{B}_{tz}^{3})
$$
\n(8.77)

However in this case none of the linear combinations $(A^1 \pm iA^2)$ or $(B^1 \pm iB^2)$ is a charge eigenstate of $i\dot{S}_l^1$ under the adjoint representation in the complexified E_6 algebra. In any case in order to identify a candidate for the $SU(2)_L$ gauge symmetry of the Standard Model a *real* $SU(2)$ subalgebra of the *real* form of E_6 is required. Such a compact real form of $SU(2)$ can be obtained from a combination of the type 2 and 3 rotation generators with:

$$
\{J^1,J^2,J^3\}=\{\sqrt{2}(\dot{R}_{\text{2d}}^2+\dot{R}_{\text{2d}}^3),\sqrt{2}(\dot{R}_{\text{2d}}^2+\dot{R}_{\text{2d}}^3),2(\dot{R}_{\text{2d}}^2+\dot{R}_{\text{2d}}^3)\}
$$

However again here a complex linear combination of J^1 and J^2 fails to form a charge eigenstate under $i\dot{S}^1_l$ ¹. It can also be noted that the third generator J^3 is in fact equal to $-2\dot{R}_{\rm at}^1$, by equation 6.46, which is a generator of the type 1 SU(2)¹ rotation subgroup and hence not even independent of the external Lorentz symmetry $SL(2,\mathbb{C})^1$ in terms of the vector space of generators. A similar observation applies to $A³$ and $B³$ in equation 8.77, and indeed was also noted for the gauge field $\tilde{Z}_{\mu}(x)$ associated with \dot{R}_{x}^{1} for the mock $SU(2)^2 \times U(1)^2$ theory before equation 8.61. These observations are not surprising since the Dynkin analysis for the Lie algebra of E_6 in section 7.3 suggests that it is not possible to append any SU(2) subgroup alongside an $SL(2,\mathbb{C})\times SU(3)\times U(1)\subset$ E⁶ decomposition, as recalled near the opening of subsection 8.3.1.

However, of the possible $SU(2)$ structures examined within the E_6 algebra, which in some sense are complementary to the type 1 Lorentz subgroup $SL(2,\mathbb{C})^1$, the subgroups $SU(2)^2$ and $SU(2)^3$ are the most promising in terms of properties *resembling* the $SU(2)_L$ gauge symmetry of the Standard Model, as has been described in this section. These observations supplement the identification of the subgroup $SL(2,\mathbb{C})^1 \times$ $\text{SU}(3)_c \times \text{U}(1)_Q \subset E_6$ in equation 8.25 which exhibits properties correlating closely with features of the Standard Model as described in sections 8.1 and 8.2. These observations also helped motivate the detailed study of the subgroup $SU(2)^2 \times U(1)^2$ in this section in an attempt to account for aspects of electroweak theory within the scope of the E_6 action on the form of $L(v_{27}) = 1$ in the present theory.

While a number of features of this mock electroweak theory resemble those of the Standard Model the lack of a complete match, together with the knowledge that the full Standard Model external and internal symmetry cannot be accommodated within E_6 , now motivates the consideration of a higher-dimensional form of temporal flow, with a higher degree of symmetry, with the goal of incorporating the physical $SU(2)_L$ gauge symmetry. The aim will be to retain the significant traits of electroweak theory as identified in this section, within the breaking of the E_6 symmetry of $L(v_{27}) = 1$ over the external spacetime M_4 , in developing a higher-dimensional expression. As a further feature in reconstructing the full details of the Standard Model it will be necessary to explain how a set of Weyl spinors might be obtained from the $a \in \mathbb{O} \subset h_3\mathbb{O}$ components listed in equation 8.28 for the ν -lepton and u -quark states. This will be the topic of section 9.1. In section 9.2 an explicit higher-dimensional form of $L(\mathbf{v}) = 1$ will be presented resulting in the identification of both left and right-handed Weyl spinors. Finally, bearing in mind the need to incorporate three generations of fermions, the possibility of a further expansion will be described in section 9.3, with the features of the Standard Model so far identified in the context of the present theory then summarised.

Chapter 9

Further Dimensions

9.1 Expanding h_3 ^O and Further Weyl Spinors

In aiming towards the identification of a physical $SU(2)_L$ symmetry acting on doublets of $SL(2,\mathbb{C})^1$ Weyl spinors in the present theory we first recall how the symmetry E_6 acting on the space $h_3\mathbb{O}$ relates to lower-dimensional forms of temporal flow expressed as $L(v) = 1$. In particular the E_6 symmetry of the cubic form $\det(\mathcal{X}) = L(v_{27}) = 1$, with $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$, may be contrasted with the case of taking the full symmetry of $L(\mathbf{v}) = 1$ to be the group $Spin^+(1,9)$ acting on the space $h_2\mathbb{O}$, intermediate in size between $h_2\mathbb{C}$ and h₃ \mathbb{O} , such that the quadratic form $\det(X)$, with $X = \binom{p - \bar{a}}{a - m} \in \mathbb{h}_2 \mathbb{O}$, is preserved as the full form of temporal flow. With $Spin^+(1,9)$ being the double cover of $SO^+(1,9)$ acting on the form $L(\mathbf{v}_{10}) = 1$ this is essentially the model described in section 5.1 as depicted in figure 5.1. In this case there is an $SL(2,\mathbb{C}) \subset Spin^+(1,9)$ subgroup, based on the choice of an imaginary octonion unit such as $q = i$ for equation 6.19, which acts as the external symmetry of 4-dimensional spacetime upon the subspace h₂ $\mathbb{C} \subset h_2\mathbb{O}$ with the two-sided action of equation 6.22 similarly as for the 10-dimensional case. This breaks the set of 45 generators of $SL(2, \mathbb{O}) \equiv$ Spin⁺(1,9) acting on the space h₂ \mathbb{O} to an internal $\text{Stab}_2(TM_4)$ set of symmetry operations which here consist purely of transverse rotations amongst the remaining six imaginary units of the $a \in \mathbb{O}$ component of h₂ \mathbb{O} . This is again sufficient to contain $SU(3)_c \times U(1)_Q$ as an internal symmetry group. Indeed it can be seen from the Dynkin diagram of figure 7.2(b), by removing the central node with the most connections, that the Lie algebra so(10) has a breaking pattern to $sl(2,\mathbb{C}) \times su(3) \times u(1)$.

However in the present theory we are not restricted to the consideration of extra *spatial* dimensions, which might lead to the study of such a $Spin⁺(1, 9)$ symmetry of 10-dimensional spacetime. Here we are dealing with a higher-dimensional form of temporal flow, allowing the structure of the above paragraph to be augmented to the group E_6 acting as the symmetry of a cubic form on the 27-dimensional space h₃ \mathbb{O} . This larger structure incorporates three interlocking Spin⁺(1,9) actions, with associated representations on three spinor spaces $\theta^a \in \mathbb{O}^2$, for $a = 1, 2$ or 3, identified in the additional components as described in section 6.4. Within this struc-

ture the components of $\theta^1 = \begin{pmatrix} \frac{c}{b} \end{pmatrix}$ and a within h₃^O transform under the internal $\text{SU}(3)_c \times \text{U}(1)_Q \subset \text{Stab}(TM_4)$ symmetry as a generation of leptons and quarks, with the appropriate fractional charges, as summarised in equations 8.26 and 8.28 of section 8.2. The three-way embedding of $SL(2,\mathbb{O}) \subset SL(3,\mathbb{O})$ is analogous to the empirical observation of three generations of leptons and quarks, although it remains to be seen whether these features do actually correlate.

Further, in augmenting the 2×2 matrices in h₂ \mathbb{O} , upon which the symmetry of 10-dimensional spacetime may be represented, to the 3×3 matrices of h₃ \mathbb{O} , with the structure of a temporal symmetry, Weyl spinor states are identified in the θ^1 components of the additional column of this matrix, as listed in equation 8.13, under the external $SL(2,\mathbb{C})^1$ symmetry. This is analogous to the motivation of the original Kaluza-Klein theories [11, 12] in which the 4×4 metric $g_{\mu\nu}$ describing the gravitational field is augmented to the case of a 5-dimensional spacetime such that the four components $A_{\mu} = g_{\mu 5}$ in the extra column of the 5 \times 5 metric describe the electromagnetic field. (This structure was later generalised to incorporate the non-Abelian case in a larger spacetime with gauge field components ω^{α}_{a} included in the metric, as described for equation 4.5 in section 4.1). In the present theory we identify fermion states in the extra temporal components, rather than gauge bosons in the additional metric components.

For the case of E₆ acting on $L(v_{27}) = 1$ not only is the above set of four Weyl spinors under the external $SL(2,\mathbb{C})^1$ symmetry identified in the θ^1 components, as described in section 8.1, but they are also seen to be aligned with the internal $\text{SU}(3)_c \times \text{U}(1)_Q$ transformation properties of the electron and a triplet of d-quark states, as described in section 8.2. In the present theory rather than generalising to a higher-dimensional spacetime here the augmentation is applied to a multi-dimensional form of temporal flow, the concept of which is further compared and contrasted with the Kaluza-Klein approach in chapter 5. These temporal structures are not restricted to a quadratic, or even cubic, form and in principle may be extended to a homogeneous polynomial form of arbitrary order.

In the previous section we assessed the possibility of identifying the structure of electroweak theory in the breaking of the E_6 symmetry on the components of h₃ \mathbb{O} . For example the gauge field components corresponding to \tilde{W}^{\pm} and \tilde{Z}^{0} gauge bosons, identified in a mock electroweak theory based on the $SU(2)^2 \times U(1)^2 \subset E_6$ symmetry, impinge on the external h₂C $\equiv TM_4$ components as described in equation 8.54, which led to equation 8.71, providing a possible mechanism for identifying gauge boson mass terms analogous to the standard Higgs sector. In addition the physical W^{\pm} states of the Standard Model act as charge raising and lowering transformations in interactions with left-handed doublets of leptons and quarks, that is $\binom{\nu}{e}$ $\binom{u}{e}$ _L and $\binom{u}{d}$ $\binom{u}{d}_L$ respectively for the first generation of fermions as listed in equation 7.36, where each component $\{\nu, e, u, d\}$ is a left-handed Weyl spinor under the external $SL(2, \mathbb{C})$ symmetry.

One aim of the present theory has been to derive the spectrum of particle states of the Standard Model, and in particular the above doublets of left-handed fermions, from the components of $h_3\mathbb{O}$ under the broken E_6 action. Towards this end the action of the $\tilde{W}^{\pm}_{\mu}(x)$ fields associated with both the $\mathrm{SU}(2)^2$ and $\mathrm{SU}(2)^3$ transformations on the e-lepton and d-quark states, which have already been associated with the components of $\theta^1 = \begin{pmatrix} \frac{c}{b} \end{pmatrix}$ through equation 8.26, should serve as a useful guide. As can be seen from equations 8.29 and 8.31 the $SU(2)^2$ and $SU(2)^3$ actions mix the components of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix} \in \mathbb{O}^2$ with the $a \in \mathbb{O}$ component. Further, the subcomponents of a transform in the same way as those of b and c under the internal $SU(3)_c$ symmetry, containing a colour singlet and a colour triplet, and the corresponding elements of a have the correct electromagnetic charges of 0 and $\frac{2}{3}$ under the U(1)_Q generator \dot{S}_l^1 to described the ν lepton and u -quark respectively as can be seen in equations 8.27 and 8.28 and reviewed above. However, as described in table 8.2 under the external Lorentz transformations of $SL(2,\mathbb{C})^1$ the 'leptonic' components of a transform as part of the vector v_4 while the 'quark' components $a(6)$ are scalars; and hence these components appear to be unsuitable to describe fermion states. In this section we focus on the possible means of constructing these further required spinors.

To see how such spinors may potentially arise and account for the ν -lepton and u-quark states the 10 real components of $X \in h_2\mathbb{O}$, embedded in the type 1 location of h_3 O as depicted in equations 6.26, 6.29 and 6.32, may be provisionally composed in terms of the 16 real components of a new object $\theta_X^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $(\bar{r}_s) \in \mathbb{O}^2$, with $r, s \in \mathbb{O}$ and

$$
X = \theta_X^1 \theta_X^{1^{\dagger}} \tag{9.1}
$$

Hence the vector X is considered to be the square of the spinor θ_X^1 , in the form as originally presented in equation 6.20. The compatibility relationship between the vector and spinor actions for the octonion case in equation 6.23 also applies here since the 2×2 transformation matrices $M \in SL(2, \mathbb{O})^1$ are required to have this property:

$$
M X M^{\dagger} = M (\theta_X^1 \theta_X^{1\dagger}) M^{\dagger} = (M \theta_X^1) (M \theta_X^1)^{\dagger}
$$
\n(9.2)

In particular this shows that under the Lorentz transformations via $M = S \in SL(2, \mathbb{C})^1$ the components of $\theta_{X}^{1} = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ \bar{s}) decompose into a set of four Weyl spinors, as is the case for $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ under the *same* transformations as described in equations 8.10–8.13. This hence shows how in principle further left-handed Weyl spinors may be indeed be identified within the E_6 action on $L(v_{27}) = 1$ by opening up further dimensions through the decomposition of equation 9.1.

In terms of the real p, m and octonion a, r, s components equation 9.1 can be written in more detail as:

$$
X = \begin{pmatrix} p & \bar{a} \\ a & m \end{pmatrix} = \theta_x^1 \theta_x^{1^{\dagger}} = \begin{pmatrix} \bar{r} \\ s \end{pmatrix} \begin{pmatrix} r & \bar{s} \end{pmatrix} = \begin{pmatrix} \bar{r}r & \bar{r}\bar{s} \\ sr & s\bar{s} \end{pmatrix}
$$
(9.3)

that is with $p = |r|^2$, $m = |s|^2$ and $a = sr$ (and as may be compared with equation 6.20). The fact that there are 16 real components of θ_X^1 given the original 10 real components of X is compatible with the underlying conceptual motivation of the present theory for which an *n*-dimensional form of temporal flow, such as $L(v_{27}) = 1$, is derived given the original 1-dimensional progression of time, and represents what is essentially a further generalisation and extension of this idea to a still higher-dimensional structure. This structure provides a means to identify a set of Weyl spinors from the external $SL(2,\mathbb{C})^1$ Lorentz action on X which might in principle be associated with physical particle states.
We shall consider how these new spinors identified within the components of θ_{X}^{1} may correlate with the first generation *v*-lepton and *u*-quark states, in a similar way that the e-lepton and d-quark states were identified within the components of θ^1 according to equation 8.26. These fermion states corresponding to $SL(2,\mathbb{C})^1$ Weyl spinors will be required to be mutually oriented within the components of θ_X^1 and θ^1 with respect to \tilde{W}^{\pm} gauge bosons which mix the corresponding leptonic or quark states, raising or lowering the electromagnetic charge of the fermion state by one unit. This SU(2) mixing of Weyl spinors *between* the components of θ_X^1 and θ^1 should be analogous to the $SU(3)_c$ mixing of the $\theta_{i,j,k}$ Weyl spinors within the θ^1 components as described on the left-hand side of table 8.7.

The close relationship between the 10-dimensional vector $X \in h_2\mathbb{O}$ and 10dimensional vectors of the form $\theta\theta^{\dagger} \in h_2\mathbb{O}$ is also exhibited by the 10-dimensional Lorentz inner product in the final term of equation 6.27 for $\det(\mathcal{X})$ with $\mathcal{X} \in h_3\mathbb{O}$. The 10-dimensional type 1 Lorentz transformations $SL(2, \mathbb{O})^1$ leave both terms of $det(\mathcal{X})$, that is both $\det(X)n$ and $2X \cdot (\theta^1 \theta^{1\dagger})$, invariant. However the subgroups $SU(2)^{2,3} \subset E_6$ mix the components of these two terms, as well as mixing components between X and θ ¹, and it is these properties which might be studied in order to describe for example a $u \leftrightarrow d$ -quark interaction in terms of $\theta_x^1 \leftrightarrow \theta^1$ components.

Substituting the $\theta_X^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $\binom{\bar{r}}{s}$ components r and s in place of p, m and a from equation 9.3 into the expression for $\det(\mathcal{X})$ in equation 6.28 leads directly to:

$$
\det(\mathcal{X}) = |r|^2 |s|^2 n - |r|^2 |b|^2 - |s|^2 |c|^2 - n|sr|^2 + 2\text{Re}(\bar{r}\bar{s}\bar{b}\bar{c})
$$
\n(9.4)

Since for any r, s in the division algebra \mathbb{O} we have $|s||r| = |sr|$ the first and fourth terms above cancel, leaving a quartic expression in $r, s, b, c \in \mathbb{O}$. Hence in principle equation 9.4 describes a homogeneous form $L(v) = 1$ with 32 dimensions, namely the real components of $\{r, s, b, c\} \in \mathbb{Q}$, with a symmetry group deriving from the action of E_6 on $L(\mathbf{v}_{27}) = 1$.

However, one significant difference between any elements $X \in h_2\mathbb{O}$ and $\theta \in$ \mathbb{O}^2 with $\theta\theta^{\dagger} \in h_2\mathbb{O}$ is that in the former case $\det(X) \in \mathbb{R}$ may take arbitrary real values while in the latter case we necessarily have $\det(\theta \theta^{\dagger}) = 0$, as was described in equation 6.21. This can be seen here from the right-hand side of equation 9.3 for which $\det(\theta_x^1 \theta_x^{1^{\dagger}}) = |r|^2 |s|^2 - |r|^2 |s|^2 = 0$, for any $\theta_x^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ (\bar{r}_s) , and accounts for the cancellation of the two quintic terms in equation 9.4. It also clearly implies that the decomposition of X as suggested in equations 9.1 and 9.3 is not possible for the general case if $\det(X) \neq 0$.

This apparent incompatibility may be remedied by further generalising equations 9.1 and 9.3 by introducing an additional spinor $\phi_{X}^{1} = \begin{pmatrix} \bar{r}' \\ s' \end{pmatrix}$ $(\bar{r}'_{s'}) \in \mathbb{O}^2$, with identical transformation properties as the original $\theta_{\rm x}^1$ in equation 9.2, such that:

$$
X = \theta_X^1 \theta_X^{1\dagger} + \phi_X^1 \phi_X^{1\dagger} \tag{9.5}
$$

with
$$
MX M^{\dagger} = M(\theta_X^1 \theta_X^{1^{\dagger}}) M^{\dagger} + M(\phi_X^1 \phi_X^{1^{\dagger}}) M^{\dagger}
$$

$$
= (M\theta_X^1)(M\theta_X^1)^{\dagger} + (M\phi_X^1)(M\phi_X^1)^{\dagger}
$$
(9.6)

This introduces a further 16 real parameters in ϕ_X^1 which transform as a further set of four Weyl spinors under the Lorentz actions with $M = S \in SL(2, \mathbb{C})^1$. The value of $\det(\theta_X^1 \theta_X^{1 \dagger} + \phi_X^1 \phi_X^{1 \dagger}) \in \mathbb{R}$ may now be compatible with the determinant of X in the general case. This is analogous to the case of the 4-dimensional Lorentz vector decomposition in equation 7.32, with the spinor substructure of the vector X potentially providing a source of microscopic physical structure, as suggested after equation 7.32 for the 4-vector field $v_4(x)$. The form of X in equation 6.1 and det(X) in equation 6.27 when substituting in equation 9.5 become:

$$
\mathcal{X} = \begin{pmatrix} |r|^2 + |r'|^2 & \bar{r}\bar{s} + \bar{r'}\bar{s'} & c \\ sr + s'r' & |s|^2 + |s'|^2 & \bar{b} \\ \bar{c} & b & n \end{pmatrix} \in \mathbf{h}_3 \mathbb{O} \tag{9.7}
$$

$$
\det(\mathcal{X}) = \det(\theta_X^1 \theta_X^{1\dot{+}} + \phi_X^1 \phi_X^{1\dot{+}})n + 2(\theta_X^1 \theta_X^{1\dot{+}} + \phi_X^1 \phi_X^{1\dot{+}}) \cdot (\theta^1 \theta^{1\dot{+}})
$$
(9.8)

Here the first part of the expression for $\det(\mathcal{X})$ contains quintic terms, which now do not cancel in general as they did in equation 9.4, while the second part contains further quartic terms. This expression hence represents an *inhomogeneous* polynomial form, and hence deviates from the form of $L(\mathbf{v}) = 1$ in equation 2.9 of section 2.1 on incorporating a further higher-dimensional dissolving of specific components, such as those of X above. The potential physical consequences of such a mathematical possibility in relation to the original homogeneous form of $L(v_{27}) = 1$ requires further clarification. While such inhomogeneous expressions may be explored upon examination they appear indeed inconsistent with the underlying conceptual basis employed in deriving equation 2.9, as relating to infinitesimal intervals of temporal flow δs . On the other hand expressions such as equation 9.8 may represent an intermediate step towards the derivation of a higher-dimensional homogeneous form, such as a purely quintic expression for $L(v) = 1$, as will be proposed hypothetically in section 9.3 in the light also of the physical motivation described below.

The need to generalise from equation 9.1 motivated the introduction of a combination of spinors, θ_X^1 and ϕ_X^1 , in equation 9.5. This latter expression $X = \theta_X^1 \theta_X^{1\bar{1}} + \phi_X^1 \phi_X^{1\bar{1}}$, with θ_X^1 and ϕ_X^1 having identical transformation properties under the external and internal symmetry actions, suggests that for example the first generation u-quark and the second generation *c*-quark states might be accommodated in the θ_X^1 and ϕ_X^1 components respectively.

As described in the previous section the SU(2)^{2,3} \subset SL(2, $\mathbb{C})^{2,3}$ actions indicated in equation 8.31 transform the octonion components $(a) \leftrightarrow (b, c)$ in a seemingly asymmetric way. Hence, with $a = sr$ from equation 9.3, a combination of $\dot{\Sigma}^{(2)\pm}$ and $\dot{\Sigma}^{(3)\pm}$, of equations 8.37 and 8.68, appear to be needed in order to transform any of the four Weyl spinors located in the components of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ into the doublet partner located within the corresponding components of $\theta_X^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $\frac{\bar{r}}{s}$.

Together the observations of the above two paragraphs suggest the possibility of a Cabibbo-like mixing between the first two generations of quarks. Empirically the gauge action $SU(2)_L$ mixes the quark states $(u) \leftrightarrow (d \cos \theta_c + s \sin \theta_c)$, where s denotes the strange quark and θ_c is the Cabibbo angle which may be generalised to the full CKM matrix for three generations, as described towards the end of section 7.2. In the Standard Model the coupling between the first and third quark generations is very small. In the present theory quark exchanges via the \tilde{W}^{\pm} bosons associated with the $SU(2)^{2,3}$ symmetry may open up a full set of possible states associated for example with

 $X = \theta_X^1 \theta_X^{1\bar{1}} + \phi_X^1 \phi_X^{1\bar{1}} + \psi_X^1 \psi_X^{1\bar{1}}$, further augmenting equation 9.5 and lifting the degeneracy to a complete set of three generations interacting via a CKM-like mixing, although a similar degeneracy will also need to be identified relating to the θ^1 components. In principle this offers a possible means of accommodating three generations of fermions into the theory which may not relate directly to the existence of the three types of $SL(2,\mathbb{O})$ embedded within $SL(3,\mathbb{O})$ as described in equations 6.32–6.35.

The full study of these phenomena, as discussed in the previous subsection, will require the identification of each of the physical mass states, as observed in the laboratory, in relation to the components of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ and $\theta^1_X = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $\binom{\bar{r}}{s}$ for example. Around equation 8.76 it was proposed that fermion masses will relate to the degree of coupling with the scalar magnitude $|v_4| = h$, and in particular the 'vacuum value' of h in the projection of $v_4 \in TM_4$, by analogy with Higgs phenomena in the Standard Model. Since terms containing both the $a \in \mathbb{O}$ and the $v^0 = h$ components of $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ do not appear in $\det(\mathcal{X})$ in equation 8.76 an explicit higher-dimensional form of $L(\mathbf{v}) = 1$, such as introduced in the following section, may be needed for further study of possible mass terms. On the other hand composition with the scalar field $n(x)$, such as for the |a| term in equation 8.76, might also provide a source of fermion mass terms.

With the possible generalisation of equation 9.5 and the above weak interactions in mind it is also necessary to determine the internal $SU(3)_c \times U(1)_Q$ symmetry transformations of the θ_X^1 components, with similar transformations implied for ϕ_X^1 . The $SO(8) \subset SL(3, \mathbb{O})$ subgroup can be generated by the composition of 3×3 matrices $\mathcal{M}^{(1)}$ of type 1 in the form of equation 6.49 based on the 2 \times 2 matrices $M = \begin{pmatrix} q & 0 \\ 0 & \bar{q} \end{pmatrix}$ $\begin{pmatrix} q & 0 \\ 0 & \overline{q} \end{pmatrix}$. Acting via the conjugation $X \to M X M^{\dagger}$ on X in equation 9.3 the component $a \in \mathbb{Q}$ transforms under the vector representation of SO(8) while the θ_X^1 components $s \in \mathbb{O}$ and $r \in \mathbb{O}$ transform individually via the spinor and dual spinor representations of SO(8) as can be seen via equation 9.2.

These three 8-dimensional representations are mutually related through triality maps described in the opening of section 6.1 and around equation 6.50 – the triality structure in the present mathematical context is also described in more detail in ([38] pp.77–80 and 120–126). Further, elements of the $G_2 \subset SO(8)$ octonion automorphism subgroup transform the vector, spinor and dual spinor, here represented by the octonions a, s and r respectively, in precisely the same way via symmetric, left and right multiplication by the same sequence of octonions. This property of termed 'strong triality' in $([38] \text{ p.123}).$

Hence under the colour gauge symmetry $SU(3)_c \subset G_2 \subset SO(8)$ each of the octonion components of $\theta_X^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ \overline{s}) transform in the same way as the component a, and hence also in the same way as the octonion components of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$. This means that the four Weyl spinors, obtained from the reduction of θ_X^1 under the external $SL(2,\mathbb{C})^1$ action, transform as a leptonic singlet and quark triplet under the internal $SU(3)_c$, just as is the case for θ^1 as summarised in equation 8.19. There then remains the question of how the $U(1)_Q$ charges for the θ_X^1 Weyl spinors compare to those for the θ^1 Weyl spinors deduced for equation 8.26.

The electromagnetic U(1)_Q generator \dot{S}_l^1 is contained in the group SO(7) ⊂ $SO(8) \subset SL(2, \mathbb{O})^1$, but unlike the $SU(3)_c$ generators $\{\dot{A}_q, \dot{G}_l\}$ it is not contained within the subgroup $G_2 \subset SO(7)$. Although $\dot{S}_l^1 = \dot{S}_l^1$ the simpler unnested single group action S_l^1 of equation 6.43 is not used here since it is not constructed as a

'compatible' action in the sense of equation 6.23 or 9.2. We hence employ S_l^1 as a member of the preferred basis incorporated into the E_6 Lie algebra composition as discussed following equation 6.43.

The group action of $S_l^1(\alpha)$ on the components of $X = \binom{p}{a}$ and $\theta_X^1 = \binom{\bar{r}}{s}$ $\binom{\bar{r}}{s}$ may be determined from table 6.2 and equation 6.39 together with table 6.1 and equation 6.24 as the type 1 nested compositions:

$$
X \rightarrow R_{\mathbf{\mathcal{U}},i}^{1}(\alpha) \circ R_{\mathbf{\mathcal{Y}},j}^{1}(\alpha) \circ R_{\mathbf{\mathcal{U}},k}^{1}(\alpha) X
$$
\n
$$
\begin{pmatrix} p & \bar{a} \\ a & m \end{pmatrix} \rightarrow M_{\mathbf{\mathcal{U}},i2}(M_{\mathbf{\mathcal{Y}},i1}(M_{\mathbf{\mathcal{Y}},j2}(M_{\mathbf{\mathcal{Y}},j1}(M_{\mathbf{\mathcal{U}},k2}(M_{\mathbf{\mathcal{U}},k1}\left(\begin{matrix} p & \bar{a} \\ a & m \end{matrix}\right)M_{\mathbf{\mathcal{U}},k1}^{\dagger})M_{\mathbf{\mathcal{U}},k2}^{\dagger})M_{\mathbf{\mathcal{Y}},j1}^{\dagger})M_{\mathbf{\mathcal{Y}},j2}^{\dagger})M_{\mathbf{\mathcal{Y}},i1}^{\dagger})M_{\mathbf{\mathcal{Y}},j2}^{\dagger})M_{\mathbf{\mathcal{Y}},i1}^{\dagger})M_{\mathbf{\mathcal{Y}},i2}^{\dagger})M_{\mathbf{\mathcal{Y}},i2}^{\dagger})M_{\mathbf{\mathcal{Y}},i2}^{\dagger}M_{\mathbf{\mathcal{Y}},i2}^{\dagger}(M_{\mathbf{\mathcal{Y}},j1}(M_{\mathbf{\mathcal{U}},k2}(M_{\mathbf{\mathcal{Y}},k1}\left(\begin{matrix} \bar{r} \\ s \end{matrix}\right))))
$$
\n
$$
\begin{pmatrix} \bar{r} \\ s \end{pmatrix} \rightarrow M_{\mathbf{\mathcal{Y}},i2}(M_{\mathbf{\mathcal{Y}},i1}(M_{\mathbf{\mathcal{Y}},j2}(M_{\mathbf{\mathcal{Y}},j1}(M_{\mathbf{\mathcal{U}},k2}(M_{\mathbf{\mathcal{U}},k1}\left(\begin{matrix} \bar{r} \\ s \end{matrix}\right))))))
$$
\n
$$
(9.9)
$$

where the expression for the $\theta_{\rm x}^1$ transformation is a consequence of equation 9.1 or 9.5 together with the compatibility of the $S_l^1(\alpha)$ action as defined in equation 6.23 or 9.2. The following notation for the 6 nested actions, with factors of ± 1 accumulated into the initial negative signs, is introduced to simplify subsequent equations:

$$
N_6(\equiv -(i\ell \cos \frac{\alpha}{2} + i \sin \frac{\alpha}{2})(i\ell((j\ell \cos \frac{\alpha}{2} + j \sin \frac{\alpha}{2})(j\ell((k\ell \cos \frac{\alpha}{2} + k \sin \frac{\alpha}{2})(k\ell \cos \frac{\alpha}{2} + k \sin \frac{\alpha}{2}))
$$

$$
N_6^{\dagger} \equiv -k\ell((k\ell \cos \frac{\alpha}{2} + k \sin \frac{\alpha}{2}))j\ell((j\ell \cos \frac{\alpha}{2} + j \sin \frac{\alpha}{2}))j\ell((l\ell \cos \frac{\alpha}{2} + i \sin \frac{\alpha}{2}))
$$

From equations 9.9 the three octonion components a, s and r then transform under $S_l^1(\alpha)$ in the manner (as may be compared with equation 6.50):

$$
a \rightarrow N_6(a)N_6^{\dagger}
$$

\n
$$
s \rightarrow N_6(s))))
$$

\n
$$
\bar{r} \rightarrow N_6(\bar{r}))))
$$

\nhence $r \rightarrow (((((r)N_6^{\dagger})))$ (9.10)

While here we have symmetric, left and right multiplication on a, s and r respectively by the same sequence of octonions, as described by ' N_6 (' and ') N_6^{\dagger} ', these three actions are not mutually related by triality since they do not describe the same transformation on \mathbb{O} . Indeed since the action $S_l^1(\alpha)$ is not part of the $G_2 \subset SO(8)$ subgroup it does not exhibit the property of 'strong triality', but rather participates in the SO(8) triality structure collectively when further generators are considered.

Taken at face value equations 9.10 imply that the electric charge identified with \dot{S}_l^1 for the \bar{r}, s components here hence differs from that for the a component. This is an undesirable feature which means that the $\binom{0}{2}$ $_{2/3}^{0}$ charge structure observed for the component parts of a under \dot{S}_l^1 in equations 8.27 and 8.28, as sought for ν -lepton and u-quark fermion states, has apparently been lost for the set of $SL(2,\mathbb{C})^1$ Weyl

spinors in $\theta_X^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ \bar{s}). In fact, as expected from the compatibility of the S_l^1 group action, the U(1)_Q transformations of the spinor $\theta_{X}^{1} = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $\binom{\bar{r}}{s}$ are identical to those of the spinor $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ and hence both spinors possess the same \dot{S}_l^1 charge values of 1 and $\frac{1}{3}$, as described for equation 8.26, which have been associated with the e-lepton and d-quark states. Indeed, as implied in the discussion around equation 9.2, the components of θ_{X}^{1} transform in precisely the same way as those of θ^1 under the action of the subgroup $\mathrm{SL}(2,\mathbb{C})^1 \times \mathrm{SU}(3)_{c} \times \mathrm{U}(1)_{Q} \subset \mathrm{SL}(2,\mathbb{O})^1$ owing to the compatibility requirement of all $M \in SL(2,\mathbb{O})$ group transformations.

A possible solution would be to maintain the same $S_l^1(\alpha)$ action on X while redefining the transformation properties of θ_X^1 under the $\mathrm{U}(1)_Q$ subgroup. That is, with $X \to \theta_X^1 \theta_X^{1\dagger}$ in equation 9.1 or 9.5, on expanding the 10-dimensional space for X to the 16-dimensional space for θ_x^1 there is a degree of redundancy in the transformation properties of θ_X^1 under $U(1)_Q$ provided X transforms in the same way. While equation 9.9 represents the simplest assumption for the action of $S_l^1(\alpha)$ on the components of θ_x^1 , based on the notion of compatibility in equation 9.2, in principle there may be further choices such as:

$$
X \rightarrow N_6 \left(\mathbf{1}_2 X \mathbf{1}_2 \right) N_6^{\dagger}
$$

= $N_6 \left(\mathbf{1}_2 \theta_X^1 \theta_X^{1 \dagger} \mathbf{1}_2 \right) N_6^{\dagger}$
with $\theta_X^1 \rightarrow N_6 \left(\mathbf{1}_2 \theta_X^1 \right) N_6^{\dagger}$ (9.11)

rather than $\theta_X^1 \to N_6(1_2 \theta_X^1)$, although with care needed to take into account the nonassociative properties of octonion composition. On employing equation 9.11 in place of equation 9.10 the action of the U(1)_Q symmetry $S_l^1(\alpha)$ on $a, s, r \in \mathbb{O}$ would be expressed uniformly as:

$$
a \rightarrow N_6(a)N_6^{\dagger} \ns \rightarrow N_6(s)N_6^{\dagger} \nr \rightarrow N_6(r)N_6^{\dagger}
$$
\n(9.12)

Hence in this case the $\binom{0}{2}$ $2/3$ charge structure of a under \dot{S}_l^1 would also apply to the $r, s \in \mathbb{O}$ components of θ_x^1 . These $U(1)_Q$ charges for θ_x^1 are here *contrived* by inserting further nested $N_6^{(\dagger)}$ actions for $\theta_X^{(1)}$ in the appropriate places for equation 9.11. However the introduction of the 16 real component object $\theta_{X}^{1} = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $\binom{\bar{r}}{s}$ itself, obtained from the 10 real component object X in equation 9.1, is contrived and defined in order to construct a possible set of doublet partners for the $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ components under the action of \tilde{W}^{\pm} charge raising and lowering operators. With physical states associated with definite representations under the Lorentz symmetry of 4-dimensional spacetime M_4 , such as the above Weyl spinors within θ^1 and θ^1 _x, the purpose here is to demonstrate the mathematical possibility of recovering weak interactions between fermions, such as those mediated via W^{\pm} gauge bosons, from within the present theory. A mathematical justification for transformations such as those of equation 9.12 might ultimately be sought within a natural higher-dimensional homogeneous form $L(v) = 1$.

Hence $\theta_X^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $\binom{\bar{r}}{s}$ provisionally describes a possible mathematical construction which possesses the appropriate transformation properties under the external symmetry $SL(2,\mathbb{C})^1$ and internal symmetry $SU(3)_c \times U(1)_Q$ to represent the neutral ν -leptons

and charge- $\frac{2}{3}$ u-quarks. These latter states are related to the charge-1 e-leptons and charge- $\frac{1}{3}$ d-quarks of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ via \tilde{W}^{\pm} interactions. The relation between the fermion states and the unit charge raising and lowering action of the \tilde{W}^{\pm} oriented with respect to the θ_X^1 components serves to mutually motivate and aid the determination of both the Weyl spinor states and the internal gauge symmetry. From this point of view, with $X \in h_2\mathbb{O}$ composed in the form of equation 9.5, the W^{\pm} and \tilde{Z}^0 may derive from a weak $SU(2) \times U(1)$ action which is less directly related to the $SU(2)^{2,3} \times U(1)^{2,3} \subset E_6$ subgroups than suggested in the previous section.

To conclude the above discussion, while equation 9.5 describes a possible way to include the required further Weyl spinor states there are several questions which remain to be resolved – these include the means by which equation 9.8 might be incorporated into a higher-dimensional *homogeneous* form of $L(\mathbf{v}) = 1$ with a larger symmetry group incorporating an appropriate electroweak $SU(2)_L \times U(1)_Y$ subgroup action and the means by which the electromagnetic charges for the ν -lepton and ν quark states may be retained from the \dot{S}^1_l action on the components of the original $a \in \mathbb{O}$ component of h₃ \mathbb{O} . Further, while here we are working at the level of the basic group and representation structure, a leading question in the full theory will be to understand the nature of physical particle states in general, and in particular for the gauge bosons, three generations of fermions and also a Higgs state as empirically observed.

The above identification of fermions by opening up the 10-dimensional vector $X \in h_2\mathbb{O}$ according to equation 9.5 can similarly be applied to the 4-dimensional Lorentz vector $h_2 \in h_2\mathbb{C}$, for the subspace $h_2\mathbb{C} \subset h_2\mathbb{O}$, according to the decomposition of equation 7.32. This latter Weyl spinor substructure of the vector $h_2 = \chi \chi^{\dagger} + \phi \phi^{\dagger}$ in terms of the *spinors* χ , ϕ has some analogy with composite Higgs and technicolor models in which fermion states are combined in *scalar* condensates in the vacuum, hence replacing the fundamental scalar Higgs of the Standard Model, as reviewed in subsection 8.3.3. Here opening up the $h_2 \in h_2 \mathbb{C}$ components to form spinors in this way incorporates the $a_{1,l}$ components of equations 8.27 and 8.28, leaving the set of $SL(2,\mathbb{C})$ ¹ Lorentz scalars in $a(6)$ which transform under the internal $SU(3)_c \times U(1)_Q$ symmetry as a colour triplet of u-quarks. In principle each of the three scalars $a_{\underline{i},i}$, $a_{\underline{i},j}$ and $a_{\underline{u},k}$ of equation 8.28 might be composed in terms of a suitable scalar product of Weyl spinors of the form $\chi^{\dagger}\chi$ with the aim of describing the fermion nature of u-quarks under the external symmetry.

These spinor decompositions involve $\mathbb C$ or $\mathbb H$ subalgebras of $a \in \mathbb O$. Hence, in comparison with equation 9.11, an extra intermediate factor of the form $e^{-l\beta}e^{l\beta}$ might be inserted for the decomposed vector $h_2 \to \chi \chi^{\dagger}$ in augmenting the $S^1_l(\alpha)$ action, to ensure the charge neutrality of the candidate neutrino states, avoiding any complication due to the non-associative nature of the octonions. With the possibility of a similar insertion for the u-quark states in principle the $U(1)_Q$ charges under S_l^1 of $\binom{0}{2}$ $\binom{0}{2/3}$, as originally found for the $\binom{a_{1,l}}{a(6)}$ components in equation 8.28, might be maintained under the spinor decomposition of these components which might hence indeed be associated with $\binom{\nu}{n}$ u_{μ}^{ν}) fermion states. Again, while such a structure might be mathematically contrived as a proof of principle, ultimately the aim will be to account for the external and internal symmetry properties of all Standard Model fermion states in a natural manner in the components of a higher-dimensional homogeneous form of $L(\boldsymbol{v})=1.$

In keeping the $\binom{\nu}{u}$ $\binom{\nu}{u}$ particle type interpretations aligned with the $\binom{a_{1,l}}{a(6)}$ components in this way, based on the spinor decomposition of $h_2 \in h_2\mathbb{C}$ (unlike the case for the collective decomposition $X \to \theta_X^1 \theta_X^{1^{\dagger}} \in h_2 \mathbb{O}$ as originally considered in equation 9.1), also suggests that exchanges with the corresponding doublet partners $\binom{e}{d}$ $\binom{e}{d}$ in the components of θ^1 might be mediated by \tilde{W}^{\pm} states closely associated with the $SU(2)^{2,3} \subset E_6$ actions as described in section 8.3 for the mock electroweak theory. Based on the $\{1, l\}$ base units these $SU(2)^{2,3}$ actions preserve the 4-way decomposition of octonion components as listed in equation 8.30 for the transformations between the $a, b, c \in \mathbb{O}$ components of h₃ \mathbb{O} of the kind described in equations 8.29 and 8.31. This is consistent with an electroweak $SU(2)$ gauge symmetry action on independent lepton and quark doublets as accommodated respectively within the $\{1, l\}$ and $(\{\mathbf{\mathit{u}},\mathit{i}\},\{\mathit{\mathit{j}},\mathit{j}\},\{\mathit{kl},\mathit{k}\})$ components of both a and $\theta^1 = \begin{pmatrix} c \\ \mathit{b} \end{pmatrix}$.

In particular the neutrino state is associated with the $a_{1,l}$ components, which also form part of the 4-vector $h_2 \in h_2 \mathbb{C}$ as projected onto the external spacetime TM_4 . In the present theory the scalar degree of freedom $|h_2| = \sqrt{\det(h_2)}$, or an alternative scalar combination of the spinor components χ, ϕ in the implicit substructure of h_2 described above, will provide a candidate for the origin of the observed Higgs particle as described in subsection 8.3.3. This apparent inconsistency with the degrees of freedom of $h_2 \in h_2 \mathbb{C}$ seemingly required to play a double role as the correlate of both the neutrino and the Higgs will be resolved in the following section.

The approach of the present theory is to gently coax the known properties of the Standard Model out of the symmetry breaking structures of forms of $L(v) = 1$ over the base manifold M_4 , with an awareness of the known empirical features while being conscious of not contriving them ultimately for the complete theory. However the possibility of contriving an augmented structure based on the components of $a \in h_3\mathbb{O}$ under the E_6 action transforming as a *v*-lepton and *u*-quark under $SL(2,\mathbb{C})^1 \times SU(3)_c \times U(1)_Q \subset$ E_6 , as described in this section, and with the further possibility of incorporating a second and third generation through additional spinors such as described for equation 9.5, is at least consistent with the possibility that a higher-dimensional form of $L(\mathbf{v}) = 1$, for example with an E_8 symmetry as will be considered in section 9.3, might naturally contain these structures. Similarly the empirical properties of left and right-handed spinors, as we recap below, will contribute to the motivation for the study of an E_7 symmetry of a higher-dimensional form of time in the following section.

In the Standard Model Lagrangian each fermion kinetic term, such as equation 7.39, or interaction term, such as equation 7.77, contains either left or right-handed fermion states, while the Yukawa or Dirac mass terms combine opposite chiralities, as for example in equations 7.69 or 7.74. In all cases the operators $P_L = \frac{1}{2}(1 - \gamma^5)$ or $P_R=\frac{1}{2}$ $\frac{1}{2}(1+\gamma^5)$, of equations 7.11 and 7.12, may be used to project out the respective left or right-handed chirality states from a 4-component Dirac spinor.

The factors of $P_L = \frac{1}{2}$ $\frac{1}{2}(1-\gamma^5)$ which appear in all fermion terms involving the $SU(2)_L$ gauge symmetry, such as in equations 7.77 and 7.78, are placed in the Lagrangian by hand in order to replicate the parity violating phenomena observed empirically for the weak interaction. This parity violation is maximal for the case of interactions via W^{\pm} gauge bosons but non-maximal for Z^0 interactions, which is associated with a linear combination of $SU(2)_L$ and $U(1)_Y$ generators, equation 7.49, as described in section 7.2. It is this asymmetry in the chiral structure, with different weak isospin transformations for left- and right-handed fields, implying that no fermion state transforms under the complex conjugate representation of that of any other fermion, which necessitates the introduction of Yukawa couplings to the Higgs field, as for equation 7.69, in order to include fermion mass terms in the Lagrangian.

In the present theory we have described how the components of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix}$ form the set of four left-handed Weyl spinors of equation 8.13 under the external Lorentz symmetry $SL(2, \mathbb{C})^1$. In this section a similar decomposition has also been identified for the components of $X = \begin{pmatrix} p & \bar{a} \\ a & m \end{pmatrix}$, for example via the spinor $\theta_X^1 = \begin{pmatrix} \bar{r} \\ s \end{pmatrix}$ $(\bar{r}_s) \in \mathbb{O}^2$ as introduced in equation 9.1. Hence the projection operator $P_L = \frac{1}{2}(1-\gamma^5)$ has not been introduced since only *left*-handed Weyl spinors under $SL(2,\mathbb{C})^1$ have so far been considered. There then remains the question of how right-handed Weyl spinor counterparts may identified within this framework, and related to the above left-handed components in a single 4-component Dirac spinor $\psi(x)$ to describe, for example, a physical electron state.

Here we began with the cubic form $\det(\mathcal{X}) = 1$, with $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$, as a 27dimensional expression of temporal flow $L(\mathbf{v}_{27}) = 1$. Determinant preserving E₆ transformations were then considered on this space, with for example $\mathcal{X} \to \mathcal{M}\mathcal{X}\mathcal{M}^{\dagger}$ for the 3×3 matrices of equations 6.29 and 6.32 incorporating any of the 2×2 matrices of category 1 or 2 in table 6.1, for the unnested case. With $\mathcal M$ representing the matrix M with each entry replaced by its octonion conjugate, as described in equation 6.7, transformations of the form:

$$
\mathcal{X} \to \overline{\mathcal{M}} \, \mathcal{X} \, \overline{\mathcal{M}}^{\dagger} \tag{9.13}
$$

clearly also leave the value of $\det(\mathcal{X})$ invariant. With E_6 a symmetry of the form of time $L(v_{27}) = 1$ the two possible representations 27 and $\overline{27}$ are equally valid while only one of them has been used so far.

Correspondingly the set of six actions with $\overline{\mathcal{M}}$, for $\mathcal{M} \in SL(2,\mathbb{C})^1$, provides an alternative choice for the type 1 Lorentz transformations acting on the vector components $v^4 \in h_2 \mathbb{C} \subset h_3 \mathbb{O}$, with $h_2 \to \overline{S} h_2 \overline{S}^{\dagger}$ in place of equation 7.31. In turn these transformations are represented on θ^1 as a set of four *right*-handed Weyl spinors. The group $SL(2,\mathbb{C})$, as for the full group E_6 , has complex representations, and the actions of $S \in SL(2, \mathbb{C})^1$ and \overline{S} describe distinct sets of transformations of θ^1 , as explained in section 7.1. This means that the left and right-handed transformations are not equivalent to each other but are instead mutually related as described in equations 7.26 and 7.27. In the context of the present theory both $S \in SL(2,\mathbb{C})^1$ and \overline{S} act as symmetry transformations leaving the form $L(v_{27}) = 1$ invariant, and hence both the left and right Weyl spinor compositions of θ^1 should in principle play a role.

The apparent asymmetry in the choice of the E_6 27 or $\overline{27}$ representation to express $L(v_{27}) = 1$, with a corresponding choice of left or right-handed representations of $SL(2,\mathbb{C})^1 \subset E_6$, the need to clearly identify both left- and right-handed fermions ψ_L and ψ_R , in particular with reference to an $SU(2)_L$ gauge symmetry, and the existence of a homogeneous quartic form as a candidate for a higher-dimensional temporal flow in the form $L(v_{56}) = 1$ all point to consideration of the group E_7 as a symmetry of time, as will be described in the following section.

9.2 E⁷ Symmetry and the Freudenthal Triple System

The introduction of further dimensions in the previous section and the observation of the quartic expression of equation 9.4, with the extension to equation 9.8 including quintic terms also, suggests the possibility of a higher-dimensional expression for the flow of time generalising beyond the cubic form $L(v_{27}) = 1$ described in chapter 6. A higher-dimensional *homogeneous* polynomial form is desired, in conformity with the derivation of equation 2.9 in chapter 2. While the determinant preserving symmetry of the space $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ describes the lowest-dimensional non-trivial representation of E_6 the smallest non-trivial representation of the exceptional Lie group E_7 is 56-dimensional and may be constructed in terms the elements x of the Freudenthal triple system $F(h_3\mathbb{O})$ ([59, 60, 61], [1] p.48).

In the above references and related publications these mathematical structures are applied in two very different contexts – namely the classification of black hole solutions in string theory and the entanglement of qubits in quantum information theory – with a correspondence between these applications identified through the mathematical forms they share. Neither application is relevant for the present discussion. While much of the literature describes a more general algebraic framework or particular cases involving for example the 'split octonions' \mathbb{O}_s or takes an underlying field of integers \mathbb{Z} , here we are interested in the octonion \mathbb{O} case over an underlying field of real numbers R as we summarise in the following.

In order to describe the Freudenthal triple system $F(h_3\mathbb{O})$ it is useful to first introduce further definitions regarding the exceptional Jordan algebra $h_3\mathbb{O}$ itself. The structure group $Str(h_3\mathbb{O})$ leaves the cubic norm $\det(\mathcal{X})$ of equations 6.27 and 6.28 invariant up to a real scalar factor, that is:

$$
Str(h_3 \mathbb{O}) = \{ g \in GL(h_3 \mathbb{O}) \mid \det(\sigma_g(\mathcal{X})) = \lambda(g) \det(\mathcal{X}), \ \forall \mathcal{X} \in h_3 \mathbb{O} \}
$$
(9.14)

with $\lambda \in \mathbb{R}$ depending only on g. The norm preserving subgroup with $\lambda = 1$ is identified as the reduced structure group $Str_0(h_3\mathbb{O}) \equiv SL(3,\mathbb{O})$. This latter symmetry corresponds to the 27-dimensional representation of $E_{6(-26)}$ as described in detail in chapter 6.

A trace bilinear map may be defined for any elements $\mathcal{X}, \mathcal{Y} \in \text{h}_3\mathbb{O}$ of the Jordan algebra, mapping $h_3 \mathbb{O} \times h_3 \mathbb{O} \to \mathbb{R}$ with:

$$
(\mathcal{X}, \mathcal{Y}) = \text{tr}(\mathcal{X} \circ \mathcal{Y}) \tag{9.15}
$$

where on the right-hand side the \circ denotes the Jordan algebra product of equation 6.2. An *adjoint* s^* for any transformation $s(\mathcal{X})$ with $s \in \mathbb{E}_6$ may be defined with respect to the above trace bilinear form such that:

$$
(s(\mathcal{X}), \mathcal{Y}) = (\mathcal{X}, s^*(\mathcal{Y})) \qquad \forall \mathcal{X}, \mathcal{Y} \in \mathbf{h}_3 \mathbb{O}
$$
\n
$$
(9.16)
$$

Along with the Jordan product there is a second natural composition for the elements of $h_3\mathbb{O}$ which is called the *Freudenthal product* and may be defined by:

$$
\mathcal{X} \wedge \mathcal{Y} = \mathcal{X} \circ \mathcal{Y} - \frac{1}{2} \big(tr(\mathcal{X}) \mathcal{Y} + tr(\mathcal{Y}) \mathcal{X} \big) + \frac{1}{2} \big(tr(\mathcal{X}) tr(\mathcal{Y}) - tr(\mathcal{X} \circ \mathcal{Y}) \big) \mathbf{1}_3 \in \mathbf{h}_3 \mathbb{O} \quad (9.17)
$$

For any $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ a quadratic adjoint map $\mathrm{h}_3\mathbb{O} \to \mathrm{h}_3\mathbb{O}$ can be defined in terms of the Freudenthal product as:

$$
\mathcal{X}^{\sharp} = \mathcal{X} \wedge \mathcal{X} \tag{9.18}
$$

or explicitly:
$$
\mathcal{X}^{\sharp} = \mathcal{X}^2 - \text{tr}(\mathcal{X})\mathcal{X} + \frac{1}{2}[\text{tr}(\mathcal{X})^2 - \text{tr}(\mathcal{X}^2)]\mathbf{1}_3
$$
 (9.19)

This 'sharp' operation satisfies the relations $(\mathcal{X}^{\sharp})^{\sharp} = \det(\mathcal{X})\mathcal{X}$ and $\mathcal{X} \circ \mathcal{X}^{\sharp} = \det(\mathcal{X})\mathbf{1}_3$. The linearisation of the quadratic adjoint is written as:

$$
\mathcal{X} \times \mathcal{Y} = (\mathcal{X} + \mathcal{Y})^{\sharp} - \mathcal{X}^{\sharp} - \mathcal{Y}^{\sharp}
$$
\n(9.20)

$$
\equiv 2\mathcal{X} \wedge \mathcal{Y} \tag{9.21}
$$

For the elements of $h_3\mathbb{O}$ the quadratic adjoint is in fact the classical adjoint, that is the transposed cofactors of $X \in \mathfrak{h}_3\mathbb{O}$ which, for the components of X presented in equation 6.1 or 9.25 below, can be written explicitly as the 3×3 matrix:

$$
\mathcal{X}^{\sharp} = \begin{pmatrix} mn - |b|^2 & cb - n\bar{a} & \bar{a}\bar{b} - mc \\ \bar{b}\bar{c} - na & pn - |c|^2 & ac - p\bar{b} \\ ba - m\bar{c} & \bar{c}\bar{a} - pb & pm - |a|^2 \end{pmatrix} \in \mathbf{h}_3 \mathbb{O}
$$
(9.22)

The vector space $F(h_3\mathbb{O})$ has 56 real components and may be introduced according to Freudenthal's construction with the vector space composition (which may be compared with the further decomposition of equation 8.1):

$$
F(\mathbf{h}_3 \mathbb{O}) \cong \mathbf{h}_3 \mathbb{O} \oplus \mathbf{h}_3 \mathbb{O} \oplus \mathbb{R} \oplus \mathbb{R} \tag{9.23}
$$

Correspondingly elements $x \in F$, with $F = F(h_3 \mathbb{O})$, are generally written in the form of a ' 2×2 matrix' as:

$$
x = \begin{pmatrix} \alpha & \mathcal{X} \\ \mathcal{Y} & \beta \end{pmatrix}, \quad \text{with } \mathcal{X}, \mathcal{Y} \in \mathbf{h}_3 \mathbb{O}, \quad \alpha, \beta \in \mathbb{R} \tag{9.24}
$$

and
$$
\mathcal{X} = \begin{pmatrix} p & \bar{a} & c \\ a & m & \bar{b} \\ \bar{c} & b & n \end{pmatrix}, \qquad \mathcal{Y} = \begin{pmatrix} P & \bar{A} & C \\ A & M & \bar{B} \\ \bar{C} & B & N \end{pmatrix}
$$
(9.25)

here with the real P, M, N and octonion A, B, C components of $\mathcal Y$ distinguished from the lower case counterpart components of X . A non-degenerate bilinear antisymmetric quadratic form mapping $F \times F \to \mathbb{R}$ may be defined on this space which acts on $x = \begin{pmatrix} \alpha & \lambda \\ \nu & \beta \end{pmatrix}$ \mathcal{Y} β $\mathcal{y} = \begin{pmatrix} \gamma & \mathcal{W} \\ \mathcal{Z} & \delta \end{pmatrix}$ $Z \delta$ $\big) \in F$ as:

$$
\{x, y\} = \alpha \delta - \beta \gamma + (\mathcal{X}, \mathcal{Z}) - (\mathcal{Y}, \mathcal{W}) \tag{9.26}
$$

Of more significance for the present theory there is also a homogeneous quartic norm $q: F \to \mathbb{R}$ defined on the components of $x \in F$ as follows:

$$
q(x) = -2[\alpha\beta - (\mathcal{X}, \mathcal{Y})]^2 - 8[\alpha \det(\mathcal{X}) + \beta \det(\mathcal{Y}) - (\mathcal{X}^{\sharp}, \mathcal{Y}^{\sharp})]
$$
(9.27)

where all the necessary definitions contained within this expression are inherited from those for the Jordan algebra $h_3\mathbb{O}$ as described above. The quadratic and quartic forms of equations 9.26 and 9.27 may be used in turn to define a trilinear mapping of the space $F \times F \times F \to F$, which is the triple product by which the 'Freudenthal triple system' gains its name. When written out explicitly in terms of the real and octonion components of equations 9.24 and 9.25 there are a large number of quartic terms in $q(x)$. In fact, via equations 9.15 and 9.22, and cross-checking with ([60] equation 9.51), we have for equation 9.27:

$$
q(x) = -2 \left[\alpha \beta - pP - mM - nN - 2(\langle a, A \rangle + \langle b, B \rangle + \langle c, C \rangle) \right]^2
$$

\n
$$
-8 \left[\beta P M N + \alpha p m n - p P m M - p P n N - n N m M
$$

\n
$$
+ (p m - \beta N) |A|^2 + (P M - \alpha n) |a|^2
$$

\n
$$
+ (m n - \beta P) |B|^2 + (M N - \alpha p) |b|^2
$$

\n
$$
+ (n p - \beta M) |C|^2 + (N P - \alpha m) |c|^2
$$

\n
$$
+ 2 \beta \text{ Re}(\overline{A} \overline{B} \overline{C}) + 2 \alpha \text{ Re}(\overline{a} \overline{b} \overline{c})
$$

\n
$$
- |a|^2 |A|^2 - |b|^2 |B|^2 - |c|^2 |C|^2
$$

\n
$$
- (c b - n \overline{a}) (\overline{B} \overline{C} - N A) - (C B - N \overline{A}) (\overline{b} \overline{c} - n a)
$$

\n
$$
- (ac - p \overline{b}) (\overline{C} \overline{A} - P B) - (A C - P \overline{B}) (\overline{c} \overline{a} - p b)
$$

\n
$$
- (ba - m \overline{c}) (\overline{A} \overline{B} - M C) - (B A - M \overline{C}) (\overline{a} \overline{b} - mc) \qquad (9.28)
$$

where the inner product $\langle a, A \rangle = \frac{1}{2}$ $\frac{1}{2}(a\overline{A}+A\overline{a})$, which has the property $\langle a, A \rangle = \langle \overline{a}, \overline{A} \rangle$, was defined in equation 6.10. Equations 9.27 and 9.28 for the quartic form $q(x)$ are the analogue of equations 6.27 and 6.28 respectively for the cubic form $\det(\mathcal{X})$. Clearly there are many more terms for the above quartic from in equation 9.28 as an extension from the cubic form of equation 6.28.

The group Inv(F) of all invertible transformations σ in F preserving the above quartic norm with $q(\sigma(x)) = q(x)$, as well as the bilinear form of equation 9.26 with $\{\sigma(x), \sigma(y)\} = \{x, y\}$, is also denoted $Aut(F)$ since it in turn forms the automorphism group of the trilinear product defined for the Freudenthal triple system. This group is found to be the non-compact real form $E_{7(-25)}$ of the exceptional Lie group E_7 . Hence, in particular, under this symmetry group the invariance of the quartic form $q(x)$, as a homogeneous polynomial, describes a possible 56-dimensional form of temporal flow which may be denoted $L(v_{56}) = 1$. The possible physical implications of this form and the accompanying E_7 symmetry will be assessed in the remainder of this section and summarised in the following one.

The symmetry of the cubic form $L(v_{27}) = 1$, in the form of $\det(\mathcal{X})$ or $\det(\mathcal{Y})$, is contained within this structure as can be seen from equation 9.27. In fact the elements $\mathcal X$ and $\mathcal Y$, with 54 real components in total, may be considered to represent a 'complexification' of the space $h_3\mathbb{O}$, with both the 27-dimensional representation of E_6 and its complex conjugate contained within the E_7 action on $q(x)$. Including the actions of the subgroup $E_6 \subset E_7$ on the elements $x \to s(x) \in F$ the transformations of full symmetry $E_7 \equiv Inv(F)$ may be categorised in terms of four sets. With $s \in E_6$, $\lambda \in \mathbb{R}$ and $C, D \in \text{h}_3\mathbb{O}$ these are [59, 60, 61]:

$$
T(s): \begin{pmatrix} \alpha & \mathcal{X} \\ \mathcal{Y} & \beta \end{pmatrix} \rightarrow \begin{pmatrix} \alpha & s(\mathcal{X}) \\ s^{*-1}(\mathcal{Y}) & \beta \end{pmatrix}
$$
(9.29)

$$
\lambda : \begin{pmatrix} \alpha & \mathcal{X} \\ \mathcal{Y} & \beta \end{pmatrix} \rightarrow \begin{pmatrix} \lambda^{-1} \alpha & \lambda^{\frac{1}{3}} \mathcal{X} \\ \lambda^{-\frac{1}{3}} \mathcal{Y} & \lambda \beta \end{pmatrix}
$$
(9.30)

$$
\phi(C): \begin{pmatrix} \alpha & \mathcal{X} \\ \mathcal{Y} & \beta \end{pmatrix} \rightarrow \begin{pmatrix} \alpha + (\mathcal{Y}, C) + (\mathcal{X}, C^{\sharp}) + \beta \det(C) & \mathcal{X} + \beta C \\ \mathcal{Y} + \mathcal{X} \times C + \beta C^{\sharp} & \beta \end{pmatrix} \quad (9.31)
$$

$$
\psi(D): \begin{pmatrix} \alpha & \mathcal{X} \\ \mathcal{Y} & \beta \end{pmatrix} \rightarrow \begin{pmatrix} \alpha & \mathcal{X} + \mathcal{Y} \times D + \alpha D^{\sharp} \\ \mathcal{Y} + \alpha D & \beta + (\mathcal{X}, D) + (\mathcal{Y}, D^{\sharp}) + \alpha \det(D) \end{pmatrix}
$$
(9.32)

where s^* is the adjoint of $s \in E_6$ as defined in equation 9.16. The set of actions s^{*-1} in equation 9.29 is equivalent to the complex conjugate of the representation defined by the set of actions $s \in E_6$ on h₃ \mathbb{O} . Under the subgroup $E_{6(-26)} \subset E_{7(-25)}$ the space F decomposes into the reducible representation ([60] equations 9.45 and 9.46):

$$
56_{E_7} \rightarrow (27 + \overline{27} + 1 + 1)_{E_6} \tag{9.33}
$$

compatible with the structure of equation 9.23 (and can be compared with the further reduction under Spin⁺(1,9) in equation 8.3). The 78 actions of E_6 combined with the single dilation action λ of equation 9.30 applied to an h₃ $\mathbb{O} \subset F$ subspace together form the 79-dimensional group $Str(h_3\mathbb{O})$ as defined in equation 9.14. The 27 independent actions of $\phi(C)$ together with the further 27 for $\psi(D)$ in equations 9.31 and 9.32 further augment the E_6 symmetry to complete the full $(78 + 1 + 27 + 27) = 133$ -dimensional exceptional Lie group E7. (Building up the symmetry structure this way is analogous to augmenting the F_4 algebra by the D^B maps in equation 6.4 to complete the full E_6 symmetry.) In addition to the continuous actions of equations 9.29–9.32 a discrete symmetry $\tau := \phi(-1_3)\psi(1_3)\phi(-1_3)$ such that:

$$
\tau : \left(\begin{array}{cc} \alpha & \mathcal{X} \\ \mathcal{Y} & \beta \end{array} \right) \rightarrow \left(\begin{array}{cc} -\beta & -\mathcal{Y} \\ \mathcal{X} & \alpha \end{array} \right) \tag{9.34}
$$

with $\tau^2(x) = -x$, may also be defined. Since $\psi(C) = \tau \phi(-C) \tau^{-1}$ the set of actions ϕ and ψ are conjugate with respect to τ . Between equations 9.29 and 9.34 the further relationship $\tau T(s) = T(s^{*-1}) \tau$ is also found.

At the Lie algebra level the actions $\dot{s} \in L(E_6)$ may be divided into the 14 elements of $L(G_2)$, denoted D^G , and the action of the 64 tracefree octonion matrices x_0 , denoted D^S in equation 6.5. The latter set further divides into the 26 boosts with Hermitian x_0 and 38 rotations with anti-Hermitian x_0 . Such decompositions were also discussed in the opening three paragraphs of section 6.5. The subgroup G_2 itself may also be obtained through sequences of nested rotations as described in section 6.4. The

dual representation of $L(E_6)$ may be obtained by defining the action $\dot{s}'(\mathcal{X})$ for each $\dot{s} \in L(E_6)$ such that ([62] equation 4):

$$
(\dot{s}(\mathcal{X}), \mathcal{Y}) = -(\mathcal{X}, \dot{s}'(\mathcal{Y})) \qquad \forall \mathcal{X}, \mathcal{Y} \in h_3 \mathbb{O}
$$
\n(9.35)

which may be contrasted with equation 9.16 at the group level. For the $L(E_6)$ maps $\mathcal{X} \to x_0 \mathcal{X} + \mathcal{X} x_0^{\dagger}$ the dual transformations correspond to:

$$
x_0' = x_0 \qquad \text{for rotations} \tag{9.36}
$$

$$
x'_0 = -x_0 \qquad \text{for boosts} \tag{9.37}
$$

that is with $x'_0 = -x_0^{\dagger}$ in general ([62] equation 5). It also follows that $\dot{s}' = \dot{s}$ for the $L(G_2)$ actions derived from transverse rotations. Applied to the subalgebra $L(E_6)$ $L(E_7)$ acting on the elements of the Freudenthal triple system these 78 generators form the first of the four sets of E_7 actions at the Lie algebra level (corresponding to equations 9.29–9.32 at the group level) which may be listed as the infinitesimal transformations of $x = \begin{pmatrix} \alpha & \lambda \\ \nu & \beta \end{pmatrix}$ $\mathcal{Y} \beta$ $(62]$ section 2):

$$
T(\dot{s}): \qquad \left(\begin{array}{cc} 0 & \dot{s}(\mathcal{X}) \\ \dot{s}'(\mathcal{Y}) & 0 \end{array}\right) \tag{9.38}
$$

$$
\dot{\lambda} : \begin{pmatrix} -\dot{\lambda}\alpha & \frac{1}{3}\dot{\lambda}\mathcal{X} \\ -\frac{1}{3}\dot{\lambda}\mathcal{Y} & \dot{\lambda}\beta \end{pmatrix}
$$
 (9.39)

$$
\phi(\dot{C}) : \qquad \begin{pmatrix} (\mathcal{Y}, \dot{C}) & \beta \dot{C} \\ \mathcal{X} \times \dot{C} & 0 \end{pmatrix} \tag{9.40}
$$

$$
\psi(\dot{D}): \qquad \begin{pmatrix} 0 & \mathcal{Y} \times \dot{D} \\ \alpha \dot{D} & (\mathcal{X}, \dot{D}) \end{pmatrix} \tag{9.41}
$$

with $\dot{s} \in L(E_6)$, $\dot{\lambda} \in \mathbb{R}$ and $\dot{C}, \dot{D} \in h_3\mathbb{O}$. Higher-order terms such as $C^{\sharp} = C \wedge C$ and the cubic norm $\det(C)$ appear for the finite group actions of equations 9.31 and 9.32.

Having extended beyond the $L(E_6)$ subalgebra to the full $L(E_7)$ we next focus on the generators of the 4-dimensional spacetime Lorentz subgroup $SL(2,\mathbb{C})^1 \subset E_6 \subset$ E_7 of type 1 as studied in section 8.1. As for all E_6 transformations for the actions of the Lorentz subalgebra $sl(2,\mathbb{C})^1 \subset L(\mathbb{E}_6)$ the dual transformations \dot{s}' in equation 9.38 have identical rotation generators to \dot{s} while the boosts are reversed, by equations 9.36 and 9.37. As was described for equations 7.24 and 7.25 of section 7.1 reversing the sign of the boosts, there parametrised by b_a , is precisely the operation which interchanges between the L and R representations of $SL(2, \mathbb{C})$.

Hence while the components of θ_l in θ^1 within $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$, defined in equation 8.11, transform as a *left*-handed Weyl spinor under $SL(2,\mathbb{C})^1$ the corresponding components of $\theta_{\mathcal{L}} = \begin{pmatrix} C_1 + C_8 l \\ B_1 - B_8 l \end{pmatrix}$ B_1-B_8l) within the θ^1 component of $\mathcal{Y} \in h_3\mathbb{O}$, extracted from equation 9.25, transform as a *right*-handed Weyl spinor under the same $SL(2,\mathbb{C})^1 \subset$ $E_6 \subset E_7$ action. (The subscript ' \mathcal{L} ' on $\theta_{\mathcal{L}}$ denotes both the use of the imaginary unit

l and the identification of the 'leptonic' components of $\theta^1 = \begin{pmatrix} C \\ B \end{pmatrix}$ in \mathcal{Y} , as will be seen below. In general the superscript '1' is not appended to components such as θ_l and $\theta_{\mathcal{L}}$ since they are unambiguously extracted from 'type 1' θ^1 components, while a superscript is included for the 'type 2' or 'type 3' case as for θ_l^2 in equation 8.51 for example). Considered as an action of 2×2 matrices $S \in SL(2, \mathbb{C})^1$ on the 2-component Weyl spinors θ_l and $\theta_{\mathcal{L}}$, extracted from the corresponding θ^1 components of X and Y respectively, and using equation 7.27, the action of equation 9.29 may be summarised as:

$$
\left(\begin{array}{c}\n\theta_l \\
\theta_{\mathcal{L}}\n\end{array}\right) \rightarrow \left(\begin{array}{cc} S & 0 \\
0 & S^{\dagger^{-1}}\n\end{array}\right) \left(\begin{array}{c}\n\theta_l \\
\theta_{\mathcal{L}}\n\end{array}\right) \tag{9.42}
$$

This is precisely the Lorentz transformation of a 4-component Dirac spinor ψ as described in equations 7.15 and 7.29. Alternatively the above expression could be obtained directly at the group level from equation 9.29 using the definition of the adjoint s^{*} in equation 9.16 applied directly to the SL(2, $\mathbb{C})^1 \subset \mathbb{E}_6$ group transformations.

As explained in section 8.1 the components of θ^1 within $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ under the action of $SL(2,\mathbb{C})^1$ actually decompose into a set of four left-handed Weyl spinors $\{\theta_l, \theta_i, \theta_j, \theta_k\}$ as listed in equation 8.13. Hence equation 9.29 contains both the original representation of $SL(2,\mathbb{C})^1$ on X, which contains the set of four left-handed Weyl spinors in the θ^1 components, simultaneously with an equivalent of the complex conjugate representation on \mathcal{Y} , which hence contains a corresponding set of four right-handed Weyl spinors, which may be denoted $\{\theta_{\mathcal{L}}, \theta_{I}, \theta_{I}, \theta_{K}\} \subset \mathcal{Y}$. Correspondingly a set of four 4-component Dirac spinors have hence been identified with:

$$
\psi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix} = \begin{pmatrix} \theta_l \\ \theta_c \end{pmatrix}, \quad \begin{pmatrix} \theta_i \\ \theta_I \end{pmatrix}, \quad \begin{pmatrix} \theta_j \\ \theta_J \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} \theta_k \\ \theta_K \end{pmatrix} \tag{9.43}
$$

with
$$
\psi = \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix} \rightarrow \begin{pmatrix} S & 0 \\ 0 & S^{\dagger^{-1}} \end{pmatrix} \begin{pmatrix} \psi_L \\ \psi_R \end{pmatrix}
$$
 (9.44)

under $S \in SL(2,\mathbb{C})^1 \subset E_6 \subset E_7$ transformations in each case.

The above analysis applied to the θ^1 components of X and Y similarly applies for the left-handed $SL(2, \mathbb{C})^1$ Weyl spinors contained within θ_X^1 under the decomposition of equation 9.1 or 9.5. In this case a corresponding set of four right-handed spinors are found in the components of θ_Y^1 obtained in turn under a decomposition which may be denoted $Y = \theta_Y^1 \theta_Y^{1^{\dagger}}$ for the $h_2 \mathbb{O} \subset h_3 \mathbb{O}$ components of \mathcal{Y} . A similar observation applies for the alternative spinor decomposition of Y beginning with the h₂ $\mathbb{C} \subset h_2\mathbb{O}$ subspace as described towards the latter part of the previous section.

The internal $SU(3)_c \times U(1)_Q$ symmetry, described in section 8.2, is composed as a subgroup of E_6 purely out of the subset of rotations. Hence, by the discussion around equation 9.36 above, these actions are identical on the components of $\mathcal X$ and $\mathcal Y$ in equation 9.29. Hence in turn the $SU(3)_c$ action on the components of X, including upon the θ^1 components as detailed in table 8.7 and summarised together with the $U(1)_O$ action in equation 8.26, is identical for the corresponding components of \mathcal{Y} , and the corresponding $U(1)_Q$ charges for the respective subcomponents of equation 9.25 are also the same. Hence the ψ_L and ψ_R components carry matching $SU(3)_c \times U(1)_Q$ transformation properties for the set of four Dirac spinors in equation 9.43 (justifying the identification of both θ_l and $\theta_{\mathcal{L}}$ as leptonic components). Similarly the SU(2)^{2,3} \times $U(1)^{2,3} \subset E_6$ rotations, for the mock electroweak theory described in section 8.3, also act on the X and Y components of $x \in F(h_3\mathbb{O})$ in the same way.

While the total number of dimensions has been increased from 27 to 56 it remains the case that only a single set of 4 dimensions will describe the external spacetime. This can be chosen as an h₂ $\mathbb{C} \subset h_3 \mathbb{O}$ subset of components $v_4 \subset \mathcal{X}$, under an SL(2, C) ⊂ E₆ action, *or* as an h₂C ⊂ h₃^O subset of components $v_4 \subset \mathcal{Y}$, transforming under the complex conjugate representation, but not both. Here we choose $v_4 \equiv h_2 \in h_2 \mathbb{C}$ as embedded within the $Y = \begin{pmatrix} P & \bar{A} \\ A & M \end{pmatrix} \in h_2 \mathbb{O}$ components of $\mathcal Y$ in equation 9.25 to represent external spacetime, with Lorentz transformations hence described by:

$$
h_2 \to h_2' = S^{\dagger^{-1}} h_2 S^{-1} \tag{9.45}
$$

rather than equation 7.31, under the action of $S \in SL(2,\mathbb{C})^1 \subset E_6$. The complex subspace with base units $\{1, l\}$ still underlies both the $SL(2, \mathbb{C})^1$ subgroup and the subspace for the vectors $h_2 \in h_2 \mathbb{C}$. These h_2 components of $\mathcal Y$ will also now be taken to form the 'vector-Higgs' correlated with the phenomena of the Standard Model Higgs sector and Yukawa couplings, as was described for the original case of $L(v_{27}) = 1$ in subsection 8.3.3. Here for the case of $L(v_{56}) = 1$ this now implies that none of the 27 components of $\mathcal{X} \in \mathrm{h}_3\mathbb{O} \subset F(\mathrm{h}_3\mathbb{O})$ are identified with components of the external spacetime vectors $v_4 \in TM_4$.

In particular this means that in addition to the d-quark and charged lepton components of left-handed Weyl spinors in $\theta^1 \subset \mathcal{X}$, potentially both u-quark and neutral lepton left-handed Weyl spinors might be identified in the X components of $\mathcal X$ as described in the previous section. The $a \in \mathbb O$ component of $\mathcal X$ has the correct $(0, \frac{2}{3})$ $\frac{2}{3}$) charge structure to describe (*v*-lepton, *u*-quark) particle states, as seen in equations 8.27 and 8.28, and is now free to accommodate both states. However while the corresponding imaginary $A(6)$ components of $\mathcal Y$ also have an \dot{S}^1_l charge of $\frac{2}{3}$, the $A_{1,l} = (A_1 + A_8l)$ part of $A \in \mathbb{O}$ in \mathcal{Y} is *occupied* by the above components $h_2 \in h_2\mathbb{C}$, representing the vector-Higgs and external spacetime, as depicted in equation 9.46.

$$
\left(\begin{array}{c}\n\alpha \\
\alpha \\
\beta\n\end{array}\begin{bmatrix}\nX \sim \theta_X^1 \theta_X^{1^{\dagger}} & \theta^1 \\
\theta^{1^{\dagger}} & n\n\end{bmatrix}\right) \sim \left(\begin{array}{c}\n\alpha \\
\alpha \\
\beta\n\end{array}\begin{bmatrix}\n\alpha \\
\alpha_L^1 & d_L \\
\alpha_L^2 & d_L\n\end{bmatrix}\right) \right)
$$
\n
$$
\left(\begin{array}{c}\n\alpha \\ \alpha_L^2 & d_L \\
\alpha_R^2 & d_R \\
\theta^2\n\end{array}\begin{bmatrix}\n\alpha \\ \alpha_L^2 & d_L \\
\alpha_R^2 & d_R \\
\theta^2\n\end{bmatrix}\right) \qquad (9.46)
$$

This provisionally provides an explanation for the existence of the left-handed neutrino ν_L while the corresponding right-handed state ν_R is prohibited, at least at the level of the basic symmetry structures, as a feature of the breakdown of leftright symmetry through the identification of external spacetime in the breaking of the full symmetry of $L(v_{56}) = 1$. This observation is accompanied by the caveat concerning the Weyl spinor composition of the components of $X \subset \mathcal{X}$ and $Y \subset \mathcal{Y}$. With this in mind, and hence with quote marks placed on the ν_L , u_L and u_R states, the relation between the component structure for elements of $x \in F(h_3\mathbb{O})$, in the form of equations 9.24 and 9.25, and the first generation of Standard Model fermions is summarised in equation 9.46.

As described in the previous section, in order to obtain left-handed Weyl spinors in the components of $X = \begin{pmatrix} p & \bar{a} \\ a & m \end{pmatrix}$ a further decomposition is required, as for example in equation 9.1 or 9.5; with a similar decomposition of $Y = \begin{pmatrix} P & \bar{A} \\ A & M \end{pmatrix}$, as for example $Y = \theta_Y^1 \theta_Y^{1\dagger}$ with $\theta_Y^1 = \begin{pmatrix} \bar{R} \\ S \end{pmatrix} \in \mathbb{O}^2$, also required to obtain the corresponding righthanded spinors within the components of $Y \subset \mathcal{Y}$. With $a = sr$ in equation 9.3 or $a = sr + s'r'$ in equation 9.7 for the $a \in \mathbb{O}$ component of X, and similarly with $A = SR$ for example for the $A \in \mathbb{O}$ component of Y, this decomposition is related to the octonion property of triality for SO(8) transformations, as described near the opening of section 6.1 and around equation 6.50. In fact, with the E_6 'rotations' acting in the same way on the $\mathcal X$ and $\mathcal Y$ components by equation 9.36 and following the discussion before equation 9.9 in the previous section, the triality symmetry implies that each of $a, s, r, A, S, R \in \mathbb{O}$ transform in precisely the same way under the action of any $SU(3)_c \subset SO(8)$ transformation.

The Lorentz spinor structure under the external $SL(2,\mathbb{C})^1$ symmetry may also be obtained under an alternative decomposition of X and Y based on the $h_2\mathbb{C} \subset h_2\mathbb{O}$ subspaces, as for example in equation 7.32, as also described in the previous section. This possibility may also be relate to the structure of the technicolor models reviewed subsection 8.3.3. With the external 4-vector $h_2 \in h_2 \mathbb{C}$ accommodated within the Y components and left-handed neutrino ν_L to be accommodated in the X components ultimately a *different* decomposition of the X and Y components may be involved in consistently accounting for the corresponding empirically observed phenomena. These phenomena require the correct matching of the internal $SU(3)_c \times U(1)_Q$ action to the observed fermion multiplets of equation 7.36. Indeed, as also described in the previous section, some care is needed in order to maintain the \dot{S}_l^1 charge structure correlating with ν -lepton and u -quark states in the spinor decomposition. Ideally a yet higherdimensional form of $L(v) = 1$ may prove the best guide for uncovering this structure in a mathematically natural manner.

While further components are needed to unfold the full spinor structure, under the enlargement of the symmetry group from E_6 to E_7 on the temporal form $L(v_{56}) = 1$ we next consider the possible identification of an internal $SU(2)_L$ action within the E₇ symmetry structure. The Dynkin diagram for the rank-7 Lie algebra E_7 is compared with that for the rank-6 Lie algebra E_6 in figure 9.1.

Figure 9.1: The Dynkin diagrams for the (a) $L(E_6)$, (b) $L(E_7)$ and (c) $L(E_8)$ Lie algebras, which may be contrasted with those for the subalgebras listed in figure 7.2.

Unlike the case for E_6 , the Lie algebra E_7 does contain a rank-6 subgroup corresponding to the combined external Lorentz symmetry and internal gauge symmetry of the Standard Model, that is:

$$
SL(2,\mathbb{C}) \times SU(3) \times SU(2) \times U(1) \quad \subset \quad E_7 \tag{9.47}
$$

The description of the internal symmetry, defined in section 8.2 as the stability group of the external h₂ $\mathbb{C} \equiv TM_4$ spacetime components and adapted here with respect to the external components of $h_2 \subset \mathcal{Y}$, will be augmented beyond the 31 E₆ generators of table 8.3 to a complete set for $\text{Stab}_7(TM_4) \subset E_7$. These will include for example the actions $\psi(D)$ of equation 9.41 for which $D(h_2) = 0$ as well as any linear combination of the four sets of E_7 generators in equations 9.38–9.41 which sum to zero on the four projected components of $h_2 \subset \mathcal{Y}$ in equation 9.46. An SU(2) ⊂ Stab₇(TM₄) ⊂ E_7 subgroup, independent of the $SL(2,\mathbb{C})^1 \times SU(3)_{c}$ symmetry, acting upon the lefthanded spinors of $\mathcal X$ and, together with the identification of a further U(1) action, completing the E⁷ decomposition of equation 9.47 may be considered as a candidate for the $SU(2)_L \times U(1)_Y$ gauge symmetry of the Standard Model. Indeed such an $\text{SU}(2) \subset \text{Stab}_7(TM_4) \subset \text{E}_7$, having not been identified within the E_6 generators, being internal to the components $h_2 \subset \mathcal{Y}$ whilst acting freely on X and hence constructed asymmetrically in terms of $\phi(C)$ and $\psi(D)$, would be expected to have an asymmetric action on the left and right-handed spinors identified in equation 9.46.

Empirically it is the gauge bosons of an $SU(2)_L$ gauge symmetry which mediate interactions within doublets of quarks $\binom{u}{d}$ $\binom{u}{d}_L$ and leptons $\binom{\nu}{e}$ $\binom{\nu}{c}_L$. Hence the identification of such an internal symmetry within the present theory may be a valuable guide to the full identification of left-handed u-quark and ν -lepton states in equation 9.46 given that we have already identified left-handed d-quark and e-lepton states within the θ ¹ components of X. With a different action on the X and Y components of $x = \begin{pmatrix} \alpha & \lambda \\ \gamma & \beta \end{pmatrix}$ \mathcal{Y} β in principle the identification of such an $SU(2)_L \subset E_7$ gauge symmetry is free to act on the left-handed doublets derived for example from the components of $\binom{\theta_X^1}{\theta_1^1}_{L_1}$ identified within X , without impinging upon the external spacetime components of Y . This hence provides a free channel for charged weak transitions within the leptonic $\binom{\nu}{e}$ $\binom{\nu}{e}$ and quark $\binom{u}{d}$ $\frac{d}{dL}$ doublets which may be extracted from equation 9.46. The analysis of such an $SU(2)_L^2$ action relating to W^{\pm} gauge boson interactions, consistent with the appropriate $SU(3)_c \times U(1)_Q$ transformations and charges for the left-handed states, may also clarify the structure of left-handed spinors themselves within the X components. More generally, guided by standard electroweak theory, the identification of the ν lepton and u -quark left-handed spinors in the components of X will be mutually related to a determination of the composition of an internal $SU(2)_L \times U(1)_Y \subset E_7$ symmetry action itself.

Towards this end, and in contrast with the opening of section 8.2, an internal symmetry might be defined as any group G consistent with the subgroup decomposition $SL(2,\mathbb{C})^1 \times \underline{G} \subset E_7$ for which the set of $SL(2,\mathbb{C})^1$ spinors transform under the trivial or fundamental representations of G . That is, while the external $SL(2,\mathbb{C})^1$ symmetry partitions the components of $L(\hat{\mathbf{v}}) = 1$ into irreducible pieces, including the spinors $\theta_{l,i,j,k}$ of equation 8.13 and table 8.2 each composed of four real components, the internal symmetry G respects this partitioning in treating the Weyl spinors as individual components of a representation of \mathcal{G} . This definition excludes for example

the SU(2) generated by $\dot{G}_q + 2\dot{S}_q^1$ for $q = i, j$ and k which, as described in the opening of subsection 8.3.1, does not transform the spinors $\theta_{l,i,j,k}$ as a fundamental representation, but does still include the internal $SU(3)_c \times U(1)_Q$ symmetry as identified in section 8.2, with the actions on the spinors described in table 8.7 and equation 8.23 as summarised, via equation 8.24, in equation 8.26. The question then regards the uniqueness of this $SU(3)_c \times U(1)_Q$ action or the existence of further internal symmetry groups which possess a similarly tidy action on the spinors.

At the same time the action of $SU(2)_L \subset E_7$ might still be expected to be closely related to the subgroups $SU(2)^{2,3} \times U(1)^{2,3} \subset E_6$ acting on the components of X, since the latter have desirable properties in relation to electroweak theory as described in the 'mock electroweak theory' of section 8.3. These include the \dot{S}_l^1 charges for the \tilde{W}^{\pm} and \tilde{Z}^{0} gauge bosons, for example for $\dot{\Sigma}^{(2)\pm}$ in equation 8.39, and the similarity of the linear dependencies for the corresponding E_6 generators, as seen for example in equation 8.47, to the structure of equation 8.48 for the Standard Model. In attempting to fit an $SU(2)_L \times U(1)_Y$ symmetry into the E_6 analysis the generator \dot{S}_l^2 was also found to provide the correct hypercharges for the left-handed fermion states in $\mathcal X$ as described following equation 8.49. An $SU(2)_L \times U(1)_Y \subset E_7$ symmetry action will differ for the X and Y components of $x \in F(h_3\mathbb{O})$, with for example presumably $Q = \frac{Y}{2}$ $rac{Y}{2}$ required for the right-handed spinors in $\mathcal Y$ as singlets of $SU(2)_L$.

A quantitative test of the E_7 symmetry breaking structure might be found in a calculation of the electroweak mixing angle θ_W , following a similar derivation that led to $\sin^2 \theta_{M^2} = \frac{3}{4}$ in equation 8.63 for the mock electroweak theory within the E_6 structure. As described in section 8.2 the relative coupling of the $U(1)_O$ gauge symmetry to the fermions, in terms of the fractional charges of the quarks, already matches the observed values. The relative value of the internal $SU(3)_c$ coupling to the spinor components of equation 9.46, in comparison with the electroweak couplings, with respect to a normalised $L(E_7)$ Killing form could also in principle be calculated.

The explicit structure of the E₆ symmetry actions on the cubic form of $\mathcal{X} \in$ h3O, obtained by generalisation of Lorentz transformations on quadratic forms [38, 39, 40, 41, could ideally be further generalised to obtain the structure of the E_7 symmetry actions on the quartic form of $x \in F(h_3\mathbb{O})$. This would involve an additional $133 - 78 = 1 + 27 + 27 = 55$ generators from equations 9.39–9.41 now expressed as tangent vectors to the 56-dimensional space of $F(h_3\mathbb{O})$. In principle the application of equation 6.55 for the full set of E₇ actions on $x \in F(h_3\mathbb{O})$ could in turn be used to determine the full 133×133 $L(E_7)$ table, building upon the 78 \times 78 $L(E_6)$ table in [38]. Another approach to such a construction might be based on the identification of $L(E_{7(-25)})$ with the Lie algebra of the symplectic group Sp(6, ①) as described in [62].

An $SU(2)_L$ action might then be sought using the new generators, either solely or in combination with the original 78 E_6 generators, acting on a set of left-handed Weyl spinors, via a spinor decomposition of X, identified within the components of X in equation 9.46, as guided by the nature of electroweak interactions for the fermions. Since the $SL(2,\mathbb{C})$ ¹ Lorentz symmetry and $SU(3)_c \times U(1)_Q$ internal symmetry have already been identified in sections 8.1 and 8.2 within the E_6 actions of equation 9.38 it may be possible to use this as a starting point to more directly construct an $SU(2)_L$ symmetry with appropriate properties out of the further generators listed in equations 9.38–9.41. That is in seeking a particular $SU(2)_L \subset E_7$ action as represented on the components of X in equation 9.46 the full 133×133 Lie algebra table for E₇ may not be required.

On the other hand the study of the complete algebra, and the subalgebras it contains, may be necessary to both identify the actions $SU(2)_L \times U(1)_Y$ corresponding to electroweak theory and to determine the weak mixing angle $\sin^2 \theta_W$ for the present theory. Even in this case a 'quantisation' of the theory to describe the phenomena of 'running coupling' may be necessary in order to make comparison with the value of $\sin^2 \theta_W \simeq 0.23$ as empirically determined at the energy scale of M_Z , as alluded to shortly after equation 8.66 in subsection 8.3.2. This full picture may also be needed to include the $SU(3)_c$ interactions in this comparison, given the differing behaviour of the running coupling associated with each of the three components of $SU(3)_c \times SU(2)_L \times$ $U(1)_Y$ in the Standard Model as sketched in figure 11.10.

In constructing an $SU(2)_L \subset E_7$ action with an appropriate action on the components of X in equation 9.46, including upon the four $\theta_{l,i,j,k}$ left-handed Weyl spinors, as part of an $SL(2,\mathbb{C})^1 \times SU(3)_c \times SU(2)_L \times U(1)_Y$ subgroup decomposition, as an exemplification of equation 9.47, the $SU(2)_L$ action might also be found to act non-trivially on the $\mathcal Y$ components of equation 9.46 and in particular impact upon the external $h_2 \in h_2 \mathbb{C} \equiv TM_4$ components. This is analogous to the $D(1)_B \subset E_6$ action in the decomposition of equation 8.35 which, although independent of $SL(2,\mathbb{C})^1$ in the Lie algebra, with the generator of equation 13.5, clearly impacts upon the external spacetime components.

In the present theory it is proposed that some of the differing properties of the internal gauge interactions associated with $SU(2)_L$ compared with $SU(3)_c \times U(1)_O$ arise since the latter forms a subgroup of Stab $(TM_4) \subset E_6$, and even of Stab₂ $(TM_4) \subset$ $SL(2, \mathbb{O})$ considered as a subgroup of a 10-dimensional spacetime symmetry as described in the opening of section 9.1, while the former is only to be identified as a subgroup of E_7 , acting on a quartic form of temporal flow, such that $SU(2)_L$ is not a subgroup of $\text{Stab}_7(TM_4)$. This structure is further proposed to be closely related to the phenomena of electroweak symmetry breaking in the Standard Model, based on the study of the mock electroweak theory described for the $SU(2)^{2,3} \times U(1)^{2,3}$ subgroups of E_6 , acting on a cubic form, as described in section 8.3.

These features of a higher-dimensional temporal form $L(\hat{\mathbf{v}}) = 1$ of cubic or higher polynomial order are distinct from those of a quadratic spacetime form. For the model considered in section 5.1 with the quadratic form $L(\mathbf{v}_{10}) = 1$, representing a 10-dimensional form of time which can also be interpreted as a higher-dimensional spacetime structure, the external $SO^+(1,3)$ and internal $SO(6)$ components of the broken full SO⁺(1,9) symmetry act *independently* on the external \overline{v}_4 and internal \underline{v}_6 components of temporal flow v_{10} , respectively, as depicted in figure 5.1(b). On extension to the cubic form of time $L(v_{27}) = 1$ the external SO⁺(1,3) symmetry was found to also act on the extra 'internal' dimensions of $\theta = \begin{pmatrix} c \\ b \end{pmatrix}$, identifying a set of four Weyl spinors, as described in section 8.1 and contrasted with the 10-dimensional spacetime case at the end of that section. Here we make the complementary observation that a component of the *internal* symmetry \mathcal{G} , in the subgroup decomposition $\text{SL}(2,\mathbb{C})\times \underline{G} \subset \hat{G}$ with $\hat{G} = \text{E}_6$ or E_7 , can itself act on the projected external $v_4 \in TM_4$ spacetime components. This possibility, for a cubic or higher form of temporal flow, is proposed to underlie the origin of mass for the corresponding gauge bosons.

With significant physical properties deriving from the combined action of the external and internal symmetry on both the external and internal temporal components the present theory deviates significantly from models based on a higher-dimensional spacetime. In particular these observations mark a departure from the resemblance with Kaluza-Klein theories, as reviewed in chapter 4 and incorporated into the geometric structures of the present theory in section 5.1, which may assist in the aim of deriving a relation between the external and internal geometry, in the form of equation 5.20, in the context of the present theory alone.

Similarly as for the proposed $SU(2)_L \times U(1)_Y \subset E_7$ subgroup the $SU(2)^{2,3} \times$ $U(1)^{2,3} \subset E_6$ actions are *not* contained within Stab(TM_4) $\subset E_6$. However in section 8.3 the impingement of these actions on the $h_2 \in \mathrm{h}_2\mathbb{C}$ components of X were seen to be analogous in structure to the Higgs mechanism of electroweak symmetry breaking and the origin of the masses for the W^{\pm} and Z^{0} gauge bosons. The object h_2 , now accommodated in the $\mathcal Y$ components in equation 9.46, together with the properties of its components, is now considered as the 'vector-Higgs', providing the source for the empirically observed Higgs phenomena.

The corresponding components ' $h_2 \subset X \subset \mathcal{X}$, in the complementary $h_2 \mathbb{C} \subset$ $h_2\mathbb{O} \subset h_3\mathbb{O}$ subspace of $x \in F(h_3\mathbb{O})$ can be opened up by a spinor decomposition, as described in the previous section, to account for the ν_L fermion state. On the other hand while the vector-Higgs $h_2 \subset Y \subset Y$ could be interpreted to be composed of spinors, in the form of equation 7.32 and by analogy with technicolor models for example, the physical expression of these components is directly in terms of a tangent vector $v_4 \in TM_4$ in the external 4-dimensional spacetime. Hence in particular a righthanded neutrino ν_R cannot be accommodated in the $\mathcal Y$ components.

The two kinds of interaction for the $SU(2)_L \times U(1)_Y$ gauge fields on the X and Y components of equation 9.46 contain analogous structures to the Standard Model Lagrangian terms respectively for the weak interactions of left-handed fermions, such as equations 7.39 and 7.40, and weak coupling to the Higgs field, such as equations 7.51 and 7.52 – with the same mixing angle θ_W applying in both cases as described after equation 7.62. For the present theory the $SU(2)_L \times U(1)_Y$ symmetry breaks to $U(1)_Q$ as generated by $\dot{S}_l^1 \in L(\text{E}_6) \subset L(\text{E}_7)$ which acts upon X and Y in the same way. In the case of the X components the $S_l^1 \equiv U(1)_Q$ action misses the ν_L components of equation 9.46 accounting for the charge neutrality of the neutrino, which is described in the Standard Model in terms of equations 7.39–7.46. In the case of the $\mathcal Y$ components the $S_l^1 \equiv U(1)_Q$ action misses the h_2 components of equation 9.46 here potentially accounting for the massless nature of the photon, as suggested for the gauge field $\tilde{A}_{\mu}(x)$ for the mock electroweak theory in equation 8.71, and as constructed for the Standard Model in equation 7.61.

These two different aspects of electroweak theory may hence here be described together in terms of the broken E_7 action on the X and Y components of $L(\mathbf{v}_{56}) =$ $q(x) = 1$ in equation 9.46. While the $SU(2)_L \times U(1)_Y \subset E_7$ action may differ on the $\mathcal X$ and $\mathcal Y$ components, involving the asymmetric actions of equations 9.31 and 9.32, a unique mixing angle θ_W and surviving $U(1)_O$ symmetry with the same action on X and \mathcal{Y} should result from the symmetry breaking over the external $h_2 \subset \mathcal{Y}$ components. In principle a complementary $\mathrm{SU}(2)_R \times \mathrm{U}(1)'_Y \subset \mathrm{E}_7$ subgroup might also be identified, by reversing the contributions from equations 9.31 and 9.32, however such a symmetry,

acting on right-handed doublets of quarks $\binom{u}{d}$ $\binom{u}{d}_R$, may be heavily suppressed due to a larger impact on the external $h_2 \subset \mathcal{Y}$ components.

In the quantum theory the propagators for the gauge fields will attain a finite mass through interaction with the external h_2 components. Naturally a 'quantisation' scheme and particle concept will be needed in order to assess the properties of the particle content of the theory (as will be developed in chapter 11), with all physical particle states transforming under well-defined representations of both the external and internal symmetry.

In identifying an $SU(2)_L \times U(1)_Y \subset E_7$ symmetry it will be desirable to maintain the features of the electroweak theory studied in section 8.3, in particular with a degree of impingement on the $h_2 \subset \mathcal{Y}$ components accounting for the corresponding gauge boson masses. This is counter to the provisional assumption in the opening of section 8.2 that an internal symmetry should belong to the stability group Stab $(TM₄)$ of the external spacetime components $h_2 \in h_2 \mathbb{C} \equiv TM_4$. Rather here in a decomposition such as equation 9.47 the emphasis is upon defining internal subgroups through the structure of their well-defined representations on the external $SL(2,\mathbb{C})^1$ spinors.

The internal $SU(3)_c \times U(1)_Q$ symmetry may also be motivated in this way, as a component of the E⁷ decomposition with well-defined representations on the spinor components, as seen in equation 8.26 for example. In this case the fact that it also happens that $SU(3)_c \times U(1)_Q \subset Stab_7(TM_4)$ is responsible for the fact that the gauge bosons associated with QCD and QED happen to be massless. It may also be the case that an internal $SU(2)_L \times U(1)_Y \subset E_7$ symmetry might also impinge on any of the scalars α, β, n and N of equation 9.46, all of which are invariant under the $\text{SU}(3)_c \times \text{U}(1)_Q$ action. The possible physical consequences of these scalar components remains to be seen, whether in terms of masses for the gauge bosons and fermions or other effects (as will be considered in chapter 13). It also remains to be seen whether the action of an $SU(2)_L \times U(1)_Y \subset E_7$ on the $h_2 \subset Y$ components may be more closely analogous to the action of the Standard Model group $SU(2)_L \times U(1)_Y$ on the Higgs complex doublet ϕ than was the case for the mock electroweak theory.

In the Standard Model fermion masses are introduced through Yukawa couplings to the Higgs field, as described in the Lagrangian of equation 7.69. In the present theory there is neither a fundamental scalar Higgs field nor an explicit Lagrangian, however amongst the long list of quartic terms in the expression for $q(x) \equiv L(v_{56}) = 1$ in equation 9.28 the top line includes the terms:

$$
q(x) \sim (\alpha \beta - ph - mh - nN) \left(\langle b, B \rangle + \langle c, C \rangle \right) \tag{9.48}
$$

with the 'vacuum value' $P = M = v^0 = h$ substituted in using equation 8.72 applied to the 'vector-Higgs' $h_2 \subset Y \subset Y \in h_3\mathbb{O}$ components of $x \in F(h_3\mathbb{O})$. Terms of this form contain both left-handed $\binom{c}{b} \subset \mathcal{X}$ and right-handed $\binom{C}{B} \subset \mathcal{Y}$ components and in this sense are reminiscent of the Standard Model Lagrangian mass terms deriving from equation 7.69. The terms of equation 9.48, in potentially contributing to the fermion masses in the full theory, supersede the cubic terms such as $h(b\bar{b} + c\bar{c})$ obtained from the form $\det(\mathcal{X}) \equiv L(\mathbf{v}_{27}) = 1$ as described for equation 8.76. In both cases the full set of terms transform under the broken symmetry of $L(\hat{v}) = 1$ rather than under the symmetry of a Lagrangian. The introduction of non-standard mass terms, as extracted from equation 9.28, is not unprecedented as can be seen by comparison with the quartic term of equation 8.75 for the technicolor model described in subsection 8.3.3.

A key part of developing the present theory will be the identification of the empirically observed properties of the neutrino sector. Of particular interest will be to identify a description of neutrino oscillations, and contrast that structure with the CKM mixing in the quark sector. These structures are also expected to relate closely to the identification of fermion masses. The low value of the left-handed neutrino mass may correlate in this theory with the lack of a right-handed counterpart in the components of $\mathcal Y$. Again with reference to equations 9.24, 9.25 and 9.46, and based on the structure of equation 8.28, the neutrino is associated with the $a_{1,l}$ components of \mathcal{X} . As possible contributions to the particle masses, in addition to equation 9.48 above, equation 9.28 contains the quartic terms:

$$
q(x) \sim PM|a|^2 + MN|b|^2 + NP|c|^2
$$

= $h^2|a|^2 + hN(|b|^2 + |c|^2)$ (9.49)

where the second line again follows on substituting the vacuum value $P = M = v^0 = h$. Hence, as well as relating to the lack of right-handed $A_{1,l}$ neutrino components in $\mathcal{Y},$ the low mass of the neutrino in comparison with the electron might depend upon the magnitude of h in comparison with that of the scalar field $N(x)$. In any case the quartic 'mass terms' for the neutrino state do differ from those for the charged lepton, and in a somewhat more complicated manner than suggested by the terms of equation 9.49 alone. Although the u-quarks have both left and right-handed components there are also differences between such quartic terms for the u and d quark states identified in equation 9.46; in this case required to account for the smaller empirical mass difference with $\frac{m_u}{m_d} \sim 0.5$ [44]. The proximity of the u and d quark masses may in fact be related to the attainment of a stable vacuum value for $L(\mathbf{v}_4) = h^2$, as will be discussed in section 13.2, and which may also correlate with the low mass for the neutrino.

With the ambiguity over the possible *mathematical* ways in which to decompose the components of $X, Y \in h_2\mathbb{O}$ into a set of spinors, as described in the previous section, ultimately an understanding of the nature of the empirically observed particle states as originating out of the present theory may be required in order to motivate a natural physical choice for such a decomposition. The full identification of scalar, spinor and gauge boson particle states will require consideration of a means of 'quantisation' for the present theory, and we pursue that direction in the following two chapters.

The scheme in equation 9.46 accounts for one family of quarks and leptons with the appropriate transformations under the internal $SU(3)_c \times U(1)_Q$ symmetry and external $SL(2,\mathbb{C})^1$ symmetry, within the above caveat for the u-quark and vlepton fermion states. In addition the particle states yet to be identified include the second and third generation of fermions, as related through CKM mixing in the case of the quark states, and their relation to the massive gauge bosons associated with electroweak theory in the Standard Model. In the following section we speculate on the possible nature of a yet higher-dimensional form of temporal flow in principle capable of accommodating these phenomena.

9.3 E₈ Symmetry and the Standard Model

The extension from E_6 acting on $L(v_{27}) = 1$ to E_7 acting on $L(v_{56}) = 1$ can be considered as a continuation of the progression to higher-dimensional forms of temporal flow which began with the $SL(2,\mathbb{C})$ Lorentz symmetry of the quadratic form $L(\mathbf{v}_4) = 1$ on 4-dimensional spacetime. This progression, the first stages of which were also described in the opening of section 9.1, is summarised here in table 9.1.

	form	dimensions	space	symmetry	$#$ generators
		$L(\mathbf{v}_4) = 1$ quadratic 4 spacetime	$\boldsymbol{v}_4\in\mathrm{h}_2\mathbb{C}$	$SL(2,\mathbb{C})$	
		$L(X) = 1$ quadratic 10 spacetime	$X\in\mathrm{h}_2\mathbb{O}$	$SL(2,\mathbb{O})$	45
		$L(\mathcal{X}) = 1$ cubic 27 temporal $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$		$E_{6(-26)}$	78
$L(x) = 1$		quartic 56 temporal $x \in F(h_3\mathbb{O})$		$E_{7(-25)}$	133

Table 9.1: Four-dimensional spacetime, as a form of temporal flow itself, may be embedded in a progression of higher-dimensional temporal forms.

The highest dimensional form of temporal flow $L(v_{56}) = 1$ has a symmetry breaking pattern to $E_{6(-26)} \subset E_{7(-25)}$ with the representations of equation 9.33 as exhibited by the structure of equation 9.29. This is analogous to the further breaking pattern of E_6 to $SL(2, \mathbb{O}) \equiv Spin^+(1, 9)$, as described by the representations of equations 8.1–8.3, which is also implied in the structure of left-hand side of equation 9.46. The $SL(2, \mathbb{O})$ symmetry of 10-dimensional spacetime is an intermediate stage on the way down to the Lorentz $SL(2, \mathbb{C})$ symmetry which further decomposes the representation space into a Lorentz 4-vector, Weyl spinors and Lorentz scalars, as described in table 8.2 and now applied to both X and $\mathcal{Y} \in \mathrm{h}_3\mathbb{O}$, with the external Lorentz 4vector $v_4 \in TM_4$ accommodated within the Y components in equation 9.46, where two further Lorentz scalar components α and β are also identified.

Apart from the three additional scalars, N, α and β in equation 9.46, the increase in dimension from 27 to 56 does not contain any redundancy in terms of comparison with the structures of the Standard Model. Most of the additional 29 dimensions are interpreted as an augmentation from 2-component Weyl spinors to 4-component Dirac spinors, together with a separation in the identification of the lefthanded neutrino state and the external spacetime $h_2 \mathbb{C} \equiv TM_4$ components.

At the level of the Dynkin diagrams of figures $9.1(a)$ and (b) the E_7 algebra marks a minimal extension from E_6 , but one which together with the 56 representation as described in the previous section may be sufficient to account for much of the structure of Standard Model symmetries and particle spectrum with little further augmentation. This additional augmentation is needed to account for the identification of the *u*-quark and ν -lepton spinor components which will require a further decomposition of the $X \subset \mathcal{X}$ and $Y \subset \mathcal{Y}$ components, with for example $X = \theta_X^1 \theta_X^{1 \top} + \phi_X^1 \phi_X^{1 \top} + \dots$ as described in equations 9.1 and 9.5 of section 9.1. The features of these two means of augmenting the form of temporal flow to a higher dimension, as described in the two previous sections, will need to be combined in the complete theory. One possible

means of achieving this will be described in the present section.

In terms of the dimension of the underlying space, as listed for the sequence of forms $L(v) = 1$ in table 9.1, we first note that a further expansion from 56 to ~80 real components would be sufficient incorporate Weyl spinors for the ν_L , u_L and u_R states of equations 9.46. This is deduced by observing that $a \in \mathbb{O}$ of equation 8.28 has 8 real components while a set of four Weyl spinors requires a total of 16 real components, or alternatively by noting that the decomposition of the form $X = \theta_X^1 \theta_X^{1\bar{1}}$ involves an augmentation from 10 to 16 real components. With a complete generation of Standard Model fermions then accounted for the second and third generations might also be directly incorporated under a further augmentation from 80 to \sim 240 real components.

Given the progression to larger symmetry groups summarised in table 9.1 from a mathematical point of view it is also natural to consider whether the Lie group E_8 , as the largest exceptional Lie group, represented on a quintic homogeneous form $L(v) = 1$, may mark one further and final possible step in this sequence. (While we refer to such a hypothetic 'quintic' form, essentially an order greater than quartic is implied). With the smallest non-trivial representation of E_8 being 248-dimensional, this possibility is particularly worth consideration in light of the observations of the previous paragraph. In a similar way that extending the symmetry from E_6 to E_7 led to the incorporation of right-handed as well as left-handed fermion states, ideally a further extension to E_8 would subsume both the E_7 symmetry of the structure in equation 9.46 and explicitly incorporate also the u-quark and ν -lepton spinor states and a full three generations of fermions all under a higher-dimensional form of $L(\mathbf{v}) = 1$ with an E_8 symmetry.

The smallest non-trivial representation of E_6 is the 27 which can be expressed as the symmetry of the cubic form $L(v_{27}) = 1$, while for E₇ the 56 representation, again the lowest-dimensional non-trivial representation, can be expressed as the symmetry of the quartic form $L(v_{56}) = 1$. However the 248 representation for E₈ is expressed in terms of the adjoint representation on the 248-dimensional E_8 Lie algebra itself, with no clear interpretation in terms of a symmetry of a form of temporal flow $L(\mathbf{v}) = 1$. Indeed the Lie algebra E_8 can be essentially introduced in terms of its action on itself, and constructed in purely algebraic terms which may involve the octonions [1], with the absence of any geometric motivation or application which might be related to a form $L(v) = 1$.

The group E_8 can be defined as the symmetry group of a 57-dimensional manifold based on $F(h_3\mathbb{O})+\mathbb{R}$, known as the 'extended Freudenthal triple system' equipped with extra geometric structure [61], with $E_7 \subset E_8$ now identified as a subgroup. These mathematical objects have some connection with the structures of M-theory [60, 61], and indeed E_8 features heavily in some branches of theoretical physics as for example in $E_8 \times E_8$ heterotic string theory. There is also some debate in the literature concerning whether or not the structure of the E_8 Lie algebra alone is large enough to fully describe the Standard Model together with gravity (see for example [63]).

For the present theory with an E_8 symmetry acting on a *hypothetical* form of temporal flow, which may be denoted $L(v_{248}) = 1$, three generations of fermions together with a vector-Higgs could be accommodated within the 248 temporal components, as outlined above, while the external Lorentz group and internal gauge symmetries could all in principle be identified within the E_8 group actions. Without the need to employ a supersymmetry there are no SUSY states or set of 'mirror' particles of any kind, although it is of course possible that new physics might be predicted with consequences that might be tested. However in the present theory the primary guiding principles are driven by conceptual ideas, rather than taking a fundamental motivation from a notion of *mathematical* elegance. Hence here it is the possible forms of temporal flow $L(v) = 1$, together with their symmetries, which lead the development of the theory, and this may or may not involve the Lie group E_8 . The progression towards higher dimensions in table 9.1 does however strongly hint towards consideration of E_8 , hence allowing this one lead from the perspective of mathematical beauty, we consider the possible marriage of this 'aesthetically pleasing mathematics' with the underlying conceptual form of the present theory.

The fact that the smallest non-trivial representation of the 248-dimensional E_8 Lie algebra is expressed as the adjoint representation does not itself preclude the possibility that a 248 representation may also be identified in terms of the symmetries of a quintic form $L(v_{248}) = 1$. By comparison for example the smallest non-trivial faithful representation of $SO(3)$ is the adjoint representation on the 3-dimensional $SO(3)$ Lie algebra, but in this case there is also a fundamental 3 representation preserving the magnitude of vectors $v_3 \in \mathbb{R}^3$ in a 3-dimensional Euclidean space. Indeed the SO(3) symmetry of the 3-dimensional form $L(v_3) = 1$ of equation 2.14 was the example of a symmetry of a multi-dimensional form of temporal flow with which we began in section 2.2. In this case the two representations of $SO(3)$ are closely related since the bilinear Killing form on the elements of the so(3) Lie algebra has the same structure as the quadratic scalar product in the space \mathbb{R}^3 as used in forming the magnitude $|v_3|$.

A quintic form underlying $L(v_{248}) = 1$, invariant under an E₈ symmetry action, may not be as closely related to the adjoint representation and the $L(E_8)$ algebra structure, unless such a quintic form (or more generally a homogeneous polynomial form of order greater than four) might be related to the bilinear Killing form on $L(E_8)$ in some way. Further, given the progression from the cubic polynomial form $\det(\mathcal{X})$ of equation 6.28 as an expression of $L(v_{27}) = 1$ with an E₆ symmetry to the terms of the quartic form $q(x)$ of equation 9.28 underlying the form $L(v_{56}) = 1$ with an E_7 symmetry, a possible quintic form for $L(v_{248}) = 1$ with an E_8 symmetry may be a considerably more complicated mathematical object still. Hence it is perhaps conceivable that such a structure has not been identified through purely algebraic means, even over fifty years after the corresponding E_6 and E_7 structures were first realised. On the other hand if such a mathematical structure does exist, namely a quintic form $L(v_{248}) = 1$ with an E₈ symmetry, then as for the other forms of table 9.1 it would naturally apply for the present theory, based on multi-dimensional forms of temporal flow, and further physical consequences would be expected to be uncovered in this further progression.

In reference [64], as an example of a more geometrical approach, all of the classical Lie groups are accounted for as isometry groups of bilinear or sesquilinear forms and the first four exceptional Lie groups, G_2 , F_4 , E_6 and E_7 , are described as isometry groups constructed for cubic or quartic forms, but with E_8 essentially absent from the discussion. More generally little reference has been identified in the literature in which a 248-dimensional representation of E_8 is described in terms of an action on a quintic, or any other homogeneous polynomial, form. However in [65, 66] a polynomial of degree eight which is invariant as a 248-dimensional representation of the compact real form of E_8 is described, and is closely related to an invariant polynomial for the real form $E_{8(8)}$. For the present theory it is then an open question whether an octic form with an E_8 symmetry might contain the quartic form with E_7 symmetry. Such a natural extension consistent with the form of temporal flow $L(\mathbf{v}) = 1$ may also be required to have a symmetry described by a non-compact real form of E_8 in order that temporal causality may be respected for physical structures identified on the base manifold M_4 , with a local $SO^+(1,3) \subset E_8$ symmetry, as will be discussed in section 13.3.

Considering the possible real forms of E_8 more generally, a suitable candidate would be $E_{8(-24)}$ since the following maximal subgroups involving the exceptional Lie groups are well known (see for example [67]):

$$
E_{7(-25)} \times SU(1,1) \quad \subset \quad E_{8(-24)}
$$

\n
$$
E_{6(-26)} \times SO(1,1) \quad \subset \quad E_{7(-25)}
$$
\n(9.50)

This suggests the employment of the chain of non-compact real forms $E_{6(-26)} \rightarrow$ $E_{7(-25)} \rightarrow E_{8(-24)}$ as symmetry groups for the corresponding forms of the sequence $L(v_{27}) = 1 \rightarrow L(v_{56}) = 1 \rightarrow L(v_{248}) = 1$, where the first two stages have been described here in chapter 6 and section 9.2 respectively, while the third form remains hypothetical. As for the structure of the first two stages it seems likely that a construction of the final form in this progression will involve the algebraic structure of the octonions in a significant way.

At the level of the complex Lie algebras the corresponding Dynkin diagrams for E_6 , E_7 and E_8 are displayed alongside each other in figure 9.1. The Lie group generated by the rank-8 $L(E_8)$ algebra is large enough to contain a rank-8 decomposition of the form:

$$
SL(2,\mathbb{C}) \times SU(3) \times SU(2) \times SU(2) \times U(1) \times U(1) \subset E_8
$$
\n(9.51)

as can be shown by straightforward analysis of the Dynkin diagrams involved. Hence while the degrees of freedom of the components of v_{248} , as an extension from $v_{56} \equiv$ $x \in F(h_3\mathbb{O})$ of equation 9.46, are sufficient to contain a full three generations of Standard Model fermions and a vector-Higgs, the E_8 symmetry group is comfortably large enough to describe the external Lorentz symmetry together with the internal $\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y$ gauge group.

While the higher-dimensional extensions of section 9.1 were *contrived*, for example via equation 9.5 leading to the inhomogeneous expression of equation 9.8, in order to describe the further necessary spinors and generations for the Standard Model, ideally these structures will be found to arise naturally within a homogeneous form $L(v_{248}) = 1$ under an E₈ symmetry broken over an external 4-dimensional spacetime M_4 . This natural structure should include a full set of $SU(3)_c \times U(1)_Q$ transformations and charges aligned with the $SL(2,\mathbb{C})^1$ spinors, completing the structure identified within the E_6 action in equations 8.26 and 8.28, and in particular supplying a mathematical justification for the electromagnetic charges of the ν -lepton and u -quark spinor states through a $U(1)_Q$ action which might be related to the form of equation 9.12 for example.

Towards the end of the previous section an $SU(2)_R \times U(1)'_Y$ subgroup was considered as a possible complementary alternative to $SU(2)_L \times U(1)_Y$ within a decomposition of E_7 in the form of equation 9.47, however the E_8 breaking structure of equation 9.51 can in principle accommodate both subgroups together. In general a decomposition of E_8 in the form of equation 9.51, arising from the symmetry breaking through a choice of $SL(2,\mathbb{C})^1$ on the external spacetime TM_4 , will contain internal symmetry groups acting on the set of spinors which do not belong to the stability group Stab₈(TM₄) ⊂ E₈. This may include for example an 'SU(2)_R', acting asymmetrically on the X and Y components of the $v_{56} \subset v_{248}$ subspace in equation 9.46 or other gauge groups with a significant impingement on the vector-Higgs $v_4 \equiv h_2 \in h_2 \mathbb{C} \equiv TM_4$ components and hence associated with very massive gauge bosons, which are hence as yet unobserved as are the corresponding gauge interactions. An internal $SU(2)_L \times U(1)_Y$ is anticipated which is also broken through a degree of impingement on the vector-Higgs components, resulting in the empirically observed massive W^{\pm} and Z^{0} gauge bosons and associated electroweak phenomena, as a progression from the 'mock electroweak theory' described in section 8.3.

Given the projection of $v_4 \in TM_4$ with an external $SL(2, \mathbb{C})^1$ symmetry and the set of fermions identified in the residual v_{248} components transforming under the internal $SU(3)_c \times U(1)_Q$ symmetry, the further internal $SU(2)_L$ symmetry action will be sought as acting on doublets of quark $\binom{u}{d}$ $\binom{u}{d}_L$ and lepton $\binom{\nu}{e}$ $\binom{\nu}{e}_L$ left-handed Weyl spinors, and not only for the first but also the second and third generation of Standard Model fermions. Masses for the fermions are anticipated to arise through interaction with the vector-Higgs as expressed in the quintic terms of $L(v_{248}) = 1$ as an extension from the quartic terms such as those of equations 9.48 and 9.49 for the $L(v_{56}) = 1$ case. A misalignment between the $SU(2)_L$ weak doublet states and the mass eigenstates for the quark sector is expected to give rise to the phenomena of CKM mixing, as described for the Standard Model towards the end of section 7.2, with a related consideration leading to the phenomena of neutrino oscillations in the lepton sector.

It would be possible to attempt to embed the structures of the Standard Model, as alluded to above, into the components of a quintic form $L(\mathbf{v}_{248}) = 1$ with an E₈ symmetry if the latter structure was already known and described in the literature. This would continue the approach adopted for the E_6 symmetry of $L(v_{27}) = 1$ and E_7 on $L(v_{56}) = 1$, as based on the corresponding mathematical structures originally discovered in the 1950s [34] and 1960s [35] respectively, for which the consequences of symmetry breaking over M_4 have been studied here in chapter 8 and section 9.2.

Alternatively the mathematical structure of E_8 acting on a quintic form underlying $L(v_{248}) = 1$, if it exists, might itself be *constructed* through its application in the present theory as a form of temporal flow based on a knowledge of the empirical properties of the Standard Model. That is, continuing the progression of table 9.1 through the Standard Model structure identified in the components of $F(h_3\mathbb{O})$ under the broken E_7 symmetry in equation 9.46, and using the need to identify spinor components for the ν -lepton and u -quarks, together with three generations of fermions oriented under an $SU(2)_L$ action and relating to CKM mixing, all in a structural correspondence with the Standard Model, might lead to the identification of a suitable underlying 248-dimensional space. The study of this mathematical structure, incorporating the subspaces of h₃ \mathbb{O} and $F(\text{h}_3\mathbb{O})$ under the subgroups E_6 and E_7 respectively,

may lead to the identification of an E₈ symmetry represented on the form $L(v_{248}) = 1$, which might then be rigorously studied as an objective mathematical entity in its own right. Such an interplay between the development of physical theories and mathematical structures has a long history of stimulating mutually beneficial progress, as for the parallel development of gauge theories and the structure fibre bundles reviewed in chapter 3.

Essentially this is the approach we have set out to follow in section 9.1 in attempting to open up further components to account for further spinors and further generations through augmentation such as that in equation 9.5. The aim is then to combine that form of extension with the augmentation to the action of E_7 on $F(h_3\mathbb{O})$ described in section 9.2 in seeking an E⁸ action on a 248-dimensional space such that a homogeneous quintic, or higher order, norm $L(v_{248}) = 1$ is invariant. As well as aiming to incorporate the essential structure of the Standard Model, in progressing in this way it is also possible that new features will appear for E_8 acting on $L(v_{248}) = 1$ as the full form of temporal flow.

For the subgroup action of $SL(2,\mathbb{O})^1 \subset E_6 \subset E_7$ on the components of $F(h_3\mathbb{O})$ in equation 9.46 the subspace elements $X \in h_2\mathbb{O}$ and $Y \in h_2\mathbb{O}$ transform as 10dimensional spacetime vectors, and need to be opened up to identify a spinor substructure as discussed in section 9.1. However under the corresponding $SL(2,\mathbb{O})^1 \subset E_8$ subgroup on the hypothetical form $L(v_{248}) = 1$ the object θ^1 (denoted θ in equation 8.2) might be directly identified along with further Majorana-Weyl spinors, including θ_X^1 and ϕ_X^1 of equation 9.5, all as naturally occurring representations within the enlarged structure and without any direct vector representations in the components of v_{248} . Further, as for the four-way decomposition of θ^1 in equation 8.13 the Majorana-Weyl spinor θ_x^1 , for example, will decompose into a set of four left-handed Weyl spinors $\{\theta_{Xl}, \theta_{Xi}, \theta_{Xj}, \theta_{Xk}\}\$ under the Lorentz subgroup $SL(2, \mathbb{C})^1 \subset SL(2, \mathbb{O})^1$; with an internal SU(2)_L \subset E₈ symmetry action sought on doublets of the SL(2, \mathbb{C})¹ Weyl spinors identified within $\begin{pmatrix} \theta_X^1 \\ \theta^1 \end{pmatrix}$.

In this case in place of *decomposing* a vector into spinor representations, as for equation 9.5, for the $SL(2,\mathbb{C})^1 \subset E_8$ action the need is rather to *construct* a 4component vector v_4 to be locally associated with the external tangent space TM_4 . This may be achieved by going the other way and composing together right-handed spinors for example (such as effectively associated with a subset of the Y components in equation 9.46), under $SL(2, \mathbb{C})^1 \subset E_8$ to form the 4-vector:

$$
h_2 = \theta_{Y\mathcal{L}}(\theta_{Y\mathcal{L}})^{\dagger} + \phi_{Y\mathcal{L}}(\phi_{Y\mathcal{L}})^{\dagger} \tag{9.52}
$$

This is essentially equation 7.32, interpreted as composing the right-hand side to form the left-hand side rather than as a decomposition of the latter. Here the Weyl spinors are fused together through the projection of the full temporal flow onto the external spacetime M_4 as an arena for perception in the world, with the local Lorentz symmetry acting on the 4-vectors $h_2 \equiv v_4 \in TM_4$ which also forms the vector-Higgs in the present theory. While this structure is analogous to the formation of a scalar Higgs in technicolor models, as a condensate of a set of proposed techniquarks interacting under an $SU(N)_{tc}$ gauge symmetry as reviewed in subsection 8.3.3, in the present theory it is the identification of the geometric form of an external spacetime as an innate feature of perception, as described in section 2.2, which necessarily draws together

spinor components into a 4-vector composition. This 4-vector h_2 under $SL(2,\mathbb{C})^1 \subset$ $SL(2,\mathbb{O})^1 \subset E_6 \subset E_7$ can be seen directly in the components of the various forms of $L(v) = 1$ for the progression in table 9.1, as shown explicitly for example in the components of $F(h_3\mathbb{O})$ on the right-hand side of equation 9.46, now considered as intermediate stages on the way to the full form $L(v_{248}) = 1$.

The fusing of $U(1)_Q$ charge neutral Weyl spinors $\theta_{Y\mathcal{L}}$ and $\phi_{Y\mathcal{L}}$ (perhaps also with a third spinor $\psi_{Y\mathcal{L}}$ to form the external spacetime vector $h_2 \equiv v_4 \in TM_4$ in equation 9.52, through the requirement of perception on the extended manifold M_4 , is consistent with absence of physical particle states corresponding to the right-handed neutrino (for all three generations). On the other hand the complementary θ_{Xl} , ϕ_{Xl} and ψ_{Xl} spinors under $SL(2,\mathbb{C})^1 \subset E_8$ remain free as a full set of three generations of left-handed neutrinos. This analysis is similar to that described for equation 9.46 under the E_7 symmetry, with $h_2 \in TM_4$ accommodated in the $\mathcal Y$ components and the ν_L -neutrino derived from the X components, for the first generation only.

Although provisional, this discussion for the hypothetical action of E_8 on $L(v_{248}) = 1$ describes a possible marriage of a full form of temporal flow, deduced on the basis of mathematical elegance as a further progression from the sequence in table 9.1, together with the basic conceptual framework of the present theory, with a knowledge of the Standard Model structure presiding over the union.

Whether or not E_8 will ultimately feature in a significant way for the present theory remains to be seen. Here the primary focus is upon homogeneous forms of temporal flow expressed as $L(v) = 1$, as derived in section 2.1, and if it happens that E_8 does not form a symmetry group of such an object then it seems unlikely that this largest exceptional group will play an important role in this theory. However such homogeneous forms are known for Lie groups as large as E_7 , as we have described in this paper. We end this chapter with a summary of the Standard Model features, based on the gauge symmetry $SU(3)_c \times SU(2)_L \times U(1)_Y$, which have been identified up to this stage within the breaking of known symmetries of $L(v) = 1$ forms through the extraction of an external Lorentz symmetry. These are described in relation to the progression of higher-dimensional forms of temporal flow listed in table 9.1.

- $L(v_4) = 1$: External spacetime Lorentz symmetry SL(2, C) acting on $v_4 \in TM_4$. The Lorentz transformations on 4-dimensional spacetime are subsequently identified with the subgroup $SL(2, \mathbb{C})^1 \subset E_6$ within the larger symmetry, as generated by the basis elements $\{\dot{B}_{1z}^1, \dot{R}_{1x}^1, \dot{B}_{1x}^1, \dot{B}_{1z}^1, \dot{R}_{1z}^1, \dot{R}_{z1}^1\}$ for sl $(2,\mathbb{C})^1$ of equation 6.57.
- $L(X) = 1$: Internal symmetry $SU(3)_c \times U(1)_Q$ actions may be identified in Stab₂(TM_4) ⊂ SL(2, \mathbb{O}). In the context of the subsequent E₆ action this symmetry is generated by the basis elements $\{\dot{A}_q, \dot{G}_l\} + \dot{S}_l^1$ acting on the components $X = \begin{pmatrix} p & \bar{a} \\ a & m \end{pmatrix} \in h_2 \mathbb{O} \subset h_3 \mathbb{O}$, with the transformations of the $a \in \mathbb{O}$ components as described in equation 8.28. (This form is closely related to the $L(\mathbf{v}_{10}) = 1$ model of figure 5.1).
- $L(\mathcal{X}) = 1$: The additional $\theta^1 = \begin{pmatrix} \frac{c}{b} \end{pmatrix} \in \mathbb{Q}^2 \subset h_3 \mathbb{O}$ components (θ in equation 6.26) transform under the external $SL(2,\mathbb{C})^1$ as 4 left-handed Weyl spinors $\theta_l, \theta_i, \theta_j, \theta_k$ as subspaces of \mathbb{O}^2 (equation 8.13). These spinors neatly dovetail with the corresponding internal $SU(3)_c \times U(1)_Q \subset Stab(TM_4) \subset E_6$ actions, as deduced from

table 8.7 and equation 8.23 and summarised in equation 8.26, hence identifying a charged lepton singlet and d-quark triplet.

Although the group E_6 , acting on $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$, is not large enough to contain an additional internal SU(2) symmetry a number of the more esoteric properties of $SU(2)_L \times U(1)_Y$ electroweak theory are reflected in the action of the type 2 subgroup $SU(2)^2 \times U(1)^2 \subset E_6$ generated by $\{\dot{R}_{\underline{z}1}^2, \dot{R}_{\underline{z}2}^2, \dot{R}_{\underline{z}1}^2\} + \dot{S}_l^2$, and similarly for the corresponding type 3 case, which complement the type 1 external $SL(2,\mathbb{C})^1$ actions, as described in section 8.3. These properties include the doublet actions of equations 8.29 and 8.31, a 'mock electroweak' symmetry breaking pattern leading to the mixing angle deduced for equation 8.63 and the potential origin of gauge boson masses for the broken $SU(2)^2 \times U(1)^2$ generators deriving from an impingement on the external spacetime components as described for equation 8.71. Fermion mass terms are similarly considered to arise through interactions with the projected external $v_4 \in TM_4$ components under $L(v_{27}) = 1$ as described for equation 8.76. The vector $v_4 \in TM_4$ itself is considered to constitute a 'vector-Higgs', with the degree of freedom of the magnitude $|v_4|$ provides a candidate for the empirically observed scalar Higgs.

In addition to the $SL(2,\mathbb{C})^1$ spinors identified from the components of θ^1 a corresponding set of 4 left-handed Weyl spinors may be identified within the components of θ_X^1 for example, upon introducing the decomposition $X = \theta_X^1 \theta_X^{1\dagger}$ of equation 9.1. These further Weyl spinors can be interpreted as the components of a neutrino and triplet of u -quarks, although care is needed to maintain the necessary electromagnetic charges of 0 and $\frac{2}{3}$ as explained around equation 9.12, and further some of these components coincide with the external $v_4 \in TM_4$, which has provisionally been associated with the above vector-Higgs.

• $L(x) = 1$: Containing now the E₆ 27 and $\overline{27}$ representations, equation 9.33, the external Lorentz symmetry $SL(2,\mathbb{C})^1 \subset E_7$ can be taken to act on the $v_4 \equiv$ $h_2 \subset \mathcal{Y} \in \mathrm{h}_3\mathbb{O} \subset F(\mathrm{h}_3\mathbb{O})$ components, which continue to both represent external spacetime and also account for the Higgs sector, as depicted in equation 9.46. The left-handed electron and d-quark Weyl spinors of the $L(\mathcal{X}) = 1$ case above now have right-handed counterparts, combining in 4-component Dirac spinors, as described in equations 9.43 and 9.44. A left-handed neutrino (along with a set of u_L -quark spinors) might now be identified by expanding the $X \in h_2 \mathbb{O} \subset$ $h_3\mathbb{O} \subset F(h_3\mathbb{O})$ components, while a right-handed counterpart may be excluded by the external $h_2 \in TM_4$ components of $\mathcal Y$ (while a set of u_R -quark spinors remain), as also indicated in equation 9.46.

The internal $SU(3)_c \times U(1)_Q \subset E_7$ symmetry acts on the X and Y components of equation 9.46 in the same way. Further internal symmetries may be sought which also act on the set of spinors in the shape of trivial or fundamental representations. In particular an internal $SU(2)_L \subset E₇$, with an asymmetric action on the $\mathcal X$ and $\mathcal Y$ components, might now be accommodated within the larger group. An explicitly left-right asymmetric coupling to fermion doublets for an internal symmetry $SU(2)_L \times U(1)_Y \subset E_7$ may be possible for this structure, with further analysis of the E_7 algebra required. The $U(1)_Q$ action, surviving the mock electroweak symmetry breaking over TM_4 , is identical on the $\mathcal Y$ and corresponding X components, accounting for the massless nature of the photon in the first case and the charge neutrality of the left-handed neutrino in the second case.

In augmenting the full symmetry from E_6 to E_7 , and hence embedding the Lorentz symmetry in the latter, there is a two-way choice regarding the embedding of the external spacetime $h_2 \mathbb{C} \subset F(h_3 \mathbb{O})$ in either the X or Y components, with the latter option taken in equation 9.46 as described in the text and alluded to above. The necessary asymmetry in this choice is then ultimately responsible for the left-right asymmetry observed for physical phenomena, and in particular leads to the parity violating properties of the weak interaction.

The Lie group E_7 does not have complex representations and is hence unsuitable as a unification group for the purely internal symmetry structure of the Standard Model, as mentioned in section 7.3. However the external Lorentz symmetry does have complex representations and including the $SL(2,\mathbb{C})$ action within the structure of the E_7 action on $F(h_3\mathbb{O})$ in this asymmetric way in turn implies a left-right asymmetry for the action of the residual internal symmetry. In the present theory this mechanism provides the source of parity violating phenomena (rather than such phenomena arising from the complementary actions of $SU(2)^{2,3}$ with respect to $SU(2)^{1}$ in terms of non-commutative quaternion subalgebras, as had been briefly considered in subsection 8.3.1 shortly after equation 8.31 as guided by [56] for example). In addition to the $SU(2)_L$ action the hypercharge symmetry $U(1)_Y$ also remains to be specifically identified, although the latter derives from the $\dot{S}_l^{2,3}$ $\ell_i^{2,3}$ generators for the mock electroweak theory as described for equation 8.47 for example.

Further, the *three* possible embeddings of $SL(2, \mathbb{O})$ acting on $h_2\mathbb{O}$ and \mathbb{O}^2 according to equations 6.32–6.35 within the structure of the E_6 action on $h_3\mathbb{O}$ may relate to the empirical observation of three generations of fermions. The embedding of a choice of Lorentz symmetry $SL(2, \mathbb{C})^1 \subset E_6$ acting on the type 1 subset $h_2 \mathbb{C} \subset h_3 \mathbb{O}$ breaks the discrete three-way symmetry between the type 1, 2 and 3 actions described in equations 6.32–6.35, hence also breaking the continuous type transformation symmetry. This in turn will lift the degeneracy of the three generations of fermions and may be related to the phenomena of CKM mixing between the quark states. However in order to explicitly accommodate three generations of Standard Model fermions the extension to E_7 symmetry on $L(\mathbf{v}_{56}) = 1$ may need to be further augmented to an E_8 action on $L(\mathbf{v}_{248}) = 1$, incorporating a spinor expansion of the original components in a form such as equation 9.5 with also a third term $\psi^1_x \psi^{1^{\dagger}}_x$. The possibility of this further extension to E_8 , which is currently hypothetical, has been the main topic of this section.

The above observations, through to the E_7 action on $F(h_3\mathbb{O})$, currently mark the point of closest approach between the present theory and the empirical world of elementary particle phenomena recorded in high energy physics experiments. A possible extension to an E_8 action of $L(v_{248}) = 1$ is suggested partly on aesthetic mathematical grounds and partly through the known structure of the Standard Model itself considered in the context of the present theory.

Here we have largely only considered a somewhat 'static' picture based on the structures of the forms $L(v) = 1$ and the corresponding symmetry groups, with emphasis on the explicit structure of the Lie group E_6 acting on the space $h_3\mathbb{O}$. For this case in addition to the terms arising from the expansion of equation 8.76 the constant value of $L(v_{27}) = 1$ will be expressed 'dynamically' on an extended spacetime manifold M_4 as the zero covariant derivative $D_{\mu}L(v_{27})=0$. The terms of the latter expression resulting from the symmetry breaking contain the internal gauge fields $Y_\mu(x)$, as was described for the $L(\mathbf{v}_{10}) = 1$ model in equation 5.51 – which includes an interaction between the gauge field Y_μ and the internal \underline{v}_6 components. The cubic temporal form $L(v_{27}) = 1$ does not have an interpretation as a higher-dimensional spacetime form and in this case, through the terms of $D_{\mu}L(v_{27}) = 0$, an internal gauge field can also impinge upon the *external* 4-dimensional spacetime components of $v_4 \in TM_4$. It is through this impingement that massive gauge bosons are anticipated to arise as described for the mock electroweak theory in subsection 8.3.3, with the field $v_4(x)$ termed the vector-Higgs through association with Higgs phenomena. In the full theory the masses for the W^{\pm} and Z^{0} gauge bosons of the Standard Model might be identified in this manner, while the fermion masses may arise through the composition of fermion components with the vector-Higgs under the full form $L(\hat{\mathbf{v}}) = 1$.

Within the expansion of $D_{\mu}L(v_{27})=0$ there are also terms of the form $h(bY_{\mu}\bar{b}+$ $cY_{\mu}\bar{c}$, by comparison with equation 8.76, with similar terms deriving from the quartic norm in the E₇ case, describing a coupling between the gauge field $Y_\mu(x)$ and the fermion components within h₃ \mathbb{O} . In this way the internal gauge field $Y_\mu(x)$, taking values for example in the $SU(3)_c$ Lie algebra, will mix the components of the Weyl spinors, such as those of the set $\{\theta_i, \theta_j, \theta_k\}$ in equation 8.19, creating the possibility of field interactions. Ultimately the consequences of the mutual couplings of all fields in the terms of $L(\hat{v}) = 1$ and $D_{\mu}L(\hat{v}) = 0$ will need to be assessed for the full form of temporal flow.

The initial dynamical equations for this theory derived from the relation between the geometry of the external spacetime and the curvature of the internal gauge fields, as deduced in section 5.1 and culminating in equation 5.20, as guided by the structure of Kaluza-Klein theories. Hence the gauge fields, such as $A_\mu(x), W_\mu^{\pm}(x) \dots$, in being closely related to the spacetime geometry $G^{\mu\nu} = f(A, W, \ldots)$ in the form of equation 5.31, seem to take some priority over possible fermion states which may be identified in turn through the field interactions, as will be described in chapter 11. That is given for example an initial $W^{\pm}_{\mu}(x)$ field in turn fermion fields $\psi(x)$ within doublets such as $\binom{\nu}{e}$ $\binom{u}{e}$ _L or $\binom{u}{d}$ $\binom{u}{d}_L$ will be drawn into relation with the external spacetime geometry $G^{\mu\nu} = f(\tilde{A}, W, \tilde{\psi}, \ldots)$ from the components of $x \in F(h_3\mathbb{O})$ via interactions with a gauge fields as an example of the generalisation described for equation 5.32 in section 5.2. In section 13.1 a direct relation between the spacetime geometry and the magnitude of the vector-Higgs field with $G^{\mu\nu} = f(\mathbf{v}_4)$ will also be derived, leading to a further and more direct link with fermions through the terms of $L(\hat{\mathbf{v}}) = 1$.

An understanding of the empirical consequences of all of the possible field interactions, and the phenomena of high energy physics in general, will require a full dynamical and quantum expression of the theory. This will include an understanding of how macroscopic 'mass' as central to general relativity through the field equation $G^{\mu\nu} = -\kappa T^{\mu\nu}$ is related to particle 'mass' as observed in the laboratory, and an exposition of a unified conceptual basis for describing both gravitational and quantum phenomena more generally. Within this unified framework the concept and the nature of physical elementary particles themselves might be addressed. A quantised theory dynamically expressed on the spacetime manifold M_4 will also be required in order to

deduce the empirical particle spectrum as well as to express kinematic quantities such as the masses of the particle states, for fermions as well as gauge and the Higgs bosons.

In quantum field theory (QFT) the particle masses feature in 'propagators' while charges and coupling constants appear in interaction 'vertex' terms. Both of these objects are intrinsic to calculations of cross-sections via the transition amplitude \mathcal{M}_{fi} , as will be described in the following chapter. The propagator factors in calculations of process probabilities contain various kinematic quantities with the dimension of mass. For example the Feynman propagator for the scalar Higgs field has a particularly simple form, $i/(p^2 - M_H^2)$ where p is the 4-momentum, which may provide a guide for the role of mass terms for the present theory. Here the effective incorporation of finite mass into the propagators for massive gauge bosons is expected to be related to that for technicolor models as described between equations 8.73 and 8.76 in subsection 8.3.3.

Similarly while equations 8.26 and 8.28 describe the correct $U(1)_Q$ charge structure for a generation of leptons and quarks, it will need to be understood how these 'charges' enter into cross-section calculations and hence actually account for the electromagnetic charge structure as observed for particle states in high energy physics (HEP) experiments. The interpretation of such QFT calculations within the context of the present theory will need to be addressed before the concepts of charge and mass can be fully comprehended here. The nature of field interactions and the concept of particles themselves will also need to be addressed in the course of this study, as we explore in the following two chapters.

Rather than beginning with fields or particles which are then postulated to have various properties and forms of interaction, in the present theory we begin essentially with a composition of, or coupling between, components of the full form $L(\hat{v}) = 1$ together with the generators of the symmetry transformations. Here 'masses' and 'charges' originate in the terms of the expressions for $L(\hat{v}) = 1$ and $D_{\mu}L(\hat{v}) = 0$. Only once these mathematical relations are expressed in terms of dynamical equations over the manifold M_4 , with spacetime geometry $G^{\mu\nu}(x) = f(Y, \hat{v})$ in the notation of equation 5.32, might particle states themselves be identified as a phenomenon arising out of the field interactions. In turn the observable characteristics of such particle phenomena might be determined.

The particle properties, including masses and mixing parameters, although arising from the underlying interactions of the fields, are not necessarily expected to be literally read off directly from the E_6 or E_7 symmetry breaking level. Indeed some particle characteristics, such as their behaviour under CPT transformations and the identification of antiparticles necessarily requires a theory expressed in an extended spacetime. In dealing with the bare $F(h_3\mathbb{O})$ components together with the algebraic form of the E_7 symmetry actions it can only be expected to uncover a shadow of the full variety of Standard Model phenomena at this level. However it is also desirable that this shadow should possess identifiable features, such as the correct fractional charges and a left-right asymmetry, that may plausibly underlie the empirical data. The dynamic aspects of the theory and a quantisation scheme will need to be developed in order to make more rigorous comparisons with the full variety of laboratory phenomena.

In the meantime, a collection of general properties of the Standard Model have already been identified in the study of the breaking of the E_6 symmetry of $L(\mathbf{v}_{27}) = 1$ in chapter 8 and E_7 symmetry of $L(v_{56}) = 1$ as presented in section 9.2. In particular the structure of the E₇ symmetry on the components of $F(\text{h}_3\textcircled{1})$ when broken over TM_4 makes significant contact with the Standard Model, as also summarised in this section, and further inroads may be possible by further exploring this structure. However the aim here is to avoid the possibility of contriving the appearance of Standard Model properties, but rather to be primarily guided by the development of the theory itself, albeit very much in the light of known empirical phenomena. A number of features, including the identification of u-quark and ν -lepton spinors and their $SU(2)_L$ interactions with the d-quarks and e-leptons respectively, the 'Yukawa couplings' and origin of mass, the structure of three generations of fermions and the mixing between them, remain to be better understood.

The progression towards higher-dimensional forms of time listed in table 9.1, together with the need to fill out the empirical picture, hints at the possibility of uncovering an E₈ symmetry action on a quintic form $L(v_{248}) = 1$ as the final 'Russian doll' in the sequence of enveloping symmetries of time, as we have described in this section. However, as well as extensions to higher dimensions a quantised theory and an understanding of physical particle states, as considered in the following two chapters, will be needed to identify further details of the Standard Model from within the present theory for a thorough comparison with and testing against the empirical data. Until then the extent to which the E_7 stage is sufficient or otherwise to account for the Standard Model will not be completely clear.

In this regard the main question concerns the identification of the structure of particle-like states within the theory before returning to further assess the correspondence between the present theory and empirical data, and then progress towards making predictions which may be tested. Before comprehending the particle concept it will be necessary to understand how in the present theory quantum phenomena arise together with the mathematical structures of quantum field theory which are intrinsic to calculations of high energy physics processes. Hence in the following chapter we begin by reviewing the standard machinery of QFT as applied for HEP experiments.

Chapter 10

Particle Physics

10.1 High Energy Physics Experiments

The concept of particle phenomena as observed in HEP experiments in the context of the theory presented in this paper will be examined here and in the following chapter. In this theory the world appears in our experience necessarily within the geometrical confines imposed in order for it to actually be perceived through the flow of time, with the geometrical conditions for the perceived 4-dimensional spacetime world projected out of a general higher-dimensional progression in time. The arbitrary nature inherent in a degenerate set of possible geometric solutions manifests itself as quantum and particle phenomena – such as observed in the detector apparatus of high energy physics (HEP) experiments, and through which we interact with and experience the world in general. This perspective, introduced in this section, will be described more thoroughly in the next chapter.

The phenomena of particles are observed in the laboratory in the limit of near 'vacuum' conditions as elementary transitions of the world as recorded in detector components. Similar phenomena will be manifest more generally in a curved spacetime associated with an arbitrary distribution of matter, however in the flat spacetime limit of the near vacuum, approximating the laboratory environment as considered here, these phenomena may be simpler to categorise. The 'particles' observed in HEP experiments are states of matter that arise in this simplifying limit, rather than the fundamental 'building blocks' of matter itself.

It is the aim of experimental high energy physics – employing huge and technologically complex macroscopic physical structures in the form of 'particle' accelerators, colliders and detectors coupled with sophisticated computer software and data analysis (see for example $[68]$) – to detect and analyse the most delicate and minimal transitions of the state of the perceived physical world. In this way the nature and properties of the elementary particles ascribed to such transitions are empirically determined – for example, the relative degree of interaction between particular gauge boson and quark fields in the case of [68]. In the present theory internal symmetries and fermion states have been identified at the level of the broken E_7 symmetry action on the multidimensional temporal form $L(v_{56}) = 1$, as described in section 9.2, and will relate to the components of the corresponding gauge fields and quark or lepton fields respectively, subject to dynamical constraints in extended spacetime. While significant contact has been made with structures of the Standard Model, as summarised in equation 9.46 and section 9.3, it will be necessary to identify in detail the mathematical correlate of HEP phenomena within the present framework in order to establish a closer relation between the theoretical and experimental environment and hence further assess the validity of the theory.

It should be kept in mind that the events recorded in a high energy physics experiments are not actively made to happen by physicists, rather the complete experimental apparatus is designed and built to passively make highly refined observations of the course of nature. The most elementary and minute transitions of the physical world, expressed for example in terms of gauge or fermion fields, are isolated and amplified through such experiments as exemplified in figure 10.1. Such a process, or 'event', may involve 'jets' of many final state particles as displayed in figure 10.1 or could be as simple as that sketched in figure 10.2 in the following section.

Figure 10.1: The most delicate changes of the macroscopic state of physical structures such as HEP detectors are interpreted in terms of 'particle tracks' composed out of a series of such minimal detectable transitions, here exemplified in an event recorded by the SLD collaboration [69].

All 'material' objects, such as particle detectors, are apparently infused with and seemingly 'composed of' field transitions. The environment of a HEP experiment is such that a particular series, or chain, of macroscopic transitions of the apparatus can be reconstructed, via amplified signals and computer algorithms, as a particle track. At the elementary microscopic level the particular components of equation 9.46 involved, as developed so far up to the action of E_7 on $L(v_{56}) = 1$ for the present theory, will determine the particle type, for example an electron or d-quark, with properties such as the observed bending of an electron track in a magnetic field or the manifestation of quarks in hadronic states determined by the coupling to the gauge field components. Similarly, the appearance of a set of such particle tracks, as seen in figure 10.1, is
correlated through the higher-order interactions with other fields, such as that of the $Z⁰$ gauge boson field, hence making connection with mathematical calculations in the corresponding theoretical framework.

Although the higher-order interactions may be complicated empirically the unique properties of elementary particles, such as the masses of the electron and muon for example, are independent of the external material environment (for example with the particle production and detection apparatus made of copper, silicon or other elements) as far as we can observe (excepting cases such as an 'effective mass' in a solid state device for example). These properties are measured to be the same in all the variety of experiments that have been set up to induce them, and also as they have been observed for a range of particle states in natural events such as cosmic ray showers. This robustness arises presumably since there is a universal 'vacuum' limit. Hence, although ordinary matter is complex, we expect to be able to isolate the robust and invariant quantities that describe the observed particle properties in the appropriate limit for theoretical calculations.

The eventual aim will be to calculate the effects seen in particle physics experiments in terms of transitions between the fields to determine the properties of the observed elementary particles. These include their masses and spins which categorise the particle transformations under the Poincaré symmetry of 4-dimensional Minkowski spacetime, assuming an approximately flat base manifold M_4 . An 'electron', for example, will be associated with particular field transformations under both a spinor representation of the global external Lorentz symmetry over M_4 and a particular representation of the internal symmetry of the local gauge group, with for example unit charge relative to other particle states under the $U(1)_{Q}$ action of electromagnetism.

Part of the defining notion of a particle is its local nature. A particle is an entity, whether in experiment or in theory, which causally connects and relates two spacetime events or interactions. In HEP experiments the chain of interactions can be traced from the production of the initial particle beams, through interactions with guiding magnets and accelerating components, into the interaction region of the collider and out into a spray of detector hits and signals to be recorded and analysed. Knowledge of the spacetime location of the directly detected interactions allows the reconstruction of kinematic quantities, such as the invariant mass or electric charge, of the particles ascribed to these observations.

The ultimate ambition here will be to describe what the 'in' and 'out' particle states in HEP experiments actually are, physically understood and mathematically expressed, as well as to account for the process taking place in the spacetime volume of the interaction region. In the spirit of this theory these phenomena, as for all physical processes in spacetime, will be 'enveloped' by the structure of the spacetime geometry as related to the other fields through $G^{\mu\nu} = f(Y, \hat{v})$ of equation 5.32, as described in section 5.2. It will be necessary to understand the precise general form of the right-hand side of this expression to address the question of what an elementary particle, such as an electron, is. For completeness this question will also include the physical nature of particle states such as quarks which are not observed to propagate macroscopically as independent objects in spacetime.

In standard field theory an independent flat spacetime background is given as an arena upon which fields may be arbitrarily added. Gauge invariance of a Lagrangian function composed of the fields is then postulated as a means to introduce interactions between fields, as described in sections 3.5 and 7.2. This construction is transferred to the corresponding quantum field theory (QFT) in which the gauge transformations mix internal components of the field operators such as $\phi(x)$. The field itself may be quantised by applying canonical commutation relations, by analogy with non-relativistic quantum mechanics, to the infinite degrees of freedom of the field, and particle creation and annihilation operators $a^{\dagger}(\boldsymbol{p})$ and $a(\boldsymbol{p})$ identified, as will be reviewed later in this chapter.

In contrast, in the present theory all elementary structures arise out of the interplay of multi-dimensional forms of the flow and symmetry of time expressed in $L(v) = 1$. The higher-dimensional mathematical form of temporal flow $L(v_{56}) = 1$ gives rise to the components of fields locally in interaction when perceived in physical 4-dimensional extended spacetime M_4 in a manner consistent with the underlying fundamental ordered one-dimensional flow of time. With the action of E_7 on $v_{56} \in$ $F(h_3\mathbb{O})$ broken over M_4 and the derivative $D_{\mu}L(v_{56})=0$ in turn fragmented through this 4-dimensional projection the physical manifestation of a degeneracy of causally linked exchanges between fields describing multiple solutions under $G^{\mu\nu}(x)$ will be identified as the origin of indeterministic interactions. It is these interactions which give rise to apparent particle phenomena, such as quarks and leptons, as objects of study in high energy physics experiments.

Hence the aim is then to understand how such *discrete* particle phenomena arise out of the fundamental elements of the theory, without needing to impose creation and annihilation operators, or using similar ad hoc quantisation techniques, to describe this particle-like behaviour. Rather the mathematical structures of the present theory are intended to match the physical structure of the world down to the most elementary level. Here particles should be derived as a phenomenon arising out of the possibility of multiple field solutions under $G^{\mu\nu}(x)$ on M_4 .

The principle goal of the following chapter will be to consider how the new theory describes the phenomena observed in high energy physics experiments, yet without the conceptual problems – for example regarding the particle interpretation – of quantum field theory. In particular this essentially means to be able to match the cross-section calculations for particle interactions in QFT except with both an underlying motivation for the nature of probabilities in these processes and a clearer understanding of the particle concept itself.

Quantum field theory, although incomplete, provides a set of pragmatic tools and strategies which have achieved great empirical success, and hence much of the mathematical machinery is expected to remain of importance. The preliminary and general nature of QFT allows for the successful elements to be extracted for comparison with the present theory. It is the agreement between calculations based on scattering matrix amplitudes in QFT and cross-sections measured in the laboratory that needs to be accounted for in the context of the present theory, and hence in the remainder of this chapter we review some of the standard textbook material on the structure of such calculations for reference in the following chapter.

10.2 Cross-section Calculation

In this chapter we consider how quantum field theory (see for example [10, 70, 71]) is employed in practice to calculate cross-sections for processes observed in high energy physics experiments, for example in proton machines such as the LHC, but in particular for the kind of events detected in electron-positron colliders as depicted in figure 10.1. The cross-section $\sigma(e^+e^- \to X)$ for a particular process quantitatively represents the likelihood for the production of the final state X . The description of this final state in general combines a particular collection of outgoing particles, or of 'jets' containing a spray of particles as for the event in figure 10.1, together with a particular range of kinematic or geometric characteristics.

The aim here will be to present the cross-section for such processes and then strip down this expression to identify how the basic structure of QFT is used to calculate the probability of such events. In the following chapter we describe how such calculations might be reconstructed in the context of the present theory. Given the cross-section $\sigma(e^+e^- \to X)$ the predicted event rate R (for N events per t seconds) is simply:

$$
R \equiv \frac{dN}{dt} = L\sigma \tag{10.1}
$$

which also *defines* the luminosity value L at which the machine is operating while producing the events. In practice 'bunches' of incoming particles are directed through the interaction region of the experiment, with bunches of $n_-\$ electrons facing oncoming bunches of n_{+} positrons (where the apparent number n_{+} of particles per bunch can be closely estimated from the total charge or energy carried by the bunch). With the effective two-dimensional overlap, normal to the beam direction, of the opposing bunches given by the area A and the rate of bunch crossings given by the frequency f , in the laboratory centre-of-mass frame, the luminosity is simply:

$$
L = \frac{fn_+n_-}{A} \tag{10.2}
$$

If this luminosity L, in units of cm⁻²s⁻¹, is known in addition to the cross-section σ , in units of cm^2 , then the rate of *detection* of the corresponding events will be R in equation 10.1 multiplied by the total efficiency ε for the experimental apparatus to observe such events. In practice L itself is measured using the detection rate ϵR for a process for which σ in equation 10.1 is both well-known and sufficiently high to achieve a small statistical uncertainty for L . In quantum electrodynamics (QED) the crosssection $\sigma(e^+e^- \to \mu^+\mu^-)$, for the process depicted below in figure 10.2 and described subsequently, is one of the simplest to calculate and is well known. It was used as a reference point for e^+e^- colliders in the 1970s in order to measure the cross-section for hadronic final state production relative to muon pairs as a function of centre-of-mass energy.

The approach taken in this chapter is to begin with *observable* quantities in HEP experiments, writing down the general expression for the cross-section as below, and then show how this is related to calculations in QFT through computation of the S-matrix. This in turn will lead to consideration of the elementary interaction terms in the Lagrangian and a description of the procedure of calculation aided by Feynman diagrams and rules. We begin then with the differential cross-section $d\sigma(e^+e^- \to X)$ for a general process at an e^+e^- collider experiment (see for example [70] p.106):

$$
d\sigma = \frac{1}{4E_1E_2|\mathbf{v}_1 - \mathbf{v}_2|} |\mathcal{M}_{fi}|^2 (2\pi)^4 \delta^4 \left(\sum_f p_f - \sum_i p_i\right) \prod_f \frac{d^3 \mathbf{p}_f}{(2\pi)^3 2E_f} \qquad (10.3)
$$

where $E_{1,2}$ and $v_{1,2}$ are the energy and 3-velocity of the particles in the two opposing incoming beams, E_f and p_f are the energy and 3-momentum for each final state particle and p_i and p_f are the 4-momenta of each initial and final state particle $(i = 1, 2$ and $f = 1, \ldots, N_f$, all in the centre-of-mass frame. A further combinatoric factor may be needed, for example to account for initial or final state particle spins for an unpolarised cross-section, as for equation 10.11 below, or a factor of $1/n!$ for a total cross-section with n identical particles in the final state.

The only non-kinematic quantity in equation 10.3 is the transition amplitude \mathcal{M}_{fi} (where here the subscript f i labels the overall process) which contains the dynamics of the transformation between the initial and final particle states. The relativistic state normalisation of equations 10.17 and 10.18 below will be employed and is consistent with the Lorentz invariance of \mathcal{M}_{fi} as constructed in the following section. Everything to the right of $|\mathcal{M}_{fi}|^2$ in equation 10.3 is the 'Lorentz invariant phase space' term dΦ for the final state. The only factor on the right-hand side of equation 10.3 which is not Lorentz invariant is the initial state flux factor $(4E_1E_2|\mathbf{v}_1-\mathbf{v}_2|)^{-1}$, however this term is invariant under Lorentz boosts along the beam direction. Indeed $d\sigma$, on the left-hand side of this equation, transforms as a two-dimensional cross-sectional area under Lorentz transformations. When composed with the luminosity L of equation 10.2 in equation 10.1 the event rate R exhibits a simple special relativistic time-dilation effect under a change of Lorentz frame, as for any physical 'clock'.

The cross-section σ can be considered as the effective cross-sectional area within scattering range of each particle in the beam, or as the number of scattering events per unit time, per unit volume, per unit flux density of the incoming beams. Indeed the above cross-section formula can be calculated by considering the interaction to take place over a finite time period T in a finite spatial volume V , which contains purely free fields in the limits $t \to \pm \infty$ relative to the interaction time around $t = 0$. Factors of T and V cancel in the final result of equation 10.3. Alternatively, a more detailed approach may be followed in which the incoming states are modelled as wave packets localised in space ([70] pp.102–106). In this case the final result for $d\sigma$ is independent of the shape of the wave packets.

For either way of deriving this formula the transition amplitude \mathcal{M}_{fi} itself in equation 10.3 is calculated for the idealised case of 'in' and 'out' plane wave states of definite momentum extending throughout spacetime. The resemblance of these states to the concept of a particle is somewhat limited due to the absence of localisation, however their use in the determination of \mathcal{M}_{fi} , and in turn the cross-section for particle interactions, may be followed pragmatically.

The transition amplitude is determined by the matrix element between the initial e^+e^- free field state represented by $|p_1, p_2\rangle$ in for $t \to -\infty$ and a particular final state $|q_1, q_2 \dots q_{N_f} \rangle$ out for $t \to +\infty$, in the respective 'in' and 'out' Fock space bases for the incoming and outgoing particles states. While neither of these two bases are simply related to a further Fock basis for interacting fields, since they both represent the free-field case they are isomorphic to each other. This isomorphism is described by the unitary operator S, connecting the 'in' and 'out' bases such that $|P\rangle_{\text{in}} = S|P\rangle_{\text{out}}$ with P denoting any state. Unitarity is required here to conserve probabilities, with the transition probability being determined by the squared modulus of the amplitude, that is $|\mathcal{M}_{fi}|^2$, by a basic *postulate* of quantum theory, as discussed further below.

This situation can be expressed in a single 'interaction picture' basis I with an initial state $|i\rangle_I$ evolving in time from $t = -\infty$, through interactions as described by the S-matrix, to be measured in the final state $|f\rangle_I$ at $t = +\infty$ with a probability determined by the matrix element:

$$
S_{fi} = \frac{\text{out}}{\f_i}\rangle_{\text{in}} = \frac{\text{out}}{\f_i}S_i\rangle_{\text{out}} = \frac{\text{in}}{\f_i}S_i\rangle_{\text{in}} \equiv \frac{I}{f_i}S_i\rangle_I \tag{10.4}
$$

where we subsequently drop the subscripts I since the interaction picture, described further in the following section, will be used throughout. The S-matrix can be written:

$$
S = \mathbf{1} + iT \tag{10.5}
$$

where 1 represents the trivial identity operation and iT , with the conventional $i = \sqrt{-1}$ factor, represents the non-trivial interaction part of the S-matrix. It is this latter part $iT = S - 1$ which is of most interest and its matrix element between the initial and final states can be written, with $p_I = \sum_i p_i$ and $p_F = \sum_f p_f$, as:

$$
\langle f|iT|i\rangle = (2\pi)^4 \delta^4(p_F - p_I) i\mathcal{M}_{fi}
$$
 (10.6)

which isolates the transition amplitude \mathcal{M}_{fi} . Expressions for $i\mathcal{M}_{fi}$ will later be associated with Feynman diagrams which in turn may be obtained directly from the Lagrangian for the field theory. Hence the transition amplitude is identified from the matrix element in equation 10.6 by factoring out an ever-present total 4-momentum conserving delta function. Such delta functions arise as a consequence of treating the external particles as idealised states of definite momentum.

In deriving the expression for the cross-section a factor of $|\langle f|S|i\rangle|^2$ is incorporated which hence contributes two factors of $(2\pi)^4 \delta^4(p_F - p_I)$; one of which may be interpreted as the spacetime interaction volume VT and cancels with other factors of V and T in the final result of equation 10.3. In this expression for the differential cross-section the surviving delta function is included in the Lorentz invariant phase space $d\Phi$ when composed with the final factor of $\prod_f d^3\mathbf{p}_f/((2\pi)^3 2E_f)$.

This latter object is a statistical factor representing the density of final states in 'small' regions of phase space between p_f and $p_f + d^3 p_f$ for each outgoing particle. These regions are constrained by the delta function for the total 4-momentum when integrating over the final state degrees of freedom of the differential cross-section. The factors of $1/E_f$ arise in the phase space from the relativistic state normalisation of equation 10.17. The first factor in equation 10.3 arises in a related way and represents the flux density for the incoming colliding particle beams.

The overall expression is such that the cross-section σ essentially represents the probability of individual particle on particle interactions and is hence correctly normalised for equations 10.1 and 10.2. Bearing in mind these latter equations together with equation 10.3 the total differential event rate can be written:

$$
dR = \left(\frac{f n_+ n_-}{A} \cdot \frac{1}{4E_1 E_2 |\mathbf{v}_1 - \mathbf{v}_2|}\right) \cdot |\mathcal{M}_{fi}|^2 \cdot \left((2\pi)^4 \delta^4 (p_F - p_I) \prod_f \frac{d^3 p_f}{(2\pi)^3 2E_f}\right)
$$
\n(10.7)

as a composition of three parts. The factor in the first brackets contributes to the likelihood of events occurring given the properties of the incoming beams from a purely statistical point of view. In a similar way the Lorentz invariant phase space $d\Phi$ in the final set of large brackets represents the range of possible outgoing state configurations as a further natural statistical factor. These two factors hence arise out of consideration of the basic classical laws of probability, essentially with the probability simply being proportional to the sum of the 'number of ways' that something can happen. A further combinatoric factor is possible, such as a sum over outgoing particle spin states, as alluded to after equation 10.3. Observations made in the experiment depend also on the efficiency ε of the detector, as alluded to after equation 10.2. Further, classical statistical methods are used to analyse the data to complete the measurements of physical quantities with the results presented along with their statistical and systematic uncertainties.

The point of this discussion is to highlight the contrast between this list of classical probabilistic factors and the middle term $|\mathcal{M}_{fi}|^2$ of equation 10.7 with which they are composed and which has rather different characteristics. Historically this final factor originated from non-relativistic quantum mechanics for which the transition probability for a state described by the normalised wavefunction $\Psi(x, t)$ to be measured in the normalised eigenstate $\Phi_i(x,t)$ is represented by the squared modulus $|A|^2$ of the amplitude $A = \langle \Phi_i(\mathbf{x}, t)|\Psi(\mathbf{x}, t)\rangle$, that is the overlap integral

$$
A = \int \Phi_i^*(\mathbf{x}, t) \Psi(\mathbf{x}, t) d^3 \mathbf{x}
$$
 (10.8)

This construction of a *probability* is a postulate of quantum theory, apparently quite different to the notion of probability as being a measure of the 'number of ways' that something can happen, as encountered in all non-quantum walks of life. This form of quantum probability was itself originally introduced to represent the likelihood for locating a particle at the spatial position x by the value of $|\Psi(x,t)|^2$, and dates from the formative years of quantum theory in the mid 1920s.

As an example the production of muon pairs in the process $e^+e^- \to \mu^+\mu^-$, as depicted in figure 10.2, will be considered. The cross-section formula of equation 10.3 simplifies for this case of scattering to a two-particle final state. The δ^4 function constrains $|\mathbf{p}_f|$ in the centre-of-mass frame to the same fixed value for each outgoing particle and, taking the approximation that all particle masses are sufficiently below the centre-of-mass energy \sqrt{s} and hence can be neglected, the differential cross-section reduces to:

$$
\frac{d\sigma}{d\Omega} = \frac{|\mathcal{M}_{fi}|^2}{64\,\pi^2\,s} \tag{10.9}
$$

where Ω is the solid angle within which the μ^- is produced. For the unpolarised process $e^+e^- \to \mu^+\mu^-$ there is a further combinatoric factor corresponding to an average over

Figure 10.2: A schematic diagram for the transition from an e^+e^- incoming state to $a \mu^+ \mu^-$ outgoing state. In the text the purely QED process is considered.

the initial electron spin states and sum over the final muon spin states, with $|\mathcal{M}_{fi}|^2$ above then replaced by:

$$
\frac{1}{4} \sum_{\text{spins}} |\mathcal{M}_{fi}|^2 = e^4 (1 + \cos^2 \theta) \tag{10.10}
$$

This is for the lowest non-trivial order of perturbation in the QFT, for which the unpolarised differential cross-section is hence given by ([70] pp.8 and 137):

$$
\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4s} (1 + \cos^2 \theta) \tag{10.11}
$$

with fine structure constant $\alpha = e^2/4\pi \simeq 1/137$, where e is the charge of the electron, conventionally taken to be negative. In equation 10.11 s is the square of the centre-ofmass energy and θ is the polar angle of the final state μ^{-} , as depicted in figure 10.2. In deriving equation 10.11 it is assumed not only that $s \gg m_{\mu^-}^2$, and hence the lepton masses are neglected, but also that s is sufficiently below M_Z^2 , so that a contribution from the weak interaction can also be neglected. In particular this means that the centre-of-mass energy is assumed to be somewhat lower than that for the experiment in figure 10.1, which operated on the Z^0 resonance. In this case, for a purely QED process, the lowest-order calculation can be associated with the Feynman diagram of figure 10.3 featuring an intermediate 'virtual photon'.

On integrating over the solid angle the total cross-section is found to be:

$$
\sigma(e^+e^- \to \mu^+\mu^-) = \frac{4\pi\alpha^2}{3s} \tag{10.12}
$$

This cross-section, based on the leading order process depicted by the Feynman diagram in figure 10.3 agrees with observations in HEP experiments to within about 10%. Most of this discrepancy is accounted for by the next order in perturbation theory ([70] p.8), with excellent agreement between the data and theory for a more thorough calculation.

Figure 10.3: Feynman diagram for the process $e^+e^- \rightarrow \mu^+\mu^-$ to lowest order in QED perturbation theory. In such diagrams the external lines on the left-hand side represent incoming particle states, while those on the right-hand side represent outgoing particles. (The direction of the arrows on the external lines is explained under 'item 3' in the discussion of Feynman diagrams in section 10.5, while the causal structure of the two vertices will also be discussed later, for example alongside figures 10.5(b) and 10.6 in section 10.4.)

The $\cos^2 \theta$ angular dependence in equation 10.11 arises in $|\mathcal{M}_{fi}|^2$ from the spin- $\frac{1}{2}$ property of the initial and final state particles. The actual calculations involving interaction processes in QED are made significantly more complicated by the presence of Lorentz spinor and vector fields, with the derivation of the right-hand side of equation 10.10 for example being non-trivial. Since we are here interested in the probability interpretation of the transition amplitude \mathcal{M}_{fi} in the following section we consider in detail a simpler, but closely analogous, model based on interacting scalar fields in order to extract the essential mathematical structure that is used in the calculation of such probabilities in a more transparent manner.

10.3 Transition Amplitudes

For the remainder of this chapter we consider a scalar model for an interacting field theory with three scalar fields, including one real field $\phi(x)$ and two complex fields $\mathcal{X}(x)$ and $\mathcal{Y}(x)$, that is with a total of five real field components, with the quanta of the complex fields being interpreted as charged particles. (The analogy with the HEP process described in the previous section being constructed here may be briefly summarised by comparing the Feynman diagrams in figures 10.3 and 10.4 for the respective lowest-order calculations). Both real and complex free fields can be expressed in terms of a corresponding annihilation and creation operator expansion which for the fields $\phi(x)$ and $\mathcal{X}(x)$ can be written as:

$$
\hat{\phi}(x) = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} \left(a(\mathbf{p}) e^{-i\mathbf{p}\cdot x} + a^{\dagger}(\mathbf{p}) e^{+i\mathbf{p}\cdot x} \right) \tag{10.13}
$$

$$
\hat{\mathcal{X}}(x) = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} \left(b_{\mathcal{X}}(\mathbf{p}) e^{-ip \cdot x} + d_{\mathcal{X}}^{\dagger}(\mathbf{p}) e^{+ip \cdot x} \right) \tag{10.14}
$$

$$
\hat{\mathcal{X}}^{\dagger}(x) = \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} \left(d_{\mathcal{X}}(\mathbf{p}) e^{-ip \cdot x} + b_{\mathcal{X}}^{\dagger}(\mathbf{p}) e^{+ip \cdot x} \right) \tag{10.15}
$$

with $p^0 = \omega_{\bf p} = \pm \sqrt{{\bf p}^2 + m^2}$ in all three expressions. The mass m for each field will be associated with the corresponding particle states which are identified in the following. In QFT the Fourier field coefficients such as $a(\boldsymbol{p})$ and $a^{\dagger}(\boldsymbol{p})$ in equation 10.13 are taken to be linear operators acting on the Fock space of particle states. The 'quantisation' of the free field is completed by imposing commutation relations on these operators:

$$
[a(\mathbf{p}), a^{\dagger}(\mathbf{p}')] = (2\pi)^3 \delta^3(\mathbf{p} - \mathbf{p}')[a(\mathbf{p}), a(\mathbf{p}')] = 0, \qquad [a^{\dagger}(\mathbf{p}), a^{\dagger}(\mathbf{p}')] = 0
$$
\n(10.16)

By imposing these relations, largely by analogy with the quantum mechanical simple harmonic oscillator, the spectrum of states possesses a ladder structure with $a^{\dagger}(\boldsymbol{p})$ interpreted as *creating* a particle of momentum p and $a(p)$ annihilating such a state. Hence in turn the $e^{\pm ip\cdot x}$ Fourier modes of equation 10.13 are associated with particle quanta of mass m the creation or annihilation of which are attributed to the free scalar field $\phi(x)$. This structure marks an attempt to achieve direct contact with the concept of particles by modelling their discrete nature, although the associated Fourier modes are clearly not localised in space. With the vacuum represented by the state $|0\rangle$ in the Fock space the annihilation operator acts as $a(p)|0\rangle = 0$ while a single particle state $|p\rangle$ is created as:

$$
|\mathbf{p}\rangle = \sqrt{2\omega_{\mathbf{p}}} \, a^{\dagger}(\mathbf{p})|0\rangle \tag{10.17}
$$

such that, given the vacuum normalisation $\langle 0|0 \rangle = 1$, we have:

$$
\langle p|q\rangle = 2\omega_p (2\pi)^3 \delta^3(p-q) \tag{10.18}
$$

which is Lorentz invariant, justifying the choice of normalisation factor employed in equation 10.17.

Analogous relations to equations 10.16 hold for each pair of operators, namely $b_{\mathcal{X}}(p), b_{\mathcal{X}}^{\dagger}(p)$ and $d_{\mathcal{X}}(p), d_{\mathcal{X}}^{\dagger}(p)$, for the $\hat{\mathcal{X}}(x)$ field of equations 10.14 and 10.15. These two pairs of operators, with the corresponding two sets of commutators, are interpreted as generating two types of particle states, with $b_{\mathcal{X}}^{\dagger}(\boldsymbol{p})$ and $b_{\mathcal{X}}(\boldsymbol{p})$ respectively creating and annihilating \mathcal{X}^- particles, and similarly with $d_{\mathcal{X}}^{\dagger}(\mathbf{p})$ and $d_{\mathcal{X}}(\mathbf{p})$ for \mathcal{X}^+ antiparticles. (The U(1) charges associated with the particles and antiparticles are actually $+1$ and −1 respectively, however the charge unit is chosen to be negative. This is by analogy with the convention adopted for the electron field, with $e < 0$ as described after equation 10.11, with negatively charged particles and positively charged antiparticles, that is positrons).

The field $\hat{y}(x)$ and its conjugate $\hat{y}^{\dagger}(x)$ can be similarly expanded in terms of corresponding creation and annihilation operators by direct analogy with equations 10.14 and 10.15 and $\hat{\mathcal{V}}^{\pm}$ particle states similarly described. The normalisation of these single particle states follows the convention of equation 10.17 and hence we define the creation operators:

$$
\hat{B}_{\mathcal{X}}^{\dagger}(\boldsymbol{p}) = \sqrt{2\omega_{\boldsymbol{p}}} \, b_{\mathcal{X}}^{\dagger}(\boldsymbol{p}) \qquad \text{with} \qquad \hat{B}_{\mathcal{X}}^{\dagger}(\boldsymbol{p}) |0\rangle = |\boldsymbol{p}_{\mathcal{X}}-\rangle \tag{10.19}
$$

$$
\hat{D}_{\mathcal{X}}^{\dagger}(\mathbf{p}) = \sqrt{2\omega_{\mathbf{p}}} d_{\mathcal{X}}^{\dagger}(\mathbf{p}) \quad \text{with} \quad \hat{D}_{\mathcal{X}}^{\dagger}(\mathbf{p})|0\rangle = |\mathbf{p}_{\mathcal{X}^{+}}\rangle \tag{10.20}
$$

$$
\hat{B}_{\mathcal{Y}}^{\dagger}(\mathbf{p}) = \sqrt{2\omega_{\mathbf{p}}} b_{\mathcal{Y}}^{\dagger}(\mathbf{p}) \quad \text{with} \quad \hat{B}_{\mathcal{Y}}^{\dagger}(\mathbf{p}) |0\rangle = |\mathbf{p}_{\mathcal{Y}}-\rangle \tag{10.21}
$$
\n
$$
\hat{D}_{\mathcal{Y}}^{\dagger}(\mathbf{p}) = \sqrt{2\omega_{\mathbf{p}}} d_{\mathcal{Y}}^{\dagger}(\mathbf{p}) \quad \text{with} \quad \hat{D}_{\mathcal{Y}}^{\dagger}(\mathbf{p}) |0\rangle = |\mathbf{p}_{\mathcal{Y}}-\rangle \tag{10.22}
$$

$$
\hat{D}_{\mathcal{Y}}^{\dagger}(\boldsymbol{p}) = \sqrt{2\omega_{\boldsymbol{p}}} d_{\mathcal{Y}}^{\dagger}(\boldsymbol{p}) \quad \text{with} \quad \hat{D}_{\mathcal{Y}}^{\dagger}(\boldsymbol{p}) |0\rangle = |\boldsymbol{p}_{\mathcal{Y}^+}\rangle \quad (10.22)
$$

with corresponding conjugate annihilation operators. These may be considered as subcomponents of the operator fields $\mathcal{X}(x)$ and $\mathcal{Y}(x)$, as for the operator in equation 10.17 with respect to the field $\phi(x)$. We stress that here in sections 10.2–10.5 we are describing the standard constructions of a quantum field theory (as described in more much detail in [10, 70, 71] for example) and in the following chapter we shall need to describe how the corresponding elements arise in the context of the new theory presented in this paper.

The Lagrangian for the model under consideration here consists of three free field parts, each of which is essentially a Klein-Gordon Lagrangian, for the fields $\phi(x)$, $\hat{\mathcal{X}}(x)$ and $\hat{\mathcal{Y}}(x)$, with mass parameters m_{ϕ} , $m_{\mathcal{X}}$ and $m_{\mathcal{Y}}$ respectively, together with an interaction part \mathcal{L}_{int} consisting of polynomial functions of the fields:

$$
\mathcal{L} = \mathcal{L}_{\phi} + \mathcal{L}_{\mathcal{X}} + \mathcal{L}_{\mathcal{Y}} + \mathcal{L}_{int}
$$

with
$$
\mathcal{L}_{\phi} = \frac{1}{2} \partial_{\mu} \hat{\phi} \, \partial^{\mu} \hat{\phi} - \frac{1}{2} m_{\phi}^{2} \, \hat{\phi}^{2}
$$

$$
\mathcal{L}_{\mathcal{X}} = \partial_{\mu} \hat{\mathcal{X}}^{\dagger} \, \partial^{\mu} \hat{\mathcal{X}} - m_{\mathcal{X}}^{2} \, \hat{\mathcal{X}}^{\dagger} \hat{\mathcal{X}}
$$

$$
\mathcal{L}_{\mathcal{Y}} = \partial_{\mu} \hat{\mathcal{Y}}^{\dagger} \, \partial^{\mu} \hat{\mathcal{Y}} - m_{\mathcal{Y}}^{2} \, \hat{\mathcal{Y}}^{\dagger} \hat{\mathcal{Y}}
$$

and
$$
\mathcal{L}_{int} = -g \hat{\phi} \, \hat{\mathcal{X}}^{\dagger} \hat{\mathcal{X}} - g \hat{\phi} \hat{\mathcal{Y}}^{\dagger} \hat{\mathcal{Y}}
$$
(10.23)

where g is the interaction coupling constant. Since the Lagrangian must be a real function a complex field appears in each term symmetrically with its conjugate field; for example $\mathcal{L}_{\mathcal{X}}$ contains the mass term $-m_{\mathcal{X}}^2 \hat{\mathcal{X}}^{\dagger} \hat{\mathcal{X}}$. It is the invariance of this total Lagrangian under the global U(1) symmetry with $\hat{\mathcal{X}} \to e^{i\alpha} \hat{\mathcal{X}}$ and $\hat{\mathcal{X}}^{\dagger} \to e^{-i\alpha} \hat{\mathcal{X}}^{\dagger}$ (and similarly for the complex $\mathcal{Y}(x)$ field) that implies a conserved U(1) charge as described above, consistent with Noether's theorem as briefly reviewed alongside equation 3.100 in section 3.5.

The simple QFT model described here is not a gauge theory and in equation 10.23 the interaction terms are added by hand. By contrast in QED or scalar electrodynamics the coupling of the charged fields to the electromagnetic field $A_u(x)$ is induced by the requirement of a *local* $U(1)$ gauge invariance of the Lagrangian, as also described in section 3.5 and exemplified in the final term of equation 3.96, although an arbitrary coupling constant can still be employed. In the Standard Model non-Abelian gauge theories are also incorporated through such expressions as for example in equations 7.39 and 7.40 of section 7.2. In all cases such Lagrangian terms imply interactions since the fields mutually influence one another in equations derived from the principle of extremal action. Here with the additional interaction terms of equation 10.23 the Euler-Lagrange equations of motion from equation 3.89, derived by varying $\hat{\phi}(x)$, $\hat{\mathcal{X}}^{\dagger}(x)$, $\hat{\mathcal{X}}(x)$, $\hat{\mathcal{Y}}^{\dagger}(x)$ and $\hat{\mathcal{Y}}(x)$ respectively as five independent fields subject to the constraint $\delta \int \mathcal{L} d^4x = 0$ (in a flat spacetime) are non-linear in the fields:

$$
(\Box + m_{\phi}^2)\hat{\phi}(x) = -g\hat{\mathcal{X}}^{\dagger}\hat{\mathcal{X}} - g\hat{\mathcal{Y}}^{\dagger}\hat{\mathcal{Y}} \qquad (10.24)
$$

$$
(\Box + m_{\mathcal{X}}^2)\hat{\mathcal{X}}(x) = -g\hat{\phi}\hat{\mathcal{X}} \quad \text{and with} \quad \hat{\mathcal{X}} \to \hat{\mathcal{X}}^{\dagger} \quad (10.25)
$$

 $(\Box + m_{\mathcal{Y}}^2)\hat{\mathcal{Y}}(x) = -g\hat{\phi}\hat{\mathcal{Y}}$ and with $\hat{\mathcal{Y}} \to \hat{\mathcal{Y}}^{\dagger}$ (10.26)

and impossible to solve exactly. Neglecting the \mathcal{L}_{int} terms in equation 10.23, that is in the limit for the coupling $g \to 0$, each of equations 10.24–10.26 reduces to the free Klein-Gordon equation for which fields of the form in equations 10.13–10.15 provide exact general solutions.

Equations 10.24–10.26 correspond to the 'Heisenberg picture' in which all of the time dependence is ascribed to the operator fields, while for the 'Schrödinger' picture' the time dependence would apply purely to the states. In all cases in quantum theory the time evolution is determined by the Hamiltonian operator H which may be expressed as the sum of a free field part H_0 and in interaction part H_{int} . In the 'interaction picture' the time dependence of all operators is determined by H_0 only, with the corresponding evolution of free operator fields such as $\phi(x)$ then readily handled (as for equation 10.13 as a solution of equation 10.24 with $g = 0$) while H_{int} governs the evolution of the states. In the interaction picture the aim is to express the transition amplitude, and hence the scattering probability, purely in terms of free fields. (This structure will be significant for making a link with the conceptual picture of the present theory, as will be discussed in 'item 3)' of section 11.2 for example.)

For the model QFT under consideration here the evolution of the states is closely related to the interaction terms of equation 10.23. Indeed if there are no time derivatives in the Lagrangian density \mathcal{L}_{int} the interaction Hamiltonian H_{int} can be written simply as:

$$
H_{\rm int} = \int d^3 x \, \mathcal{H}_{\rm int} = -\int d^3 x \, \mathcal{L}_{\rm int} \tag{10.27}
$$

In the interaction picture the initial state $|i\rangle$ evolves according to the unitary operator U into the state $|\Psi(t)\rangle \equiv U(t, -\infty)|i\rangle$ at time t, with the equation of motion:

$$
i\frac{d}{dt}|\Psi(t)\rangle = H_{\text{int}}(t)|\Psi(t)\rangle
$$
\n(10.28)

with the Hamiltonian $H_{\text{int}}(t)$ defining the time evolution. The scattering amplitude is obtained from the overlap of the state $|\Psi(t)\rangle$ evolved to $t = +\infty$ with the given final state $|f\rangle$, that is the matrix element:

$$
S_{fi} = \langle f|U(+\infty, -\infty)|i\rangle \tag{10.29}
$$

In the interaction picture the 'initial value problem' for $U(t, -\infty)$ is posed by the initial condition $U(-\infty, -\infty) = 1$ together with the equation of motion obtained directly from equation 10.28:

$$
i\frac{d}{dt}U(t, -\infty) = H_{\text{int}}(t)U(t, -\infty)
$$
\n(10.30)

As an Hermitian operator the Hamiltonian H acts as the infinitesimal generator of a one-parameter unitary group. This unitary symmetry is employed in QFT to model the conservation of probability in scattering processes. A solution to equation 10.30, which might naively be expected to take the form $U(t, -\infty) \sim e^{-itH_{\text{int}}}$, when taking into account the time dependence and operator action can be obtained by iteration (and checked by direct substitution into equation 10.30) and then restructured using the time-ordered product T of operators. Considering the evolution for any time interval from t_0 to t it is found that:

$$
U(t, t_0) =
$$

\n
$$
\mathbf{1} + (-i) \int_{t_0}^t dt_1 H_{int}(t_1) + (-i)^2 \int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 H_{int}(t_1) H_{int}(t_2)
$$

\n
$$
+ (-i)^3 \int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 \int_{t_0}^{t_2} dt_3 H_{int}(t_1) H_{int}(t_2) H_{int}(t_3) + \dots \qquad (10.31)
$$

\n
$$
= \mathbf{1} - i \int_{t_0}^t dt_1 T[H_{int}(t_1)] - \frac{1}{2} \int_{t_0}^t dt_1 \int_{t_0}^t dt_2 T[H_{int}(t_1) H_{int}(t_2)]
$$

$$
+ \frac{i}{3!} \int_{t_0}^t dt_1 \int_{t_0}^t dt_2 \int_{t_0}^t dt_3 T[H_{\rm int}(t_1)H_{\rm int}(t_2)H_{\rm int}(t_3)] + \dots \qquad (10.32)
$$

$$
= \sum_{n=0}^{\infty} \frac{(-i)^n}{n!} \int_{t_0}^t dt_1 \int_{t_0}^t dt_2 \dots \int_{t_0}^t dt_n T[H_{\rm int}(t_1) H_{\rm int}(t_2) \dots H_{\rm int}(t_n)] \tag{10.33}
$$

$$
= T[\exp(-i\int_{t_0}^t dt' H_{\rm int}(t'))]
$$
\n(10.34)

The factor of $\frac{1}{2}$ appears on the right-hand side in equation 10.32 since the extended integral does everything that is needed for the corresponding term in equation 10.31 twice. This generalises to the factor of $1/n!$ in equation 10.33 for the corresponding combinatorial over-counting for the higher-order terms. The final expression above is a useful shorthand notation for equation 10.33. The S-matrix, as introduced in equation 10.4, is then defined, on taking $t_0 = -\infty$ and $t = +\infty$, as the unitary operator:

$$
S = U(+\infty, -\infty) = Te^{-i\int_{-\infty}^{+\infty} dt \, H_{\rm int}(t)} \tag{10.35}
$$

which appeared in equation 10.29 for the transition amplitude for a particular process. Hence the S-matrix contains the information needed to calculate the probability of scattering from one plane wave state to another. In the interaction picture the basis for the external plane wave states is expressed in terms of the same sets of annihilation and creation operators which provide the coefficients of the Fourier expansions of the fields $\phi(x)$, $\mathcal{X}(x)$ and $\mathcal{Y}(x)$, of equations 10.13–10.15 for example, through which in turn H_{int} and hence the S-matrix is expressed in equation 10.35, containing all the information about the interaction.

For the case of $\mathcal{L}_{\text{int}} = 0$ the interaction Hamiltonian is zero and trivially $S = 1$. In the general case with $\mathcal{L}_{int} \neq 0$ equations 10.29 and 10.35 together describe a timeordered chain of field operations between the initial and final states. This time ordering is explicit in equation 10.31 owing to the temporal limits for each integral and the order of the interaction Hamiltonian operators in the integrand. Essentially the Smatrix represents everything that can happen at all intermediate times between the initial and final free states according to the \mathcal{L}_{int} terms. As will be described below, when the calculation is restructured with a more symmetric set of temporal limits for each integral in equation 10.32 the time ordering with T ensures causal relations are maintained through this chain, with Hamiltonian field operators acting in the correct sequence with intermediate field states first being created before being annihilated.

Given this iterative solution for $U(t, t_0)$ described in equations 10.31–10.34, for $(t, t_0) = (+\infty, -\infty)$, the assumption of *perturbation theory* is that the first few terms provide a good approximation to the exact full expression. This may be possible if the magnitude of the first few terms in equation 10.29 decreases (or if there are cancellations between large terms) with increasing order n , as defined in equation 10.33, which will generally be the case if the coupling constant, such as q in equation 10.23 or α in equation 10.11 for the case of QED, is sufficiently small. Even in this case many terms will lead to divergent integrals in QFT which will need to be accounted for by renormalisation. However, even this does not imply that the expression for $U(+\infty, -\infty)$, and in turn S_{fi} , will converge for large n. Nevertheless the first few terms of perturbation theory do lead to calculations that have a well-defined meaning in that they generate quantities that can be compared with experiment, as is the case for muon pair production in the Standard Model as described towards the end of the previous section.

By analogy with the real process $e^+e^- \to \mu^+\mu^-$ here for the scalar field model we consider the scattering process $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$, that is, using equations 10.19– 10.22, between:

the initial state
$$
\mathcal{X}^+ \mathcal{X}^-
$$
: $|i\rangle = \hat{D}_{\mathcal{X}}^{\dagger}(\mathbf{p}_2) \hat{B}_{\mathcal{X}}^{\dagger}(\mathbf{p}_1) |0\rangle$ at $t = -\infty$
and final state $\mathcal{Y}^+ \mathcal{Y}^-$: $|f\rangle = \hat{D}_{\mathcal{Y}}^{\dagger}(\mathbf{q}_2) \hat{B}_{\mathcal{Y}}^{\dagger}(\mathbf{q}_1) |0\rangle$ at $t = +\infty$

Hence for the process $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$ under consideration in the scalar field model the transition amplitude of equation 10.29, via equation 10.35, can be written:

$$
S_{fi} = \langle 0 | \hat{B}_{\mathcal{Y}}(\boldsymbol{q}_1) \hat{D}_{\mathcal{Y}}(\boldsymbol{q}_2) T[\exp\big(-i \int_{-\infty}^{+\infty} dt \, H_{\rm int}(t)\big)] \, \hat{D}_{\mathcal{X}}^{\dagger}(\boldsymbol{p}_2) \hat{B}_{\mathcal{X}}^{\dagger}(\boldsymbol{p}_1) |0\rangle \qquad (10.36)
$$

where H_{int} is expressed in terms of a polynomial in the interaction picture operator fields $\phi(x)$, $\mathcal{X}(x)$ and $\mathcal{Y}(x)$. These are free-fields evolving simply under H_0 and can be expanded in terms of creation and annihilation operators, that is by substituting the free fields of equations 10.13–10.15 (as well as for $\hat{y}(x)$ and $\hat{y}^{\dagger}(x)$) into equations 10.23 and 10.27 in turn, hence linking the initial and final states in equation 10.36. The general problem then in the interaction picture is to evaluate terms of the form:

$$
\int dt_1, dt_2 \dots dt_n T[H_{\rm int}(t_1) H_{\rm int}(t_2) \dots H_{\rm int}(t_n)] \tag{10.37}
$$

between the external particle Fock states. This calculation can be somewhat simplified by noting that these terms, together with the initial and final state creation operators in equation 10.36, are sandwiched between vacuum states which have the property $a(p)|0\rangle = 0$ and $\langle 0|a^{\dagger}(p) = 0$ for an arbitrary annihilation operator $a(p)$ and its conjugate. Hence the goal is to use the commutation relations for such operators, for example equation 10.16, to extract the residual non-zero terms from equation 10.36. This is achieved by decomposing the time-ordered product into a combination of normalordered terms and contractions, which takes a simple form for the product of two field values: ^p–—–^q

$$
T(\hat{\phi}(x)\hat{\phi}(y)) = \quad : \hat{\phi}(x)\hat{\phi}(y): + \hat{\phi}(x)\hat{\phi}(y) \tag{10.38}
$$

Here the final term is the contraction which can be defined as the difference between the time-ordered product and the normal-ordered product of the field values. The normal-ordered product, denoted by the colon braces : \hat{F} :, is defined such that all annihilation operators are placed to the right of all creation operators in each term, and hence $\langle 0 | : \hat{F} : |0 \rangle = 0$, that is the vacuum expectation value (v.e.v.) for the normal-ordered product of any collection \hat{F} of fields is zero. The contracted product in equation 10.38 is a scalar multiple of the identity operator 1, as can be shown by considering the case for $x^0 > y^0$ and for $x^0 < y^0$. For example:

$$
\hat{\phi}(x)\hat{\phi}(y) = T(\hat{\phi}(x)\hat{\phi}(y)) - \hat{\phi}(x)\hat{\phi}(y); \text{ which for the } x^{0} > y^{0} \text{ case:}
$$
\n
$$
= \int \frac{d^{3}p}{(2\pi)^{3}} \frac{d^{3}q}{(2\pi)^{3}} \frac{1}{\sqrt{2\omega_{p}}} \frac{1}{\sqrt{2\omega_{q}}} \left\{ (a(p)e^{-ip\cdot x} + a^{\dagger}(p)e^{+ip\cdot x}) (a(q)e^{-iq\cdot y} + a^{\dagger}(q)e^{+iq\cdot y}) - (a(p)a(q)e^{+ip\cdot x}e^{-iq\cdot y} + a^{\dagger}(q)a(p)e^{-ip\cdot x}e^{+iq\cdot y} + a^{\dagger}(p)a(q)e^{+ip\cdot x}e^{-iq\cdot y} + a^{\dagger}(p)a(q)e^{+ip\cdot x}e^{-iq\cdot y} + a^{\dagger}(p)a(q)e^{+ip\cdot x}e^{+iq\cdot y} \right\}
$$
\n
$$
= \int \frac{d^{3}p}{(2\pi)^{3}} \frac{d^{3}q}{(2\pi)^{3}} \frac{1}{\sqrt{2\omega_{p}}} \frac{1}{\sqrt{2\omega_{q}}} \left(a(p)a^{\dagger}(q) e^{-ip\cdot x} e^{+iq\cdot y} - a^{\dagger}(q)a(p) e^{-ip\cdot x} e^{+iq\cdot y} \right)
$$
\n
$$
= \int \frac{d^{3}p}{(2\pi)^{3}} \frac{d^{3}q}{(2\pi)^{3}} \frac{1}{\sqrt{2\omega_{p}}} \frac{1}{\sqrt{2\omega_{q}}} [a(p), a^{\dagger}(q)] e^{-ip\cdot x} e^{+iq\cdot y}
$$
\n
$$
= \int \frac{d^{3}p}{(2\pi)^{3}} \frac{d^{3}q}{(2\pi)^{3}} \frac{1}{\sqrt{2\omega_{p}}} \frac{1}{\sqrt{2\omega_{q}}} (2\pi)^{3} \delta^{3}(p-q) e^{-ip\cdot x} e^{+iq\cdot y}
$$
\n
$$
= \int \frac{d^{3}p}{(2\pi)^{3}} \frac{1}{2\omega_{p}} e^{-ip\cdot (x-y)} \qquad (\text{for } x^{0} > y^{0}) \qquad (10.39)
$$

which is a scalar quantity, and with $e^{-ip \cdot (x-y)}$ replaced by $e^{+ip \cdot (x-y)}$ in the concluding line found for the case $x^0 < y^0$. Hence taking the v.e.v. of equation 10.38, with the normalisation $\langle 0|0 \rangle = 1$, shows that:

$$
\langle 0|T(\hat{\phi}(x)\hat{\phi}(y))|0\rangle = \hat{\phi}(x)\hat{\phi}(y)
$$
\n(10.40)

which is an object also known as the 'Feynman propagator' for the field $\hat{\phi}(x)$. The complete contractions for the fields of equations 10.13–10.15 can be written as:

$$
\widehat{\phi}(x)\widehat{\phi}(y) = i \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot (x-y)}}{k^2 - m_\phi^2 + i\varepsilon}
$$
(10.41)

$$
\widehat{\dot{\mathcal{X}}(x)\dot{\mathcal{X}}^{\dagger}(y)} = i \int \frac{d^4k}{(2\pi)^4} \, \frac{e^{-ik \cdot (x-y)}}{k^2 - m_{\mathcal{X}}^2 + i\varepsilon} \tag{10.42}
$$

as will be explained in the following section, see for example equation 10.71, where the role of ε will also be described. While the above functions are identical the latter case can be interpreted as representing \mathcal{X}^- particle propagation for $x^0 > y^0$ and \mathcal{X}^+ antiparticle propagation for $x^0 \, \langle y^0 \rangle$, since in the latter case the antiparticle creation operator d^{\dagger}_{λ} $\chi^{\dagger}(\boldsymbol{p})$ of equation 10.14 acts first at the earlier time x^0 .

The generalisation of equation 10.38 for higher-order compositions of fields, in particular for those occurring in equation 10.37, is given by Wick's theorem. This expresses the T-product as a sum of terms involving permutations of normal-ordered products composed with contracted field pairs. Many of these terms vanish when taking the v.e.v. due to their normal-ordered part, leaving residual terms expressible as a product of pair-wise contractions, that is Feynman propagators.

However, the terms in the Wick expansion of the T-ordered product in equation 10.36 do not act on the vacuum directly due to the operators for the initial and final states and hence it is necessary to consider also the more trivial contractions such as (by substituting in for example equations 10.14 and 10.19):

$$
\hat{\mathcal{X}}(x)\hat{B}_{\mathcal{X}}^{\dagger}(\boldsymbol{p}) = \langle 0|\hat{\mathcal{X}}(x)\hat{B}_{\mathcal{X}}^{\dagger}(\boldsymbol{p})|0\rangle \n= \langle 0|\int \frac{d^{3}\boldsymbol{q}}{(2\pi)^{3}}\frac{1}{\sqrt{2\omega_{\boldsymbol{q}}}}\left(b_{\mathcal{X}}(\boldsymbol{q})e^{-iq\cdot x} + d_{\mathcal{X}}^{\dagger}(\boldsymbol{q})e^{+iq\cdot x}\right)\sqrt{2\omega_{\boldsymbol{p}}}b_{\mathcal{X}}^{\dagger}(\boldsymbol{p})|0\rangle \n= \langle 0|\int \frac{d^{3}\boldsymbol{q}}{(2\pi)^{3}}\sqrt{\frac{\omega_{\boldsymbol{p}}}{\omega_{\boldsymbol{q}}}}e^{-iq\cdot x}(2\pi)^{3}\delta^{3}(\boldsymbol{q}-\boldsymbol{p})|0\rangle \n= \langle 0|e^{-ip\cdot x}|0\rangle = e^{-ip\cdot x}
$$
\n(10.43)

$$
\hat{D}_{\mathcal{Y}}(\boldsymbol{q})\hat{\mathcal{Y}}(x) = \langle 0|\hat{D}_{\mathcal{Y}}(\boldsymbol{q})\hat{\mathcal{Y}}(x)|0\rangle = e^{+iq\cdot x}
$$
\n(10.44)

which can be interpreted as the position space representation of the one-particle wavefunctions for the respective initial and final single particle states. These have a simple form since there is no time dependence for the operators $\hat{B}^{\dagger}_{\mathcal{X}}(p)$ and $\hat{D}_{\mathcal{Y}}(q)$ and the order of products in these two expressions is given explicitly with creation operators for initial state particles acting first and those for final state particles acting last in temporal order.

For example, substituting $\int dt H_{\text{int}}(t) = -\int d^4x \mathcal{L}_{\text{int}}(x)$ from equation 10.27, with the interaction Lagrangian \mathcal{L}_{int} of equation 10.23, into equation 10.36 the lowestorder non-trivial term in the perturbative expansion, corresponding to $n = 2$ in equation 10.33, for S_{fi} will include a contribution from the expression:

$$
S_{fi}|_{n=2} = -\frac{g^2}{2} \langle 0|\hat{B}_{\mathcal{Y}}(\boldsymbol{q}_1)\hat{D}_{\mathcal{Y}}(\boldsymbol{q}_2) T\Big(\int d^4x \,\hat{\phi}(x)\hat{\mathcal{X}}^{\dagger}(x)\hat{\mathcal{X}}(x)\int d^4y \,\hat{\phi}(y)\hat{\mathcal{Y}}^{\dagger}(y)\hat{\mathcal{Y}}(y)\Big) D_{\mathcal{X}}^{\dagger}(\boldsymbol{p}_2) B_{\mathcal{X}}^{\dagger}(\boldsymbol{p}_1)|0\rangle
$$

=
$$
-\frac{g^2}{2} \int d^4x \,d^4y \,\hat{B}_{\mathcal{Y}}(\boldsymbol{q}_1)\hat{\mathcal{Y}}^{\dagger}(y) \,\hat{D}_{\mathcal{Y}}(\boldsymbol{q}_2)\hat{\mathcal{Y}}(y) \,\hat{\phi}(x)\hat{\phi}(y) \,\mathcal{X}^{\dagger}(x) D_{\mathcal{X}}^{\dagger}(\boldsymbol{p}_2) \,\mathcal{X}(x) B_{\mathcal{X}}^{\dagger}(\boldsymbol{p}_1) \tag{10.45}
$$

As an alternative to expressions such as equation 10.45 the operators creating the initial and final states, such as B_{λ}^{\dagger} $\mathcal{L}^{\dagger}(\mathbf{p}_1)$ and the Hermitian conjugate of $\hat{D}_{\mathcal{Y}}^{\dagger}(\mathbf{q}_2)$, can also be expressed in terms of functions of free fields, such as $\mathcal{X}(x)$ or $\mathcal{Y}(x)$. In this case an additional Feynman propagator is introduced for each external particle state as expressed in the LSZ reduction formula ([70] p.227). In this form each contribution in the expansion of the scattering amplitude is expressed as the Fourier transform of the v.e.v. of a T-product of free fields, that is of a Green's function (or correlation

function). This full LSZ expression may be needed for example for a consistent treatment of ultraviolet divergences in higher perturbative orders. Here we deal essentially with the 'truncated' Green's function, describing the internal interactions, in order to abstract out the general structure needed to calculate the transition amplitude, as required to make connection with the present theory in the following chapter.

Each non-zero term in the transition amplitude can be represented by a Feynman diagram. In practice QFT calculations of such terms begin with the corresponding Feynman diagrams as constructed from a small set of rules. For example the lowest-order non-trivial term described in equation 10.45 corresponds to the diagram in figure 10.4.

Figure 10.4: Feynman diagram for the process $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$ to lowest order in perturbation theory in the scalar model; closely analogous to the diagram for the QED process $e^+e^- \rightarrow \mu^+\mu^-$ shown in figure 10.3 for which the general comments in the caption apply also here.

More generally the essence of the transition amplitude calculation can be distilled out into a collection of Feynman rules and diagrams as will be described in section 10.5 and table 10.1 for the scalar model. These may be obtained either from the canonical quantisation route, as described above (taking care to handle fermion state operator anticommutators correctly in the case of the Standard Model) or the path integral approach to QFT. Here we are interested in the origin of the Feynman rules, which may be written down from the Lagrangian density for a particular model, for comparison with the present theory. From this point of view the approach of canonical quantisation will prove to be more illuminating, in particular through the intermediate stage of equation 10.31 as will be described in the following chapter. On the other hand the formalism of the path integral, while pragmatically serving as a valuable calculational tool for QFT, seems to provide less in the way of relevant conceptual insight for the present theory.

By substituting the contractions in the form of equations 10.41–10.44 the leading-order term of the transition amplitude expressed in equation 10.45 can be written out explicitly as (with the integrals covering all terms to the right of the integral signs):

$$
S_{fi}|_{n=2} = -\frac{g^2}{2} \int d^4x \, d^4y \, e^{+iq_1 \cdot y} \, e^{+iq_2 \cdot y} \, i \int \frac{d^4k}{(2\pi)^4} \, \frac{e^{-ik \cdot (x-y)}}{k^2 - m_\phi^2 + i\varepsilon} \, e^{-ip_2 \cdot x} \, e^{-ip_1 \cdot x}
$$
\n
$$
= -\frac{g^2}{2} \, i \int d^4x \, d^4y \, \frac{d^4k}{(2\pi)^4} \, \frac{1}{k^2 - m_\phi^2 + i\varepsilon} \, e^{i(k+q_1+q_2)\cdot y} \, e^{-i(k+p_1+p_2)\cdot x}
$$
\n
$$
= -\frac{g^2}{2} \, i \int d^4y \, \frac{d^4k}{(2\pi)^4} \, \frac{1}{k^2 - m_\phi^2 + i\varepsilon} \, e^{i(k+q_1+q_2)\cdot y} \, (2\pi)^4 \, \delta^4(k+p_1+p_2)
$$
\n
$$
= -\frac{g^2}{2} \, i \int d^4y \, \frac{1}{(-p_1-p_2)^2 - m_\phi^2 + i\varepsilon} \, e^{i(q_1+q_2-p_1-p_2)\cdot y}
$$
\n
$$
= -\frac{g^2}{2} \, \frac{i}{(p_1+p_2)^2 - m_\phi^2 + i\varepsilon} \, (2\pi)^4 \, \delta^4(q_1+q_2-p_1-p_2) \tag{10.46}
$$

where the three integrals over d^4x , d^4k and d^4y have been carried out in the third, fourth and fifth lines above respectively. The final expression is relatively simple and explicitly shows how such terms of the matrix element S_{fi} are functions of the coupling g, the particle masses and the momentum variables. Indeed since HEP experiments generally prepare initial particles in momentum states and measure the final particles also in particular momentum states such calculations are simplified by beginning with momentum space Feynman rules, as will be described in section 10.5. In this case the scattering matrix is calculated in terms of momentum space Green's functions which are related to the corresponding position space functions, such as equation 10.41, by a Fourier transform (see also equation 10.72 in the following section).

In explicit calculations the final integral over position space always leads to an overall 4-momentum conserving delta function, as for the bottom line in equation 10.46. This is factored out and not included in the definition of the transition amplitude \mathcal{M}_{fi} as was described for equation 10.6, and hence this delta function is also not included in the Feynman rules for $i\mathcal{M}_{fi}$. Further, in equation 10.45 only complementary halves of \mathcal{L}_{int} from equation 10.23 have been employed under each integral. The reverse choice corresponds to swapping the coordinate labels x and y on the two vertices of the Feynman diagram in figure 10.4. Hence the complete expression for $S_{fi}|_{n=2}$ based on equations 10.45 and 10.46 will contain a further equivalent contribution with the dummy variables x and y interchanged. More generally an amplitude $i\mathcal{M}_{fi}$ will be associated with each topologically distinct Feynman diagram, with the permutation of n! ways of associating the *n* interactions with *n* vertices for an nth order diagram cancelling the $\frac{1}{n!}$ factor in the expansion of equation 10.33. This cancellation is generally incorporated into the Feynman rules for a quantum field theory, including the case of the model QFT considered here as will be described in the opening of section 10.5 (see the discussion of 'rule 6' following table 10.1).

With the above observations on mind, and by reference to equations 10.4– 10.6, the transition amplitude for this leading-order term can be extracted from equation 10.46 (now including also the $x \leftrightarrow y$ case) for the Feynman diagram of figure 10.4 (drawn *without* the explicit x, y labels) as:

$$
\mathcal{M}_{fi} = -g^2 \frac{1}{(p_1 + p_2)^2 - m_{\phi}^2 + i\varepsilon} \tag{10.47}
$$

and hence
$$
|\mathcal{M}_{fi}|^2 = \frac{g^4}{s^2}
$$
 (10.48)

where for the second equation it has been assumed that $s = (p_1+p_2)^2 \gg m_{\phi}^2$, and also ε has been set to zero as will be explained in the following section. The differential crosssection for $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$ scattering to lowest non-trivial order is then obtained by substituting this transition amplitude into equation 10.9 for this two-particle final state to find $\frac{d\sigma}{d\Omega} = \frac{g^4}{64\pi^2}$ $\frac{g^4}{64\pi^2s^3}$.

The purpose of this section has been to show explicitly how such transition amplitudes, featuring in the general cross-section and hence event rate formulae of equations 10.3 and 10.7, are calculated. In the case of muon production the contribution from the lowest-order transition amplitude in equation 10.10 is rather different to the analogous case for the scalar model in equation 10.48. In the case of $e^+e^- \to \mu^+\mu^$ the coupling $e = \sqrt{4\pi\alpha}$ is dimensionless, unlike the case for g in the scalar model, and (combined with the kinematic normalisation factors for the Dirac spinor and electromagnetic fields) this leads to an absence of s in equation 10.10, while for equation 10.48 there is no θ dependence since the model deals with scalar fields only. However, underlying these differences the essential elements of quantum field theory going into these calculations are very similar. In the following section we explore further the basic ingredients and structure of the transition amplitude in the context of the scalar field model.

10.4 Propagators and Causality

Central to the calculation of the amplitude in equation 10.36, via Wick's theorem for the general T-ordered product of several fields, is the Feynman propagator. This was introduced for the scalar field $\phi(x)$ in equations 10.38–10.41 and is generally denoted by the symbol Δ_F ('delta F') with a conventional factor of i (or by $D_F \equiv i\Delta_F$ as for [70]) in the expression:

$$
i\Delta_F(x-y) = \langle 0|T(\hat{\phi}(x)\hat{\phi}(y))|0\rangle \tag{10.49}
$$

$$
= \langle 0 | \theta(x^{0} - y^{0}) \hat{\phi}(x) \hat{\phi}(y) + \theta(y^{0} - x^{0}) \hat{\phi}(y) \hat{\phi}(x) | 0 \rangle \qquad (10.50)
$$

The θ -function takes the value $\theta(t) = 1$ for $t > 0$ and $\theta(t) = 0$ for $t < 0$ (with the value $\theta(0) = \frac{1}{2}$ less significant since $\theta(t)$ is generally used under a time integral; see also the discussion of equation 10.64 below) and explicitly expresses the time ordering of the field product. The Hamiltonian H_{int} is composed of a product of free fields in the interaction picture with the scalar field $\phi(x)$ having the Fourier expansion of equation 10.13. The field $\phi(x)$ can be constructed as a sum of positive and negative frequency parts, $\hat{\phi}(x) = \hat{\phi}^+(x) + \hat{\phi}^-(x)$, with $a(\mathbf{p})$ and $a^{\dagger}(\mathbf{p})$ operator coefficients respectively:

$$
\hat{\phi}^+(x) = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2\omega_p}} a(p) e^{-ip \cdot x}
$$
\n(10.51)

$$
\hat{\phi}^-(x) = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2\omega_p}} a^{\dagger}(p) e^{+ip \cdot x}
$$
\n(10.52)

The $e^{-ip\cdot x}$ components are termed 'positive frequency' since as wavefunctions they would represent states of positive energy under the quantum mechanical operator $H \equiv$ $E = i\hbar\partial/\partial t$ (as implied for the same operator in equation 11.51 of section 11.4 we generally employ natural units with $\hbar = 1$ and $c = 1$ in this paper). Similarly the $e^{+ip\cdot x}$ modes are termed 'negative frequency'. Hence decomposing $\hat{\phi}(x)$ into a sum of the positive and negative frequency parts, with $\hat{\phi}^+(x)|0\rangle = 0$ and $\langle 0|\hat{\phi}^-(x) = 0$, equation 10.50 for the scalar Feynman propagator can be written:

$$
i\Delta_F(x-y) =
$$

$$
\theta(x^0 - y^0) \langle 0 | \hat{\phi}^+(x) \hat{\phi}^-(y) | 0 \rangle + \theta(y^0 - x^0) \langle 0 | \hat{\phi}^+(y) \hat{\phi}^-(x) | 0 \rangle \tag{10.53}
$$

$$
= \theta(x^{0} - y^{0}) \langle 0 | [\hat{\phi}^{+}(x), \hat{\phi}^{-}(y)] | 0 \rangle + \theta(y^{0} - x^{0}) \langle 0 | [\hat{\phi}^{+}(y), \hat{\phi}^{-}(x)] | 0 \rangle \qquad (10.54)
$$

$$
= \theta(x^{0} - y^{0}) i\Delta^{+}(x - y) + \theta(y^{0} - x^{0}) i\Delta^{+}(y - x)
$$
\n(10.55)

In the final line above the function $\Delta^+(x-y)$ can be defined in terms of the commutator of the positive and negative frequency parts of the field and then written out explicitly using equations 10.51 and 10.52:

$$
i\Delta^{+}(x-y) = [\hat{\phi}^{+}(x), \hat{\phi}^{-}(y)] \tag{10.56}
$$

$$
= \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} \int \frac{d^3 \mathbf{q}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{q}}}} \left[a(\mathbf{p}), a^{\dagger}(\mathbf{q}) \right] e^{-ip \cdot x} e^{+iq \cdot y} \qquad (10.57)
$$

$$
= \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2\omega_p} e^{-ip \cdot x} e^{+ip \cdot y} \tag{10.58}
$$

where the constraint on the energy components, such as $p^0 = +\omega_p = +\sqrt{p^2 + m^2}$, is understood in these expressions, and equation 10.16 has been used in the final line – which agrees with equation 10.39 for the $x^0 > y^0$ case as expected. Again here, since $\Delta^+(x-y)$ is simply a function rather than an operator, the vacuum normalisation $\langle 0|0 \rangle = 1$ has been used to factor out the vacuum states in equation 10.54 above to obtain equation 10.55. Integrals of the form $\int \frac{d^3p}{(2\pi)^3}$ $\frac{d^3\bm{p}}{(2\pi)^3} \frac{f(p)}{2\omega_{\bm{p}}}$ $\frac{J(p)}{2\omega_p}$ are Lorentz invariant provided $f(p)$ is a general Lorentz invariant function ([70] p.23, equation 2.40), and hence from equation 10.58 it can be seen that the function $\Delta^+(x-y)$ is Lorentz invariant. Together with the function:

$$
i\Delta^{-}(x-y) = [\hat{\phi}^{-}(x), \hat{\phi}^{+}(y)] = -[\hat{\phi}^{+}(y), \hat{\phi}^{-}(x)] = -i\Delta^{+}(y-x)
$$
(10.59)

these can be written in the manifestly Lorentz invariant form:

$$
i\Delta^{\pm}(x-y) = \pm \int \frac{d^4p}{(2\pi)^4} e^{-ip \cdot (x-y)} \theta(\pm p^0) 2\pi \delta(p^2 - m^2)
$$
 (10.60)

The objects θ and δ are 'generalised functions', or 'distributions', which typically only make full mathematical sense when composed with regular functions in an integrand. A representation of the θ -function will be given below. In one dimension the Dirac δ -function can be defined by the property:

$$
\int dx f(x)\,\delta(x-x') = f(x')\tag{10.61}
$$

which is essentially to substitute the value $x = x'$ into any function $f(x)$. The onedimensional δ-function can be represented by the following expression, which has the subsequent properties (while generally in the text denoting four-parameter objects, x and k each represent a single real variable in equations $10.61-10.63$:

$$
\delta(x - x') = \frac{1}{2\pi} \int_{-\infty}^{+\infty} dk \, e^{\pm ik(x - x')} \tag{10.62}
$$
\n
$$
dx \, \delta(x - x') = 1,
$$

 $with$

and
$$
\int dx f(x) \delta(g(x)) = \sum_{i} \frac{f(a_i)}{|g'(a_i)|}
$$
 with $g(x) = 0$ for $x = \{a_1, a_2...\}$

i.e.
$$
\delta(g(x)) = \sum_{i} \frac{\delta(x - a_i)}{|g'(a_i)|}
$$

e.g.
$$
\delta(x^2 - a^2) = \frac{1}{2a} (\delta(x - a) + \delta(x + a))|_{a \ge 0}
$$
(10.63)

The final expression above can be substituted into equation 10.60 and the p^0 integral performed to show that it is equivalent to the expression for $\Delta^+(x-y)$ in equation 10.58 and to that for $\Delta^-(x-y)$ via equation 10.59 for the $p^0 < 0$ case.

The expression for $\Delta^+(x-y)$ in equation 10.60 describes the positive energy and 'on-mass-shell' momentum space overlap integral of the plane waves, or wavefunctions, $e^{-ip\cdot x}$ and $(e^{-ip\cdot y})^*$. In quantum theory the probability for a particle originating at the spacetime location y to be found at the location x is represented precisely by this amplitude (which via a Fourier transform is analogous to the wavefunction transition amplitude of equation 10.8). In quantum field theory the form of this amplitude $i\Delta^+(x-y) = \langle 0|\phi^+(x), \phi^-(y)|0\rangle$, from equations 10.53 and 10.55, indeed suggests the propagation of a particle *created* at y and *annihilated* at x. Since the spacetime locations x and y are arbitrary x may be either later or earlier than y.

The Feynman propagator can be expressed either in terms of operators acting on the vacuum state, equations 10.49 and 10.50, or in terms of plane waves as described in equations 10.55 and 10.58, with the bridge between these forms of $\Delta_F(x-y)$ provided by the intermediate equations. In either case a temporal ordering is introduced via the θ -functions.

For $x^0 > y^0$ the Feynman propagator is simply $\Delta_F(x-y) = \Delta^+(x-y)$, from equation 10.55, and hence represents the amplitude for a positive energy particle to propagate forward in time from y to x. On the other hand the 'negative energy' part in equation 10.60, with $p^0 < 0$ and $\theta(-p^0) = 1$, represents a propagation from x to y in the $x^0 < y^0$ part of $\Delta_F(x-y)$ and in QFT is interpreted as an *antiparticle* of *positive* energy carried forward in time from x to y . As described following equation 10.42 for

the complex scalar field case and for $x^0 < y^0$ the operator $\hat{\mathcal{X}}$ acts before $\hat{\mathcal{X}}^{\dagger}$ with $d^{\dagger}_{\hat{\mathcal{Z}}}$ creating an antiparticle; while for the real scalar field $\hat{\phi}$ there is no distinction between particle and antiparticle states. Hence $\Delta_F(x-y)$ can be consistently interpreted as only representing propagation forwards in time. Further, from equation 10.55 $\Delta_F(x-y)$ is clearly symmetric in x and y , as is the above interpretation.

In actual calculations all spacetime location variables, such as $\{x, y\}$ for the propagator $\Delta_F(x-y)$, will appear under an integral, such as the $\int d^4x d^4y$ in the first line of equation 10.46, over all spacetime (including regions outside the light cone with $(x - y)^2 < 0$) hence showing explicitly how all possible time orderings are included equally. These integrals essentially represent a Fourier transform to momentum space, allowing for a simplification of the calculations in terms of the momentum space Feynman rules as will be presented in the following section.

Hence the Feynman propagator $\Delta_F(x-y)$ combines wave-like functions $e^{\pm ip\cdot x}$ and particle-like operators $a^{(\dagger)}(\mathbf{p})$ of the field $\hat{\phi}(x)$ with structures of *causality* through the θ -functions, for example in equation 10.53; – apparently elements required to describe the dynamics of exchanges between fields in an interacting theory. It is represented pictorially by an internal line in a Feynman diagram such as figure 10.5(b). Such

Figure 10.5: (a) The function $\Delta^+(x-y)$ represented as the creation, propagation and annihilation of a particle state from y to x in spacetime. (b) The internal line Feynman propagator between two spacetime points, representing equation 10.55. No time ordering is implied in either diagram.

diagrams do not represent literal particle trajectories but should merely be interpreted as mnemonic symbols for mathematical terms such as $\Delta_F(x-y)$ which form the basis of perturbative calculations for an interacting QFT. Indeed the form of $\Delta_F(x-y)$ results from the *restructuring* of the S-matrix calculation of equation 10.31, which describes an explicitly causal chain of operator actions, to the form of equation 10.32 with θ -functions implicitly introduced to impose the apparent time ordering required for mathematical consistency with the first equation.

Hence with the Feynman propagator $\Delta_F(x-y)$ employed to aid *calculation* in this way there need not be any direct physical interpretation of this object. However, due to the time ordering, the Feynman propagator can be considered to represent the internal part of both 'processes' depicted in figure 10.6 below, in which a specific time direction is indicated. While the latter diagram in particular represents a purely mathematical element of the calculation both of these 'processes' are implied in a single Feynman diagram, such as figure 10.5(b), for which there is no explicit temporal direction relating the two vertices.

Figure 10.6: The two terms in equation 10.53 for the Feynman propagator $\Delta_F(x-y)$ describe respectively the two internal ϕ field 'processes' depicted here. In (a) an internal particle state propagates from y to x while in (b) an internal antiparticle propagates from x to y , however with no distinction between particle and antiparticle states for a real scalar field such as $\hat{\phi}$. In (b) ϕ , \mathcal{Y}^+ and \mathcal{Y}^- particle states are created out of the vacuum at x.

The propagator $\Delta_F(x-y)$ depends only on the 4-vector difference $(x-y)$. The functions $\Delta^{\pm}(x-y)$, and hence also $\Delta_F(x-y)$, are non-zero outside the light cone region, $(x-y)^2 < 0$, where they decay exponentially. While the $\Delta^{\pm}(x-y)$ are Lorentz invariant the function $\theta(x^0 - y^0)$ is not Lorentz invariant for spacelike separations outside the light cone. However the combination of both terms in equation 10.55 is Lorentz invariant.

The generalised function $\theta(t)$ itself can be expressed in the Fourier, or integral, representation as:

$$
\theta(t) = \lim_{\eta \to 0^+} \frac{i}{2\pi} \int_{-\infty}^{+\infty} \frac{e^{-ist}}{s + i\eta} ds \qquad (10.64)
$$

which as a *distribution* is differentiable everywhere (unlike the closely related Heaviside function $H(t)$ defined with $H(t) = 1$ for $t \ge 0$ and $H(t) = 0$ for $t < 0$). In fact:

$$
\frac{d\theta(t)}{dt} = \lim_{\eta \to 0^+} \frac{i}{2\pi} \int \frac{-is \, e^{-ist}}{s + i\eta} \, ds = \frac{1}{2\pi} \int e^{-ist} ds = \delta(t) \tag{10.65}
$$

from the representation of the δ -function in equation 10.62. The substitution of the θ-function into equation 10.55 for $\Delta_F(x-y)$ is aided by first making the change of integration variable $s \to k^0 - \omega$, with finite real constant ω , in equation 10.64 so that:

$$
\theta(t) = \lim_{\eta \to 0^{+}} \frac{i}{2\pi} \int_{-\infty}^{+\infty} \frac{e^{-i(k^{0}-\omega)t}}{k^{0}-\omega+i\eta} dk^{0}
$$

$$
= \lim_{\eta \to 0^{+}} \frac{i}{2\pi} e^{+i\omega t} \int_{-\infty}^{+\infty} \frac{e^{-ik^{0}t}}{k^{0}-\omega+i\eta} dk^{0}
$$
and hence:
$$
\theta(t) e^{-i\omega t} = \lim_{\eta \to 0^{+}} \frac{i}{2\pi} \int_{-\infty}^{+\infty} \frac{e^{-ik^{0}t}}{k^{0}-\omega+i\eta} dk^{0}
$$
(10.66)

This expression for the θ -function, along with equation 10.58 for the function $\Delta^+(x-y)$, can be substituted into equation 10.55 for the Feynman propagator as follows:

$$
\Delta_F(x-y) = \theta(x^0 - y^0) \Delta^+(x-y) + \theta(y^0 - x^0) \Delta^+(y-x)
$$
(10.67)

$$
= \theta(x^0 - y^0) (-i) \int \frac{d^3p}{(2\pi)^3} \frac{1}{2\omega_p} e^{+ip \cdot (x-y)} e^{-ip^0 \cdot (x^0 - y^0)}
$$

$$
+ \theta(y^0 - x^0) (-i) \int \frac{d^3p}{(2\pi)^3} \frac{1}{2\omega_p} e^{+ip \cdot (y-x)} e^{-ip^0 \cdot (y^0 - x^0)} \Big|_{p^0 = +\omega_p = +\sqrt{p^2 + m^2}}
$$

Since $\{x, y\}$ are fixed for each value of $\Delta_F(x - y)$ the θ -function can be moved inside the d^3p integral and with $p^0 = +\omega_p$, which is constant for each value of the 3-vector p , equation 10.66 above may be substituted into the square brackets below:

$$
\Delta_F(x-y) = (-i) \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} e^{+i\mathbf{p}\cdot(\mathbf{x}-\mathbf{y})} \left[\theta(x^0 - y^0) e^{-i\omega_{\mathbf{p}}\cdot(x^0 - y^0)} \right] \n+ (-i) \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} e^{+i\mathbf{p}\cdot(\mathbf{y}-\mathbf{x})} \left[\theta(y^0 - x^0) e^{-i\omega_{\mathbf{p}}\cdot(y^0 - x^0)} \right] \n= (-i) \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} e^{+i\mathbf{p}\cdot(\mathbf{x}-\mathbf{y})} \left[\lim_{\eta \to 0^+} \frac{i}{2\pi} \int \frac{e^{-ik^0(x^0 - y^0)}}{k^0 - \omega_{\mathbf{p}} + i\eta} dk^0 \right] \n+ (-i) \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} e^{+i\mathbf{p}\cdot(\mathbf{y}-\mathbf{x})} \left[\lim_{\eta \to 0^+} \frac{i}{2\pi} \int \frac{e^{-ik^0(y^0 - x^0)}}{k^0 - \omega_{\mathbf{p}} + i\eta} dk^0 \right] (10.69)
$$

Hence the 3-momentum integral has been enlarged to a 4-parameter integral by including the full unrestricted range of the k^0 variable associated with the θ -function integral. That is while the p^0 component of the 4-vector p is constrained to the value $\omega_p = +\sqrt{p^2 + m^2}$, the free k⁰ integration variable is introduced from equation 10.66. In relabelling the 3-momentum p by the 3-vector k the above final expression is seen to take the form of an apparent 4-momentum integral:

$$
\Delta_F(x - y) = \lim_{\eta \to 0^+} \int \frac{d^4k}{(2\pi)^4 2\omega_k} \left[\frac{e^{-ik \cdot (x - y)}}{k^0 - \omega_k + i\eta} + \frac{e^{+ik \cdot (x - y)}}{k^0 - \omega_k + i\eta} \right] \qquad (10.70)
$$

\n
$$
= \lim_{\eta \to 0^+} \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot (x - y)} \left[\frac{1}{2\omega_k} \left(\frac{1}{k^0 - \omega_k + i\eta} + \frac{1}{-k^0 - \omega_k + i\eta} \right) \right]
$$

\n
$$
= \lim_{\eta \to 0^+} \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot (x - y)} \left[\frac{\omega_k - i\eta}{\omega_k((k^0)^2 - \omega_k^2 + 2i\omega_k\eta - \eta^2)} \right]
$$

\n
$$
= \lim_{\varepsilon \to 0^+} \int \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot (x - y)}}{k^2 - m^2 + i\varepsilon} \qquad (10.71)
$$

Here the second line is obtained by reversing the sign of all 4 integration variables in the second term in square brackets in equation 10.70. The third and final lines follow after some straightforward algebra, with the new limiting parameter $\varepsilon \simeq +2\omega_{\mathbf{k}}\eta$ introduced, and with the limit $\varepsilon \to 0^+$ for the integral understood even if not explicitly stated. Through substituting $\omega_k^2 = k^2 + m^2$ (see equation 10.67) into the third line, and with $k^2 = (k^0)^2 - k^2$ in the final line, k is treated as a Lorentz 4-vector. This is the expression for the Feynman propagator scalar function quoted in equation 10.41 (with a factor of i from equation 10.49). This function of the spacetime difference $(x - y)$ may also be written:

$$
\Delta_F(x - y) = \int \frac{d^4k}{(2\pi)^4} e^{-ik \cdot (x - y)} \widetilde{\Delta}_F(k)
$$

with
$$
\widetilde{\Delta}_F(k) = \frac{1}{k^2 - m^2 + i\varepsilon}
$$
 (10.72)

being the momentum space representation of the Feynman propagator, obtained as the coefficients in the Fourier decomposition of the position space function.

Unlike the 4-momentum integral expression for $\Delta^{\pm}(x-y)$ in equation 10.60, for the Feynman propagator in equation 10.71 there is no 'mass-shell' condition with a $\delta(k^2 - m^2)$ function, and with 4 independent 'momentum' variables the Feynman propagator represents 'states' which are generally 'off-shell'. This situation motivates the term 'virtual particle' in referring to the 'propagating entity'. On the other hand $\Delta_F(x-y)$ is constructed in equation 10.67 out of elements which are on-shell with energy $\omega_p = \pm \sqrt{p^2 + m^2}$, with the off-shell interpretation for the full expression arising through the incorporation of the θ -functions.

Equation 10.71 follows from the structure of $\Delta^+(x-y)$, which is found through $[a(\mathbf{p}), a^{\dagger}(\mathbf{p}')]$ commutators appearing for example in the expansion of terms in equation 10.31 between vacuum states to determine a scattering amplitude, together with the θ -functions, which are deployed when the calculation is reorganised with the time ordering T of equation 10.32. Hence the notion of 'virtual particle states' may be considered to be a purely mathematical construction arising from this reworking of the calculation.

In equation 10.64 the θ -function is defined by a contour integration in the complex plane. This involves a combination of Cauchy's theorem and the residue theorem – respectively for integration contours surrounding a region of the integrand function which is regular or containing singularities, together with Jordan's lemma for the vanishing of particular e^{-ist} contour integrals depending on the sign of the real parameter t in the complex s-plane. The result is that $\theta(t)$ can be expressed in equation 10.64 with the horizontal integration contour C of figure 10.7(a), in which the pole in the integrand at $s = -i\eta$ is also shown.

The single pole in the integrand function for $\theta(t)$ carries over into two poles in the complex plane (since there are two θ -functions in equation 10.68 leading to equation 10.70) for the integrand in equation 10.71 for $\Delta_F(x-y)$. In this latter equation (which was derived from equation 10.69) it is understood that the k^0 integration should be carried out first following the straight contour C along the real axis in figure 10.7(b). Using Cauchy's theorem this contour integral can be 'analytically continued' by a 90^0 counterclockwise rotation to the imaginary k^0 axis without encountering any poles. Under this 'Wick rotation' to Euclidean 4-space (with k^0 replaced by $k^4 = i k^0$ to form a Euclidean 4-vector with k) the parameter η (and hence ε in equation 10.71) may be discarded.

Alternatively equation 10.71 and the real k^0 integration in figure 10.7(b) is equivalent setting $\varepsilon = 0$ and performing the resulting integral:

$$
\Delta_F(x-y) = \int_{C_F} \frac{d^4k}{(2\pi)^4} \frac{e^{-ik \cdot (x-y)}}{k^2 - m^2}
$$
\n(10.73)

Figure 10.7: Integration contours (a) in the complex s-plane for $\theta(t)$ defined in equation 10.64 and (b) in the complex k^0 -plane for the Feynman propagator $\Delta_F(x-y)$ in equation 10.71. The single pole in the first case and pair of poles in the second case are also indicated.

following the contour C_F with an implied limit of infinitesimal detours below the first then above the second pole on the real axis as displayed by the thick line in figure 10.8. Although these expressions are equivalent equation 10.71 is generally quoted in preference to equation 10.73 since the $i\varepsilon$ term in the former case serves to explicitly indicate the side on which the contour avoids the poles.

Figure 10.8: The six functions $\Delta^{\pm}(x-y)$, $\Delta(x-y)$ and $\Delta_{F,R,A}(x-y)$ described in the text can be defined by the integration along six different contours $(C^{\pm}, C \text{ and } C_{F,R,A})$ respectively) in the complex k^0 -plane for the same integrand function presented in equation 10.73.

Maintaining the same integrand while adapting the contour C_F employed in equation 10.73 in a total of six different ways leads to the expression of a total of six different functions, all related to $\Delta_F(x-y)$, and each then defined here in a related mathematical form. However, the primary importance is given to the C_F contour and the Feynman propagator in QFT since this object arises prominently in

the calculation of scattering amplitudes. The three contours C_F , C_R and C_A hug the real axis in figure 10.8 with the integral determined in the limit of vanishingly small detours around the poles. However these integrals do *include* these infinitesimal detours are not the Cauchy principle values of the integrals which 'hop over' the poles in this limit and would then be identical for ' C_F ', ' C_R ' and ' C_A '.

The three remaining contours C, C^+ and C^- can be taken anywhere in the complex plane, so long as they navigate around the poles with the topology indicated in figure 10.8. These contour integrals in the complex k^0 -plane simply have the values of $-2\pi i$ times the residues enclosed, with a negative sign relative to the residue theorem which is based on anticlockwise circulating contours. It is again understood that this complex k^0 integral is performed first in equation 10.73 for the respective contours, before the remaining real $\int d^3k$, in defining the Δ , Δ^+ and Δ^- functions.

Here the outer contour C , encompassing both poles in figure 10.8, represents the Lorentz invariant singular function $\Delta(x - y)$. This function can be introduced in the discussion of causality relating to field interactions and defined directly in terms of the field commutator:

$$
i\Delta(x - y) = [\hat{\phi}(x), \hat{\phi}(y)]
$$

= $[\hat{\phi}^+(x), \hat{\phi}^-(y)] + [\hat{\phi}^-(x), \hat{\phi}^+(y)]$ (10.74)

$$
= i\Delta^{+}(x-y) + i\Delta^{-}(x-y)
$$
 (10.75)

$$
= \int \frac{d^4k}{(2\pi)^4} \,\varepsilon(k^0) \, 2\pi \, \delta(k^2 - m^2) e^{-ik \cdot (x-y)} \tag{10.76}
$$

using equations 10.59 and 10.60 and with $\varepsilon(k^0) = +1, 0, -1$ for $k^0 > 0, k^0 = 0, k^0 < 0$ respectively. Equation 10.75 is consistent with the residue theorem with the integral contour C in figure 10.8 enclosing both poles, which are separately enclosed by C^+ and C^- . With $\Delta(x-y) = 0$ for $(x-y)^2 < 0$, unlike the case for the individual $\Delta^{\pm}(x-y)$ components, this function represents causality in field interactions through equation 10.74, in the sense that it implies $\hat{\phi}(x)$ and $\hat{\phi}(y)$ operate independently of each other outside the light cone. Each of these three functions satisfies the Klein-Gordon equation:

$$
(\Box_x + m^2) \Delta^{(\pm)}(x - y) = 0 \tag{10.77}
$$

where the differential operator \Box_x acts on the spacetime variables corresponding to x, and m in the above is understood to be the mass m_{ϕ} associated with the scalar field $\hat{\phi}(x)$. In the spatial plane $x^0 - y^0 = 0$ the function $\Delta(x - y)$ also satisfies the time derivative equation $\partial_0 \Delta(\mathbf{x} - \mathbf{y}, 0) = -i\delta^3(\mathbf{x} - \mathbf{y})$ which, via equation 10.74, and the conjugate field $\hat{\pi}(x) = \partial_0 \phi(x)$, is consistent with the equal-time field commutation relation:

$$
[\hat{\phi}(\boldsymbol{x},t),\hat{\pi}(\boldsymbol{y},t)] = i\,\delta^3(\boldsymbol{x}-\boldsymbol{y})\tag{10.78}
$$

Here we have arrived at this expression by employing the commutation relation $[a(\mathbf{p}), a^{\dagger}(\mathbf{q})] = (2\pi)^3 \delta^3(\mathbf{p} - \mathbf{q})$ in order obtain equation 10.76 from equation 10.74 via equation 10.57. However the 'canonical' commutation relation of equation 10.78 may be postulated ahead of equation 10.16 as the field quantisation rule, as a generalisation from the non-relativistic quantum mechanical relation $[\hat{x}^a, \hat{p}^b] = i\hbar \delta^{ab}$ for $a, b = \{1, 2, 3\}$ in the three spatial dimensions.

In contrast to the three $C^{(\pm)}$ contours for the three $\Delta^{(\pm)}$ functions in figure 10.8 the three remaining contour integrals essentially follow the real k^0 axis, differing only in their means of bypassing the two poles as described above. Although figure 10.8 provides a neat mathematical way of summarising these six functions it is important to understand their conceptual meaning and the relationships between them.

In particular the two functions $\Delta_R(x-y)$ and $\Delta_A(x-y)$ are the 'retarded' and 'advanced' parts of the Lorentz invariant singular function $\Delta(x - y)$, that is:

$$
\Delta_R(x - y) = \theta(x^0 - y^0) \Delta(x - y) \qquad (= 0 \text{ for } x^0 < y^0) \qquad (10.79)
$$

$$
\Delta_A(x - y) = -\theta(y^0 - x^0) \Delta(x - y) \qquad (= 0 \text{ for } x^0 > y^0) \qquad (10.80)
$$

Both of these functions of course vanish outside the light cone since $\Delta(x-y)$ does. The function $\Delta_R(x-y)$ also vanishes for $x^0 < y^0$ into the past while $\Delta_A(x-y)$ vanishes into the future. In solutions for a classical theory both retarded and advanced waves can be identified, with the latter then being eliminated on the grounds of causality. Bearing in mind the antiparticle interpretation described earlier in this section, the retarded and advanced functions are of comparable significance in quantum field theory. These two functions, along with the Feynman propagator $\Delta_F(x-y)$, are Green's functions which satisfy the inhomogeneous Klein-Gordon equation:

$$
(\Box_x + m^2) \Delta_{F,R,A}(x - y) = -\delta^4(x - y)
$$
\n(10.81)

The conventional factor of i introduced in equation 10.49 is chosen so that such a factor is absent in the above equation. The choice of detours around the poles for the contour integration in figure 10.8 reflects different choices of boundary conditions for solutions to the differential equation 10.81, such as the vanishing of the functions into the past or the future described in equations 10.79 and 10.80. The relation of the Feynman propagator to the retarded and advanced Green's functions can be seen from figure 10.8 to be:

$$
\Delta_F(x - y) = \Delta_R(x - y, \theta(k^0)) + \Delta_A(x - y, \theta(-k^0))
$$
\n(10.82)

That is, with the $\theta(\pm k^0)$ -functions understood to be attached to the integrand in the right-hand side of equation 10.73, the contour integral for the Feynman propagator Δ_F follows the advanced contour C_A in the negative frequency $k^0 < 0$ half-plane and the retarded contour C_R for positive frequency $k^0 > 0$. Alternatively, on attaching the $\theta(\pm k^0)$ to the integrand of equation 10.76 which is then substituted into equations 10.79 and 10.80 and in turn into equation 10.82 the resulting expression is found to be identical to equation 10.55, with the latter expressing $\Delta_F(x-y)$ in terms of the $\Delta^{\pm}(x-y)$ functions.

Retarded and advanced propagators are employed in quantum field theory to study solutions to the equations of motion. For example, with regard to the scalar model of section 10.3, expressions such as:

$$
\hat{\phi}(x) = \int d^4y \,\Delta_R(x-y) \, g\hat{\mathcal{X}}^\dagger(y)\hat{\mathcal{X}}(y) \tag{10.83}
$$

may be considered. The retarded propagator $\Delta_R(x-y)$ satisfies equation 10.81, which applied to equation 10.83 yields:

$$
\left(\Box_x + m^2\right)\hat{\phi}(x) = -g\hat{\mathcal{X}}^\dagger(x)\hat{\mathcal{X}}(x) \tag{10.84}
$$

as an equation of motion for the quantum field $\hat{\phi}(x)$ with source term $-g\hat{\mathcal{X}}^{\dagger}(x)\hat{\mathcal{X}}(x)$. This is equation 10.24, for the two fields $\phi(x)$ and $\mathcal{X}(x)$ of the scalar model, which in the previous section was derived from the Lagrangian of equation 10.23. This method of obtaining solutions to equations of motion via Green's functions was originally employed for classical field theories. For the classical case the right-hand side of equation 10.84 may act as a source of disturbance generating a wave motion for the corresponding classical field $\phi(x)$ on the left-hand side, while in the quantum case the right-hand side may act as a source for the production of particles of the quantum field $\phi(x)$.

10.5 Feynman Rules and Optical Theorem

The various systematic procedures involved in calculating a given transition amplitude for a given interacting quantum field theory can be conveniently summarised in a small set of rules, which are most simply expressed in the momentum space representation, obtained in turn for the Feynman propagators in their Fourier expansions. The Feynman rules associate mathematical elements of the calculation with graphical elements in a diagram representing a particular contribution to the transition amplitude. These rules are written down here for the scalar model with the interaction Lagrangian of equation 10.23 in table 10.1. These rules resemble those for the simpler interacting field theory based on a single scalar field $\hat{\phi}(x)$ with the interaction Lagrangian $\mathcal{L}_{int} = -\frac{\lambda}{4!} \hat{\phi}^4$ ([70] p.115), which is often presented as a model QFT.

Representing possible terms in the transition amplitude by the possible topologies of graphical diagrams greatly assists the bookkeeping involved in the calculation. While terms in the expansion of $S = Te^{-i\int dt H_{int}(t)}$ of equation 10.35 can be pictured this way the internal lines should not be literally interpreted as representing trajectories of 'virtual particles', indeed there is no reference to location at all in the momentum space Feynman rules. Rather the topology of the diagrams describes the structure of possible mathematical terms. Here we make some further comments on these rules 1–6 as listed in table 10.1:

- 1. Each line, whether internal or external, is associated with a particular field type. The direction of an arrow on a line can be used to distinguish a particle from an antiparticle when relevant, as described in 'item 3' below. The propagator term is $i\Delta_F(k)$ from equation 10.72, where the factor of i follows from the convention of equation 10.49.
- 2. The coupling g is added by hand in equation 10.23 and hence for the interaction Hamiltonian in equation 10.27. The factor of $-i$ originates from equation 10.34 and in turn from the evolution equation 10.30.
- 3. The factors in these first three items are multiplied together. The external lines can be labelled with the on-mass-shell 4-momentum k , with an arrow on the line following the momentum transfer (into or out of the terminating vertex) for a particle and in the opposite direction for an antiparticle (with a similar convention for internal lines), as depicted in figures 10.3 and 10.4.

Table 10.1: The Feynman rules in momentum space for the scalar model, relating mathematical terms and instructions to the elements of a Feynman diagram, each of which contributes to a transition amplitude $i\mathcal{M}_{fi}$.

- 4. The momentum conservation for each vertex arises from the $\int d^4x$ over spacetime associated with each of n factors of $\mathcal{L}_{int}(x)$ in the nth order of perturbation, with the x-dependence in the integrand purely in terms of the form $e^{i(\sum_a k_a)x}$, with the $\sum_{a} k_a$ summing over all lines connected to the vertex. This is seen for example for $n = 2$ in equation 10.46, where all the various factors for the $S_{fi}|_{n=2}$ term of equation 10.45 are composed.
- 5. The loop integrals over $\int d^4k$ tend to diverge leading to the need for renormalisation, as will be discussed below for figure 10.9 and also in section 11.3. The loop integral includes the full independent range $-\infty < k^0 < +\infty$, arising originally from equations 10.66–10.71 as described in the previous section. In other quantum field theories there may also be a discrete sum over field indices such as spin.
- 6. This factor is simply the exponential expansion coefficient of equations 10.33 and 10.34 multiplied by $n!$ from the number of ways the dummy integration variables $\{x, y, \ldots\}$ can label the *n* vertices of the Feynman diagram. In other theories there may also be symmetry factors for permutations of identical particles, as for example in the $\hat{\phi}^4$ theory ([70] p.93).

Bearing in mind equations 10.5 and 10.6 each Feynman diagram corresponds to a contribution to the S-matrix without the overall factor of $(2\pi)^4 \delta^4(p_F - p_I)$, that is the

transition amplitude $i\mathcal{M}_{fi}$. Since the amplitude appears as $|\mathcal{M}_{fi}|^2$ in the cross-section calculation of equation 10.3 the overall factor of i is sometimes neglected.

The above rules can be applied to the Feynman diagram of figure 10.4, representing the lowest-order term for the process $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$. Reading off the Feynman rules in table 10.1 for this diagram we find directly:

$$
i\mathcal{M}_{fi} = -g^2 \frac{i}{(p_1 + p_2)^2 - m_{\phi}^2 + i\varepsilon}
$$
 (10.85)

This is the same expression for the transition amplitude as obtained in equation 10.47 by explicit calculation, as it should be. The Feynman rules, as applied above, strip out the essence of such calculations.

We recall here that the transition probability is obtained from the square of the absolute value of the transition amplitude, by a basic postulate of quantum theory, as discussed around equation 10.8. The transition amplitude itself is strictly composed of all of the terms in the expansion of equation 10.35, of which only the lowest-order nontrivial term for $n = 2$ has been accounted for in equation 10.85. It is an assumption of perturbation theory that the subsequent inclusion of terms of higher order into the sum gives a rapidly improving approximation to physical quantities such that very few orders are needed in practice. One aim of the following chapter is to understand how this procedure works in the context of the theory presented in this paper, but here we first explore a next-to-leading order term in the standard QFT approach for the scalar model. One of several contributions to the transition amplitude for $n = 4$ is described by the Feynman diagram in figure 10.9.

Figure 10.9: Feynman diagram for the process $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$ for a possible higherorder perturbation. At this 'next-to-leading order' level an unconstrained internal loop momentum r appears, depicted here for the \mathcal{X} field.

In this case reading off the instructions from table 10.1 'rule 5' is invoked for the freedom in the internal loop momentum r which is not constrained by the application of 'rule 4', leading to the amplitude contribution:

$$
i\mathcal{M}_{fi} = g^4 \left(\frac{i}{(p_1+p_2)^2 - m_{\phi}^2 + i\varepsilon}\right)^2 \int \frac{d^4r}{(2\pi)^4} \frac{i}{r^2 - m_{\mathcal{X}}^2 + i\varepsilon} \frac{i}{(p_1+p_2-r)^2 - m_{\mathcal{X}}^2 + i\varepsilon} \tag{10.86}
$$

In QFT such loop momentum integrals are frequently divergent, as is the case here and for similar terms in the $\hat{\phi}^4$ scalar model, giving infinite and hence meaningless answers if taken at face value. This leads to the need for a program of 'renormalisation' in order to extract useful results out of these calculations.

In practice the divergent internal loop integrals are first made finite by introducing a parameter to smooth the integrand or act as a cut-off to the integration range, a process known as 'regularisation'. The theory is then renormalised, essentially by calibration against an empirical input, before the regularising parameters are eliminated. The aim is to achieve finite predictive quantities in this way for comparison with further physical measurements, such as the observation of 'running coupling' which is a consequence of renormalisation as will be described in section 11.3.

In the natural units we are adopting, with $\hbar = 1$ and $c = 1$, any physical quantity can be expressed in units of mass, that is with dimension M^D , where the mass dimension M is reciprocal to that of length and time, that is $M^1 \equiv L^{-1} \equiv T^{-1}$. The success of the renormalisation procedure generally depends upon the power of the mass dimension D for the coupling parameter itself. Since $\int \mathcal{L} d^4x$ represents the 'action' which is a dimensionless quantity with $D = 0$, the Lagrangian density $\mathcal L$ has dimension $D = 4$, which is also consistent with equation 10.27 since the Hamiltonian H has dimension $D = 1$. If the coupling parameter in the interaction Lagrangian has $D \geq 0$ such a theory is probably renormalisable, whereas theories with $D < 0$, such as gravitation for which Newton's constant G_N has $D = -2$, are non-renormalisable.

Hence the renormalisation procedure works for quantum field theories with dimensionless coupling constants, such as QED and the Standard Model in general and also the scalar model with $\mathcal{L}_{int} = -\frac{\lambda}{4!} \hat{\phi}^4$. For the scalar model considered here with $\mathcal{L}_{\text{int}} = -g\hat{\phi}\hat{\mathcal{X}}^{\dagger}\hat{\mathcal{X}} - g\hat{\phi}\hat{\mathcal{Y}}^{\dagger}\hat{\mathcal{Y}}$ the full Lagrangian of equation 10.23 implies that the coupling g has dimension $D = +1$, and hence the theory can be renormalised. Such a theory with $D > 0$ may even be 'super-renormalisable' and contain no infinities at all after some order of perturbation.

Since a cross-section σ has the dimension L^2 the right-hand side of equation 10.3 must also have the overall dimension $D = -2$, which is also the dimension of the initial state flux factor in this equation. For a two-particle final state the Lorentz invariant phase space $d\Phi$ is dimensionless, implying that the amplitude \mathcal{M}_{fi} itself should also have $D = 0$ in this case. This is consistent with the dimensionless coupling e of QED in equation 10.10 and with the coupling g having $D = 1$ for the scalar model in equations 10.48, 10.85 and 10.86. More generally the dimension of the transition amplitude \mathcal{M}_{fi} will depend upon the multiplicity of the final state and the conventions employed for initial and final state normalisation, consistent with the composition of factors forming the cross-section having the appropriate net dimension, as is the case for equation 10.3.

Higher-order corrections, as appearing for the internal propagator for the field ϕ of figure 10.4 when dressed as in figure 10.9, will also be important for the external particle states. This applies also for calculations in QED and Standard Model QFT calculations in general. Although the theory begins by describing free field states it is not possible in the physical world to decouple the electron field from the electromagnetic field (or the \mathcal{X} field from the ϕ field in the scalar model) since they are intrinsic elements of a single interacting system.

Any parameters, such as masses m_{ϕ} and m_{χ} in the model here, ascribed to a free field will be unphysical and unmeasurable. Instead a finite set of fundamental physical parameters can be operationally defined as those quantities which are directly measurable in the laboratory. The self-interaction effects for the observed particle states are absorbed into these measured parameters, with the Fock space of initial and final states (in the interaction picture basis) assumed to represent precisely the observed masses and charges of physically produced or detected particles in matrix element $\langle f|S|i\rangle$ calculations. These renormalised parameters obey the fundamental conservation laws of external and internal symmetries in collision processes. The physical renormalised mass is not the same object as the 'bare' mass parameter appearing in the Lagrangian of the theory.

As well as the obvious necessity to 'tame the infinities' for calculations of physical quantities, the finite results obtained must also respect the basic requirement of probability conservation, namely that the total probability for something to happen must always be equal to 1. This fundamental principle translates in quantum theory into the unitarity of the S-matrix, with the restrictions of this condition having implications for the relationship between physical quantities such as the cross-section σ and the structure of the transition amplitude \mathcal{M}_{fi} as will be described here.

The unitarity of the S-matrix of equation 10.35, that is the property $SS^{\dagger} =$ $S^{\dagger}S = 1$, together with the definition of the operator $T = i(1 - S)$ in equation 10.5, hence with $T^{\dagger} = -i(1 - S^{\dagger})$, implies that:

$$
TT^{\dagger} = T^{\dagger}T = i(T^{\dagger} - T) \qquad (10.87)
$$

and therefore:
$$
\langle f|TT^{\dagger}|i\rangle = i\langle f|T^{\dagger}|i\rangle - i\langle f|T|i\rangle
$$
 (10.88)

Inserting a sum over a complete set of intermediate states $|m\rangle$ the left-hand side of this expression can be written as:

$$
\langle f|TT^{\dagger}|i\rangle = \sum_{m} \left(\prod_{j=l}^{r_m} \int \frac{d^3 \mathbf{k}_j}{(2\pi)^3 2E_j} \right) \langle f|T|m\rangle \langle m|T^{\dagger}|i\rangle \tag{10.89}
$$

where r_m is the number of particles in each state $|m\rangle$ and $\frac{d^3 k_j}{(2\pi)^3 2}$ $\frac{a}{(2\pi)^3 2E_j}$ is the invariant phase space element for the particle state normalisation adopted, as described in section 10.2 and required here for the insertion of the unit operator 1 between T and T^{\dagger} . The two terms on the right-hand side of equation 10.88 can be written as:

$$
i\langle f|T|i\rangle = i\mathcal{M}_{fi} (2\pi)^4 \,\delta^4(p_F - p_I) \tag{10.90}
$$

$$
i\langle f|T^{\dagger}|i\rangle = i\mathcal{M}_{if}^{*}(2\pi)^{4}\delta^{4}(p_{F}-p_{I})
$$
\n(10.91)

These are obtained directly from equation 10.6, which can also be applied to the righthand side of equation 10.89 and hence substituted into equation 10.88 along with equations 10.90 and 10.91 to find:

$$
\sum_{m} \mathcal{M}_{fm} (2\pi)^4 \delta^4(p_F - p_M) \mathcal{M}_{im}^* \left[(2\pi)^4 \delta^4(p_M - p_I) \prod_{j=l}^{r_m} \int \frac{d^3 \mathbf{k}_j}{(2\pi)^3 2E_j} \right]
$$

= $(i\mathcal{M}_{if}^* - i\mathcal{M}_{fi}) (2\pi)^4 \delta^4(p_F - p_I)$ (10.92)

This is a non-linear relationship between transition amplitudes, with a product on the left and a sum on the right-hand side, resulting from the unitarity of the S-matrix. Given the second δ -function on the left-hand side the first one $\delta^4(p_F - p_M)$ may be

replaced by $\delta^4(p_F - p_I)$, which hence cancels with the δ -function on the right-hand side. The term in square brackets is simply the Lorentz invariant phase space $d\Phi$, as described for equations 10.3 and 10.7, here for the intermediate states, and hence equation 10.92 can be written simply as:

$$
\sum_{m} \left(\mathcal{M}_{fm} \mathcal{M}_{im}^* \int d\Phi \right) = i(\mathcal{M}_{if}^* - \mathcal{M}_{fi}) \tag{10.93}
$$

Considering a two-particle initial state and setting $|f\rangle = |i\rangle$, corresponding to elastic forward scattering at a HEP collider with the final state being identical to the initial state, and by comparison with equation 10.3, the left-hand side above is then identical to the *total* cross-section for the transition from an initial state $|i\rangle$ to any state $|m\rangle$, up to an initial state flux factor, which again relates to the state normalisation. That is, with $|f\rangle = |i\rangle$ and since $|\mathcal{M}_{im}| = |\mathcal{M}_{mi}|$, equation 10.93 becomes:

$$
\sum_{m} \left(|\mathcal{M}_{mi}|^2 \int d\Phi \right) = 2 \operatorname{Im}(\mathcal{M}_{ii}) \tag{10.94}
$$

$$
\equiv 4E_1E_2|\mathbf{v}_1-\mathbf{v}_2|\sigma_{\text{tot}} = 2\,\text{Im}(\mathcal{M}_{ii}) \tag{10.95}
$$

where equation 10.3, with an implied integration over the phase space for each final state to obtain the total cross-section σ_{tot} , has been substituted in for the left-hand side in the second line. (Here $\text{Im}(\mathcal{M}_{ii})$ is of course a real number, as for the standard definition of the imaginary part of a complex number, in contrast to the definition of the imaginary part of an octonion as described immediately before equation 6.10). The flux factor can be expressed in terms of the total centre-of-mass energy E_T (= \sqrt{s}) and the momentum of either initial particle in the centre-of-mass frame $|p_i|$ (noting however that this factor is not fully Lorentz invariant, as described after equation 10.3), such that the total cross-section can finally be written as:

$$
\sigma_{\text{tot}}(i \to \text{anything}) = \frac{\text{Im}(\mathcal{M}_{ii})}{2E_T|\mathbf{p}_i|} \tag{10.96}
$$

This relationship, along with its derivation, is a form of the 'optical theorem' $(70 \text{ p.}231, \text{ equation } 7.50)$. It is a consequence of the S-matrix unitarity condition in scattering experiments, which in turn expresses basic properties of the laws of probability, and has further implications for observable quantities. Here it shows how the total cross-section for the production of any final state is directly related to the imaginary part of the forward scattering amplitude, up to the normalisation factor in equation 10.96. By equations 10.5 and 10.6 the imaginary part of \mathcal{M}_{ii} corresponds to the non-trivial real part of $\langle i|S|i\rangle$, with many intermediate processes contributing. The significance of this result in the context of the present paper is that it demonstrates a linear relationship between a cross-section, that is the likelihood of an event occurring, and an amplitude.

The generalised optical theorem as expressed in equation 10.93 can also be applied to the case of a single particle initial state. On again setting $|f\rangle = |i\rangle$ in this case an expression for the total decay rate Γ can be identified as:

$$
\Gamma(i \to \text{anything}) = \frac{\text{Im}(\mathcal{M}_{ii})}{m_i} \tag{10.97}
$$

where m_i is the mass of the initial state particle. For a single particle the tree level contribution to \mathcal{M}_{ii} is just the propagator $\Delta_F(k)$ of equation 10.72. For $\varepsilon \to 0$ this function is real except when the particle is on-shell, with the consequence that Im(1/(k² – m² + iε)) ~ δ(k² – m²).

This observation can be generalised for higher-order perturbations. In fact the application of the optical theorem in a quantum field theory can also be demonstrated in terms of Feynman diagrams, where it can also be proved to all orders of perturbation theory by applying 'cutting rules' ([70] pp.232–236, [72] pp.183–196). An example obtained by relabelling the Feynman diagram in figure 10.9 to represent an amplitude for the forward scattering process $\mathcal{X}^+\mathcal{X}^ \rightarrow$ $\mathcal{X}^+\mathcal{X}^-$, with identical incoming and outgoing particles and momenta, via two $\hat{\phi}$ field propagators and a $\hat{\mathcal{Y}}$ field internal loop is shown here in figure 10.10.

Figure 10.10: A Feynman diagram for the forward scattering process $\mathcal{X}^+ \mathcal{X}^ \to \mathcal{X}^+ \mathcal{X}^-$, with a 'cut line' drawn through the intermediate loop propagators of the \hat{y} field.

By careful analysis of the singularities that occur when internal propagators go on-mass-shell under internal loop momenta integrals, twice the imaginary part of the amplitude can be obtained by summing over the 'cutting' possibilities (only one for the diagram in figure 10.10, shown by the vertical dashed line) and replacing the term in the Feynman rule for each propagator that may be simultaneously put on-shell by the cut as:

$$
\frac{i}{k^2 - m^2 + i\varepsilon} \to 2\pi i \,\delta(k^2 - m^2) \tag{10.98}
$$

(with the sign and factors of 2 and i depending on the conventions adopted) before performing the $\int d^4r$ over the loop 4-momentum. Hence the imaginary part of a loop amplitude is obtained by placing the intermediate states on-shell together, as may have been expected from the optical theorem itself since the final states for cross-sections and decay rates, equations 10.96 and 10.97 respectively, consist of on-shell particles. Each way of placing intermediate states on-shell together, as for figure 10.10, is called a 'cut' after Cutkosky, with the above cutting rules providing a method to compute the imaginary part of a transition amplitude in general.

The cutting rules for obtaining the imaginary part of the transition amplitude for a given Feynman diagram can be derived by summing over sets of replacements of each Feynman propagator Δ_F by either Δ_F , Δ_F^* , Δ^+ or Δ^- in the Feynman rules. This calculational tool involves a sum over permutations of selected vertices which determine

the kind of replacement for each Δ_F (see for example [72] p.186). Indeed it can be seen that replacing Δ_F of equation 10.71 with Δ^{\pm} from equation 10.60 incorporates the substitution of equation 10.98 together with the introduction of a factor of $\theta(\pm k^0)$. This latter factor relates to the time ordering of the corresponding vertices and the resulting interpretation as an apparent particle or antiparticle propagating forwards in time between the two vertices. While in equation 10.55 or 10.67 the Feynman propagator was constructed out of two Δ^{\pm} components, here it is taken apart again and a single on-shell part retained.

This on-mass-shell condition is expressed by the δ -function in equation 10.60. On performing the k^0 integral this constraint leads to the form of equation 10.58 which in the present context represents the phase space factor for real final state external particles, on-mass-shell and with positive energy, in cross-section or decay rate calculations. The remaining propagators Δ_F , for example for the two internal ϕ field lines in figure 10.10 are unchanged, representing their usual (non-physical) aid to calculation as described in the previous two sections.

For example for the Feynman diagram in figure 10.10, by adapting equation 10.86 and applying the substitutions from equation 10.98 to the basic Feynman rules of table 10.1 we obtain:

$$
2 \operatorname{Im} \left(\mathcal{M}(\mathcal{X}^+ \mathcal{X}^- \to \mathcal{X}^+ \mathcal{X}^-) \right)
$$

= $g^4 \left(\frac{i}{(p_1 + p_2)^2 - m_\phi^2 + i \varepsilon} \right)^2 \int \frac{d^4 r}{(2\pi)^4} 2\pi i \delta(r^2 - m_\mathcal{Y}^2) 2\pi i \delta((k_1 - r)^2 - m_\mathcal{Y}^2)$
(10.99)

The latter integral can be more easily performed under the substitution of the original 4-momenta q_1 and q_2 , as indicated in figure 10.10, in place of the integral over r, by including a 4-momentum constraint in:

$$
\int \frac{d^4r}{(2\pi)^4} \equiv \int \frac{d^4q_1}{(2\pi)^4} \int \frac{d^4q_2}{(2\pi)^4} (2\pi)^4 \delta^4(q_1+q_2-k_1)
$$

This leads to:

$$
2 \operatorname{Im} \left(\mathcal{M} (\mathcal{X}^+ \mathcal{X}^- \to \mathcal{X}^+ \mathcal{X}^-) \right)
$$

= $-g^4 \left(\frac{i}{(p_1 + p_2)^2 - m_\phi^2 + i \varepsilon} \right)^2 \int \frac{d^4 q_1}{(2\pi)^4} \int \frac{d^4 q_2}{(2\pi)^4}$

$$
2\pi \delta (q_1^2 - m_\mathcal{Y}^2) 2\pi \delta (q_2^2 - m_\mathcal{Y}^2) (2\pi)^4 \delta^4 (q_1 + q_2 - k_1)
$$

= $-g^4 \left(\frac{i}{(p_1 + p_2)^2 - m_\phi^2 + i \varepsilon} \right)^2 \int \frac{d^3 q_1}{(2\pi)^3} \frac{1}{2\omega_{q_1}} \int \frac{d^3 q_2}{(2\pi)^3} \frac{1}{2\omega_{q_2}} (2\pi)^4 \delta^4 (q_1 + q_2 - k_1)$
(10.100)

by applying equation 10.63 to obtain the bottom line with $q_{1,2}^0 > 0$. Here the initial $q_{1,2}^0$ part of the $\int \frac{d^4q_{1,2}}{(2\pi)^4}$ $\frac{d^4q_{1,2}}{(2\pi)^4}$ integrals over $\delta(q_{1,2}^2 - m_{\mathcal{Y}}^2)$ place the momenta $q_{1,2}$ on-shell resulting in integrals of the form $\int \frac{d^3 q_{1,2}}{(2\pi)^3}$ $\frac{d^o\bm{q}_{1,2}}{(2\pi)^3}\frac{1}{2\omega_{\bm{q}}}$ $\frac{1}{2\omega_{q_{1,2}}}$, that is over the relativistic phase space. Together

with the overall $(2\pi)^4 \delta^4(p_F - p_I)$ for 4-momentum conservation implied in the final delta function this identifies the Lorentz invariant phase space factor $d\Phi$ for a two-body y^+y^- final state in equation 10.100. The remaining factor, before the first \int sign, can be identified with $|\mathcal{M}_{fi}|^2$ for the scattering amplitude \mathcal{M}_{fi} of equation 10.85 for the process $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$ at the level of the Feynman diagram depicted in figure 10.4, and hence (swapping the two sides of equation 10.100):

$$
|\mathcal{M}(\mathcal{X}^+\mathcal{X}^- \to \mathcal{Y}^+\mathcal{Y}^-)|^2 \int d\Phi = 2 \operatorname{Im} \left(\mathcal{M}(\mathcal{X}^+\mathcal{X}^- \to \mathcal{X}^+\mathcal{X}^-) \right)
$$
(10.101)

This equation verifies the optical theorem relation of equation 10.94 for the $\mathcal{Y}^+ \mathcal{Y}^-$ final state contribution to the total cross-section in $\mathcal{X}^+ \mathcal{X}^-$ collisions for the Feynman diagram analysis at this level of perturbation theory. A further contribution for a $\mathcal{X}^+ \mathcal{X}^-$ final state can be obtained in a very similar manner based on an intermediate $\mathcal{X}^+ \mathcal{X}^-$ loop, in place of the $\mathcal{Y}^+ \mathcal{Y}^-$ loop, in figure 10.10. The above argument applies to arbitrary loop diagrams and this Feynman diagram approach based on the cutting rules can be used to prove the optical theorem to all orders of perturbation theory ([70] pp.235–236), with care for combinatoric factors and the consistency of the conventions used in general.

Our main point here has been to review the relation between a physical crosssection and an expression linear in a component of an amplitude, namely the imaginary part of the forward scattering amplitude, both in terms of the total cross-section in equation 10.96 and at the level of individual processes as implied in equation 10.101. These expressions relate to the optical theorem and the unitarity constraint which in turn represents the basic property that the total probability must always equal one. This structure will provide a means to connect calculations of the likelihood of scattering processes for the present theory with the techniques of quantum field theory, as we shall describe in section 11.2. In the meantime, in the following section, we assess the nature of basic field interactions in the context of the present theory.
Chapter 11

A Novel Conception of HEP

Processes

11.1 Degeneracy of Spacetime Solutions

In this chapter we consider how the probabilistic nature of quantum phenomena arises in the context of the present theory, and in particular in the environment of laboratory experiments. The main goals will be to relate the calculation of cross-sections, for example, for the present theory with the corresponding formalism of QFT and to address the related question concerning the nature of particle phenomena generally. Here the probability for a particular process will be a measure of the degeneracy of field states describing the mathematical form of a particular 4-dimensional geometry, that is through the symmetry of possible local reinterpretations of fields such as the gauge field $Y_\mu(x)$ or fermion field $\psi(x)$ (denoted without 'hats', since these are not quantum field operators here) under the same Einstein tensor $G^{\mu\nu}(x)$. The spacetime geometry is locally completely insensitive to reinterpretations of the underlying fields, that is exchanges between components of the fields $\delta Y(x) \leftrightarrow \delta \psi(x)$, which leave $G^{\mu\nu}(x)$ locally unchanged, while the geometric contracted Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$ remains globally valid.

The field interactions proceed by a kind of 'Chinese whispers' of field indistinguishability, as a degenerate mathematical possibility underlying the spacetime geometry. This leads directly to the indeterminate nature characteristic of quantum phenomena. Probabilities, in the form of cross-sections and decay rates, will then arise in proportion to the sum of the 'number of ways' in which such underlying field descriptions are possible.

We begin however by considering a particular case for which $G^{\mu\nu}(x)$ is a function of a single internal gauge field. Based on the breaking of the full symmetry of $L(\hat{v}) = 1$ into external and internal parts over an extended base manifold M_4 a relationship between the external geometry described by $G^{\mu\nu}(x)$ and internal gauge fields $Y_\mu(x)$ over $x \in M_4$ was developed in chapters 2–5. Structures are identified analogous to those of Kaluza-Klein theory in leading to equation 4.16-4.17, which is conjectured to arise out of the geometric constraints of the present theory culminating in equation 5.20, for which the practical normalisation convention $\chi = \frac{\kappa}{2}$ may be adopted. For the present theory with an external linear connection $\Gamma^{\rho}{}_{\mu\nu}(x)$ and an internal gauge field $A_\mu(x)$, deriving from an internal Abelian U(1) gauge symmetry, the relation between the external Riemannian curvature and internal gauge curvature is expressed in equation 5.22, via the above connection with classical Kaluza-Klein theory, and as reproduced here:

$$
-\frac{1}{\kappa}G^{\mu\nu} = +F^{\mu}_{\ \rho}F^{\rho\nu} + \frac{1}{4}g^{\mu\nu}F_{\rho\sigma}F^{\rho\sigma}
$$
 (11.1)

Under the Bianchi identity $G^{\mu\nu}_{;\mu} = 0$ this relation *implies* the source-free homogeneous Maxwell equation 5.30, as explained in section 5.2. Here, within the context of the present theory the generator of the internal $U(1)_Q$ gauge symmetry of electromagnetism is identified with the element \dot{S}^1_l within the set of E₆ Lie algebra actions as described in section 8.2. The electromagnetic gauge field $A_\mu(x)$, associated with the U(1)_Q generator, is in turn identified with the field $\tilde{A}_{\mu}(x)$ as described in and following equation 8.58. We next consider the form of the free field $A_\mu(x)$ as a solution of Maxwell's equation, which can be written in terms of the gauge field itself as:

$$
\Box A^{\mu}(x) = 0 \tag{11.2}
$$

The energy-momentum tensor for the electromagnetic field can be obtained directly through the definition $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$ with the geometry $G^{\mu\nu}(x)$ determined in terms of the electromagnetic field tensor $F^{\mu\nu}$ according to equation 11.1 above. (A normalisation convention setting $\kappa = 8\pi G_N = -1$ might also be adopted in order to emphasise the equivalence of both sides in this definition $T^{\mu\nu} := G^{\mu\nu}$. Here the electromagnetic gauge field itself is analysed under the assumption of an approximately flat spacetime. A real field $A^{\mu}(x)$ can be expressed in terms of Fourier components, which in terms of trigonometric functions and a single 4-vector k takes the form:

$$
A^{\mu}(x) = A_c^{\mu}(\mathbf{k}) \cos k \cdot x + A_s^{\mu}(\mathbf{k}) \sin k \cdot x \tag{11.3}
$$
\n
$$
\begin{bmatrix} A_c^{\mu} & \cdots & A_s^{\mu} \end{bmatrix} \tag{11.3}
$$

$$
\equiv \frac{A_c^{\mu}}{\cos\left(\tan^{-1}\frac{A_s^{\mu}}{A_c^{\mu}}\right)}\cos\left(k \cdot x - \tan^{-1}\frac{A_s^{\mu}}{A_c^{\mu}}\right) \quad (11.4)
$$

$$
= A^{\mu}(\mathbf{k}) e^{-ik \cdot x} + A^{\mu *}(\mathbf{k}) e^{+ik \cdot x}
$$
 (11.5)

$$
\equiv C^{\frac{1}{2}} \varepsilon_r^{\mu}(\mathbf{k}) A_r(\mathbf{k}) e^{-ik \cdot x} + C^{\frac{1}{2}} \varepsilon_r^{\mu}(\mathbf{k}) A_r^*(\mathbf{k}) e^{+ik \cdot x}
$$
(11.6)

In equation 11.4 the gauge field is expressed in terms of a single cosine function, that is in the form $A^{\mu}(x) = A^{\mu} \cos(k \cdot x + \lambda)$, with no sum implied over the index $\mu = 0, 1, 2, 3$, using the trigonometric identity $\cos \alpha \cos \beta \pm \sin \alpha \sin \beta = \cos(\alpha \mp \beta)$. Equation 11.5 follows from equation 11.3 with $A^{\mu}(\mathbf{k}) = \frac{1}{2}(A_c^{\mu}(\mathbf{k}) + iA_s^{\mu}(\mathbf{k}))$, and with $A^{\mu}(\mathbf{k})$ expressed as $C^{\frac{1}{2}} \varepsilon_r^{\mu}(\mathbf{k}) A_r(\mathbf{k})$ in the final line.

Hence for each 4-vector k each of the four vector components of $A^{\mu}(x)$ can be associated with the real coefficients $A_c^{\mu}(\mathbf{k})$ and $A_s^{\mu}(\mathbf{k})$ or the complex coefficients $A^{\mu}(\mathbf{k})$ and $A^{\mu*}(\mathbf{k})$, either pair of which can be considered to be independent in terms of possible field interactions as described in the following section. With all physical phenomena being invariant under spacetime translations, and hence with no preferred set of coordinates $\{x\}$, and hence with $A_s^{\mu}(\mathbf{k}) \neq 0$ in general, equation 11.4 forms a relatively cumbersome expression for the free field and will not be employed further.

The constant coefficient $C^{\frac{1}{2}}$ is introduced with the square root in equation 11.6 since each factor of $A^{\mu}(x)$ appears quadratically in the expression for $G^{\mu\nu}$ in equation 11.1. In line with textbook solutions to Maxwell's equations, and anticipating the quantum field analysis, four polarisation vectors $\varepsilon_r^{\mu}(\mathbf{k})$ are introduced, representing a 4-vector object for each of $r = 0, 1, 2, 3$ (see for example [71] section 5.1). These provide a constant basis for the 4-vector $A^{\mu}(\mathbf{k})$, analogous to the variable tetrad components $e^{\mu}_{a}(x)$ describing four vector fields for $a = 0, 1, 2, 3$ as a basis for tangent 4-vectors to the manifold M_4 , as employed in section 5.3 for example. A standard choice of basis is such that the polarisation vectors are real and orthogonal, with respect to the Minkowski metric $\eta_{\mu\nu} = \text{diag}(1, -1, -1, -1)$, that is:

$$
\varepsilon_r(\mathbf{k}) \cdot \varepsilon_s(\mathbf{k}) = \varepsilon_{r\,\mu}(\mathbf{k}) \varepsilon_s^{\mu}(\mathbf{k}) = \begin{cases} \delta_{rs} & r = 0 \\ -\delta_{rs} & r = 1, 2, 3 \end{cases}
$$
(11.7)

More specifically a standard basis in a given reference frame can be taken with:

$$
\varepsilon_0^{\mu}(\mathbf{k}) = (1, 0, 0, 0) \tag{11.8}
$$

$$
\varepsilon_r^{\mu}(\mathbf{k}) = (0, \varepsilon_r(\mathbf{k})) \qquad r = 1, 2, 3 \tag{11.9}
$$

with
$$
\mathbf{k} \cdot \boldsymbol{\varepsilon}_r(\mathbf{k}) = 0
$$
 $r = 1, 2$ (11.10)

and
$$
\varepsilon_3(\mathbf{k}) = \mathbf{k}/|\mathbf{k}|
$$
 (11.11)

The $r = 0$ case is the scalar, or timelike, polarisation vector, while $r = 1, 2$ represent transverse polarisation vectors and the case for $r = 3$ is called the longitudinal polarisation vector. (With respect to a 3D scalar product with metric $\delta_{ij} = \text{diag}(+1, +1, +1)$) the 3-vector parts of the transverse polarisation vectors satisfy $\varepsilon_r(\mathbf{k}) \cdot \varepsilon_s(\mathbf{k}) = +\delta_{rs}$ for $r = 1, 2, 3$ according to equation 11.7; while equation 11.10 for $r = 1, 2$ is also valid for the 4D scalar product with $k \cdot \varepsilon_r(\mathbf{k}) = k_\mu \varepsilon_r^\mu(\mathbf{k}) = 0$.

The full set of four polarisation states $\varepsilon_r^{\mu}(\mathbf{k})$ for $r = 0, 1, 2, 3$ provides a Lorentz covariant description for the 4-component vector field $A^{\mu}(x)$ and suggests the possibility of four kinds of photon states corresponding to these four degrees of freedom. However, in the standard theory, the requirement of gauge invariance, which allows some field excitations to be transformed to zero, together with the massless condition $k^2 = 0$ for the free electromagnetic field, result in there being only *two* physical photon states corresponding to the transverse polarisation states ε_1^{μ} $\frac{\mu}{1}(\boldsymbol{k})$ and ε_2^{μ} $\frac{\mu}{2}(\boldsymbol{k}).$

Maxwell's equation in the form of equation 11.2 is obtained from equation 5.30 under the Lorenz gauge condition $\partial_{\mu}A^{\mu} = 0$, as described for the inhomogeneous case of equations 3.91 and 3.92 in section 3.5. In turn equation 11.6 forms a solution of the free field case of equation 11.2 provided that the massless condition $k^2 = 0$ holds. Further the Lorenz gauge condition itself, applied to equation 11.6, requires that $k_{\mu} \varepsilon^{\mu} = 0$ and hence, from equations 11.8–11.11, the transverse polarisation states are clearly permitted.

When substituted into equation 11.1 the gauge field $A^{\mu}(x)$ of equation 11.6, with $A_r(\mathbf{k}) \neq 0$ for either $r = 1$ or $r = 2$ only, yields a large number of terms, most of which are zero due to the conditions $k \cdot \varepsilon_{1,2}(k) = 0$, for these transverse states, and $k^2 = 0$ (for example no $F_{\rho\sigma}F^{\rho\sigma}$ terms remain) leading to:

$$
T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu} = +C k^{\mu} k^{\nu} \left(2|A_r|^2 - A_r^2 e^{-2ik \cdot x} - A_r^{*2} e^{+2ik \cdot x} \right)
$$

= +2C k^{\mu} k^{\nu} \left(|A_r|^2 - \left(\text{Re}(A_r^2) \cos 2k \cdot x + \text{Im}(A_r^2) \sin 2k \cdot x \right) \right)
= +2C k^{\mu} k^{\nu} |A_r|^2 \left(1 + \cos(2k \cdot x + \alpha) \right) (11.12)

The form of the final line, with $\alpha \in \mathbb{R}$, follows by a similar argument that led to equation 11.4, in terms of a single real cosine function. Taking the 4-vector $k = (k^0, 0, 0, k^0)$, representing the propagation of the electromagnetic wave in the direction of the x^3 coordinate (with transverse polarisation vectors ε_1^{μ} $\ell_1^{\mu}(\mathbf{k}) = (0, 1, 0, 0)$ and ε_2^{μ} $\frac{\mu}{2}(\mathbf{k}) = (0, 0, 1, 0)$ for example), the variable part of $T^{\mu\nu}(x)$ is sketched alongside that for the gauge field $A^{\mu}(x)$ in figure 11.1 as projected onto the spatial coordinate x^3 on M_4 .

Figure 11.1: The energy-momentum tensor $T^{\mu\nu}(x) := -\frac{1}{\kappa} G^{\mu\nu}(x)$ is modulated by a non-negative cosine function as depicted above, corresponding to an electromagnetic vector field $A^{\mu}(x)$ in the form of a plane wave.

Although the field function $A^{\mu}(x)$ is presented in the Lorenz gauge, the form of the Einstein tensor $G^{\mu\nu(x)}$ is gauge invariant. Indeed this was one of the conditions used to derive the relation between the external and internal geometry, as described for example in the discussion following equation 5.13 in section 5.1, in leading to equation 11.1 itself. Hence in turn the energy-momentum tensor $T^{\mu\nu}(x)$, of equation 11.12 and figure 11.1, is naturally gauge invariant.

As for the electric E_i and magnetic B_i field components of the gauge invariant electromagnetic field tensor F in equation 5.23, the geometry of the Einstein tensor $G^{\mu\nu}(x)$ in equation 11.12 represents an unambiguous physical feature associated with the electromagnetic wave. Further, as discussed shortly after equation 5.28 the scalar curvature R associated with any electromagnetic field vanishes, with the Ricci tensor $R^{\mu\nu}$ hence identified with the Einstein tensor $G^{\mu\nu}$. Hence equation 11.12 describes a 'wave of Ricci curvature', which is complementary to the usual notion of a gravitational wave, with the latter composed of purely Weyl curvature in the Ricci vacuum as described after equation 5.44, also in section 5.2.

Indeed under the assumption of an approximately flat spacetime, as employed for equation 11.14 below, and with $R^{\mu\nu} = G^{\mu\nu}$ given by equation 11.12 it can be seen from equation 5.44 that $K^{\rho\sigma\mu} = 0$, that is the source of Weyl curvature vanishes for this geometry. As implied in the discussion before equation 3.69 the Weyl curvature vanishes for any conformally flat geometry, and hence a metric of the form:

$$
g_{\mu\nu}(x) = (1 + \beta \cos 2k \cdot x)\eta_{\mu\nu} \tag{11.13}
$$

for a small value of $\beta \in \mathbb{R}$, provides a candidate solution underlying an Einstein tensor in the form of equation 11.12. Indeed, assuming the Levi-Civita connection of equation 3.53, via equations 3.73 and 3.74 it can be seen that the scalar curvature R vanishes for such a metric if $k^2 = 0$, and that the resulting $G^{\mu\nu}(x) = 4\beta k^{\mu}k^{\nu}\cos 2k \cdot x$, to first order in β , exhibits a corresponding oscillatory behaviour, although more work is needed to obtain the precise form for a metric underlying the Einstein tensor of equation 11.12.

The physical spacetime curvature described by $G^{\mu\nu}(x)$ in equation 11.12 is assumed to be very small. As described in section 5.2, alongside equation 5.22, while the Einstein tensor is theoretically directly related to the internal gauge field, in the form of equation 11.12 for example, in units appropriate for laboratory measurements the Einstein equation can be written $G^{\mu\nu} = -\kappa T^{\mu\nu}$, where the normalisation constant κ is a very small number. Hence while the energy-momentum carried by the electromagnetic wave may be readily detected the distortion of the spacetime geometry away from Minkowski flatness is extremely small and utterly unobservable via any direct means. In turn the plane wave description of equations 11.3–11.6, modelled on the flat spacetime case, can be used to a very good approximation. The divergence of the Einstein tensor in equation 11.12 can be expressed in terms of the energy-momentum tensor in this approximately flat spacetime limit with $T^{\mu\nu}_{;\mu} \to T^{\mu\nu}_{;\mu}$ in Cartesian coordinates, as described in the opening of section 5.2. Consistent with the Bianchi identity $G^{\mu\nu}_{;\mu} = 0$ this object can be seen to vanish, due to the condition $k^2 = 0$, as would be expected:

$$
T^{\mu\nu}_{\ \ ,\mu} = 2C \, k^{\mu} k^{\nu} \, |A_r|^2 \Big(-2k_{\mu} \sin(2k \cdot x + \alpha) \Big) = 0 \tag{11.14}
$$

In the context of the present theory while the polarisation requirement $k_{\mu} \varepsilon^{\mu} = 0$ can again be seen to be a consequence of imposing the Lorenz gauge condition on such plane wave solutions for the field $A^{\mu}(x)$, the 'momentum' requirement $k^2 = 0$ is a consequence of the necessity for the free field solution to satisfy the geometric Bianchi identity as for equation 11.14. That is as a plane wave the free field is necessarily 'massless' in order to identify a consistent solution for $G^{\mu\nu}(x)$ in the form of equation 11.1, that hence might occur in nature. Equivalently the requirement $k^2 = 0$ could be considered to be a consequence of the homogeneous Maxwell equation 11.2, which itself is a direct consequence of the Bianchi identity $G^{\mu\nu}_{\ \ ;\mu} = 0$ given equation 11.1, as described for equation 5.30. Equation 11.14 also of course directly implies energy-momentum conservation, $T^{\mu\nu}_{\mu\nu} = 0$, for the gauge field described in equation 11.6 as employed in equation 11.12.

Recalling that in general relativity the components $T^{\mu\nu}(x)$ represent the energymomentum *density*, the field $A^{\mu}(x)$ in a spatial volume V carries 4-momentum P^{μ} which may be expressed as:

$$
P^{\mu} = \int_{V} d^{3}x \, T^{\mu 0} \tag{11.15}
$$

In the present theory the energy-momentum is always fundamentally determined by the Einstein tensor $G^{\mu\nu}(x)$ through the Einstein equation $T^{\mu\nu} := G^{\mu\nu}$. As also described

in the opening paragraphs of section 5.2 this is in contrast to the Lagrangian approach for which an energy-momentum tensor $t^{\mu\nu}$ can be defined giving rise to a conserved 4-momentum, in the form of equation 11.15, as described in equation 3.102 and the subsequent discussion of section 3.5.

The components P^{μ} are locally four conserved quantities which transform amongst each other covariantly as a 4-vector under Lorentz transformations. Setting $|A_r|^2 = 1$, with the real coefficient $C^{\frac{1}{2}}$ taking care of the field normalisation in equation 11.6, and substituting the top line of equation 11.12 into the above expression, taking into account the vanishing of the integral of the $e^{\pm i k \cdot x}$ terms for suitably defined boundary conditions for the volume V , yields:

$$
P^{\mu} = \int_{V} d^{3}x \, 2C \, k^{0} k^{\mu} = 2 \, V \, C \, k^{0} k^{\mu} \tag{11.16}
$$

Hence by setting the coefficient $C = \frac{1}{2V k^0}$ the 4-vector k^{μ} in the Fourier component can be identified with the 4-momentum P^{μ} of the field in the volume V. Such an object with field values localised to within the volume V might naively be considered to represent a 'particle', although a less simplistic particle concept that emerges in the present theory will be described in section 11.3.

Hence in turn $C^{\frac{1}{2}} = \frac{1}{\sqrt{2L}}$ $\frac{1}{2V k^0}$ is the normalisation required in equation 11.6, given the transverse polarisation vectors described in equations 11.7–11.11, taking $|A_r| = 1$ and considering the field in the volume V_1 to possess 4-momentum $P^{\mu} \equiv k^{\mu}$. The origin of this field normalisation factor $C^{\frac{1}{2}}$ here therefore is in the interpretation of the Einstein tensor as the energy-momentum, and in particular with $G^{00}(x)$ as the energy density in the local reference frame $T^{00} := -\frac{1}{\kappa} G^{00} \sim C k^0 k^0 \propto \frac{k^0}{V}$ relating the k⁰ component of the Fourier expansion of the $A^{\mu}(x)$ gauge field directly to the physical energy P^0 carried by the field (that is, the classical Hamiltonian H). This construction is independent of the spatial volume V which, being arbitrary within the choice of the boundary conditions, should cancel in all calculations of physical quantities when interactions are considered, as it does for cross-section calculations in QFT as described in the discussion following equation 10.3.

In the present theory the energy of a real field, such as $A^{\mu}(x)$, is obtained directly by substitution of the field into the appropriate expression for the right-hand side of $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$. Taking a complex-valued expression for the field $A^{\mu}(x)$ in the form of the first term on the right-hand side of equation 11.6, for example, leads to the subsequent expression for the energy-momentum tensor:

$$
A^{\mu}(x) = C^{\frac{1}{2}} \varepsilon_r^{\mu}(\mathbf{k}) A_r(\mathbf{k}) e^{-ik \cdot x}
$$
 (11.17)

$$
\Rightarrow T^{\mu\nu} = -C k^{\mu} k^{\nu} A_r^2 e^{-2ik \cdot x} \tag{11.18}
$$

which, while consistent with $T^{\mu\nu}_{,\mu} = 0$, is a *complex* tensor and hence does not represent a real energy-momentum tensor $T^{\mu\nu}$, or a real geometric tensor $G^{\mu\nu}$. In addition here $A^{\mu}(x)$ is required in any case to be real in order to represent the real components of a $U(1)_Q$ Lie algebra-valued vector field, that is a classical macroscopic gauge field.

Alternatively, the first term on the right-hand side of equation 11.3, for example, is real and does, alone, produce a real energy-momentum tensor:

$$
A^{\mu}(x) = C^{\frac{1}{2}} \varepsilon_r^{\mu}(\mathbf{k}) A_{cr}(\mathbf{k}) \cos k \cdot x \qquad (11.19)
$$

$$
\Rightarrow T^{\mu\nu} = +C k^{\mu} k^{\nu} A_{cr}^2 \sin^2 k \cdot x \tag{11.20}
$$

as a special case of equation 11.12 (with $A_r = \frac{1}{2}A_{cr} \in \mathbb{R}$).

All three expressions for $A^{\mu}(x)$ in equations 11.3 (or 11.6), 11.17 and 11.19 also necessarily satisfy Maxwell's equation $\Box A^{\mu} = 0$ since this is implicit in the identity $G^{\mu\nu}{}_{;\mu} = 0$ when applied to equation 11.1 as was described in equation 5.29 and the subsequent discussion as reviewed above. In all cases a solution with A^{μ} dependent upon the Fourier mode 4-vector k expressed in the Lorenz gauge requires a polarisation vector with $k_{\mu} \varepsilon^{\mu} = 0$, and with $G^{\mu\nu}$ in the form of equation 11.1 the geometric Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$ implies $k^2 = 0$.

For the standard treatment of a massive vector field with $k^2 = m^2 \neq 0$, as for the case of a massive gauge vector boson, the plane wave expansion in the form of any of equations 11.3–11.6 can again be employed, and the Lorenz gauge condition again implies $k_{\mu} \varepsilon^{\mu} = 0$ for the polarisation vector. In this case however the remaining gauge freedom, subject to $\partial_{\mu}A^{\mu} = 0$, cannot be used to uncover a cancellation between the scalar and longitudinal components of polarisation. Hence for massive gauge bosons there are three possible states, with the longitudinal degree of freedom appended to the two transverse polarisation states. In this case Maxwell's equation is replaced by an expression incorporating a mass term:

$$
(\Box + m^2)A^{\mu} = 0 \tag{11.21}
$$

In the context of the present theory on substituting the free field in the form of equation 11.6 into the expression $G^{\mu\nu} = f(A)$ of equation 11.1 the Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$ in the form of equation 11.14 is no longer satisfied for this new case with $k^2 \neq 0$. This suggests that the direct relationship between the Einstein tensor and a gauge field of the form of equation 5.20 and 5.31, as employed for equation 11.1, no longer holds, but rather a more general expression is to be sought, as suggested by the form of equation 5.32 in section 5.2. In the present context this latter expression $G^{\mu\nu} = f(A, \hat{\boldsymbol{v}})$ implicitly incorporates the consequences of *interactions* between the gauge field $A^{\mu}(x)$ and components of the temporal flow under the full form $L(\hat{\boldsymbol{v}}) = 1$.

Indeed in subsection 8.3.3 it has been suggested that in the present theory gauge boson masses arise through an impingement of the corresponding internal symmetry on the external vector $h_2 \equiv v_4 \in TM_4$ of equation 8.72, which forms the components of a 'vector-Higgs'. This argument was constructed in part by analogy with technicolor models, with longitudinal components for massive gauge bosons obtained when the propagators are corrected for the field interactions, as described following equation 8.73. In the context of the present theory while equation 11.1 together with the identity $G^{\mu\nu}_{\ \ ;\mu} = 0$ implies equation 11.2, the form of $G^{\mu\nu} = f(A, \hat{\boldsymbol{v}})$ under the same identity is expected to be consistent with equation 11.21.

The $A^{\mu}(x)$ gauge field associated with electromagnetism is in fact massless. In the present theory in terms of the corresponding $U(1)_Q \subset E_6$ internal symmetry this property is attributed to the fact that the $U(1)_Q$ generator \dot{S}^1_l does not impact upon the external $v_4 \in TM_4$ components of v_{27} , as presented for example in equation 8.71. However there are interactions between $A^{\mu}(x)$ and other temporal components which suggest that the free field expansion and equation 11.1 will not represent the full picture.

Indeed, in the present theory the gauge field $A^{\mu}(x)$ and the associated internal $U(1)_Q$ symmetry are not considered as basic entities in themselves, rather they are introduced since they act on components within the form $L(\hat{v}) = 1$. These latter components include the Dirac spinors ψ of equations 9.43 and 9.46, as identified in the components of v_{56} in the extension to the E₇ symmetry of the full form $L(v_{56}) = 1$ in section 9.2, which unlike the v_4 components do transform non-trivially under the $U(1)_Q$ action. In principle these temporal components provide a greater freedom for building the spacetime geometry, now with an underlying degeneracy of possible $\delta A(x) \leftrightarrow \delta \psi(x)$ field 'redescriptions', always consistent with the Bianchi identity $G^{\mu\nu}_{;\mu} = 0$ for the external spacetime.

Objects transforming as a 4-vector can be constructed out of Dirac spinors in the form of $\overline{\psi}\gamma^{\mu}\psi$ via the conjugate field $\overline{\psi} = \psi^{\dagger}\gamma^{0}$ as introduced for equation 3.96 in section 3.5. As for the standard theory the relationship between ψ and $\overline{\psi}$ is expected to relate to the dynamics of fermions and antifermions for physical particle states propagating in spacetime. For the present theory 'interactions' between the vector field $A^{\mu}(x)$ and a fermion field $\psi(x)$ take the form of vector field 'redescriptions' as provisionally sketched in figure 11.2.

Figure 11.2: Field redescriptions: (a) The same function of spacetime is associated with the field $\overline{\psi}(x)\gamma^{\mu}\psi(x)$ before time $t_2 \in T$ and with the field $A^{\mu}(x)$ at later times, here in the spatial volume V. (b) The field $A^{\mu}(x)$ in the spacetime volume VT is redescribed as the field $\overline{\psi}(x)\gamma^{\mu}\psi(x)$ from time $t_1 \in T$.

While the field function is relabelled $A^{\mu}(x) \to \overline{\psi}(x) \gamma^{\mu} \psi(x)$ at time $x^{0} = t_{1}$ in figure 11.2(b), the *function* form itself is independent of the choice of t_1 , as is the local geometric structure $G^{\mu\nu} = f(A, \psi)$ in spacetime. Figure 11.2(b) does not represent the field $A^{\mu}(x)$ 'turning into' the field $\overline{\psi}(x)\gamma^{\mu}\psi(x)$ at time t_1 , rather this possible redescription is everywhere *implicit* in $A^{\mu}(x)$ as a function on M_4 . For example the plane wave described by $A^{\mu}(x)$ in figure 11.1 might be redescribed in terms of the field $\overline{\psi}(x)\gamma^{\mu}\psi(x)$ at any time t_1 with the external form of $T^{\mu\nu} := G^{\mu\nu}$ remaining unchanged throughout the spacetime volume VT .

This notion of field indistinguishability is closely analogous in spirit to the 'arithmetic indistinguishability' of the multi-dimensional form $L(\mathbf{v}) = 1$ from the original one-dimensional temporal flow within which the form $L(\mathbf{v}) = 1$ is ever implicit, as described in section 2.1. That is one-dimensional time s innately contains possible 'redescriptions', such as $s^2 = (x^1)^2 + (x^2)^2 + (x^3)^2$, which may potentially be interpreted as geometric or spatial structures. Further, when projected onto the base manifold M_4 in the full theory the form $L(\hat{\boldsymbol{v}}) = 1$ will provide constraints on possible

field redescriptions, represented by $A^{\mu}(x) \leftrightarrow \overline{\psi}(x) \gamma^{\mu} \psi(x)$ here, as will be described for equation 11.33 below for example.

Although locally indistinguishable, the possibility of local field redescriptions such as depicted in figures $11.2(a)$ and (b) will lead to *globally* distinguished and observable phenomena on M_4 . This includes the possible outcomes of a 'Schrödinger's cat' type experiment, as will be described in section 11.4. This is possible since different field descriptions point towards a different set of subsequent possible field redescriptions propagating in the broader spacetime environment, always under the constraint $G^{\mu\nu}{}_{;\mu} = 0$. The relative probability for a specific observable effect will depend directly upon the degeneracy of the local underlying possible field descriptions, as we shall explore in the following section.

In the full theory the spacetime geometry with metric $g_{\mu\nu}(x)$ and Einstein tensor $G^{\mu\nu}(x)$ are continuous and smooth over M_4 and have 'surveillance' over the other fields, as described in section 5.2 in the discussion shortly before equation 5.44 for example. As well as shaping the equations of motion for macroscopic fields and entities this surveillance will also constrain the form of microscopic field interactions and exchanges. While the original form of $T^{\mu\nu} := G^{\mu\nu} = f(A)$ is expected to be associated with photon states in some way, further redescriptions of a form suggested by the sketches of figure 11.2 will ultimately introduce matter terms $T^{\mu\nu}$ primarily associated with the spinor ψ field components, to be associated with electron states for example.

The precise mathematical form of the field redescriptions remains to be fully understood. However this structure, as pictured in figure 11.2, brings to mind Huygen's principle for the description of a field at a later time as propagated from earlier times and the form of the retarded propagator, closely relating to $\Delta_R(x-y)$ of equation 10.79 for the scalar case. Here however, rather than fields propagating through a pre-existing spacetime background, the 4-dimensional spacetime M_4 with geometry $G^{\mu\nu} = f(A, \psi)$ is constructed in terms of the fields. In any case we provisionally represent the structures of figure $11.2(a)$ and (b) respectively by the mathematical relations:

$$
A^{\mu}(x) = \int d^4y D_r^{\mu\nu}(x-y) \overline{\psi}(y) \gamma_{\nu}\psi(y) \qquad (11.22)
$$

$$
A^{\mu}(x) = \int d^4y D_a^{\mu\nu}(x-y) \overline{\psi}(y) \gamma_{\nu}\psi(y) \qquad (11.23)
$$

The role of the functions $D_r^{\mu\nu}(x-y)$ and $D_a^{\mu\nu}(x-y)$ is hence to provide a more rigorous account of the field exchanges depicted graphically and somewhat naively in figure 11.2. Here, by analogy with the case of standard electrodynamics, the 'redescription function' $D_r^{\mu\nu}(x-y)$ is analogous to the retarded propagator $D_R^{\mu\nu}$ $R^{\mu\nu}(x-y)$ for a vector field. Such a redescription may also take place 'into the past', that is by analogy with the advanced propagator $D_A^{\mu\nu}$ $A^{\mu\nu}(x-y)$, equivalent to the field exchange $A^{\mu}(x) \to \overline{\psi}(x)\gamma^{\mu}\psi(x)$ forward in time, as described in terms of $D_a^{\mu\nu}(x-y)$ in equation 11.23 and depicted in figure 11.2(b).

Hence we provisionally identify the functions $D_{r,a}^{\mu\nu}(x-y)$ in equations 11.22 and 11.23 with the propagators $D_{R,A}^{\mu\nu}(x-y)$. The vector retarded propagator for the massless gauge boson case can be defined in terms of the corresponding scalar propagator $\Delta_R(x-y)$ as:

$$
D_R^{\mu\nu}(x - y) = \lim_{m \to 0} [-g^{\mu\nu} \Delta_R(x - y)] \tag{11.24}
$$

with
$$
\Box_x D_R^{\mu\nu}(x-y) = g^{\mu\nu} \delta^4(x-y) \qquad (11.25)
$$

hence following from equation 10.81, and with a similar construction for the advanced propagator. In turn equation 11.22 implies:

$$
\Box A^{\mu} = \overline{\psi}\gamma^{\mu}\psi \tag{11.26}
$$

that is Maxwell's equation $F^{\mu\nu}_{;\mu} = j^{\nu}$ of equation 3.91 for the inhomogeneous case with source current $j^{\mu} = \overline{\psi} \gamma^{\mu} \psi$.

This relation, deriving from equation 11.22, is incompatible with the classical expression of equation 11.1, which led to $\Box A^{\mu} = 0$. This generalisation from equation 11.2 with the addition of the source term j^{μ} on the right-hand side is analogous to the extension with a mass term m^2 on the left-hand side in equation 11.21, in both cases arising out of interactions between the gauge field $A^{\mu}(x)$ and components of the temporal flow under $L(\hat{v}) = 1$. In both cases this involves opening up a more general relation between the spacetime geometry and the internal fields, and correspondingly more general properties of matter described by the energy-momentum tensor $T^{\mu\nu} := G^{\mu\nu} = f(Y, \hat{v})$, as outlined in the discussion around equation 5.32 in section 5.2. For the generalisation to $G^{\mu\nu} = f(A, \psi)$ considered here the evolution of the both fields, $A^{\mu}(x)$ and $\psi(x)$, will in turn be shaped in conformity with the Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$ (which led to the source-free Maxwell equation for the electromagnetic field $A^{\mu}(x)$ alone in equation 11.2).

From equations 10.76 and 10.79 the scalar retarded propagator as appearing in equation 11.24 can be written as:

$$
\Delta_R(x-y) = -i\theta(x^0 - y^0) \int \frac{d^4k}{(2\pi)^4} \,\varepsilon(k^0) \, 2\pi \, \delta(k^2 - m^2) e^{-ik \cdot (x-y)} \tag{11.27}
$$

This function contains similar features to those required for $D_r^{\mu\nu}(x-y)$ in equation 11.22, including a θ-function for the temporal ordering of the field redescription, which takes place at time t_2 in figure 11.2(a). Further, this propagator is employed to obtain field solutions in the form of equation 10.83, which also applies for classical fields as described at the end of section 10.4, and which is closely analogous to equation 11.22 for the field redescription above.

As used above in deriving equation 11.26 the retarded propagator $\Delta_R(x-y)$ of equation 11.27 satisfies equation 10.81. Indeed in QFT the propagators $\Delta_{F,R,A}(x - y)$ may be introduced as inverse functions for the operator $(\Box_x + m^2)$ in equation 10.81, with appropriate boundary conditions, motivated by the search for solutions to differential equations of motion for the fields, such as equation 10.84. These equations of motion are themselves derived via the Euler-Lagrange equation 3.89 given an original postulated Lagrangian as the starting point, which led for example to equation 10.24 (incorporating equation 10.84) for the scalar model. A very similar situation applies for the QFT employed for the Standard Model in particle physics.

In the standard theory the complete Lagrangian, including the interaction terms, is subject to the Euler-Lagrange equation collectively. For example the combined Maxwell and Dirac Lagrangian, given by equation 3.96 for the internal $U(1)_Q$

case, under variation of the gauge field $A^{\mu}(x)$ and its spacetime derivatives $\partial_{\nu}A^{\mu}(x)$ leads, in the Lorenz gauge, directly to:

$$
\Box A^{\mu} = \overline{\psi} \gamma^{\mu} \psi =: j^{\mu} \tag{11.28}
$$

as implied in equations 3.97 and 3.98. In order to arrive at this expression the variation of *both* the $F_{\mu\nu}F^{\mu\nu}$ and $j^{\mu}A_{\mu}$ parts implied in equation 3.96 are mutually related by appearing in the same Lagrangian object under a single Euler-Lagrange equation.

By contrast the form of the redescription propagator $D_r^{\mu\nu}(x-y)$ of equation 11.22 is not motivated on the grounds of finding solutions for equations of motion such as equation 11.28, but rather in the present theory it is conceptually motivated on the grounds of a degeneracy of field solutions under the construction of the spacetime geometry $G^{\mu\nu} = f(A, \psi)$ over M_4 . In fact here there is no similar direct expression with an explicit source term for the microscopic case, as there is in the standard theory with equation 11.28 above. In the present theory simple differential equations such as equation 11.26 arise as a consequence of the possibility of mutual field redescriptions at the microscopic level. However, apparent source terms in these expressions might be identified which are reminiscent of those seen in the field equations of motion for the Standard Model. A generalisation of the gauge-fermion field interactions described in equations 11.22–11.26 for non-Abelian gauge symmetries for comparison with the general case of equations 3.97 and 3.98 in section 3.5 could also be considered.

Here, rather than an interaction Lagrangian or Hamiltonian relating the different fields as for QFT, the form of temporal flow $L(\hat{v}) = 1$ places mutual constraints on field values and provides selection rules for possible 'transitions' linking possible initial, intermediate and final states. Here the field interaction terms appear not in a single Lagrangian function but rather through a range of constraint equations, which may be provisionally listed as:

$$
L(\hat{\mathbf{v}}) = 1; \qquad D_{\mu}L(\hat{\mathbf{v}}) = 0; \qquad G^{\mu\nu} = f(Y); \qquad G^{\mu\nu}_{;\mu} = 0 \tag{11.29}
$$

Of these $L(\hat{v}) = 1$, as a scalar invariant, is perhaps most closely related to a standard Lagrangian, however in being constrained to the fixed scalar value 1 further field interactions are implied in the terms of $D_{\mu}L(\hat{v}) = 0$. The third of these constraints is the relation between the external geometry and internal degrees of freedom consisting purely of gauge fields, that is equation 5.20, and relates closely to Kaluza-Klein theories as described in section 5.1. Together with $G^{\mu\nu}_{;\mu}=0$ further geometric structures such as the Bianchi identity $DF = 0$ for the internal gauge fields constrain the equations of motion.

Underlying the more general spacetime geometry $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, it is the possibility of gauge-fermion field redescriptions such as expressed in equations 11.22 and 11.23 as considered here for the Abelian case, consistent with the selection rules of equations 11.29, which leads to the identification of the current $j^{\mu} := \overline{\psi} \gamma^{\mu} \psi$ in equation 11.26, which is identical in form to equation 11.28. In addition to the vector field transitions described above, spinor field redescriptions may also be considered with for example:

$$
\psi(x) = \int d^4y S_r(x - y)A(y)\psi(y) \tag{11.30}
$$

where $A = \gamma^{\mu} A_{\mu}$, by analogy with equation 11.22. As for the vector case in figure 11.2 this field redescription is also possible for the reverse temporal ordering. The spinor redescription function $S_r(x-y)$ is here closely related to the spinor retarded propagator which may be expressed as $S_R(x - y) = (\mathrm{i}\partial_x + m)\Delta_R(x - y)$ in terms of the scalar propagator of equation 11.27, which satisfies the relation $(i\partial_x - m)S_R(x-y) = \delta^4(x-y)$ $([70] \text{ p.63}).$

In a similar way that equation 11.22 led to equation 11.26, that is Maxwell's equation with a source term, here equation 11.30 leads to the Dirac equation, also with a source term, assuming that the properties of the spinor redescription function $S_r(x-y)$ are similar to the propagator $S_R(x-y)$. In this case the action of $(i\partial_x - m)$ on both sides of equation 11.30 results in:

$$
(i\partial \hspace{-8pt}/ - m)\psi = A\psi \tag{11.31}
$$

This is the Dirac equation that was obtained in section 3.5 via the Dirac Lagrangian in leading to equation 3.99, here with the convention for the gauge covariant derivative $D_{\mu} = \partial_{\mu} + iA_{\mu}.$

In the context of the present theory the mass m terms in these equations will also be introduced through field interactions. In the case of equation 11.21 for a massive gauge field the mass arises from the impact of the gauge symmetry upon the components external vector-Higgs $v_4 \in TM_4$, as recalled in the discussion after equation 11.21, as introduced in the terms of $D_{\mu}L(\hat{\mathbf{v}}) = 0$ of equation 11.29. Mass terms for fermions on the other hand will be incorporated through the constraint of $L(\hat{\mathbf{v}}) = 1$ itself in equation 11.29, in the form of Yukawa-like couplings between the fermion components and the same vector-Higgs, as described for equation 8.76 in subsection 8.3.3 in the case of the form $L(v_{27}) = 1$ and for equation 9.48 in section 9.2 in the case of the form $L(v_{56}) = 1$. Both for gauge bosons and fermions the interaction mass terms will correct the form of the corresponding Feynman propagators in the quantum theory. However, in focussing on the gauge-fermion interactions in the following we neglect the mass terms and hence equation 11.31 reduces to simply (within a conventional factor of i :

$$
\partial \psi = A\psi \tag{11.32}
$$

In the present theory field redescriptions occur if permitted by the constraint equations 11.29, which effectively provide interaction selection rules. For the case of an electromagnetic gauge field $A^{\mu}(x)$ associated with the internal $U(1)_Q$ symmetry generated by $\dot{S}_l^1 \in L(\text{E}_6)$, as described for example in equations 8.22–8.24 of section 8.2, interactions between the gauge and fermion fields can be identified in the expression $D_{\mu}L(v_{27}) = 0$, here taking the conserved quantity $L(v_{27}) = 1$ as the full form of temporal flow. This is analogous to the expression for $D_{\mu}L(v_{10}) = 0$ in equation 5.51, as described towards the end of section 5.4, for the $SO^+(1, 9)$ model, while here for the E_6 symmetry of the form $L(\mathbf{v}_{27}) = 1$ of equation 6.28 the expression $D_\mu L(\mathbf{v}_{27}) = 0$ includes terms of the form:

$$
D_{\mu}L(v_{27}) = \ldots + pb(\partial_{\mu}\bar{b} + \dot{s}_{f}A_{\mu}\bar{b}) + m\bar{c}(\partial_{\mu}c + \dot{s}_{f}A_{\mu}c) + \ldots = 0 \qquad (11.33)
$$

$$
= \ldots + h \theta^{1^{\dagger}} D_{\mu} \theta^{1} + \ldots = 0 \tag{11.34}
$$

In the first line \dot{s}_f carries the \dot{S}_l^1 charges of the corresponding fermion components and in the second line the values $p = m = v^0 = h$ via equation 8.72 and the spinor

 $\theta^1 = \begin{pmatrix} \frac{c}{b} \end{pmatrix}$ of equation 6.26 have been substituted in. The spinor θ^1 decomposes into the four Weyl spinors $\theta_{l,i,j,k}$ of equation 8.13 under the external $SL(2,\mathbb{C})^1$ symmetry, each of which is augmented to a Dirac spinor ψ of equation 9.43 upon extension to the E_7 symmetry of $L(v_{56}) = 1$. The Dirac spinor for the 'electron' field for example will consist of the 4-component object:

$$
\psi = \begin{pmatrix} c_1 + c_8l \\ b_1 - b_8l \\ C_1 + C_8l \\ B_1 - B_8l \end{pmatrix}
$$
\n(11.35)

in the notation of equation 9.25. Having identified $\psi(x)$ in the components of $F(h_3\mathbb{O})$ its conjugate $\overline{\psi}$ can also be constructed, and both fields expanded in terms of plane waves with complex coefficients, as was the case for the electromagnetic wave in equation 11.6. The nature of particle and antiparticle states will ultimately need to be addressed in relation to such field expansions, although here we deal directly with the fields and their mutual exchanges.

Hence in generalising from equation 11.34 for the E_7 symmetry case the expression $D_{\mu}L(v_{56}) = 0$ will contain terms incorporating factors of the form $\psi^{\dagger}D_{\mu}\psi$ involving a juxtaposition of gauge and fermion fields, with the latter identified in the components of $F(h_3\mathbb{O})$. In the present theory field exchanges in the form of equation 11.30, with the ensuing equations of motion such as equation 11.32, are required to be compatible with the constraints such as $D_{\mu}L(v_{56})=0$.

The precise means of implementing these constraints remains to be well understood, although the terms identified are analogous to the form of those found in the Standard Model Lagrangian. Further, the mutual redescriptions of the field functions are considered to be discrete, as suggested by the provisional picture of figure 11.2, which is reminiscent of the actions of the creation and annihilation operators in the expansion of quantum fields which appear through an interaction Lagrangian in expressions such as equation 10.45 in a quantum field theory.

Here equations of motion such as equation 11.32, derived from the field redescription of equation 11.30, must be filtered through the selection rules such as equation 11.34, deriving from equations 11.29, with a corresponding range of charges. This is one factor leading to differences in the likelihood of a particular process to occur. Specifically the relative factors of \dot{s}_f for different fermion components in equation 11.33 will relate to the relative number of ways in which such a process may be channelled via equation 11.30, which takes the same form for all such processes, as will be described further after figure 11.5 in the following section. Hence the factors of \dot{s}_f , obtained from the components of \dot{S}_l^1 in equations 8.22–8.24, with $|\dot{s}_f| = 1$ and $|\dot{s}_f| = \frac{1}{3}$ $\frac{1}{3}$ provisionally associated with charged leptons and d-type quarks respectively in section 8.2 provide a factor of three in the relative interaction strength between these fermion states and the electromagnetic field, that is with an apparent 'fractional charge' of $\frac{1}{3}$ for the *d*-quark relative to the unit electron charge.

For both equation 11.33 in the present theory and equation 10.23 in the model quantum field theory an interaction is mediated since changes is one field influence another field through their mutual composition in these expressions, with the constraint of $D_{\mu}L(\hat{v})=0$ in the former case and through the Euler-Lagrange equation of motion derived form the total Lagrangian in the latter case. In the present theory equations of motion with field interactions are induced through consistency with equations 11.29 rather than directly as Euler-Lagrange equations of motion from a Lagrangian with interaction terms.

Interactions between gauge and fermion fields arise for the Standard Model through the Lagrangian approach by requiring the invariance of the total Lagrangian $\mathcal L$ under local internal symmetry transformations, such as with the gauge group $U(1)_{\mathcal Q}$ in the case of electromagnetism. This implies an equivalence or indistinguishability between for example a photon and an e^+e^- pair, with $A^{\mu} \leftrightarrow \overline{\psi}\gamma^{\mu}\psi$, or between an electron and an electron-photon pair, with $\psi \leftrightarrow A\psi$; which implies the possibility of physical interactions between the fields. Similarly in the present theory it is the property of invariance of the form $L(\hat{\mathbf{v}}) = 1$ with respect to the internal symmetry, dynamically expressed over M_4 through terms such as those of equation 11.33, that allows interchanges between gauge and fermion field components corresponding to a multitude of possible solutions for the geometric form $G^{\mu\nu} = f(A, \psi)$ in 4-dimensional spacetime.

In the present theory it is the possibility of such multiple solutions with coupling between the $A^{\mu}(x)$ and $\psi(x)$ fields implied in $D_{\mu}L(\hat{v})=0$ terms that leads to the identification of the current $j^{\mu} := \overline{\psi} \gamma^{\mu} \psi$ in equation 11.26. The fields A^{μ} and $\overline{\psi} \gamma^{\mu} \psi$ mutually appear in the field redescriptions of equations 11.22 and 11.23 which are also subject to the selection rules implied in $D_{\mu}L(\hat{v}) = 0$ and incorporated into a world geometry, with the form of $G^{\mu\nu}(x)$ generalised from equation 11.1 but always with $G^{\mu\nu}{}_{;\mu}=0$ as a further constraining identity.

The further constraint $G^{\mu\nu} = f(Y)$ listed in equations 11.29, referring to the direct relation between the external and internal geometry as expressed in equation 5.20, itself will generalise to incorporate gauge-gauge field exchanges for the case of a non-Abelian internal symmetry. That is, for a gauge field $Y^{\mu}(x)$ associated with a non-Abelian internal gauge symmetry with:

$$
-\frac{1}{\kappa}G^{\mu\nu} = F^{\alpha\mu}_{\quad\rho}F_{\alpha}^{\ \rho\nu} + \frac{1}{4}g^{\mu\nu}F^{\alpha}_{\ \rho\sigma}F_{\alpha}^{\ \rho\sigma} \tag{11.36}
$$

and
$$
F^{\alpha}_{\ \mu\nu} = \partial_{\mu} Y^{\alpha}_{\ \nu} - \partial_{\nu} Y^{\alpha}_{\ \mu} + c^{\alpha}_{\ \beta\gamma} Y^{\beta}_{\ \mu} A^{\gamma}_{\ \nu}
$$
 (11.37)

with the latter from equation 3.38, there will be possible gauge field redescriptions consistent with the cubic and quartic terms of $G^{\mu\nu} = f(Y)$, namely:

$$
\partial Y \, YY \quad \text{terms} \quad \Rightarrow \quad Y \leftrightarrow YY \quad \text{exchanges} \tag{11.38}
$$
\n
$$
Y \, Y \, Y \quad \text{terms} \quad \Rightarrow \quad YY \leftrightarrow YY \quad \text{exchanges}
$$

Mutual gauge field redescriptions channelled through these constraints will augment the form of equation 11.36, similarly as for equations 11.22 and 11.23 and again under the identity $G^{\mu\nu}_{;\mu} = 0$, allowing for further possible solutions for the extended spacetime geometry.

In the Standard Model such cubic and quartic gauge field interaction terms for a non-Abelian gauge field similarly appear through terms quadratic in the curvature tensor F , in this case via a Lagrangian in the form of equation 3.94. In a quantum field theory for describing particle phenomena, such as for the Standard Model, there are certain constraints placed on the form of the Lagrangian. In general all possible terms which are allowed by gauge invariance and other symmetries of the theory should be included, but there should be no terms involving coupling constants with negative dimension D , in order to construct a renormalisable theory, as described after equation 10.86 in section 10.5. For QCD (quantum chromodynamics) in addition to equation 3.94 the Lagrangian term:

$$
\mathcal{L} = \frac{\alpha_s}{4\pi} \theta F_\alpha^{\ \mu\nu}{}^*F^\alpha_{\ \mu\nu} \qquad \text{with} \qquad {}^*F^\alpha_{\ \mu\nu} = \frac{1}{2} \varepsilon_{\mu\nu\rho\sigma} F^{\alpha\rho\sigma} \tag{11.39}
$$

is also admitted. Here $\alpha_s = \frac{g_s^2}{4\pi}$ is the strong coupling while the index α corresponds to the Lie algebra values and $*F^{\alpha}_{\mu\nu}$ is the dual field strength tensor, as originally introduced for the electromagnetic field in equation 5.24. The θ -parameter is sometimes considered as the $19th$ parameter of the Standard Model along with the 18 others (as summarised later in table 15.2 of section 15.2). However this Lagrangian term implies CP violation for strong interactions, contradicting empirical observations, unless the θ -parameter is unnaturally very small. This is the 'strong CP problem' in the Standard Model, which indicates that the Lagrangian approach may contain too many terms, leading to effects not seen in nature.

In the present theory gauge field interactions have a different origin. The expression for $G^{\mu\nu}$ in terms of the gauge field strength $F^{\alpha}_{\ \mu\nu}$ as implied in equation 11.29 in the form of 11.36 can be rewritten in a form similar to equation 5.27, with a term quadratic in the dual field strength. However, as noted after equation 5.28, there is no term of the form in equation 11.39 and hence the strong CP problem is potentially sidestepped in this Lagrangian-free theory.

Regardless of the nature of the underlying gauge or fermion field content, the object $G^{\mu\nu}(x)$, describing the spacetime geometry of M_4 , is a real-valued tensor, while the identity $G^{\mu\nu}_{;\mu} = 0$ is a real-valued vector. Similarly the constraints $L(\hat{\boldsymbol{v}}) = 1$ and $D_{\mu}L(\hat{v})=0$ are a real-valued scalar and real-valued vector respectively. The collection of these objects, as listed in equations 11.29 (with $G^{\mu\nu} = f(Y)$ interpreted as a local constraint), is analogous to the collection of terms in a single real-valued scalar Lagrangian, and in the present theory they will also be interrelated through the full dynamics. However, as for a real-valued Lagrangian, the components of fields underlying these objects may be mathematically analysed into complex-valued parts, such as the Fourier modes for the electromagnetic field in equations 11.5 and 11.6.

More generally, as described in subsection 2.2.3 and equation 2.30, a real-valued gauge field $Y(x)$ on M_4 was originally obtained as the pull-back of the Maurer-Cartan 1-form defined on the manifold of an unbroken symmetry group \ddot{G} . Subsequently internal gauge fields deriving from the symmetry breaking over the base manifold were considered, as appearing in the final term of equation 2.47 for example. Although only the complete real field $Y(x)$ represents a *macroscopic* gauge field (as discussed after equation 11.18), the functional form of the gauge field $Y^{\alpha}_{\mu}(x)$ may be analysed into complex Fourier components. Similarly, the 56 real components of a vector $v_{56} \in F(h_3\mathbb{O})$ under $L(\mathbf{v}_{56}) = 1$, including the various fermion subcomponents $\psi(x) \subset \mathbf{v}_{56}(x)$, when expressed as functions over M_4 may be analysed into complex Fourier modes. Further,

in principle such *complex* $e^{\pm ik\cdot x}$ Fourier mode components of the fields $Y(x)$ and $\psi(x)$, or a hybrid combination, might be composed at the microscopic level to form real expressions for objects such as $G^{\mu\nu}(x)$ and $D_{\mu}L(\mathbf{v}_{56}(x)) = 0$ over M_4 .

While a crucial observation for the present theory is that the spacetime associated with any field propagation is *not* flat, as pictured in figure 11.1 with $G^{\mu\nu} = -\kappa T^{\mu\nu}$ for example, here the geometry is assumed to be sufficiently close to flat in order to employ such a plane wave expansion in essentially Cartesian coordinates, as described before equation 11.14. The field redescription functions, featuring in equations 11.22, 11.23 and 11.30 for example, are closely related to the retarded propagator $\Delta_R(x-y)$ of equation 11.27. This latter function itself is expressed as an integral over $e^{-ik \cdot (x-y)}$ Fourier modes, suggesting that in turn the exchanges and interactions between the components of fields such as $Y(x)$ and $\psi(x)$ might also be most conveniently analysed in terms of $e^{\pm i k \cdot x}$ Fourier modes, as is the case for the field expansions in quantum field theory. That is the association of the functions $D_{r,a}^{\mu\nu}(x-y)$ with the propagators $D_{R,A}^{\mu\nu}(x-y)$ as provisionally suggested after equations 11.22 and 11.23, as for the association of the function $S_r(x-y)$ in equation 11.30 with the propagator $S_R(x-y)$, may involve analysis of the corresponding field structures in terms of complex-valued components. These structures in the present theory will be linked with the cross-section calculations of QFT in the following section.

In all cases the mutual field exchanges are required to be consistent with the full set of constraints of equations 11.29, with the geometric condition $G^{\mu\nu}_{;\mu} = 0$ in 4-dimensional spacetime implying 4-momentum conservation through the definition of energy-momentum as $T^{\mu\nu} := G^{\mu\nu}$. The underlying one-dimensional form of temporal progression is reflected in the structure of a causal sequence of field redescriptions, as expressed by the θ -function in $\Delta_R(x-y)$ of equation 11.27, while the δ -function in that expression relates to the appropriate matching of Fourier modes for the general case, for which a finite mass m may result from further field interactions. Each possible field redescription itself, for individual Fourier modes such as $e^{-ik \cdot x}$, may provisionally be associated by analogy with QFT with an element of a Feynman diagram, namely a vertex diagram of the kind listed in 'rule 2' of table 10.1, as depicted in the examples of figure 11.3.

Figure 11.3: Three Feynman vertex diagrams correlated with the possible field exchanges (a) $A \leftrightarrow \overline{\psi}\psi$, (b) $\psi \leftrightarrow A\psi$ and (c) $Y \leftrightarrow YY$, as associated with equations 11.23, 11.30 (strictly with S_r replaced by S_a here, since the implied time ordering is from left to right in these diagrams) and the cubic terms of equation 11.38 respectively; with a 4-way gauge field vertex also possible for the quartic terms of the latter equation.

The field redescription of equation 11.30, associated with figure 11.3(b), is

directly suggested by the form of the terms of $D_{\mu}L(v_{27}) = 0$ in equations 11.33 and 11.34 via equation 11.32, although a higher-dimensional full form such as $L(v_{56}) = 1$ will be needed for more explicit details. More generally the juxtaposition of a gauge field and quadratic fermion field factor in the terms of $D_{\mu}L(v_{56}) = 0$ may lead to interactions between this combination of fields with various spacetime orientations, while sharing the same vertex topology, resulting in the exchange of figure $11.3(a)$ for example.

Similarly, as well as identifying particular particle states in the components of $F(h_3\mathbb{O})$ the distinction between particles and antiparticles, together with their different dynamic behaviour, will require a full consideration of the fields under the symmetries of extended 4-dimensional spacetime. The provisional correlation of the combination of the fermion field ψ and its conjugate $\overline{\psi}$ with the combination of a fermion and antifermion pair, as discussed after equation 11.35, will be dependent upon the temporal orientation of the field components on the extended manifold M_4 .

The association between terms of the constraints in equation 11.29 and the form on an interaction Lagrangian, as emphasised by the Feynman vertices of figure 11.3, raises the question of how calculations for quantities such as cross-sections, as measured in HEP experiments, might be determined in the present theory and how such calculations might be related to the Feynman rules more generally. This will form the topic of the following section.

11.2 Determination of Process Probability

Here we make a provisional connection between the calculation of physical quantities such as cross-sections, as described in the previous chapter, and the notion of a degeneracy of field redescriptions underlying the corresponding processes, as introduced in the previous section. Since such calculations in quantum field theory have achieved great success in comparison with empirical HEP observations a relation between the present theory and the mathematical structures and tools of QFT will be desirable.

First we consider as an example a field sequence $\overline{\psi}\gamma^{\mu}\psi \to \overline{\varphi}\gamma^{\mu}\varphi$, where $\psi, \varphi \subset \overline{\varphi}$ $v_{56} \in F(h_3\mathbb{O})$ denote fermion components, with the interaction taking place in the spatial volume V over a time period T via an intermediate $A^{\mu}(x)$ field state. This situation is depicted in figure 11.4 which essentially consists of a juxtaposition of figures 11.2(a) and (b) where the initial and final fermion types may differ in general.

In this section we consider interactions at the level of such field exchanges. As alluded to at the end of the previous section the structure of physical particle states in spacetime, including both particle and antiparticle states, is yet to be identified in this theory. Further, the inclusion of the second and third generation fermions may require a further extension of the full form $L(\hat{v}) = 1$, as suggested for example in section 9.3. However a field state such as $\overline{\psi}\gamma^{\mu}\psi$ is provisionally considered to represent fermion pairs such as e^+e^- or $\mu^+\mu^-$ leptons or $d\bar{d}$ or $t\bar{t}$ quarks for example. Hence the field sequence in figure 11.4 mimics a HEP collision process such as $e^+e^- \rightarrow \mu^+\mu^-$. In the following section the physical nature of the actual incoming and outgoing particle states observed in HEP phenomena will be considered.

In the analogous QFT calculation the initial and final 'particle' states are rep-

Figure 11.4: The transition from an initial $\overline{\psi}\gamma^{\mu}\psi$ field state to a final $\overline{\varphi}\gamma^{\mu}\varphi$ state via an intermediate description of the field function in terms of a mathematically equivalent $A^{\mu}(x)$ field state.

resented by complex plane waves, that is Fourier modes of the form $e^{\pm ik\cdot x}$, as discussed for equations 10.43 and 10.44 for example. This is similar to the picture initially considered here in figure 11.4 with the field functions in spacetime expanded in terms of Fourier modes such as those of equation 11.6. However, in the present theory the incoming, interacting and outgoing field states conform everywhere to an expression of the spacetime geometry described by the real tensor $G^{\mu\nu}(x) = f(A, \psi, \varphi)$.

For the case of d discrete intervals of time Δt_i during which the field exchanges between $t = 0$ and $t = T$ in figure 11.4 may occur the total number of ways N in which the overall transition may proceed is simply:

$$
N = \sum_{i=1}^{d} \left(R(t_1 \in \Delta t_i : A \to \overline{\varphi} \varphi) \sum_{j=1}^{i-1} R(t_2 \in \Delta t_j : \overline{\psi} \psi \to A) \right)
$$
(11.40)

with R denoting 'redescription' such that $R(\Delta t_i : A \to \overline{\varphi}\varphi) \equiv 1$ simply expresses the fact that the corresponding field exchange $A^{\mu} \leftrightarrow \overline{\varphi} \gamma^{\mu} \varphi$ is allowed during the time interval Δt_i . More generally $R(t)$ will take the value 1 if the corresponding field exchange is allowed, according to the constraint equations 11.29 as described in the previous section, and 0 if it is not.

For the process with incoming field state $\overline{\psi}\gamma^{\mu}\psi$ the total field function is already distributed everywhere in V from time $t = 0$ in figure 11.4, and as a function in spacetime it is indistinguishable from that of the outgoing $\overline{\varphi}\gamma^{\mu}\varphi$ field state at $t=T$. The field redescription applies *everywhere* in V simultaneously at any time such as t_2 or t_1 since this simply involves a reinterpretation of the *same* field function, with nothing physically changing in VT . Hence from the point of view of the spacetime geometry and $G^{\mu\nu}(x)$ alone it would be possible to link the states $\overline{\psi}\gamma^{\mu}\psi$ and $\overline{\varphi}\gamma^{\mu}\varphi$ directly, without an intermediate $A^{\mu}(x)$ field description. This is prevented in the present theory by the absence of selection rule being provided by constraints such as $D_{\mu}L(\hat{v}) = 0$ which determine whether $R(t) = 1$ or $R(t) = 0$ for a particular field redescription.

This is closely analogous to the Lagrangian approach in QFT as described for example for the scalar model where the absence of a coupling term of the form $\hat{\mathcal{X}}^{\dagger}\hat{\mathcal{X}}\hat{\mathcal{Y}}^{\dagger}\hat{\mathcal{Y}}$ in the interaction Lagrangian of equation 10.23, and hence via equation 10.27 in $H_{\text{int}}(t)$, implies that the collision process $\mathcal{X}^+ \mathcal{X}^- \to \mathcal{Y}^+ \mathcal{Y}^-$ requires an intermediate ϕ state as depicted in the Feynman diagram of figure 10.4. Similarly the lack of a direct electronmuon coupling in the Standard Model Lagrangian leads to consideration of scattering processes via an intermediate photon, such as depicted in figure 10.3, which will be seen to be analogous to figure 11.4 for the present theory.

Taking equation 11.40 to the continuum limit, as implied in figure 11.4, a measure of the total degeneracy D can then be expressed as:

$$
D(T,0) = \int_0^T dt_1 \int_0^{t_1} dt_2 R(t_1)R(t_2)
$$
 (11.41)

The structure of this equation has some similarity to the second-order term in the expansion of the time evolution operator $U(t,t_0)$ in quantum field theory. In the interaction picture, with interaction Hamiltonian H_{int} , the operator U satisfies the differential equation 10.30, as described in section 10.3, with the iterative solution for $U(t, t_0)$ displayed in equation 10.31.

In equation 10.31 the factors of the Hamiltonian operator H_{int} in each term naturally stand in time order, with the earliest to the right and latest to the left, in virtue of the time integration limits. As explained in section 10.3 this expansion of the time evolution operator $U(t, t_0)$ can be written in the familiar more compact form of equations 10.34, via equations 10.32 and 10.33, using the T-product of operators which *imposes* time ordering over a broadened, and more symmetric, range of time integrals. In particular the second-order term in equation 10.31 can be replaced by that in equation 10.32 since:

$$
\int_{t_0}^t dt_1 \int_{t_0}^{t_1} dt_2 H_{\rm int}(t_1) H_{\rm int}(t_2) \equiv \frac{1}{2} \int_{t_0}^t dt_1 \int_{t_0}^t dt_2 T\big(H_{\rm int}(t_1) H_{\rm int}(t_2)\big) \qquad (11.42)
$$

It is the similarity between equation 11.41, as a measure of the degeneracy or number of ways in which to describe the field transition sequence $\overline{\psi}\gamma^{\mu}\psi \to A^{\mu} \to \overline{\varphi}\gamma^{\mu}\varphi$, and the left-hand side of equation 11.42 that provides a further preliminary entry point for the present theory into the workings of QFT. In a similar way that the field exchange of figure 11.2(b) has been provisionally associated with the Feynman vertex diagram of figure 11.3(a), the $A^{\mu}(x)$ internal field stage of figure 11.4 might be associated with the Feynman propagator corresponding to the internal line of the Feynman diagram in figure 10.3 for example, via the relation between equations 11.41 and 11.42 described above.

Equations 10.31 and 10.32 are matched on a term by term basis and hence the terms of the perturbative expansion of equation 10.34 match those of equation 10.31. In turn the higher-order terms of equation 10.31 can be associated with higher-order sequences of field redescriptions, such as depicted in figure 11.6 below. In the Feynman rules for the mathematical elements associated with a Feynman diagram at order n in perturbation theory the factor of $1/n!$ in equation 10.33 cancels against a factor of n! from the possible vertex permutations, as described shortly after equation 10.46 and summarised for 'rule 6' in the opening of section 10.5. Hence in the Feynman rules for the second-order term correlated with the right-hand side of equation 11.42 the factor of $\frac{1}{2}$ does not appear.

Via the above associations the field exchange sequence described in figure 11.4 is analogous to the Feynman diagram in figure 10.4 for the corresponding scalar model QFT calculation. While a possible physical interpretation of the Feynman propagator $\Delta_F(x-y)$ in terms of 'virtual particles' is conceptually dubious, as discussed in section 10.4 (for example after equation 10.72), this object is a key part of calculations in QFT and we return to this propagator – which in the scalar field case may be expressed for the internal field operator $\phi(x)$ in canonical QFT by the equation:

$$
i\Delta_F(x-y) = \langle 0|T(\hat{\phi}(x)\hat{\phi}(y))|0\rangle \tag{11.43}
$$

as we began with equation 10.49 in section 10.4. This object arose when the transition amplitude calculation was restructured with the time evolution operator $U(\infty, -\infty)$ in the form of equation 10.32 placed between vacuum states, in particular for the secondorder term. This object hence consists of terms implicitly containing time-ordered field products, such as $T(\phi(x)\phi(y))$ in the right-hand side of equation 11.42.

The time ordering implies that $\Delta_F(x-y)$ consists of two parts, associated with $\theta(x^0 - y^0)$ and $\theta(y^0 - x^0)$, as described in equations 10.50–10.55 and as represented by the two diagrams in figure 10.6. From the point of view of the concept of field redescriptions in the present theory the first diagram, figure 10.6(a), can be physically motivated as representing the field redescription causal sequence such as $\psi \gamma^\mu \psi \to A^\mu \to \overline{\varphi} \gamma^\mu \varphi$ as depicted in figure 11.4 while the second diagram, figure $10.6(b)$, represents a figment of the mathematical restructuring of the calculation, leading in turn to the notion of intermediate 'virtual particle' states.

Nevertheless, via the above chain of argument each case of an intermediate $A^{\mu}(x)$ field state, as depicted in figure 11.4, may be *provisionally* associated with a corresponding Feynman propagator $D_F^{\mu\nu}$ $_{F}^{\mu\nu}(x-y)$. That is, intermediate field redescriptions such as that in figure 11.4 may be associated with the 'virtual particle' states as represented by the internal line in figure 10.4, and in Feynman diagrams in general, considered as a restructuring of a calculation which is here fundamentally based on an underlying conceptual notion of a degeneracy of field descriptions.

Associating a Feynman propagator with each intermediate causal redescription, such as that with the field $A^{\mu}(x)$ in figure 11.4 as described above, supplements the set of interaction vertices associated with the constraints of equations 11.29, as exemplified in figure 11.3. Hence with propagators identified in addition to the vertices these objects may be combined to form Feynman diagrams more generally. Beginning from the idea that the probability of an observable process is a measure of the number of ways in which it can occur, summing over all possible intermediate field redescriptions as for example in equation 11.41, the aim is to effectively reproduce a full set of Feynman rules, for comparison with the Standard Model version of table 10.1, and further to use this relation in order to make calculations of empirical quantities such as cross-sections.

Regarding the Feynman vertices the key to understanding how $D_{\mu}L(v_{27})=0$ terms, for example, are to be used in place of a Lagrangian here may be found in the coupling strength, which is put in by hand in the Lagrangian case. In equation 11.33 the value of \dot{s}_f for the leptonic states is 3 times larger than for the quark states, as determined in section 8.2 and noted in the previous section. The question then is how this mathematical factor of 3 corresponds to an empirical factor of 3 in 'electric charge' with an underlying explanation in terms of the degeneracy for the number of ways a process can occur. Consider the processes described by the Feynman diagrams in figures $11.5(a)$ and (b), either of which may be correlated with, while not literally representing, the field sequence depicted in figure 11.4 as described above.

Figure 11.5: Feynman diagrams for the electromagnetic processes (a) $e^+e^- \rightarrow \mu^+\mu^$ and (b) $e^+e^- \rightarrow d\bar{d}$, together with a 'higher-order correction' via (c) a radiated photon and (d) a gluon exchange between the final state quarks respectively.

In the calculation of the degeneracy for a process, as initially described for figure 11.4, the number of possibilities depends upon the total time T available for the process, as can be seen in equation 11.41. For a quantum field theory, the spacetime volume factor VT for an interaction cancels in cross-section and decay rate calculations, as described in section 10.2 following equation 10.3, essentially since the effective values of V and T in external spacetime are the same for all possible processes. A similar cancellation might be expected for calculations based on field degeneracies in the present theory. On the other hand, unlike the case for the common external dimensions of the interaction, here for the present theory, the effective 'charge volume' C in the internal space dimension varies from process to process, as indicated by the differing values of \dot{s}_f in equation 11.33, and does not cancel in such calculations. In QFT these three spaces are closely related, as seen for example in the CPT theorem, while in the present theory they are mutually related through the structure and symmetries of the underlying temporal flow in the form $L(\hat{\mathbf{v}}) = 1$. A more precise expression for the way in which the relative charges channel the relative likelihood for different field exchanges, and indeed a fuller understanding of the relation of the present theory to the Lagrangian approach in general, requires further study, as was also discussed after equation 11.35.

Given the 'virtual photon' mediating both processes in figures 11.5(a) and (b) further internal degeneracy, as for example in figure 11.6 below, will be essentially the same for both cases and not effect the relative rates. That is the branching fractions or relative cross-sections for competing processes will depend on the differences in the number of ways, and this may be dominated by the factors of $|\dot{s}_f| = 1$ or $|\dot{s}_f| = \frac{1}{3}$ 3 associated with the final state vertex in figures $11.5(a)$ and (b) respectively. Differences may also arise due to the mass of the final state particles (upon which the final state phase space depends), relating to further possible field interactions with the components of the vector-Higgs $h_2 \equiv v_4 \in TM_4$, and more generally due to higher-order field exchange possibilities, such as those represented in figures $11.5(c)$ and (d) ; as will be further discussed in the following section.

In the full theory the possible Feynman diagrams will generalise corresponding to the range of gauge fields and further interactions identified for a full set of internal symmetries as studied in chapters 8 and 9, and which show a significant resemblance to the structures of the Standard Model of particle physics. For example, in figure 11.5(d) an $SU(3)_c$ gauge field exchange is included. There are eight internal $SU(3)_c$ generators, as described in section 8.2 and listed down the left-hand side of table 8.7. Unlike the $U(1)_Q$ action in equation 11.33 these mix the components of $\theta^1 = \begin{pmatrix} c \\ b \end{pmatrix} \in \mathbb{O}^2$ between different Weyl spinors hence introducing interactions between the corresponding quark states. The identification of an $SU(2)_L \subset E_7$ (or within a larger symmetry of time such as E₈), also mediating between the external $SL(2,\mathbb{C})^1$ Weyl spinors in $F(h_3\mathbb{O})$ (or within a higher-dimensional form of time such as $L(v_{248}) = 1$) will provide a further internal gauge symmetry action central to an understanding of electroweak theory within the present theory.

The measure of degeneracy in equation 11.41 can be generalised to higherorder sequences of A^{μ}, ψ, φ field exchanges which mirror the general expansion to higher-order perturbations for QFT in equation 10.31; with the Hamiltonian operator $H_{\text{int}}(t)$ in the latter case replaced by the 'redescription parameter' $R(t)$, as determined by the constraints of equations 11.29, in the former case. The temporal sequence of figure 11.6 provides an example of the ways in which the causal sequence of figure 11.4 may be generalised for nested sequences of field indistinguishability to arbitrary high order.

$$
\left[\left|\nabla\gamma^{\mu}\psi\right|\right] = A^{\mu} \left[\left|\nabla\gamma^{\mu}\psi\right|\right] = A^{\mu} \left[\left|\nabla\gamma^{\mu}\psi\right|\right] = 0
$$

Figure 11.6: The transition from an initial $\overline{\psi}\gamma^{\mu}\psi$ state to a final $\overline{\varphi}\gamma^{\mu}\varphi$ state via an intermediate description of the field function in terms of a sequence mathematically equivalent $A^{\mu}(x) \rightarrow \overline{\psi}(x) \gamma^{\mu} \psi(x) \rightarrow A^{\mu}(x)$ field states.

The corresponding degeneracy for the chain of field interpretations in figure 11.6, as an augmentation of equation 11.41, is expressed as :

$$
D(T,0) = \int_0^T dt_1 \int_0^{t_1} dt_2 \int_0^{t_2} dt_3 \int_0^{t_3} dt_4 R(t_1: A \to \overline{\varphi}\varphi) R(t_2: \overline{\psi}\psi \to A)
$$

$$
R(t_3: A \to \overline{\psi}\psi) R(t_4: \overline{\psi}\psi \to A)
$$
(11.44)

While the sequence of field descriptions pictured in figure 11.4 can be correlated with the Feynman diagram of figure 10.4, via equations 11.41 and 11.42, the higher-order process of figure 11.6 is similarly analogous to the form of figure 10.9, representing the T-ordered expression for this fourth-order term for the scalar QFT model. A similar correspondence may be identified between field sequences for the present theory and Feynman diagrams in QED, as depicted in figure 11.11 in the following section for example. These figures represent steps in the direction of connecting the structures of the present theory with Feynman diagrams and rules more generally.

Intuitively the extra sums over the two additional intermediate times, labelled t_3 and t_2 in figure 11.6 and equation 11.44, will lead to a relative 'infinity' of new ways in which the overall event may proceed from the initial to the final state. However the degeneracy measure D for both equations 11.41 and 11.44 is actually finite. On the other hand the intermediate state composed of ψ and $\overline{\psi}$ between t_3 and t_2 involves two field contributions simultaneously, each of which may be expanded into Fourier modes independently with a combined product of the form $\sim e^{-i(p_1+p_2)\cdot x}$ which, although the total $p_1 + p_2$ is constrained, leads to a further infinity in the degeneracy of the internal 4-momentum. In this case the integral sum over p_1 is unlimited, unlike the situation for the time integrals, and is expected to be reflected in the divergent momentum loop integrals, as for example in equation 10.86 for the scalar model, in the correspondence with QFT calculations. For the present theory, as for QFT, such divergences might be expected to cancel when observable quantities such as branching ratios are appropriately normalised, as will be described in the following section, with such observables ultimately dominated by the charges involved in the final interaction of the sequence as discussed above.

In the present theory the world geometry is necessarily described by the real tensor $G^{\mu\nu}(x)$ which itself in principle may be composed out of the real or complex components of fields, such as $A^{\mu}(x)$ and $\psi(x)$. Regarding the degeneracy count itself it is an open question concerning whether there is a unique or optimal way in which possible field redescriptions should be counted, consistent with the constraint equations 11.29. This question concerns both the domain of the field functions, as a patchwork of regions in spacetime or in momentum space for example, and also the form of the field functions. Here we are analysing the degeneracy count in terms of complex Fourier modes on the base manifold M_4 . In this sense each $e^{\pm i k \cdot x}$ component is not considered as an independent physical field, rather this decomposition provides a mathematical means of identifying a set of mutually independent field solutions which may be summed over.

In describing the transitions between fields such as $\psi(x)$, $A^{\mu}(x)$ and $\varphi(x)$ it is possible that linear combinations of real sine and cosine expansion terms, rather than *complex* $e^{\pm i k \cdot x}$ parts, might be employed to preserve the identity of *real*, and hence physical, fields under the spacetime geometry $G^{\mu\nu}(x)$ subject to the constraint equations 11.29 everywhere. For example considering the real Fourier components $A_c^{\mu}(\mathbf{k})$ and $A_s^{\mu}(\mathbf{k})$ of the field in equation 11.3 to be exchanged independently maintains a real condition for the field $A^{\mu}(x)$ which at every stage may compose an intermediate, but physical, gauge field coupled to the fermion fields consistent with $D_{\mu}L(\hat{\mathbf{v}})=0$.

However here we have described field interactions such as $A^{\mu} \leftrightarrow \overline{\psi} \gamma^{\mu} \psi$ in terms of the indistinguishability of complex Fourier modes of the fields, as expanded for example in equations 11.5 and 11.6 for the electromagnetic field, in part since this provides a closer link with the framework of QFT. Indeed, as alluded to towards the end of the previous section, many of the tools involved in QFT, such as the various propagators and the δ -function of equation 10.62 and the θ -function of equation 10.64, are conveniently expressed in terms of complex Fourier modes. Further, complex components of gauge fields have already been considered with regard to the charged gauge bosons $\tilde{W}^{(2)\pm}_{\mu}(x)$ of equation 8.67 in section 8.3, by analogy with the standard electroweak gauge fields $W^{\pm}_{\mu}(x)$ of equation 7.57 in section 7.2, which relate to the corresponding physical interactions with Lorentz spinors. Hence here the field redescriptions will be analysed in terms complex Fourier modes in the determination of a real measure or count of the degeneracy of field solutions.

As described in the previous section both parts of equation 11.6 are required to identify a field state carrying real energy-momentum, which in the present theory is determined by the form of the field under $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$. Hence transitions between fields must necessarily link *both* of the $e^{\pm ik\cdot x}$ parts with the *external* 4-momentum k, as identified through equation 11.16, matched under an everywhere real $T^{\mu\nu}$:= $-\frac{1}{\kappa}G^{\mu\nu}$ energy-momentum tensor, subject to the identity $G^{\mu\nu}_{\;\;;\mu} = 0$, and also with the internal representations of the field components matching under the constraints of equation 11.29 in general, with the form $L(\hat{v}) = 1$ broken over the base manifold.

With $A^{\mu}(\mathbf{k}) = \frac{1}{2}(A_c^{\mu}(\mathbf{k}) + iA_s^{\mu}(\mathbf{k})) \in \mathbb{C}$ a general complex number in equation 11.5 transitions in the field $A^{\mu}(x)$ can be considered to take place treating $A^{\mu}(k)$ and $A^{\mu*}(k)$ as independent degrees of freedom in terms of possible exchanges with complex Fourier modes of the fermion fields. This implies the possibility of intermediate *complex* fields such as $A^{\mu}(x)$ and $\psi(x)$ while hybrid combinations of these gauge and fermion fields mutually form under *real* objects such as $G^{\mu\nu}(x)$ and $D_{\mu}L(\hat{v}) = 0$.

Hence the temporal sequence of redescriptions should be considered independently for the complex $e^{-ik\cdot x}$ and $e^{+ik\cdot x}$ parts such that, for example, the processes represented in figures 11.4 and 11.6 may be generalised for this independence, as depicted for example in figure 11.7.

Figure 11.7: The transition from an initial $\overline{\psi}\gamma^{\mu}\psi$ state to a final $\overline{\varphi}\gamma^{\mu}\varphi$ state generalised for an intermediate description of the field function in terms of the complex Fourier modes $e^{-ik \cdot x}$ and $e^{+ik \cdot x}$ independently in time.

With the need to account for both sets of possible sequences as exemplified in figure 11.7 the probability P for the overall process $\overline{\psi}\psi \to \overline{\varphi}\varphi$ is proportional to $D_+ \times D_-,$ where D_+ represents the degeneracy of ways via $e^{+ik\cdot x}$ exchanges and D_-

that for the $e^{-ik\cdot x}$ mode exchanges, each of which has a structure similar to that in equation 11.41 or 11.44. Alternatively the process probability could be expressed in terms of the degeneracies D_c and D_s representing the number of field exchanges relating to the cosine and sine Fourier modes as alluded to above, with for example $A_c^{\mu}(\mathbf{k})$ and $A_s^{\mu}(\mathbf{k})$ of equation 11.3 independent, and with $P \propto D_+D_- \equiv D_cD_s$. In this case all fields are real-valued and hence can be interpreted as physical entities at all times, however here we pursue the equivalent calculation based on the complex decomposition.

Earlier in this section we have described a correlation between the form of a degeneracy count $D(T, 0)$ and the anatomy of a Feynman diagram, with for example figure 11.4 compared with figure 11.5(a) or (b), via the structure of the expansion of the QFT operator $U(t, t_0)$ of equations 10.31–10.34. Here the underlying physical basis of probability calculations is found in the field degeneracies, with the use of Tordered products in QFT, via the θ -functions, simply implementing a restructuring of the calculations. Hence in turn the representation of the Feynman propagator in figure 10.6 should not be interpreted as two possible physical processes. On the other hand the fact that the underlying field redescriptions are free to take place independently for *both* the $e^{-ik \cdot x}$ and $e^{+ik \cdot x}$ field components, as depicted for example for the process $\overline{\psi}\gamma^{\mu}\psi \to \overline{\varphi}\gamma^{\mu}\varphi$ in figure 11.7, does extend the range of possible field redescriptions and hence will have physically observable consequences. With both sets of field redescriptions for the $e^{\pm ik\cdot x}$ Fourier modes required to link the initial and final states the process probability takes the form $P \propto D_+ D_-$, and we hence now wish to determine a correlate for this product in QFT.

In figure 11.7 the field states at $t = 0$ and $t = T$ (and hence also for $t \to \pm \infty$) represent real external particle states, that is on-mass-shell particles. This suggests that the diagram in figure 11.7 can be 'unfolded' to represent an extension of a linear degeneracy count, having the same basic structure as figure 11.4 or 11.6, but with a 'fold line' denoting on-shell states. The corresponding unfolded diagram is depicted in figure 11.8(a). This field sequence correlates with the structure of the Feynman diagram of figure 11.8(b), with the fold line mapped to the cut line – for which the propagators are simultaneously placed on-mass-shell, as originally described for figure 10.10.

According to the 'cutting rules', as also described in section 10.5, the imaginary part of the transition amplitude associated with a Feynman diagram is obtained by summing over the cutting possibilities. These involve adapting the Feynman rules for each possibility by placing the 'cut' virtual states on-mass-shell – and hence open to interpretation as external particle states – via equation 10.98, which essentially replaces each corresponding Feynman propagator Δ_F by one of the Δ^{\pm} function components described in equations 10.55–10.60. From the unfolding of figure 11.7 the initial and final field states in figure 11.8(a) are equivalent, and hence the cutting rules applied to the corresponding figure 11.8(b) yields the imaginary part of the forward scattering amplitude for the e^+e^- initial state $|i\rangle$, namely in fact $2\text{Im}(\mathcal{M}_{ii})$, as contributed by placing the cut line on the intermediate $\mu^+\mu^-$ state for this Feynman diagram.

The important observation of the present theory is that $\text{Im}(\mathcal{M}_{ii})$ is a real number, and hence might be directly compared with event probabilities with contributions of the form $P \propto D_+ D_-$ based on a count of the 'number of ways' in which an

Figure 11.8: (a) The unfolding of figure 11.7, with a corresponding 'fold line' and reparametrised time intervals. (b) A correlated Feynman diagram for the forward scattering process $e^+e^- \rightarrow e^+e^-$, with a 'cut line' drawn through the intermediate loop propagators of the muon field.

observed process might arise. Adding all possible contributions for all possible final states, exemplified by the process in figure 11.7, then correlates, via the generalisation of figure 11.8, with the imaginary part of the forward scattering amplitude for all possible Feynman diagrams for the full perturbative expansion. The resulting real number Im(\mathcal{M}_{ii}) is in turn directly related to the total cross-section σ , equation 10.96, via the optical theorem as described in section 10.5. Hence we arrive at a provisional relationship between a degeneracy count and a physical observable.

As described towards the end of section 10.5 the optical theorem can be proven to all orders of perturbation through the analysis of Feynman diagrams. The cut pictured in figure 11.8(b) represents one contribution to $\text{Im}(\mathcal{M}_{ii})$ for this diagram, with a second contribution provided by placing the cut instead through the $d\bar{d}$ fermion loop. Hence by the above discussion the determination of $\text{Im}(\mathcal{M}_{ii})$ via the cutting rules for this diagram correlates with a sum of a $D_{+}D_{-}$ field sequence for both a $\mu^{+}\mu^{-}$ final state and a $d\bar{d}$ final state. Similarly for equation 10.101 the imaginary part of the Feynman diagram of figure 10.10 was determined corresponding to opening up a $\mathcal{Y}^+ \mathcal{Y}^-$ final state, with a further contribution to $\text{Im}(\mathcal{M}_{ii})$ at this order of perturbation obtained by replacing the loop in figure 10.10 with a $\mathcal{X}^+ \mathcal{X}^-$ state, as described after equation 10.101.

The Feynman diagram with the cut of figure 11.8(b), in placing the $\mu^+\mu^-$ pair on-mass-shell and via the optical theorem, contributes to the cross-section $\sigma(e^+e^-\to e^-e^-)$ $\mu^+\mu^-$). However the structure of Im(\mathcal{M}_{ii}), in summing over the cuts, generally incorporates a collection of final states from which individual cross-sections for particular processes need to be untangled, as they are for the sum on the left-hand side of equation 10.94 for example. Also, as alluded to in the caption comments, the fold-line in figure 11.8(a) should in principle be constrained to the 'half-time' point to accurately represent the degeneracy count of figure 11.7. Further, we have considered the degeneracy count, based on particular sequences of fields leading to a particular final state, to represent a measure of the probability $P \propto D_+D_-$ for a particular process. However, in order to actually represent a probability this count needs to be determined relative to the total degeneracy for all possible final states, which will provide the overall normalisation and which so far we have not taken into account. In looking to address these points we recap how a particular final state is extracted and an event probability determined in the context of all possible outcomes in the framework of a QFT, with the aim of establishing a more precise link with similar calculations for the present theory.

As described in section 10.3 in QFT the initial state $|i\rangle$ evolves through a period of field interactions into the state $|\Psi(\infty)\rangle = S|i\rangle$ according to the S-matrix of equation 10.35. This evolution is governed at each moment by the equation of motion expressed in equation 10.28 in which the interaction Hamiltonian $H_{\text{int}}(t)$ contains all possible field interactions. Hence $|\Psi(\infty)\rangle$ in turn contains all possible final states. Since $H_{\text{int}}(t)$ is Hermitian the evolution of the state in equation 10.28 is a unitary transformation, and hence if the initial state normalisation is chosen with $\langle i|i\rangle = 1$ this is preserved such that $\langle \Psi(t)|\Psi(t)\rangle = 1$ at any time t. On inserting a sum over a complete orthonormal set of similarly normalised final states $|f\rangle$ we have $\sum_{f} \langle \Psi(t) | f \rangle \langle f | \Psi(t) \rangle = 1$, and in particular in the aftermath of the interaction, we have:

$$
\sum_{f} |\langle f | \Psi(\infty) \rangle|^2 = 1 \tag{11.45}
$$

as a mathematical identity. Hence the objects $|\langle f|\Psi(\infty)\rangle|^2$, in the sense of consisting of a set of positive real numbers that sum to unity, do have the property of representing probabilities, and in a structure which implicitly contains information about all possible final states.

A relationship between the degeneracy $D(T, 0)$ of equation 11.41 and the second order term of $U(t, t_0)$ of equation 11.42 was described for a particular field sequence leading to a particular final state, as pictured in figure 11.4. However, in general a degeneracy count associated with all terms of the entire S-matrix is desired in order to express everything that can happen, according to the field redescriptions permitted by the constraints of equations 11.29 in place of an interaction Hamiltonian, and hence incorporate all possible outcomes. This suggests a 'complexification' of the probability calculation based on the degeneracy count such that the unitarity constraint, that is $SS^{\dagger} = 1$ in QFT, might effectively be employed to normalise the total probability for any outcome to unity.

The subcomponent degeneracy counts D_+ and D_- , originally considered to provide a measure of the probability $P \propto D_+ D_-,$ are each real numbers. The probability for any process is a positive real number $P \in \mathbb{R}$ from 0 to 1, as for any probability, and as for the square root of this quantity $p = \sqrt{P}$. However in principle it may be possible to consider a complexification of the underlying calculation, represented by $p \to \tilde{p} \in \mathbb{C}$, such that $P = \tilde{p}^*\tilde{p}$. This is considered to be essentially the case in quantum theory where unitary symmetry is used to model the properties of probabilities, and in the case of QFT the role of the above complex quantity \tilde{p} is played by the transition amplitude \mathcal{M}_{fi} .

Specifically, the likelihood of an event in QFT is proportional to the squared

modulus of the transition amplitude, as extracted from the terms of equation 11.45 via equation 10.6, and as introduced in equation 10.3. With the cross-section for a HEP process, for example, linked to the imaginary part of the forward scattering amplitude via the optical theorem expression of equation 10.96 and this latter object, as the real number Im(\mathcal{M}_{ii}), correlated with a degeneracy count $D_+D_-\,$, as described for figure 11.8, we have the following chain of associations:

$$
P \propto D_+ D_- \sim \text{Im}(\mathcal{M}_{ii}) \sim |\mathcal{M}_{fi}|^2 \tag{11.46}
$$

Here, in order for calculations in the present theory to converge with the formalism of QFT, the process probability on the left-hand side is linked with the QFT calculation on the right-hand side via the mediation of $\text{Im}(\mathcal{M}_{ii})$. The provisional connection on the side of the present theory with $D_{+}D_{-}$ has been described above and the connection through the optical theorem with $|\mathcal{M}_{fi}|^2$ on the side of QFT was described in section 10.5.

While the structure of QFT on right-hand side of equation 11.46 exhibits the basic property of probability conservation, via equation 11.45, the input from the present theory on the left-hand side provides an explanation of the underlying physical nature of the probabilities in terms of the relative degeneracy of the field redescriptions involved – that is the 'number of ways' in which the event may happen. Essentially the progression from left to right in equation 11.46 represents a complexification of the calculation in order to employ unitarity to gather a normalised expression of the degeneracy count from which particular final states might be extracted with a combined probability of unity.

The fact that the degeneracy count for field redescription sequences may be correlated with Feynman diagrams, as described for figure 11.8, together with the fact that the optical theorem can be demonstrated order by order in perturbation theory via the analysis of Feynman diagrams, as described in section 10.5, suggests that the structure of equation 11.46 might be explored further for low orders of perturbation. Indeed the assumption of perturbation theory, provided the coupling constant is sufficiently small, is that only the first few terms of the expansion of the S-matrix of equation 10.35 are required for precise calculations.

In principle here it might be possible to work backwards from QFT Feynman rules, such as those in table 10.1 based on the Fourier expansions of quantum fields such as $\phi(x)$ in the interaction picture, via the construction of the Feynman propagator $\Delta_F(x-y)$ as implied in the right-hand side of equation 11.42, and use the analogy between the left-hand side of that expression and equation 11.41 to make a detailed connection with the present theory. This connection, employing also the optical theorem, should also provide a guide for deducing a more rigorous mathematical expression for the underlying conceptual picture of the present theory, with the spacetime geometry $G^{\mu\nu}(x)$ constructed in terms of fields such as $A^{\mu}(x)$ and $\psi(x)$ as one of many possible solutions.

On understanding the parallels between QFT and the present theory and making the connection from the right-hand side of equation 11.46 the aim would be to extract from the constraints of equations 11.29 effective Lagrangian terms within the framework of the QFT formalism, expressed in the flat spacetime of special relativity. On importing aspects of the present theory into QFT in this way, with field redescriptions expressed in terms of the algebra of creation and annihilation operators, the aim would be to follow through calculations such as cross-sections using the familiar machinery of QFT.

In this section we have largely considered the alternative route beginning with the provisional picture described in figures 11.4, 11.6 and 11.7 for the present theory leading to the simple relation $P \propto D_+D_-$ for a process probability, with $D_+, D_- \in \mathbb{R}$. Through comparing the structure of figures 11.7 and 11.8(b), via figure 11.8(a), and making the association $D_+D_-\sim \text{Im}\mathcal{M}_{ii}$ this calculation might be 'complexified' as guided by the optical theorem of QFT. In particular a unitarity constraint could be employed to effectively normalise the process probability calculation for all possible outcomes, as expressed in terms of an amplitude $\mathcal{M}_{fi} \in \mathbb{C}$. This complex transition amplitude may then in turn be determined as described in the previous chapter, and in particular in terms of the Feynman propagator $\Delta_F(x-y)$ and Feynman rules, such as those of table 10.1.

This approach is anchored in left-hand side of equation 11.46, with the aim of first motivating all development from the perspective of the present theory in itself. On establishing a link with the framework of QFT various techniques, such as the employment of unitarity in probability calculations, might be extracted from QFT and adapted for use in the framework of the present theory. It may also be possible learn from the relation of QFT to phenomena in condensed matter physics, as we allude to in the following section. Here the aim is to understand the nature of physical particle states and determine cross-sections and other observable quantities within the environment of the present theory, for which the spacetime geometry accompanying empirical phenomena is not flat. However in a suitable limit the present theory may approximate to the form of a QFT in flat spacetime.

The plausibility of either approach, from the left or right side of equation 11.46, rests on the identification of connections between the present theory and QFT which straddle the parallel development of the theories. Such a correspondence will be summarised in points 1) to 7) below. The ultimate aim here would be to comprehend and follow through a complete calculation in the present theory, without any arbitrary reference to standard QFT, and to establish a direct connection with HEP empirical phenomena. However, using the canonical approach to QFT as a close guide is a reasonable strategy since it has been used widely and successfully in practice to obtain results for comparison with experiment.

In the present theory there have been two distinct considerations:

(a) The nature of field redescriptions and an understanding of the permitted elementary exchanges, such as depicted in figure 11.2, according to the various equations of constraint in the theory. This was the topic of the previous section.

The fields such as $A^{\mu}(x)$ and $\psi(x)$ are not introduced onto a pre-existing 4dimensional manifold M_4 , rather spacetime itself, with the spacetime geometry $G^{\mu\nu}(x) = f(A, \psi)$, is shaped by the possibilities of the field descriptions. Hence figure 11.2 should not be interpreted too literally but rather a more dynamical mathematical expression of field redescriptions is desired. This might take the form of equations 11.22 or 11.23 (or 11.30 for the spinor case) in terms of retarded or advanced Green's functions, provided these expressions are compatible with constraints deriving from equations 11.29.

(b) The calculation of the probability of observable processes, for example in HEP experiments, based on a count of the possible internal field degeneracies underlying the process, as depicted for example in figure 11.7. This has been the topic of the present section.

Again here the sequence of $A^{\mu}(x)$ and $\psi(x)$ fields in figure 11.7, superposed as if upon a pre-existing spacetime, presents a somewhat naive and mechanical picture for the degeneracy count. A more conceptually and mathematically rigorous expression of this count may be required to describe the multiplicity of ways in which the geometry of spacetime $G^{\mu\nu} = f(A, \psi)$ may be fabricated out of these fields.

One of the initial aims has been to establish a correspondence between the basic elements of the present theory and those of calculations in QFT. In QFT the construction of the transition amplitude \mathcal{M}_{fi} generally breaks down into very simple elements as described by the Feynman rules, as listed in table 10.1 of section 10.5 for the scalar model. Hence the goal is to explain how the 'number of ways' approach of figures 11.4, 11.6 and 11.7 leads to the Feynman rules which determine the quantity \mathcal{M}_{fi} , and understand why $|\mathcal{M}_{fi}|^2$ should determine the probability for various processes as expressed in cross-section or decay rate calculations.

The parallels identified between the present theory and QFT are listed here. The first six items below loosely correlate with the respective Feynman rules of table 10.1 and the subsequent discussion in section 10.5.

1) The number of ways a series of field redescriptions may unfold through a onedimensional temporal progression with degeneracy D , with terms such as equations 11.41 and 11.44, is analogous to the perturbative expansion of the time evolution operator $U(t, t_0)$ of equation 10.31 in QFT. The 'number of ways' integral sum is naturally normalised by the linear uniform flow of time, with 'one way' for each equal discrete temporal interval Δt_i in equation 11.40 taken to the continuum limit $\Delta t_i \rightarrow 0$ for equation 11.41. This symmetry between equal time intervals implies a flat prior probability distribution as a basis for a Bayesian statistical approach. It then needs to be understood how the propagator Δ_F of equation 11.43, taking the form of equation 10.72, arises as an effective momentum space prior probability distribution when the calculation is restructured as for QFT.

As simply a set of real parameters in the expansion of a field into Fourier modes the variables $k \in \mathbb{R}^4$, which may be interpreted as 4-momentum under $T^{\mu\nu}$:= $-\frac{1}{\kappa}G^{\mu\nu}$, as described in the previous section for $G^{\mu\nu} = f(A)$ in leading from equation 11.6 to equation 11.16, may also appear in factors relating to process probabilities as a result of calculations based on underlying field degeneracy. This is the case for cross-section calculations in QFT with factors of the Feynman propagator $\tilde{\Delta}_F(k) = 1/(k^2 - m^2 + i\varepsilon)$ effectively appearing as a *weight* factor, as for example in equation 10.47. Hence in the restructuring of process calculations for the present theory, via the introduction of T-ordering in equation 10.32 and the resulting Feynman propagators, such prior probability distributions should also appear through this connection with QFT.

2) The redescription expansion is moderated by the need for consistency with the constraint equations. These include the higher-dimensional form of temporal flow $L(\hat{\mathbf{v}}) = 1$ with $D_{\mu}L(\hat{\mathbf{v}}(x)) = 0$ and the original form of the external geometry $G^{\mu\nu}(x) = f(Y)$ with $G^{\mu\nu}_{;\mu} = 0$ throughout; as listed in equations 11.29 and all effectively acting as selection rules for field interactions. Collectively these constraints are analogous to a Lagrangian, including in particular the \mathcal{L}_{int} terms in QFT as associated with the vertices in Feynman diagrams. For the present theory the 'number of ways' a process may occur is taken to be proportional to the couplings implicit in the constraints, such as the factors of \dot{s}_f in the terms of $D_{\mu}L(v_{27})=0$ in equation 11.33, as also discussed after figure 11.5.

In QFT the structures correlating with (a) and (b), listed above for the present theory, are seemingly inextricably linked. The interaction Lagrangian, which is closely associated with the selection rules provided by $D_uL(\hat{v}) = 0$ for example in (a), appears explicitly in the S-matrix, through equations 10.27 and 10.35, which is used in the determination of event probabilities for item (b) above. That is in QFT the mathematical structure of possible field interactions is embedded in the structure of event probability calculations. Effectively this is achieved through the mechanism of 'quantisation' itself, with the expansion of the fields in terms of creation and annihilation operators, which essentially converts a classical composition of fields in an interaction term into a selection rule for contributions to the S-matrix.

In calculations of the transition amplitude \mathcal{M}_{fi} the commutation relations, such as equations 10.16, ensure the correct matching and avoid unwanted cross-terms in compositions of the interaction Lagrangian or Hamiltonian $H_{int}(t)$ in the terms of equation 10.31 and its time-ordered form in equations 10.32–10.34. The sequences of creation and annihilation operators placed between vacuum states also ensures causality in QFT calculations in the sense that any intermediate state must always be created before it is annihilated to yield a non-zero matrix element S_{fi} . Sequences of creation and annihilation operators from the interaction Lagrangian embedded in S_{fi} ultimately determine relative probabilities in the context of all possible processes.

A similar method of 'quantisation' might be employed in the present theory in order to incorporate the constraints of equations 11.29 as selection rules for chains of field redescriptions between initial and final states in a degeneracy count, through the structure of $R(t)$ in equations 11.41 and 11.44 for example.

3) A free field solution for $A^{\mu}(x)$ under $G^{\mu\nu} = f(A)$ in the form of equation 11.1 may be expanded in terms of $e^{\pm i k \cdot x}$ Fourier modes as described in equation 11.6, as consistent with Maxwell's equations under $G^{\mu\nu}{}_{;\mu} = 0$. Exchanges between fields such as $A^{\mu} \leftrightarrow \overline{\psi} \gamma^{\mu} \psi$ are considered in terms of the complex Fourier modes of the fields. Similarly for QFT calculations as presented in chapter 10 using the interaction picture, as discussed after equation 10.26, between the initial and final plane waves of the form $e^{\pm ik\cdot x}$ the state evolution is mediated by an expansion of free fields of the form in equations 10.13–10.15, which are solutions of the Klein-Gordon equation for the scalar model.

In the canonical quantisation approach to QFT, as described in chapter 10, anni-

hilation and creation operators, such as $a(p)$ and $a^{\dagger}(p)$, are associated with the Fourier modes $e^{-ip\cdot x}$ and $e^{+ip\cdot x}$ of the field respectively, as seen in equation 10.51 for $\hat{\phi}^+(x)$, equation 10.52 for $\hat{\phi}^-(x)$ and equation 10.13 for the complete free scalar field $\hat{\phi}(x)$. In a QFT calculation the complex plane waves of the form $e^{\pm i k \cdot x}$ representing the incoming and outgoing particle states are linked by a chain of creation and annihilation operators for a variety of fields to determine the transition amplitude as described for example in equation 10.36. This structure, employed throughout the calculations in the interaction picture, provides a close analogy with the present theory.

The quantum field $\phi(x)$ of equation 10.13 does not represent a solution of the equations of motion given an interaction, nor does it represent a physical entity in any context. Rather this expansion $\phi(x)$ carries the potential for all possible transitions for the corresponding classical field in terms of Fourier components. This is the interpretation in the present theory, for which such quantum field expansions might be employed in the construction of chains of field redescriptions, expressed in terms of complex Fourier modes and employed in a degeneracy count for any process.

4) The geometric constraint $G^{\mu\nu}_{;\mu} = 0$ over the external 4-dimensional spacetime, with energy-momentum $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$, implies the conservation of 4-momentum for all possible field redescriptions (in the flat spacetime limit considered here, as discussed before equation 11.14). In QFT calculations the time integral $\int dt$ over the interaction Hamiltonian H_{int} is replaced by a manifestly Lorentz invariant spacetime integral $\int d^4x$ over the interaction Lagrangian density \mathcal{L}_{int} via equation 10.27 which, as seen for example in the lines of equations 10.46, leads to the constraint of 4-momentum conservation for each interaction vertex as expressed by the δ^4 -functions.

Whether spacetime integrals, as a generalisation of purely temporal integrals, might feature in a generalisation of the field redescription degeneracy count for solutions underlying a particular geometry $G^{\mu\nu}(x)$ is open to consideration. However here the field exchanges have been considered to take place purely through a temporal progression, consistent with the notion of a fundamental one-dimensional progression in time that underpins the conceptual basis of the whole theory. In any case, in the present theory 4-momentum conservation is ensured through the prevailing relation $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$ and the identity $G^{\mu\nu}_{;\mu} = 0$ which hold throughout spacetime and in particular for local exchanges of the underlying fields. For such exchanges applied to the Fourier modes such as $A^{\mu}(x) \sim e^{-ik \cdot x}$ and $\overline{\psi}(x) \gamma^{\mu} \psi(x) \sim e^{-ip_1 \cdot x} e^{-ip_2 \cdot x}$ for example the 4-momentum conservation in a $A^{\mu} \leftrightarrow \overline{\psi} \gamma^{\mu} \psi$ field redescription takes the form of the mutual condition $k = p_1 + p_2$. This is essentially implied in the requirement that *locally* the spacetime geometry $G^{\mu\nu}(x)$ itself is unchanged for such an underlying field redescription.

5) In the present theory an infinity in the degeneracy count occurs when for example the intermediate $A^{\mu}(x)$ field state in figure 11.4 is augmented for a further intermediate redescription in terms of a pair of fields, such as $A^{\mu} \to \overline{\psi} \gamma^{\mu} \psi \to A^{\mu}$ as shown in figure 11.6. Here the degeneracy count of equation 11.44 will be further augmented as the field $A^{\mu}(x) \sim e^{-ik \cdot x}$ is replaced by the field $\overline{\psi}(x) \gamma^{\mu} \psi(x) \sim$ $e^{-ip_1 \cdot x}e^{-ip_2 \cdot x}$ up to a mutual freedom in the share of the total 4-momentum between p_1 and p_2 , accounting for an infinite degeneracy of solutions, as described after equation 11.44.

This is closely analogous to the ambiguity in the 4-momentum carried by an internal loop in a Feynman diagram, such as that in figure 10.9 leading to the divergent momentum integral $\int d^4r$ in equation 10.86, and as frequently encountered in QFT. In both cases a means of 'renormalisation' is required in order to obtain a finite calculation. By matching such infinities in the present theory with the analogous quantities in QFT a similar program of renormalisation might be obtained for the present theory, although with a different interpretation as will be described in the following section. Indeed, the degeneracy count for any given process in any case stands in need of a 'normalisation' with respect to the count of the number of ways in which anything can happen.

- 6) Various combinatoric factors due to permutations of interactions for higher-order field redescriptions, or symmetries between identical particle states, will need to be assessed for the present theory and related to the corresponding factors based on the analysis of Feynman diagrams in QFT. Discrete sums over field degrees of freedom such as spin in QFT also reflect the number of ways a process may occur.
- 7) The need to match both the $e^{-ik\cdot x}$ and $e^{+ik\cdot x}$ complex Fourier modes of the fields, through independent chains of degeneracies D_+ and D_- , underlying a real expression of $L(\hat{\bm{v}}) = 1$ and $G^{\mu\nu} = f(Y, \hat{\bm{v}})$, means that an overall event probability is of the form $P \propto D_+ \times D_-$ as described for figure 11.7 (rather than $P \propto D$ alone from 'item 1)' above). For practical calculations it is the relative ratios of the degeneracies for the range of possible processes that is needed to obtain actual probabilities with $\sum_{F} P_{F} = 1$, for a sum over all possible final states F arising from an initial state interaction, including the case for which the final state is identical to the initial state.

The calculation of D_+D_- is correlated with the determination of Im(\mathcal{M}_{ii}) in QFT, as described for figure 11.8, which via a complexification of the calculation and the optical theorem is then closely related to the amplitude squared $|\mathcal{M}_{fi}|^2$ in QFT as described for equation 11.46. Expressed this way the unitary symmetry applying to the complex amplitudes \mathcal{M}_{fi} models the conservation of the total probability, implicitly normalising the degeneracy count for all possible processes. The fact that renormalisation is required in QFT shows that this application of unitarity is only partially successful, and does not necessarily automatically normalise the degeneracy count completely. Indeed even for a renormalisable QFT finite calculations might not be achievable at a very high order of perturbation, and in general a more watertight method of normalisation might be sought for the present theory.

For a complete calculation in this theory, putting all of the pieces together, the actual value of the probability P_F for a process yielding the final state F is determined by the relative, rather than absolute, number of ways in which it can occur, essentially as is the case for the probabilities of classical physics. For example degeneracy counts over the infinite possibilities in the timing of field redescriptions, such as those in equation 11.41, may be independent of the choice of the external fields, as for example in figure 11.4 which may describe the leptonic or quark final states for figure $11.5(a)$ or (b). More generally the infinities in the count of the number of ways will be in common for a range of competing processes and will cancel in the calculation of physical quantities such that the total probability for any outcome will necessarily satisfy the requirement $\sum_{F} P_{F} = 1$. Some care will then be needed in this theory to deal with infinities that arise in the stages of such calculations. However, since all probabilities are normalised by the total degeneracy for any process the bound $0 \leq P_F \leq 1$ will apply trivially. A relative infinity of ways to produce one particular final state F will result in a probability $P_F = 1$, which may be problematic in terms of comparison with the corresponding empirical value, but it is not possible for the theory to yield a nonsensical infinity for the calculated value.

The calculation of probabilities via a complexification may prove an effective technique to apply for the present theory, once the relation between the underlying real number measure of degeneracy and the QFT calculation through equation 11.46 has been fully understood. In this translation of the calculation a 'unitarity' condition will model probability conservation, consistent with kinematic factors appearing through the propagators, as described for 'item 1)' above, provided the ultimate expression for the probability is a dimensionless quantity.

While the seven points listed above express a close parallel between structures in the present theory and perturbative calculations in QFT, as well as obtaining the Feynman rules for \mathcal{M}_{fi} the full cross-section expression is needed for comparison with empirical data. For QFT the structure of the cross-section σ was introduced in equation 10.3 as a product of three factors, namely the amplitude squared $|\mathcal{M}_{fi}|^2$ together with the initial state flux factor and the final state Lorentz invariant phase space $d\Phi$. Various normalisation factors such as the volume V and time interval T of the interaction cancel in forming this expression.

The probability for a process, whether expressed in terms of a degeneracy count or not, should be a dimensionless quantity, as is the transition amplitude \mathcal{M}_{fi} for the two-body final states considered in chapter 10. In general \mathcal{M}_{fi} need not be a dimensionless quantity provided the cross-section has the dimension of a length squared, as for example in equation 10.12, and as described in the discussion following equation 10.86.

The present theory may involve a different breakdown across the three factors composing the expression for the cross-section, compared with that displayed for example in equation 10.7, with the form of the appropriate normalisation for all three factors, including those for the initial state flux factor and final state phase space, possibly differing also from the QFT case. For the present theory, as for QFT, it is ultimately the calculated cross-section that is required to be of the appropriate form in the context of equation 10.1. The normalisation of factors required for consistency with the cross-section having the mass dimension $D = -2$ will be closely correlated with the normalisation employed to obtain dimensionless probabilities that sum to unity.

As described for equation 10.7 in section 10.2 the event rate is proportional to the initial state luminosity and flux factors and final state phase space, as would be expected based on a classical notion of probability. In this section we have argued for the replacement or interpretation of the central term $|\mathcal{M}_{fi}|^2$ in this expression in the form of a quantity representing an underlying measure of the degeneracy of ways in which the process may occur, in terms of sequences of field redescriptions, and hence constituting a further purely statistical factor having essentially the same character as a classical probability.

Explicitly, in this section we have considered field interactions in terms of possible field redescriptions, involving the $e^{\pm i k \cdot x}$ Fourier modes of the fields, as expressed for example in equations 11.22 and 11.30 of the previous section, causally linked together to mediate observable processes, conceived as a field sequence such as depicted in figure 11.4 or 11.6 and combined as for figure 11.7, and as allowed by the form of constraint equations 11.29 such as $D_{\mu}L(\hat{v}) = 0$ and $G^{\mu\nu}_{;\mu} = 0$. This however leads to a picture of the extended spacetime geometry $G^{\mu\nu}(x)$ itself constructed as one solution out of a myriad of possible ways based on local field description degeneracy, again subject to the constraint equations, not only for HEP processes but everywhere throughout the 4-dimensional spacetime world.

This implies a conception of HEP phenomena, such as an $e^+e^- \to \mu^+\mu^-$ event, supported by the underlying field exchanges which seamlessly also support the macroscopic physical world including the detector apparatus itself. In turn the physics of quantum mechanics is seamlessly connected to the world of classical physics. In the section 11.4 we further explore this conceptual picture within which quantum and particle phenomena are found alongside macroscopic objects and gravitation in a unified framework.

While the above seven points provide a useful guide into the workings of such calculations ultimately a stand-alone approach within the present theory may be desired. In this way the aim is to achieve explicit calculations for comparison with HEP processes such as e^+e^- collisions for the full theory. To make a detailed comparison between the present theory and HEP data ultimately the particle concept, and in particular the nature of the 'in' and 'out' states at a collider experiment, will need to be understood within the context of the present theory. This will require an understanding of the nature of physical particle states propagating in spacetime in general, relating to a fully 'renormalised' expression of field exchanges, rather than representing particle states in the form of simple $e^{-ik\cdot x}$ plane waves as for QFT. This direction will be explored in the following section.

11.3 Renormalisation and Particle States

The relation $G^{\mu\nu} = f(A)$ derives from the internal $U(1)_Q \subset E_7$ action within the isochronal symmetry of $L(v_{56}) = 1$, and is expressed explicitly in equation 11.1 as determined through the analogy with Kaluza-Klein theory as described in section 5.1. In deriving directly from the basic structure of the theory, through equation 2.30 applied to the full symmetry group, the internal gauge field component $A^{\mu}(x)$ itself, which appears in expressions such as $D_{\mu}L(v_{56}) = 0$ for the broken full symmetry, can be considered as a 'bare' or elementary field at the 'microscopic' level from the point of view of QFT. This same field, implicit in equation 11.1 and hence satisfying the relation $\Box A^{\mu} = 0$ of equation 11.2, in the Lorenz gauge, is also essentially the classical gauge potential of Maxwell's electrodynamics of 1864, associated with directly observable laboratory effects. In this sense, again from the point of view of QFT, the gauge field $A^{\mu}(x)$ can be considered as a 'dressed' or renormalised field at the 'macroscopic' level. The question then arises as to how these two views of the same field $A^{\mu}(x)$ are consistent.

For a non-Abelian gauge field $Y^{\mu}(x)$ the relation $G^{\mu\nu} = f(Y)$ of equations 11.29, that is the classical field expression of equation 5.20, contains self-interaction terms as described in equations 11.36–11.38. Hence, even from the perspective of gauge fields alone, the macroscopic form for $G^{\mu\nu} = f(Y)$ will be necessarily shaped and corrected as a consequence of the multiple solutions for the spacetime geometry, as built upon a degeneracy of underlying gauge field redescriptions, with the constraint $G^{\mu\nu}_{\mu\mu} = 0$ holding throughout the base manifold. However, in the full theory the gauge field $A^{\mu}(x)$, associated with the internal Abelian U(1)_Q gauge group, is also not free since it couples to fermions through the constraints of equations 11.29, as seen in the terms of equation 11.33 for example. Through field exchanges as considered for equation 11.22 Maxwell's equation is modified to the form of equation 11.26, with a source term deriving from the fermion components. Hence it is necessary to consider the macroscopic form of the spacetime geometry $G^{\mu\nu}(x) = f(A, \psi)$, and understand how this relates to the original classical expression for $G^{\mu\nu}(x) = f(A)$ and also to empirical phenomena.

Empirically electromagnetic waves are observed to propagate 'at the speed of light' effectively according to Maxwell's equation $\Box A^{\mu} = 0$, with solutions such as that in equation 11.6 for the transverse polarisation states $r = 1$ or 2 and with $k^2 = 0$. Hence the overall form of the function $G^{\mu\nu}(x)$, on the left-hand side of equation 11.12 with $G^{\mu\nu}_{\mu\nu} = 0$ implied in equation 11.14, appears to be completely transparent to underlying exchanges of indistinguishable fields, with possible intermediate stages similar to those of figure 11.6 or 11.7, which percolate down through higher orders with the spacetime geometry $G^{\mu\nu}(x)$ always preserved over the possible field redescriptions. That is, unlike the general case, the underlying gauge-fermion field redescriptions appear to make little or no impression on the spacetime geometry associated with an electromagnetic wave – with $G^{\mu\nu}(x) = f(A, \psi) \simeq f(A)$ which takes the shape of $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu}$ as depicted in figure 11.1 for example.

In QED these higher-order solutions are described in terms of photon selfenergy contributions, as shown for example in the Feynman diagrams of figure 11.9.

Figure 11.9: A series of possible Feynman diagrams which 'dress' the original 'bare' photon propagator, which itself corresponds to the first diagram alone.

The particles observed in experiments correspond to renormalised states of the fields. The quanta of the electromagnetic field are massless, even though the higherorder corrections to the photon propagator in figure 11.9 contain virtual particles such
as e^+e^- and $d\bar{d}$ pairs. In QED the preservation of the bare photon mass $m_\gamma = 0$, and hence the equation of motion $\Box A^{\mu}(x) = 0$, for the renormalised field is explained in terms of Ward identities (see for example [70]). This observation in QED is analogous to the transparency of the geometry $G^{\mu\nu} = f(A)$ to higher-order microscopic field redescriptions in the present theory, maintaining the macroscopic field condition $k^2 =$ $m^2 = 0$, and a correlated mathematical explanation might be sought here.

In the standard theory of QED the behaviour of the field $A^{\mu}(x)$ deviates from that in classical electrodynamics due to the properties of low energy e^+e^- pairs. In HEP experiments an effective internal structure of the photon is manifested in 'twophoton collisions', such as the process $\gamma\gamma \to c\bar{c}$ induced and observed at e^+e^- colliders. In such experiments the photon expresses itself in revealing the internal structure of its dressed state. Equivalent empirical effects are expected to arise from the principles of the present theory, with the internal structure of matter composed of endless possible internal 'bare' field redescriptions. Here for example solutions for $G^{\mu\nu} = f(A, \psi)$ may take the effective macroscopic form of an electromagnetic field alone, such as the wave solution in equation 11.6, while implicitly containing a myriad of possible field components and hence carrying the potential for the associated interactions as seen for example in two-photon collisions.

The mathematical divergences associated with higher-order loop diagrams in QED are tamed by accepting the non-physical nature of quantities such as 'mass' and 'charge' in the bare Lagrangian and instead aligning the physical parameters of the renormalised theory with empirical values of mass and charge, as described briefly following equation 10.86. The effect of combining an empirically measured generic coupling parameter g with quantum corrections determined in theory, through the machinery of renormalisation in QFT, leads to the observable phenomenon of 'running coupling' in which the parameter q is found to depend on the energy scale E as described by the 'renormalisation group equation':

$$
\frac{d}{d\ln E} g(E) = \beta(g(E))\tag{11.47}
$$

The function β depends upon the particular theory. In the Standard Model β is positive for the U(1)_Y gauge group and negative for the non-Abelian internal symmetries resulting in the running coupling shown qualitatively in figure 11.10.

The energy dependence of the coupling g, representing the general case in equation 11.47, is independent of the bare Lagrangian parameters, and also independent of the regularisation method and parameters used to temporarily suppress the divergences in the process of renormalisation.

As described in section 10.5 generally a quantum field theory is renormalisable, and finite results may be obtained for comparison with experiment, if the coupling parameter q is of mass dimension M^D with $D \geq 0$. All of the couplings for the Standard Model, such as g and g' in equation 7.40, have $D=0$ and the corresponding QFT is just renormalisable. Even here though for the renormalised Standard Model divergences remain in the sense that the expansion series for equation 10.35 in equation 10.29 does not generally converge at higher orders, although the problem does not become apparent until terms of approximately order 137 in the case of QED for example ([26] p.681, this is the point alluded to at the end of 'item 7)' in the previous section), far beyond the first few orders needed for calculations in practice.

Figure 11.10: The running coupling g' , g and $g_s = \sqrt{4\pi\alpha_s}$ respectively for the U(1)_Y, $SU(2)_L$ and strong $SU(3)_c$ gauge interactions in the Standard Model. Extrapolated from their laboratory values over a number of orders of magnitude in energy scale, via equation 11.47 with conventional β functions, the three parameters mutually intersect, although not simultaneously, at around 10^{14} – 10^{16} GeV ([70] p.787).

The structure and tools of QFT have a broad scope of applications and do not necessarily describe the fields or particle states of a 'fundamental' theory. An effective quantum field theory is one which is only valid as a physical theory below a certain energy threshold and describes particle states appropriate within that energy range. Such an effective QFT, for example a theory for nucleon-pion scattering, is necessarily an approximation to nature, with different physics and new particle states observed at higher energy. The interpretation of particles associated with an effective field theory, such as nucleon and pion states, as 'fundamental' particles is hence unsatisfactory.

Renormalisable QFTs such as the Standard Model are also considered to be low energy effective field theories. The form of the renormalisation group equation, and contact with empirical observations, is insensitive to high-energy, short-distance phenomena, which are also unknown. Hence QFT provides a *phenomenological* framework for particle physics with fields in the Standard Model Lagrangian transforming under the $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge group describing the types of particles that are observed in high energy physics experiments. The theory applies over a wide energy range and provides a unifying framework incorporating weak and strong, in addition to electromagnetic, interactions. The corresponding quanta of the Standard Model quantum fields describe the particle states of leptons, quarks, gauge bosons and the Higgs, all of which from an empirical point of view appear to be elementary. However this is not a conclusion that can be drawn from the QFT for the Standard Model itself.

A more fundamental theory is needed to ascertain the true elementary structures of nature. The renormalisation for the QFT of the Standard Model has had great pragmatic success in particle physics but, as well as being insensitive to the method by which divergences are 'cut-off', in general has very little to say regarding the structure of an ultimate high energy theory. Hence the results of the Standard Model renormalised QFT are plausibly consistent with an underlying theory for which interaction probabilities are fundamentally expressed in terms of a degeneracy count of possible redescriptions of the underlying field function as proposed in this paper. The present theory aims to describe the actual nature and behaviour of physical entities down to arbitrarily short distances and up to any energy scale.

Indeed the present theory is intended to be a fundamental, rather than an effective, theory, in contrast with the Lagrangian approach, as has already been emphasised in section 5.2 and as will be discussed further in section 15.2. The present theory is also completely 'renormalisable' in an essentially trivial way since probabilities are constructed simply in terms of the relative 'number of ways' field solutions may be obtained. These involve nested sequences going down through higher orders of field exchanges, as depicted for example in figure $11.11(a)$, which itself represents a higher-order extension from the form of figure 11.6 for a single $e^{-ik \cdot x}$ field component sequence, here depicted alongside the associated Feynman diagram.

Figure 11.11: (a) A higher-order sequence of field exchanges for the process $e^+e^- \rightarrow$ $\mu^+\mu^-$ together with (b) the correlated Feynman diagram with internal loops. This figure is similar to figure 11.8, except here with differing initial and final states and without a fold or cut line.

Even considering the degrees of freedom of the field redescription timings t_i the sum of possible ways is infinite, as described following equation 11.44. Further, the internal 4-momentum freedom for the $d\bar{d}$ field state, for example, in figure 11.11(a) translates into the divergent momentum integral for the corresponding $d\bar{d}$ virtual particle loop in the Feynman diagram of figure $11.11(b)$, as described in 'item 5)' of the previous section. For yet higher orders this structure implies a nested product of infinite sums and integrals which would appear to more and more dominate calculations for more and more 'dressed' diagrams. However it is conceivable that such infinite degeneracy counts largely cancel, resulting in a non-trivial finite calculation of cross-sections or branching ratios.

For example, by relabelling the final state, figure $11.11(a)$ can be considered

to represent a field sequence underlying either an $e^+e^- \to \mu^+\mu^-$ or $e^+e^- \to d\bar{d}$ event, amongst other possibilities. Since the intermediate redescriptions in figure 11.11(a) are applicable for both processes $e^+e^- \to \mu^+\mu^-$ and $e^+e^- \to d\bar{d}$ the relative 'branching ratio' to obtain the final state $d\bar{d}$ is simply:

$$
BR(e^+e^- \to d\bar{d}) = \frac{\text{sum of ways for } d\bar{d}}{\text{sum of ways for } \{d\bar{d} \text{ or } \mu^+\mu^-\}} \to \text{``}\left(\frac{\infty}{\infty}\right),\tag{11.48}
$$

For either final state there is an infinite degeneracy of intermediate states owing to the implied unconstrained 4-momenta for example. These infinities clearly cancel in calculations such as equation 11.48 since there is a similar, in fact here identical, 'degree of divergence' in each case. Indeed generally in forming measurable branching fractions cancellation between common factors will provide the main source of normalisation. Further normalisation factors will be involved in deriving event rates and cross-sections such as $\sigma(e^+e^- \to \mu^+\mu^-)$, as described towards the end of the previous section.

A similar situation arises in QFT with for example the $4th$ order Feynman diagram in figure 10.9 together with the same diagram with the final state relabelled by $\mathcal{X}^+\mathcal{X}^-$, for the processes $\mathcal{X}^+\mathcal{X}^- \to \mathcal{Y}^+\mathcal{Y}^-$ and $\mathcal{X}^+\mathcal{X}^- \to \mathcal{X}^+\mathcal{X}^-$ respectively, both of which contain loops with infinite degrees of freedom in terms of the corresponding momentum integrals. In QFT the methods of renormalisation lead to finite crosssections and branching fractions for comparison with the empirical data. In fact in QFT tree level diagrams, such as figure 10.4, already give a good approximation for the rates of such processes, provided the coupling constant is sufficiently small. This is the case for QED in which the cross-section calculation based on the tree-level diagram in figure 10.3 gives a good approximation for the process $e^+e^- \rightarrow \mu^+\mu^-$ as described towards the end of section 10.2.

For the present theory based fundamentally on degeneracy counts the interpretation of equation 11.48 may be contrasted with the case of Newtonian calculus in which the ratio $\frac{\delta y \to 0}{\delta x \to 0}$ has a well defined meaning and value since δy and δx tend to zero in a related manner through a continuous function $y = f(x)$. Here, in a similar and yet complementary situation, the limit of the ratio $\sum_{n=1}^{\infty}$ $\sum_{x\to\infty} y\to\infty$ in equation 11.48 gives a finite and well defined result due to the close relationship between the divergence in the numerator and that in the denominator.

This a very literal notion of (re)normalisation in calculating probabilities. It is analogous to everyday cases such as the probability of hitting the '20' on a dartboard. There are an infinite number of ways in which the point of the dart can land on the surface of the 20 segment. However this infinity is normalised by the infinite number of ways of landing in any other region such that the total probability is finite and approximately $\frac{1}{20}$ (for a suitably random dart thrower). Alternatively the sum over points may be quantified as a finite integral over surface area, rather like the finite integral over possible field redescription times t_i in equation 11.41 for figure 11.4 as a measure of the sum of ways to describe the underlying field function.

The above analogy demonstrates the close association of classical and quantum probabilities in the present theory as will be discussed further in the following section. The cancellation in equation 11.48 not only applies for the infinite degeneracy of field redescription times t_i but also for the unrestricted internal momentum freedom, implicit in the $d\bar{d}$ internal state of figure 11.11(a) for example, which is also infinite in terms of a real-valued $\int d^4k$ measure.

In practice calculations of branching fractions and cross-sections may be much more readily performed by noting the symmetry of the system (analogous for example to the equal sizes of the twenty segments on a circular dartboard in the metaphor described above). In the case of QFT unitary symmetry, in calculations based on complex amplitudes, is applied to model the conservation of probability; and yields successful results when supplemented by the techniques of renormalisation. However these calculations, founded on postulated complex-valued entities, miss the physical meaning of the infinities as a real-valued degeneracy in the number of ways a process can occur. Hence in the present theory renormalisation based on a real degeneracy count is expected to be closely related to QFT renormalisation based on complex objects, such as amplitudes and propagators, with similar conclusions except with finite results necessarily to all orders in the present theory.

While generating finite results when normalised for specific processes it is plausible that the sums and integrals over the myriad of continuous possibilities, such as for figure $11.11(a)$, and for an endless range of higher-order field sequences, may have residual effects such as the dependency on the energy scale of physically measurable interaction strengths as described by the running coupling in figure 11.10. Underlying differences in branching ratios such as equation 11.48 will then depend directly upon differences in the 'bare' couplings associated with the field redescriptions such as $A^{\mu} \leftrightarrow \psi \gamma^{\mu} \psi$ for example. These include the \dot{s}_f real coefficient factors of magnitude 1 or $\frac{1}{3}$ for the U(1)_Q coupling in equation 11.33, applied for the outer layer of field exchanges, that is in the external vertices of the Feynman diagram as for example in figure 11.5(a) and (b) and as described there in the subsequent text, and will apply here for the final field redescription at time $t = t_1$ in figure 11.11(a). (In QED there are Ward identities which both preserve the bare value of the photon mass $m_{\gamma} = 0$, as alluded to earlier in this section with reference to figure 11.9, and also which preserve the ratios of charges through renormalisation, and again a correspondence might be sought with the structures of the present theory.)

The field redescriptions underlying the many solution possibilities are profusely diffused throughout spacetime, from the temporal origin of the universe in the Big Bang, shaping the initial conditions for the evolution of the cosmos as considered in the following two chapters, to the quantum effects observed in laboratory experiments such as that represented in figure 10.1 and described further in the following section. As well as the photon 'self-energy' contributions of figure 11.9 the field redescriptions 'dress' the initial and final state particles for an event observed at a collider experiment. These higher-order solutions include the final state processes suggested by the Feynman diagrams in figures 11.5(c) and (d). Since the $d\bar{d}$ fields undergo strong $\text{SU}(3)_c$ interactions, producing an observed final state $\pi^+\pi^-$ pair for example, *objectively* it might be expected that many more spacetime world solutions with a $d\bar{d}$ compared with a $\mu^+\mu^-$ final state might be identified, in the context of a grand ensemble of all possible $G^{\mu\nu}(x) = f(Y, \hat{v})$ solutions on M_4 . This consideration would suggest that the branching ratio of equation 11.48 should effectively be unity, owing to the apparent relative infinity of ways to produce a $d\bar{d}$ rather than a $\mu^+\mu^-$ final state. As pointed out in the discussion after 'item 7)' in the previous section such a conclusion for the present theory, although being internally consistent, would appear to be drastically incompatible with empirical phenomena.

However laboratory phenomena, as for all observations, subjectively evolve progressively in time. At the time t_1 of the final field redescription in figure 11.11(a) the likelihood of a field exchange will depend upon the $U(1)_Q$ coupling \dot{s}_f regardless of what has happened before or what can happen after. Hence the charge value of 1 or $\frac{1}{3}$ will dominate the cross-section. Subsequent field redescriptions and interactions for the final state produced, as represented in terms of the Feynman diagrams in figures $11.5(c)$ and (d) for example, will not affect the branching fraction calculation for equation 11.48 other than through their implications for a final state phase space factor, which in the present theory correlates with the range of spacetime geometries associated with a particular set of final state particles.

More generally out of the grand ensemble of all possible $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ solutions it might be expected that a typical world would be dominated by strong $SU(3)_c$ interactions and corresponding forms of matter, since a relatively much larger range of field redescriptions are possible, via the set of eight self-interacting gluons, compared with other kinds of Standard Model interactions. However we do not apprehend a full 4-dimensional universe all together in its full temporal extent as a given object, rather we subjectively sample a possible world progressively through time. The corresponding progressive accumulation of probabilities selects a type of possible $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ solution which is extremely rare in the context of the full ensemble, with a sparser more open form of matter shaped by a more democratic contribution from the components of $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge interactions.

That is the $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ solution that we observe is selected with all probabilities oriented with respect to an underlying one-dimensional temporal flow from the past to the future, moulding the matter content and laws of physics for such a universe, with the structure of causality built into the world we perceive. As for perception of the world in space and time itself, this subjective causal aspect of observations is a further necessary a priori structure through which we experience the world, as will be discussed further in chapter 14.

While the accumulation of probabilities along a causal path through a choice of world solutions shapes the macroscopic properties of matter on the large scale, the probabilities locally determine the relative likelihood to achieve different outcomes such as for example the event $e^+e^- \rightarrow d\bar{d}$ or $e^+e^- \rightarrow \mu^+\mu^-$ in a HEP experiment as described above. Once the final state particles, such as $\pi^{+}\pi^{-}$ or $\mu^{+}\mu^{-}$, are formed and propagate through spacetime to the extent that the macroscopic shape of $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ diverges the relative degeneracy count of field redescriptions for different processes under the same $G^{\mu\nu}(x)$ geometry no longer applies. That is a branching ratio such as equation 11.48 is determined by the relative number of world solutions effectively within a local finite spacetime volume (similarly as represented by VT for the QFT calculations described in section 10.2) with a common local geometry described by $G^{\mu\nu}(x)$, regardless of what can happen after the final states form.

In quantum mechanics the causal sequence of probabilities is reflected in the evolution of the wavefunction Ψ in equation 11.51 below as punctuated by apparent 'collapses' of the wavefunction, as will be discussed in the following section. As will also be described further in the next section the underlying statistical origin of quantum phenomena in the present theory is very similar in nature to that for a classical statistical system, with outcomes essentially determined by the 'number of ways' in which something can happen. The causal accumulation of probability, that is the temporal ordering property as described above for quantum phenomena in the present theory, naturally also applies for systems of classical physics. In the classical world the temporal ordering of probabilities underlies the second law of thermodynamics for example, which will be considered in relation to the very early universe in section 13.2.

The statistical approach underlying quantum phenomena in the present theory, fundamentally based on a real-valued degeneracy of field possibilities, has a microscopic structure analogous to that studied in the classical physics of critical phenomena. There the forces and behaviour of basic elements of condensed matter systems, such as magnets or fluids, are sufficiently well known to be modelled and parametrised. There is also a close relationship between such systems and quantum field theory at the phenomenological level – in fact a correspondence can be identified between renormalisation in QFT and the theory of critical phenomena which leads to a principle of universality for statistical fluctuations, which is equivalent to the cut-off independence in QFT ([70] p.268). However, although the empirical tests in HEP have been very successful, in the case of the QFT for the Standard Model the short-distance physics, only provisionally represented by field parameters in the bare Lagrangian, is essentially unknown, as alluded to earlier in this section.

Potentially the present theory extends the analogy between HEP phenomena and critical phenomena conceptually as well as mathematically, with the microscopic world being 'modelled' on the idea of underlying field redescriptions. This makes a closer relation to the theory of condensed matter systems than for standard QFT, with the latter founded pragmatically on calculations based on complex transition amplitudes.

In principle the present theory reaches down without limit into the microworld revealing an internal structure in terms of nested multiple field solutions continuing indefinitely in almost fractal-like manner, analogous to the perturbative expansion of the QFT time-evolution operator expressed as an infinite series in equation 10.31. On the other hand the scope of the theory in principle also feeds upwards and seamlessly into the phenomena of condensed matter physics itself, with magnetic and fluid properties emerging at the macroscopic level, and into the realm of classical physics and classical probabilities, as alluded to above and described further in the following section.

Out of the construction of the spacetime geometry over sequences of field degeneracies, of arbitrary high order, it is suggested that the phenomena of particle states themselves arise, apparently propagating through field configurations in spacetime in the fully 'renormalised' theory and mutually interacting, accounting for the phenomena observed in HEP experiments. This picture of particle states brings to mind the excitations of 'phonons' in the medium of a solid state device, with here the colourful variety of Standard Model particle types arising out of the variety of underlying internal field interactions allowed by the broken form of $L(\mathbf{v}_{56}) = 1$ and $D_{\mu}L(\mathbf{v}_{56}) = 0$ and the constraint equations 11.29 on M_4 in general.

As for the Standard Model, in the present theory particle masses arise through the interactions of the corresponding field with a 'Higgs' field. Here a vector-Higgs field is associated with the components of $h_2 \equiv v_4 \subset v_{56}$ of equation 9.46 projected onto the local tangent space TM_4 , with the effective Higgs phenomenology provisionally identified as described in subsection 8.3.3. As well as the selection of the external TM_4 subspace component of $F(\mathbf{h}_3\mathbb{O})$ here 'spontaneous symmetry breaking' is also realised in terms of a particular choice of vector field $v_4(x)$ which may 'point' in an arbitrary direction at any given location $x \in M_4$. This structure may be closely relate to the statistical methods employed in spontaneous symmetry breaking for critical phenomena, as for example associated with the properties of ferromagnetism. Recalling that the Higgs mechanism was developed from the early 1960s through analogy with spontaneous symmetry breaking phenomena as originally conceived in condensed matter physics this observation sees the Higgs concept returning to familiar territory.

Low energy effective phenomena might also arise and be related to the Standard Model, which itself may considered to be an effective field theory as discussed after figure 11.10. In this case while some components of $v_{56} \in F(h_3\mathbb{O})$ such as the Dirac spinors ψ might correlate directly with elementary fermion states, the vector-Higgs v_4 components may correlate less directly with the empirically observed scalar Higgs particle. In the Standard Model this latter state is itself treated as a 'fundamental particle' in the effective theory with symmetry breaking modelled by a scalar Higgs field ϕ in the contrived potential of equation 7.53 as described in section 7.2. In the present theory the degree of freedom $|v_4|$ is considered as a candidate for a field underlying the observed Standard Model scalar Higgs particle, which hence does not correspond to a fundamental scalar field in the components of $L(v_{56}) = 1$ projected over M_4 . While the scalar condensates of technicolor models, described shortly before equation 8.73 in subsection 8.3.3, are analogous to BCS pairs of electrons bound through interactions with phonons in solid state devices, a different relation to condensed matter systems might be sought for the present theory since here technicolor gluons are not required to bind the scalar Higgs together.

While the microworld is infused with field function redescriptions, such as $A^{\mu} \leftrightarrow \overline{\psi} \gamma^{\mu} \psi$, in the multiple solutions under $G^{\mu\nu}(x)$ physically transmitted real particle states, such as electrons and photons, as detected in HEP experiments propagate over macroscopic distances with measurable and regular properties such as mass m , charge e, spin s and average lifetime τ . These features, which define the particle types, are regular and reproducible and hence must to some degree arise as the properties of self-sufficient discrete entities, in the sense of being generally independent of the conditions under which they are produced and the environment within which they are observed. Such real propagating particle states are associated with a distinct impression in the spacetime geometry, that is the form of $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, itself. In propagating over macroscopic distances particle states, such as photons and electrons, are revealed through their observable apparent interactions between each other and with the elements of macroscopic apparatus.

An electron state in the e^- beam of a particle accelerator for example is in constant interaction with the electromagnetic fields produced by the accelerating, bending and focussing components of the machine, via the elementary field exchanges depicted in figure 11.12(a). Even for a freely propagating electron interactions with an electromagnetic field $A^{\mu}(x)$ are present in terms of internal 'self-energy' possibilities, similar to those for the photon in figure 11.9, as shown here for a free electron state in figure 11.12(b).

Both situations depicted in figure 11.12 are submerged within a saturation of multiple solution possibilities for $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ such that the empirically observed

Figure 11.12: Exchanges between the electron field, described in terms of the components of the fermion field $\psi(x)$, and the electromagnetic gauge field $A^{\mu}(x)$, for (a) an interaction with experimental equipment via an external photon and (b) a self-energy contribution for a free electron in terms of an internal photon.

electron state emerges out of this myriad of interactions as an apparently robust and reproducible discrete entity. Such a particle entity may be guided and to some degree localised, propagating in a 4-dimensional spacetime expression of the underlying 1-dimensional temporal flow with properties shaped out of the full higher-dimensional form $L(\hat{v}) = 1$. The particle states exhibit probabilistic behaviour, of the form modelled by quantum theory, as inherited from the probabilistic nature of the underlying degenerate set of possible field configurations, as described in the previous section.

For the case of the plane wave electromagnetic field $A^{\mu}(x)$ of equation 11.6 and explicit form of the spacetime geometry was derived in equation 11.12, with $G^{\mu\nu} \sim$ $\kappa \frac{k^{\mu}k^{\nu}}{Vk^0}$ via the coefficient C extracted from equation 11.16. This geometry for the field in a spatial volume V was provisionally associated with a 'photon' of 4-momentum $P^{\mu} = k^{\mu}$ and $k^2 = 0$. In this naive picture the photon propagates as a kind of 'microscopic gravity wave', as suggested by equation 11.12, consisting of purely Ricci curvature as described after figure 11.1 and suggesting a metric $g_{\mu\nu}(x)$ of a form similar to equation 11.13. The 4-momentum carried by such a 'particle state' is naturally 'quantised' in the sense that the parameter $k \in \mathbb{R}^4$ in the $e^{\pm i k \cdot x}$ Fourier modes appears in the expression $T^{\mu\nu} := -\frac{1}{\kappa} G^{\mu\nu} = f(A)$, that is equation 11.12, since $G^{\mu\nu}(x)$ is a function of the spacetime derivatives of the gauge field $A^{\mu}(x)$.

The actual nature of physical particle structure is expected to be rather more elaborate than initially suggested by this picture of plane waves in a given volume, which was initially motivated in part by analogy with QFT as recapped at the end of the previous section. For the present theory, unlike the external states the intermediate states of field redescriptions, over which the form of the local spacetime geometry $G^{\mu\nu}(x)$ is unchanged, may however indeed involve independent complex wave components. As described in section 11.2 a hybrid set of $e^{-ik\cdot x}$ and $e^{+ik\cdot x}$ mode field exchanges in such interactions, as depicted in figure 11.7, correlates with the apparent 'amplitude squared' rule for the associated interaction probabilities. On the other hand it is in the nature of a 'particle' to possess properties quite distinct from plane waves.

Whether or not considered in terms of wave packets or within a volume V a 'particle' here is also not considered to be a 'point-like' entity, but rather a state of fields as a function on M_4 dynamically prescribed through the conditions of $G^{\mu\nu}$ = $f(Y, \hat{v})$ and $L(\hat{v}) = 1$. Higher energy particle transitions may be possible in ever smaller effective volumes V , correlated with higher 4-momentum k , without limit, with an apparent 'size' or structure never observable for the initial and final state

'entities' in processes such as $e^+e^- \rightarrow \mu^+\mu^-$. Indeed such 'particle interactions' are manifestations of field redescriptions which effectively apply throughout a finite volume V simultaneously, as indicated in and described for figure 11.4 for example, without reference to any point-like particle structure at all.

The fact that particle phenomena become apparent for interactions on very short distance scales, relative to macroscopic laboratory equipment, may be due to there being a greater likelihood for field functions to be indistinguishable for small spacetime volumes. On the other hand the idealised case of transition amplitude \mathcal{M}_{fi} calculations in QFT effectively considers plane waves defined in the limit $V \to \infty$, with factors of V cancelling for observable quantities such as cross-sections. In the present theory the role of an apparent volume V with regards to particle interactions, and the discrete 'quantised' nature of particle states and interactions more generally, requires further understanding.

The factor of $\frac{1}{A}$ in the event rate formula of equation 10.7, from the expression for the luminosity in equation 10.2, makes intuitive sense when picturing the incoming beam components as 'bunches of particles'. However, here the question is how a greater intensity of field interactions, apparently corresponding a smaller area A , increases the production probability for final state particles, with the particle concept *itself* deriving from the underlying field interactions. The relation of the macroscopic to the microscopic world through a program resembling renormalisation will be key to addressing these questions.

As described in section 5.2, and reviewed in the opening of the following section, the generalisation from the classical solution $G^{\mu\nu} = f(A)$, closely relating to Kaluza-Klein theory, will modify the macroscopic form of $G^{\mu\nu}(x)$ in way that incorporates the charge density $\sigma(x)$ in the current $J^{\mu} = \sigma u^{\mu}$ of equation 5.40, the material density $\rho(x)$ in $-\frac{1}{\kappa}G^{\mu\nu} = \rho u^{\mu}u^{\nu}$ from equation 5.39 and the structure of matter $T^{\mu\nu} := -\frac{1}{\kappa}G^{\mu\nu}$ more generally.

From this point of view elementary particle states, such as the electrons and muons observed in HEP experiments, can be considered as quantum transitions within the macroscopic world, which is geometrically described by $G^{\mu\nu} = f(Y, \hat{v})$. With gravitation encompassing quantum phenomena this describes an environment one layer outside the traditional approach to QFT for which the particle states are simply given as the initial and final states of particle interactions. We hence return to the conception of the physical world as described section 10.1 for the experiment depicted in figure 10.1 for example. In the meantime, in chapter 10, we have dismantled the QFT cross-section calculation in order to identify a correspondence with the present theory, as summarised in points (1) –7)' of the previous section; with the ultimate aim of reassembling such calculations in light of the present theory and fully accounting for the observed particle phenomena.

Together with the observations of chapters 8 and 9 for the breaking of higherdimensional forms of $L(\hat{\mathbf{v}}) = 1$ we may hope to gain some insight into the reasons for the observed properties of the various particle types without having to merely write them in by hand based on empirical findings. The abstract Fock space of QFT is not required, with creation and annihilation of particles through mutual exchanges now being firmly grounded in the field state of the macroscopic world. Such a state may consist in the physical components of experimental apparatus themselves, which exhibit essentially classical behaviour, providing a framework to make firm calculations and predictions for the properties of the apparent particle transitions recorded.

An electron state is then consistent both with the idea that nested multiple field solutions, generalising from figure 11.12(b), continue indefinitely down on the microscopic level together with the spacetime geometry satisfying $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ and $G^{\mu\nu}{}_{;\mu}=0$ at the macroscopic level, with a form of 'renormalisation' relating the two levels. In the case of the electromagnetic field the massless 'renormalised' field has a close resemblance to the bare field $A^{\mu}(x)$ of equation 11.6, as described in the opening of this section. Further, from the perspective of Kaluza-Klein theory, the external geometry is directly related to the internal gauge fields through $G^{\mu\nu} = f(Y)$ in the form of equation 5.20 for example. On the other hand, in the absence of an expression of the form $G^{\mu\nu} = f(\psi)$, the physical fermion particle states of an 'electron field' for example appear to have a somewhat more distant resemblance to the bare $\psi \subset v_{56}$ subcomponents with which they were originally identified through the action of $E₇$ on the components of $F(\mathrm{h}_3\mathbb{O})$ broken over TM_4 , as summarised in equation 9.46.

That is, rather than being described directly by the $\psi(x) \subset v_{56} \in F(h_3\mathbb{O})$ field components projected onto M_4 the form of $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ for an e^- particle state observed in HEP experiments will be shaped through interactions with other fields, such as $A^{\mu}(x)$, resulting in a 'renormalised' or 'dressed' state. This was suggested towards the end of section 9.3 where it was also hinted however that fermion particle states might be identified more directly through interactions of a spinor $\psi(x)$ field and the vector-Higgs $v_4(x)$, initially shaping a geometry more simply of the form $G^{\mu\nu} = f(\hat{\mathbf{v}})$, as will be described in section 13.1.

The $\psi \leftrightarrow v_4$ field exchanges between the spinor and vector-Higgs fields, consistent with the constraint $L(\hat{v}) = 1$ of equations 11.29, are also proposed to give rise to the generation of 'mass' for the fermion states. The corresponding interaction terms, as described for equation 9.48, are reminiscent of Yukawa couplings of a fermion field to the Higgs field in equation 7.69 for the Standard Model. However while such a 'bare mass' might be identified at the level of $L(\hat{v}) = 1$, the physical mass, and indeed the concept of mass itself, as an observable quantity is only defined for the macroscopic dressed state as described in terms of the energy-momentum $T^{\mu\nu} := G^{\mu\nu}$. For a free electron state in the complete theory the aim will be to identify the corresponding macroscopic form of $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, and to understand how $k^2 = m_e^2$ arises as a robust observable quantity for such a state, as deriving from the underlying interactions of the 'bare' fields. Further light will be shed on the nature and origin of mass in the presentation of cosmology in the context of the present theory, in particular towards the end of section 13.1 and opening of section 13.2.

As well as carrying energy-momentum density in $T^{\mu\nu} := G^{\mu\nu}$ particle phenomena are observed through the transfer of discrete values of 4-momenta k , ever enveloped within a spacetime geometry and consistent with $G^{\mu\nu}_{;\mu} = 0$, such that the total initial and total final momenta match in processes such as $e^+e^- \rightarrow \mu^+\mu^-$. A full understanding of the nature of such interactions, as provisionally described in sections 11.1 and 11.2, is of course intimately related to an understanding of the nature of the particle concept itself. This may require a full exploration of the relation between the present theory, quantum field theory and condensed matter physics as alluded to in this section.

In general terms to understand what is an electron state or what is a muon state, as observed in HEP processes such as displayed in figure 10.1 or 10.2, it is necessary to think of the full 4-dimensional spacetime picture in relation to the underlying field component redescriptions. This will be described further for figure 11.13 in the following section and connects to the broader question concerning the incorporation of a theory accounting for the quantum properties of fields and particles alongside general relativity in a consistent framework, in the form of the theory presented in this paper. The conception of particle phenomena for the present theory will also be discussed further alongside figure 15.2 in section 15.2 of the concluding chapter, with particle states correlated with the emergence of discrete topologies for geometric solutions for $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ in the near vacuum limit.

11.4 Quantum Phenomena and Gravitation Unified

In the present theory we begin with a 1-dimensional temporal progression and hence need to *build* a 4-dimensional spacetime M_4 with geometry $G^{\mu\nu}(x)$ out of the structure and symmetries of the underlying multi-dimensional form of temporal flow $L(\hat{v}) = 1$. The degeneracy of possible solutions for the ways in which this may be achieved results in the indeterminacy of empirical observations in our world and other apparent 'quantum' phenomena, as studied for example in HEP experiments.

Beginning with the electromagnetic field $A^{\mu}(x)$ in section 11.1 the possibility of alternative solutions involving the fermion field $\psi(x)$ underlying the spacetime geometry were expressed in terms of the field redescriptions of equations 11.22 and 11.23. These equations satisfy equation 11.26 in which the current $j^{\mu} = \overline{\psi} \gamma^{\mu} \psi$ may be considered as a source term. Such 'microscopic' field redescriptions via the mutual exchanges $A^{\mu} \leftrightarrow j^{\mu}$ are incorporated into the spacetime geometry, generalising from the classical relation $G^{\mu\nu} = f(A)$ of equation 11.1 as originally derived through association with Kaluza-Klein theory in section 5.1.

The observable world is awash with the interchanges between the $A^{\mu}(x)$ and $\psi(x)$ fields, together with higher-order redescriptions through which the fields may interact, saturating the world, as described alongside figure 11.12 in the previous section. This gives rise to a rather fluid mathematical creation of matter as perceived through these exchanges and hence the properties and forms of the 'macroscopic' material world are conditioned by them. This describes the general relativistic limit pertaining to tangible physical objects that take shape on M_4 over the collective contribution of the internal fields, such that the apparent composition of the Einstein tensor may be written simply as:

$$
G^{\mu\nu} = -\kappa T^{\mu\nu}(Y, \hat{\boldsymbol{v}}) \tag{11.49}
$$

This is equation 5.32 of section 5.2 expressed in a form which emphasises the implicit field composition of the material world. The effective energy-momentum tensor $T^{\mu\nu}$ on M⁴ may take different forms in terms of the apparent macroscopic matter distribution on the manifold, but it must always be fundamentally composed out of the interplay of the underlying fields, mutually subject to the constraint equations 11.29. For example the energy-momentum tensor might describe a perfect fluid and the Einstein tensor will be macroscopically composed as described in equation 5.37, as we shall consider for the cosmological scales of the universe, alongside equations 12.2 and 12.3, in the following chapter. The general form of that equation for the structure of $G^{\mu\nu}(x)$ incorporates macroscopic 'pressure' $p(x)$ terms and defines the scalar field $\rho(x)$ which in general relativity is identified with the familiar notion of 'matter density'. For the case of a pressureless perfect fluid we have:

$$
-\frac{1}{\kappa}G^{\mu\nu} =: T_{\epsilon}^{\mu\nu} = \rho u^{\mu} u^{\nu}
$$
\n(11.50)

that is equation 5.39, with $T_{\epsilon}^{\mu\nu}$ interpreted as the effective energy-momentum tensor for a pressureless fluid. In the original formulation of general relativity such an energy-momentum tensor, through the above field equation, would be interpreted as the 'source of curvature' on the manifold. This correspondence with general relativity was explored in more detail in section 5.2 where it was described how equation 11.50 leads to the geodesic equation of motion for this form of matter, that is equation 5.36, owing to the Bianchi identity $G^{\mu\nu}_{\mu\nu} = 0$, without the need to introduce the geodesic constraint as an additional postulate of the theory.

The extension of the classical field relation of equation 11.1, which implies the homogeneous Maxwell equation $\Box A^{\mu} = 0$ of equation 11.2, as shown for equation 5.30, with the inclusion of the charged matter term $\rho u^{\mu} u^{\nu}$ on the right-hand side of equation 5.41 is an example of a break away from the pure Kaluza-Klein relation of $G^{\mu\nu} = f(A)$. This deviation from a free electromagnetic field alone is here considered at the level of macroscopic phenomena, which overlays the microscopic field interactions which led to equations 11.21 and 11.26 in section 11.1 and as described in the opening of the previous section. Applying $G^{\mu\nu}{}_{;\mu}=0$ to the full expression in equation 5.41 led to the incorporation of a charged current J^{μ} , with $\Box A^{\mu} =: J^{\mu} = \sigma u^{\mu}$ as defined in equation 5.40, and to the identification of the Lorentz force law of equation 5.43 as a deviation from the purely gravitational geodesic flow. Here the charged current J^{μ} corresponds to that observed in macroscopic classical experiments, typically for the non-relativistic limit such as performed by Faraday in the $19th$ century. Hence in addition to the apparent matter density ρ in equation 11.50 effective macroscopic phenomena also involve the charge density σ in $J^{\mu} = \sigma u^{\mu}$. Both the macroscopic and microscopic currents are conserved, with $J^{\mu}_{;\mu} = 0$ as described following equation 5.43 in section 5.2 also applying for $j^{\mu} = \overline{\psi} \gamma^{\mu} \psi$ of equation 11.26 (as originally expressed for equation 3.101 in section 3.5 for the Lagrangian approach).

Within this limiting case of general relativity described above, that is neglecting explicit quantum phenomena, if the approximation of a flat spacetime for which $G^{\mu\nu}(x) \simeq 0$ may be assumed then the theory of special relativity will apply to the laws of physics. Further beyond that limit the motion of bodies for physical systems in which the relative velocities are small compared with the speed of light may be analysed using classical Newtonian mechanics. Local energy-momentum conservation in all physical processes is ensured under the Bianchi identity $G^{\mu\nu}{}_{;\mu}=0$, since the energymomentum tensor $T^{\mu\nu}$ is identified with the spacetime geometry $G^{\mu\nu}$, regardless of the magnitude of the spacetime curvature. As described in the opening of section 5.2 this observation applies in particular in approaching the flat spacetime limit with $T^{\mu\nu}_{\mu\mu} = 0$, and will also apply for the further limit of the non-relativistic case; with the corresponding energy-momentum conservation also encompassing all underlying quantum phenomena in all cases.

Microscopic transitions of internal fields such as $A^{\mu} \leftrightarrow \overline{\psi} \gamma^{\mu} \psi$, and quantum processes in general, may be recorded in macroscopic devices, generally in the form of amplified electronic signals. All such macroscopic equipment is also itself composed over field interactions in the form of equation 11.49 and effectively described by an appropriate classical energy-momentum tensor $T^{\mu\nu}$, for the solid state devices typically employed, and at a basic level a tiny 'detector recoil' will accompany any production or detection of particle states as a consequence of 4-momentum conservation. The equations governing the evolution and interactions of the microscopic world hence merge into the equations of motion for 'classical' objects, such as described by geodesic trajectories or the Lorentz force law. This framework will then shed some light on a key question concerning the relation of quantum mechanics to the world of classical physics.

In the previous section it has been outlined how empirically observed particle states, such as electrons and muons, might be identified in parallel with a program of 'renormalisation' for the present theory, and merge seamlessly into the state of the macroscopic environment. In figure 11.13 a typical high energy physics process, as described in section 10.2 and already depicted in figure 10.2, is contrasted with a typical experiment involving non-relativistic quantum theory.

Figure 11.13: (a) The process $e^+e^- \rightarrow \mu^+\mu^-$ as observed in HEP experiments for which the cross-section can be calculated in QFT. (b) The double-slit experiment in which a single electron is detected on the screen according to a probability distribution determined in non-relativistic quantum mechanics.

In figure 11.13(a) a particular event is detected, a final state $\mu^+\mu^-$ pair at an angle θ , presumably mediated by one of many possible intermediate sequences of field states such as represented in figure 11.7. In figure 11.13(b) an electron is detected at a particular location A out of a continuum of possibilities including $B, C \dots$ In standard quantum theory both of these processes are assumed to take place against a flat background of space and time, which for figure $11.13(a)$ is Minkowskian and for figure 11.13(b) is Newtonian. In the present theory however the base manifold curvature, although smooth, is finite and non-flat essentially everywhere in 4-dimensional spacetime, with both processes depicted in figure 11.13 representing particular features of a global $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}}) \neq 0$ solution.

Events of the kind sketched in figure 11.13(a) are readily observed by experiments of the kind depicted in figure 10.1 for example. In this case both the macroscopic

SLD detector and microscopic $e^+e^- \rightarrow \mu^+\mu^-$ interaction are uniformly enveloped within a particular solution for $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. The interaction region of such an experiment for such an event will locally have a spacetime geometry $G^{\mu\nu}$ of a similar form to that for $T^{\mu\nu} := G^{\mu\nu}$ pictured in figure 11.1 and as represented by the wavy lines in figure $11.13(a)$. As described in section 11.1 the associated metric solution $g_{\mu\nu}(x)$ will have properties closely relating to the metric of equation 11.13; and the underlying field redescriptions, as represented for example by figure 11.7 in section 11.2, will necessarily respect this external physical geometric form.

Similarly the system described in figure 11.13(b) will be enveloped within a particular $G^{\mu\nu} = f(Y, \hat{v})$ solution, with a non-flat metric $g_{\mu\nu}(x)$ description. For a sufficiently high intensity source with a stable interference pattern observed on the screen a wave-like solution for $G^{\mu\nu}(x)$ will permeate the spaces between the elements of apparatus. The lower intensity case, with a single electron detected on the screen as indicated in the figure, will correspond to a different 4-dimensional world solution for $G^{\mu\nu}(x)$. While both wave-like and particle-like solutions are shaped by an enveloping geometry with $G^{\mu\nu}{}_{;\mu}=0$ uniformly throughout space and time, the underlying indeterministic character of the field redescriptions become evident as discrete particle phenomena emerge at low intensity. In all cases the direct identification of $-\kappa T^{\mu\nu} := G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ implies that the field equation of general relativity is faithfully reproduced, even for the case of a single particle state exhibiting the underlying quantum behaviour.

While a solution $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ envelopes the full 4-dimensional system depicted in figure 11.13(b), including the macroscopic apparatus, the indeterminacy of the single particle process lies in the perfect symmetry of possible field solutions underlying the smooth function $G^{\mu\nu}(x)$ locally at the source S, which is the same for any possible outcome. This situation is then very similar to that in figure 11.13(a), with the source S corresponding to the interaction region, as represented by the rectangular box, and with the range of outcomes $A, B, C \ldots$ corresponding to the angular range $0 < \theta < \pi$. The comparison is even more direct if the intermediate double-slit screen is removed from the apparatus in figure 11.13(b).

In all cases the full 4-dimensional solution $G^{\mu\nu} = f(Y, \hat{v})$ encompassing the entire system is intrinsically shaped through a 1-dimensional causal accumulation of probabilistic outcomes wherever the geometry $G^{\mu\nu}(x)$ is locally expressible in terms of a degeneracy of underlying field functions. The inclusion of the double-slit screen in figure 11.13(b) is accompanied by a more complicated spectrum of single particle solutions, as might be expected since the full system is more complicated. In this case the relative probabilities, while depending crucially on the underlying field degeneracy at S, turns out to be weighted by the interference pattern as shown.

The spacetime curvature itself is far too small to be directly detectable, for example by geodesic deviation, although the fact that $G^{\mu\nu}(x)$ is non-zero in these laboratory experiments is crucial in the present theory. As described above the local spacetime curvature associated with the interaction region in figure $11.13(a)$ will be analogous to that for the free electromagnetic wave as derived in equation 11.12 and pictured in figure 11.1. This curvature will naturally be higher in cases of higher energy density such as at the interaction region of the LHC, where it remains also far too small to be observable.

With $T^{\mu\nu} := G^{\mu\nu}$ the spacetime curvature is also indirectly made apparent through the presence of energy-momentum. For example, since energy-momentum is everywhere conserved in line with the identity $G^{\mu\nu}{}_{;\mu} = 0$, a small recoil of the electron source S in figure 11.13(b) will causally precede the detection of an electron at A. Indeed, in principle 'elementary' particles might be observed with detectors in a way analogous to 'Brownian motion' with macroscopic matter 'recoiling' against the elementary transitions of the fields within which it is immersed, bringing out the properties of both the particles and material objects themselves.

Considering a thought experiment with a very lightweight source S situated at a very long distance from the detector screen in figure 11.13(b) in principle an observation of the momentum recoil of S could precede the detection of the signal at A (and for apparatus consisting of the source and screen alone the prediction of the hit location on the screen would be very direct). The total momentum of the system, including the source, double-slit screen and detection screen, will be conserved. The same quantum interference pattern would still appear on the screen given a large number of such events.

With the momentum recoil of the macroscopic source S too small to be measurable in practice for the process depicted in figure 11.13(b) the first and only sign of the event will be through the amplification of an electronic signal at A. Pragmatically the possible observable outcomes can be represented in terms of an electron wavefunction $\Psi(x)$ evolving according to Schrödinger's equation until collapsing to zero at $B, C \ldots$ at the moment when the electron is detected at A . In the present theory such a description in terms of an apparently non-local action of wavefunction collapse represents our knowledge of the state of the system rather than its underlying physical evolution.

In standard non-relativistic quantum mechanics the basic principle of the conservation of energy and momentum is considered to hold together with the constraint that no signals may be transmitted faster than light. For the case of the experiment depicted in figure 11.13(b) this leads to the question concerning the location of the 'conserved energy' during the intermediate period between the emission of a particle of a given energy at S and the later detection of a particle of the same energy at A. The corresponding energy-momentum cannot be carried by the wavefunction for example, due to the discontinuous nature of the wavefunction collapse.

In the present theory the 'energy-momentum' is distributed throughout in terms of the 4-dimensional geometry $T^{\mu\nu} := G^{\mu\nu}$. Energy-momentum conservation is everywhere implied in the identity $G^{\mu\nu}_{\mu\nu} = 0$, with nothing being transmitted faster than the speed of light – as defined by the light cone structure which arises through the projection of the full form $L(\hat{v}) = 1$ onto the manifold M_4 as described in section 5.3. Since a solution $G^{\mu\nu}(x) = f(Y, \hat{v})$ primarily describes the shape of a particular spacetime geometry it may have a highly counter-intuitive distribution when *interpreted* through $T^{\mu\nu} := G^{\mu\nu}$ as an apparent flow of '*matter'* through space. Some forms of geometry do possess a form with a natural interpretation in terms of energy-momentum, as expressed for the macroscopic example in equation 11.50 for a pressureless fluid. However, more generally rather more arbitrary geometries are permitted, provided $G^{\mu\nu}{}_{;\mu} = 0$, and some form of continuous geometry $G^{\mu\nu}(x)$ will be associated with the single particle process depicted in figure 11.13(b).

For the present theory the question concerns the manner in which everywhere

continuous solutions for a geometry $G^{\mu\nu}(x)$ can be apparently channelled in certain discrete and localised ways which give the impression of 'particle' transitions. That is, locally the energy-momentum $T^{\mu\nu} := G^{\mu\nu}$ can be interpreted as the emission or detection of a discrete particle, for example at S or A respectively in figure 11.13(b). The answer presumably lies in the nature of the *fixed* constraint equations 11.29, such as $D_{\mu}L(\hat{v})=0$, which channel the underlying field redescriptions in a limited number of ways and in turn determine the properties of the particle transitions which emerge, as alluded to in the previous section.

All empirical phenomena, whether naturally occurring or constructed in physical experiments such as that in figure 10.1, will be enveloped under a geometry $G^{\mu\nu}(x) = f(Y, \hat{v})$. Since in the laboratory setting this non-trivial geometry, that is any deviation from Minkowski spacetime, is completely unobservable the consequences of this perspective may be pursued by considering more extreme cases, such as the thought experiment described earlier for figure 11.13(b) with a very lightweight source S far removed from the detector screen, as well as by analysing phenomena physically realised in practice. Any beam of electromagnetic radiation carries energy and is hence associated with spacetime curvature as for the case of standard general relativity and as depicted in figure 11.1 for example. In a further thought experiment intense beams of light, for example produced by lasers, could in principle be configured such that the tiny geodesic deviation of a suitable test body projected through the curved spacetime associated with the laser beam and over a large distance might be observed, without any photons of the beam being detected.

This situation may be contrasted with the empirical observation of the deflection of light itself in the gravitational field of the sun, as first reported just a few years after the formulation of general relativity. In these cases, for both the above thought and practical experiments, there is an 'interaction' between light and gravity without the detection of any photons or the need to appeal to any properties associated with quantum theory. For the above thought experiment similar observations would hold for an intense beam of particles such as electrons in place of the lasers, and leads to the conclusion that the electron field associated with the event of detecting even single electron in figure 11.13(b) will indeed be accompanied by a small, although utterly undetectable, spacetime curvature.

As described in section 11.2 the spacetime curvature $G^{\mu\nu}(x)$ is always a real valued tensor field but may be constructed out of a hybrid of complex components, such as the $e^{\pm ik\cdot x}$ Fourier modes, of the underlying fields such as $A^{\mu}(x)$ and $\psi(x)$ as depicted in figure 11.7 for example. The matching of both the $e^{-ik \cdot x}$ and $e^{+ik \cdot x}$ parts coming together into a real-valued function for a single field such as $\psi(x)$ may correlate with detection events, such as at A in figure 11.13(b), through which apparently propagating particle states, such as electrons, are revealed. More generally the concept and nature of elementary particles needs to be fully addressed, as was discussed in the previous section and will be further elaborated in section 15.2.

While the understanding of the nature of particles as observed in the laboratory requires further work, it is clear in the present theory that there are no 'graviton' states since the gravitational field itself is *not* quantised in any sense. In fact general relativity provides a classical description of the geometry of the external perceptual framework of the world which fully accounts for the phenomena of gravitation. There is no given flat 4-dimensional spacetime manifold and hence no 'force' of gravity as an apparent empirical addition on top of such a Minkowski spacetime. In turn there is no place for gravitons as 'carriers' of such a gravitational force and no compelling motivation to consider any form of quantisation of gravity.

In the present theory 'quantisation' is a phenomenon that applies to the fields underlying spacetime solutions of the form $G^{\mu\nu} = -\kappa T^{\mu\nu}(Y, \hat{\boldsymbol{v}})$, that is equation 11.49 above. Only the right-hand 'matter' side of this expression is effectively quantised, as a consequence of the degeneracy of field redescriptions, involving interchanges of gauge and fermion fields for example, which underlie the solution. To attempt to impose 'quantisation rules' on the left-hand 'geometry' side of this expression would be to quantise the same object $G^{\mu\nu} \equiv T^{\mu\nu}$ twice in two different ways. The external geometry $G^{\mu\nu}(x)$ itself implicitly incorporates a choice of $T^{\mu\nu}(Y,\hat{\boldsymbol{v}})$ and effectively the identification of this 4-dimensional spacetime geometry is itself the mechanism of quantisation for all non-gravitational fields. That is, the possibility of multiple solutions of the form $G^{\mu\nu} = f(Y, \hat{v})$ underlying the external geometric framework for perception of objects in the world is the reason why the fields implicit in the energymomentum tensor $T^{\mu\nu} := G^{\mu\nu}$ are quantised, with no similar argument applying to the degrees of freedom of the spacetime geometry itself.

Quantum field theory is however formulated against a flat spacetime background and we may also consider the corresponding limit for the present theory. For the respective theories of general relativity and quantum fields the geometry of the spacetime manifold and that of the internal gauge fields are independent constructions. In the present unifying theory the relation between them is identified through a larger, all encompassing, symmetry group \hat{G} for the full general form of the flow of time $L(\hat{\boldsymbol{v}}) = 1$, linking the external and internal forces of nature, as for example seen in equations 5.20 and 11.1. Their distinctive, complementary, features arise in the breaking of the full symmetry group over the base manifold M_4 , as depicted in figure 5.1 for the $L(\mathbf{v}_{10}) = 1$ model. Considering the full forms $L(\mathbf{v}_{27}) = 1$ and $L(\mathbf{v}_{56}) = 1$ in turn the surviving local gauge symmetry and resulting field interactions in this theory have been compared with corresponding features of the Standard Model in chapters 8 and 9. The properties of these interactions will be drawn out and made apparent through discrete particle phenomena, which themselves can only be fully explored in the present theory when the associated minute deviations from a flat geometry are fully embraced, as described above and in the previous section.

In a curved spacetime there are generally no preferred choices of Lorentz frames and through the local freedom in $l(x) \in SO^+(1,3)$ general relativity can be interpreted as having some relation to gauge theory, as alluded to towards the end of section 3.4. However, in the limit of a flat linear connection, with $\Gamma(x) \to 0$ in a Minkowski coordinate system, it is also meaningful to define a global external gauge, that is a global basis for components of a tangent vector field in TM_4 , with a single choice of $l \in SO^+(1,3)$. While the equations of physics are gauge covariant we expect them to take a particularly simple form when such a natural global gauge is possible. This is the case in special relativity and also in Newtonian physics for which a Galilean reference frame is typically preferred. In the limit of a flat spacetime the freedom of local symmetry in $l(x) \in SO^+(1,3)$ has effectively been *broken* to the much more restricted freedom of a global symmetry on M4.

In fact, in the spirit of the present theory as introduced in section 2.2, the requirement of perception implies that the local Lorentz symmetry freedom of the local reference frames as a function of $x \in M_4$ acts globally over macroscopic scales to a good approximation and hence is essentially broken down from a local to a merely global symmetry, and hence with far fewer degrees of freedom. This is the reverse of the usual case seen in gauge field theories in which a global symmetry is generalised to become a local symmetry leading to the interactions described in the Standard Model of particle physics for example.

It is this assumption of what is essentially a hole in the full symmetry \ddot{G} of $L(\hat{\mathbf{v}}) = 1$ carved out by the global $SO^+(1,3)$ symmetry on M_4 that allows the deployment of a global Minkowski coordinate frame on the base manifold that in turn allows the expansion of each field as a sum over the linearly independent functions of a Fourier series on the base manifold, as described for the electromagnetic field for example in equations 11.3–11.6. The question concerning the relation between the interactions of such fields in the present theory in this limit and calculations performed in quantum theory has been considered in the previous sections of this chapter and is further elaborated in the following.

Generally in physics there are numerous examples in which observable quantities parametrised by real numbers are analysed through expressions involving the algebra of complex numbers. To take a simple example an oscillating quantity such as the electric current in a wire of the form $I = I_0 \cos \omega t \in \mathbb{R}$ can be expressed as $I = I_0 \text{Re}(e^{i\omega t}) \in \mathbb{R}$. The straightforward mathematical properties of objects such as $e^{i\omega t} \in \mathbb{C}$, under multiplication and differentiation for example, may then be exploited in calculations before the underlying physically real (in the sense of 'existing') part is extracted in terms of the mathematically real (in the sense of \mathbb{R}) part at the end of the calculation.

In quantum theory complex analysis is used directly from the foundations. Via either canonical quantisation or the path integral approach as the starting point for QFT, Feynman rules and the complex transition amplitude \mathcal{M}_{fi} may be constructed on the way to extracting real-valued cross-sections or decay rates at the end of a calculation. Similarly the postulates of non-relativistic quantum mechanics are couched in terms of complex mathematical objects from the beginning – with a complex wavefunction $\Psi(x)$ or state vector in a Hilbert space completely defining the dynamical state of a quantum system and empirical predictions obtained in terms of the real eigenvalues of Hermitian operators.

These structures for quantum theory appear quite distinct from other applications of complex analysis in physics which, as for the example of the electric current $I = I_0 \text{Re}(e^{i\omega t})$ above, *begin* with real-valued quantities. In this sense, by comparison, quantum theory appears to hang in the air, apparently lacking a more tangible conceptual foundation. The present theory aims to supply such an underlying physical basis for quantum theory in terms of the relative frequency of possible solutions for fabricating the 4-dimensional spacetime M_4 itself, with the geometry $G^{\mu\nu}(x) = f(Y, \hat{\boldsymbol{v}})$, as provisionally expressed in terms of the probability $P \propto D_+ D_-$ of equation 11.46. Linked to the QFT probability $|\mathcal{M}_{fi}|^2$ via the structure of the real-valued quantity $\text{Im}(\mathcal{M}_{ii})$ and the optical theorem, as discussed for figure 11.8 and summarised in points '1)–7)' of section 11.2, the aim is to build the theory up from beneath QFT through a complexification of the underlying probability computation, which is based on the degeneracy of solutions, on the way adopting some the mathematical machinery of QFT itself.

Historically QFT was developed in the late 1920s on the coat-tails of the original quantum mechanics by promoting the wavefunction to an operator field (which was sometimes called 'second quantisation', although there is still only *one* quantisation). By Fourier analysing the vector potential $A^{\mu}(x)$, as a free-field solution of Maxwell's equations, into normal modes and applying a quantum mechanical harmonic oscillator treatment to each mode independently photons, as quanta of the electromagnetic field, were the first 'particle states' to be studied in a QFT. Since in the present theory we began by making contact with QFT in the environment of HEP experiments the connection in the other direction, with quantum mechanics arising as a limit of QFT, should also be considered. For example, the retarded propagator or Green's function of QFT, which enters the present theory as described for equation 11.22, can be taken to the non-relativistic limit in which it is found to be identical to the transition amplitude for single particle transitions in quantum mechanics. (The possibility of such a connection can be inferred from the relation of the function $\Delta_R(x-y)$ to $\Delta^+(x-y)$ through equations 10.75 and 10.79 and the structure of $\Delta^+(x-y)$ in equations 10.58 and 10.60 in comparison with equation 10.8 as discussed after equation 10.63).

While the path integral approach has not been found useful for establishing the link between the present theory and QFT, as alluded to before equation 10.46 in section 10.3, the relationship between QFT and QM is perhaps most readily seen in terms of the path integral approach for which the same basic postulates apply in both cases. The transition amplitude K is treated as a fundamental object and identified with the sum over 'all possible paths' of a phase factor $e^{iA/\hbar}$, where A is the classical action associated with the path (see for example [10] chapter 8). The mathematical properties of this phase factor are exploited in the structure of the theory, with the transition probability postulated to take the form $P = |K|^2$ such that the basic law of probability conservation is upheld. Both QFT, with a mathematical formalism for generating expressions associated with Feynman diagrams through higher-order functional derivatives, and the single particle theory of QM for the non-relativistic case, which can be generalised with higher-order Green's functions to describe multiparticle systems, may be derived from the path integral approach. The two cases of spontaneous symmetry breaking, in condensed matter physics and the Higgs sector of the Standard Model, alluded to in the previous section may also be described in very similar ways mathematically using the path integral approach, which is otherwise here seen as a useful formal method of performing calculations rather than relating to the conceptual basis for the present theory.

In the present theory objects such as the Schrödinger wavefunction $\Psi(x)$ in QM and transition amplitudes, such as \mathcal{M}_{fi} for QFT, are also considered as mathematical constructions for pragmatic use in the relevant calculations of real observable quantities such as event probabilities. As complex representations none of these mathematical objects directly represent physical entities such as fields or particle states, although complex Fourier modes of the fields have been employed in the degeneracy count as represented for example in figure 11.7.

For the present theory the fundamental objects in spacetime are the real-valued

fields directly drawn out from the components of $L(\hat{\mathbf{v}}) = 1$, and its symmetry actions, over the base manifold M_4 , as described in the opening of this section. The energymomentum possessed by such fields is strictly defined through $T^{\mu\nu} := G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, as described throughout this chapter, in terms of the 4-dimensional spacetime geometry description via the Einstein tensor effectively composed of the underlying fields. Field interactions and transitions follow from the degeneracy of possible solutions. This definition of energy-momentum in $T^{\mu\nu}$ is independent of the field content, applying to the quantum as well as classical physics case, with 4-momentum conservation corresponding in all cases to the identity $G^{\mu\nu}{}_{;\mu} = 0$.

While in the present theory we begin with the form $L(\hat{\mathbf{v}}) = 1$ and then identify the spacetime geometry over an extended manifold M_4 , in QFT the starting point is a flat spacetime manifold itself. From this point of view in standard quantum theory the presence of energy-momentum $T^{\mu\nu} \neq 0$ alongside the flat spacetime assumption $G^{\mu\nu} = 0$ not only directly contradicts the central field equation of general relativity but also evades any possibility of a unifying theory of quantum mechanics with gravitation. That is, since in QFT a flat 4-dimensional spacetime is a given background arena the conceptual origin of indeterminate quantum phenomena as a degeneracy of solutions for the underlying spacetime structure itself is entirely missed.

Hence in quantum theory wavefunctions and amplitudes are introduced at the outset and unitary symmetry imposed in order to model the probabilities of such phenomena. This approach dates back to matrix mechanics, presented by Heisenberg in 1925, in resorting to a mathematical framework aimed at coherently linking observable phenomena without an underlying conceptual and physical motivation as the basis. On the one hand the present theory provides such an underlying basis for quantum phenomena in terms of a degeneracy of field solutions for the spacetime geometry, and on the other hand it should also be able to account for the original quantum mechanics of Heisenberg and Schrödinger in the non-relativistic limit.

Although the curvature of the spacetime geometry $G^{\mu\nu} = -\kappa T^{\mu\nu} \neq 0$ is unobservably small by many orders of magnitude on the scale of atomic or HEP phenomena the standard equations of quantum mechanics and QFT, in assuming a flat spacetime background, do depend on the existence of a smooth continuum of spacetime points $x \in M_4$ with the structure of a global Minkowski metric η_{ab} on the manifold, since this is required to give meaning to the location of wavefunction or operator field values in these theories. In the present theory this continuum takes the full metric form $g_{\mu\nu}(x)$ of general relativity, which is determined by the fields themselves, describing a spacetime which is only approximately flat.

In the mathematical formalism of QFT the points x of an independent flat spacetime background are mapped into operators $x \to \hat{\phi}(x)$ which are defined by their action on the states of the system. Quantisation rules are imposed on the dynamical degrees of freedom of the field $\phi(x)$ itself, as described for equations 10.13 and 10.16 in section 10.3, while the spacetime location x is simply a parametrisation for the field in terms of a set of real number coordinates $\{x\}$. In non-relativistic quantum mechanics the operator \hat{x}^a , appearing in the discussion below equation 10.78, represents the spatial location of a particle state, while there is no operator corresponding to time. However in QFT, which is invariant under the transformations of special relativity, there are no operators corresponding to either time or space. Since these quantities are clearly 'observables' quantum theory as it stands is not a universal theory, rather extended spacetime provides an 'external' background arena for QFT, as it does for classical mechanics in the non-relativistic limit.

General relativity is the theory of external space and time and is itself not a theory standing in need of quantisation, either on empirical or necessary theoretical grounds. While some approaches to 'quantum gravity' seek to include gravitation and spacetime geometry *within* the framework of an extended quantum theory, here in the present theory the phenomena of quantisation arise beneath the surveillance of gravitation, with the geometric degrees of freedom associated with general relativity hence outside the domain of quantum theory. As a consequence, for example, there are no gravitons for this theory, as discussed earlier in this section.

The concept of time plays a central role both in relativity theory and in quantum mechanics. In general relativity the proper time, with the interval $d\tau$ of equation 5.48 for infinitesimal displacements, can be used to parametrise a series of events on the manifold, such as those that map out the spacetime trajectory of a physical object with 4-velocity flow $u^{\mu} = dx^{\mu}/d\tau$ as described in section 5.3. In quantum mechanics the temporal evolution of a state is determined by the Hamiltonian operator H (as introduced before equation 10.27), which also describes the energy of the state, with the wavefunction $\Psi(t)$ for example in the time-dependent Schrödinger equation satisfying:

$$
i\frac{\partial}{\partial t}\Psi = H\Psi \tag{11.51}
$$

(as exemplified in equation 10.28 for the evolution of the state vector in QFT, and alluded to after equation 10.52 with $H \equiv \hat{E}$ in quantum mechanics). One of the main difficulties with background dependent approaches to quantum gravity that apply the superposition principle of quantum theory to spacetime geometries, or make quantum transitions from one to another, is that, owing to the principle of general covariance in general relativity, there is no well defined way to map points in one spacetime to those of another. Labelling the points with coordinates does not help since under general covariance coordinates are of no physical significance, as described in section 3.4. This, in particular, means that there is no unique way to specify a map from a temporal derivative on one spacetime manifold to a temporal derivative on another. Hence the temporal evolution of a quantum state in equation 11.51 cannot be transferred in any meaningful way between different spacetimes. This absence of a well defined independent temporal parameter is known as the 'problem of time' in quantum gravity.

It is a problem which does not arise within the present theory since here gravity itself is not quantised and there is no 'superposition of spacetimes'. As for classical general relativity, here the emphasis is on complete four-dimensional solutions for the spacetime geometry satisfying $G^{\mu\nu} = f(Y, \hat{\mathbf{v}})$, with indeterminacy and the probabilistic nature of quantum phenomena inherent fundamentally in the degeneracy of many possible field solutions which underlie the world geometry. Hence there is no difficulty in identifying a universal one-dimensional time parameter (such as the proper time in the local frame of any given observer) and the 'problem of time' does not arise here, as it does for theories which place the temporal evolution as conceived in a quantum theory at the forefront. Rather here quantum effects arise underneath gravity, and can be consistently parametrised in terms of coordinates on the single spacetime background of perception, with the ready availability of unambiguous temporal derivatives.

The structure of quantum theory is hence fused within the structure of general relativity, with 4-dimensional spacetime infused throughout with a 1-dimensional causal progression in time as employed for example in equation 11.51.

Geometric structures, including the causal structure of spacetime, are described by degrees of freedom expressed in the metric $g_{\mu\nu}(x)$ and tetrad $e^{\mu}_{a}(x)$ fields, consistent with the Riemann tensor $R^{\rho}_{\sigma\mu\nu}(x)$, on a single spacetime manifold M_4 . Indeed, this manifold is itself constructed out of the more fundamental underlying notion of progression in temporal flow s as expressed through $L(\hat{v}) = 1$. The projection onto the manifold results in relative time dilation phenomena as described in section 5.3. In the 4-dimensional continuum each observer carries a clock which provides a time parameter which may be applied in quantum experiments in the laboratory or for observations in general within the spacetime arena; with temporal parameters for mutual observers simply related by relativistic transformations.

The probabilistic nature in terms of what can happen as the outcome of a laboratory experiment involving quantum phenomena motivates the construction of the quantum state or wavefunction locally parametrised through a 1-dimensional progression in time according to equation 11.51 for each local observer and the employment of the associated quantum theoretical tools.

In the present theory we begin with *real* fields such as $A^{\mu}(x)$ of equations 11.3– 11.6, with a complex decomposition into parts such as $e^{-ik \cdot x}$ which seem to resemble a complex wavefunction $\Psi(x, t)$. Indeed, similarly as applied to the positive frequency modes of the quantum field component $\hat{\phi}^+(x)$ as described after equations 10.51 and 10.52, the Hamiltonian operator H of equation 11.51 can be applied to the complex component of the classical electromagnetic field in equation 11.17 resulting in:

$$
HA^{\mu}(x) = k^{0}A^{\mu}(x) \tag{11.52}
$$

with eigenvalue k^0 . This is identical to the energy of the normalised electromagnetic field within the volume V as described following equation 11.16, which was extracted through the relations $-\kappa T^{\mu\nu} := G^{\mu\nu} = f(A)$. Equation 11.52 exemplifies how the 'operator plus wavefunction' description can offer a concise way to extract properties such as the 4-momentum from the field as a useful tool for calculations. Although the real-valued field $A^{\mu}(x)$ of equation 11.6 precisely describes the actual field, if it may be reconstructed uniquely as the 'realification' of a complex component such as equation 11.17 (that is, by adding that equation to its complex conjugate) then the latter in principle carries all the information concerning the real physical field, and also satisfies the same equation of motion as the real field as described after equation 11.20.

However, in quantum mechanics the wavefunction $\Psi(x)$ represents a single observable particle, applying to a physical electron state in figure 11.13(b) for example, and it remains to be seen explicitly how such particle states may be described in terms of underlying fields and their interactions in the present theory. Hence an understanding of field renormalisation and the nature of observable particles, as discussed in the previous section, will need to be further developed in order to establish the full connection between the mathematical objects of the present theory and the pragmatic devices of quantum theory.

While representing a single particle a wavefunction $\Psi(x, t)$ is in general a continuous function of the spatial coordinates with intrinsically non-local properties in terms of the temporal evolution as a measurement is made and therefore exhibits a non-particle-like structure itself. In a measurement of position the quantum particle is observed to be in a particular spatial location which is determined by the wavefunction, to the extent that the squared modulus of this complex function $|\Psi(x, t)|^2$ determines the probability to detect the particle at that location, as alluded to after equation 10.8, with the wavefunction immediately 'collapsing' to that measured point. While carrying information concerning various physical quantities, when combined with the appropriate quantum mechanical operator, the wavefunction $\Psi(\boldsymbol{x})$, unlike the field $A^{\mu}(x)$ does not itself *carry* energy or any other physical attribute and hence there is no *physical* discontinuity for the experiment of figure 11.13(b) when the electron is observed at a particular location. Rather, in the interpretation of the present theory, the energy is contained in components of $T^{\mu\nu} := G^{\mu\nu}$ which is continuous everywhere in these experiments, as described earlier in this section.

From this point of view the quantum mechanical wavefunction reflects our best knowledge of the range of world solutions our current empirical situation is consistent with; it evolves in a determined way U through passage of laboratory time, as governed by equation 11.51 for a given Hamiltonian operator, in such a way as to yield probabilistic predictions for which particular solution state we shall find ourselves observing at the time of the next measurement (see for example [26] p.592). Since the wavefunction is a non-physical entity, the so-called 'collapse' or 'reduction' R of the wavefunction merely represents the change in our knowledge when such an observation is made, and is not itself a constituent property of the physical world.

Hence the apparent conceptual difficulties concerning 'wavefunction collapse' are a somewhat artificial feature of quantum mechanics since the change in evolution law from the unitary U for the wavefunction Ψ , describing a superposition of states, to reduction R selecting an eigenstate Ψ_i in the measurement, is just a pragmatic device for calculation (similarly as for the employment of the transition amplitude \mathcal{M}_{fi} in QFT) and does not directly describe the behaviour of a physical entity, such as represented by the gauge field $A^{\mu}(x)$ for example.

From the perspective of the subjective laboratory view with a sequence of events seemingly evolving in time upon a given 4-dimensional background manifold some quantum phenomena appear mysterious, such as the 'spooky action at a distance' as predicted and observed for Einstein-Podolski-Rosen (EPR) experiments. Such experiments demonstrate that quantum phenomena cannot be accounted for by an underlying theory which is both local and deterministic, as constructed in terms of 'hidden variables' for example. In the present theory however the phenomena of EPR correlations and quantum entanglement in general are all sown into the fabric of the full 4-dimensional spacetime solutions under the geometry $G^{\mu\nu}(x)$. These observations are hence in principle naturally accounted for without any non-local interactions or behaviour and without hidden variables but with indeterminacy fully embraced as a manifestation of the local degeneracy of possible fields, including the gauge field $A^{\mu}(x)$ and fermion field $\psi(x)$, necessarily featuring in solutions for $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. Here the underlying fields such as $A^{\mu}(x)$ are intimately involved in the *construction* of the spacetime geometry, rather than introduced separately as classical waves spreading out over a *pre-existing* M_4 background.

The local causality in the present theory incorporates the restriction that sig-

nals cannot propagate faster than the speed of light, with special relativity holding locally as for general relativity. (In principle a form of the 'equivalence principle', as described in section 3.4, might be adopted, but the employment of a 'torsion-free' external geometry is a simplifying and provisional assumption both for general relativity and for the present theory, as discussed in section 5.3 and also section 13.3). Here 'causality' means of course that the range of probabilities for possible future states, and not the actual future state itself, is determined locally by the present state.

Although the local redescriptions of the fields such as depicted in figures 11.4, 11.6 and 11.7 are arbitrary within the constraints of equations 11.29 the overall theory is 'deterministic' in the sense that all possible worlds, all solutions, potentially exist. On the other hand events in the single solution of our world do necessarily appear indeterministic – 'God does play dice' from the point of view of observations in our universe.

In the case of Schrödinger's famous thought experiment the outcome can only be to perceive an alive or a dead cat ([26] p.808), while an entity described by the quantum state '|alive \rangle + \langle dead \rangle ' cannot be observed. The present framework incorporates a theory of perception through which each of the two possible macroscopic states corresponds to a separate $G^{\mu\nu}(x)$ world, each necessarily observed subject to $G^{\mu\nu}$ $\mathcal{L}'_{;\mu}(x) = 0$ and constructed out of the flow of time with $L(\hat{\boldsymbol{v}}) = 1$, such that we cannot perceive both large scale states simultaneously since they describe different worlds. The more practical experiments with an electron being detected at $A, B, C \dots$ in figure 11.13(b), or the muon detected at an angle θ figure 11.13(a), are associated with a spectrum of different worlds.

Whatever the relative probability of the two alternative outcomes as determined by the apparatus of a 'Schrödinger's cat' type experiment, it is possible to consider two sets of worlds each of which consists of a 'coarse-grained' ensemble characterised by one of the two possible outcomes. More generally we inhabit one of a much larger ensemble of possible worlds, each distinguished by the resolution of a vast number of locally indeterministic processes intrinsic to the 4-dimensional world solutions. With the range of worlds resulting from the many ways to construct $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ over a 4-dimensional base manifold each solution, each universe, is as real as ours. (This statement carries the caveat that each universe should support observers, in the sense described in chapter 14).

The availability of 'many solutions' for $G^{\mu\nu}(x)$ in spacetime responsible for the indeterminacy in such experiments is reminiscent of the 'many worlds' interpretation of quantum mechanics. However, here the theory has many solutions by nature, this feature is not an interpretation of the theory. In the many worlds interpretation of quantum mechanics the wavefunction is taken literally as a real entity with the above observations of both 'an alive and a dead cat' effectively interpreted as a bifurcation of our world as one of many such divisions in a 'branching universe'. In the present theory the other worlds might be thought of existing 'out there' in a realm of possible mathematical solutions, unlike the more intimate picture of the many worlds interpretation.

Here there is also no essential observer participation in 'wavefunction collapse' in the sense of the 'many minds' interpretation of quantum mechanics, rather the wavefunction, as a non-physical entity, is our own pragmatic construction employed to predict the likelihood of future events. On the other hand in the present theory the observer does have an innate role in shaping the overall theory through the subjective nature of perception on the base manifold, which implies the breaking the full $L(\hat{\mathbf{v}}) = 1$ symmetry and the ensuing physical structures. This perspective is influenced by the Kantian philosophy concerning the a priori nature of perception in the form of space, time and causality, as will be further elaborated in chapter 14 and in particular in the opening paragraphs of section 14.2.

During the early history of quantum mechanics the meaning of the formalism in terms of the 'Copenhagen interpretation', was a natural, pragmatic and provisional way of addressing the conceptual difficulties raised. This also marked a relatively conservative break away from the world of classical mechanics, combining the quantum with the classical aspects of the world in a way that upheld the classical behaviour of experimental apparatus and the classical notion that physics exclusively studies the properties of a single universe, although now, however, one with an intrinsic element of uncertainty. While the postulates and mathematical structure of quantum theory has remained essentially intact and unchanged since the 1920s, the debate over the interpretation of the theory continues into the $21st$ century.

The main difficulty with the Copenhagen interpretation is the 'measurement problem' concerning the grey area of interface between classical apparatus and the quantum system under investigation and the nature of the apparent 'wavefunction collapse'. This issue is highlighted by the 'Schrödinger's cat' thought experiment and helped motivate the later many worlds interpretation alluded to above. In the present theory the measurement problem is resolved through the seamless employment of a classical notion of probability, defined in terms of the number of ways an event can happen, all the way down from the macroscopic apparatus to the underlying microscopic field redescriptions. This theory hence unifies the notion of probability for the classical and quantum domains, as applies for example to the experiments depicted in figure 11.13.

As well as having a common underlying origin the meaning of the probability of an outcome for a quantum process (involving for example an experiment in figure 11.13 or the fate of Schrödinger's cat) and for a classical process (such as the roll of a dice or the toss of a coin) is subjectively the same, in terms of for example how we might make choices dependent upon such outcomes. In both the quantum and classical cases the outcome probability is calculated based on our knowledge of the set-up of the system before the experiment is performed. However there is also a significant objective difference in the nature of quantum and classical chance even in the context of the present theory. The difference is that in quantum theory the outcome is fundamentally unknowable in advance, whereas for the classical case the probability merely represents the practical limitations of our knowledge and our ignorance of the precise details of the initial conditions. The actual outcome of such classical experiments would in principle be calculable and fully determined if we could gather sufficient data and muster the necessary computational power (the improving accuracy of weather forecasting with improving technology provides an example). On the other hand, although in the many solutions there are many worlds and essentially everything that *can* happen *does* happen in some universe, quantum phenomena from our perspective in our world are objectively and inherently indeterministic.

For a given observed event, for a process such as $e^+e^- \rightarrow \mu^+\mu^-$ pictured in figure 11.13(a), the question can be asked whether a *particular* sequence of field exchanges actually mediates the process between the initial and final states. In terms of the field sequence $\overline{\psi}\gamma^{\mu}\psi \to \overline{\varphi}\gamma^{\mu}\varphi$ in figure 11.6 for example this corresponds to the question of whether there are particular values for t_4, t_3, t_2 and t_1 , whether the intermediate $\overline{\psi}\gamma^{\mu}\psi$ field state represents a $\mu^+\mu^-$, $d\bar{d}$ or other fermion pair between t_3 and t_2 , and the value of the corresponding unconstrained internal 4-momentum degrees of freedom. In turn there is an endless list of possible field sequences, with field exchanges separated by intervals of time down to $\delta t \to 0$.

These possibilities are not observable, but it is precisely the fact that they signify distinct descriptions of the overall process that contributes to the total probability to observe the event which is statistically measurable. In a similar way that one particular outcome of many possibilities is observed, such as a $\mu^+\mu^-$ or $\tau^+\tau^-$ final state at an angle θ to the incoming e^- beam in figure 11.13(a), from a philosophical point of view it is consistent to think of the internal process as following one particular sequence, such as via a $\mu^+\mu^-$ or $d\bar{d}$ internal fermion state in figure 11.6 for example, with particular values for the continuous degrees of freedom described above. (Although since there is an endless number of infinitely nested possible field redescriptions, as alluded to in the previous section, the idea of singling out 'one' such sequence may be poorly defined). This is again analogous to the classical case in which the outcome of the roll of a dice, for example, is the result of one particular dynamical path taken by the dice out of an infinite range of possibilities $-$ a path which although not predictable is, however, observable to within practical limits of precision for the classical system.

This interpretation is of course required to also be consistent with all observations of quantum phenomena. These include interference effects, such as described in figure 11.13(b), apparently well accounted for in terms of a superposition of wavefunctions, which in turn feature in the course of the calculations involving complex number algebra, but which don't individually generally represent a particular 'way' in which a process occurs. It will be necessary to trace a path from the many solutions picture of degeneracy in the present theory to the QFT Feynman rules for cross-section calculations based on the amplitude \mathcal{M}_{fi} , through equation 11.46 as described in section 11.2, and further to the postulates of quantum mechanics and the construction of the wavefunction $\Psi(x)$ for the non-relativistic limit, in order to see how such phenomena (and their quantum mechanical description) are compatible with the present theory.

The QFT calculation for the event rate at an e^+e^- collider, for processes such as depicted in figure 11.13(a), was presented in equation 10.7 and described in section 10.2. A doubling of the incoming luminosity, for example by doubling the bunch crossing frequency f in equation 10.2, or a doubling of the available final state phase space, in the final term of equation 10.7, leads to a direct doubling of the observed event rate. On the other hand on *adding* new intermediate processes interference between the complex amplitudes \mathcal{M}_{fi} may lead to a *reduction* of the event rate. Indeed, in practice the phenomenology predicted as a result of adding new hypothetical processes in such a calculation is sometimes studied in order to explain the observation of a lower than expected cross-section. However, according to the basic principles of the present theory the addition of new processes will only add to the 'number of ways' through which to bridge an initial to a final state and always serve to increase cross-sections and decay rates.

The question then may be asked how apparent interference phenomena arise in the present theory with probabilities based on degeneracy counts which always accumulate in a positive sense. However, it should be noted that there is no one-toone correspondence between components of the degeneracy count D and contributions to the transition amplitude \mathcal{M}_{fi} . Rather these two means of calculating the *total* probability are collectively related by a correspondence of the form of equation 11.46, which in particular implies a mechanism for normalising the degeneracy count through a complexification of the calculation.

Interference phenomena in quantum theory are more explicitly presented in the experiment of figure 11.13(b). As alluded to above this system can be analysed in terms of two wavefunctions, each emanating from one of the two intermediate slits, and added together to form the pattern of constructive and destructive interference generating the probability distribution for events observed on the final screen. Again there is no direct analogue of the 'superposition of wavefunctions' in the present theory, and again there is no one-to-one correspondence between wavefunctions and elements of a degeneracy count.

In the present theory such a degeneracy count is also not based on the 'number of ways' in which an electron, as a particle state, could pass through the slits, but rather on the number of underlying field solutions for $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ given the degeneracy of field redescriptions underlying the common geometry $G^{\mu\nu}(x)$ for the source S. Particle phenomena themselves arise as an apparent feature of these solutions. In fact, strictly speaking it is the phenomena of particle emission or detection, for example from the source S or at the point A on the screen in figure 11.13(b), that emerge in these solutions, with no continuous trajectory of a particle-like entity ever observed. Only the particle-like interactions are ever actually directly recorded.

Even for the events of sophisticated experiments such as depicted in figure 10.1 the apparent 'tracks' of particles are reconstructed from a series individual detector hits, in particular in a tracking chamber. 'Joining the dots' in this way creates an illusion of continuous particle trajectories, as was presumed for the incoming and outgoing particle states sketched in figure 10.2 for example. The theory is hence required to explain how field solutions for $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ incorporate apparent particle emission and detection phenomena, which in many cases create the impression of intermediate particle trajectories – as an interpretation in part based on a close analogy with the properties of classical particles. Since the effective local field interaction volume can be arbitrarily small the associated elementary particle states have no apparent size, consistent with a point-like interpretation.

In conclusion, for the present theory particle effects and the probabilistic nature of quantum phenomena generally arise out of the merging of two necessary features of the world. On the one hand the world we inhabit must be perceivable, as expressed mathematically in terms of geometric structures on an extended manifold such as $M₄$. On the other hand all such mathematical structures derive from a fundamentally one-dimensional temporal progression which may be expressed in terms of a general multi-dimensional form $L(\hat{v}) = 1$ together with its symmetries. Resolving these two requirements in a compatible manner leads to the equations of motion and physical properties of the tangible material world as perceived in spacetime and incorporating the phenomena of 'quantum mechanical' transitions deriving from the degeneracy of underlying field solutions.

While the underlying field components of $L(\hat{\mathbf{v}}) = 1$ on M_4 are in principle subject to the *full* symmetry degrees of freedom of the multi-dimensional form of time the geometrical interpretation needed to support the perceptual frame of the world requires the identification of a Riemannian geometry on the base manifold of the appropriate mathematical form with a lower symmetry. Here, as for quantum theory in general, symmetry rather than scale is the key to quantum processes; although (as discussed shortly after figure 11.12 in the previous section) with a higher degree of field symmetry more likely to be encountered on a 'microscopic' scale quantum phenomena are most frequently associated with such dimensions. The spirit of the principles of quantum mechanics is hence preserved in this new theory in unification with gravitation, with the identification $-\kappa T^{\mu\nu} := G^{\mu\nu}$ expressing the field equation of general relativity.

The similar nature of the interplay between the larger symmetry and the broken symmetry in the present theory to the situation in quantum mechanics can be exemplified by the Zeeman effect. The energy levels of the hydrogen atom are split by the presence of a uniform magnetic field, as a preferred direction in 3-dimensional space *reducing* or breaking the rotational symmetry of the system from $SO(3)$ to $SO(2)$. Passing a beam of electrons through a magnetic field configured to select a certain spin state provides a further example. Generally, in all cases of a measurement of a quantum mechanical system a structure of lower symmetry, such as the configuration of laboratory equipment, is imposed upon the intrinsically higher symmetry of the unobserved state.

In the present theory quantum phenomena arise through the unavoidable a priori imposition of the lower symmetry of 4-dimensional spacetime upon the general flow of time as a prerequisite for perception and observation in the world itself. Through our innate faculty to organise and interpret our experiences in the world through a coherent global geometrical manifold M_4 (playing the part of the directional magnetic field in the analogy with the Zeeman effect) the full E_7 symmetry of $L(\mathbf{v}_{56}) = 1$ is broken down to the local external symmetry $SO^+(1,3)$ together with the internal gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y$ (which, as the surviving symmetries, collectively play the part of SO(2) in the Zeeman analogy). However, while in the Zeeman effect the magnetic field direction is a particular choice of experimental setup, in perception the Lorentz frame, within an approximately global $SO^+(1,3)$ symmetry, is a *necessary* form for all physical experience of the world and hence applies to all experiments and observations.

Further, while the SO(2) symmetry of the uni-directional magnetic field imposed on a hydrogen atom with SO(3) symmetry results in a discrete splitting of the atomic energy levels, the surviving $SO^+(1,3) \times SU(3)_c \times SU(2)_L \times U(1)_Y$ external and gauge symmetry of the 4-dimensional perceptual field imposed over the full flow of time $L(\mathbf{v}_{56}) = 1$ with an E₇ symmetry will be correlated with a discrete set of possible transitions of the microscopic world which determines the spectrum of elementary particles. (Strictly speaking the 'surviving symmetry' is $SO^+(1,3)\times SU(3)_c \times U(1)_Q$ since the electroweak symmetry $SU(2)_L \times U(1)_Y$ is itself broken down to $U(1)_Q$ through its action on the external spacetime components of the 'vector-Higgs' $h_2 \equiv v_4 \in TM_4$. As

described in section 9.2 the electroweak symmetry is also yet to be explicitly identified in terms of E_7 generators). In general the resulting phenomena will be exhibited in the observed properties of particles in HEP experiments as well as in the non-relativistic limit of quantum mechanics itself, as exemplified in figures 11.13(a) and (b) respectively.

While the physical structures of both gravitational and quantum theory are ever present in nature it is possible to consider the limiting cases of the present theory as applicable to the corresponding empirical observations. The limit in which classical general relativity emerges on the one hand and a complementary limit through which an apparent quantum field theory emerges on the other hand can be described in terms of two significant symmetries for our world with the external Lorentz group $\overline{H} = SO^{+}(1,3)$ (in the notation of section 2.3) as a subgroup of $\overline{G} = \mathbb{E}_7$, with the latter being the symmetry of the full form of temporal flow $L(\mathbf{v}_{56}) = 1$ as described in section 9.2. These alternative limits can be characterised by the role of the linear connection $\Gamma(x)$ on the spacetime manifold M_4 , as described in table 11.1.

Symmetry	GR limit	QFT limit
$\overline{H} = SO^+(1,3)$	local symmetry on M_4	global symmetry on M_4
	generally $\Gamma(x) \neq 0$	can take $\Gamma(x) = 0$ exactly
$\hat{G} = \mathrm{E}_7$	effective macroscopic matter	local $E_7/SO^+(1,3)$ symmetry
	$T^{\mu\nu}(x) := G^{\mu\nu}(x) = f(Y, \hat{\mathbf{v}})$	\Rightarrow Y(x) gauge fields

Table 11.1: Limits in which general relativity and quantum field theory arise. The employment of a local or global freedom for Lorentz frames with $l(x) \in SO^+(1,3)$ was also discussed earlier in this section in relation to gauge theory.

The fact that GR and QFT emerge as almost exclusive complementary limits is not surprising given the notorious incompatibility of the respective mathematical theories and difficulties in uniting them under a single framework. However there is necessarily a trace of overlap even in the limiting cases. In the GR limit quantum effects are always locally present underneath the effective energy-momentum tensor which describes the apparent matter distribution, with macroscopic material properties shaped by the underlying quantum world. Similarly in the QFT limit particle interactions are clearly associated with regions of matter density and hence a minute but finite spacetime curvature is involved, which is a critical observation from the perspective of the present theory.

As well as shedding light on the respective limits, the present theory may also address conceptual problems for physical systems where both gravitational and quantum effects are significant. For example the difficulties seen in some approaches to quantum gravity such as the 'problem of time', as described earlier in this section, and the non-renormalisable nature of quantised gravity, as implied in the discussion following equation 10.86 in section 10.5, are avoided here since gravity itself is not quantised. While one aim of the present theory is to explore particle physics phenomena in the flat spacetime limit with Riemann curvature tensor components $R^{\rho}_{\sigma\mu\nu}(x) \rightarrow 0$, as an approximation to laboratory conditions to test the theory, the case for 'quantum transitions' and 'particle effects' for $R^{\rho}{}_{\sigma\mu\nu}(x) \neq 0$, and in general for a *highly* curved spacetime, is intended to be fully accounted for in this inclusive theory.

The general form of the relation $T^{\mu\nu}(x) := G^{\mu\nu}(x) = f(Y, \hat{\boldsymbol{v}})$ in table 11.1 will be applicable even in locations of the universe with extreme spacetime curvature, such as in the vicinity black holes and during the 'Big Bang' epoch. For example, an environment in which both gravitational and quantum effects are expected to be significant arises for the phenomenon of the emission of Hawking radiation (1974) in the highly curved spacetime in the proximity of a black hole, and similarly for the Unruh effect (1976) in which an observer undergoing a uniform high acceleration in the 'vacuum' of a flat spacetime can detect thermal radiation. Quantum and particle effects should be calculable with the present theory in such environments, and also for Big Bang cosmology – which will be discussed in the following two chapters.

In QFT the Fock space representation is generally only valid for free fields in flat spacetime. The Fourier expansion of the field $\ddot{\phi}(x)$ in equation 10.13 relies on the Poincaré symmetry of flat spacetime for the preferred basis of normal modes $e^{\pm ip\cdot x}$ and a corresponding preferred vacuum state $|0\rangle$. Particle excitations are built upon this ground state via the operators $a^{\dagger}(\boldsymbol{p})$ and $a(\boldsymbol{p})$. In flat Minkowski spacetime only global inertial frames of reference are used for which the particle content of a state, implied in the Fourier components, agrees for all observers.

This construction is not possible in curved spacetime for which the reference frames of global coordinate systems are necessarily non-inertial. For QFT in curved spacetime there is generally no unique set of normal modes, which results in different inequivalent expressions of a particular QFT without a unique vacuum state, and the particle interpretation in turn becomes ambiguous. Hence in general there is no objective possibility of identifying either a vacuum or specific particle state for QFT in general relativity. However, interference between normal modes expressed in different general coordinate systems has the physical consequence that real particles may be created by gravitational fields.

Indeed physical particle states produced by gravitational fields or, equivalently, by accelerated observers are in principle detectable and hence do represent real objective phenomena which in principle should be consistently accounted for in a complete theory. Similarly the particle states observed in high energy physics experiments are empirical objective entities. In all cases the detection of particle effects hinges on the nature of particle or field interactions, without which the particles could not be observed. In the present theory it remains then to fully understand the nature of particle phenomena, and their apparent physical interactions in general, as emerging out of the underlying interactions of fields, as represented by a degeneracy of redescriptions, as we began to address in the previous section and will further consider in section 15.2 in the discussion of figure 15.2.

In the present theory the use of the Fourier transform expansion in equation 11.6 is merely an effective approximation that arises in the limit of a flat Minkowski spacetime, and in which the apparent particle effects might most simply be analysed. Elementary particles are not fundamental entities out of which the world is built, they are a robust phenomenon that arises in the flat spacetime (and near vacuum) limit, as alluded to in the opening of section 10.1 and as studied in experiments such as depicted in figure 10.1. The properties of 'particles' may be less robust in highly curved spacetime, and more difficult to calculate than in the fixed limit of flat background manifold, but there is no fundamental conceptual difficulty.

The field redescription $\overline{\psi}\gamma^{\mu}\psi \to A^{\mu} \to \overline{\varphi}\gamma^{\mu}\varphi$ of figure 11.4 is presumed to be locally enveloped in a spacetime geometry $G^{\mu\nu}(x)$ which takes a form resembling that of $T^{\mu\nu} := G^{\mu\nu}$ in figure 11.1. If such an interaction takes place in the prevailing environment of a highly curved spacetime, for example in the proximity of a black hole, then to a certain extent all of the fields, $\psi(x)$, $\varphi(x)$ and $A^{\mu}(x)$, will be 'bent the same way' and hence processes such as depicted in figure 11.4 might be largely unaffected. Similar underlying field redescriptions in combination with immense gravitational tidal forces might then provide a description of black hole evaporation in the context of the present theory.

The question can also be asked concerning the nature of phenomena for yet more extreme spacetime curvature, such as in the region of a black hole 'singularity' or generally corresponding to a yet higher scale of energy. In the context of figure 11.10 the GUT scale, at around 10^{15} GeV, in marking a point of gauge coupling unification ought to be of significance for the present theory in terms of the phenomena of the internal forces, while the external gravitational field will be treated in the same manner as for the low energy phenomena. Further, in the present theory gravity itself is not 'quantised', there are no 'graviton' particles, and the Planck scale at around $10^{19} \,\mathrm{GeV}$ may just be a dimensional quirk with no particular significance. Hence arbitrarily high energy densities and arbitrarily high spacetime curvature might be considered in the present theory in a continuous manner without limit.

In summary, from the point of view of the present theory the postulate in quantum theory that an event probability is determined by the square of the absolute value of an 'amplitude', with unitary symmetry imposed to ensure the structure is consistent with the basic laws of probability, should be considered as a *provisional* construction standing in need of an underlying conceptual basis and physical explanation. Such an explanation would be preferred in place of any theoretical 'postulate', and here it lies in the idea of the natural degeneracy inherent in the number of ways local field solutions may be found for $G^{\mu\nu} = f(Y, \hat{v})$ for processes such as those observed in figure 11.13 and more generally.

This is the key to combining general relativity and quantum phenomena in a single complete and unified theory. Indeed, given the prohibitive conceptual and mathematical difficulties encountered in attempting to unify these two pillars of $20th$ theoretical physics it seems likely that a significant concept or postulate on at least one side must yield some ground. Here the definition of probability in terms of amplitudes in quantum theory seems a reasonable place for this, with the amplitudes and wavefunctions of quantum theory then representing calculational tools employed in an intermediate complexification of a computation. This approach has been exemplified by unravelling the QFT event rate calculation of equation 10.7 and making the case for replacing the contribution from the amplitude \mathcal{M}_{fi} by a quantity based on a degeneracy count D via the associations of equation 11.46.

This foundation also unifies the notion of probability with the classical concept in the sense of essentially referring to the 'number of ways' that a process can occur given a particular initial state or situation. However, while classical probabilities concern the number of ways that things can happen in spacetime M_4 , quantum probabilities concern the more fundamental question of the number of ways in which the spacetime manifold M_4 itself can be constructed with a world geometry described by $G^{\mu\nu}(x) = f(Y, \hat{\boldsymbol{v}})$. Further, in principle this approach to quantum phenomena also leads to a clarification of the meaning of 'renormalisation' as discussed for equation 11.48 in the previous section.

For theories which postulate extra spatial dimensions, such as the Kaluza-Klein theories described in chapter 4, our 4-dimensional spacetime world is contained within the larger space, for example as a 4-dimensional *brane* embedded within the higher-dimensional *bulk* manifold or with the extra dimensions being 'compactified', as discussed in section 5.4. For the present theory founded on one-dimensional temporal flow the extended physical world is perceived through the structure and symmetries of the multi-dimensional form $L(\hat{v}) = 1$, with the degeneracy of solutions for constructing such a 4-dimensional world underlying the phenomena of quantum theory while the external spacetime geometry itself conforms with the structure of general relativity.

As well as combining general relativity and quantum theory in a consistent framework within which the two theories are separately preserved in essence, the complete conceptual theory is based on sound intuitive principles, founded upon the ever pervading multi-dimensional form of temporal flow $L(\hat{v}) = 1$ rather than upon seemingly arbitrary, mysterious or purely pragmatic assumptions. The theory should of course also be able to make predictions and be found to be in full agreement with all observations. Such a correspondence has been initiated in chapters 8 and 9 with regards to comparison with the Standard Model of particle physics. Further, on incorporating all physical scales, including that of HEP phenomena, in principle the present theory is expected to be profusely testable.

All the underlying fields in nature, which underlie for example electron and photon particle states, are in continual interaction through mutual indistinguishability under the external geometry $G^{\mu\nu}(x)$ – from the interaction region of an HEP experiment such as that in figure 10.1, to atoms and molecules, through to biological organisms, planets, stars and galaxies, with the underlying processes moulding a smooth and continuous geometry $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ with all the quantum phenomena embedded within and in turn, through $T^{\mu\nu} := G^{\mu\nu}$, shaping the structure and apparent phenomena of the material world.

As well as the extreme environment of a highly curved spacetime alluded to above, the complementary question concerning the nature of the 'vacuum state' can also be considered. Even in the apparent vacuum, away from tangible physical matter, in general a form of $G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ must be present throughout M_4 in order to describe the spacetime geometry. This structure might in principle implicitly include a form of effective 'vacuum energy', incorporated into the spacetime geometry and describing the effects of a cosmological constant Λ , at least to a good approximation, and hence in turn accounting for observations of the large scale structure of the universe. Indeed, more generally, as well as terrestrial laboratory phenomena the present theory has also been developed with the cosmological scale in mind, and hence in the following two chapters we review aspects of cosmology in the context of the new theory.

Chapter 12

Cosmology

12.1 The Large Scale Structure of the Universe

While the previous chapter focussed on the application of the present theory to the smallest observable scales, regarding in particular the quantum field and particle phenomena studied in high energy physics experiments, here we return to consider general relativity and gravitation, continuing the thread from sections 5.2 and 5.3 in the light of the intermediate chapters, as applied up to the largest empirically accessible scale of the observable universe and beyond. In the context of the large scale structure of 4-dimensional spacetime the right-hand side of equation 5.32 can generally be considered to describe the effective macroscopic form of apparent matter terms, with $-\kappa T^{\mu\nu} := G^{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ for this equation, that is in the GR limit as summarised in table 11.1, with the practical normalisation factor of $-\kappa$ inserted. However an understanding of the impact upon the spacetime geometry of the underlying microscopic fields and their interactions will also be directly relevant both for the universe at the present epoch as well as in its much earlier history. Indeed since the energy density in the early universe reaches and surpasses that attainable in high energy physics experiments, the environment of the immediate aftermath of the Big Bang may itself provide a possible test arena for theoretical particle physics through any imprint which the corresponding phenomena may leave in the structure of the cosmos which is still observable today.

In the following two sections we review some of the main features of standard textbook cosmology, as deduced from and motivated by empirical observations. In the following chapter we then collect and describe a series of observations concerning the present theory which, at a qualitative level at least, correlate with a number of aspects of modern cosmology. Without making a quantitative argument in terms of cosmological parameters these aspects include the dark sector of implied matter and energy in the universe and the origin and nature of the Big Bang and the very early universe itself.

The rather direct application of the conceptual scheme described in the previous chapters to the cosmological scale will first be outlined briefly in this section. This

application is possible since the general picture of the standard cosmological model of the evolution of the universe according to Einstein's field equation of general relativity, given broad underlying assumptions concerning the large scale structure of spacetime, is naturally compatible with the present framework.

Based on the translation symmetry represented in figure 2.2 we described in sections 2.1 and 2.2 how the perceptual background of a flat SO(3) symmetric spatial manifold M_3 could be effectively derived through the structure and symmetries of the flow of time expressed in the form $L(v_3) = 1$ of equation 2.14. On extending this model to the case of a full $\tilde{G} = SO(5)$ symmetry of $L(v_5) = 1$ projected over M_3 , as described for figure 2.7 in section 2.3, a finite external (and also internal) curvature was obtained. Subsequently a $\hat{G} = SO^+(1,9)$ model for the case of a 4-dimensional spacetime M_4 as pictured in figure 5.1 was described in section 5.1, again introducing minor distortions from a flat geometry corresponding to the effects of general relativity. These geometric distortions are presumed to be undetectable in everyday experience – that is out of the degrees of freedom of the full symmetry group of $L(\hat{\boldsymbol{v}}) = 1$ projected onto the base manifold M_4 we require the local $SO^+(1,3) \subset G$ subgroup to be broken down to an approximately global symmetry of the 4-dimensional spacetime manifold, incorporating an approximately Euclidean 3-dimensional space, forming the backdrop for our perception of physical objects in the world.

This requirement is borne out by our observations of the world around us on the scale of the solar system for which the non-Euclidean effects of general, as well as special, relativity are indeed imperceptible. The non-Euclidean effects such as the deflection of starlight passing close to the sun are well beyond the reach of casual observation. On the other hand local observations such as the accelerating fall of an apple from a tree might at first sight be ascribed to a 'force of gravity' active within a flat arena of space and time, rather than to an effect of a curved spacetime arena itself. The apparent flatness of the local geometry both from the point of view of our everyday experience of the world and also for most scientific experiments accounts for the fact that the existence of a non-Euclidean element of 3-dimensional space combined with 1-dimensional time was not recognised, through centuries of scientific developments, until the early 1900s.

Carrying the same principle of our innate requirement of perception in the world to the largest scale in which we encompass everything in our observable universe it seems natural to ask how it could be possible for our existence and experiences to influence in any way the shape or form of the universe over regions measuring billions of light-years across. However, a central point of the work presented in this paper is that here we consider the whole universe to be the physical manifestation that is created through and within the possibility of our experiencing it and is therefore shaped by the necessary form of that possibility, as we shall describe further in chapter 14. The initial naive picture that hence comes to mind is then based upon the assumption of an approximately Euclidean background extending to the largest observable scale, neglecting the (generally imperceptible) local variations from flatness, with the flow of time propagating through a 4-dimensional manifold as depicted in figure 12.1. This picture represents the largest scale realisation, for our own 4-dimensional universe, of the general idea introduced in figure 2.3 of section 2.2 for the model 3-dimensional world.

Figure 12.1: Propagation of galaxies, clusters of galaxies and large scale physical structures through the M_4 spacetime manifold, with the temporal dimension directed from left to right and one spatial dimension suppressed.

We further recall that in the full theory the components of the 4-dimensional vector field $v_4(x)$ on M_4 are considered to be locally embedded in a higher-dimensional form of temporal flow $L(v_{27}) = 1$ via the space of $v_{27} \equiv \mathcal{X} \in \mathfrak{h}_3\mathbb{O}$ matrices through equations 8.4 and 8.5 of section 8.1, and in turn within the form $L(v_{56}) = 1$ via the elements $x \in F(h_3\mathbb{O})$ in the form of equation 9.46 as described in section 9.2. From a purely mathematical point of view the intermediate 4-dimensional case is readily bypassed in generalising from a 1-dimensional temporal progression to higher-dimensional forms, here represented by a 27-dimensional and on to a 56-dimensional form of time with a full $\hat{G} = \mathbb{E}_6$ and $\hat{G} = \mathbb{E}_7$ symmetry respectively. However, in order to *physically* experience or perceive any structures implicit within the general form of time a lowerdimensional part, with mathematical properties isomorphic to the geometrical forms required to define the perception of objects in the world, is projected out, or syphoned off, from the full general form of temporal flow.

In our world this has been taken to be achieved through extracting $v_4 \equiv h_2 \in$ $h_2\mathbb{C} \subset h_3\mathbb{O} \subset F(h_3\mathbb{O}),$ with the quartic form $L(\mathbf{v}_{56}) = 1$ for $\mathbf{v}_{56} \in F(h_3\mathbb{O})$ having an E₇ symmetry, and projecting the vector component $v_4 \text{ }\subset v_{56}$ onto TM₄. The form $L(\mathbf{v}_4) = \det(\mathbf{h}_2) = h^2$ has the symmetry group $SL(2,\mathbb{C}) \subset E_7$ which, as described in section 7.1, is the double cover of the external Lorentz group acting on Lorentz 4-vectors. While the representations of the complementary internal symmetry upon the components of $L(v_{27}) = 1$ and $L(v_{56}) = 1$ are reminiscent of Standard Model properties, as described in chapter 8 and section 9.2 respectively, an extension for example to an E_8 symmetry of a form $L(v_{248}) = 1$, as outlined hypothetically in
section 9.3, may be needed to fully incorporate the structure of the Standard Model.

In principle this projection, on employing the associated 4-dimensional translation symmetry of the form $L(v_{56}) = 1$, opens out the local Lorentz subgroup into an approximately global symmetry on the M_4 manifold, breaking the full local E_7 symmetry while at the same time actually generating the spacetime manifold itself. That is the M⁴ manifold is created in the act of the symmetry breaking projection from $v_{56} \in F(h_3 \mathbb{O}) \to v_4 \in TM_4 \equiv h_2 \mathbb{C}$, with $v^a(x)$ for $a = 0, 1, 2, 3$ being the components of the Lorentz tangent vector field in a local Minkowski coordinate frame on the manifold as described in section 5.3. In terms of the initial picture, deriving from the translation symmetry as described for figure 2.2, the metric $g_{\mu\nu}(x) = \text{diag}(1, -1, -1, -1)$ may in fact be adopted globally. This continues to be the case to a good approximation in practice even when planets, stars and galaxies are incorporated as depicted in figure 12.1. However, while maintaining the approximation of neglecting the local distortions correlated with matter in these forms, it will not be possible to adopt a global Minkowski frame when considering the overall cosmological point of view.

That is, while compatible with an approximately flat $SO^+(1,3)$ frame locally, on the scale of the solar system for example, on larger spacetime scales through to the vast arena of the universe studied in cosmology there is no longer a necessity for the geometric form of spacetime to describe a perceptual frame even in approximation. Further, extrapolating beyond our possible experience or observation of the world the 4 dimensional geometrical interpretation may itself at some point break down altogether. This may apply to extreme regions such as black holes at any epoch and the structure of the very early history of the universe and the Big Bang. The effects of particle physics studied in an Earth-bound laboratory, and in the previous chapters, are also likely to play a significant role when extrapolated to the extreme conditions of a highly non-Euclidean spacetime geometry, as alluded to in the previous section.

In the following section the standard cosmological model and a range of possible large scale metric solutions will be reviewed, before turning to the very early universe in section 12.3. This will provide a basis for the perspective of the present theory to be presented in the following chapter.

12.2 The Standard Model of Cosmology

While the Standard Model for particle physics has been constructed in recent decades in parallel with the findings of high energy physics experiments, the underlying tools of quantum field theory were originally developed in the 1920s through to the 1940s. The framework for cosmological models was originally developed over a similar period following soon after the publication of general relativity in 1915 and through to the 1930s, although again here the 'standard model of cosmology' has only become established in recent decades in the light of the empirical data revealed with modern observational technology. In this section we examine the picture of the cosmos and the standard cosmological model that has emerged out of this work (see for example [5, 73]).

The standard approach incorporates general relativity, as reviewed in sections 3.3 and 3.4, for which the empirical observation that spacetime curvature is strongly correlated with the presence of matter is expressed through the field equation $G^{\mu\nu} = -\kappa T^{\mu\nu}$. This equation postulates the equality of the Einstein tensor $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2} R g^{\mu\nu}$ with the energy-momentum tensor $T^{\mu\nu}$, to within a constant of proportionality. This approach is here summarised in terms of three quotes from [74]:

- We wish to relate the curvature of spacetime to the presence of matter, since gravity appears in the neighbourhood of matter' ([74] p.232). The proportionality constant is determined for weak fields by comparison with Newton's theory of gravity and found to be $\kappa = 8\pi G_N$, where G_N is Newton's constant, as described for equation 3.75.
- 'It will be assumed that the metric in a nearly empty universe is nearly Minkowski' ([74] p.229). Essentially this implies that a flat spacetime arena M_4 is presupposed before the introduction of matter. Within relatively local portions of the universe a flat Minkowski spacetime can act as a boundary condition in regions sufficiently far from matter, as for the example of the Schwarzschild solution in equation 5.49.
- '...the vanishing of the divergence of $G^{\mu\nu}$ as a mathematical identity implies the vanishing of the divergence of $T^{\mu\nu}$ ([74] p.232). That is in light of the contracted Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$ this conclusion follows immediately given that the Einstein field equation is assumed to hold.

The divergence-free nature of $T^{\mu\nu}$ can be interpreted as the conservation of energy and momentum in the limit of an approximately flat spacetime, as described in the opening of section 5.2, since for a suitable choice of coordinates with linear connection $\Gamma \to 0$ we have $T^{\mu\nu}_{\mu} = 0$. On the other hand, this equation for the conservation of energy and momentum is often cited as a starting point and then expressed in a general curved spacetime as $T^{\mu\nu}_{;\mu} = 0$, and it is this observation that then justifies the introduction of $T^{\mu\nu}$ on the right-hand side of the field equation itself, with the Einstein tensor on the left-hand side, since it happens to be also the case that $G^{\mu\nu}_{;\mu} = 0$ as the contracted Bianchi identity. Consistent with this requirement an additional divergence-free term may be postulated, associated with a 'cosmological constant' Λ, as may be necessary to account for the empirically observed evolution of the universe, yielding the full standard field equation as quoted in equation 3.84 and reproduced here (where we turn here to a convention of generally using lower indices in such expressions):

$$
G_{\mu\nu} + \Lambda g_{\mu\nu} = -\kappa T_{\mu\nu}.
$$
\n(12.1)

In 1922 Aleksandr Friedmann made two classes of assumptions in order to obtain solutions for the spacetime structure of the universe as a whole. The first class required that the gravitational field should satisfy the equation 12.1, that is the Einstein field equation including the cosmological constant term (Friedmann considered the case for both arbitrary Λ as well as $\Lambda = 0$, with matter represented as a pressureless fluid with energy-momentum tensor $T_{\mu\nu} = \rho u_{\mu} u_{\nu}$ where ρ is the proper density of matter. In 1927 Georges Lemaître, working independently of Friedmann, considered the more general case by including a spatially isotropic pressure term and hence treating matter as a perfect fluid with an energy-momentum tensor in the form of equation 5.37, that is:

$$
T_{\mu\nu} = (\rho + p)u_{\mu}u_{\nu} - pg_{\mu\nu}
$$
\n(12.2)

where p is the pressure and here u_{μ} represents the 4-velocity of the flow of galaxies, as depicted for example in figure 12.1. This energy-momentum is then substituted into Einstein's equation 12.1 to give:

$$
G_{\mu\nu} + \Lambda g_{\mu\nu} = -\kappa (\rho + p) u_{\mu} u_{\nu} + \kappa p g_{\mu\nu}.
$$
 (12.3)

We now know that the contribution of radiation pressure to the evolutionary dynamics of the universe is most significant for around the first 10,000 years of its history, with the contribution of the matter density becoming comparable around 50,000 years after the Big Bang and subsequently increasingly dominating over the radiation term. Hence the idealisation of Friedmann, treating the flow of galaxies as a dust or pressureless fluid with $p = 0$, makes a very good approximation for modelling the cosmic evolution, particularly since the epoch of the 'decoupling' of matter from radiation around 372,000 years [44] after the Big Bang, still relatively early in the 13.8 billion year history of the universe.

The second class of assumptions made by Friedmann in order to obtain a solution concern the nature of preferred coordinate systems and more direct restrictions on the form of the metric deriving from symmetries imposed on the spacetime. Based on the picture of galaxies pursuing non-intersecting world lines, for which figure 12.1 represents only a particular special case, 3-dimensional spacelike hypersurfaces, orthogonal to and parametrised by a global timelike coordinate t , are assumed to have a uniform t-dependent 3-dimensional scalar curvature $R_3(t)$ independent of the location on a given 3-dimensional spatial surface. The unambiguous cosmic time t is taken to be the proper time as measured for any given galaxy. The 'Copernican view', that here on Earth we do not inhabit a central or preferred location of the universe, is subsumed into the 'cosmological principle' which asserts that at any given cosmic time t the universe on large scales is spatially homogeneous and isotropic about any location.

From an observational point of view at the present epoch the assumption of homogeneity may be justified by the smallness of fluctuations in the distribution of galactic clusters on scales larger than a few 100 Mpc (megaparsecs, where 1 parsec is around 3.26 light-years) in an observable universe with distance scales of up to the order of the Hubble radius:

$$
R_H := c/H_0 \simeq 3000h^{-1} \text{ Mpc}
$$
 (12.4)

with $h \approx 0.7$ and the Hubble constant H_0 defined below for equation 12.13. Similarly, the assumption of isotropy may be justified by the evenness of the cosmic microwave background (CMB) radiation to within of order 1 part in $10⁵$ as observed over the full coverage of the sky from the Earth.

The mathematical basis for the assumptions of the cosmological principle was studied thoroughly by H.P. Robertson and independently by A.G. Walker in the 1930s. The 3-dimensional hypersurfaces for constant t are everywhere orthogonal to a congruence of geodesics given by the integral curves of the vector field $\partial/\partial t$. For each solution the hypersurface curvature $R_3(t)$, while it can vary with time, remains always either positive (3-sphere), negative (hyperboloid) or zero (for a spatially flat universe). The Robertson-Walker line element is the most general spacetime metric compatible with homogeneity and isotropy and can be expressed in terms of intervals of proper time τ as:

$$
d\tau^2 = dt^2 - a^2(t) \left[\frac{dr^2}{(1 - kr^2)} + r^2 (d\theta^2 + \sin^2\theta d\phi^2) \right]
$$
 (12.5)

where the parameters $a(t)$ and k will be described below. With the world line of any given idealised galaxy expressed in terms of constant 3-dimensional spatial spherical coordinates $\{r, \theta, \phi\}$ the full 4-dimensional set $\{t, r, \theta, \phi\}$ describes a comoving coordinate system with the cosmic time parameter t equivalent to the proper time τ elapsed for the galaxy.

As for any metric for 4-dimensional spacetime here the convention is to take the components of $g_{\mu\nu}(x)$ to have the dimension of length squared, that is $[g_{\mu\nu}] = [d\tau^2] =$ L^2 (with the dimension of length L equivalent to that of time T since $c = 1$, while in the notation of the discussion following equation 10.86 the mass M dimension of $g_{\mu\nu}$ is $D = -2$). In turn the components of the metric inverse have the dimension $[g^{\mu\nu}] = L^{-2}$. (We note that in the present theory the internal Killing metric components, such as $g_{\alpha\beta}$ in equation 4.4 for the case of a Kaluza-Klein metric, are not interpreted as representing a physical length in a higher-dimensional space in the present theory, however here we are dealing with purely external metric components in 4-dimensional spacetime). This convention is consistent with the principle of general covariance in general relativity, as described in section 3.4, which implies that in general no physical significance can be attached to a set of coordinates, which consists of numerical parameters of dimension L^0 . As implied in the name, only when the manifold is endowed with a 'metric' are lengths defined. In fact all parameters on the right-hand side of equation 12.5, including the scale factor $a(t)$, can be considered to be dimensionless quantities. Since an implicit factor of $g_{00} = 1$ carrying the dimension L^2 accompanies the dt^2 term in equation 12.5, the cosmic time coordinate t may be *interpreted* as having the dimension of $T \equiv L$. A similar interpretation might be applied to spatial coordinates in certain cases, in particular for Euclidean coordinates $\{x, y, z\}$ in the limit of a flat spacetime with Minkowski metric.

The sign of the dimensionless real number k in equation 12.5 indicates the sign of the 3-space curvature. For $k = 0$ the spatial hypersurfaces are flat, although even in this case the 4-dimensional curvature will generally be finite. For non-zero values of k the coordinate r may be redefined as $r \to r/|k|^{\frac{1}{2}}$, and the scale factor as $a \to a|k|^{\frac{1}{2}}$, such that the thus normalised values of $k = +1, -1$ and 0 represent the positive, negative and zero spatial curvature solutions respectively.

The simplifying assumptions of the cosmological principle have hence reduced the 10 parameters of the unknown metric $g_{\mu\nu}(x)$ down to a single real parameter $a(t)$ along with a discrete set of three possible values for k in equation 12.5. Together $a(t)$ and k characterise the Robertson-Walker line element which itself represents a trial solution for cosmological models. The specific form of the line element will be determined by the dynamics provided by equation 12.3, which depends in turn on the choice of cosmological constant Λ and the 'equation of state' relating ρ and p in the energymomentum tensor on the right-hand side of Einstein's field equation. Equation 12.2, with ρ and p functions of t only, is in fact the most general form of energy-momentum tensor consistent with the requirements of homogeneity and isotropy as expressed in the cosmological principle, which is also respected by the Λ term in equation 12.3, with $g_{\mu\nu}(x)$ in the form of equation 12.5. The resulting differential equations in the single independent variable t may be solved for a, ρ and p, each of which is a function of t only owing to the homogeneity assumption.

The Einstein tensor is constructed from the Riemann curvature tensor in terms of the components of the Ricci tensor as $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}$, as introduced after equation 3.71 and via equation 3.74; the Riemann tensor is a function of the linear connection Γ as expressed in equation 3.73, and the torsion-free Levi-Civita connection of equation 3.53 is employed as also described in section 3.3. The components of the metric tensor $g_{\mu\nu}$ implied in equation 12.5 are:

$$
g_{00} = 1,
$$
 $g_{11} = \frac{-a^2}{(1 - kr^2)},$ $g_{22} = -a^2r^2,$ $g_{33} = -a^2r^2\sin^2\theta$ (12.6)

These can be substituted into the above chain of relations, via the linear connection, to determine the components of the Ricci tensor $R_{\mu\nu}$ and scalar curvature $R = g^{\mu\nu} R_{\mu\nu}$ as (see for example [75] p.151):

$$
R_{00} = 3\frac{\ddot{a}}{a}
$$
\n
$$
R_{11} = -(a\ddot{a} + 2\dot{a}^2 + 2k)/(1 - kr^2)
$$
\n
$$
R_{22} = -(a\ddot{a} + 2\dot{a}^2 + 2k)r^2
$$
\n
$$
R_{33} = -(a\ddot{a} + 2\dot{a}^2 + 2k)r^2 \sin^2 \theta
$$
\n
$$
R = 6\left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2}\right)
$$
\n(12.8)

with both $g_{\mu\nu} = 0$ and $R_{\mu\nu} = 0$ for $\mu \neq \nu$, and the notation $\dot{a} = da/dt$ and $\ddot{a} = d^2a/dt^2$ has been employed. Further following the standard procedure and completing the chain of relations from the metric $g_{\mu\nu}$ to the Einstein tensor $G_{\mu\nu}$ the above expressions for $R_{\mu\nu}$ and R are substituted into the field equation 12.3, which includes the cosmological term and energy-momentum in the form of a perfect fluid, with components of the galactic flow 4-velocity $u_{\mu} = g_{\mu\nu} \frac{dx^{\nu}}{d\tau} = (1, 0, 0, 0)$ in the comoving coordinates, to find for the G_{00} and G_{11} components respectively:

$$
\frac{\dot{a}^2}{a^2} + \frac{k}{a^2} - \frac{1}{3}\Lambda = \frac{\kappa}{3}\rho \tag{12.9}
$$

$$
2\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{k}{a^2} - \Lambda = -\kappa p \tag{12.10}
$$

Only the above two independent non-trivial equations result since the equations for the G_{22} and G_{33} components are each identical to that for G_{11} in equation 12.10, due to the symmetries of the cosmological principle, while the set of six equations for $G_{\mu\nu}$ with $\mu \neq \nu$ are identically zero on both sides. In equations 12.9 and 12.10 the parameters a, ρ and p are functions of the cosmic time t while Λ , κ and k are constants.

Multiplying equation 12.9 by a^3 , differentiating the full resulting expression with respect to t and replacing the left-hand side by equation 12.10 multiplied by $\dot{a}a^2$ leads to the relations:

$$
\frac{d}{dt}(\rho a^3) = -3p\dot{a}a^2 = -p\frac{d}{dt}a^3
$$
\nthat is:

\n
$$
\frac{d}{da}(\rho a^3) = -3p a^2
$$
\n(12.11)

These equations may also be derived directly from the identity $T^{\mu\nu}_{;\mu} = 0$ for the perfect fluid energy-momentum tensor of equation 12.2 given the metric components of equation 12.6 (see for example [75] pp.152–153, with the same result holding if a Λ $\frac{\Lambda}{\kappa}g_{\mu\nu}$ term is included in $T_{\mu\nu}$ since $\left(\frac{\Lambda}{\kappa}\right)$ $\frac{\tilde{\Lambda}}{\kappa} g^{\mu\nu} \Big|_{;\mu} = 0$. Alternatively the constraint $T^{\mu\nu}_{;\mu} = 0$ can be combined with equation 12.9 in order to derive equation 12.10.

This apparent redundancy between the Einstein field equation and the expression $T^{\mu\nu}_{;\mu} = 0$ is expected since, as alluded to above, the form of the field equation $G_{\mu\nu} = -\kappa T_{\mu\nu}$ can itself be motivated by the divergence-free identity which applies to both sides and contains equivalent information. In fact in defining $-\kappa T_{\mu\nu} := G_{\mu\nu}$, which is the interpretation implied in the third bullet point near the opening of this section, the identity $T^{\mu\nu}_{;\mu} = 0$ is simply a *copy* of the contracted Bianchi identity $G^{\mu\nu}{}_{;\mu}=0$ which is an intrinsic property of the Einstein tensor (see also for example [6] p.729).

The apparent 'conservation law' $T^{\mu\nu}_{;\mu} = 0$ can not be directly interpreted as the 'conservation of total energy-momentum' unless the 4-dimensional geometry involved is asymptotically a flat Minkowski spacetime, which is not generally the case for the Robertson-Walker line element of equation 12.5. In general a suitably flat spacetime can be identified for local regions of the universe, as suggested in the second bullet point near the opening of this section, and more specifically leads to energymomentum conservation when applied in the laboratory setting, such as for the QFT limit summarised in table 11.1 towards the end of section 11.4.

As described above in comparison with the contribution from radiation pressure the universe has been matter dominated since a relatively short time after the Big Bang. Hence considering the pressure-free case of dust with $p = 0$ equation 12.11 implies that ρa^3 is constant in time. In this case the matter density at any epoch can be written as $\rho = \rho_0 \frac{a_0^3}{a^3}$, where a subscript '0' on a quantity such as the density ρ_0 or scale factor a_0 denotes the present day value at cosmic time $t = t_0$. Generally the boundary condition $a(0) = 0$ will be employed, with the cosmic time $t = 0$ designating the origin of the universe at the 'Big Bang'. For such a cosmology the present cosmic time t_0 hence denotes the current ago of the universe.

The physical spatial distance $d(t)$ between any two galaxies at a given cosmic time t is simply $d(t) = a(t)\Delta\Sigma$ where $\Delta\Sigma$ is the comoving 'coordinate distance' between the galaxies (when interpreted with care for the length L dimension as introduced through the metric as described above, similarly as discussed for the interpretation of the Schwarzschild solution around equation 5.50). The physical speed of one of the galaxies relative to the other is $v = \frac{d}{dt}d(t)$ which leads directly to the expression:

$$
v = \frac{\dot{a}(t)}{a(t)}d(t) = H(t)d(t)
$$
\n(12.12)

where $H(t) := \frac{\dot{a}(t)}{a(t)}$ is the Hubble parameter, which generally varies with time. Equation 12.12 expresses Hubble's law which states that at any given epoch t the relative speed between any two galaxies on the corresponding spatial hypersurface is directly proportional to the distance $d(t)$ between them, with the constant of proportionality given by the Hubble parameter $H(t)$ at that cosmic time. Hubble's law is a direct consequence of the form of the Robertson-Walker line element of equation 12.5 with variable $a(t)$ and says nothing about the actual dynamics, that is the function $a(t)$ itself, for the universe. At the present epoch the Hubble parameter is called the Hubble constant (since it is the same everywhere in space) $H_0 = H(t_0) = \frac{\dot{a}_0}{a_0}$ and is empirically found to take the value:

$$
H_0 = 100h \, \text{km s}^{-1} \text{Mpc}^{-1} \tag{12.13}
$$

with $h = 0.673 \pm 0.012$ [44], as employed above in equation 12.4.

The functional form of $a(t)$ itself may be determined from equation 12.9, which is also called the Friedmann equation. While considering the case with $p = 0$ if the cosmological constant is also neglected by setting $\Lambda = 0$ the Friedmann equation for any cosmic time t may be written as:

$$
H^2 + \frac{k}{a^2} = \frac{\kappa}{3}\rho \tag{12.14}
$$

The particular value of $\rho = \rho_c = \frac{3H^2}{\kappa}$ $\frac{H^2}{\kappa}$ is called the 'critical density' and corresponds to a solution with $k = 0$, that is a universe which is spatially flat at any epoch. This solution is known as the Einstein-de Sitter model and describes an ever-expanding universe with scale factor $a(t) \propto t^{\frac{2}{3}}$, as listed in table 12.1 (in contrast the radiation dominated case with $p \neq 0$ and equation of state $p = \frac{1}{3}$ $\frac{1}{3}\rho$ results in a dynamics with $a(t) \propto t^{\frac{1}{2}}$ for $k = 0$, as also listed in the table).

FLRW model: $G_{\mu\nu} = -\Lambda g_{\mu\nu}$			Matter $\rho \neq 0$ Radiation $p \neq 0$ $R_{\mu\nu} = \lambda(t)v_{\mu}v_{\nu}$	
$p = \epsilon \rho : \epsilon =$				
	$a(t) \propto \int e^{\sqrt{\frac{\Lambda}{3}}(t-t_0)} a_0$	$t^{\frac{2}{3}}$	$t\overline{2}$	$t\bar{3}$
$\rho(t) \propto$	constant	$t^{-2} \sim a^{-3}$	$t^{-2} \sim a^{-4}$	$t^{-2} \sim a^{-6}$

Table 12.1: Four FLRW (Friedmann-Lemaître-Robertson-Walker) cosmological models for $k=0$ with an energy-momentum $T_{\mu\nu} = -\frac{1}{\kappa}G_{\mu\nu}$ in the form of equation 12.2, with p and ρ related via the equation of state $p = \epsilon \rho$, corresponding to a universe dominated by a cosmological term (see below), matter, radiation and through $R_{\mu\nu} = \lambda(t)v_{\mu}v_{\nu}$ (for section 13.1) respectively. The evolution of the scale factor $a(t)$ and effective matter density $\rho(t)$ are obtained as solutions for equations 12.9 and 12.10.

Equation 12.14 can be rearranged in the form:

$$
\frac{k}{a^2} = H^2(\Omega_M - 1) \tag{12.15}
$$

on introducing the matter density parameter $\Omega_M = \frac{\rho}{\rho}$ $\frac{\rho}{\rho_c} = \frac{\kappa \rho}{3H^2}$. For $\Omega_M > 1$ the spatial curvature is positive, $k = +1$, and the evolution equation for $a(t)$, that is the Friedmann equation 12.9 with $\Lambda = 0$, shows that the universe will inevitably collapse back down to the condition $a = 0$, while for $\Omega_M < 1$ the spatial curvature is negative, $k = -1$, and the evolution equation for $a(t)$ shows that the universe will expand forever as for the $\Omega_M = 1$ case, with the latter then representing the critical value upon which the ultimate destiny of the universe depends. While the Einstein-de Sitter universe with $\Omega_M = 1$ describes the unique spatially flat case with $\rho = \rho_c = \frac{3H^2}{\kappa}$ $\frac{H^2}{\kappa}$ for a given $H(t)$ at any cosmic time t, for the spatially non-flat cases with $k = +1$ and $k = -1$ there is a continuous range of solutions with $\rho > \rho_c$ and $\rho < \rho_c$ respectively, for any given values of ρ_c and t.

At the present epoch the density parameter for ordinary baryonic matter alone, which is largely readily visible in the form of galaxies of stars and clouds of dust and gas, is observed to have a value of $\Omega_{B_0} = 0.050 \pm 0.002$ [44] which, being much less than unity, would imply that we inhabit a universe with spatial curvature $k = -1$ if such matter were the sole source of gravitation. The contribution of 'dark matter', which is needed to explain the rotation dynamics of stars within galaxies as well as the dynamics of individual galaxies within clusters, is found to be larger with density parameter $\Omega_{D_0} = 0.265 \pm 0.011$ [44], implying a total matter density parameter at the present epoch of $\Omega_{M_0} \simeq 0.315$. However this total is still inconsistent with independent observations, namely of the angular anisotropy of the CMB radiation, which suggest that the universe is very close to being spatially flat with $k = 0$.

Since Ω_{M_0} falls well short of the total value needed to account for the observed spatial flatness, and since this quantity is only sensitive to gravitating matter associated with clustering up to the largest scales probed, a significant contribution from relativistic particles or a vacuum energy term is implied. With a negligible contribution from the CMB radiation itself of $\Omega_{R_0} \simeq 5.5 \times 10^{-5}$ [44] (and with an even smaller contribution predicted for relic neutrinos from the Big Bang) we continue to assume $p = 0$ but allow the cosmological constant Λ to take a finite value in equation 12.9, which can be divided by $H^2 = \frac{\dot{a}^2}{g^2}$ $\frac{a^2}{a^2}$ and rearranged in the form:

$$
\Omega_M + \Omega_\Lambda = 1 + \frac{k}{\dot{a}^2} \tag{12.16}
$$

with
$$
\Omega_M = \frac{\kappa \rho}{3H^2}
$$
 and $\Omega_{\Lambda} := \frac{\Lambda}{3H^2}$ (12.17)

Analysis of the Hubble diagram for distant supernovae of type SN Ia independently determines a value of $\Omega_{\Lambda_0} = 0.685 \pm 0.017$ [44]. Hence, as can be seen from equation 12.16, the empirical observations of $\Omega_{M_0} + \Omega_{\Lambda_0} \simeq 1.000$ and of spatial flatness consistent with $k = 0$ from the CMB anisotropy are in excellent agreement.

However while these observations are mutually consistent it appears coincidental that the contributions from Ω_{M_0} and Ω_{Λ_0} are of the same order of magnitude at the present epoch. In particular from equation 12.17 and the empirical values of the density parameters the present overall matter density in the universe is approximately half that of the vacuum energy, with $\rho_0 \simeq \frac{1}{2}\Lambda/\kappa$. Since ρ was much larger in the earlier universe and is ever decreasing into the future, and since the matter density ρ describes matter which is heavily clumped into clusters of galaxies and the stars within, while Λ/κ (which may be generically termed 'dark energy') is of an apparently very different nature, both constant in time and uniformly distributed in space, the approximate coincidence of their present average values, within a factor of two, is notable. It is also observed that within the Ω_{M_0} contribution itself the dark matter content is around five times that of the baryonic matter, which is assumed to be a feature largely independent of cosmic time. An understanding of the origin of the above empirical observations will require a theoretical understanding of the nature of the dark sector itself.

A solution for the large scale cosmic geometry must also be consistent with equation 12.10, which can be employed to further analyse the dynamics. Substituting

 \dot{a}^2 $\frac{\dot{a}^2}{a^2}+\frac{k}{a^2}$ $\frac{k}{a^2}$ from equation 12.9 into this second dynamic equation leads directly to the relation:

$$
\frac{\ddot{a}}{a} = -\kappa \left(\frac{\rho + 3p}{6} \right) + \frac{\Lambda}{3}
$$
\n(12.18)

An era of accelerating expansion of the universe, that is with $\ddot{a}(t) > 0$, provides a formal definition of cosmic 'inflation'. From the above equation it can be seen that this is the case for $(\rho + 3p) < 0$ for $\Lambda = 0$, or for $\Lambda > 0$ if ρ and p are relatively small, or some combination of these factors. The dynamics can be described in terms of the 'deceleration parameter', defined as $q := -\frac{a\ddot{a}}{\dot{a}^2}$ $\frac{aa}{a^2}$. Taking the case $p = 0$, using equation 12.17 and dividing equation 12.18 by H^2 the deceleration parameter is found to be related to the density parameters as:

$$
q = \frac{\Omega_M}{2} - \Omega_\Lambda \tag{12.19}
$$

At the present epoch, as for the previous several billion years, with the value of $\frac{\Omega_M}{2}$ < Ω_{Λ} and $q < 0$ the expansion of the universe is accelerating, and at an increasing rate. In contrast during the first few billion years of cosmic evolution the values were such that $\frac{\Omega_M}{2} > \Omega_{\Lambda}$ with $q > 0$ and the rate of expansion of the universe was, temporarily, slowing down – as sketched in figure 12.2 .

Figure 12.2: A qualitative depiction of the evolution of the cosmological scale factor $a(t)$ as a function of cosmic time t, up to and beyond the present epoch t_0 .

In the future as the matter density $\rho(t)$ and the value of Ω_M decrease with the expanding universe the cosmological constant Λ will increasingly dominate the large scale evolution of the cosmos. With $p = 0$, $k = 0$, $\Lambda > 0$ and taking the limit $\rho \to 0$ equation 12.9 becomes simply $\dot{a}^2 = \frac{1}{3}\Lambda a^2$. Hence if such a cosmic epoch begins at time $t = t_\Lambda$ the scale factor increases as $a(t) = \exp\left(\sqrt{\frac{1}{3}\Lambda} (t - t_\Lambda)\right) a(t_\Lambda)$, which is also consistent with equation 12.18. For a cosmology entirely determined by a cosmological constant then $t_\Lambda = 0$ and, setting $a(0) = 1$ for this scenario, this describes the de Sitter model with line element:

$$
d\tau^2 = dt^2 - e^{At} d\Sigma_{k=0}^2
$$
 (12.20)

where $A = 2\sqrt{\frac{1}{3}\Lambda}$ is a constant and $d\Sigma_{k=0}^2$ represents the 3-dimensional spatial part of the line element in equation 12.5 for the spatially flat case with $a^2(t) = e^{At}$. Since if $\Lambda = 0$ ordinary matter on the right-hand side of equation 12.1 does not yield a solution in the form of equation 12.20 this special case for FLRW cosmology with an exponential expansion factor was originally considered to represent a matterless vacuum with Einstein equation $G_{\mu\nu} + \Lambda g_{\mu\nu} = 0$, which is equivalent to the Ricci tensor being constrained to the form $R_{\mu\nu} = \Lambda g_{\mu\nu}$ with constant Λ .

However, since the 'vacuum' Einstein equation can be written with the cosmological term on the right-hand side as $G_{\mu\nu} = -\Lambda g_{\mu\nu} \equiv -\kappa T_{\mu\nu}(\Lambda)$ the exponential expansion observed for our universe at the present epoch is generally attributed to 'vacuum energy' or 'dark energy', in contrast to 'dark matter' and as alluded to in the discussion following equation 12.17 above. By direct comparison with equation 12.2 the object $T_{\mu\nu}(\Lambda)$ may be interpreted as a non-standard form of energy-momentum for a 'fluid' with an energy density $\rho_{\Lambda} = \Lambda/\kappa$ and pressure $p_{\Lambda} = -\rho_{\Lambda} = -\Lambda/\kappa$ which are constant in time as well as space even as the universe evolves. This substitution, replacing Λ with effective values of ρ_{Λ} and p_{Λ} , can also be applied directly in the evolution equations 12.9 and 12.10, from which the accelerating expansion may be deduced via equation 12.18 since $(\rho_{\Lambda} + 3p_{\Lambda}) < 0$ for $\Lambda > 0$ (consistent of course with employing $Λ$ itself directly in equation 12.18). The quantities $ρ_Λ$ and $p_Λ$ remain constant in time even if an energy-momentum tensor with $T_{\mu\nu}\neq 0$ for ordinary matter is included in the field equation 12.1, as is the case in equation 12.3.

The above de Sitter model of equation 12.20 was introduced in 1917 and originally thought to represent a static solution until it was shown how test particles would fly apart from each other in such a universe. In the same year a truly static universe model was proposed by Einstein, also with $p = 0$ and $\Lambda > 0$ as for the de Sitter model but in this case with a finite matter density ρ tuned to solve equations 12.9 and 12.10 with the constraint $\dot{a} = \ddot{a} = 0$. The solution for the Einstein model requires a positive curvature $k = +1$ and a constant density $\rho = \frac{2}{\kappa} \Lambda$ for ordinary matter fixed for all time as the universe neither expands nor contracts.

From an observational point of view an initial data set of measurements of significant redshifts for a number of nebulae was observed by V.M. Slipher as early 1917, that is the same year the above models were proposed. In the early 1920s the brightest nebulae were resolved into stars, including those of the Cepheid type allowing Edwin Hubble to estimate their distances out to several million light-years. At this time it was established that the nebulae are in fact further distant galaxies comparable in size to our own and the visible scale of the cosmos was vastly augmented. That galaxies are receding away from our own Milky Way with velocities proportional to their distance from us, consistent with equation 12.12, was first discovered by Hubble in 1929.

Following the empirical conclusion from the 1920s that the universe is expanding and Eddington's theoretical observation in 1930 that the static Einstein model is unstable a policy of dropping the cosmological constant term Λ was generally adopted. This led in particular to the Einstein-de Sitter model of 1932, with $\Lambda = 0, k = 0, p = 0$ and $\rho = \rho_c = \frac{3H^2}{\kappa}$ $\frac{H^2}{\kappa}$ evolving in time, as described above following equation 12.14. As described alongside equations 12.16 and 12.17 observations in cosmology dating from the 1990s have resulted in the reintroduction of a $\Lambda > 0$ term, which is now incorporated into the standard model of cosmology. (The present domination of this term over the matter density, composed of both baryonic and dark matter, with $\rho_0 \simeq \frac{1}{2\epsilon}$ $\frac{1}{2\kappa}\Lambda$ contrasts with above finely balanced Einstein universe for which $\rho = \frac{2}{\kappa} \Lambda$). While the impact of the cosmological constant on the more recent evolution of the universe is clearly visible in figure 12.2 the much earlier radiation dominated period, while also forming a key part of the standard model, in spanning a period of less than 50,000 years after the Big Bang is far too brief to feature on the linear scale adopted in this figure. In the following section we motivate and review some of the theoretical ideas applied to the yet far earlier universe.

12.3 Inflationary Theory

The redshift z observed for distant galaxies by Hubble, and through to present day observations now extending out across several billion light-years, is defined by the relation:

$$
1 + z \equiv \frac{\lambda_0}{\lambda_e} = \frac{a(t_0)}{a(t_e)}
$$
\n(12.21)

where t_e is the cosmic time of the emission of radiation from a distant galaxy with wavelength λ_e (which can be deduced from well-known patterns of spectral lines) and t_0 is the present cosmic time at which we detect the radiation and measure the wavelength to be λ_0 in our galaxy. With the value of $z_0 = 0$ for the present epoch and adopting the convention $a(t_0) = 1$ there is a simple relationship between the redshift z at an earlier epoch and the corresponding scale factor $a(t_e)$ at that time. Since for our universe $a(t)$ is an ever increasing function of time, as can be seen in figure 12.2, the value of the redshift z can be used to label the earlier epochs of our expanding universe. Hubble's observations of a positive redshift are explained via equation 12.21 by the simple fact that $a(t_e)$ was smaller in the past, while an increasing value of $a(t)$ at any given time t implies a positive value for the Hubble parameter defined in equation 12.12.

This cosmological redshift does not arise from the Doppler effect, which only strictly applies in a local or extended *flat* region of spacetime, but from the passage of light through a curved 4-dimensional spacetime. As a further example the non-Euclidean geometry of spacetime also accounts for the gravitational redshift or blueshift resulting from the propagation of light away from or towards, respectively, a massive object, which is also an effect of general relativity.

Although the 4-dimensional geometry of our universe is far from (pseudo-) Euclidean, the observations described in the previous section indicate that the 3 dimensional spatial hypersurfaces at any given cosmic epoch t appear to be very flat. While the hypothetical Einstein universe was found by Eddington to be balanced precariously, as alluded to at the end of the previous section, there is also an apparent instability concerning the state of the universe we actually observe. If the value of the total density parameter Ω is not exactly equal to one in a matter or radiation dominated universe, such as we have described for the first few billion years of our own cosmos, this value will diverge away from unity as the universe evolves. Taking equation 12.15, generically replacing Ω_M by a density parameter Ω and with $H^2 = \frac{\kappa \rho}{3\Omega}$,

as for the form of equations 12.17 for example, leads directly to the relation:

$$
\left(\Omega^{-1} - 1\right)\rho a^2 = -\frac{3k}{\kappa} \tag{12.22}
$$

from which different conclusions may be reached depending on the equation of state for the apparent matter density ρ , given that the right-hand side of this expression is a constant. In particular for an expanding FLRW universe that is matter dominated or radiation dominated the quantity ρa^2 decreases with cosmic time in proportion to a^{-1} or a^{-2} respectively, as can be seen from the columns of table 12.1. Hence it can be seen from equation 12.22 that a value of $\Omega \neq 1$ will diverge further from unity as such a universe evolves.

That is any small deviation of the density parameter Ω from the value of unity at an earlier epoch with a large redshift z will have been greatly amplified by the present day, such that in going back to the extreme case of the Planck epoch of $t \approx 10^{-43}$ seconds after the Big Bang an apparent fine tuning of the density parameter to about 1 part in 10^{60} is required in order to be consistent with the present day observation of spatial flatness for the universe $([5]$ p.323). The need for an explanation of this precise tuning of the initial spatial flatness condition arising out of the Big Bang is known as the 'flatness problem'.

That the 4-dimensional geometry can be highly curved even for a spatially flat cosmology with $k = 0$ is particularly evident in the early universe. For the matter dominated case it can be seen by substituting terms from equations 12.9 and 12.10 into equation 12.8 that the scalar curvature of the spacetime is simply $R = \kappa \rho$ (as would be expected from the paragraph following equation 5.35). Hence as $t \to 0$, in principle to an epoch even earlier than the Planck time, with the scale factor $a(t) \rightarrow 0$ while $\rho \rightarrow \infty$ the scalar curvature R diverges to infinity at what is referred to as the 'initial singularity' at $t = 0$. It is sometimes noted that the standard cosmological model hence predicts the paradoxical origins of the universe in such an initial singularity, studied by S.W. Hawking, G.F.R. Ellis and R. Penrose around 1965–70, a point at which general relativity, which governs the model itself, breaks down. However any conclusions drawn from the structure of gravitation at the Planck scale are inevitably uncertain given the as yet unknown role of quantum phenomena in such an extreme environment. As described in section 11.4 for the present theory gravitation itself is not quantised and hence in principle the Planck scale will be of less significance and not represent a barrier to further extrapolation to arbitrarily early times, as will be considered in section 13.2.

It is also informative to write the Robertson-Walker line element of equation 12.5 with the cosmic time coordinate t transformed to a conformal time parameter $\eta = \int_0^t$ dt′ $\frac{dt'}{a(t')}$ as:

$$
d\tau^2 = a^2(\eta) \left[d\eta^2 - d\Sigma_k^2 \right] \tag{12.23}
$$

where $d\Sigma_k^2$ represents the spatial part of the line element inside the square brackets of equation 12.5 with $k = +1, 0$ or -1 . Hence by adopting the conformal time coordinate η in equation 12.23 the scale factor $a(\eta)$ can be seen as a special case of a conformal transformation, which more generally takes the form $g_{\mu\nu}(x) \to f(x)g_{\mu\nu}(x)$ where $f(x)$ is an arbitrary real function of spacetime (a very different example of which was considered in equation 11.13). For the case $k = 0$ the Robertson-Walker line element is hence related to a flat 4-dimensional spacetime via a conformal transformation. It can also be shown, using a further suitable coordinate transformation, that the geometry for each of the $k = \pm 1$ cases is also conformally flat ([5] p.71). Hence for all FLRW models the 4-dimensional geometry of the universe, with the metric of equation 12.5, is conformally flat, which implies the vanishing of the Weyl curvature tensor, introduced before equation 3.69, that is $C_{\rho\sigma\mu\nu}(x) = 0$, even though components of the Ricci curvature $R_{\mu\nu}(x)$ part of the Riemann tensor may attain arbitrarily large values in the very early universe.

The initial singularity of the Big Bang is a spacelike boundary of spacetime in our distant past, represented by the horizontal wiggly line in the conformal diagram of figure 12.3. In such a diagram all null-rays, that is with proper time line element $d\tau^2 = 0$, are drawn at 45° and hence the causal properties of the spacetime are made apparent. The vertical axis of such a diagram is linear in the conformal time η with the horizontal axis representing comoving coordinate distances $\Delta\Sigma$, consistent with equation 12.23. In figure 12.3 epochs on the vertical axis are labelled by the cosmic time t , although of course not to scale, and intervals of the horizontal axis at any given epoch can be converted to physical proper distances $a(t)\Delta\Sigma$, as described before equation 12.12. A ray of light emitted at time $t = t_e$ and reaching us now at $t \equiv t_0$ will have travelled the comoving coordinate distance

$$
\Sigma_p(t_0, t_e) = \Delta \eta = \int_{t_e}^{t_0} \frac{dt'}{a(t')}
$$
 (12.24)

where $\delta t'/a(t')$ is the coordinate distance traversed in a small interval of cosmic time $\delta t'$. Hence any signal emitted beyond this distance at time t_e will not have been able to reach us yet and hence in turn $\Sigma_p(t_0, t_e)$ is termed the 'particle horizon'. For any given t_e the particle horizon grows with time $t = t_0$ from the perspective of the observer b in figure 12.3. The particle horizon can also be defined as the proper distance $R_p(t_0, t_e) = a(t_0) \Sigma_p(t_0, t_e)$ on the 3-dimensional spatial hypersurface at the time t_0 , that is $R_p(t_0, t_e) = a(t_0) \Delta \eta$, where $\Delta \eta$ is the conformal time elapsed between $t = t_e$ and $t = t_0$.

At the present epoch t_0 the largest particle horizon $R_p(t_0)$ corresponds to signals emitted at the time of the Big Bang. Setting $t_e = 0$ the integral in equation 12.24 converges provided the equation of state is such that $\rho(t)$ decreases at least as fast as $a^{-2}(t)$, as it does for a matter or radiation dominated universe as seen in table 12.1. For various cosmological models the particle horizon, obtained from equation 12.24, is generally greater than t_0 itself since $a(t)$ tends to be smaller for earlier times $t < t_0$. For a radiation dominated universe the particle horizon has a value of $R_p(t_0) = 2t_0$, while for the matter dominated case the value is $R_p(t_0) = 3t_0$. (For the Einstein-de Sitter model with $k = 0$ the age of the universe can be expressed as $t_0 = \frac{2}{3H}$ $\frac{2}{3H_0}$ and the particle horizon is twice the Hubble radius, $R_p = 3t_0 = \frac{2}{H}$ $\frac{2}{H_0} = 2R_H$, with the latter defined in equation 12.4.)

For our own universe the present particle horizon is determined to be $R_p(t_0) \simeq$ 46 billion light-years, which is greater than $3t_0$, where $t_0 \approx 13.8$ billion years is the age of the universe, owing to increasing impact of the cosmological Λ term at the present epoch. The proper distance $R_p(t_0)$ represents the edge of the observable universe in terms the present distance to objects on the 3-dimensional spatial hypersurface at

Figure 12.3: Conformal diagram depicting the past light cone from our present location b at cosmic time $t = t_0$ extending back to the Big Bang singularity at $t = 0$. The recombination era $t = t_{\text{rec}} \left(\simeq t_0/37,000 \right)$ is also indicated. As a plot of conformal time η versus comoving coordinate distance Σ the diagram is a 2-dimensional representation of a 4-dimensional spacetime.

cosmic time $t = t_0$, not of course as we actually might see them via light emitted in the distant past. While $R_p(t_0, t_e) = a(t_0) \Sigma_p(t_0, t_e)$ is the present particle horizon for observing events from time $t = t_e$, the proper distance to such an event on the horizon at the time of signal emission was $a(t_e)\Sigma_p(t_0,t_e)$. For comparing particle horizons at different epochs comoving coordinate distances $\Delta\Sigma$, that is intervals of the horizontal axis in conformal diagrams, will be move convenient, as we describe in the following.

As well as the Big Bang at $t = 0$ and the present era $t = t_0$ the time of 'recombination' $t = t_{\text{rec}}$ is also labelled in figure 12.3. This is the epoch around 372,000 years after the Big Bang, with a redshift of $z \approx 1090$ and as the temperature dropped below around 4,000 K, during which the residual electrons, which had not annihilated with positrons, combined with protons and other light nuclei to form neutral atoms, mainly hydrogen and helium (the name 'recombination' is somewhat inaccurate as this process is an initial combination of such objects, unless thought of as a return to charge neutral states in a different form to that at $t = 0$). Since there is only an extremely small interaction between an external electromagnetic field and neutral atoms this also marks the era of decoupling between radiation and matter alluded to after equation 12.3. Photons from this decoupling epoch have effectively been propagating freely since $t = t_{\text{rec}}$, relatively early in the 13.8 billion year history of the universe, until detected in the present as the observed CMB radiation now redshifted to a temperature below 3 K.

From our perspective photons composing the CMB radiation were emitted from anywhere on the 2-sphere of our past light cone in 4-dimensional spacetime at the time $t = t_{\text{rec}}$. Two points u and v on the continuous surface of a 2-sphere can be arbitrarily close together, unlike the points u and v in figure 12.3 on the past light cone of this 2dimensional representation of spacetime. If the comoving coordinate distance between u and v at $t = t_{\text{rec}}$ is greater than twice the particle horizon $\Sigma_p(t_{\text{rec}})$ then the two spacetime points have never been in causal contact. Hence from our perspective b , with both u and v observed on our particle horizon $\Sigma_p(t_0, t_{\text{rec}})$, there is no reason to expect a homogeneity of physical quantities such as the CMB temperature as measured and compared for such regions u and v which have not been in causal contact with each other. In fact the particle horizon at the recombination era $\Sigma_p(t_{\text{rec}})$ only subtends of order 1[°] in the sky from our present perspective *b* on Earth. The difficulty in contriving an assumption of homogeneity as an initial condition of the hot Big Bang to account for the observed uniformity of the CMB temperature to within 1 part in 10^5 over all angles of the sky is known as the 'horizon problem'.

In place of postulating homogeneous initial conditions across causally separated spatial regions of the very early universe the only means by which the temperatures at u and v might be related through a process of thermalisation is to arrange for the possibility of causal contact in their past. This requires a mechanism through which the Big Bang epoch effectively retreats back further below the recombination era in the conformal diagram of figure 12.3, as demonstrated in figure 12.4. This in turn can be achieved by a sufficient rescaling of proper spatial distances with $a(t_{\text{inf}}-\epsilon) \ll a(t_{\text{inf}}+\epsilon)$, where ϵ may be a very short time interval, effectively 'miniaturising' 3-dimensional space during the epoch $t < t_{\text{inf}}$. In this case the horizontal displacements in figure 12.4 labelled by comoving coordinate intervals $\Delta\Sigma$ now represent much shorter physical proper distances $a(t)\Delta\Sigma$ for $t < t_{\text{inf}}$ and a given null-ray propagating for a given cosmic time interval Δt occupies a somewhat larger portion of the vertical axis which is linear in conformal time intervals $\Delta \eta \sim \frac{\Delta t}{a(t)}$. Hence the Big Bang epoch at $t = 0$ is pushed back in the conformal diagram to accommodate this rescaling. Hence in turn the comoving coordinate distance traversed by null-rays in a fixed cosmic time interval during this early epoch before $t = t_{\text{rec}}$ can in principle comfortably encompass the present particle horizon $\Sigma_p(t_0, t_{\text{rec}})$ at $t = t_0$ for signals emitted at $t = t_{\text{rec}}$ (see for example [26] pp.744–747).

Figure 12.4: Conformal diagram depicting a similar cosmic history as figure 12.3 with the same three values of $t = 0$, $t = t_{\text{rec}}$ and $t = t_0$ but with the addition of a further epoch $t = t_{\text{inf}}$ during which the scale factor $a(t)$ is 'inflated' by an enormous degree in a short period of cosmic time.

Evolving forwards in time from the Big Bang the rapid expansion of the universe scale factor $a(t)$ at the epoch t_{inf} , which in principle solves the horizon problem, is termed 'inflation', as a particular case of an accelerating expansion described generically after equation 12.18. The question then still remains regarding the physical mechanism behind such a radical transformation of the spacetime geometry at that very early epoch. Guided by the de Sitter model with the line element of equation 12.20 describing an exponential expansion with scale factor $a(t) \propto \exp\left(\sqrt{\frac{1}{3}\Lambda} t\right)$ one way to achieve inflation is with a very large, but only temporarily active, cosmological term of the form $\Lambda g_{\mu\nu}$ in Einstein's field equation 12.1.

On introducing a new scalar field $\varphi(x)$ (which is unrelated to the scalar Higgs field $\phi(x)$ of the Standard Model of particle physics described in section 7.2) a false vacuum state obtained for a suitable potential $V(\varphi)$ can model the effect of a cosmological term via an energy-momentum tensor $T_{\mu\nu}$ with a term of the form $V(\varphi)g_{\mu\nu}$ (such as in equation 12.25 below). That is, a potential $V(\varphi, T)$, as a function of the field $\varphi(x)$ and temperature $T(x)$, may be contrived such that the high temperature vacuum state $\varphi = 0$ becomes a 'false vacuum' as the universe achieves a 'supercooled' condition below a certain critical temperature T_c in the very early universe. The phase transition to the new true vacuum state with $\varphi \neq 0$ for $T < T_c$ may involve either quantum mechanical tunnelling through an intermediate potential barrier ('old inflation') or a gradual roll down a potential slope ('new inflation'). In either case the potential function $V(\varphi, T)$ may be suitably contrived in order that the true vacuum is not immediately attained and the energy of the false vacuum state dominates the cosmological evolution equations for a brief period of time. This cosmic time period of $t_{\text{inf}} \sim 10^{-35} - 10^{-32}$ seconds can be long enough for the scale factor $a(t)$ to increase by a factor of $\sim 10^{30}$ or more, effectively solving the horizon problem by the rapid inflation of a small homogeneous region of the very early universe (see for example [5] chapter 11).

While the de Sitter model of equation 12.20 assumes a spatially flat universe with $k = 0$, the evolution of the scale factor $a(t)$ resulting from a cosmological term $\Lambda q_{\mu\nu}$ in the field equation can also be determined for the cases of $k = \pm 1$ with spatial curvature. It is found that for $k = +1, 0$ and -1 the scale factor evolves as $a(t) \propto \cosh\left(\sqrt{\frac{\Lambda}{3}}\right)$ $\frac{\Lambda}{3} t$), $\exp\left(\sqrt{\frac{\Lambda}{3}}\right)$ $\frac{\Lambda}{3} t$) and $\sinh\left(\sqrt{\frac{\Lambda}{3}}\right)$ $\frac{\Lambda}{3} t$ \setminus respectively, and hence the $k = \pm 1$ solutions in time converge towards the de Sitter solution with $k = 0$ and constant Hubble parameter $H(t) = \sqrt{\frac{\Lambda}{3}}$ $\frac{\Delta}{3}$ ([5] p.326). This convergence towards a density parameter Ω of unity can also be seen from equation 12.22 since the equation of state for a cosmological term implies that $\rho a^2 \propto a^2$, as can be seen from table 12.1, which hence rapidly increases during inflation, driving $\Omega \to 1$. Hence during inflation solutions for $a(t)$ with finite spatial curvature rapidly approach the purely exponential expansion solution with $k = 0$, that is the de Sitter model for a flat universe with the $\Lambda g_{\mu\nu}$ term simulated by the energy of the false vacuum during the inflationary period. The brief inflationary era t_{inf} described in the previous paragraph, and depicted in figure 12.4, is sufficient to suppress any non-zero spatial curvature by a factor of around 10⁶⁰ or more, hence in principle solving the flatness problem described after equation 12.22, in addition to solving the horizon problem.

Inflationary theory was initially proposed by Alan Guth in 1980, precisely

to address the horizon problem while also accounting for the flatness problem. In fact the strong bias towards spatial flatness is sometimes considered to have been a successful *prediction* of the theory. The hypothetical period of inflation at t_{inf} drives the total density parameter Ω extremely close to unity in the early universe such that the subsequent radiation dominated era of thousands of years and matter dominated era of billions of years have been insufficient to prise the value of Ω away from the value of one, as described following equation 12.22, to any measurable degree. The more recent and increasingly dominant effect of the apparently presently active cosmological term $\Lambda g_{\mu\nu}$ is again tending to bind the density parameter yet closer to unity, although this effect has thus far been too weak to account for the observation of spatial flatness without the much earlier and much more dramatic inflationary epoch.

However unlike the cosmological term $\Lambda g_{\mu\nu}$ which accounts for the present day relatively pedestrian accelerating expansion of the universe the much earlier period of rapid inflation is required to terminate, and such a change in conditions is generally ascribed to a phase transition as introduced above. The original 'old inflation' model employed a first order phase transition via quantum tunnelling from the false to the true vacuum once the temperature had dropped sufficiently to allow penetration through the potential barrier. However the quantum nature of the transition results in bubble formation and corresponding large inhomogeneities that are not observed. This 'graceful exit problem' can be solved by 'new inflation' which ends via a transition from the false vacuum at a local maximum in the potential at $\varphi = 0$, that is through a second order phase transition, which proceeds more nearly simultaneously throughout the universe. An almost flat potential around $\varphi = 0$ can result in a 'slow roll' down to the true vacuum at the potential minimum, still allowing sufficient time for a dramatic inflationary expansion.

Amongst a range of inflationary models proposed 'chaotic inflation' in principle also solves the graceful exit problem. In this model the potential of the scalar field can take a much simpler form such as $V(\varphi) = m^2 \varphi^2$ or $V(\varphi) = \lambda \varphi^4$ with a single minimum at $\varphi = 0$. Under a large range of possible initial conditions in the primordial chaos in some regions the value of $\varphi(x)$ may be far from the minimum. Such a value, required to be essentially uniform over a region of space of order the present day Hubble radius, can stimulate an inflationary period. A large inflation factor is possible provided that the constant λ for example is chosen such that the potential function is sufficiently shallow to allow a sufficiently delayed roll down to the true vacuum value at $\varphi = 0$. As the true vacuum is attained and inflation ends our observable universe is contained within a single bubble, one of many resulting from the initial chaotic conditions. Even if the scalar field φ begins with a value close to the minimum at zero quantum fluctuations can drive this value further from the minimum resulting in a self-sustaining 'stochastic inflation', or even motivating consideration of an 'eternal inflation' model.

For any of the above inflationary models an energy-momentum tensor can be derived from a standard Lagrangian for a scalar field, namely $\mathcal{L} = \frac{1}{2}$ $\frac{1}{2}\partial_{\mu}\varphi\partial^{\mu}\varphi-V(\varphi)$ including a kinetic as well as the potential term, via Noether's theorem as described for equation 3.102 (some care is needed with the interpretation of translation invariance since here we are clearly *not* dealing with a globally flat Minkowski spacetime, however equation 3.102 may be applied for sufficiently small spacetime regions by the strong equivalence principle described in section 3.4 and then generalised for the result below on replacing $\eta_{\mu\nu}$ by $g_{\mu\nu}$) leading directly to ([5] p.329):

$$
T_{\mu\nu} = \partial_{\mu}\varphi \,\partial_{\nu}\varphi - \frac{1}{2}\partial_{\rho}\varphi \partial^{\rho}\varphi \,g_{\mu\nu} + V(\varphi)g_{\mu\nu} \tag{12.25}
$$

In addition to the cosmological term for a temporarily finite (and uniform at least over the spatial extent of the observable universe) value $V(\varphi) \equiv \frac{\Lambda}{\kappa}$, with an effective equation of state $p_{\Lambda} = -\rho_{\Lambda} = -V(\varphi)$, driving the exponential expansion, there are also kinetic terms in the derivatives of the scalar field $\varphi(x)$. An equation of motion for $\varphi(x)$ can be derived as the Euler-Lagrange equation for the stationarity of the action $S = \int \mathcal{L}\sqrt{|g|}d^4x$ which, since the metric $g_{\mu\nu}(x)$ incorporates the scale factor $a(t)$, is found to include a Hubble drag term of the form $H\dot{\varphi}$ ([5] p.331).

If after the Planck time the universe is initially radiation dominated then as the temperature drops below the critical temperature T_c inflation begins to dominate and the radiation is rapidly redshifted. During the vacuum driven expansion the universe is essentially devoid of matter and radiation, with the scalar field φ completely dominating towards the end of inflation, however any coupling between φ and matter fields leads to a further drag term in the equation of motion for φ . As the minimum of $V(\varphi)$ is approached the dynamic equations drive rapid oscillations, which are dampened by the drag terms. This in turn fuels a reheating in the post-inflation era as the vacuum energy is converted into interacting particles, including the familiar states of the Standard Model. This period of transition to essentially zero vacuum energy, in which the energy is transferred from the scalar field φ to ordinary matter and radiation via their mutual interactions, may also be the time during which any mechanism that generates an asymmetry between matter and antimatter, as still manifestly observed today, may act. The origin of dark matter might also turn out to be associated with the termination of inflation. This epoch then merges into the beginning of the radiation and then matter dominated FLRW periods of the standard cosmological model as described in the previous section, with the initial conditions set by the inflationary expansion.

In de Sitter spacetime, as for that of inflation, the event horizon (which is distinct from the particle horizon) is of finite size, as for the case of back holes. This means that the conditions for producing Hawking radiation, as alluded to towards the end of section 11.4, are also present during inflation. In turn the possibility arises that quantum fluctuations can become frozen into residual classical deformations in the latter stages of inflation. In turn these classical fluctuations will modulate the density of the radiation and matter produced at the end of inflation, seeding the evolution of large scale structure as eventually manifested in galactic formations. Similar fluctuations during the inflationary epoch are also predicted to generate a background of gravitational waves which still propagate today and which, although being much more difficult detect, are in principle observable through the large scale CMB anisotropies which may provide a signature for the metric distortions of the gravity waves.

A significant degree of fine tuning is required for any model of inflation based on the properties of a postulated scalar field $\varphi(x)$ in order to achieve a match with a range of empirical observations, which is somewhat unsatisfactory since inflationary theory was designed to avoid the necessary fine tuning as initially implied by the horizon problem and flatness problem. There is also no understanding of the origin of the vast difference between the magnitude of the effective cosmological term associated with inflation due to $V(\varphi)$, which is of $O(10^{-10})$ in natural units, and apparent cosmological constant Λ of the present epoch, which is of $O(10^{-120})$ in natural units. Indeed the unaccounted for magnitude of the latter number itself, the 'cosmological constant problem' is one of the biggest puzzles in physics, as already alluded to briefly at the end of section 4.1.

A further significant issue regarding the standard model of cosmology, which is not addressed by inflation, relates to the origin of the very special conditions of the Big Bang in that the entropy of the early universe must have apparently been extremely low, despite the high degree of thermalisation achieved for the degrees of freedom of the electromagnetic field. The degrees of freedom of the gravitational field may be described by the Weyl tensor ([26] section 28.8), although both the Ricci curvature and Weyl curvature parts of the Riemann tensor exhibit the effects of gravity. The Weyl curvature and its distorting tidal effect tend to increase as matter gravitationally clumps into dense regions, diverging to infinity in the neighbourhood of a black hole. The entropy associated with a black hole is correspondingly extremely high, attaining values much higher than that associated with ordinary thermal entropy. On the other hand, as described following equation 12.23, in the idealisation of the FLRW cosmological models the spacetime is conformally flat with zero Weyl curvature. This suggests that if the universe originates in a state very close to an FLRW model the initially low entropy may correlate with the very low Weyl curvature, both of which then tend to increase as matter progressively clumps together as the universe evolves.

More generally the 'Weyl Curvature Hypothesis', proposed by Roger Penrose in 1979 ([26] section 28.8), asserts that $C_{\rho\sigma\mu\nu}(x) = 0$, or is at least very close to zero, as a constraint on the initial singularity. Hence the universe shares at least this property, of conformal flatness, with the FLRW models in the early stages. (In principle this constraint might be further augmented by the condition $k = 0$ as the universe evolves into a spatially flat model due to a subsequent period of inflation). This hypothesis of zero Weyl curvature for the initial singularity of the Big Bang is then in stark contrast to the situation for the terminal singularities of black holes as alluded to above.

This very special condition of the Big Bang represents an enormous constraint of low entropy on the initial conditions which in turn provides a suitable point of departure for the second law of thermodynamics. Gravitation, in comparison to all other fields, hence appears to have had a very special status, aloof from thermalisation in the Big Bang, with the second law of thermodynamics only later exercised through the gravitational degrees of freedom. While inflation, as described for figure 12.4, provides the breathing space for ordinary matter, including the electromagnetic field, to reach thermal equilibrium in the aftermath of the hot Big Bang, the question remains to explain why gravitation should apparently be treated in such a radically different manner to the other forces of nature. The theory presented in this paper may shed some light on these questions since, as discussed in the previous chapter, here the gravitational field itself is not quantised and is hence different from all other fields in this respect.

Further, while for a range of given initial conditions inflationary theory can solve the horizon problem, which was introduced in figure 12.3, by opening up a suitable spacetime volume to allow points such as u and v to exhibit the same temperature through thermalisation, as described for figure 12.4, the structure of these diagrams indicates that there may be a more fundamental difficulty with this picture. Namely, since *any* two different points such as x and y on the spacelike surface of the initial singularity at $t = 0$, as indicated in figure 12.3 for example, have clearly never been in causal contact with each other it is difficult to conceive how the Big Bang could be effectively 'triggered' simultaneously across this potentially infinite 3-dimensional hypersurface. This observation applies even if initial properties, such as the temperature, are very different at x and y . It also applies in exactly the same way for figure 12.4 and inflation is of no help in addressing this 'start-up problem'.

On the other hand if the Big Bang can be considered as a 'spacelike event', encompassing the points of a large region of the initial spatial hypersurface, then there seems no reason to suppose that the simultaneous 'cause' of the Big Bang at points such as x and y in figure 12.3 might not also 'cause' them to have the same properties such as temperature. Indeed, the notion of a simultaneous start-up along the $t = 0$ spacelike surface which endows different points with *different* properties, implying the application of a range of possible start-up conditions and resulting in an uneven temperature distribution, seems somewhat more contrived. That is, it seems any two points like x and y on the initial singularity must be related in order for the universe to start-up at both, and any solution to this problem may well itself entail a high degree of homogeneity in the very early universe and solve the horizon problem without the need for inflation. A source of later fluctuations and inhomogeneity will then still be needed to account for the origin and formation of the galactic structures seen today.

However, even without the issue of the uncertain role of quantum phenomena under the extreme gravitational conditions of the very early universe, care is needed in the extrapolation to the earliest epoch. For most FLRW models as the cosmic time approaches the moment of the Big Bang $t \to 0$ the scale factor also approaches zero $a(t) \rightarrow 0$, indeed the boundary condition $a(0) = 0$ is adopted for various dynamic solutions, as described before equation 12.12. In this limit any finite comoving coordinate distance $\Delta\Sigma$ corresponds to a vanishing physical proper distance $a(0)\Delta\Sigma = 0$. With the horizontal axes in figures 12.3 and 12.4 representing coordinate distances this naive analysis implies that the observable universe at present came from a physically vanishingly small region of the initial singularity. In turn the initial singularity, represented by the wiggly line in these figures, might perhaps be interpreted as a fully causally connected entity in the limit $t = 0$, amending the strict causal structure of the conformal diagrams in this extreme case.

However, since the spacelike coordinate distances are unlimited in magnitude even as $a(t) \to 0$ any proper distance in the limit $t \to 0$ is obtained as the product of one number in principle approaching infinity with another approaching zero, a situation which approaches the meaningless. Hence rather than speculating upon 'how many angels can dance on the head of a pin', what is really needed is a more complete understanding of what *happens* in the Big Bang, what *causes* it to happen and even why there should be a universe at all.

Chapter 13

A Novel Perspective on Cosmological Structure

13.1 The Dark Sector

Within the context of the present theory the external geometric structure of the world is intimately associated with a subjective perceptual requirement, forged out of a multi-dimensional form of temporal flow expressed as $L(\hat{v}) = 1$, rather than being an apparently arbitrary feature of an objective universe independent of the need for perception. Indeed the specific identification of 3-dimensional spatial expanses with an approximately global SO(3) symmetry would seem to be a somewhat redundant and unnecessary feature of such an inanimate mathematical entity. On the other hand the extent of spatial flatness for the observable universe, as described in the previous chapter, goes far beyond that utilised for perception by sentient beings on the planet Earth.

Further, given the observed Hubble constant of equation 12.13 at the present epoch, in a period of 100 years the fractional change in the scale factor is $\Delta a_0/a_0 \simeq$ 0.7×10^{-8} . Hence on the scale of a human lifetime the Robertson-Walker line element of equation 12.5, for the $k = 0$ case, describes a flat Minkowskian spacetime to within 1 part in 10^8 , with the expanding universe seemingly hanging suspended as a vast spatial expanse through a given human interval of cosmic time. For the horizontal time axis representing a duration of 100 years figure 12.1 would then represent an accurate snapshot of our universe at the present epoch.

However the breakdown of global Lorentz symmetry beyond our 100 year thick slice of the universe is readily observed in the cosmological redshift. This redshift, defined in equation 12.21 and as first observed by Hubble and others and now probing distant galaxies reaching back over billions of years in cosmic time, uncovers the non-Euclidean geometry of the cosmos as summarised by the evolution of the scale factor $a(t)$ depicted in figure 12.2.

The question then is the extent to which the present theory might account for the observations of such large scale structure in cosmology, and the phenomena of the dark sector more generally as summarised in section 12.2, as we shall explore in this section. In section 12.3 it was described how the origin of spatial flatness and the cosmological principle of homogeneity and isotropy, beyond the pragmatism of assumptions employed for FLRW models, can in principle be accounted for by the theory of inflation in the very early universe. In the following section the evolution of the very early universe and the nature of the Big Bang itself will be considered here within the context of the projection of spacetime out of the general form of temporal flow for the present theory. In section 13.3 the extent to which cosmological and other physical parameters might be explicitly constrained by the theory will also be considered.

The pure flow of time s, underlying the multi-dimensional form of temporal flow through $L(\hat{v}) = 1$, is directly related to the proper time τ elapsed from the point of view of any timelike trajectory through spacetime, as described in section 5.3. Time dilation effects for τ , as implied in the metric $g_{\mu\nu}(x)$ such as that for the Schwarzschild solution of equation 5.49, are directly equivalent to those for s. A similar observation applies for the Robertson-Walker metric of equations 12.5 and 12.6 and hence for an idealised galaxy based observer, with constant comoving coordinates $\{r, \theta, \phi\}$, the fundamental time parameter s in being to proportional to τ is in turn equivalent to the cosmic time parameter t . Only in this special case under the assumptions of an FLRW model might s be associated with a preferred universal temporal parameter, namely the cosmic time t , for observers attached to idealised galaxies in the context of such a model.

However, the fundamental temporal flow s itself does not represent a unique universal parameter. In the context of large scale structure a local parameter s, subject to each observer, depends upon the relative motion of the observer with respect to a galaxy or the relative finite peculiar velocity of the galaxy itself, in precisely the same way as the proper time τ in special relativity. Similarly the parameter s will depend upon the location of the observer with respect to a local source of gravity, such as any massive body or even a black hole, again exactly as for the proper time τ , in this case as for general relativity.

The relative time dilations for a community of N observers, each of whom is associated with a personal flow of pure time $s_I \equiv \tau_I$ (for $I = 1...N$, generalising from the case of 'twin A' and 'twin B' described at the end of section 5.3), distributed anywhere in the universe dovetail together in a mutually consistent manner. The particular temporal parameter s_I for a given observer describes the 'fundamental' flow of time underlying the mathematical structure of the multi-dimensional form $L(\hat{\mathbf{v}}) = 1$ through which the physical processes of the universe unfold from the perspective of that observer. In this sense each s_I is a universal temporal parameter, as noted in the discussion of the 'problem of time' in section 11.4 following equation 11.51.

Locally the flow of time $s \equiv \tau$ parametrises the evolution of fields, such as a gauge field $Y(x)$ or fermion field $\psi(x)$ and microscopic quantum phenomena generally, as well as the dynamics of macroscopic entities, such as a dust cloud described by the energy-momentum tensor $T_{\mu\nu} = \rho u_{\mu} u_{\nu}$ or classical matter generally. Either quantum or classical processes may be utilised in the construction of a physical clock which may in turn be employed to *measure* the proper time τ itself and hence observe time dilation effects. With the microscopic quantum properties of matter underlying, and in harmony with, the macroscopic geometry of gravitational phenomena there is no 'problem of time' in this picture, as described in section 11.4, with gravity itself not quantised.

In general relativity local coordinates can always be found such that for any 4-dimensional metric, such as that in equation 12.5, the line element can be expressed through a local Minkowski metric with $d\tau^2 = \eta_{ab} dx^a dx^b$. In the present theory such a local structure *derives* from a 4-dimensional form of temporal flow $ds^2 = \frac{\eta_{ab}}{h^2} dx^a dx^b$, that is equation 5.47 which is equivalent to equation 5.46, that is the expression:

$$
L(\mathbf{v}_4) = h^2 \tag{13.1}
$$

This latter structure is embedded within a higher-dimensional form such as $L(\mathbf{v}_{27}) = 1$ or $L(v_{56}) = 1$ as described in chapter 8 and section 9.2 respectively. It is the higherdimensional form which both sets the normalisation for the temporal flow s and gives rise to a range of many possible solutions for an extended 4-dimensional spacetime, with geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ as described for equation 5.32, incorporating quantum phenomena in the degeneracy of solutions as described in chapter 11.

Hence with the geometry $G_{\mu\nu}(x)$ and the spacetime manifold M_4 itself together drawn out of the structures implicit in $L(\hat{v}) = 1$, with solutions such that $G_{\mu\nu} =$ $f(Y, \hat{v}) \neq 0$ in general, there is no presumption of taking a flat background manifold as a starting point or expectation of obtaining such a Minkowskian spacetime geometry. With the external curvature related to the internal curvature as the symmetries of $L(\hat{v}) = 1$ are projected over M_4 , as conjectured in section 5.1 in comparison with Kaluza-Klein theory, there is a solution with both zero external and zero internal curvature, as implied in equation 5.20 for example with $G_{\mu\nu} = f(Y) = 0$. Even in this case the assumption, as applied in section 5.3, that the value of $L(\mathbf{v}_4) = h^2(x)$ of equation 13.1, as projected out of $L(\hat{\mathbf{v}}) = 1$, is constant throughout spacetime is required to obtain a flat spacetime manifold. The consequences of a variation in the value of $h(x)$, as alluded to at the end of section 5.3, will be considered shortly and will contribute, along with the freedom of the gauge fields and quantum transitions, to a solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ which is non-zero in the general case.

Our a priori predisposition to mentally project a flat background of space and time onto the world in order to perceive objects in it will be consistent with the above mathematical structure provided an effective assumption of $G_{\mu\nu}(x) = 0$ is a sufficiently good approximation at least for the region of the world we locally inhabit. As discussed in section 12.1 this means for example that the local observation of a falling apple can be accounted for in terms of a 'force of gravity' superposed upon an apparently flat arena of space and time, which in practice is both as precise as and much simpler than a full explanation in terms of spacetime curvature. On the global cosmological scale the observed accelerating expansion of the universe not only contradicts the assumption of a flat 'vacuum' geometry, but is also counter-intuitive given the terrestrial bias of associating gravity with a universal force of attraction.

In the present theory the question does not concern what needs to be added to a flat background manifold to produce the effects of terrestrial gravity or the introduction of an apparent vacuum energy to account for the accelerating expansion of the universe,

but rather, in all cases involving gravitation, to ask what is the form of $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ in general. This observation applies to both everyday material objects such as apples and trees and also in the apparent absence of tangible matter in the case of the dark sector for cosmology. This approach in the present theory can be summarised in the following three points (which may be contrasted respectively with the three points listed near the opening of section 12.2 for the standard theory):

- Rather than beginning with a flat spacetime $G_{\mu\nu} = 0$ and then introducing terms such as $T_{\mu\nu}$ or $\Lambda g_{\mu\nu}$ through Einstein's field equation 12.1 as an apparent source of curvature, with matter in some sense actively perturbing the otherwise flat geometry, here the energy-momentum tensor is defined through the Einstein equation, that is $-\kappa T_{\mu\nu} := G_{\mu\nu}$, with the external geometry itself determined through the relation $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ out of the underlying flow of time in the form $L(\hat{v}) = 1$ (as for the example of equation 11.12 and figure 11.1).
- Hence there is no flat spacetime background, acting as a boundary condition, as an apparent consequence of the absence of matter. Originating from our apparently innate bias to conceive of such a flat spacetime as a given entity, this assumption in part underlies the apparent mystery of the cosmological constant, requiring the term $\Lambda g_{\mu\nu}$ to be added to the field equation in a seemingly ad hoc manner to account for the empirical observation.
- On the third point quoted from [74] in section 12.2, a similar interpretation applies here. The identity $T^{\mu\nu}_{;\mu} = 0$ follows trivially from the definition of $T_{\mu\nu} := G_{\mu\nu}$ given the geometric Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$. Indeed, the reverse interpretation of the Einstein equation with $G_{\mu\nu} := T_{\mu\nu}$ implying that matter somehow causes spacetime curvature is more problematic since an independent justification is then required for the relation $T^{\mu\nu}_{;\mu} = 0$ in a general curved spacetime, while the identity $G^{\mu\nu}_{;\mu} = 0$ does not require any such external support.

Regarding the accelerating expansion of the universe the question then boils down to what in the structure of $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ might account for this observation. Ultimately a full understanding will be required for the general macroscopic form for $G_{\mu\nu}(x)$ constructed over a degeneracy of underlying local field exchanges $\delta Y \leftrightarrow \delta \hat{v}$, in principle incorporating some of the machinery of a quantum field theory as described in chapters 10 and 11. Both the matter density ρ and radiation pressure p, for equation 12.2 substituted into equation 12.1 to obtain equation 12.3, represent possible macroscopic forms of $G_{\mu\nu}(x)$ which, while also entailing classical thermodynamic phenomena, are dependent upon the statistical range of possible exchanges for the microscopic fields. Arising out of the degeneracy of possible field solutions the conceptual origin of quantum and particle phenomena in the present theory differs to that in standard QFT as described in chapter 11. Correspondingly the notion of a 'vacuum state' is also different. Indeed the failure of calculations of the value for the vacuum energy in QFT to match the empirical value for Λ (typically by 120 orders of magnitude, as discussed towards the end of the previous section, see also for example [70] pp.790– 791) provides a further argument for the need to reassess the underlying structure of QFT itself, in particular in relation to the theory of gravitation. The possibility of addressing the cosmological constant problem within the context of the present theory was raised at the end of section 11.4.

While a number of features of the broken $E₇$ action on the components of $F(\text{h}_3\textcircled{0})$ projected over M_4 explicitly match features of the Standard Model of particle physics, as described for equation 9.46 and summarised in the bullet points in section 9.3, in this chapter we shall describe more qualitatively potential connections between features of the present theory and those of the standard cosmological model and theories of the very early universe.

As alluded to above a correlation between the external curvature and internal gauge fields $Y(x)$, expressed generically as $G_{\mu\nu} = f(Y)$, via the action integral of equation 5.18, was described in section 5.1 through a comparison with the framework of Kaluza-Klein theory. Further, towards the end of section 11.3 it was implied that an external geometry of a form which might ideally be expressed as $G_{\mu\nu} = f(\psi)$, corresponding for example to the electron field $\psi(x)$ for figure 11.13(a) and (b) in section 11.4, may arise from the fermion components within the space $F(h_3\mathbb{O})$ for the 56-dimensional vectors under $L(v_{56}) = 1$ through interactions with the gauge fields or more directly via an expression of the form $G_{\mu\nu} = f(\hat{\mathbf{v}})$. Similarly, without yet having a fully developed quantised theory, the possible physical manifestation of further components in the space $F(h_3\mathbb{O})$ may be considered.

In addition to the Lorentz vector v_4 and Lorentz spinor components of an element of $F(h_3\mathbb{O})$, transforming under the external subgroup $SL(2,\mathbb{C})^1 \subset E_6 \subset E_7$, identified in equation 9.46 there are four Lorentz scalar components α, β, n and N which may also contribute to shaping the external geometry through $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. In principle any of these four scalars, or even the scalar magnitude $|v_4|$ projected onto M_4 , could contribute to the macroscopic geometry. Each of the Lorentz scalars α, β, n, N and $|v_4|$ also transform trivially under the internal $SU(3)_c \times U(1)_Q$ gauge groups identified in section 8.2 and incorporated into the E⁷ symmetry in section 9.2. Hence, while the specific nature of $SU(2)_L \times U(1)_Y$ actions on these, or any other, components of $F(h_3\mathbb{O})$ is not yet known, in lacking both strong and electromagnetic interactions any of these scalar fields might contribute to the dark sector in cosmology.

For example a constant value for a scalar field such as α, β, n, N or $|v_4|$ projected over spacetime might be associated with the constancy of the scalar Λ for an effective cosmological constant term $\Lambda g_{\mu\nu}$ in the field equation 12.1 deriving from at least one of these fields. Interactions between α, β, n, N and $|v_4|$ implied under the terms of the constraint $L(\mathbf{v}_{56}) = 1$ may underlie empirically observed gravitational effects, in particular with the first four of these scalar fields coupled to the vector-Higgs v_4 in this way. Similar interactions under $L(v_{56}) = 1$ also relate to the fermion masses as described for equation 9.48, and in particular for the low neutrino mass alongside equation 9.49.

The identification of these scalars in the components of the full form $L(\mathbf{v}_{56}) = 1$ projected over M_4 is analogous to the appearance of a multiplet of scalar fields deriving from the components of a higher-dimensional metric in some forms of Kaluza-Klein theory, via a non-Killing metric Φ on the gauge group G as alluded to towards the end of section 4.3, as the geometry is 'reduced' over a 4-dimensional spacetime. In the present theory there is no higher-dimensional physical metric but, as for the scalars of Kaluza-Klein theories, here also scalar fields deriving from the breaking of the full form of temporal flow $L(\mathbf{v}_{56}) = 1$ may have implications for cosmology.

While ordinary matter, subject to the Standard Model internal gauge symmetry $\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y$, clumps together with an energy density $\rho(x)$ an essential requirement for a cosmological term is that, while locally having a much lower energy density than ordinary matter, it should have a largely even effect over cosmological scales in the apparent 'vacuum' of spacetime. While here not making a quantitative or specific argument for the $\Lambda g_{\mu\nu}$ term in the field equation the presence of a number a scalar fields in the theory, any of which may impact upon the external geometry, provides a source for investigation.

If a scalar field deriving from a component such as N in $F(h_3\mathbb{O})$ does give rise to a geometry of the form $G_{\mu\nu} = -\Lambda g_{\mu\nu}$ the effective energy density $T_{\mu\nu} := -\frac{1}{\kappa} G_{\mu\nu}$ in the form of a perfect fluid with constant energy density $\rho_{\Lambda} = \frac{\Lambda}{6}$ $\frac{\Lambda}{\kappa}$, and equation of state $p_{\Lambda} = -\rho_{\Lambda}$, might appear as a form of 'dark energy' arising as an apparent vacuum state, as described after equation 12.20 in section 12.2. The dynamical implications of such a term, as implied in equation 12.20 and summarised for the first FLRW model listed in table 12.1 in section 12.2, are well known to qualitatively match the empirical observation of the accelerating expansion of the universe at the present epoch.

Regarding the projection of the components of $v_{56} \in F(h_3\mathbb{O})$ onto the base manifold, and again without here making a rigorous argument, a symmetric rank-2 energy-momentum tensor *could* be constructed as $T_{\mu\nu} = \frac{1}{\kappa}$ $\frac{1}{\kappa} \lambda v_{\mu} v_{\nu}$, where λ is a real constant and $v_\mu(x) = g_{\mu\nu}v^\nu$ are the components of the Lorentz 4-vector $v_4 \subset v_{56} \in F(h_3\mathbb{O})$ projected onto TM₄ with magnitude $|v_4| = h$. This proposal is also motivated by analogy with the energy-momentum for dust $T_{\mu\nu} = \rho u_{\mu} u_{\nu}$, as contained in equation 12.2 for $p = 0$, with the 4-velocity u (with $|u| = 1$) representing the motion of idealised galaxies in the FLRW models. The field equation for $T_{\mu\nu} = \frac{1}{\kappa}$ $\frac{1}{\kappa} \lambda v_{\mu} v_{\nu}$ can be written as $G_{\mu\nu} + \lambda v_{\mu}v_{\nu} = 0$, which has a similar appearance to the field equation $G_{\mu\nu} + \Lambda g_{\mu\nu} = 0$ for the de Sitter model. On assuming the timelike flow of v_4 to be aligned with the galactic flow, as is the case for the 4-velocity u , the components of v_4 are simply $v_{\mu} = h u_{\mu}$, which are numerically the same as $v^{\mu} = (h, 0, 0, 0)$ on employing the metric of equations 12.5 and 12.6 and the comoving coordinates $\{t, r, \theta, \phi\}$. The substitution of $T_{\mu\nu} = \frac{1}{\kappa}$ $\frac{1}{6} \lambda v_{\mu} v_{\nu}$ into the field equations 12.9 and 12.10 then leads to the identical situation as the matter dominated case except here with an apparent matter density $\rho = \frac{1}{\kappa} \lambda h^2$. For constant $h(x)$ these equations do not lead to a solution unless $\lambda(t)$ is allowed to vary with cosmic time as for the parameter $\rho(t)$, in which case this model is identical to the matter dominated case as listed in the middle column of table 12.1.

Alternatively, since the full geometry is described by the Riemann tensor (which for example is also directly correlated with the internal curvature through relations on a bundle space such as equations 5.2 and 5.13 in the manner of a Kaluza-Klein theory) the Ricci tensor, defined with components $R_{\mu\nu} = R^{\sigma}{}_{\mu\nu\sigma}$, might be considered to be geometrically more fundamental than the Einstein tensor in terms of having a direct link with the underlying fields such as $N(x)$ or $v_4(x)$ deriving from the components of $F(h_3\mathbb{O})$ in equation 9.46. For the case of a constant scalar N giving rise to a cosmological constant Λ postulating the relation $R_{\mu\nu} = \Lambda g_{\mu\nu}$ implies directly that $G_{\mu\nu} := R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = -\Lambda g_{\mu\nu}$, which is identical to the case of the first model in table 12.1 already considered above. On the other hand postulating the relation $R_{\mu\nu} = \lambda v_{\mu}v_{\nu}$ as a possible vacuum limit does lead to a new scenario. Substituting this expression, with $v_4 = (h, 0, 0, 0)$ again aligned with the comoving coordinates, into the R_{00} component obtained from the Robertson-Walker line element in equation 12.7 leads immediately to the relation $3\frac{\ddot{a}}{a} = \lambda h^2$. This expression describes an exponentially expanding universe for constant $\lambda > 0$ and provides some of the motivation for originally considering a $\lambda v_{\mu}v_{\nu}$ term in the field equations.

However, a solution is of course required to be consistent with all components of the field equation. The fundamental role of the Einstein tensor is essentially due to the contracted Bianchi identity $G^{\mu\nu}_{;\mu} = 0$. The relation $R_{\mu\nu} = \lambda v_{\mu}v_{\nu}$ implies in turn $G_{\mu\nu} := R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = \lambda v_{\mu} v_{\nu} - \frac{1}{2}$ $\frac{1}{2}\lambda h^2 g_{\mu\nu}$ which via the definition $T_{\mu\nu} := -\frac{1}{\kappa} G_{\mu\nu}$ leads to an effective energy-momentum tensor in the form of equation 12.2 for this model with the equation of state $p_{\lambda} = \rho_{\lambda} = -\frac{\lambda h^2}{3\kappa}$. This contrasts with the de Sitter model with a Λ term for which $p_{\Lambda} = -\rho_{\Lambda} = -\frac{\Lambda}{\kappa}$ $\frac{\Lambda}{\kappa}$, as reviewed above. However the differing signs means that for the case of $R_{\mu\nu} = \lambda(t)v_{\mu}v_{\nu}$ a solution for equations 12.9 and 12.10 is only possible if λ is negative (that is $p_{\lambda} = \rho_{\lambda} > 0$) and allowed to vary in time, with the result listed in the final column of table 12.1 in section 12.2. Hence rather than being able to account for an accelerating expansion this hypothesis describes a more extreme deceleration than either the matter or radiation dominated models. Only the first case listed in table 12.1 describes an accelerating expansion, with $(\rho + 3p) < 0$ as discussed following equation 12.18.

The redundancy between equations 12.9, 12.10 and the expression $T^{\mu\nu}_{;\mu} = 0$ was highlighted by equation 12.11, and similarly here the identity $T^{\mu\nu}_{;\mu} = 0$ itself for $T_{\mu\nu} = \frac{1}{\kappa} \lambda v_{\mu} v_{\nu}$ or $T_{\mu\nu} = -\frac{1}{\kappa} (\lambda v_{\mu} v_{\nu} - \frac{1}{2} \lambda h^2 g_{\mu\nu})$ prohibits a constant value for λ . However in principle a full solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ may involve a range of contributions individually in the form of those in table 12.1 as well as others besides. In this case there will be a string of terms effectively composing the energy-momentum tensor which *collectively* are required to satisfy $T^{\mu\nu}_{;\mu} = 0$, a relation which in general may no longer hold for a particular contribution. This is very similar to the situation as described for equation 5.41 in section 5.2 for which a synthesis of charged matter and the electromagnetic field led to the Lorentz force law under the constraint $T^{\mu\nu}_{;\mu} = 0$. Here effectively a synthesis of several terms may arise within $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ on the cosmological scale.

The 4-vector v_4 in a $\lambda v_\mu v_\nu$ term could also be considered to have non-zero spatial components which might in principle relate to the formation of large scale structure in the universe and open up possibilities not available for a purely scalar degree of freedom in a $\Lambda g_{\mu\nu}$ term. However this in turn would imply the complication of loosening the FLRW assumptions of homogeneity and isotropy in the definition of the metric in equation 12.5. Even within those assumptions the possibilities with finite spatial curvature $k = \pm 1$, as well the purely $k = 0$ solutions of table 12.1, might be further considered. More generally, if the general structure of $-\kappa T_{\mu\nu} := G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$ on the largest scales of the universe can be established it will be a case of refitting the cosmological data with the parameters of the new model.

However, unlike the need to provisionally postulate explicit terms such as $\Lambda g_{\mu\nu}$ or $\lambda v_{\mu}v_{\nu}$ in the Einstein equation, as potentially effectively arising from a Lorentz scalar such as N or the Lorentz vector v_4 in the components of $v_{56} \in F(h_3\mathbb{O})$ projected over M_4 , there is a much more direct and intrinsic way in which this projection can shape the 4-dimensional spacetime geometry. We describe this observation, and its possible implications for the large scale structure of the universe, for the remainder of this section.

Earlier in this section, as for the discussion in section 5.3, the gravitational time dilation effects for $s \equiv \tau$ have been considered to result entirely from the metric $g_{\mu\nu}(x)$ as might be obtained through the Einstein equation 12.1, such as the case of the Schwarzschild solution of equation 5.49, that is essentially for cases with a known form of energy-momentum tensor. So far a constant magnitude has been assumed for $|v_4|^2 = L(v_4) = h^2$ in equation 13.1 in the projection of $v_4 \subset v_{56}$ onto TM_4 . However in principle all fields on M_4 may vary, within the necessary constraints such as $L(v_{56}) = 1$, similarly as the internal gauge field $Y(x)$ can vary under the constraint that the action integral of equation 5.18 should remain stationary, that is $\delta \tilde{I} = 0$, for example. Since the components of $v_4 \in TM_4$ represent the injection of the pure temporal flow s into the base manifold M_4 any variation in $|v_4|$ will itself have some impact on the spacetime geometry. Here we begin by considering this impact upon an otherwise flat manifold.

Hence we first return to the translation symmetry of the form $L(\mathbf{v}_4) = h^2$ under the four degrees of freedom $\{x^0, x^1, x^2, x^3\} \in \mathbb{R}^4$, as originally depicted for the 3-dimensional case in figure 2.2. Here the constant vector field $v_4 = (h, 0, 0, 0)$, in conformity with a constant value for h and with $v^0 = dx^0/ds = h$, is aligned with the global Lorentz frame as represented in figure 13.1 (a) . This first figure depicts the uniform translation symmetry implicit in the form $L(\mathbf{v}_4)$ as described in equation 2.13, which contrasts with the case in figure 13.1(b) in which the magnitude $h(x)$ of the 4vector $v_4(x)$ is free to vary.

Figure 13.1: The vector field v_4 subject to $L(v_4) = h^2(x)$ for (a) the original translation symmetry over $\mathbb{R}^4 \equiv M_4$ with constant $h(x)$ and global Lorentz symmetry and (b) with $h(x)$ variable and only local Lorentz symmetry. In both cases the flow v_4 is aligned to the timelike coordinate x^0 while x^i with $i = 1, 2, 3$ represents the three spacelike coordinates.

For the present theory the local metric $g_{\mu\nu}(x)$ is projected from the form $L(\mathbf{v}_4) = \eta_{ab}v^a v^b = h^2$, framing the local injection of temporal flow into the base manifold. However with the local coordinate x^0 of figure 13.1 representing the fundamental flow of time according to the relation $\delta s = \delta x^0/h$, the expression $L(\mathbf{v}_4) = h^2$, subject to the constraint $L(\hat{v}) = 1$, also sets the scale for temporal flow in the local frame. That is, the x^0 coordinate representation of time will vary with the value of h. With $\delta s = \delta x^0/h$ the pure time s will effectively flow more slowly in regions of large h, corresponding to the vectors v_4 with a larger magnitude in figure 13.1(b), and more quickly in spacetime regions with a lower value of h . More generally the relation $L(v_4) = h^2$ can be rearranged in the form $ds^2 = \frac{\eta_{ab}}{h^2} dx^a dx^b$, that is the final relation in equation 5.47, with the spacetime metric extracted as:

$$
g_{\mu\nu}(x) = \frac{1}{h^2(x)} \eta_{\mu\nu}
$$
\n(13.2)

when expressed in the global coordinates on the extended manifold M_4 (see also the discussion of equation 13.3 below). This physical metric $g_{\mu\nu}$, related to flat spacetime through the conformal transformation $\eta_{\mu\nu} \to \eta_{\mu\nu}/h^2(x)$, describes a non-Euclidean manifold incorporating time dilation effects. As for general relativity, while general coordinate systems are arbitrary and unphysical, local inertial frames with $g_{\mu\nu}(x) =$ $\eta_{\mu\nu}$ and $\partial_{\sigma}g_{\mu\nu}=0$ do have physical significance. Such an inertial frame may be identified globally for figure 13.1(a) but only locally for figure 13.1(b). By the strong equivalence principle the laws of physics according to special relativity apply in a local inertial reference frame. As described in section 3.4 the weak equivalence principle is sufficient to incorporate the notion that all gravitational effects can be transformed away in a sufficiently small spacetime volume, and can be interpreted as implying that the torsion vanishes.

While the unphysical nature of coordinate systems in general relativity is encapsulated under general covariance, as also described in section 3.4, any coordinate system can be represented by the parameter space grid of figure $3.6(a)$. In the special case of Minkowski spacetime such a coordinate grid can be mapped onto the 4-dimensional manifold such that the metric has constant components $g_{\mu\nu}(x) = \eta_{\mu\nu}$, as is the case for the spacetime underlying the constant vector flow depicted in figure 13.1(a). In this case a unique family of coordinate charts are identified through the parameter space of translation symmetry of the form $L(\mathbf{v}_4) = h^2$, as described in equation 2.13, and related to each other via global Lorentz transformations. On the other hand in projecting the coordinate grid of figure 3.6(a) onto the spacetime underlying figure 13.1(b) the simplest expression for the metric takes the form of equation 13.2.

For the metric of either figure 13.1(a) or (b) obtained in this way through the underlying injection of temporal flow s into the spacetime manifold, as for the case of a metric determined as a solution to Einstein's equation as considered in section 5.3, the proper time τ recorded by physical clocks is again tied to the fundamental flow of time s. This is the case since the laws of physics, including those utilised by the structure of clocks, unfold through the underlying temporal flow s and hence the proper time $\tau \equiv s$ exhibits the equivalent time dilation effects due to variation in $L(\nu_4) = h^2$, as was the case for other sources of temporal dilation. The question then concerns the more specific nature of this relation between τ and s, as originally discussed at the end of section 5.3 and earlier in this section.

The fundamental temporal flow s is modelled by the real line and hence can be represented by the values of a pure real number $s \in \mathbb{R}$, intervals of which can be expressed in terms of a set of real parameters of arbitrarily high dimension, as described for equation 2.4, which is an essential observation for the present paper. On the other hand the proper time τ represents intervals of 4-dimensional spacetime on the manifold M_4 and is expressed by a real number associated with the dimension of length L (which

is equivalent to the dimension of time T since natural units are employed, and in a sense it would be more appropriate to use T as we are ultimately dealing with multidimensional forms of temporal flow). Hence the constant factor γ relating the pure 1-dimensional temporal flow s to a corresponding empirically observable progression in proper time $\tau = \gamma s$ is one which carries the dimension of length L. Hence in turn proper time intervals for the spacetime geometry underlying figure $13.1(a)$ or (b) can be expressed through:

$$
d\tau^2 = \gamma^2 ds^2 = \frac{\gamma^2}{h^2} \eta_{\mu\nu} dx^{\mu} dx^{\nu}
$$
 (13.3)

Here then the metric $g_{\mu\nu} = \frac{\gamma^2}{h^2} \eta_{\mu\nu}$ explicitly carries the dimension of L^2 , as was described for the general case in the discussion following equation 12.5 in section 12.2. Since the underlying temporal flow s is not directly observed, and the scale of the real line parametrising s is in any case arbitrary, once empirical units, such as metres, are chosen for τ the coordinate parameters can in turn be chosen such that the metric takes a convenient form. For example in the case of figure $13.1(a)$ pseudo-Euclidean coordinates can be chosen such that $g_{\mu\nu}(x) = \eta_{\mu\nu}$ everywhere.

While the value of the factor γ is of no meaning, its significance lies in representing a constant relation $\tau = \gamma s$. In practice setting $\gamma = 1$ can be interpreted as choosing the arbitrary scale of $s \in \mathbb{R}$ to match the practical parametrisation of the proper time τ . In this case the basic metric $g_{\mu\nu}$ from equation 13.3 is that of equation 13.2. With $h(x)$ varying the constancy of γ in equation 13.3 implies that in general it is not possible to find any coordinates such that $g_{\mu\nu}(x) = \eta_{\mu\nu}$ globally for the scenario in figure 13.1(b), although this relation is always possible locally, as also suggested by the equivalence principle.

Hence variation in the value of $h(x)$ on M_4 directly modifies the effective metric $g_{\mu\nu}(x)$, warping the spacetime geometry that underlies the vector field in figure 13.1(b) for example. Assuming the geometry to be described in terms of a torsion-free linear connection the corresponding Levi-Civita connection Γ can be constructed as a function of the metric of equation 13.2 via equation 3.53 and in turn the components of the full Riemannian curvature tensor $R^{\rho}_{\ \sigma\mu\nu}$ of equation 3.73 computed. In turn the Einstein tensor, for the conformal geometry $g_{\mu\nu} = \theta(x)\eta_{\mu\nu}$ with a real scalar field $\theta(x) = h^{-2}(x)$, is found explicitly (and cross-checked with a related calculation in [9] pp.42 and 76) to take the form:

$$
G_{\mu\nu} = -\frac{3}{2}\theta^{-2}\partial_{\mu}\theta\partial_{\nu}\theta + \frac{3}{4}\theta^{-2}\partial_{\rho}\theta\partial^{\rho}\theta g_{\mu\nu} + \theta^{-1}\partial_{\mu}\partial_{\nu}\theta - \theta^{-1}\Box\theta g_{\mu\nu}
$$
(13.4)

A similar expression, with a different set of coefficients, is obtained as a function of h under the substitution $\theta \to h^{-2}$, as for any other scalar field related to θ by a simple power expression. The derivation of this expression for $G_{\mu\nu}$ follows the same chain of relations that led to the form of G_{00} and G_{11} , appearing alongside the corresponding $\Lambda g_{\mu\nu}$ terms on the left-hand side of equations 12.9 and 12.10 respectively, given the metric form of equation 12.6 and via the Ricci tensor $R_{\mu\nu}$ and scalar curvature R. However here equation 13.4 represents a direct warping of the spacetime geometry due to the variation in $L(\mathbf{v}_4) = h^2$ which implies equation 13.2, without the need to employ further assumptions regarding the form of an energy-momentum tensor in Einstein's equation in order to extract a solution.

Hence this construction can be contrasted with the usual determination of a metric $g_{\mu\nu}$ in general relativity. There the metric is extracted as a solution to the set of second order differential equations contained in the Einstein field equation 12.1 under assumptions of symmetry regarding both the matter distribution and the form of the metric itself. This was the approach taken for the Schwarzschild solution of equation 5.49 and also for the cosmological models based on the Robertson-Walker line element of equation 12.5. Here in contrast the form of the metric $g_{\mu\nu} = \frac{1}{h^2} \eta_{\mu\nu}$ implies a linear connection Γ and Riemannian curvature $\mathbf R$ and hence leads to the Einstein tensor $G_{\mu\nu} = f(\mathbf{v}_4)$ as a *consequence* of the variation in $L(\mathbf{v}_4) = h^2(x)$ under the constraint $L(\hat{\mathbf{v}}) = 1$, rather than as an equation to solve for the metric.

In practice the distribution $h(x)$ might be constrained by observations of the corresponding gravitational effects, in a similar way that the constant k and scale factor $a(t)$ of the Robertson-Walker line element of equations 12.5 and 12.6 are determined through empirical observations, found to be consistent with $k = 0$ and deducing the structure depicted in figure 12.2 for example. In this sense the procedure to constrain the actual function $h(x)$ is very similar to the standard approach for general relativity, that is by matching equation 13.4 with empirical observations. On the other hand in this case it may also prove possible to calculate both the typical value of $h(x)$, and the typical range of variation in this magnitude, as constrained for example under the relation $L(\hat{\mathbf{v}}) = 1$, within the theory itself.

In addition to the warped spacetime $G_{\mu\nu} = f(\hat{\mathbf{v}})$ of equation 13.4 geometries of the form $G_{\mu\nu} = f(Y)$, relating the external curvature to the internal gauge fields as described in section 5.1, are also possible. The combined general expression $G_{\mu\nu}$ = $f(Y, \hat{v})$ can be interpreted to incorporate a contribution from the gauge fields $Y(x)$ which determine the metric via the differential field equations $G_{\mu\nu} = f(Y)$, while the geometry $G_{\mu\nu} = f(\hat{\boldsymbol{v}})$ concerns the direct impact of $L(\boldsymbol{v}_4) = h^2(x)$ on the metric in the form of equation 13.2. As described in chapter 11 ordinary matter exhibiting quantum phenomena will arise out of an underlying degeneracy of solutions for $G_{\mu\nu} = f(Y, \hat{v})$ given the field exchanges such as $\delta Y \leftrightarrow \delta \hat{v}$ allowed according to the selection rules summarised in equations 11.29.

Since the fermion components $\psi(x)$ in $F(h_3\mathbb{O})$ are correlated with the components of $v_4 \equiv h_2$ in $F(h_3 \mathbb{O})$ under the constraint $L(v_{56}) = 1$ as described for equation 9.48, fermion terms may explicitly appear through field exchanges of the form $\delta v_4 \leftrightarrow \delta \psi$. That is, these interactions may directly give rise to the effective geometry $G_{\mu\nu} = f(\psi)$ alluded to earlier in this section and towards the end of section 11.3, and as applied for the external geometric structure associated with the fermion fields underlying the processes depicted in figures 11.13(a) and (b) for example. In general the full set of microscopic field redescription possibilities, consistent with the constraint equations, will need to be taken into account to determine the form of macroscopic geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ as shaped through a local degeneracy of field solutions as described in chapter 11.

The gravitational time dilation effect, that is the relative slowing of time in the vicinity of a massive object, described by the Schwarzschild solution for the metric in equations 5.49 and 5.50, can be ascribed to the presence of the massive object itself in general relativity. Accordingly the situation for regions in figure 13.1(b) with relatively large values of $h = |\mathbf{v}_4|$, and hence a relative slowing of the flow of time, might be 'reverse engineered' to identify an apparent presence of 'matter' in such a region. That is, for the geometry $G_{\mu\nu} = f(\mathbf{v}_4)$ in equation 13.4 it is possible to define an associated energy-momentum tensor through $-\kappa T_{\mu\nu}^D := G_{\mu\nu} = f(\mathbf{v}_4)$. Here $T_{\mu\nu}^D$ does not then represent 'ordinary matter' which is built upon a degeneracy of gauge and fermion field solutions for the geometry $G_{\mu\nu}(x)$ over M_4 , and in particular made 'visible' through the $U(1)_Q$ electromagnetic interactions, but rather an underlying warping of spacetime geometry itself. Hence, while not describing baryonic matter, the implicit energymomentum $T_{\mu\nu}^D$ is a candidate for the 'dark matter' of the universe.

With $T_{\mu\nu}^D$ defined in this way for the Einstein tensor $G_{\mu\nu}$ of equation 13.4 deriving directly from the metric $g_{\mu\nu}$ of equation 13.2, this in fact follows the procedure for obtaining solutions for Einstein's equation by cataloguing $(g_{\mu\nu}(x), T_{\mu\nu}(x))$ pairs as outlined towards the end of section 5.2. Here however it is the form of the metric that is physically motivated and not arbitrary while the resulting energy-momentum tensor need not necessarily correspond to any known form of matter.

The term 'dark matter' implies a kind of 'phantom source' of gravitation, which is only detectable through its manifestation as a structure of spacetime geometry, and indeed the above definition of $T_{\mu\nu}^D$ essentially describes a purely gravitational phenomenon. In general a structure described in the relation $-\kappa T_{\mu\nu} := G_{\mu\nu}$ may or may not be detectable as 'matter' and may or may not be detectable as 'gravity'. For example ordinary baryonic matter in the form of stars or planets is both visible as matter $T_{\mu\nu}$ and evident as gravity $G_{\mu\nu}$. On the other hand baryonic matter in the form of tables and chairs, while clearly exhibiting a number of properties of matter, does not give rise to any detectable gravitational effects. Contrary to that situation 'dark matter' in the form of $G_{\mu\nu} = f(\mathbf{v}_4)$ might produce very significant gravitational phenomena without being associated with any apparent material effects at all. For example, as alluded to above, the variation in $h(x)$ might be determined through observations of galactic rotation curves and gravitational lensing effects rather than an explicit empirical detection of a 'dark matter' distribution (as would be possible for example for a cloud of dust on a galactic scale). A fourth case is conceivable in which a definite mathematical form of $-\kappa T_{\mu\nu} := G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ has evaded detection both as a material and a gravitational entity.

Since the material effects of the local ordinary matter distribution present themselves more immediately than the corresponding gravitational phenomena, historically the sense that universal gravity is a property to be associated with matter was a natural point of view to adopt. In turn the Einstein equation $G_{\mu\nu} = -\kappa T_{\mu\nu}$, influenced by the Newtonian gravity which arises in the appropriate limit, was initially interpreted to imply that in some sense matter 'causes' the curvature of spacetime . That interpretation is considered to be a 'reverse engineering' from the perspective adopted in this paper in which the energy-momentum tensor is simply defined through $-\kappa T_{\mu\nu} := G_{\mu\nu}$, with the spacetime geometry determined primarily as a solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ subject to the constraint equations (see also the discussion in the opening paragraphs of section 5.2).

In terms of the spacetime solution in the particular region of the early universe, through the mutual gravitation of dark matter and ordinary baryonic matter the effects of $G_{\mu\nu} = f(\mathbf{v}_4)$, as a network of creases in the underlying fabric of spacetime, might have guided the formation of galaxies and galactic clusters. The properties of these structures are then visible today through the motions of galaxies within clusters and the rotation curves of stars within galaxies, all still in mutual gravitational interaction with the dark matter. This interplay between baryonic and dark matter is depicted in figure 13.2, where the final stage labelled (e) corresponds to the kind of structures observed through to the present epoch as also represented in figure 12.1.

Figure 13.2: (c) Fluctuations in the magnitude $h(x)$ of the vector field v_4 , represented by the vertical arrows (as for figure 13.1), in the early universe (d) gravitationally merge along with baryonic matter, represented as points of dust, as the universe evolves leading to (e) the formation of large scale galactic structures as observed through to the present epoch. (Earlier epochs will be represented in figure $13.3(a)$, (b) in the following section).

In the standard cosmological model it is known that the dark matter cannot be baryonic due to the abundances of the light elements resulting from nucleosynthesis in the early universe. Dark matter composed of relic particles from the Big Bang must also be weakly interacting in order to have evaded direct detection. In the case of 'cold dark matter' (CDM) the relic particles have a low thermal velocity leading to a hierarchical formation of structure through the merger of smaller initial units beginning in the early universe. This description is consistent with the picture in figure 13.2, except that for 'dark matter' in the form of variations in $|v_4|$ there are seemingly no associated 'particle' phenomena at all.

Since $v_4(x)$ is a 4-vector field, as well as fluctuations in $|v_4|$, ascribed to the temporal component v^0 in figures 13.1(b) and 13.2, in principle there may be variations in the spatial components v^i also (as suggested earlier in this section for the case in the final column of table 12.1 for a spacetime geometry incorporating a $\lambda v_{\mu}v_{\nu}$ term in the Einstein field equation) which could be pictured as a horizontal component for the vectors in these figures. Such spatial fluctuations are counter to the assumption of strict homogeneity and isotropy, as indeed are variations in the magnitude $h(x)$, but they could potentially be a factor in the observed peculiar motions of galaxies and clusters of galaxies and might even be associated with a 'dark flow' if observations of such phenomena were to be established.

Further, while fluctuations towards higher values of $|v_4|$, that is a larger value for $L(\mathbf{v}_4) = h^2$, correspond to regions of spacetime with an apparent slowing of the flow of time $\tau \equiv s$, and hence associated with 'dark matter', regions with a smaller value for $L(\mathbf{v}_4) = h^2$ will have a complementary spacetime geometry with a faster rate of temporal flow and in principle the opposite gravitational effect. Such regions may hence tend to open up cosmic 'voids' between the galactic clusters and play an important role in the structural evolution process represented in figure 13.2. On the largest observable scales such a gravitational repulsion might also be a factor in the composition of the apparent 'dark energy'.

It is a very familiar idea that a 2-dimensional surface embedded within a 3 dimensional space will generally have an intrinsic curvature, such as the surface of a ball for example. Here we are considering the embedding of 4-dimensional spacetime within the structures of a higher-dimensional form of temporal flow $L(\hat{\mathbf{v}}) = 1$, and it is again natural to expect that in general a finite intrinsic curvature for the 4 dimensional manifold might result. It is hence proposed that such intrinsic curvature for the spacetime geometry, closely correlated with variation in the component values of the projected 4-vector $v_4 \,\subset \hat{v}$ onto TM₄, constitutes at least a significant factor in accounting for the observed phenomena of the 'dark sector' in cosmology.

In general relativity spacetime curvature might be considered to account for the origin of mass in general by interpreting Einstein's field equation essentially as a definition of energy-momentum $-\kappa T_{\mu\nu} := G_{\mu\nu}$, as we have in this paper and as reviewed above. This is in contrast with the Standard Model of particle physics in which the Higgs field $\phi(x)$ and the Higgs mechanism of spontaneous symmetry breaking, as described in section 7.2, is responsible for the origin of mass through field interactions for a theory framed in a flat spacetime background.

In the present theory with a continuous variation in the magnitude of the underlying field $v_4(x)$ on M_4 , implying time dilation effects and shaping the spacetime geometry, an apparent 'mass' might be associated with this field through $-\kappa T_{\mu\nu}^D :=$ $G_{\mu\nu} = f(\nu_4)$ as described above, and hence the field $\nu_4(x)$ can be considered as the source of this apparent mass. In subsection 8.3.3, and in particular in the discussion around equation 8.76, and similarly around equation 9.48 in section 9.2, the same field $v_4(x)$ has been associated with Higgs phenomena in conveying masses to the fermion and gauge boson fields via possible $\delta v_4 \leftrightarrow \delta \psi$ and $\delta v_4 \leftrightarrow \delta Y$ interactions respectively, compatible with the constraints summarised in equations 11.29.

Combining these observations suggests that the physical mechanism through which a $\psi(x)$ or $Y(x)$ field interaction with the 'vector-Higgs' field $v_4(x)$ results in a 'mass' for the fermions or gauge bosons respectively is through the effect on the local geometry due to the projection of the field $v_4(x) \in TM_4$ out of $\hat{v}(x)$ under the full form $L(\hat{v}) = 1$. Such a δv_4 interaction may locally correspond to a further geometric effect on top of the continuous $v_4(x)$ variation. In the present theory such external gravitational effects will be compatible with underlying quantum effects through the above field redescriptions of the form $\delta v_4 \leftrightarrow \delta \psi$ and $\delta v_4 \leftrightarrow \delta Y$, which add to the list of possible interaction vertices of figure 11.3 in the correspondence with Feynman rules for a quantum field theory. In the context of QFT the relation between the 'bare mass' associated with these field interactions and the measured mass of physical particle states, as considered near the end of section 11.3, will depend upon this impact on the external spacetime geometry.

In this theory while both dark matter and Higgs phenomena are directly associated with the field $v_4(x)$ the dark matter is of course not *composed* of Higgs particles. Rather the Higgs interactions via *discrete* $\delta|\mathbf{v}_4|$ exchanges are closely associated with visible baryonic matter in the universe, and as observed in high energy physics experiments, while dark matter relates to a continuous variation in the underlying field $v_4(x)$. Even if $L(v_4) = h^2(x)$ was constant on large scales, hence with no dark matter phenomena, 'strongly coupled' field interactions with the vector-Higgs field v_4 and Higgs particles would still be observable in the laboratory. This observation is compatible with the apparently 'weakly interacting' effects of dark matter as a gravitational phenomenon that arises through variations in $h(x)$ on the galactic scale. That is, while the spacetime geometry resulting from the continuous $v_4(x)$ variation need not itself be associated directly with any quantum or particle phenomena, the scalar $\delta |\mathbf{v}_4|$ interactions of the same everywhere pervading field may give rise to the detected Higgs particle states.

More generally the question remains to understand whether specific quantum or particle effects might be associated with the dark sector, and how the geometric phenomena arising from the injection of the temporal flow into the spacetime manifold relate to the properties of the familiar Standard Model particle states. Together these phenomena shape the world geometry as described collectively under $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. The external geometry will involve the conformal transformation of equation 13.2, which generates Ricci curvature and leads to equation 13.4, together with more general solutions over a degeneracy of underlying internal field exchanges, resulting in a combination and interplay of both Ricci and Weyl curvature in general.

Dark matter is empirically observed to be associated with galactic clusters, and hence the value of $h(x)$ is expected to be larger in such regions and lower in inter-galactic space, as sketched in figure $13.2(e)$. Given that copious photons of light and other Standard Model particles can be detected on Earth after being transmitted through such regions, in travelling from distant galaxies, it appears that the properties of such particles must be physically robust for small variations of $h(x)$ to some degree.

The development of the full physical form and consequences of the expression $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ will require a greater understanding of the incorporation of quantum phenomena as introduced in chapter 11. The full implications of the theory, derived either via direct calculation or simulation, for particle physics as well as cosmology will depend both on the degree of variation of $L(\mathbf{v}_4) = h^2(x)$ and the typical value of $h(x)$ itself at the present epoch. With material properties and the laws of physics likely to have some dependence on the value of $h(x)$ it may be that the Standard Model of particle physics requires a certain apparent 'tuning' of this parameter in order to allow the formation of ordinary baryonic matter itself. This raises the question more generally of the possible uniqueness, or otherwise, of the 'physical constants' as observed in the world, both for Standard Model of particle physics and in terms of the cosmological parameters, as we shall discuss in section 13.3.

In the following section we first consider the possibility that both the average value and the fluctuations in $h(x)$ may have been very different in the very early universe, leading to a 'phase transition' to an average value of $h(x)$ compatible with the nature and properties of Standard Model interactions in particle physics and which has remained stable to the present day. In this scenario the phase transition may mark a point of convergence upon the familiar laws of physics in 4-dimensional spacetime more generally. These may include the second law of thermodynamics expressed in terms of the degrees of freedom of familiar interacting particles which are themselves produced in the phase transition.

In summary, the 'novel perspective' in the title of this chapter refers to the manner in which the intrinsic geometry of the 4-dimensional spacetime backdrop for cosmology is shaped through the projection of the extended M_4 manifold out of the full multi-dimensional form of progression in time $L(\hat{v}) = 1$. While the present theory based on general forms of time is very simple there are a number of features such as the projection of the vector-Higgs field v_4 onto TM_4 , generating the conformal transformation of equation 13.2 parametrised for example by the dilation symmetries described in the opening of the following section, and a set of elementary scalar fields α, β, n and N, as described earlier in this section, which potentially correlate with large scale cosmological phenomena in particular associated with the dark sector. These features identified within the components of $v_{56} \in F(h_3\mathbb{O})$ for the 56-dimensional form $L(v_{56}) = 1$ are complementary to the features identified for the Standard Model of particle physics as summarised in equation 9.46 and in section 9.3. Since the known phenomena of HEP cannot account for the dark sector in cosmology new features, such as identified in this section, are indeed required to account for the cosmological parameters. There then remains the question concerning the degree to which the mathematical structures described in this section might compare quantitatively with empirical observations of the large scale physical structure of the universe.

A first step will be to seek a guide through a comparison between the approach of the present theory and geometric models aimed at accounting for the dark sector of cosmology in the existing literature. While papers involving conformal gravity (see for example [76, 77]) may account for elements of the dark sector in geometric terms, such models appear somewhat different to the approach described in this section. In replacing the Einstein-Hilbert action by a conformally invariant action based on the Weyl tensor these papers do however implicitly incorporate geometric transformations of the kind in equation 13.2 and hence may relate to the structures of the present theory. The present theory both aims to avoid the employment a Lagrangian formalism and does not propose a 'modified gravity' of any kind. In fact here the Einstein equation is identified as a fundamental feature embedded within the definition of energy-momentum through the expression $-\kappa T_{\mu\nu} := G_{\mu\nu} = f(Y, \hat{v})$, which also provides the interpretation of the Einstein equation in the context of the present theory. While one aim of this paper has been to avoid 'postulating' a Lagrangian of any form, the Einstein-Hilbert action of equation 3.79 for general relativity, for the vacuum case with $\mathcal{L} = 0$ and $\Lambda = 0$, has been adapted in section 5.1 to facilitate a provisional connection between the external and internal geometry arising from the symmetries of $L(\hat{\mathbf{v}}) = 1$ broken
over the base manifold M_4 , as described for equations 5.18 and 5.19 and guided by Kaluza-Klein theory.

Further, rather than devising a scheme tailored to match empirical observations, here we begin with an underlying conceptual motivation and foundation for the theory. Once this theory has been sufficiently developed a full cosmological model might be established leading for example to a calculation for the density parameters Ω_B , Ω_D and Ω_{Λ} , as introduced following equation 12.15 in the previous chapter. In principle the cosmological data itself might then be refit within the context of the theory in order to test these ideas quantitatively.

13.2 The Very Early Universe

The highest-dimensional form of temporal flow considered in any detail in this paper is the form $L(v_{56}) = 1$ with E_7 symmetry, as introduced in section 9.2. If any of the four scalar components $\{\alpha, \beta, n, N\}$ of $v_{56} \in F(h_3\mathbb{O})$ in equation 9.46 is found to be associated with an effective cosmological term $\Lambda g_{\mu\nu}$ in equation 12.1 then the magnitude of this component will be correlated with the magnitude of the acceleration of the cosmic expansion, which may be arbitrarily small. Similarly the magnitude and variation of the projected $v_4 \in TM_4$ components may directly correlate with the properties of dark matter or dark energy, while as a 'vector-Higgs' the field v_4 also generates mass terms for the fermions and gauge bosons and underlies Higgs phenomena in general, as also reviewed in the previous section. Interactions between the scalars $\{\alpha, \beta, n, N\}$ and v_4 might also generate massive weakly-interacting scalar states as a possible contribution to the dark sector.

One means of varying $|v_4|$ can be described via a simple dilation symmetry as a subgroup of $SO^+(1,9)$ for the model described in section 5.1 for a 10-dimensional form $L(\mathbf{v}_{10}) = 1$ projected over M_4 as pictured in figure 5.1. This dilation symmetry acts on the components of v_{10} such that the magnitudes of the external \overline{v}_4 and internal \underline{v}_6 vectors are traded subject to the constraint $L(\mathbf{v}_{10}) = |\overline{\mathbf{v}}_4|^2 + |\underline{\mathbf{v}}_6|^2 = 1$. This variation in $h = |\mathbf{v}_4|$ can also be described in terms of a one-parameter subgroup denoted $D(1)_X \subset SL(2,\mathbb{O})$ acting as a dilation symmetry on the components of $v_{10} \equiv X \in h_2\mathbb{O}$ in equation 6.16 preserving $L(\mathbf{v}_{10}) = \det(X) = 1$.

In terms of the largest form of temporal identified another possibility for 'tuning' the magnitude of the v_4 components under the constraint of the 56-dimensional form $L(v_{56}) = 1$ lies in the dilation symmetry, which will be denoted $D(1)$ _λ \subset E₇, as parametrised by $\lambda \in \mathbb{R}$ and introduced in equation 9.30. This symmetry acts upon all 27 components of $\mathcal{Y} \in \mathrm{h}_{3}\mathbb{O}$ of the 56-dimensional space $F(\mathrm{h}_{3}\mathbb{O})$ in a uniform way, and not only on the $v_4 \subset \mathcal{Y}$ subset of components in equation 9.46.

As an intermediate case a further dilation of components can be identified within the E₆ symmetry on the h₃ \mathbb{O} subspaces of $F(\mathrm{h}_3 \mathbb{O})$ and will be denoted $D(1)_B$, as generated by the linear combination of boosts $\dot{B}_{tz}^1 + 2\dot{B}_{tz}^2$ and as introduced in equation 8.35. From table 6.6 this E_6 generator as a vector field in the tangent space $Th_3\mathbb{O}$ has the form:

$$
\dot{B}_{\mathbf{L}}^1 + 2\dot{B}_{\mathbf{L}}^2 = \begin{pmatrix} +p & +\bar{a} & -\frac{1}{2}c \\ +a & +m & -\frac{1}{2}\bar{b} \\ -\frac{1}{2}\bar{c} & -\frac{1}{2}b & -2n \end{pmatrix}
$$
(13.5)

Hence under the action of $D(1)_B$ on the components of $\mathcal{X} \in \mathcal{H}_3\mathbb{O}$ in equation 6.26 the 10 components of X are inflated while the remaining 17 components of θ and n are deflated, or vice versa. The consistency of this $D(1)_B$ action with the preservation of $L(v_{27}) = det(\mathcal{X})$ can be seen directly from the form of $det(\mathcal{X})$ in equation 6.27 together with the generator coefficients in equation 13.5. For this particular linear combination of boosts the rank-6 E_6 Lie algebra contains a rank-6 subalgebra decomposition, which in terms of the corresponding Lie groups can be written as:

$$
SL(2,\mathbb{C})^1 \times SU(3)_c \times U(1)_Q \times D(1)_B \subset E_6 \tag{13.6}
$$

As mentioned for the same decomposition in equation 8.35 the mathematical structure of this Lie subalgebra is described in [38] (the first example in Appendix D, p.187). In the present paper the subgroup $SL(2,\mathbb{C})^1$ has been identified with the external symmetry of spacetime M_4 in section 8.1, and $SU(3)_c \times U(1)_Q$ as the internal symmetry subgroup within Stab $(TM₄)$ in section 8.2, in each case with a corresponding physical interpretation. Clearly $D(1)_B$ is not a subgroup of Stab(TM₄) due to the action on the $v_4 \text{ }\subset X$ components in equation 13.5, however since $D(1)_B$ is independent of $SL(2,\mathbb{C})^1 \times SU(3)_c \times U(1)_Q$, as described in equation 13.6, it may also be of physical significance.

Regardless of the means of varying the 4-vector magnitude $|v_4| = h$, whether via $D(1)_X$, $D(1)_B$, $D(1)_\lambda$ or otherwise, the geometric impact of relatively high values of $L(\mathbf{v}_4) = h^2(x)$ projected out of the full form $L(\hat{\mathbf{v}}) = 1$ over extended regions of the cosmos is considered to form a candidate for the effects of dark matter, as described in the previous section. As well as these small variations in $h(x)$, the implications of a much larger time-dependent evolution in the scalar value $h(t)$, as averaged over the 3dimensional spatial hypersurfaces as a function of cosmic time t, can be considered for the very early universe, as we describe in this section. For example we shall consider the progression from a value approaching zero $h(t) \to 0$ for $t \to 0$ towards a stable average value $h(t_v) = h_0$ in a period of time associated with the epoch of the Big Bang, with $h_0 = |\mathbf{v}_4|$ hence also denoting the present day average value. On adopting a normalisation factor of $\gamma = 1$ (a further natural option would be to set $\gamma = h_0$) relating the fundamental temporal flow s with the proper time τ , as described for equation 13.3, the basic metric deformation can be described by equation 13.2. An interval of proper time $d\tau$ can then be expressed as:

$$
d\tau^2 = ds^2 = \frac{1}{h^2(t)} \left[dt^2 - d\Sigma^2 \right] \tag{13.7}
$$

where $d\Sigma$ represents a Euclidean 3-dimensional spatial element. While this line element has the form of a conformal transformation dependent upon $h(t)$, similarly as for the case of equation 12.23 for the FLRW models with conformal time parameter η , we continue to think of t as the 'cosmic time' parameter. Given that the value of $h(t)$ is only considered to differ from the present value h_0 for $t < t_v$ in the very early universe, and that elementary physical structures will be unfamiliar during that epoch as the nature of the projection of $L(\hat{v}) = 1$ over that region of M_4 is correspondingly also different, the form of any 'physical clock' and the measure of time itself will need further consideration for this earliest era. Hence in the above mathematical expression we keep track of the coordinate time t in place of defining a new temporal parameter.

The identification of $D(1)_B \subset E_6$ alongside other subgroups in equation 13.6 is analogous to the proposed subgroup $SU(2)_L \times U(1)_Y \subset E_7$, as a candidate for the gauge symmetry underlying the left-handed weak interactions, as described in equation 9.47 and the subsequent discussion. In the case of $SU(2)_L \times U(1)_Y$ the fact that this symmetry is not independent of $Stab(TM_4)$ (while it is independent of $SL(2,\mathbb{C})^1 \times SU(3)_{c}$ leads to the phenomena of 'electroweak symmetry breaking' through the interaction of the SU(2)_L \times U(1)_Y gauge fields and the $v_4 \in TM_4$ vector-Higgs field, breaking the symmetry down to $U(1)_Q$. For the future development of the present theory it will be important to gain an understanding of the interplay between electroweak symmetry breaking (as well as the role of the unification scale described for figure 11.10) and the breaking of the $D_B(1)$ symmetry action (or other dilation symmetry) in the very early universe.

A 'gauge field' associated with the $D(1)_B$ dilation symmetry, through interaction with the components of v_{56} , might itself drive an inflationary effect in the very early universe. The very different physical environment associated with a very different magnitude of $v_4 \,\subset\, v_{56}$ projection onto TM₄ in the very early universe might also in principle incorporate some of the effects of a cosmological inflation. In either case, following the inflationary epoch the $D(1)_B$ symmetry action would effectively be broken in a 'phase transition' as the value of $|v_4|$ is stabilised and the parameters of the Standard Model of particle physics are established. That is, with field interactions leading to a mutual stabilisation of both the value of $|v_4| = h_0$ and the Standard Model parameters.

In this theory, with the field v_4 also closely associated with Higgs phenomena, these parameters include the masses of the fermions through couplings implied in the constraint $L(v_{56}) = 1$, as recalled towards the end of the previous section. The D(1)_B symmetry, as generated by the E_6 Lie algebra element of equation 13.5, applies to both X and $\mathcal{Y} \in \mathrm{h}_3\mathbb{O}$ in equation 9.46 and with a uniform action on all components of $Y \subset h_2\mathbb{O}$, not just the $v_4 \in h_2\mathbb{C}$ subspace. The X and Y components in equation 9.46 carry the u-quark and ν -lepton states while the d-quark and e-lepton states reside in the θ^1_{λ} $\frac{1}{\chi}$ and $\theta_{\mathcal{Y}}^1$ y components. Hence the stabilisation of the magnitude of $v_4 \in TM_4$ in the projection out of $L(v_{56}) = 1$, as the $D(1)_B$ symmetry is broken in the early universe, will establish the observed masses for the u-quark and ν -lepton states in comparison with those for the d-quark and e-lepton states. The 'tuning' to these values may be automatic if there is some mechanism underlying the stability for the corresponding value of $L(\mathbf{v}_4) = h^2$ (see also the discussion at the end of section 9.2).

As described in section 9.3 a yet higher-dimensional form of temporal flow may be required to fully identify the *u*-quark and *v*-lepton states as $SL(2,\mathbb{C})^1$ fermions as well as to identify the second and third generation of Standard Model fermions. The structure of such a higher-dimensional symmetry of time, possibly involving an E_8 symmetry on a form of temporal flow $L(v_{248}) = 1$ of greater than quartic order, may hence be needed to address the question of the stability of $L(\mathbf{v}_4) = h^2(t)$.

As well as the fermions the masses for the W^{\pm} and Z^{0} gauge bosons from the $SU(2)_L \times U(1)_Y$ sector of the theory and the Higgs mass itself will also be established at the epoch of this phase transition. Fermion pairs such as e^+e^- might be produced via the decay of a heavy gauge boson, as associated with the Feynman vertex of figure 11.3(a), or via other underlying field exchanges of the form $\delta Y \leftrightarrow \delta \psi$ generalising from a geometric solution $G_{\mu\nu} = f(Y)$ for the spacetime geometry. However the energy density of the early universe in the form of $-\kappa T_{\mu\nu} := G_{\mu\nu} = f(\mathbf{v}_4)$ might also be converted into fermion states through underlying $\delta v_4 \leftrightarrow \delta \psi$ field exchanges, under $L(\hat{\mathbf{v}}) = 1$ as implied in the previous section.

As well as the production of such fermion pairs new interactions, for example under the full form $L(v_{248}) = 1$, may be significant in the high energy density environment of the very early universe. Such a form, involving quintic or higher order field composition terms, might involve the production of leptons and quarks in the same interactions, with potentially a mechanism for creating an asymmetry between matter and antimatter acting during this very early epoch through to the phase transition. A further possibility might involve gauge bosons, for example from a 'beyond the Standard Model' $SU(2) \subset E_8$ subgroup of the full symmetry of time (in principle identified through an explicit symmetry breaking decomposition in the form of equation 9.51 acting on the components of $L(v_{248}) = 1$, which might mediate interactions between leptons and quarks in an analogous manner to the ' X and Y ' gauge bosons of an SU(5) GUT model. The origin of this imbalance of matter over antimatter remains to be understood, but the underlying asymmetry in the directed flow of time, the parity asymmetry arising from in the choice of the $v_4 \in TM_4$ projection out of the components of $F(h_3\textcircled{0})$ in equation 9.46, a mechanism for combined lepton plus quark production or even CP violation in the quark sector, all in the context of the present theory, may play a part here.

In the standard theory by the time the temperature of the universe has cooled to 10^{12} K at around 10^{-4} seconds after the Big Bang quarks and gluons no longer form a component of a relatively weakly interacting plasma, along with leptons and photons, but become confined in hadronic states. A proton to photon ratio of around 10^{-9} to one is established at this epoch, with a negligible contribution from antiprotons. Similarly most of the initial electrons and positrons mutually annihilate leaving a residual e^- contribution, balancing the residual p^+ states, leading to the much later recombination era as electrons combine with nuclei forming neutral atoms around 372,000 years after the Big Bang, marking the origin of the CMB radiation as observed today and as described in section 12.3. An understanding of the origin of the imbalance between matter and antimatter states in the very early universe, accounting for the predominance of 'matter' states as still observed today, within the context of the present theory may also aid in the identification of the mathematical structure of the currently hypothetical E_8 action on the form $L(\mathbf{v}_{248}) = 1$, augmenting the input from the required Standard Model properties as discussed in section 9.3.

For the case of $\mathcal{X} \in \mathrm{h}_3\mathbb{O}$ with constant $L(v_{27}) = \det(\mathcal{X}) = 1$ and given a very small initial value of $|v_4|$ for a projected $v_4 \subset \mathcal{X}$ a very large value for the scalar n is permitted, as can be seen from equation 6.27. Similarly, in the context of the form $L(v_{56}) = 1$ and equation 9.46, with $v_4 \in TM_4$ projected from the Y components, a very large value of the scalar field $N(x)$ may be achieved if the dilation symmetry $D(1)_B$, with the generator of equation 13.5, is involved in obtaining a very small value of $|v_4|$. Further, if this latter value is obtained via the dilation symmetry $D(1)_\lambda$ of equation 9.30 then a very large value for either the scalar field $n(x)$ or $\beta(x)$ from equation 9.46 can result. Hence if a cosmological term $\Lambda g_{\mu\nu}$ is derived for equation 12.1 in the early universe, with the scalar Λ closely related to any of the scalars N, n or β then a temporary but rapid inflationary dynamics might be obtained.

The action of a dilation symmetry $D(1)_B$ or $D(1)_\lambda$, or some combination, may slide the value of $|v_4| = h \approx 0$, corresponding to an initial unstable 'inflationary state' in the very early universe, towards a preferred solution under $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ with a stable value for $|v_4| = h_0$ via interactions with other fields on M_4 . In the very early universe $v_4(x) \simeq 0$, correlated with a very large value for a scalar field such as $N(x)$ or $\beta(x)$, might account for inflationary phenomena, with the largest inflation driven for example by $N \to \infty$ as $|v_4| \to 0$ for $t \to 0$. After the magnitude of v_4 has grown in time the same field with small variations around $|v_4| = h_0$ might account for dark matter effects, as described in the previous section, while a residual, now small value for $N(x)$ or $\beta(x)$ might account for the dark energy term $\Lambda g_{\mu\nu}$ at the present epoch, as a greatly suppressed remnant of the early inflationary era. A field such as $N(x)$ driving inflation in the very early universe must be weakly coupled at the present epoch in order to have evaded detection in the laboratory. On the other hand the stable and complementary scalar field $h_0(x) = |v_4|$ is associated with the vector-Higgs field v_4 , giving rise to phenomena which are evident in experiments.

At the time of the phase transition $t = t_v$ the energy of the vector-Higgs field v_4 is transferred to fermion and gauge particle states under the external geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ solution. In addition to the stable value of $|\boldsymbol{v}_4| = h_0$ the masses of the fermions will be established under terms of $L(\hat{\mathbf{v}}) = 1$, as described for the case of $L(v_{56}) = 1$ in equations 9.48 and 9.49 at the end of section 9.2. Hence the low value of the cosmological constant Λ may be correlated with the low value of the neutrino mass, and the pattern of fermion masses more generally, according to the balance between the stable scalar values for $|v_4| = h_0, n, N, \alpha$ and β in equation 9.46, although again a full form such as $L(v_{248}) = 1$ may be required for the full picture.

Under the assumptions applied for FLRW models, as described in section 12.2, while for $t > t_v$ a radiation dominated solution for the line element of equation 12.5 initially emerges, for $t < t_v$ both the scale factor $a(t)$ of that equation and the conformal factor $h(t) = |\mathbf{v}_4|$ of equation 13.7 combine together to form the line element:

$$
d\tau^2 = ds^2 = \frac{1}{h^2(t)} \left[dt^2 - a^2(t) d\Sigma^2 \right]
$$
 (13.8)

In general it will be necessary to solve the 4-dimensional geometry $G_{\mu\nu} = f(Y, \hat{v})$ to determine the dynamical form of $g_{\mu\nu}(x)$ both for $t > t_v$ for the evolution from a radiation to a matter dominated universe and on to the era of dark energy dominance and potentially also for an 'inflationary' epoch for $t < t_v$, both in principle involving an evolution of the scale factor $a(t)$ driven by an effective $\Lambda g_{\mu\nu}$ term in equation 12.1 induced by a scalar component such as n, N, α or β . As described above such a scalar field might take a very large value in the early universe as balanced against $h(t) \rightarrow 0$ as $t \to 0$ under the constraint $L(\hat{\boldsymbol{v}}) = 1$. As was described for equation 13.7 the parameter t is considered to represent 'cosmic time' rather than 'conformal time', even for $t < t_v$. This convention is further justified here with the scale factor $a(t)$ incorporated into the more complete expression in equation 13.8.

However, for $t < t_v$ in addition to a possible $\Lambda g_{\mu\nu}$ term there are new features that arise in the present theory. The initial low value for $|v_4| = h(t)$ in equation 13.8 implies a relatively rapid flow of the fundamental time s through the spacetime manifold as parametrised by the cosmic time t . This property is complementary to the relative slowing of time in the later universe associated with small fluctuations to relatively high values of $|v_4|$ distributed in space, with the resulting gravitational effects ascribed to apparent regions of 'dark matter' as described for figures 13.1 and 13.2. The complementary case with much lower values of $|v_4|$ in the very early universe may imply an effective expansion of spacetime (which might also apply to a smaller degree in regions of the later universe correlating with 'voids' between galactic clusters, as discussed after figure 13.2 also in the previous section).

In the case of the very early universe the conformal scaling via the factor $h^{-2}(t)$ in equation 13.8 as $h(t)$ becomes smaller for $t \to 0$ means that intervals of the 'comoving coordinates' $\{t, r, \theta, \phi\}$ represent greater physical spacetime volumes as $t \to 0$. For this conformal geometry an infinite spacetime volume may be inscribed within a finite coordinate boundary (as might be represented for example by the 'Circle Limit' woodcuts of M.C. Escher described in [26] pp.33–34).

Allowing an infinite passage of time in the past in this way with $\tau \equiv s \to -\infty$ as $t \to 0$ may itself not help solve the 'horizon problem' since spatial volumes are dilated by the same factor. That is, the conformal diagram of figure 12.3, representing the causal structure for the evolution of the universe since the 'initial singularity' at $t = 0$, is unchanged by variation in $h(t)$ alone. As for the standard approach to solving the horizon problem it appears necessary to 'miniaturise' physical spatial displacements relative to temporal intervals at the earliest epoch via for example an inflationary dynamics of the scale factor $a(t)$, as has been applied for figure 12.4. Hence although in the present theory much more time may be available in the very early universe both $h(t)$ and the scale factor $a(t)$, which will be mutually correlated in the dynamics, play significant roles in equation 13.8.

Nevertheless, in the present theory the magnitude of any inflationary effect, in terms of the increase in the scale factor $a(t)$, and its period of duration may be somewhat different than in the original theory of inflation. Here the non-uniformity in the way that the underlying flow of time s is injected into the spacetime manifold as $h(t) = |\mathbf{v}_4|$ evolves may have consequences which partially, or even totally, remove the need for a rapid 'inflation'. A much smaller value for $h(t)$ in the very early universe relative to the present day value will also mean that the properties of physical structures are likely to be very different, compared with those of the Standard Model for example. These differing structures may also in principle imply uniform characteristics, such as 'temperature', across the initial singularity, with little if any time required to attain the high degree of 'thermal equilibrium' as observed today for the CMB radiation across the full spatial extent of the observable universe.

In assuming the vector field $v_4(x)$ to take the particular unstable value in the very early universe with $v_4(x) \to 0$ as $t \to 0$ the dynamics of the cosmic evolution, described globally through $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, will change with the phase transition at the time at which the stable value $|v_4| = h_0$ emerges at $t = t_v$. This potentially abrupt change in the nature of the dynamics may be accompanied by a reduction in symmetry, in particular regarding the effective breaking of the dilation symmetry, composed of a combination of the groups $D(1)_X$, $D(1)_B$ and $D(1)_\lambda$ described in the opening of this section for example. In the original inflation theory there is no relation between the postulated scalar inflaton field $\varphi(x)$ and the Standard Model scalar Higgs field $\phi(x)$ of particle physics. In the present theory 'inflation' in the early universe is correlated with a very small value for $v_4(x)$, while Higgs phenomena derive from the present stable vector field with $|v_4| = h_0$, with the two values of the same field v_4 related through the action of the dilation symmetries in the very early universe.

There are however some models in the literature for which inflation in the very early universe is correlated with the Standard Model Higgs field, and the properties of conformal transformations, in some way, as for example in [78, 79, 80]. These references typically incorporate a coupling between the Higgs and gravitational fields by postulating a new interaction term in a Lagrangian of the form $\mathcal{L} \sim \xi \phi^{\dagger} \phi R$, where ϕ is the scalar Higgs field, R is the scalar curvature and ξ is a new coupling parameter. In the theory presented in this paper however the Higgs sector is more intimately associated with gravity since variations in the magnitude $h(x) = |\mathbf{v}_4|$ of the vector-Higgs field $v_4(x)$ directly impact upon the external spacetime geometry via a change in the metric of the form described by equation 13.2.

The picture of the very early universe in the present theory does also have close parallels with the original inflationary models described in section 12.3. The scalar magnitude $h = |\mathbf{v}_4|$ for the initial projection of $\mathbf{v}_4 \in TM_4$ out of the components of \hat{v} in the very early universe is analogous to the initial value of the scalar field φ in inflationary theory. For example in 'old' or 'new' inflation the initial value $\varphi = 0$ becomes a 'false vacuum' as the potential $V(\varphi, T)$ is modified with the dropping cosmic temperature T until subsequently a stable condition with $\varphi = \varphi_0 \neq 0$ is achieved. In the present theory the consequences can be considered for a state with $h = |\mathbf{v}_4| \simeq 0$ in the very early universe followed by a continual range of projections of $L(\hat{v}) = 1$ over M_4 until the stable value $h = |\mathbf{v}_4| = h_0 \neq 0$ is achieved, particularly in terms of the form of the geometric solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$.

In the present theory interactions between the components of the vector field $v_4(x)$ and for example the fermion field $\psi(x)$ under the $L(v_{56}) = 1$ terms, together with gauge fields $Y(x)$ via terms in the expansion of $D_{\mu}L(v_{56})=0$, may compose a thermal system incorporating an effective temperature dependent potential $V(h,T)$. Such interactions will also generate 'drag terms' in the dynamics of the evolution of $h(t)$ leading to damping effects accompanying a possible period of oscillations about the potential minimum as energy is transferred to Standard Model particle states created and 'reheated' as the point of stabilisation with $h(t) = h_0$ at time $t = t_v$ is approached. This describes the 'phase transition' at the end of an inflationary period, leaving a residual dark energy contribution, arising for example from a much reduced and stable value for the scalar field $N(x)$, in addition to the Standard Model particle spectrum as observed today. A radiation dominated FLRW cosmology emerges at this time $t = t_v$ out of the 'Big Bang' with the initial conditions of the standard cosmological model having been set.

In beginning with $L(\mathbf{v}_4) = h^2 \simeq 0$ and converging towards $L(\mathbf{v}_4) = h_0^2$ at time $t = t_v$ via interactions under the terms of the full form of $L(\hat{\boldsymbol{v}}) = 1$ this picture is closely analogous to the second order phase transition of the 'new inflation' model described in section 12.3, involving a 'slow roll' down an effective potential slope $V(h,T)$ on the way to achieving the stable value. For the present theory the 'potential energy' associated with the original unstable value of $h \approx 0$, following the analogy of the 'false vacuum' state in the new inflation model, might itself effectively provide a direct source of an inflationary expansion.

The variation of $|v_4| = h(x)$ alone modifies an otherwise flat Minkowski spacetime via the conformal transformation $g_{\mu\nu}(x) = \theta(x)\eta_{\mu\nu}$, with $\theta(x) = h^{-2}(x)$ from equation 13.2. Via the Levi-Civita connection $\Gamma(x)$ this results in the Einstein tensor $G_{\mu\nu}(x)$ explicitly presented in equation 13.4. The shaping of the geometry $G_{\mu\nu} = f(\theta)$ from beneath in this way contrasts with the geometry $G_{\mu\nu} = -\kappa T_{\mu\nu}(\varphi)$ deriving from the energy-momentum source $T_{\mu\nu}(\varphi)$ of equation 12.25, which in turn was derived from a Lagrangian for the postulated scalar field $\varphi(x)$ in the original inflation theory. Despite this difference in origin there is a close similarity between the kinetic terms in the field $\varphi(x)$ in equation 12.25 and the first two terms in the field $\theta(x) = h^{-2}(x)$ in equation 13.4, suggesting the possibility of a similar field dynamics for the two models.

For inflationary theory, in addition to the kinetic term drag terms are also introduced into the Lagrangian, as described following equation 12.25, and relate to the physical phenomenon of post-inflation reheating during which the energy of the false vacuum is converted into interacting particles. A similar effect may arise in the present theory, with an effective potential $V(h,T)$ and 'drag terms' for the new theory deriving from interaction terms implicit in the form $L(\hat{v}) = 1$ as described above. The favoured minimum in $V(h, T)$ (which may be largely independent of the effective temperature T) will correspond to the stable value $|v_4| = h_0$, without the need to *contrive* an appropriate form for the potential $V(\varphi, T)$ as is the case for inflationary theory, since all the couplings of the present theory are effectively implied within the constraint equations 11.29.

An alternative proposal for the present theory features initial conditions with $|v_4| \gg h_0$ with potentially large fluctuations in the components of the field $v_4(x)$, perhaps accompanied by a large value for the scalar field $\alpha(x)$ from equation 9.30 providing the source of an inflationary $\Lambda g_{\mu\nu}$ term in the very early universe. Amongst the many possible solutions for $G_{\mu\nu} = f(Y, \hat{v})$ in principle arbitrarily extreme spacetime geometries may occur, but without necessarily being supported throughout the full expanse of the manifold M_4 . Such an extreme structure may describe the initial geometry in the Big Bang, where the conditions may even be somewhat 'chaotic' as an extended spacetime solution is first shaken out of the mathematical possibilities implied in the form $L(\hat{\boldsymbol{v}}) = 1$ and its symmetries, assuming the present universe to have evolved from such a state. With potentially a large range of possibilities for $|v_4(x)| \gg h_0$ the initial value for $v_4(t)$ for $t \to 0$ may be required to be fairly uniform over a spatial extent of order the Hubble radius, as for the initial value of the scalar field $|\varphi| > |\varphi_0|$ in models of 'chaotic inflation', as also described in section 12.3.

However, while there may be a range of possible 'false vacuum' initial conditions for the projection of $v_4 \,\subset \,\hat{v}$ onto TM₄ in the present theory, the 'post-inflationary' stable value of $|v_4| = h_0$, as coordinated with the parameters of the Standard Model of particle physics, may still be uniquely determined. The possible range of initial values for $|v_4| \gg h_0$ in principle implies a range of inflationary effects and a corresponding

range of properties for the later evolution of the universe, some of which may be compatible with the present day universe as actually observed, and in particular with both the horizon and flatness problems resolved as for standard inflationary theory.

This raises the question of the uniqueness of the present theory, which will be discussed more generally in the following section. For the case of $h(t) = |\mathbf{v}_4| \to 0$ as $t \to 0$ the scalar function $\theta = h^{-2}$ diverges. Hence even in this case, if θ is interpreted as the inflationary field, the present theory may be interpreted in a manner analogous to chaotic inflation, with θ effectively taking a broad range of large values in the 'primordial chaos' of the very early universe. Although these options, in relation to chaotic inflation, might be considered further, here we explore in more detail the implications of taking $h(t) = |\mathbf{v}_4| \to 0$ as $t \to 0$ as a potentially unique starting point.

Hence this general structure involving a transition from the initial condition with $v_4(x) \rightarrow 0$, as depicted in figure 13.3(a), towards the stable state with $|v_4| = h_0$, with a value which may also be determined uniquely, and as represented in figure 13.3 at stage (c), may be essentially unambiguous. The state immediately emerging from the phase transition in figure $13.3(c)$, along with ordinary matter represented by the sprinkling of points of dust, was also represented in figure $13.2(c)$, although with the spatial fluctuations of the field $v_4(x)$ neglected in the earlier figure.

Figure 13.3: (a) Beginning with $v_4(x) \approx 0$ at the temporal origin of 4-dimensional spacetime, (b) the value of $|v_4| = h$ grows, with potentially large fluctuations in both magnitude and direction, until the phase transition with (c) a stable value attained for $v_4(x)$ with small fluctuations about the components $(h_0, 0, 0, 0)$ for v_4 in the comoving cosmological frame $\{t, r, \theta, \phi\}$. (Later epochs are depicted in figure 13.2(d),(e) in the previous section).

The picture of the phase transition between (b) and (c) in figure 13.3 is analogous to the that associated with the property of ferromagnetism in a piece of iron.

The atoms in the iron can be considered as forming a lattice of a very large number of randomly oriented magnets for temperatures $T > T_c$ above the critical value. This is similar to the situation in figure 13.3(b), except that the atomic magnets would be represented by 3-dimensional spatial vectors of a uniform constant magnitude. Upon cooling to a temperature $T < T_c$ it is energetically favourable for neighbouring magnetic vectors to align, with an analogous phenomena applying for the vector field $v_4(x)$ as the stable value $|v_4| = h_0$ is attained as depicted in figure 13.3(c), with small fluctuations about the average 4-vector value of $v_4(x)$ greatly exaggerated in the diagram.

For the present theory the symmetry breaking in the phase transition, both in terms of the actions of the dilation symmetry for the magnitude of $v_4(x)$ and fluctuations in the orientation of this 4-vector, further suggests a close relationship between $v_4(x)$ and the Standard Model Higgs field. Indeed the stable vacuum value $v_4 = (h_0, 0, 0, 0)$ is precisely the same 4-vector as that in equation 8.72 (where a pseudo-Euclidean basis $\{t, x, y, z\}$ for \mathbf{v}_4 was employed), with $v^0 = h_0$, as described in subsection 8.3.3. The 'vacuum symmetry' is broken as the vector-Higgs field $v_4(x)$ takes a magnitude and particular direction in spacetime which, on average, is presumed to be essentially aligned with the preferred cosmological frame parametrised by comoving coordinates $\{t, r, \theta, \phi\}.$

That is, the comoving cosmological frame is aligned with the average distribution of visible matter, which in turn is presumed to have been formed and evolved in line with the underlying flow $v_4(x)$ through the spacetime manifold M_4 . The degree of correlation between local fluctuations in the flow v_4 and peculiar motions on a galactic scale at the present epoch is an open question. While the laws of physics are locally Lorentz invariant actual physical structures clearly are not, and this also applies to the large scale structure of the universe. For example a directional relative blueshift and redshift for the detected CMB radiation depends upon the local choice of Lorentz frame for the observer. In our case these shifts are due to our local motion within our galaxy, and can be readily corrected for in the CMB maps.

As for the inflationary theories described in section 12.3 quantum fluctuations and potentially Hawking radiation in the inflationary epoch generate inhomogeneities in the very early universe which may become frozen as classical fluctuations in energy density at the end of inflation, ultimately seeding the formation of galactic structures. In the present theory these quantum effects include interactions between $v_4(x)$ and other fields, such as those for the fermions $\psi(x)$ and gauge bosons $Y(x)$ as well as scalar fields such as $N(x)$, all subject to the constraint equations 11.29 in forming the overall external geometric solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ in spacetime. Fluctuations in the value of $h(x) = |\mathbf{v}_4(x)|$ directly impact upon the spacetime geometry, as described for equation 13.2, and hence in particular may generate large scale structure when amplified as the scale factor $a(t)$ rapidly grows.

Fluctuations in the spatial components of $v_4(x)$ could also in principle have a large effect during the evolution of the very early universe as represented by the stage of figure 13.3(b), particularly for the case of a pre-inflation spatially 'miniaturised' world, with a relatively very small value of $a(t)$, as described for figure 12.4. A calculation of how such fluctuations might stir up the primordial geometry would involve taking into account all components of the field $v_4(x)$ to determine $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, rather than just the magnitude $|\mathbf{v}_4| = h = \theta^{-\frac{1}{2}}$ as was the case for equation 13.4. Such a cosmological model, with fluctuations in both the magnitude and direction of $v_4(x)$ impacting upon the large scale structure, would differ from the forms derived in equations 12.9 and 12.10, for G_{00} and G_{11} respectively, for which homogeneity and isotropy were assumed, unless a statistical average is taken for the large scale structure conforming to those assumptions.

As the vector field $v_4(x)$ stabilises through the phase transition to the stage depicted in figure 13.3(c) small residual variations in the components of $v_4(x)$ might still remain, and be found to be finely grained on the scale of the observable universe. This residual fingerprint of the earlier fluctuations is the network of creases in the fabric of spacetime as has been described in the previous section for the same epoch as also depicted in figure $13.2(c)$. At this point, and throughout the remaining evolution of the cosmos, this small residual variation in the components of $v_4(x)$ might account for the phenomena of dark matter, and even a dark flow, as also suggested in the previous section. That is, with sufficient deviation from the assumptions of uniformity of the cosmological principle these residual variations might seed the early formation of galaxies and clusters of galaxies through gravitational merging into the courser structures observed at the present epoch. This cosmic imprint in the underlying spacetime geometry, arising from fluctuations in the very early universe, is interpreted as a manifestation of 'cold dark matter' in particular, as was described in the previous section for structures observed through to the present epoch as depicted in figure $13.2(e)$.

As noted above, on top of these geometric effects of a continuous variation in the field $v_4(x)$ in the present theory the same field is responsible for the Higgs sector in particle physics through interactions or exchanges with other fields. More generally, throughout the history of the universe quantum transitions, in the form of $\delta v_4 \leftrightarrow \delta \psi$ or $\delta \hat{v} \leftrightarrow \delta Y$ field exchanges underlying the multiple possible solutions for the spacetime geometry $G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$, with the constraints of equations 11.29 applying everywhere on M_4 , will shape the evolution of the cosmos, including the epoch of the very early universe. This shaping includes both the impact of observable fluctuations as described above as well as the physical implications arising from the statistical average of the microscopic interactions.

As described in chapter 11 the direct association of the likelihood of an observable quantum event with the 'number of ways' in which the same empirical effect can be achieved, quantified in terms of the degeneracy of underlying field solutions for the same external local geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, unifies the quantum process notion of probability itself with the classical concept. While a time-ordered accumulation of probabilities in the quantum case is relevant for cross-section calculations, in the case of classical phenomena it gives rise to the second law of thermodynamics as quantified by an ever increasing value of entropy for any evolving thermodynamic state. In the present theory all such thermodynamic phenomena are played out in time and do not themselves drive an 'arrow of time', as will be clear in the following chapter.

As alluded to above the structure of the very early universe may allow sufficient breathing space for the thermalisation of the particle degrees of freedom in the epoch before the phase transition in figure 13.3, as is the case for the pre-inflation environment in figure 12.4 as described in section 12.3. However some care is needed in applying the principles of thermodynamics and statistical mechanics, familiar from their application in the flat spacetime environment of the laboratory for example, in the potentially highly curved and dynamic spacetime of the very early universe. Even basic notions such as 'temperature' or a 'black body spectrum' may be hard to define in such an extreme environment. The approach may be justified to some extent by applying thermodynamics within small spacetime regions which approximate to local inertial frames, and adopting the strong equivalence principle, given a sufficient number of 'particles' and 'particle interactions', or underlying field exchanges, within such a region to apply statistical methods. Further, the properties of the 'particles' and fields themselves in the era before the phase transition may be very different to the familiar Standard Model particles and fields that emerge out of the Big Bang.

It is also noted that the universe, and in particular the structure of the very early universe, is a *single* system for empirical study. Hence thermodynamic arguments, which consider an *ensemble* of systems each of which might form a small component within the universe, as for example employed for laboratory experiments, may not apply for the potentially unique system composing the precursor to and immediate aftermath of the Big Bang. That is the observable universe today may have evolved from state in the very early universe which is too small or simple to incorporate a statistical average, and which might in fact be dominated by the effect of a single 'fluctuation'.

Further, as was described towards the end of section 12.3, beyond the 'horizon problem' there is apparently a 'start-up problem' in the need to choreograph a vast number of spacelike separated 'bangs' along the initial singularity, either in figure 12.3 or 12.4, in order to effectively simultaneously trigger the 'Big Bang' itself at cosmic time $t = 0$. Analogous to a synchronised display of fireworks there might be range of 'temperatures' across the range of 'bangs' creating inhomogeneous initial conditions. However the Big Bang is not such a terrestrial event and there seems no reason why it should not be in the nature of the start-up to generate essentially homogeneous thermodynamic conditions over the entire spacelike hypersurface at $t = 0$, upon which local fluctuations may be identified in terms field exchanges underlying the multiple solutions for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. In particular the different nature of gravity, associated with the smooth geometry $G_{\mu\nu}(x)$, compared with the quantum phenomena associated with the internal field interactions, may play an important role in this structure.

In the immediate aftermath of the phase transition of figure $13.3(c)$ with the 'vacuum energy' being converted into Standard Model particles through transitions of the form $\delta v_4 \leftrightarrow \delta \psi$ under $L(v_{56}) = 1$, with familiar microscopic quantum properties, many more degrees of freedom may open up. The entropy content of the observable universe emerging from this epoch will depend on the reheating effects of the drag terms implicit in $L(\hat{v}) = 1$ combined with the kinetic terms implied in equation 13.4, which were discussed earlier in this section and similarly as described following equation 12.25 for inflationary theory. However, as also noted in section 12.3 the gravitational field appears to have had a very special role in the Big Bang and very early universe in being aloof from the thermalisation process.

The strong equivalence principle (as reviewed in section 3.4 and adopted above) in part demonstrates how the characteristics of gravity fundamentally differ from the other forces of nature. The properties of local inertial frames are key to the structure of general relativity, with all physical phenomena other than gravity behaving in such a frame as if gravity were completely absent, while gravity itself is described by the geometry of the extended spacetime. The differences between gravity and other physical phenomena will be significant for addressing issues for the early universe, including also the 'flatness problem' as well as the 'horizon problem' and an understanding of the role of entropy.

Within the present theory the special status of the external gravitational field further derives from the fact that it is of a quite different, 'unquantised' nature in comparison with the internal gauge $Y(x)$ and fermion $\psi(x)$ fields. The external geometry, described for example in terms of the metric components $g_{\mu\nu}(x)$ or linear connection $\Gamma(x)$, does not partake in the statistical physics of the internal fields which lies beneath continuous geometric solutions of the form $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. In the expression $-\kappa T_{\mu\nu} := G_{\mu\nu}$ the right-hand side describes the smooth external geometry, with all quantum mechanical properties of matter implicitly underlying the energy-momentum tensor on the left-hand side. Such quantum phenomena, based on an degeneracy of field solutions, can generally be described to a good approximation within local inertial frames, as was the case in sections 11.1 and 11.2. While the electromagnetic field, for example, exhibits thermal properties through the underlying field interactions the external gravitational field, being aloof from such interactions, has a very different relation with thermodynamic phenomena, and also, being unquantised, does not directly partake in quantum fluctuations.

The phenomenon of Hawking radiation, as discussed towards the end of section 11.4, arises for quantised fields in the classical curved spacetime of a black hole exterior, with the consequence for example that a black hole with mass $O(10^6)$ kg will evaporate in approximately one second. Such phenomena involve the quantum mechanical description of the vacuum but the gravitational field itself is not quantised, and lead to a study of the thermodynamic and entropy properties of black holes. Similar properties of the vacuum may arise for the present theory, since gravity is not quantised here, and also be important in the study of the thermodynamic and entropy properties of the very early universe, in particular during the inflationary period.

For the case in which the initial geometry is dominated by variation in the value of $|v_4| = h(x)$, with a metric of the form $g_{\mu\nu}(x) = h^{-2}(x)\eta_{\mu\nu}$ in equation 13.2, the spacetime geometry of equation 13.4 is conformally flat, even for arbitrarily large variations in the scalar field $h(x)$. For such a geometry the Weyl curvature tensor vanishes, $C_{\rho\sigma\mu\nu}(x) = 0$, consistent with the proposal of the Weyl curvature hypothesis as motivated and described towards the end of section 12.3. Hence this observation may account for the 'cosmological problem' concerning the extraordinarily special state of the Big Bang to 1 part in $10^{10^{123}}$ (according to [26] p.777) as required for the low entropy initial conditions which underlie the subsequent evolution of the cosmos consistent with second law of thermodynamics.

Through interactions and fluctuations of the form $\delta v_4 \leftrightarrow \delta \psi$, in particular with the transfer of energy from the vacuum to Standard Model particle states towards the end of the inflationary period corresponding to figure $13.3(c)$, a non-conformally flat geometry will emerge incorporating Weyl curvature, and hence the propagation of gravitational waves for example, as well as Ricci curvature. As described in section 12.3 (with reference to [26] section 28.8) the entropy of the gravitational field might be expressed in terms of the degrees of freedom of the Weyl curvature and hence contribute to the increase in entropy from this time. As also described in section 12.3, following equation 12.23, all FLRW models are consistent with $C_{\rho\sigma\mu\nu}(x) = 0$ but require something like an initial period of inflation to explain why observations are consistent with $k = 0$, that is with spatial flatness. Similarly for the present theory an inflationary evolution for $a(t)$ in equation 13.8 in the very early universe may relate to this observation.

As described in section 11.3, and depicted in figure 11.10, the three coupling parameters of the Standard Model gauge group $SU(3)_c \times SU(2)_L \times U(1)_Y$ approximately converge at an energy scale of $O(10^{15})$ GeV. This unification scale will mark a significant threshold in the early universe, and it will be important to understand how it relates to the epoch of the phase transition in figure 13.3(c) for the present theory. The interplay between the dilation symmetry, such as $D(1)_B$, and electroweak symmetry, together with the nature of their breaking, will also be key, as described shortly after equation 13.7 with reference to equation 13.6. The electroweak symmetry $SU(2)_L \times U(1)_Y$ is broken by its action on $v_4 \in TM_4$, with the stable value for the magnitude $|v_4| = h_0$ arising out of the Big Bang at $t = t_v$. For $t < t_v$, and in particular for $t \to 0$ with $|v_4| \ll h_0$ via the action of the dilation symmetry, the properties of the electroweak symmetry and the Higgs sector more generally will be somewhat different, for example with regards to the pattern of particle masses. To address the complete symmetry breaking picture it will be required to explicitly identify the electroweak symmetry $SU(2)_L \times U(1)_Y$ within the full E_7 , or E_8 , symmetry of the full form $L(\hat{v}) = 1$, in relation to the dilation symmetries, such as $D(1)_B$, and the $SL(2,\mathbb{C})^1 \times SU(3)_c \times U(1)_Q$ symmetry already identified, completing the development of these structures described in chapters 8 and 9.

As also alluded to towards the end of section 11.4 for a theory of 'quantum gravity', with the degrees of freedom of the gravitational field quantised, significant effects are expected at the Planck energy scale $E_P = \left(\frac{c^3 \hbar}{G_N}\right)^2$ $G_{\!N}$ $\int_{0}^{\frac{1}{2}} \simeq 1.2 \times 10^{19} \text{ GeV}$. For any description of the very early universe in the context of such a theory all classical field concepts in turn fail at epochs earlier than the Planck time $t_P = \left(\frac{G_N \hbar}{c^5}\right)^2$ $\left(\frac{C_{N}\hbar}{c^{5}}\right)^{\frac{1}{2}}\simeq 5\times 10^{-44}$ seconds. However the energy scale E_P is considered to be of no special significance for the present theory and the time scale t_P , representing for example the extremely early universe, in principle presents no barrier for this theory. Hence the nature of the universe down through epochs at arbitrary cosmic times $t < t_P$ might be studied within the context of the present theory. This leads essentially to two broad possibilities as depicted in figures 13.4(a) and (b). In these diagrams $t = t_v$ (presumably with $t_v \gg t_P$) denotes the epoch of the phase transition at which there is a convergence to the average value $L(\mathbf{v}_4) = |\mathbf{v}_4|^2 = h_0^2$, as represented in figure 13.3(c). For either figure 13.4(a) or (b) the epoch of the 'Big Bang' can be identified with the time $t = t_v$ or more generally with the period from $t = 0$ to $t = t_v$ and the state emerging at that latter time.

In the first case for figure 13.4(a) the time $t = 0$ can be considered to be the moment at which an extended 4-dimensional spacetime world first emerges out of the forms of the pure temporal flow s as identified through the geometric relation $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. This is the point in time at which extended and potentially infinite 3-dimensional spatial hypersurfaces may be identified as an offshoot out of the multidimensional form of temporal flow $L(\hat{v}) = 1$ and a spacetime geometry with metric $g_{\mu\nu}(x)$ established, although with a significant deviation from flatness possible both

Figure 13.4: Two scenarios for the relation between the temporal origin of the universe and the fundamental flow of time with (a) $s \to -\infty$ for $t < 0$ and (b) $s \to -\infty$ for positive values of $t \to 0$. The width of each figure for $t > 0$ represents the spatial scale factor $a(t)$ as a function cosmic time t, neither of which are drawn to scale.

for the 4-dimensional curvature and for the 3-dimensional hypersurfaces. Considering a time $t > 0$ in figure 13.4(a) and retracing the temporal flow backwards the time $t = 0$ marks the point at which the geometrical interpretation in terms of a 4-dimensional extended manifold, supported by the mathematical structure and symmetries of the form $L(\hat{\mathbf{v}}) = 1$, completely breaks down.

Before $t = 0$ in figure 13.4(a) the parameter t no longer represents a coordinate on the manifold M_4 , while the fundamental flow of time $s \to -\infty$ continuous without any limit as expressible through a general form $L(\hat{v}) = 1$, as always, but without any projection of $v_4 \in TM_4$ components onto an extended manifold. Here, as depicted for example in figure 13.3(a), we have considered the case with $v_4(x) \simeq 0$ in the very early universe. In the context of figure 13.4(a) beginning with $v_4(x) = 0$ at $t = 0$ with $|v_4(x)| = h(x)$ generally growing with $t > 0$ in the very early universe, as depicted in figure 13.3(b), the time $t = 0$ could be considered as the epoch at which a fragment of temporal flow under $L(\hat{v}) = 1$ is 'syphoned off' into the thereby created spacetime manifold M_4 . However it is also conceivable that this point of spacetime creation at $t = 0$ can be accompanied by arbitrary values for $v_4(x) > 0$, in principle even with $|v_4| \gg h_0.$

The width in both figures 13.4(a) and (b) represents the spatial scale factor $a(t)$ of equation 13.8, under the presumption of a solution with $a(t) \to 0$ as $t \to 0$ and some form of inflationary expansion leading up to $t = t_v$, not drawn to scale. The behaviour of the ratio $\frac{a(t)}{h(t)}$, and in particular whether this fraction tends towards zero, infinity or is finite as $t \to 0$, will be significant for understanding the nature of the geometry of the manifold M_4 in this limit, according to the spacetime structure described by equation 13.8. The geometry in this limit will also be important in relation to the horizon problem and the 'start-up problem' as discussed for figures 12.3 and 12.4 in section 12.3. This might be best approached via a redefined cosmic time parameter such as \bar{t} with $\delta \bar{t} = \frac{\delta t}{h(t)}$ $\frac{\partial t}{h(t)}$ and with the line element of equation 13.8 correspondingly replaced by:

$$
d\tau^2 = d\bar{t}^2 - \frac{a^2(t)}{h^2(t)} d\Sigma^2
$$
\n(13.9)

As has been discussed earlier, care is needed for the meaning of 'cosmic time' for the epoch $t < t_v$, whether parametrised by t or \overline{t} , since physical clocks will be of a somewhat different nature for the very early universe, and indeed do not exist in any form for $t < 0$. In any case a more complete theory is required to avoid the dangers of speculating on the number of angels that might be accommodated upon the head of a pin, as noted at the end of section 12.3.

The above comments also apply for the scenario depicted in figure 13.4(b), for which necessarily $v_4(x) \rightarrow 0$ at the spacelike edge of the manifold M_4 in the past at $t = 0$. For this second picture the relation between the flow of the fundamental time parameter s and the 'cosmic time' coordinate t is sketched in figure 13.5.

Figure 13.5: For the scenario depicted in figure 13.4(b) the projection $v_4(x) \in TM_4$ converges to zero for $t \to 0$ in the very early universe. While $t = 0$ marks a coordinate boundary to the 4-dimensional spacetime M_4 the range of the fundamental temporal flow $-\infty < s < +\infty$ is tucked away and entirely contained within this manifold. Adopting the approximate components $(h, 0, 0, 0)$ for v_4 in the comoving frame the phase transition $t = t_v$ marks the point at which $v^0 = |\mathbf{v}_4(x)| = h(t) = h_0$ stabilises.

For this scenario if $v_4(x) \in TM_4$ converges to zero in an appropriate manner as $t \to 0$ then as $s \to -\infty$ the spacelike hypersurface at $t = 0$, potentially an 'initial singularity' as $a(t) \rightarrow 0$, is never attained and all of the fundamental flow of time $-\infty < s < +\infty$ is absorbed into the extended spacetime M_4 of the universe. With $s \to -\infty$ without limit at the temporal coordinate origin on M_4 the structure for $t < t_v$ in figure 13.4(b) and 13.5 might be pictured poetically as the bottomless waterfall at the end of time. With the familiar structures of the Standard Model of particle physics emerging in the phase transition, from this epoch and for all times $t > t_v$ the fundamental time flow s is equivalent to both the proper time τ and also the cosmic time t for idealised observers in the context of an FLRW cosmological model, as described near the opening of section 13.1.

For the case of the scenario depicted in figure 13.4(b) the present day universe is, in a sense, infinitely old in terms of the fundamental time parameter s. However for the picture in *both* figure $13.4(a)$ and (b) the physical and mathematical structures can be traced back to arbitrarily early times for $s \to -\infty$, with the difference being that for (a) physical structures are no longer defined for $t < 0$ while for (b) parameter values $t \leq 0$ are outside the domain of the underlying temporal flow s. In both cases physical structures relating to the Standard Model of particle physics arise out of the Big Bang at $t = t_v$. This is the point in time at which we can effectively 'start the clock' with $s \equiv \tau \equiv t$, as might be measured through familiar physical processes, now determined to stretch back through around 13.8 billion years of cosmic evolution. Such an apparent temporal origin for the laws of physics in our 4-dimensional world may be necessary for consistency with an environment supporting biological life at the present epoch. Here we refer in particular to the second law of thermodynamics which implies the universe is still evolving away from the particularly low entropy state conceivably corresponding to the nature or uniformity of the gravitational field in the very early universe, as described above.

For either scenario depicted in figure 13.4 the cosmic evolution itself is a feature of the full macroscopic 4-dimensional spacetime $G_{\mu\nu} = f(Y, \hat{v})$, as shaped by microscopic field interactions in the form of the local degeneracies of fields underlying the possible solutions, consistent with the constraint equations 11.29, as described in chapter 11. As discussed in section 11.4 in combining gravitation with quantum theory the notion of a 4-dimensional spacetime solution of general relativity takes precedence over the 1-dimensional propagation of an apparent quantum state, with the latter described in terms of a local time coordinate, hence also circumventing the 'problem of time' encountered by some approaches to quantum gravity. As also concluded in section 11.4 the nature of probability in quantum processes is essentially the same as that for classical systems, at heart formulated in terms of the 'number of ways' that an empirical effect may be produced.

On the large scale, with many underlying degrees of freedom, the interplay of both quantum and classical statistical phenomena will contribute to the shaping of the cosmological solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. This solution will also incorporate macroscopic contributions to the geometry in the form of $G_{\mu\nu} = f(Y)$ of equation 5.20, by comparison with Kaluza-Klein theory as described in section 5.1, and of the form $G_{\mu\nu} = f(\hat{\boldsymbol{v}})$ of equation 13.4 from variations of $|\boldsymbol{v}_4| = h(x) = \theta^{-\frac{1}{2}}(x)$ in the projection of $L(\hat{\mathbf{v}}) = 1$ onto M_4 , as described in this chapter. A correspondence with the techniques of 'renormalisation' in quantum field theory might in principle be developed in order to study the relation between the macroscopic external geometry and the underlying 'bare' fields, as has been described in section 11.3.

The question then concerns how the combination of all of the above geometrical and statistical factors in determining a solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, with $T_{\mu\nu} := G_{\mu\nu}$ providing the interpretation of equation 12.1, might collectively account for the observed cosmic evolution, compatible in approximation with the assumptions of the FLRW models and the metric form of equations 12.5 and 12.6, together with the large scale galactic structures. While observations of the latter structures require an apparent 'dark matter' component, on the largest scale the solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ is required to account for the apparent effects of 'dark energy', for example in the form of an effective cosmological term $\Lambda g_{\mu\nu}$ in the Einstein field equation. As for the earlier inflationary epoch, the modern era parameter Λ may not be entirely constant, but with any variation such that $(\Lambda g^{\mu\nu})_{;\mu} \neq 0$ exactly compensated by an apparent effective energy-momentum tensor with $T_{\epsilon;\mu}^{\mu\nu} \neq 0$ consistent with $G^{\mu\nu}{}_{;\mu} = 0$ and equation 12.1. This possibility was alluded to in the previous section in the discussion regarding table 12.1, and with reference to a similar observation for equation 5.41. In the present theory the total energy-momentum tensor $T_{\mu\nu} := G_{\mu\nu}$ is defined to incorporate any possible 'dark energy' cosmological term, and indeed the full solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$.

In describing the overall cosmological evolution in the spirit of the FLRW models the metric of the line element in equation 12.5 or 13.8 underlying the full 4 dimensional solution $G_{\mu\nu} = f(Y, \hat{v})$ will incorporate the expansion of the universe, including that of the present day, in terms of the scale factor $a(t)$. The perspective adopted here is not that the universe is expanding now because it was expanding in the past, analogous to the kinematic propagation of the flight of a cannonball from one moment to the next along its trajectory, in either case raising the question of how it was set in motion in the first place. Rather here the very early universe is conceived of as one particular region of the full four-dimensional spacetime manifold M_4 , which happens to exhibit properties such as $a(t) \to 0$ and $h(t) \to 0$ as the coordinate parameter $t \to 0$, consistent with the overall $G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$ external geometry solution.

This is analogous to thinking of the Earth as being in orbit around the sun at the present day not as a kinematic consequence of the fact that it was in orbit one year ago or a billion years ago but since the 4-dimensional spacetime trajectory, featuring an approximately elliptical orbit, exists as a geodesic solution for a 4-dimensional Schwarzschild spacetime. In fact since the Bianchi identity $G^{\mu\nu}{}_{;\mu} = 0$ implies geodesic motion, as described for equation 5.36 in section 5.2, the full spacetime geometry of an entire planetary system can be conceived of as a particular 4-dimensional solution for $G_{\mu\nu}(x)$. The idealised Schwarzschild solution itself describes an infinite and eternal 4dimensional spacetime with $G_{\mu\nu}(x) = 0$ everywhere, except for the point at the centre of spherical spatial symmetry, with the metric of equation 5.49. While the components of this Schwarzschild metric are constant in time but vary as a function of the radial coordinate r via the factors of $(1 - 2G_N M/r)$, the geometry of the Robertson-Walker metric for an FLRW cosmological solution is independent of the spatial coordinates but varies with the time coordinate through the scale factor $a(t)$. Both cases represent full 4-dimensional spacetime geometries.

In the present theory both $a(t)$ and $h(t)$ in the line element of equation 13.8 shape the geometry for the very early universe with $t < t_v$, with a correlated evolution of these parameters associated with a period of inflation. The comparison, earlier in this section, with the 'new inflation' model represents an analogy for the present theory, however the 'slow roll' down from $h(t) \simeq 0$ for $t \to 0$ to the stable average value $h(t_v) = h_0$ may or may not end with a series of 'oscillations' as the minimum of the effective potential $V(h,T)$ is achieved. In any case, given the correlation between $a(t)$ and $h(t)$, it is conceivable that spatial regions with residual small positive fluctuations $h(x) > h_0$ may have 'inflated' a little longer leaving a value of the scale factor $a(x)$ also slightly larger than the average value at the end of inflation.

For the large scale evolution of the observable universe for any time $t > t_v$, with quantities averaged over each 3-dimensional spatial hypersurface, the value $h(t)$ = h_0 remains constant and stable while $a(t)$ continues to increase, parametrising the expansion of the universe as sketched in figure 12.2. However on the local scale of galaxies and galactic clusters it is the correlated distribution in space of $a(x)$ and $h(x)$, initially established at $t \simeq t_v$, that might be associated with dark matter. That is, evolving forward to the present day, the effects of dark matter might be attributed to regions with small fluctuations of $h(x) > h_0$ together with a correlated spatial profile in $a(x)$, rather than simply the conformal scaling alone of equation 13.2 as suggested following figure 13.1 in the previous section. In this way, generalising from equation 13.8 for spacetime variation of h and a , the line element takes the form:

$$
d\tau^2 = \frac{1}{h^2(x)}dt^2 - \frac{a^2(x)}{h^2(x)}d\Sigma^2
$$
 (13.10)

This structure opens up a greater degree of independence between the temporal and spatial components of the metric, with for example $g_{00}(x)$ relatively low and $g_{ii}(x)$ for $i = 1, 2, 3$ relatively high in spatial regions where both $h(x) > h_0$ and $\frac{a(x)}{h(x)}$ are relatively high, which in this sense is more reminiscent of the Schwarzschild solution of equation 5.49, and which also may have geometric properties more characteristic of a distribution of an apparent form of 'matter' than variation of $h(x)$ alone.

While, given an initially flat spacetime, the purely conformal action of $h(x)$ only generates Ricci curvature, the metric of equation 13.10 will generate both Ricci and Weyl curvature contributions extended throughout the spacetime manifold M_4 , both in regions of galactic clusters and the voids between. Having the variation of both $h(x)$ and $a(x)$ in equation 13.10 increases the potential to match the observations of galactic motions and rotation curves, together with gravitational lensing effects, as a candidate for dark matter in interaction with the distribution of ordinary baryonic matter. On the yet larger scale of cosmological evolution these contributions to the dynamics of the universe might also be compared with the measured density parameters Ω_D and Ω_B , in addition to Ω_{Λ} , as introduced in section 12.2, as part of a global fit to the cosmological data.

In summary, the large scale structure and cosmological evolution of the universe are to be identified generally as aspects of a full 4-dimensional solution for the spacetime geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. There is no presupposition of a flat spacetime manifold. In projecting an extended 4-dimensional spacetime M_4 out of the full multi-dimensional form $L(\hat{v}) = 1$ of the fundamental temporal flow s large scale geometric distortions might be expected, which in turn may correlate with the observations ascribed to inflation, dark energy and dark matter, as reviewed in the previous chapter. There remains, of course, the need for a more complete theory and a much more thorough analysis, but in the meantime the possible variation of the magnitude of the projected 4-vector $v_4 \in TM_4$ and the identification of several scalar fields α, β, n and N from the components of $L(v_{56}) = 1$ indicates the potential for the application of the present theory to these cosmological questions.

The above discussion applies for the geometry of the 4-dimensional spacetime manifold M_4 whether in the context of the scenario depicted in figure 13.4(a) or (b). However, compared with the first scenario of figure $13.4(a)$ that in figure $13.4(b)$ is more symmetric in time in the sense that both the limit for $s \to -\infty$ as well as for $s \to +\infty$ is incorporated within the 4-dimensional spacetime solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, as depicted in figure 13.6.

Figure 13.6: As parametrised by the fundamental temporal flow s the spacetime manifold underlying the physical universe can be of infinite extent without boundary in time as well as in space for the scenario of figures 13.4(b) and 13.5.

We inhabit a region of this eternal and infinite spacetime located within the period of several tens of billions of years following the phase transition at $t = t_v$ during which complex physical structures supporting biological life can be found, as represented in $13.6(e)$ and corresponding to the epoch of figure $13.2(e)$. It may be that both the far future through to $s \to +\infty$ as well as the far past with $s \to -\infty$ may become progressively less structurally varied and eventful compared with the present epoch. For $s \to +\infty$ the universe may evolve into a relatively uneventful interplay between slowly evaporating massive black holes and thermal radiation, as depicted in figure 13.6(f), while for $s \to -\infty$ there may be an equally uneventful asymptotic progression with $|v_4| \to 0$, as depicted in figures 13.3(a) and 13.6(a). In this picture a physical understanding of the structure of the universe for both $s \to +\infty$ and $s \to -\infty$ may be equally open to study.

On the other hand there then remains the question concerning the reason why

the universe should exist at all. In the context of the scenario in figure $13.4(a)$, as for the standard cosmological models discussed in section 12.2, in tracing the cosmological history back through the epoch of the very early universe this question can be phrased in terms of the cause of the Big Bang and the nature of the temporal origin of the universe itself. However, with everything, including the Big Bang happening in time, and with all physical structures in the universe for the present theory built entirely upon the notion of the one-dimensional flow of time s , there will still remain the question of the foundation of this apparently fundamental temporal entity itself, a question which applies equally for the scenario in figure $13.4(b)$. This will form the topic for the following chapter. In the meantime, in the following section, we consider the extent to which the properties and laws of physics of the universe, as depicted for example in figure 13.6, might or might not be unique within the conceptual notions and mathematical constraints of the present theory.

13.3 Uniqueness

In this section we consider several topics concerning the extent to which the particular properties as empirically observed for the universe might be either necessarily determined or down to chance, within the context of the present theory, beginning with the values of the large scale cosmological parameters. Without a full understanding of their underlying origin, the fact that the density parameters are observed to take the values $\Omega_{B_0} = 0.050 \pm 0.002$, $\Omega_{D_0} = 0.265 \pm 0.011$ and $\Omega_{\Lambda_0} = 0.685 \pm 0.017$ [44], as reviewed in section 12.2, mutually within an order of magnitude or so of each other at the present epoch, given the apparent possibility for each to range over many orders of magnitude, is striking. On the other hand given a universe dominated by either a cosmological constant Λ or matter density ρ term the Friedmann equation 12.9, particularly for the $k = 0$ case with $H^2 = \frac{1}{3}(\Lambda + \kappa \rho)$, shows that the Hubble parameter is essentially determined by Λ or ρ respectively, and is clearly not an independent observable.

At the present epoch for our universe, which is consistent with $k = 0$ and with the cosmological term beginning to dominate, it is then to be expected that $\Lambda \sim R_H^{-2}$ are of the same order of magnitude, where R_H is the Hubble radius introduced in equation 12.4. This observation is a direct consequence of the field equation 12.1 which leads to the dynamical solution for the metric structure of equation 12.5, including the case of a Λ dominated universe. If the history of the scale factor $a(t)$ is such that H_0^{-1} approximates the current age of the universe, which is the case for our universe with the cosmic evolution sketched in figure 12.2, then R_H will be of the same order as the scale of the observable universe hence in turn relating $\Lambda^{-\frac{1}{2}}$ to this scale given the dominance of the Λ term at the present epoch.

While the constant Λ in equation 12.1, considered as a geometrical effect, has the length dimension of L^{-2} the equivalent 'vacuum energy density' $\rho_{\Lambda} = \Lambda/\kappa$ has the dimension $ML^{-3} \equiv L^{-4}$ and may be directly compared with the mass density ρ for both ordinary and dark matter. It should be noted though that on substituting \dot{a}^2 $\frac{\dot{a}^2}{a^2}+\frac{k}{a^2}$ $\frac{k}{a^2}$ from equation 12.9 into equation 12.10, for the Λ dominated case, it is the extra factor of $-\Lambda$ in the second equation which leads to a positive value for \ddot{a} in the case of positive vacuum energy density $\rho_{\Lambda} > 0$. This difference can be interpreted as a consequence of the effective 'equation of state' for dark energy, with $p_{\Lambda} = -\rho_{\Lambda}$, as also implied in equations 12.18 and 12.19.

While the Λ term is beginning to dominate, the present day values of ρ_{Λ} and ρ_M (with the latter composed of both baryonic and dark matter together) still have a comparable impact on the large scale cosmological dynamic equations. The value of $\rho_{\Lambda} \simeq 5.8 \times 10^{-27}$ kg m⁻³ is apparently uniform in space and time, and hence the same locally as well as globally, and can be compared with the global value of $\rho_M \simeq$ 2.6×10^{-27} kg m⁻³, which includes a contribution from $\rho_B \simeq 0.4 \times 10^{-27}$ kg m⁻³, at the present epoch. However the value of ρ_B changes significantly with the cosmic epoch while local values for density of ordinary baryonic matter, such as for the planet Earth with $\rho_{B_E} \simeq 5,500 \text{ kg m}^{-3}$, are much more stable in time. The magnitude of the stable terrestrial ratio of $\rho_{B_E}/\rho_{\Lambda} \simeq 10^{30}$ then provides a measure of the apparently very different nature of ordinary matter and dark energy.

Another well known apparently natural 'large number' in physics concerns the order of magnitude of the Standard Model couplings of particle physics in comparison to the strength of the gravitational interaction. For example the ratio of the classical electrostatic force between an electron and a proton to the classical gravitational force between them has a value of $O(10^{39})$ to one. This empirical observation was also alluded to near the opening of section 5.2 in motivating the need to introduce practical normalisation factors in studying the implications of equation 5.20 in the laboratory environment. In the present theory, with general relativity and the Standard Model relating to the external and internal structures of $L(\mathbf{v}_{56}) = 1$ respectively, the relative strengths of the corresponding interactions in general will be related to the identification and interpretation of equation 5.20, which in turn is related to the geometric structures of Kaluza-Klein theory. The fact that the gravitational field is not 'quantised', and hence does not exhibit the running coupling of figure 11.10 for example, further distinguishes gravity from the Standard model forces in the present theory.

The differing strengths of gravitational and internal gauge forces should also be connected in some way with the relative magnitudes of the components, such as those of the vector v_4 or spinors ψ , within $v_{56} \in F(h_3\mathbb{O})$ of equation 9.46 in the symmetry breaking projection over M_4 . With the forms of matter and dark energy also relating to structures within $L(v_{56}) = 1$ and its symmetries these relative magnitudes for the components of v_{56} may also determine the widely differing local values of ρ_{B_E} and ρ_{Λ} , with the value of Λ possibly relating to the value of a scalar field such as $N(x)$, $n(x)$ or $\beta(x)$ projected out of the components of $F(h_3\mathbb{O})$. Hence the symmetry breaking pattern of E₇ on $L(\mathbf{v}_{56}) = 1$ down to an external SL $(2, \mathbb{C})^1$ acting on $\mathbf{v}_4 \in TM_4$ together with the internal structures and the details of the projection of the components of $v_{56}(x)$ over M_4 may underlie the empirical observation of both of the above large numbers.

It is the relative weakness of gravity that allows structures to form on large scales, from the formation of stable planetary bodies through to clusters of galaxies. On the other hand the relative strength of the internal forces shapes the smaller scale structures from terrestrial geology down through biological and chemical systems to the elements of particle physics. Immersed in the relatively small scale biological structures our perspective is one of a spacetime which is flat to a very good approximation upon which an apparent 'force of gravity' is observed to determine the motion of material

objects such as apples and cannonballs, as described in section 12.1 and before the bullet points in section 13.1.

For all of the reasons of the above paragraph a world in which the elementary interactions of the Standard Model of particle physics are of a much greater strength than that of gravitation is 'anthropically' favoured. Such a preference may correlate with a certain value, or range of values, for the magnitude $L(\mathbf{v}_4) = h^2(x)$ in the projection of the $v_4 \text{ }\subset v_{56}$ components onto TM_4 , and in turn underlie the empirical observation of $\rho_{B_E} \gg \rho_{\Lambda}$ locally on Earth and for concentrations of baryonic matter generally. With the global density ρ_M of the combination of ordinary and dark matter (assuming 'dark matter' to behave in a similar manner to baryonic matter in this respect) declining from a potentially divergent value in the initial singularity and seemingly asymptotically approaching zero in the future, the observation that $\rho_{M_0} \sim \rho_{\Lambda}$ are of the same order at the present epoch, an apparently arbitrary point in cosmic time, appears to be essentially coincidental.

This determination of $\rho_{M_0} \sim \rho_{\Lambda}$ has some analogy with the observation that $r_m \sim r_s$ at the present epoch, where r_m and r_s are the apparent sizes of the moon and the sun respectively as viewed from the Earth. The value of r_m has been declining since the formation of the Earth-moon system as the average distance between these two bodies increases by $O(1 \text{ cm})$ every year due to the nature of the gravitational interaction between the two bodies. Hence the present situation in which the moon is apparently just large enough to create a total solar eclipse is largely coincidental. However there are anthropic arguments, with the distance of the Earth from the sun being in the 'habitable zone' (not too near and too hot while also not too far and too cold) and similarly for the distance of moon from the Earth resulting in a magnitude of tides which may have aided the early development of biological life, which make such an apparent coincidence much more likely. Similarly there may be underlying anthropic reasons involving the nature of cosmological evolution which make the observation of $\rho_{M_0} \sim \rho_{\Lambda}$ more probable during a cosmic epoch supporting biological life.

In summary, in the present theory the observation of $\rho_{B_E} \gg \rho_{\Lambda}$ is expected to be correlated with the observation that Standard Model forces are far greater in strength than the gravitational force. Indeed the cosmological term $\Lambda g_{\mu\nu}$ might be considered effectively as a geometric perturbation within general relativity as the large scale external spacetime structure M_4 is identified through the projection of $v_4 \in TM_4$ out of $L(\hat{v}) = 1$, rather than an internal effect underlying the solution $G_{\mu\nu} = f(Y, \hat{v})$. The relation between Λ and the Hubble radius, described near the opening of this section, may also hint at a geometric origin for the cosmological term. As well as the great difference in strength, the rather different nature of gravitational compared with internal gauge forces is further emphasised in the present theory by the fact that the degrees of the freedom of the gravitational field, describing the external spacetime geometry, are not quantised here.

With dark matter associated with the external geometric consequences of a variation in the magnitude $h(x) = |v_4|$, as described in the previous two sections, here the dark sector in general is associated with locally 'weakly interacting' general relativistic effects. In the context of a solution for the full 4-dimensional cosmological geometry 'density parameters' such as Ω_D and Ω_{Λ} may not have the same meaning as for the standard theory, since for example the above candidate for 'dark matter'

may not evolve in time in the same way as the baryonic matter density and the above origin for 'dark energy' may not imply a constant value for Λ . In any case the present observation of $\Omega_{D_0} \sim \Omega_{\Lambda_0}$ may be a consequence of a correlated geometric origin for the associated empirical effects, collectively arising from the warping of the manifold M_4 in the projection out of $L(\hat{\mathbf{v}}) = 1$, while the proximity of Ω_{B_0} to these values may in part be due to an element of coincidence as described above in the analogy with the apparent size of the moon and the sun.

In developing the present theory further gravitational or material effects may be derived in studying the general structure of $G_{\mu\nu} = f(Y, \hat{v})$ beyond those of the empirically observed baryonic matter and dark sector. It would seem to require a significant coincidence if all such effects are of a measurable magnitude and hence observable at the present epoch. If there are physical consequences of the relation $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ which have not yet been detected, and which may be beyond the reach of any practical observation, this itself would partly account for the apparent coincidence of $\rho_{M_0} \sim \rho_{\Lambda}$. That is, these two latter quantities may form a subset of effects which collectively comprise a list of mutual contributions to $G_{\mu\nu}(x)$ at present, with a range of other potential terms having much lower density parameters and hence remaining undetected. For example if the empirically deduced cosmological term itself had been just one order of magnitude smaller it would have been far harder to detect. On the other hand while a contribution to the cosmic evolution of the form $R_{\mu\nu} = \lambda(t)v_{\mu}v_{\nu}$ (as described in section 13.1 and listed in the final column in table 12.1) has not been observed such a term, with a sufficiently low value of $\lambda(t)$, might in principle form part of the large scale spacetime solution. With a larger range of such contributions it is more likely for any two of them, such as ρ_{M_0} and ρ_{Λ} , to take similar values and be mutually observable.

In chapter 11 the degeneracy of multiple possible local field solutions underlying the spacetime geometry $G_{\mu\nu} = f(Y, \hat{v})$ was described as the origin of indeterministic quantum phenomena in general. However in terms of constructing a solution there may also be a degeneracy in terms of the average projected values of the components of for example $v_4(x)$ and $\psi(x)$ out of $F(h_3\mathbb{O})$ globally over M_4 . In this case there may be only a small certain range of values which lead to physical properties of matter capable of supporting life as we know it. Even with this degree of anthropic selection to 'dial in' certain ratios of the components of $v_{56} \in F(\mathrm{h}_3\mathbb{O})$, via the dilation symmetries described in the opening of section 13.2 for example, since only a small number of 'free' parameters are involved in the projection of $v_4 \in TM_4$ under the fixed structures of $L(v_{56}) = 1$ the theory would still be highly constrained, and hence in principle still capable of making predictions which might be tested. In section 13.2 the point of view was adopted that the interactions under the constraints of the theory are such that a unique stable value of $|v_4| = h_0$ is achieved, resulting in a phase transition in the very early universe, implying an even greater degree of predictability for the theory.

For the scenario described in figures 13.4(b), 13.5 and 13.6 at the end of the previous section a unique asymptotic condition with $h(t) = |\mathbf{v}_4| \to 0$ as $t \to 0$ has also been presumed. The ensuing progression from $h(t) \to 0$ to the stable value $h(t_n) = h_0$ was compared with models of 'new inflation'. It is also possible to consider a range of starting conditions for $h(t) < h_0$ as $t \to 0$ and even a broad range of values $h(t) > h_0$ for $t \to 0$, as might be associated with the scenario depicted in figure 13.4(a), and

evoking a comparison with models of 'chaotic inflation'. In turn a range of long term cosmological conditions might emerge out of the subsequent phase transition, even given the same stable value for $h(t_v) = h_0$, and hence in principle with a degree of anthropic selection implied for our own habitable universe.

This raises the question of the degree of uniqueness regarding other aspects of the theory. With the general form of the function $L(\mathbf{v})$ determined, as described in section 2.1, it is a well defined mathematical problem to identify particular forms and then consider the reasons why certain of these may be significant for the physical world. Two such significant forms that we have identified are $L(v_{56})$ with an E₇ symmetry acting on elements of $F(h_3\mathbb{O})$ and $L(\mathbf{v}_4)$ with the Lorentz symmetry acting on the 4dimensional tangent space TM_4 on the base manifold. With respect to the larger form symmetry breaking over M_4 identifies the smaller form via the chain $L(\mathbf{v}_4) \to L(\mathbf{v}_{10}) \to$ $L(v_{27}) \rightarrow L(v_{56})$, rather like a sequence of Russian dolls, with a corresponding chain of subgroups $SO^+(1,3) \subset SL(2,0) \subset E_6 \subset E_7$ as summarised in table 9.1 in the opening of section 9.3.

Alternatively a progression of forms $L(v_4) \rightarrow L(v_9) \rightarrow L(v_{27}) \rightarrow L(v_{56})$ aligned with the subgroup chain $SO^+(1,3) \subset SL(3,\mathbb{C}) \subset E_6 \subset E_7$ might be considered by expanding the Lorentz symmetry action of $SL(2,\mathbb{C})$ on $v_9 \equiv \mathcal{X} \in h_3\mathbb{C}$ in equation 7.35 to an $SL(3,\mathbb{C})$ symmetry of the 9-dimensional form $L(\mathbf{v}_9) = \det(\mathcal{X}) = 1$. The $sl(3,\mathbb{C})$ Lie algebra basis of equation 8.33 explicitly demonstrates how this structure is naturally embedded within the $SL(3, \mathbb{O}) \equiv E_6$ action on the 27-dimensional space $h_3\mathbb{O}$ at the next stage of the sequence.

At either end of this chain it may be asked why these two particular forms are selected out of a large array of possibilities – why the projection should be onto a 4-dimensional spacetime manifold and why the highest-dimensional form $L(\hat{v}) = 1$ should be represented by a quartic expression in 56 dimensions with an E_7 symmetry. It could be considered whether further worlds, different to our own and of course not observable by us, could be created out of other possible mathematical forms of $L(\mathbf{v})$. That is, whether the forms $L(\mathbf{v}_4) = h^2$ and $L(\mathbf{v}_{56}) = 1$ are largely identified as choices that agree with our world, or whether either or both of these are determined by physical stability or mathematical symmetry arguments for example.

By extension from the 3-dimensional model world of section 2.2 and figure 2.3 with an SO(3) symmetry one way to construct a 4-dimensional world would be to embed 3-dimensional spatial hypersurfaces within a 4-dimensional base manifold with local tangent vectors $v'_4(x)$ satisfying the form $L(v'_4) = (v^1)^2 + (v^2)^2 + (v^3)^2 + (v^4)^2 = 1$ (appending one dimension to the model case of equation 2.14) with an SO(4) symmetry. However while geometric curvature and even particle trajectories might be defined in such a world, given the local $SO(4)$ symmetry on M_4 there is no consistent definition and distinction of a 'temporal' direction compared with 'spatial' displacements. In principle one of the four dimensions could be arbitrarily declared to represent an apparent temporal component, however due to the nature of the symmetry between the four components the causal structure on M_4 would not be well defined.

However given that the 4-dimensional manifold arises as a multi-dimensional manifestation of the ordered 1-dimensional flow of time itself and the necessity for the temporal causal structure to be retained on the manifold, the form $L(\mathbf{v}_4) = (v^0)^2 (v¹)² - (v²)² - (v³)² = 1$ of equation 5.1 is naturally preferred. As described in

section 5.3 the metric of Lorentz signature implied in this form locally defines a 'light cone' structure on the extended manifold, which hence distinguishes timelike from spacelike directions on M_4 . The symmetry preserving this form $L(\mathbf{v}_4) = 1$ is the noncompact Lorentz group $SO^+(1,3)$, which is also the group which preserves the causality structure on a Minkowski spacetime [81]. The local symmetry of the 3-dimensional spatial hypersurfaces is identified as the $SO(3)$ subgroup of the Lorentz group. These observations concerning causality appear decisive in favour of the 'pseudo-Euclidean' form $L(\mathbf{v}_4) = 1$ over the above 'Euclidean' alternative $L(\mathbf{v}'_4) = 1$.

A more accurate model for chapter 2 would have involved the 3-dimensional Lorentz group group $SO^+(1, 2)$ acting on the form $L(v_3) = (v^0)^2 - (v^1)^2 - (v^2)^2$ projected onto the tangent space of M_3 , as a subgroup of the full symmetry $G = SO^+(1, 4)$ acting on a 5-dimensional Lorentzian form for example. This would be necessary to identify timelike and spacelike vectors and temporal causality on the base manifold M_3 . However dealing with the simplified Euclidean model in chapter 2 was sufficient to demonstrate the relation between the external and internal symmetry in the present theory, with the same conceptual ideas applying for the case of the real world with the 4-dimensional Lorentz group on the base space M_4 as described in section 5.1, leading to the connection with Kaluza-Klein theory as also discussed in that section.

As well as the local Lorentz symmetry, which also holds to a good approximation on for example the scale of the solar system in the case of our world, we may also consider whether the base manifold M_n is required to have $n = four$ spacetime dimensions. If we attempt to construct another possible world using similar reasoning to that presented in this paper then we would expect something similar to general relativity, that is gravitation, to arise out of the geometrical properties on the base manifold of the world, independent of its dimension. One important factor may be that while for a 4-dimensional spacetime base manifold robust, stable planetary orbits around a massive object, such as a star, are to be found in the solutions to the equations of gravitation, this does not arise for other dimensions of base space.

This was shown to be the case for the motion of a body near a massive gravitating object, as the source of the Schwarzschild solution for general relativity, in an n-dimensional spacetime by F.R. Tangherlini in 1963. While for $n = 4$ the metric solution takes the form of equation 5.49 the functional form of $g_{\mu\nu}(x)$ depends on the value of n. Although theoretically a circular orbit may be permitted in some cases for $n > 4$, the slightest perturbation, for example from the impact of a 'meteor' or the gravitational influence of a third body, would cause the 'planet' to wander out of orbit and into a path forever receding to larger distances or spiralling inwardly until colliding with the central 'star'. The same conclusion, that a stable orbit is only possible for $m = 3$ spatial dimensions, was also found by Paul Ehrenfest in 1917 for Newton's theory of gravity in which the gravitational potential is determined as a solution of the m-dimensional Poisson equation (which was introduced for the $m = 3$ case above equation 3.75 in section 3.4).

Clearly the stability of the elliptical orbit of the Earth around the sun is necessary for life on our planet in our world, although this does not imply that the equivalent stability is absolutely necessary for life in another world with $n \neq 4$ spacetime dimensions. For example the 'chemistry' in such a world would be vastly different from our own and the relative time scale for the development of life structures to the time scale of planetary motions may also be vastly different – potentially allowing a civilisation to evolve out of the primordial chemical soup stirring on the planet in the time it takes to glance past a star, even assuming such an encounter with a low entropy source in the form of stellar 'nuclear' energy is necessary.

Having then decided upon the 4-dimensional Lorentz group on the M_4 manifold to break the full symmetry there are still issues concerning the degree to which assumptions made about the form of the linear connection $\Gamma(x)$ on M_4 are necessary. In the present theory the base manifold M_4 derives from the projected form $L(\mathbf{v}_4)$ and hence regarding the local geometry there are a range of local coordinate systems at any point on M_4 for any of which the metric has a Minkowski form $g_{\mu\nu}(x) = \delta^a_{\mu} \delta^b_{\nu} \eta_{ab}$. In turn, in deriving from an $SO^+(1,3)$ -valued connection form the linear connection $\Gamma(x)$ will be metric compatible, as also discussed in section 5.3.

As for general relativity, in a 4-dimensional spacetime there is enough freedom in general coordinate transformations to set $\partial_{\lambda}g_{\mu\nu}(x) = 0$ at any given point $x \in$ M_4 . However, as reviewed in section 3.4 (and also following equation 13.2) general relativity goes further by asserting the 'equivalence principle' – according to which the gravitational field can be transformed away at any given point, that is, a local inertial frame can be constructed such that not only $g_{\mu\nu}(x) = \delta^a_{\mu} \delta^b_{\nu} \eta_{ab}$ and $\partial_{\lambda} g_{\mu\nu}(x) = 0$ but also $\Gamma(x) = 0$ for any given $x \in M_4$. This means for example that there exists everywhere a local coordinate system in which a geodesic trajectory as described in equation 3.76 for the 4-vector **u**, with components $u^{\mu} = dx^{\mu}/d\tau$, takes the simple form $du/d\tau = 0$. Since the torsion tensor T, with the components of equation 3.60, must be zero in all coordinate systems if it is zero in any of them, such a linear connection $\Gamma(x)$ is necessarily torsion-free.

In the present theory an extended frame of reference throughout which both $\partial_{\lambda}g_{\mu\nu}(x) \simeq 0$ and $\Gamma(x) \simeq 0$ is preferred for the anthropic purpose of framing an environment for perception. This is certainly consistent with the existence of local coordinates such that the equivalence principle holds with both $\partial_{\lambda}g_{\mu\nu}(x) = 0$ and $\Gamma(x) = 0$ at any given $x \in M_4$, and taking the torsion to be zero may be a very good approximation. However given the mathematical basis for what we are taking as the act of perception it seems perhaps artificial to impose the extra restriction on the connection that it should necessarily be torsion-free or, further, require that the strong equivalence principle in general should hold. It may be that there is a non-vanishing contribution to physical phenomena from torsion which has so far been beyond the reach of observation – for example any contribution to the connection coefficients $\Gamma^{\lambda}{}_{\mu\nu}(x)$ asymmetric in the $\{\mu,\nu\}$ indices would have no effect on the simple geodesic motion of equation 3.76 – and neglecting it has therefore been of no consequence.

This is also the case in general relativity where setting the torsion equal to zero acts as a simplifying assumption. Both in general relativity and the present theory the linear connection Γ is a metric connection with $\nabla g = 0$, but this does not imply that the torsion should vanish. In the Einstein-Cartan version of general relativity the more general geometry with finite torsion is considered (with extra dimensions such a generalisation is also significant for the Kaluza-Klein theories reviewed in section 4.2). In this case while the spacetime curvature is still related to the energy-momentum of matter through the Einstein equation the torsion is a function of the spin current of matter. Unlike curvature the torsion does not propagate in the matter-free vacuum

and the two theories are identical in such an environment. Further, given that the spin density is small for ordinary matter in the universe the two theories have been experimentally indistinguishable, and this itself justifies adopting $T = 0$ as a simplifying assumption.

In the present theory it is an open question whether the linear connection Γ is necessarily symmetric and torsion-free, and if so to explain why this is the case. More generally the question regards whether the spacetime geometry and forms of matter consistent with $G_{\mu\nu} = f(Y, \hat{v})$ contain the structures of torsion and a spin current. In the meantime as for general relativity the assumption $T = 0$ may be adopted to simplify some of the mathematical expressions, with in particular the Levi-Civita connection of equation 3.53 hence being employed. This is analogous to adopting the simplifying conditions which underlie the Robertson-Walker line element of equation 12.5 in order to study models for the evolution of the universe as a whole, even though the assumptions of the cosmological principle clearly do not hold exactly. The degree to which the large scale structure deviates from the conditions of homogeneity and isotropy may itself not be a uniquely restricted property of the universe.

Even for the $T = 0$ case, in constructing the external geometry in terms of the internal fields, as well as the 10 components of the Einstein tensor in the form of $G_{\mu\nu}$ = $f(Y, \hat{v})$ itself the full 20 independent components of the Riemann curvature tensor may also explicitly depend on those fields with $R^{\rho}_{\sigma\mu\nu} = f(Y, \hat{\mathbf{v}})$. This will include the Weyl tensor components $C^{\rho}_{\sigma\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ and hence the identity $C_{\sigma\mu} = C^{\rho}_{\sigma\mu\rho} = 0$, as described towards the end of section 3.3, as well as the Bianchi identity $G^{\mu\nu}_{;\mu} = 0$, will also apply implicitly for the internal fields. The 10 degrees of freedom of $C^{\rho}_{\sigma\mu\nu}(x)$ are still considered to represent the 'vacuum' in the sense that they complement the 10 degrees of freedom of $-\kappa T_{\mu\nu} := G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu}$ in the decomposition of equation 3.69 for the full Riemann tensor $R^{\rho}_{\sigma\mu\nu} = f(\tilde{Y}, \tilde{\boldsymbol{v}})$.

In classical general relativity while matter is identified with the content of the energy-momentum tensor $T_{\mu\nu}$ the vacuum geometry with $G_{\mu\nu} = 0$ and $C^{\rho}{}_{\sigma\mu\nu} \neq 0$ still carries energy, in the form of gravity waves for example, as also discussed after equation 5.44 in section 5.2. Hence energy can propagate through the 'vacuum' of spacetime even when not expressed in terms of any underlying internal fields. In contrast it is also possible in the present theory that there may be fields on M_4 , for example from some of the components of $F(h_3\mathbb{O})$ underlying the form $L(v_{56}) = 1$, at least in some regions of spacetime, that may not directly contribute to the spacetime structure of $G_{\mu\nu} = f(Y, \hat{v})$ at all, and hence which do not carry energy-momentum.

While for general relativity the Einstein equation 3.84 can be derived from the Einstein-Hilbert action of equation 3.79, it can be shown, as demonstrated by Cartan, Weyl and others, that the most general divergence-free symmetric 2-index tensor constructed from the metric and its derivatives up to second order is a linear combination of $G_{\mu\nu}$ and $g_{\mu\nu}$ (see for example [82] appendix II). This consideration itself leads to Einstein's equation 3.84 and 12.1, with Λ a free parameter, as essentially the only admissible field equation for a geometric theory of gravity consistent with a divergencefree energy-momentum tensor on the right-hand side. Regardless of the method of derivation the significance of the Einstein equation derives largely from the contracted Bianchi identity $G^{\mu\nu}_{;\mu} = 0$, which then necessarily applies to the energy-momentum tensor. On the other hand symmetries in the apparent distribution of matter can be

employed to assist the search for solutions, with for example equation 12.2, with ρ and p being functions of cosmic time only, being the most general energy-momentum tensor consistent with the assumptions of the cosmological principle, as described in section 12.2.

In the present theory energy-momentum is defined through $T_{\mu\nu} := G_{\mu\nu}$. While $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ incorporates ordinary matter, a possible $\Lambda g_{\mu\nu}$ term, dark matter phenomena and the structure of the very early universe collectively into an apparent $T_{\mu\nu}(x)$ there may be further geometric or material phenomena, arising out of the internal fields or their interactions, which have not yet been detected. This possibility was alluded to earlier in this section in the discussion of the observation of $\rho_{M_0} \sim \rho_{\Lambda}$, as exemplified by a potential contribution originating from a term of the form $R_{\mu\nu} = \lambda(t)v_{\mu}v_{\nu}$, and now incorporates also the possibility of finite torsion. The potential for new phenomena will be of particular interest if yet higher-dimensional forms of $L(\hat{v}) = 1$ are considered, with a corresponding larger symmetry, which may also be needed to fully account for known Standard Model particle phenomena.

The Lie group E_6 was originally selected as a candidate symmetry for the full form of $L(\hat{\mathbf{v}}) = 1$, in the context of the present theory, in part since it is already of well known interest in relation to the observed gauge groups of elementary particle theory, as reviewed in section 7.3. However, it was primarily chosen for detailed study as it acts on a relatively high dimensional vector space, with 27 dimensions compared with the four on the base manifold M_4 , and stands out as exhibiting particularly rich mathematical structures, involving for example the triality symmetry of the octonions and three interlocking actions of $SL(2, \mathbb{O})$ as described in chapter 6, through which to channel the temporal flow via the components of v_{27} under the constraint of the 27-dimensional cubic form $L(v_{27}) = 1$. Expressed in terms of the octonions, which themselves form the largest of the normed division algebras, this form of $L(\hat{v}) = 1$ provides a unique structure. The existence of elaborate mathematical properties within the substructures of the E_6 symmetry acting on $h_3\mathbb{O}$ matrices is perhaps the reason why E_6 stands out as a kind of significant mathematical resonance amongst other possible symmetries of temporal forms in yet higher dimensions. By comparison for example higher-dimensional spacetime symmetries $SO^+(1,m)$, acting on quadratic Lorentzian forms with an arbitrarily large number m of spatial dimensions, arguably exhibit a somewhat less elaborate structure.

In section 9.2 the analysis was extended to the smallest non-trivial representation of E_7 realised as an action on the 56-dimensional space $F(h_3\mathbb{O})$ preserving a certain quartic form $L(v_{56}) = 1$ and incorporating the octonions in two independent h3O subspaces. Building upon the properties identified for the symmetry breaking of the E_6 action on h₃^{σ} described in chapter 8, the structure of the broken E_7 action on $F(h_3\mathbb{O})$ when projected over M_4 has a number of properties reminiscent of the Standard Model, as summarised in equation 9.46. However it is still very much an open question as to which other symmetry groups should perhaps be considered and what observable effects they may have on our own world. These effects might be manifested in particle physics phenomena through the prediction of additional states, or the determination of the properties of known states, which might be observed in high energy physics experiments.

The hypothetical extension to an E_8 symmetry on a 248-dimensional form

 $L(v_{248}) = 1$, as described in section 9.3, would in principle be large enough to incorporate the full set of known Standard Model states, including all three generations of the fermions. Given that E_8 is the largest exceptional Lie algebra, terminating the chain of Dynkin diagrams depicted in figure 9.1 of section 9.2, such a form of temporal flow might uniquely complete the sequence of extensions listed in table 9.1 of section 9.3. As a continuation of that sequence, and also in particular to contain the non-compact Lorentz group $SO^+(1,3)$ as the local symmetry of the causal structure on M_4 as discussed earlier in this section, this may involve the non-compact real form $E_{8(-24)}$ as described for equation 9.50.

The present theory is based on the observation that the one-dimensional progression in time, via the elementary arithmetic properties of the real line \mathbb{R} , can be expressed in terms of variables in an arbitrary number of dimensions. In principle the same observation might be applied to each of the n real components underlying an *n*-dimensional form of temporal flow $L(\mathbf{v}_n) = 1$. For the case of the orthogonal group $O(n)$ in the limit $n \to \infty$ certain properties related to the octonions make various calculations more tractable. In his study of the homotopy groups of the topological group $O(\infty)$ in 1957 Raoul Bott discovered the isomorphism $\pi_{i+8}(O(\infty)) \cong \pi_i(O(\infty))$. Such period 8 structures, which are also seen for Clifford algebras and known generally as 'Bott periodicity', are all closely related to the 8-dimensional octonions. Similar periodicity structures may become relevant in the exploration of higher-dimensional forms of $L(\hat{\boldsymbol{v}}) = 1$, for which octonion elements explicitly feature, and might even be important for calculations relating to the degeneracy of solutions underlying $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, involving a higher-order nesting of field redescriptions, which underlie quantum and particle phenomena.

The progression towards higher-dimensional forms of $L(\hat{\mathbf{v}}) = 1$ described above may uncover a uniquely determined mathematical structure. Given also that the 4 dimensional Lorentzian form $L(v^4) = h^2$ projected into M_4 may necessarily provide the means of breaking the higher symmetry the laws of physics observed in our universe might in turn be uniquely determined. Even in this case our universe does not represent the unique manifestation of such a world, but rather one of a vast number of possible solutions for the external geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, built upon an underlying degeneracy of local internal field descriptions as expounded in chapter 11. While events at a HEP experiment, such as depicted in figure 10.1, exhibit the intrinsic structure of quantum uncertainty, the spectrum and properties of the particles identified in the laboratory may be unique. On the other hand in principle there might still be solutions for multiple universes with a range of large scale cosmological structures depending on the nature of the overall $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ solution, in particular with regard to the apparent conditions in the very early universe.

It nevertheless will be required to carve out of the full form $L(\hat{\mathbf{v}}) = 1$ a universe like ours, such as depicted in figure 13.6 and described at the end of the previous section, incorporating all of the observed large scale structure and the phenomena of the Big Bang. Regardless of the degree of uniqueness of such a world, in being constructed out of the multi-dimensional forms of temporal flow, it derives in turn from the priority of one-dimensional temporal flow as the underlying basis of the universe. Hence the conceptual question remains regarding the origin of this one-dimensional structure itself, as we consider in the following chapter.

Chapter 14

The Origin of Time

14.1 Two Loose Ends in the Theoretical Sciences

The aim of theoretical physics at a fundamental level could be described as a program to uncover the basic scientific principles of the world, the consequences of which encompass all empirical phenomena. From the objective point of view the existence of the universe, and its matter content, began with the Big Bang and evolved according to equations of motion, as governed by the fundamental principles, for billions of years as the matter condensed into galaxies, stars and planets, some of which are conducive for biological life, until eventually conscious observers such as ourselves in turn evolved, with the ability to contemplate the world and the cosmos around us. Two of the most pressing kinds of questions raised by this picture concern the nature of (1) the Big Bang and (2) conscious life:

- (1) What can we say about the universe before the Big Bang? How and why does the Big Bang occur? How is spacetime itself created? Can the 'initial singularity' be avoided? What determines the particular initial conditions? How is matter created and what determines its properties? Why are the laws of physics the way they are?
- (2) Given that a material universe is created and set in motion subject to the physical laws, how is it possible to mould the conscious experiences of observers, aware of themselves and the world around them, out of inert, lifeless, material substance of a seemingly qualitatively entirely different nature?

It seems inevitable that any physical theory must be founded on a 'loose end' concerning the basic elements of the theory. This is the case whether these basic entities consist of particles, fields, strings, spacetime, extra dimensions, or some combination of these or further concepts, and is generally justified on the grounds that 'one has to start somewhere'. A similar argument could be made for the present theory founded on the concept of time. This paper has presented the mathematical development of this theory, beginning with the general form of temporal flow $L(\mathbf{v}) = 1$ as deduced

for equation 2.9, through the construction of a physical world in space and time for comparison with observations, leading to a discussion of the possible uniqueness of this structure in the previous section – which addresses some of the points of item (1) above. However, no matter how far progress might be made with the elucidation of empirical phenomena the theory is incomplete so long as there remains the question regarding the origin of temporal flow itself, as represented by the loose end on the left-hand side of figure 14.1.

Figure 14.1: Beginning with the notion of one-dimensional progression in time, via the general mathematical form $L(v) = 1$, both an extended spacetime manifold and the physical bodies perceived within it are in turn derived. The two loose ends concern respectively the origin of time itself and the subjective experience of the observer.

With the basic entity having such a simple structure, namely a one-dimensional ordered flow of time modelled by the real line \mathbb{R} , this first loose end is particularly striking for the present theory. By comparison a theory founded for example upon the basic entities of a set of fields in spacetime begins with a great deal of structure, and can to a large extent be considered as a study of the phenomenology of fields in spacetime. However here since the fundamental temporal flow, represented by the real line, cannot be readily decomposed into simpler elements it is very natural to raise the question of its origin, and in turn there is a greater sense of obligation to address the issue of a foundation for the present theory.

Given a description of the physical world, whether founded on the notion of time or other basic concepts, containing bodies which can be observed, the second loose end, as depicted on the right-hand side of figure 14.1, regards the question of how it is possible for an entity to be *aware* of an observation. This question concerns the issue of how 'we', as beings conscious of observations and thoughts, are embedded within the structures of the world. The physical structure of the organic brain is closely associated with this latter loose end as an apparent vehicle for self-reference capable of encoding subjective experiences within the physical world. In this section we consider how such a structure might be modelled or explained in terms of mathematical or physical elements, before returning to the first loose end of figure 14.1.

The idea that conscious experience can arise out of physical structures on the spacetime manifold M_4 should not be too controversial since it is essentially implied in most approaches to fundamental physics. If based on a quantum field theory, as applied in the Standard Model of particle physics for example, all properties of matter ultimately arise from the properties of the basic fields and their mutual interactions. Hence the microscopic properties of matter underlie the structure of macroscopic objects in the world including both inanimate objects such as rocks and pencils as well as biological structures such as flowers and brains. The self-reflective, self-conscious activity of the human brain must therefore be supported by the underlying elements of the theory and the structures which they generate in spacetime. This is essentially the case for any physical theory, since it is evident that beings conscious of experience arise in the same world as described by the theory. Both aspects of this world, that is the subjective mental phenomena as well as the objective material phenomena, are then in principle amenable to theoretical analysis.

On the practical side, since the early history of computing, with devices designed or constructed first of mechanical and later electronic components, comparisons have been drawn between 'artificial intelligence' and the workings of the naturally occurring physical structure of the brain. Indeed, the design of a computer as envisaged by Alan Turing in the 1930s and 1940s was based on modelling the action of the human mind with the ambition to 'build a brain' out of electronic components. This came with the significant advance in the design whereby programs as well as data could be stored in symbolic form, allowing both to be modified and manipulated by the 'universal machine'. On the more philosophical side Turing demonstrated that there are questions involving the performance of a universal machine which are intrinsically 'non-computable' for the device. Turing also came to the conclusion that the actions of a human brain are 'computable'; with such thought processes then being amplified through the actions of the human body.

The notion of computability for physical devices has a close parallel in the field of pure mathematics, regarding in particular the demonstration by Kurt Gödel a few years earlier that propositions can be constructed in an arithmetical calculus which are intrinsically unprovable within the calculus. It is this latter analysis we consider here in order to then describe a model for a self-referencing subjective state.

Proposition VI of Gödel's 1931 paper, On Formally Undecidable Propositions of Principia Mathematica and Related Systems I [83] can be paraphrased: 'All consistent axiomatic formulations of number theory include undecidable propositions'; that is, there are true statements of number theory which its methods of proof are too weak to demonstrate. The argument can be applied to any calculus (that is a formal system consisting of a set of axioms and rules of inference) powerful enough to express the basic arithmetic (with addition and multiplication) of the natural numbers $(0, 1, 2, \ldots)$. Hence any such formal system is 'incomplete'. The essential idea employed by Gödel was to find a way to use mathematical reasoning to explore mathematical reasoning itself (see for example [84, 85]).

Following a chain of deductions which begins with a construction known as 'Gödel numbering' a formula called G (after Gödel) is derived which is the mirror image within the arithmetical calculus of the meta-mathematical statement that: 'The formula G is not demonstrable'. Gödel was able to show that if in fact G is demonstrable then its formal contradictory $\sim G$ (i.e. 'not G') would also be demonstrable, leading to an obvious inconsistency. He proved that if the formal system is consistent then G is formally *undecidable*; that is, neither G nor $\sim G$ can be deduced from the axioms and rules of the calculus.

It can however be seen by meta-mathematical reasoning that G is in fact a true proposition of the calculus. Hence G is a true arithmetical formula and in fact expresses a certain property of all natural numbers. Hence an arithmetical truth has been discovered which can not be deduced formally from the axioms and rules of inference of the calculus. Any calculus incorporating arithmetic is incomplete in this way (in the original historical context this signalled the demise of David Hilbert's challenge to prove the contrary). Although we are free to simply add G as an extra axiom for the formal system, in this case a different true undecidable arithmetical formula G' could be constructed from the augmented calculus. Again, adding G' as an axiom we would still be able to construct a true undecidable G'' , and so on; that is, for any augmented set of axioms and rules it will always be possible to construct further undecidable propositions – the calculus is 'essentially incomplete'.

The essential points of Gödel's theorem for our purposes are summarised here:

- The symbols, axioms, rules, theorems and general expressions of a calculus or formal system capable of expressing arithmetic can be mapped onto a subset of the integers by Gödel numbering.
- Meta-mathematical statements about expressions of the calculus are associated with a mirror image within the arithmetic itself.
- Assuming that the calculus is consistent, formulas such as G can be constructed which can be shown to be true while being formally undecidable – it is not possible to prove either G or \sim G within the calculus.
- Augmenting the calculus with new axioms such as G leads to a new calculus for which new undecidable formulas such as G' can be found; completeness of arithmetic can not be achieved, it is 'essentially incomplete'.
- The consistency of the calculus can not be proved from within the system, but it can be demonstrated by meta-mathematical reasoning outside the system.

We next ask how the above considerations may be of relevance in the theoretical sciences and in particular in relation to the theory investigated in this paper. The general mathematical form $L(\mathbf{v}) = 1$ was derived in equations 2.1–2.9 on considering the notion of progression in time to have a structure isomorphic to the algebra of the real numbers \mathbb{R} , including the basic arithmetic operations of $+$ and \times . Since the natural numbers N are embedded as a subset of the real numbers the mathematical calculus concerned with $L(v) = 1$ is certainly sufficient to express the usual rules of arithmetic for the non-negative integers. Further, in developing this physical theory certain mathematical structures arising from the forms and symmetries of $L(\mathbf{v}) = 1$ have been taken to be isomorphic to the structures that we perceive in the physical world.

It is a world in which we find both natural and manufactured machines and devices which are in some cases capable of expressing statements about mathematics, and in particular about the kind of mathematical calculus that underlies the world. Since the physical world can be expressed in mathematical terms capable of describing the behaviour of objects and devices in the world exhibiting for example structures (such as the human brain) powerful enough to perform arithmetic operations and support states of self-reference, then it seems that 'formally undecidable propositions' must inevitably arise in the application of these mathematical structures. We may then consider the possibility that the manifestation of such mathematical phenomena in the world is in the form of our own conscious experience of being in an 'undecided state', with the above list of five points correlated with the corresponding list below:

- There is a necessary isomorphism between the physical structure of everything in the material world, including brains, and mathematical structures expressible in the calculus underlying the expression $L(\mathbf{v}) = 1$.
- The human brain is capable of making meta-mathematical statements about structures deriving from the mathematics of $L(\mathbf{v}) = 1$, which therefore necessarily have a mirror image in structures deriving from the $L(\mathbf{v}) = 1$ calculus itself.
- We experience questions we can ask of ourselves in making a choice, such as "Shall I pick up the pen or the pencil in front of me?" as being undecidable (that is, we cannot predict our own future actions).
- In making a choice, for example in picking up the pencil, we find ourselves in a new state for which a further horizon of similarly undecidable questions perpetually arise.
- Our experiences are organised and synthesised into a self-consistent and coherent awareness of the world.

This is indeed, of course, very far from being a definitive analysis of the phenomenon of our conscious experience in the world. The intention here is rather merely to consider the close analogy with the elements that go into the construction of Gödel's theorems. That there may be a more significant relation between these two cases is suggested by their close structural similarity, the fact that they are both grounded in mathematical considerations involving self-reference and the fact that potentially highly complicated mathematical expressions arise in both cases. We observe further that in considering a choice it is precisely our 'undecided' state that we are aware of.

For this preliminary discussion of this phenomenon in the context of the present theory we proceed with the following simple experiment. For clarity of exposition the discussion is presented in terms of my experiences in the world, where my and I refer to any individual, such as the person currently reading this text. I can place, for example, a pen and a pencil on the table in front of me and allow myself to deliberate for several seconds over the question "shall I pick up the pen or the pencil?", while filtering out other thoughts as far as possible. In performing such an experiment the experience is one of initially having an awareness of being in an 'undecided' state, in which I may 'change my mind' several times almost as if compelled along on a wave of reasoning guided by practical or aesthetic judgements concerning, for example, the utility of the pencil or the colour of the pen, and then, quite suddenly, as if I have to let go, I find myself in the 'decided' state of having chosen the pencil and hold it in my hand (in fact, the more casually or lazily I make the choice the more it feels determined by the rational course of the world, including subconscious processes, with my conscious deliberation being a kind of internally reflecting resistance to that flow). That we can readily do this kind of 'thought experiment' and attempt to observe what happens when the choice is made serves to emphasise just how central the phenomenon of conscious decision making is in the world. A general physical theory of the world should then ideally have something to say about this phenomenon or be able to offer a good reason why it does not.

Here we comment on the fundamental difference between questions we can ask of the kind "will the apple fall off the tree?" and of the kind "shall I pick up the apple?". The former question about the external world, not involving self-reference, is 'undecided' to the extent that we lack the relevant knowledge about the physical state of the objects concerned – we simply await the resolution of the question as carried externally in the inertia of the world (and with a similar interpretation applying for the outcome of indeterministic quantum processes, as depicted in figure 11.13(b) for example). For the latter question regarding whether or not to pick up the apple, in attempting to predict our own future action based on our internal thoughts we are conscious of falling over ourselves in search of the answer until we experience the resolution.

To proceed further we consider a self-referential mathematical system R which is assumed to be correlated with a physical brain state. For such a given formal system R in principle a large number n of undecidable propositions G_i , with $i = 1 \ldots n$, might be formulated; as represented in figure $14.2(a)$. If any one of these is taken to be absorbed into the mathematical structure as a new axiom then a new formal system R' with a new horizon of internally undecidable propositions G'_{i} arises, as depicted in figures $14.2(b)$ and (c).

Figure 14.2: Expansion of a formal system R as the 'undecidable' proposition G_i is incorporated as a new axiom.

From the subjective point of view the system R represents a self-reflective conscious state of mind, which is in constant interaction with the subconscious mind and the world beyond, which are also represented by mathematical structures and provide a reservoir of information and data in the environment E which might enter conscious thought as represented by figure 14.2(b), effectively corresponding to a realisation of the truth of G_i . The subjective correlate of a single step $R \to R'$ is considered to be the experience of making a choice. The overall mathematical structure with the progression $R \to R' \to R''$ drawn out through the interaction between the conscious state R and the broader environment E is outlined in figure 14.3.

The essential feature of figures 14.2 and 14.3 is that any change in the system R, due to the interaction between R and the mathematical forms of E, results in a *progression.* The all-encompassing mathematical environment E in figure 14.3 can be thought of as a static sculpture of mathematical objects. Within this structure the self-referential systems with $\dots R \subset R' \subset R'' \dots$ carve out a one-dimensional ordered progression. A given self-referential state R' within this sequence absorbs the state R accompanied by one of its undecidable propositions G_i , now included as an axiom within R' , with respect to which R represents the 'past'. Similarly R' is itself in turn contained within R'' with the latter state incorporating a resolution of an undecided

Figure 14.3: In terms of mathematical objects the formal system R corresponds to a subset of a larger environment E which provides the source of information and data for the progression depicted in figure 14.2.

proposition of the state R' , that is one of the G'_{i} , and hence representing the 'future'. Along with this terminology, with for example the self-reflective state R' corresponding to the 'present' experience, incorporating R in the past and drawn towards R'' in the future, the structure represented in figures 14.2 and 14.3 is postulated as a model for subjective *temporalisation*.

To follow the above analogy with Gödel's theorem closely then would be to say that our experienced state of being undecided finds resolution by absorption into a new state in which a particular choice, or corresponding new 'axiom', is included. The possibilities to incorporate further new axioms in the attempt to resolve a perpetual state of undecidability leads to an ordered progression (incorporating $\dots G, G', G'' \dots$ into $\ldots R$, R' , $R'' \ldots$ respectively) which is therefore structurally identical to, and proposed as the origin of, our experience of temporality.

It is important to emphasise here that it is the mutual association of the R states in this ordered series that has itself a temporal structure. It is not a question of being situated at R' for example and asking how it is possible to move on to R'' , since for something to move presupposes an already existing flow of time with respect to which the motion takes place. Rather it is the unambiguously ascending logical order of this series itself which, having a structure that can be mapped onto and modelled by the one-dimensional ordered real line R, reveals the form of time itself. We have 'time' already in its pure and simplest essence as an ordered progression in this abstract series.

From the objective point of view a state R corresponds to a limited physical system in the world, correlated in particular with features of a physical brain. Via physical processes new data can be introduced through the interaction between, effectively, the conscious brain and the subconscious brain, as well as with the rest of the physical world, as will be described further below and in the following section. Future actions are not fully determined by or contained within the self-reflective state R itself. The physical process of the subconscious intervening in the conscious deliberation, as modelled by the progression from R to $R + G_i$ and attainment of the new state R' depicted in figure 14.2, correlates with the subjective experience of 'letting go' after a period of 'falling over oneself' in debating whether to pick up the pen or the pencil. From an internal subjective point of view the conscious mind is ignorant of the choice

until it is made and the individual finds himself holding the pencil rather than the pen for example, contributing to his sense of temporalisation. The passage of time and conscious experience more generally may feel somewhat mysterious since we do not generally perceive the objective structures and interactions represented in figure 14.3, only their internal subjective correlate.

Evidently our thoughts are not really as clear cut or 'binary' as suggested in the example above when confronted with a simple choice such as "Shall I pick up the pen or the pencil in front of me?". It is not that we are really considering an isolated possible future state corresponding to each alternative G_i in figure 14.2(a). Rather, there is an enormous ensemble of possible future states which may be divided into two sets, each with a vast range of members:

A) {I shall pick up the pen + X} B) {I shall pick up the pencil + Y } (14.1)

where X and Y each refer to possible features of a state of mind in addition to whether or not I hold the pen or pencil respectively. The idea here is not so much that individual undecidable propositions G uniquely correspond to simple thoughts or actions such as 'I pick up the pen'. Rather it is to be considered that there is a vat of an enormously large number of correlated G-like statements G_i (with $i = 1...n$ and n an extremely large number) relevant to a particular brain state. A subset of the G_i will incorporate the statement 'I pick up the pen' amongst other actions, others will incorporate the statement 'I pick up the pencil' amongst other actions, while still further subsets of the G_i will represent the cases 'I pick up both' or 'I do not pick up anything'. Each of these ghostly undecidable propositions G_i points towards a possible extension of my self-reflective state. Such extended systems draw us in and as we progress from one state to another, augmented, state our sense of temporality is created.

In the course of this dynamical stream of temporalisation I shall find myself coming into a state of picking up the pen or pencil, depending on the choice of the possible G_i . This set of potential G_i is itself of course very dynamic, as represented for example by the set G'_{i} in figure 14.2(c), and evolves in turn with the incorporation of new axioms, or choices, and new information into my system, corresponding to the ever evolving set of my possible future actions.

That the nature of subjective awareness may be correlated with the mathematical notion of the undecidable in self-referencing systems opens the door to a more thorough investigation. However, technically, in the context of the physical world, it may be that 'computability', rather than the closely related notion of 'decidability', is a more directly relevant concept to employ, since we know that the laws of physics in our world are such that 'computing machines' (both artificial and organic) are supported. That is, we are directly dealing with the states of such 'devices' in the physical world rather than with abstract mathematical symbols in a formal system, although there is a close structural parallel between the two cases. The discussion has been framed in terms of 'decidability' partly due to the similarity of the language used to express the experience of making a choice; that is, in making a choice we are primarily conscious of being in an undecided state. On the other hand given this coincidence of language terms some caution is needed in order to avoid being misled into taking the connection too literally.

It is indeed very much open to question how far to take the analogy between the structures pertaining to Gödel's theorem and the subjective process of decision making, although there is some degree of correspondence as indicated by the two sets of bullet points listed earlier in this section. With contradictory 'undecided' propositions from sets A and B being simultaneously entertained in equation 14.1, corresponding for example to G_i and G_j with $i \neq j$, this structure does seem to have some important differences also with the above mathematical correlate, since for Gödel's theorem the 'undecidable' describes the relation between G_i and $\sim G_i$, with the proposition G_i representing an unprovable but true statement. Although this implies that to some degree Gödel's construction should be taken metaphorically here, the employment of a mathematical framework with self-referential structures is still very relevant.

As well as the subjective interpretation the structure in figure 14.3 must also correlate with a physical manifestation. From this objective perspective the laws of physics must support a kind of inertia in the substructure of the physical brain, corresponding to the subconscious mind, that carries the subject into just one of the array of 'true' states either in set A or in set B of equation 14.1; that is into a new structure of self-reference such that the other options (in particular those in set B or set A respectively) become manifestly false propositions. The wiring of the subconscious mind in this sense will govern to a large degree the patterns of behaviour of an individual.

Naturally, we are taking this to be a phenomenon that our thoughts are thoroughly and continuously saturated with, rather than a discrete set of deliberations such as "hmmm, shall I pick up the pen or pencil?". That is, many of our 'choices' in this sense are simply the train of thoughts at the forefront of our mind that continually bubble up even when we are not *trying* to think. Most of these thoughts are not directly accompanied by an external bodily action such as picking up an object or not. For example each process of 'changing my mind', as described for the thought experiment shortly after the second set of bullet points above, is also a choice, even when not accompanied by a decisive external action.

An analogy between our thought processes and the mathematical structures underlying Gödel's theorem has been elucidated by other authors. In the preface to reference [85] (p.7) Hofstadter refers to elementary expositions involving a selfreferencing loop leading to undecidable propositions, such as G considered above, as containing:

. . . only the most bare-bones strange loop, and it resides in a system whose complexity is pathetic, relative to that of an organic brain. Moreover, a formal system is static; it doesn't change or grow over time. A formal system does not live in a society of other formal systems, mirroring them inside itself, and being mirrored in turn inside its "friends". . . . there is no counterpart to time, no counterpart to development, let alone to birth and death.

For Hofstadter, it is the self-referential and mirroring properties of the brain, giving rise to abstract structures similar to the 'strange loops' encountered in demonstrating Gödel's theorem, that is central to the emergence of an animated conscious 'I' from the inanimate particles of matter of the brain. As suggested by Hofstadter, for the present theory also, a mathematical structure somewhat more complicated than that required to demonstrate Gödel's theorem might be needed to account for these phenomena.

In this paper, we consider that the possibility for such systems to change and grow is not only something that objectively takes place in time; but moreover it is the ordered nature implicit in such a series of potentially related states that describes temporalisation itself. It is the possible existence of this ordered progression of systems which, through its simple structural isomorphism to an ordered one-dimensional mathematical series (that can be mapped onto the real line \mathbb{R}), itself corresponds to our immediate experience of time. The resolution of undecidable propositions from one system to another corresponds to the progression of choices we find ourselves making, with varying degrees of awareness. From the subjective point of view these choices are not deterministic, in the usual sense of the word, since they are not something that happens in time; rather they are the *generators of temporality* itself.

The progression depicted in figure 14.3 only has one direction. This underlies our experience of an apparent 'arrow of time' which corresponds simply to the one-way nature of this process (always with the possibility of losing knowledge of the world as our memory becomes frayed at the edges, it being supported by an imperfect physical device and following behind in the wake of our new experiences). The phrase 'arrow of time' is somewhat misleading since it implies the possibility of time having the opposite sense, that is flowing in the 'other' direction, effectively as if an empirical time parameter t could be seen to be reversed with $t \to -t$. However this is not the case as the progression in figure 14.3 possesses only a single direction, which may be associated with $+t \in \mathbb{R}$. The sequence of self-reflective states R, subjectively experienced as the flow of time, creates the inertia of the external world carrying physical objects. These objects include, for example, the components of a clock which can be used to measure the 'time' t. The fact that we can imagine, or even construct, a physical clock to 'run backwards', or for example watch scenes of a movie played backwards, creates the illusion of an alternative possible sense for time. However, we can only detect that a physical clock is running backwards since it operates relative to the fundamental underlying ordered progression of time.

The purely mathematical structure of figure 14.3, encapsulating the experience of a 1-dimensional temporal progression, can itself be encoded within the physical structures of a 4-dimensional spacetime world as depicted in figure 14.4. Here the structures in M_4 can be considered to represent a *static* 4-dimensional physical sculpture, as a manifestation of the static mathematical sculpture described for figure 14.3, within which a chain of states, having a one-to-one isomorphic correspondence with a self-reflective experience of a one-dimensional temporal flow, is embedded.

This origin of our experience of 1-dimensional time is analogous to the origin of our perception of 3-dimensional space. In general an abstract mathematical structure might be interpreted in several possible ways, whether geometrical or not. The arena for spatial perception arises out of a possible interpretation of the mathematical structure and symmetries of the multi-dimensional form $L(\mathbf{v}) = 1$, and in particular the properties of the components v^i , with $i = 1, 2, 3$, of the form $L(v_4) = (v^0)^2 - (v^1)^2 - (v^2)^2 - (v^3)^2$ with an SO(3) \subset SO⁺(1,3) symmetry, in terms of extended geometrical forms, as was described more generally in chapter 2. This incorporates the perception of physical objects in an extended 3-dimensional space, as

Figure 14.4: A representation of the progression of the self-reflecting state of figure 14.3 as translated into an extended physical environment M_4 .

represented for example on the hypersurface planes in figure 2.3. The structures of that figure, interpreted as a 4-dimensional spacetime, can be superposed on the manifold M_4 of figure 14.4, within which more complicated mathematical structures also arise out of the full mathematical form $L(\hat{v}) = 1$ when projected onto the 4-dimensional base manifold. These further mathematical objects, described for example in terms of fields on M_4 , incorporate series of self-referring elements which have a structural isomorphism with an experience of a directed 1-dimensional flow in time.

For the world sketched in figure 14.4 the mathematical structures in spacetime hence have both the necessary mathematical properties to give rise to perception of objects in space, that is in a 3-dimensional geometrical volume (represented by 2 dimensional planes in figure 2.3), as well as the experience of events in time, in a direction geometrically 'orthogonal' to the 3-dimensional spatial hypersurfaces on the manifold M_4 , which possesses the local $SO^+(1,3)$ symmetry of the form $L(\mathbf{v}_4) = h^2$ of equation 5.46. Hence both spatial and temporal forms of perception are encoded in the mathematical structures of the world. That is, we consider not only that the structures obtained from $\hat{G}/SO^+(1,3)$ symmetry breaking for $L(\hat{\mathbf{v}})=1$ projected over M_4 can be equivalent to the geometrical shapes we experience in space, but that they also incorporate self-referential mathematical structures which may be isomorphic to the self-reflecting progression that we experience subjectively as the flow of time.

In turn this one-dimensional temporalisation itself provides the source of dynamical laws through the breaking of the multi-dimensional form of temporal flow $L(\hat{v}) = 1$ over the 4-dimensional spacetime, generating the physical laws on M_4 with which physical structures in general, and those depicted in figure 14.4 in particular, must be compatible. Hence the progression $R \to R' \to R''$ identified within the mathematical environment E in figure 14.3 must be consistent with the seemingly *inevitable* progression $R \to R' \to R''$ of states in the physical spacetime environment M_4 in figure 14.4 described as an apparent consequence of the laws of physics.

The inertia of the physical world conforming to these laws of nature carries with it both the subconscious and conscious components of the brain and with them a 'decision' already shaped in the former is swept into a new self-reflecting state of the latter, for which an 'undecided state of mind' is now experienced as being resolved. More generally, information and data in the broader environment of M_4 in figure 14.4, as labelled by E in figure 14.3, which implicitly includes both the subconscious element and anything else distinguished from the self-reflecting R state considered, can in principle contribute to the progression.

An analogy can be made between the self-reflective system R in the extended environment M_4 and a thermodynamic system B embedded within the same larger environment on the manifold M_4 . While interactions between R and the further structures in M_4 lead inevitably to the progression of the self-reflective state $R \to R' \to R''$, for example in terms an increase in subjective 'information', the interaction between B and the broader environment leads inexorably to an increase in the total entropy S. In fact *both* the incorporation of a Gödel statement G_i into R, as depicted in figure 14.2, as well as the case of increasing entropy might be considered as analogies for the phenomenon of temporalisation. The first example may also carry some elements above a mere metaphor, while for the second example an increase in entropy will accompany the physical process underlying the subjective experience of time.

For a sufficiently complex system such as a human brain the complete immersion of the self-reflective state within the wider environment might effectively generate a continuous temporalisation. Indeed, while for figures 14.2–14.4 a series of discrete steps has been described, subjectively we generally experience a continuous flow of time without any gaps or jumps. For example, while watching a ball roll along a table, essentially obeying Newton's first law of motion, we observe a smooth progression relative to our internal sense of temporality. It is this continuous subjective experience of the flow of time, as modelled by the one-dimensional real line R, that forms the basic entity of the present theory.

Indeed, although subjective experience in general exhibits a correlation with objective phenomena it is not explicitly described by the latter phenomena. For example, the sensation of 'green' is associated with radiation from an interval of the electromagnetic spectrum with a wavelength of around 500 nm in physical interaction with the cells of the human eye and the resulting neural activity in the brain. However the subjective experience of 'green' is not explicitly contained in the description at any level of detail of these objective physical processes. Similarly here, the subjective experience of a continuous flow in time is associated with the physical structures implied in figure 14.4. However it is not necessarily the case that a continuous sequence needs to be identified in a physical system based on the progression $R \to R' \to R''$ in order for it to underlie a subjective experience of time which can be accurately modelled by the continuous real line R.

The irreversibility of conscious choices, the origin of the 'arrow of time', is echoed in the irreversibility of many physical systems which are all governed by equations derived from the general mathematical form of progression in time. For example the second law of thermodynamics itself arising as a statistical consequence of a progression of states, as alluded to in section 13.2. An essential difference is that while entropy increase is solely something which happens in time, the physical progression

 $R \to R' \to R''$ of figure 14.4 is directly correlated with a subjective experience of time which *drives* the temporal flow itself. In addition to systems of classical physics, quantum phenomena are also subject to the underlying ordered flow of time which is infused into the base manifold M_4 . Calculations of probabilities and cross-sections for quantum processes depend on the accumulation of the possible field degeneracies conforming to a causal sequence in time, building upon time-ordered expressions as described for equations 11.41 and 11.44 and more generally in sections 11.2 and 11.3.

In addition to the fundamental temporal progression itself there are a large number of apparently one-dimensional quantities which may be constructed out of the physical structures on M_4 . As well as the example of entropy S these include the temperature T of a body or even the 'time' t recorded by a mechanical clock. However each of these quantities correlates solely with the objective collective actions of molecular motions in spacetime and each defines a measurable property of the fourdimensional world. Even the time t recorded by the clock is not a 1-dimensional geometric entity, but rather signifies a certain coincidence between the hands of the clock and the numerals on its dial in 3-dimensional space.

In fact no purely 1-dimensional phenomenon can be objectively inscribed within a 4-dimensional spacetime without reference to the extended M_4 manifold or physical processes within it. While the self-reflective physical structure in figure 14.4 is similarly diffused in spacetime, objectively in terms of the firing of brain neurons for example, the subjective experience of time is of a different character. Unlike a physical quantity such as S , the mental process of experiencing time is a *purely* 1-dimensional phenomenon and in this subjective sense it is not located within spacetime. The progression $R \to R' \to R''$ is subjectively fused in mind into a purely 1-dimensional experience of a qualitatively different nature to, and hence distinguished from, the 4-dimensional spacetime arena.

This one-dimensional structure is the origin of time in the world, in the form of subjective temporalisation, and provides the foundation which underlies the general mathematical form of temporal flow $L(v) = 1$ and the physical laws in spacetime itself. It is through attempting to address the second loose end on the right-hand side of figure 14.1 that a source has been identified for the first loose end on the left-hand side.

14.2 A Universal Foundation

For a description of the universe in terms of a purely objective theory a 4-dimensional background arena for events in spacetime, as for the case of general relativity, might be postulated as a fundamental entity or perhaps derived from a higher-dimensional spacetime. This is consistent with the observation that all physical events in the world have *both* a spatial *and* a temporal location in the universe. For the present theory it is noted, however, that while we observe such events in spacetime our subjective experience in the world is more fundamentally temporal than spatial. While many experiences are accompanied by a sense of both time and space, all appear to exhibit a temporal aspect while some, such as the experience of listening to a piece of music or of simply thinking itself, lack any accompanying sense of an extended spatial arena. This observation, along with the simplification of founding a theory on one dimension rather than four, provided a source of motivation for the present theory.

The sensation of time that accompanies all subjective experiences may be modelled mathematically by a continuous interval of the real line \mathbb{R} , which is precisely the same one-dimensional structure of temporal flow considered objectively as a presence which underlies the structure of the entire universe. That is, innate within the expression for this temporal flow in the multi-dimensional form $L(v) = 1$ of equation 2.9, along with its symmetries, the form of the physical universe throughout an expanse of both time and space is supported. As discussed towards the end of the previous section, both temporal causality and a spatial geometry, deriving from the form $L(v) = 1$, are infused throughout the manifold M_4 .

At the mathematical level the unfolding of this structure is analogous to some degree to the properties of the Mandelbrot set, in that a highly complex pattern is identified through a very simple mathematical expression. A further analogy we consider here is the simple differential equation $\frac{\partial^2 y}{\partial x^2} + y = 0$ with the possible solution $y = \sin x$. This sine wave is typically represented as a graph in the 2D plane incorporating for example a horizontal axis for values of $-\pi \leq x \leq +\pi$. However innate in the expression $\frac{\partial^2 y}{\partial x^2} + y = 0$ the actual mathematical solution is of course present throughout the infinite real line for $-\infty < x < +\infty$, even though we typically only picture a small portion of this solution. Similarly while the extended 'spatial' arena corresponding to the translation symmetry of $L(v) = 1$ is pictured for a finite region in figure 2.2 this purely mathematical structure is of infinite extent in all n dimensions. This observation still applies when the construction of the spacetime arena is generalised for the geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, as one of many possible solutions involving differential equations in, and a degeneracy of, the underlying fields. As for the sine wave above, this purely mathematical solution has no limit for the coordinate parameters on the base manifold (including in fact the particular solution for $G_{\mu\nu}(x)$ of equation 11.12 which is itself described by a simple sine wave function as represented over an interval of x^3 in figure 11.1), while as for the Mandelbrot set the structure which emerges in general may be highly complex.

These mathematical patterns and structures on M_4 arise through the projection of the full form $L(\hat{v}) = 1$ onto the base manifold and the associated symmetry breaking. Through an innate subjective interpretation of certain entities on M_4 there arises for us a vivid impression of material phenomena which appear to be detached and hovering outside us in an apparently spatial expanse. This entire *perceived* world is however mathematically enfolded within the one-dimensional subjective temporal progression through which everything in the world is observed. (This description is very much influenced by the notion of the *a priori* necessity for both temporal and spatial forms of experience, and their mutual relation, as elucidated by Immanuel Kant). The observation that spatial structures through the form $L(\mathbf{v}) = 1$ can be implicitly enfolded within the experienced one-dimensional flow of time that accompanies all of our perceptions in the world completes the initial motivation for the present theory described in the opening paragraph of this section.

The form $L(v) = 1$ itself is derived from within the notion of a 'moment of time', divided into infinitesimal intervals, as described in section 2.1. From the mathematical point of view the solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ over the manifold M_4 is a structure implicit within the full form $L(\hat{v}) = 1$ and its symmetries, describing the geometry of an infinite expanse of 4-dimensional spacetime (as for example implied for figure 13.6) which does *not* 'take time' to unfold across the cosmos, rather it underlies all cosmic structure, similarly as the solution $y = \sin x$ is implicit within the expression $\frac{\partial^2 y}{\partial x^2} + y = 0$ across the full extent of the x-axis. This mathematical structure logically precedes the laws of physics and the properties of physical objects perceived in the world. As a manifestation of the underlying mathematical structures these physical properties include causal relations in general, incorporating for example the dynamic evolution of the fields, on M_4 . The causal and spatial relations between physical events unfolding in the world, which do 'take time', create the sense of a world *outside* accommodating all of the apparent material phenomena. All physical structures are subject to the laws of physics, which derive from the underlying mathematical forms, which apply for example to the phenomena depicted in figures 14.3 and 14.4 which in turn have both an objective and a subjective interpretation.

The logical precedence of the elements of the theory described above is unpacked in the following sequence:

- (1) The objective starting point of the theory is one-dimensional progression in time with a mathematical structure isomorphic to an interval of the real line R.
- (2) From the basic arithmetic properties of $\mathbb R$ a general multi-dimensional flow in time subject to the constraint $L(\mathbf{v}) = 1$ can be derived.
- (3) The identification of extended geometrical structures from the form and symmetries of $L(v) = 1$ provides a basis for the necessary arena for perception, that is a subjective experience of a spatial expanse.
- (4) In breaking the symmetry of the full form $L(\hat{\mathbf{v}}) = 1$ over the M_4 base manifold the properties of material phenomena are sculptured and made visible in conformity with the resulting laws of physics.
- (5) The material objects in the world include the complex structures of physical devices, such as brains, capable of performing mathematical operations and encoding a progression of states of self-reference governed by the physical laws.
- (6) The sequence of self-referential states, linked through a contiguous resolution of their associated 'undecidable propositions', correlate with subjective thoughts and experiences, ever accompanied by the sense of an ordered flow in time.
- (7) The subjective temporalisation may be modelled by an interval of the real line R having a one-dimensional mathematical structure identical to that in item (1).

The first four points listed above form the main thrust of this paper from the opening chapters through to and including chapter 13, while the remainder of the above chain has been the topic of the present chapter. In this paper the self-reflective structures depicted in figure 14.4 and discussed in the previous section are proposed as the means through which subjective experiences arise, although this may be a vast simplification, or even largely a metaphor, for the actual mechanism. In any case, the existence of a sequence such as that described in the latter four points above, beginning with an empirically observed physical world and leading to self-reflective conscious experience in the world, is incontrovertible to the extent that it is evident that the presence of conscious beings is amongst the known phenomena of the physical world. This observation applies for any physical theory, as discussed shortly after figure 14.1, although the details of the theoretical mechanism that underlies the subjective thought process remains open to investigation.

For any physical theory built upon essentially any postulated entities, such as fields or particles and a background of spacetime, the universe can be described in mathematical terms as a 'static' 4-dimensional object, for example in the form of a spacetime diagram for the entire cosmos, which includes within it the full history of each human brain and all other material entities. However this is clearly not the way we see the universe, rather the 4-dimensional spacetime structure of the brain must prescribe our subjective perception of the universe as dynamically evolving through a progression in time. Hence for any such physical theory the above segment of argument in points (5) and (6) can still be applied, however there then remains dangling the prominent loose end that there is no apparent justification for the origin and properties of the initially postulated physical entities themselves, other than that they may be contrived pragmatically, for example in terms of a Lagrangian function in spacetime, to match the empirical data from observations and experiments.

On the other hand the key observation for the present theory is that the final link, item (7) in the above chain, representing the fact that temporalisation is contained as an ever present feature of subjective experience in the world, reconnects the chain to the initial link of item (1) at the top. Hence not only is a mechanism for the origin of time conceivable, providing a foundation for the left-hand loose end of figure 14.1, but this temporalisation itself arises through self-reflecting structures, identified in the physical world itself, which account for our subjective experiences in general and the right-hand loose end of figure 14.1. The chain then naturally closes into the cycle depicted in figure 14.5.

It is a feature of the present theory that the two loose ends of figure 14.1 can be mutually tied up in this way. From a mathematical point of view each of the six stages in figure 14.5 is contained within the previous stage, supplying a foundation for all of the structures of the theory. This system can then be considered to establish a 'universal foundation' for the present theory. The entire system is self-supporting in the sense that whenever we ask "where does X come from?", where X can be time, space, matter, conscious beings, or anything at all, the question can be answered in terms of something else *within* the system. Without the need for any external foundation or assumptions and entire structure in figure 14.5 hence detaches itself and floats free.

This figure does not, of course, express an impossible cyclic chain of cause and effect relating the six stages in a temporal sense. Indeed time itself is contained as one link within this cycle hence incorporating also the notion of temporal causality within this structure, and in particular for the physical laws in node (4). Rather each connection between neighbouring stages has the logical nature of a structural isomorphism, more precisely in the sense that the properties of node $(i + 1, \text{ mod } 6)$ are contained within the structure of node (i), with the net effect of expressing a self-contained and consistent mathematical and physical system. While providing a chain of concepts for the benefit of deliberation the six nodes of figure 14.5 can be

Figure 14.5: A self-contained 'time cycle' leading from the notion of progression in time, through the general mathematical form $L(v) = 1$ and perception of physical structures in spacetime, to self-reflective entities incorporating experience of a progression in time and hence completing the cycle.

considered to collapse down to a single entity. This entity contains an entire universe created through the temporalisation represented by the structure in figure 14.3 which itself is entirely enveloped within the same physical world.

In the opening of this section it was noted that while empirically everything happens in *spacetime* from the subjective point of view *time* is a more fundamental mode of experience than space. This observation provides part of the original motivation for basing the present investigations on a general form of temporal flow $L(v) = 1$, together with its symmetries, rather than upon a 4-dimensional, or even higher-dimensional, spacetime structure. The further observation here that it is in the nature of time itself to provide the link connecting the two loose ends in the theoretical sciences described in the previous section and in figure 14.1 adds further circumstantial support for this approach.

While the means of supporting spatial perception arises from a very direct interpretation of the geometric forms implicit in the mathematical properties of $L(v) = 1$, as described in chapter 2 and corresponding to nodes $(2) \rightarrow (3)$ of figure 14.5, the means of generating temporal experience arises from the far more complex mathematical structures represented in figure 14.3, as described in the previous section and corresponding to nodes $(5) \rightarrow (6)$ of figure 14.5. The figure as a whole can be seen as an interplay between 1-dimensional and multi-dimensional forms of time. The underlying mechanism for obtaining an extended 4-dimensional world out of 1-dimensional temporal flow, summarised in the upper half of figure 14.5, differs from the far more complex structures required to identify a purely 1-dimensional entity out of the 4-dimensional physical world, as summarised in the lower half of the figure. (As discussed at the end of the previous section a simple physical clock, for example, does not possess any intrinsically 1-dimensional geometric structure). It is the very different nature of the mechanism for obtaining multi-dimensional forms and extended spacetime from 1-dimensional time on the one hand and for identifying temporal flow itself out of the higher-dimensional and spacetime structures on the other hand, as required for subjective perception and experience, that opens up the non-trivial system of figure 14.5.

The contrast between the *objective* features of temporal progression, identified as the simplest element of figure 14.5, and the subjective experience of progression in time arising out of the *most complex* structures in this system, while both aspects of time, in nodes (1) and node (6) respectively, share the identical structure of an interval of the real line, underlies the enigmatic quality of the concept of time itself. References to the seemingly more philosophical nature of time in the physics literature are rare but not entirely absent. Near the beginning of the introduction to his $Space-Time-Matter$ Hermann Weyl writes ([82] p.1):

Since the human mind first wakened from slumber, and was allowed to give itself free rein, it has never ceased to feel the profoundly mysterious nature of time-consciousness, of the progression of the world in time, – of Becoming. It is one of those ultimate metaphysical problems which philosophy has striven to elucidate and unravel at every stage of its history.

While the upper half of figure 14.5, that is the chain of nodes $(1)-(4)$, represents the development of the theory within the traditional scope of physics, the entire scheme, including the lower half of the figure, is fully incorporated within the sphere of scientific study more generally. Indeed experiments are performed, dating for example from those conducted by the neurologist Benjamin Libet in the early 1980s, concerned with the relation between the physical brain and conscious actions, that is essentially nodes (5) and (6) respectively in figure 14.5. In such experiments physical cerebral activity is found to precede a conscious awareness of intention typically by around 300 milliseconds or more.

For the present theory a conscious intention, or choice, is associated with the origin of temporalisation, as described for figures 14.2 and 14.3. This leads to the multi-dimensional form of temporal flow $L(\hat{v}) = 1$ through which derives the mathematical structure underlying the entire physical universe on M_4 , incorporating its full eternal temporal extent both into the past and into the future. This physical universe includes in particular the brain state 300 milliseconds before the conscious choice was experienced, and indeed at any other time, as embedded within M_4 as represented in figure 14.4. Hence the overall scheme presented here is fully consistent with the experimental findings of Libet and others. More generally the full cycle of figure 14.5, including all of the nodes and links, is fully amenable to theoretical and scientific investigation.

As a preliminary discussion the remarks made here on the origin of our temporal experience and the phenomenon of consciousness, together with their mutual association, are necessarily somewhat speculative. However, it is meaningful to formulate such questions, the worldview presented in this paper provides a new arena through which the construction of figure 14.5 seems inevitable, and this provides a firm mathematical basis for a possible scientific enquiry into the nature of subjective phenomena compatible with the basic structure of the present theory.

Most fields of scientific study are rooted in node (4) of figure 14.5, in that the natural starting point for any scientific investigation is observation of the physical world around us. For the physical sciences the general aim is to deduce the basis of the underlying structure of the world, extrapolating inwards as for example in the direction of nodes $(3) \rightarrow (2)$ for the present theory, while on the other hand the biological sciences, for example, also study the world at face value and extrapolate outwards, which might include the properties of nodes $(5) \rightarrow (6)$ in figure 14.5. However, unlike the case for other scientific theories in general, in the present theory it is natural to extrapolate one step further, both inwards and outwards, to establish the final link in node (1) and hence complete the circuit. Here the overall structure of figure 14.5 then has the shape of providing an answer to the general question "why is there something rather than nothing?", rather than merely displacing it.

From the mathematical perspective while an exposition of the structures in figure 14.5 could begin with any given node the simple mathematical structure of time, as a 1-dimensional progression modelled by the real line R, provides a convenient entry point into the study of the whole system. In particular the unique properties of a real interval provide an unambiguous structure upon which to develop the theory, as will be emphasised later in this section. If the properties of the real line R are considered to define the axioms, which in general underlie all the expressions which may be derived in a formal system, then the self-contained structure of figure 14.5 might be thought of as a mathematical system which 'contains its own axioms'.

From this point of view as a single entity of self-creation the time cycle in figure 14.5 can be considered firstly as a purely mathematical structure which can be described in terms of the six nodes displayed with each one mathematically identical to, or contained within, the previous node of the chain. This picture can then be 'coloured in' with both the objective material features of a physical world and the subjective experienced aspects of self-reflective thoughts and perceptions.

The subjective experiences, as much as the objective material phenomena observed, are an irreducible component of this system. Indeed it is the experience of time, as well as of space, that generates necessary links in the time cycle of figure 14.5. Such a world cannot exist unless it is experienced. The two loose ends, left exposed in many conceptual worldviews, relating to the origin of conscious experience and the origin of the material world are interwoven into one coherent system. Here the emphasis does not weigh heavily upon a pre-existing material content of any kind, but rather takes an overall more balanced view within which 'matter' is identified with a form of experience shaped in 'mind'.

The apparent distinction between mind and body arises in part since the spatially distributed matter we experience appears to exist out there, however here the concepts of 'mind' and 'matter' are intimately intertwined within one system. We have no need to postulate two wholly different kinds of substance and ponder how they interact, such as through the pineal gland in the brain in the worldview of René Descartes. Rather mind and matter are different aspects of the same self-contained

system: the conscious mind being bound with the structures of mental activity and temporality embedded in the physical world, while spatially extended matter itself is carved out of the multi-dimensional algebraic properties of time. Hence both sides of the philosophical dichotomy between mind and matter are accounted for and the points of view of both the idealist and the materialist democratically amalgamated into this structure. We are not spirits haunting Earthly bodies, and neither are we machines in search of a soul.

While forming components of one overarching framework both the objective structure of the physical world and the subjective forms of experience in the world can be described in terms of theoretical elements, and each is sufficiently distinctive and well defined to seemingly take on a 'life of its own'. From the point of view of the present theory the materialist is grounded in node (4) of figure 14.5 and can construct a relatively short, physically motivated, argument to account, via node (5), for the realm of the idealist in node (6). On the other hand the idealist, based upon the subjective experiences of node (6), is required to make a more lengthy detour, via the conceptual and mathematical structures of nodes (1), (2) and (3), in order to arrive at the materialist's realm in node (4). This asymmetry in the apparent directness of mutual explanatory power perhaps in part accounts for the predominance of the materialist, 'a spade is a spade', philosophy that has underpinned most progress in the history of science, in addition to its practical utility.

Outside the present chapter of this paper, as for the vast majority of work in theoretical physics in general, the focus has been with the study of a mathematical description or model of the physical material world, here through equations such as $L(\hat{\mathbf{v}}) = 1$ and $G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$. However mental phenomena, such as our awareness of the physical world and decision making actions, are also very much a feature of the universe and in principle equally amenable to theoretical analysis, as discussed above.

To recap, in the present theory the mathematical structure described in figure 14.3 models our conscious self-reflective state and ever present feeling of not knowing for sure quite what we shall do in the next moment. This perpetual uncertainty as to our own thoughts or actions resolves momentarily in a choice G_i opening up a new horizon of uncertainty, as represented in the progression of figure 14.2. The self-reflective state is inexorably drawn through the series $\dots R \to R' \to R'' \dots$ of figure 14.3 correlating with an internal experience of a sequence of thoughts, aspects of which have a complex mathematical representation, but in all cases associated with a subjective experience of a simple one-dimensional temporal flow.

Within this structure the term *freewill*, as used without hesitation in everyday language, is identified as this 'experience of choice' as one feature of the overall system of figure 14.5. Everything that happens objectively in the physical world follows in the wake of this subjective temporalisation phenomenon. The historical philosophical debate concerning 'freewill versus determinism' becomes more strictly a question of 'freewill versus the laws of physics' in the context of modern day science. The laws of physics include 'indeterministic' quantum phenomena as a feature of the objective world which in the present theory are *not* correlated with the subjective act of making a conscious choice. Indeed the intrinsically random transitions of quantum effects are of a wholly different nature to rational decision making or freewill. On the other hand quantum properties are a major component of the laws of physics, and it is this full

package of physical laws which determine all physical structures. These include the physical state of the brain which evolves in time according to the laws of physics, exhibiting properties which do correlate with the interaction between the conscious and subconscious mind as implied in figure 14.4 and hence providing the vehicle to carry self-reflective experiences.

The traditional philosophical difficulty in reconciling freewill and the laws of physics derives from the observation that an apparently independent objective world evolving according to a set of deterministic laws (together with random quantum phenomena) seems to leave no room for the notion of freewill. However, here in the present theory, since the physical world is brought into being through a subjective temporalisation sufficient breathing space opens up for the concept of freewill – not as a secondary phenomenon on top of a given physical world, but as an irreducible feature in dynamic interplay with it, as summarised in the time cycle of figure 14.5. An element of the philosophical confusion concerning these issues arises as there is considerable ambiguity in the meaning of the term 'freewill' in itself. The present theory provides a context within which the notion of freewill might be more precisely defined. Within the system of figure 14.5 the means by which the world is experienced in mind is as important as the empirical forms of matter, with freewill being a property of the former while the laws of physics are a property of the latter.

It seems of course counter-intuitive to suggest that the great expanse and 'weight' of the entire physical universe might be created through and carried in a single waking moment of thought. However, as described near the opening of this section, the mathematical structures underlying a solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ innate in the form of temporal flow $L(\hat{\mathbf{v}}) = 1$ are perfectly 'weightless' and infinitely delicate, effortlessly supporting an entire cosmic history throughout the full expanse of the physical manifestation of the universe. If the laws of physics in this spacetime are compatible with the local evolution of a physical brain as depicted in figure 14.4 which encodes the self-reflective sequence of figure 14.3 which in turn represents a subjective experience of a one-dimensional temporal progression isomorphic to the ordered real line in node (1) of figure 14.5 then the circuit closes and the experiencing being locates himself at a particular place in a particular world (in this chapter in this context pronouns such as 'himself' or 'his' refer to a non-gender-specific being in any world). This spacetime location will be within the habitable epoch of the cosmological evolution as depicted in figure $13.6(e)$, and most likely upon a planet orbiting within the habitable zone of a suitable solar system as considered in section 13.3, for the case of our own universe. The poets have more readily conceived of such a world, as for example in the often quoted opening lines from William Blake's Auguries of Innocence of 1803:

> To see a world in a grain of sand And a heaven in a wild flower, Hold infinity in the palm of your hand And eternity in an hour.

Here, not limited by poetic licence, we require only a moment rather than an hour through which the entire universe may be perceived. The contention of the present theory sees the world and the heavens, together with an infinite expanse of space and an eternal temporal duration all held within a moment of time. The completion of this picture is depicted in figure 14.5 with the experience of time itself contained within the structures of the physical universe.

While the entire physical universe is created through the experience of a single temporal moment, the moment itself is not unique. The circuit of figure 14.5 can be closed by any one of a large number of possible local structures representing the progression of figure 14.3, each embedded within the physical world and each associated with a moment of experienced time. Indeed if the physical world is capable of supporting such a structure at all then in principle there may be many examples. This generalisation is depicted in figure 14.6.

Figure 14.6: Rather than the single experience of time represented by node (6) in figure 14.5 in general a large number of physical structures on M_4 may generate moments of temporalisation, each represented by one of the small boxes labelled by A or B here and each of which completes the circuit of a time cycle for the same physical world of node (4).

In particular the set of temporal moments labelled by the series $A_1 \ldots A_i$ in figure 14.6 might form a contiguous structure in the sense of the embedding of the progression of figure 14.4 within the physical world on M_4 . The corresponding temporal experience correlated with this structure is the sensation of a 'sliding now' for a particular individual. Similarly while the 'spark' that creates the universe from within can be any moment A_i associated with such an individual, it could also belong to any other being, such as the temporal experience represented by B in figure 14.6. In general the experiences of a community of beings A, B, C, \ldots may be inscribed within the same manifestation of a physical world, as depicted in figure 14.7.

Figure 14.7: A depiction of the trajectories of several self-reflecting 'beings' A, B and C experiencing life in the M_4 spacetime manifold.

Any structure of temporalisation, such as those represented by each 'X' in figure 14.7 can take the place of node (6) in figure 14.5 and complete the circuit which also incorporates the physical world itself which a community of beings such as A, B and C cohabit. Each of these individuals observes a time-ordered progression of states of material entities, from stars and planets to tables and chairs and other individuals, distributed in a 4-dimensional spacetime as originally envisaged in figure 2.3. While a given observer A experiences the subjective freewill of his own self-reflective state, and an internal temporalisation as represented in figure 14.3, from his perspective both the subconscious as well as the conscious elements of the brains of the other beings B and C are unambiguously seen to partake seamlessly in the physical flow of events in the world. That is, the behaviour of the other, progressing in parallel and as represented for example in figure 14.4, conforms to the basic laws of physics exactly as any other physical entity such as the tables and chairs carried along in the inertia of the world. With a perfectly reciprocal account given from the internal subjective point of view of B or C the mutually consistent perspectives of all individuals dovetail together within the common physical world.

In conformity with this symmetry between A, B and C in terms of a perspective on freewill and the laws of physics each observer carries a personal experienced fundamental time parameter s. This temporal flow s is equivalent to the proper time τ recorded by a physical clock in the frame of the individual, as related by the constant factor γ described for equation 13.3. The progression in time $s \equiv \tau$ for any given individual is related to that associated with each of the other observers through the dilation effects of both special and general relativity in spacetime, as described for the 'twins' A and B towards the end of section 5.3 and generalised near the opening of section 13.1, again in a completely reciprocal manner.

For any individual the seemingly vast potential arena for the flow of time in the universe at large contrasts sharply with the observation that we experience time at an apparently brief moment. The concept of 'now' can be identified subjectively with the 'present moment', which consists of a small duration rather than a point in time. This leads us to pose the question – given such a vast expanse of time – "why is it now!?"; as opposed to, say, some time last week. This question is particularly challenging for theories of the world which posit an initial extensive and objective spacetime arena upon which the laws of physics are mathematically constructed from an independent perspective outside spacetime, such that the physical laws governing all phenomena have perfect symmetry with respect to translation of location in either space or time. Within such a framework 'now' is generally conceived objectively as a point in time, as a mathematical point of the real line. While we have a wide choice over where to make an observation the fact that we necessarily observe the world as it is now, at this particular point in time, appears to explicitly break the time translation symmetry.

The problem disappears when we consider the meaning of 'now' within the theory presented in this paper. Our self-referencing awareness involves the physical structure of a small region of the world which is sufficiently complex to support 'undecided states', but further complex structure carried in the physical world, in particular that of the subconscious brain, holds the resolution to such states and draws conscious awareness into the wider world in the process of temporalisation. Beyond the brain we find also the human body, the habitable environment and the entire physical world unfolding through the physical realisation of temporality creating a *situation* in which the conscious being exists. Since every situation is an experience and every experience is an experience now the logical meaning of the word 'now' in this system is entirely redundant (although, of course, it has a practical purpose in everyday language). The fact that it is *now*, rather than some time last week, is simply that I am this experience, whereas the situation for a particular individual at a particular time last week is that experience. The apparent problem is then largely an issue of the assumptions made in the use of language regarding the identity of an individual (the 'I am') as something more attached to a bodily form than to an experience.

The fact that it feels like 'now' comes from the fact that the world exists 'all at once' – that we can conceive of a past and future progression within which we place ourselves in the present, now. However, past and future are not periods of a pre-existing external and independent world-time; rather the past and future refer to locations within the universe with respect to the perceiving being whom experiences the situation – it is a description of the experience which partitions a self-reflecting conscious state into a concrete past and an uncertain future as a necessary structural form of a thinking being. (This aspect of the worldview being described here is philosophically close to the standpoint of existentialism, and is influenced in part by the philosophy of Jean-Paul Sartre).

I have to experience the world *now* in a similar way that I also find myself here at a particular spatial location in the world. While the 'body' of the whole world is created through the structure of our being, here is where my eyes, and other

sense organs of the human body, locate me spatially relative to other physical objects in the world. To necessarily exist here and now is simply the statement of having to be the centre of reference for an experience in a world. This central vantagepoint is essentially the location of the physical manifestation of the associated thought processes, as represented in figure 14.4, within the extended spacetime manifold M_4 .

From the perspective of any individual such as A the universe created through any given experienced moment, such as A_3 in figure 14.6, not only mutually supports the contiguous moments of the 'sliding now' and into the span of the current day, but also the moments of yesterday, and the past in general, and those of tomorrow, and the future in general. As well as the spacetime separation between moments experienced by A and B each individual is also separated from himself in time, corresponding to the moments marked ' X ' on the trajectory of A in figure 14.7 for example. The identical universes generated from A's experienced moments on Monday, Tuesday, Wednesday and so on resonate together into a single life history. Any moment of temporalisation not only brings the corresponding present self into being but also the physical structure for all the past and future 'selves' in the life history of the same individual. This system is hence comfortably compatible with the experimental findings of Libet and others as noted in the discussion following figure 14.5. The mutual relations between any of $\ldots A_1, A_2 \ldots A_i \ldots$ from a single life history dovetail together, as with the moments of any other beings, such as B and C , in the same physical world.

The exhaustive spacetime coverage of the universe created through each temporalised moment A_i together with the inertia of the derived material processes in the world maintains the physical manifestation of any individual during non-waking hours or through different shades of consciousness. In this way historically separated waking moments are seamlessly stitched together over periods of years alongside those of other beings immersed in the same world.

The existence of different shades of consciousness, such as the experience of dreams, suggests that a rigid geometric framework in space may not be essential for some forms of perception, although dream sequences are closely associated with waking experiences. The question regards whether spatial perception is required in some form in order to complete the circuit of the time cycle in figure 14.5. As discussed in section 12.1 and section 13.1 (before the bullet points) our a piori imposition of an extended 3-dimensional frame for our perceptions in the world does not perfectly match the non-Euclidean geometry of the world – which we however effectively interpret as being flat while certain phenomena are ascribed to an apparent force of gravity. We very rarely perceive solely events within a local inertial reference frame, such as within an orbiting spacecraft, however such an idealised limiting geometry is not required in order for us to be able to interpret and organise our perceptions of the world in a manner compatible with the presumption of a flat Euclidean frame of reference.

In addition to providing a spatial orientation for vivid conscious experiences of the world, with material objects obeying physical laws of motion within the perceptual framework, the general laws of physics themselves, which shape all material properties, arise from the projection of the full form of temporal flow $L(\hat{\boldsymbol{v}}) = 1$ onto the base manifold M_4 . Complex mathematical and physical structures which arise in this breaking of $L(\hat{\boldsymbol{v}}) = 1$ and its full symmetry over M_4 accommodate the mechanism for self-reflective conscious experience itself, as described for figures 14.3 and 14.4. The physical laws deriving from the symmetry breaking hence not only maintain objective material objects in the world but also images of the same objects which can be maintained in our subjective thoughts even while the object is not being directly perceived (as for example in a dream or in a waking moment in which we simply look away from the physical object while still thinking about it).

Hence the laws of physics derived from the symmetry breaking of $L(\hat{\boldsymbol{v}}) = 1$ over the base manifold M_4 give rise to both the structure of conscious self-reflective states and the material phenomena, perceived against the M_4 background, which constitute objects of consciousness. This then describes the primary requirement of the symmetry breaking of $L(\hat{v}) = 1$ in order to complete the time cycle in figure 14.5, that is to open up structures that may be presented as objects of conscious experience together with the self-reflective elements capable of contemplating such objects. In our world these structures are obtained through the projection of $L(\hat{v}) = 1$ over a locally approximately flat 4-dimensional spacetime M_4 which incorporates a 3-dimensional spatial arena for the perception of objects.

In principle we can enquire what it might be like to be immersed in a highly curved spacetime environment of a different world. Without the support of an effectively Euclidean spatial orientation it would be harder to organise our incoming sensory data and difficult to predict the physical consequences of our own actions and to engage in such a world generally. The likelihood of errors of judgement in this respect is much lower in the local environment of an apparently flat spacetime combined with the very regular patterns of motion deriving from Newtonian gravity, as we encounter in our own world.

It seems very natural to us that space ought to have Euclidean properties, as witnessed by the historical perseverance of the geometrical laws of Euclid formulated in ancient Greece, which until the early $20th$ century were assumed to describe the real world. While applying to an excellent approximation in the local environment of the Earth and solar system, the assumption of a flat spacetime geometry breaks down for large scale cosmological structures. In general any manifold with two or more dimensions can exhibit arbitrarily large curvature at any point, as is the case for our 4-dimensional universe for which the curvature diverges in the proximity of the initial singularity or a black hole. However the curvature of any 1-dimensional manifold is trivially zero and the geometry necessarily 'Euclidean'. Hence an interval of the 1-dimensional real line \mathbb{R} , as a unique and robust structure, and as a parametrisation of the subjective experience of temporal flow, provides an unambiguous starting point from which the present theory has been developed in this paper.

This discussion raises the questions considered in section 13.3 regarding whether or not the symmetry of $L(\hat{v}) = 1$ is uniquely required to be broken over a 4-dimensional spacetime M_4 and whether structures identified in the symmetry breaking are required to be compatible with the notion of perception as conceived in our world. Whether a complete time cycle of the kind in figure 14.5 incorporating self-reflecting beings without an a priori spatial perception of any form could exist, or even whether there are conscious organisms within our own world that completely lack any spatial awareness, may be difficult questions to address. Such self-reflecting creatures may still necessarily require an M_4 base space to break the full $L(\hat{v}) = 1$ form in order to physically exist (as do all non-sentient biological life forms in our world), yet without employing a subjective spatial interpretation of the 3-dimensional structures on M_4 . In a similar way we require the extra dimensions of the form $L(\hat{\mathbf{v}}) = 1$ in order to physically exist ourselves, yet without our being directly aware of them.

Here we recall that the term 'perception' is being employed not just in the narrow sense of that which we are visually aware of in the moment. It refers more generally to an organising faculty for all the data about the world that enters and our thoughts through all of our senses. This data is accumulated both directly, for example through the experience of vision or touch, as well as indirectly, for example via intermediate objects, tools of experimentation or the accounts of other people. This data concerns aspects of the world in spacetime ranging from our immediate locality, down to the minute microscopic scales explored in HEP experiments, out to regions very remote from us and through to the limit of observations relating to the structure and evolution of the cosmos. Perception is a form of knowledge that encompasses everything we can understand about the world in space and time, in principle anything associated with nodes (3), (4) and (5) in figure 14.5.

Moulded by this form of perception physical structures as we experience them exhibit the effortless complexity inherent in the breaking of $L(\hat{v}) = 1$ over the infinite expanse of M_4 as depicted for example in figure 13.6. As alluded to in the opening of this section such a structure is analogous to the endless delicacy of the Mandelbrot set, which arises from the iteration of a simple mathematical expression in the complex plane. In both cases an inexhaustible variety of fine detail can be observed wherever we choose to 'zoom in' and examine for example biological forms in the physical world or the spiralling patterns of the Mandelbrot set. At the shortest physical distances probed the properties of elementary particles emerge over an underlying fractal-like structure of field solutions, as described in section 11.3, while at the other end of the scale, throughout the expanse of the observable universe, we perceive the manifestation of the laws of physics in the swirling patterns of galaxies and galactic clusters. The observations of cosmology, on this largest scale, are contained within the physical world of node (4) in figure 14.5, which in turn provides a context for understanding the 'cause' of the Big Bang and the origin of the universe more generally within the 'system of the world' presented in this paper.

14.3 A Context for Cosmology

The big picture for the present theory, as represented by the time cycle of figure 14.5 which sees the conscious observer in a dynamic interplay with the entire physical universe as an irreducible, integral component of the world, offers a very different perspective to the 'Copernican view', which sees humanity playing a far less significant role in the cosmos. The fact that the physical manifestation of humanity represents a tiny contribution to the total matter content of the Earth, which itself is in orbit around a far more massive sun, which in turn is one of countless stars distributed through the galactic structures of the universe all serves to cement the Copernican worldview concerning our apparent insignificance in the grand scheme of things. This is a misconception of the nature of the cosmos from the point of view of the present theory. On the other hand here there are potentially a vast number of subjective experiences which may complete the circuit for any physical universe, as described for figure 14.6, and each objective physical universe is one of a potentially vast number solutions of the form $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ capable of supporting self-reflective temporalising beings.

In any case the 'cosmological principle', as described in section 12.2, is valid for our universe in being sufficiently consistent with empirical observations to provide a valuable aid in finding solutions for the spacetime geometry on the largest scales observable. Such an entire solution for a physical universe, represented in a spacetime of unlimited 4-dimensional extent as depicted in figure 13.6 and with a geometry expressed as $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ in the present theory, is created as a mathematical possibility *within* the system of figure 14.5. The nature of this geometric solution is very much in the spirit of general relativity for which a spacetime geometry such as the Schwarzschild solution of equation 5.49, although typically employed to determine a planetary orbit about a star, represents an infinite 4-dimensional spacetime.

Here the possibility of the overall mathematical solution represented in figure 14.5 is the reason why the universe exists. Our local perspective of observing the flow of cause and effect in the everyday physical world leads by analogy to the presumption that the universe itself must have been created either by an event in time or by an event coinciding with the beginning of time. For any creation event in time the question then ever remains regarding the cause of that event while for a creation event at $t = 0$ the nature of an event without an apparent cause is certainly no less problematic from a conceptual point of view. In either case there are an array of further conceptual difficulties regarding the origin our own universe, such as the 'start-up problem', as described towards the end of section 12.3.

In the present theory the creation of the universe is not something that 'happens' in the Big Bang, or temporally before it, rather the very early universe and the Big Bang correspond to a certain region of the spacetime geometry at a particular epoch of the full 4-dimensional solution. This early epoch is beyond the horizon of our direct experience but its existence depends upon the self-reflective temporalising experience that arises in the history of the universe, as does everything in the cosmos. All the physical structure and conditions of the universe, including that for all future as well as past epochs and throughout the vast spatial expanse both within and beyond our observational reach at any epoch, are brought into being through the nature of a temporalising entity, which in turn is supported within the physical world, as depicted here in figure 14.8.

All experience in general is played out through a moment in time, including our perception of the physical world, with the structure of the universe being mathematically described by a solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ as conforming to the full form of temporal flow $L(\hat{v}) = 1$. Hence here the structure of the entire universe is derived mathematically through a moment of time, typically conceived as a duration of order one second as represented by a small one-dimensional interval ∆s, as the window through which it is perceived for example by the observer in the centre of figure 14.8. This contrasts with the standard cosmological models for which the entire observable universe evolves physically from a vanishingly small 3-dimensional spacelike hypersurface of size $a(t)\Delta\Sigma \to 0$ for $t\to 0$, as described for figures 12.3 and 12.4 in section 12.3, corresponding to the point at the base of figure 14.8.

Figure 14.8: The physical universe contains its own means of creation as perceived through the window of an interval of pure time ∆s subjectively experienced by the observer within the world. This picture is in contrast to standard cosmology for which the observable universe evolves from a vanishingly small spatial extent $a(t)\Delta\Sigma$ at $t=0$. Here the scenario described for figure 13.4(b) has been depicted.

For the standard approach all of the field content, particle properties and physical laws in general need to be added onto the spacetime in order to determine the evolution of the universe from the initial spacelike state, which is presumed to exhibit suitable initial conditions. However for the present theory all of the fields and physical laws derive from the structure and symmetries of $L(\hat{v}) = 1$ through the necessary projection onto M_4 in framing our perception of the world, including the Standard Model particle properties as identified in chapters 8 and 9. Here the apparent 'initial conditions' for $t \to 0$ represent a particular region of the full spacetime solution as required in order that the overall solution contains a habitable epoch such that the circuit in figure 14.5 closes, and hence in principle the initial conditions might appear improbable from the standard perspective.

Indeed, as described in the previous two sections, developing the theory from an interval of one-dimensional temporal flow Δs brings with it the possibility of constructing a universal foundation as represented in figure 14.5, while beginning with a spacelike hypersurface $\Delta\Sigma$ at $t=0$ leaves questions open concerning not only the source of physical laws and the nature of the initial conditions but also the origin of spacetime itself. Further the existence of the temporal moment is evident, in fact we are perhaps more intimately familiar with our experience of it than of anything else in the world, while the hypothetical initial spacelike state of the universe is an extremely remote entity. Hence overall, the notion of the present theory that everything is 'perceived through a moment of time' is perhaps not less reasonable than the standard picture for which everything 'evolves from a point of space'.

The system constructed in figure 14.5, for which figure 14.8 represents a particular manifestation such as our own world, can be considered as being centred fundamentally upon addressing the question of how it is possible to have subjective experiences of a world. As described in the previous section such experiences always take place here and now in the world, with everything else we can say about the universe, whether at some distance in space or extrapolated through time into the future or the past, necessarily consistent with the fact that we experience the world here in the present moment. The environment we experience in the present incorporates, amongst other things, observations based on a geometrical spacetime manifold; in particular we are able to perceive a world since it is cast against an approximately flat spatial background. However, there is no reason to expect the mathematical preservation of such an approximately flat pseudo-Euclidean spacetime indefinitely into the past as we extrapolate beyond the horizon of our direct physical experience of the world. The geometry of the very early universe for example, being beyond our immediate perception, with a potentially extreme spacetime curvature, is not required to be compatible with our a priori imposition of a flat framework of space and time within which to organise our impressions of the world and plot our actions within it.

Hence neither an approximation to spatial flatness nor any other constraint on the 3 or 4-dimensional geometry is required for the early universe regions of figures 13.4 or 14.8, in particular in approaching $t \to 0$. In fact at earlier times there remains no requirement for the identification of a 3-dimensional spatial or 4-dimensional spacetime manifold structure of any kind, as is the case for the scenario depicted in figure 13.4(a). However while the identification of the manifold M_4 itself, together with the projection $v_4 \in TM_4$, may break down at an epoch before the Big Bang the general form of temporal flow $L(\hat{v}) = 1$ remains ever valid for any value of the fundamental temporal parameter, even for $s \to -\infty$ as described for figure 13.4(a) in section 13.2.

From this point of view while the universe can be considered to be infinitely old, in terms of the value $s \to -\infty$, the cosmic time $t = 0$ can be considered to be the point in time at which a 4-dimensional spacetime manifold M_4 unfolds from the form $L(\hat{v}) = 1$, as depicted in figure 13.4(a). The familiar laws of physics in 4-dimensional spacetime, including the second law of thermodynamics expressed in terms of the degrees of freedom of Standard Model particle states and the gravitational field, may first be collectively applied as they emerge from the Big Bang at $t = t_v$; for either the scenario of figure $13.4(a)$ or (b) as also described in section 13.2. While we do not directly interact with the very early universe we are intimately connected with it not only through observations in cosmology but also through the need for the conditions of both stellar and biological evolution to arise and be consistent with the support of self-reflective beings at the present epoch.

In order to achieve this in addition to the microscopic field and particle interactions underlying the macroscopic gravitational structure $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, as empirically observed in the high energy physics laboratory and the cosmos respectively, at an intermediate scale the laws of physics implicit in this solution must necessarily be compatible with the development of the structures of molecular biology, such as DNA, which underpin the evolution of life. The complex biological structures implicit in figure 14.4, correlated with the subjective experience of temporalisation, must themselves arise in the material dynamics of the universe in a manner consistent with the laws of physics in 4-dimensional spacetime. That is, the physical universe we observe must support not only the formation and history of the solar system but also the evolution of biological life on Earth and the birth and development of specific self-reflective beings as manifested in human form and as represented in the centre of figure 14.8 for our world.

It could be asked: if the whole universe is brought into being through an experience of it here and now, why does it appear that biological evolution, leading up to the human race was necessary? Why not have readily formed humans along with the Earth and our local environment suddenly appearing, along with the identification of the M_4 manifold itself dating from the 'cosmic time' $t = 0$ just a few centuries or even a few minutes ago? However, the full extent of our spacetime world, including everything causally related to us from the past, must conform to the form of our perception in spacetime through the breaking of $L(\hat{v}) = 1$ over M_4 and the consequential laws of physics as implicit in the solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. The flow of the world in our past and into the future must obey these laws and also be consistent with our biological form as observers in the present.

Such an overall solution might be much more likely achieved through a very simple initial state followed by a prolonged cosmic and biological evolution as shaped by the laws of physics, rather than the apparently more direct route via a highly improbable 'initial state', in the form for example of a 'snapshot' of the universe taken a few minutes ago, which may in any case be prohibited through contradiction with the necessary laws of physics. This would still be the case even if the 'snapshot' only met the minimal requirement of preserving the complex form of the local environment, in which case the large scale cosmos would also most probably look very different to our universe. At the other extreme the universe, as an extended spacetime manifold, may not have a temporal origin at all in the sense that arbitrarily early times with fundamental time parameter $s \to -\infty$ might be contained within M_4 . This is the seemingly more natural scenario depicted in figure 13.4(b) for which the 'initial state' corresponds to the asymptotic conditions as $s \to -\infty$ and $t \to 0$, as also described for figure 13.5.

On the other hand the conditions in the universe observable today, even neglecting the consequences of the cosmic expansion, could not have prevailed indefinitely into the past. The laws of physics, in particular the second law of thermodynamics, demonstrate that it is not possible to sustain an everlasting immortal species on the Earth, and itself implies a necessarily finite lapse of time into the past to an apparent origin for our physical universe, which is also consistent with the observed expansion of the universe. Hence human life forms must have been moulded out of the state of the physical world at an apparent temporal origin of the 4-dimensional universe, that is the time at which the familiar laws of physics were established, culminating in a physical evolutionary process which in our case involves the processes of genetic mutations and natural selection. This apparent temporal origin must itself have an explanation in terms of the overall theory, and is here associated with the phase transition at the end of the Big Bang, that is at $t = t_v$ in the scenario of figure 13.4(a) or (b) and as also indicated in figure 14.8.

That 'there was evolution' is a statement from our perspective within the universe, which itself can be considered from an outside perspective as an 'atemporal' static 4-dimensional entity, about the world as a whole and the structure it must have for us to exist here and now in 4-dimensional spacetime. To 'visualise' the whole universe it is convenient to return to the three-dimensional spacetime analogy and combine the content of figures 14.4, 14.7 and 14.8. Through the circuit of figures 14.5 and 14.6 life draws itself into being out of the 'mathematical vacuum'. While the laws of nature on our spacetime manifold are carved out of the general form of progression in time, the actual physical forms we encounter in the universe, whether in our present or uncovered from our past, are moulded to conform with the possibility of our own bodily existence and conscious experience within it.

Many features of the world that we observe, such as our existence within a community of beings (the experiences of whom mutually dovetail together as described for figure 14.7) rather than finding ourselves in isolation, are the way they are since the world in which we find ourselves situated must accommodate a physical sequence of events, including for example an evolutionary and social history, leading up to the form of each individual experience.

All matter of the universe is brought into existence through our experience and perception of it as being mathematically, and hence physically, connected to the necessity that the experience itself exists. Hence all of our body organs, blood vessels and so on, as well as the human brain, necessarily come into being through the mechanisms and processes that give rise to life, in terms of its physical parts, along with the entire biological world, through the logical and rational requirement that we must be physically sustained within the world which we experience. The seemingly great improbability of life in terms of the complexity of biological structures such as sensory organs and the nervous system is essentially irrelevant. If such a biological system is physically possible at all and represents a self-reflective temporalising structure within the mathematical system of figure 14.5 then it will draw itself into being and *exist* as the realisation of an underlying mathematical necessity.

The constraints on the form of such a mathematical solution will be all the more stringent if there are essentially no free parameters in the breaking of the full form of temporal flow $L(\hat{\boldsymbol{v}}) = 1$ over a base manifold M_n (for an *n*-dimensional world solution). However, as for any mathematical problem, whether or not a solution actually exists does not depend upon the apparent difficulty of the problem. Whether or not a degree of tuning is possible for the symmetry breaking parameters (and whether or not $n = 4$), as considered in section 13.3, and regardless of the extent to which the apparent 'initial conditions' of the universe might be constrained, life will find a way if any solution for the structure in figure 14.5 exists, no matter how difficult or how remote the possibility of such a solution might seem to us.

Conscious life draws itself into being, through a self-supporting system, within the constraints of the mathematical form of the physical world it engages with. This is not necessarily a straightforward feat to achieve, in the sense of the non-trivial mathematical and physical structures required. Indeed, the fact that our ability to physically experience the world relies on the support of a human body which is enormously complex on the scale of the fundamental laws of physics (gravitational and quantum particle) is itself evidence of the difficulties of embedding the physical manifestation of a conscious life within a physical world constructed within the constraints imposed by the underlying mathematical progression in time experienced by the conscious beings themselves.

As a solution to the cycle of figure 14.5 the physical world of node (4) can be described in the mathematical terms of the full 4-dimensional spacetime structure $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ as moulded in conformity with the simple state of a moment of time in node (1). Alternatively the structure of the universe can be described effectively in the physical terms of a dynamic evolution from an apparently initial state at $t = 0$, as discussed for figure 14.8. Expressed in this latter way the physical development of the organic form of conscious beings out of a comparatively far simpler physical state at the apparent temporal origin of the world, according to precisely determined laws of physics, not surprisingly requires a relatively long period of biological evolution on a planet such as the Earth in a stable orbit around a star such as the sun. Hence the fact that we find ourselves in a universe at a spacetime location such that the sun is 149 million kilometres away in space and the Big Bang is 13.8 billion years away from us in our temporal past, as depicted in figure 14.8, have similar explanations: both are required of our physical environment in order that we, conscious beings, can consistently exist here and now. The observed vastness of the cosmos that surrounds us beyond the solar system is in some sense a byproduct arising from the non-triviality of realising a solution for the overall structure in figure 14.5, albeit a byproduct which is entirely 'weightless' from a mathematical point view as described towards the opening of section 14.2.

Analogous observations would apply to worlds other than our own, drawn into existence as a solution for the general form of figure 14.5, insomuch as it would seem surprising for a 'simple' solution to exist. The question concerning the uniqueness of our world, as considered in section 13.3, requires consideration of other worlds that could be created by and through other self-reflective beings. Since the 1950s philosophers in this world have sometimes enquired "what is it like to be a bat?", which is very difficult to answer since, amongst other things, bats and humans have different forms of sense perception. This kind of question becomes yet much harder if we attempt to enquire "what is it like to experience a different possible world to our own?" Here we refer to a different world with different laws of physics and perhaps even a base manifold with an intrinsically highly non-Euclidean geometry or a different dimensionality to ours.

All the varieties of other possible worlds with different laws of physics still have significant features in common, assuming they fall within the general framework described in this paper, involving a multi-dimensional form of temporal flow. The full form $L(\hat{v}) = 1$ may in a strict sense represent the greatest possible dissolving of the temporal flow via an infinite dimensional channelling through $\hat{v} \in \mathbb{R}^{\infty}$, as alluded to towards the end of section 13.3, and hence be unique for all worlds. Symmetries such as E_6 acting on the form $L(v_{27}) = 1$ with $v_{27} \equiv \mathcal{X} \in \mathrm{h}_3\mathbb{O}$ or E_7 acting on the form $L(v_{56}) = 1$ with $v_{56} \equiv x \in F(h_3\mathbb{O})$ may also represent significant mathematical resonances which dominate the actual physics observed in any universe. The physical laws themselves are then effectively determined in breaking the full symmetry, for example through the identification of a subgroup acting on the subspace of vectors $v_n \subset$ \hat{v} projected onto the tangent space of an *n*-dimensional base manifold M_n . This smaller space, together with the symmetry group for $L(\mathbf{v}_n)$, is broken out of the larger space and symmetry group of $L(\hat{v}) = 1$ in the formation of a global background manifold which acts as a geometrical reference frame for events perceived in the world. This background provides the relief against which apparent material objects are brought to the attention of the self-reflective beings through the laws of physics resulting from the breaking of the full symmetry group G .

Whether there is only one such kind of world, of which our own would then be a particular manifestation, or several, which might even be catalogued, is likely to be difficult to determine (perhaps even much more so than categorising all possible biological life forms given the laws of physics within our own universe, whether on the Earth or elsewhere). Certainly for any world to be possible in this framework is equivalent to the statement that it must actually exist, and in this case our variety of universe would not be entirely unique. However, we would not be able to communicate with other worlds, or the creatures living within them, and there is no question of interference with the internal consistency of our own world.

While the existence of other worlds with different laws of physics is an open question, there will be, according to this theory, many possible solutions for a geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ on a 4-dimensional spacetime manifold M_4 , apart from our own world, which share the same laws of physics and will also internally support conscious life under circumstances similar to those in which we find ourselves on Earth. The notion of 'many worlds' as an *interpretation* of quantum mechanics is distinct from, although implicit within, the overall framework presented in this paper, where here we are referring to the 'many solutions' embedded within the theory, as discussed in section 11.4. Different solutions for $G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$ involve $\delta Y \leftrightarrow \delta \hat{\mathbf{v}}$ field exchanges in principle anywhere on the spacetime manifold M_4 , even back to the Big Bang epoch for cosmic time $0 < t < t_v$, and when considered from a dynamic point of view our universe in some sense might be considered to have 'branched' from another possible solution at each quantum event.

Such quantum transitions, which are indeterministic from the perspective of a single universe, taking place in the very early universe might serve to seed the eventual formation of stars, galaxies and large scale structure generally, as alluded to in sections 12.3 and 13.2. That is, in part due to the causal temporal accumulation of such probabilistic events, the impact of a quantum fluctuation for $t < t_v$ on the overall structure of the universe might generally be far more dramatic than a similar 'branching' resulting from a 'Schrödinger's cat' type experiment performed at the present epoch. While many solutions for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ might be considered to be mutually related by such 'branching' events, in the present theory each solution is primarily interpreted as an independent full 4-dimensional spacetime solution in its own right.

Much of modern science adopts an essentially materialist worldview in line with our Newtonian heritage. From this objective point of view with the universe seen as a fundamentally material phenomenon, created in the Big Bang as an inanimate physical entity with the various seemingly arbitrary parameters of cosmology and particle physics, it appears extremely fortunate for us that such a world can both support biological life and lead to the development of our own society, culminating in our own personal human form, through a series of chance events. In particular life itself, as we know it, would be impossible given a small change in any of a range of the empirically measured physical parameters.

As usually presented this means that, for example, the laws of physics are required to be such that the chemical elements necessary for life on Earth could be manufactured in the hot Big Bang – which successfully accounts for the relative abundances of the light nuclei, D, 3 He, 4 He and 7 Li, cooked up from a hot soup of protons and neutrons in the first few minutes – together with the much later generation of the heavier elements through stellar nucleosynthesis. The latter stage is possible thanks to a seemingly fortuitous energy level of the carbon nucleus that allows the three-body reaction 3^4 He \rightarrow ¹²C to proceed at a reasonable rate. In 1953, in a famous case of anthropic principle reasoning, the necessity of this carbon resonance was predicted by Fred Hoyle, to account for our own presence in the world as a carbon-based life form dependent on the heavy elements. Given this motivation the resonance was then experimentally observed shortly afterwards.

However, for the present theory the universe, through the structure of figure 14.5, is born out the intimate interplay between conscious beings and the physical world. The complexity of the resulting physical structures within such a solution creates the illusion of the fortuity of our own existence. Due to the non-trivial nature of solutions achieving a completion of the time cycle in figure 14.5 any possible physical world is likely to appear highly complex, as discussed above. Hence beings in any such world are likely to require a number of parameters to describe empirical findings in their world, as for the Standard Model of particle physics in our world for example. Hence in turn, with the physical support for known biological life forms apparently collapsing under a hypothetical change in the empirical parameters, beings in such a world might consider themselves lucky. Given the familiarity of our own world as a starting point we can readily conceive of many ways in which a physical world could not support life, through small perturbations to the properties of our own universe,

but it is much harder to conceive of very different worlds with very different solutions for supporting the structures of conscious life.

Hence here there is a major contrast between the present theory and various forms of the anthropic principle, which are generally subject to criticism due to their lack of predictive power. For example based on the anthropic principle a theory may postulate the existence of a very large ensemble of different universes with different initial conditions, physical constants or laws of nature – then the fact that our universe is necessarily a member of the ensemble in which the structures for life can form necessarily greatly restricts the possible structure of the physical laws and conditions that we can observe. However the potentially vast range of physical properties for the worlds of the whole ensemble, most of which are presumably not observed by any being, is in no way limited by this principle.

Here the present theory is 'anthropic' to a more extreme extent in the sense that the only worlds that *exist at all* are those that can be brought into being through a conscious, temporalising observer, and in this case we may hope to discover the opposite conclusion that the laws of physics are necessarily determined, or at least highly constrained, by this requirement (although naturally there will always be the more trivial anthropic matter of the local selection of a habitable environment, such as the Earth, within such a world). That is, rather than postulating a large ensemble of typically inanimate physical universes with a range of parameters, one of which happens to provide a suitable environment for ourselves, we draw our own world into existence, sculpting the physical contents of the world out of the possibilities inherent in perceiving a world through the forms of $L(v) = 1$ as a solution for figure 14.5.

The number of parameters needed to describe the projection of $v_4 \in TM_4$ out of the components of $\mathcal{X} \in h_3 \mathbb{O}$ or $x \in F(h_3 \mathbb{O})$, involving for example the dilation symmetries described in the opening of section 13.2, is much less than the number of parameters needed to describe the empirical data as observed in particle physics and cosmology. Hence the present theory in principle will be highly predictive even if there are some possibilities for the variation of certain parameters within the mathematical constraints of the theory, that is with an anthropic degree of tuning for the parameters involved in the projection of $v_4 \in TM_4$ out of $L(\hat{v}) = 1$. However, in chapter 13 we have generally presumed that interactions between the fields as determined by the constraint equations 11.29 result in a fixed and stable value around $|v_4| = h_0$ emerging from the phase transition at $t = t_v$ in the very early universe, as described for figure 13.3(c). The potential uniqueness of the laws of physics and particle properties arising at this time suggests the present theory should be profusely testable.

Here the perspective is to consider the 'early' universe to be an object of study as a limiting extrapolation from our present experience in the world rather than as an objective self-sufficient physical state that happens to be the causal origin leading up to the present conditions in our world, as has been described for figure 14.8. With a solution for the time cycle structure of figure 14.5 taking priority and founding the theory, not only is nothing needed as a temporal antecedent of the Big Bang to cause the universe to exist, but the particular conditions of the Big Bang and early universe are shaped by the overall consistency of the solution within the structural constraints implied in figure 14.5. These constraints on the apparent 'initial conditions' of the early universe are ultimately manifested in the physical and biological processes required to

support self-reflective life forms at the present epoch, as described above.

Here we take, possibly rather indirect, measurements of cosmological structure including that for the earliest epochs of the universe, as for laboratory experiments in particle physics, as being extensions of our world experience – quantitatively differing from the nature of everyday experience in the world more generally, but in all cases subject to the same laws of physics and all within the same system. From the basic experiences of thinking, listening to music, walking down the street and watching an apple fall from a tree to performing experiments and studying the structures of biology, chemistry and physics on all scales, there is a continuity from the notion of experience through to, and incorporating, the practice of experimental and empirical observations. In the present theory both the notions of scientific observations and subjective experiences more generally are drawn together and unified as particular manifestations of experience in time.

If the present theory were to be founded on a purely objective notion of onedimensional temporal flow, as modelled by the real line R, the sequence for figure 14.5 could still be constructed linking the nodes $(1), (2) \dots (6)$ but without the final link between nodes (6) and (1). For such a theory the subjectively experienced time of node (6) would hence be derived from a long chain of non-trivial steps $(1), (2) \dots (6)$ beneath which the fundamental objective temporal entity of node (1) would be very much hidden from our immediate view of the physical world, and would not be an entity we might directly perceive. However, this unnatural duplication of the concept of time in nodes (1) and (6) is avoided through the actual perspective of the present theory which has been developed *beginning with* the notion of subjective experienced time. Indeed temporal flow is not something that we 'see' in the world, as is the case even for the elementary forms of space, rather it is an innate characteristic of our engagement in the world. That is temporal flow is not a property of the world which we need to set out to discover, as for the 'hidden' structures of material phenomena or particle interactions of node (4) for example. Rather we do directly perceive the underlying temporal flow of node (1) since it is identified with our immediate experience of time in node (6), hence in turn completing cycle of figure 14.5.

The overall system of figure 14.5 is perhaps best understood by thinking through the cycle of six nodes and links in turn, beginning from any point, but the structure can be contracted down in a number of ways including a more minimal scheme describing an interplay between experience and the empirical, or essentially between subjective temporal flow and the objective laws of physics as associated with nodes (1) and (4) respectively. Ultimately the full set of six nodes coincide as six facets of the internal structure of the possibility of conscious experience, conceived as a unified whole, essentially adopting the philosophical outlook of existentialism as alluded to in the previous section. As discussed in the previous section, from this point of view the possibility of an experience is a more fundamental concept than the individuals who believe they have them, and with the laws of physics, which shape both the physical individual and his environment, also determined through the constraints on the possible forms an experience can take within the system of figure 14.5.

From the philosophical perspective of materialism, which is grounded largely in node (4) of figure 14.5, the 'problem of consciousness' arises since the concept of subjective experiences seems to be of a qualitatively different nature to anything studied in the realm of the physical world. While from this point of view consciousness appears mysterious and beyond the reach of the physical sciences, it nevertheless remains the case that conscious experience is a very real phenomenon of the world, and indeed it is the feature of the world with which we are most intimately familiar. Hence an inclusive scientific theory should either have something to say regarding the nature of consciousness or provide a good explanation as to why it should not, as suggested shortly before figure 14.2. On this basis the speculative structure of figures 14.2–14.4 has been studied here in section 14.1. One possible justification for not addressing this question regards the complexity of the human brain, being beyond the current scope of an exhaustive scientific understanding.

On the other hand the nature of subjective experience can be very simple, as exemplified by the 'thought experiment' involving picking up a pen or pencil as also described in section 14.1. This suggests that the broad objective physical correlate of such experiences might also be described in terms far simpler than those required to give an account of the detailed structure of the brain. Together with the practical experiments of Libet and others discussed in section 14.2 it is clear that the phenomena of conscious thought are in any case open to study. Indeed research into consciousness is a scientific field of study in its own right, although one which is not traditionally closely linked with physics. It's relevance for the present theory lies in the close relationship between the nature of consciousness and the structures proposed to complete the cycle of figure 14.5. In return the perspective of the present theory, in which consciousness is closely associated with temporalisation and related to the physical world through figure 14.5, might in principle be of value for the corresponding area of study in neurology, for which a firmly materialist standpoint is commonly adopted.

It is suggested here that consciousness is not something that can be fully explained as a phenomenon arising solely within a pre-existing physical world, as would be required from a purely materialist perspective. Subjective experiences cannot be directly described in terms of objective matter, but rather correlate with certain mathematical structures which underlie the physical world within the context of the system depicted in figure 14.5. On the other hand the content of the physical world is not fully contained *within* the horizon of our conscious observations, as might be the case for the pure idealist. We can conceive of an infinite expanse of the physical world in space and time beyond the horizon of our direct experience as supported by the full mathematical solution for $G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$ implied in the structure of figure 14.5, which itself provides the context for all the structures of cosmology. Conscious experience is an irreducible feature of the world and the means by which a mathematically possible universe is realised through the intimate interplay between the subjective and objective aspects of figure 14.5.

Much of the apparent mystery of 'consciousness' owes to the fact that nothing exists without its support and hence it is impossible to step back and isolate the phenomenon 'in itself'. Everything that exists or happens does so within the context of consciousness, even our awareness of a discussion of consciousness itself, with the phenomena of the physical world ultimately inseparable from the experiences of temporalising beings. A theory which, on the contrary, attempts to construct a notion of consciousness entirely within the limits of a given independent physical world, implying that such a world can 'exist' even in the absence of such sentient beings, is necessarily

dealing with an incomplete system. Rather, while also supporting the physical correlate of conscious mental phenomena, the physical world is itself engulfed within the sphere of conscious experience, as implied in the relations depicted in figure 14.5.

For the above materialist worldview in addition to the difficulty in constructing an explanation of consciousness upon a given physical world, as alluded to also for the right-hand end of figure 14.1 and discussed more generally in section 14.1, on the other hand there remains the second major loose end regarding a foundation for the physical theory itself.

An appeal to 'beautiful mathematics' is often made either explicitly or implicitly as a significant motivating force in theoretical physics, promoting a sense that nature 'ought' to make use of aesthetically pleasing mathematical structures. While some successes may be cited, notably for example regarding the Dirac equation for a fermion field (quoted here in equations 3.99 and 11.31 with a gauge field interaction included), the achievements of this approach, in terms of discovering empirical phenomena that match a beautiful mathematical theory (applied in particle physics or cosmology), have been particularly limited in recent decades. This approach also has serious philosophical difficulties, regarding not least the highly subjective notion of 'beautiful mathematics' itself and the means through which physical entities in the world should relate to the mathematical components of the theory.

Alternatively an objective physical theory might be founded upon a conceptual idea regarding the nature of an inanimate physical world, which will subsequently be formulated and developed in mathematical terms in order to derive testable consequences for the theory. Examples of this approach include the description of gravitation in terms of a curvature of 4-dimensional spacetime in general relativity, or the properties of discrete particle-like entities interacting in a flat spacetime. However it is difficult to conceive of any physical concept which does not itself stand in apparent need of a further underlying explanation. Progress may be proposed, for example with gravitation and the geometry of our world in 4-dimensions arising out of a more fundamental higher-dimensional spacetime or with particle phenomena deriving from a field theory, but at some point the basic physical entities, together with perhaps a Lagrangian formalism or a quantisation procedure, is essentially 'postulated' as an apparently necessary starting point.

The foundations of such a theory can be justified provisionally on the grounds that 'one has to start somewhere', as alluded to in the opening of section 14.1, provided the theory satisfies a criterion of empirical success. Based upon that success we learn what a more fundamental theory should effectively look like in a certain environment or under certain limiting conditions, such as those for general relativity or quantum field theory as described for table 11.1 in section 11.4. Whether an objective physical theory is founded chiefly upon mathematical, conceptual or empirical grounds (and in practice in some combination) the foundational loose end is generally accompanied by questions concerning the nature of the origin of the universe in the Big Bang, which is needed in order 'to get the ball rolling' in the first place, as summarised in point (1) in the opening of section 14.1.

The approach of the present theory, with respect to the two loose ends of figure 14.1, is to fully embrace the subjective element of our engagement in the world. With all experience in the world having a temporal aspect the theory is founded purely on the notion of a one-dimensional flow of 'time' as a necessary component of both the subjective and objective world. Since time is a feature of the world, which we experience directly without any intermediate interpretation, this offers an extremely conservative starting point for a theory. To be aware of anything at all is to experience an irreducible moment in time, as a basic aspect of thought and experience generally. With all thinking having a necessarily temporal dimension we have essentially retreated to the minimal observation that, with a twist on the famous words of Descartes, 'I think therefore I temporalise'. This provides the mathematical basis for a full physical theory which supports the entire structure of the universe as perceived in an experience itself. In its simplicity this starting point is largely devoid of any arbitrary aspects, unlike the case for most theories which are motivated on mathematical or conceptual grounds which are purely objective.

Through the dual subjective and objective nature of time, both modelled on the same mathematical real line, this theory can ultimately also supply its own foundation, tying up the two loose ends of figure 14.1 in the shape of figure 14.5. Although here the theory is motivated from the direction of a conceptual argument, based upon temporal flow, rather than from the direction of 'beautiful mathematics', the mathematical structure represented in figure 14.5 has itself a degree of elegance in its simplicity and self-contained nature. However instead of beginning with mathematical beauty together with the presumption of its necessary application to the physical world, here the realisation of the mathematically elegant structure described in figure 14.5 contains its own inevitability, in that it incorporates both self-reflective intelligent entities and its own foundation.

Further, this structure provides a context within which an entire universe, as depicted for example in figures 13.6 and 14.8 and supported by a spacetime manifold M_4 of infinite extent, forms part of the overall solution, hence incorporating all features of the physical world, from the microscopic to cosmological scales, including the Big Bang and events in the arbitrarily distant past. Although the subjective aspects are necessary to conceive of the whole system and help motivate the initial foundation of the theory in terms of the flow of time, the structures contained within nodes (1) – (4) of figure 14.5 can be essentially treated as an objective physical theory, as has been the case for the large majority of the work presented in this paper, which may be measured against observation in the empirical world as for any other theory.

While the simplicity and elegance of the mathematical structure of nodes (1) – (4) might itself be considered, in order to fully justify the present theory not only on conceptual grounds but also from the perspective of the mathematical elegance of figure 14.5 as a whole a more rigorous mathematical account of the lower half of the chain through nodes $(4) \rightarrow (5) \rightarrow (6) \rightarrow (1)$ might be desirable. However, all elements of the cycle are open to such an exploration, and in section 14.1 we described a possible approach to uncovering a mathematical correlate of self-reflective subjective thoughts and decision making.

There we also noted a close analogy between the mathematical structures relating to Gödel's notion of decidability and the properties of physical devices relating to Turing's notion of computability. Following Turing and the ambition to develop artificial intelligence it is conceivable to attempt to build a machine exhibiting the properties self-reflective conscious experiences and creative thought. The design of such a machine might include a complicated arrangement of malleable and adaptable electronic, and even biological, components capable of internal development, as well as an array of sensory input devices and means of interacting with the environment. Given the design on paper, for the machine to actually 'exist' it would then need to be built, requiring the physical assembly of the necessary technological components. Only when manufactured in this way could we declare, in the words of Dr. Frankenstein, that "it's alive!".

If the machine could think and have experiences in a similar way that we do, it might also ask itself how the physical universe and its place in the world came into being, and might also be drawn to a conclusion in the form of the system described in figure 14.5. For the case of this artificial intelligence the full physical environment must include not only *our* biological evolution but also the particular human inventors and technicians with the ability to design and construct the machine.

On the other hand if we consider directly the purely mathematical construction of self-reflecting elements relating to Gödel's theorem or a similar theoretical structure, rather than taking the computing route of Turing, the conclusion is somewhat different. In this case we might design a particular mathematical system capable of describing self-reflective states and which also contains its own foundation as sketched in figure 14.5. This mathematical structure, as for any logically possible mathematical construction, is in principle a free creation for our mind to think about abstractly and objectively from an independent point of view. While we can discover such a logically coherent structure in this case any ambition to *build* such an entity would be meaningless (unless it could be mapped onto the design of a practical machine as described above). However, since the kind of structure depicted in figure 14.5 has the characteristic that it contains thoughts and experiences of internal elements all within the same structure together with its own foundation it is in the nature of this mathematical system to spontaneously realise its own existence, detached from any external support. The contention here then is that our own experiences in our own universe are a particular manifestation of precisely such a self-illuminating world.

Chapter 15

Towards a Complete Theory

15.1 Summary and Future Directions

The underlying unifying principle for the theory is simply the observation that everything takes place through progression in time. Based upon this principle in this paper we have explored the extent to which the empirical phenomena of the physical world might be accounted for. In the previous chapter we have described how physical structures in the world might themselves inscribe subjective experience of progression in time and hence act as the source of temporalisation itself. Regarding the general structure of the theory, we first summarise here the main novel ideas presented as the foundation for the physical world as described in detail in the preceding chapters.

The mathematical possibility of a multi-dimensional flow in time is expressed through the general mathematical form of progression in time $L(\mathbf{v}) = 1$ as derived for equation 2.9. The creation of an extended spacetime manifold out of the flow of time is possible through an innate subjective interpretation of a subset of the algebraic structures incorporated within $L(v) = 1$ in terms of a geometrical representation. This spatialisation of the world is considered a subjective phenomenon insofar as it is through it that experience of a physical world by sentient beings is possible. The description of the geometry of the resulting extended external spacetime is identified with that for general relativity, as applying for all physical scales.

Since the extended frame for perception is constructed out of a substructure of the full form of temporal flow described by $L(\hat{\mathbf{v}}) = 1$ a natural mechanism for breaking the higher, unifying, symmetry of time arises. Non-gravitational fields and interactions are induced on the spacetime manifold through the residual components of the full form and symmetry of $L(\hat{v}) = 1$. The possibility of a degeneracy of solutions for the external spacetime geometry underlies the phenomena of quantum theory and particle physics. The breaking of explicit full symmetry groups for candidate forms for $L(\hat{v}) = 1$ over the 4-dimensional spacetime base space is found to yield structures closely correlating with features of the Standard Model of particle physics.

A significant novel feature of this theory is that the spacetime manifold is not postulated as a starting point, rather it is grounded as a possible structure within the
multi-dimensional flow of time, arising out of the translation symmetry inherent in the form $L(\hat{v}) = 1$. In other background-free theories one main difficulty is to explain the origin of such an extended spacetime structure. Hence most theories employ a pre-existing 4-dimensional manifold, or a higher-dimensional spacetime arena in which to embed the former, and then introduce fields or other mathematical entities upon the manifold. Since here we extended the symmetry group of $L(\mathbf{v}_4) = 1$ to act on a higher-dimensional form of time, absorbing the 4-dimensional one, these ideas could also be considered as a theory with extra dimensions. However, here they are not extra dimensions of a spacetime, although algebraic forms or symmetries which also have such a geometrical interpretation (including the temporal form $L(v_{10}) = 1$ with the symmetry $SO^+(1,9)$ considered in figure 5.1 for the model of section 5.1) may happen to arise in the mathematics. On the other hand this theory can also be conceived as a rather more economical approach with fewer dimensions, in that the world emerges from a one-dimensional progression of time.

The physical theory presented in this paper, based on the notion of a fundamental underlying progression in time taking the general form $L(\mathbf{v}) = 1$, has progressed along four main fronts, as depicted in figure 15.1. In this concluding chapter these theoretical developments are summarised along with a discussion of how they are mutually related and might be combined together in progressing towards a complete theory.

$$
(2)
$$

\n E_7 Action on $F(h_3\mathbb{O})$
\n $L(\mathbf{v}_{56}) = 1, D_{\mu}L(\mathbf{v}_{56}) = 0$
\n(Standard Model)
\n $G_{\mu\nu} = f(Y), G^{\mu\nu}_{;\mu} = 0, \text{ on } M_4$
\n $L(\mathbf{v}) = 1 \longrightarrow \text{ Many Solutions}$
\n(Kaluza-Klein Theory)
\n \downarrow
\n $L(\mathbf{v}) = 1 \longrightarrow \text{ Many Solutions}$
\n $\delta Y \leftrightarrow \delta \mathbf{v}_{56}$ Redescriptions
\n(Quantum Field Theory)
\n \downarrow
\n $L(\mathbf{q})$
\nLarge Scale Structure
\n $G_{\mu\nu} = f(\mathbf{v}_{56})$ and Generalised
\n(Standard Cosmology)

Figure 15.1: Developed from the original underlying notion of the primary role of temporal flow these four areas of progress $(1), (2), (3)$ and (4) have been described in detail in this paper in chapters 2–5, 6–9, 10–11 and 12–13 respectively. (In each case the main guide from established physical theory is appended parenthetically).

The four fronts of the theory described in figure 15.1 contain aspects of the interplay between the various forms of the flow of time considered, from one-dimensional temporal causality itself up to the largest form $L(v_{56}) = 1$, the full symmetry of which is broken over the base manifold M_4 . Individually these four fronts exhibit the following principal features:

- (1) Motivated by the notion of perception over a 4-dimensional base manifold M_4 four extended external dimensions are initially identified through translation symmetries of the full form $L(\hat{\mathbf{v}}) = 1$. Subgroups of 'rotational' symmetries of $L(\hat{\mathbf{v}}) = 1$ imply the identification of gauge fields on M_4 relating to both the external and internal geometry and the unifying framework of a principle fibre bundle for general relativity and classical gauge theory can be constructed. With the external and internal geometry correlated as the full symmetry of $L(\hat{v}) = 1$ is broken in the projection over M_4 this structure, with the four external dimensions identified as above rather than with the 'extra' dimensions being 'compactified', is reminiscent of non-Abelian Kaluza-Klein theories.
- (2) Motivated by its mathematically rich structure out of the infinite possible forms of $L(v) = 1$, a 56-dimensional form of temporal progression $L(v_{56}) = 1$ with a high degree of symmetry is identified through the action of the group E_7 in preserving a quartic form defined on the space $F(h_3\mathbb{O})$, containing the determinant preserving action of E_6 on the space h₃ σ . When broken over the external M_4 base manifold the residual internal gauge group contains features of the symmetry $SU(3)_c \times SU(2)_L \times U(1)_Y$ acting upon components of $F(h_3\mathbb{O})$, including subspaces identified as spinors under the local external Lorentz symmetry $SL(2,\mathbb{C})$ with charges under an internal $U(1)_Q$ symmetry, which are reminiscent of the Standard Model of particle physics.
- (3) Conforming with the underlying one-dimensional causal flow of time the degeneracy of field solutions for the world geometry $G_{\mu\nu}(x)$, consistent with the broken form of temporal flow expressed dynamically on the base manifold via expressions such as $D_{\mu}L(v_{56})=0$, selection rules for exchanges between gauge $Y(x)$ and spinor $\psi(x)$ fields may be obtained. This leads to interaction phenomena with a mathematical structure reminiscent of calculations employing the time evolution operator $U(t, t_0)$ in a quantum field theory based upon a given Lagrangian.
- (4) In constructing the base manifold M_4 out of the full form $L(\mathbf{v}_{56}) = 1$ and its symmetries variation in the magnitude of the projected subspace vectors $v_4(x) \in$ TM_4 , with $|\mathbf{v}_4|^2 = L(\mathbf{v}_4) = h^2(x)$, itself generates a non-flat external geometry. The general solution for the 4-dimensional geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ might also incorporate a cosmological term in principle deriving from the scalar components of $F(h_3\mathbb{O})$. Collectively the resulting large scale structure of the cosmos may correlate with the observed phenomena of the dark sector and properties of the very early universe, that is in a manner reminiscent of the standard cosmological model and inflationary theory.

Hence the theory represents new directions of research in fundamental physics branching into several areas. At the same time the main part of this work sits comfortably within the existing infrastructure of theoretical and experimental physics. The mathematical framework has been adopted entirely from that used in much of contemporary theoretical physics, with the novel input more in the nature of the overall conceptual picture.

The essential theoretical ingredients to account for the Standard Model of particle physics and large scale cosmological structure, while sidestepping the Lagrangian formalism and also providing a conceptual basis for the 'quantisation' of the fields, are in principle all found in the structures of the present theory. All four of the above fronts are directly related to consideration of the basic idea expressed in the general form of temporal flow $L(\mathbf{v}) = 1$, and are mutually related to each other. The immediate future direction and main aim for further study on each front is first summarised here:

- (1) Use the mutual relationship between the external and internal curvature in originating from symmetries of the same full form $L(\hat{\mathbf{v}}) = 1$ projected over M_4 , described in terms of the differential geometry of the structure of a fibre bundle, to derive the relation $G_{\mu\nu} = f(Y)$ in the form of equation 5.20 without any explicit application of an action integral such as equation 5.18 as adapted from Kaluza-Klein theory.
- (2) Determine a higher-dimensional form of temporal flow and corresponding symmetry to build upon the features of the Standard Model identified in the action of E_7 on $L(\mathbf{v}_{56}) = 1$ when broken over M_4 as summarised in equation 9.46. For example a presently hypothetical E₈ action on a full form $L(v_{248}) = 1$ might be sought, the structure of which will be guided by fields and interactions of the Standard Model Lagrangian yet to be accounted for.
- (3) Use a statistical approach to HEP phenomena with probabilities based upon field degeneracy, building upon the relationship with quantum field theory described for equation 11.46 and possibly employing the analogy between the properties of condensed matter systems and QFT, to develop the theory through to the calculation of cross-sections and the identification and conceptual understanding of particle states without imposing quantisation rules.
- (4) Build upon the geometry $G_{\mu\nu}(x)$ of equation 13.4, deriving from a variation of the magnitude $L(\mathbf{v}_4) = h^2(x)$, to a full general form $G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$ incorporating also scalar fields and applied for the large scale structure of the universe, in order to make a more quantitative comparison between the present theory and empirical observations in cosmology; with one aim being to deduce which scenario, such as that in figure $13.4(a)$ or (b), applies for the very early universe.

The main prediction of the theory at present is a mathematical one concerning the existence of an E₈ symmetry acting upon a quintic or higher order form $L(v_{248}) = 1$ as alluded to in front (2) above. This structure, as an extension from the E_7 action on $L(v_{56}) = 1$, when broken over M_4 should incorporate further Standard Model properties such as three generations of fermions, as motivated in detail in section 9.3. More generally the overall aim is to fuse the above four areas together in a full unified theory, and assess the consequences and possible predictions of the theory that can be further compared with and tested against empirical data from HEP experiments, cosmology and other observations. We begin here by observing the following relations between the four theoretical branches summarised in figure 15.1.

 $(1+3)$ The key motivation for front (1) is the identification of a smooth external geometry $G_{\mu\nu}(x)$ on M_4 as an arena for perception in the world. Since there is no similar requirement regarding the need for a 'smooth' internal geometry of gauge fields it would be more natural to *begin* with the structure of fronts $(1+3)$ combined, as implied in the relation $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ as a possible solution for the world geometry on M_4 . A finely fragmented and fractal-like structure of field exchanges $\delta Y \leftrightarrow \delta \hat{v}$ underlies the smooth external spacetime arena, with $G^{\mu\nu}{}_{;\mu}=0$ maintained as a geometric identity. In this way the degeneracy of many possible solutions brings the phenomena of general relativity and quantum theory together at the same time in the process of identifying the base manifold itself, rather than beginning with a 'classical theory' of the form $G_{\mu\nu} = f(Y)$ which is then 'quantised'.

The relation between the initial theoretical 'bare' fields and empirically observed 'dressed' fields was also described in the opening of section 11.3. Indeed, a geometrical relation of the form $G_{\mu\nu} = f(Y)$ might still be identifiable for macroscopic fields, such as the empirically observed electromagnetic field. Out of the complete framework the standard theories alluded to parenthetically for fronts (1) and (3) in figure 15.1 may be shown to emerge in the appropriate limits: namely Kaluza-Klein theory in a curved spacetime as an example of the macroscopic field limit of general relativity and QFT in the limit of a flat spacetime for microscopic fields, as described for table 11.1 in section 11.4.

(2+4) In the present theory the phenomena of electroweak symmetry breaking and in particular the masses of particle states observed in the laboratory arise out of interactions between the components of the vector-Higgs field $v_4(x)$ and other fields such as the fermions $\psi(x)$ identified in the components of $F(h_3\mathbb{O})$ through the terms of the quartic form $L(v_{56}) = 1$. On the other hand cosmological structure depends on variation in the magnitude $|v_4| = h(x)$ as $v_4 \in TM_4$ is projected out of the full form $L(v_{56}) = 1$ over M_4 , which itself provides a geometric explanation of the origin of mass in terms of an effective energy-momentum tensor defined in $-\kappa T_{\mu\nu} := G_{\mu\nu} = f(\mathbf{v}_{56}) \neq 0$. Hence these two notions of mass are intimately related via the field $v_4(x)$.

The dilation symmetries, acting on the components of $F(h_3\mathbb{O})$ as discussed in the opening of section 13.2, change the value of $|v_4|$ and may be significant in relation to the mechanism of electroweak symmetry breaking in the very early universe. The physics of the very early universe may also guide the identification of a higher-dimensional form of time, such as the hypothetical $L(v_{248}) = 1$ with E₈ symmetry. In particular the mechanism for generating a matter-antimatter asymmetry might be determined by interaction terms implicit in the form $L(v_{248}) = 1$ or involve a further internal gauge field deriving from the E_8 action, as also discussed in section 13.2. Hence the structure of the full form $L(\hat{\bm{v}}) = 1$ is closely linked with an understanding of significant questions in cosmology.

 $(1+2)$ In equation 6.3 of chapter 6 the generators of the symmetry of a 27-dimensional form of $L(v_{27}) = 1$ were introduced as operators that annihilate the cubic norm $\det(\mathcal{X})$ with $v_{27} \equiv \mathcal{X} \in \mathrm{h}_3\mathbb{O}$. A complete basis for this 78-dimensional Lie algebra of E_6 , as represented by vectors of the tangent space $R \in Th_3\mathbb{O}$, is listed in tables 6.6 and 6.7 at the end of section 6.5. Such a 'static' generator can be pulled back to a Lie algebra valued 1-form $Y_\mu(x)$ on M_4 , as initially described in subsection 2.2.3, and appears in 'dynamic' expressions on the base manifold. Kaluza-Klein models based on fibres identified with homogeneous spaces were reviewed in section 4.3, and might provide additional insight in comparison with the closely related theories constructed on principle fibre bundles described in sections 4.1 and 4.2.

With regards to the model described for figure 5.1 in section 5.1, with the full symmetry group $SO^+(1,9)$ acting on the form $L(\mathbf{v}_{10}) = 1$ over M_4 , the structure of the Lie algebra for $SO^+(1,9)$ can itself be expressed in terms of vector fields on the space of 10-dimensional vectors $v_{10} \equiv X \in h_2\mathbb{O}$ with $\det(X) = 1$, based on the opening of section 6.3. With $h_2 \mathbb{O} \subset h_3 \mathbb{O}$ embedded as a subspace a close connection is made with the above case for E_6 acting upon the homogeneous space composed of vectors $v_{27} \equiv \mathcal{X} \in \mathrm{h}_3\mathbb{O}$ of unit determinant. The E_7 action on $F(h_3\mathbb{O})$, broken over the 4-dimensional base space M_4 , represents a higherdimensional extension of this structure, while the full form of $L(\hat{\mathbf{v}}) = 1$ that provides the actual setting for a description of the real world is open to further investigation. Hence branch (1) relates to branch (2) of figure 15.1 essentially in the choice of $L(\hat{v}) = 1$ and the corresponding full symmetry group over the base manifold M_4 .

 $(2+3)$ Taking the example of the E₇ case, the generators of the internal symmetry action $R \in TF(h_3\mathbb{O})$ give rise to the gauge fields $Y_\mu(x)$ on the base space while the components of $\mathbf{v}_{56} \in F(\mathbf{h}_3 \mathbb{O})$ are also intimately related to the base manifold through the translation symmetry over $x \in M_4$ as originally described for figure 2.2. Hence since $F(h_3\mathbb{O})$ forms the representation space of E₇ the gauge fields $Y_\mu(x)$ naturally couple with components of $v_{56}(x)$, including the spinor fields $\psi(x)$. The dynamics of the interaction between the components of $v_{56} \in F(h_3\mathbb{O})$ and the gauge fields, under the constant form $L(v_{56}) = 1$, is subject to the constraint $D_{\mu}L(v_{56}) = 0$, expressed through the covariant derivative $D_{\mu} \sim \partial_{\mu} + Y_{\mu}$ (as for the E_6 example in equation 11.33). In this way interaction terms similar in form to those introduced for \mathcal{L}_{int} in the Lagrangian approach for the Standard Model are identified. Arising from symmetry breaking over the base manifold M_4 the possible $\delta Y \leftrightarrow \delta \psi$ exchanges of field components are also constrained by the set of degenerate solutions under the same local external geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$.

All observed fermion states interact with at least one gauge boson via terms of $D_{\mu}L(v_{56}) = 0$, as applied for the electron self-energy interaction in figure 11.12(b) for example. Hence the external geometric structures relating to the $\psi(x)$ components will be shaped by the bare gauge fields such as $A_\mu(x)$ with which they interact. With the bare gauge fields subject to $G_{\mu\nu} = f(Y)$ from the isochronal Kaluza-Klein relation the physical fermion particle states will emerge through modifications to the geometry $G_{\mu\nu}(x)$ due to $\delta Y \leftrightarrow \delta \psi$ interactions. In turn the question of the form of $G_{\mu\nu} = f(\psi)$ for electron, muon and further particle states might be considered. This form of solution should also extrapolate to the non-relativistic limit, such as for the implied electron state linking S and A in figure 11.13(b) for example.

(3+4) Given also the non-trivial geometry $G_{\mu\nu} = f(\boldsymbol{v}_{56})$ from $L(\boldsymbol{v}_4) = h^2(x)$ variation the implications of further field interactions of the form $\delta v_4 \leftrightarrow \delta \psi$ under the constraint $L(\mathbf{v}_{56}) = 1$ will also contribute to the form of $G_{\mu\nu} = f(\psi)$. These interactions with the vector-Higgs field $v_4(x)$ are expected to relate to the origin of fermion masses, with the details giving rise to the mass difference between the electron and d-quark states for example. In order to investigate the mass differences between the three generations of fermions, such as between the electron and muon, a higher-dimensional form such as $L(v_{248}) = 1$ may be required. The equality of the empirically observed electric charge across the generations may relate to the role of 'Ward identities' in the QFT limit.

With the relation $G_{\mu\nu} = f(\mathbf{v}_{56})$ generalised for multiple solution field exchanges under the form $G_{\mu\nu} = f(Y, v_{56})$ essentially all matter $T_{\mu\nu} := G_{\mu\nu}$ is expected to be associated with quantum phenomena, with the variety material forms observed in the universe shaped according to the probabilistic nature of the underlying field composition. The relative probabilities of local solutions for $G_{\mu\nu} = f(Y, v_{56})$ are determined through a 'number of ways' statistical count of the underlying field redescriptions, essentially as for the determination of probabilities for classical systems. This leads to a unified approach to quantum and classical thermodynamic properties, which in particular will be significant for studying the evolution from $t = 0$ to the phase transition at $t = t_v$, as the stable value $L(\mathbf{v}_4) = h_0^2$ is attained in the very early universe, as described for figure 13.3. This may also mark an epoch of fermion production via $\delta v_4 \leftrightarrow \delta \psi$ exchanges as the properties of the Standard Model of particle physics emerge in the phase transition.

 $(4+1)$ While we have considered beginning with the classical geometric relations $G_{\mu\nu} =$ $f(\boldsymbol{v}_{56})$ or $G_{\mu\nu} = f(Y)$ more generally these two means of obtaining finite external curvature will be combined in a general solution for $G_{\mu\nu} = f(Y, \hat{v})$. In the full theory field interactions of the form $\delta Y \leftrightarrow \delta v_4$, resulting from the action of the corresponding gauge symmetry on the external components $v_4 \in TM_4$, will relate closely to the identification of gauge boson masses and the phenomena of electroweak symmetry breaking generally.

In principle the theory might rather begin with the full general form of $G_{\mu\nu}$ = $f(Y, \hat{v})$, fully incorporating quantum phenomena and completing the program described for fronts $(1+3)$ combined above, as will be required to fully account for both the large scale structure in cosmology and the phenomena observed in the HEP laboratory. While the pure 'bare' forms of the relations $G_{\mu\nu} = f(\mathbf{v}_{56})$ or $G_{\mu\nu} = f(Y)$ may not be found in nature, due to the possibility of underlying field interactions, each of these relations may play a role in an appropriate classical field limit.

Hence the aim from the developments in figure 15.1 is to generalise from (1) the geometric structure of gravitational and gauge fields deriving from the isochronal symmetry of $L(v_{56}) = 1$ to incorporate interactions with the field components of (2) v_{56} itself subject to the dynamic relation $D_{\mu}L(v_{56}) = 0$ derived from the action of the full symmetry of E_7 on $F(h_3\mathbb{O})$ broken over M_4 , taking into account the impact of (4) variation in the projected value of $|v_4| = h(x)$, to arrive at a general form of solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ over (3) a degeneracy of 'quantum' field redescriptions underlying

an external geometry with $G^{\mu\nu}{}_{;\mu} = 0$ everywhere, which itself provides one of the constraint equations 11.29.

Collectively progress on fronts (1) , (3) and (4) of figure 15.1 can be considered together under the ambition of accounting for the empirical properties of a quantum field theory *without* applying standard quantisation rules for the present theory. These three fronts all relate to the identification of a smooth geometry $G_{\mu\nu}(x)$ constructed in terms of fields extended on the spacetime manifold M_4 , the identification of which, as the background for perception in the world, itself motivates this construction. This area of research, guided by the analogy between QFT and condensed matter systems, might proceed based on a provisional assumption for the full symmetry of the full form $L(\hat{\mathbf{v}}) = 1$ such as the E₆ case.

In fact for this purpose a yet simpler, but non-trivial, model could be considered based on $\hat{G} = SL(3,\mathbb{C})$ as the full symmetry of time acting upon elements $\hat{\mathbf{v}} = \mathbf{v}_9 \in h_3\mathbb{C}$ such that $L(v_9) = det(v_9) = 1$ is invariant. This structure incorporates a subgroup action $SL(2,\mathbb{C}) \subset SL(3,\mathbb{C})$ on the subcomponents of $v_4 \equiv h_2 \in h_2\mathbb{C}$, identified with the external tangent space TM_4 , as described for equation 7.35 at the end of section 7.1. The structure of the resulting symmetry breaking to $SL(2,\mathbb{C}) \times U(1) \subset SL(3,\mathbb{C})$ over the base manifold M_4 may be sufficient to study a model accommodating both general relativity together with a form of quantum electrodynamics deriving from the internal $U(1)$ symmetry. On generalising from the complex space $\mathbb C$ to the octonions $\mathbb O$ the symmetry action SL(3, $\mathbb C$) is itself contained as a subgroup of SL(3, $\mathbb O$) \equiv E₆ as explicitly demonstrated by the generator composition of equations 8.32 and 8.33 in subsection 8.3.1. In this way the form $L(v_9) = 1$ naturally takes its place in the progression $L(v_4) \to L(v_9) \to L(v_{27}) \to L(v_{56})$ discussed in section 13.3.

Independently of combining the above three fronts, that is (1) , (3) and (4) , further progress may be made on the structure of front (2) itself which, although the subspace of vectors $v_4 \in h_2\mathbb{C}$ is associated with the external spacetime, considers the symmetry structure of $L(\hat{v}) = 1$ without explicitly projecting the components into fields over M_4 . This further study concerns, for example, the explicit identification of an internal $SU(2)_L \times U(1)_Y \subset E_7$ subgroup together with a determination of $\sin^2 \theta_W$ and the study of electroweak properties within the theory based on the form $L(v_{56}) = 1$. However the larger ambition for front (2) will be the identification of the full general form of temporal flow, involving for example an E_8 symmetry of the currently hypothetical form $L(v_{248}) = 1$. The progression of table table 9.1 and the known structure of equation 9.50 together with the general discussion of section 9.3 strongly hints towards the real form $E_{8(-24)}$ as a candidate to be sought for such a full symmetry.

A more thorough understanding of quantum phenomena in spacetime and a determination of the full form of $L(\hat{\mathbf{v}}) = 1$ are hence the two main branches to be pursued en route to the formation of a complete theory incorporating all four fronts of figure 15.1, with the aim to account both for cosmological observations and the properties of the Standard Model of particle physics through the structure of $L(\hat{v}) = 1$, and without introducing a Lagrangian or any other arbitrary postulates for any point of the theory.

15.2 Reconstructing HEP Phenomena

For contrast with the present theory the general recipe for constructing a standard field theory is summarised in the following three stages. This involves in particular employing a Lagrangian, such as equation 3.96 or as described in section 7.2 for the Standard Model, to introduce interactions into the theory in order to describe the phenomena observed in HEP experiments.

- (a) Together with the Lorentz group for the external spacetime symmetry, a gauge group is selected, generally motivated on empirical grounds, to describe the internal symmetry of the model. The field content of the theory, in terms of the field transformation properties as a choice of the representations of the symmetry groups, is also determined in order to comply with the findings of experiments.
- (b) A scalar Lagrangian as a function of the fields is written down, invariant under the symmetries of the theory, with various caveats on the general form of the terms – for example to ensure the renormalisability of the quantum version of the theory. The Lagrangian function is used in conjunction with the principle of extremal action to determine the equations of motion for the fields.
- (c) The classical theory can be quantised for example by introducing field operators $\phi(x)$, commutation relations and a Fock space of particle states such as $|p\rangle$ as reviewed in the opening of section 10.3. The framework of QFT is built upon a flat spacetime background as a given entity.

From the point of view taken here the introduction of a scalar Lagrangian function in item (b) above is conceptually a particularly poorly motivated aspect of the standard theory. The roots of the Lagrangian approach originate historically in the study of classical mechanics for non-relativistic material bodies, reproducing Newton's Laws of Motion in a more general framework. Later, further pragmatic progress and empirical success was achieved in generalising this framework to incorporate field theories and also to derive relativistic field equations in the Minkowski spacetime of special relativity. The Lagrangian approach is also employed for the quantised fields of QFT in a flat spacetime on the one hand, and in general relativity, with the geometric $R\sqrt{|g|}$ Lagrangian term based on the Ricci scalar R for example in equation 3.79, in a curved spacetime on the other hand.

However there is no underlying conceptual justification for the invention of such a scalar field, the integral of which over a set of spacetime coordinates should be stationary under field variations, either for a classical or quantum theory. In the QFT for the Standard Model it is the empirical observation of the effects of local gauge groups through their representations on apparent particle multiplets that guides the construction the Lagrangian, taylored to generate the desired equations of motion. That the Lagrangian framework should remain valid for a unified theory of quantum phenomena and gravitation is a further assumption built upon an uncertain foundation.

By contrast with the Lagrangian approach, in the present theory a fundamental scalar function which is not only stationary but constrained to a particular scalar value is readily identified, that is $L(\mathbf{v}_{56}) = 1$. Although general empirical features, such as the required rank of a unification group as described in section 7.3, serve as a useful guide for the study of E_6 and E_7 as a symmetry of time, here empirical details of the Standard Model are uncovered in the structure of the external and internal broken symmetry action on the components of the spaces h₃ \mathbb{O} and $F(\mathrm{h}_3\mathbb{O})$, as described in chapters 8 and 9. Further, the equations of motion for the fields on M_4 can be derived purely as a consequence of the constraints of the theory, which are summarised in equations 11.29. For example Maxwell's equation 11.26 and the Dirac equation 11.31 result from the degeneracy of field solutions subject to the constraints, as described in section 11.1. Hence in contrast to the recipe for a standard field theory listed above in (a) –(c), the necessary ingredients arise naturally in the present framework as listed below:

- (A) All the main symmetries considered must form a group or subgroup of a symmetry of time, that is of the equation $L(v) = 1$. The Lorentz group is motivated by its pseudo-Euclidean structure as required for external perception, while the internal gauge groups are identified in the breaking of the higher, richer, symmetry such as E_7 over the base manifold M_4 . The representations are already essentially determined since the Lorentz and $E₇$ groups are selected by their actions upon the vector spaces $h_2\mathbb{C}$ and $F(h_3\mathbb{O})$ respectively, with the broken internal gauge groups acting upon multiplets of $SL(2,\mathbb{C})^1 \subset E_7$ Weyl spinors.
- (B) Equations of motion are constrained by the fundamental requirement $L(\mathbf{v}_{56}) = 1$ which further implies $D_{\mu}L(v_{56})=0$, as listed in equations 11.29. Further constraints on the equations of motion for the fields are governed by the relation $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, consistent with the Bianchi identities for the external and internal symmetries. This structure over M_4 naturally arises as required to frame a world of physical perception, in a geometrical space and time, out of the general form of temporal flow. Field 'interactions' are implied at the outset in the form of the above expressions over the base manifold, in terms of gauge $Y(x)$ and spinor $\psi(x)$ fields for example, leading to expressions such as equation 11.33.
- (C) In the present theory the phenomena of quantisation correspond to the degeneracy of the multiple solutions implied in the expression $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, consistent with $L(v_{56}) = 1$, as has been summarised in the previous section. That is, the fields are intrinsically involved in creating the non-trivial geometry $G_{\mu\nu}(x)$ of the base manifold itself. It then remains to be described how the particle phenomena seen in HEP experiments, in particular the nature of the initial and final particle states, arise out of these field exchanges in spacetime.

The non-gravitational fields on M_4 derive from the symmetries and components of the 'extra dimensions' of temporal flow, in a manner analogous to the employment of the additional degrees of freedom in theories based on extra spacetime dimensions such as Kaluza-Klein theories. Here the equations of motion are simply equations for the variation of the mathematical structures which arise as projected onto the 4 dimensional base manifold and parametrised by the underlying 1-dimensional temporal flow. They are *not* equations of motion for some other body or entity introduced independently of time itself.

The field and particle content of the theory will be determined by the choice of the full and external forms of temporal flow, here taken to be $L(v_{56}) = 1$ and

 $L(\mathbf{v}_4) = h^2$ on M_4 with their respective symmetries of E₇ and SL $(2,\mathbb{C})^1$ (with the latter originally identified as a subgroup of E_6 as described for equation 8.6). The mathematical and conceptual limitations on the choice of these significant forms and the component normalisation such as h^2 , and hence the observed field and particle properties induced through the symmetry breaking, were considered in the section 13.3. There questions were raised concerning the uniqueness of the present theory and the extent to which it is constrained given, for example, the possibility of further higherdimensional forms of temporal flow.

Here, with E_7 taken to describe the symmetry of the full 56-dimensional form of temporal flow, for the complete theory the full set of broken $L(v_{56}) = 1$ and $D_{\mu}L(v_{56}) = 0$ terms may be written out. All empirical effects must then be consistent with these equations together with the local geometrical forms $G_{\mu\nu} = f(Y)$ and $G_{\mu\nu} = f(\mathbf{v}_{56})$, the latter of which augments the set in equation 11.29, as combined globally under the solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ together with the identity $G^{\mu\nu}_{;\mu} = 0$ framing the spacetime manifold. Hence the set of possible field couplings, as expressed through causal sequences of degenerate field redescriptions, must conform to this set of equations. These equations, essentially acting as selection rules, are listed in the left-hand column of table 15.1 alongside examples of possible terms and the associated field interactions or empirical effects in the remaining columns.

Equations	Terms	Field Interactions and Phenomena
$L(v_{56})=1$	$\sim vv\psi\psi$	Yukawa-type couplings for fermion masses
		involving vector-Higgs v_4 components
$D_{\mu}L(v_{56})=0$ $\sim vv\overline{\psi}Y\psi$		gauge-fermion interactions for internal forces
		also gauge- v_4 coupling for Z^0, W^{\pm} masses
$G_{\mu\nu} = f({\bf v}_{56})$	equation 13.4	significant for geometry of dark sector
		and evolution of the very early universe
$G_{\mu\nu}=f(Y)$	$\sim FF$	with $F = dY + \frac{1}{2}[Y, Y]$, equation 3.37, have
		gauge field cubic and quartic self-coupling
$G^{\mu\nu}_{;\mu} = 0$	$T^{\mu\nu}_{\ \ \mu}(Y,\hat{\boldsymbol{v}})=0$	conservation of energy-momentum and
		constraint on field equations of motion

Table 15.1: The set of constraints in the first column determine the field interactions and associated field equations of motion, in place of an imposed Lagrangian.

The interactions described in the right-hand column bare a close resemblance to those placed by hand in the Standard Model Lagrangian, however the corresponding field terms in table 15.1 arise naturally in the present theory. Collectively the constraints in table 15.1 expressed over the spacetime manifold M_4 replace the need to introduce a scalar Lagrangian function. With respect to local internal symmetry transformations all of the equations in table 15.1 are gauge invariant while they transform

covariantly under external Lorentz transformations as scalar, vector or tensor representations. This latter feature, as well as the fact that there are several equations, distinguishes this theory from the scalar Lagrangian approach, and indeed the present theory will need to be fully worked out independently of the standard framework.

Given a sufficient understanding of how field degeneracy in the present theory relates to quantum phenomena it may be possible to deduce effective Lagrangian terms from the constraints of the equations listed in table 15.1 and import these structures into the framework of a QFT employing a Lagrangian approach. This substitution of fields and interactions derived from the present theory into the standard procedure summarised in items (a), (b) and (c) above might be provisionally followed all the way through to standard QFT calculations such as cross-sections. However, the alternative approach, with the emphasis on a complete understanding of the present theory, would be much preferred in the long term, with the formalism of a QFT Lagrangian later identified in a suitable limit of the complete theory.

For the present theory the meaning of quantisation itself is to be found in the degeneracy of field solutions, without following a standard QFT approach such as attaching creation $a^{\dagger}(\boldsymbol{p})$ and annihilation $a(\boldsymbol{q})$ operators to the field components and applying canonical commutation rules. However in the process of calculation the field couplings arising from the equations in table 15.1 may be associated with vertex diagrams, as was described for a few cases in figure 11.3, as one part of the correspondence with Feynman rules described more completely in section 11.2. That is, while the present theory is constructed on a firm conceptual foundation, the empirical successes of QFT suggests that a complexification of a calculation and the employment of the mathematical tools of QFT, such as amplitudes and unitary evolution, might also be applied pragmatically here. Hence the optimal approach may be to straddle both perspectives – pursuing the development of the present theory while incorporating calculational tools from QFT.

Between the macroscopic structure of the external geometry $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ and the internal microscopic field interaction exchanges, consistent with the equation $D_{\mu}L(v_{56}) = 0$ for example, nested layers of multiple solutions will shape the physical manifestation of the theory in a way reminiscent of 'renormalisation' techniques in QFT. While the particle concept and HEP calculations may be motivated from within the present theory mathematical tools extracted and adapted from QFT will play an important role in the development of the complete theory and the establishment of a detailed comparison with empirical measurements.

Since the physical couplings and masses measured for HEP phenomena correspond to renormalised states it isn't expected that the full features of the Standard Model should be seen directly in the bare broken terms of E_7 on $F(h_3\mathbb{O})$ for example. In QFT the bare Standard Model Lagrangian, with the Higgs field added in a relatively unnatural way, does mimic the processes of HEP to some extent. For the present theory, intended as an underlying fundamental theory, the fact that a number of features placed by hand into the Standard Model Lagrangian have already been reproduced, as summarised in section 9.3, suggests that further specific details of empirical phenomena might be uncovered for the complete theory. These empirical details include in particular the 18 free parameters of the Standard Model as summarised in table 15.2.

SM Parameters	#	Origin in present theory
Fermion Masses	9	$\psi \leftrightarrow v_4$ coupling in $L(\hat{v}) = 1$ terms
		equations 8.76 and 9.48
Gauge Couplings	3	$\psi \leftrightarrow Y$ coupling in $D_{\mu}L(\hat{\boldsymbol{v}}) = 0$ terms
		equations 11.33 and 11.34
Higgs Potential	$\overline{2}$	$v_4 \in TM_4$ projected from full \hat{v}
		equation 8.72, with $ \mathbf{v}_4 = h_0$ stable
Quark Mixing CKM	4	mass and gauge couplings for 3 generations
		may require 'E ₈ on $L(v_{248}) = 1$ '

Table 15.2: The 18 parameters of the Standard Model and their correspondence in the present theory. All essentially originate as couplings implied in $L(\hat{v}) = 1$ as exemplified in the above equation references, including further parameters for the neutrino sector.

The QCD θ -parameter, introduced in equation 11.39 and which is consistent with zero empirically, is not included in the table since the corresponding field interaction terms do not arise in the present theory, as described in section 11.1. On the other hand the new structures presented in this paper may imply new kinds of interaction terms which do have empirical consequences. As well as identifying new processes the present theory may be tested through its ability to reproduce the details of known phenomena through the interactions listed in table 15.1.

As noted in that table, these include observations of the large scale structure in cosmology, which may relate to variation in the magnitude $|v_4|$ under $G_{\mu\nu} = f(v_{56})$. In addition to accounting for the Standard Model particle properties the complete theory would aim to provide a match for the cosmological data, including the density parameters Ω_B , Ω_D and Ω_{Λ} introduced in section 12.2, and the structure of the cosmic evolution generally. In particular the Lorentz scalar components α, β, n and N of $F(\text{h}_3\textcircled{1})$ in equation 9.46, which also transform trivially under the $SU(3)_c \times U(1)_Q \subset E_7$ gauge group while effectively acquiring mass through interactions with the vector-Higgs v_4 under the terms of $L(v_{56}) = 1$, may contribute to the dark sector in cosmology, as discussed in section 13.1.

Other known phenomena are not explicitly expressed in table 15.1. An example is provided by the CKM quark mixing parameters alluded to in table 15.2, which can be expressed explicitly in the Standard Model Lagrangian as described for equation 7.78. In the Standard Model the phenomena of CKM mixing arise for the three generations of quarks due to the mismatch between the Yukawa and gauge couplings, as described towards the end of section 7.2. While fermion masses and gauge couplings arise in the present theory as indicated in the upper half of table 15.2, the further necessary ingredient of three generations required for CKM mixing may require a further extension to for example an E_8 symmetry acting upon the hypothetical form $L(v_{248}) = 1$ as discussed in section 9.3.

Further parameters for three generations of neutrino masses and corresponding mixing phenomena are also needed as a known extension to the Standard Model, and are presumed to have a similar origin as described above for the quark sector in the present theory. As also suggested in section 9.3 the $SU(2)_L$ internal symmetry may play an essential role in distinguishing three generations of fermion states. It will also be required to identify neutrino and u-type quark states that transform as $SL(2,\mathbb{C})^1$ Weyl spinors and hence form $SU(2)_L$ doublet partners with charged lepton and d-type quark Weyl spinors respectively, which may also involve the identification of a full E_8 symmetry action on $L(v_{248}) = 1$.

The phenomena of electroweak symmetry breaking arise since the $SU(2)_L \times$ $U(1)_Y$ symmetry action itself also impinges on the components of the external vector-Higgs field $v_4 \in TM_4$. These interactions of the $SU(2)_L \times U(1)_Y$ gauge fields account for the massive nature of the Z^0 and W^{\pm} gauge bosons as described in subsection 8.3.3. That is the masses of all particles, fermions and gauge bosons, are here postulated to originate through field interactions with the components of $v_4(x) \in TM_4$ rather than with a fundamental scalar Higgs field. The large mass of the Z^0 and W^{\pm} bosons, of the same order as that of the empirically observed Higgs boson, will need to be understood in the context of the present theory. Indeed, the Higgs particle state itself will also need to be identified within this theory, echoing the empirical search for the Higgs which concluded successfully in 2012 at the Large Hadron Collider.

In the Standard Model the masses for the $Z⁰$, W^{\pm} and Higgs boson can be expressed in terms of gauge coupling and Higgs parameters of the left-hand side table 15.2 as described in section 7.2. The scalar Higgs field ϕ exhibits self-coupling, with terms such as $\phi^{\dagger} \phi$ and $(\phi^{\dagger} \phi)^2$ in the Lagrangian potential of equation 7.53, as contrived to break the symmetry of the vacuum. Within the new approach the scalar Higgs is provisionally identified with the magnitude $h(x)$ of the vector-Higgs $v_4(x)$ as projected onto TM_4 such that the relation $L(\mathbf{v}_4) = |\mathbf{v}_4|^2 = h^2$ is directly identified within the full form $L(\hat{v}) = 1$. Cubic and quartic field couplings, within the terms of $L(\mathbf{v}_{27}) = 1$ and $L(\mathbf{v}_{56}) = 1$ respectively, involving the components of \mathbf{v}_4 (coupled with combinations of the four scalar fields from the α , β , n and N components of $F(\mathbf{h}_3\mathbb{O})$ for example, as can be seen in equation 9.28), generate an effective potential $V(h,T)$, which may be dependent upon an apparent temperature T , as described in section 13.2. For the new approach yet further possible interactions will arise for higher-order field exchanges or a higher-dimensional full form of time. An initial unstable value of $h(t)$ has been considered for the extreme spacetime environment of the very early universe as discussed in section 13.2 in relation to inflationary theory, with the stable value $h(t) = h_0$ achieved at cosmic time $t = t_v$ marking a phase transition.

In chapters 6–9 of this paper the emphasis has been on the identification of known Standard Model properties from within the structure of the present theory, as summarised in the four bullet points and further discussion in section 9.3. The further ambition is to develop the theory to the point of making new empirical predictions that might be tested in existing and future laboratory experiments in particle physics as well as through observations in cosmology. Such theoretical predictions could be worked out concurrently with the running of the LHC in time to anticipate new effects that may appear in the data analysis. The predictions might also influence the design specifications for the future International Linear Collider.

For the present theory in addition to breaking the full symmetry of $L(\hat{\mathbf{v}}) = 1$ through the choice of the projected vector $v_4 \in TM_4$, with the stable value of $|v_4| = h_0$, symmetry breaking is also exhibited through the choice of particular components for the vector-Higgs v_4 in the local tangent space on the 4-dimensional manifold. This choice, represented in figure $13.3(c)$ with exaggerated fluctuations about the mean value, is analogous to the choice of component contributions for the Standard Model scalar Higgs vacuum value in equation 7.54. However, due to the difference in underlying structure, differences between the Standard Model Higgs phenomena and predictions of the present theory might be observable in the laboratory environment.

In considering the hypothetical structure of an E_8 action on a form $L(v_{248}) = 1$ the possibility of identifying the external spacetime vector $h_2 \equiv v_4 \in TM_4$ by fusing together a set of two or three right-handed spinors $\{\theta_{Y\mathcal{L}}, \phi_{Y\mathcal{L}}, \psi_{Y\mathcal{L}}\} \in \mathbb{C}^2$ under $SL(2,\mathbb{C})^1 \subset E_8$ was described alongside equation 9.52 in section 9.3. In turn there are a number of ways of identifying scalars from the components of the above three spinors, including the scalar magnitude $|v_4| = h$. This in principle opens up the possibility of identifying additional Higgs-like states, beyond the earlier possible scalar states that might be associated with the α , β , n and N components of $F(\mathrm{h}_3\mathbb{O})$ for the E₇ case. In addition to a direct search for such scalar states at the LHC an e^+e^- collider tuned to operate as a 'Higgs factory' might be sensitive to some of the observable consequences. Since the employment of the three spinors in this way corresponds to the empirical absence of a set of three generations of right-handed neutrinos, these structures may also impact upon the neutrino sector in a manner beyond the Standard Model.

Considered in general terms the extension to an E_8 symmetry itself also suggests the possibility of new gauge bosons beyond the Standard Model deriving from the extra $SU(2) \times U(1)$ that is appended to the familiar Standard Model symmetry in the rank-8 decomposition of equation 9.51 in section 9.3. However, the first objective is a mathematical one in identifying the predicted E_8 action on a quintic or higher order form $L(v_{248}) = 1$ itself, as highlighted in the previous section, and to assess the further extent to which known Standard Model properties might be recovered before considering additional empirical consequences in great detail. In the meantime the general manner in which particle states might be described from a conceptual point of view can be further elaborated as we now consider.

Under the assumption of a global flat spacetime in the laboratory the Lorentz symmetry may be augmented to the 10-parameter Poincaré group and particle states classified by their mass m and spin s (or helicity h for $m = 0$) according to the values of (m^2) and $(m^2)s(s+1)$ (for $m \neq 0$) they take respectively for the two Casimir operators $P_{\mu}P^{\mu}$ and $W_{\mu}W^{\mu}$, where W^{μ} is the Pauli-Lubanski vector. This applies to all particle states, including hadrons composed of quarks and the Higgs scalar which is presumed to be composed out of the collection of non-scalar field components of the vector-Higgs field v_4 in the present theory, as recalled above (with an analogous construction for technicolor models reviewed in subsection 8.3.3).

The four Weyl spinors of equation 8.13 identified in the components of θ^1 in section 8.1 relate to projected components of the larger Dirac spinors, which in turn can be identified within the components of $F(h_3\mathbb{O})$ in equation 9.46 under the action of $SL(2,\mathbb{C})^1 \subset E_7$. The fermions of the Standard Model are Dirac spinors, with differing properties for the projected left and right-handed Weyl spinor parts as reviewed in chapter 7. These different properties arise here through the necessarily asymmetric embedding of the vector-Higgs $v_4 \in TM_4$ with respect to the $\mathcal{X}, \mathcal{Y} \in H_3\mathbb{O}$ subspaces of $F(\text{h}_3\textcircled{1})$ and the resulting asymmetric action of an internal $SU(2)_L \subset E_7$ symmetry on these components in equation 9.46.

As described in section 8.2 alongside the $U(1)_Q$ symmetry of electromagnetism the broken E_6 symmetry on the space $h_3\mathbb{O}$ also includes $SU(3)_c$ as a pure internal symmetry, to be associated with massless gauge bosons, the gluons of QCD, in the Standard Model. In subsection 8.3.2 it was described how this $U(1)_O$ symmetry survives the breaking of an $SU(2)^2 \times U(1)^2 \subset E_6$ symmetry in a 'mock electroweak theory', as a provisional guide towards the identification of an $SU(2)_L \times U(1)_Y$ symmetry within E_7 or E_8 acting on the full temporal form $L(\hat{\boldsymbol{v}}) = 1$ in the complete theory.

Combining the above external properties under the Poincaré symmetry with full set of internal quantum numbers according to the transformation properties under $SU(3)_c \times SU(2)_L \times U(1)_Y$ will lead to a classification of particle states for a more thorough comparison with the Standard Model framework. That the enormous wealth of experimental data in high energy physics all points to a concise and simple table of a relatively small number of elementary particles, the fermions and bosons, as summarised in the Standard Model of particle physics with the 18 parameters of table 15.2 above, further motivates the aim to determine such particle properties in the present theory by taking a mathematical limit or approximation that mirrors the physical conditions to be found in such laboratory experiments.

In order to make contact with terrestrial laboratory experiments in HEP it will be necessary to proceed from the ideas presented in this paper through practical calculations for processes such as those in figures 10.1 and 11.13 and beyond to more general, and even novel, applications. In the particular case of figure 10.1 out of the general solutions $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ over M_4 the emergence of the initial e^+ and e^- states, an intermediate Z^0 boson and the final state particles will need to be described. Out of the annihilation of the particle and antiparticle in the centre-of-mass system in figure $11.13(a)$ a large number of field transmutations are possible, whether through a photon or a Z^0 boson state, allowing a large number of possible $\delta \psi \leftrightarrow \delta Y$ field exchanges and further states to be produced. These include the leptonic final state depicted in figure 11.13(a) as well as the hadronic jets seen in figure 10.1, resulting from quark pair production, together with all the particle states within the jets. These and further particle phenomena need to be accounted for within the structure and constraints of the present theory, as has been described in chapter 11.

One way of approaching the nature of particle states might be to consider the simple decay process $Z^0 \to e^+e^-$ via $\delta Y \leftrightarrow \delta \psi$ field exchanges resulting in the propagation of two independent fermions. This would also require an understanding of the Z^0 gauge boson mass in terms of $\delta Y \leftrightarrow \delta v_4$ interactions, incorporated into a solution $G_{\mu\nu} = f(Y)$ for a massive gauge field with $k^2 = m^2 \neq 0$, possessing a third polarisation state ε_3^{μ} $\frac{\mu}{3}$, and which satisfies equation 11.21. Similarly a Higgs decay process such as $H \to e^+e^-$ could be studied directly in terms of $\delta v_4 \leftrightarrow \delta \psi$ field exchanges, closely relating to the mechanism for fermion production during the phase transition at $t = t_v$ in the very early universe described in section 13.2.

On the other hand a purely QED process might be considered with the electromagnetic field $A^{\mu}(x)$ interacting with fermions. Since the photon is massless a possible approach would be to take a superposition, or sum, of electromagnetic fields, each in the form of equation 11.6, mimicking the situation of a two-photon collision and hence able to produce fermion pairs, as alluded to near the opening of section 11.3. The nature of a single intermediate photon state, effectively with $k^2 \neq 0$, in the centre-ofmass frame of an e^+e^- collider might also be considered. The production of fermions would be required to proceed through field redescriptions of the form $A^{\mu} \leftrightarrow \overline{\psi} \gamma^{\mu} \psi$, as initially discussed for figure 11.2, consistent with the constraint equations 11.29 under a geometric solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$.

A consistent normalisation of the fields will be required in field exchanges of the form $A^{\mu} \leftrightarrow \overline{\psi} \gamma^{\mu} \psi$, under the local geometry $G_{\mu\nu} = f(Y, \hat{\mathfrak{v}})$ with $G^{\mu\nu}{}_{;\mu} = 0$, linking external and intermediate field states. This will relate the $C^{\frac{1}{2}}$ coefficient and polarisation vectors $\varepsilon_r^{\mu}(\mathbf{k})$ for the electromagnetic field, as introduced in equation 11.6, to the spinor coefficients for a Dirac field $\psi(x)$. In the standard theory there are four independent solutions to the free Dirac equation labelled by the 4-component coefficients $u^{1,2}(p)$ and $v^{1,2}(p)$, with for example $\psi(x) = u^{1}(p)e^{-ip\cdot x}$ which may be normalised by kinematic factors of energy and mass (see for example [70] sections 3.3 and 5.2). Similarly in the present theory the coefficients of the electron field $\psi(x)$ for example will contain energy p^0 and mass m factors which will need to match those for the normalisation coefficients of the electromagnetic field A^{μ} in Fourier mode expansion exchanges between the fields under $G_{\mu\nu} = f(A, \psi)$. In all cases such 'kinematic factors' arise from 'numerical parameters' such as $p \in \mathbb{R}^4$ in the Fourier modes $e^{\pm ip\cdot x}$ themselves. As well as being mutually compatible these normalisation factors will ultimately translate into the appropriate dimensions for cross-section calculations, as described towards the end of section 11.2

In the environment of HEP experiments it is generally assumed that the spacetime is flat and a Minkowski coordinate system employed such that the external Lorentz connection has components $A^a{}_{b\mu}(x) = 0$, corresponding to a linear connection $\Gamma(x) = 0$ by equation 3.51. Transforming under the global Lorentz symmetry the components of the 4-component Dirac spinors $\psi(x)$ are normalised as alluded to above. The Lorentz connection $A^a_{b\mu}(x)$ acts on a Dirac spinor $\psi(x)$ through the associated spinor connection as a representation of the Lorentz symmetry. This structure can also be applied to the more general case of a curved spacetime, employing a spinor bundle over M_4 to express the dynamics of the Lorentz connection, with $A^a{}_{b\mu}(x) \neq 0$ in general, on the base manifold in relation to spinor fields. As described towards the end of the previous section, a starting point might be to develop a minimal model based on the full symmetry $G = SL(3, \mathbb{C})$ acting on $v_9 \in I_3 \mathbb{C}$ leaving the form $L(v_9) = 1$ invariant. For this model fermion states derive from the Weyl spinor ψ_L in equation 7.35 in interaction with an internal U(1)-valued gauge field, in principle describing a model for QED.

As well as classifying particle states such as gauge bosons and fermions in a representation space according to their transformation properties under the external and internal symmetry groups and their possible interactions, the structure of tangible physical particles in spacetime as detected in experiments can also be investigated. Physical particles evidently transfer energy and momentum, which can be described by the tensor $T_{\mu\nu}(x)$ and is presumed to be conserved in 4-dimensional spacetime. In the present theory energy-momentum is defined by the relation $T_{\mu\nu} := G_{\mu\nu}$ (within a practical normalisation factor of $-\kappa$), and hence the transfer of a finite amount of energy must necessarily be associated with $G_{\mu\nu} \neq 0$ and hence a non-flat spacetime, while the identity $G^{\mu\nu}_{;\mu} = 0$ also ensures energy-momentum conservation throughout. In turn this tangible spacetime form of a particle is expressed in terms of the underlying fields as a solution for $G_{\mu\nu} = f(Y, \hat{v})$. This smooth external geometry represents a macroscopic 'dressed' or 'renormalised' object constructed out of the underlying microscopic 'bare' field exchanges. Representing the electron beam in a HEP accelerator for example, observable properties associated with the energy-momentum for the electron field are carried by the tensor:

$$
T_{\mu\nu} := G_{\mu\nu} = f(Y, \hat{\mathbf{v}})
$$
\n^(15.1)

This is equation 5.32 of section 5.2, where a particular vector space representing the full temporal flow \hat{v} may be substituted in. The expression $G_{\mu\nu} = f(Y, v_{56})$ implies an underlying innumerably nested sequence of indistinguishable field descriptions under $G_{\mu\nu}(x)$. This geometry is entirely constructed out of field components derived from $L(v_{56}) = 1$ and the corresponding E₇ symmetry actions. However, in this theory, it seems quite possible that some components of the fundamental form $L(\mathbf{v}_{56}) = 1$ and the gauge fields may exist on M_4 without contributing to the geometry field $G_{\mu\nu}$. With $T_{\mu\nu} := G_{\mu\nu}$ this would imply that not all fields in spacetime have energy-momentum in the sense of $T_{\mu\nu}\neq 0$. This possibility was discussed in section 13.3 and compared to the case of gravity waves which, while associated with a geometry with $G_{\mu\nu} = 0$, carry energy via a finite Weyl curvature as described after equation 5.44 in section 5.2. Here we consider the measurable phenomena of HEP particle types and properties to be determined by the mutual constraints of equations 11.29 applied to the underlying fields and conveyed via energy-momentum in the form of the generalised expression of equation 15.1, as originally employed for the special case of the free electromagnetic field leading to figure 11.1 in section 11.1.

While a significant correlation between the structures of the present theory and calculations in QFT has been identified as described in sections 11.1 and 11.2, a key question remains regarding the precise conceptual form and mathematical expression of the nature of field quantisation. One major aspect concerns whether the projected field components themselves are effectively fragmented into discrete elements distributed over spacetime and related via $\delta Y(x)$ and $\delta \hat{v}(x)$ differences, as has typically been conceived as the theory has developed, with the components of the external gravitational field composing the only smooth and continuous functions on M_4 . An alternative view might see all fields smooth and continuous on M_4 , with discrete exchanges only in the local contributions to $G_{\mu\nu}(x)$ in equation 15.1 consistent with equations 11.29, considered as 'excitations' of the fields and giving rise to observations of apparent quantum phenomena. A full understanding of this description of such quantum phenomena in the context of the present theory is one of the two main branches to be pursued as summarised at the end of the previous section.

With *all* physical entities described by equation 15.1, subject to constraints such as $L(\hat{v}) = 1$, this includes solutions that incorporate the phenomena of apparent particle effects, as discussed in section 11.3. These solutions must describe the discrete emission and detection of the *same* conserved 4-momentum p with $p^2 = m^2$ and conserved charges, arising from the internal field constraints, giving the rather mechanical impression of an intermediate 'classical particle' or projectile of some form. As discussed in section 11.4 the 'particle tracks' that we construct by joining up detector hits, as depicted in figure 10.1 for example, reinforces this illusion of an independent particle-like entity pursuing a continuous trajectory.

One way to approach the nature of the actual physical structure underlying such particle-like phenomena is to begin by considering a general state of macroscopic matter described by $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, as represented by the 'bulky' geometry of figure 15.2(a), which might represent for example the matter content $T_{\mu\nu} := G_{\mu\nu}$ of ordinary 'table and chairs'. Subsequently a progression down to a more minimal field content underlying a solution of $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ can be considered, down to a stage that does not simply gradually fade away towards $G_{\mu\nu}(x) = 0$, but rather solutions for geometric structure emerge that take on the shape of a discrete set of topologies due to the discrete constraints on the underlying fields. In this case a somewhat 'tubular' structure might arise as the vacuum limit is approached, as represented in figure 15.2(b). These near vacuum conditions correspond for example to the environment created in HEP experiments as described near the opening of section 10.1.

Figure 15.2: Representations of 4-dimensional solutions for $G_{\mu\nu} = f(Y, \hat{v})$ for (a) the general case of ordinary extended matter (b) the discrete structure emerging as permitted by the underlying field constraints as the vacuum state is approached.

The pattern of inner lines in figure 15.2(b) are analogous to the contours on a map representing the altitude of a continuous physical terrain, with the geometry $G_{\mu\nu}(x)$ being perfectly smooth and continuous, as also for figure 15.2(a) and all other cases. Hence this geometry might more accurately be represented by a continuous shading. Considered as a full 4-dimensional spacetime solution the contour tubes in the near vacuum region in figure 15.2(b) connect and are continuous with macroscopic entities such as HEP accelerators and detectors, as represented by the outer structure in the same figure. The inner structure in figure $15.2(b)$, with time directed from left to right, might represent for example the overall particle interaction process $e^+e^- \rightarrow$ $\mu^+\mu^-$, via an intermediate γ or Z^0 state, which is typically pictured in terms of particle trajectories in 3-dimensional space as depicted in figure 11.13(a) and described in section 11.4.

While shaped by the discrete enveloping topology the spacetime geometry for such a process will also be modulated by a wave-like structure of a form similar to equation 11.12 and figure 11.1, corresponding to a particular 4-momentum transfer. As also described in section 11.1 the spacetime metric $g_{\mu\nu}(x)$ itself associated with this modulation is presumed to take a form similar to equation 11.13. In the overall solution of equation 15.1 for such a process the left-hand side $T_{\mu\nu} := G_{\mu\nu}$ of the equation describes both the kinematic properties of the interaction via the energymomentum tensor $T_{\mu\nu}(x)$ and the smooth external geometry $G_{\mu\nu}(x)$ as for general relativity. Through the right-hand side ' $f(Y, \hat{v})$ ' of the same equation all quantum properties are sown into this structure in the form of an underlying set of discrete field redescriptions of the form $\delta Y \leftrightarrow \delta \hat{v}$, subject to the constraints such as $L(\hat{v}) = 1$, which determine in turn the possible set of discrete particle types and interactions that can be observed in HEP experiments.

That is, while $G_{\mu\nu}(x)$ is perfectly smooth and continuous there is both a discrete set of apparent particle types and a discrete set of possible topologies, corresponding for example to n-particle final states, that may be obtained for the near-vacuum solutions. This structure hence provides a coherent conception of the nature and properties of particle states observed in the laboratory. For example a continuous range of conserved momenta is available for the apparent emission and detection of a fermion state within the discrete constraint $p^2 = m^2$, corresponding to an apparent particle mass m which arises from the underlying interactions between the particular fermion field $\psi(x)$ and the vector-Higgs field $v_4(x)$.

The metric $g_{\mu\nu}(x)$ for the external geometry depicted in figure 15.2(b) represents a particular solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ on the macroscopic scale of HEP experiments, similarly as the Schwarzschild metric of equation 5.49 represents a particular macroscopic solution on a much larger scale. Unlike the large scale case, for which the precise trajectory of planetary orbits and the deflection of light passing near the sun is observable, it is clearly not possible in practice to send 'test particles' through the laboratory environment of figure 15.2(b) in order to map out the spacetime curvature (although such a project can be readily conceived in terms of a thought experiment, as for that involving geodesic deviation due to the geometry of intense beams of light as described in section 11.4).

However, crucially for the present theory, this non-trivial external geometry with metric $g_{\mu\nu}(x)$ is a physical characteristic of a possible solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ and the test of this proposal, which will require all elements of the full theory, will rest on the ability to identify HEP processes which are actually observed and to predict new phenomena. This will involve both the determination of the internal quantum numbers of the apparent particle types, as implied in the underlying field structure $f(Y, \hat{v})$ for such a process, and in particular the apparent kinematic constraints on the 4-momentum p transferred, where with $p^2 = m^2$ and $T_{\mu\nu} := G_{\mu\nu}$ the invariant mass m provides a direct characterisation of the external geometry itself.

Within the field constraints more generally a range of topologies which are rather more complicated than that depicted in figure 15.2(b) will arise. For example the process recorded in figure 10.1 is identified as an $e^+e^- \to Z^0 \to b\bar{b}$ event in the analysis of [68]. Such a process typically involves 'particle tracks', as shown in the event picture, each of which apparently emanates from one a sequence of vertices, each of which in turn is associated with the Z^0 boson itself or a B or D hadron in a subsequent decay chain. With generally five such decay vertices for each such event mutually separated by typically a few millimetres, within the volume of the detector for which the closest devices are a few centimetres from the interaction point, the topology of the apparent particle-like structure described by the solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ will be relatively intricate for these processes.

Yet other forms of solutions for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ may appear less 'particle-like' as for the case of an e^- state apparently simultaneously 'passing through both slits' in the experiment depicted in figure 11.13(b). The overall geometry $G_{\mu\nu}(x)$ for the set-up of figure 11.13(b) for the case of a high intensity electron beam, with the full interference pattern clearly observed on the final screen, will be of a macroscopic form as described for figure $15.2(a)$ above. As the intensity is turned down, corresponding to a transition towards a near vacuum solution as exemplified in figure 15.2(b), an overall geometry will emerge incorporating the transfer of an apparent single e^- particle from the source S to the detector hit A in figure 11.13(b) in continuity with the structure of the apparatus of the double-slit experiment. The geometry of such a solution serves to emphasise the fact that a 'particle' should not be considered as a kind of localised entity in the form of an 'energy-knot' propagating in 3-dimensional space (see for example [82] pp.202–204), but rather as an apparent phenomenon associated with a particular kind of smooth extended 4-dimensional solution for $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ constructed over the underlying field possibilities. Similar 4-dimensional spacetime solutions will also incorporate the phenomena of quantum entanglement and EPR experiments as discussed in section 11.4. In many cases however a solution for $G_{\mu\nu} = f(Y, \hat{\mathbf{v}})$ will take a form consistent with the notion of a localised propagating particle-like entity.

Although in the present theory there are also no fundamental 'string-like' objects, there may be some relation to string theory (for which there are also no fundamental particle entities) in that diagrams with a similar topology to that of the inner structure in figure 15.2(b) also appear in relation to string theory calculations. Here however rather than describing the trajectory and interactions of a set of closed strings the tubular contours in figure 15.2(b) purely represent the structure of an extended 4-dimensional geometry. In string theory such a diagram correlates with the 'tree level' process as represented by the Feynman diagram of figure 10.3 for example, while for the present theory figure 15.2(b) represents the full physical process with arbitrarily nested field exchanges implied under the solution $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$. However, although the inner structure of figure 15.2(b) in relating to a process such as $e^+e^- \rightarrow \mu^+\mu^$ has a very different physical and conceptual meaning to analogous diagrams featuring in string theory, some of the mathematical properties of topological structures in 4-dimensional spacetime might be jointly applicable.

Rather than the phenomena of a discrete spectrum of particles being determined by the vibrations and tension of hypothetical strings, here such phenomena are generated by the possibility of actual underlying field redescriptions subject to the constraints of equations 11.29. As considered above a practical starting point may be to identify QED processes involving electron-photon interactions, such as with Bhabha or Compton scattering events, in this unified framework alongside general relativity. This study might begin with a model based on the full symmetry $SL(3,\mathbb{C})$ for the form $L(v_9) = 1$ before generalising to the octonion case with a full $SL(3, \mathbb{O}) \equiv E_6$ symmetry acting on the form $L(v_{27}) = 1$. The action of the internal $U(1)_Q \subset E_6$ symmetry generated by \dot{S}_l^1 on the spinor components of $Th_3\mathbb{O}$, as seen for example in equations 11.33 and 11.34 in the terms of the field constraint equation $D_{\mu}L(v_{27}) = 0$, gives rise to the phenomena of electrodynamics. The precise manner in which the factors of $|\dot{s}_f| = 1$ or $|\dot{s}_f| = \frac{1}{3}$ in this expression translate into the corresponding factor of three in charge ratio for physical renormalised particle states, as discussed for figure 11.5 in section 11.2 in the context of cross-section calculations, will need to be determined alongside the full understanding of the structure of quantum phenomena and particle states themselves.

The above QED phenomena will generalise for the complete internal symmetry identified in the breaking of the E_6 symmetry over the extended external M_4 manifold, and then further with the full symmetry of time identified as E_7 or even E_8 on the full form of temporal flow $L(\hat{v}) = 1$. The insight gained from the U(1)_Q case might then be extended for the remaining internal generators to identify further features of the Standard Model and beyond as they arise naturally out of the complete theory. It is likely that the full package will be required with all the features of figure 15.1 combined together, and the full set of possible fields and field interactions incorporated, in order to determine specific quantities such as the electron mass and the full set of Standard Model parameters as summarised in table 15.2, including the neutrino sector, generally.

In conclusion, the field and particle content of the present theory, in terms of figure 15.1, includes the external gravitational and internal gauge fields which arise from the symmetries of $L(\hat{\mathbf{v}}) = 1$ and are mutually related as described for 'front (1)', together with the fermion and 'vector-Higgs' fields identified from the $F(h_3\mathbb{O})$ components studied for 'front (2)'. Consistent with the gauge invariance of the constraint equations the non-gravitational fields mutually interact to form combinations under possible solutions $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$ for the world geometry on M_4 as described for 'front (3)', taking into account the intrinsic warping of the spacetime geometry due to variation in $|v_4(x)|$ and the role of the scalar field components as studied for 'front (4)'. In order to develop this theory further and establish full contact with the results of HEP experiments, as well as with empirical observations in cosmology and physical phenomena more generally, the four fronts of figure 15.1 will need to be further developed and combined as provisionally outlined in the previous section.

15.3 Concluding Remarks

While emphasising the possibilities for progressing outwards from the structure of figure 15.1 the present theory is based upon the multi-dimensional form of temporal flow $L(\hat{\boldsymbol{v}}) = 1$, at the centre of the figure, which in turn derives from the simple structure of one-dimensional progression in time as described in section 2.1. With both the familiar four dimensions of the extended spacetime manifold M_4 and the 'extra dimensions', which are associated with the properties of physical objects in spacetime, deriving from a single temporal dimension the question concerning the origin of time itself is inevitable. A naive further reduction down to 'zero dimensions' together with a contrived argument to generate one dimension is not considered here to be of any great value. On the other hand the observation that the arithmetic properties of multiple dimensions are implicit within the arithmetic structure of the real line R, as described in section 2.1, provides a natural and major motivation for the present theory. A second founding motivation for the entire theory is the apparent necessity for any and every subjective experience, including our observations of the physical world, to take place in time. This conception of the theory itself implies a subjective nature for the origin of time and leads to the conclusions described in chapter 14, and in particular

to the 'universal foundation' for the theory depicted in figure 14.5.

This overall structure can be considered as a *system* rather than just a *theory* $(in the usual sense of the word) - it is intended not merely to represent the world by$ a model, but rather it aims to describe the way the world actually is, and how it is possible for it to be. This is in a similar spirit that a biologist, for example, might describe the system of a living organism – although finding such a metaphor for the whole system is particularly problematic due to its unique and all-embracing nature.

It is a system founded upon general experience of living in the world as well as upon knowledge gained from the high energy physics laboratory together with cosmology and from scientific observations in general. Indeed all such experiments and observations are just a refined and specialised form of our experience in the world. While the primary aim has been to demonstrate a unified theory that can account for a wealth of scientific data, and thereby also provide a means of verification of the ideas, it has also been considered desirable to incorporate the nature of experience itself in the world. This leads to a unification not only of experimental findings but also of science as a whole with our experiences of the world in general. Hence although much of the presentation has involved scientific knowledge, from particle physics to cosmology, the overall conceptual scheme arrived at is that of a world which one can feel oneself to be immersed or engaged within while walking down the street.

While the physical laws and structures of the 4-dimensional world are carved out of the general flow of time, as filtered by the spacetime form of perception, the actual physical objects we encounter, such as complex organic life forms, are moulded to conform with the possibility of our actual existence in the world. The apparent stability of the perceived physical forms – from inter-galactic structures to the insect world on Earth – gives the illusion of a robust universe, independent of conscious life, constructed upon an independently existing material substratum, a notion upon which the early development of science also built its foundations. It is an illusion which continues to yield enormous practical advances in navigating our way around the physical world.

Both time and space are direct forms of subjective experience of mathematical structures in the world, through which the physical world itself is created and sustained as incorporated in node (4) of figure 14.5. Although a more rigorous mathematical description of all aspects of this structure is to be sought this does not imply that the system of the world is itself fundamentally a 'mathematical object'. Rather, as is the case in general, mathematics provides a precise and concise means of describing and elaborating both physical and abstract structures. It is conceivable though that there may be essential properties of complex entities in the physical world such as the structure of the human brain which cannot be transcribed into a mathematical language which is both precise and concise enough for an exhaustive and practical description. Such a physical entity is of course 'still there' even if it cannot be succinctly expressed in mathematical terms, in which case a mathematical approximation to nature might still be employed for practical purposes.

For the present theory mathematics offers a precise, quantitative language for the scientific study of the conceptual, organic interplay between the physical world and conscious observer as represented in figure 14.5. However, while there is considerable scope for further mathematical development of the theory the time cycle structure can be conceptually and logically coherent even if it may be humanly difficult to comprehend or develop a precise mathematical description of certain elements, such as for nodes (5) and (6) of figure 14.5, or if such an element does not directly correlate with a mathematical expression in a sense that we might recognise from familiar textbook maths. These elements of the theory may be correspondingly harder to both investigate in full detail as well as model in mathematical terms. Regardless of these practical difficulties the fact remains ultimately that we do 'see' the world through a one-dimensional progression in time (in a similar sense that we see some objects as 'green' as described towards the end of section 14.1). This continuous temporal progression is inseparably fused together with all subjective experience as a fundamental characteristic of all experiences.

Taking the 1-dimensional flow of time in node (1) of figure 14.5 to be modelled accurately by an interval of the mathematical real line R can itself be considered as a provisional assumption. This can be justified since experience of a moment of time has the very simple structure of a continuous one-dimensional progression which may be uniquely and unambiguously represented by the properties of the real line. This assumption may be further justified by empirical tests of the consequences of the theory derived, via nodes (2) and (3) , for node (4) of figure 14.5.

Again, further stepping around the cycle in this figure, the employment of tractable mathematical language may fall short of providing an accurate and unambiguous account of the full nature of the self-reflective structures R which are central to nodes (5) and (6) of figure 14.5. The use of the mathematical structures relating to Gödel's theorem and undecidable propositions G in section 14.1 marked a provisional attempt to model such a structure, although in a manner that seems far too simplistic.

However, further mathematical development of this aspect of the theory is both desirable and possible, with the aim of identifying a more precise description of the progression of self-reflective physical states, as crudely represented in figure 14.4, in mathematical terms. This may involve a degree of approximation based on a statistical approach to the phenomena of systems composed of many parts, by analogy for example with the thermodynamic properties of entropy. Even if such a mathematical structure remains somewhat elusive the conceptual ideas regarding the notion of subjective temporalisation might in principle be tested to some extent against empirical findings in the field of neuroscience. Some of the ideas presented might also be of relevance in the field of artificial intelligence (as initially discussed at the end of section 14.3) featuring for example the design of a device as a 4-dimensional entity in spacetime incorporating a structure of internal temporalisation – that is a machine not just programmed to do things in time but also capable of internally representing a potentially subjective temporal structure itself.

In contrast to these more speculative elements of the theory the full mathematical expression of the upper half of figure 14.5, beginning with the objective flow of time modelled by an interval of the one-dimensional real line $\mathbb R$ and leading via the multi-dimensional form of temporal flow $L(\hat{v}) = 1$ and its symmetries to the extended spacetime arena of the physical world as constructed through one of a myriad of solutions for the expression $G_{\mu\nu} = f(Y, \hat{\boldsymbol{v}})$, is in principle highly testable and has also been by far the main focus of the present theory. Amidst the resulting quantum phenomena the full theory can be applied to the observations of HEP experiments as modelled by the techniques of QFT and expressed in the form of the Standard Model of particle physics, and here arising from the E⁷ symmetry of a 56-dimensional form of time. The external theory of general relativity, describing gravitational phenomena, is here unified with the internal theories of gauge fields and particle physics through the projection of the form $L(v_{56}) = 1$ and breaking of its symmetry in the identification the spacetime manifold M_4 as an arena for perception in the world.

Through these ideas the present theory also incorporates the subjective way in which we experience an apparently classical world of Newtonian material objects. Although having its origins in the fundamental notion of progression in time and perception in space the theory has developed with large scale cosmology and the Standard Model of laboratory particle phenomena in mind, resting heavily upon knowledge accumulated by the experimental and theoretical communities over recent decades to draw out the system of the world presented in this paper. The theory is expected to be profusely testable in terms of determining the extent to which the known form of the physical world can be ascertained from the basic conceptual ideas of the theory in addition to making new predictions for as yet unobserved phenomena which might be discovered. Indeed the properties already deduced from the theory, in matching a number of features of the Standard Model mark a first success for the theory. This success is summarised in section 9.3 where further progress is proposed in seeking an E_8 symmetry of an appropriate form $L(v_{248}) = 1$ as a mathematical prediction of the theory.

The other principle area for study in the next stage of developing the theory is towards a more detailed understanding of the application of statistical methods and renormalisation techniques for the present theory in relation to QFT. The phenomena of 'running coupling' will be of relevance here and the extrapolation of the three gauge couplings from the laboratory energy scale may encounter 'new physics' in terms of new interactions or states identified in the theory on the way up to the GUT scale. Consistency with the unification of the gauge couplings hence will also provide a test of this theory. The Planck scale seems to be of no great significance for the present theory since gravity is not quantised.

Returning again to figure 15.1, with the theory developed from the notion of a multi-dimensional form of time $L(v) = 1$, front (1) has shown how a Kaluza-Klein related unification between gravitational and internal gauge fields can arise naturally out of an underlying isochronal symmetry, rather than an isometry, for a world perceived over a 4-dimensional spacetime manifold. The results presented for front (2) already establish a substantial connection with empirical data in the form of several basic features of the Standard Model. Within the same framework, generalised for multiple solutions, front (3) has described how the calculational tools of quantum field theory might be incorporated, again originating out of the basic principles of this new theory. In addition to accounting for small scale laboratory phenomena, culminating in the particle concept described for figure $15.2(b)$, the large scale structure of cosmology is also addressed in front (4), including the remote reaches of the very early universe, leading to the conception of the cosmos summarised in figure 14.8. While well defined areas of further development have been identified the progress made and properties uncovered in all directions, together with the simplicity inherent in the founding notion of the flow of time, add to the overall plausibility of the theory.

Bibliography

- [1] John C. Baez, 'The Octonions', Bull. Am. Math. Soc. 39, 145–205 (2002) [arXiv:math/0105155 [math-ra]].
- [2] Shoshichi Kobayashi and Katsumi Nomizu, 'Foundations of Differential Geometry', Volume I, Wiley-Interscience (1963), Wiley Classics Library (1996).
- [3] Yvonne Choquet-Bruhat and Cécile DeWitt-Morette, 'Analysis, Manifolds and Physics. Part I: Basics', revised edition, North-Holland (1982).
- [4] Marián Fecko, 'Differential Geometry and Lie Groups for Physicists', Cambridge University Press (2011).
- [5] John A. Peacock, 'Cosmological Physics', Cambridge University Press, (1999).
- [6] Charles W. Misner, Kip S. Thorne and John Archibald Wheeler, 'Gravitation', W. H. Freeman and Company (1973).
- [7] Carlo Rovelli, 'Quantum Gravity', Cambridge University Press (2004).
- [8] Ryoyu Utiyama, 'Invariant Theoretical Interpretation of Interaction', Phys. Rev. 101, 1597 (1956).
- [9] S.W. Hawking and G.F.R. Ellis 'The Large Scale Structure of Space-Time', Cambridge University Press (1973).
- [10] Michio Kaku, 'Quantum Field Theory: A Modern Introduction', Oxford University Press, (1993).
- [11] Theodor Kaluza, 'On the Problem of Unity in Physics', Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.), 966 (1921).
- [12] Oskar Klein, 'Quantum Theory and Five-Dimensional Relativity', Z. Phys. 37, 895 (1926).
- [13] Y. M. Cho, 'Higher-Dimensional Unifications of Gravitation and Gauge Theories', J. Math. Phys. 16 (10), 2029 (1975).
- [14] R. Kerner, 'Generalization of the Kaluza-Klein Theory for an Arbitrary Non-Abelian Gauge Group', Annales Poincaré Phys. Theor. 9 (2), 143–152 (1968).
- [15] L. N. Chang, K. I. Macrae and F. Mansouri, 'Geometrical Approach to Local Gauge and Supergauge Invariance: Local Gauge Theories and Supersymmetric Strings', Phys. Rev. D 13, 235 (1976).
- [16] C. A. Orzalesi, 'Multidimensional Unified Theories', Fortsch. Phys. 29, 413–440 (1981).
- [17] W. Kopczyński, 'Metric-Affine Unification of Gravity and Gauge Theories', Acta Phys. Polon. B 10, 365 (1979).
- [18] C. A. Orzalesi and M. Pauri, 'Spontaneous Compactification, Gauge Symmetry and the Vanishing of the Cosmological Constant', Phys. Lett. B 107 (3), 186 (1981). C. A. Orzalesi and M. Pauri, 'Geodesic Motion in Multidimensional Unified Gauge Theories', Nuovo Cim. B 68 (2), 193–202 (1982).
- [19] M. W. Kalinowski, 'Vanishing of the Cosmological Constant in Non-Abelian Kaluza-Klein Theories', Int. J. Theor. Phys. 22 (5), 385 (1983).
- [20] M. O. Katanaev, 'Linear Connection in Theories of Kaluza-Klein Type', Theor. Math. Phys. 56, 795–798 (1984) [Teor. Mat. Fiz. 56 (2), 246–250 (1983)].
- [21] P. S. Wesson and J. Ponce de Leon, 'Kaluza-Klein Equations, Einstein's Equations, and an Effective Energy-momentum Tensor', J. Math. Phys. 33 (11), 3883 (1992).
- [22] R. Percacci and S. Randjbar-Daemi, 'Kaluza-Klein Theories on Bundles With Homogeneous Fibers. 1', J. Math. Phys. 24 (4), 807–814 (1983).
- [23] J. F. Luciani, 'Space-Time Geometry and Symmetry Breaking', Nucl. Phys. B 135, 111–130 (1978). G. Domokos and S. Kövesi-Domokos, 'Gauge Fields on Coset Spaces', Nuovo Cim. A 44 (2), 318–330 (1978). M. A. Awada, 'Kaluza-Klein Theory over Coset Spaces', Phys. Lett. B 127 (6), 415–418 (1983). D. Bailin and A. Love, 'Kaluza-Klein Theories', Rept. Prog. Phys. 50, 1087–1170 (1987). M. J. Duff, 'Kaluza-Klein Theory in Perspective', in *Stockholm 1994, The Oskar Klein centenary* 22-35 (1994) [arXiv:hep-th/9410046].
- [24] C.-C. Chiang, S.-C. Lee, S.-L. Lou and G. Marmo, 'Curvature Tensor for Kaluza-Klein Theories with Homogeneous Fibers', Phys. Rev. D 32 (6), 1364–1368 (1985).
- [25] F. Mansouri and L. N. Chang, 'Gravitation as a Gauge Theory', Phys. Rev. D 13 (12), 3192–3200 (1976).
- [26] Roger Penrose, 'The Road to Reality (A Complete Guide to the Laws of the Universe)', Jonathan Cape (2004).
- [27] J. L. Synge, 'Relativity: The General Theory', North-Holland Publishing Company, Amsterdam (1966).
- [28] F. Mansouri, 'Superunified Theories Based on the Geometry of Local (Super-) Gauge Invariance', Phys. Rev. D 16 (8), 2456–2467 (1977).
- [29] E. Witten, 'Search for a Realistic Kaluza-Klein Theory', Nucl. Phys. B 186, 412 (1981).
- [30] M. J. Duff, B. E. W. Nilsson and C. N. Pope, 'Kaluza-Klein Supergravity', Phys. Rept. 130 (1&2), 1–142 (1986).
- [31] J. Ponce de Leon, 'The Effective Energy-Momentum Tensor in Kaluza-Klein Gravity with Large Extra Dimensions and Off-Diagonal Metrics', Int. J. Mod. Phys. D 11, 1355–1380 (2002) [arXiv:gr-qc/0105120].
- [32] A. Gamba, 'Peculiarities of the Eight-Dimensional Space', J. Math. Phys. 8 (4), 775–781 (1967).
- [33] Jörg Schray and Corinne A. Manogue, 'Octonionic Representations of Clifford Algebras and Triality', Found. Phys. 26, 17 (1996) [arXiv:hep-th/9407179].
- [34] Claude Chevalley and Richard D. Schafer, 'The Exceptional Simple Lie algebras F⁴ and E6', Proc. Nat. Acad. Sci. USA 36, 137–141 (1950).
- [35] Hans Freudenthal, 'Lie Groups in the Foundations of Geometry', Adv. Math. 1, 145–190 (1964).
- [36] A. Sudbery, 'Division Algebras, (Pseudo)Orthogonal Groups and Spinors', J. Phys. A: Math. Gen. 17, 939–955 (1984).
- [37] Corinne A. Manogue and Jörg Schray, 'Finite Lorentz Transformations, Automorphisms, and Division Algebras', J. Math. Phys. 34, 3746–3767 (1993) [arXiv:hepth/9302044].
- [38] Aaron D. Wangberg, 'The Structure of E6', PhD. Thesis (Advisor: Tevian Dray), Oregon State University (2007) arXiv:0711.3447 [math.RA], with 'The Multiplication Table of E_6 ', i.e. the commutation table for sl(3, $\circled{0}$), available at http://course1.winona.edu/awangberg/E6/E6_color_commutation_poster.pdf
- [39] Corinne A. Manogue and Tevian Dray, 'Octonions, E6, and Particle Physics', J. Phys. Conf. Ser. 254, 012005 (2010) [arXiv:0911.2253v2 [math.RA]].
- [40] Tevian Dray and Corinne A. Manogue, 'Octonionic Cayley Spinors and E6', arXiv:0911.2255v2 [math.RA] (2010).
- [41] Aaron Wangberg and Tevian Dray, 'E₆, the Group: The Structure of $SL(3, \mathbb{O})$ ', J. Algebra Appl. 14 (6), 1550091 (2015) [arXiv:1212.3182 [math.RA]].
- [42] T. Teubner, 'The Standard Model', Proceedings of the RAL School for Experimental High Energy Physics Students, Oxford, September (2009). www.ppd.stfc.ac.uk/PPD/resources/pdf/StandardModel09.pdf
- [43] Howard Georgi and S. L. Glashow, 'Unity of All Elementary-Particle Forces', Phys. Rev. Lett. 32 (8), 438–441 (1974).
- [44] K. A. Olive et al. (Particle Data Group Collaboration), 'Review of Particle Physics', Chin. Phys. C 38 (9), 090001 (2014).
- [45] Scott Willenbrock, 'Symmetries of the Standard Model', arXiv:hep-ph/0410370 (2008).
- [46] P. Sikivie, L. Susskind, M. B. Voloshin and V. I. Zakharov, 'Isospin Breaking in Technicolor Models', Nucl. Phys. B 173, 189–207 (1980).
- [47] Katsusada Morita, 'Algebraic Gauge Theory of Quarks and Leptons', Prog. Theor. Phys. 68 (6), 2159–2175 (1982).
- [48] Geoffery Dixon, 'Division Algebras, (1,9)-Space-time, Matter-antimatter Mixing', arXiv:hep-th/9303039 (1993).
- [49] Feza Gürsey and Chia-Hsiung Tze, 'On the Role of Division, Jordan, and Related Algebras in Particle Physics', World Scientific, Singapore (1996).
- [50] Tevian Dray and Corinne A. Manogue, 'Quaternionic Spin', in *Ablamowicz, R. (ed.), Fauser, B. (ed.): Clifford algebras and their applications in mathematical physics, vol. 1* 21-38 (1999) [arXiv:hep-th/9910010].
- [51] Moshe Carmeli, 'SL(2, C) Invariance and the Gravitational Field', Annals Phys. 71, 603–610 (1972).
- [52] M. Carmeli and S. Malin, 'Reformulation of General Relativity as a Gauge Theory', Annals Phys. 103, 208–232 (1977).
- [53] F. G¨ursey, P. Ramond and P. Sikivie, 'A Universal Gauge Theory Model based on E_6 ', Phys. Lett. B 60 (2), 177–180 (1976).
- [54] Howard Georgi, 'Lie Algebras in Particle Physics', Perseus Books (1999).
- [55] M. Günaydin and F. Gürsey, 'Quark Statistics and Octonions', Phys. Rev. D 9, 3387–3391 (1974).
- [56] Katsusada Morita, 'Gauge Theories over Quaternions and Weinberg-Salam Theory', Prog. Theor. Phys. 65 (6), 2071–2074 (1981).
- [57] L. Susskind, 'Dynamics of Spontaneous Symmetry Breaking in the Weinberg-Salam Theory', Phys. Rev. D 20, 2619 (1979) [SLAC-PUB-2142].
- [58] E. Farhi and L. Susskind, 'Technicolour', Phys. Rept. 74 (3), 277–321 (1981).
- [59] S. Krutelevich, 'Jordan Algebras, Exceptional Groups, and Higher Composition Laws', arXiv:math/0411104 [math.NT] (2004).
- [60] L. Borsten, D. Dahanayake, M. J. Duff, H. Ebrahim and W. Rubens, 'Black Holes, Qubits and Octonions', Phys. Rept. 471, 113–219 (2009) [arXiv:0809.4685 [hep-th]].
- [61] Michael Rios, 'Jordan C^{*}-Algebras and Supergravity', arXiv:1005.3514 [hep-th] $(2010).$
- [62] Tevian Dray, Corinne A. Manogue, Robert A. Wilson, 'A Symplectic Representation of E7', arXiv:1311.0341 [math.RA] (2013).
- [63] J. Distler and S. Garibaldi, 'There is no "Theory of Everything" inside E8', Commun. Math. Phys. 298, 419 (2010) [arXiv:0905.2658 [math.RT]]. A. Garrett Lisi, 'An Explicit Embedding of Gravity and the Standard Model in E8', arXiv:1006.4908 [gr-qc] (2010).
- [64] Michael Aschbacher, 'Some Multilinear Forms with Large Isometry Groups', Geometriae Dedicata 25 (1–3), 417–465 (1988).
- [65] Martin Cederwall and Jakob Palmkvist, 'The Octic E_8 Invariant', J. Math. Phys. 48, 073505 (2007) [arXiv:hep-th/0702024].
- [66] Skip Garibaldi and Robert M. Guralnick, 'Simple Groups Stabilizing Polynomials', Forum of Mathematics, Pi 3, e3 (2015) [arXiv:1309.6611v3 [math.GR]].
- [67] P. Ramond, 'Algebraic Dreams', arXiv:hep-th/0112261 (2001).
- [68] K. Abe *et al.* (SLD Collaboration), 'Measurement of the branching ratios of the Z^0 into heavy quarks', Phys. Rev. D 71, 112004 (2005) [SLAC-PUB-9941] [arXiv:hep $ex/0503005$.
- [69] SLD Collaboration, http://www-sld.slac.stanford.edu/sldwww/sld.html
- [70] Michael E. Peskin and Daniel V. Schroeder, 'An Introduction to Quantum Field Theory', Westview Press (1995).
- [71] Franz Mandl and Graham Shaw, 'Quantum Field Theory', Wiley (1999), second edition, Wiley (2010).
- [72] Martinus J. G. Veltman, 'Diagrammatica: The Path to Feynman Rules', Cambridge Lect. Notes Phys. 4, 1 (1994).
- [73] J. V. Narlikar, 'An Introduction to Cosmology', Cambridge University Press (2002).
- [74] H. P. Robertson and Thomas W. Noonan, 'Relativity and Cosmology', W.B. Saunders Company (1968).
- [75] J. Foster and J. D. Nightingale, 'A Short Course in General Relativity', Longman Scientific & Technical (1986).
- [76] P. D. Mannheim, 'Alternatives to Dark Matter and Dark Energy', Prog. Part. Nucl. Phys. 56, 340 (2006) [arXiv:astro-ph/0505266].
- [77] R. K. Nesbet, 'Conformal Gravity: Dark Matter and Dark Energy', Entropy 15, 162 (2013) [arXiv:1208.4972 [physics.gen-ph]].
- [78] J. García-Bellido, J. Rubio, M. Shaposhnikov and D. Zenhäusern, 'Higgs-Dilaton Cosmology: From the Early to the Late Universe', Phys. Rev. D 84, 123504 (2011) [arXiv:1107.2163 [hep-ph]].
- [79] F. Bezrukov, 'The Higgs Field as an Inflaton', Class. Quant. Grav. 30, 214001 (2013) [arXiv:1307.0708 [hep-ph]].
- [80] P. Brax and A. C. Davis, 'Conformal Inflation Coupled to Matter', JCAP 1405, 019 (2014) [arXiv:1401.7281 [astro-ph.CO]].
- [81] E. C. Zeeman, 'Causality Implies the Lorentz Group', J. Math. Phys. 5, 490–493 (1964).
- [82] Hermann Weyl, 'Space Time Matter', translated by Henry L. Brose, Dover Publications (1952), (fourth edition, first published in 1922).
- [83] Kurt Gödel, 'Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I', Monatshefte für Mathematik und Physik 38, 173–198 (1931).
- [84] Ernest Nagel and James R. Newman, 'Gödel's Proof', Routledge & Kegan Paul Ltd (1958), New York University Press (2008).
- [85] Douglas R. Hofstadter, 'G¨odel, Escher, Bach: an Eternal Golden Braid', Basic Books (1999) (first published in 1979).