

Enhanced Information Exclusion Relations

Yunlong Xiao^{1, 2}, Naihuan Jing^{3, 1*}, and Xianqing Li-Jost²

¹School of Mathematics, South China University of Technology, Guangzhou 510640, China

²Max Planck Institute for Mathematics in the Sciences, Leipzig 04103, Germany

³Department of Mathematics, North Carolina State University, Raleigh, NC 27695, USA

*Corresponding author: jing@ncsu.edu

ABSTRACT

In Hall's reformulation of the uncertainty principle, the entropic uncertainty relation occupies a core position and provides the first nontrivial bound for the information exclusion principle. Based upon recent developments on the uncertainty relation, we present new bounds for the information exclusion relation using majorization theory and combinatoric techniques, which reveal further characteristic properties of the overlap matrix between the measurements.

Introduction

Mutual information is a measurement of correlations and plays a central role in communication theory¹⁻³. While the entropy describes uncertainties of measurements⁴⁻⁸, mutual information quantifies bits of gained information. Furthermore, information is a more natural quantifier than entropy except in applications like transmission over quantum channels⁹. The sum of information corresponding to measurements of position and momentum is bounded by the quantity $\log 2\Delta X\Delta P_X/\hbar$ for a quantum system with uncertainties for complementary observables ΔX and ΔP_X , and this is equivalent to the Heisenberg uncertainty principle¹⁰. Both the uncertainty relation and information exclusion relation¹¹⁻¹³ have been used to study the complementarity of observables such as position and momentum. The standard deviation has also been employed to quantify uncertainties, and it has been recognized later that the entropy seems more suitable in studying certain aspects of uncertainties.

As one of the well-known entropic uncertainty relations, Maassen and Uffink's formulation⁸ states that

$$H(M_1) + H(M_2) \geq -\log c_{max}, \quad (1)$$

where $H(M_k) = H(M_k, \rho) = -\sum_j p_j^k \log_2 p_j^k$ with $p_j^k = \langle u_j^k | \rho | u_j^k \rangle$ ($k = 1, 2; j = 1, 2, \dots, d$) for a given density matrix ρ of dimension d , and $c_{max} = \max_{i_1, i_2} c(u_{i_1}^1, u_{i_2}^2)$, and $c(u_{i_1}^1, u_{i_2}^2) = |\langle u_{i_1}^1 | u_{i_2}^2 \rangle|^2$ for two orthonormal bases $M_1 = \{|u_{i_1}^1\rangle\}$ and $M_2 = \{|u_{i_2}^2\rangle\}$ of d -dimensional Hilbert space \mathcal{H} .

Hall¹¹ generalized Eq.(1) to give the first bound of the *Information Exclusion Relation* on accessible information about a quantum system represented by an ensemble of states. Let M_1 and M_2 be as above on system A , and let B be another classical register (which may be related to A), then

$$I(M_1 : B) + I(M_2 : B) \leq r_H, \quad (2)$$

where $r_H = \log_2(d^2 c_{max})$ and $I(M_i : B) = H(M_i) - H(M_i|B)$ is the *mutual information*¹⁴ corresponding to the measurement M_i on system A . Here $H(M_i|B)$ is the conditional entropy relative to the subsystem B . Moreover, if system B is quantum memory, then $H(M_i|B) = H(\rho_{M_i B}) - H(\rho_B)$ with $\rho_{M_i B} = (\mathcal{M}_i \otimes I)(\rho_{AB})$, while $\mathcal{M}_i(\cdot) = \sum_{k_i} |u_{k_i}^i\rangle \langle u_{k_i}^i | (\cdot) | u_{k_i}^i\rangle \langle u_{k_i}^i |$. Eq. (2) depicts that it is impossible to probe the register B to reach complete information about observables M_1 and M_2 if the maximal overlap c_{max} between measurements is small. Unlike the entropic uncertainty relations, the bound r_H is far from being tight. Grudka *et al.*¹⁵ conjectured a stronger information exclusion relation based on numerical evidence (proved analytically in some special cases)

$$I(M_1 : B) + I(M_2 : B) \leq r_G, \quad (3)$$

where $r_G = \log_2 \left(d \cdot \left[\sum_{d \text{ largest}} c(u_{i_1}^1, u_{i_2}^2) \right] \right)$. As the sum runs over the d largest $c(u_{i_1}^1, u_{i_2}^2)$, we get $r_G \leq r_H$, so Eq. (3) is an improvement of Eq. (2). Recently Coles and Piani¹⁶ obtained a new information exclusion relation stronger than Eq. (3), which can also be strengthened to the case of quantum memory¹⁷

$$I(M_1 : B) + I(M_2 : B) \leq r_{CP} + H(A|B), \quad (4)$$

where $r_{CP} = \min\{r_{CP}(M_1, M_2), r_{CP}(M_2, M_1)\}$, $r_{CP}(M_1, M_2) = \log_2 \left(d \sum_{i_1, i_2} \max c(u_{i_1}^1, u_{i_2}^2) \right)$, and $H(A|B) = H(\rho_{AB}) - H(\rho_B)$ is the conditional von Neumann entropy with $H(\rho) = -\text{Tr}(\rho \log_2 \rho)$ the von Neumann entropy, while ρ_B represents the reduced state of the quantum state ρ_{AB} on subsystem B . It is clear that $r_{CP} \leq r_G$.

As pointed out in Ref.¹¹, the general information exclusion principle should have the form

$$\sum_{m=1}^N I(M_m : B) \leq r(M_1, M_2, \dots, M_N, B), \quad (5)$$

for observables M_1, M_2, \dots, M_N , where $r(M_1, M_2, \dots, M_N, B)$ is a nontrivial quantum bound. Such a quantum bound is recently given by Zhang *et al.*¹⁸ for the information exclusion principle of multi-measurements in the presence of quantum memory. However, almost all available bounds are not tight even for the case of two observables.

Our goal in this paper is to give a general approach for the information and exclusion principle using new bounds for two and multiple observables of quantum systems of any finite dimension by generalizing Coles-Piani's uncertainty relation and using majorization techniques. In particular, all our results can be reduced to the case without the presence of quantum memory.

The close relationship between the information exclusion relation and the uncertainty principle has promoted mutual developments. In the applications of the uncertainty relation to the former, there have been usually two available methods: either through subtraction of the uncertainty relation in the presence of quantum memory or utilizing the concavity property of the entropy together with combinatorial techniques or certain symmetry. Our second goal in this work is to analyze these two methods and in particular, we will show that the second method together with a special combinatorial scheme enables us to find tighter bounds for the information exclusion principle. The underlined reason for effectiveness is due to the special composition of the mutual information. We will take full advantage of this phenomenon and apply a distinguished symmetry of cyclic permutations to derive new bounds, which would have been difficult to obtain without consideration of mutual information.

We also remark that the recent result¹⁹ for the sum of entropies is valid in the absence of quantum side information and cannot be extended to the cases with quantum memory by simply adding the conditional entropy between the measured particle and quantum memory. To resolve this difficulty, we use a different method in this paper to generalize the results of Ref.¹⁹ in Lemma 1 and Theorem 2 to allow for quantum memory.

Results

We first consider the information exclusion principle for two observables, and then generalize it to multi-observable cases. After that we will show that our information exclusion relation gives a tighter bound, and the bound not only involves the d largest $c(u_{i_1}^1, u_{i_2}^2)$ but also contains all the overlaps $c(u_{i_1}^1, u_{i_2}^2)$ between bases of measurements.

We start with a qubit system to show our idea. The bound offered by Coles and Piani for two measurements does not improve the previous bounds for qubit systems. To see these, set $c_{i_1 i_2} = c(u_{i_1}^1, u_{i_2}^2)$ for brevity, then the unitarity of overlaps between measurements implies that $c_{11} + c_{12} = 1$, $c_{11} + c_{21} = 1$, $c_{21} + c_{22} = 1$ and $c_{12} + c_{21} = 1$. Assuming $c_{11} > c_{12}$, then $c_{11} = c_{22} > c_{12} = c_{21}$, thus

$$\begin{aligned} r_H &= \log_2(d^2 c_{max}) = \log_2(4c_{11}), \\ r_G &= \log_2\left(d \sum_{d \text{ largest}} c(u_{i_1}^1, u_{i_2}^2)\right) = \log_2(2(c_{11} + c_{22})), \\ r_{CP} &= \min\{r_{CP}(M_1, M_2), r_{CP}(M_2, M_1)\} = \log_2(2(c_{11} + c_{22})), \end{aligned} \quad (6)$$

hence we get $r_H = r_G = r_{CP} = \log_2(4c_{11})$ which says that the bounds of Hall, Grudka *et al.*, and Coles and Piani coincide with each other in this case.

Our first result already strengthens the bound in this case. Recall the implicit bound from the tensor-product majorization relation²⁰⁻²² is of the form

$$H(M_1|B) + H(M_2|B) \geq -\frac{1}{2} \omega \mathfrak{B} + H(A) - 2H(B), \quad (7)$$

where the vectors $\mathfrak{B} = (\log_2(\omega \cdot \mathfrak{A}_{i_1 i_2}))^\downarrow$ and $\mathfrak{A}_{i_1 i_2} = (c(u_{i_1}^1, u_{i_1}^2) c(u_{i_2}^1, u_{i_2}^2))_{ij}^\downarrow$ are of size d^2 . The symbol \downarrow means re-arranging the components in descending order. The majorization vector bound ω for probability tensor distributions $(p_{i_1}^1 p_{i_2}^2)_{i_1 i_2}$ of state ρ is the d^2 -dimensional vector $\omega = (\Omega_1, \Omega_1 - \Omega_2, \dots, \Omega_d - \Omega_{d-1}, 0, \dots, 0)$, where

$$\Omega_k = \max_{\rho} \sum_{|\{(i_1, i_2)\}|=k} p_{i_1}^1 p_{i_2}^2.$$




Figure 1. First comparison with Hall's bound. The upper orange curve (our bound $2 + \frac{1}{2}\omega\mathfrak{B}$) is tighter than the lower blue one (Hall's bound r_H) almost everywhere.




Figure 2. First comparison with Hall's bound. The difference $r_H - 2 - \frac{1}{2}\omega\mathfrak{B}$ of our bound from Hall's bound r_H for $a \in [0.5, 1]$ is shown.

The bound means that

$$(p_{i_1}^1 p_{i_2}^2)_{i_1 i_2} \prec \omega,$$

for any density matrix ρ and \prec is defined by comparing the corresponding partial sums of the decreasingly rearranged vectors. Therefore ω only depends on $c_{i_1 i_2}$ ²⁰. We remark that the quantity $H(A) - 2H(B)$ assumes a similar role as that of $H(A|B)$, which will be clarified in Theorem 2. As for more general case of N measurements, this quantity is replaced by $(N-1)H(A) - NH(B)$ in the place of $NH(A|B)$. A proof of this relation will be given in the section of Methods. The following is our first improved information exclusion relation in a new form.

Theorem 1. For any bipartite state ρ_{AB} , let M_1 and M_2 be two measurements on system A , and B the quantum memory correlated to A , then

$$I(M_1 : B) + I(M_2 : B) \leq 2 + \frac{1}{2}\omega\mathfrak{B} + 2H(B) - H(A), \quad (8)$$

where ω is the majorization bound and \mathfrak{B} is defined in the paragraph under Eq. (7).

See Methods for a proof of Theorem 1.

Eq. (8) gives an implicit bound for the information exclusion relation, and it is tighter than $\log_2(4c_{max}) + 2H(B) - H(A)$ as our bound not only involves the maximal overlap between M_1 and M_2 , but also contains the second largest element based on the construction of the universal uncertainty relation ω ^{21,22}. Majorization approach^{21,22} has been widely used in improving the lower bound of entropic uncertainty relation. The application in the information exclusion relation offers a new aspect of the majorization method. The new lower bound not only can be used for arbitrary nonnegative Schur-concave function²³ such as Rényi entropy and Tsallis entropy²⁴, but also provides insights to the relation among all the overlaps between measurements, which explains why it offers a better bound for both entropic uncertainty relations and information exclusion relations. We also remark that the new bound is still weaker than the one based on the optimal entropic uncertainty relation for qubits²⁵.




Figure 3. Comparison of our bound with that of Coles and Piani. Our bound $2 + \frac{1}{2}\omega\mathfrak{B}$ (lower in green) is better than Coles-Piani's bound r_{CP} (upper in purple) everywhere.

As an example, we consider the measurements $M_1 = \{(1, 0), (0, 1)\}$ and $M_2 = \{(\sqrt{a}, e^{i\phi}\sqrt{1-a}), (\sqrt{1-a}, -e^{i\phi}\sqrt{a})\}$. Our bound and $\log_2 4c_{max}$ for $\phi = \pi/2$ with respect to a are shown in FIG. 1.

FIG. 1 shows that our bound for qubit is better than the previous bounds $r_H = r_G = r_{CP}$ almost everywhere. Using symmetry we only consider a in $[\frac{1}{2}, 1]$. The common term $2H(B) - H(A)$ is omitted in the comparison. Further analysis of the bounds is given in FIG. 2.

Theorem 1 holds for any bipartite system and can be used for arbitrary two measurements M_i ($i = 1, 2$). For example, consider the qutrit state and a family of unitary matrices $U(\theta) = M(\theta)O_3M(\theta)^\dagger$ ^{16,20} where

$$M(\theta) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & \sin \theta \\ 0 & -\sin \theta & \cos \theta \end{pmatrix},$$

$$O_3 = \frac{1}{\sqrt{6}} \begin{pmatrix} \sqrt{2} & \sqrt{2} & \sqrt{2} \\ \sqrt{3} & 0 & -\sqrt{3} \\ 1 & -2 & 1 \end{pmatrix}. \quad (9)$$

Upon the same matrix $U(\theta)$, comparison between our bound $2 + \frac{1}{2}\omega\mathfrak{B}$ and Coles-Piani's bound r_{CP} is depicted in FIG. 3.

In order to generalize the information exclusion relation to multi-measurements, we recall that the universal bound of tensor products of two probability distribution vectors can be computed by optimization over minors of the overlap matrix^{21,22}. More generally for the multi-tensor product $(p_{i_1}^1 p_{i_2}^2 \cdots p_{i_N}^N)$ corresponding to measurement M_m on a fixed quantum state, there exists similarly a universal upper bound ω : $(p_{i_1}^1 p_{i_2}^2 \cdots p_{i_N}^N) \prec \omega$. Then we have the following lemma, which generalizes Eq. (7).

Lemma 1. For any bipartite state ρ_{AB} , let M_m ($m = 1, 2, \dots, N$) be N measurements on system A , and B the quantum memory correlated to A , then the following entropic uncertainty relation holds,

$$\sum_{m=1}^N H(M_m|B) \geq -\frac{1}{N}\omega\mathfrak{B} + (N-1)H(A) - NH(B), \quad (10)$$

where ω is the d^N -dimensional majorization bound for the N measurements M_m and \mathfrak{B} is the d^N -dimensional vector $(\log(\omega \cdot \mathfrak{A}_{i_1, i_2, \dots, i_N}))^\dagger$ defined as follows. For each multi-index (i_1, i_2, \dots, i_N) , the d^N -dimensional vector $\mathfrak{A}_{i_1, i_2, \dots, i_N}$ has entries of the form $c(1, 2, \dots, N)c(2, 3, \dots, 1) \cdots c(N, 1, \dots, N-1)$ sorted in decreasing order with respect to the indices (i_1, i_2, \dots, i_N) where $c(1, 2, \dots, N) = \sum_{i_2, \dots, i_{N-1}} \max_{i_1} c(u_{i_1}^1, u_{i_2}^2) \cdots c(u_{i_{N-1}}^{N-1}, u_{i_N}^N)$.

See Methods for a proof of Lemma 1.

We remark that the *admixture bound* introduced in Ref.¹⁹ was based upon the majorization theory with the help of the action of the symmetric group, and it was shown that the bound outperforms previous results. However, the admixture bound cannot be extended to the entropic uncertainty relations in the presence of quantum memory for multiple measurements directly. Here we first use a new method to generalize the results of Ref.¹⁹ to allow for the quantum side information by mixing properties of the conditional entropy and Holevo inequality in Lemma 1. Moreover, by combining Lemma 1 with properties of the entropy we are able to give an enhanced information exclusion relation (see Theorem 2 for details).




Figure 4. Comparison of our bound with that of Zhang et al. Our bound r_x in the bottom is tighter than the top curve of Zhang's bound \mathcal{U}_1 .

The following theorem is obtained by subtracting entropic uncertainty relation from the above result.

Theorem 2. For any bipartite state ρ_{AB} , let M_m ($m = 1, 2, \dots, N$) be N measurements on system A , and B the quantum memory correlated to A , then

$$\sum_{m=1}^N I(M_m : B) \leq \log_2 d^N + \frac{1}{N} \omega \mathfrak{B} + NH(B) - (N-1)H(A) := r_x, \quad (11)$$

where $\frac{1}{N} \omega \mathfrak{B}$ is defined in Eq. (10).

See Methods for a proof of Theorem 2.

Throughout this paper, we take $NH(B) - (N-1)H(A)$ instead of $-(N-1)H(A|B)$ as the variable that quantifies the amount of entanglement between measured particle and quantum memory since $NH(B) - (N-1)H(A)$ can outperform $-(N-1)H(A|B)$ numerically to some extent for entropic uncertainty relations.

Our new bound for multi-measurements offers an improvement than the bound recently given in Ref.¹⁸. Let us recall the information exclusion relation bound¹⁸ for multi-measurements (state-independent):

$$\sum_{m=1}^N I(M_m : B) \leq \min \left\{ \widetilde{\mathcal{U}}_1, \widetilde{\mathcal{U}}_2, \widetilde{\mathcal{U}}_3 \right\} \quad (12)$$

with the bounds $\widetilde{\mathcal{U}}_1$, $\widetilde{\mathcal{U}}_2$ and $\widetilde{\mathcal{U}}_3$ are defined as follows:

$$\begin{aligned} \widetilde{\mathcal{U}}_1 &= N \log_2 d + NH(B) - (N-1)H(A) + \min_{(i_1 \dots i_N) \in \mathfrak{S}_N} \left\{ \log \max_{i_N} \left\{ \sum_{i_2 \dots i_{N-1}} \max_{i_1} \prod_{n=1}^{N-1} c(u_{i_n}^n, u_{i_{n+1}}^{n+1}) \right\} \right\}, \\ \widetilde{\mathcal{U}}_2 &= (N-1) \log_2 d + NH(B) - (N-1)H(A) + \min_{(i_1 \dots i_N) \in \mathfrak{S}_N} \left\{ \log \sum_{i_2 \dots i_N} \max_{i_1} \prod_{n=1}^{N-1} c(u_{i_n}^n, u_{i_{n+1}}^{n+1}) \right\}, \\ \widetilde{\mathcal{U}}_3 &= N \log_2 d + \frac{N}{2} (2H(B) - H(A)) + \frac{1}{|I_2|} \sum_{(k,l) \in I_2} \left\{ \min \left\{ \log \max_{i_k} c(u_{i_k}^k, u_{i_l}^l), \log \max_{i_l} c(u_{i_k}^k, u_{i_l}^l) \right\} \right\}. \end{aligned}$$

Here the first two maxima are taken over all permutations $(i_1 i_2 \dots i_N) : j \rightarrow i_j$, and the third is over all possible subsets $I_2 = \{(k_1, l_1), \dots, (k_{|I_2|}, l_{|I_2|})\}$ such that $(k_1, l_1, \dots, k_{|I_2|}, l_{|I_2|})$ is a $|I_2|$ -permutation $1, \dots, N$. For example, $(12), (23), \dots, (N-1, N), (N1)$ is a 2-permutation of $1, \dots, N$, while $(12), (13), \dots, (N-1, N), (N, 1)$ is an $(N-1)$ -permutation of $1, \dots, N$. Clearly, $\widetilde{\mathcal{U}}_3$ is the average value of all potential two-measurement combinations.

Using the permutation symmetry, we have the following Theorem which improves the bound $\widetilde{\mathcal{U}}_3$.

Theorem 3. Let ρ_{AB} be the bipartite density matrix with measurements M_m ($m = 1, 2, \dots, N$) on the system A with a quantum




Figure 5. Comparison of our two bounds via combinatorial and majorization methods: the top curve is r_{opt} (combinatorial), while the lower curve is r_x (majorization).

memory B as in Theorem 2, then

$$\sum_{m=1}^N I(M_m : B) \leq N \log_2 d + \frac{N}{2} (2H(B) - H(A)) + \frac{1}{|I_L|} \sum_{(k_1, k_2, \dots, k_L) \in I_L} \left\{ \min_{(k_1, k_2, \dots, k_L)} \left\{ \log \max_{i_L} \sum_{k_2, \dots, k_{L-1}} \max_{k_1} \prod_{n=1}^{L-1} c(u_{k_n}^n, u_{k_{n+1}}^{n+1}) \right\} \right\} := r_{opt}, \quad (13)$$

where the minimum is over all L -permutations of $1, \dots, N$ for $L = 2, \dots, N$.

In the above we have explained that the bound $\widetilde{\mathcal{U}}_3$ is obtained by taking the minimum over all possible 2-permutations of $1, 2, \dots, N$, naturally our new bound r_{opt} in Theorem 3 is sharper than $\widetilde{\mathcal{U}}_3$ as we have considered all possible multi-permutations of $1, 2, \dots, N$.

Now we compare $\widetilde{\mathcal{U}}_1$ with r_x . As an example in three-dimensional space, one chooses three measurements as follows²⁶ :

$$\begin{aligned} u_1^1 &= (1, 0, 0), u_2^1 = (0, 1, 0), u_3^1 = (0, 0, 1); \\ u_1^2 &= \left(\frac{1}{\sqrt{2}}, 0, -\frac{1}{\sqrt{2}} \right), u_2^2 = (0, 1, 0), u_3^2 = \left(\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}} \right); \\ u_1^3 &= (\sqrt{a}, e^{i\phi} \sqrt{1-a}, 0), u_2^3 = (\sqrt{1-a}, -e^{i\phi} \sqrt{a}, 0), u_3^3 = (0, 0, 1). \end{aligned}$$

FIG 4 shows the comparison when a changes and $\phi = \pi/2$, where it is clear that r_x is better than $\widetilde{\mathcal{U}}_1$.

The relationship between r_{opt} and r_x is sketched in FIG. 5. In this case r_x is better than r_{opt} for three measurements of dimension three, therefore $\min\{r_{opt}, r_x\} = \min\{r_x\}$. Rigorous proof that r_x is always better than r_{opt} is nontrivial, since all the possible combinations of measurements less than N must be considered.

On the other hand, we can give a bound better than $\widetilde{\mathcal{U}}_2$. Recall that the concavity has been utilized in the formation of $\widetilde{\mathcal{U}}_2$, together with all possible combinations we will get following lemma (in order to simplify the process, we first consider cases of three measurements, then generalize it to multiple measurements).

Lemma 2. For any bipartite state ρ_{AB} , let M_1, M_2, M_3 be three measurements on system A in the presence of quantum memory B , then

$$\sum_{m=1}^3 I(M_m : B) \leq -2H(A|B) + \sum_{\text{cyclic perm}} \log_2 \left[\sum_{i_3} \left(\sum_{i_2} \max_{i_1} c(u_{i_1}^1, u_{i_2}^2) c(u_{i_2}^2, u_{i_3}^3) \right)^{\frac{1}{3}} \right], \quad (14)$$

where the sum is over the three cyclic permutations of $1, 2, 3$.

See Methods for a proof of Lemma 2.

Observe that the right hand side of Eq. (14) adds the sum of three terms

$$\frac{1}{3} \sum_{i_3} p_{i_3}^3 \log_2 \left[\sum_{i_2} \max_{i_1} c(1, 2, 3) \right], \quad \frac{1}{3} \sum_{i_1} p_{i_1}^1 \log_2 \left[\sum_{i_3} \max_{i_2} c(2, 3, 1) \right], \quad \frac{1}{3} \sum_{i_2} p_{i_2}^2 \log_2 \left[\sum_{i_1} \max_{i_3} c(3, 1, 2) \right].$$

Naturally, we can also add $\frac{1}{2} \sum_{i_3} p_{i_3}^3 \log_2 [\sum_{i_2} \max_{i_1} c(1, 2, 3)]$ and $\frac{1}{2} \sum_{i_1} p_{i_1}^4 \log_2 [\sum_{i_3} \max_{i_2} c(2, 3, 1)]$. By the same method, consider all possible combination and denote the minimal as r_{3y} . Similar for N -measurements, set the minimal bound under the concavity of logarithm function as r_{Ny} , moreover let $r_y = \min_m \{r_{my}\}$ ($1 \leq m \leq N$), hence $r_y \leq \widetilde{\mathcal{U}}_2$, finally we get

Theorem 4. For any bipartite state ρ_{AB} , let M_m ($m = 1, 2, \dots, N$) be N measurements on system A , and B the quantum memory correlated to A , then

$$\sum_{m=1}^N I(M_m : B) \leq \min\{r_x, r_y\} \quad (15)$$

with $\frac{1}{N} \omega \mathfrak{B}$ the same in Eq. (10). Since $\min\{r_x, r_y\} \leq \min\{\widetilde{\mathcal{U}}_1, \widetilde{\mathcal{U}}_2, \widetilde{\mathcal{U}}_{opt}\}$ and all figures have shown our newly construct bound $\min\{r_x, r_y\}$ is tighter. Noted that there is no clear relation between r_x and r_y , while the bound r_y cannot be obtained by simply subtracting the bound of entropic uncertainty relations in the presence of quantum memory. Moreover, if r_y outperforms r_x , then we can utilize r_y to achieve new bound for entropic uncertainty relations stronger than $-\frac{1}{N} \omega \mathfrak{B}$.

Conclusions

We have derived new bounds of the information exclusion relation for multi-measurements in the presence of quantum memory. The bounds are shown to be tighter than recently available bounds by detailed illustrations. Our bound is obtained by utilizing the concavity of the entropy function. The procedure has taken into account of all possible permutations of the measurements, thus offers a significant improvement than previous results which had only considered part of 2-permutations or combinations. Moreover, we have shown that majorization of the probability distributions for multi-measurements offers better bounds. In summary, we have formulated a systematic method of finding tighter bounds by combining the symmetry principle with majorization theory, all of which have been made easier in the context of mutual information. We remark that the new bounds can be easily computed by numerical computation.

Methods

Proof of Theorem 1. Recall that the quantum relative entropy $D(\rho||\sigma) = \text{Tr}(\rho \log_2 \rho) - \text{Tr}(\rho \log_2 \sigma)$ satisfies that $D(\rho||\sigma) \geq D(\tau\rho||\tau\sigma) \geq 0$ under any quantum channel τ . Denote by $\rho_{AB} \rightarrow \rho_{M_1 B}$ the quantum channel $\rho_{AB} \rightarrow \sum_i |u_i^1\rangle\langle u_i^1| \rho_{AB} |u_i^1\rangle\langle u_i^1|$, which is also $\rho_{M_1 B} = \sum_i |u_i^1\rangle\langle u_i^1| \otimes \text{Tr}_A(\rho_{AB} |u_i^1\rangle\langle u_i^1|)$. Note that both $M^i = \{|u_i^1\rangle\}$ ($i = 1, 2$) are measurements on system A , we have that for a bipartite state ρ_{AB}

$$\begin{aligned} H(M_1|B) - H(A|B) &= H(\rho_{M_1 B}) - H(\rho_{AB}) = \text{Tr}(\rho_{AB} \log_2 \rho_{AB}) - \text{Tr}(\rho_{M_1 B} \log_2 \rho_{M_1 B}) \\ &= D(\rho_{AB} || \sum_i |u_i^1\rangle\langle u_i^1| \otimes \text{Tr}_A(\rho_{AB} |u_i^1\rangle\langle u_i^1|)). \end{aligned}$$

Note that $\text{Tr}_B \text{Tr}_A(\rho_{AB} |u_i^1\rangle\langle u_i^1|) = p_i^1$, the probability distribution of the reduced state ρ_A under the measurement M_1 , so $\sigma_{B_i} = \text{Tr}_A(\rho_{AB} |u_i^1\rangle\langle u_i^1|) / p_i^1$ is a density matrix on the system B . Then the last expression can be written as

$$\begin{aligned} &D(\rho_{AB} || \sum_i p_i^1 |u_i^1\rangle\langle u_i^1| \otimes \sigma_{B_i}) \\ &\geq D(\rho_{M_2 B} || \sum_{i_1, i_2} p_{i_1}^1 C_{i_1 i_2} |u_{i_2}^2\rangle\langle u_{i_2}^2| \otimes \sigma_{B_{i_1}}). \end{aligned}$$

If system B is a classical register, then we can obtain

$$H(M_1) + H(M_2) \geq H(A) - \sum_{i_2} p_{i_2}^2 \log \sum_{i_1} p_{i_1}^1 c(u_{i_1}^1, u_{i_2}^2), \quad (16)$$

by swapping the indices i_1 and i_2 , we get that

$$H(M_2) + H(M_1) \geq H(A) - \sum_{i_1} p_{i_1}^1 \log \sum_{i_2} p_{i_2}^2 c(u_{i_2}^2, u_{i_1}^1). \quad (17)$$

Their combination implies that

$$H(M_1) + H(M_2) \geq H(A) - \frac{1}{2} \left(\sum_{i_2} p_{i_2}^2 \log \sum_{i_1} p_{i_1}^1 c(u_{i_1}^1, u_{i_2}^2) + \sum_{i_1} p_{i_1}^1 \log \sum_{i_2} p_{i_2}^2 c(u_{i_2}^2, u_{i_1}^1) \right), \quad (18)$$

thus it follows from Ref.²⁷ that

$$H(M_1|B) + H(M_2|B) \geq H(A) - 2H(B) - \frac{1}{2} \left(\sum_{i_2} p_{i_2}^2 \log \sum_{i_1} p_{i_1}^1 c(u_{i_1}^1, u_{i_2}^2) + \sum_{i_1} p_{i_1}^1 \log \sum_{i_2} p_{i_2}^2 c(u_{i_2}^2, u_{i_1}^1) \right), \quad (19)$$

hence

$$\begin{aligned} I(M_1|B) + I(M_2|B) &= H(M_1) + H(M_2) - (H(M_1|B) + H(M_2|B)) \\ &\leq H(M_1) + H(M_2) + \frac{1}{2} \left(\sum_{i_2} p_{i_2}^2 \log \sum_{i_1} p_{i_1}^1 c(u_{i_1}^1, u_{i_2}^2) + \sum_{i_1} p_{i_1}^1 \log \sum_{i_2} p_{i_2}^2 c(u_{i_2}^2, u_{i_1}^1) \right) + 2H(B) - H(A) \\ &\leq H(M_1) + H(M_2) + \frac{1}{2} \sum_{i_1, i_2} p_{i_1}^1 p_{i_2}^2 \log_2 \left(\sum_{i, j} p_i^1 p_j^2 c(u_i^1, u_{i_2}^2) c(u_j^2, u_{i_1}^1) \right) + 2H(B) - H(A) \\ &\leq 2 + \frac{1}{2} \omega \mathfrak{B} + 2H(B) - H(A), \end{aligned} \quad (20)$$

where the last inequality has used $H(M_i) \leq \log_2 d$ ($i = 1, 2$) and the vector \mathfrak{B} of length d^2 , whose entries $\mathfrak{B}_{i_1 i_2} = \log_2(\omega \cdot \mathfrak{A}_{i_1 i_2})$ are arranged in decreasing order with respect to (i_1, i_2) . Here the vector \mathfrak{A} is defined by $\mathfrak{A}_{i_1 i_2} = c(u_{i_1}^1, u_{i_2}^2) c(u_{i_2}^2, u_{i_1}^1)$ for each (i_1, i_2) and also sorted in decreasing order. Note that the extra term $2H(B) - H(A)$ is another quantity appearing on the right-hand side that describes the amount of entanglement between the measured particle and quantum memory besides $-H(A|B)$.

We now derive the *information exclusion relation* for qubits in the form of $I(M_1 : B) + I(M_2 : B) \leq 2 + \frac{1}{2} \omega \mathfrak{B} + 2H(B) - H(A)$, and this completes the proof.

Proof of Lemma 1. Suppose we are given N measurements M_1, \dots, M_N with orthonormal bases $\{|u_{i_j}^j\rangle\}$. To simplify presentation we denote that

$$c_{i_1, \dots, i_N}^{1, \dots, N} = c(u_{i_1}^1, u_{i_2}^2) c(u_{i_2}^2, u_{i_3}^3) \cdots c(u_{i_{N-1}}^{N-1}, u_{i_N}^N).$$

Then we have that²⁶

$$(1 - N)H(A) + \sum_{m=1}^N H(M_m) \geq -\text{Tr}(\rho \log \sum_{i_1, i_2, \dots, i_N} p_{i_1}^1 c_{i_1, \dots, i_N}^{1, \dots, N} |u_{i_N}^N\rangle \langle u_{i_N}^N|) = -\sum_{i_N} p_{i_N}^N \log \sum_{i_1, i_2, \dots, i_{N-1}} p_{i_1}^1 c_{i_1, \dots, i_N}^{1, \dots, N}. \quad (21)$$

Then consider the action of the cyclic group of N permutations on indices $1, 2, \dots, N$, and taking the average gives the following inequality:

$$\sum_{m=1}^N H(M_m) \geq -\frac{1}{N} \omega \mathfrak{B} + (N - 1)H(A), \quad (22)$$

where the notations are the same as appeared in Eq. (10). Thus it follows from Ref.²⁷ that

$$\sum_{m=1}^N H(M_m|B) \geq -\frac{1}{N} \omega \mathfrak{B} + (N - 1)H(A) - NH(B). \quad (23)$$

The proof is finished.

Proof of Theorem 2. Similar to the proof of Theorem 1, due to $I(M_m : B) = H(M_m) - H(M_m|B)$, thus we get

$$\begin{aligned} \sum_{m=1}^N I(M_m : B) &= \sum_{m=1}^N H(M_m) - \sum_{m=1}^N H(M_m|B) \\ &\leq \sum_{m=1}^N H(M_m) + \frac{1}{N} \omega \mathfrak{B} + NH(B) - (N - 1)H(A) \\ &\leq \log_2 d^N + \frac{1}{N} \omega \mathfrak{B} + NH(B) - (N - 1)H(A), \end{aligned} \quad (24)$$

with the product $\frac{1}{N} \omega \mathfrak{B}$ the same in Eq. (10).

Proof of Lemma 2. First recall that for $\sum_{i_3} p_{i_3}^3 \log_2 [\sum_{i_2} \max_{i_1} c(u_{i_1}^1, u_{i_2}^2) c(u_{i_2}^2, u_{i_3}^3)]$ we have

$$\begin{aligned} & H(M_3) + \sum_{i_3} p_{i_3}^3 \log_2 [\sum_{i_2} \max_{i_1} c(u_{i_1}^1, u_{i_2}^2) c(u_{i_2}^2, u_{i_3}^3)] \\ &= \sum_{i_3} p_{i_3}^3 \log_2 \frac{1}{p_{i_3}} + \sum_{i_3} p_{i_3}^3 \log_2 [\sum_{i_2} \max_{i_1} c(u_{i_1}^1, u_{i_2}^2) c(u_{i_2}^2, u_{i_3}^3)] \\ &\leq \log_2 \sum_{i_3} [\sum_{i_2} \max_{i_1} c(u_{i_1}^1, u_{i_2}^2) c(u_{i_2}^2, u_{i_3}^3)], \end{aligned}$$

where we have used concavity of log. By the same method we then get

$$\begin{aligned} \sum_{m=1}^3 I(M_m : B) &\leq \sum_{m=1}^3 H(M_m) + 3H(B) - 2H(A) + \frac{1}{3} \sum_{i_3} p_{i_3}^3 \log_2 [\sum_{i_2} \max_{i_1} c(1, 2, 3)] \\ &\quad + \frac{1}{3} \sum_{i_2} p_{i_2}^2 \log_2 [\sum_{i_1} \max_{i_3} c(3, 1, 2)] + \frac{1}{3} \sum_{i_1} p_{i_1}^1 \log_2 [\sum_{i_3} \max_{i_2} c(2, 3, 1)] \\ &= \sum_{m=1}^3 H(M_m) + 3H(B) - 2H(A) + \sum_{i_3} p_{i_3}^3 \log_2 [\sum_{i_2} \max_{i_1} c(1, 2, 3)]^{\frac{1}{3}} \\ &\quad + \sum_{i_2} p_{i_2}^2 \log_2 [\sum_{i_1} \max_{i_3} c(3, 1, 2)]^{\frac{1}{3}} + \sum_{i_1} p_{i_1}^1 \log_2 [\sum_{i_3} \max_{i_2} c(2, 3, 1)]^{\frac{1}{3}} \\ &\leq 3H(B) - 2H(A) + \log_2 \sum_{i_3} [\sum_{i_2} \max_{i_1} c(1, 2, 3)]^{\frac{1}{3}} + \log_2 \sum_{i_2} [\sum_{i_1} \max_{i_3} c(3, 1, 2)]^{\frac{1}{3}} + \log_2 \sum_{i_1} [\sum_{i_3} \max_{i_2} c(2, 3, 1)]^{\frac{1}{3}} \\ &= 3H(B) - 2H(A) \\ &\quad + \log_2 \left\{ \left[\sum_{i_3} \left(\sum_{i_2} \max_{i_1} c(1, 2, 3) \right)^{\frac{1}{3}} \right] \left[\sum_{i_2} \left(\sum_{i_1} \max_{i_3} c(3, 1, 2) \right)^{\frac{1}{3}} \right] \left[\sum_{i_1} \left(\sum_{i_3} \max_{i_2} c(2, 3, 1) \right)^{\frac{1}{3}} \right] \right\}, \quad (25) \end{aligned}$$

with $c(1, 2, 3)$, $c(2, 3, 1)$ and $c(3, 1, 2)$ the same as in Eq. (14) and this completes the proof.

References

1. Cover, T. M. & Thomas, J. A. *Elements of information theory*. 2nd ed. (Wiley, New York, 2005).
2. Holevo, A. S. *Bounds for the quantity of information transmitted by a quantum communication channel*. Probl. Inf. Trans. **9**, 177 (1973).
3. Yuen, A. S. & Ozawa, M. *Ultimate information carrying limit of quantum systems*. Phys. Rev. Lett. **70**, 363 (1993).
4. Wehner, S. & Winter, A. *Entropic uncertainty relations a survey*. New J. Phys. **12**, 025009 (2010).
5. Bosyk, G. M., Portesi, M. & Plastino, A. *Collision entropy and optimal uncertainty*. Phys. Rev. A **85**, 012108 (2012).
6. Białynicki-Birula, I. & Mycielski, J. *Uncertainty relations for information entropy in wave mechanics*. Comm. Math. Phys. **44**, 129 (1975).
7. Deutsch, D. *Uncertainty in quantum measurements*. Phys. Rev. Lett. **50**, 631 (1983).
8. Maassen, H. & Uffink, J. B. M. *Generalized entropic uncertainty relations*. Phys. Rev. Lett. **60**, 1103 (1988).
9. Dupuis, F., Szehr, O. & Tomamichel, M. *A decoupling approach to classical data transmission over quantum channels*. IEEE Trans. on Inf. Theory **60**(3), 1562 (2014).
10. Heisenberg, W. *Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik*. Z. Phys. **43**, 172 (1927).
11. Hall, M. J. W. *Information exclusion principle for complementary observables*. Phys. Rev. Lett. **74**, 3307 (1995).
12. Hall, M. J. W. *Quantum information and correlation bounds*. Phys. Rev. A **55**, 100 (1997).
13. Coles, P. J., Yu, L., Gheorghiu, V. & Griffiths, R. B. *Information-theoretic treatment of tripartite systems and quantum channels*. Phys. Rev. A **83**, 062338 (2011).

14. Reza, F. M., *An introduction to information theory*. (McGraw-Hill, New York, 1961).
15. Grudka, A., Horodecki, M., Horodecki, P., Horodecki, R., Kłobus, W. & Pankowski, Ł. *Conjectured strong complementary correlations tradeoff*. Phys. Rev. A **88**, 032106 (2013).
16. Coles, P. J. & Piani, M. *Improved entropic uncertainty relations and information exclusion relations*. Phys. Rev. A **89**, 022112 (2014).
17. Berta, M., Christandl, M., Colbeck, R., Renes, J. M. & Renner, R. *The uncertainty principle in the presence of quantum memory*. Nat. Phys. **6**, 659 (2010).
18. Zhang, J., Zhang, Y. & Yu, C.-S. *Entropic uncertainty relation and information exclusion relation for multiple measurements in the presence of quantum memory*. Scientific Reports **5**, 11701 (2015).
19. Xiao, Y., Jing, N., Fei, S.-M., Li, T., Li-Jost, X., Ma, T. & Wang, Z.-X. *Strong entropic uncertainty relations for multiple measurements*. Phys. Rev. A **93**, 042125 (2016).
20. Rudnicki, Ł., Puchała, Z. & Życzkowski, K. *Strong majorization entropic uncertainty relations*. Phys. Rev. A **89**, 052115 (2014).
21. Friedland, S., Gheorghiu, V. & Gour, G. *Universal uncertainty relations*. Phys. Rev. Lett. **111**, 230401 (2013).
22. Puchała, Z., Rudnicki, Ł. & Życzkowski, K. *Majorization entropic uncertainty relations*. J. Phys. A **46**, 272002 (2013).
23. Marshall, A. W., Olkin, I. & Arnold, B. C. *Inequalities: theory of majorization and its applications*. 2nd ed. Springer Series in Statistics. (Springer, New York, 2011).
24. Havrda, J. & Charvát, F. *Quantification method of classification processes. Concept of structural α -entropy*. Kybernetika **3**, 30 (1967).
25. Ghirardi, G., Marinatto, L. & Romano, R. *An optimal entropic uncertainty relation in a two-dimensional Hilbert space*. Phys. Lett. A **317**, 32 (2003).
26. Liu, S., Mu, L.-Z. & Fan, H. *Entropic uncertainty relations for multiple measurements*. Phys. Rev. A **91**, 042133 (2015).
27. Nielsen, M. A. & Chuang, I. L.. *Quantum computation and quantum information*. (Cambridge Univ. Press, Cambridge, 2000).

Acknowledgments

We would like to thank Tao Li for help on computer diagrams. This work is supported by National Natural Science Foundation of China (grant nos. 11271138, 11531004), China Scholarship Council and Simons Foundation grant no. 198129.

Author contributions statement

Y. X. and N. J. analyzed and wrote the manuscript. They reviewed the paper together with X. L.-J.

Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Xiao, Y., Jing, N. and Li-Jost, X. Enhanced Information Exclusion Relations. Sci. Rep. (2016), Article no. —