# The rotation - Lithium depletion correlation in the $\beta$ Pictoris association and LDB age determination

S. Messina<sup>1</sup>, A.C. Lanzafame<sup>2,1</sup>, G.A. Feiden<sup>3</sup>, M. Millward<sup>4</sup>, S. Desidera<sup>5</sup>, A. Buccino<sup>6</sup>, I. Curtis<sup>7</sup>, E. Jofré<sup>8,9</sup>, P. Kehusmaa<sup>10</sup>, B.J. Medhi<sup>11</sup>, B. Monard<sup>12</sup>, R. Petrucci<sup>8,9</sup>

<sup>1</sup> INAF-Catania Astrophysical Observatory, via S.Sofia, 78 I-95123 Catania, Italy e-mail: sergio.messina@oact.inaf.it

- <sup>3</sup> Department of Physics & Astronomy, Uppsala University, Box 516, SE-751 20, Uppsala, Sweden e-mail: gregory.feiden@physics.uu.se
- <sup>4</sup> York Creek Observatory, Georgetown, Tasmania, Australia
- e-mail: mervyn.millward@yorkcreek.net
- <sup>5</sup> INAF-Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122 Padova, Italy e-mail: silvano.desidera@oapd.inaf.it
- <sup>6</sup> Instituto de Astronomía y Física del Espacio (IAFE-CONICET), Buenos Aires, Argentina e-mail: abuccino@iafe.uba.ar
- <sup>7</sup> IAU Minor Planet Center code D79, 2 Yandra Street, Vale Park, South Australia 5081 Australia e-mail: Ivan.Curtis@keyworks.com.au
- <sup>8</sup> Observatorio Astronómico de Córdoba, Laprida 854, X5000BGR, Córdoba, Argentina e-mail: jofre.emiliano@gmail.com; romina@oac.unc.edu.ar
- <sup>9</sup> Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina
- <sup>10</sup> Harlingten Atacama Observatory, San Pedro de Atacama, Chile. e-mail: petri@kehusmaa-astro.com
- <sup>11</sup> Aryabhatta Research Institute of Observational Sciences, Manora Peak, Nainital 263129, India e-mail: biman@aries.res.in
- <sup>12</sup> Klein Karoo Observatory, Western Cape, South Africa e-mail: bmonard@mweb.co.za

#### ABSTRACT

*Context.* There is evidence in the 125-Myr Pleiades cluster, and more recently in the 5-Myr NGC 2264 cluster, that rotation plays a key role in the Lithium (Li) depletion processes among low-mass stars. Fast rotators appear to be less Li-depleted than equal-mass slow rotators.

Aims. We intend to explore the existence of a Li depletion - rotation connection among the  $\beta$  Pictoris members at an age of about 24 Myr, and to use such correlation either to confirm or to improve the age estimate based on the Lithium Depletion Boundary (LDB) modeling.

*Methods.* We have photometrically monitored all the known members of the  $\beta$  Pictoris association with at least one Lithium equivalent width (Li EW) measurement from the literature.

*Results.* We measured the rotation periods of 30 members for the first time and retrieved from the literature the rotation periods for other 36 members, building a catalogue of 66 members with measured rotation period and Li EW.

Conclusions. We find that in the  $0.3 < M < 0.8 M_{\odot}$  range, there is a strong correlation between rotation and Li EW. For higher mass stars, no significant correlation is found. For very low mass stars in the Li depletion onset, at about  $0.1 M_{\odot}$ , data are too few to infer a significant correlation. The observed Li EWs are compared with those predicted by the Dartmouth stellar evolutionary models that incorporate the effects of magnetic fields. After decorrelating the Li EW from the rotation period, we find that the hot side of the LDB is fitted well by Li EW values corresponding to an age of  $25\pm3$  Myr in good agreement with independent estimates from the literature.

Key words. Stars: activity - Stars: late-type - Stars: rotation - Stars: starspots - Stars: individual: beta Pictoris association - Stars: abundances

#### 1. Introduction

In recent years much attention has been payed to the  $24\pm3\,\text{Myr}$  (Bell et al. 2015) young  $\beta$  Pictoris stellar association. Several studies have allowed to increase sig-

nificantly the number of confirmed members and to discover many more candidate members. Just to mention the most recent works, we refer the readers to Lépine & Simon (2009); Kiss et al. (2011); Schlieder et al. (2010, 2012); Shkolnik et al. (2012); Malo et al. (2013, 2014a,b). The first comprehensive search for the rotation periods of the low-mass members of  $\beta$  Pictoris was carried out by

<sup>&</sup>lt;sup>2</sup> Universitá di Catania, Dipartimento di Fisica e Astronomia, Sezione Astrofisica, via S. Sofia 78, I-95123 Catania, Italy

e-mail: a.lanzafame@unict.it

Send offprint requests to: Sergio Messina

Messina et al. (2010, 2011) who retrieved a total of 38 low-mass members (i.e., spectral types from late F to M) from the earlier compilations of Zuckerman & Song (2004), Torres et al. (2006), and Kiss et al. (2011). Their study provided the rotation periods of 33 out of 38 members.

In the light of the mentioned studies, we have again explored up to the most recent literature and, at this time, we could finally compile a new list of 117 stars among members and candidate members with spectral types later than about F3V. Then, we have started a new rotational study on this enlarged sample. To get the photometric rotation periods of our targets, we used our own observations, archive data, and also we made use of periods from the literature.

As result of our photometric investigation, we obtained the rotation periods of 112 out of 117 stars. Specifically, we measured for the first time the rotation periods of 56 stars. For another 27 stars, we could confirm with our analysis of new or archived data the values reported in the literature. For 29 stars we adopted the literature values. For the remaining 5 stars, our periodogram analysis did not provide the rotation period. The results of this investigation are presented in the catalogue of photometric rotation period of the  $\beta$  Pictoris association members (Messina et al. 2016; Paper I) where we describe the photometric observations newly obtained, their reduction and analysis, and a detailed discussion of our results obtained for each individual star.

In this paper (Paper II), we focus on a sub-sample consisting of 66 members for which we know the rotation period and have one measurement at least (from the literature) of the Lithium equivalent width (EW). This sub-sample is to date the largest of any known young loose association. We will make use of it to investigate the correlation between rotation and Li depletion and to compare with earlier results the age of the  $\beta$  Pictoris association obtained by the modeling of the Lithium Depletion Boundary (LDB), after decorrelating the Li EW from rotation.

#### 2. Sample description

The sample under analysis consists of 66 members with spectral type later than F3V with one measurement, at least, of the Li EW (see Table 1). This sample, which has significantly increased with respect to earlier studies, makes a new investigation on the age of the  $\beta$  Pictoris association with the LDB modeling method necessary. In fact, earlier studies have made use of a smaller number of association members. Mentuch et al. (2008) made use of a sample of 23 members; Macdonald & Mullan (2010) used a sample of only 10 members. Binks & Jeffries (2014) and Malo et al. (2014b) used, respectively, about 40 and 34 members.

Moreover, we also now know the rotation periods of all these 66 stars, which put us in the position to get more accurate results and to deeply explore the mechanism of Li depletion. More specifically, we measured for the first time the rotation periods of 30 out of 66 members. These new measurements will be presented in the mentioned catalogue containing the photometric rotation periods of all 112 members/candidate members of the association. For another 16 members, the rotation periods were taken from Messina et al. (2010, 2011). The remaining 20 rotation periods were taken from different literature sources.

This sample represents, to date, one of the three largest

samples ever used to investigate the correlation between rotation period and Li depletion. The other samples consist of members of the Pleiades and of the M34 stellar open clusters investigated by Gondoin (2014) using the rotation periods from Hartman et al. (2010), and of members of the NGC 2264 open cluster whose results on the Lithiumrotation connection are presented by Bouvier et al. (2016). The earlier larger sample of Pleiades members analysed by Soderblom et al. (1993) made use of the projected rotational velocity  $(v \sin i)$ , which is less accurate owing to the uncertainty arising from the unknown sin *i*.

#### 3. Li distribution

In Fig. 1, we plot the distribution of Li EW for all 66 members versus the  $V-K_s$  color. This color index is measured using the  $K_s$  magnitude from the 2MASS project (Cutri et al. 2003) and the brightest (presumably unspotted) V magnitude from the long photometric time series available in the All Sky Automated Survey (ASAS) archive (Pojmanski 2002). This choice allows us to reduce the impact of photometric variability on the V-band magnitude. Among different color indexes, the  $V-K_s$  turned out to be the most homogeneous and that with the largest range of values, allowing an accurate representation of different masses. We applied no color correction for interstellar absorption since the members under study have distances in the range from  $10\,\mathrm{pc}$  to  $80\,\mathrm{pc}$ , therefore the interstellar reddening can be considered negligible. The average uncertainty on this color is  $\sigma_{V-K_s} = 0.03 \text{ mag}$ . There are only 8 stars that belong to close binaries that are not resolved neither in V nor in  $K_s$  magnitudes and consist of components differing by no more than 2 spectral subclasses. For these stars the measured  $V-K_s$  for the primary component, the one with measurement of Li EW, resulted to be redder than the expected value for the spectral type. In these cases we adopted the  $V-K_s$  values from Pecaut & Mamajek (2013), and corresponding to the spectral type of the primary component. The Li EW measurements are retrieved from Mentuch et al. (2008), da Silva et al. (2009), Kiss et al. (2011), Binks & Jeffries (2014), Moór et al. (2013), Malo et al. (2014b), Reid et al. (2002), Torres et al. (2006), and Desidera et al. (2015). The average uncertainty associated to these measurements is generally not larger than  $\sim 40 \,\mathrm{m}$ Å. The error bars associated to the measurements (in both axes) are equal to the size of the symbols plotted in Fig. 1. In a few cases more than one measurement is available. The range of values of the Li EW in these cases is plotted with a dotted line. Almost all measurements (for a total of 20 members) in the color range  $4.5 < V-K_s < 5.3 \text{ mag}$  are upper limits. Different symbols are used to distinguish the different nature of the considered members (see the legend in Fig. 1). Detailed information on each member and the criterion to distinguish close from wide components is given in Messina et al. (Paper III, in prep.).

In Fig. 1, we note three important features. First, we note a very well defined LDB that will be used for an estimate of the age. Second, in the color range  $3.2 < V-K_s < 4.5$  mag, we note, even by just a visual inspection, a scatter in the Li EW significantly larger than in the range of bluer members. We will investigate the role played by the rotation in producing this scattered depletion. Finally, we note that in the range  $2.5 < V-K_s < 3.0$  mag, we have only three stars



Fig. 1. Distribution of Li EW versus  $V-K_s$  color for confirmed members of the  $\beta$  Pictoris association. The meaning of symbols is given in the legend. Dotted vertical lines indicate the range of values for stars with multiple Li EW measurements. All circled symbols indicate stars hosting debris discs.

that all have Li EWs significantly lower than the average. This color range corresponds to K2–K3 spectral types and is reminiscent of the Li dip observed in more evolved mainsequence stars (see, e.g., Balachandran 1995). One of these stars is HIP 84586, a SB2 spectroscopic binary whose tidal locking between the two components may have altered the internal mixing and, consequently, enhanced the Li depletion rate with respect to single stars. The other two stars are TYC 6878 0195 1 and HIP 11437A, both components of wide binaries that are expected to have evolved, rotationally, as single stars. We only note that HIP 11437A hosts a debris disc.

### 4. The rotation - Li depletion correlation

We first analyse the possible correlation between rotation and Li EW to investigate which role the rotation may play in producing the observed Li EW dispersion among stars with similar masses.

This information will be used for a more accurate modeling of the LBD to infer the association age.

We analyse separately the following three sub-groups: blue stars with  $0.5 \leq V-K_s < 3.4 \text{ mag}$  totaling 19 members, red stars with  $3.4 \leq V-K_s \leq 4.5 \text{ mag}$  totaling 21 members, and very red stars with  $5.4 < V-K_s < 5.9 \text{ mag}$ totaling 6 members that correspond to the onset of the Li depletion. Stars with  $V-K_s > 5.9 \text{ mag}$  have not undergone any Li depletion. In the blue stars sample, we observe a trend of increasing Li EW with increasing color (decreasing mass); in the red stars sample, we observe the opposite trend; in the very red star sample we observe again a trend

of increasing Li EW with increasing color. As it is shown in the top panel of Fig. 2, linear fits are enough for our purposes, i.e. approximating the mass dependence to outline the correlation with the rotation period. In the middle panel we plot the residuals with respect to the linear fits. Only one star (HIP 11437A), represented by a square and significantly deviating from the trend, was excluded from the fit. Positive residuals indicate stars that are more Li rich than the average, whereas negative residuals indicate stars that are more Li-depleted than the average. In the case of blue stars, we find a peak-to-peak amplitude of the dispersion  $\sigma = 90 \text{ mÅ}$  of the residuals from the fit, whereas in the case of red stars the peak-to-peak amplitude of the dispersion is about twice as large,  $\sigma = 180 \text{ mÅ}$ . Finally, in the case of very red stars we find  $\sigma = 125 \,\mathrm{mA}$ . The dotted lines represent peak-to-peak amplitudes of dispersions in the three color ranges. Now, to probe any dependence of the residual dispersion on the rotation period, we computed the Pearson linear correlation coefficients (r) and significance levels<sup>1</sup> ( $\alpha$ ) between the residuals and the rotation period as well as the Spearman's rank correlation (Spearman 1904). In the case of blue stars we find r =-0.19 with a significance level  $\alpha < 50\%$ , i.e. no dependence exists of the residuals on rotation (from Spearman's rank correlation we found  $\rho = -0.04$  with very small significance p-value = 0.77). On the contrary, for red stars we found

<sup>&</sup>lt;sup>1</sup> The significance level  $\alpha$  represents the probability of observing a value of the correlation coefficient larger than r for a random sample having the same number of observations and degrees of freedom (Bevington 1969)



Fig. 2. Top panel: distribution of Li EW versus V–K<sub>s</sub> color in three color ranges:  $0.5 < V-K_s < 3.4 \text{ mag}$  (19 stars),  $3.4 < V-K_s < 4.5 \text{ mag}$  (21 stars), and  $5.4 < V-K_s < 5.9 \text{ mag}$  (6 stars) of  $\beta$  Pictoris members with known rotation period and with overplotted (solid lines) the linear fits. *Middle panel*: residuals from the fit with horizontal dotted lines indicating the  $1\sigma$  dispersion. *Bottom panels*: distribution of Li EW residuals versus rotation period in the three color ranges. Solid lines are linear fits, r is the Pearson linear correlation coefficient and  $\alpha$  the significance level.

r = -0.71 with very high significance level  $\alpha > 99.9\%$ (similarly, from rank correlation we found  $\rho = -0.56$  with high significance p-value = 0.01). Finally, for very red stars, we find r = +0.13 with a significance level  $\alpha \sim 80\%$ (from Spearman's rank correlation we found  $\rho = 0.29$  with very small significance p-value = 0.58). Therefore, since in this regime of very red stars we have no enough data, we can not asses the existence or not of a rotation - Li depletion connection.

In the case of red stars  $(3.2 \leq V-K_s \leq 4.5 \text{ mag})$  we can state that the Li depletion is significantly correlated to rotation that seems to maintain fast rotators less depleted than slow rotators according to Eq. (1):

$$\delta EW(Li) = 101 \pm 22 - 17.7 \pm 3.4 \times P \tag{1}$$

where the Li EW is in mÅ and the rotation period P in days. The average uncertainty on the rotation period ( $\Delta P = 0.7\%$ ) is negligible with respect to that on the Li EW. After decorrelating the Li EW from rotation of red stars, the dispersion turns to be reduced by about a factor of 2,  $\sigma = 50$  mÅ and becomes to be comparable to that measured among blue stars, where no dependence on rotation was observed, and among very red stars. This residual dispersion is larger than the uncertainty associated to the Li EW measurements. Therefore, apart from mass and rotation, other causes must play a role in producing the observed dispersion. The intrinsic variability of the Li EW is certainly one cause. In fact, we know that spotted stars, as all our targets are, exhibit a rotational modulation of the Li EW. Specifically, the EW is larger at the rotation phases when the spots are best in view, and, consequently, the average surface temperature is lower. In our sample 26 stars have two EW measurements. We find that the average difference between these multiple measurements is  $<\Delta EW>$  = 26 mÅ . We note only two extreme cases: TX Psa and 2MASS J05082729-2101444 both very red stars with  $\Delta EW>$  100 mÅ.

Generally, the rotational Li variability in young stars is not larger than about 5% (see Appendix in Bouvier et al. 2016). This corresponds in our case to a range from  $\sim$ 5 to 25 mÅ in absolute values, which is as the same order of the intrinsic variability we found.

#### 4.1. Observed versus predicted LBD

It is interesting to make a comparison between the measured distribution of Li EWs and the predictions of evolutionary models. In the following, we first consider the evolutionary models of Baraffe et al. (2015) for solar abundances. Solar abundances have been measured in a few members, like  $\beta$  Pic and PZ Tel, and generally they were adopted for the whole association (see, e.g., Mentuch et al. 2008). The surface Lithium abundance is first derived from the ratio of surface Lithium abundance to initial abundance provided with the models using an initial abundance of Li<sub>0</sub>



Fig. 3. Left panel: distribution of Li EWs versus V–K<sub>s</sub> color for  $\beta$  Pic members. Solid lines represent the Li EWs predicted by the models of Baraffe et al. (2015) for ages in the range from 5 to 25 Myr. Right panel: same as in the left panel, but with models arbitrarily shifted by 0.7 mag towards redder colors to match with the observations.

=  $1 \times 10^{-9}$ . Then, to make the comparison with the observations, we transformed the model Li abundance into Li EW. For this purpose, we have used the curves of growth from Zapatero Osorio et al. (2002). They are valid in the effective temperature range 2600 < T<sub>eff</sub> < 4100 K and for 1.0 < A(Li) < 3.4.

In the left panel of Fig. 3, we plot the measured Li EW and compare this distribution with the Li EW derived from Baraffe et al. (2015) models. We note a strong mismatch between observations and model predictions. The models for ages in the 10-25 Myr range, predict a substantial/complete depletion in the  $V-K_s$  color range 3.8-4.5 mag, whereas the Li gap is observed at colors that are redder by about 0.7 mag. In fact, a better match is achieved if we arbitrarily shift the model colors by  $\Delta(V-K_s) =$  $+0.7 \,\mathrm{mag}$  (see right panel of Fig. 3). Moreover, assuming that the fast rotators have rotation-unaffected Li abundances we have maintained their observed Li EW, whereas for the slower rotators we have increased the Li EW according to Equation (1). As it is shown in the right panel of Fig. 3, we find that, even applying this  $V-K_s$  color shift and the rotation decorrelation, we can infer different ages by looking at different part of the observed Li depletion since the slope of the observed depletion is different than that predicted by the models.



Fig. 4. Absolute magnitude versus  $V-K_s$  color of members in the Li gap. Solid lines represent the isochrones corresponding to ages from 5 (top) to 25 Myr (bottom).

We note that the same mismatch between observations and models is found when different colors (e.g., V–I, J–H, or J-K) or the effective temperatures are used to make the comparison. The colors of the stars in the Li depletion gap  $4.4 < V-K_s < 5.4$  mag have been accurately checked. For a few stars (either single or resolved components in wide binaries) the  $V-K_s$  is measured for each component. For other stars (components of unresolved binaries) the  $V-K_s$  is derived from the spectral type of the primary component, which is the one with measured Li. In the first case the error on V-K<sub>s</sub> is  $\pm 0.03$  mag, in the latter case is  $\pm 0.15$  mag. In Fig. 4 we compare the absolute magnitudes of the members in the Li gap with isochrones for ages from 5 to 25 Myr taken from Baraffe et al. (2015). We note that within the gap  $(4.4 < V-K_s < 5.4 \text{ mag})$  the stars seem to be either more luminous or redder than expected for the most recent age estimation of 24 Myr by Bell et al. (2015). A better agreement is observed at bluer colors where stars are within the 10–25 Myr isochrones. If we correct the magnitudes of close components for binarity, we get a better agreement, but do not solve the overluminosity. This indicates that, in order to reproduce the data, the predicted Baraffe et al. (2015) colours should be shifted redward.

All stars in the gap have a prominent magnetic activity as it is inferred from photometric variability, X-ray emission and Near-UV/Far-UV excess. Therefore, models including the effect of magnetic fields should be more adequate to describe the Li depletion pattern. For this reason, we also made use of the Dartmouth stellar evolutionary models that incorporate the effects of magnetic fields (Feiden 2016); (Mann et al. 2016). In these models, inhibition of convection by magnetic fields cools the stellar surface temperature thereby slowing the contraction rate of young stars. Stars have a larger radius and a higher luminosity at a given age, as a result

These models were computed for solar metallicity and for an equipartition magnetic field strength in the range  $2500 < Bf_{eq} < 3000 G$ . Model effective temperatures were transformed into V–K<sub>s</sub> colors using the empirical T<sub>eff</sub>-V–K<sub>s</sub> relation from Pecaut & Mamajek (2013) valid for young 5–30 Myr stars. Li abundances were transformed into Li EW as done in the case of the Baraffe et al. models.



Fig. 5. Left panel: distribution of Li EWs versus  $V-K_s$  color for  $\beta$  Pic members. Solid lines represent the Li EWs predicted by the Dartmouth models for ages in the range from 20 to 30 Myr. Right panel: same as in the left panel, but Li EW decorrelated from rotation, assuming that fast rotators have rotation-unaffected Li EW.

In the left panel of Fig. 5, we plot the measured Li EW and compare this distribution with the Li EW derived from the Dartmouth et al. models. In this case, no arbitrary color shift is needed to match with the hot side of the Li gap. A better match is achieved in the right panel of Fig. 5, when the Li EW are decorrelated from rotation, assuming again that the fast rotators have rotation-unaffected Li abundances.

The reduced chi-squares<sup>2</sup> computed from the fit to the hot boundary of the Li gap have their minimum (see Fig. 6) for an age of  $25\pm3$  Myr. The uncertainty formally represents the age interval by which the  $\chi^2_{\nu}$  increases by one unity from the minimum value. The relatively large values of the  $\chi^2_{\nu}$  indicate either that the EW scatter arising from the intrinsic variability (rotational modulation) is larger than our estimate of 26 mÅ or the uncertainties associated to the measurements are in several case underestimated to some level.

Our age estimation is in good agreement with respect to the more recent age estimate of  $23\pm4$  Myr and  $24\pm3$  Myr by Mamajek & Bell (2014) and Bell et al. (2015), respectively, and significantly lower than the Macdonald & Mullan (2010) estimate.

#### 5. Conclusions

We have retrieved from the literature the measured Li EW of 66 members of the young  $\beta$  Pictoris association. We have carried out a photometric monitoring of these members that allowed us to measure for the first time the rotation periods of 30 members. For other 16 members we retrieved the rotation periods from Messina et al. (2010, 2011), and for the remaining members we used different sources in the literature.

We have explored the existence of a connection between rotation and Li depletion. After removing the mass dependence of the Li EW, using linear fits to EW versus V-K<sub>s</sub>, we found that for stars with  $0.5 \leq V-K_s \leq$ 3.4 mag, roughly corresponding to masses M > 0.8 M<sub>☉</sub>, no significant correlation is found between Li EW and rotation period. On the contrary, in the color range 3.4  $\leq$  V–K<sub>s</sub>  $\leq$  4.5 mag, roughly corresponding to masses 0.3 < M < 0.8 M $_{\odot}$ , we find a strong correlation between the Li EW and the rotation period, where fast rotators are much less Li-depleted than slow rotators. Finally, in the color range 5.4 < V–K<sub>s</sub> < 5.9 mag, roughly corresponding to masses M  $\sim$  0.1 M $_{\odot}$ , we have some hint for an inverted correlation where fast rotators are more depleted than slow rotators. However, this correlation is currently based on only 7 stars and the significance level is high (80%), but this is due to just one point that could be an outlier for whatever reason.

Interestingly, the dispersion in the  $3.4 \leq V-K_s \leq 4.5 \text{ mag}$ has a peak-to-peak amplitude that amounts to 180 mÅwhich is about a factor of 2 larger than that measured in the same mass range in the 5-Myr NGC 2264 open cluster (Bouvier et al. 2016), and about a factor 2 smaller than that measured in the 125-Myr Pleiades open cluster (Soderblom et al. 1993). Therefore, we note that the effect of rotation on the Li depletion is also age dependent and increases with age.

A comparison with the Baraffe et al. (2015) models, shows a mismatch of about  $0.7 \,\mathrm{mag}$  in the V-K<sub>s</sub> color between the observed and the predicted color range where the Li gap falls. Models predict the Li gap at bluer colors than observed. On the contrary, the Dartmouth models that incorporate the effects of magnetic fields provide a good match with the hot side of the Li depletion gap, although some mismatch on the cool side of the Li gap remains. Comparing the Dartmouth models with the hot side of the Li depletion gap, we infer an age of  $25\pm3$  Myr, which is in very good agreement with the most recent age estimates for the  $\beta$  Pictoris association. However, the relatively large values of the reduced chi-squares suggest that either the intrinsic Li EW variability is still underestimated or some other factor, apart from rotation, plays a relevant role in producing the observed scatter among stars with similar mass.

Acknowledgements. Research on stellar activity at INAF- Catania Astrophysical Observatory is supported by MIUR (Ministero dell'Istruzione, dell'Universitá e della Ricerca). This research has made use of the Simbad database, operated at CDS (Strasbourg, France). SM thanks Jerome Bouvier for useful discussion and the anony-

<sup>&</sup>lt;sup>2</sup> To compute the  $\chi^2_{\nu}$  we used the uncertainty  $\sigma = \sqrt{\sigma_0^2 + \sigma_v^2}$ where  $\sigma_0$  is the measurement error and  $\sigma_v$  is the average intrinsic variability.

Table 1. Properties of the 66 members of the  $\beta$  Pictoris association studied in this work

Target		RA		DEC			Sp.T	V	V-Ks	Р	Li EW
0	(hh, mm, ss)		n, ss)	$(^{\circ},  ',  '')$			*	(mag)	(mag)	(d)	(mÅ)
HIP 560	00	06	50.08	-23	06	27.20	F3V	6.15	0.910	0.224	87.0
TYC 1186-0706-1	00	23	34.66	+20	14	28.75	K7.5V+M5	10.96	3.623	7.900	338.0
GJ 2006A	00	27	50.23	-32	33	06.42	M3.5Ve	12.87	4.858	3.990	29.0
GJ 2000B	00	27	00.30 05 40	-32	33	23.80	M5.5Ve	13.10	0.044 6.050	4.910	20.0
2MASS J01112542+1526214A 2MASS J01351303 0712517	01	25	20.42 13.03	$^{+15}_{-07}$	20 12	21.00 51.77	$M_{0} + M_{0}$ $M_{4} = V$	14.40 13.22	0.202 5.502	0.911	593.0 46.7
TVC 1208-0468-1	01	37	39.42	$^{-07}_{\pm 18}$	35	32.91	K3+K5	9.85	3.114	2 803	440.0
HIP 10679	02	17	24.74	+28	44	30.43	G2V	7.75	1.488	0.777	160.0
HIP 10680	02	17	25.28	+28	44	42.16	F5V	6.95	1.163	0.240	140.0
HIP 11437A	02	27	29.25	+30	58	24.60	K4	10.12	3.040	12.500	248.0
HIP 11437B	02	27	28.05	+30	58	40.53	M1	12.44	4.219	4.660	220.0
HIP 12545	02	41	25.90	+05	59	18.00	K6Ve	10.37	3.301	4.830	450.0
GJ3305	04	37	37.30	-02	29	28.00	M1+M?	10.59	4.177	4.890	140.0
2MASS J04435686+3723033	04	43	56.87	+37	23	03.30	M3Ve+M5?	12.98	4.179	4.288	194.0
HIP 23200 HIP 23300	04	00	34.83 47.10	$+01 \\ -57$	47	25.00	MOVo	10.05	3.990 3.756	4.430	270.0
HIP 23418A	05	01	58 80	$\pm 09$	59	20.00	M3V	10.00 11.45	4.780	1 220	53.0
BD -211074A	05	06	49.90	-21	35	09.00	M1.5V	10.29	4.350	9.300	20.0
BD-211074B	05	06	49.90	-21	35	09.00	M2.5V	11.67	4.643	5.400	20.0
2MASS J05082729-2101444	05	08	27.30	-21	01	44.40	M5.6V	14.70	5.867	0.280	618.0
2MASS J05241914-1601153	05	24	19.15	-16	01	15.30	M4.5 + M5	13.50	5.603	0.401	217.0
HIP 25486	05	27	04.76	-11	54	03.47	F7V	6.22	1.294	0.966	191.0
2MASS J05335981-0221325	05	33	59.81	-02	21	32.50	M2.9V	12.42	4.725	7.250	49.0
2MASS J06131330-2742054	06	13	13.31	-27	42	05.50	M3.V:	12.09	5.230	16.9	28.0
HIP 29964	06	18	28.20	-72	02	41.00	K4Ve	9.80	2.986	2.670	420.0
1 WA 22 HID 76620	10	11	20.89	-03	54 49	26.50	MD KOV	13.99	0.301	0.830	202.0
2MASS 116430128-1754274	16	- JO - / 3	01.20	-37 -17	42 54	27.00 27.50	MOG	12 50	2.110	4.270 5.140	292.0
HIP 84586	17	17	25.50	-66	57	04.00	G5IV+K5IV	7.23	2.528	1.680	250.0
HD 155555C	17	17	31.29	-66	57	05.49	M3.5Ve	12.71	5.081	4.430	20.0
TYC 872822621	17	29	55.10	-54	15	49.00	K1V	9.55	2.186	1.830	360.0
GSC 08350-01924	17	29	20.67	-50	14	53.00	M3V	12.86	4.766	1.982	50.0
V4046Sgr	18	14	10.50	-32	47	33.00	K5+K7	10.44	3.191	2.420	440.0
2MASS J18151564-4927472	18	15	15.64	-49	27	47.20	M3V	12.86	4.780	0.447	46.0
HIP 89829	18	19	52.20	-29	16	33.00	GIV	8.89	1.837	0.571	290.0
2MASS J18202275-1011131A	18	20	22.74	-10	51	13.62	K5Ve+K7Ve	10.63	3.350	4.650	530.0 400.0
TYC 907724891 TYC 907307621	18	40	57.02 52.60	-62	10	36.00	M1Ve	9.30	3.204 3.946	5.340	490.0 332.0
HD 173167	18	48	06.36	-62	13	47.02	F5V	7.28	1.144	0.250	107.0
TYC 740800541	18	50	44.50	-31	47	47.00	K8Ve	11.20	3.660	1.075	492.0
HIP 92680	18	53	05.90	-50	10	50.00	K8Ve	8.29	1.924	0.944	287.0
TYC 687210111	18	58	04.20	-29	53	05.00	M0Ve	11.78	3.762	0.503	483.0
2MASS J19102820-2319486	19	10	28.21	-23	19	48.60	M4V	13.20	4.985	3.640	55.0
TYC 687801951	19	11	44.70	-26	04	09.00	K4Ve	10.27	2.904	5.650	320.0
2MASS J19233820-4606316	19	23	38.20	-46	06	31.60	MOV	11.87	3.598	3.242	422.0
1 Y C 744311021 2MASS 110560204 2207186AP	19	00 56	04.37	-32	07	37.71	MU.UV MAV	11.80	3.954 5.116	11.300	110.0
2MASS J19500294-5207180AB	20	01	37.18	-32 -33	13	14.01	M1	13.23 12.25	4 056	12700	100.0
2MASS J20015710-3515155	20	05	56.41	-32	16	59.15	M2:	11.96	4.022	8.368	140.0
HD 191089	20	09	05.21	-26	13	26.52	F5V	7.18	1.104	0.488	91.0
2MASS J20100002-2801410AB	20	10	00.03	-28	01	41.10	M2.5 + M3.5	12.80	4.640	0.470	46.0
2MASS J20333759-2556521	20	33	37.59	-25	56	52.20	M4.5	14.87	5.993	0.710	504.0
HIP 102141A	20	41	51.20	-32	26	07.00	M4Ve	10.36	5.416	1.191	20.0
HIP 102141B	20	41	51.10	-32	26	10.00	M4Ve	10.36	5.416	0.781	20.0
2MASS J20434114-2433534	20	43	41.14	-24	33	53.19	M3.7+M4.1	12.73	4.971	1.610	28.0
HIP 102409	20	45	09.50	-31	20	27.00	MIVe	8.73	4.201	4.860	80.0
HIP 103311 TVC 624002001	20	00 56	47.07	-17	10	51.04 54.00	F8V K6Vo I M2	10.69	1.539	0.300	147.0
2MASS 121100535-1010573	20	10	02.70	-17	10	57.40	M2V	10.02 11.54	1 3/4	3.410 3.710	420.0
2MASS J21100030-1010073	21	10	31.48	-27	10	57.80	M4.5V	14.90	5.600	0.650	502.0
TYC 9486-927-1	21	$25^{-5}$	27.49	-81	38	27.68	M2V	11.70	4.360	0.542	104.0
TYC 221113091	22	00	41.59	+27	15	13.60	M0V	11.39	3.666	1.109	40.0
TYC 934004371	22	42	48.90	-71	42	21.00	K7Ve	10.60	3.706	4.460	440.0
HIP 112312	22	44	58.00	-33	15	02.00	M4Ve	12.10	5.168	2.370	30.0
TX Psa	22	45	00.05	-33	15	25.80	M4.5Ve	13.36	5.567	1.080	450.0
TYC 583206661	23	32	30.90	-12	15	52.00	M0Ve	10.54	3.971	5.680	185.0

mous Referee for useful comments that helped us to improve this paper.

## References

- Balachandran, S. 1995, in The Light Element Abundances, ed.
- P. Crane Baraffe, I., Homeier, D., Allard, F., & Chabrier, G. 2015, A&A, 577, A42
- Bell, C. P. M., Mamajek, E. E., & Naylor, T. 2015, MNRAS, 454, 593 Bevington, P. R. 1969, Data reduction and error analysis for the physical sciences

Binks, A. S. & Jeffries, R. D. 2014, MNRAS, 438, L11

- Bouvier, J., Lanzafame, A. C., Venuti, L., et al. 2016, A&A, 590, A78
  Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, 2MASS All Sky Catalog of point sources.
- da Silva, L., Torres, C. A. O., de La Reza, R., et al. 2009, A&A, 508, 833
- Desidera, S., Covino, E., Messina, S., et al. 2015, A&A, 573, A126 Feiden, G. A. 2016, ArXiv: 1604.08036
- Gondoin, P. 2014, A&A, 566, A72
- Hartman, J. D., Bakos, G. Á., Kovács, G., & Noyes, R. W. 2010, MNRAS, 408, 475
- Kiss, L. L., Moór, A., Szalai, T., et al. 2011, MNRAS, 411, 117



Fig. 6. Reduced chi-squares versus age obtained from the residuals from the fit to the hot boundary of the Li depletion gap.

Lépine, S. & Simon, M. 2009, AJ, 137, 3632

- Macdonald, J. & Mullan, D. J. 2010, ApJ, 723, 1599
- Malo, L., Artigau, É., Doyon, R., et al. 2014a, ApJ, 788, 81
- Malo, L., Doyon, R., Feiden, G. A., et al. 2014b, ApJ, 792, 37
- Malo, L., Doyon, R., Lafrenière, D., et al. 2013, ApJ, 762, 88
- Mamajek, E. E. & Bell, C. P. M. 2014, MNRAS, 445, 2169
- Mann, A. W., Newton, E. R., Rizzuto, A. C., et al. 2016, ArXiv:1604.06165
- Mentuch, E., Brandeker, A., van Kerkwijk, M. H., Jayawardhana, R., & Hauschildt, P. H. 2008, ApJ, 689, 1127
- Messina, S., Desidera, S., Lanzafame, A. C., Turatto, M., & Guinan, E. F. 2011, A&A, 532, A10
- Messina, S., Desidera, S., Turatto, M., Lanzafame, A. C., & Guinan, E. F. 2010, A&A, 520, A15
- Moór, A., Szabó, G. M., Kiss, L. L., et al. 2013, MNRAS, 435, 1376
- Pecaut, M. J. & Mamajek, E. E. 2013, ApJS, 208, 9
- Pojmanski, G. 2002, Acta Astron., 52, 397
- Reid, I. N., Kirkpatrick, J. D., Liebert, J., et al. 2002, AJ, 124, 519
- Schlieder, J. E., Lépine, S., & Simon, M. 2010, AJ, 140, 119
- Schlieder, J. E., Lépine, S., & Simon, M. 2012, AJ, 143, 80
- Shkolnik, E. L., Anglada-Escudé, G., Liu, M. C., et al. 2012, ApJ, 758, 56
- Soderblom, D. R., Jones, B. F., Balachandran, S., et al. 1993, AJ, 106, 1059
- Spearman, C. 1904, The proof and measurement of association between two things, Vol. 15, 72–101
- Torres, C. A. O., Quast, G. R., da Silva, L., et al. 2006, A&A, 460, 695
- Zapatero Osorio, M. R., Béjar, V. J. S., Pavlenko, Y., et al. 2002, A&A, 384, 937
- Zuckerman, B. & Song, I. 2004, ARA&A, 42, 685