

# AN EXPLICIT VOLUME FORMULA FOR THE LINK $7_3^2(\alpha, \alpha)$ CONE-MANIFOLDS

JI-YOUNG HAM, JOONGUL LEE, ALEXANDER MEDNYKH\*, AND ALEKSEY RASSKAZOV

ABSTRACT. We calculate the volume of the  $7_3^2$  link cone-manifolds using the Schläfli formula. As an application, we give the volume of the cyclic coverings branched over the link.

## 1. INTRODUCTION

Let us denote the link complement of  $7_3^2$  in Rolfsen's link table by  $X$ . Note that it is a hyperbolic knot. Hence by Mostow-Prasad rigidity theorem,  $X$  has a unique hyperbolic structure. Let  $\rho_\infty$  be the holonomy representation from  $\pi_1(X)$  to  $\mathrm{PSL}(2, \mathbb{C})$  and denote  $\rho_\infty(\pi_1(X))$  by  $\Gamma$ , a Kleinian group.  $X$  is a  $(\mathrm{PSL}(2, \mathbb{C}), \mathbb{H}^3)$ -manifold and can be identified with  $\mathbb{H}^3/\Gamma$ . Thurston's orbifold theorem guarantees an orbifold,  $X(\alpha) = X(\alpha, \alpha)$ , with underlying space  $S^3$  and with the link  $7_3^2$  as the singular locus of the cone-angle  $\alpha = 2\pi/k$  for some nonzero integer  $k$ , can be identified with  $\mathbb{H}^3/\Gamma'$  for some  $\Gamma' \in \mathrm{PSL}(2, \mathbb{C})$ ; the hyperbolic structure of  $X$  is deformed to the hyperbolic structure of  $X(\alpha)$ . For the intermediate angles whose multiples are not  $2\pi$  and not bigger than  $\pi$ , Kojima [10] showed that the hyperbolic structure of  $X(\alpha)$  can be obtained uniquely by deforming nearby orbifold structures. Note that there exists an angle  $\alpha_0 \in [\frac{2\pi}{3}, \pi)$  for the link  $7_3^2$  such that  $X(\alpha)$  is hyperbolic for  $\alpha \in (0, \alpha_0)$ , Euclidean for  $\alpha = \alpha_0$ , and spherical for  $\alpha \in (\alpha_0, \pi]$  [19, 8, 10, 20]. For further knowledge of cone-manifolds a reader can consult [1, 7].


Even though we have wide discussions on orbifolds, it seems to us we have a little in regard to cone-manifolds. Explicit volume formulae for hyperbolic cone-manifolds of knots and links are known a little. The volume formulae for hyperbolic cone-manifolds of the knot  $4_1$  [8, 10, 11, 15], the knot  $5_2$  [13], the link  $5_1^2$  [16], the link  $6_2^2$  [17], and the link  $6_3^2$  [2] have been computed. In [9] a method of calculating the volumes of two-bridge knot cone-manifolds was introduced but without explicit formulae. In [7, 6], explicit volume formulae of cone-manifolds for the hyperbolic twist knot and for the knot with Conway notation  $C(2n, 3)$  are computed. Similar methods are used for computing Chern-Simons invariants of orbifolds for the twist knot and  $C(2n, 3)$  knot in [5, 4].

The main purpose of the paper is to find an explicit and efficient volume formula of hyperbolic cone-manifolds for the link  $7_3^2$ . The following theorem gives the volume formula for  $X(\alpha)$ .

**Theorem 1.1.** *Let  $X(\alpha)$ ,  $0 \leq \alpha < \alpha_0$  be the hyperbolic cone-manifold with underlying space  $S^3$  and with singular set the link  $7_3^2$  of cone-angle  $\alpha$ .  $X(0)$  denotes  $X$ . Then the volume of  $X(\alpha)$  is given by the following formula*

---

\*The author was funded by the Russian Science Foundation (grant 16-41-02006).

FIGURE 1. Link  $7_3^2$  in Rolfsen's link table

$$\text{Vol}(X(\alpha)) = \int_{\alpha}^{\pi} 2 \log \left| \frac{A - iV}{A + iV} \right| d\alpha,$$

where for  $A = \cot \frac{\alpha}{2}$ ,  $V$  ( $\text{Re}(V) \leq 0$  and  $\text{Im}(V) \geq 0$  is the largest) is a zero of the Riley-Mednykh polynomial  $P = P(V, A)$  for the link  $7_3^2$  given below.

$$\begin{aligned} P &= 8V^5 + 8A^2V^4 + (8A^4 + 16A^2 - 8)V^3 + (4A^6 + 8A^4 - 12A^2)V^2 \\ &+ (A^8 + 4A^6 - 2A^4 - 12A^2 + 1)V - 4A^6 - 8A^4 + 4A^2. \end{aligned}$$

The following corollary gives the hyperbolic volume of the  $k$ -fold strictly-cyclic covering [12, 18] over the link  $7_3^2$ ,  $M_k(X)$ , for  $k \geq 3$ .

**Corollary 1.2.** *The volume of  $M_k(X)$  is given by the following formula*

$$\text{Vol}(M_k(X)) = k \int_{\frac{2\pi}{k}}^{\pi} 2 \log \left| \frac{A - iV}{A + iV} \right| d\alpha,$$

where for  $A = \cot \frac{\alpha}{2}$ ,  $V$  ( $\text{Re}(V) \leq 0$  and  $\text{Im}(V) \geq 0$  is the largest) is a zero of the Riley-Mednykh polynomial  $P = P(V, A)$  for the link  $7_3^2$ .

In Section 2, we present the fundamental group  $\pi_1(X)$  of  $X$  with slope  $9/16$ . In Section 3, we give the defining equation of the representation variety of  $\pi_1(X)$ . In Section 4, we compute the longitude of the link  $7_3^2$  using the Pythagorean theorem. And in Section 5, we give the proof of Theorem 1.1 using the Schläfli formula.

## 2. LINK $7_3^2$

Link  $7_3^2$  is presented in Figure 1. It is the same as  $W_3$  from [2]. The slope of this link is  $7/16$ . The link with slope  $9/16$  is the mirror of the link  $7_3^2$ . Since the volume of the link with slope  $7/16$  is the same as the volume of link with slope  $9/16$ , in the rest of the paper, the link with slope  $9/16$  is used.

The following fundamental group of  $X$  is stated in [2] with slope  $7/16$ .

**Proposition 2.1.**

$$\pi_1(X) = \langle s, t \mid s w s^{-1} w^{-1} = 1 \rangle,$$

where  $w = s^{-1}[s, t]^2[s, t^{-1}]^2$ .

### 3. $(\mathrm{PSL}(2, \mathbb{C}), \mathbb{H}^3)$ STRUCTURE OF $X(\alpha)$

Let  $R = \mathrm{Hom}(\pi_1(X), \mathrm{SL}(2, \mathbb{C}))$ . Given a set of generators,  $s, t$ , of the fundamental group for  $\pi_1(X)$ , we define a set  $R(\pi_1(X)) \subset \mathrm{SL}(2, \mathbb{C})^2 \subset \mathbb{C}^8$  to be the set of all points  $(h(s), h(t))$ , where  $h$  is a representation of  $\pi_1(X)$  into  $\mathrm{SL}(2, \mathbb{C})$ . Since the defining relation of  $\pi_1(X)$  gives the defining equation of  $R(\pi_1(X))$  [21],  $R(\pi_1(X))$  is an affine algebraic set in  $\mathbb{C}^8$ .  $R(\pi_1(X))$  is well-defined up to isomorphisms which arise from changing the set of generators. We say elements in  $R$  which differ by conjugations in  $\mathrm{SL}(2, \mathbb{C})$  are *equivalent*. A point on the variety gives the  $(\mathrm{PSL}(2, \mathbb{C}), \mathbb{H}^3)$  structure of  $X(\alpha)$ .

Let

$$h(s) = \begin{bmatrix} \cos \frac{\alpha}{2} & ie^{\frac{\rho}{2}} \sin \frac{\alpha}{2} \\ ie^{-\frac{\rho}{2}} \sin \frac{\alpha}{2} & \cos \frac{\alpha}{2} \end{bmatrix}, \quad h(t) = \begin{bmatrix} \cos \frac{\alpha}{2} & ie^{-\frac{\rho}{2}} \sin \frac{\alpha}{2} \\ ie^{\frac{\rho}{2}} \sin \frac{\alpha}{2} & \cos \frac{\alpha}{2} \end{bmatrix}.$$

Then  $h$  becomes a representation if and only if  $A = \cot \frac{\alpha}{2}$  and  $V = \cosh \rho$  satisfies a polynomial equation [21, 14]. We call the defining polynomial of the algebraic set  $\{(V, A)\}$  as the *Riley-Mednykh polynomial* for the link  $7_3^2$ . Throughout the paper,  $h$  can be sometimes any representation and sometimes the unique hyperbolic representation.

Given the fundamental group of  $X$ ,

$$\pi_1(X) = \langle s, t \mid sws^{-1}w^{-1} = 1 \rangle,$$

where  $w = s^{-1}[s, t]^2[s, t^{-1}]^2$ , let  $S = h(s)$ ,  $T = h(t)$  and  $W = h(w)$ . Then the trace of  $S$  and the trace of  $T$  are both  $2 \cos \frac{\alpha}{2}$ .

**Lemma 3.1.** *For  $n \in \mathrm{SL}(2, \mathbb{C})$  which satisfies  $nS = S^{-1}n$ ,  $nT = T^{-1}n$ , and  $n^2 = -I$ ,*

$$SWS^{-1}W^{-1} = -(SWn)^2.$$

*Proof.*

$$\begin{aligned} (SWn)^2 &= SWnSWn = SWS^{-1}n(S^{-1}(STS^{-1}T^{-1})^2(ST^{-1}S^{-1}T)^2)n \\ &= SWS^{-1}(S(S^{-1}T^{-1}ST)^2(S^{-1}TST^{-1})^2)n^2 = -SWS^{-1}W^{-1}. \end{aligned}$$

□

From the structure of the algebraic set of  $R(\pi_1(X))$  with coordinates  $h(s)$  and  $h(t)$  we have the defining equation of  $R(\pi_1(X))$ . The following theorem is stated in [2, Proposition 4] with slope  $7/16$ .

**Theorem 3.2.**  *$h$  is a representation of  $\pi_1(X)$  if  $V$  is a root of the following Riley-Mednykh polynomial  $P = P(V, A)$  which is given below.*

$$\begin{aligned} P &= 8V^5 + 8A^2V^4 + (8A^4 + 16A^2 - 8)V^3 + (4A^6 + 8A^4 - 12A^2)V^2 \\ &\quad + (A^8 + 4A^6 - 2A^4 - 12A^2 + 1)V - 4A^6 - 8A^4 + 4A^2. \end{aligned}$$

*Proof.* Note that  $SWS^{-1}W^{-1} = I$ , which gives the defining equations of  $R(\pi_1(X))$ , is equivalent to  $(SWn)^2 = -I$  in  $\mathrm{SL}(2, \mathbb{C})$  by Lemma 3.1 and  $(SWn)^2 = -I$  in  $\mathrm{SL}(2, \mathbb{C})$  is equivalent to  $\mathrm{tr}(SWn) = 0$ .

We can find two  $n$ 's in  $\mathrm{SL}(2, \mathbb{C})$  which satisfies  $nS = S^{-1}n$  and  $n^2 = -I$  by direct computations. The existence and the uniqueness of the isometry (the involution) which is represented by  $n$  are shown in [3, p. 46]. Since two  $n$ 's give the same element in  $\mathrm{PSL}(2, \mathbb{C})$ , we use one of them. Hence, we may assume

$$n = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix},$$

$$S = \begin{bmatrix} \cos \frac{\alpha}{2} & ie^{\frac{\rho}{2}} \sin \frac{\alpha}{2} \\ ie^{-\frac{\rho}{2}} \sin \frac{\alpha}{2} & \cos \frac{\alpha}{2} \end{bmatrix}, \quad T = \begin{bmatrix} \cos \frac{\alpha}{2} & ie^{-\frac{\rho}{2}} \sin \frac{\alpha}{2} \\ ie^{\frac{\rho}{2}} \sin \frac{\alpha}{2} & \cos \frac{\alpha}{2} \end{bmatrix}.$$

Recall that  $P$  is the defining polynomial of the algebraic set  $\{(V, A)\}$  and the defining polynomial of  $R(\pi_1(X))$  corresponding to our choice of  $h(s)$  and  $h(t)$ . By direct computation  $P$  is a factor of  $\mathrm{tr}(SWn) = -4i \sinh \rho (2V^2 + A^4 + 2A^2 - 1)P$ . As in [2],  $P$  can not be  $\sinh \rho$  or have only real roots. Also,  $P$  can not have only purely imaginary roots similarly.  $P$  in the theorem is the only factor of  $\mathrm{tr}(SWn)$  which is different from  $\sinh \rho$  and has roots which are not real or purely imaginary.  $P$  is the Riley-Mednykh polynomial.  $\square$

#### 4. LONGITUDE

Let  $l_s = ws$  and  $l_t = (t^{-1}[t, s]^2[t, s^{-1}]^2)t$ . Then  $l_s$  and  $l_t$  are the longitudes which are null-homologous in  $X$ . Let  $L_S = h(l_s)$  and Let  $L_T = h(l_t)$ .

**Lemma 4.1.**  $\mathrm{tr}(S^{-1}L_T) = \mathrm{tr}(S)$  and  $\mathrm{tr}(T^{-1}L_S) = \mathrm{tr}(T)$ .

*Proof.* Since

$$\begin{aligned} S^{-1}L_T &= S^{-1}(T^{-1}(TST^{-1}S^{-1}TST^{-1}S^{-1} \cdot TS^{-1}T^{-1}STS^{-1}T^{-1}S)T) \\ &= (T^{-1}S^{-1}TST^{-1}S^{-1}T)(S^{-1})(T^{-1}S^{-1}TST^{-1}S^{-1}T)^{-1}, \end{aligned}$$

$$\mathrm{tr}(S^{-1}L_T) = \mathrm{tr}(S^{-1}) = \mathrm{tr}(S).$$

The second statement can be obtained in a similar way.  $\square$

**Definition.** The *complex length* of the longitude  $l$  ( $l_s$  or  $l_t$ ) of the link  $7_3^2$  is the complex number  $\gamma_\alpha$  modulo  $4\pi\mathbb{Z}$  satisfying

$$\mathrm{tr}(h(l)) = 2 \cosh \frac{\gamma_\alpha}{2}.$$

Note that  $l_\alpha = |\mathrm{Re}(\gamma_\alpha)|$  is the real length of the longitude of the cone-manifold  $X(\alpha)$ .

By sending common fixed points of  $T$  and  $L_T = h(l_t)$  to 0 and  $\infty$ , we have

$$T = \begin{bmatrix} e^{\frac{i\alpha}{2}} & 0 \\ 0 & e^{-\frac{i\alpha}{2}} \end{bmatrix}, \quad L_T = \begin{bmatrix} e^{\frac{\gamma_\alpha}{2}} & 0 \\ 0 & e^{-\frac{\gamma_\alpha}{2}} \end{bmatrix},$$

and the following normalized line matrices of  $T$  (resp.  $L_T$ ) which share the fixed points with  $T$  (resp.  $L_T$ ).

$$\begin{aligned} l(T) &\equiv \frac{T - T^{-1}}{2i \sinh \frac{i\alpha}{2}} \\ &= \frac{1}{i(e^{\frac{i\alpha}{2}} - e^{-\frac{i\alpha}{2}})} \begin{bmatrix} e^{\frac{i\alpha}{2}} - e^{-\frac{i\alpha}{2}} & 0 \\ 0 & e^{-\frac{i\alpha}{2}} - e^{\frac{i\alpha}{2}} \end{bmatrix} \\ &= \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}, \end{aligned}$$

$$\begin{aligned} l(L_T) &\equiv \frac{L_T - L_T^{-1}}{2i \sinh \frac{\gamma\alpha}{2}} \\ &= \frac{1}{i(e^{\frac{\gamma\alpha}{2}} - e^{-\frac{\gamma\alpha}{2}})} \begin{bmatrix} e^{\frac{\gamma\alpha}{2}} - e^{-\frac{\gamma\alpha}{2}} & 0 \\ 0 & e^{-\frac{\gamma\alpha}{2}} - e^{\frac{\gamma\alpha}{2}} \end{bmatrix} \\ &= \begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}, \end{aligned}$$

which give the orientations of axes of  $T$  and  $L_T$ .

Now, we are ready to prove the following theorem which gives Theorem 4.3. Recall that  $\gamma_\alpha$  modulo  $4\pi\mathbb{Z}$  is the *complex length* of the longitude  $l_s$  or  $l_t$  of  $X(\alpha)$ . The following theorem is a particular case of Proposition 5 from [2].

**Theorem 4.2.** (*Pythagorean Theorem*) [2] *Let  $X(\alpha)$  be a hyperbolic cone-manifold and let  $\rho$  be the complex distance between the oriented axes  $S$  and  $T$ . Then we have*

$$i \cosh \rho = \cot \frac{\alpha}{2} \coth \left( \frac{\gamma\alpha}{4} \right).$$

*Proof.*

$$\begin{aligned}
\cosh \rho &= -\frac{\operatorname{tr}(l(S)l(T))}{2} \\
&= -\frac{\operatorname{tr}(l(S)l(L_T))}{2} \\
&= \frac{\operatorname{tr}((S - S^{-1})(L_T - L_T^{-1}))}{8 \sinh \frac{i\alpha}{2} \sinh \frac{\gamma\alpha}{2}} \\
&= \frac{\operatorname{tr}(SL_T - S^{-1}L_T - SL_T^{-1} + (L_T S)^{-1})}{8 \sinh \frac{i\alpha}{2} \sinh \frac{\gamma\alpha}{2}} \\
&= \frac{2(\operatorname{tr}(SL_T) - \operatorname{tr}(S^{-1}L_T))}{8 \sinh \frac{i\alpha}{2} \sinh \frac{\gamma\alpha}{2}} \\
&= \frac{\operatorname{tr}(S)\operatorname{tr}(L_T) - 2\operatorname{tr}(S^{-1}L_T)}{4 \sinh \frac{i\alpha}{2} \sinh \frac{\gamma\alpha}{2}} \\
&= \frac{\operatorname{tr}(S)\operatorname{tr}(L_T) - 2\operatorname{tr}(S)}{4 \sinh \frac{i\alpha}{2} \sinh \frac{\gamma\alpha}{2}} \\
&= \frac{\operatorname{tr}(S)(\operatorname{tr}(L_T) - 2)}{4 \sinh \frac{i\alpha}{2} \sinh \frac{\gamma\alpha}{2}} \\
&= \frac{2 \cos \frac{\alpha}{2} (2 \cosh \frac{\gamma\alpha}{2} - 2)}{4i \sin \frac{\alpha}{2} \sinh \frac{\gamma\alpha}{2}} \\
&= -i \cot \frac{\alpha}{2} \tanh\left(\frac{\gamma\alpha}{4}\right).
\end{aligned}$$

where the first equality comes from [3, p. 68], the sixth equality comes from the Cayley-Hamilton theorem, and the seventh equality comes from Lemma 4.1. Therefore, we have

$$i \cosh \rho = \cot \frac{\alpha}{2} \coth\left(\frac{\gamma\alpha}{4}\right).$$

□

Pythagorean theorem 4.2 gives the following theorem which relates the eigenvalues of  $h(l)$  and  $V = \cosh \rho$  for  $A = \cot \frac{\alpha}{2}$ .

**Theorem 4.3.** *Recall that  $l$  is the longitude. By conjugating if necessary, we may assume  $h(l)$  is upper triangular. Let  $L = h(l)_{11}$ . Let  $A = \cot \frac{\alpha}{2}$ . Then the following formulae show that there is a one to one correspondence between the the eigenvalues of  $h(l)$  and  $V = \cosh \rho$ :*

$$iV = A \frac{L - 1}{L + 1} \text{ and } L = \frac{A - iV}{A + iV}.$$

*Proof.* By Theorem 4.2,

$$\begin{aligned}
iV &= i \cosh \rho \\
&= \cot \frac{\alpha}{2} \tanh\left(\frac{\gamma\alpha}{4}\right) \\
&= \cot \frac{\alpha}{2} \frac{\sinh\left(\frac{\gamma\alpha}{4}\right)}{\cosh\left(\frac{\gamma\alpha}{4}\right)} \\
&= \cot \frac{\alpha}{2} \frac{e^{\frac{\gamma\alpha}{4}} - e^{-\frac{\gamma\alpha}{4}}}{e^{\frac{\gamma\alpha}{4}} + e^{-\frac{\gamma\alpha}{4}}} \\
&= \cot \frac{\alpha}{2} \frac{e^{\frac{\gamma\alpha}{2}} - 1}{e^{\frac{\gamma\alpha}{2}} + 1} \\
&= A \frac{L - 1}{L + 1}.
\end{aligned}$$

If we solve the above equation,

$$iV = A \frac{L - 1}{L + 1},$$

for  $L$ , we have

$$L = \frac{A - iV}{A + iV}.$$

□

## 5. PROOF OF THEOREM 1.1

According to [19, 8, 10, 20], there exists an angle  $\alpha_0 \in [\frac{2\pi}{3}, \pi)$  such that  $X(\alpha)$  is hyperbolic for  $\alpha \in (0, \alpha_0)$ , Euclidean for  $\alpha = \alpha_0$ , and spherical for  $\alpha \in (\alpha_0, \pi)$ . Denote by  $D(X(\alpha))$  the discriminant of  $P(V, A)$  over  $V$ . Then  $\alpha_0$  is the only zero of  $D(X(\alpha))$  in  $[\frac{2\pi}{3}, \pi)$ .

From Theorem 4.3, we have the following equality,

$$(1) \quad |L|^2 = \left| \frac{A - iV}{A + iV} \right|^2 = \frac{|A|^2 + |V|^2 + 2A \operatorname{Im} V}{|A|^2 + |V|^2 - 2A \operatorname{Im} V}.$$

For the volume, we choose  $L$  with  $|L| \geq 1$  and hence we have  $\operatorname{Im}(V) \geq 0$  by Equality (1). The component of  $V$  with  $\operatorname{Im}(V) \geq 0$  which becomes real at  $\alpha_0$  has negative real part. On the geometric component which gives the unique hyperbolic structure, we have the

volume of a hyperbolic cone-manifold  $X(\alpha)$  for  $0 \leq \alpha < \alpha_0$ :

$$\begin{aligned} \text{Vol}(X(\alpha)) &= - \int_{\alpha_0}^{\alpha} 2 \left( \frac{l_{\alpha}}{2} \right) d\alpha \\ &= - \int_{\alpha_0}^{\alpha} 2 \log |L| d\alpha \\ &= - \int_{\pi}^{\alpha} 2 \log |L| d\alpha \\ &= \int_{\alpha}^{\pi} 2 \log |L| d\alpha \\ &= \int_{\alpha}^{\pi} 2 \log \left| \frac{A - iV}{A + iV} \right| d\alpha, \end{aligned}$$

where the first equality comes from the Schläfli formula for cone-manifolds (Theorem 3.20 of [1]), the second equality comes from the fact that  $l_{\alpha} = |Re(\gamma_{\alpha})|$  is the real length of the one longitude of  $X(\alpha)$ , the third equality comes from the fact that  $\log |L| = 0$  for  $\alpha_0 < \alpha \leq \pi$  by Equality (1) since all  $V$ 's are real for  $\alpha_0 < \alpha \leq \pi$ , and  $\alpha_0 \in [\frac{2\pi}{3}, \pi)$  is the zero of the discriminant  $D(X(\alpha))$ . Numerical calculations give us the following value for  $\alpha_0$ :  $\alpha_0 \approx 2.83003$ .

## REFERENCES

- [1] Daryl Cooper, Craig D. Hodgson, and Steven P. Kerckhoff. *Three-dimensional orbifolds and cone-manifolds*, volume 5 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2000. With a postface by Sadayoshi Kojima.
- [2] D. Derevnin, A. Mednykh, and M. Mulazzani. Volumes for twist link cone-manifolds. *Bol. Soc. Mat. Mexicana (3)*, 10(Special Issue):129–145, 2004.
- [3] Werner Fenchel. *Elementary geometry in hyperbolic space*, volume 11 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1989. With an editorial by Heinz Bauer.
- [4] Ji-Young Ham and Joongul Lee. Explicit formulae for Chern-Simons invariants of the hyperbolic orbifolds of the knot with Conway's notation  $C(2n, 3)$ . [www.math.snu.ac.kr/~jyham](http://www.math.snu.ac.kr/~jyham), 2016. Preprint.
- [5] Ji-Young Ham and Joongul Lee. Explicit formulae for Chern-Simons invariants of the twist knot orbifolds and edge polynomials of twist knots. *Matematicheskii Sbornik*, 207 (9), 2016.
- [6] Ji-Young Ham and Joongul Lee. The volume of hyperbolic cone-manifolds of the knot with Conway's notation  $C(2n, 3)$ . *J. Knot Theory Ramifications*, 25(6):1650030, 9, 2016.
- [7] Ji-Young Ham, Alexander Mednykh, and Vladimir Petrov. Trigonometric identities and volumes of the hyperbolic twist knot cone-manifolds. *J. Knot Theory Ramifications*, 23(12):1450064, 16, 2014.
- [8] Hugh Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. On a remarkable polyhedron geometrizing the figure eight knot cone manifolds. *J. Math. Sci. Univ. Tokyo*, 2(3):501–561, 1995.
- [9] Hugh M. Hilden, María Teresa Lozano, and José María Montesinos-Amilibia. Volumes and Chern-Simons invariants of cyclic coverings over rational knots. In *Topology and Teichmüller spaces (Katinkulta, 1995)*, pages 31–55. World Sci. Publ., River Edge, NJ, 1996.
- [10] Sadayoshi Kojima. Deformations of hyperbolic 3-cone-manifolds. *J. Differential Geom.*, 49(3):469–516, 1998.
- [11] Sadayoshi Kojima. Hyperbolic 3-manifolds singular along knots. *Chaos Solitons Fractals*, 9(4-5):765–777, 1998. Knot theory and its applications.
- [12] John P. Mayberry and Kunio Murasugi. Torsion-groups of abelian coverings of links. *Trans. Amer. Math. Soc.*, 271(1):143–173, 1982.



- [13] Alexander Mednykh. The volumes of cone-manifolds and polyhedra. <http://mathlab.snu.ac.kr/~top/workshop01.pdf>, 2007. Lecture Notes, Seoul National University.
- [14] Alexander Mednykh and Aleksey Rasskazov. On the structure of the canonical fundamental set for the 2-bridge link orbifolds. [www.mathematik.uni-bielefeld.de/sfb343/preprints/pr98062.ps.gz](http://www.mathematik.uni-bielefeld.de/sfb343/preprints/pr98062.ps.gz), 1998. Universitt Bielefeld, Sonderforschungsbereich 343, Discrete Strukturen in der Mathematik, Preprint, 98062.
- [15] Alexander Mednykh and Alexey Rasskazov. Volumes and degeneration of cone-structures on the figure-eight knot. *Tokyo J. Math.*, 29(2):445–464, 2006.
- [16] Alexander Mednykh and Andrei Vesnin. On the volume of hyperbolic Whitehead link cone-manifolds. *SCIENTIA, Series A: Sci. Ser. A Math. Sci. (N.S.)*, 8:1–11, 2002.
- [17] Alexander D. Mednykh. Trigonometric identities and geometrical inequalities for links and knots. In *Proceedings of the Third Asian Mathematical Conference, 2000 (Diliman)*, pages 352–368. World Sci. Publ., River Edge, NJ, 2002.
- [18] Michele Mulazzani and Andrei Vesnin. The many faces of cyclic branched coverings of 2-bridge knots and links. *Atti Sem. Mat. Fis. Univ. Modena*, 49(suppl.):177–215, 2001. Dedicated to the memory of Professor M. Pezzana (Italian).
- [19] Joan Porti. Spherical cone structures on 2-bridge knots and links. *Kobe J. Math.*, 21(1-2):61–70, 2004.
- [20] Joan Porti and Hartmut Weiss. Deforming Euclidean cone 3-manifolds. *Geom. Topol.*, 11:1507–1538, 2007.
- [21] Robert Riley. Nonabelian representations of 2-bridge knot groups. *Quart. J. Math. Oxford Ser. (2)*, 35(138):191–208, 1984.

DEPARTMENT OF SCIENCE, HONGIK UNIVERSITY, 94 WAUSAN-RO, MAPO-GU, SEOUL, 04066, KOREA.

*E-mail address:* [jiyoungnam1@gmail.com](mailto:jiyoungnam1@gmail.com)

DEPARTMENT OF MATHEMATICS EDUCATION, HONGIK UNIVERSITY, 94 WAUSAN-RO, MAPO-GU, SEOUL, 04066, KOREA.

*E-mail address:* [jglee@hongik.ac.kr](mailto:jglee@hongik.ac.kr)

SOBOLEV INSTITUTE OF MATHEMATICS, PR. KOTYUGA 4, NOVOSIBIRSK 630090, NOVOSIBIRSK STATE UNIVERSITY, PIROGOVA 2, NOVOSIBIRSK 630090, RUSSIA.

*E-mail address:* [mednykh@math.nsc.ru](mailto:mednykh@math.nsc.ru)

WEBSTER INTERNATIONAL UNIVERSITY, 146 MOO 5, TAMBON SAM PHRAYA, CHA-AM, PHETCHABURI 76120, TAILAND

*E-mail address:* [arasskazov69@webster.edu](mailto:arasskazov69@webster.edu)