LOGARITHMIC COEFFICIENTS OF SOME CLOSE-TO-CONVEX FUNCTIONS

MD FIROZ ALI AND A. VASUDEVARAO

ABSTRACT. The logarithmic coefficients γ_n of an analytic and univalent function f in the unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ with the normalization f(0) = 0 = f'(0) - 1 are defined by $\log \frac{f(z)}{z} = 2 \sum_{n=1}^{\infty} \gamma_n z^n$. In the present paper, we consider close-to-convex functions (with argument 0) with respect to odd starlike functions and determine the sharp upper bound of $|\gamma_n|$, n = 1, 2, 3 for such functions f.

1. INTRODUCTION

Let \mathcal{A} denote the class of analytic functions f in the unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ normalized by f(0) = 0 = f'(0) - 1. Any function f in \mathcal{A} has the following power series representation

(1.1)
$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

The class of univalent (i.e. one-to-one) functions in \mathcal{A} is denoted by \mathcal{S} . A function $f \in \mathcal{A}$ is called starlike (convex respectively) if $f(\mathbb{D})$ is starlike with respect to the origin (convex respectively). Let \mathcal{S}^* and \mathcal{C} denote the class of starlike and convex functions in \mathcal{S} respectively. It is well-known that a function $f \in \mathcal{A}$ is in \mathcal{S}^* if and only if Re (zf'(z)/f(z)) > 0 for $z \in \mathbb{D}$. Similarly, a function $f \in \mathcal{A}$ is in \mathcal{C} if and only if Re (1 + zf''(z)/f'(z)) > 0 for $z \in \mathbb{D}$. From the above it is easy to see that $f \in \mathcal{C}$ if and only if $zf' \in \mathcal{S}^*$. Given $\alpha \in (-\pi/2, \pi/2)$ and $g \in \mathcal{S}^*$, a function $f \in \mathcal{A}$ is said to be close-to-convex with argument α and with respect to g if

Let $\mathcal{K}_{\alpha}(g)$ denote the class of all such functions. Let

$$\mathcal{K}(g) := \bigcup_{\alpha \in (-\pi/2, \pi/2)} \mathcal{K}_{\alpha}(g) \text{ and } \mathcal{K}_{\alpha} := \bigcup_{g \in \mathcal{S}^*} \mathcal{K}_{\alpha}(g)$$

be the classes of functions called close-to-convex functions with respect to g and close-to-convex functions with argument α , respectively. The class

$$\mathcal{K} := \bigcup_{\alpha \in (-\pi/2, \pi/2)} \mathcal{K}_{\alpha} = \bigcup_{g \in \mathcal{S}^*} \mathcal{K}(g)$$

File: FV_ADM_2016.tex, printed: 2021-8-23, 20.15

²⁰¹⁰ Mathematics Subject Classification. Primary 30C45, 30C55.

Key words and phrases. Analytic, univalent, starlike, convex, close-to-convex functions, logarithmic coefficient.

is the class of all close-to-convex functions. It is well-known that every close-toconvex function is univalent in \mathbb{D} (see [5]). Geometrically, $f \in \mathcal{K}$ means that the complement of the image-domain $f(\mathbb{D})$ is the union of non-intersecting half-lines.

For the function $f \in S$, the logarithmic coefficients $\gamma_n (n = 1, 2, ...)$ are defined by

(1.3)
$$\log \frac{f(z)}{z} = 2 \sum_{n=1}^{\infty} \gamma_n z^n, \quad z \in \mathbb{D}.$$

I.E. Bazilevich first noticed that the logarithmic coefficients are very essential in the coefficient problem of univalent functions. He estimated (see [2, 3]) in depending upon the positive Hayman constant (see [10]) how close the coefficients γ_n (n = 1, 2, ...) of the functions of class S are to the relative logarithmic coefficients of the Koebe function $k(z) = z/(1-z)^2$. He also estimated the value $\sum_{n=1}^{\infty} n |\gamma_n|^2 r^{2n}$ which after multiplication by π is equal to the area of the image of the disk |z| < r < 1 under the function $\frac{1}{2} \log(f(z)/z)$ for $f \in S$. The celebrated de Branges' inequalities (the former Milin conjecture) for univalent functions f state that

$$\sum_{k=1}^{n} (n-k+1) |\gamma_k|^2 \le \sum_{k=1}^{n} \frac{n+1-k}{k}, \quad n = 1, 2, \dots,$$

with equality if and only if $f(z) = e^{-i\theta}k(e^{i\theta}z), \theta \in \mathbb{R}$ (see [4]). De Branges [4] used this inequality to prove the celebrated Bieberbach conjecture. Moreover, the de Branges' inequalities have also been the source of many other interesting inequalities involving logarithmic coefficients of $f \in S$ such as (see [6])

$$\sum_{k=1}^{\infty} |\gamma_k|^2 \le \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}.$$

More attention has been given to the results of an average sense (see [5, 6, 13]) than the exact upper bounds for $|\gamma_n|$ for functions in the class \mathcal{S} . Very few exact upper bounds for $|\gamma_n|$ seem have been established. For the Koebe function $k(z) = z/(1-z)^2$, the logarithmic coefficients are $\gamma_n = 1/n$. Since the Koebe function k(z) plays the role of extremal function for most of the extremal problems in the class \mathcal{S} , it is expected that $|\gamma_n| \leq \frac{1}{n}$ holds for functions in \mathcal{S} . But this is not true in general, even in order of magnitude [5, Theorem 8.4]. Indeed, there exists a bounded function f in the class \mathcal{S} with logarithmic coefficients $\gamma_n \neq O(n^{-0.83})$ (see [5, Theorem 8.4]). By differentiating (1.3) and equating coefficients we obtain

(1.4)
$$\gamma_1 = \frac{1}{2}a_2$$

(1.5)
$$\gamma_2 = \frac{1}{2}(a_3 - \frac{1}{2}a_2^2)$$

(1.6)
$$\gamma_3 = \frac{1}{2}(a_4 - a_2a_3 + \frac{1}{3}a_2^3).$$

If $f \in S$ then $|\gamma_1| \leq 1$ follows from (1.4). Using Fekete-Szegö inequality [5, Theorem 3.8] in (1.5), it is easy to obtain the following sharp estimate

$$|\gamma_2| \le \frac{1}{2}(1+2e^{-2}) = 0.635\dots$$

For $n \geq 3$, the problem seems much harder, and no significant upper bound for $|\gamma_n|$ when $f \in S$ appear to be known.

For functions in the class S^* , using the analytic characterization Re (zf'(z)/f(z)) > 0 for $z \in \mathbb{D}$ it is easy to prove that $|\gamma_n| \leq \frac{1}{n}$ for $n \geq 1$ and equality holds for the Koebe function $k(z) = z/(1-z)^2$. The inequality $|\gamma_n| \leq \frac{1}{n}$ for $n \geq 2$ also holds for functions in the class \mathcal{K} was claimed in a paper of Elhosh [7]. However, Girela [8] pointed out an error in the proof of Elhosh [7] and, hence, the result is not substantiated. Indeed, Girela proved that for each $n \geq 2$, there exists a function $f \in \mathcal{K}$ such that $|\gamma_n| > \frac{1}{n}$. Recently, it has been proved [14] that $|\gamma_3| \leq \frac{7}{12}$ for functions in \mathcal{K}_0 (close-to-convex functions with argument 0) with the additional assumption that the second coefficient of the corresponding starlike function g is real. But this estimate is not sharp as pointed out in [1] where the authors proved that $|\gamma_3| \leq \frac{1}{18}(3 + 4\sqrt{2}) = 0.4809$ for functions in \mathcal{K}_0 without assuming the additional assumption that the second coefficient of the corresponding starlike function g is real. In the same paper, the authors also determined the sharp upper bound $|\gamma_3| \leq \frac{1}{243}(28 + 19\sqrt{19}) = 0.4560$ for close-to-convex functions with argument 0 and with respect to the Koebe function and conjectured that this upper bound is also true for the whole class \mathcal{K}_0 .

Let S_2^* denote the class of odd starlike functions and \mathcal{F} denote the class of close-to-convex functions with argument 0 and with respect to odd starlike functions. That is,

$$\mathcal{F} = \left\{ f \in \mathcal{A} : \operatorname{Re} \frac{zf'(z)}{g(z)} > 0, \ z \in \mathbb{D}, \text{ for some } g \in \mathcal{S}_2^* \right\}.$$

It is important to note that the class \mathcal{F} is rotationally invariant. In the present article, we determine the sharp upper bound of $|\gamma_n|$, n = 1, 2, 3 for functions in \mathcal{F} .

2. Main Results

Let \mathcal{P} denote the class of analytic functions P of the form

(2.1)
$$P(z) = 1 + \sum_{n=1}^{\infty} c_n z^n$$

such that $\operatorname{Re} P(z) > 0$ in \mathbb{D} . Functions in \mathcal{P} are sometimes called Carathéodory functions. To prove our main results, we need some preliminary lemmas.

Lemma 2.1. [5, p. 41] For a function $P \in \mathcal{P}$ of the form (2.1), the sharp inequality $|c_n| \leq 2$ holds for each $n \geq 1$. Equality holds for the function P(z) = (1+z)/(1-z).

Lemma 2.2. [12] Let $P \in \mathcal{P}$ be of the form (2.1) and μ be a complex number. Then

(2.2)
$$|c_2 - \mu c_1^2| \le 2 \max\{1, |2\mu - 1|\}.$$

The result is sharp for the functions given by $P(z) = \frac{1+z^2}{1-z^2}$ and $P(z) = \frac{1+z}{1-z}$.

Lemma 2.3. [11] Let $P \in \mathcal{P}$ be of the form (2.1). Then there exist $x, t \in \mathbb{C}$ with $|x| \leq 1$ and $|t| \leq 1$ such that

$$2c_2 = c_1^2 + x(4 - c_1^2) \quad and 4c_3 = c_1^3 + 2(4 - c_1^2)c_1x - c_1(4 - c_1^2)x^2 + 2(4 - c_1^2)(1 - |x|^2)t.$$

Theorem 2.1. Let $f \in \mathcal{F}$ be of the form (1.1). Then $|\gamma_1| \leq \frac{1}{2}$, $|\gamma_2| \leq \frac{1}{2}$ and $|\gamma_3| \leq \frac{1}{972}(95 + 23\sqrt{46})$. The inequalities are sharp.

Proof. Let $f \in \mathcal{F}$ be of the form (1.1). Then there exists an odd starlike function $g(z) = z + \sum_{n=1}^{\infty} b_{2n+1} z^{2n+1}$ and a Carathéodory function $P \in \mathcal{P}$ of the form (2.1) such that

(2.3)
$$zf'(z) = g(z)P(z).$$

Comparing the coefficients on the both sides of (2.3) gives

(2.4)
$$a_2 = \frac{1}{2}c_1, \ a_3 = \frac{1}{3}(b_3 + c_2) \text{ and } a_4 = \frac{1}{4}(b_3c_1 + c_3).$$

Substituting a_2, a_3 and a_4 given by (2.4) in (1.4), (1.5) and (1.6) and then further simplification gives

(2.5)
$$\gamma_1 = \frac{1}{2}a_2 = \frac{1}{4}c_1$$

(2.6)
$$\gamma_2 = \frac{1}{2} \left(a_3 - \frac{1}{2} a_2^2 \right) = \frac{1}{6} b_3 + \frac{1}{6} \left(c_2 - \frac{3}{8} c_1^2 \right)$$

(2.7)
$$2\gamma_3 = a_4 - a_2 a_3 + \frac{1}{3}a_2^3 = \frac{1}{24} \left(2c_1 b_3 + c_1^3 - 4c_1 c_2 + 6c_3 \right).$$

In view of Lemma 2.1, it follows from (2.5) that $|\gamma_1| \leq \frac{1}{2}$ and equality holds for a function f defined by zf'(z) = g(z)P(z), where $g(z) = z/(1-z^2)$ and P(z) = (1+z)/(1-z). Since g is an odd starlike function, $|b_3| \leq 1$ (see [9, Chaptar 4, Theorem 3, page 35]). Using Lemma 2.2, it follows from (2.6) that

$$|\gamma_2| \le \frac{1}{6}|b_3| + \frac{1}{6}\left|c_2 - \frac{3}{8}c_1^2\right| \le \frac{1}{6} + \frac{1}{3} = \frac{1}{2}$$

and equality holds for a function f defined by zf'(z) = g(z)P(z), where $g(z) = z/(1-z^2)$ and $P(z) = (1+z^2)/(1-z^2)$.

Writing c_2 and c_3 in terms of c_1 with the help of Lemma 2.3, it follows from (2.7) that

(2.8)
$$48\gamma_3 = 2c_1b_3 + \frac{1}{2}c_1^3 + c_1x(4-c_1^2) - \frac{3}{2}c_1x^2(4-c_1^2) + 3(4-c_1^2)(1-|x|^2)t,$$

where $|x| \leq 1$ and $|t| \leq 1$. Since the class \mathcal{F} is invariant under rotation, without loss of generality we can assume that $c_1 = c$, where $0 \leq c \leq 2$. Taking modulus on both the sides of (2.8) and then applying triangle inequality and $|b_3| \leq 1$, it follows that

$$48|\gamma_3| \le 2c + \left|\frac{1}{2}c^3 + cx(4-c^2) - \frac{3}{2}cx^2(4-c^2)\right| + 3(4-c^2)(1-|x|^2),$$

where we have also used the fact $|t| \leq 1$. Let $x = re^{i\theta}$ where $0 \leq r \leq 1$ and $0 \leq \theta \leq 2\pi$. For simplicity, by writing $\cos \theta = p$ we obtain

(2.9)
$$48|\gamma_3| \le \psi(c,r) + |\phi(c,r,p)| =: F(c,r,p),$$

where $\psi(c, r) = 2c + 3(4 - c^2)(1 - r^2)$ and

$$\phi(c,r,p) = \left(\frac{1}{4}c^6 + c^2r^2(4-c^2)^2 + \frac{9}{4}c^2r^4(4-c^2)^2 + c^4(4-c^2)rp\right)^{1/2}$$
$$-\frac{3}{2}c^4r^2(4-c^2)(2p^2-1) - 3c^2(4-c^2)r^3p\right)^{1/2}.$$

Thus we need to find the maximum value of F(c, r, p) over the rectangular cube $R := [0, 2] \times [0, 1] \times [-1, 1].$

By elementary calculus it is easy to verify the following:

$$\max_{0 \le r \le 1} \psi(0, r) = \psi(0, 0) = 12, \quad \max_{0 \le r \le 1} \psi(2, r) = 4,$$
$$\max_{0 \le c \le 2} \psi(c, 0) = \psi\left(\frac{1}{3}, 0\right) = \frac{37}{3}, \quad \max_{0 \le c \le 2} \psi(c, 1) = \psi(2, 1) = 4 \quad \text{and}$$
$$\max_{(c,r) \in [0,2] \times [0,1]} \psi(c, r) = \psi\left(\frac{1}{3}, 0\right) = \frac{37}{3}.$$

We first find the maximum value of F(c, r, p) on the boundary of R, i.e on the six faces of the rectangular cube R.

On the face c = 0, we have $F(0, r, p) = \psi(0, r)$ for $(r, p) \in R_1 := [0, 1] \times [-1, 1]$. Thus

$$\max_{(r,p)\in R_1} F(0,r,p) = \max_{0\le r\le 1} \psi(0,r) = \psi(0,0) = 12.$$

On the face c = 2, we have F(2, r, p) = 8 for $(r, p) \in R_1$.

On the face r = 0, we have $F(c, 0, p) = 2c + 3(4 - c^2) + \frac{1}{2}c^3$ for $(c, p) \in R_2 := [0, 2] \times [-1, 1]$. Note that F(c, 0, p) is independent of p. Thus, by using elementary calculus it is easy to see that

$$\max_{(c,p)\in R_2} F(c,0,p) = F\left(\frac{2}{3}(3-\sqrt{6}),0,p\right) = \frac{8}{9}\left(9+\sqrt{6}\right) = 12.3546.$$

On the face r = 1, we have $F(c, 1, p) = \psi(c, 1) + |\phi(c, 1, p)|$ for $(c, p) \in R_2$. We first prove that $\phi(c, 1, p) \neq 0$ in the interior of R_2 . On the contrary, if $\phi(c, 1, p) = 0$ in the interior of R_2 then

$$|\phi(c,1,p)|^2 = \left|\frac{1}{2}c^3 + ce^{i\theta}(4-c^2) - \frac{3}{2}ce^{2i\theta}(4-c^2)\right|^2 = 0$$

and hence

$$\frac{1}{2}c^3 + cp(4-c^2) - \frac{3}{2}c(4-c^2)(2p^2-1) = 0 \text{ and}$$
$$c(4-c^2)\sin\theta - \frac{3}{2}c(4-c^2)\sin2\theta = 0.$$

On further simplification, this reduces to

$$\frac{1}{2}c^2 + p(4-c^2) - \frac{3}{2}(4-c^2)(2p^2-1) = 0 \text{ and } 1-3p = 0,$$

which is equivalent to p = 1/3 and $c^2 = 6$. This contradicts the range of $c \in (0, 2)$. Thus $\phi(c, 1, p) \neq 0$ in the interior of R_2 . Next, we find the maximum value F(c, 1, p) in the interior of R_2 . Suppose that F(c, 1, p) has the maximum at an interior point of R_2 . Then at such point $\frac{\partial F(c, 1, p)}{\partial c} = 0$ and $\frac{\partial F(c, 1, p)}{\partial p} = 0$. From $\frac{\partial F(c, 1, p)}{\partial p} = 0$ (for points in the interior of R_2), a straight forward calculation gives

(2.10)
$$p = \frac{2(c^2 - 3)}{3c^2}$$

Substituting the value of p as given in (2.10) in the relation $\frac{\partial F(c,1,p)}{\partial c} = 0$ and further simplification gives

$$2c - 3c^3 + \sqrt{6(c^2 + 2)} = 0.$$

Taking the last term on the right hand side and squaring on both the sides yields

(2.11)
$$9c^6 - 12c^4 - 2c^2 - 12 = 0.$$

This equation has exactly one root in (0, 2) which can be shown using the wellknown Strum theorem for isolating real roots and hence for the sake of brevity we omit the details. By solving the equation (2.11) numerically, we obtain the approximate root 1.3584 in (0, 2) and the corresponding value of p obtained from (2.10) is -0.4172. Thus the extremum points of F(c, 1, p) in the interior of R_2 lie in a small neighborhood of the points $A_1 = (1.3584, 1, -0.4172)$ (on the plane r = 1). Clearly $F(A_1) = 9.3689$. Since the function F(c, 1, p) is uniformly continuous on R_2 , the value of F(c, 1, p) would not vary too much in the neighborhood of the point A_1 .

Next we find the maximum value of F(c, 1, p) on the boundary of R_2 . Clearly, F(0, 1, p) = 0, F(2, 1, p) = 8,

$$F(c, 1, -1) = \begin{cases} 2c + c(10 - 3c^2) & \text{for } 0 \le c \le \sqrt{\frac{10}{3}} \\ 2c - c(10 - 3c^2) & \text{for } \sqrt{\frac{10}{3}} < c \le 2 \end{cases}$$

and

$$F(c, 1, 1) = \begin{cases} 2c + c(2 - c^2) & \text{for } 0 \le c \le \sqrt{2} \\ 2c - c(2 - c^2) & \text{for } \sqrt{2} < c \le 2. \end{cases}$$

By using elementary calculus we find that

$$\max_{0 \le c \le 2} F(c, 1, -1) = F\left(\frac{2\sqrt{3}}{3}, 1, -1\right) = \frac{16\sqrt{3}}{3} = 9.2376 \quad \text{and}$$
$$\max_{0 \le c \le 2} F(c, 1, 1) = F\left(\frac{2\sqrt{3}}{3}, 1, 1\right) = \frac{16\sqrt{3}}{9} = 3.0792.$$

Therefore

$$\max_{(c,p)\in R_2} F(c,1,p) \approx 9.3689.$$

On the face p = -1,

$$F(c, r, -1) = \begin{cases} \psi(c, r) + \eta_1(c, r) & \text{for} \quad \eta_1(c, r) \ge 0\\ \psi(c, r) - \eta_1(c, r) & \text{for} \quad \eta_1(c, r) < 0, \end{cases}$$

where $\eta_1(c,r) = c^3(3r^2 + 2r + 1) - 4cr(3r + 2)$ and $(c,r) \in R_3 := [0,2] \times [0,1]$. To find the maximum value of F(c,r,-1) in the interior of R_3 we need to solve the pair of equations $\frac{\partial F(c,r,-1)}{\partial c} = 0$ and $\frac{\partial F(c,r,-1)}{\partial r} = 0$ in the interior of R_3 . But it is important to note that $\frac{\partial F(c,r,-1)}{\partial c}$ and $\frac{\partial F(c,r,-1)}{\partial r}$ may not exist at points in $S_1 = \{(c,r) \in R_3 : \eta_1(c,r) = 0\}$. Solving these pair of equations, we find that

$$\max_{(c,r)\in \operatorname{int} R_3\setminus S_1} F(c,r,-1) = F\left(\frac{1}{3}(\sqrt{82}-8), \frac{1}{57}(\sqrt{82}-5), -1\right)$$
$$= \frac{4}{81}(41\sqrt{82}-121) = 12.359.$$

Now we find the maximum value of F(c, r, -1) on the boundary of R_3 and on the set S_1 . Note that

$$\max_{(c,r)\in S_1} F(c,r,-1) \le \max_{(c,r)\in R_3} \psi(c,r) = \frac{37}{3} = 12.33.$$

On the other hand by using elementary calculus, as before, we find that

$$\max_{0 \le r \le 1} F(0, r, -1) = \max_{0 \le r \le 1} 12(1 - r^2) = F(0, 0, -1) = 12, \quad \max_{0 \le r \le 1} F(2, r, -1) = 8,$$

$$\max_{0 \le c \le 2} F(c, 0, -1) = \max_{(c, p) \in R_2} F(c, 0, p) = F\left(\frac{2}{3}(3 - \sqrt{6}), 0, -1\right) = \frac{8}{9}\left(9 + \sqrt{6}\right) = 12.3546$$

and
$$\max_{0 \le c \le 2} F(c, 1, -1) = F\left(\frac{2\sqrt{3}}{3}, 1, -1\right) = \frac{16\sqrt{3}}{3} = 9.2376.$$

Hence, by combining the above cases we obtain

$$\max_{(c,r)\in R_3} F(c,r,-1) = F\left(\frac{1}{3}(\sqrt{82}-8), \frac{1}{57}(\sqrt{82}-5), -1\right)$$
$$= \frac{4}{81}(41\sqrt{82}-121) = 12.359.$$

On the face p = 1,

$$F(c, r, 1) = \begin{cases} \psi(c, r) + \eta_2(c, r) & \text{for} \quad \eta_2(c, r) \ge 0\\ \psi(c, r) - \eta_2(c, r) & \text{for} \quad \eta_2(c, r) < 0, \end{cases}$$

where $\eta_2(c,r) = c^3(3r^2 - 2r + 1) - 4cr(3r - 2)$ for $(c,r) \in R_3$. To find the maximum value of F(c,r,1) in the interior of R_3 we need to solve the pair of equations $\frac{\partial F(c,r,1)}{\partial c} = 0$ and $\frac{\partial F(c,r,1)}{\partial r} = 0$ in the interior of R_3 . But it is important to note that $\frac{\partial F(c,r,1)}{\partial c}$ and $\frac{\partial F(c,r,1)}{\partial r}$ may not exist at points in $S_2 = \{(c,r) \in R_3 : \eta_2(c,r) = 0\}$. Solving these pair of equations, we find that

$$\max_{(c,r)\in \operatorname{int} R_3\setminus S_2} F(c,r,1) = F\left(\frac{1}{3}(8-\sqrt{46}),\frac{1}{75}(11-\sqrt{46}),1\right)$$
$$= \frac{4}{81}(95+23\sqrt{46}) = 12.3947.$$

Now, we find the maximum value of F(c, r, 1) on the boundary of R_3 and on the set S_2 . By noting that (see earlier cases)

$$\max_{\substack{(c,r)\in S_2}} F(c,r,1) \le \max_{\substack{(c,r)\in R_3}} \psi(c,r) = \frac{37}{3} = 12.33,$$

$$\max_{\substack{0\le r\le 1}} F(0,r,1) = 12, \quad \max_{\substack{0\le r\le 1}} F(2,r,1) = 8,$$

$$\max_{\substack{0\le c\le 2}} F(c,0,1) = \frac{8}{9} \left(9 + \sqrt{6}\right) = 12.3546,$$

$$\max_{\substack{0\le c\le 2}} F(c,1,1) = \frac{16\sqrt{3}}{9} = 3.0792$$

and combining all the cases, we find that

$$\max_{(c,r)\in R_3} F(c,r,1) = F\left(\frac{1}{3}(8-\sqrt{46}),\frac{1}{75}(11-\sqrt{46}),1\right)$$
$$= \frac{4}{81}(95+23\sqrt{46}) = 12.3947.$$

Let $S'=\{(c,r,p)\in R: \phi(c,r,p)=0\}.$ Then

$$\max_{(c,r,p)\in S'} F(c,r,p) \le \max_{(c,r)\in R_3} \psi(c,r) = \psi\left(0,\frac{1}{3}\right) = \frac{37}{3} = 12.33.$$

We prove that F(c, r, p) has no maximum value at any interior point of $R \setminus S'$. Suppose that F(c, r, p) has a maximum value at an interior point of $R \setminus S'$. Then at such point $\frac{\partial F(c,r,p)}{\partial c} = 0$, $\frac{\partial F(c,r,p)}{\partial r} = 0$ and $\frac{\partial F(c,r,p)}{\partial p} = 0$. Note that $\frac{\partial F(c,r,p)}{\partial c}$, $\frac{\partial F(c,r,p)}{\partial r}$ and $\frac{\partial F(c,r,p)}{\partial p}$ may not exist at points in S'. From $\frac{\partial F(c,r,p)}{\partial p} = 0$ (for points in the interior of $R \setminus S'$), a straight forward but laborious calculation gives

(2.12)
$$p = \frac{3c^2r^2 + c^2 - 12r^2}{6c^2r}.$$

Substituting the value of p as given in (2.12) in $\frac{\partial F(c,r,p)}{\partial r} = 0$ and simplifying, we obtain

$$(4 - c^2)r(\sqrt{6(c^2 + 2)} - 6) = 0.$$

This equation has no solution in the interior of $R \setminus S'$ and hence F(c, r, p) has no maximum in the interior of $R \setminus S'$.

Thus combining all the above cases we find that

$$\max_{(c,r,p)\in R} F(c,r,p) = F\left(\frac{1}{3}(8-\sqrt{46}), \frac{1}{75}(11-\sqrt{46}), 1\right)$$
$$= \frac{4}{81}(95+23\sqrt{46}) = 12.3947$$

and hence from (2.9) we obtain

(2.13)
$$|\gamma_3| \le \frac{1}{972}(95 + 23\sqrt{46}) = 0.2582.$$

We now show that the inequality (2.13) is sharp. An examination of the proof shows that equality holds in (2.13) if we choose $b_3 = 1$, $c_1 = c = \frac{1}{3}(8 - \sqrt{46})$, $x = \frac{1}{75}(11 - \sqrt{46})$ and t = 1 in (2.8). For such values of c_1 , x and t, Lemma 2.3 gives $c_2 = \frac{1}{27}(134 - 19\sqrt{46})$ and $c_3 = \frac{2}{243}(721 - 71\sqrt{46})$. A function $P \in \mathcal{P}$ having the first three coefficients c_1, c_2 and c_3 as above is given by (2.14)

$$P(z) = (1 - 2\lambda)\frac{1 + z}{1 - z} + \lambda \frac{1 + uz}{1 - uz} + \lambda \frac{1 + \overline{u}z}{1 - \overline{u}z}$$

= $1 + \frac{1}{3}(8 - \sqrt{46})z + \frac{1}{27}(134 - 19\sqrt{46})z^2 + \frac{2}{243}(721 - 71\sqrt{46})z^3 + \cdots,$

where $\lambda = \frac{1}{10}(-4 + \sqrt{46})$ and $u = \alpha + i\sqrt{1 - \alpha^2}$ with $\alpha = \frac{1}{18}(-1 - \sqrt{46})$. Hence the equality holds in (2.13) for a function f defined by zf'(z) = g(z)P(z), where $g(z) = z/(1 - z^2)$ and P(z) is given by (2.14). This completes the proof. \Box

Acknowledgement: The authors thank Prof. K.-J. Wirths for useful discussion and suggestions. The first author thank University Grants Commission for the financial support through UGC-SRF Fellowship. The second author thank SERB (DST) for financial support.

References

- MD FIROZ ALI and A. VASUDEVARAO, On logarithmic coefficients of some close-to-convex functions, arXiv:1606.05162.
- [2] I.E. BAZILEVICH, Coefficient dispersion of univalent functions, Mat. Sb. 68(110) (1965), 549– 560.
- [3] I.E. BAZILEVICH, On a univalence criterion for regular functions and the dispertion of their derivatives, *Mat. Sb.* 74(116) (1967), 133–146.
- [4] L. DE BRANGES, A proof of the Bieberbach conjecture. Acta Math. 154 (1985), no. 1-2, 137–152.
- [5] P. L. DUREN, Univalent functions (Grundlehren der mathematischen Wissenschaften 259, New York, Berlin, Heidelberg, Tokyo), Springer-Verlag, 1983.
- [6] P. L. DUREN and Y. J. LEUNG, Logarithmic coefficients of univalent functions, J. Analyse Math. 36 (1979), 36–43.
- [7] M. M. ELHOSH, On the logarithmic coefficients of close-to-convex functions, J. Austral. Math. Soc. Ser. A 60 (1996), 1–6.
- [8] D. GIRELA, Logarithmic coefficients of univalent functions, Ann. Acad. Sci. Fenn. Math. 25 (2000), 337–350.
- [9] A. W. GOODMAN, Univalent Functions, Vols. I and II. Mariner Publishing Co. Tampa, Florida, 1983.
- [10] W.K. HAYMAN, Multivalent Functions, Cambridge University Press, 1958.
- [11] R.J. LIBERA and E.J. ZŁOTKIEWICZ, Early coefficients of the inverse of a regular convex function, Proc. Amer. Math. Soc. 85 (2) (1982) 225–230.
- [12] W. MA and D. MINDA, A unified treatment of some special classes of univalent functions, Proceedings of the Conference on Complex Analysis, Tianjin, 1992 (Int. Press), 157–169.
- [13] O. ROTH, A sharp inequality for the logarithmic coefficients of univalent functions, Proc. Amer. Math. Soc. 135(7) (2007), 2051–2054.
- [14] D.K. THOMAS, On the logarithmic coefficients of close to convex functions, Proc. Amer. Math. Soc. 144 (2016), 1681–1687.

MD FIROZ ALI, DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY KHARAG-PUR, KHARAGPUR-721 302, WEST BENGAL, INDIA.

E-mail address: ali.firoz89@gmail.com

A. VASUDEVARAO, DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY KHARAGPUR, KHARAGPUR-721 302, WEST BENGAL, INDIA.

E-mail address: alluvasu@maths.iitkgp.ernet.in