Burstiness and tie reinforcement in time varying social networks

Enrico Ubaldi,^{1, 2, 3} Alessandro Vezzani,^{2, 4} Márton Karsai,⁵ Nicola Perra,⁶ and Raffaella Burioni^{2, 3}

¹Institute for Scientific Interchange Foundation, 10126 Torino, Italy

²Dipartimento di Fisica e Scienza della Terra, Università di Parma,

Parco Area delle Scienze 7/A, 43124 Parma, Italy

³INFN, Gruppo Collegato di Parma, Parco Area delle Scienze 7/A, 43124 Parma, Italy

⁴Centro S3, CNR-Istituto di Nanoscienze, Via Campi 213A, 41125 Modena Italy

⁵ Univ de Lyon, ENS de Lyon, INRIA, CNRS, UMR 5668, IXXI, 69364 Lyon, France

⁶Centre for Business Network Analysis, University of Greenwich, Park Row, London SE10 9LS, United Kingdom

(Dated: August 24, 2021)

We introduce a time-varying network model accounting for burstiness and tie reinforcement observed in social networks. The analytical solution indicates a non-trivial phase diagram determined by the competition of the leading terms of the two processes. We test our results against numerical simulations, and compare the analytical predictions with an empirical dataset finding good agreements between them. The presented framework can be used to classify the dynamical features of real social networks and to gather new insights about the effects of social dynamics on ongoing spreading processes.

The recent availability of longitudinal and timeresolved datasets capturing social behaviours has induced a paradigm shift in the way we study, describe, and model the interactions between individuals. It moved the focus from static, time-aggregated, representations to time-varying, dynamical, characterisations of social networks [1–4]. Thinking in terms of time-varying systems allows to overcome the limitations arising from the depiction of social ties as fixed and immutable in time [2, 3]. Indeed, it allows to capture a set of complex dynamics driving the evolution of links [5–11] and to uncover the effects of such dynamics on processes unfolding on the networks' fabrics [12–21] (see Ref. [3] for a recent review).

While social networks are shaped by a multitude of processes [22], two particular mechanisms have been found to play central roles in their emergence and evolution [13, 14, 23–27]. The first is social activity, i.e. the propensity of individuals to be engaged in social act per unit time. Observations in a range of real datasets, capturing different types of social dynamics, show that activity is heterogeneously distributed among people [27–30]. Furthermore, the activity of single individuals evolves through *bursty* temporal patterns, inducing heterogeneous inter-event time distributions [31–39]. In other words, not only individuals show heterogeneous propensities to be socially active, but their activation is bursty as well.

The second important mechanism determines the *allocation of social ties*, i.e. the selection process driving the creation or renewal of a particular connection. Intuitively, social tie allocation is not random. In fact, empirical observations show that people tend to distribute a large fraction of their social acts towards already existing strong ties, while allocating a smaller amount of interactions to create new social relationships or to maintain weak ties [23–25, 27, 40, 41]. In other words, in time some connections are frequently reinforced by repeated

interactions, while others are not. Although the study of these mechanisms has been the focus of a range of works [13, 14, 23–27], a general modeling framework is still missing. Such a framework would allow for the analytical characterization on how the interplay of heterogeneous activity patterns and tie allocation mechanisms shape the evolution of social networks, and in turn the processes taking place on their fabrics.

Here we introduce a model of time-varying networks that allows to regulate the relative strength of burstiness and tie reinforcement simultaneously. We analytically solve the asymptotic behaviour of the model and find a non-trivial phase diagram determined by the interplay of the two processes under investigation. In particular, we observe two different dynamical regimes, one in which burstiness governs the evolution of the network, and vice versa another in which the dynamics is completely determined by the process of ties allocation. Interestingly, if the reinforcement of previously activated connections is sufficiently strong, burstiness can be sub-leading the network evolution even in the presence of large inter-event time fluctuations. The theoretical results are validated considering a real world social network, with analytical predictions fitting the empirical observations. Consequently, the proposed framework can be also used to classify the temporal features of real networks, which could provide new insights on the effects that social mechanisms have on spreading processes unfolding on social networks.

In the framework of activity driven networks [24, 27, 28], a network \mathcal{G} is formed by N nodes and each node i is assigned with an activity a_i drawn from an arbitrary distribution F(a). The activity sets the activation rate of node i, i.e. the probability $a_i\Delta t$ that a node active in time interval Δt . In general, F(a) is chosen to be a broad function reflecting the shape of the corresponding distribution in empirical observations [24, 27, 28]. Typically



Figure 1. In the phase diagram we report the delimiting lines of the different scaling regions as found in Eq. (6). Evidencing the **Str**ong **B**urstiness **R**egime (StrBR), the **W**eak **B**urstiness **R**egime (WBR), and the **Sup**pressed **B**urstiness **R**egime (SupBR). We also show the parameters value of the simulations presented in Fig. 2 (yellow tags), and in Fig. 3 (white and blue tags).

a power law distribution i.e. $F(a) \sim a^{-(\nu+1)}$ is observed for large activity.

The inter-event time τ_i , i.e. the time between two subsequent activations of the agent *i*, is directly connected with the agent activity, since $a_i = 1/\langle \tau_i \rangle$. All activitydriven models proposed so far considered a Poissonian distributed τ_i [42]. However, in social systems the interevent time distribution of a single agent is strongly heterogeneous and usually spans over several orders of magnitude. In order to capture this bursty nature of human dynamics, we impose that the inter-event time τ_i for node *i* is drawn from a power-law distribution $\Psi(\tau_i)$:

$$\Psi(\tau_i) = \frac{\alpha}{\xi_i^{-\alpha}} \tau_i^{-(1+\alpha)}, \ \tau_i \in [\xi_i, +\infty), \tag{1}$$

where the exponent α characterizes the distribution and ξ_i is a lower time cutoff. The latter represents the characteristic timescale for the node *i*, i.e. $\xi_i \sim 1/a_i$, as the γ -th moment of the distribution $\Psi(\tau_i)$ reads $\langle \tau_i^{\gamma} \rangle \sim \xi_i^{\gamma}$. If also the ξ_i are heterogeneously distributed as

$$\Phi(\xi_i) \propto \xi_i^{\nu-1},\tag{2}$$

for small ξ_i , as a consequence we obtain a network in which the corresponding activity potential a_i is broadly distributed. In particular the activity distribution behaves as $F(a_i) \propto a_i^{-(\nu+1)}$ for large a_i . We note that, instead of introducing an agent dependent cut-off, different definitions are possible, e.g. considering a distribution of waiting times $\Psi(\tau_i) = \alpha \delta_i^{\alpha} / (\delta_i + \tau_i)^{1+\alpha}$, where $a_i \sim 1/\delta_i$ since $\langle \tau_i^{\gamma} \rangle \sim \delta_i^{\gamma}$. Our model, therefore, belongs to a novel class of activity driven networks, where the agent time scale is set by a parameter in the waiting time distribution.

When a node is active, it initiates interactions with other nodes in the network. This way the degree k_i of a node *i*, defined as the number of connected peers of *i* up to time *t*, is evolving in time. To model this evolution we build on the latest development of the model in which the



Figure 2. (color online) The scaling of the P(k, t) function for (a) $\beta = 0.7$, $\alpha = 1.35$ (WBR region) and (b) $\beta = 0.7$, $\alpha = 1.6$ (SupBR regime). In each plot we consider logarithmically spaced times between $t = 10^4$ and $t = 10^7$ averaged over 10^5 representation of the single agent evolution. The curves referring to the longest time $t = 10^7$ are shown in solid thick line, while shorter times are shown in dashed lines. A comparison with a normal distribution (black dashed lines) evidence a good agreement with the SupBR data (b) while it completely misses the WBR case (a). Insets plot the $\langle k(t) \rangle$ and the corresponding analytical prediction of Eq. (7) (green solid lines).

selection of ties is driven by a reinforcement process [24, 27]. In particular, if at time t a node i of degree k_i is active it will contact a new, randomly chosen node with probability $p_i(k_i)$. Instead, with probability $1 - p_i(k_i)$ it reinforces a tie by contacting a node randomly chosen amongst the k_i already contacted agents. The form of $p_i(k_i)$ has been measured and characterized [27] in several datasets as:

$$p_i(k_i) = \left(1 + \frac{k_i}{c_i}\right)^{-\beta_i},\tag{3}$$

where β_i measures the strength of the reinforcement process, and c_i sets the characteristic number of ties that an individual is able to maintain before the reinforcement process takes place. Though simple, the reinforcement mechanism provides a reliable description of real world datasets and allows for analytical tractability.

In our simulations, initially, for each node i we set the integrated degree $k_i = 0$ and assign a lower cut-off ξ_i according to distribution $\Phi(\xi_i)$ in Eq. (2). The evolution starts by extracting, for each node i, a time τ_i at which the node will get active for the first time. We then activate one node at a time accordingly to their next activation time. When active, an agent i selects a



Figure 3. (color online). Numerical simulation of the full multi agent dynamics. (a) the rescaled P(k, t) distribution for ξ_i distributed with $\nu = 1.4$, $\alpha = 0.5$, $\beta = 0.75$. Curves refer to ten logarithmically spaced times between $t = 10^5$ and $t = 10^8$ (dashed lines, the longest time is shown in solid line). The data correspond to the StrBR regime. In the inset we show the growth of average degree for different activity classes (symbols) rescaled as $t \to t(a + \langle a \rangle)$. The analytical prediction of Eq. (7) is shown for comparison (blue solid line). (b) the P(k,t) distribution for $\nu = 1.2$, $\beta = 0.5$ and $\alpha = 2.2$, at seven logarithmically spaced times between $10^4 \leq t \leq 10^6$. Inset compares the degree distribution at the final simulation time (circles) with the analytical prediction (solid line) of Eq.(8).

randomly chosen other agent in the network with probability $p(k_i) = (1 + k_i/c)^{-\beta}$; in this case the value of k_i is increased by one both for the connecting and the connected nodes. Otherwise, with probability $1 - p_i(k_i)$, the agent *i* interacts with a randomly chosen neighbor node which has been already connected to *i*. For simplicity we fix β and *c* constant for all nodes. After each iteration the interaction of node *i* is removed and a new activation time is selected by an inter-event time τ_i drawn from the distribution in Eq. (1).

In the following we apply a single agent approximation, in which agents can only attach to other nodes and never get contacted. In this case we can write the master equation (ME) describing their degree evolution, and analytically solve it in the asymptotic limit of large times. In this case the activity is fixed to the value $a_0 = \xi_0^{-1}$ where ξ_0 is the characteristic time of the considered agent. In particular, let us define Q(k,t) as the probability for an agent active at time t to have degree k right after t. The ME then reads as

$$Q(k,t) = \int_{\xi_0}^{\infty} \frac{Q(k-1,t-t')}{t'^{\alpha+1}} \frac{c^{\beta}}{(c+k-1)^{\beta}} + \int_{\xi_0}^{\infty} \frac{Q(k,t-t')}{t'^{\alpha+1}} \left(1 - \frac{c^{\beta}}{(c+k)^{\beta}}\right) + \delta(k,0)\delta(t,0).$$
(4)

The first term accounts for the probability that the agent gets active and contacts a new node, while the second term represents the probability of connecting to an already contacted neighbor. We evaluate the probability P(k,t) for a node to have degree k at the time t, by integrating Eq. (4) over all the possible waiting time values, i.e.

$$P(k,t) = \int_{\xi_0}^{\infty} Q(k,t-t') \int_{t'}^{\infty} \frac{1}{\tau^{\alpha}} dt d\tau.$$
 (5)

In the asymptotic regime P(k,t) is (see Supporting Materials for details):

$$P(k,t) \simeq \begin{cases} \frac{1}{(ta_0)^{\frac{\alpha}{1+\beta}}} f_{\alpha\beta} \left(A'_{\alpha,\beta} \frac{k}{(ta_0)^{\frac{\alpha}{1+\beta}}} \right) & \text{if } \alpha < 1, \\ \frac{1}{(ta_0)^{\frac{1}{\alpha} - \frac{1}{(1+\beta)}}} f_{\alpha\beta} \left(A'_{\alpha,\beta} \frac{k-v(ta_0)^{1/(1+\beta)}}{(ta_0)^{\frac{1}{\alpha} - \frac{1}{(1+\beta)}}} \right) & \text{if } 1 < \alpha < \frac{2\beta+2}{2\beta+1}, \\ \frac{1}{(ta_0)^{\frac{1}{2(1+\beta)}}} \exp\left[-A_{\beta} \frac{\left(k-C_{\beta}(ta_0)^{\frac{1}{1+\beta}} \right)^2}{(ta_0)^{1/(1+\beta)}} \right] & \text{if } \alpha > \frac{2\beta+2}{2\beta+1}, \end{cases}$$
(6)

where $f_{\alpha\beta}(x)$ is a non-Gaussian scaling function (see [43]), v is the drift velocity of the peak of the distribution and $A_{\alpha,\beta}$, A_{β} , C_{β} are constants depending on α and β , respectively. degree $\langle k(t) \rangle$ is:

$$\langle k(t) \rangle \propto \begin{cases} t^{\alpha/(1+\beta)} & \text{if } \alpha < 1, \\ t^{1/(1+\beta)} & \text{if } \alpha > 1. \end{cases}$$
(7)

This solution leads to a dynamical phase diagram of the model, summarized in Fig. 1. For $\alpha < 1$, in the **Str**ong **B**urstiness **R**egime (StrBR) burstiness strongly affects the dynamics. Here the scaling function $f_{\alpha\beta}(x)$ is

As a consequence of Eq. (6), the growth of the average

not Gaussian and the exponent $\alpha/(1+\beta)$, leading the growth of $\langle k(t) \rangle$, depends both on the burstiness and on the reinforcement exponents, β and α respectively. On the other hand, for $\alpha > (2\beta + 2)/(2\beta + 1)$, we have a **Sup**pressed **B**urstiness **R**egime (SupBR), where the dynamics is independent of α . In particular, the reinforcement driven behavior, described in reference [27], is fully recovered with a Gaussian scaling function and a connectivity growing as $t^{1/(1+\beta)}$. Finally, for 1 < $\alpha < (2\beta + 2)/(2\beta + 1)$ the average connectivity grows as $t^{1/(1+\beta)}$ just as in the systems with suppressed burstiness. In this regime, the scaling function is not Gaussian and its scaling length depends on the burstiness exponent α . We call this phase the Weak Burstiness Regime (WBR). The non trivial dependence on β and α of the transition line between WBR and SupBR highlights the complex interplay between burstiness and tie reinforcement occurring for $1 < \alpha < 2$. Fig. 2 shows that the curve $\alpha = (2\beta + 2)/(2\beta + 1)$ marks a transition from a Gaussian to a non Gaussian scaling function, providing a numerical support to the analytical asymptotic results. In particular, in Fig. 2 (a) panel the left tail of the curve is slowly increasing with time, and the asymmetric distribution cannot be fitted with a Gaussian. On the other hand, in Fig. 2 (b) we observe the opposite behavior: the long time curve is slowly converging to a normal PDF.

Interestingly, the single-agent model provides a qualitatively correct description even of the multi-agent case, where different agents display different activities (see Fig. 3). In particular, one can suppose that the degree distribution Eq. (6) holds for each node of the system if one replaces $a_0 t$ with $(a + \langle a \rangle)t$. In this case at and $\langle a \rangle t$ represent the contribution to the growth of the degree induced by the node activity and by the activity of the rest of the network respectively (see [27] and [44] for an explicit calculation in the case without burstiness). Simulations evidence that the approximation works quite well, however, larger evolution times are required for observing the asymptotic scaling behavior. The same hypothesis allows to evaluate the degree distribution among different agents. In particular, if the activity a is distributed according to Eq. (2), at a given time t the degrees, for large k, are distributed as:

$$\rho(k) \propto \begin{cases} k^{-[\nu(1+\beta)/\alpha+1]} & \text{if } \alpha < 1, \\ k^{-[\nu(1+\beta)+1]} & \text{if } \alpha > 1. \end{cases}$$
(8)

As shown in Fig. 3(b) inset, the simulation results are well described by the asymptotic behavior in Eq. (8).

To motivate our model with real world observations, we checked how the proposed scaling picture corresponds to rescaled empirical P(k,t) functions measured in a Twitter Mentions Network (TMN) datasets (for details see SM). First of all, the system appears with broad activity and degree distributions (see Fig. 4(a)) with exponents satisfying the relations obtained in Eq. 8. More-



Figure 4. (color online). Twitter mentions dataset. (a) The activity distribution F(a) (blue squares) fitted as $F(a) \propto a^{-(1+\nu)}$ (green solid line) with $\nu = 1.25$ and the degree distribution $\rho(k)$ (red circles) with the predicted behavior (blue solid line) of Eq. (8). In the inset we plot the waiting-time distribution $\Psi(\tau_i)$ (blue circles) and the fit $\Psi(\tau_i) \propto \tau_i^{-(1+\alpha)}$ (red solid line) giving $\alpha = 0.95$. (b) The distribution P(a, k, t) for a selected activity class, the degree k is rescaled dividing by $t^{\alpha/(1+\beta)}$ where $\beta = 0.47$ has been found in [27] and α was evaluated in the upper panel. Inset shows the average degree growth $\langle k(t) \rangle$ for different activity classes a (symbols) rescaling time as $t \to t(a + \langle a \rangle)$. The predicted behavior of Eq. (7) is shown for comparison (green solid line).

over, the inter-event time distribution approximately follows a power-law (see Fig. 4(a) inset). Notice that, given the measured value of the exponent $\alpha \sim 0.95$, we expect the TMN system to fall in the StrBR region. This is verified in Fig. 4(b) where indeed the P(k,t) distributions at different evolution times are not Gaussian and seem to apply to the expected dynamical scaling. The proposed framework allows to classify the dynamical features of real social networks and thus anticipate their effects on spreading processes taking place on their fabrics.

In summary, we introduced a new model, which is able to capture two key aspects driving the evolution of social networks: burstiness and tie reinforcement. We solved the ME of the model in the large time regime and explored analytically a complex phase space, where changes in the relative importance between the two mechanisms are linked to different degree distributions and emerging dynamics. Interestingly, the presented framework is able to predict observables of real networks and provides a new way to classify their dynamical features. Starting from the analytic result, an interesting further improvement is the introduction of a dynamic on the network population, encoding the fact that in real world dataset agents typically can enter or exit from the system during the network evolution.

Our results also have the potential to provide new insights for the characterization of spreading processes unfolding on social networks. In particular, the proposed modeling framework could help to clarify the role of burstiness on contagion phenomena, which is currently subject of a heated debate. The model can potentially be extended further by including other social processes such as the presence of communities or ageing of nodes.

- [1] C. T. Butts, Sociological Methodology **38**, 155 (2008).
- [2] P. Holme and J. Saramäki, Physics Reports 519, 97 (2012),.
- [3] P. Holme, The European Physical Journal B 88, 1 (2015).
- [4] B. Gonçalves and N. Perra, Social phenomena: From data analysis to models (Springer, 2015).
- [5] S. Liu, N. Perra, M. Karsai, and A. Vespignani, Phys. Rev. Lett. **112**, 118702 (2014).
- [6] L. Isella, J. Stehlé, A. Barrat, C. Cattuto, J.-F. Pinton, and W. Van den Broeck, Journal of theoretical biology 271, 166 (2011).
- [7] P. Grindrod, M. C. Parsons, D. J. Higham, and E. Estrada, Physical Review E 83, 046120 (2011).
- [8] S. Praprotnik and V. Batagelj, Ars Mathematica Contemporanea 11 (2015).
- [9] G. Ghoshal and P. Holme, Physica A: Statistical Mechanics and its Applications 364, 603 (2006).
- [10] J. Saramäki and E. Moro, The European Physical Journal B 88, 1 (2015).
- [11] J. Saramäki, E. A. Leicht, E. López, S. G. Roberts, F. Reed-Tsochas, and R. I. Dunbar, Proceedings of the National Academy of Sciences 111, 942 (2014).
- [12] S.-Y. Liu, A. Baronchelli, and N. Perra, Phys. Rev. E 87, 032805 (2013).
- [13] G. Miritello, E. Moro, and R. Lara, Phys. Rev. E 83, 045102 (2011).
- [14] M. Starnini and R. Pastor-Satorras, Phys. Rev. E 89, 032807 (2014).
- [15] E. Valdano, L. Ferreri, C. Poletto, and V. Colizza, Physical Review X 5, 021005 (2015).
- [16] L. E. Rocha and V. D. Blondel, PLoS Comput Biol 9, e1002974 (2013).
- [17] I. Scholtes, N. Wider, R. Pfitzner, A. Garas, C. J. Tessone, and F. Schweitzer, Nature communications 5 (2014).
- [18] D. Han, M. Sun, and D. Li, Physica A: Statistical Mechanics and its Applications 432, 354 (2015).
- [19] M. Morris and M. Kretzschmar, Aids 11, 641 (1997).
- [20] L. E. Rocha and N. Masuda, New Journal of Physics 16,

063023 (2014).

- [21] M. Kivelä, R. K. Pan, K. Kaski, J. Kertész, J. Saramäki, and M. Karsai, Journal of Statistical Mechanics: Theory and Experiment **2012**, P03005 (2012).
- [22] M. O. Jackson et al., Social and economic networks, Vol. 3 (Princeton university press Princeton, 2008).
- [23] J.-P. Onnela, J. Saramäki, J. Hyvönen, G. Szabó, D. Lazer, K. Kaski, J. Kertész, A.-L. Barabási, Proceedings the Naand of tional Academy of Sciences 104, 7332 (2007), http://www.pnas.org/content/104/18/7332.full.pdf+html.
- [24] M. Karsai, N. Perra, and A. Vespignani, Sci. Rep. 4, 4001 (2014).
- [25] G. Miritello, R. Lara, M. Cebrian, and E. Moro, Sci. Rep. 3 (2013).
- [26] B. Gonçalves, N. Perra, and A. Vespignani, PLoS ONE 6, e22656 (2011).
- [27] E. Ubaldi, N. Perra, M. Karsai, A. Vezzani, R. Burioni, and A. Vespignani, arXiv preprint arXiv:1509.04563 (2015).
- [28] N. Perra, B. Gonçalves, R. Pastor-Satorras, and A. Vespignani, Sci. Rep. 2 (2012).
- [29] M. Vincenzo Tomasello, N. Perra, C. Juan Tessone, M. Karsai, and F. Schweitzer, Sci. Rep. 4 (2014),.
- [30] B. Ribeiro, N. Perra, and A. Baronchelli, Sci. Rep. 3 (2013).
- [31] R. D. Malmgren, D. B. Stouffer, A. E. Motter, and L. A. N. Amaral, Proceedings of the National Academy of Sciences 105, 18153 (2008), http://www.pnas.org/content/105/47/18153.full.pdf.
- [32] A. Moinet, M. Starnini, and R. Pastor-Satorras, Phys. Rev. Lett. **114**, 108701 (2015).
- [33] K.-I. Goh and A.-L. Barabási, EPL (Europhysics Letters) 81, 48002 (2008).
- [34] A. Vázquez, J. a. G. Oliveira, Z. Dezsö, K.-I. Goh, I. Kondor, and A.-L. Barabási, Phys. Rev. E 73, 036127 (2006).
- [35] A.-L. Barabasi, Nature **435**, 207 (2005).
- [36] M. Karsai, K. Kaski, A.-L. Barabási, and J. Kertész, Sci. Rep. 2 (2012).
- [37] H.-H. Jo, M. Karsai, J. Kertész, and K. Kaski, New Journal of Physics 14, 013055 (2012).
- [38] M. Karsai, K. Kaski, and J. Kertész, PLoS ONE 7, e40612 (2012).
- [39] M. Karsai, M. Kivelä, R. K. Pan, K. Kaski, J. Kertész, A.-L. Barabási, and J. Saramäki, Phys. Rev. E 83, 025102 (2011).
- [40] M. Granovetter, Sociological Theory 1, 201 (1983).
- [41] L. Weng, M. Karsai, N. Perra, F. Menczer, and A. Flammini, arXiv preprint arXiv:1505.02399 (2015).
- [42] With exception for [32] which however does not consider the reinforcement process.
- [43] R. Burioni, G. Gradenigo, A. Sarracino, A. Vezzani, and A. Vulpiani, Journal of Statistical Mechanics: Theory and Experiment **2013**, P09022 (2013).
- [44] M. Starnini and R. Pastor-Satorras, Phys. Rev. E 87, 062807 (2013).

Supplemental Material for "Burstiness and tie reinforcement in time varying social networks".

Enrico Ubaldi,^{1, 2, 3} Alessandro Vezzani,^{2, 4} Márton Karsai,⁵ Nicola Perra,⁶ and Raffaella Burioni^{2, 3}

¹Institute for Scientific Interchange Foundation, 10126 Torino, Italy

²Dipartimento di Fisica e Scienza della Terra, Università di Parma,

Parco Area delle Scienze 7/A, 43124 Parma, Italy

³INFN, Gruppo Collegato di Parma, Parco Area delle Scienze 7/A, 43124 Parma, Italy

⁴Centro S3, CNR-Istituto di Nanoscienze, Via Campi 213A, 41125 Modena Italy

⁵Univ de Lyon, ENS de Lyon, INRIA, CNRS, UMR 5668, IXXI, 69364 Lyon, France

⁶Centre for Business Network Analysis, University of Greenwich, Park Row, London SE10 9LS, United Kingdom

I. DATASET

The analyzed dataset is the Twitter fire-hose (i.e. all the citations done from all the users) from January the 1st to September the 31st of 2008.

The dataset consists of 536, 210 nodes performing about 160M events and connected by 2.6M edges.

Since the data are daily aggregated we infer the inter-event time distribution for $\tau_i \leq 24$ h by assuming the events done by a node within a single day to be homogeneously distributed during the 24 hours of the day. As we are measuring the α exponent leading the $\Psi(\tau_i) \propto \tau_i^{-(1+\alpha)}$ in the right tail of the distribution this assumption does not change the resulting α .

To measure the reinforcement process and specifically the $\beta = 0.47$ exponent we measure the attachment probability on nodes featuring similar stories, i.e. with a comparable activity a_i (i.e. the number of events actually engaged by the node *i*) and final degree k_i (see [1] for details).

We also checked that sorting the nodes accordingly to their average inter-event time ξ_i and final degree k_i instead of by their activity does not change the measured value of β that reads $\beta = 0.50$ instead of $\beta = 0.47$.

We also considered two other datasets at our disposal (Mobile Phone Calls network and the American Physical Society co-authorship network) but they both fall in the SupBR region as we show in Fig. 1. Given this result we did not further analyzed these datasets as they are already well described by the previous version of the model.



FIG. 1: The waiting-time distribution $\Psi(\tau_i)$ for (a) Mobile phone network (circles) and (b) the co-authorship network of the Physical Review B journal (squares). We also show the fitting curve of the right tail $\Psi(\tau_i) \propto \tau_i^{-(1+\alpha)}$ giving $\alpha \sim 1.45$ for the mobile calls dataset and $\alpha \sim 2.1$ for the PRB one. Given these results, and provided that the (minimum) value of β found in the mobile phone call dataset is $\beta_{\min} = 1.2$ we conclude that both the system are above the $\alpha = (2\beta + 2)/(2\beta + 1)$ curve, thus falling in the SupBR regime.

II. MASTER EQUATION

A.
$$P(k,t)$$

We start from the probability Q(k,t) that an agent makes a call at time t and afterwards its connectivity is k. In the single agent approximation we have:

$$Q(k,t) = \tilde{\mathcal{C}} \int_{\xi_0}^{+\infty} \frac{Q(k-1,t-t')}{t'^{\alpha+1}} \frac{c^{\beta}}{(c+k-1)^{\beta}} dt' + \tilde{\mathcal{C}} \int_{\xi_0}^{+\infty} \frac{Q(k,t-t')}{t'^{\alpha+1}} \left(1 - \frac{c^{\beta}}{(c+k)^{\beta}}\right) dt' + \delta(k,0)\delta(t,0), \quad (1)$$

where α is the exponent driving the inter-event time distribution $P(t) = \tilde{\mathcal{C}}t^{-(\alpha+1)}$ and $\tilde{\mathcal{C}}$ is the normalization constant $\tilde{\mathcal{C}} = \alpha \xi_0^{\alpha}$.

To obtain the probability distribution P(k,t) that the agent has degree k at time t we must integrate Eq. (1) so that:

$$P(k,t) = \int_{\xi_0}^t dt' Q(k,t-t') \int_{t'}^{+\infty} d\xi \frac{1}{\xi^{\alpha+1}}.$$
(2)

Let us perform the Fourier Transform of Eq. (1) in time, by sending the integration variable $t \to (t - t')$ we get:

$$\tilde{Q}(k,\omega) = \tilde{\mathcal{C}}\left[\frac{c^{\beta}}{(c+k-1)^{\beta}}\tilde{Q}(k-1,\omega)\int_{\xi_0}^{\infty}\frac{e^{i\omega t'}}{t'^{\alpha+1}}dt' + \left(1-\frac{c^{\beta}}{(c+k)^{\beta}}\right)\tilde{Q}(k,\omega)\int_{\xi_0}^{\infty}\frac{e^{i\omega t'}}{t'^{\alpha+1}}dt' + \right] + \delta(k,0).$$
(3)

By taking the limit $k \to \infty$ of Eq. (4) we end up with

$$\tilde{Q}(k,\omega) = \tilde{\mathcal{C}}\left[\left(\frac{c}{k}\right)^{\beta} \left[\tilde{Q}(k-1,\omega) - \tilde{Q}(k,\omega)\right] \int_{\xi_0}^{\infty} \frac{e^{i\omega t'}}{t'^{\alpha+1}} dt' + \tilde{Q}(k,\omega) \int_{\xi_0}^{\infty} \frac{e^{i\omega t'}}{t'^{\alpha+1}} dt' + \right] + \delta(k,0).$$

$$\tag{4}$$

B. Integral Contributions

The issue is now to compute the integral appearing in Eq. (4). There are three intervals of the exponent α leading to three different results. In all the three cases we will perform the integral by taking the $\omega \to 0$ limit, i.e. take into account only the long-time, asymptotic region of the solution.

• $0 < \alpha < 1$: in this regime we can take the $\omega \to 0$ limit and expand the exponential term $e^{i\omega t} \sim 1 + i\omega t + \mathcal{O}(t^2)$. However, the first term proportional to t diverges as $\alpha < 1$. We can use the following trick to estimate the first diverging contribution of the integral:

$$\tilde{\mathcal{C}}\frac{\partial}{\partial\omega}\int_{\xi_0}^{\infty}\frac{e^{i\omega t'}}{t'^{\alpha+1}}dt' = i\tilde{\mathcal{C}}\int_{\xi_0}^{\infty}\frac{e^{i\omega t'}}{t'^{\alpha}}dt',\tag{5}$$

and by defining $|\omega|t' = x$ we get

$$i\tilde{\mathcal{C}}\int_{|\omega|\xi_{0}}^{\infty} \frac{e^{ix\operatorname{sign}(\omega)}}{\left(\frac{x}{|\omega|}\right)^{\alpha}} \frac{dt'}{|\omega|} = i\tilde{\mathcal{C}}|\omega|^{\alpha-1} \int_{|\omega|\xi_{0}}^{\infty} \frac{e^{ix\operatorname{sign}(w)}}{x^{\alpha}} dt' = i\tilde{\mathcal{C}}|\omega|^{\alpha-1} \int_{|\omega|\xi_{0}}^{\infty} \frac{\cos(x\operatorname{sign}(\omega)) + i\sin(x\operatorname{sign}(\omega))}{x^{\alpha}} dt' \sim (6)$$
$$i\tilde{\mathcal{C}}|\omega|^{\alpha-1} \left[\frac{(\xi_{0}|\omega|)^{1-\alpha}}{\alpha-1} + \Gamma(1-\alpha) \left[\cos\left(-\frac{\pi\alpha}{2}\right) + i\operatorname{sign}(\omega)\sin\left(-\frac{\pi\alpha}{2}\right)\right]\right].$$

Now we can integrate the last term of Eq. (6) in $d\omega$ to get the leading term of the integral in the $\omega \to 0$ limit

$$\tilde{\mathcal{C}}\int_{\xi_0}^{\infty} \frac{e^{i\omega t'}}{t'^{\alpha+1}} dt' \sim 1 - (\xi_0 \omega)^{\alpha} \frac{\alpha}{\alpha} \Gamma(1-\alpha) \left[\cos\left(-\frac{\pi\alpha}{2}\right) + i\operatorname{sign}(\omega) \sin\left(-\frac{\pi\alpha}{2}\right) \right] = 1 - (\xi_o \omega)^{\alpha} A_{\alpha}, \tag{7}$$

where the 1 comes from the first term of the exponential expansion and where we dropped the terms $\propto \omega$ as they die out faster than the ones $\propto \omega^{\alpha}$. We also encoded the complex constant multiplying the $(\xi_0 \omega)^{\alpha}$ term in the A_{α} symbol.

• $1 < \alpha < 2$: in this case we can apply the same procedure of the previous case but expanding the series to the second order, thus deriving twice with respect to ω . The leading terms of the integral of Eq. (4) are:

$$\tilde{\mathcal{C}} \int_{\xi_0}^{\infty} \frac{e^{i\omega t'}}{t'^{\alpha+1}} dt' \sim 1 + i(\omega\xi_0) \frac{\alpha}{(\alpha-1)} + \frac{(\xi_0\omega)^{\alpha}}{(\alpha-1)} \Gamma(2-\alpha) \left[\cos\left(\frac{\pi\alpha}{2}\right) - i\operatorname{sign}(\omega)\sin\left(\frac{\pi\alpha}{2}\right) \right],\tag{8}$$

and we define $C_{\alpha} = \frac{1}{(\alpha-1)} \Gamma(2-\alpha) \left[\cos\left(\frac{\pi\alpha}{2}\right) - i \operatorname{sign}(\omega) \sin\left(\frac{\pi\alpha}{2}\right) \right].$

• $\alpha > 2$: in this case it is sufficient to consider the first three orders of the integral expansion so that:

$$\tilde{\mathcal{C}} \int_{\xi_0}^{\infty} \frac{e^{i\omega t'}}{t'^{\alpha+1}} dt' \sim 1 + i(\omega\xi_0) \frac{\alpha}{(\alpha-1)} - (\omega\xi_0)^2 \frac{\alpha}{(\alpha-2)}.$$
(9)

C. Asymptotic solution $\alpha < 1$

Using Eq. (7) we can rewrite Eq. (4) as

$$A_{\alpha}(\xi_{0}\omega)^{\alpha}Q(k,w) + \left(\frac{c}{k}\right)^{\beta}\frac{\partial Q(k,\omega)}{\partial k} = \delta(k,0).$$
⁽¹⁰⁾

To solve the equation we introduce the variable $h = k^{1+\beta}$ so that:

$$A_{\alpha}(\xi_0|\omega|)^{\alpha}Q(h,\omega) + c^{\beta}(1+\beta)\frac{\partial Q(h,\omega)}{\partial h} = \delta(h,0).$$
(11)

We Fourier transform this equation in space sending $h \to q$ getting

$$A_{\alpha}(\xi_0|\omega|)^{\alpha}Q(q,\omega) + ic^{\beta}(1+\beta)qQ(q,\omega) = 1,$$
(12)

so that:

$$Q(q,\omega) = \frac{1}{A_{\alpha}(\xi_0|\omega|)^{\alpha} + iqc^{\beta}(1+\beta)}.$$
(13)

Now let us introduce the variable h also in Eq. (2) calling $R(h,\omega) = P(h^{1/(1+\beta)},\omega)$. We then perform the FT of Eq. (2) with respect to h calling q the transformed variable and getting:

$$R(q,\omega) = \tilde{Q}(q,\omega) \int e^{i\omega t'} \int_{t'}^{\infty} \frac{\tilde{\mathcal{C}}}{\xi^{\alpha+1}} dt' d\xi.$$
(14)

The integral on the r.h.s. of the last equation is the same as the one in Eq. (7) with $\alpha \rightarrow \alpha - 1$ so that:

$$R(q,\omega) = \frac{B_{\alpha}|\omega|^{\alpha-1}}{A_{\alpha}(\xi_0|\omega|)^{\alpha} + ic^{\beta}(1+\beta)q},$$
(15)

where $B_{\alpha} = A_{\alpha-1}\xi_0^{\alpha}$. This equation is the same of Eqq. (8) and subsequent in reference [2] therefore we have

$$P(h,t) = \frac{1}{c^{\beta}(\beta+1)(t/\xi_0)^{\alpha}} f_{\alpha}\left(\frac{h}{c^{\beta}(\beta+1)(t/\xi_0)^{\alpha}}\right)$$
(16)

where f_{α} is the Lévy function. Reintroducing the degree variable $k = h^{1/(1+\beta)}$ we find:

$$P(k,t) = \frac{k^{\beta}}{c^{\beta}(t/\xi_{0})^{\alpha}} f_{\alpha} \left(\frac{k^{\beta+1}}{c^{\beta}(\beta+1)(t/\xi_{0})^{\alpha}} \right),$$
(17)

that can be rewritten as:

$$P(k,t) = \frac{1}{t^{\alpha/(1+\beta)}} \tilde{f}_{\alpha\beta} \left(\frac{k}{(t/\xi_0)^{\alpha/(1+\beta)}} \right) = \frac{1}{t^{\alpha/(1+\beta)}} \tilde{f}_{\alpha\beta} \left(\tilde{k} \right), \tag{18}$$

where $\tilde{k} = k/(c^{\beta}(\beta+1)(t/\xi_0)^{\alpha})^{1/(1+\beta)}$ and $\tilde{f}_{\alpha\beta}(\tilde{k})$ is an unknown function of \tilde{k} .

Equations (18) and (17) states that the peak of the P(k,t) distribution (i.e. the average degree) grows as $t^{\alpha/(1+\beta)}$. We can show these last findings in an alternative way. Prompted by the results of the numerical simulations we suppose a scaling form of the P(k,t) that reads:

$$P(k,t) \simeq \frac{1}{t^{\gamma}} P\left(\frac{k^{1+\beta}}{t^{\gamma}}\right) \xrightarrow{h=k^{1+\beta}} P(h,t) \simeq \frac{1}{t^{\gamma}} P\left(\frac{h}{t^{\gamma}}\right).$$
(19)

If we now compute the space and time Fourier transform of Eq. (19) calling q the transformed variable of h we find

$$P(q,\omega) = \int \int e^{i\omega t + iqh} \frac{1}{t^{\gamma}} P\left(\frac{h}{t^{\gamma}}\right) dt dh = \int \int e^{it' + ih'} \frac{1}{\omega} G\left(\frac{h'/q}{(t'/\omega)^{\gamma}}\right) dt' dh' = \frac{1}{\omega} \tilde{G}\left(\frac{q}{\omega^{\gamma}}\right), \tag{20}$$

where we renamed $\omega t \to t'$ and $qh \to h'$.

By comparing the last term of the last line of Eq. (20) with Eq. (15) we find that $\gamma = \alpha$ so that we recover the results shown in Eq. (18).

The resulting scaling regime reads:

$$P(h,t) = \frac{1}{c^{\beta}(\beta+1)(t/\xi_0)^{\alpha}} f_{\alpha\beta} \left(\frac{h}{c^{\beta}(\beta+1)(t/\xi_0)^{\alpha}}\right)$$
(21)

where $f_{\alpha\beta}$ is a non Gaussian scaling function. Reintroducing the connectivity k we get:

$$P(k,t) = \frac{k^{\beta}}{c^{\beta}(t/\xi_0)^{\alpha}} f_{\alpha} \left(\frac{k^{\beta+1}}{c^{\beta}(\beta+1)(t/\xi_0)^{\alpha}} \right) = \left(\frac{t}{\xi_0} \right)^{\frac{-\alpha}{\beta+1}} \tilde{f}_{\alpha\beta} \left(\frac{k}{\left(\frac{t}{\xi_0} \right)^{\alpha/(\beta+1)}} \right), \tag{22}$$

where we grouped the $k/(t/\xi_0)^{\alpha/(\beta+1)}$ terms and where $\tilde{f}_{\beta\alpha}(\tilde{k})$ is an unknown, non-Gaussian scaling function of $k/\left(\frac{t}{\xi_0}\right)^{\alpha/(\beta+1)}$.

Moreover the overall time scaling of the average degree of an activity class reads:

$$k \sim c^{\beta/(1+\beta)} (at)^{\alpha/(1+\beta)} \tag{23}$$

where we used the fact that ξ_0 is the inverse of the activity *a*.

D. Asymptotic solution $1 < \alpha < 2$

First of all let us compute the functional form of the Fourier transformed P(k, t). As in the previous case, we find that the solution follows the:

$$\frac{1}{t^{\gamma}}P\left(\frac{k^{1+\beta}-t}{t^{\gamma}}\right) = \frac{1}{t^{\gamma}}P\left(\frac{h-t}{t^{\gamma}}\right),\tag{24}$$

where $h = k^{1+\beta}$. If we now follow the same procedure as in Equations (19) and following, we find:

$$\int e^{i\omega t + iqh} \frac{1}{t^{\gamma}} P\left(\frac{h - t}{t^{\gamma}}\right) dt dh = \left[\epsilon = h - t\right] = \int e^{i\omega t + iq(\epsilon + t)} \frac{1}{t^{\gamma}} P\left(\frac{\epsilon}{t^{\gamma}}\right) dt dh = \left[(\omega + q)t = t'; q\epsilon = q'\right] = \int e^{it' + i\epsilon'} \left(\frac{\omega + q}{t'}\right)^{\gamma} P\left(\frac{\epsilon'}{q} \left(\frac{\omega + q}{t}\right)^{\gamma}\right) \frac{dt'}{\omega + q} \frac{d\epsilon'}{q} = \frac{1}{\omega + q} g\left(\frac{q}{(\omega + q)^{\gamma}}\right),$$
(25)

so that, calling $\tilde{\omega} = \omega + q$ we have $P(q, \tilde{\omega}) = \frac{1}{\tilde{\omega}} g\left(\frac{q}{\tilde{\omega}\gamma}\right)$. Using Eq. (8) we can rewrite Eq. (4) as

$$Q(k,\omega) = -\left(\frac{c}{k}\right)^{\beta} \frac{\partial Q}{\partial k} + \frac{1}{2} \left(\frac{c}{k}\right)^{\beta} \frac{\partial^2 Q}{\partial k^2} + i\omega \langle \xi \rangle Q(k,\omega) + Q(k,\omega) + \omega^{\alpha} A_{\alpha} Q(k,\omega) + \frac{\beta c^{\beta}}{k^{\beta+1}} Q(k,\omega) + \delta(k,0), \quad (26)$$

where $\langle \xi \rangle$ is the first momentum of the inter event time distribution (i.e. its mean value) and the second-last term comes from the expansion of the $(c/(c+k-1))^{\beta}$ term. We then introduce as in the previous case the $h = k^{\beta+1}$ variable, so that $1/k^{\beta} \cdot \partial/\partial k \to \beta \partial/\partial h$ and $1/k^{\beta} \cdot \partial^2/\partial k^2 \sim (1+\beta)^2 h^{\beta/(\beta+1)} \cdot \partial^2/\partial h^2$. We then Fourier transform with respect to space sending $h \to q$ getting:

$$-i\omega\langle\xi\rangle P(q,\omega) - A_{\alpha}\omega^{\alpha}P(q,\omega) = -i\beta c^{\beta}qP(q,\omega) + \frac{1}{2}(1+\beta)^{2}\int e^{iqh}h^{\beta/(\beta+1)}\frac{\partial^{2}P(h,\omega)}{\partial h^{2}}dh + \beta c^{\beta}\int h^{-1}e^{iqh}P(h,\omega)dh + 1,$$
(27)

where $P(h,\omega) = Q(h^{1/(1+\beta)},\omega) = Q(k,\omega)$. Now let us focus on the integral containing the second-derivative term. We can rewrite it as:

$$\int e^{iqh} h^{\beta/(1+\beta)} \frac{\partial^2}{\partial h^2} \int e^{-i\omega t} P(h,t) dh dt = \iint e^{iqh'} e^{-i\omega' t} (h' + \beta c^\beta t)^{\beta/(1+\beta)} \frac{\partial^2}{\partial h'^2} P((h' + \beta c^\beta t), t) dh' dt,$$
(28)

where we introduced $\omega' = \omega - \beta c^{\beta} q$ and $h' = h - \beta c^{\beta} t$. Note that we can approximate $(h' + \beta c^{\beta} t) \simeq \beta c^{\beta} t$, as we expect $h' \ll t$. If we now integrate by parts Eq. (28) and re-write $P(h' + \beta c^{\beta} t, t)$ as $\int e^{i\tilde{\omega}t + i\tilde{q}h'} P(\tilde{q}, \tilde{\omega}) d\tilde{q} d\tilde{\omega}$ we get:

$$-q^{2} \iint dh' dt e^{iqh'} e^{-i\omega't} (\beta c^{\beta} t)^{\beta/(1+\beta)} \int e^{i\tilde{\omega}t + i\tilde{q}h'} P(\tilde{q}, \tilde{\omega}) d\tilde{q} d\tilde{\omega} = -q^{2} \iint dt d\tilde{\omega} e^{-i\omega't + i\tilde{\omega}t} (\beta c^{\beta} t)^{\beta/(1+\beta)} P(q, \tilde{\omega}), \quad (29)$$

where we used the fact that the integrals in dq and $d\tilde{q}$ give us a $\delta(q, \tilde{q})$. Now we insert our prediction on the $P(q, \tilde{\omega})$ of Eq. (25). In addiction we call $\omega' t = t'$ and, once the substitution is done, $\tilde{\omega}/\omega' = y$ getting:

$$-q^{2} \iint dy dt' e^{-it'+iyt'} (\beta c^{\beta} t'/\omega')^{\beta/(1+\beta)} \tilde{\omega} f\left(\frac{q}{\tilde{\omega}\gamma}\right) = -\frac{q^{2}}{\omega'^{\beta/(1+\beta)}} H\left(\frac{q}{\tilde{\omega}\gamma}\right), \tag{30}$$

where H(x) is an unknown scaling function.

Now, making the $\omega \to \omega'$ substitution and putting the result of Eq. (30) in Eq. (27) we get:

$$-i\omega'\langle\xi\rangle f\left(\frac{q}{\omega^{\gamma}}\right) - A_{\alpha}(\omega' + \beta c^{\beta}q)^{\alpha} f\left(\frac{q}{\omega^{\gamma}}\right) + \frac{1}{2}(1+\beta)^{2}\left(\frac{q}{\omega'^{\frac{1}{2}\beta/(1+\beta)}}\right)^{2} H\left(\frac{q}{\tilde{\omega}^{\gamma}}\right) = 1.$$
(31)

Now we collect an ω' term from all the members to isolate the leading order finding

$$-i\langle\xi\rangle f\left(\frac{q}{\omega\gamma}\right) - A_{\alpha}\left(\omega'^{1-1/\alpha} + \beta c^{\beta}\frac{q}{\omega'^{1/\alpha}}\right)^{\alpha} f\left(\frac{q}{\omega\gamma}\right) + \frac{1}{2}(1+\beta)^{2}\left(\frac{q}{\omega'^{\frac{1}{2}\left(\frac{\beta}{(1+\beta)}+1\right)}}\right)^{2} H\left(\frac{q}{\tilde{\omega}\gamma}\right).$$
(32)

Now if the term proportional to A_{α} is the leading one we find that the exponent $\gamma = 1/\alpha$, otherwise, if the term proportional to $H(q/\omega'^{\gamma})$ leads the evolution we find $\gamma = (2\beta + 1)/(2\beta + 2)$. The separation between the two cases is set at the point where the two exponents are equal, e.g. the A_{α} term wins if

$$\alpha < \frac{2\beta + 2}{2\beta + 1}.\tag{33}$$

In the other case the term proportional to $H(q/\tilde{\omega}^{\gamma})$ wins and we recover the calculation of the previous work (see also Section II E).

Given these results, to show the scaling form of the P(k,t) distribution let us recall the assumed scaling form of the P(k,t) of Eq. (24). As already said the drift of the peak of the distribution goes as $\langle k(t) \rangle \propto t^{1/(1+\beta)}$, so that the variable inside the distribution function reads:

$$\frac{k^{1+\beta} - vt}{t^{\gamma}}.$$
(34)

If we rewrite $k^{\beta+1} = (k - t^{1/(1+\beta)} + t^{1/(1+\beta)})^{1+\beta}$, we can introduce the variable $x = k - t^{1/(1+\beta)} \ll 1$ and expand Eq. (34) in x:

$$k^{1+\beta} \simeq t + (1+\beta)xt^{\beta/(1+\beta)} + \mathcal{O}(x^2).$$
 (35)

By substituting Eq. (35) in Eq. (34) we find the distribution P(k, t) to scale as:

$$\frac{1}{t^{\gamma-\frac{\beta}{1+\beta}}}P\left(\frac{x}{t^{\gamma-\frac{\beta}{1+\beta}}}\right),\tag{36}$$

where $x = k - t^{1/(1+\beta)}$. Then in the Gaussian scaling $\gamma = \frac{2\beta+1}{2(\beta+1)}$ so that the scaling form reads:

$$\frac{1}{t^{\frac{1}{2(1+\beta)}}}P\left(\frac{x}{t^{\frac{1}{2(1+\beta)}}}\right),\tag{37}$$

while in the other case $\gamma = \frac{1}{\alpha}$ so that the scaling form reads:

$$\frac{1}{t^{\frac{1}{\alpha} - \frac{\beta}{1+\beta}}} P\left(\frac{x}{t^{\frac{1}{\alpha} - \frac{\beta}{1+\beta}}}\right).$$
(38)

As a last remark, we stress that, regardless of the α and β exponent, the peak of the distribution, i.e. the average value $\langle k(t) \rangle$ grows as

$$\langle k(t) \rangle \propto k^{\frac{1}{1+\beta}}.$$
(39)

This is in good agreement with numerical simulations and also allows for the prediction of the degree distribution $\rho(k)$ as we will show in Section III.

E. Asymptotic solution $\alpha > 2$

Using Eq. (9) we can rewrite Eq. (4) as

$$Q(k,\omega) = -\left(\frac{c}{k}\right)^{\beta} \frac{\partial Q}{\partial k} + \frac{1}{2} \left(\frac{c}{k}\right)^{\beta} \frac{\partial^2 Q}{\partial k^2} + i\omega \langle \xi \rangle Q(k,\omega) + Q(k,\omega) + \omega^2 \langle \xi^2 \rangle Q(k,\omega) + \frac{\beta c^{\beta}}{k^{\beta+1}} Q(k,\omega) + \delta(k,0), \quad (40)$$

where $\langle \xi^2 \rangle$ is the second moment of the P(t) inter-event time distribution. The calculations in this case can be done in the direct space and we recover the results of the previous work, in particular we find the P(k,t) to scale as a Gaussian with mean value growing as $(t/\xi_0)^{1/(1+\beta)}$ and variance growing as $t^{1/[2(1+\beta)]}$:

$$P(k,t) \simeq \frac{1}{(t/\xi_0)^{\frac{1}{2(1+\beta)}}} \exp\left[-A_\beta \frac{\left(k - C_\beta(t/\xi_0)^{\frac{1}{1+\beta}}\right)^2}{(t/\xi_0)^{1/(1+\beta)}}\right].$$
(41)

This results has the same functional form of the one found in Section IID for the $\alpha > (2\beta + 2)/(2\beta + 1)$ case.

III. DEGREE DISTRIBUTION $\rho(k)$

The degree distribution can be evaluated recalling the scaling form accordingly to the different phases of the system. In particular we found:

$$P(k,t) \simeq \begin{cases} \frac{1}{(t/\xi_0)^{\frac{\alpha}{1+\beta}}} f_{\alpha\beta} \left(A'_{\alpha,\beta} \frac{k}{(t/\xi_0)^{\frac{\alpha}{1+\beta}}} \right) & \text{if } \alpha < 1, \\ \frac{1}{(t/\xi_0)^{\frac{1}{\alpha} - \frac{1}{(1+\beta)}}} f_{\alpha\beta} \left(A'_{\alpha,\beta} \frac{k - v(t/\xi_0)^{1/(1+\beta)}}{(t/\xi_0)^{\frac{1}{\alpha} - \frac{1}{(1+\beta)}}} \right) & \text{if } 1 < \alpha < \frac{2\beta + 2}{2\beta + 1}, \\ \frac{1}{(t/\xi_0)^{\frac{1}{2(1+\beta)}}} \exp\left[-A_{\beta} \frac{\left(k - C_{\beta}(t/\xi_0)^{\frac{1}{1+\beta}} \right)^2}{(t/\xi_0)^{1/(1+\beta)}} \right] & \text{if } \alpha > \frac{2\beta + 2}{2\beta + 1}, \end{cases}$$
(42)

so that we can evaluate, at fixed time t, the $\rho(k)$ distribution as:

$$\rho(k) = \int_{\xi_{\min}}^{\xi_{\max}} F(\xi_0) P(k, t) d\xi_0,$$
(43)

where $F(\xi)$ is the distribution of the lower-cut off ξ , i.e. the lower cut-off of the inter-event time distribution for each agent. If $F(\xi) = \delta(\xi - \xi_0)$ the $\rho(k)$ is trivially the P(k,t). If we instead let the ξ of the system be distributed as

 $F(\xi) \propto \xi^{\nu-1}$ we can evaluate the degree distribution $\rho(k)$. Let us recall that such a distribution of the lower cut-off ξ corresponds to an activity distribution going as:

$$F(a) \propto a^{-(\nu+1)},\tag{44}$$

where we used the fact that $a \propto \xi^{-1}$.

For the $\alpha < 1$ case we get:

$$\rho(k) \propto \int \xi^{\nu-1} \xi^{-\alpha/(1+\beta)} f_{\alpha\beta} \left(\frac{k}{(t/\xi)^{\alpha/(1+\beta)}} \right) d\xi = \int \xi^{\nu-1} \xi^{-\alpha/(1+\beta)} f'_{\alpha\beta} \left(\xi k^{(1+\beta)/\alpha} \right) d\xi = [\xi' = \xi k^{(1+\beta)/\alpha}] = \int k^{-1} \left(\frac{\xi'}{k^{(1+\beta)/\alpha}} \right)^{\nu-1} f'_{\alpha\beta} \left(\xi'^{\alpha/(1+\beta)} \right) \frac{d\xi'}{k^{(1+\beta)/\alpha}} = \mathcal{C}k^{-\left[\frac{1+\beta}{\alpha}\nu+1\right]},$$
(45)

where C is a constant with respect to k.

For $\alpha > 1$ we can use the scaling variable $x = k - t^{1/(1+\beta)}$ as a $\delta(k - t^{1/(1+\beta)})$ so that we recover the result of the previous work:

$$\rho(k) \propto k^{-[(1+\beta)\nu+1]}.$$
(46)

- Enrico Ubaldi, Nicola Perra, Márton Karsai, Alessandro Vezzani, Raffaella Burioni, and Alessandro Vespignani. Asymptotic theory for the dynamic of networks with heterogenous activity and social ties allocation. arXiv preprint arXiv:1509.04563, 2015.
- [2] R Burioni, G Gradenigo, A Sarracino, A Vezzani, and A Vulpiani. Rare events and scaling properties in field-induced anomalous dynamics. Journal of Statistical Mechanics: Theory and Experiment, 2013(09):P09022, 2013.