

Commuting differential operators of rank 2 with rational coefficients

Vardan Oganessian

Department of Geometry and Topology, Faculty of Mechanics and Mathematics, Lomonosov Moscow State University, Moscow, 119991 Russia.

E-mail address: vardan.o@mail.ru

Abstract. In this paper we find new self-adjoint commuting operators of rank 2 with rational coefficients and prove that any elliptic and hyperelliptic curves of genus 2 are spectral curves of commuting operators with rational coefficients. Also the case when curves of genus 3 are spectral curves of commuting operators with rational coefficients is studied.

Introduction

The commutativity condition of two differential operators

$$L_n = \sum_{i=0}^n u_i(x) \partial_x^i, \quad L_m = \sum_{i=0}^m v_i(x) \partial_x^i$$

is equivalent to a very complicated system of non-linear differential equations. If two differential operators commute, then there is a nonzero polynomial $R(z, w)$ such that $R(L_n, L_m) = 0$ (see [1]). The curve Γ defined by $R(z, w) = 0$ is called the *spectral curve*. The genus of the curve $R(z, w) = 0$ is called the genus of commuting pair. If

$$L_n \psi = z \psi, \quad L_m \psi = w \psi,$$

then $(z, w) \in \Gamma$. For almost all $(z, w) \in \Gamma$ the dimension of the space of common eigenfunctions ψ is the same. The dimension of the space of common eigenfunctions for generic $P = (z, w) \in \Gamma$ is called the *rank*. The rank is a common divisor of m and n .

This research was supported by grant RSF 16-11-10260 and was done at the faculty of Mechanics and Mathematics, department of Geometry and Topology of Moscow State University

Commutative rings of such operators were classified by Krichever [3]. The ring is determined by the spectral curve and some additional spectral data. If the rank equals 1, then there are explicit formulas for coefficients of commutative operators in terms of Riemann theta-functions (see [2]).

If the rank equals 1, then common eigenfunction $\psi(x, P)$ is Baker-Akhiezer function, where $P \in \Gamma$. Baker-Akhiezer function is a function with the following properties.

- 1) Function $\psi(x, P)$ has one essential singularity at a fixed point $q \in \Gamma$

$$\psi(x, P) = e^{kx} \left(1 + \frac{\xi_1(x)}{k} + \frac{\xi_2(x)}{k^2} + \dots \right),$$

where k^{-1} is a local parameter in a neighborhood of q .

- 2) Function ψ has simple poles at some points $\gamma_1, \dots, \gamma_g$, where g is the genus of Γ .

The set $\{\Gamma, q, \gamma_1, \dots, \gamma_g\}$ is called the spectral data. If we take the spectral data where $D = \gamma_1 + \dots + \gamma_g$ is a non-special divisor, then there is a unique function $\psi(x, P)$ satisfying the conditions 1) and 2). Let us mention that if the spectral curve is singular, then in general ψ is not a function on the spectral curve, but ψ is a section of a torsion-free sheaf on $\Gamma \setminus \{q\}$ (see [13]).

The case when rank is greater than 1 is much more difficult. The first examples of commuting ordinary differential operators of the nontrivial rank 2 and the nontrivial genus $g = 1$ were constructed by Dixmier [12] for the nonsingular elliptic spectral curve $w^2 = z^3 - \alpha$, where α is an arbitrary nonzero constant:

$$L = (\partial_x^2 + x^3 + \alpha)^2 + 2x,$$

$$M = (\partial_x^2 + x^3 + \alpha)^3 + 3x\partial_x^2 + 3\partial_x + 3x(x^2 + \alpha).$$

Operators L and M is the commuting pair of the Dixmier operators of rank 2, genus 1.

The general classification of commuting ordinary differential operators of rank greater than 1 was obtained by Krichever [3]. The general form of commuting operators of rank 2 for an arbitrary elliptic spectral curve was found by Krichever and Novikov [4]. The general form of operators of rank 3 for an arbitrary elliptic spectral curve was found by Mokhov [5], [6]. Moreover, examples of commuting ordinary differential operators of arbitrary

genus and arbitrary rank with polynomial coefficients were constructed in [8], [7].

Mironov in [9] constructed examples of commuting operators

$$L = (\partial_x^2 + A_3x^3 + A_2x^2 + A_1x + A_0)^2 + g(g+1)A_3x,$$

$$M^2 = L^{2g+1} + a_{2g}L^{2g} + \dots + a_1L + a_0,$$

where a_i are some constants and $A_i, A_3 \neq 0$, are arbitrary constants. Operators L and M are commuting operators of rank 2, genus g .

Let us suppose that function \mathcal{P} satisfies the equation

$$(\mathcal{P}'(x))^2 = g_2\mathcal{P}^2(x) + g_1\mathcal{P}(x) + g_0, \quad g_2 \neq 0,$$

where g_1 and g_2 are arbitrary constants. In [10] it was proved by Mironov that operators L_1 and M_1 ,

$$L_1 = (\partial_x^2 + \alpha_1\mathcal{P}(x) + \alpha_0)^2 + \alpha_1g_2g(g+1)\mathcal{P}(x), \quad \alpha_1 \neq 0,$$

$$M_1^2 = L_1^{2g+1} + a_{2g}L_1^{2g} + \dots + a_1L_1 + a_0,$$

where a_i are some constants, α_i are arbitrary constants, is a commuting pair of rank 2, genus g .

Let $\wp(x)$ be the Weierstrass elliptic function satisfying the equation $\wp'(x) = 4\wp^3(x) + g_2\wp(x) + g_3$. Mironov proved in [10] that operators L_2 and M_2 ,

$$L_2 = (\partial_x^2 + \alpha_1\wp(x) + \alpha_0)^2 + s_1\wp(x) + s_2\wp^2(x),$$

$$M_2^2 = L_2^{2g+1} + b_{2g}L_2^{2g} + \dots + b_1L_2 + b_0,$$

where b_i are some constants, $\alpha_1 = \frac{1}{4} - 2g^2 - 2g$, $s_1 = \frac{1}{4}g(g+1)(16\alpha_0 + 5g_2)$, $s_2 = -4g(g+2)(g^2 - 1)$ and α_0 is an arbitrary constant, are also a commuting pair of rank 2, genus g .

Examples of commuting operators of rank two with trigonometric coefficients were found in [14].

Let us consider

$$L_1 = (\partial_x^2 + A_6x^6 + A_2x^2)^2 + 16g(g+1)A_6x^4,$$

$$L_2 = (\partial_x^2 + A_4x^4 + A_2x^2 + A_0)^2 + 4g(g+1)A_4x^2,$$

where $g \in \mathbb{N}$, $A_6 \neq 0$, $A_4 \neq 0$, A_2, A_0 are arbitrary constants. Operators L_1 and L_2 commute with some differential operators M_1 and M_2 of order $4g+2$

respectively (see [15], [16]). The spectral curves of these operators have the form $w^2 = z^{2m+1} + a_{2m}z^{2m} + \dots + a_1z + a_0$. Common eigenfunction of operators L_1, M_1 and L_2, M_2 in some special cases were found in [18]. Moreover, the following theorems are proved in [15], [16].

- 1) If $L = (\partial_x^2 + A_n x^n + A_{n-1} x^{n-1} + \dots + A_0)^2 + B_k x^k + B_{k-1} x^{k-1} + \dots + B_0$, where $n > 3, n \in \mathbb{N}, A_n \neq 0, B_k \neq 0$, commutes with a differential operator M of order $4g + 2$ and M, L are operators of rank 2, then $k = n - 2$ and $B_k = (n - 2)^2 m(m + 1) A_n$ for some $m \in \mathbb{N}$.
- 2) For $n > 6$ operator $L = (\partial_x^2 + A_n x^n)^2 + B_{n-2} x^{n-2}$ does not commute with any differential operator M of order $4g + 2$, where M and L could be a pair of rank 2.
- 3) If $n = 5$, then $L = (\partial_x^2 + Ax^5)^2 + 18Ax^3, A \neq 0$, commutes with a differential operator M of order $4g + 2$ for all g and M, L are operators of rank 2.
- 4) The operator $(\partial_x^2 + Ax^5)^2 + 9m(m + 1)Ax^3, A \neq 0$ does not commute with any differential operator M of order $4g + 2$, for $m > 1$, where M and L could be a pair of rank 2.

Commuting operators L_4 and M , where $L_4 = \partial_x^4 + u(x)$ and $M^2 = L_4^{2g+1} + \beta_{2g} L_4^{2g} + \dots + \beta_0$ were studied in [17].

Mironov and Zhegllov (see [11]) proved that for arbitrary integer m and arbitrary spectral curve Γ given by equation $w^2 = z^3 + c_2 z^2 + c_1 z + c_0$ there are polynomials

$$V_m = \alpha_{m+2} x^{m+2} + \dots + \alpha_0, \quad W_m = \beta_m x^m + \dots + \beta_0$$

such that operator

$$L_{4,m} = (\partial_x^2 + V_m)^2 + W_m$$

commutes with a six order operator $L_{6,m}$. And the spectral curve $L_{4,m}, L_{6,m}$ coincides with Γ . Also, Mironov and Zhegllov studied in [11] automorphisms of the first Weyl algebra.

Theorem 1. *The operator*

$$L = (\partial_x^2 + A_6 x^6 + A_2 x^2 + \frac{A_{-2}}{x^2} + \frac{A_{-6}}{x^6})^2 + 16g(g + 1)A_6 x^4 + B_0, \quad (1)$$

where $g \in \mathbb{N}, A_6 \neq 0$ and A_2, A_{-2}, A_{-6}, B_0 are arbitrary constants, commutes with a differential operator M of order $4g + 2$. The spectral curve has the

form $w^2 = z^{2m+1} + a_{2m}z^{2m} + \dots + a_1z + a_0$. The operators L and M are operators of rank 2.

Theorem 2. *The operator*

$$L = (\partial_x^2 + A_4x^4 + A_2x^2 + A_0 + \frac{A_{-2}}{x^2})^2 + 4g(g+1)A_4x^2 + B_0, \quad (2)$$

where $g \in \mathbb{N}$, $A_4 \neq 0$ and A_2, A_0, A_{-2}, B_0 are arbitrary constants, commutes with a differential operator M of order $4g + 2$. The spectral curve has the form $w^2 = z^{2m+1} + a_{2m}z^{2m} + \dots + a_1z + a_0$. The operators L and M are operators of rank 2.

Theorem 3.

1) *Any elliptic curve*

$$w^2 = z^3 + \beta_2z^2 + \beta_1z + \beta_0,$$

where β_i are arbitrary constants, is spectral curve of commuting operators from Theorem 1.

2) *Any hyperelliptic curve of the form*

$$w^2 = z^5 + \beta_4z^4 + \beta_3z^3 + \beta_2z^2 + \beta_1z + \beta_0,$$

where β_i are arbitrary constants, is spectral curve of commuting operators from Theorems 1 or 2.

3) *And any hyperelliptic curve of the form*

$$w^2 = z^7 + \beta_6z^6 + \beta_5z^5 + \beta_4z^4 + \beta_3z^3 + \beta_2z^2 + F_1z + F_0,$$

where β_i are arbitrary constant and $F_1 = F_1(\beta_6, \dots, \beta_2)$ and $F_0 = F_0(\beta_6, \dots, \beta_2)$ are some constants, is spectral curve of commuting operators from Theorems 1 or 2.

Note that Theorem 3 doesn't say that any hyperelliptic curve of genus 2 is spectral curve of operators only from Theorem 1 or only from Theorem 2. But any hyperelliptic curve of genus 2 is spectral curve of operators from Theorem 1 or Theorem 2.

Acknowledgments

The author wishes to express gratitude to Professor O. I. Mokhov for advices and help in writing this paper.

Commuting operators of rank 2

Let us consider the operator

$$L = (\partial_x^2 + V(x))^2 + W(x) \quad (3)$$

It is proved in [9] that the operator L commutes with an operator M of order $4g+2$, operators L, M are operators of rank 2 and the spectral curve of L, M has the form

$$w^2 = z^{2g+1} + \beta_{2g}z^{2g} + \dots + \beta_0$$

if and only if there exists a polynomial

$$Q = z^g + a_1(x)z^{g-1} + a_2(x)z^{g-2} + \dots + a_{g-1}(x)z + a_g(x)$$

that the following relation is satisfied

$$Q^{(5)} + 4VQ''' + 6V'Q'' + 2Q'(2z - 2W + V'') - 2QW' \equiv 0, \quad (4)$$

where Q' means $\partial_x Q$. The spectral curve of commuting pair L, M has the form

$$4w^2 = 4F(z) = 4(z-W)Q^2 - 4V(Q')^2 + (Q'')^2 - 2Q'Q''' + 2Q(2V'Q' + 4VQ'' + Q^{(4)}). \quad (5)$$

Using (4), we have

$$4Q'z \equiv -Q^{(5)} - 4VQ''' - 6V'Q'' - 2Q'V'' + 2QW' + 4Q'W.$$

We get the following system.

$$\begin{cases} a_1 = W/2 + C_1 \\ 4a'_2 = -a_1^{(5)} - 4Va_1''' - 6V'a_1'' - 2a_1'V'' + 2a_1W' + 4a_1'W \\ \dots \\ 4a'_{i+1} = -a_i^{(5)} - 4Va_i''' - 6V'a_i'' - 2a_i'V'' + 2a_iW' + 4a_i'W \\ \dots \\ 4a'_g = -a_{g-1}^{(5)} - 4Va_{g-1}''' - 6V'a_{g-1}'' - 2a'_{g-1}V'' + 2a_{g-1}W' + 4a'_{g-1}W \\ 0 = -a_g^{(5)} - 4Va_g''' - 6V'a_g'' - 2a_g'V'' + 2a_gW' + 4a_g'W \end{cases}$$

It is clear from the system that the result in [9] can be formulated in the following way. Let $a_1 = W/2 + C_1$, where C_1 is an arbitrary constant. Define a_i by recursion

$$a_{i+1} = C_{i+1} + \frac{1}{4} \int (-a_i^{(5)} - 4V a_i''' - 6V' a_i'' - 2a_i' V'' + 2a_i W' + 4a_i' W) dx \quad (6)$$

We see that (3) commutes with an operator of order $4g+2$ and these operators are operators of rank 2 if and only if there exist constants C_1, \dots, C_g such that $a_{g+1} \equiv \text{const}$.

For example,

$$a_2 = -\frac{1}{8}W^{(4)} - \frac{1}{2}VW'' - \frac{1}{4}V'W' + \frac{3}{8}W^2 + \frac{1}{2}WC_1 + 2C_2,$$

where C_2 is a constant, which appears after integration. In general, a_i contains i constants, i.e., $a_i(x) = a_i(x; C_1, \dots, C_i)$.

Proof of Theorem 1.

The operator L has the form

$$L = (\partial_x^2 + A_6 x^6 + A_2 x^2 + \frac{A_{-2}}{x^2} + \frac{A_{-6}}{x^6})^2 + 16A_6 g(g+1)x^4 + B_0. \quad (7)$$

Let us prove that we can choose constants C_1, C_2, \dots, C_g in such a way that $a_{g+1} = \text{const}$. Calculation shows that

$$a_1 = C_1 + \frac{B_0}{2} + 8A_6 g(g+1)x^4.$$

We see from (6) that a_{i+1} is polynomial in x and linear in a_i . Let us apply (6) to x^{4k} . Assuming that $a_i = x^{4k}$ we get

$$a_{i+1} = C_{i+1} - 16A_{-6}k(k-1)x^{4k-8} - k(4k-2)(3+4A_{-2}-16k+16k^2)x^{4k-4} + (B_0 - 16A_2k^2)x^{4k} + \frac{8A_6(g-k)(g+k+1)(2k+1)}{k+1}x^{4k+4}.$$

Let us calculate calculate a_2 and a_3

$$a_2 = \tilde{C}_2 + 4A_6(2C_1 + 3B_0 - 32A_2)g(g+1)x^4 + 96A_6^2g(g+1)(g-1)(g+2)x^8, \quad (8)$$

$$\begin{aligned}
a_3 = & \tilde{C}_3 + 4A_6g(1+g)((16A_2 - B_0)(32A_2 - 3B_0 - 2C_1) + \\
& + 2(\tilde{C}_2 - 144A_6(35 + 4A_{-2})(g-1)(g+2)))x^4 - \\
& - 48A_6^2(160A_2 - 5B_0 - 2C_1)g(g+1)(g-1)(g+2)x^8 + \\
& + 1280A_6^3(g-2)(g-1)g(g+1)(g+2)(g+3)x^{12},
\end{aligned} \tag{9}$$

where \tilde{C}_i are constants and depend on C_i .

If $a_i = \tilde{C}_i + K_i^1x^4 + K_i^2x^8 + \dots + K_i^ix^{4i}$, then

$$\begin{aligned}
a_{i+1} = & \tilde{C}_{i+1} + (8A_6g(g+1)\tilde{C}_i + (B_0 - 16A_2)K_i^1 - 2 \cdot 6(35 + 4A_{-2})K_i^2 - 16 \cdot 2 \cdot 3A_{-6}K_i^3)x^4 + \\
& + (const \cdot K_i^1 + const \cdot K_i^2 - const \cdot K_i^3 - const \cdot K_i^4)x^8 + \dots + \\
& + \frac{8A_6(2i+1)(g-i)(g+i+1)K_i^ix^{4i+4}}{i+1}.
\end{aligned}$$

Note that all coefficients, except the leading term, contain constants C_1, C_2, \dots . And K_i^{i-1} contains only C_1 , K_i^{i-2} contains only C_1 and C_2 , K_i^{i-3} only C_1, C_2, C_3 etc.

So, the term $const \cdot x^{4(g+1)}$ in a_{g+1} vanishes because it is multiplied by $const \cdot (g-g)(g+g+1) = 0$. Therefore, we must choose the constants C_1, C_2, \dots to vanish $K_{g+1}^m = 0$ for all m . We always can do it because the leading term in a_{g+1} equals $const \cdot x^{4g}$ and depends only on C_1 , penultimate on C_1 and C_2 etc.

Theorem 1 is proved.

Proof of Theorem 2.

The operator L has the form

$$L = (\partial_x^2 + A_4x^4 + A_2x^2 + A_0 + \frac{A_{-2}}{x^2})^2 + 4g(g+1)A_4x^2. \tag{10}$$

As in the previous proof, we want to find constants C_1, C_2, \dots, C_g such that $a_{g+1} = const$. Calculation shows that

$$a_1 = C_1 + \frac{B_0}{2} + 2A_4g(g+1)x^2.$$

We mentioned before that a_{i+1} is polynomial in x and linear in a_i . Assume that $a_i = x^{2k}$, then from (6) we have

$$a_{i+1} = C_{i+1} - k(k-1)(3 + 4A_{-2} + 4k(k-2))x^{2k-4} - 2A_0k(2k-1)x^{2k-2} + (B_0 - 4A_2k^2)x^{2k} + \frac{2A_4(2k+1)(g-k)(g+k+1)x^{2k+2}}{k+1}. \quad (11)$$

Let us calculate a_2 and a_3

$$a_2 = \widetilde{C}_2 + A_4(2C_1 + 3B_0 - 8A_2)g(g+1)x^2 + 6A_4^2g(g+1)(g-1)(g+2)x^4, \quad (12)$$

$$a_3 = \widetilde{C}_3 + A_4g(g+1) \left((4A_2 - B_0)(8A_2 - 3B_0 - 2C_1) + 2\widetilde{C}_2 - 72A_0A_4(g-1)(g+2) \right) x^2 - 3A_4^2(40A_2 - 5B_0 - 2C_1)g(g+1)(g-1)(g+2)x^4 + 20A_4^3g(g+1)(g-1)(g+2)(g-2)(g+3)x^6, \quad (13)$$

where \widetilde{C}_i are constants and depend on C_i .

If $a_i = \widetilde{C}_i + K_i^1x^2 + K_i^2x^4 + \dots + K_i^ix^{2i}$, then

$$a_{i+1} = \widetilde{C}_{i+1} + \left(2A_4g(g+1)\widetilde{C}_i - (B_0 - 4A_2)K_i^1 - 12A_0K_i^2 - 6(15 + 4A_{-2})K_i^3 \right) x^2 + (const \cdot K_i^1 - const \cdot K_i^2 - const \cdot K_i^3 - const \cdot K_i^4) x^4 + \dots + \frac{2A_4(2i+1)}{i+1}(g-i)(g+i+1)K_i^ix^{2i+2}.$$

As in the proof of the previous Theorem all terms except the leading term contain constants C_1, C_2, \dots . And K_i^{i-1} contains only C_1 , K_i^{i-2} contains only C_1, C_2 and K_i^{i-3} only C_1, C_2, C_3 etc.

The term $const \cdot x^{2(g+1)}$ in a_{g+1} vanishes because it is multiplied by $\frac{2A_4(2g+1)}{g+1}(g-g)(g+g+1) = 0$. So we must choose constants C_1, C_2, \dots in such a way that $A_{g+1}^i = 0$ for all i . We always can do it because the leading term in a_{g+1} depends only on C_1 , penultimate on C_1 and C_2 etc.

Theorem 2 is proved.

Proof of Theorem 3.

1) From Theorem 1 we see that the operator

$$L = \left(\partial_x^2 + A_6 x^6 + A_2 x^2 + \frac{A_{-2}}{x^2} + \frac{A_{-6}}{x^6} \right)^2 + 32A_6 x^4 + B_0$$

commutes with an operator M of order 6. From (8) we see that if $g = 1$, then we must take $C_1 = \frac{32A_2 - 3B_0}{2}$. Using formula (5), where $Q = z + a_1(x)$ we find the spectral curve of operators L and M

$$w^2 = z^3 + \beta_2 z^2 + \beta_1 z + \beta_0,$$

where

$$\begin{aligned} \beta_2 &= (32A_2 - 3B_0), \\ \beta_1 &= (256A_2^2 - 64A_2B_0 + 3B_0^2 + 64A_6(4A_{-2} + 3)), \\ \beta_0 &= -(16A_2 - B_0)((16A_2 - B_0)B_0 - 64A_6(4A_{-2} + 3)) - 4096A_6^2A_{-6}. \end{aligned}$$

So, we must solve the system of equations in $A_6, A_2, A_{-2}, A_{-6}, B_0$

$$\begin{cases} \beta_2 = q_2 \\ \beta_1 = q_1 \\ \beta_0 = q_0 \end{cases}$$

where q_2, q_1 and q_0 are arbitrary constants. The solution is

$$\begin{aligned} A_2 &= \frac{3B_0 + q_2}{32}, \\ A_{-2} &= \frac{3B_0^2 - 768A_6 + 4q_1 + 2B_0q_2 - q_2^2}{1024A_6}, \\ A_{-6} &= \frac{B_0^3 - 8q_0 + 4B_0q_1 + B_0^2q_2 + 4q_1q_2 - B_0q_2^2 - q_2^3}{32768A_6^2}, \end{aligned}$$

where $A_6 \neq 0, B_0$ are arbitrary constants.

2) From Theorem 2 we know that the operator

$$L = \left(\partial_x^2 + A_4 x^4 + A_2 x^2 + A_0 + \frac{A_{-2}}{x^2} \right)^2 + 24A_4 x^2 + B_0$$

commutes with an operator M of order 10. Again, using formula (5), (12) and (13) we find the spectral curve of operators L and M

$$w^2 = z^5 + \beta_4 z^4 + \beta_3 z^3 + \beta_2 z^2 + \beta_1 z + \beta_0,$$

where

$$\beta_4 = 40A_2 - 5B_0,$$

$$\beta_3 = 2(264A_2^2 + 168A_0A_4 - 80A_2B_0 + 5B_0^2),$$

$$\beta_2 = 2(1280A_2^3 - 792A_2^2B_0 - 504A_0A_4B_0 - 5B_0^3 + 24A_2(156A_0A_4 + 5B_0^2) + 72A_4^2(13 + 12A_{-2})),$$

$$\begin{aligned} \beta_1 = & 4096A_2^4 + 27648A_0^2A_4^2 - 5120A_2^3B_0 - 3744A_4^2B_0 + 1008A_0A_4B_0^2 + 5B_0^4 + \\ & + 144A_2^2(256A_0A_4 + 11B_0^2) - 3456A_4^2B_0A_{-2} + 32A_2(72A_4^2(17 + 12A_{-2}) - 468A_0A_4B_0 - 5B_0^3), \end{aligned}$$

$$\begin{aligned} \beta_0 = & 27648A_0^2A_4^2(4A_2 - B_0) + 48A_0A_4(1024A_2^3 - 768A_2^2B_0 + 156A_2B_0^2 - 7B_0^3 + \\ & + 1728A_4^2(3 + 4A_{-2})) - (16A_2 - B_0)(256A_2^3B_0 - 144A_2^2B_0^2 + \\ & + 24A_2(B_0^3 - 384A_4^2) + B_0(144A_4^2(13 + 12A_{-2})) - B_0^3). \end{aligned}$$

So, we again must solve the system of equations in $A_4, A_2, A_0, A_{-2}, B_0$

$$\begin{cases} \beta_4 = q_4 \\ \beta_3 = q_3 \\ \beta_2 = q_2 \\ \beta_1 = q_1 \\ \beta_0 = q_0 \end{cases} \quad (14)$$

where q_4, q_3, q_2, q_1, q_0 are arbitrary constants. From the first equation we find that

$$B_0 = \frac{40A_2 - q_4}{5}.$$

Then we obtain that

$$A_0 = \frac{560A_2^2 + 5q_3 - 2q_4^2}{1680A_4}$$

$$A_{-2} = \frac{11200A_2^3 - 327600A_4^2 + 175q_2 + 300A_2q_3 - 105q_3q_4 - 120A_2q_4^2 + 28q_4^3}{302400A_4^2}$$

$$A_2 = \frac{6125q_1 - 1500q_3^2 - 2450q_2q_4 + 1935q_3q_4^2 - 387q_4^4}{56448000A_4^2}$$

Hence the last equation of (14) has the form

$$-\frac{2304}{35}A_4^2(5q_3 - 2q_4^2) +$$

$$+ \frac{(-6125q_1 + 1500q_3^2 + 2450q_2q_4 - 1935q_3q_4^2 + 387q_4^4)^2}{8643600000A_4^2} +$$

$$+ \frac{625(7q_1 - 12q_3^2)q_4 + 5175q_3q_4^3 - 828q_4^5 + 125q_2(100q_3 - 47q_4^2)}{21875} =$$

$$= A_4^2h_2 + \frac{h_{-2}}{A_4^2} + h_0 = q_0,$$

where h_i are coefficients of the equation. We want to find solution $A_4 \neq 0$.

If $q_4 = q_3 = q_2 = q_1 = 0$ and $q_0 = 1$, then left side equals zero and right side equals 1. Hence for some q_i we have not solution.

There doesn't exist solution $A_4 \neq 0$ if and only if one of the following system is satisfied

$$\begin{cases} h_2 = 0 \\ h_{-2} = 0 \\ h_0 \neq q_0 \end{cases}, \quad \begin{cases} h_2 = 0 \\ h_{-2} \neq 0 \\ h_0 = q_0 \end{cases}, \quad \begin{cases} h_2 \neq 0 \\ h_{-2} = 0 \\ h_0 = q_0 \end{cases}. \quad (15)$$

Now let us consider operator (1) from Theorem 1. If $g = 2$, then using (5), (8) and (9), we get that the spectral curve of operators L and M from Theorem 1 has the form

$$w^2 = z^5 + \gamma_4z^4 + \gamma_3z^3 + \gamma_2z^2 + \gamma_1z + \gamma_0,$$

where

$$\gamma_4 = 160A_2 - 5B_0,$$

$$\gamma_3 = 8448A_2^2 - 640A_2B_0 + 10B_0^2 + 192A_6(213 + 28A_{-2}),$$

$$\begin{aligned} \gamma_2 = & 2(81920A_2^3 - 12672A_2^2B_0 - 5B_0^3 - 288A_6B_0(213 + 28A_{-2}) + \\ & + 96A_2(5B_0^2 + 48A_6(359 + 52A_{-2})) + 55296A_6^2A_{-6}), \end{aligned}$$

$$\begin{aligned} \gamma_1 = & 1048576A_2^4 - 327680A_2^3B_0 + 5B_0^4 + 576A_6B_0^2(213 + 28A_{-2}) + \\ & + 2304A_2^2(11B_0^2 + 1024A_6(19 + 4A_{-2})) + 128A_2(55296A_6^2A_{-6} - 5B_0^3 - 144A_6B_0(359 + 52A_{-2})) + \\ & + 221184A_6^2(1890 + 496A_{-2} + 32A_{-2}^2 - B_0A_{-6}), \end{aligned}$$

$$\begin{aligned} \gamma_0 = & -1048576A_2^4B_0 - B_0^5 - 192A_6B_0^3(213 + 28A_{-2}) + \\ & + 32768A_2^3(5B_0^2 + 384A_6(3 + 4A_{-2})) - 768A_2^2B_0(11B_0^2 + 3072A_6(19 + 4A_{-2})) + \\ & + 84934656A_6^3(35 + 4A_{-2})A_{-6} + 110592A_6^2B_0(-3780 - 992A_{-2} - 64A_{-2}^2 + B_0A_{-6}) + \\ & + 32A_2(5B_0^4 - 288A_6B_0^2(359 + 52A_{-2}) - 221184A_6^2(105 + 152A_{-2} + 16A_{-2}^2 - B_0A_{-6})). \end{aligned}$$

So, the system (14) for operator (1) has the form

$$\begin{cases} \gamma_4 = q_4 \\ \gamma_3 = q_3 \\ \gamma_2 = q_2 \\ \gamma_1 = q_1 \\ \gamma_0 = q_0 \end{cases} \quad (16)$$

Solving the first, second and the third equations of (16) we get

$$\begin{aligned} B_0 &= \frac{160A_2 - q_4}{5} \\ A_{-2} &= \frac{8960A_2^2 - 204480A_6 + 5q_3 - 2q_4^2}{26880A_6} \\ A_{-6} &= \frac{716800A_2^3 + 58982400A_2A_6 + 175q_2 + 1200A_2q_3 - 105q_3q_4 - 480A_2q_4^2 + 28q_4^3}{19353600A_6^2} \end{aligned}$$

And system (16) is equivalent to the system

$$\begin{cases} g_1 = 6125q_1 \\ g_0 = 153125q_0 \end{cases} ,$$

where

$$g_1 = 1500q_3^2 - 28901376000A_2^2A_6 - 42467328000A_6^2 + 2450q_2q_4 - \\ -1935q_3q_4^2 + 387q_4^4 + 460800A_6(5q_3 - 2q_4^2)$$

$$g_0 = 87500(6144A_6(q_2 - 2048A_2(7A_2^2 - 72A_6)) + (147456A_2A_6 + q_2)q_3) - \\ -45000(3211264A_2^2A_6 + (768A_6 + q_3)(6144A_6 + q_3))q_4 - \\ -2625(1966080A_2A_6 + 11q_2)q_4^2 + 150(542720A_6 + 177q_3)q_4^3 - 3861q_4^5$$

To prove the part 2 of Theorem 3 we must show that the following systems have solution.

$$\left\{ \begin{array}{l} h_2 = 0 \\ h_{-2} = 0 \\ h_0 \neq q_0 \\ g_1 = 6125q_1 \\ g_0 = 153125q_0 \\ A_6 \neq 0 \end{array} \right. , \quad (17)$$

$$\left\{ \begin{array}{l} h_2 = 0 \\ h_{-2} \neq 0 \\ h_0 = q_0 \\ g_1 = 6125q_1 \\ g_0 = 153125q_0 \\ A_6 \neq 0 \end{array} \right. , \quad (18)$$

$$\left\{ \begin{array}{l} h_2 \neq 0 \\ h_{-2} = 0 \\ h_0 = q_0 \\ g_1 = 6125q_1 \\ g_0 = 153125q_0 \\ A_6 \neq 0 \end{array} \right. \quad (19)$$

From system (17) we get

$$\left\{ \begin{array}{l} q_3 = \frac{2q_4^2}{5} \\ q_1 = \frac{2q_2q_4}{5} - \frac{3q_4^4}{125} \\ q_0 \neq \frac{q_2q_4^2}{25} - \frac{9q_4^5}{3125} \\ A_6 = -\frac{49A_2^2}{72} \\ \frac{822083584}{3}A_2^5 - \frac{7168(25q_2 - 2q_4^3)}{75}A_2^2 + \frac{q_2q_4^2}{25} - \frac{9q_4^5}{3125} = q_0 \\ A_6 \neq 0 \end{array} \right.$$

If equation $\frac{822083584}{3}A_2^5 - \frac{7168(25q_2 - 2q_4^3)}{75}A_2^2 + \frac{q_2q_4^2}{25} - \frac{9q_4^5}{3125} = q_0$ has so-

lution $A_2 = 0$, then $q_0 = \frac{q_2q_4^2}{25} - \frac{9q_4^5}{3125}$ but it contradicts the third equation.

So, we obtain that system (17) always has solution.

Now let us consider system (18). We have

$$\left\{ \begin{array}{l} q_3 = \frac{2q_4^2}{5} \\ 125q_1 - 50q_2q_4 + 3q_4^4 \neq 0 \\ q_0 = \frac{q_4(625q_1 - 125q_2q_4 + 6q_4^4)}{3125} \\ -4718592A_2^2A_6 = \frac{125q_1 - 50q_2q_4 + 3q_4^4}{125} + \frac{339738624}{49}A_6^2 \\ A_2\left(\frac{3623878656}{7}A_6^2 + 50331648\left(\frac{125q_1 - 50q_2q_4 + 3q_4^4}{4718592 \cdot 125} + \frac{339738624}{4718592 \cdot 49}A_6^2\right)\right) = \\ \quad = -\frac{24576}{175}A_6(25q_2 - 2q_4^3) \\ A_6 \neq 0 \end{array} \right.$$

This system has solution because if $A_6 = 0$ is solution of

$$\left\{ \begin{array}{l} -4718592A_2^2A_6 = \frac{125q_1 - 50q_2q_4 + 3q_4^4}{125} + \frac{339738624}{49}A_6^2 \\ A_2\left(\frac{3623878656}{7}A_6^2 + 50331648\left(\frac{125q_1 - 50q_2q_4 + 3q_4^4}{4718592 \cdot 125} + \frac{339738624}{4718592 \cdot 49}A_6^2\right)\right) = \\ = -\frac{24576}{175}A_6(25q_2 - 2q_4^3) \end{array} \right.$$

then $125q_1 - 50q_2q_4 + 3q_4^4 = 0$ but there exists solution (A_2, A_6) for all q_i . So, we obtain that system (18) has solution.

Finally, let us consider system (19). We get

$$\left\{ \begin{array}{l} 5q_3 - 2q_4^2 \neq 0 \\ q_1 = \frac{1500q_3^2 + 2450q_2q_4 - 1935q_3q_4^2 + 387q_4^4}{6125} \\ q_0 = \frac{(100q_3 - 33q_4^2)(875q_2 - 450q_3q_4 + 117q_4^3)}{153125} \\ A_6(62720A_2^2 + 92160A_6 - 5q_3 + 2q_4^2) = 0 \\ A_6(92160A_6(1120A_2 - 3q_4) - 10035200A_2^3 + 700q_2 - \\ - 188160A_2^2q_4 - 405q_3q_4 + 106q_4^3 + 3360A_2(5q_3 - 2q_4^2)) = 0 \\ A_6 \neq 0. \end{array} \right.$$

Then

$$\left\{ \begin{array}{l} 5q_3 - 2q_4^2 \neq 0 \\ q_1 = \frac{1500q_3^2 + 2450q_2q_4 - 1935q_3q_4^2 + 387q_4^4}{6125} \\ q_0 = \frac{(100q_3 - 33q_4^2)(875q_2 - 450q_3q_4 + 117q_4^3)}{153125} \\ A_6 = \frac{-62720A_2^2 + 5q_3 - 2q_4^2}{92160} \\ (62720A_2^2 - 5q_3 + 2q_4^2)(2867200A_2^3 - 160A_2(5q_3 - 2q_4^2) - 25q_2 + 15q_3q_4 - 4q_4^3) = 0 \\ A_6 \neq 0 \end{array} \right.$$

This system has solution. So, the part 2 of Theorem 3 is proved.

3). From Theorem 2 we know that operator

$$L = \left(\partial_x^2 + ax^4 + bx^2 + c + \frac{d}{x^2} \right)^2 + 48ax^2 + e, \quad a \neq 0$$

commutes with an operator M of order 14. Using formula (5) we find the spectral curve of operators L and M

$$w^2 = z^7 + \beta_6 z^6 + \beta_5 z^5 + \beta_4 z^4 + \beta_3 z^3 + \beta_2 z^2 + \beta_1 z + \beta_0,$$

where

$$\beta_6 = -7(e - 16b),$$

$$\beta_5 = 21(224b^2 + 96ac - 32be + e^2),$$

$$\beta_4 = (92416b^3 + 3168a^2(37 + 12d) + 288ac(516b - 35e) - 23520b^2e + 1680be^2 - 35e^3),$$

$$\beta_3 = 872704b^4 - 369664b^3e + 47040b^2e^2 - 2240be^3 + 35e^4 + 1152a^2(930c^2 + 8b(749 + 264d) - 11(37 + 12d)e) + 576ac(6312b^2 - 1032be + 35e^2),$$

$$\begin{aligned} \beta_2 = & 3(1204224b^5 + 115200a^3c(341 + 108d) - 872704b^4e + 184832b^3e^2 - \\ & 15680b^2e^3 + 560be^4 - 7e^5 + \\ & 192ac(62048b^3 - 18936b^2e + 1548be^2 - 35e^3) + 192a^2(8b^2(24107 + 9972d) + \\ & 48b(1410c^2 - 749e - 264de) + 3e(-1860c^2 + 407e + 132de))). \end{aligned}$$

We must solve the system of equations in a, b, c, d, e

$$\begin{cases} \beta_6 = q_6 \\ \beta_5 = q_5 \\ \beta_4 = q_4 \\ \beta_3 = q_3 \\ \beta_2 = q_2 \end{cases} \quad (20)$$

where q_6, q_5, q_4, q_3 and q_2 are arbitrary constants. Solving the first, second,

third and fourth equations of system (20) we obtain

$$\begin{cases} e = \frac{112b - q_6}{7} \\ c = \frac{4704b^2 + 7q_5 - 3q_6^2}{14112a} \\ d = \frac{68992b^3 - 5743584a^2 + 49q_4 + 308bq_5 - 35q_5q_6 - 132bq_6^2 + 10q_6^3}{1862784a^2} \\ b = \frac{7595q_5^2 - 28812q_3 + 16464q_4q_6 - 12390q_5q_6^2 + 2655q_6^4}{17259540480a^2} \\ h_2a^2 + \frac{h_{-2}}{a^2} + h_0 = q_2 \end{cases} \quad (21)$$

where

$$\begin{aligned} h_2 &= \frac{9600}{49}(7q_5 - 3q_6^2) \\ h_{-2} &= \frac{(28812q_3 - 7595q_5^2 - 16464q_4q_6 + 12390q_5q_6^2 - 2655q_6^4)^2}{323233222201344} \\ h_0 &= \frac{22638q_3q_6 - 18375q_5^2q_6 + 14665q_5q_6^3 - 2514q_6^5 + 1029q_4(25q_5 - 17q_6^2)}{52822} \end{aligned}$$

Arguing as before we see that we have not solution $a \neq 0$ of (21) if and only if one of the following system is satisfied.

$$\begin{cases} h_2 = 0 \\ h_{-2} = 0 \\ h_0 \neq q_0 \end{cases}, \quad \begin{cases} h_2 = 0 \\ h_{-2} \neq 0 \\ h_0 = q_0 \end{cases}, \quad \begin{cases} h_2 \neq 0 \\ h_{-2} = 0 \\ h_0 = q_0 \end{cases}. \quad (22)$$

Let us consider the operator

$$L = \left(\partial_x^2 + ax^6 + bx^2 + \frac{c}{x^2} + \frac{d}{x^6} \right)^2 + 192ax^4 + e, \quad a \neq 0$$

If $g = 3$, then the spectral curve of operators L and M from Theorem 1 has the form

$$w^2 = z^7 + \gamma_6z^6 + \gamma_4z^4 + \gamma_3z^3 + \gamma_2z^2 + \gamma_1z + \gamma_0,$$

where

$$\gamma_6 = 7(64b - e),$$

$$\gamma_5 = 3(384a(501 + 28c) + 7(3584b^2 - 128be + e^2)),$$

$$\begin{aligned}
\gamma_4 &= 5914624b^3 + 2433024a^2d - 376320b^2e + 6720be^2 - 35e^3 + \\
&\quad + 1152a(16b(8227 + 516c) - 5e(501 + 28c)), \\
\gamma_3 &= 223412224b^4 - 23658496b^3e + 752640b^2e^2 - 8960be^3 + 35e^4 + \\
&\quad + 110592a^2(755475 + 87560c + 2480c^2 + 5632bd - 88de) + \\
&\quad + 2304a(128b^2(41567 + 3156c) - 32be(8227 + 516c) + 5e^2(501 + 28c)), \\
\gamma_2 &= 3(1233125376b^5 + 353894400a^3d(635 + 36c) - 223412224b^4e + \\
&\quad + 11829248b^3e^2 - 250880b^2e^3 + 2240be^4 - 7e^5 + 768a(2048b^3(77977 + 7756c) - \\
&\quad - 384b^2e(41567 + 3156c) + 48be^2(8227 + 516c) - 5e^3(501 + 28c)) + \\
&\quad + 36864a^2(425472b^2d + 32b(1840425 + 311960c + 11280c^2 - 528de) - \\
&\quad + 3e(755475 + 87560c + 2480c^2 - 44de))).
\end{aligned}$$

Solving the first, second and the third equations we get

$$\left\{ \begin{array}{l} e = \frac{448b - q_6}{7} \\ c = \frac{75264b^2 - 4040064a + 7q_5 - 3q_6^2}{225792a} \\ d = \frac{908328960ab + 4415488b^3 + 49q_5 + 1232bq_5 - 35q_5q_6 - 528bq_6^2 + 10q_6^3}{119218176a^2} \\ g_3 = q_3 \\ g_2 = q_2 \end{array} \right. \quad (23)$$

where

$$\begin{aligned}
g_3 &= 306708480ab^2 + \frac{4q_4q_6}{7} + \\
&\quad + \frac{5(1519q_5^2 - 10987147100160a^2 - 2478q_5q_6^2 + 531q_6^4 - 3354624a(7q_5 - 3q_6^2))}{28812}, \\
g_2 &= 1258291200ab^3 + \frac{920125440}{7}ab^2q_6 - \frac{2457600}{49}ab(3354624a + 7q_5 - 3q_6^2) - \\
&\quad - \frac{604293090508800a^2q_6 + 173705q_5^2q_6 - 69020q_5q_6^3 + 5991q_6^5}{739508} + \\
&\quad + \frac{2058q_4(175q_5 - 31q_6^2) - 430080a(6860q_4 - 1897q_5q_6 + 113q_6^3)}{739508}
\end{aligned}$$

As in the previous part, to prove the part 3 of Theorem 3 we must show that the following systems have solution.

$$\left\{ \begin{array}{l} h_2 = 0 \\ h_{-2} = 0 \\ h_0 \neq q_2 \\ g_3 = q_3 \\ g_2 = q_2 \\ a \neq 0 \end{array} \right. , \quad (24)$$

$$\left\{ \begin{array}{l} h_2 = 0 \\ h_{-2} \neq 0 \\ h_0 = q_2 \\ g_3 = q_3 \\ g_2 = q_2 \\ a \neq 0 \end{array} \right. , \quad (25)$$

$$\left\{ \begin{array}{l} h_2 \neq 0 \\ h_{-2} = 0 \\ h_0 = q_2 \\ g_3 = q_3 \\ g_2 = q_2 \\ A_6 \neq 0 \end{array} \right. \quad (26)$$

From system (24) we obtain

$$\left\{ \begin{array}{l} q_5 = \frac{3q_6^2}{7} \\ q_3 = \frac{196q_4q_6 - 15q_6^4}{343} \\ q_2 \neq \frac{3(98q_4q_6^2 - 9q_6^5)}{2401} \\ a = \frac{637b^2}{3960} \\ -\frac{502293069824}{121}b^5 - \frac{33280(49q_4 - 5q_6^3)}{2541}b^2 + \frac{3(98q_4q_6^2 - 9q_6^5)}{2401} = q_2 \\ a \neq 0 \end{array} \right.$$

We see that this system has solution.

From system (25) we get

$$\left\{ \begin{array}{l} q_5 = \frac{3q_6^2}{7} \\ 343q_3 - 196q_4q_6 + 15q_6^4 \neq 0 \\ q_2 = \frac{3q_6(343q_3 - 98q_4q_6 + 6q_6^4)}{2401} \\ 306708480ab^2 = q_3 + \frac{93428121600}{49}a^2 - \frac{4q_4q_6}{7} + \frac{15q_6^4}{343} \\ -\frac{943718400}{343}a(20384b + 99q_6) + \frac{20480a(77271040b^3 - 245q_4 + 8072064b^2q_6 + 25q_6^3)}{3773} \\ -\frac{q_6(343q_3 - 196q_4q_6 + 15q_6^4)}{2401} = 0 \\ a \neq 0 \end{array} \right.$$

If $a = 0$ is root of fourth equation, then $343q_3 - 196q_4q_6 + 15q_6^4 = 0$ but it contradicts the second equation. But the fourth and fifth equations have solutions (a,b) for all q_i .

System (26) has the form

$$\left\{ \begin{array}{l} 7q_5 - 3q_6^2 \neq 0 \\ q_3 = \frac{7595q_5^2 + 16464q_4q_6 - 12390q_5q_6^2 + 2655q_6^4}{28812} \\ q_2 = \frac{-173705q_5^2q_6 + 69020q_5q_6^3 - 5991q_6^5 + 2058q_4(175q_5 - 31q_6^2)}{739508} \\ a = \frac{13(526848b^2 - 7q_5 + 3q_6^2)}{42577920} \\ (526848b^2 - 7q_5 + 3q_6^2)(49(97026048b^3 + 15q_4 - 32bq_5) - 525q_5q_6 + 672bq_6^2 + 150q_6^3) = 0 \\ a \neq 0. \end{array} \right.$$

This system has solution.

Theorem 3 is proved.

References

- [1] Burchnell J.L., Chaundy T.W.: Commutative ordinary differential operators. Proc. London Math. Soc. 21 (1923), 420-440; Proc. Royal Soc. London (A) 118, 557–583 (1928)
- [2] Krichever. "Integration of nonlinear equations by the methods of algebraic geometry." Functional Anal. Appl., 11:1, 12–26 (1977)
- [3] Krichever I.M.: Commutative rings of ordinary linear differential operators. Functional Anal. Appl., 12:3, 175–185 (1978)
- [4] Krichever I.M., Novikov S.P.: Holomorphic fiberings and nonlinear equations. Finite zone solutions of rank 2. Sov. Math., Dokl. 20(4), 650–654 (1979)
- [5] Mokhov O.I.: Commuting ordinary differential operators of rank 3 corresponding to an elliptic curve. Russian Math. Surveys, 37:4, 129–130 (1982)
- [6] Mokhov O.I.: "Commuting differential operators of rank 3, and nonlinear differential equations." Math. USSR, Izvestiya, 35:3, 629–655 (1990)
- [7] Mokhov O. I.: On Commutative Subalgebras of the Weyl Algebra Related to Commuting Operators of Arbitrary Rank and Genus. Mat. Zametki, 94:2, 314–316 (2013)
- [8] Mokhov O.I.: Commuting ordinary differential operators of arbitrary genus and arbitrary rank with polynomial coefficients. American Mathematical Society Translations, Volume 234, 323-336 (2014)
- [9] Mironov A.E.: Self-adjoint commuting differential operators and commutative subalgebras of the Weyl algebra. Inventiones Mathematicae 197, no. 2, 417–431 (2014)
- [10] Mironov A.E.: Periodic and rapid decay rank two self-adjoint commuting differential operators. Amer. Math. Soc. Transl. Ser. 2, V. 234, 309–322 (2014)

- [11] Mironov A.E., Zheglov A.B.: Commuting ordinary differential operators with polynomial coefficients and automorphisms of the first Weyl algebra. *International Mathematics Research Notices* Vol. 2016 2974–2993 (2016)
- [12] Dixmier J.: Sur les algebres de Weyl. *Bulletin de la Societe Mathematique de France* 96, 209–242 (1968)
- [13] Mumford D.: An algebro-geometric construction of commuting operators and of solutions to the Toda lattice equation, Korteweg deVries equation and related nonlinear equation. *Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., Kyoto, 1977)*, pp. 115–153, Kinokuniya Book Store, Tokyo, (1978)
- [14] Davletshina V.N.: Commuting differential operators of rank 2 with trigonometric coefficients. *Siberian Mathematical Journal*, 56, 405–410 (2015)
- [15] Oganessian V.S: Commuting differential operators of rank 2 and arbitrary genus g with polynomial coefficients. *Russian Math. Surveys*, 70 (1), 165–167 (2015)
- [16] Oganessian V.S.: Commuting differential operators of rank 2 with polynomial coefficients. *Functional Analysis and Its Applications*, 50:1, 54–61 (2016)
- [17] Oganessian V.S.: Explicit characterization of some commuting differential operators of rank 2. *International Mathematics Research Notices*, (2016) doi:10.1093/imrn/rnw085
- [18] Oganessian V.S.: Common eigenfunctions of commuting differential operators of rank 2. *Mathematical Notes*, 99:2, 308–311 (2016)