

On Hypohamiltonian Snarks and a Theorem of Fiorini

Jan Goedgebeur *

*Department of Applied Mathematics, Computer Science & Statistics, Ghent University,
Krijgslaan 281-S9, 9000 Ghent, Belgium*

Carol T. Zamfirescu †

*Department of Applied Mathematics, Computer Science & Statistics, Ghent University,
Krijgslaan 281-S9, 9000 Ghent, Belgium*

In memory of the first author's little girl Ella.

Received dd mmmm yyyy, accepted dd mmmmm yyyy, published online dd mmmmm yyyy

Abstract

We discuss an omission in the statement and proof of Fiorini's 1983 theorem on hypohamiltonian snarks and present a version of this theorem which is more general in several ways. Using Fiorini's erroneous result, Steffen showed that hypohamiltonian snarks exist for some $n \geq 10$ and each even $n \geq 92$. We rectify Steffen's proof by providing a correct demonstration of a technical lemma on flower snarks, which might be of separate interest. We then strengthen Steffen's theorem to the strongest possible form by determining all orders for which hypohamiltonian snarks exists. This also strengthens a result of Máčajová and Škoviera. Finally, we verify a conjecture of Steffen on hypohamiltonian snarks up to 36 vertices.

Keywords: hypohamiltonian, snark, irreducible snark, dot product

Math. Subj. Class.: 05C10, 05C38, 05C45, 05C85.

1 Introduction

A graph G is *hypohamiltonian* if G itself is non-hamiltonian, but for every vertex v in G , the graph $G - v$ is hamiltonian. A *snark* shall be a cubic cyclically 4-edge-connected graph with chromatic index 4 (i.e. four colours are required in any proper edge-colouring) and girth at least 5. We refer for notions not defined here to [21] and [6].

Motivated by similarities between the family of all snarks and the family of all cubic hypohamiltonian graphs regarding the orders for which such graphs exist, Fiorini [7] studied the hypohamiltonian properties surrounding Isaacs' so-called "flower snarks" [11] (defined rigorously below). The a priori surprising interplay between snarks and hypohamiltonian graphs has been investigated extensively—we now give an overview. Early contributions

*The first author is supported by a Postdoctoral Fellowship of the Research Foundation Flanders (FWO).

†The second author is a PhD fellow at Ghent University on the BOF (Special Research Fund) scholarship 01DI1015.

E-mail addresses: jan.goedgebeur@ugent.be (Jan Goedgebeur), czamfirescu@gmail.com (Carol T. Zamfirescu)

include Fiorini’s aforementioned paper [7], in which he showed that there exist infinitely many hypohamiltonian snarks. (In fact, according to Máčajová and Škoviera [16], it was later discovered that a family of hypohamiltonian graphs constructed by Gutt [10] includes Isaacs’ snarks, thus including Fiorini’s result.)

Skupień showed that there exist exponentially many hypohamiltonian snarks [18], and Steffen [21] proved that there exist hypohamiltonian snarks of order n for every even $n \geq 92$ (and certain $n < 92$)—we will come back to this result in Section 3. For more references and connections to other problems, see e.g. [16, 18, 3, 22]. Hypohamiltonian snarks have also been studied in connection with the famous Cycle Double Cover Conjecture [3] and Sabidussi’s Compatibility Conjecture [8].

The smallest snark, as well as the smallest hypohamiltonian graph, is the famous Petersen graph. Steffen [20] showed that every cubic hypohamiltonian graph with chromatic index 4 is *bicritical*, i.e. the graph itself is not 3-edge-colourable but deleting any two distinct vertices yields a 3-edge-colourable graph. Nedela and Škoviera [17] proved that every cubic bicritical graph is cyclically 4-edge-connected and has girth at least 5. Therefore, every cubic hypohamiltonian graph with chromatic index 4 must be a snark.

This article is organised as follows. In Section 2 we discuss the omission in Fiorini’s theorem on hypohamiltonian snarks [7] and its consequences and state a more general version of this theorem. In Section 3 we first rectify a proof of Steffen on the orders for which hypohamiltonian snarks exist which relied on Fiorini’s theorem—this erratum is based on giving a correct proof of a technical lemma concerning flower snarks, which may be of separate interest. We then prove a strengthening of Steffen’s theorem, which is best possible, as all orders for which hypohamiltonian snarks exist are determined. Our result is stronger than a theorem of Máčajová and Škoviera [15] in the sense that our result implies theirs, while the inverse implication does not hold. Finally, in Section 4 we comment upon and verify a conjecture of Steffen on hypohamiltonian snarks [22] for small hypohamiltonian snarks.

2 Fiorini’s Theorem Revisited

We call two edges independent if they have no common vertices. Let G and H be disjoint connected graphs on at least 6 vertices. Consider $G' = G - \{ab, cd\}$, where ab and cd are independent edges in G , $H' = H - \{x, y\}$, where x and y are adjacent cubic vertices in H , and let a', b' and c', d' be the other neighbours of x and y in H , respectively. Then the *dot product* $G \cdot H$ is defined as the graph

$$(V(G) \cup V(H'), E(G') \cup E(H') \cup \{aa', bb', cc', dd'\}).$$

Two remarks are in order. (1) Under the above conditions, the dot product may be disconnected. (2) In fact, there are two ways to form the dot product for a specific ab, cd, xy . One also has the possibility to add the edges $\{ac', bd', ca', db'\}$. However, in this paper we will always construct the dot product by adding the former set of edges, summarising this in the following statements as “ a, b, c, d are joined by edges to the neighbours of x and y , respectively”.

According to Skupień [19], the dot product was introduced by Adel’son-Vel’skiĭ and Titov [1], and later and independently by Isaacs [11]. Its original purpose was to obtain new snarks by combining known snarks. Isaacs was the first to explicitly construct infinitely many snarks [11]. Fiorini then proved that the dot product can also be used to combine

two hypohamiltonian snarks into a new one. Unfortunately, Fiorini's argument is incorrect. We shall discuss this omission within this section, and correct the proof of a lemma of Steffen [21] which depended on Fiorini's result in Section 3.

In a graph G , a pair (v, w) of vertices is *good* in G if there exists a hamiltonian path in G with end-vertices v and w . Two pairs of vertices $((v, w), (x, y))$ are *good* in G if there exist two disjoint paths which together span G , and which have end-vertices v and w , and x and y , respectively.

Theorem 2.1 (Fiorini, Theorem 3 in [7]). *Let G be a hypohamiltonian snark having two independent edges ab and cd for which*

- (i) *each of (a, c) , (a, d) , (b, c) , (b, d) , $((a, b), (c, d))$ is good in G ;*
- (ii) *for each vertex v , exactly one of (a, b) , (c, d) is good in $G - v$.*

If H is a hypohamiltonian snark with adjacent vertices x and y , then the dot product $G \cdot H$ is also a hypohamiltonian snark, where ab and cd are deleted from G , x and y are deleted from H , and vertices a, b, c, d are joined by edges to the neighbours of x and y , respectively.

Cavicchioli et al. [5] point out the omissions in Theorem 2.1: in order for the proof to work, the given vertex pairs need to be good in the graph *minus the edges ab, cd* . They restate the theorem and give a new proof.

Theorem 2.2 (Cavicchioli et al., Theorem 3.2 in [5]). *Let G be a hypohamiltonian snark having two independent edges ab and cd for which*

- (i) *each of (a, c) , (a, d) , (b, c) , (b, d) , $((a, b), (c, d))$ is good in $G - \{ab, cd\}$;*
- (ii) *for each vertex v , each of (a, b) , (c, d) is good in $G - \{v, ab, cd\}$.*

If H is a hypohamiltonian snark with adjacent vertices x and y , then the dot product $G \cdot H$ is also a hypohamiltonian snark, where ab and cd are deleted from G , x and y are deleted from H , and vertices a, b, c, d are joined by edges to the neighbours of x and y , respectively.

In above statements, the fact that the dot product of snarks is itself a snark had already been shown [1, 11], so indeed only the hypohamiltonicity was to be proven.

We point out that the hypotheses in Theorem 2.2 are unattainable for $v \in \{a, b, c, d\}$, which is tied to the fact that the requirements in (ii) are stronger than what is needed to prove the statement.

In [9], the following (second) restatement of Theorem 2.1 is given. Note that in [9] the graphs are required to be cubic and below we do not state this requirement—we do however need the two vertices which are removed to be cubic. This allows us to use exactly the same proof as in [9].

Theorem 2.3 (Goedgebeur and Zamfirescu). *Let G be a non-hamiltonian graph having two independent edges ab and cd for which*

- (i) *each of (a, c) , (a, d) , (b, c) , (b, d) , $((a, b), (c, d))$ is good in $G - \{ab, cd\}$;*
- (ii) *for each vertex v , at least one of (a, b) and (c, d) is good in $G - \{v, ab, cd\}$.*

If H is a hypohamiltonian graph with cubic adjacent vertices x and y , then the dot product $G \cdot H$ is also a hypohamiltonian graph, where ab and cd are deleted from G , x and y are deleted from H , and vertices a, b, c, d are joined by edges to the neighbours of x and y , respectively.

If G and H are planar, and ab and cd lie on the same facial cycle, then the dot product can be applied such that $G \cdot H$ is planar, as well. If g (h) is the girth of G (H), then the girth of $G \cdot H$ is at least $\min\{g, h\}$. If G and H are cubic, then so is $G \cdot H$.

Note that the fact that G is non-hamiltonian together with condition (ii) implies that G must be hypohamiltonian.

In the following, we will call the pair of edges ab, cd from the statement of Theorem 2.3 *suitable*. The Petersen graph is the smallest snark, and the two Blanuša snarks on 18 vertices are the second-smallest snarks. All three graphs are also hypohamiltonian. Due the huge automorphism group of the Petersen graph, it can be verified by hand that it does not contain a pair of suitable edges. Although both Blanuša snarks are dot products of two Petersen graphs, the Petersen graph does not contain a pair of suitable edges. Thus, in a certain sense, Theorem 2.3 is *not* “if and only if”, i.e. there exist dot products whose factors do not contain suitable edges.

Let us end this section with a remark which may prove to be useful in other applications. Throughout its statement and proof, we use the notation from Theorem 2.3.

Observation 2.4. We have that $G \cdot H + ab$, $G \cdot H + cd$, and $G \cdot H + ab + cd$ are hypohamiltonian, as well.

Proof. Put $N(x) = \{a', b', y\}$ and $N(y) = \{c', d', x\}$ such that the unique neighbour of a' (b', c', d') in G is a (b, c, d). Assume $G \cdot H + ab + cd$ contains a hamiltonian cycle \mathfrak{h} . Thus, at least one of ab and cd lies in \mathfrak{h} , say ab . We treat $H - \{x, y\}$ as a subgraph of $G \cdot H$. If $aa', bb' \in E(\mathfrak{h})$, then $\mathfrak{h} \cap H \cup a'xb' \cup c'yd'$ gives a hamiltonian cycle in H , a contradiction. If $aa', bb' \notin E(\mathfrak{h})$, then the cycle $\mathfrak{h} \cap G + cd$ yields a contradiction. So w.l.o.g. $aa' \in E(\mathfrak{h})$ and $bb' \notin E(\mathfrak{h})$. This implies the existence of a hamiltonian path in $H - \{x, y\}$ with end-vertices a' and $u \in \{c', d'\}$. But this path together with $uyxa'$ is a hamiltonian cycle in H , a contradiction. It follows that $G \cdot H + ab$ and $G \cdot H + cd$ are non-hamiltonian, as well. \square

3 On a Theorem of Steffen on Hypohamiltonian Snarks

3.1 Rectifying Steffen’s proof

A snark is *irreducible* if the removal of every edge-cut which is not the set of all edges incident with a vertex yields a 3-edge-colourable graph. Steffen’s article [21] is motivated by the following problem.

Problem 3.1 (Nedela and Škoviera [17]). For which even number $n \geq 10$ does there exist an irreducible snark of order n ? In particular, does there exist an irreducible snark of each sufficiently large order?

Steffen settled the second question of Problem 3.1 by giving the following main result from [21].

Theorem 3.2 (Steffen, Theorem 2.5 in [21]). *There is a hypohamiltonian snark of order n*

- (1) for each $n \in \{m : m \geq 64 \text{ and } m \equiv 0 \pmod{8}\}$
- (2) for each $n \in \{10, 18\} \cup \{m : m \geq 98 \text{ and } m \equiv 2 \pmod{8}\}$
- (3) for each $n \in \{m : m \geq 20 \text{ and } m \equiv 4 \pmod{8}\}$
- (4) for each $n \in \{30\} \cup \{m : m \geq 54 \text{ and } m \equiv 6 \pmod{8}\}$
- (5) for each even $n \geq 92$.

Isaacs' *flower snark* J_{2k+1} , see [11], is the graph

$$\left(\{a_i, b_i, c_i, d_i\}_{i=0}^{2k}, \{b_i a_i, b_i c_i, b_i d_i, a_i a_{i+1}, c_i d_{i+1}, d_i c_{i+1}\}_{i=0}^{2k} \right),$$

where addition in the indices is performed modulo $2k + 1$.

However, the proof of [21, Lemma 2.3], which is essential for the proof of the theorem, is erroneous, since it uses Theorem 2.1 (and it does not, by sheer coincidence, work with Theorem 2.3). We here give a correct proof of that lemma.




Figure 1: The flower snark J_9 . The suitable edges b_0c_0 and b_4c_4 are marked in bold red.

Lemma 3.3 (Steffen, Lemma 2.3 in [21]). *The flower snarks J_9 , J_{11} , and J_{13} satisfy the conditions of Theorem 2.3.*

Proof. In [21], in each of the graphs J_9 , J_{11} , and J_{13} , the suitable edges were chosen to be b_0c_0 and b_4c_4 . However, in [21], for various vertices v , the hamiltonian paths did not satisfy condition (ii) from Theorem 2.3, as the paths used one of the edges b_0c_0 or b_4c_4 . This was for instance the case for $v \in \{a_0, a_8, d_0\}$ in J_9 , for $v \in \{a_0, a_{10}, d_0\}$ in J_{11} and for $v \in \{a_0, c_1, c_{12}, d_0\}$ in J_{13} , see Claims 6, 7, and 8 in the Appendix of [21].

We will now prove that b_0c_0 and b_4c_4 are indeed suitable edges for Theorem 2.3 for J_9 , J_{11} , and J_{13} . For J_9 the proof is given below (and partially in the Appendix), while the technical details of the proofs for J_{11} and J_{13} can be found in the Appendix. The mapping between the a_i, b_i, c_i, d_i (used by Steffen) and the vertex numbers used in the proof is shown in Figures 1–3. We use numbers as labels in the proof to make it easier to read these graphs using a computer for verifying the results.

Proof that b_0c_0 and b_4c_4 are suitable edges for J_9 .

Figure 1 shows the flower snark J_9 . In J_9 , b_0c_0 and b_4c_4 correspond to the edges $(0, 26)$ and $(11, 12)$, respectively.

$(0, 11), (0, 12), (26, 11)$ and $(26, 12)$ are good in $J_9 - \{(0, 26), (11, 12)\}$ due to the following hamiltonian paths, respectively:

- 11, 10, 5, 6, 7, 8, 9, 30, 29, 28, 27, 35, 24, 23, 22, 17, 16, 15, 32, 31, 12, 13, 14, 19, 18, 33, 34, 21, 20, 25, 26, 4, 3, 2, 1, 0
- 12, 13, 14, 15, 16, 11, 10, 5, 4, 26, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 30, 9, 8, 7, 6, 29, 28, 3, 2, 1, 23, 24, 35, 27, 0
- 11, 10, 5, 4, 3, 2, 1, 0, 27, 28, 29, 6, 7, 8, 9, 30, 31, 12, 13, 14, 19, 18, 17, 16, 15, 32, 33, 34, 35, 24, 23, 22, 21, 20, 25, 26
- 12, 13, 8, 7, 2, 3, 4, 5, 6, 29, 28, 27, 0, 1, 23, 22, 17, 18, 33, 32, 31, 30, 9, 10, 11, 16, 15, 14, 19, 20, 21, 34, 35, 24, 25, 26

$((0, 26), (11, 12))$ is good in $J_9 - \{(0, 26), (11, 12)\}$ due to the following two disjoint paths with end-vertices 0 and 26, and 11 and 12, respectively, which together span J_9 .

- 26, 25, 20, 19, 14, 13, 8, 7, 2, 1, 0
- 12, 31, 32, 15, 16, 17, 18, 33, 34, 21, 22, 23, 24, 35, 27, 28, 3, 4, 5, 6, 29, 30, 9, 10, 11

We showed by computer that at least one of $(0, 26)$ or $(11, 12)$ is good in $J_9 - \{v, (0, 26), (11, 12)\}$ for every $v \in V(J_9)$. In each case we verified that the path found by the computer is indeed a valid hamiltonian path in the graph. Below we explicitly show this for $v = 0$. The hamiltonian paths for the other values of v can be found in the Appendix.

$v = 0$: 12, 13, 14, 15, 32, 31, 30, 29, 6, 5, 10, 9, 8, 7, 2, 1, 23, 24, 25, 26, 4, 3, 28, 27, 35, 34, 33, 18, 19, 20, 21, 22, 17, 16, 11 □

Since Steffen's statement of Lemma 3.3 remains intact, the proof and statement of his main result, reproduced above as Theorem 3.2, are correct as given in [21]. Even though we prove a stronger version of Steffen's theorem in the next section, we think it is important to fix the proof of Lemma 3.3 as there may be others who rely on this lemma, or might want to rely on it in the future.

3.2 Orders of hypohamiltonian snarks

We shall now prove a strengthening of Steffen's theorem, which in a sense is strongest possible since we will determine all orders for which hypohamiltonian snarks exist. We emphasise that our proof's mechanism contains significantly fewer "moving parts" than Máčajová and Škoviera's [15], and, as mentioned in the introduction, our theorem also strengthens their result. We do need the following two easily verifiable lemmas.

Lemma 3.4. *The second Blanuša snark B_2 shown in Figure 4 has a pair of suitable edges.*

Proof. Figure 4 shows the second Blanuša snark B_2 . By computer we determined that B_2 has exactly three pairs of suitable edges: $((6, 8), (10, 16)), ((3, 9), (12, 17))$ and $((4, 7), (13, 15))$. We will now prove by hand that $((6, 8), (10, 16))$ is a suitable edge pair.

$(6, 10), (6, 16), (8, 10)$ and $(8, 16)$ are good in $B_2 - \{(6, 8), (10, 16)\}$ due to the following hamiltonian paths, respectively:

- 10, 11, 12, 17, 16, 15, 13, 14, 0, 1, 5, 4, 7, 8, 9, 3, 2, 6
- 16, 15, 9, 8, 7, 17, 12, 13, 14, 10, 11, 1, 0, 2, 3, 4, 5, 6




Figure 2: The flower snark J_{11} . The suitable edges b_0c_0 and b_4c_4 are marked in bold red.

- 10, 11, 1, 0, 14, 13, 12, 17, 16, 15, 9, 3, 2, 6, 5, 4, 7, 8
- 16, 15, 9, 3, 4, 5, 6, 2, 0, 1, 11, 10, 14, 13, 12, 17, 7, 8

$((6, 8), (10, 16))$ is good in $B_2 - \{(6, 8), (10, 16)\}$ due to the following two disjoint paths with end-vertices 6 and 8, and 10 and 16, respectively, which together span B_2 .

- 8, 7, 4, 5, 6
- 16, 17, 12, 11, 1, 0, 2, 3, 9, 15, 13, 14, 10

We now prove that at least one of $(6, 8)$ or $(10, 16)$ is good in $B_2 - \{v, (6, 8), (10, 16)\}$ for every $v \in V(B_2)$. By symmetry we only need to prove this for $v = 0, 2, 4, 6, 7, 8$.

- $v = 0$: 8, 7, 4, 5, 1, 11, 10, 14, 13, 12, 17, 16, 15, 9, 3, 2, 6
- $v = 2$: 8, 9, 3, 4, 7, 17, 16, 15, 13, 12, 11, 10, 14, 0, 1, 5, 6
- $v = 4$: 16, 15, 13, 14, 0, 1, 5, 6, 2, 3, 9, 8, 7, 17, 12, 11, 10
- $v = 6$: 16, 15, 9, 8, 7, 17, 12, 13, 14, 0, 2, 3, 4, 5, 1, 11, 10
- $v = 7$: 8, 9, 15, 16, 17, 12, 13, 14, 10, 11, 1, 0, 2, 3, 4, 5, 6
- $v = 8$: 16, 15, 9, 3, 2, 6, 5, 4, 7, 17, 12, 13, 14, 0, 1, 11, 10

□




Figure 3: The flower snark J_{13} . The suitable edges b_0c_0 and b_4c_4 are marked in bold red.




Figure 4: The second Blanuša snark. It has 18 vertices. The suitable edges $(6, 8)$ and $(10, 16)$ are marked in bold red.

Lemma 3.5. *The first Loupekine snark L_1 shown in Figure 5 has a pair of suitable edges.*

Proof. Figure 5 shows the first Loupekine snark L_1 . By computer we determined that L_1 has exactly six pairs of suitable edges: $((0, 1), (17, 20)), ((0, 2), (8, 17)), ((1, 5), (14, 20)), ((2, 3), (8, 10)), ((3, 4), (10, 12))$ and $((4, 5), (12, 14))$. We will now prove by hand that $((3, 4), (10, 12))$ is a suitable edge pair.

$(3, 10), (3, 12), (4, 10)$ and $(4, 12)$ are good in $L_1 - \{(3, 4), (10, 12)\}$ due to the following hamiltonian paths, respectively:

- 10, 7, 9, 6, 8, 17, 19, 21, 16, 13, 11, 0, 1, 5, 4, 18, 20, 14, 12, 15, 2, 3
- 12, 14, 20, 18, 4, 5, 7, 10, 8, 17, 19, 21, 16, 9, 6, 1, 0, 11, 13, 15, 2, 3
- 10, 7, 5, 1, 0, 11, 13, 16, 9, 6, 8, 17, 20, 14, 12, 15, 2, 3, 19, 21, 18, 4
- 12, 14, 11, 0, 1, 6, 9, 16, 13, 15, 2, 3, 19, 21, 18, 20, 17, 8, 10, 7, 5, 4

$((3, 4), (10, 12))$ is good in $L_1 - \{(3, 4), (10, 12)\}$ due to the following two disjoint paths with end-vertices 3 and 4, and 10 and 12, respectively, which together span L_1 .

- 4, 5, 1, 6, 8, 17, 20, 18, 21, 19, 3
- 12, 14, 11, 0, 2, 15, 13, 16, 9, 7, 10

We now prove that at least one of $(3, 4)$ or $(10, 12)$ is good in $L_1 - \{v, (3, 4), (10, 12)\}$ for every $v \in V(L_1)$. By symmetry we only need to prove this for $v = 1, 4, 5, 6, 7, 8, 9, 10, 16, 17, 18, 21$.

- $v = 1$: 12, 15, 13, 16, 9, 6, 8, 17, 20, 14, 11, 0, 2, 3, 19, 21, 18, 4, 5, 7, 10
- $v = 4$: 12, 14, 20, 18, 21, 16, 9, 6, 8, 17, 19, 3, 2, 15, 13, 11, 0, 1, 5, 7, 10
- $v = 5$: 4, 18, 20, 17, 8, 10, 7, 9, 6, 1, 0, 2, 15, 12, 14, 11, 13, 16, 21, 19, 3
- $v = 6$: 4, 5, 1, 0, 2, 15, 12, 14, 11, 13, 16, 9, 7, 10, 8, 17, 20, 18, 21, 19, 3
- $v = 7$: 12, 14, 20, 17, 19, 3, 2, 15, 13, 11, 0, 1, 5, 4, 18, 21, 16, 9, 6, 8, 10
- $v = 8$: 12, 14, 20, 17, 19, 3, 2, 15, 13, 11, 0, 1, 6, 9, 16, 21, 18, 4, 5, 7, 10
- $v = 9$: 4, 5, 7, 10, 8, 6, 1, 0, 2, 15, 12, 14, 11, 13, 16, 21, 18, 20, 17, 19, 3
- $v = 10$: 4, 5, 7, 9, 16, 13, 11, 0, 1, 6, 8, 17, 19, 21, 18, 20, 14, 12, 15, 2, 3
- $v = 16$: 4, 5, 1, 6, 9, 7, 10, 8, 17, 19, 21, 18, 20, 14, 12, 15, 13, 11, 0, 2, 3
- $v = 17$: 4, 5, 1, 6, 8, 10, 7, 9, 16, 13, 11, 0, 2, 15, 12, 14, 20, 18, 21, 19, 3
- $v = 18$: 4, 5, 1, 6, 8, 10, 7, 9, 16, 21, 19, 17, 20, 14, 12, 15, 13, 11, 0, 2, 3
- $v = 21$: 4, 18, 20, 14, 12, 15, 2, 0, 11, 13, 16, 9, 6, 1, 5, 7, 10, 8, 17, 19, 3

□




Figure 5: The first Loupekine snark L_1 . It has 22 vertices. The suitable edges $(3, 4)$ and $(10, 12)$ are marked in bold red.

The following is the strongest form of Steffen's Theorem 3.2.

Theorem 3.6. *There exists a hypohamiltonian snark of order n iff $n \in \{10, 18, 20, 22\}$ or n is even and $n \geq 26$.*

Proof. For $n = 10$, it is well-known that the Petersen graph is hypohamiltonian and it is also well-known that no snarks exist of order 12, 14 or 16.

In Lemma 3.4 we showed that the second Blanuša snark B_2 (which has order 18) contains a pair of suitable edges. In [3] it was proven that hypohamiltonian snarks exist for all even orders from 18 to 36 with the exception of 24 (see Table 1). Let S_n denote a hypohamiltonian snark of order n . Using Theorem 2.3, we form the dot product $B_2 \cdot S_n$ for $n \in \{18, 20, 22, 26, 28, 30, 32\}$ and obtain hypohamiltonian snarks of all even orders between 34 and 48 with the exception of 40 (recall that the dot product of two snarks is a snark).

By Lemma 3.5 we know that the first Loupekine snark L_1 (which has order 22) contains a pair of suitable edges. Applying Theorem 2.3 to this snark and a 22-vertex hypohamiltonian snark, we obtain a hypohamiltonian snark of order 40.

We form the dot product $B_2 \cdot S_n$ for all even $n \in \{34, \dots, 48\}$ and obtain hypohamiltonian snarks of all even orders from 50 to 64. This may now be iterated ad infinitum, and the proof is complete. \square

In [15] Máčajová and Škoviera proved the following theorem (which fully settles Problem 3.1).

Theorem 3.7 (Máčajová and Škoviera, Theorems A and B in [15]). *There exists an irreducible snark of order n iff $n \in \{10, 18, 20, 22\}$ or n is even and $n \geq 26$.*

Nedela and Škoviera [17] proved that a snark is irreducible if and only if it is bicritical, and Steffen [20] showed that every hypohamiltonian snark is bicritical—while the converse is not true, as will be shown in Table 1.

In [4] Carneiro, da Silva, and McKay determined all 4-vertex-critical snarks up to 36 vertices and Škoviera [23] showed that a snark is 4-vertex-critical if and only if it is irreducible. (A graph G without k -flow is k -vertex-critical if, for every pair of vertices (u, v) of G , identifying u and v yields a graph that has a k -flow (see [4] for more details)).

The number of hypohamiltonian snarks on $n \leq 36$ vertices was determined in [3] and can be found in Table 1 together with the number of irreducible snarks from [4]. These graphs can also be downloaded from the *House of Graphs* [2] at <http://hog.grinvin.org/Snarks>.

As can be seen from Table 1, there are a significant number of irreducible snarks which are not hypohamiltonian. The smallest such snarks have order 26. So Theorem 3.6 implies Theorem 3.7, while the inverse implication is not true.

Order	irreducible	hypo.	hypo. and $\lambda_c = 4$	hypo. and $\lambda_c \geq 5$
10	1	1	0	1
18	2	2	2	0
20	1	1	0	1
22	2	2	0	2
24	0	0	0	0
26	111	95	87	8
28	33	31	30	1
30	115	104	93	11
32	13	13	0	13
34	40 328	31 198	29 701	1 497
36	13 720	10 838	10 374	464
38	?	?	$\geq 51\,431$?
40	?	?	$\geq 8\,820$?
42	?	?	$\geq 20\,575\,458$?
44	?	?	$\geq 8\,242\,146$?

Table 1: Number of irreducible and hypohamiltonian snarks (see [4, Table 1] and [3, Table 2]). λ_c stands for cyclic edge-connectivity. The counts of cases indicated with a ' \geq ' are possibly incomplete; all other cases are complete.

The hypohamiltonian snarks on $n \geq 34$ vertices constructed by the dot product in the proof of Theorem 3.6 clearly all have cyclic edge-connectivity 4. By combining this with Table 1 we obtain:

Corollary 3.8. *There exists a hypohamiltonian snark of order n and cyclic edge-connectivity 4 iff $n \in \{18, 26, 28, 30\}$ or n is even and $n \geq 34$.*

As already mentioned, every hypohamiltonian snark is irreducible, thus Corollary 3.8 implies [15, Theorem A]. For higher cyclic edge-connectivity, the following is known.

Theorem 3.9 (Máčajová and Škoviera [16]). *There exists a hypohamiltonian snark of order n and cyclic connectivity 5 for each even $n \geq 140$, and there exists a hypohamiltonian snark of order n and cyclic connectivity 6 for each even $n \geq 166$.*

If we relax the requirements from hypohamiltonicity to irreducibility, more is known:

Theorem 3.10 (Máčajová and Škoviera [15]). *There exists an irreducible snark of order n and cyclic connectivity 5 iff $n \in \{10, 20, 22, 26\}$ or n is even and $n \geq 30$, and there exists an irreducible snark of order n and cyclic connectivity 6 for each $n \equiv 4 \pmod{8}$ with $n \geq 28$, and for each even $n \geq 210$.*

Note that as every hypohamiltonian graph is irreducible, Theorem 3.9 also implies that $n \geq 210$ can be improved to $n \geq 166$ in Theorem 3.10.

The smallest hypohamiltonian snark of cyclic edge-connectivity 5 has order 10 and is the Petersen graph, and the second-smallest such graph has order 20. The flower snark J_7 of order 28 is the smallest cyclically 6-edge-connected hypohamiltonian snark. We conclude this section with the following two problems motivated by Theorem 3.10 and results of Kochol [12, 14].

Problem 3.11 (Máčajová and Škoviera [15]). Construct a cyclically 6-edge-connected snark (irreducible or not) of order smaller than 118 and different from any of Isaacs' snarks.

Problem 3.12. Determine all orders for which cyclically 6-edge-connected snarks exist.

4 On a Conjecture of Steffen on Hypohamiltonian Snarks

Consider a cubic graph G . We denote with $\mu_k(G)$ the minimum number of edges not contained in the union of k 1-factors of G , for every possible combination of k 1-factors. If $\mu_3(G) = 0$, then G is 3-edge-colourable. In [22], Steffen made the following conjecture on hypohamiltonian snarks.

Conjecture 4.1 (Steffen, Conjecture 4.1 in [22]). *If G is a hypohamiltonian snark, then $\mu_3(G) = 3$.*

If true, this conjecture would have significant consequences, e.g. by Theorem 2.14 from [22], it would imply that every hypohamiltonian snark has a Berge-cover (a bridgeless cubic graph G has a Berge-cover if $\mu_5(G) = 0$).

We wrote a computer program for computing $\mu_3(G)$ and tested Conjecture 4.1 on the complete lists of hypohamiltonian snarks up to 36 vertices. This leads to the following observation.

Observation 4.2. There are no counterexamples to Conjecture 4.1 among the hypohamiltonian snarks with at most 36 vertices.

The authors of [3] noted a huge increase (from 13 to 31 198) in the number of hypohamiltonian snarks from order 32 to 34, see Table 1. Using a computer, we were able to determine that 29 365 out of the 29 701 hypohamiltonian snarks with cyclic edge-connectivity 4 on 34 vertices can be obtained by performing a dot product on a hypohamiltonian snark on 26 vertices and the Petersen graph. We also determined that the remaining hypohamiltonian snarks with cyclic edge-connectivity 4 on 34 vertices are obtained by performing a dot product on the Blanuša snarks. Intriguingly, our computations show that some hypohamiltonian snarks can be obtained by performing a dot product on a hypohamiltonian snark on 26 vertices and the Petersen graph, as well as by performing a dot product on the Blanuša snarks.

There is also a (slightly less dramatic) increase in the cyclically 5-edge-connected case—these are obviously not dot products—and we believe it to be due to more general graph products, for instance “superposition” introduced by Kochol [13]. It would be interesting to further explore these transformations in order to fully understand these sudden increases and decreases in numbers.

Using a computer, we determined that all hypohamiltonian snarks with cyclic edge-connectivity 4 up to 36 vertices can be obtained by performing a dot product on two hypohamiltonian snarks. This leads us to pose the following question.

Problem 4.3. Is every hypohamiltonian snark with cyclic edge-connectivity 4 a dot product of two hypohamiltonian snarks?

Recall that in Theorem 2.3 the graphs G and H are hypohamiltonian, but the theorem is *not* “if and only if”, since although the Petersen graph does not contain a pair of suitable edges, the Blanuša snarks (which are dot products of two Petersen graphs) are hypohamiltonian. We believe the answer to this problem to be “no” due to the following observation. In order to cover all cases, we would need to add to condition (ii) of Theorem 2.3 the possibility of $((a, b), (c, d))$ being good in $G - \{v, ab, cd\}$. However, we would then need to require from H that it contains a 2-factor containing exactly two (necessarily odd) cycles. Although we were unable to find a counter-example, we believe that there exist hypohamiltonian snarks which do not possess such a 2-factor.

We also determined all hypohamiltonian snarks up to 44 vertices which can be obtained by performing a dot product on two hypohamiltonian snarks. The counts of these snarks can be found in the fourth column of Table 1. We also verified Conjecture 4.1 on these snarks.

Observation 4.4. There are no counterexamples to Conjecture 4.1 among the hypohamiltonian snarks with at most 44 vertices which are a dot product of two hypohamiltonian snarks.

Acknowledgements

We thank Nico Van Cleemput for providing us with a script which greatly enhanced the quality of our figures. We would also like to thank Martin Škoviera for informing us about the equivalence of irreducible and vertex-critical graphs. Finally, we also wish to thank Eckhard Steffen for useful suggestions.

References

- [1] G. M. Adel'son-Vel'skiĭ and V. K. Titov, On edge 4-chromatic cubic graphs (in Russian), in: *Proc. Seminar of 1971 at Moscow Univ., Voprosy Kibernetiki [vol. not numbered]*, 1973 pp. 5–14.
- [2] G. Brinkmann, K. Coolsaet, J. Goedgebeur and H. Mélot, House of Graphs: a database of interesting graphs, *Discrete Applied Mathematics* **161** (2013), 311–314, available at <http://hog.grinvin.org/>.
- [3] G. Brinkmann, J. Goedgebeur, J. Häggglund and K. Markström, Generation and properties of snarks, *Journal of Combinatorial Theory, Series B* **103** (2013), 468–488.
- [4] A. Carneiro, C. da Silva and B. McKay, A faster test for 4-flow-criticality in snarks, *LAGOS'15 - VIII Latin-American Algorithms, Graphs and Optimization Symposium, Electronic Notes in Discrete Mathematics* **50** (2015), 193–198.

- [5] A. Caviccholi, T. E. Murgolo, B. Ruini and F. Spaggiari, Special classes of snarks, *Acta Applicandae Mathematica* **76** (2003), 57–88.
- [6] R. Diestel, Graph Theory, *Graduate Texts in Mathematics* **173** (2010).
- [7] S. Fiorini, Hypohamiltonian snarks, in: *Graphs and Other Combinatorial Topics (ed.: M. Fiedler)*, Teubner, volume 59, 1983 pp. 70–75.
- [8] H. Fleischner and R. Häggkvist, Circuit double covers in special types of cubic graphs, *Discrete Mathematics* **309** (2009), 5724–5728.
- [9] J. Goedgebeur and C. T. Zamfirescu, New results on hypohamiltonian and almost hypohamiltonian graphs, *arXiv preprint arXiv:1606.06577* (2016).
- [10] S. Gutt, Infinite families of hypohamiltonian graphs, *Académie Royale de Belgique. Bulletin de la Classe des Sciences. 5e Série* **63** (1977), 432–440.
- [11] R. Isaacs, Infinite families of nontrivial trivalent graphs which are not Tait colorable, *American Mathematical Monthly* **82** (1975), 221–239.
- [12] M. Kochol, A cyclically 6-edge-connected snark of order 118, *Discrete Mathematics* **161** (1996), 297–300.
- [13] M. Kochol, Snarks without small cycles, *Journal of Combinatorial Theory, Series B* **67** (1996), 34–47.
- [14] M. Kochol, Reduction of the 5-flow conjecture to cyclically 6-edge-connected snarks, *Journal of Combinatorial Theory, Series B* **90** (2004), 139–145.
- [15] E. Máčajová and M. Škoviera, Irreducible snarks of given order and cyclic connectivity, *Discrete Mathematics* **306** (2006), 779–791.
- [16] E. Máčajová and M. Škoviera, Constructing hypohamiltonian snarks with cyclic connectivity 5 and 6, *Electronic Journal of Combinatorics* **14** (2007), #R18.
- [17] R. Nedela and M. Škoviera, Decompositions and reductions of snarks, *Journal of Graph Theory* **22** (1996), 253–279.
- [18] Z. Skupień, Exponentially many hypohamiltonian snarks, in: *6th Czech-Slovak International Symposium on Combinatorics, Graph Theory, Algorithms and Applications, Electronic Notes in Discrete Mathematics*, volume 28, 2007 pp. 417–424.
- [19] Z. Skupień, Multi-compositions in exponential counting of hypohamiltonian snarks, in: *Convexity and Discrete Geometry Including Graph Theory, Springer Proc. in Math. & Stat. (eds.: K. Adiprasito, I. Bárány, C. Vîlcu)*, Springer, volume 148, 2016 pp. 47–58.
- [20] E. Steffen, Classification and characterizations of snarks, *Discrete Math.* **188** (1998), 183–203.
- [21] E. Steffen, On bicritical snarks, *Mathematica Slovaca* **51** (2001), 141–150.
- [22] E. Steffen, 1-factor and cycle covers of cubic graphs, *Journal of Graph Theory* **78** (2015), 195–206.
- [23] M. Škoviera, Flow-critical and critical snarks coincide, *In preparation* (2016).

Appendix

Below we give the technical details which were left out in the proof of Lemma 3.3.

Proof that b_0c_0 and b_4c_4 are suitable edges for J_9 (continued)

We will now prove that at least one of $(0, 26)$ or $(11, 12)$ is good in $J_9 - \{v, (0, 26), (11, 12)\}$ for every $v \in V(J_9)$ except for $v = 0$, which we have already shown above in the proof of Lemma 3.3.

- $v = 1$: 26, 4, 5, 6, 7, 2, 3, 28, 29, 30, 31, 12, 13, 8, 9, 10, 11, 16, 17, 18, 33, 32, 15, 14, 19, 20, 25, 24, 23, 22, 21, 34, 35, 27, 0
- $v = 2$: 12, 31, 32, 15, 14, 13, 8, 7, 6, 5, 10, 9, 30, 29, 28, 3, 4, 26, 25, 24, 23, 1, 0, 27, 35, 34, 33, 18, 19, 20, 21, 22, 17, 16, 11
- $v = 3$: 26, 4, 5, 6, 7, 2, 1, 23, 22, 21, 20, 25, 24, 35, 34, 33, 32, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 31, 30, 29, 28, 27, 0
- $v = 4$: 26, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 30, 29, 28, 3, 2, 1, 23, 24, 35, 27, 0
- $v = 5$: 26, 4, 3, 2, 1, 23, 22, 21, 20, 25, 24, 35, 34, 33, 32, 15, 14, 19, 18, 17, 16, 11, 10, 9, 30, 31, 12, 13, 8, 7, 6, 29, 28, 27, 0
- $v = 6$: 26, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 5, 4, 3, 2, 7, 8, 9, 30, 29, 28, 27, 35, 24, 23, 1, 0
- $v = 7$: 26, 4, 5, 6, 29, 30, 31, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 32, 33, 34, 35, 24, 25, 20, 21, 22, 23, 1, 2, 3, 28, 27, 0
- $v = 8$: 26, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 9, 30, 29, 28, 3, 4, 5, 6, 7, 2, 1, 23, 24, 35, 27, 0
- $v = 9$: 26, 4, 3, 2, 1, 23, 22, 21, 20, 25, 24, 35, 34, 33, 32, 15, 14, 19, 18, 17, 16, 11, 10, 5, 6, 7, 8, 13, 12, 31, 30, 29, 28, 27, 0
- $v = 10$: 12, 13, 14, 15, 32, 31, 30, 9, 8, 7, 2, 3, 28, 29, 6, 5, 4, 26, 25, 24, 23, 1, 0, 27, 35, 34, 33, 18, 19, 20, 21, 22, 17, 16, 11
- $v = 11$: 26, 4, 3, 2, 1, 23, 22, 21, 34, 35, 24, 25, 20, 19, 14, 15, 16, 17, 18, 33, 32, 31, 12, 13, 8, 7, 6, 5, 10, 9, 30, 29, 28, 27, 0
- $v = 12$: 26, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 30, 9, 8, 13, 14, 15, 16, 11, 10, 5, 4, 3, 2, 7, 6, 29, 28, 27, 35, 24, 23, 1, 0
- $v = 13$: 12, 31, 30, 29, 6, 5, 10, 9, 8, 7, 2, 1, 0, 27, 28, 3, 4, 26, 25, 20, 21, 22, 23, 24, 35, 34, 33, 32, 15, 14, 19, 18, 17, 16, 11
- $v = 14$: 26, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 15, 16, 11, 10, 9, 8, 13, 12, 31, 30, 29, 28, 3, 4, 5, 6, 7, 2, 1, 23, 24, 35, 27, 0
- $v = 15$: 26, 4, 3, 2, 1, 23, 22, 21, 20, 25, 24, 35, 34, 33, 32, 31, 12, 13, 14, 19, 18, 17, 16, 11, 10, 5, 6, 7, 8, 9, 30, 29, 28, 27, 0
- $v = 16$: 12, 13, 8, 7, 2, 3, 28, 29, 6, 5, 4, 26, 25, 24, 23, 1, 0, 27, 35, 34, 33, 18, 17, 22, 21, 20, 19, 14, 15, 32, 31, 30, 9, 10, 11
- $v = 17$: 26, 4, 3, 2, 1, 23, 22, 21, 20, 25, 24, 35, 34, 33, 18, 19, 14, 13, 12, 31, 32, 15, 16, 11, 10, 5, 6, 7, 8, 9, 30, 29, 28, 27, 0
- $v = 18$: 26, 25, 20, 19, 14, 15, 32, 33, 34, 21, 22, 17, 16, 11, 10, 9, 8, 13, 12, 31, 30, 29, 28, 3, 4, 5, 6, 7, 2, 1, 23, 24, 35, 27, 0
- $v = 19$: 26, 4, 3, 28, 27, 35, 34, 21, 20, 25, 24, 23, 22, 17, 18, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 5, 6, 29, 30, 9, 8, 7, 2, 1, 0
- $v = 20$: 26, 25, 24, 35, 27, 28, 29, 6, 5, 4, 3, 2, 7, 8, 13, 12, 31, 30, 9, 10, 11, 16, 17, 18, 19, 14, 15, 32, 33, 34, 21, 22, 23, 1, 0
- $v = 21$: 26, 4, 3, 2, 1, 23, 22, 17, 18, 19, 20, 25, 24, 35, 34, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 30, 29, 28, 27, 0

- $v = 22$: 26, 25, 20, 21, 34, 33, 32, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 31, 30, 29, 28, 3, 4, 5, 6, 7, 2, 1, 23, 24, 35, 27, 0
- $v = 23$: 26, 4, 3, 28, 27, 35, 24, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 5, 6, 29, 30, 9, 8, 7, 2, 1, 0
- $v = 24$: 26, 25, 20, 19, 14, 15, 32, 33, 18, 17, 16, 11, 10, 9, 8, 13, 12, 31, 30, 29, 28, 3, 4, 5, 6, 7, 2, 1, 23, 22, 21, 34, 35, 27, 0
- $v = 25$: 26, 4, 3, 2, 1, 23, 24, 35, 34, 33, 18, 17, 22, 21, 20, 19, 14, 13, 12, 31, 32, 15, 16, 11, 10, 5, 6, 7, 8, 9, 30, 29, 28, 27, 0
- $v = 26$: 12, 13, 8, 7, 2, 3, 4, 5, 6, 29, 28, 27, 0, 1, 23, 22, 21, 34, 35, 24, 25, 20, 19, 14, 15, 16, 17, 18, 33, 32, 31, 30, 9, 10, 11
- $v = 27$: 26, 4, 5, 6, 7, 2, 3, 28, 29, 30, 31, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 32, 33, 34, 35, 24, 25, 20, 21, 22, 23, 1, 0
- $v = 28$: 12, 13, 14, 15, 32, 31, 30, 29, 6, 5, 10, 9, 8, 7, 2, 3, 4, 26, 25, 24, 23, 1, 0, 27, 35, 34, 33, 18, 19, 20, 21, 22, 17, 16, 11
- $v = 29$: 26, 4, 5, 6, 7, 8, 13, 12, 31, 30, 9, 10, 11, 16, 17, 18, 19, 14, 15, 32, 33, 34, 35, 24, 25, 20, 21, 22, 23, 1, 2, 3, 28, 27, 0
- $v = 30$: 26, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 9, 8, 7, 2, 3, 4, 5, 6, 29, 28, 27, 35, 24, 23, 1, 0
- $v = 31$: 12, 13, 8, 7, 6, 5, 10, 9, 30, 29, 28, 27, 0, 1, 2, 3, 4, 26, 25, 20, 21, 22, 23, 24, 35, 34, 33, 32, 15, 14, 19, 18, 17, 16, 11
- $v = 32$: 26, 25, 24, 23, 1, 2, 7, 6, 5, 4, 3, 28, 29, 30, 31, 12, 13, 8, 9, 10, 11, 16, 15, 14, 19, 20, 21, 22, 17, 18, 33, 34, 35, 27, 0
- $v = 33$: 26, 4, 3, 28, 27, 35, 34, 21, 20, 25, 24, 23, 22, 17, 18, 19, 14, 13, 12, 31, 32, 15, 16, 11, 10, 5, 6, 29, 30, 9, 8, 7, 2, 1, 0
- $v = 34$: 26, 25, 24, 35, 27, 28, 29, 6, 5, 4, 3, 2, 7, 8, 13, 12, 31, 30, 9, 10, 11, 16, 17, 18, 33, 32, 15, 14, 19, 20, 21, 22, 23, 1, 0
- $v = 35$: 26, 4, 3, 2, 1, 23, 24, 25, 20, 19, 18, 17, 22, 21, 34, 33, 32, 31, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 30, 29, 28, 27, 0

Proof that b_0c_0 and b_4c_4 are suitable edges for J_{11}

Figure 2 shows the flower snark J_{11} and here b_0c_0 and b_4c_4 correspond to the edges $(0, 32)$ and $(11, 12)$, respectively.

$(0, 11)$, $(0, 12)$, $(32, 11)$ and $(32, 12)$ are good in $J_{11} - \{(0, 32), (11, 12)\}$ due to the following hamiltonian paths, respectively:

- 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 43, 30, 29, 28, 23, 22, 17, 16, 15, 38, 37, 12, 13, 14, 19, 18, 39, 40, 21, 20, 25, 24, 41, 42, 27, 26, 31, 32, 4, 3, 2, 1, 0
- 12, 13, 14, 15, 16, 11, 10, 5, 4, 32, 31, 26, 25, 24, 41, 42, 27, 28, 23, 22, 17, 18, 19, 20, 21, 40, 39, 38, 37, 36, 9, 8, 7, 6, 35, 34, 3, 2, 1, 29, 30, 43, 33, 0
- 11, 10, 5, 4, 3, 2, 1, 0, 33, 34, 35, 6, 7, 8, 9, 36, 37, 12, 13, 14, 19, 18, 17, 16, 15, 38, 39, 40, 41, 24, 25, 20, 21, 22, 23, 28, 29, 30, 43, 42, 27, 26, 31, 32
- 12, 13, 8, 7, 2, 3, 4, 5, 6, 35, 34, 33, 0, 1, 29, 28, 23, 22, 17, 18, 39, 38, 37, 36, 9, 10, 11, 16, 15, 14, 19, 20, 21, 40, 41, 24, 25, 26, 27, 42, 43, 30, 31, 32

$(0, 32)$, $(11, 12)$ is good in $J_{11} - \{(0, 32), (11, 12)\}$ due to the following two disjoint paths with end-vertices 0 and 32, and 11 and 12, respectively, which together span J_{11} .

- 32, 31, 26, 25, 20, 19, 14, 13, 8, 7, 2, 1, 0

- 12, 37, 38, 15, 16, 17, 18, 39, 40, 21, 22, 23, 24, 41, 42, 27, 28, 29, 30, 43, 33, 34, 3, 4, 5, 6, 35, 36, 9, 10, 11

The following hamiltonian paths show that at least one of $(0, 32)$ or $(11, 12)$ is good in $J_{11} - \{v, (0, 32), (11, 12)\}$ for every $v \in V(J_{11})$.

- $v = 0$: 12, 13, 14, 15, 38, 37, 36, 35, 6, 5, 10, 9, 8, 7, 2, 1, 29, 30, 31, 32, 4, 3, 34, 33, 43, 42, 41, 24, 23, 28, 27, 26, 25, 20, 19, 18, 39, 40, 21, 22, 17, 16, 11
- $v = 1$: 32, 4, 5, 6, 7, 2, 3, 34, 35, 36, 37, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 38, 39, 40, 41, 24, 23, 22, 21, 20, 25, 26, 31, 30, 29, 28, 27, 42, 43, 33, 0
- $v = 2$: 12, 37, 38, 15, 14, 13, 8, 7, 6, 5, 10, 9, 36, 35, 34, 3, 4, 32, 31, 30, 29, 1, 0, 33, 43, 42, 41, 24, 23, 28, 27, 26, 25, 20, 19, 18, 39, 40, 21, 22, 17, 16, 11
- $v = 3$: 32, 4, 5, 6, 7, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 33, 0
- $v = 4$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 3, 2, 1, 29, 30, 43, 33, 0
- $v = 5$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11, 10, 9, 36, 37, 12, 13, 8, 7, 6, 35, 34, 33, 0
- $v = 6$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 5, 4, 3, 2, 7, 8, 9, 36, 35, 34, 33, 43, 30, 29, 1, 0
- $v = 7$: 32, 4, 5, 6, 35, 36, 37, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 38, 39, 40, 41, 24, 23, 22, 21, 20, 25, 26, 31, 30, 29, 28, 27, 42, 43, 33, 34, 3, 2, 1, 0
- $v = 8$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 9, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 9$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11, 10, 5, 6, 7, 8, 13, 12, 37, 36, 35, 34, 33, 0
- $v = 10$: 12, 13, 14, 15, 38, 37, 36, 9, 8, 7, 2, 3, 34, 35, 6, 5, 4, 32, 31, 30, 29, 1, 0, 33, 43, 42, 41, 24, 23, 28, 27, 26, 25, 20, 19, 18, 39, 40, 21, 22, 17, 16, 11
- $v = 11$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 20, 25, 24, 23, 22, 17, 16, 15, 14, 19, 18, 39, 38, 37, 12, 13, 8, 7, 6, 5, 10, 9, 36, 35, 34, 33, 0
- $v = 12$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 8, 9, 10, 11, 16, 15, 38, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 13$: 12, 37, 36, 35, 6, 5, 10, 9, 8, 7, 2, 1, 0, 33, 34, 3, 4, 32, 31, 26, 27, 28, 29, 30, 43, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11
- $v = 14$: 32, 31, 26, 25, 24, 41, 42, 27, 28, 23, 22, 17, 18, 19, 20, 21, 40, 39, 38, 15, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 15$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 14, 13, 12, 37, 38, 39, 18, 17, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 16$: 12, 13, 8, 7, 2, 3, 34, 35, 6, 5, 4, 32, 31, 30, 29, 1, 0, 33, 43, 42, 41, 24, 25, 26, 27, 28, 23, 22, 17, 18, 39, 40, 21, 20, 19, 14, 15, 38, 37, 36, 9, 10, 11
- $v = 17$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 18, 39, 38, 37, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 18$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 38, 15, 14, 19, 20, 21, 22, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 19$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 20, 25, 24, 23, 22, 17, 18, 39, 38, 37, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 20$: 32, 31, 26, 25, 24, 41, 42, 27, 28, 23, 22, 21, 40, 39, 38, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 21$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 39, 18, 17, 22, 23, 24, 25, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0

- $v = 22$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 21, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 23$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 24, 25, 20, 19, 18, 17, 22, 21, 40, 39, 38, 37, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 24$: 32, 31, 26, 25, 20, 21, 22, 23, 28, 27, 42, 41, 40, 39, 38, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 25$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 24, 23, 22, 17, 18, 19, 20, 21, 40, 39, 38, 37, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 26$: 32, 31, 30, 29, 1, 2, 7, 6, 5, 4, 3, 34, 35, 36, 37, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 38, 39, 40, 41, 24, 25, 20, 21, 22, 23, 28, 27, 42, 43, 33, 0
- $v = 27$: 32, 4, 3, 2, 1, 29, 28, 23, 24, 25, 26, 31, 30, 43, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 28$: 32, 31, 26, 27, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 29$: 32, 4, 3, 34, 33, 43, 30, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 5, 6, 35, 36, 9, 8, 7, 2, 1, 0
- $v = 30$: 32, 31, 26, 25, 20, 21, 22, 23, 24, 41, 40, 39, 38, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 28, 27, 42, 43, 33, 0
- $v = 31$: 32, 4, 3, 2, 1, 29, 30, 43, 42, 41, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 40, 39, 38, 37, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 32$: 12, 13, 8, 7, 2, 3, 4, 5, 6, 35, 34, 33, 0, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 20, 25, 24, 23, 22, 17, 16, 15, 14, 19, 18, 39, 38, 37, 36, 9, 10, 11
- $v = 33$: 32, 4, 5, 6, 7, 2, 3, 34, 35, 36, 37, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 38, 39, 40, 41, 24, 23, 22, 21, 20, 25, 26, 31, 30, 43, 42, 27, 28, 29, 1, 0
- $v = 34$: 12, 13, 14, 15, 38, 37, 36, 35, 6, 5, 10, 9, 8, 7, 2, 3, 4, 32, 31, 30, 29, 1, 0, 33, 43, 42, 41, 24, 23, 28, 27, 26, 25, 20, 19, 18, 39, 40, 21, 22, 17, 16, 11
- $v = 35$: 32, 4, 5, 6, 7, 8, 13, 12, 37, 36, 9, 10, 11, 16, 17, 18, 19, 14, 15, 38, 39, 40, 41, 24, 23, 22, 21, 20, 25, 26, 31, 30, 29, 28, 27, 42, 43, 33, 34, 3, 2, 1, 0
- $v = 36$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 9, 8, 7, 2, 3, 4, 5, 6, 35, 34, 33, 43, 30, 29, 1, 0
- $v = 37$: 12, 13, 8, 7, 6, 5, 10, 9, 36, 35, 34, 33, 0, 1, 2, 3, 4, 32, 31, 26, 27, 28, 29, 30, 43, 42, 41, 40, 21, 22, 23, 24, 25, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11
- $v = 38$: 32, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 15, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 39$: 32, 4, 3, 2, 1, 29, 28, 27, 26, 31, 30, 43, 42, 41, 40, 21, 20, 25, 24, 23, 22, 17, 18, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 40$: 32, 31, 26, 25, 24, 41, 42, 27, 28, 23, 22, 21, 20, 19, 14, 15, 38, 39, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 41$: 32, 4, 3, 2, 1, 29, 28, 27, 42, 43, 30, 31, 26, 25, 24, 23, 22, 17, 18, 19, 20, 21, 40, 39, 38, 37, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0
- $v = 42$: 32, 31, 26, 27, 28, 23, 22, 21, 20, 25, 24, 41, 40, 39, 38, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 37, 36, 35, 34, 3, 4, 5, 6, 7, 2, 1, 29, 30, 43, 33, 0
- $v = 43$: 32, 4, 3, 2, 1, 29, 30, 31, 26, 25, 24, 23, 28, 27, 42, 41, 40, 39, 18, 17, 22, 21, 20, 19, 14, 13, 12, 37, 38, 15, 16, 11, 10, 5, 6, 7, 8, 9, 36, 35, 34, 33, 0

Proof that b_0c_0 and b_4c_4 are suitable edges for J_{13}

Figure 3 shows the flower snark J_{13} and here b_0c_0 and b_4c_4 correspond to the edges $(0, 38)$ and $(11, 12)$, respectively.

$(0, 11)$, $(0, 12)$, $(38, 11)$ and $(38, 12)$ are good in $J_{13} - \{(0, 38), (11, 12)\}$ due to the following hamiltonian paths, respectively:

- 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 51, 36, 35, 34, 29, 28, 23, 22, 17, 16, 15, 44, 43, 12, 13, 14, 19, 18, 45, 46, 21, 20, 25, 24, 47, 48, 27, 26, 31, 30, 49, 50, 33, 32, 37, 38, 4, 3, 2, 1, 0
- 12, 13, 14, 15, 16, 11, 10, 5, 4, 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 42, 9, 8, 7, 6, 41, 40, 3, 2, 1, 35, 36, 51, 39, 0
- 11, 10, 5, 4, 3, 2, 1, 0, 39, 40, 41, 6, 7, 8, 9, 42, 43, 12, 13, 14, 19, 18, 17, 16, 15, 44, 45, 46, 47, 24, 23, 22, 21, 20, 25, 26, 31, 30, 29, 28, 27, 48, 49, 50, 51, 36, 35, 34, 33, 32, 37, 38
- 12, 13, 8, 7, 2, 3, 4, 5, 6, 41, 40, 39, 0, 1, 35, 34, 29, 28, 23, 22, 17, 18, 45, 44, 43, 42, 9, 10, 11, 16, 15, 14, 19, 20, 21, 46, 47, 24, 25, 26, 27, 48, 49, 30, 31, 32, 33, 50, 51, 36, 37, 38

$(0, 38), (11, 12)$ is good in $J_{13} - \{(0, 38), (11, 12)\}$ due to the following two disjoint paths with end-vertices 0 and 38, and 11 and 12, respectively, which together span J_{13} .

- 38, 37, 32, 31, 26, 25, 20, 19, 14, 13, 8, 7, 2, 1, 0
- 12, 43, 44, 15, 16, 17, 18, 45, 46, 21, 22, 23, 24, 47, 48, 27, 28, 29, 30, 49, 50, 33, 34, 35, 36, 51, 39, 40, 3, 4, 5, 6, 41, 42, 9, 10, 11

The following hamiltonian paths show that at least one of $(0, 38)$ or $(11, 12)$ is good in $J_{13} - \{v, (0, 38), (11, 12)\}$ for every $v \in V(J_{13})$.

- $v = 0$: 12, 13, 14, 15, 44, 43, 42, 41, 6, 5, 10, 9, 8, 7, 2, 1, 35, 36, 37, 38, 4, 3, 40, 39, 51, 50, 49, 30, 29, 34, 33, 32, 31, 26, 25, 24, 23, 28, 27, 48, 47, 46, 45, 18, 19, 20, 21, 22, 17, 16, 11
- $v = 1$: 38, 4, 5, 6, 7, 2, 3, 40, 41, 42, 43, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 44, 45, 46, 47, 24, 23, 22, 21, 20, 25, 26, 31, 30, 49, 48, 27, 28, 29, 34, 35, 36, 37, 32, 33, 50, 51, 39, 0
- $v = 2$: 12, 43, 44, 15, 14, 13, 8, 7, 6, 5, 10, 9, 42, 41, 40, 3, 4, 38, 37, 36, 35, 1, 0, 39, 51, 50, 49, 30, 29, 34, 33, 32, 31, 26, 25, 24, 23, 28, 27, 48, 47, 46, 45, 18, 19, 20, 21, 22, 17, 16, 11
- $v = 3$: 38, 4, 5, 6, 7, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 39, 0
- $v = 4$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 3, 2, 1, 35, 36, 51, 39, 0
- $v = 5$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 42, 43, 12, 13, 8, 7, 6, 41, 40, 39, 0
- $v = 6$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 4, 3, 2, 7, 8, 9, 42, 41, 40, 39, 51, 36, 35, 1, 0
- $v = 7$: 38, 4, 5, 6, 41, 42, 43, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 44, 45, 46, 47, 24, 23, 22, 21, 20, 25, 26, 31, 30, 29, 28, 27, 48, 49, 50, 51, 36, 37, 32, 33, 34, 35, 1, 2, 3, 40, 39, 0
- $v = 8$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 9, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 9$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 5, 6, 7, 8, 13, 12, 43, 42, 41, 40, 39, 0
- $v = 10$: 12, 13, 14, 15, 44, 43, 42, 9, 8, 7, 2, 3, 40, 41, 6, 5, 4, 38, 37, 36, 35, 1, 0, 39, 51, 50, 49, 30, 29, 34, 33, 32, 31, 26, 25, 24, 23, 28, 27, 48, 47, 46, 45, 18, 19, 20, 21, 22, 17, 16, 11
- $v = 11$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 46, 47, 24, 25, 20, 19, 14, 15, 16, 17, 18, 45, 44, 43, 12, 13, 8, 7, 6, 5, 10, 9, 42, 41, 40, 39, 0
- $v = 12$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 42, 9, 8, 13, 14, 15, 16, 11, 10, 5, 4, 3, 2, 7, 6, 41, 40, 39, 51, 36, 35, 1, 0
- $v = 13$: 12, 43, 42, 41, 6, 5, 10, 9, 8, 7, 2, 1, 0, 39, 40, 3, 4, 38, 37, 32, 33, 34, 35, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11
- $v = 14$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 15, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0

- $v = 15$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 43, 12, 13, 14, 19, 18, 17, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 16$: 12, 13, 8, 7, 2, 3, 40, 41, 6, 5, 4, 38, 37, 36, 35, 1, 0, 39, 51, 50, 49, 30, 29, 34, 33, 32, 31, 26, 25, 24, 23, 28, 27, 48, 47, 46, 45, 18, 17, 22, 21, 20, 19, 14, 15, 44, 43, 42, 9, 10, 11
- $v = 17$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 18, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 18$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 14, 15, 44, 45, 46, 21, 22, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 19$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 28, 29, 30, 31, 26, 25, 20, 21, 46, 47, 24, 23, 22, 17, 18, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 20$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 25, 26, 27, 28, 23, 22, 21, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 21$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 17, 18, 19, 20, 25, 24, 47, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 22$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 21, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 23$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 28, 29, 30, 31, 26, 25, 24, 47, 46, 45, 18, 17, 22, 21, 20, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 24$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 46, 21, 22, 23, 28, 27, 26, 25, 20, 19, 14, 15, 44, 45, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 25$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 24, 47, 46, 45, 18, 17, 22, 21, 20, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 26$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 27, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 27$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 47, 24, 23, 28, 29, 30, 31, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 28$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 27, 26, 25, 20, 21, 22, 23, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 29$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 30, 31, 26, 25, 24, 23, 28, 27, 48, 47, 46, 45, 18, 17, 22, 21, 20, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 30$: 38, 37, 32, 31, 26, 27, 28, 29, 34, 33, 50, 49, 48, 47, 46, 21, 22, 23, 24, 25, 20, 19, 14, 15, 44, 45, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 31$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 30, 29, 28, 23, 24, 25, 26, 27, 48, 47, 46, 45, 18, 17, 22, 21, 20, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 32$: 38, 37, 36, 35, 1, 2, 7, 6, 5, 4, 3, 40, 41, 42, 43, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 44, 45, 46, 47, 24, 23, 22, 21, 20, 25, 26, 31, 30, 49, 48, 27, 28, 29, 34, 33, 50, 51, 39, 0
- $v = 33$: 38, 4, 3, 2, 1, 35, 34, 29, 30, 31, 32, 37, 36, 51, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 34$: 38, 37, 32, 33, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 35$: 38, 4, 3, 40, 39, 51, 36, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 41, 42, 9, 8, 7, 2, 1, 0
- $v = 36$: 38, 37, 32, 31, 26, 27, 28, 29, 30, 49, 48, 47, 46, 21, 22, 23, 24, 25, 20, 19, 14, 15, 44, 45, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 34, 33, 50, 51, 39, 0
- $v = 37$: 38, 4, 3, 2, 1, 35, 36, 51, 50, 49, 30, 29, 34, 33, 32, 31, 26, 25, 24, 23, 28, 27, 48, 47, 46, 45, 18, 17, 22, 21, 20, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 38$: 12, 13, 8, 7, 2, 3, 4, 5, 6, 41, 40, 39, 0, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 46, 47, 24, 25, 20, 19, 14, 15, 16, 17, 18, 45, 44, 43, 42, 9, 10, 11
- $v = 39$: 38, 4, 5, 6, 7, 2, 3, 40, 41, 42, 43, 12, 13, 8, 9, 10, 11, 16, 17, 18, 19, 14, 15, 44, 45, 46, 47, 24, 23, 22, 21, 20, 25, 26, 31, 30, 29, 28, 27, 48, 49, 50, 51, 36, 37, 32, 33, 34, 35, 1, 0

- $v = 40$: 12, 13, 14, 15, 44, 43, 42, 41, 6, 5, 10, 9, 8, 7, 2, 3, 4, 38, 37, 36, 35, 1, 0, 39, 51, 50, 49, 30, 29, 34, 33, 32, 31, 26, 25, 24, 23, 28, 27, 48, 47, 46, 45, 18, 19, 20, 21, 22, 17, 16, 11
- $v = 41$: 38, 4, 5, 6, 7, 8, 13, 12, 43, 42, 9, 10, 11, 16, 17, 18, 19, 14, 15, 44, 45, 46, 47, 24, 23, 22, 21, 20, 25, 26, 31, 30, 29, 28, 27, 48, 49, 50, 51, 36, 37, 32, 33, 34, 35, 1, 2, 3, 40, 39, 0
- $v = 42$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 9, 8, 7, 2, 3, 4, 5, 6, 41, 40, 39, 51, 36, 35, 1, 0
- $v = 43$: 12, 13, 8, 7, 6, 5, 10, 9, 42, 41, 40, 39, 0, 1, 2, 3, 4, 38, 37, 32, 33, 34, 35, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 22, 21, 20, 25, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11
- $v = 44$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 25, 26, 27, 28, 23, 22, 17, 18, 45, 46, 21, 20, 19, 14, 15, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 45$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 28, 29, 30, 31, 26, 25, 20, 21, 46, 47, 24, 23, 22, 17, 18, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 46$: 38, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 25, 26, 27, 28, 23, 22, 21, 20, 19, 14, 15, 44, 45, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 47$: 38, 4, 3, 2, 1, 35, 34, 33, 32, 37, 36, 51, 50, 49, 48, 27, 26, 31, 30, 29, 28, 23, 24, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 48$: 38, 37, 32, 31, 30, 49, 50, 33, 34, 29, 28, 27, 26, 25, 20, 21, 22, 23, 24, 47, 46, 45, 44, 15, 14, 19, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 49$: 38, 4, 3, 2, 1, 35, 34, 33, 50, 51, 36, 37, 32, 31, 30, 29, 28, 23, 24, 25, 26, 27, 48, 47, 46, 45, 18, 17, 22, 21, 20, 19, 14, 13, 12, 43, 44, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0
- $v = 50$: 38, 37, 32, 33, 34, 29, 28, 27, 26, 31, 30, 49, 48, 47, 46, 21, 22, 23, 24, 25, 20, 19, 14, 15, 44, 45, 18, 17, 16, 11, 10, 9, 8, 13, 12, 43, 42, 41, 40, 3, 4, 5, 6, 7, 2, 1, 35, 36, 51, 39, 0
- $v = 51$: 38, 4, 3, 2, 1, 35, 36, 37, 32, 31, 30, 29, 34, 33, 50, 49, 48, 47, 24, 23, 28, 27, 26, 25, 20, 19, 18, 17, 22, 21, 46, 45, 44, 43, 12, 13, 14, 15, 16, 11, 10, 5, 6, 7, 8, 9, 42, 41, 40, 39, 0