

FUNCTIONAL INEQUALITIES INVOLVING NUMERICAL DIFFERENTIATION FORMULAS OF ORDER TWO

TOMASZ SZOSTOK

ABSTRACT. We write expressions connected with numerical differentiation formulas of order 2 in the form of Stieltjes integral, then we use Ohlin lemma and Levin-Stechkin theorem to study inequalities connected with these expressions. In particular, we present a new proof of the inequality

$$(1) \quad f\left(\frac{x+y}{2}\right) \leq \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt \leq \frac{1}{y-x} \int_x^y f(t) dt$$

satisfied by every convex function $f : \mathbb{R} \rightarrow \mathbb{R}$ and we obtain extensions of (1). Then we deal with nonsymmetric inequalities of a similar form.

1. INTRODUCTION

Writing the celebrated Hermite-Hadamard inequality

$$(2) \quad f\left(\frac{x+y}{2}\right) \leq \frac{1}{y-x} \int_x^y f(t) dt \leq \frac{f(x) + f(y)}{2}$$

in the form

$$(3) \quad f\left(\frac{x+y}{2}\right) \leq \frac{F(y) - F(x)}{y-x} \leq \frac{f(x) + f(y)}{2}$$

we can see that (2) is, in fact, an inequality involving two very simple quadrature operators and a very simple differentiation formula. In papers [11] and [12] the quadrature operators occurring in (2) were replaced by more general ones whereas in [9] the middle term from (2) was replaced by more general formulas used in numerical differentiation. Thus inequalities involving expressions of the form

$$\frac{\sum_{i=1}^n a_i F(\alpha_i x + \beta_i y)}{y-x}$$

2010 *Mathematics Subject Classification.* 26A51, 26D10, 39B62.

Key words and phrases. Hermite-Hadamard inequality, differentiation formulas, convex functions.

where $\sum_{i=1}^n a_i = 0$, $\alpha_i + \beta_i = 1$ and $F' = f$ were considered. In the current paper we deal with inequalities for expressions of the form

$$(4) \quad \frac{\sum_{i=1}^n a_i \Phi(\alpha_i x + \beta_i y)}{(y-x)^2}$$

(where $\Phi'' = f$) which are used to approximate the second order derivative of F and, surprisingly, we discover a connection between our approach and the inequality (1).

First we make the following simple observation.

Remark 1. Let $f, F, \Phi : [x, y] \rightarrow \mathbb{R}$ be such that $\Phi' = F, F' = f$, let $n_i, m_i \in \mathbb{N} \cup \{0\}, i = 1, 2, 3$ $a_{i,j} \in \mathbb{R}, \alpha_{i,j}, \beta_{i,j} \in [0, 1], \alpha_{i,j} + \beta_{i,j} = 1, i = 1, 2, 3; j = 1, \dots, n_i$ $b_{i,j} \in \mathbb{R}, \gamma_{i,j}, \delta_{i,j} \in [0, 1], \gamma_{i,j} + \delta_{i,j} = 1, i = 1, 2, 3; j = 1, \dots, m_i$. If the inequality

$$(5) \quad \sum_{i=1}^{n_1} a_{1,i} f(\alpha_{1,i} x + \beta_{1,i} y) + \frac{\sum_{i=1}^{n_2} a_{2,i} F(\alpha_{2,i} x + \beta_{2,i} y)}{y-x} \\ + \frac{\sum_{i=1}^{n_3} a_{3,i} \Phi(\alpha_{3,i} x + \beta_{3,i} y)}{(y-x)^2} \leq \sum_{i=1}^{m_1} b_{1,i} f(\gamma_{1,i} x + \delta_{1,i} y) \\ + \frac{\sum_{i=1}^{m_2} b_{2,i} F(\gamma_{2,i} x + \delta_{2,i} y)}{y-x} + \frac{\sum_{i=1}^{m_3} b_{3,i} \Phi(\gamma_{3,i} x + \delta_{3,i} y)}{(y-x)^2}$$

is satisfied for $x = 0, y = 1$ and for all continuous and convex functions $f : [0, 1] \rightarrow \mathbb{R}$ then it is satisfied for all $x, y \in \mathbb{R}, x < y$ and for each continuous and convex function $f : [x, y] \rightarrow \mathbb{R}$. To see this it is enough to observe that expressions from (5) remain unchanged if we replace $f : [x, y] \rightarrow \mathbb{R}$ by $\varphi : [0, 1] \rightarrow \mathbb{R}$ given by $\varphi(t) := f(x + t(y-x))$.

The simplest expression used to approximate the second order derivative of f is of the form

$$f''\left(\frac{x+y}{2}\right) \approx \frac{f(x) - 2f\left(\frac{x+y}{2}\right) + f(y)}{\left(\frac{y-x}{2}\right)^2}$$

Remark 2. From numerical analysis it is known that

$$f''\left(\frac{x+y}{2}\right) = \frac{f(x) - 2f\left(\frac{x+y}{2}\right) + f(y)}{\left(\frac{y-x}{2}\right)^2} - \frac{\left(\frac{y-x}{2}\right)^2}{12} f^{(4)}(\xi).$$

This means that for convex g and for G such that $G''' = g$ we have

$$g\left(\frac{x+y}{2}\right) \leq \frac{G(x) - 2G\left(\frac{x+y}{2}\right) + G(y)}{\left(\frac{y-x}{2}\right)^2}.$$

In this paper we shall obtain some inequalities for convex functions which do not follow from numerical differentiation results.

In order to get such results we shall use Stieltjes integral. In paper [10] it was observed that the classical Hermite-Hadamard inequality (2) easily follows from the following Ohlin lemma

Lemma 1. (Ohlin [8]) *Let X_1, X_2 be two random variables such that $EX_1 = EX_2$ and let F_1, F_2 be their distribution functions. If F_1, F_2 satisfy for some x_0 the following inequalities*

$$F_1(x) \leq F_2(x) \text{ if } x < x_0 \text{ and } F_1(x) \geq F_2(x) \text{ if } x > x_0$$

then

$$(6) \quad Ef(X_1) \leq Ef(X_2)$$

for all continuous and convex functions $f : \mathbb{R} \rightarrow \mathbb{R}$.

Ohlin lemma was used also in paper [11]. However in the present approach (similarly as in [9] and [12]) we are going to use a more general result from [6], (see also [7] Theorem 4.2.7). In this theorem we use the notations from [7].

Theorem 1. (Levin, Stechkin) *Let $F_1, F_2 : [a, b] \rightarrow \mathbb{R}$ be two functions with bounded variation such that $F_1(a) = F_2(a)$. Then, in order that*

$$\int_a^b f(x)dF_1(x) \leq \int_a^b f(x)dF_2(x)$$

for all continuous and convex functions $f : [a, b] \rightarrow \mathbb{R}$ it is necessary and sufficient that F_1 and F_2 verify the following three conditions:

$$(7) \quad F_1(b) = F_2(b),$$

$$(8) \quad \int_a^x F_1(t)dt \leq \int_a^x F_2(t)dt, \quad x \in (a, b),$$

and

$$(9) \quad \int_a^b F_1(t)dt = \int_a^b F_2(t)dt.$$

Remark 3. As it easy to see, if measures occurring in Ohlin lemma are concentrated on the interval $[x, y]$ then this lemma is an easy consequence of Theorem 1. However Theorem 1 is more general for two reasons: it allows functions F_1, F_2 to have more crossing points than one and functions F_1, F_2 do not have to be cumulative distribution functions. Therefore we shall use this theorem even if functions F_1, F_2 have exactly one crossing point.

Let now $f : [x, y] \rightarrow \mathbb{R}$ be any function and let $F, \Phi : [x, y] \rightarrow \mathbb{R}$ be such that $F' = f$ and $\Phi' = F$. We need to write the expression

$$(10) \quad \frac{\Phi(x) - 2\Phi\left(\frac{x+y}{2}\right) + \Phi(y)}{\left(\frac{y-x}{2}\right)^2}$$

in the form

$$\int_x^y f dF_1$$

for some F_1 . In the next proposition we show that it is possible – here for the sake of simplicity we shall work on the interval $[0, 1]$.

Proposition 1. *Let $f : [0, 1] \rightarrow \mathbb{R}$ be any function and let $\Phi : [0, 1] \rightarrow \mathbb{R}$ be such that $\Phi'' = f$. Then we have*

$$4\left(\Phi(0) - 2\Phi\left(\frac{1}{2}\right) + \Phi(1)\right) = \int_x^y f dF_1$$

where $F_1 : [0, 1] \rightarrow \mathbb{R}$ is given by

$$(11) \quad F_1(t) := \begin{cases} 2x^2 & x \leq \frac{1}{2}, \\ -2x^2 + 4x - 1 & x > \frac{1}{2}. \end{cases}$$

Proof. Let $F : [0, 1] \rightarrow \mathbb{R}$ be such that $\Phi' = F$. Now, to prove this proposition it is enough to do the following calculations

$$\begin{aligned} \int_0^1 f dF_1 &= \int_0^{\frac{1}{2}} 4xf(x)dx + \int_{\frac{1}{2}}^1 (-4x + 4)f(x)dx \\ &= 2F\left(\frac{1}{2}\right) - 0 \cdot F(0) - \int_0^{\frac{1}{2}} 4F(x)dx - 0 \cdot F(1) - 2F\left(\frac{1}{2}\right) + \int_{\frac{1}{2}}^1 4F(x)dx \\ &= 4\Phi(1) - 8\Phi\left(\frac{1}{2}\right) + 4\Phi(0). \end{aligned}$$

□

Remark 4. Observe that if Φ and f are such as in Proposition 1 then the following equality is satisfied

$$\frac{\Phi(x) - 2\Phi\left(\frac{x+y}{2}\right) + \Phi(y)}{\left(\frac{y-x}{2}\right)^2} = \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt.$$

After this observation it turns out that inequalities involving the expression (10) were considered in the paper of Dragomir [3] where (among others) the following inequalities were obtained

$$(12) \quad f\left(\frac{x+y}{2}\right) \leq \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt \leq \frac{1}{y-x} \int_x^y f(t)dt.$$

As we already know (Remark 2) the first one of the above inequalities may be obtained using the numerical analysis results.

Now the inequalities from the Dragomir's paper easily follow from Ohlin lemma but there are many possibilities of generalizations and modifications of inequalities (12). These generalizations will be discussed in the following chapters.

2. THE SYMMETRIC CASE

We start with the following remark.

Remark 5. Let $F_*(t) = at^2 + bt + c$ for some $a, b, c \in \mathbb{R}, a \neq 0$. It is impossible to obtain inequalities involving $\int_x^y f dF_*$ and any of the expressions: $\frac{1}{y-x} \int_x^y f(t)dt, f\left(\frac{x+y}{2}\right), \frac{f(x)+f(y)}{2}$ which were satisfied for all convex functions $f : [x, y] \rightarrow \mathbb{R}$. Indeed, suppose for example that we have

$$\int_x^y f dF_* \leq \frac{1}{y-x} \int_x^y f(t)dt$$

for all convex $f : [x, y] \rightarrow \mathbb{R}$. Without loss of generality we may assume that $F_*(x) = 0$, then from Theorem 1 we have $F_*(y) = 1$ Also from Theorem 1 we get

$$\int_x^y F_*(t)dt = \int_x^y F_0 dt$$

where $F_0(t) = \frac{t-x}{y-x}, t \in [x, y]$ which is impossible, since F_* is either strictly convex or concave.

This remark means that in order to get some new inequalities of the Hermite-Hadamard type we have to integrate with respect to functions constructed with use of (at least) two quadratic functions, as it was the case in Proposition 1 Now we may present the main result of this section.

Theorem 2. *Let x, y be some real numbers such that $x < y$ and let $a \in \mathbb{R}$. Let $f, F, \Phi : [x, y] \rightarrow \mathbb{R}$ be any functions such that $F' = f$ and $\Phi' = F$ and let $T_a f(x, y)$ be defined by the formula*

$$T_a f(x, y) = \left(1 - \frac{a}{2}\right) \frac{F(y) - F(x)}{y-x} + 2a \frac{\Phi(x) - 2\Phi\left(\frac{x+y}{2}\right) + \Phi(y)}{(y-x)^2}.$$

*Then the following inequalities hold for all convex functions f :
if $a \geq 0$ then*

$$(13) \quad T_a f(x, y) \leq \frac{1}{y-x} \int_x^y f(t)dt,$$

if $a \leq 0$ then

$$(14) \quad T_a f(x, y) \geq \frac{1}{y-x} \int_x^y f(t) dt,$$

if $a \leq 2$ then

$$(15) \quad f\left(\frac{x+y}{2}\right) \leq T_a f(x, y),$$

if $a \geq 6$ then

$$(16) \quad T_a f(x, y) \leq f\left(\frac{x+y}{2}\right),$$

if $a \geq -6$ then

$$(17) \quad T_a f(x, y) \leq \frac{f(x) + f(y)}{2},$$

Furthermore:

if $a \in (2, 6)$ then the expressions $T_a f(x, y)$, $f\left(\frac{x+y}{2}\right)$ are not comparable in the class of convex functions

if $a < -6$ then expressions $T_a f(x, y)$, $\frac{f(x)+f(y)}{2}$ are not comparable in the class of convex functions.

Proof. In view of Remark 1 we may restrict ourselves to the case $x = 0, y = 1$. Take $a \in \mathbb{R}$, let $f : [0, 1] \rightarrow \mathbb{R}$ be any convex function and let $F, \Phi : [0, 1] \rightarrow \mathbb{R}$ be such that $F' = f, \Phi' = F$. Define $F_1 : [0, 1] \rightarrow \mathbb{R}$ by the formula

$$(18) \quad F_1(t) := \begin{cases} at^2 + \left(1 - \frac{a}{2}\right)t & t < \frac{1}{2}, \\ -at^2 + \left(1 + \frac{3a}{2}\right)t - \frac{a}{2} & t \geq \frac{1}{2}. \end{cases}$$

First we shall prove that $T_a f(0, 1) = \int_0^1 f dF_1$. Indeed, we have

$$\begin{aligned} \int_0^1 f dF_1 &= \int_0^{\frac{1}{2}} f dF_1 + \int_{\frac{1}{2}}^1 f dF_1 = \int_0^{\frac{1}{2}} f(t) \left(2at + 1 - \frac{a}{2}\right) dt \\ &+ \int_{\frac{1}{2}}^1 f(t) \left(-2at + 1 + \frac{3a}{2}\right) dt = F\left(\frac{1}{2}\right) \left(\frac{a}{2} + 1\right) - F(0) \left(1 - \frac{a}{2}\right) \\ &- \int_0^{\frac{1}{2}} F(t) 2adt + F(1) \left(1 - \frac{a}{2}\right) - F\left(\frac{1}{2}\right) \left(\frac{a}{2} + 1\right) + \int_{\frac{1}{2}}^1 F(t) 2adt \\ &= \left(1 - \frac{a}{2}\right) (F(1) - F(0)) + 2a \left(\Phi(0) - 2\Phi\left(\frac{1}{2}\right) + \Phi(1)\right) \end{aligned}$$

Now let $F_2(t) = t, t \in [0, 1]$. Then functions F_1, F_2 have exactly one crossing point (at $\frac{1}{2}$) and

$$\int_0^1 F_1(t)dt = \frac{1}{2} = \int_0^1 tdt.$$

Moreover if $a > 0$ then function F_1 is convex on the interval $(0, \frac{1}{2})$ and concave on $(\frac{1}{2}, 1)$. Therefore it follows from Ohlin lemma that for $a > 0$ we have

$$(19) \quad \int_0^1 f dF_1 \leq \int_0^1 f dF_2$$

which, in view of Remark 1, yields (13) and for $a < 0$ the opposite inequality is satisfied which gives (14).

Take

$$(20) \quad F_3(t) := \begin{cases} 0 & t \leq \frac{1}{2}, \\ 1 & t > \frac{1}{2}. \end{cases}$$

It is easy to check that for $a \leq 2$ we have $F_1(t) \geq F_3(t)$ for $t \in [0, \frac{1}{2}]$, and $F_1(t) \leq F_3(t)$ for $t \in [\frac{1}{2}, 1]$ and this means that from Ohlin lemma we get (15).

Suppose that $a > 2$. Then there are three crossing points of functions F_1 and $F_3 : x_0, \frac{1}{2}, x_1$, where $x_0 \in (0, \frac{1}{2}), x_1 \in (\frac{1}{2}, 1)$ Function

$$\varphi(s) := \int_0^s F_3(t) - F_1(t)dt, \quad s \in [0, 1]$$

is increasing on intervals $[0, x_0], [\frac{1}{2}, x_1]$ and decreasing on $[x_0, \frac{1}{2}]$ and on $[x_1, 1]$. This means that φ takes its absolute minimum at $\frac{1}{2}$. As it is easy to calculate $\varphi(\frac{1}{2}) \geq 0$ if $a \geq 6$ which, in view of Theorem 1, gives us (16).

To see that for $a \in (2, 6)$ expressions $T_a f(x, y)$ and $f(\frac{x+y}{2})$ are not comparable in the class of convex functions it is enough to observe that in this case $\varphi(x_0) > 0$ and $\varphi(\frac{1}{2}) < 0$.

Now let

$$(21) \quad F_4(t) := \begin{cases} 0 & t = 0, \\ \frac{1}{2} & t \in (0, 1), \\ 1 & t = 1. \end{cases}$$

Similarly as before, if $a \geq -2$ then we have $F_1(t) \geq F_4(t)$ for $t \in [0, \frac{1}{2}]$ $F_1(t) \leq F_4(t)$ for $t \in [\frac{1}{2}, 1]$ i.e. there is only one crossing point of these functions and (17) is obvious. However, for $a \in (-2, -6]$ we have

$$(22) \quad \int_0^{\frac{1}{2}} F_1(t)dt \leq \frac{1}{4} = \int_0^{\frac{1}{2}} F_4(t)dt$$

and therefore, in view of Theorem 1 we still have (17). In the case $a < -6$ inequality (22) is no longer true which means that expressions $T_a f(x, y)$ and $\frac{f(x)+f(y)}{2}$ are not comparable in the class of convex functions. \square

This theorem provides us with a full description of inequalities which may be obtained using Stieltjes integral with respect to a function of the form (18). Some of the obtained inequalities are already known. For example from (13) and (14) we obtain the inequality

$$\frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt \leq \frac{1}{y-x} \int_x^y f(t) dt$$

whereas from (15) for $a = 2$ we get the inequality

$$f\left(\frac{x+y}{2}\right) \leq \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt.$$

However inequalities obtained for "critical" values of a i.e. $-6, 6$. are here particularly interesting. In the following corollary we explicitly write these inequalities.

Corollary 1. *For every convex function $f : [x, y] \rightarrow \mathbb{R}$ the following inequalities are satisfied*

$$(23) \quad 3 \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt \leq \frac{2}{y-x} \int_x^y f(t) dt + f\left(\frac{x+y}{2}\right)$$

and

$$(24) \quad \frac{4}{y-x} \int_x^y f(t) dt \leq 3 \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt + \frac{f(x)+f(y)}{2}$$

Remark 6. In the paper [4] S.S. Dragomir and I. Gomm obtained the following inequality

$$(25) \quad 3 \int_x^y f(t) dt \leq 2 \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt + \frac{f(x)+f(y)}{2}.$$

Inequality (24) from Corollary 1 is stronger than (25). Moreover, as it was observed in Theorem 2 inequalities (23) and (24) cannot be improved i.e. the inequality

$$\frac{1}{y-x} \int_x^y f(t) dt \leq \lambda \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) ds dt + (1-\lambda) \frac{f(x)+f(y)}{2}$$

for $\lambda > \frac{3}{4}$ is not satisfied by every convex function $f : [x, y] \rightarrow \mathbb{R}$ and the inequality

$$\frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) dsdt \leq \gamma \frac{1}{y-x} \int_x^y f(t)dt + (1-\gamma)f\left(\frac{x+y}{2}\right)$$

with $\gamma < \frac{2}{3}$ is not true for all convex functions $f : [x, y] \rightarrow \mathbb{R}$.

In Corollary 1 we obtained inequalities for the triples:

$$\frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) dsdt, \int_x^y f(t)dt, \frac{f(x)+f(y)}{2}$$

and

$$\frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) dsdt, \int_x^y f(t)dt, f\left(\frac{x+y}{2}\right).$$

In the next remark we present an analogous result for expressions

$$\frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) dsdt, \frac{f(x)+f(y)}{2}, f\left(\frac{x+y}{2}\right).$$

Remark 7. Using functions: F_1 defined by (11) and F_5 given by

$$(26) \quad F_5(t) := \begin{cases} 0 & t = 0, \\ \frac{1}{6} & t \in \left(0, \frac{1}{2}\right) \\ \frac{5}{6} & t \in \left[\frac{1}{2}, 1\right) \\ 1 & t = 1, \end{cases}$$

we can see that

$$\frac{1}{6}f(x) + \frac{2}{3}f\left(\frac{x+y}{2}\right) + \frac{1}{6}f(y) \geq \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) dsdt$$

for all convex functions $f : [x, y] \rightarrow \mathbb{R}$.

Moreover it is easy to see that the above inequality cannot be strengthened which means that the inequality

$$af(x) + bf\left(\frac{x+y}{2}\right) + af(y) \geq \frac{1}{(y-x)^2} \int_x^y \int_x^y f\left(\frac{s+t}{2}\right) dsdt$$

where $a, b \geq 0, 2a + b = 1$ is not satisfied by all convex functions f if $a < \frac{1}{6}$.

3. THE NON-SYMMETRIC CASE

In this part of the paper we shall obtain inequalities for $f(\alpha x + (1-\alpha)y)$ and for $\alpha f(x) + (1-\alpha)f(y)$ where α is not necessarily equal to $\frac{1}{2}$.

Now, in contrast to the symmetric case (Remark 5), it is possible to prove inequalities using just one quadratic function but before we

do this we shall present a nonsymmetric version of Hermite-Hadamard inequality involving only the primitive function of f .

Proposition 2. *Let x, y be some real numbers such that $x < y$ and let $\alpha \in [0, 1]$. Let $f : [x, y] \rightarrow \mathbb{R}$, be a convex function, let $F : [x, y] \rightarrow \mathbb{R}$ be such that $F' = f$. If $S_\alpha^1 f(x, y)$ is defined by*

$$S_\alpha^1 f(x, y) := \frac{\frac{-\alpha}{1-\alpha}F(x) + \frac{2\alpha-1}{\alpha(1-\alpha)}F(\alpha x + (1-\alpha)y) + \frac{1-\alpha}{\alpha}F(y)}{y-x}$$

then the following inequality is satisfied:

$$(27) \quad f(\alpha x + (1-\alpha)y) \leq S_\alpha^1 f(x, y) \leq \alpha f(x) + (1-\alpha)f(y).$$

Proof. As usually, the proof will be done on the interval $[0, 1]$. Define functions $F_6, F_7, F_8 : [0, 1] \rightarrow \mathbb{R}$ by the following formulas:

$$(28) \quad F_6(t) := \begin{cases} 0 & t \leq 1-\alpha, \\ 1 & t > 1-\alpha, \end{cases} ,$$

$$(29) \quad F_7(t) := \begin{cases} 0 & t = 0, \\ \alpha & t \in (0, 1), \\ 1 & t = 1, \end{cases} ,$$

and

$$(30) \quad F_8(t) := \begin{cases} \frac{\alpha}{1-\alpha}t & t \in [0, 1-\alpha), \\ \frac{1-\alpha}{\alpha}t + \frac{2\alpha-1}{\alpha} & t \in [1-\alpha, 1]. \end{cases} ,$$

We have:

$$\begin{aligned} \int_0^1 F_6(t)dt &= \int_0^1 F_7(t)dt = \int_0^1 F_8(t)dt = \alpha, \\ \int_0^1 f dF_6 &= f(1-\alpha) \\ \int_0^1 f dF_7 &= \alpha f(0) + (1-\alpha)f(1) \\ \int_0^1 f dF_8 &= S_\alpha^1 f(0, 1). \end{aligned}$$

Moreover both of the pairs (F_6, F_8) and (F_8, F_7) has only one crossing point. Thus it suffices to use Theorem 1 to obtain inequalities (27). \square

Theorem 3. *Let x, y be some real numbers such that $x < y$ and let $\alpha \in [0, 1]$. Let $f : [x, y] \rightarrow \mathbb{R}$, be a convex functions, let F be such that $F' = f$ and let Φ satisfy $\Phi' = F$. If $S_\alpha^2 f(x, y)$ is defined by*

$$S_\alpha^2 f(x, y) := \frac{(4-6\alpha)F(y) + (2-6\alpha)F(x)}{y-x} - \frac{(6-12\alpha)(\Phi(y) - \Phi(x))}{(y-x)^2}$$

then the following conditions hold true:

$$(31) \quad S_{\alpha}^2 f(x, y) \leq \alpha f(x) + (1 - \alpha)f(y),$$

if $\alpha \in [\frac{1}{3}, \frac{2}{3}]$ then

$$(32) \quad S_{\alpha}^2 f(x, y) \geq f(\alpha x + (1 - \alpha)y),$$

if $\alpha \in [0, 1] \setminus [\frac{1}{3}, \frac{2}{3}]$ then expressions $S_{\alpha}^2 f(x, y)$ and $f(\alpha x + (1 - \alpha)y)$ are incomparable in the class of convex functions,

if $\alpha \in (0, \frac{1}{3}] \cup [\frac{2}{3}, 1)$ then

$$(33) \quad S_{\alpha}^2 f(x, y) \leq S_{\alpha}^1 f(x, y)$$

and if $\alpha \in (\frac{1}{3}, \frac{1}{2}) \cup (\frac{1}{2}, \frac{2}{3})$ then $S_{\alpha}^1 f(x, y)$ and $S_{\alpha}^2 f(x, y)$ are incomparable in the class of convex functions.

Proof. Take

$$F_9(t) = (3 - 6\alpha)t^2 + (6\alpha - 2)t, \quad t \in [0, 1]$$

and let F_6, F_7, F_8 be defined so as in Proposition 2. Then we have

$$(34) \quad \begin{aligned} \int_0^1 f dF_1 &= \int_0^1 ((6 - 12\alpha)t + 6\alpha - 2)f(t) dt \\ &= F(1)(6 - 12\alpha) - \int_0^1 (6 - 12\alpha)F(t) dt + (6\alpha - 2)(F(1) - F(0)) \\ &= (4 - 6\alpha)F(1) + (2 - 6\alpha)F(0) - (6 - 12\alpha)(\Phi(1) - \Phi(0)) \end{aligned}$$

and

$$\int_0^1 F_9(t) dt = \int_0^1 F_8(t) dt = \int_0^1 F_7(t) dt = \int_0^1 F_6(t) dt.$$

It is easy to see that functions F_9, F_7 have exactly one crossing point thus we have

$$\int_0^1 f dF_9 \leq \int_0^1 f dF_7$$

which gives us (31).

Now, assume that $\alpha \in [\frac{1}{3}, \frac{2}{3}]$ then function F_9 is increasing and, consequently,

$$F_9(t) \geq F_6(t), \quad t \in [0, 1 - \alpha], \quad \text{and} \quad F_9(t) \leq F_6(t), \quad t \in (1 - \alpha, 1].$$

Thus for every convex function f we have

$$\int_0^1 f dF_6 \leq \int_0^1 f dF_9$$

which yields (32).

Let now $\alpha < \frac{1}{3}$. Then function F_9 is decreasing on some interval $[0, d]$ and increasing on $[d, 1]$. Observe that from the equality,

$$\int_0^1 F_9(t)dt = \int_0^1 F_6(t)dt$$

we know that functions F_9, F_6 must have a crossing point in the interval $(0, 1 - \alpha)$, further these functions cross also at the point $1 - \alpha$. Thus there are two crossing points of F_9, F_6 in view of Lemma 2 from [9] this means that expressions

$$\int_0^1 f dF_6, \int_0^1 f dF_9$$

are incomparable in the class of convex functions (as claimed). The reasoning in the case $\alpha > \frac{2}{3}$ is similar.

Now we shall prove the inequality (33). If $\alpha \in (0, \frac{1}{3}] \cup [\frac{2}{3}, 1)$ then $F_9'(0) \leq \frac{\alpha}{1-\alpha}$ and $F_9'(1) \leq \frac{1-\alpha}{\alpha}$ this means that functions F_9 and F_8 have only one crossing point and, therefore we have (33).

If on the other hand $\alpha \in (\frac{1}{3}, \frac{1}{2})$ then $F_9'(0) < \frac{\alpha}{1-\alpha}$ and $F_9'(1) > \frac{1-\alpha}{\alpha}$ and, consequently functions F_9, F_8 have two crossing points. Similarly as before from Lemma 2, [9] we know that for $\alpha \in (\frac{1}{3}, \frac{1}{2})$, $S_\alpha^1 f(x, y)$ and $S_\alpha^2 f(x, y)$ are incomparable in the class of convex functions, as claimed. It is easy to see that in the case $\alpha \in (\frac{1}{2}, \frac{2}{3})$ functions F_9, F_8 have again two crossing points which finishes the proof. \square

4. CONCLUDING REMARKS AND EXAMPLES

In the previous sections we made an exhaustive study of two types of inequalities. Now we briefly describe the possible extensions of our results.

Remark 8. In order to obtain inequalities involving expressions of the form $\frac{a_1\Phi(x)+a_2\Phi(\alpha x+(1-\alpha)y)+a_3\Phi(y)}{(y-x)^2}$ functions of the form

$$F_1(t) := \begin{cases} ax^2 + (1-\alpha)x & t \in [0, \alpha] \\ cx^2 + (1-c\alpha-c)x + c\alpha & t \in [\alpha, 1] \end{cases}$$

where $c = \left(-\frac{\alpha}{1-\alpha}\right)^3$ must be used. Since the description of all possible cases in Theorem 2 was already quite complicated, we shall not present these inequalities in details here.

Remark 9. It is possible to use methods developed in this paper to get inequalities involving longer expressions of the form (4). In order to

do that it is necessary to use more than two quadratic functions. For example considering function

$$(35) \quad F_1(t) := \begin{cases} 4t^2 & t \leq \frac{1}{4}, \\ -4t^2 + 4t - \frac{1}{2} & t \in \left(\frac{1}{4}, \frac{1}{2}\right] \\ 4t^2 - 4t + \frac{3}{2} & t \in \left(\frac{1}{2}, \frac{3}{4}\right] \\ -4t^2 + 8t - 3 & t > \frac{3}{4} \end{cases}$$

and using Levin-Stechkin theorem, we get the following inequality

$$\frac{8\Phi(x) - 16\Phi\left(\frac{3x+y}{4}\right) + 16\Phi\left(\frac{x+y}{2}\right) - 16\Phi\left(\frac{x+3y}{4}\right) + 8\Phi(y)}{(y-x)^2} \leq \frac{1}{y-x} \int_x^y f(t)dt$$

where $f : [x, y] \rightarrow \mathbb{R}$ is any convex function and $\Phi'' = f$.

Remark 10. We have

$$\int_0^1 t^2 dF_9(t) = \frac{5}{6} - \alpha$$

and

$$\int_0^1 t^2 dF_6(t) = (1 - \alpha)^2.$$

This means that for two values of $\alpha : \frac{3-\sqrt{3}}{6}$ and $\frac{3+\sqrt{3}}{6}$ we have $\int_0^1 t^2 dF_9(t) = \int_0^1 t^2 dF_6(t)$ Moreover, as it was mentioned in the proof of Theorem 3, functions F_9, F_6 have in this case two crossing points. This means that, from [2] (see also [10]) we get that the inequalities:

$$S_{\frac{3-\sqrt{3}}{6}}^2(x, y) \geq f\left(\frac{3-\sqrt{3}}{6}x + \frac{3+\sqrt{3}}{6}y\right)$$

and

$$S_{\frac{3+\sqrt{3}}{6}}^2(x, y) \leq f\left(\frac{3+\sqrt{3}}{6}x + \frac{3-\sqrt{3}}{6}y\right)$$

are satisfied by all 2-convex functions $f : [x, y] \rightarrow \mathbb{R}$

Remark 11. It is easy to see that all inequalities obtained in this paper in fact characterize convex functions (or 2-convex functions). This is a consequence of results contained in paper [1].

REFERENCES

- [1] M. Bessenyei, Zs. Páles, *Characterization of higher order monotonicity via integral inequalities*, Proc. Roy. Soc. Edinburgh Sect. A 140 (2010), no. 4, 723–736.
- [2] M. Denuit, C. Lefevre, M. Shaked, *The s -convex orders among real random variables, with applications*, Math. Inequal. Appl. 1 (1998), no. 4, 585–613.
- [3] S.S. Dragomir, *Two mappings in connection to Hadamard's inequalities* J. Math. Anal. Appl. **167** (1992), no. 1, 49–156.
- [4] S. S. Dragomir, I. Gomm, *Some new bounds for two mappings related to the Hermite-Hadamard inequality for convex functions* Numer. Algebra Control Optim. **2** (2012), no. 2, 271–278.
- [5] S.S. Dragomir, C.E.M. Pearce, *Selected Topics on Hermite–Hadamard Inequalities and Applications*, 2000.
- [6] V.I. Levin, S.B. Stečkin; *Inequalities*. Amer. Math. Soc. Transl. (2) 14 1960 1–29.
- [7] C. P. Niculescu, L-E. Persson, *Convex functions and their applications. A contemporary approach*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 23. Springer, New York, 2006.
- [8] J. Ohlin, *On a class of measures of dispersion with application to optimal reinsurance*, ASTIN Bulletin **5** (1969), 249–66.
- [9] A. Olbryś, T. Szostok, *Inequalities of the Hermite-Hadamard type involving numerical differentiation formulas*, Results Math., **67** (2015), 403–416.
- [10] T. Rajba, *On The Ohlin lemma for Hermite-Hadamard-Fejer type inequalities*, Math. Ineq. Appl. **17**, Number 2 (2014), 557–571.
- [11] T. Szostok *Ohlin's lemma and some inequalities of the Hermite-Hadamard type*, Aequationes Math. **89** (2015) 915–926
- [12] T. Szostok *Levin Steckin theorem and inequalities of the Hermite-Hadamard type*, submitted

INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, BANKOWA 14, 40-007
KATOWICE

E-mail address: `tszostok@math.us.edu.pl`