

# van der Waals Interactions in Hadron Resonance Gas: From Nuclear Matter to Lattice QCD

Volodymyr Vovchenko,<sup>1,2,3</sup> Mark I. Gorenstein,<sup>4,1</sup> and Horst Stoecker<sup>1,2,5</sup>

<sup>1</sup> *Frankfurt Institute for Advanced Studies, Goethe Universität Frankfurt, D-60438 Frankfurt am Main, Germany*

<sup>2</sup> *Institut für Theoretische Physik, Goethe Universität Frankfurt, D-60438 Frankfurt am Main, Germany*

<sup>3</sup> *Department of Physics, Taras Shevchenko National University of Kiev, 03022 Kiev, Ukraine*

<sup>4</sup> *Bogolyubov Institute for Theoretical Physics, 03680 Kiev, Ukraine*

<sup>5</sup> *GSI Helmholtzzentrum für Schwerionenforschung GmbH, D-64291 Darmstadt, Germany*

An extension of the ideal hadron resonance gas (HRG) model is constructed which includes the attractive and repulsive van der Waals (VDW) interactions between baryons. This VDW-HRG model yields the nuclear liquid-gas transition at low temperatures and high baryon densities. The VDW parameters  $a$  and  $b$  are fixed by the ground state properties of nuclear matter, and the temperature dependence of various thermodynamic observables at zero chemical potential are calculated within VDW-HRG model. Compared to the ideal HRG model, the inclusion of VDW interactions between baryons leads to a qualitatively different behavior of second and higher moments of fluctuations of conserved charges, in particular in the so-called crossover region  $T \sim 140 - 190$  MeV. For many observables this behavior resembles closely the results obtained from lattice QCD simulations. This hadronic model also predicts nontrivial behavior of net-baryon fluctuations in the region of phase diagram probed by heavy-ion collision experiments. These results imply that VDW interactions play a crucial role in thermodynamics of hadron gas. Thus, the commonly performed comparisons of the ideal HRG model with the lattice and heavy-ion data may lead to misconceptions and misleading conclusions.

PACS numbers: 24.10.Pa, 25.75.Gz

Keywords: hadron resonance gas, van der Waals interactions, conserved charges fluctuations

The thermodynamic properties of strongly interacting matter at zero chemical potential and finite temperature have been computed using Monte Carlo simulations in lattice QCD [1, 2]. A crossover is observed [3] in the temperature range of 140 – 190 MeV. At lower temperatures,  $T \sim 100 - 150$  MeV, QCD exhibits features similar to simple ideal hadron resonance gas (IHRG) which successfully reproduces many lattice observables [4–7]. In the crossover region, however, the agreement between IHRG and lattice QCD deteriorates. The breakdown of the IHRG model especially concerns the higher order fluctuations and correlations of conserved charges [8], resulting in statements that hadrons melt quickly and are basically absent at  $T > 160$  MeV [9]. In this Letter, it is shown that these conclusions are inconclusive. van der Waals (VDW) interactions between baryons play a crucial role for the thermodynamics of hadron fluid at sufficiently high temperatures. As a result, the qualitative features of the thermodynamics of interacting HRG appear to be close to lattice results in the crossover region. The results also have important phenomenological relevance for heavy-ion collision experiments where measurements of conserved charges fluctuations have been suggested as probes for chemical freeze-out [10, 11] or the QCD critical point [12].

The IHRG model does not capture the VDW nature of nucleon-nucleon interaction and, thus, fails to describe the properties of nuclear matter at small temperatures and large baryon densities. This shortcoming of this common HRG model is usually considered to be of minor significance when applied to ultra-relativistic heavy-ion collisions or to lattice data, although recently possible relevance for fluctuations was pointed out [13]. The repulsive part of VDW interactions had often been included into HRG by means of an excluded-volume (EV) procedure [14], usually assuming identical EV interactions between all hadron pairs [15]. The grand canonical ensemble formulation of the full VDW equation with both attractive and repulsive interactions, and including quantum statistics, was developed in Refs. [16–18] for single-component systems. In these works, the basic features of nuclear matter have been successfully described by the VDW equation with Fermi statistics for nucleons. The VDW parameters  $a$  and  $b$  were uniquely fixed by reproducing the saturation density  $n_0 = 0.16 \text{ fm}^{-3}$  and binding energy  $E/A = -16 \text{ MeV}$  of the ground state of nuclear matter. For nucleons the values  $a = 329 \text{ MeV fm}^3$  and  $b = 3.42 \text{ fm}^3$  were obtained in Ref. [18], and also later in Ref. [19]. The resulting model predicts a liquid-gas first-order phase transition in symmetric nuclear mat-

ter with a critical point located at  $T_c \simeq 19.7$  MeV and  $\mu_c \simeq 908$  MeV ( $n_c \simeq 0.07$  fm<sup>3</sup> =  $0.45 n_0$ ).

In the following a minimal extension of IHRG model, which includes the VDW interactions between (anti)baryons, is described. We refer to this model as VDW-HRG and it is based on the following assumptions:

1. VDW interactions are assumed to exist between all pairs of baryons and between all pairs of antibaryons. The VDW parameters  $a$  and  $b$  for all (anti)baryons are assumed to be equal to those of nucleons, as obtained from the fit to the ground state of nuclear matter.

2. The baryon-antibaryon, meson-meson, and meson-(anti)baryon VDW interactions are neglected.

In a sense the present VDW-HRG model is a “minimal-interaction” extension of the IHRG model, which describes the basic properties of nuclear matter. Whether significant VDW interactions exist between hadron pairs other than (anti)baryons is not clearly established. For instance, it has been argued that short-range interactions between baryons and antibaryons may be dominated by annihilation processes and not by repulsion [20], and this is our motivation to exclude VDW terms for them in this study. The presence of significant mesonic eigenvolumes, comparable to those of baryons, leads to significant suppression of thermodynamic functions in the crossover region at  $\mu_B = 0$ , which is at odds with lattice data (see Refs. [20, 21]). The attractive interactions involving mesons, on the other hand, normally lead to resonance formation [22], which are already included in HRG by construction. For these reasons we neglect the meson-related VDW interactions in this study. The VDW-HRG consists of three sub-systems: Non-interacting mesons, VDW baryons, and VDW antibaryons. The total pressure reads

$$p(T, \boldsymbol{\mu}) = P_M(T, \boldsymbol{\mu}) + P_B(T, \boldsymbol{\mu}) + P_{\bar{B}}(T, \boldsymbol{\mu}), \quad (1)$$

with

$$P_M(T, \boldsymbol{\mu}) = \sum_{j \in M} p_j^{\text{id}}(T, \mu_j) \quad (2)$$

$$P_B(T, \boldsymbol{\mu}) = \sum_{j \in B} p_j^{\text{id}}(T, \mu_j^{B*}) - a n_B^2 \quad (3)$$

$$P_{\bar{B}}(T, \boldsymbol{\mu}) = \sum_{j \in \bar{B}} p_j^{\text{id}}(T, \mu_j^{\bar{B}*}) - a n_{\bar{B}}^2, \quad (4)$$

where  $M$  stands for mesons,  $B$  for baryons, and  $\bar{B}$  for antibaryons,  $p_j^{\text{id}}$  is the Fermi or Bose ideal gas pressure,  $\boldsymbol{\mu} = (\mu_B, \mu_S, \mu_Q)$  are the chemical potentials which regulate the average values of net baryon

number  $B$ , strangeness  $S$ , electric charge  $Q$ ,  $\mu_j^{B(\bar{B})*} = \mu_j - b P_{B(\bar{B})} - a b n_{B(\bar{B})}^2 + 2 a n_{B(\bar{B})}$ , and  $n_B$  and  $n_{\bar{B}}$  are, respectively, total densities of baryons and antibaryons.

The calculation of mesonic pressure  $P_M(T, \boldsymbol{\mu})$  is straightforward. The shifted chemical potentials  $\mu_j^{B(\bar{B})*}$  of (anti)baryons depend explicitly on (anti)baryon pressure  $P_{B(\bar{B})}$  and on total (anti)baryon density  $n_{B(\bar{B})}$ . By taking the derivatives of  $P_{B(\bar{B})}$  with respect to the baryochemical potential one obtains additional equations for  $n_{B(\bar{B})}(T, \boldsymbol{\mu})$

$$n_{B(\bar{B})} = (1 - b n_{B(\bar{B})}) \sum_{j \in B(\bar{B})} n_j^{\text{id}}(T, \mu_j^{B(\bar{B})*}). \quad (5)$$

At given  $T$  and  $\boldsymbol{\mu}$ , Eqs. (3)-(5) are solved numerically, giving  $P_{B(\bar{B})}(T, \boldsymbol{\mu})$  and  $n_{B(\bar{B})}(T, \boldsymbol{\mu})$ . Entropy density is calculated as  $s = (\partial p / \partial T)_\mu$ , and energy density is obtained from Gibbs relation.

The present calculations include all established strange and non-strange hadrons which are listed in the Particle Data Tables [23], with exception of  $\sigma$  and  $\kappa$  mesons [24, 25]. The finite widths of the resonances are included by means of an additional mass integration over their Breit-Wigner shapes. We employ the HRG code used in [26, 27], modified to include the VDW interactions between (anti)baryons. The temperature dependence of the scaled pressure  $p/T^4$ , energy density  $\varepsilon/T^4$ , and the speed of sound squared  $c_s^2 = dp/d\varepsilon$  calculated at  $\boldsymbol{\mu} = 0$  within IHRG and VDW-HRG models is compared to the lattice data in Fig. 1. To clarify the role of attractive and repulsive interactions we also show the calculations, denoted as EV-HRG, where the VDW attraction was “switched off”, i.e.  $a = 0$ . Since the matter is meson dominated at  $\mu_B = 0$ , and the mesons are modeled as non-interacting, no significant suppression of thermodynamic functions is seen, in contrast to earlier studies [20, 21], where constant EV interactions between all hadrons were assumed. The energy density is somewhat below the lattice data at  $T > 160$  MeV for VDW-HRG. This may be explained by missing heavy Hagedorn states which add a significant contribution to the energy density [21]. The temperature dependence of the speed of sound squared,  $c_s^2$ , is consistent with lattice data and shows a minimum at  $T \sim 155 - 160$  MeV, in contrast to the IHRG where  $c_s^2$  decreases slowly and monotonically.

In addition to the thermodynamical functions, the VDW-HRG model allows us to calculate the fluctua-




Figure 1: The temperature dependence of (a) scaled pressure and energy density, and of (b) the square of the speed of sound at zero chemical potential, as calculated within IHRG (dashed black lines), EV-HRG with  $b = 3.42 \text{ fm}^3$  (dash-dotted red lines), and VDW-HRG with  $a = 329 \text{ MeV fm}^3$  and  $b = 3.42 \text{ fm}^3$  (solid blue lines). Lattice QCD results of Wuppertal-Budapest [1] and HotQCD [2] collaborations are shown, respectively, by symbols and green bands.

tions of conserved charges:

$$\chi_{lmn}^{BSQ} = \frac{\partial^{l+m+n} p/T^4}{\partial(\mu_B/T)^l \partial(\mu_S/T)^m \partial(\mu_Q/T)^n}. \quad (6)$$

The fluctuations of the net number of light quarks  $L = (u + d)/2 = (3B + S)/2$  are also considered.

The temperature dependencies of the 2nd order susceptibilities are shown in Fig. 2. These include (a) net number of light quarks  $\chi_2^L$ , (b) net baryon number  $\chi_2^B$ , (c) net strangeness  $\chi_2^S$ , and (d) baryon-electric charge correlator  $\chi_{11}^{BQ}$ . The  $\chi_2^L$  calculated within the VDW-HRG model shows a very different behavior compared to IHRG at  $T > 160 \text{ MeV}$ , and agrees well with the lattice data [4] up to  $T = 180 \text{ MeV}$ . A qualitatively similar picture is obtained for  $\chi_2^B$ . The qualitative difference between IHRG and VDW-HRG models appears to be driven by the EV interaction terms between (anti)baryons, while the inclusion of VDW attraction leads to an improved agreement with the lattice data. The strangeness susceptibility  $\chi_2^S$  is described fairly well by the IHRG model, but appears to be underestimated by the VDW-HRG model. We have also found that the baryon-strangeness correlator (not shown in plots) is rather notably underestimated by all considered HRG models. Does this reflect the presence of hitherto undiscovered strange hadrons? The inclusion of such states was shown to improve the agreement between lattice data and IHRG [28, 29]. The correlator  $\chi_{11}^{BQ}$  between net baryon number and net electric charge has a very different temperature dependence in IHRG and VDW-HRG. In the IHRG

model the  $\chi_{11}^{BQ}$  increases rapidly at  $T > 150 \text{ MeV}$ , in stark contrast to the lattice data. In the VDW-HRG model this correlator has a broad bump with a maximum at  $T \sim 160\text{--}190 \text{ MeV}$ , showing a behavior which is in qualitative agreement to the correlator obtained on the lattice.

The higher order fluctuations are also analyzed and exhibited in Fig. 3. All considered observables show very different behavior between IHRG and VDW-HRG. The net-light number  $\chi_4^L/\chi_2^L$  monotonically increases in the IHRG model and overshoots the lattice data at  $T \sim 140 \text{ MeV}$ . The VDW-HRG model, in contrast, yields a non-monotonic behavior with a wide peak at  $T \sim 120\text{--}145 \text{ MeV}$ , resembling the lattice data [7], which peaks at slightly higher temperature. The peak in the  $T$ -dependence of net strangeness  $\chi_4^S/\chi_2^S$  is relatively well reproduced within the VDW-HRG model. In contrast, the IHRG model shows no maximum at all. It is remarkable that our model shows flavor hierarchy: The peak for net-light number  $\chi_4/\chi_2$  is at smaller temperatures as compared to the peak in net-strangeness. The same result is seen in the lattice data! It was argued that this observation is related to the flavor separation in the deconfinement transition in QCD [7]. Since the VDW-HRG model has only hadronic degrees of freedom, the present results cast doubt on this interpretation to trace back to deconfinement the observed flavor dependence in  $\chi_4/\chi_2$ , as well as the presence of the peaks themselves. Turning to higher-order fluctuations of net-baryon number: The net-baryon kurtosis




Figure 2: The temperature dependence of the 2nd order susceptibilities of conserved charges. These include (a) net number of light quarks  $\chi_2^L$ , (b) net baryon number  $\chi_2^B$ , (c) net strangeness  $\chi_2^S$ , and (d) baryon-electric charge correlator  $\chi_{11}^{BQ}$ . Calculations are done within IHRG (dashed black lines), EV-HRG (dash-dotted red lines), and VDW-HRG (solid blue lines). Lattice QCD results of the Wuppertal-Budapest [4, 7] (for  $\chi_{11}^{BQ}$  preliminary results [30, 31] are used) and HotQCD [5] collaborations are shown, respectively, by symbols and green bands.

sis,  $\chi_4^B/\chi_2^B$ , shows the expected Skellam behavior for IHRG model with values very close to unity. The VDW-HRG model, on the other hand, shows a stark decrease at  $T = 130 - 165$  MeV, i.e. in the so-called “crossover region”, even though the VDW-HRG model does not contain any transition to the quark-gluon degrees of freedom. The  $\chi_4^B/\chi_2^B$  even turns negative at  $T > 165$  MeV. This decrease of  $\chi_4^B/\chi_2^B$  is also seen on the lattice [6], although it starts at higher  $T = 145$  MeV and  $\chi_4^B/\chi_2^B$  does not become negative. Also, the temperature dependence of the sixth order cumulant ratio  $\chi_6^B/\chi_2^B$  is predicted: The VDW-HRG model exhibits very strong variations and non-monotonous behavior in the “crossover region”. Will a similarly dramatic  $T$ -dependent behavior be observed in corresponding lattice simulations?

Finally, the kurtosis of net-baryon fluctuations at

finite baryon density is explored (Fig. 3d). For simplicity it is assumed that  $\mu_S = \mu_Q = 0$ . The region of negative  $\chi_4^B/\chi_2^B$  at small  $\mu_B$  is smoothly connected to the region of the liquid-vapor phase transition in nuclear matter, and seems relevant for “chemical freeze-out” in heavy-ion collisions (see dashed line). The VDW-HRG model suggests non-monotonic behavior of  $\chi_4/\chi_2$  with respect to collision energy, in stark contrast to IHRG [32]. This implies that non-trivial fluctuations of net-baryon number in heavy-ion collisions [33–35] may simply be a manifestation of the nuclear liquid-gas phase transition (see also [13] and [36]). Since VDW interactions also affect the thermal fits [26, 37], this question demands further studies.

For many observables, the *quantitative* agreement of the VDW-HRG model calculations presented here with the lattice data in the “crossover region” is not




Figure 3: The temperature dependence of the  $\chi_4/\chi_2$  cumulant ratio for (a) net number of light/strange quarks, (b) net baryon number, and the (c)  $\chi_6/\chi_2$  for net baryon number. Calculations are done within IHRG (dashed lines), EV-HRG (dash-dotted lines), and VDW-HRG (solid lines). The lattice QCD results of the Wuppertal-Budapest collaboration [6, 7] are shown by symbols. (d) The  $\chi_4^B/\chi_2^B$  ratio in VDW-HRG in the  $T$ - $\mu_B$  plane. The possible chemical freeze-out line in heavy-ion collisions taken from [27] is sketched by the dashed line.

perfect. This is hardly surprising. Indeed, we have modeled the VDW interactions between baryons in the simplest way possible: It is assumed that the VDW interactions between all baryons are the same as those between nucleons, as obtained from nuclear matter properties at  $T = 0$ . Conceptually, the VDW-HRG model is quite different from an underlying fundamental QCD theory. Still, the presented analysis is essentially parameter-free, in the sense that no new parameters which could be adjusted to lattice data were introduced. Indeed, the two VDW parameters had been fixed by reproducing the saturation properties of nuclear matter [17, 19], independently from any lattice data. While there are other model parameters, e.g., the hadron list and its properties, they are known and fixed experimentally. It is feasible that VDW parameters are different for different baryon

pairs. In the course of calculations, it was noticed that the agreement of the VDW-HRG with the lattice data is improved by taking smaller values of the nucleon or baryon EV parameter,  $b \simeq 2 - 3 \text{ fm}^3$ . Such modification does not necessarily break down the existing agreement of our model with the properties of nuclear matter: As suggested in Ref. [38], the heavier and/or strange baryons may have smaller eigenvolumes, thus reducing the average  $b$ . The present VDW-HRG model leaves plenty of room for improvement. Owing to the expectation that possible meson-related VDW interactions are considerably weaker than the baryon-baryon VDW interactions, we do not expect major qualitative changes to the results presented in this paper for baryon number susceptibilities. This, however, may not be the case for other observables and should be carefully explored in future works.

To summarize, a minimal extension of the IHRG model is presented which includes both attractive and repulsive VDW interactions between baryons, with parameters  $a$  and  $b$  taken from previous fits to the ground state of nuclear matter. Compared to the usually used IHRG model, the VDW-HRG model shows a qualitatively different behavior of most fluctuations and correlations of conserved charges in the “crossover region” at zero chemical potential. This behavior resembles closely the lattice QCD results. These results hint towards crucial importance of the VDW interactions in the hadron gas, and indicate that commonly performed comparisons of IHRG with the lattice data may result in misleading conclusions. Particularly, our results suggest that hadrons do not melt quickly with increasing temperature, as one could conclude on the basis of the IHRG. It is feasible that the nuclear liquid-gas phase transition manifests itself into significant non-trivial fluctuations of net-baryon number in heavy-ion collisions. The influence of VDW interactions on thermal fits to hadron yield data from heavy-ion collisions is another possibility which will be explored.

*Acknowledgments.* We are grateful to Volker Koch for stimulating discussions, and acknowledge fruitful comments from Paolo Alba, Leonid Satarov, and Jan Steinheimer. We also thank Szabolcs Borsanyi for providing the preliminary data of Wuppertal-Budapest collaboration for the baryon-electric charge correlator in tabulated format. This work was supported by HIC for FAIR within the LOEWE program of the State of Hesse. V.V. acknowledges the support from HGS-HIRe for FAIR. H.St. acknowledges the support through the Judah M. Eisenberg Laureatus Chair at Goethe University. The work of M.I.G. was supported by the Program of Fundamental Research of the Department of Physics and Astronomy of National Academy of Sciences of Ukraine.

- 
- [1] S. Borsanyi, Z. Fodor, C. Hoelbling, S. D. Katz, S. Krieg, and K. K. Szabo, Phys. Lett. B **730**, 99 (2014).
- [2] A. Bazavov *et al.* [HotQCD Collaboration], Phys. Rev. D **90**, 094503 (2014).
- [3] Y. Aoki, G. Endrodi, Z. Fodor, S. D. Katz, and K. K. Szabo, Nature **443**, 675 (2006).
- [4] S. Borsanyi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti, and K. Szabo, JHEP **1201**, 138 (2012).
- [5] A. Bazavov *et al.* [HotQCD Collaboration], Phys. Rev. D **86**, 034509 (2012).
- [6] R. Bellwied, S. Borsanyi, Z. Fodor, S. D. Katz, A. Pasztor, C. Ratti, and K. K. Szabo, Phys. Rev. D **92**, 114505 (2015).
- [7] R. Bellwied, S. Borsanyi, Z. Fodor, S. D. Katz, and C. Ratti, Phys. Rev. Lett. **111**, 202302 (2013).
- [8] A. Bazavov *et al.*, Phys. Rev. Lett. **111**, 082301 (2013).
- [9] F. Karsch, Acta Phys. Polon. Supp. **7**, 117 (2014).
- [10] S. Borsanyi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti, and K. K. Szabo, Phys. Rev. Lett. **113**, 052301 (2014).
- [11] A. Bazavov *et al.*, Phys. Rev. Lett. **109**, 192302 (2012).
- [12] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, Phys. Rev. Lett. **81**, 4816 (1998).
- [13] K. Fukushima, Phys. Rev. C **91**, 044910 (2015).
- [14] D. H. Rischke, M. I. Gorenstein, H. Stoecker, and W. Greiner, Z. Phys. C **51**, 485 (1991).
- [15] P. Braun-Munzinger, I. Heppe, and J. Stachel, Phys. Lett. B **465**, 15 (1999).
- [16] V. Vovchenko, D. V. Anchishkin, and M. I. Gorenstein, J. Phys. A **48**, 305001 (2015).
- [17] V. Vovchenko, D. V. Anchishkin, and M. I. Gorenstein, Phys. Rev. C **91**, 064314 (2015).
- [18] V. Vovchenko, D. V. Anchishkin, M. I. Gorenstein, and R. V. Poberezhnyuk, Phys. Rev. C **92**, 054901 (2015).
- [19] K. Redlich and K. Zalewski, Acta Phys. Polon. B **47**, 1943 (2016), [arXiv:1605.09686 [cond-mat.quant-gas]].
- [20] A. Andronic, P. Braun-Munzinger, J. Stachel, and M. Winn, Phys. Lett. B **718**, 80 (2012).
- [21] V. Vovchenko, D. V. Anchishkin, and M. I. Gorenstein, Phys. Rev. C **91**, 024905 (2015).
- [22] R. Venugopalan and M. Prakash, Nucl. Phys. A **546**, 718 (1992).
- [23] K. A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38**, 090001 (2014).
- [24] W. Broniowski, F. Giacosa, and V. Begun, Phys. Rev. C **92**, 034905 (2015).
- [25] B. Friman, P. M. Lo, M. Marczenko, K. Redlich, and C. Sasaki, Phys. Rev. D **92**, 074003 (2015).
- [26] V. Vovchenko and H. Stoecker, J. Phys. G **44**, 055103 (2017).
- [27] V. Vovchenko, V. V. Begun, and M. I. Gorenstein, Phys. Rev. C **93**, 064906 (2016).
- [28] P. M. Lo, M. Marczenko, K. Redlich, and C. Sasaki, Phys. Rev. C **92**, 055206 (2015).
- [29] A. Bazavov *et al.*, Phys. Rev. Lett. **113**, 072001 (2014).
- [30] S. Borsányi, PoS LATTICE **2015**, 015 (2016) [arXiv:1511.06541 [hep-lat]].
- [31] Wuppertal-Budapest collaboration, in preparation.
- [32] P. Alba, W. Alberico, R. Bellwied, M. Bluhm, V. Mantovani Sarti, M. Nahrgang, and C. Ratti, Phys. Lett. B **738**, 305 (2014).
- [33] M. M. Aggarwal *et al.* [STAR Collaboration], Phys. Rev. Lett. **105**, 022302 (2010).
- [34] L. Adamczyk *et al.* [STAR Collaboration], Phys. Rev. Lett. **112**, 032302 (2014).

- [35] X. Luo [STAR Collaboration], PoS CPOD **2014**, 019 (2015) [arXiv:1503.02558 [nucl-ex]].
- [36] A. Mukherjee, J. Steinheimer, and S. Schramm, arXiv:1611.10144 [nucl-th].
- [37] L. M. Satarov, V. Vovchenko, P. Alba, M. I. Gorenstein, and H. Stoecker, Phys. Rev. C **95**, 024902 (2017).
- [38] P. Alba, V. Vovchenko, M. I. Gorenstein, and H. Stoecker, arXiv:1606.06542 [hep-ph].