

The Symmetricity of Normal Modes in Symmetric Complexes

Guang Song^{1,2,3*}

¹Department of Computer Science,

²Program of Bioinformatics and Computational Biology,

³L. H. Baker Center for Bioinformatics and Biological Statistics,

Iowa State University, Ames, IA 50011, USA

*Correspondence to: Guang Song, Tel: 515-294-1696;

Fax: 515-294-0258; E-mail: gsong@iastate.edu

July 9, 2018

Abstract

In this work, we look at the symmetry of normal modes in symmetric structures, particularly structures with cyclic symmetry. We show that normal modes of symmetric structures have different levels of symmetry, or symmetricity. One novel theoretical result of this work is that, for a ring structure with m subunits, the symmetricity of the normal modes falls into m groups of equal size, with normal modes in each group having the same symmetricity. The normal modes in each group can be computed separately, using a much smaller amount of memory and time (up to m^3 less), thus

making it applicable to larger complexes. We show that normal modes with perfect symmetry or anti-symmetry have no degeneracy while the rest of the modes have a degeneracy of two. We show also how symmetry in normal modes correlates with symmetry in structure. While a broken symmetry in structure generally leads to a loss of symmetricity in symmetric normal modes, the symmetricity of some symmetric normal modes is preserved even when symmetry in structure is broken. This work suggests a deeper reason for the existence of symmetric complexes: that they may be formed not only for structural purpose, but likely also for a dynamical reason, that certain symmetry is needed by a given complex to obtain certain symmetric motions that are functionally critical.

1 Introduction

Many proteins in cell need to form structure complexes in order to function. In E-coli, it was estimated that over 80% of proteins form structural complexes¹. Most of the structures reported in PDB form complexes too. According to a recent work by Levy et al.², about 1/2 to 2/3 of the proteins in PDB are multi-meric, i.e., containing multiple chains. Such multi-unit complexes are called oligomers or multimers. They represent the quaternary structures of proteins. An oligomer is composed of more than one peptide chains, each of which is called a subunit of the complex. Some of these complexes are composed of monomers of different types, which are called heteroligomers, or heteromers, while others are composed of identical subunits and are called homoligomers or homomers.

How proteins form quaternary complexes is an important research topic. There are a number of work that predict the quaternary structures of proteins, such as PQS³, PISA⁴, 3D-complex², PiQSi⁵, etc. Some of these are fully-automatic^{3,4}, while others are manually curated⁵. 3D-complex² provides a novel hierarchical classification of the organization of protein complexes and it uses biological assemblies of PDB as input. In their recent

study⁶, Teichmann and her co-workers delved deeper into the organizing principle of protein complexes and found that most assembly steps fall into three basic types: “dimerization, cyclization, and heteromeric subunit addition”⁶. According to this principle, protein complexes can be organized neatly into a “periodic table”,⁶ which not only provides fresh insights into the patterns and formation principle of protein complexes but also may be used to predict complexes that are not yet observed structurally.

Homomers are more abundant than heteromers. One known fact about homomers is that more of them have an even number of subunits than odd.² Levy et al.⁷ shows that most homomers have either cyclic or dihedral symmetry, while other kinds of symmetry are rare. For cyclic structures, Kidera and coworkers⁸ found that there is no preference for an even or odd number of subunits, and therefore the reason behind the preference for an even number of subunits is that complexes with dihedral symmetry always require an even number of subunits⁸. In summary, most proteins form structure complexes and a majority of these complexes are homomers and symmetric. *symmetric structures are prevalent*. Complexes with cyclic symmetry are commonly called ring structures. In the work, we focus on the normal modes of symmetric ring structures. The study can be extended to other symmetric structures and perhaps even pseudo-symmetric structures¹, of which the subunits are not identical but the overall structure is almost symmetric.

Normal modes analysis⁹⁻¹¹, as a powerful tool for studying protein vibrational dynamics near equilibrium state, has been extensively studied and applied for the last two decades using coarse-grained models¹²⁻³². Dynamics produced from NMA has provided insightful understanding of the functional mechanisms of a wide range of proteins. Deeper understandings regarding the properties of normal modes themselves also have been obtained, such as the property of low frequency modes, high-frequency modes, hot-spot residues³³, dynamics-residues²⁴, the universality of the vibrational spectrum^{34,35}, the effective degeneracy of normal modes³⁶, etc. These insightful understandings of normal modes are helpful

for a more accurate and appropriate use of them in various applications.

The topic of this study is about another property of normal modes, namely, the symmetry of the normal modes in symmetric structures. Since the structures under consideration are symmetric, There are a number of interesting questions that can be raised regarding their normal modes:

- One is about **efficiency**. Since the complex is symmetric, is there a way to take advantage of the symmetry so as to obtain the normal modes more efficiently?
- Another is about **degeneracy**. Are all the modes degenerate due to the symmetry in structure? What is the level of degeneracy?
- **Dynamics**. Since the structure is symmetric, are the modes/motions also symmetric? How to measure the level of a mode's symmetry?
- Lastly, about **function**. Are the functional processes symmetric? If so, how are they related to the intrinsic symmetric modes of symmetric ring structures?

In their seminal work, Simonson and Perahia³⁷ addressed the first issue regarding efficiency and showed that by using group theory the time for computing normal modes of symmetric structures can be greatly reduced. Using a similar approach, Vlijmen and Karplus showed that efficient normal mode analysis could be carried out for large systems with icosahedral symmetry³⁸ and successfully applied it to Icosahedral viruses³⁹. In a recent work, Matsunaga et al.⁸ demonstrated that structure symmetry has a determinant effect on the protein dynamics of circularly symmetric structure. They observed that multimers with a highly composite number of subunits (such as 6 or 12) tend to have more inter-unit fluctuations while multimers with a prime number of subunits tend to have more intra-unit fluctuations.

Our work here focuses on the *degeneracy* and *dynamics* of symmetric complexes. Our previous work showed that the functional motions of some symmetric structures are symmetric and are closely related to symmetric modes⁴⁰.

2 Methods

For symmetric ring structures of m subunits of size N (i.e., N atoms or residues in each subunit), the use of symmetry can reduce the size of the matrix that has to be diagonalized from $3mN \times 3mN$ to a series of smaller matrices of $3N \times 3N$. The most general, rigorous way to take advantage of symmetry is to use group theory^{8,37–39,41}, which utilizes the character table and irreducible representations to find symmetric coordinates^{8,37–39,41}. When expressed in symmetric coordinates, the Hessian matrix becomes block diagonal. For the special case of symmetric ring structures, the process can be understood using a mathematically simpler approach that is based on circulant matrix.

2.1 Circulant Matrix

A circular matrix has the following form^{42,43}:

$$\begin{bmatrix} c_0 & c_1 & \dots & c_{m-2} & c_{m-1} \\ c_{m-1} & c_0 & \dots & c_{m-3} & c_{m-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ c_2 & c_3 & \dots & c_0 & c_1 \\ c_1 & c_2 & \dots & c_{m-1} & c_0 \end{bmatrix} \quad (1)$$

The normalized eigenvectors of a circulant matrix are given by

$$v_j = \frac{1}{\sqrt{m}}(1, \omega_j, \omega_j^2, \dots, \omega_j^{m-1})^T, \quad j = 0, 1, \dots, m-1, \quad (2)$$

where $\omega_j = \exp\left(\frac{2\pi ij}{m}\right)$ are the m^{th} roots of unity and i is the imaginary unit.

The corresponding eigenvalues are:

$$\lambda_j = c_0 + c_1\omega_j + c_2\omega_j^2 + \dots + c_{m-1}\omega_j^{m-1}, \quad j = 0, 1, \dots, m-1. \quad (3)$$

2.2 Small Oscillations of M Identical Masses in a Hoop

For a spring-mass system of m identical masses connected with m identical springs in a hoop, the Kirchhoff matrix is a circulant matrix where $c_0 = 2$, $c_{m-1} = c_1 = -1$, and the rest of c 's are all zeros. Therefore, the eigenvalues are:

$$\lambda_j = c_0 + c_1\omega_j + c_{m-1}\omega_j^{m-1} = 2 - 2\cos\left(\frac{2\pi j}{m}\right), \quad j = 0, 1, \dots, m-1. \quad (4)$$

Except for λ_0 (which is 0) and $\lambda_{\frac{m}{2}}$ when m is even, all the modes are degenerate and have a degeneracy of 2.

2.3 Symmetric Circulant Block Matrix

A symmetric circulant block matrix has the following form:

$$H = \begin{bmatrix} A_0 & A_1 & \dots & A_{m-2} & A_{m-1} \\ A_{m-1} & A_0 & \dots & A_{m-3} & A_{m-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ A_2 & A_3 & \dots & A_0 & A_1 \\ A_1 & A_2 & \dots & A_{m-1} & A_0 \end{bmatrix} \quad (5)$$

and $A_k = A'_{m-k}$ since the matrix is symmetric. In the discussion that follows, we limit ourselves to the scenario that H is also real, as is for a Hessian matrix.

An efficient way to obtain H 's eigenvalues and eigenvectors exists and was given by Cao⁴⁴.

Part of his proof is included here for clarity. In his proof, Cao constructed a series of M matrices:

$$\tilde{M}_j = \sum_{k=0}^{m-1} \omega_m^{kj} A_k \quad (6)$$

where ω_m is equal to $e^{\frac{2\pi i}{m}}$.

Specifically,

$$\tilde{M}_0 = \sum_{k=0}^{m-1} A_k, \quad (7)$$

and when m is even,

$$\tilde{M}_{m/2} = \sum_{k=0}^{m-1} (-1)^k A_k. \quad (8)$$

According to Cao⁴⁴, define a matrix F_m as:

$$F_m = \frac{1}{\sqrt{m}} \begin{bmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & \omega_m & \omega_m^2 & \dots & \omega_m^{m-1} \\ 1 & \omega_m^2 & \omega_m^4 & \dots & \omega_m^{2(m-1)} \\ \dots & \dots & \dots & \dots & \dots \\ 1 & \omega_m^{m-1} & \omega_m^{2(m-1)} & \dots & \omega_m^{(m-1)^2} \end{bmatrix} \quad (9)$$

Assume \mathbf{u}_j is one of eigenvectors of \tilde{M}_j , then $\mathbf{w} = (0_n, 0_n, \dots, \mathbf{u}_j, \dots, 0_n)'$ will be the eigenvectors of $\text{diag}(\tilde{M}_0, \tilde{M}_1, \dots, \tilde{M}_j, \dots, \tilde{M}_{m-1})$, a diagonal block matrix, where 0_n represents a row vector with n zeros and $n = 3N$. The corresponding eigenvector \mathbf{v} of the original

matrix H in Eq. (5) can be obtained by⁴⁴,

$$\mathbf{v} = (F_m \otimes I_n) \mathbf{w} = \begin{bmatrix} \mathbf{u}_j \\ \mathbf{u}_j \omega_m^j \\ \vdots \\ \mathbf{u}_j \omega_m^{(m-1)j} \end{bmatrix} \quad (10)$$

where I_n is an identity matrix of dimension n . Now let \mathbf{u}_j be,

$$\mathbf{u}_j = \begin{bmatrix} r_1 e^{i\theta_1} \\ r_2 e^{i\theta_2} \\ \vdots \\ r_n e^{i\theta_n} \end{bmatrix} \quad (11)$$

Then,

$$\mathbf{v} = \begin{bmatrix} r_1 e^{i\theta_1} \\ \vdots \\ r_n e^{i\theta_n} \\ r_1 e^{i(\theta_1 + \phi)} \\ \vdots \\ r_n e^{i(\theta_n + \phi)} \\ \vdots \\ r_1 e^{i(\theta_1 + (m-1)\phi)} \\ \vdots \\ r_n e^{i(\theta_n + (m-1)\phi)} \end{bmatrix} = \begin{bmatrix} r_1 \cos(\theta_1) \\ \vdots \\ r_n \cos(\theta_n) \\ r_1 \cos(\theta_1 + \phi) \\ \vdots \\ r_n \cos(\theta_n + \phi) \\ \vdots \\ r_1 \cos(\theta_1 + (m-1)\phi) \\ \vdots \\ r_n \cos(\theta_n + (m-1)\phi) \end{bmatrix} + i \begin{bmatrix} r_1 \sin(\theta_1) \\ \vdots \\ r_n \sin(\theta_n) \\ r_1 \sin(\theta_1 + \phi) \\ \vdots \\ r_n \sin(\theta_n + \phi) \\ \vdots \\ r_1 \sin(\theta_1 + (m-1)\phi) \\ \vdots \\ r_n \sin(\theta_n + (m-1)\phi) \end{bmatrix} = \mathbf{p} + i\mathbf{q}, \quad (12)$$

where $\phi = 2\pi j/m$.

Since H is a real symmetric matrix, it means that both \mathbf{p} and \mathbf{q} are the eigenvectors of H , having the same eigenvalue. Next, we will show that \mathbf{p} and \mathbf{q} are orthogonal to each other, i.e., $\mathbf{p} \cdot \mathbf{q} = 0$.

Consider \mathbf{p}_u and \mathbf{p}_d that are obtained by rotating \mathbf{p} upward or downward by n elements, respectively:

$$\mathbf{p}_u = \begin{bmatrix} r_1 \cos(\theta_1 + \phi) \\ \vdots \\ r_n \cos(\theta_n + \phi) \\ \vdots \\ r_1 \cos(\theta_1 + (m-1)\phi) \\ \vdots \\ r_n \cos(\theta_n + (m-1)\phi) \\ r_1 \cos(\theta_1) \\ \vdots \\ r_n \cos(\theta_n) \end{bmatrix}, \mathbf{p}_d = \begin{bmatrix} r_1 \cos(\theta_1 + (m-1)\phi) \\ \vdots \\ r_n \cos(\theta_n + (m-1)\phi) \\ r_1 \cos(\theta_1) \\ \vdots \\ r_n \cos(\theta_n) \\ \vdots \\ r_1 \cos(\theta_1 + (m-2)\phi) \\ \vdots \\ r_n \cos(\theta_n + (m-2)\phi) \end{bmatrix} \quad (13)$$

Now it is evident that $\mathbf{p}_u = \mathbf{p} \cos(\phi) - \mathbf{q} \sin(\phi)$. $\mathbf{p}_d = \mathbf{p} \cos(\phi) + \mathbf{q} \sin(\phi)$. Since $\mathbf{p}_u \cdot \mathbf{p}_u = \mathbf{p}_d \cdot \mathbf{p}_d$, we have, $\mathbf{p} \cdot \mathbf{q} = 0$.

Since \mathbf{p} and \mathbf{q} are both eigenvectors of H and share the same eigenvalue, the normal modes that these eigenvectors represent have a degeneracy of 2, which is consistent with small oscillations of m identical masses on a hoop (see section 2.2). It is worth noting that this kind of degeneracy originates purely from symmetry in structure and is different from the kind of degeneracy of normal modes caused by structure uncertainty³⁶.

The degeneracy of 2 in normal modes can be understood also in an alternative way.

Notice that,

$$\tilde{M}_{m-j} = \sum_{k=0}^{m-1} \omega_m^{k(m-j)} A_k = \sum_{k=0}^{m-1} \bar{\omega}_m^{kj} A_k = \overline{\tilde{M}_j} \quad (14)$$

That is, M_{m-j} is a complex conjugate of M_j . It is clear both \tilde{M}_j and \tilde{M}_{m-j} are Hermitian matrices and have real eigenvalues. Furthermore, since they are complex conjugate to each other, \tilde{M}_j and \tilde{M}_{m-j} have the same set of eigenvalues. Therefore, the normal modes computed from \tilde{M}_j and \tilde{M}_{m-j} using eq. (10) have a degeneracy of 2 in general. The only exception to this is that \tilde{M}_0 (eq. (7)), or $\tilde{M}_{m/2}$ (eq. (8)) when m is even, is a real matrix. The modes computed from them have no degeneracy. Later, we will show that the modes computed from \tilde{M}_0 are perfectly symmetric modes, while modes computed from $\tilde{M}_{m/2}$ are perfectly anti-symmetric modes.

2.4 Representing Hessian Matrix as a Circulant Block Matrix

For a symmetric ring structure with m subunits, it has a m -fold cyclic symmetry (C_m). Let R represent a rotation of $2\pi/m$ degree. Representing the motions of each subunit in its local frame, the Hessian matrix becomes a symmetric circulant block matrix:

$$H = \begin{bmatrix} A_0 & A_1 & \dots & A_{m-2} & A_{m-1} \\ A_{m-1} & A_0 & \dots & A_{m-3} & A_{m-2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ A_2 & A_3 & \dots & A_0 & A_1 \\ A_1 & A_2 & \dots & A_{m-1} & A_0 \end{bmatrix} \quad (15)$$

where A_i represents the interaction between the current subunit and the i_{th} subunit down the ring (clockwise or counter-clockwise).

The eigenvectors of H are given in Eq. (10). The eigenvectors in the *global* coordinate

are:

$$\mathbf{v} = \begin{bmatrix} \mathbf{u}_j \\ (I_n \otimes R)\mathbf{u}_j\omega_m^j \\ \vdots \\ (I_n \otimes R^{n-1})\mathbf{u}_j\omega_m^{(m-1)j} \end{bmatrix} \quad (16)$$

2.5 The Symmetricity of Normal Modes

For ring structures with m identical subunits, we define the symmetricity of its modes as follows⁴⁰. For each mode \mathbf{m}_i , we perform a rotation of $2\pi/m$ along its central axis. Let \mathbf{m}'_i be the mode after the rotation. The symmetricity of mode i is defined as the overlap (or dot product) between \mathbf{m}_i and \mathbf{m}'_i , i.e.,

$$\text{symmetricity} = \mathbf{m}_i \cdot \mathbf{m}'_i \quad (17)$$

If a mode is perfectly symmetric along the central axis, the rotation has no effect and its symmetricity should be 1. That is, the mode is invariant under rotations of multiples of $2\pi/m$.

If we represent the motion of each subunit in its local frame, the effect of the rotation is the same as rotating the elements of \mathbf{m}_i by one subunit block (n elements). If we let \mathbf{m}_i be \mathbf{p} (see Eq. (12)), as \mathbf{p} indeed is an eigenvector of H , then \mathbf{m}'_i is the same as \mathbf{p}_u in Eq. (13). Consequently, since $\mathbf{p} \cdot \mathbf{p} = 1$ and $\mathbf{p} \cdot \mathbf{q} = 0$, we have,

$$\text{symmetricity} = \mathbf{m}_i \cdot \mathbf{m}'_i = \mathbf{p} \cdot \mathbf{p}_u = \cos(\phi) = \cos(2\pi j/m), \quad (18)$$

where $j = 0, 1, \dots, n-1$. This means for symmetric ring structures with m units, $1/m$ of the modes have symmetricity of $\cos(0) = 1$, $1/m$ of the modes have symmetricity of $\cos(2\pi/m)$, $1/m$ of the modes have symmetricity of $\cos(4\pi/m)$, and so on.

2.6 Construct Symmetric Complexes

The biological assembly reported in PDB⁴⁵ for symmetric ring structures generally are not exactly perfectly symmetric along the central axis. There is usually a small amount of deviation from the otherwise perfectly symmetric structure.

To construct a perfectly symmetric structure model is simple. One can pick one subunit from a given ring structure and perform on it a series of m rotations (clockwise or counter-clockwise) of $2\pi/m$ degrees along the central axis (where m is the number of subunits) until it comes back to its original conformation and collect all the intermediate conformations, which, together with the subunit's initial conformation, form a conformation of a symmetric ring structure. We term such a perfectly symmetric structure as *perfect structure* in the rest of the paper, as contrast to actual PDB structures that may or may not be perfectly symmetric.

Definition 1: Perfect Structure. A perfect structure is a ring structure, constructed or actual, that has perfect axial symmetry.

A perfect structure of p97. p97⁴⁶ is an important protein in the extended AAA (ATPases Associated with diverse cellular Activities) family. p97 is a symmetric hexamer and there are about a dozen of p97 structures deposited in PDB⁴⁰. One of them is 5ftl⁴⁷, a cryo-EM structure of p97 bound with ATP analogs. 5ftl is nearly exactly symmetric. We apply the above procedure and construct a perfect structure of p97 using chain A of 5ftl and name the new structure model *5ftl-perfect*. The root mean square distance between 5ftl and 5ftl-perfect is 0.0042 Å. Both 5ftl and 5ftl-perfect are used in our study

3 Results

The symmetricity of normal modes in ring structures describes the extent of synchronization in motion between adjacent subunits along the ring. When applied to the whole ring

structure, it can reveal the extent of cooperativity²⁸ among the subunits. In the following, we will take a close investigation of symmetricity in ring structures and its implications.

3.1 Efficient Computations of Normal Modes of Ring Structures

The circulant block Hessian matrix allows us to efficiently compute normal modes of ring structures. Particularly, Eqs. (5) to (12) show how the task of solving for the normal modes of the whole multi-mer can be divided by solving for the eigenvalues of a few matrices of a smaller size: m times smaller to be precise, where m is the number of subunits in the complex. Since solving for eigenvalues may take up to a cubic time to the size of the matrix, the computational gain can be enormous. This allows normal modes of ring structures be computed in a much smaller amount of time, using a much smaller amount of memory, thus making it possible to extend the normal mode computations to larger ring structures.

Moreover, in so doing, the normal modes are naturally grouped by their level of symmetricity. That is, the normal modes within each group have the same symmetricity (see the Methods section on symmetricity).

This natural grouping can be highly beneficial when only modes with certain symmetricity is needed. For example, if all we care about are symmetric modes (i.e., symmetricity = 1), we only need to solve one M matrix (see Methods)! The computation time will be further reduced. Indeed, for symmetric ring structures, it is likely that only symmetric or anti-symmetric (symmetricity=-1) are functionally important. For example, a previous study on p97⁴⁰ indicated that only symmetric modes contribute to the conformation changes that also are symmetric.

3.2 Perfectly Symmetric and Anti-Symmetric Modes

For symmetric ring structures with m subunits, it is evident that $1/m$ of the modes are symmetric. For those ring structures with an even number of subunits, an additional $1/m$ of the modes are anti-symmetric. The anti-symmetric modes can be computed by solving Eq. (8). Structures with an odd number of subunits do not have anti-symmetric modes. Anti-symmetric modes represent motions where half the subunits (every other subunit around the ring) synchronize perfectly and move in the opposite direction to the other half of subunits. This pattern of motions may be important for the functions of some complexes whose two subsets of subunits alternate their roles in function, such as ATP binding and hydrolysis.

It is helpful to realize that both symmetric and anti-symmetric of an m -subunit structure are symmetric modes of the same structure if structure is considered as an $m/2$ -dimer that has $m/2$ -fold axial symmetry. For example, the heat shock locus protein (HslU), which consists of 6 subunits, was found to behave as a hexamer when all six units are bound with ADP (or ATP)^{48,49}, and as a trimer of dimer when only every other subunit binds with an ADP^{49,50}.

3.3 Symmetric Modes Form a Closed Subspace

Symmetric modes can be obtained by solving Eq. 7. It is evident that symmetric modes amount to $1/m$ of the total number of modes. All the symmetric modes take the form of:

$$\mathbf{S}_j = \begin{bmatrix} \mathbf{u}_j \\ \mathbf{u}_j \\ \vdots \\ \mathbf{u}_j \end{bmatrix} \quad (19)$$

where \mathbf{u}_j is an eigenvector of \tilde{M}_0 in Eq. (7) and $j = 0, 1, \dots, 3N$, where N is number of atoms in each subunit. Therefore, it is obvious that all the symmetric modes form a closed subspace in the sense that any symmetric conformation displacement can be written as a linear combination of S_j 's. That is, for a symmetric conformational displacement \mathbf{ds} ,

$$\mathbf{ds} = \begin{bmatrix} \mathbf{d} \\ \mathbf{d} \\ \vdots \\ \mathbf{d} \end{bmatrix} \quad (20)$$

we can always have:

$$\mathbf{ds} = \sum_{i=1}^{3N} c_i \mathbf{S}_i \quad (21)$$

This is evident since the displacement of each unit, \mathbf{d} , in general can always be written as a linear combination of \mathbf{u}_j 's.

$$\mathbf{d} = \sum_{i=1}^{3N} c_i \mathbf{u}_j. \quad (22)$$

There are a couple of important implications. First, if we are interested only in symmetric conformation displacements or conformation transitions, only symmetric modes are needed and they alone provide complete information regarding how the transition may take place. A second implication is that non-symmetric modes cannot be linearly combined to give a symmetric displacement.

As we will see later, for ring structures that are not completely symmetric, symmetricity is partially broken due to degeneracy³⁶ and the number of modes with perfect symmetry (i.e., symmetricity=1) is reduced and their fraction is less than $1/m$ and consequently they no longer form a closed subspace. However, symmetric modes that are functional are generally of low frequency and are robust to degeneracy and can remain to be symmetric. These symmetric modes, though no longer forming a closed subspace, may still be sufficient to

interpret any symmetric function-related conformation changes⁴⁰.

3.4 The Symmetricity of Normal Modes

The level of symmetricity reveals the extent to which adjacent subunits along the ring are moving together, or the extent of their similarity. The theoretical prediction on symmetricity as given in Eq. 18 states that the whole set of modes can be divided into m groups of the same size and the modes in each group have the same symmetricity. The symmetricity for group j ($0 \leq j \leq m - 1$) is $\cos(\frac{2\pi j}{m})$. The angle $\frac{2\pi j}{m}$ represents the amount of phase shift between adjacent subunits.

Figure 1(A) shows the symmetricity distribution of the normal modes of p97, computed from a perfectly symmetric model of p97: 5ftl-perfect, which is constructed from PDB⁴⁵ structure 5ftl (see Methods). The computed distribution matches perfectly with the theoretical prediction: 1/6 of the modes have symmetricity of 1 or -1, and 1/3 of the modes have symmetricity of 0.5 (which equals to $\cos(\frac{2\pi}{6})$ or $\cos(\frac{5*2\pi}{6})$) or -0.5 (which equals to $\cos(\frac{2*2\pi}{6})$ or $\cos(\frac{4*2\pi}{6})$).

Now consider a unit polygon with m sides as shown in Figure 1(B), with vertices starting at (1, 0) and proceeding counterclockwise at angles in multiples of $2\pi/m$, the symmetricity of each group of normal modes mentioned above is the same as the x-coordinate values of the vertices of the polygon. Groups i and $m - i$ clearly have the same symmetricity. When n is even, there is a group with symmetricity of -1, or anti-symmetric.

Symmetric modes distribute through the whole frequency range. Since Symmetric modes are obtained by solving the Hessian matrix in Eq. 7, it is perceivable that the symmetric modes should distribute through the whole frequency range. Figure 2 shows the density of state distributions of modes with different symmetricity. It is seen that modes with different symmetricity spread evenly and have nearly the same distribution (ignoring the scaling factor).

3.5 Symmetricity Pattern is Partially Broken in Real Structures

Crystal structures reported in PDB contains a set of coordinates solved from the structure factors. These coordinates represent structural data in the asymmetric units. They may not represent the complete complex or may contain multiple copies of the same molecule. To construct the complete biological assembly of a molecule based on the given coordinate data from the asymmetric units (as well as information on space group and unit cell), software such as PISA⁴ or PQS³ is often used. There are cases where the asymmetric unit contains the whole multimer, such as the hexamer structure of HslU (PDB-id: 1DO2⁴⁹) or p97 (PDB-id:5C18⁵¹).

The recent advancement of cryo-EM technology makes possible the determination of many large structure complexes at near-atomic resolution⁵². Cryo-EM usually assumes structure symmetry (for complexes that are symmetric) and is able to produce a whole structure assembly.

For most of the homomer structures reported in PDB, the subunits are not perfectly symmetric. There exist some slight structure deviations from the otherwise perfectly symmetric structures. The effective degeneracy of normal modes³⁶ dictates that modes are degenerate under slight structure variations and may mix together with other modes with similar frequencies. Consequently, not all the modes computed from a perfectly symmetric structure will maintain their symmetricity in reality.

Figure 3 shows the symmetricity plot of p97 (pdb-id: 5ftl) as computed by ANM¹⁶. The result is the same as that in Figure 1(A) except that the original PDB structure of 5ftl is used here. This cryo-EM structure of p97 (pdb-id: 5ftl) as reported in PDB is nearly perfectly axially symmetric. The root mean square deviation between this structure and the one constructed above by selecting chain *A* of 5ftl and rotating multiples of 60 degrees is only 0.0042 Å.

From the figure it is seen that, comparing to the 2,000+ symmetric modes (i.e., sym-

metricity=1) of the perfectly symmetric structure in Figure 1(A), there are only about 200 modes in the actual structure that have a symmetricity of nearly 1 (≥ 0.98).

What are these 200 modes? Are they just a subset of the 2000+ symmetric modes of the perfectly symmetric structure? To answer this question, we find, for each mode of the perfect structure, the best matching mode in the modes of the actual structure. Figure 4 gives a scatter plot of the symmetricity of the modes of the perfect structure and the symmetricity of their best matching modes in the actual structure. The figure shows that the modes with high symmetricity (≥ 0.8) of the actual structure all match to modes with symmetricity of 1 in the perfect structure. The symmetric modes (symmetricity=1) of the perfect structure match to the modes of the actual structure with a wide-spread range of symmetricity. Of all the symmetric modes of the perfect structure, only a small percentage of them are able to preserve their symmetricity.

The next question is, what are these modes that are able to preserve their perfect symmetricity of 1 and why? What happens to the other symmetric modes? Figure 5 shows the frequency distribution of the modes that are able to preserve their perfect symmetricity of 1. Most of these modes fall into either the low frequency end (low mode indices) or the high frequency end (high mode indices).

To find out why the modes at either the low frequency end or the high frequency end can preserve their symmetricity and what happens to the other symmetric modes, we plot in Figure 6, for the 2,167 symmetric modes of the perfect structure, the symmetricity of their besting matching modes in the actual structure and the overlaps between them and their best matching modes. Not surprisingly, a strong correlation is found: the modes that are mostly preserved (or unchanged) under structure variation, as indicted by a large overlap, also preserve most of their symmetricity. Thus, to the question raised earlier, “Why are the modes at the low or high frequency end able to preserve their symmetricity?” the answer is that these modes are robust to small structural changes and remain mostly unchanged

(with a large overlap). This is thus consistent with our previous finding that modes at the low or high frequency end are less degenerate³⁶. On the other hand, the low overlaps of the other modes indicate that they have deformed greatly under the structure deviation. Figure 6 shows that non-degenerate modes are the ones that can maintain their symmetry, while degenerate modes generally cannot.

It is perceivable that for some systems symmetric modes are critical to the realization of functions. The ability to preserve their symmetry is consequently important.

3.6 Obtain Symmetric and/or Anti-Symmetric Modes from Nearly-Symmetric Structures

Practically speaking, most ring structures deposited in PDB do not have exact symmetry. The coordinates of the atoms in each subunit may be slightly off from their otherwise perfectly symmetric locations. Moreover, we do not expect symmetric ring structures such as p97 to maintain a mathematically exact symmetry while they function in cell. For such structure models that have nearly exact symmetry, how do we obtain symmetric modes efficiently? Apparently, we cannot apply Eq. (7) or (8) if the structure is not exactly symmetric.

One possible solution is to choose not to take advantage of the symmetry and compute the normal modes using the whole structure. The drawback is that this can become computationally too costly, especially for large systems. Another possible solution is to reconstruct a perfectly symmetric structure from one of the subunits by applying axial symmetric rotations and then apply Eq. (7) to compute modes. The drawback of this approach is that the modes computed by the reconstructed structure (which has a perfect symmetry) are generally somewhat different from those computed with the original (PDB) structure. Such difference reflects the degeneracy of protein normal modes³⁶.

A more ideal solution is to be able to obtain symmetric modes that are robust to the

structure differences among the subunits. Is there an efficient way to obtain the subset of symmetric modes that are robust to structure variations without solving the Hessian matrix of the whole multi-mer? To answer this question, it is helpful to realize that the subunits are highly similar to one another in structure and they only deviate slightly from the otherwise perfectly symmetric structure. In our previous work, we have shown that some normal modes, especially those at low frequency end, are robust to small structure deviations while others are not and become degenerate³⁶. For these non-degenerate modes, the slight deviation from the perfect symmetry does not disturb them.

For this reason, we determine symmetric modes that are robust to small structure deviations in the following way. First, we use each subunit in turn as the center and use Eq. (7) compute the symmetric modes. We then find the symmetric modes that are common to all subunits (modes are considered common or the same if the overlaps between them are greater than 0.99). These modes are robust to structure differences among the subunits.

Figure 7 shows the index/frequency distribution of the symmetric modes that are common to all subunits. The modes computed from each subunit are compared with the 2,167 symmetric modes of a perfectly symmetric structure of p97: 5ftl-perfect. Modes that are common to all subunits are then shown in Figure 7 as a histogram. The figure shows that the modes that are common to all subunits consist mostly of the modes at the low frequency end, representing domain motions, or modes at the high frequency end, representing localized motions. The distribution in Figure 7 is similar to that in Figure 5, which is computed using the whole structure. Indeed, Of the first 50 lowest frequency symmetric modes of the actual structure, which are more likely to be functionally important than the other modes, 47 are included among the symmetric modes common to all subunits as identified above.

There is also a drawback with this last approach. The set of symmetric modes that are common to all subunits are not exactly the same as the set of symmetric modes computed from the whole complex, as indicted above. However, it is probable that both sets capture all

the symmetric modes that are functionally important and thus their small difference is not a problem. When this is in doubt, one may also compute the modes from the whole complex (given that it is computationally feasible) and compare the two sets of modes closely.

4 Discussion

In this work, we have looked into the symmetricity of protein normal modes and its implications. Symmetricity is introduced to define how similar the motions of the adjacent subunits are in symmetric structures. We show that the symmetry in structure can be taken advantage of to compute the normal modes of symmetric ring structures most efficiently. We then present a new theoretical result on the symmetricity of the normal modes of ring structures and confirm the theoretical prediction with computational results. Lastly, we show that the symmetricity pattern is broken in real structures due to the degeneracy of protein normal modes under structure variations. The work has several important implications.

The importance of symmetric and anti-symmetric modes in biological functions. It is perceivable that some functional motions of structurally symmetric systems are symmetric. The present work shows how to obtain symmetric normal modes separately in a fraction of the time that is otherwise needed to get all the modes. Anti-symmetric modes also are likely to be functionally important, as they represent a motion pattern in which every other subunit in a ring structure synchronizes perfectly and moves in the opposite direction to the other half of the subunits. As symmetric modes, anti-symmetric modes also can be obtained separately, using Eq. (8). The heat shock protein (HslU)⁴⁹, for example, which also is a hexamer, was found to allosterically bind ADP at every other subunit^{49,50}. The behavior may be best understood using anti-symmetric motions, where every other subunit has perfectly synchronized motions. It is likely that most AAA+ proteins⁵³ that employ a threading mechanism may use symmetric motions along the central pore as their primary

function motions. A deeper understanding of the motion patterns of their symmetric modes may help us better understand, for example, how protein unfolding and degradation are carried out.

On the other hand, it is possible that some molecular systems, though having symmetric structures, may not utilize symmetric motions as their primary functional movements. This is probably the case with GroEL/GroES complex⁵⁴. A major function of GroEL/ES is to provide a conducive environment for proteins to unfold and refold. It may rely on different kinds of motions such as twisting or stretching to accomplish this purpose. GroEL/GroES is a heptamer with seven subunits, which is a primer number. It cannot function as a trimer of dimers or a dimer of trimers as the hexameric HslU can. This might be the reason why its function motions are less or even not symmetric. It is possible that multi-mers with an even (or a highly composite) number of subunits should have more symmetric functional motions than those with an odd (or even a prime) number of subunits⁸.

Conformation changes that are symmetric require symmetric modes. It is likely that in some complexes symmetric motions are critical to the functions, such as threading. Consequently, the system must have a way to preserve at least some of its symmetric modes, especially those are key to function. Not all symmetric modes are robust to structure variations: some modes are more robust to structure perturbations than the others. The former preserve their motion patterns under small structural variations while the latter are unraveled under the same condition. Understanding the structural reason why some symmetric modes are preserved while the others are not should be useful. We show that the preserved symmetric modes are those with no degeneracy³⁶, mostly at either the low frequency end or the high frequency end. Non-degenerate normal modes are more likely to be functional³⁶.

The importance of symmetry in structure to symmetric motions. Identical protein units of some types can come together to form homooligomers. Whenever this is feasible, the interactions among them often drive them naturally to form beautiful complexes

of certain symmetry. The symmetric structure is energetically favored, corresponding to at least a stable local minimum in the energy landscape. Consequently, structure deviations from the symmetry are understandably disfavored. On the other hand, our work here shows that there is probably a dynamical reason why symmetry in structure ought to be maintained. Figure 3 shows that a structural deviation as small as 0.0042\AA can significantly affect the symmetricity distribution of the normal modes. We find that scarcely any symmetric modes are left (data not shown) when structure deviation is greater than 0.1\AA for p97. This means that if symmetric motions are critical to a complex's function, then maintaining the symmetry in structure (i.e., not having a significant deviation from it) not only makes the structure look appealing but also is important to preserving its key motion patterns and thus its functions as well. Symmetry in nature is not only aesthetically pleasing but may exist also for survival.

Degeneracy. For a circularly symmetric structure with m subunits, i.e., with C_m symmetry, what should be the degree of the degeneracy of its normal modes? Since the structure is m -fold symmetric, does it mean its normal modes have a degeneracy of m ? Group theory and our work here show that for symmetric or anti-symmetric modes (whose symmetricity equals to 1 or -1), there is no degeneracy. For all the other modes, the degeneracy is 2. For any pair of degenerate modes \mathbf{p}_1 and \mathbf{p}_2 , it is helpful to know that any linear combination of \mathbf{p}_1 and \mathbf{p}_2 also represents a valid mode³⁶. Besides the degeneracy originating from symmetry, there exists also degeneracy due to small structure deviations³⁶. This latter kind of degeneracy can cause some of symmetric modes to become degenerate with other modes, thus losing their perfect symmetry. However, functionally important symmetric modes should somehow be able to preserve their symmetricity.

C_2 symmetry. Having only two subunits (a dimer), structures with C_2 symmetry is a special case of the general circularly symmetric structures with m subunits (C_m). First, structures with C_2 symmetry have only symmetric (symmetricity=1) and anti-symmetric

(symmetricity=-1) modes. This means none of their modes are degenerate! In all the modes, the motions of two subunits are either fully symmetric, or fully anti-symmetric. Being the simplest case of structures with cyclic symmetry, C_2 structures may be ideal candidates for investigating how the symmetry in normal modes may be broken when the symmetry in structure is broken under small structure deviations. It will be helpful to quantitatively characterize symmetric modes that are robust to structure perturbations and to understand why small structure perturbations do not alter them.

Extension to other symmetric structures. Here we focus only on symmetric ring structures of C_n symmetry. It would be interesting to know what normal modes are like for structures with other kinds of symmetry (such as dihedral symmetry) and how their normal mode patterns are related to function. There are many symmetric complexes and they employ a number of different kinds of symmetry. Why do proteins form such symmetric complexes? One plausible reason given is that interactions drive it. Symmetric homomers are favored since interactions between the same structures are more favored than those between different structures⁵⁵. Another reason for the existence of symmetric complexes is for structure purpose. Symmetric complexes are formed, for example, to create a channel for protein unfolding and degradation⁵³, or to create a closed chamber for protein unfolding and refolding⁵⁴, etc. Indeed, the existence of symmetry in structures in nature has inspired scientists to bio-engineer new structures with desired symmetry and geometry using nucleic acids⁵⁶ or proteins^{57,58}. Yet another reason could be for functional purpose: the structure and the symmetry in the structure are needed for achieving certain patterns of motions that are functionally critical. Regarding this no much is known. Future studies in this area could reveal interesting insights and offer inspirations. It is foreseeable that symmetry-based ideas may be employed also to design structures with desired motion patterns someday in the future.

Link to small oscillations of n identical masses on a frictionless hoop. As

pointed out earlier in the paper, there is a tight link between symmetric ring structures and a system of n identical masses connected with identical springs on a frictionless hoop, which is a commonly used an example on small oscillations in classical mechanics⁵⁹. If we model each subunit as a sphere, a ring structure will become such a system of n masses. It is known that such an oscillating system has a zero mode and the rest of the modes (except for one if n is even) have a degeneracy of 2, just as we have shown for symmetric ring structures.

Acknowledgment

Funding from National Science Foundation (CAREER award, CCF-0953517) is gratefully acknowledged.

References

- [1] D. S. Goodsell and A. J. Olson. Structural symmetry and protein function. *Annu. Rev. Biophys. Biomol. Struct.*, 29:105–153, June 2000.
- [2] E. D. Levy, J. B. Pereira-Leal, C. Chothia, and S. A. Teichmann. 3D complex: a structural classification of protein complexes. *PLoS Comput Biol*, 2(11):e155, 2006.
- [3] K. Henrick and J. M. Thornton. PQS: a protein quaternary structure file server. *Trends Biochem Sci*, 23(9):358–61, 1998.
- [4] E. Krissinel and K. Henrick. Inference of macromolecular assemblies from crystalline state. *J Mol Biol*, 372(3):774–97, 2007.
- [5] E. D. Levy. PiQSi: protein quaternary structure investigation. *Structure*, 15(11):1364–7, 2007.

- [6] S. E. Ahnert, J. A. Marsh, H. Hernandez, C. V. Robinson, and S. A. Teichmann. Principles of assembly reveal a periodic table of protein complexes. *Science*, 350(6266):aaa2245, 2015.
- [7] E. D. Levy, E. Boeri Erba, C. V. Robinson, and S. A. Teichmann. Assembly reflects evolution of protein complexes. *Nature*, 453(7199):1262–5, 2008.
- [8] Y. Matsunaga, R. Koike, M. Ota, J. R. Tame, and A. Kidera. Influence of structural symmetry on protein dynamics. *PLoS One*, 7(11):e50011, 2012.
- [9] N. Go, T. Noguti, and T. Nishikawa. Dynamics of a small globular protein in terms of low-frequency vibrational modes. *Proc. Natl. Acad. Sci. USA*, 80(12):3696–3700, June 1983.
- [10] B. Brooks and M. Karplus. Harmonic dynamics of proteins: normal modes and fluctuations in bovine pancreatic trypsin inhibitor. *Proc. Natl. Acad. Sci. USA*, 80(21):6571–6575, November 1983.
- [11] M. Levitt, C. Sander, and P. S. Stern. The normal modes of a protein: Native bovine pancreatic trypsin inhibitor. *Int. J. Quant. Chem.*, 10:181–199, 1983.
- [12] M. M. Tirion. Large amplitude elastic motions in proteins from a single-parameter, atomic analysis. *Phys. Rev. Lett.*, 77:1905–1908, 1996.
- [13] I. Bahar, A. R. Atilgan, and B. Erman. Direct evaluation of thermal fluctuations in proteins using a single-parameter harmonic potential. *Folding & Design*, 2(3):173–181, 1997.
- [14] K. Hinsen. Analysis of domain motions by approximate normal mode calculations. *Proteins*, 33(3):417–429, March 1998.

- [15] K. Hinsén, A. Thomas, and M. J. Field. Analysis of domain motions in large proteins. *Proteins*, 34(3):369–382, February 1999.
- [16] A. R. Atilgan, S. R. Durell, R. L. Jernigan, M. C. Demirel, O. Keskin, and I. Bahar. Anisotropy of fluctuation dynamics of proteins with an elastic network model. *Biophys. J.*, 80(1):505–515, January 2001.
- [17] F. Tama and Y. H. Sanejouand. Conformational change of proteins arising from normal mode calculations. *Protein Eng.*, 14(1):1–6, January 2001.
- [18] O. Keskin, I. Bahar, D. Flatow, D. G. Covell, and R. L. Jernigan. Molecular mechanisms of chaperonin GroEL-GroES function. *Biochemistry*, 41(2):491–501, January 2002.
- [19] G. Li and Q. Cui. A coarse-grained normal mode approach for macromolecules: an efficient implementation and application to Ca(2+)-ATPase. *Biophys. J.*, 83:2457–2474, Nov 2002.
- [20] C. Xu, D. Tobi, and I. Bahar. Allosteric changes in protein structure computed by a simple mechanical model: Hemoglobin $t \leftrightarrow r2$ transition. *J. Mol. Biol.*, 333(1):153–168, 2003.
- [21] J. Ma. New advances in normal mode analysis of supermolecular complexes and applications to structural refinement. *Curr. Protein Pept. Sci.*, 5:119–123, 2004.
- [22] P. Maragakis and M. Karplus. Large amplitude conformational change in proteins explored with a plastic network model: adenylate kinase. *J. Mol. Biol.*, 352:807–822, Sep 2005.
- [23] A. W. Van Wynsberghe and Q. Cui. Comparison of mode analyses at different resolutions applied to nucleic acid systems. *Biophys. J.*, 89(5):2939–2949, November 2005.

- [24] W. Zheng and B. Brooks. Identification of dynamical correlations within the myosin motor domain by the normal mode analysis of an elastic network model. *J. Mol. Biol.*, 346(3):745–759, February 2005.
- [25] O. Kurkcuoglu, R. L. Jernigan, and P. Doruker. Loop motions of triosephosphate isomerase observed with elastic networks. *Biochemistry*, 45(4):1173–1182, January 2006.
- [26] F. Tama and C. L. Brooks III. Symmetry, form, and shape: guiding principles for robustness in macromolecular machines. *Annu. Rev. Biophys. Biomol. Struct.*, 35:115–133, 2006.
- [27] T. R. Lezon, A. Sali, and I. Bahar. Global motions of the nuclear pore complex: insights from elastic network models. *PLoS Comput. Biol.*, 5(9), September 2009.
- [28] L. Yang, G. Song, and R. L. Jernigan. Protein elastic network models and the ranges of cooperativity. *Proc. Natl. Acad. Sci. USA*, 106(30):12347–12352, July 2009.
- [29] T.-L. Lin and G. Song. Generalized spring tensor models for protein fluctuation dynamics and conformation changes. *BMC Structural Biology*, 10(Suppl 1):S3+, 2010.
- [30] M. H. Kim, S. Seo, J. I. Jeong, B. J. Kim, W. K. Liu, B. S. Lim, J. B. Choi, and M. K. Kim. A mass weighted chemical elastic network model elucidates closed form domain motions in proteins. *Protein Sci.*, 22:605–613, May 2013.
- [31] H. Na and G. Song. A natural unification of GNM and ANM and the role of inter-residue forces. *Phys. Biol.*, 11(3):036002, April 2014.
- [32] H. Na and G. Song. Bridging between normal mode analysis and elastic network models. *Proteins*, 82:2157–2168, September 2014.
- [33] P. Ozbek, S. Soner, and T. Haliloglu. Hot spots in a network of functional sites. *PLoS ONE*, 8(9):e74320, 2013.

- [34] K. Hinsen and G. R. Kneller. Projection methods for the analysis of complex motions in macromolecules. *Mol. Sim.*, 23:275–292, 2000.
- [35] H. Na, G. Song, and D. Ben-Avraham. Universality of vibrational spectra of globular proteins. *Phys. Biol.*, 13(1):016008, February 2016.
- [36] H. Na and G. Song. The effective degeneracy of protein normal modes. *Phys Biol*, 13(3):036002, 2016.
- [37] T. Simonson and D. Perahia. Normal modes of symmetric protein assemblies. application to the tobacco mosaic virus protein disk. *Biophys J*, 61(2):410–27, 1992.
- [38] H. W. van Vlijmen and M. Karplus. Normal mode analysis of large systems with icosahedral symmetry: application to (Dialanine)₆₀ in full and reduced basis set. *J. Chem. Phys.*, 115(2):691698, 2001.
- [39] H. W. van Vlijmen and M. Karplus. Normal mode calculations of icosahedral viruses with full dihedral flexibility by use of molecular symmetry. *J Mol Biol*, 350(3):528–42, 2005.
- [40] H. Na and G. Song. Predicting the functional motions of p97 using symmetric normal modes. *Proteins*, 2016. Under review.
- [41] F.A. Cotton. *Chemical Applications of Group Theory*. Wiley, New York, 3rd edition, 1990.
- [42] P. J. Davis. *Circulant Matrices*. Wiley, New York, 1970.
- [43] https://en.wikipedia.org/wiki/Circulant_matrix.
- [44] Cao Z. A note on symmetric block circulant matrix. *J. Math. Res. Exposition*, 10:469–473, 1990.

- [45] H. M. Berman, J. Westbrook, Z. Feng, G. Gilliland, T. N. Bhat, H. Weissig, I. N. Shindyalov, and P. E. Bourne. The protein data bank. *Nucleic Acids Res*, 28(1):235–42, 2000.
- [46] Y. Ye, H. H. Meyer, and T. A. Rapoport. The AAA ATPase Cdc48/p97 and its partners transport proteins from the ER into the cytosol. *Nature*, 414:652–656, December 2001.
- [47] S. Banerjee, A. Bartesaghi, A. Merk, P. Rao, S. L. Bulfer, Y. Yan, N. Green, B. Mroczkowski, R. J. Neitz, P. Wipf, V. Falconieri, R. J. Deshaies, J. L.S. Milne, D. Huryn, M. Arkin, and S. Subramaniam. 2.3 Å resolution cryo-EM structure of human p97 and mechanism of allosteric inhibition. *Science*, 351:871–875, February 2016.
- [48] J. Wang, J. J. Song, M. C. Franklin, S. Kamtekar, Y. J. Im, S. H. Rho, I. S. Seong, C. S. Lee, C. H. Chung, and S. H. Eom. Crystal structures of the HslVU peptidase-ATPase complex reveal an ATP-dependent proteolysis mechanism. *Structure*, 9(2):177–84, 2001.
- [49] M. Bochtler, C. Hartmann, H. K. Song, G. P. Bourenkov, H. D. Bartunik, and R. Huber. The structures of HslU and the ATP-dependent protease HslU-HslV. *Nature*, 403(6771):800–5, 2000.
- [50] J. Wang, J.J. Song, I.S. Seong, M.C. Franklin, S. Kamtekar, S.H. Eom, and C.H. Chung. Nucleotide-dependent conformational changes in a protease-associated atpase HslU. *Structure*, 9:1107–1116, 2001.
- [51] P. Hänzelmann and H. Schindelin. Structural basis of ATP hydrolysis and intersubunit signaling in the AAA+ ATPase p97. *Structure*, 24:127–139, January 2016.
- [52] X. Bai, G. McMullan, and S. H.W Scheres. How cryo-EM is revolutionizing structural biology. *Trends in Biochemical Sciences*, 40(1):49 – 57, 2015.

- [53] P. I. Hanson and S. W. Whiteheart. AAA+ proteins: have engine, will work. *Nat. Rev. Mol. Cell Biol.*, 6:519–529, July 2005.
- [54] Z. Xu, A. L. Horwich, and P. B. Sigler. The crystal structure of the asymmetric GroEL-GroES-(ADP)₇ chaperonin complex. *Nature*, 388:741–750, August 1997.
- [55] D. B. Lukatsky, B. E. Shakhnovich, J. Mintseris, and E. I. Shakhnovich. Structural similarity enhances interaction propensity of proteins. *J Mol Biol*, 365(5):1596–606, 2007.
- [56] N. C. Seeman. Nanomaterials based on DNA. *Annu Rev Biochem*, 79:65–87, 2010.
- [57] Y. T. Lai, D. Cascio, and T. O. Yeates. Structure of a 16-nm cage designed by using protein oligomers. *Science*, 336(6085):1129, 2012.
- [58] N. P. King, W. Sheffler, M. R. Sawaya, B. S. Vollmar, J. P. Sumida, I. Andre, T. Gonen, T. O. Yeates, and D. Baker. Computational design of self-assembling protein nanomaterials with atomic level accuracy. *Science*, 336(6085):1171–4, 2012.
- [59] Herbert Goldstein. *Classical mechanics*. Addison-Wesley Press, Cambridge, Mass, 1950.

Figure Legends

Figure 1.

(A) The symmetricity distribution of the normal modes of a perfectly symmetric hexamer (p97). The structure is constructed manually using subunit A of PDB structure 5ftl (see the Methods section). (B) The symmetricity values shown in (A) are the same as the abscissa values of the vertices of a unit hexagon.

Figure 2.

Symmetric modes distribute through the whole frequency range. Frequency values are obtained by simply taking the square root of the eigenvalues (λ 's). The ANM model¹⁶ is used, whose cutoff distance is set at 13 Å and whose spring constant is set at 1. 100 bins are used.

Figure 3.

The symmetricity of an actual structure (5ftl) that is nearly perfectly symmetric. A uniform bin size of 0.02 is used and there are 100 bins. The RMSD between this structure and the manually constructed perfect structure (5ftl-perfect) is only 0.0042 Å. However, the symmetricity distribution is significantly affected and becomes more widely spread than that in Figure 1(A), though the peaks remain clearly identifiable. In total, only 237 modes maintain a very high symmetricity (greater than 0.98, in the rightmost bin). In comparison, 2,167 modes, or one sixth of the total number of the modes, have perfect symmetricity (equal to 1) in Figure 1(A).

Figure 4.

The symmetricity of the modes of a perfect structure (5ftl-perfect) and that of their best matching modes in the actual structure (5ftl).

Figure 5.



The indices of the 237 symmetric modes that preserve their symmetricity under small structure variations. Most of these modes fall into the low frequency or the high frequency end.

Figure 6.

Between the 2,167 symmetric modes of the perfect structure (5ftl-perfect) and their best matching modes in the actual structure (5ftl), modes that maintain a high overlap maintain also a high symmetricity in the actual structure (see the text).

Figure 7.

Symmetric modes that are common to all subunits of a p97 structure (pdb-id: 5ftl). Symmetric modes are computed using Eq. (7) as each subunit in turn is used as the center. The modes computed from each subunit are then compared with the symmetric modes of a perfect structure of p97: 5ftl-perfect. The indices of the symmetric modes that are common to all subunits fall mostly into either the low frequency or the high frequency end.



(B)

Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:




Figure 6:




Figure 7: