

Homoclinical structure of SICNNs with rectangular input currents

Mehmet Onur Fen^{1,*}, Fatma Tokmak Fen²

¹*Basic Sciences Unit, TED University, 06420 Ankara, Turkey*

²*Department of Mathematics, Gazi University, 06500 Ankara, Turkey*

Abstract

Shunting inhibitory cellular neural networks (SICNNs) with continuous as well as discontinuous external inputs are investigated. The descriptions of homoclinic and heteroclinic motions are provided in the functional sense for the multidimensional dynamics of SICNNs, and it is demonstrated that the networks under investigation exhibit such motions. Homoclinic and heteroclinic outputs that are asymptotic to quasi-periodic outputs are illustrated.

Keywords: Shunting inhibitory cellular neural networks; Homoclinic and heteroclinic outputs; Rectangular input currents; Stable and unstable sets; Quasi-periodic outputs

1 Introduction

Starting with the studies Poincaré [1] the investigation of homoclinic and heteroclinic motions has been one of the attractive topics among scientists who study nonlinear dynamics. The presence of homoclinic motions is an effective tool for the demonstration of chaotic behavior since any neighborhood of a structurally stable Poincaré homoclinic orbit admits nontrivial hyperbolic sets that contain a countable number of saddle periodic orbits and continuum of non-periodic Poisson stable orbits [2]-[4]. Besides, heteroclinic motions are also crucial for chaotic dynamics [5, 6].

The present study is devoted to the investigation of homoclinic and heteroclinic motions generated by shunting inhibitory cellular neural networks (SICNNs), which is a class of biologically inspired cellular neural networks introduced by Bouzerdoum and Pinter [7]. This type of neural networks have been extensively applied in psychophysics, speech, perception, robotics, adaptive pattern recognition, vision and image processing [8]-[15].

The model of SICNNs in its most original form is as follows [7]. Consider a two dimensional grid of processing cells arranged into m rows and n columns, and let C_{ij} denote the cell at the (i, j) position of the lattice, where $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$. In SICNNs, neighboring cells exert mutual

*Corresponding Author. E-mail: monur.fen@gmail.com, Tel: +90 312 585 02 17

inhibitory interactions of the shunting type. Define the r -neighborhood of the cell C_{ij} as

$$N_r(i, j) = \{C_{hl} : \max(|h - i|, |l - j|) \leq r, 1 \leq h \leq m, 1 \leq l \leq n\}.$$

The dynamics of a cell C_{ij} are described by the following nonlinear ordinary differential equation:

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{hl} \in N_r(i, j)} C_{ij}^{hl} f(x_{hl}(t))x_{ij} + \mathcal{L}_{ij}(t), \quad (1.1)$$

where x_{ij} is the activity of the cell C_{ij} ; $\mathcal{L}_{ij}(t)$ is the external input to the cell C_{ij} ; the constant $a_{ij} > 0$ represents the passive decay rate of the cell activity; $C_{ij}^{hl} \geq 0$ is the coupling strength of post-synaptic activity of the cell C_{hl} transmitted to the cell C_{ij} ; the positive and continuous function $f(x_{hl})$ represents the output or firing rate of the cell C_{hl} .

The existence and stability of periodic, almost-periodic and anti-periodic solutions of SICNNs have been widely studied in the literature [16]-[25]. Moreover, chaos in SICNNs were considered in the papers [26]-[30]. However, to the best of our knowledge, the existence of homoclinic and heteroclinic solutions of SICNNs has not been investigated before in the literature. Motivated by the lack of mathematical methods for the investigation of homoclinic and heteroclinic motions in SICNNs, we propose the results of the present study. Both continuous and discontinuous external inputs in rectangular form are utilized in our model.

The generation of homoclinic and heteroclinic motions in systems of differential equations by means of discontinuous perturbations was first realized by Akhmet [31] on the basis of functional spaces. The paper [31] was concerned with the presence of homoclinic motions embedded in the chaotic attractor of a relay system (see [32, 33] for details about relay systems). Similar results for impulsive differential equations were obtained in the studies [34, 35], and the formation of homoclinic and heteroclinic motions in economic models was considered within the scope of the paper [36]. Based on the definitions given in [31], in the present study, we provide the descriptions of homoclinic and heteroclinic motions for the multidimensional dynamics of SICNNs in the functional sense, and rigorously prove that the discontinuous inputs give rise to the generation of homoclinic and heteroclinic motions.

One of the techniques for the confirmation of chaos in neural networks is to determine homoclinic and heteroclinic motions. A geometric construction of a transversal homoclinic orbit for a nonlinear neuron model with time delays was provided by Chen [37] in order to show the existence of chaos. Moreover, Li et al. [38] proved the existence of a saddle periodic orbit in the dynamics of a small Hopfield neural network with a memristive synaptic weight, and verified the existence of a hyperchaotic invariant set via a homoclinic intersection of its stable and unstable manifolds by using the Smale-Birkhoff homoclinic theorem. Zou et al. [39], on the other hand, presented a method for finding homoclinic and heteroclinic motions in the three-cell autonomous cellular neural network, which has chaotic behavior similar to that

of the double scroll attractor in a certain parameter range. In the study [39], homoclinic motions that are asymptotic to equilibria are considered. However, in this paper, we take into account a larger class of homoclinic and heteroclinic motions, and illustrate the presence of such motions which are asymptotic to quasi-periodic ones in Section 4.

The rest of the paper is organized as follows. In the next section the model of SICNNs with rectangular input currents (RICs) is introduced, and some results concerning the bounded solutions are mentioned. Section 3 is devoted to the descriptions of homoclinic and heteroclinic motions of SICNNs as well as the main result of the present study. An illustrative example, which support the theoretical results, is provided in Section 4. Finally, some concluding remarks are given in Section 5.

2 The model

We will take into account the external input $\mathcal{L}_{ij}(t)$ in SICNN (1.1) as the sum of the continuous external input $L_{ij}(t)$ and the RIC $P_{ij}(t, \zeta)$. More precisely, the main object of investigation is the following SICNN,

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} f(x_{hl}(t))x_{ij} + L_{ij}(t) + P_{ij}(t, \zeta). \quad (2.2)$$

In model (2.2), for each $i = 1, 2, \dots, m$ and $j = 1, 2, \dots, n$, the input currents $P_{ij}(t, \zeta)$ are defined through the equations $P_{ij}(t, \zeta) = \zeta_k^{ij}$ for $t \in (\theta_k, \theta_{k+1}]$, $k \in \mathbb{Z}$, such that the sequence $\{\theta_k\}_{k \in \mathbb{Z}}$ of discontinuity moments is strictly increasing, $|\theta_k| \rightarrow \infty$ as $|k| \rightarrow \infty$, and $\zeta = \{\zeta_k\}_{k \in \mathbb{Z}}$ is a sequence generated by the map

$$\zeta_{k+1} = F(\zeta_k), \quad (2.3)$$

where $F : \Lambda \rightarrow \Lambda$ is a continuous function and Λ is a bounded subset of \mathbb{R}^{mn} . Our purpose is to prove the existence of homoclinic and heteroclinic motions in the dynamics of SICNN (2.2) in the case that the map (2.3) admits homoclinic and heteroclinic orbits.

The dynamics of SICNNs with RICs were also investigated in the paper [29]. It was found in [29] that if the discontinuity moments of the RICs change chaotically, then the network possesses chaos with infinitely many almost periodic motions in basis. Homoclinic and heteroclinic motions were not investigated in [29], and the novelty of the present study is the investigation of such motions in the dynamics of SICNNs. Another difference is that the discontinuity moments of the RICs are generated by a discrete chaotic map in [29], while the values of the RICs in the intervals of constancy are generated by a discrete map, which possesses homoclinic and heteroclinic orbits. Besides, it was mentioned in [29] that the discontinuity moments of the RICs can be comprehended as spike moments, and it is reasonable

to use networks with this type of inputs in modeling biological neural systems.

In an electrical equivalent circuit of a cell of SICNNs, the synaptic conductance of each inhibitory channel is proportional to the firing rate of the cell controlling it, and the shunting conductance of the cell is the sum of the synaptic conductances of individual inhibitory channels [7]. Since both continuous and discontinuous external inputs are utilized, one may design an electrical equivalent circuit of the network (2.2) by properly coupling appropriate pulse generating circuits [40, 41] with the one mentioned in [7].

Throughout the paper, \mathbb{R} and \mathbb{Z} will stand for the sets of real numbers and integers, respectively, and the norm $\|u\| = \max_{(i,j)} |u_{ij}|$ will be used, where $u = \{u_{ij}\} = (u_{11}, \dots, u_{1n}, \dots, u_{m1}, \dots, u_{mn}) \in \mathbb{R}^{m \times n}$.

The following conditions are required.

- (C1) There exists a positive number L_f such that $|f(s_1) - f(s_2)| \leq L_f |s_1 - s_2|$ for all $s_1, s_2 \in \mathbb{R}$;
- (C2) There exist positive numbers M_f and M_{ij} such that $\sup_{s \in \mathbb{R}} |f(s)| \leq M_f$ and $\sup_{t \in \mathbb{R}} |L_{ij}(t)| \leq M_{ij}$ for each $i = 1, 2, \dots, m, j = 1, 2, \dots, n$.

Suppose that the inequality $M_f \max_{(i,j)} \frac{\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl}}{a_{ij}} < 1$ is valid. Let us denote $\gamma = \min_{(i,j)} a_{ij}$, $\delta = \max_{(i,j)} \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl}$ and $K_0 = \frac{\max_{(i,j)} \frac{M_{ij} + M_F}{a_{ij}}}{1 - M_f \max_{(i,j)} \frac{\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl}}{a_{ij}}}$, where $M_F = \sup_{\xi \in \Lambda} \|F(\xi)\|$. It is worth noting that the number γ is positive and δ is nonnegative since the numbers a_{ij} are positive and the coupling strengths C_{ij}^{hl} are nonnegative.

The following condition is also needed.

- (C3) $\gamma - (M_f + L_f K_0) \delta > 0$.

Since the model (2.2) satisfies the Lipschitz condition in the intervals of constancy of $P_{ij}(t, \zeta)$, the local existence and uniqueness of solutions are valid, and at the discontinuity moments, one can use the continuity condition of solutions to proceed for the next interval of constancy of $P_{ij}(t, \zeta)$. Therefore, the local existence and uniqueness of solutions as well as their continuation to the maximal interval of existence are assured. The reader is referred to the books [42, 43] for more information about differential equations with discontinuous right hand side.

If the conditions (C1) – (C3) are valid, then the technique mentioned in the paper [27] can be used to prove that for each solution $\zeta = \{\zeta_k\}_{k \in \mathbb{Z}}$ of (2.3), there exists a unique bounded on \mathbb{R} solution $\phi_\zeta(t) = \{\phi_\zeta^{ij}(t)\}$, $i = 1, 2, \dots, m, j = 1, 2, \dots, n$, of (2.2), which satisfies the following relation

$$\phi_\zeta^{ij}(t) = - \int_{-\infty}^t e^{-a_{ij}(t-s)} \left[\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} f(\phi_\zeta^{hl}(s)) \phi_\zeta^{ij}(s) - L_{ij}(s) - P_{ij}(s, \zeta) \right] ds \quad (2.4)$$

such that $\sup_{t \in \mathbb{R}} \|\phi_\zeta(t)\| \leq K_0$. Moreover, under the same conditions, it can be verified in a very similar way to Theorem 2 [44] that for a fixed solution $\zeta = \{\zeta_k\}_{k \in \mathbb{Z}}$ of (2.3) all other solutions of SICNN (2.2)

converge exponentially to the unique bounded on \mathbb{R} solution $\phi_\zeta(t)$.

3 Homoclinic and heteroclinic motions

Let us describe the stable, unstable and hyperbolic sets as well as the homoclinic and heteroclinic motions for both SICNN (2.2) and the discrete map (2.3). These definitions are adapted from the papers [31, 34].

Suppose that Θ denotes the set of all sequences $\zeta = \{\zeta_k\}_{k \in \mathbb{Z}} \subset \Lambda$ obtained by (2.3). The stable set of a sequence $\zeta \in \Theta$ is defined as

$$W^s(\zeta) = \{\eta \in \Theta \mid \|\eta_k - \zeta_k\| \rightarrow 0 \text{ as } k \rightarrow \infty\},$$

and the unstable set of ζ is

$$W^u(\zeta) = \{\eta \in \Theta \mid \|\eta_k - \zeta_k\| \rightarrow 0 \text{ as } k \rightarrow -\infty\}.$$

The set Θ is called hyperbolic if for each $\zeta \in \Theta$ the stable and unstable sets of ζ contain at least one element different from ζ . A sequence $\eta \in \Theta$ is homoclinic to another sequence $\zeta \in \Theta$ if $\eta \in W^s(\zeta) \cap W^u(\zeta)$. Moreover, $\eta \in \Theta$ is heteroclinic to the sequences $\zeta^1 \in \Theta$, $\zeta^2 \in \Theta$, $\eta \neq \zeta^1$, $\eta \neq \zeta^2$, if $\eta \in W^s(\zeta^1) \cap W^u(\zeta^2)$.

On the other hand, let \mathcal{A} be the set consisting of all bounded on \mathbb{R} solutions $\phi_\zeta(t)$, $\zeta \in \Theta$, of SICNN (2.2). A bounded solution $\phi_\eta(t) \in \mathcal{A}$ belongs to the stable set $W^s(\phi_\zeta(t))$ of $\phi_\zeta(t) \in \mathcal{A}$ if $\|\phi_\eta(t) - \phi_\zeta(t)\| \rightarrow 0$ as $t \rightarrow \infty$. Besides, $\phi_\eta(t)$ is an element of the unstable set $W^u(\phi_\zeta(t))$ of $\phi_\zeta(t)$ provided that $\|\phi_\eta(t) - \phi_\zeta(t)\| \rightarrow 0$ as $t \rightarrow -\infty$.

We say that \mathcal{A} is hyperbolic if for each $\phi_\zeta(t) \in \mathcal{A}$ the sets $W^s(\phi_\zeta(t))$ and $W^u(\phi_\zeta(t))$ contain at least one element different from $\phi_\zeta(t)$. A solution $\phi_\eta(t) \in \mathcal{A}$ is homoclinic to another solution $\phi_\zeta(t) \in \mathcal{A}$ if $\phi_\eta(t) \in W^s(\phi_\zeta(t)) \cap W^u(\phi_\zeta(t))$, and $\phi_\eta(t) \in \mathcal{A}$ is heteroclinic to the bounded solutions $\phi_{\zeta^1}(t)$, $\phi_{\zeta^2}(t) \in \mathcal{A}$, $\phi_\eta(t) \neq \phi_{\zeta^1}(t)$, $\phi_\eta(t) \neq \phi_{\zeta^2}(t)$, if $\phi_\eta(t) \in W^s(\phi_{\zeta^1}(t)) \cap W^u(\phi_{\zeta^2}(t))$.

The relations between the stable and unstable sets of the network (2.2) and the discrete map (2.3) are respectively provided in Lemma 3.1 and Lemma 3.2.

Lemma 3.1 *Assume that the conditions (C1) – (C3) are valid, and let ζ and η be elements of Θ . If $\eta \in W^s(\zeta)$, then $\phi_\eta(t) \in W^s(\phi_\zeta(t))$.*

Proof. Fix an arbitrary positive number ϵ , and let α be a real number such that

$$\alpha \geq 1 + \frac{1}{\gamma - (M_f + L_f K_0) \delta}.$$

Since $\eta = \{\eta_k\}_{k \in \mathbb{Z}}$ is an element of $W^s(\zeta)$, there exists an integer k_0 such that $\|\eta_k - \zeta_k\| < \frac{\epsilon}{\alpha}$ for all $k \geq k_0$.

According to (2.4), the bounded solutions $\phi_\eta(t)$ and $\phi_\zeta(t)$ satisfy the following relation

$$\begin{aligned} \phi_\eta^{ij}(t) - \phi_\zeta^{ij}(t) &= - \int_{-\infty}^t e^{-a_{ij}(t-s)} \left[\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} f(\phi_\eta^{hl}(s)) \phi_\eta^{ij}(s) - P_{ij}(s, \eta) \right. \\ &\quad \left. - \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} f(\phi_\zeta^{hl}(s)) \phi_\zeta^{ij}(s) + P_{ij}(s, \zeta) \right] ds. \end{aligned}$$

Using the inequality $|P_{ij}(t, \eta) - P_{ij}(t, \zeta)| < \frac{\epsilon}{\alpha}$, $t \in (\theta_{k_0}, \infty)$, one can obtain on the same interval that

$$\begin{aligned} |\phi_\eta^{ij}(t) - \phi_\zeta^{ij}(t)| &\leq 2(M_f K_0 \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} + M_F) \int_{-\infty}^{\theta_{k_0}} e^{-a_{ij}(t-s)} ds \\ &+ \int_{\theta_{k_0}}^t e^{-a_{ij}(t-s)} M_f \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} |\phi_\eta^{ij}(s) - \phi_\zeta^{ij}(s)| ds \\ &+ \int_{\theta_{k_0}}^t e^{-a_{ij}(t-s)} L_f K_0 \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} |\phi_\eta^{hl}(s) - \phi_\zeta^{hl}(s)| ds + \int_{\theta_{k_0}}^t e^{-a_{ij}(t-s)} \frac{\epsilon}{\alpha} ds. \end{aligned}$$

Therefore, we have for $t > \theta_{k_0}$ that

$$\begin{aligned} \|\phi_\eta(t) - \phi_\zeta(t)\| &\leq 2 \max_{(i,j)} \frac{1}{a_{ij}} \left(M_f K_0 \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} + M_F \right) e^{-\gamma(t-\theta_{k_0})} + \frac{\epsilon}{\gamma\alpha} \left(1 - e^{-\gamma(t-\theta_{k_0})} \right) \\ &+ (M_f + L_f K_0) \max_{(i,j)} \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} \int_{\theta_{k_0}}^t e^{-\gamma(t-s)} \|\phi_\eta(s) - \phi_\zeta(s)\| ds. \end{aligned} \quad (3.5)$$

Define the function $v(t) = e^{\gamma t} \|\phi_\eta(t) - \phi_\zeta(t)\|$ and let

$$R_0 = 2 \max_{(i,j)} \frac{1}{a_{ij}} \left(M_f K_0 \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} + M_F \right) - \frac{\epsilon}{\gamma\alpha}.$$

The inequality (3.5) yields

$$v(t) \leq R_0 e^{\gamma\theta_{k_0}} + \frac{\epsilon}{\gamma\alpha} e^{\gamma t} + (M_f + L_f K_0) \delta \int_{\theta_{k_0}}^t v(s) ds, \quad t > \theta_{k_0}.$$

Applying the Gronwall's Lemma [45] to the last inequality one can verify that

$$\begin{aligned} v(t) &\leq \frac{\epsilon}{\gamma\alpha} e^{\gamma t} + R_0 e^{(M_f + L_f K_0)\delta(t-\theta_{k_0})} e^{\gamma\theta_{k_0}} \\ &\quad + \frac{\epsilon(M_f + L_f K_0)\delta}{\gamma\alpha [\gamma - (M_f + L_f K_0)\delta]} e^{\gamma t} \left(1 - e^{-[\gamma - (M_f + L_f K_0)\delta](t-\theta_{k_0})} \right). \end{aligned}$$

Multiplying both sides of the last inequality by $e^{-\gamma t}$ we obtain that

$$\|\phi_\eta(t) - \phi_\zeta(t)\| < \frac{\epsilon}{\alpha [\gamma - (M_f + L_f K_0)\delta]} + R_0 e^{-[\gamma - (M_f + L_f K_0)\delta](t-\theta_{k_0})}, \quad t > \theta_{k_0}.$$

Now, let E be a positive number such that $E \geq \frac{1}{\gamma - (M_f + L_f K_0)\delta} \ln \left(\frac{R_0 \alpha}{\epsilon} \right)$. Hence, for $t \geq E + \theta_{k_0}$, we have

$$\|\phi_\eta(t) - \phi_\zeta(t)\| < \frac{\epsilon}{\alpha} \left(1 + \frac{1}{\gamma - (M_f + L_f K_0)\delta} \right) \leq \epsilon.$$

Consequently, $\phi_\eta(t)$ belongs to $W^s(\phi_\zeta(t))$. \square

Lemma 3.2 *Assume that the conditions (C1) – (C3) are valid, and let ζ and η be elements of Θ . If $\eta \in W^u(\zeta)$, then $\phi_\eta(t) \in W^u(\phi_\zeta(t))$.*

Proof. Fix an arbitrary positive number ϵ , and let β be a real number satisfying

$$\beta > \frac{1}{\gamma} \left(1 - (M_f + L_f K_0) \max_{(i,j)} \frac{\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl}}{a_{ij}} \right)^{-1}.$$

Because the sequence $\eta = \{\eta_k\}_{k \in \mathbb{Z}}$ belongs to the unstable set $W^u(\zeta)$ of $\zeta = \{\zeta_k\}_{k \in \mathbb{Z}}$, there exists an integer k_0 such that $\|\eta_k - \zeta_k\| < \frac{\epsilon}{\beta}$ for all $k \leq k_0$. In this case, for each i and j , one can confirm that $|P_{ij}(t, \eta) - P_{ij}(t, \zeta)| < \frac{\epsilon}{\beta}$, $t \leq \theta_{k_0+1}$.

Making use of the relation

$$\begin{aligned} \phi_\eta^{ij}(t) - \phi_\zeta^{ij}(t) &= - \int_{-\infty}^t e^{-a_{ij}(t-s)} \left[\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} f(\phi_\eta^{hl}(s)) \phi_\eta^{ij}(s) - P_{ij}(s, \eta) \right. \\ &\quad \left. - \sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl} f(\phi_\zeta^{hl}(s)) \phi_\zeta^{ij}(s) + P_{ij}(s, \zeta) \right] ds, \end{aligned}$$

it can be verified for $t \leq \theta_{k_0+1}$ that

$$|\phi_\eta^{ij}(t) - \phi_\zeta^{ij}(t)| < \frac{\epsilon}{\beta a_{ij}} + (M_f + L_f K_0) \frac{\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl}}{a_{ij}} \sup_{t \leq \theta_{k_0+1}} \|\phi_\eta(t) - \phi_\zeta(t)\|.$$

The last inequality implies that

$$\sup_{t \leq \theta_{k_0+1}} \|\phi_\eta(t) - \phi_\zeta(t)\| \leq \frac{\epsilon}{\gamma \beta} \left(1 - (M_f + L_f K_0) \max_{(i,j)} \frac{\sum_{C_{hl} \in N_r(i,j)} C_{ij}^{hl}}{a_{ij}} \right)^{-1} < \epsilon.$$

Hence, $\lim_{t \rightarrow -\infty} \|\phi_\eta(t) - \phi_\zeta(t)\| = 0$. \square

The following theorem can be proved using the results of Lemma 3.1 and Lemma 3.2.

Theorem 3.1 *Under the conditions (C1) – (C3), the following assertions are valid.*

- (i) *If $\eta \in \Theta$ is homoclinic to $\zeta \in \Theta$, then $\phi_\eta(t) \in \mathcal{A}$ is homoclinic to $\phi_\zeta(t) \in \mathcal{A}$;*
- (ii) *If $\eta \in \Theta$ is heteroclinic to $\zeta^1, \zeta^2 \in \Theta$, then $\phi_\eta(t) \in \mathcal{A}$ is heteroclinic to $\phi_{\zeta^1}(t), \phi_{\zeta^2}(t) \in \mathcal{A}$;*
- (iii) *If Θ is hyperbolic, then the same is true for \mathcal{A} .*

The next section is devoted to an illustrative example.

4 An Example

Let us consider the SICNN

$$\frac{dx_{ij}}{dt} = -a_{ij}x_{ij} - \sum_{C_{hl} \in N_1(i,j)} C_{ij}^{hl} f(x_{hl}(t))x_{ij} + L_{ij}(t) + P_{ij}(t, \zeta), \quad (4.6)$$

where $i, j = 1, 2, 3$, $f(s) = \frac{\arctan(s^2)}{9}$, $\theta_k = \frac{7k}{2}$, $k \in \mathbb{Z}$, $L_{11}(t) = 0.4 \sin(2\pi t) + 0.6 \cos(\sqrt{5}t)$, $L_{12}(t) = 0.4 \cos(\sqrt{2}t) - 0.2 \cos t$, $L_{13}(t) = 0.5 \sin(3t) + 0.3 \sin(2\pi t)$, $L_{21}(t) = 0.1 \sin(3\pi t) - 0.6 \cos(\sqrt{2}t)$, $L_{22}(t) = 0.2 \sin(\pi t) + 0.1 \cos(2t)$, $L_{23}(t) = 0.1 \cos(2\sqrt{3}t) + 0.7 \cos(5\pi t)$, $L_{31}(t) = 0.4 \sin(\pi t/2) + 0.5 \cos t$, $L_{32}(t) = 0.7 \sin(3t) - 0.1 \sin(\sqrt{2}t)$, $L_{33}(t) = 0.5 \sin(\sqrt{3}t) + 0.2 \sin(5t)$,

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{pmatrix} 0.4 & 0.6 & 0.5 \\ 0.7 & 0.3 & 0.8 \\ 0.4 & 0.9 & 0.6 \end{pmatrix},$$

and

$$\begin{pmatrix} C_{ij}^{11} & C_{ij}^{12} & C_{ij}^{13} \\ C_{ij}^{21} & C_{ij}^{22} & C_{ij}^{23} \\ C_{ij}^{31} & C_{ij}^{32} & C_{ij}^{33} \end{pmatrix} = \begin{pmatrix} 0.004 & 0.006 & 0.002 \\ 0.006 & 0.005 & 0.002 \\ 0.001 & 0.007 & 0 \end{pmatrix}$$

whenever the cells C_{hl} , $h, l = 1, 2, 3$ belong to $N_1(i, j)$, i.e., for fixed h and l , the coupling strengths C_{ij}^{hl} , $i, j = 1, 2, 3$, are taken to be equal to each other in cases where C_{hl} belongs to $N_1(i, j)$.

In (4.6), for each i and j , we take the rectangular input currents as $P_{ij}(t, \zeta) = \zeta_k$, $t \in (\theta_k, \theta_{k+1}]$, $k \in \mathbb{Z}$. Here, the sequence $\zeta = \{\zeta_k\}_{k \in \mathbb{Z}}$ is a solution the logistic map

$$\zeta_{k+1} = F_\mu(\zeta_k), \quad (4.7)$$

where $F_\mu(s) = \mu s(1 - s)$. For the values of the parameter $\mu \in (0, 4]$, the unit interval $[0, 1]$ is invariant under the iterations of (4.7) [46]. The inverses of the function F_μ with ranges on the intervals $[0, 1/2]$ and $[1/2, 1]$ are $G_\mu(s) = \frac{1}{2} \left(1 - \sqrt{1 - \frac{4s}{\mu}} \right)$ and $H_\mu(s) = \frac{1}{2} \left(1 + \sqrt{1 - \frac{4s}{\mu}} \right)$, respectively.

One can calculate that

$$\sum_{C_{hl} \in N_1(1,1)} C_{11}^{hl} = 0.021, \quad \sum_{C_{hl} \in N_1(1,2)} C_{12}^{hl} = 0.025, \quad \sum_{C_{hl} \in N_1(1,3)} C_{13}^{hl} = 0.015,$$

$$\begin{aligned} \sum_{C_{hl} \in N_1(2,1)} C_{21}^{hl} &= 0.029, & \sum_{C_{hl} \in N_1(2,2)} C_{22}^{hl} &= 0.033, & \sum_{C_{hl} \in N_1(2,3)} C_{23}^{hl} &= 0.022, \\ \sum_{C_{hl} \in N_1(3,1)} C_{31}^{hl} &= 0.019, & \sum_{C_{hl} \in N_1(3,2)} C_{32}^{hl} &= 0.021, & \sum_{C_{hl} \in N_1(3,3)} C_{33}^{hl} &= 0.014. \end{aligned}$$

Conditions (C1) – (C3) are satisfied for SICNN (4.6) with $\gamma = 0.3$, $\delta = 0.33$, $L_f = \sqrt[4]{27}/18$, $M_f = \pi/18$, $M_{11} = 1$, $M_{12} = 0.6$, $M_{13} = 0.8$, $M_{21} = 0.7$, $M_{22} = 0.3$, $M_{23} = 0.8$, $M_{31} = 0.9$, $M_{32} = 0.8$, $M_{33} = 0.7$, and $K_0 = 5.0979$. Using the results of [29] one can confirm that for each periodic sequence $\zeta = \{\zeta_k\}_{k \in \mathbb{Z}}$, SICNN (4.6) possesses a unique quasi-periodic output.

In order to demonstrate the existence of a homoclinic motion in the dynamics of (4.6), let us take $\mu = 3.9$. The orbit $\eta = \{\dots, H_{3.9}^3(\eta_0), H_{3.9}^2(\eta_0), H_{3.9}(\eta_0), \eta_0, F_{3.9}(\eta_0), F_{3.9}^2(\eta_0), F_{3.9}^3(\eta_0), \dots\}$, where $\eta_0 = 1/3.9$, is homoclinic to the fixed point $\zeta^* = 2.9/3.9$ of (4.7) [47]. Let us denote by $\phi_\eta(t) = \{\phi_\eta^{ij}(t)\}$ and $\phi_{\zeta^*}(t) = \{\phi_{\zeta^*}^{ij}(t)\}$ the bounded solutions of SICNN (4.6) corresponding to η and ζ^* , respectively. According to Theorem 3.1, the output $\phi_\eta(t)$ is homoclinic to the quasi-periodic output $\phi_{\zeta^*}(t)$. The graphs of the outputs $\phi_\eta^{22}(t)$ and $\phi_{\zeta^*}^{22}(t)$ corresponding to the cell C_{22} of (4.6) are represented in Figure 1. The output $\phi_\eta^{22}(t)$ is in blue color, whereas $\phi_{\zeta^*}^{22}(t)$ is in red color. The outputs corresponding to the remaining cells, which are not just pictured here, exhibit similar homoclinic behavior. Figure 1 manifests that $\|\phi_\eta(t) - \phi_{\zeta^*}(t)\| \rightarrow 0$ as $t \rightarrow \pm\infty$, i.e., a homoclinic motion takes place in the dynamics of (4.6).




Figure 1: Homoclinic output of SICNN (4.6). The outputs $\phi_\eta^{22}(t)$ and $\phi_{\zeta^*}^{22}(t)$ corresponding to the cell C_{22} are represented in blue and red colors, respectively. The simulation confirms that $\phi_\eta(t)$ is homoclinic to the quasi-periodic output $\phi_{\zeta^*}(t)$.

Now, we will demonstrate the presence of a heteroclinic motion of (4.6). For that purpose, let us take into account the network (4.6) with the value of the parameter $\mu = 4$. It was shown in [47] that the orbit $\tilde{\eta} = \{\dots, G_4^3(\tilde{\eta}_0), G_4^2(\tilde{\eta}_0), G_4(\tilde{\eta}_0), \tilde{\eta}_0, F_4(\tilde{\eta}_0), F_4^2(\tilde{\eta}_0), F_4^3(\tilde{\eta}_0), \dots\}$, where $\tilde{\eta}_0 = 1/4$, is heteroclinic to the fixed points $\zeta^1 = 3/4$ and $\zeta^2 = 0$ of the logistic map (4.7). Let $\phi_{\tilde{\eta}}(t) = \{\phi_{\tilde{\eta}}^{ij}(t)\}$, $\phi_{\zeta^1}(t) = \{\phi_{\zeta^1}^{ij}(t)\}$ and $\phi_{\zeta^2}(t) = \{\phi_{\zeta^2}^{ij}(t)\}$ be the bounded solutions of SICNN (4.6) corresponding to the orbits $\tilde{\eta}$, ζ^1 and ζ^2 , respectively. It is worth noting that $\phi_{\zeta^1}(t)$ and $\phi_{\zeta^2}(t)$ are quasi-periodic. According to Theorem 3.1, $\phi_{\tilde{\eta}}(t)$ is heteroclinic to the solutions $\phi_{\zeta^1}(t)$ and $\phi_{\zeta^2}(t)$. The graphs of $\phi_{\tilde{\eta}}^{22}(t)$, $\phi_{\zeta^1}^{22}(t)$ and $\phi_{\zeta^2}^{22}(t)$ are depicted in Figure 2 in blue, red and green colors, respectively. Figure 2 supports the result of Theorem 3.1 such that SICNN (4.6) possesses a heteroclinic motion.




Figure 2: Heteroclinic output of SICNN (4.6). The outputs $\phi_{\bar{\eta}}^{22}(t)$, $\phi_{\zeta_1}^{22}(t)$ and $\phi_{\zeta_2}^{22}(t)$ corresponding to the cell C_{22} are represented in blue, red and green colors, respectively. It is seen in the figure that $\phi_{\bar{\eta}}(t)$ is heteroclinic to the quasi-periodic outputs $\phi_{\zeta_1}(t)$ and $\phi_{\zeta_2}(t)$.

5 Conclusions

In this paper, we theoretically prove the existence of homoclinic and heteroclinic outputs in SICNNs whose external inputs are generated by a discrete map. Continuous as well as discontinuous inputs are utilized in the model under investigation. The definitions of homoclinic and heteroclinic motions for the multidimensional dynamics of SICNNs are provided in the functional sense. The presented example reveals that quasi-periodic as well as homoclinic and heteroclinic motions can coexist in the dynamics of SICNNs with discontinuous external inputs. An advantage of the homoclinic and heteroclinic motions obtained in our study is that they can be simulated numerically. Our results can be easily adapted to other types of recurrent neural networks such as Hopfield [48] and Cohen-Grossberg [49] neural networks, and the presented technique may be useful for the mathematical investigation of chaotic behavior in brain dynamics and the rest of the nervous system.

References

- [1] Andersson KG (1994) Poincaré's discovery of homoclinic points. Arch Hist Exact Sci 48:133-147
- [2] Gonchenko SV, Shil'nikov LP, Turaev DV (1996) Dynamical phenomena in systems with structurally unstable Poincaré homoclinic orbits. Chaos 6:15-31
- [3] Shil'nikov LP (1967) On a Poincaré-Birkhoff problem. Math. USSR-Sbornik 3:353-371
- [4] Smale S (1965) Diffeomorphisms with many periodic points. In: Cairns SS (ed) Differential and Combinatorial Topology: A Symposium in Honor of Marston Morse, Princeton University Press, Princeton, NJ, pp 63-70
- [5] Bertozzi AL (1988) Heteroclinic orbits and chaotic dynamics in planar fluid flows. Siam J Math Anal 19:1271-1294
- [6] Chacón R, Bejarano JD (1995) Homoclinic and heteroclinic chaos in a triple-well oscillator. J Sound Vib 186:269-278

- [7] Bouzerdoum A, Pinter RB (1993) Shunting inhibitory cellular neural networks: Derivation and stability analysis. *IEEE Trans Circuits Syst I, Fundam Theory Appl* 40:215-221
- [8] Bouzerdoum A, Pinter RB (1990) A shunting inhibitory motion detector that can account for the functional characteristics of fly motion-sensitive interneurons. In: *Proceedings of IJCNN International Joint Conference on Neural Networks, San Diego, CA, USA*, pp 149-153
- [9] Bouzerdoum A, Nabet B, Pinter RB (1991) Analysis and analog implementation of directionally sensitive shunting inhibitory cellular neural networks. In: *Visual information processing: from neurons to chips, Proc. SPIE 1473*, pp 29-38
- [10] Bouzerdoum A, Pinter RB (1992) Nonlinear lateral inhibition applied to motion detection in the fly visual system. In: Pinter RB, Nabet B (ed) *Nonlinear Vision, Boca Raton, CRC Press*, pp 423-450
- [11] Carpenter GA, Grossberg S (1988) The ART of adaptive pattern recognition by a self-organizing neural network. *Computer* 21:77-88
- [12] Cheung HN, Bouzerdoum A, Newland W (1999) Properties of shunting inhibitory cellular neural networks for colour image enhancement. In: *Proceedings of 6th International Conference on Neural Information Processing, Perth, Western Australia*, pp 1219-1223
- [13] Fukushima K (1989) Analysis of the process of visual pattern recognition by the neocognitron. *Neural Netw* 2:413-420
- [14] Jernigan ME, McLean GF (1992) Lateral inhibition and image processing. In: Pinter RB, B. Nabet B (ed) *Nonlinear Vision, Boca Raton, FL, CRC Press*, pp 451-462
- [15] Pinter RB, Olberg RM, Warrant E (1989) Luminance adaptation of preferred object size in identified dragonfly movement detectors. In: *Proceedings of IEEE International Conference on Systems, Man and Cybernetics*, pp 682-686
- [16] Peng G, Huang L (2009) Anti-periodic solutions for shunting inhibitory cellular neural networks with continuously distributed delays. *Nonlinear Anal Real World Appl* 10:2434-2440
- [17] Chen A, Cao J (2002) Almost periodic solution of shunting inhibitory CNNs with delays. *Phys Lett A* 298:161-170
- [18] Shao J (2008) Anti-periodic solutions for shunting inhibitory cellular neural networks with time-varying delays. *Phys Lett A* 372:5011-5016
- [19] Cai M, Zhang H, Yuan Z (2008) Positive almost periodic solutions for shunting inhibitory cellular neural networks with time-varying delays. *Math Comput Simulation* 78:548-558

- [20] Meng H, Li Y (2008) New convergence behavior of shunting inhibitory cellular neural networks with time-varying coefficients. *Appl Math Lett* 21:717-721
- [21] Chen Z (2013) A shunting inhibitory cellular neural network with leakage delays and continuously distributed delays of neutral type. *Neural Comput & Applic* 23:2429-2434
- [22] Ou C (2009) Almost periodic solutions for shunting inhibitory cellular neural networks. *Nonlinear Anal Real World Appl* 10:2652-2658
- [23] Peng L, Wang W (2013) Anti-periodic solutions for shunting inhibitory cellular neural networks with time-varying delays in leakage terms. *Neurocomputing* 111:27-33
- [24] Wang P, Li B, Li Y (2015) Square-mean almost periodic solutions for impulsive stochastic shunting inhibitory cellular neural networks with delays. *Neurocomputing* 167:76-82
- [25] Liu X (2015) Exponential convergence of SICNNs with delay and oscillating coefficients in leakage terms. *Neurocomputing* 168:500-504
- [26] Gui Z, Ge W (2006) Periodic solution and chaotic strange attractor for shunting inhibitory cellular neural networks with impulses. *Chaos* 16:033116
- [27] Akhmet MU, Fen MO (2013) Shunting inhibitory cellular neural networks with chaotic external inputs. *Chaos* 23:023112
- [28] Akhmet MU, Fen MO (2015) Attraction of Li-Yorke chaos by retarded SICNNs. *Neurocomputing* 147:330-342
- [29] Akhmet M, Fen MO, Kivılcım A (2016) Li-Yorke chaos generation by SICNNs with chaotic/almost periodic postsynaptic currents. *Neurocomputing* 173:580-594
- [30] Fen MO, Akhmet M (2016) Impulsive SICNNs with chaotic postsynaptic currents. *Discrete Contin Dyn Syst Ser B* 21:1119-1148
- [31] Akhmet MU (2010) Homoclinical structure of the chaotic attractor. *Commun Nonlinear Sci Numer Simulat* 15:819-822
- [32] Akhmet MU (2009) Devaney's chaos of a relay system. *Commun Nonlinear Sci Numer Simulat* 14:1486-1493
- [33] Akhmet M, Fen MO (2016) *Replication of Chaos in Neural Networks, Economics and Physics*. Springer-Verlag, Berlin, Heidelberg
- [34] Akhmet MU (2008) Hyperbolic sets of impact systems. *Dyn Contin Discrete Impuls Syst Ser A Math. Anal.* 15 (Suppl. S1):1-2. Proceedings of the 5th International Conference on Impulsive and Hybrid Dynamical Systems and Applications, Beijing, Watan Press

- [35] Fen MO, Tokmak Fen F, Homoclinic and heteroclinic motions in hybrid systems with impacts. *Mathematica Slovaca* (accepted)
- [36] Akhmet M, Fen MO (2016) Homoclinic and heteroclinic motions in economic models with exogenous shocks. *Applied Mathematics and Nonlinear Sciences* 1:1-10
- [37] Chen SS (2009) Delayed transiently chaotic neural networks and their application. *Chaos* 19:033125
- [38] Li Q, Tang S, Zeng H, Zhou T (2014) On hyperchaos in a small memristive neural network. *Nonlinear Dyn* 78:1087-1099
- [39] Zou F, Katérle A, Nossek JA (1993) Homoclinic and heteroclinic orbits of the three-cell cellular neural networks. *IEEE Trans Circuits Syst I, Fundam Theory Appl* 40:843-848
- [40] Tyagi BK, Mehotra RK (1976) Special type of emitter-coupled rectangular pulse generator. In: *IEE-IERE Proceedings, India*, pp 223-227
- [41] Ping W, Jiali B, Hong W, Huiping W (2003) Multi-pulse Generator for electroporation. In: *Proceedings of the 25th Annual International Conference of the IEEE EMBS, Cancun, Mexico, September 17-21*, pp 2970-2973
- [42] Filippov AF (1988) *Differential Equations with Discontinuous Righthand Sides: Control Systems*. Springer, The Netherlands
- [43] Akhmet M (2011) *Nonlinear Hybrid Continuous/Discrete-Time Models*. Atlantis Press, Paris
- [44] Huang X, Cao J (2003) Almost periodic solution of shunting inhibitory cellular neural networks with time-varying delay. *Phys Lett A* 314:222-231
- [45] Corduneanu C (1971) *Principles of Differential and Integral Equations*. Allyn and Bacon, Inc., Boston
- [46] Hale J, Koçak H (1991) *Dynamics and Bifurcations*. Springer-Verlag, New York
- [47] Avrutin V, Schenke B, Gardini L (2015) Calculation of homoclinic and heteroclinic orbits in 1D maps. *Commun Nonlinear Sci Numer Simulat* 22:1201-1214
- [48] Hopfield JJ (1984) Neurons with graded response have collective computational properties like those of two-state neurons. *Proc Natl Acad Sci USA* 81:3088-3092
- [49] Cohen MA, Grossberg S (1993) Absolute stability of global pattern formation and parallel memory storage by competitive neural networks. *IEEE Trans Syst Man Cybern SMC* 13:815-826