

Dynamical Maps, Quantum Detailed Balance and Petz Recovery Map

Álvaro M. Alhambra^{1,*} and Mischa P. Woods^{1,2,†}

¹*Department of Physics and Astronomy, University College London,
Gower Street, London WC1E 6BT, United Kingdom*

²*QuTech, Delft University of Technology, Lorentzweg 1, 2611 CJ Delft, Netherlands*

Markovian master equations (formally known as quantum dynamical semigroups) can be used to describe the evolution of a quantum state ρ when in contact with a memoryless thermal bath. This approach has had much success in describing the dynamics of real-life open quantum systems in the lab. Such dynamics increase the entropy of the state ρ and the bath until both systems reach thermal equilibrium, at which point entropy production stops. Our main result is to show that the entropy production at time t is bounded by the relative entropy between the original state and the state at time $2t$. The bound puts strong constraints on how quickly a state can thermalise, and we conjecture that the factor of 2 is tight. The proof makes use of a key physically relevant property of these dynamical semigroups – detailed balance, showing that this property is intimately connected with the field of recovery maps from quantum information theory. We envisage that the connections made here between the two fields will have further applications. We also use this connection to show that a similar relation can be derived when the fixed point is not thermal.

I. INTRODUCTION

It is very often observed in nature that physical systems relax to an equilibrium state. This phenomenon, which has very evident consequences at the macroscopic scales of our everyday experience, ultimately relies on the dynamics of the microscopic components. This fact was understood in the early days of statistical mechanics, and since then a large amount of work has been produced with the aim of trying to understand how exactly physical systems reach thermal equilibrium.

Any such evolution will be ultimately generated through some reversible dynamics on a large composite system, that is effectively irreversible as seen by a smaller part of that composite system. This irreversibility means that, in a coarse-grained sense, entropy will be produced throughout the process. The entropy production can be linked to the fact that correlations between a big thermal object (a heat bath) and one smaller subsystem S are increasingly harder to access, which forces the coarse-graining of the description [1]. Intuitively, the more irreversible a process is, the more entropy is produced, and the closer a particular system will be to equilibrium.

In this work we look at a commonly used family of quantum evolutions that model the dynamics of a system weakly coupled to a thermal bath, and show explicitly how the amount of entropy produced along a particular evolution is related to how much a state changes along that evolution. These maps were first studied by Davies [2] and are a quantum generalization of the classical Glauber dynamics. They are Markovian, and have the thermal state as a fixed point of the evolution.

In the limit of a large thermal bath, the total entropy produced by such a process is given by how much the

free energy of a system decreases with time [3]. The free energy for a state $\rho_S(t)$ at time t is defined as

$$F_\beta(\rho_S(t)) = \text{Tr}[\hat{H}_S \rho_S(t)] + \frac{1}{\beta} \text{Tr}[\rho_S(t) \log \rho_S(t)], \quad (1)$$

where \hat{H}_S is the Hamiltonian of the subsystem of interest, and β^{-1} is the temperature of the bath. This is a consequence of the fact that *i)* the dynamics has the thermal state of the system $\tau_S = \frac{e^{-\beta \hat{H}_S}}{\mathcal{Z}}$ as a fixed point and *ii)* the quantum relative entropy is contractive under quantum evolutions. More explicitly, for an evolution from time $t = 0$ to t , the total amount of entropy produced is given by $F_\beta(\rho_S(0)) - F_\beta(\rho_S(t))$, which is always positive.

Our main result is Theorem 2, which states that under the condition that the interaction between system and bath is time-independent, we can lower-bound the entropy production at time t by the state at time $2t$.

This sharpens some intuitive notions, namely that if not much entropy is produced during a time interval Δt , the state will not change very much during the time interval $2\Delta t$, but if it does, then a large amount of entropy must have been produced at an earlier time, namely during the time interval Δt . This relationship between the entropy production at an earlier time to the state at a later time is not due to any memory effects, since the dynamics is purely Markovian. Moreover, as we will explain, the key ingredient is a general property of these Davies maps, namely *quantum detailed balance*.

The bound proves a physically relevant particular case of an open conjecture about general quantum maps first formulated in [4]. The strongest possible version of the conjecture is known to not be true in full generality [5], although it has been shown for particular sets such as unital maps [6], classical stochastic matrices [4], catalytic thermal operations [7] and we here show it for Davies maps. All these results relate the decrease of relative entropy with a measure of how well a given pair of states

* alvaro.alhambra.14@ucl.ac.uk

† mischa.woods@gmail.com

can be recovered through a particular *recovery map*, and are generalizations of an early result by Petz [8]. For the best results up to date on general quantum maps, see [9–12].

This line of research was very much inspired by the study of quantum conditional mutual information and *quantum Markov chains* [13], where recent results put lower bounds on the quantum conditional mutual information with measures of how well the operation of tracing out a subsystem within a tripartite state can be undone. See [14] for an early breakthrough, and [5, 15–17] for further refinements. Recovery maps have found many applications in quantum information theory in general, such as coding theorems [18], approximate error correction [19] or asymmetry [20]. They also appear in the derivation of quantum fluctuation theorems [21, 22].

Our results, inspired by findings in quantum information theory, are a consequence of the observation that if a dynamical map satisfies quantum detailed balance, a property of thermodynamical processes, then this implies that the map is its own recovery map. The connection between information theory and thermodynamics goes back a long way, to the seminal work of Landauer [23] and has furthered our understanding of both significantly. Within the current surge of information-theory approaches to quantum thermodynamics (see [24] for a review), our result provides another example of how ideas from one may find definite applications in the other.

We shall first introduce Davies maps, outline their properties, and explain their entropy production. This is followed by the statement of main result and a discussion on the bound it self. We finally conclude with some suggestions for open questions. The technical results have all been placed in the appendix, where we also include a discussion on Davies maps and of the bound in the infinitesimal time limit.

II. DAVIES MAPS AND ENTROPY PRODUCTION

Davies maps are a particular set of quantum dynamical semigroups that describe the evolution of a system on a d_S dimensional Hilbert space that is weakly interacting with a heat bath. The first rigorous derivation of their form was given in [2] (see [25, 26] for more modern treatments). As they are time-continuous quantum semigroups, their generator takes the form of a Lindbladian operator, which we define as

$$\frac{d\rho_S(t)}{dt} = \mathcal{L}(\rho_S(t)) + i\theta(\rho_S(t)), \quad (2)$$

where \mathcal{L} is called the *Lindbladian* and $\theta(\cdot) = -[H_{\text{eff}}, \cdot]$ is called the unitary part, with H_{eff} the effective Hamiltonian. The solution is a one-parameter family of CPTP maps $M_\Delta(\cdot)$, $\Delta \geq 0$ which governs the dynamics, $M_\Delta(\rho(t)) = \rho(t + \Delta)$. We will not delve into the full

details here, but instead highlight the important properties the canonical form of Davies maps, denoted $T_t(\cdot)$, possess:

- 1) They arise from the weak system-bath coupling limit
- 2) They can be written in the form $T_t(\cdot) = e^{it\theta + t\mathcal{L}}(\cdot)$, with θ and \mathcal{L} time independent
- 3) θ and \mathcal{L} commute: $\theta(\mathcal{L}(\cdot)) = \mathcal{L}(\theta(\cdot))$
- 4) They have a thermal fixed point: $T_t(\tau_S) = \tau_S$, where τ_S is the Gibbs state of the system at temperature T_S .
- 5) Their Lindbladians and unitary part satisfy *Quantum detailed balance* (QDB):

$$\langle A, \mathcal{L}^\dagger(B) \rangle_\Omega = \langle \mathcal{L}^\dagger(A), B \rangle_\Omega, \quad (3)$$

$$[H_{\text{eff}}, \Omega] = 0, \quad (4)$$

for all $A, B \in \mathbb{C}^{d_S \times d_S}$, where \mathcal{L}^\dagger is the adjoint Lindbladian and in the case of Davies maps, $\Omega = \tau_S$. The scalar product in Eq. (3) is defined as

$$\langle A, B \rangle_\Omega := \text{Tr}[\Omega^{1/2} A^\dagger \Omega^{1/2} B]. \quad (5)$$

This is sometimes referred to as *reversibility* or KMS condition. It is stronger than 4), since it has as a consequence that Ω is the fixed point, $\mathcal{L}(\Omega) = 0$.

In Appendix A we give a more detailed account of the microscopic origin of these maps, and of the form of the weak coupling limit, property 1). In the literature, there are various different definitions of QDB which are in general not equivalent. We show in Appendix C, that for maps satisfying time translation symmetry, such as Davies maps, definition 5) is equivalent to the definition of QDB in [25, 27].

In addition to the properties above, it is sometimes assumed that:

- 6) The dynamics associated with Davies maps converge to the fixed point, $\lim_{t \rightarrow \infty} T_t(\rho_S(0)) = \tau_S$.

Such convergence is guaranteed if more stringent conditions are imposed on the Davies map [28–31]. We will *not* need to assume 6) here.

Since we wish to bound the distance from the state at time t to the fixed point, we need a distance measure. For this we use the *relative entropy* $D(\rho||\sigma) = \text{Tr}[\rho(\log \rho - \log \sigma)]$. This measure is meaningful since it is non-negative, zero iff $\rho = \sigma$, and is contractive under CPTP maps. For the special case that σ is a Gibbs state, it has an interpretation in terms of a free energy,

$$D(\rho(t)||\tau_S) = \beta F_\beta(\rho_S(t)) - \log Z_S, \quad (6)$$

where $Z_S = \text{Tr}[e^{-\beta H_S}]$ is the partition function of the system, which we assume is constant. We can thus write

$$\begin{aligned} D(\rho(0)||\tau_S) - D(\rho(t)||\tau_S) &= \beta (F_\beta(\rho_S(0)) - F_\beta(\rho_S(t))) \\ &= \beta \Delta E - \Delta S, \end{aligned} \quad (7)$$

with ΔE , ΔS the changes in mean energy and entropy of the system. Due to the contractivity property of D , Eq. (7) is non-negative and non-decreasing with $t \geq 0$. The l.h.s. of Eq. (7) is referred to in the literature as the *entropy production*. The reason for this name is as follows. For a large thermal reservoir, small changes of energy (that is, heat transferred to the system) are proportional to changes of entropy in it, with proportionality constant $\frac{1}{\beta}$. Hence, we can identify the change in energy in the system with a change of entropy in the reservoir $\beta \Delta E \simeq -\Delta S_{\text{bath}}$, so that the difference in free energy of the system for a time interval Δt is equal to the total entropy generated during the interval Δt in system and bath. Therefore, Eq. (7) constitutes a natural measure of the irreversibility of the process. Furthermore, as one intuitively might expect, the entropy production only depends on the dissipative part of the dynamics, as we explain in the Section A3 of the appendix. Therefore, we will assume for simplicity that $\theta = 0$ in the next Section unless stated otherwise.

If one were to change the initial state of the environment for the maximally mixed state, then the system can only exchange entropy, but not heat/energy with it. These correspond to unital maps, in which case the free energy is replaced with the entropy gain of the system alone. In that case, a lower bound on the entropy they produce in terms of the adjoint of the unital map can be found in [6].

III. MAIN RESULTS

Our main result is a tight lower bound on the change of free energy, or entropy produced, for a finite time. For Davies maps, the following lemma holds:

Lemma 1. All Davies maps $T_t(\cdot)$, satisfy the inequality

$$F_\beta(\rho_S(0)) - F_\beta(\rho_S(t)) \geq \frac{1}{\beta} D(\rho_S(0)||\tilde{T}_t(\rho_S(t))), \quad (8)$$

where $\tilde{T}_t(\cdot)$ is the *time-reversed* map or Petz recovery map, defined as

$$\tilde{T}_t(\cdot) = \tau_S^{1/2} T_t^\dagger \left(\tau_S^{-1/2}(\cdot) \tau_S^{-1/2} \right) \tau_S^{1/2}, \quad (9)$$

with T_t^\dagger denoting the adjoint of T_t .

Proof. See Appendix A2 □

For Lemma 1 to hold, only properties 1) and 4) are required. In addition, we find that there is a striking connection between property 4) and the Petz recovery map

which we will now explain. A quantum dynamical semi-group M_t which obeys QDB has a Petz recovery map \tilde{M}_t which is equal to the map itself $\tilde{M}_t = M_t$ (See Theorem 8 in Appendix). Petz derived his famous recovery map in 1986 [8] while the first appearance of the detailed balance condition goes back at least to the work of Boltzmann in 1872 [32] and quantum detailed balance to Alicki in 1976 [33]. To the best of the authors knowledge, this connection between results from the communities of quantum information theory and quantum dynamical semi-groups was previously unknown.

The classical definition of detailed balance, in terms of the transition probabilities $p(j|i)$ of a classical Master equation, implies that, at equilibrium, a particular jump between energy levels $E_i \rightarrow E_j$ has the same total probability as the opposite jump $E_j \rightarrow E_i$, such that $p(j|i) \frac{e^{-\beta E_i}}{Z} = p(i|j) \frac{e^{-\beta E_j}}{Z}$. The condition in Eq. (3) is the most natural quantum generalization of that (although as shown in [34] different ones are also possible). In that sense, QDB can be understood as the fact that a particular thermalization process coincides with its own time-reversed map, which is defined as in Eq. (9) (for more details, see e.g. [35]).

On the other hand, the Petz recovery map $\tilde{\Gamma}(\cdot)$, given a state σ and a CPTP map $\Gamma(\cdot)$, is formally defined as [8, 36, 37]

$$\tilde{\Gamma}(\cdot) = \sigma^{1/2} \Gamma^\dagger \left(\Gamma(\sigma)^{-1/2}(\cdot) \Gamma(\sigma)^{-1/2} \right) \sigma^{1/2}. \quad (10)$$

This map is such that we have that iff $D(\rho||\sigma) = D(\Gamma(\rho)||\Gamma(\sigma))$ then $\tilde{\Gamma}(\Gamma(\rho)) = \rho$ and $\tilde{\Gamma}(\Gamma(\sigma)) = \sigma$. It appears in quantum information theory when one tries to find the best possible way to recover data after it is processed [38, 39].

We can hence rewrite Lemma 1 as

Theorem 2. All Davies maps $T_t(\cdot)$, satisfy the inequality

$$F_\beta(\rho_S(0)) - F_\beta(\rho_S(t)) \geq \frac{1}{\beta} D(\rho_S(0)||\rho_S(2t)). \quad (11)$$

Proof. See Appendix A3. □

In addition to assuming detailed balance, condition 5), we have also used condition 2). If the Lindbladian \mathcal{L} is time dependent, i.e. 2) is not satisfied, Eq. (11) holds but with $\rho_S(2t)$ replaced with $T_t(\rho(t))$.

While, as mentioned at the end of Section II, entropy production is invariant under a change in the unitary part of the dynamics, it is interesting to find the Petz recovery map when θ is not set to zero. We show in Lemma 9 in the appendix, that the Petz recovery map $\tilde{M}_t(\cdot)$ of a map $M_t(\cdot)$ satisfying QDB and for which \mathcal{L} and θ commute [property 3) of Davies maps], reverses the unitary part of the dynamics, while keeping the same dissipative part, that is

$$\tilde{M}_t(\cdot) = e^{-it\theta + t\mathcal{L}}(\cdot), \quad (12)$$




FIG. 1. An example of the inequality in Theorem 2 for a Davies map on a qubit given in [40]. The solid (blue) curve is the amount of entropy produced $\beta F_\beta(\rho(0)) - \beta F_\beta(\rho(t))$ and the dashed (purple) the lower bound $D(\rho_S(0)||\rho_S(2t))$. It can be seen how the lower bound at $t = 0$ starts at zero, and how for large times the two curves quickly converge to the total amount of entropy produced $D(\rho_S(0)||\tau_S)$.

and thus $\tilde{T}_t(T_t(\cdot)) = e^{2t\mathcal{L}}(\cdot)$. So not only is the l.h.s. of Eq. 11 invariant under a change in the unitary part of the dynamics; but so is the r.h.s.

In Fig. 1 we show a simple example of the inequality for the case of Davies maps applied on a qubit. Eq. (11) is tight at $t = 0$ and also in the large time limit, as long as condition 6) is satisfied. In this limit, the total entropy that has been produced is equal to $\frac{1}{\beta} D(\rho(0)||\tau_S)$, which both sides of the inequality approach as $\rho_S \rightarrow \tau_S$.

On the other hand, for very short times, the lower bound becomes trivial. In particular, in Appendix A 4 we show what both sides of the inequality tend to in the limit of infinitesimal time transformations. The entropy production becomes a *rate*, and the lower bound to it approaches 0. The latter can be seen in Fig. 1 by the fact that the dashed curve has a stationary point at the origin.

Non-trivial lower bounds on the rate of entropy production, in the form of log-Sobolev inequalities [41] can be used to derive bounds on the time it takes to converge to equilibrium for particular instances of Davies maps. Hence, given that Theorem 2 is completely general, and holds also for Davies maps that do not reach thermal equilibrium efficiently, the fact that the lower bound vanishes for infinitesimal times is not surprising.

Recall that the factor of 2 in Eq. (11) is a consequence of the observation that the Petz recovery map is equal to the map itself. A natural question is then, *is the factor 2 fundamental?* The following conjecture suggests it is:

Conjecture 3. [Tightness of entropy production bound] The largest constant $k \geq 0$ such that

$$F_\beta(\rho_S(0)) - F_\beta(\rho_S(t)) \geq \frac{1}{\beta} D(\rho_S(0)||\rho_S(kt)) \quad (13)$$

holds for all Davies maps is $k = 2$.

Violations of Eq. (13) have been found numerically for $k = 2.0001$, so the conjecture is highly likely to be true.




FIG. 2. Examples plots for Conjecture 3 for the Davies map for qubits from [40]. The solid (blue) curve is the amount of entropy produced $\beta F_\beta(\rho(0)) - \beta F_\beta(\rho(t))$ (l.h.s. of Eq. (13)) while the dashed lines correspond to $D(\rho(0)||\rho(kt))$ (r.h.s. of Eq. (13)) for different k . We see that when the constant k is greater than 2 the bound does not hold any more, showing that the $k = 2$ case is indeed special. For $k < 2$ the bound holds intuitively (given that it holds for $k = 2$), but results in a worse bound. This shows that the constraint set by Eq. (11) reflects a special feature of how Davies maps thermalize. Moreover, we see that a $k > 2$ would predict (incorrectly) a faster thermalisation rate, thus confirming that Eq. (11) is an implicit universal bound on the rate of thermalisation for Davies maps.

See Fig. 2 for more details. This means that Eq. (2) is likely to be the strongest constraint of its kind that Davies maps obey, and it hence sets a somewhat strong relation between how much the free energy and the systems state at a later time change during a thermalization process.

IV. BEYOND DAVIES MAPS

So far we have addressed the issue of universal bounds for Davies maps which constrain the speed of convergence to the thermal fixed point. We now turn our attention to what recent developments from quantum information theory can say about convergence of dynamical semi-groups in general. A recent advancement in quantum information is the development of *universal recoverability* maps [9, 10, 12]. By universal recoverability, it is meant that given a state σ and a CPTP map Γ , one can use the recovery map to lower bound the relative entropy difference $D(\rho||\sigma) - D(\Gamma(\rho)||\Gamma(\sigma))$ for *all* quantum states ρ . In general the lower bound takes on a complicated form (see Appendix B). However, for the case of dynamical semi-groups satisfying QDB and the following property, the bound is more explicit.

Let us assume that we have a one-parameter dynamical semi-group $M_t(\cdot)$ equipped with a fixed point Ω that satisfies a condition we call *Time-translation symmetry w.r.t. fixed point* (TTSFP),

$$\mathcal{L}(\cdot) = \Omega^{it} \mathcal{L}(\Omega^{-it}(\cdot)\Omega^{it}) \Omega^{-it} \quad \forall t \in \mathbb{R}. \quad (14)$$

This condition is satisfied for example by dynamical semigroups which arise naturally in the weak coupling limit or the low-density limit. Davies maps are one such example, but there are others [42].

The properties lead to the following result:

Theorem 4. Let the Quantum Dynamical Semigroup $M_t(\cdot)$ satisfy QDB and TTSFP. Then the following holds

$$D(\rho(0)||\Omega) - D(\rho(t)||\Omega) \geq -2 \log F(\rho, M_t(\rho(t))), \quad (15)$$

where $F(\rho, \sigma) = \text{Tr}[\sqrt{\sqrt{\sigma}\rho\sqrt{\sigma}}]$ is the quantum fidelity. Moreover, if the generators are time-independent we may write $M_t(\rho(t)) = \rho(2t)$.

It is well known that $D(\rho||\sigma) \geq -2 \log F(\rho||\sigma)$ with equality *only* for special instances. Therefore, for Davies maps, Eq. (15) is satisfied but with a weaker bound than Theorem 2.

V. CONCLUSION

We have proven a lower bound on the amount of entropy produced for a family of thermalization processes described by quantum dynamical semigroups. In order to do this, we have shown that an inequality which is not true for general CPTP maps holds for these maps. From that perspective, it would be interesting to investigate what are the features of these and other particular maps that make them obey this inequality, and whether such bounds have more interesting physical consequences. One potential application of this in open quantum systems, is to use a tightened monotonicity inequality to find when *information backflow* occurs in non-Markovian dynamics [43].

One of the important questions regarding Davies maps is how fast they converge to equilibrium. There have been several approaches to this question, mostly inspired

by their classical analogues, which include the computation of the spectral gaps [26, 34, 44] or the logarithmic-Sobolev inequalities [41, 45]. In particular we note that the latter take the form of lower bounds to the rate of entropy production of particular Davies maps, from which thermalization times can be derived. It would be interesting to know if the bound of Eq. (11), albeit true for all Davies maps no matter their thermalization times, may contain information about their asymptotic convergence. For instance, one could look at how fast is the inequality saturated in particular cases. We however leave this for future work.

One of the main features in the study of dynamical thermalisation processes, such as Davies maps, is quantum detailed balance; and we have shown that this condition can be written in terms of the Petz recovery map that appears in information theory. It is not completely clear what the physical origin of this coincidence is, and begs for a deeper understanding. The condition of detailed balance is ubiquitous in thermalization processes, and in particular, current algorithms for simulating thermal states on a quantum computer, such as the quantum Metropolis algorithm [46], obey it, which makes it all the more interesting. As such, the useful connection we establish here between the Petz recovery map and quantum detailed balance, is likely to have further implications for both thermodynamics and information theory.

ACKNOWLEDGMENTS

The authors would like to thank David Sutter, Michael Wolf, Nilanjana Datta, Mark Wilde and Toby Cubitt for helpful discussions. AA and MW acknowledge support from FQXi and EPSRC. This work was partially supported by the COST Action MP1209.

-
- [1] Massimiliano Esposito, Katja Lindenberg, and Christian Van den Broeck. Entropy production as correlation between system and reservoir. *New Journal of Physics*, 12(1):013013, 2010.
 - [2] E.B. Davies. Markovian master equations. *Commun. Math. Phys.*, 39:91 – 110, 1974.
 - [3] Herbert Spohn. Entropy production for quantum dynamical semigroups. *Journal of Mathematical Physics*, 19(5):1227–1230, 1978.
 - [4] Ke Li and Andreas Winter. Squashed entanglement, k-extendibility, quantum markov chains, and recovery maps. *arXiv preprint arXiv:1410.4184*, 2014.
 - [5] Fernando GSL Brandao, Aram W Harrow, Jonathan Oppenheim, and Sergii Strelchuk. Quantum conditional mutual information, reconstructed states, and state redistribution. *Physical review letters*, 115(5):050501, 2015.
 - [6] Francesco Buscemi, Siddhartha Das, and Mark M. Wilde. Approximate reversibility in the context of entropy gain, information gain, and complete positivity. *Phys. Rev. A*, 93:062314, Jun 2016.
 - [7] Stephanie Wehner, Mark M Wilde, and Mischa P Woods. Work and reversibility in quantum thermodynamics. *arXiv preprint arXiv:1506.08145*, 2015.
 - [8] Denes Petz. Sufficient subalgebras and the relative entropy of states of a von neumann algebra. *Communications in mathematical physics*, 105(1):123–131, 1986.
 - [9] Mark M Wilde. Recoverability in quantum information theory. In *Proc. R. Soc. A*, volume 471, page 20150338. The Royal Society, 2015.
 - [10] Marius Junge, Renato Renner, David Sutter, Mark M Wilde, and Andreas Winter. Universal recovery from a decrease of quantum relative entropy. *arXiv preprint arXiv:1509.07127*, 2015.

- [11] David Sutter, Marco Tomamichel, and Aram W Harrow. Strengthened monotonicity of relative entropy via pinched petz recovery map. *IEEE Transactions on Information Theory*, 62(5):2907–2913, 2016.
- [12] David Sutter, Mario Berta, and Marco Tomamichel. Multivariate trace inequalities. *arXiv preprint arXiv:1604.03023*, 2016.
- [13] Patrick Hayden, Richard Jozsa, Denes Petz, and Andreas Winter. Structure of states which satisfy strong subadditivity of quantum entropy with equality. *Communications in mathematical physics*, 246(2):359–374, 2004.
- [14] Omar Fawzi and Renato Renner. Quantum conditional mutual information and approximate markov chains. *Communications in Mathematical Physics*, 340(2):575–611, 2015.
- [15] David Sutter, Omar Fawzi, and Renato Renner. Universal recovery map for approximate markov chains. *Proc. R. Soc. A*, 2016, 472(2186), 2016.
- [16] Mario Berta and Marco Tomamichel. The fidelity of recovery is multiplicative. *IEEE Transactions on Information Theory*, 62(4):1758–1763, 2016.
- [17] Nilanjana Datta and Mark M Wilde. Quantum markov chains, sufficiency of quantum channels, and rényi information measures. *Journal of Physics A: Mathematical and Theoretical*, 48(50):505301, 2015.
- [18] Salman Beigi, Nilanjana Datta, and Felix Leditzky. Decoding quantum information via the Petz recovery map. *Journal of Mathematical Physics*, 57(8), 2016.
- [19] Hui Khoon Ng and Prabha Mandayam. Simple approach to approximate quantum error correction based on the transpose channel. *Physical Review A*, 81(6):062342, 2010.
- [20] Iman Marvian and Seth Lloyd. From clocks to cloners: Catalytic transformations under covariant operations and recoverability. *arXiv preprint arXiv:1608.07325*, 2016.
- [21] Johan Aberg. Fully quantum fluctuation theorems. *arXiv preprint arXiv:1601.01302*, 2016.
- [22] Álvaro M Alhambra, Lluís Masanes, Jonathan Oppenheim, and Christopher Perry. The second law of quantum thermodynamics as an equality. *arXiv preprint arXiv:1601.05799*, 2016.
- [23] R. Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5(3):183–191, July 1961.
- [24] John Goold, Marcus Huber, Arnau Riera, Lidia del Rio, and Paul Skrzypczyk. The role of quantum information in thermodynamics—a topical review. *Journal of Physics A: Mathematical and Theoretical*, 49(14):143001, 2016.
- [25] Robert Alicki and K Lendi. *Quantum Dynamical Semigroups and Applications*, volume 717. Springer Science & Business Media, 2007.
- [26] Kristan Temme. Lower bounds to the spectral gap of davis generators. *Journal of Mathematical Physics*, 54(12):122110, 2013.
- [27] Andrzej Kossakowski, Alberto Frigerio, Vittorio Gorini, and Maurizio Verri. Quantum detailed balance and kms condition. *Comm. Math. Phys.*, 57(2):97–110, 1977.
- [28] Herbert Spohn and Joel L. Lebowitz. *Irreversible Thermodynamics for Quantum Systems Weakly Coupled to Thermal Reservoirs*, pages 109–142. John Wiley & Sons, Inc., 2007.
- [29] Herbert Spohn. An algebraic condition for the approach to equilibrium of an open N-level system. *Letters in Mathematical Physics*, 2(1):33–38, 1977.
- [30] Alberto Frigerio. Quantum dynamical semigroups and approach to equilibrium. *Letters in Mathematical Physics*, 2(2):79–87, 1977.
- [31] Alberto Frigerio. Stationary states of quantum dynamical semigroups. *Communications in Mathematical Physics*, 63(3):269–276, 1978.
- [32] Ludwig Boltzmann. *Lectures on gas theory*; Berkeley. University of California Press, Berkeley, CA, 1964.
- [33] Robert Alicki. On the detailed balance condition for non-hamiltonian systems. *Reports on Mathematical Physics*, 10(2):249 – 258, 1976.
- [34] K Temme, Michael James Kastoryano, MB Ruskai, Michael Marc Wolf, and F Verstraete. The χ^2 -divergence and mixing times of quantum markov processes. *Journal of Mathematical Physics*, 51(12):122201, 2010.
- [35] Gavin E Crooks. Quantum operation time reversal. *Physical Review A*, 77(3):034101, 2008.
- [36] Dénes Petz. Sufficiency of channels over von neumann algebras. *The Quarterly Journal of Mathematics*, 39(1):97–108, 1988.
- [37] Dénes Petz. Monotonicity of quantum relative entropy revisited. *Reviews in Mathematical Physics*, 15(01):79–91, 2003.
- [38] Howard Barnum and Emanuel Knill. Reversing quantum dynamics with near-optimal quantum and classical fidelity. *Journal of Mathematical Physics*, 43(5):2097–2106, 2002.
- [39] Mark M Wilde. *Quantum information theory*. Cambridge University Press, 2013.
- [40] Wojciech Roga, Mark Fannes, and Karol Życzkowski. Davies maps for qubits and qutrits. *Reports on Mathematical Physics*, 66(3):311–329, 2010.
- [41] Michael J Kastoryano and Kristan Temme. Quantum logarithmic sobolev inequalities and rapid mixing. *Journal of Mathematical Physics*, 54(5):052202, 2013.
- [42] M. M. Wolf, J. Eisert, T. S. Cubitt, and J. I. Cirac. Assessing non-markovian quantum dynamics. *Phys. Rev. Lett.*, 101:150402, Oct 2008.
- [43] Elsi-Mari Laine, Jyrki Piilo, and Heinz-Peter Breuer. Measure for the non-markovianity of quantum processes. *Physical Review A*, 81(6):062115, 2010.
- [44] R Alicki, M Fannes, and M Horodecki. On thermalization in kitaev’s 2d model. *Journal of Physics A: Mathematical and Theoretical*, 42(6):065303, 2009.
- [45] Kristan Temme, Fernando Pastawski, and Michael J Kastoryano. Hypercontractivity of quasi-free quantum semigroups. *Journal of Physics A: Mathematical and Theoretical*, 47(40):405303, 2014.
- [46] K Temme, TJ Osborne, Karl G Vollbrecht, D Poulin, and Frank Verstraete. Quantum metropolis sampling. *Nature*, 471(7336):87–90, 2011.
- [47] E Brian Davies. Generators of dynamical semigroups. *Journal of Functional Analysis*, 34(3):421–432, 1979.
- [48] Piotr Źwikliński, Michał Studziński, Michał Horodecki, and Jonathan Oppenheim. Limitations on the evolution of quantum coherences: Towards fully quantum second laws of thermodynamics. *Physical review letters*, 115(21):210403, 2015.
- [49] Matteo Lostaglio. Elementary thermal operations and the universality of thermodynamic constraints. *arXiv preprint arXiv:1607.00394*, 2016.

- [50] Fernando G. S. L. Brandão, Michał Horodecki, Jonathan Oppenheim, Joseph M. Renes, and Robert W. Spekkens. Resource theory of quantum states out of thermal equilibrium. *Phys. Rev. Lett.*, 111:250404, Dec 2013.
- [51] F. Rellich. Perturbation theory of eigenvalue problems. *Lecture Notes reprinted by Gordon and Breach, 1968*, 1953.
- [52] T. Kato. Perturbation theory for linear operators, grundlehren. *SpringerVerlag, Berlin, New York*, 132, 1976.
- [53] M. Ohya and D. Petz. *Quantum Entropy and Its Use*. Theoretical and Mathematical Physics. Springer Berlin Heidelberg, 2004.
- [54] Heinz-Peter Breuer and Francesco Petruccione. *The theory of open quantum systems*. Oxford University Press on Demand, 2002.
- [55] Eric Carlen. Trace inequalities and quantum entropy: an introductory course. 2010.
- [56] Michael J Kastoryano and Fernando GSL Brandao. Quantum gibbs samplers: the commuting case. *Communications in Mathematical Physics*, pages 1–43.
- [57] J. Dereziński and F. Fruboes. *Fermi Golden Rule and open quantum systems, "Open Quantum Systems III Recent Developments"*. 2006.
- [58] Iman Marvian and Robert W Spekkens. Modes of asymmetry: the application of harmonic analysis to symmetric quantum dynamics and quantum reference frames. *Physical Review A*, 90(6):062110, 2014.
- [59] Matteo Lostaglio, Kamil Korzekwa, David Jennings, and Terry Rudolph. Quantum coherence, time-translation symmetry, and thermodynamics. *Physical Review X*, 5(2):021001, 2015.

Appendix A: Technical results

1. Davies maps and conditions for Lemma 1

Davies maps are derived from considering the dynamics of a state $\rho_S \in \mathcal{S}(\mathcal{H}_S)$, where \mathcal{H}_S is of finite dimension d_S , in contact with a thermal bath on an infinite dimensional Hilbert space \mathcal{H}_B . We will here specify the minimal assumptions about the bath and its interaction with the system necessary for the derivation of Lemma 5 and Lemma 1. In order to guarantee other properties, such as the existence of a fixed point or detailed balance, more subtle constraints are also necessary.

Let \hat{H}_B be a self-adjoint Hamiltonian on \mathcal{H}_B . Since we want states on \hat{H}_B to be thermodynamically stable, we assume that $Z_B = \text{Tr}[\exp(-\beta\hat{H}_B)] < \infty$ for all $\beta > 0$. \hat{H}_B must therefore have a purely discrete spectrum, which is bounded below and has no finite limit points; that is, there are only a finite number of energy levels in any finite interval ΔE . The quantum state $\rho_S \in \mathcal{S}(\mathcal{H}_S)$ with its free self-adjoint Hamiltonian \hat{H}_S of finite dimension interacts with the system via a bounded interaction term $\hat{I} \in \mathcal{B}(\mathcal{H}_S \otimes \mathcal{H}_B)$, with a parameter $\lambda > 0$ determining the interaction strength as follows

$$\hat{H}_{SB} = \hat{H}_S \otimes \mathbb{1}_B + \mathbb{1}_S \otimes \hat{H}_B + \lambda \hat{I}. \quad (\text{A1})$$

The initial state on $\mathcal{S}(\mathcal{H}_S \otimes \mathcal{H}_B)$ is assumed to be product, $\rho_S \otimes \tau_B$, with τ_B the Gibbs state at inverse temperature β . The dynamics of the system at time \tilde{t} is given by the unitary operator

$$U(\tilde{t}) := e^{-i\tilde{t}\hat{H}_{SB}} \quad (\text{A2})$$

after tracing out the environment. More precisely, by

$$\text{Tr}_B [U(\tilde{t})\rho_S \otimes \tau_B U^\dagger(\tilde{t})] \in \mathcal{S}(\mathcal{H}_S), \quad (\text{A3})$$

where U^\dagger denotes the adjoint of U .

The Davies map $T_t(\cdot)$ is defined by taking the limit that the interaction strength λ goes to zero, while the time \tilde{t} goes to infinity while maintaining $\tilde{t}\lambda^2 := t$ fixed. More concisely,

$$T_t(\cdot) = \lim_{\lambda \rightarrow 0^+} \text{Tr}_B [U(\tilde{t})(\cdot) \otimes \tau_B U^\dagger(\tilde{t})] \in \mathcal{S}(\mathcal{H}_S) \quad \text{subject to } \tilde{t}\lambda^2 = t \text{ fixed.} \quad (\text{A4})$$

It is assumed that in this limit $U(\tilde{t})$ and its inverse $U^\dagger(\tilde{t})$ are still unitary operators mapping states on $\mathcal{S}(\mathcal{H}_S \otimes \mathcal{H}_B)$ to states on $\mathcal{S}(\mathcal{H}_S \otimes \mathcal{H}_B)$. To gain more physical insight into this construction, we refer to [2, 28, 47]. We remind the reader that the conditions described in Section A 1 are not sufficient for the map $T_t(\cdot)$ to satisfy other properties, such as the convergence to a fixed point or detailed balance, more subtle constraints are also necessary. We will not go into the details of these additional conditions, since only sufficient (but perhaps not necessary) conditions are known, e.g. [2]. In other Sections, we will additionally take advantage of the known fact that Davies maps satisfy quantum detailed balance.

2. Proof and statement of Lemma 1

In order to prove the main theorem we need a lemma about Davies maps first. We show that in the weak coupling limit, correlations between the system and the environment (the bath) are not created if both start as initially uncorrelated thermal states. In order to do this, we will need to introduce a finite dimensional cut-off on \mathcal{H}_B and prove the results for the truncated space, and finally proving uniform convergence in the bath system size by removing the cut-off by taking the infinite dimensional limit. Let \hat{P}_n denote the projection onto a finite dimensional Hilbert Space $\mathcal{H}_{B,n} \subset \mathcal{H}_B$. Furthermore, assume that $\mathcal{H}_{B,1} \subset \mathcal{H}_{B,2} \subset \mathcal{H}_{B,3} \dots$ and that $\lim_{n \rightarrow \infty} \mathcal{H}_{B,n} = \mathcal{H}_B$. For concreteness (although not strictly necessary), one could let $\hat{P}_n = \sum_{k=0}^n |E_k\rangle\langle E_k|$ where $|E_0\rangle, |E_1\rangle, |E_2\rangle, \dots$ are the eigenvectors of \hat{H}_B ordered in increasing eigenvalue order.

We define the truncated self-adjoint Hamiltonians on \mathcal{H}_B as $\hat{H}_B^{(n)} = \hat{P}_n \hat{H}_B \hat{P}_n$ with a corresponding Gibbs state denoted by $\tau_{B,n} \in \mathcal{S}(\mathcal{H}_{B,n})$. Similarly, we construct unitaries on $\mathcal{H}_{B,n}$ by

$$U_n = \exp\left(-i\Delta\hat{H}_{SB}^{(n)}\right), \quad \hat{H}_{SB}^{(n)} = (\mathbb{1}_S \otimes \hat{P}_n) \hat{H}_{SB} (\mathbb{1}_S \otimes \hat{P}_n) \quad (\text{A5})$$

and define $\hat{I}_n := (\mathbb{1}_S \otimes \hat{P}_n) \hat{I} (\mathbb{1}_S \otimes \hat{P}_n)$. We recall the definition of the thermal state of the system $\tau_S \in \mathcal{S}(\mathcal{H}_S)$, which is given by

$$\tau_S = \frac{e^{\beta \hat{H}_S}}{Z_S}, \quad Z_S > 0 \quad (\text{A6})$$

for some inverse temperature $\beta > 0$

The lemma is the following:

Lemma 5 (Correlations at the fixed point). Let $\alpha > 0$, $\Delta \in \mathbb{R}$ and the constant $\tilde{Z}_{SB}^{n,\alpha} = \text{Tr}[(\tau_S \otimes \tau_{B,n})^\alpha]$. Then, for all $n \in \mathbb{N}^+$, we have the bound

$$\frac{1}{2} \|U_n(\tau_S \otimes \tau_{B,n})^\alpha U_n^\dagger - (\tau_S \otimes \tau_{B,n})^\alpha\|_1 \leq \tilde{Z}_{SB}^{n,\alpha} \beta \sqrt{\lambda \|\hat{I}_n\|}, \quad (\text{A7})$$

where $\tau_S, \tau_{B,n}$ are thermal states at inverse temperatures $\beta_S, \beta_{B,n}$ respectively, and $\|\cdot\|_1, \|\cdot\|$ is the one-norm and operator norm respectively.

Proof. The result is a consequence of mean energy conservation under the unitary transformation U_n and Pinsker's inequality.

Define the shorthand notation $\tilde{\tau}_{SB}^{n,\alpha} = U_n(\tau_S \otimes \tau_{B,n})^\alpha U_n^\dagger / \tilde{Z}_{SB}^{n,\alpha} \in \mathcal{S}(\mathcal{H}_S \otimes \mathcal{H}_{B,n})$ and $\tilde{Z}_{SB}^{n,\alpha} := \tilde{Z}_S^\alpha \tilde{Z}_B^{n,\alpha}$, $\tilde{Z}_S^\alpha := \text{Tr}[\tau_S^\alpha]$, $\tilde{Z}_B^{n,\alpha} := \text{Tr}[\tau_{B,n}^\alpha]$. By direct evaluation of the relative entropy,

$$D(\tilde{\tau}_{SB}^{n,\alpha} \| (\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha}) / \beta = \text{Tr}[\hat{H}_S \tilde{\tau}_{SB}^{n,\alpha}] + \text{Tr}[\hat{H}_B^{(n)} \tilde{\tau}_{B,n}^\alpha] - (\alpha\beta)^{-1} S(\tau_S^\alpha \otimes \tau_{B,n}^\alpha / \tilde{Z}_{SB}^{n,\alpha}) + \ln(\tilde{Z}_{SB}^{n,\alpha}), \quad (\text{A8})$$

where we have used unitary in-variance of the von Neumann entropy $S(\cdot)$. Thus since

$$0 = D((\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha} \| (\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha}) / \beta \quad (\text{A9})$$

$$= \text{Tr}[\hat{H}_S \tau_S^\alpha / \tilde{Z}_S^\alpha] + \text{Tr}[\hat{H}_B^{(n)} \tau_{B,n}^\alpha / \tilde{Z}_B^{n,\alpha}] - (\alpha\beta)^{-1} S(\tau_S^\alpha \otimes \tau_{B,n}^\alpha / \tilde{Z}_{SB}^{n,\alpha}) + \ln(\tilde{Z}_{SB}^{n,\alpha}), \quad (\text{A10})$$

we conclude

$$D(\tilde{\tau}_{SB}^{n,\alpha} \| (\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha}) / \beta = \text{Tr}[\hat{H}_S \tilde{\tau}_{SB}^{n,\alpha}] + \text{Tr}[\hat{H}_B^{(n)} \tilde{\tau}_{B,n}^\alpha] - \text{Tr}[\hat{H}_S \tau_S^\alpha / \tilde{Z}_S^\alpha] - \text{Tr}[\hat{H}_B^{(n)} \tau_{B,n}^\alpha / \tilde{Z}_B^{n,\alpha}]. \quad (\text{A11})$$

Energy conservation implies

$$\text{Tr}[\hat{H}_{SB}^{(n)} (\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha}] = \text{Tr}[\hat{H}_{SB}^{(n)} \tilde{\tau}_{SB}^{n,\alpha}]. \quad (\text{A12})$$

Combining Eqs. (A12), (A11) we achieve

$$D(\tilde{\tau}_{SB}^{n,\alpha} \| (\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha}) = \text{Tr}[\lambda \hat{I}_n (\tilde{\tau}_{SB}^{n,\alpha} - (\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha})] \beta. \quad (\text{A13})$$

Pinsker inequality states that for any two density matrices ρ, σ ,

$$D(\rho \| \sigma) \geq \frac{1}{2} \|\rho - \sigma\|_1^2. \quad (\text{A14})$$

It follows from it, and from Eq. (A13),

$$\|U_n(\tau_S \otimes \tau_{B,n})^\alpha U_n^\dagger - (\tau_S \otimes \tau_{B,n})^\alpha\|_1 \leq \tilde{Z}_{SB}^{n,\alpha} \beta \sqrt{2 \text{Tr}[\lambda \hat{I}_n (\tilde{\tau}_{SB}^{n,\alpha} - (\tau_S \otimes \tau_{B,n})^\alpha / \tilde{Z}_{SB}^{n,\alpha})]} \quad (\text{A15})$$

$$\leq 2 \tilde{Z}_{SB}^{n,\alpha} \beta \sqrt{\sup_{\rho \in \mathcal{S}(\mathcal{H}_S \otimes \mathcal{H}_{B,n})} |\text{Tr}[\hat{I}_n \rho]| \lambda} \quad (\text{A16})$$

$$\leq 2 \tilde{Z}_{SB}^{n,\alpha} \beta \sqrt{\lambda \|\hat{I}_n\|} \quad (\text{A17})$$

□

This lemma may be of independent interest, as it makes explicit the idea mentioned in previous work such as [48, 49] of how Davies maps, in the weak coupling limit, can be taken as free operations in the resource theory of athermality [50].

With it at hand, we can prove the central lemma.

Lemma 6 (Lemma 1 of main text). Assume conditions in Section A 1 hold. Then all maps $T_t(\cdot)$ satisfy the inequality

$$D((\cdot)\|\tau_S) - D(T_t(\cdot)\|\tau_S) \geq D\left((\cdot)\|\tilde{T}_t(T_t(\cdot))\right), \quad \forall t \geq 0 \quad (\text{A18})$$

where $\tilde{T}_t(\cdot)$ is the Petz recovery map corresponding to $T_t(\cdot)$,

$$\tilde{T}_t(\cdot) = \tau_S^{1/2} T_t^\dagger \left(\tau_S^{-1/2}(\cdot) \tau_S^{-1/2} \right) \tau_S^{1/2}, \quad (\text{A19})$$

with T_t^\dagger denoting the adjoint of T_t .

Proof. Had there been no interaction term (i.e. $\lambda = 0$) and the bath been finite dimensional, the proof of this Lemma would have been straightforward, using the techniques developed in [7] involving simple manipulations of the relative entropy, and the data processing inequality for finite dimensional baths. The added difficulty here will be in proving monotone convergence as the bath Hilbert space tends to infinity. To achieve this, we will use Lemma 5 and continuity arguments. We will perform the calculations for the map $\text{Tr}_B \left[e^{-i\hat{H}_{SB}}(\cdot) \otimes \rho_B e^{i\hat{H}_{SB}} \right]$ rather than $T_t(\cdot)$ itself. We will finally take the limit described in Eq. (A4) to conclude the proof.

Noting that the relative entropy between two copies is zero, followed by using its additivity and unitarity invariance properties, we find for $\rho_S \in \mathcal{S}(\mathcal{H}_S)$,

$$D(\rho_S\|\tau_S) = D(\rho_S \otimes \tau_{B,n}\|\tau_S \otimes \tau_{B,n}) = D(U_n \rho_S \otimes \tau_{B,n} U_n^\dagger \| U_n \tau_S \otimes \tau_{B,n} U_n^\dagger) \quad (\text{A20})$$

$$= D(U_n \rho_S \otimes \tau_{B,n} U_n^\dagger \| \tau_S \otimes \tau_{B,n} + \sqrt{\lambda} \hat{B}_n(\lambda)) \quad (\text{A21})$$

where $\hat{B}_n(\lambda) := (U_n \tau_S \otimes \tau_{B,n} U_n^\dagger - \tau_S \otimes \tau_{B,n}) / \sqrt{\lambda}$.

With the identity $D(\gamma_{CD}\|\zeta_{CD}) - D(\gamma_D\|\zeta_D) = D(\gamma_{CD}\|\exp(\ln \zeta_{CD} + \ln \mathbb{1}_C \otimes \zeta_D - \ln \mathbb{1}_C \otimes \zeta_D))$ for bipartite states γ_{CD} , ζ_{CD} , we have that

$$D(U_n \rho_S \otimes \tau_{B,n} U_n^\dagger \| \tau_S \otimes \tau_{B,n} + \sqrt{\lambda} \hat{B}_n(\lambda)) - D(\sigma_S \| \tau_S + \sqrt{\lambda} \text{Tr}_{B,n}[\hat{B}_n(\lambda)]) \quad (\text{A22})$$

$$= D\left(U_n \rho_S \otimes \tau_{B,n} U_n^\dagger \left\| \exp\left(\ln(\tau_S \otimes \tau_{B,n} + \sqrt{\lambda} \hat{B}_n(\lambda)) + \ln \sigma_S \otimes \mathbb{1}_{B,n} - \ln(\tau_S + \sqrt{\lambda} \text{Tr}_{B,n}[\hat{B}_n(\lambda)]) \otimes \mathbb{1}_{B,n} \right) \right\| \right) \quad (\text{A23})$$

$$\geq D\left(\rho_S \left\| \text{Tr}_{B,n} \left[U_n^\dagger \exp\left(\ln(\tau_S \otimes \tau_{B,n} + \sqrt{\lambda} \hat{B}_n(\lambda)) + \ln \sigma_S \otimes \mathbb{1}_{B,n} - \ln(\tau_S + \sqrt{\lambda} \text{Tr}_{B,n}[\hat{B}_n(\lambda)]) \otimes \mathbb{1}_{B,n} \right) U_n \right] \right\| \right) \quad (\text{A24})$$

where $\sigma_{S,n} := \text{Tr}_{B,n}[U_n \rho_S \otimes \tau_{B,n} U_n^\dagger]$ and in the last line we have used the unitarity invariance of the relative entropy followed by the data processing inequality. Plugging Eq. (A21) into Eq. (A24) followed by taking the $n \rightarrow \infty$ limit, we obtain

$$D(\rho_S\|\tau_S) - D(\sigma_S\|\tau_S + \sqrt{\lambda} \text{Tr}_B[\hat{B}(\lambda)]) \quad (\text{A25})$$

$$\geq D\left(\rho_S \left\| \text{Tr}_B \left[U^\dagger \exp\left(\ln(\tau_S \otimes \tau_B + \sqrt{\lambda} \hat{B}(\lambda)) + \ln \sigma_S \otimes \mathbb{1}_B - \ln(\tau_S + \sqrt{\lambda} \text{Tr}_B[\hat{B}(\lambda)]) \otimes \mathbb{1}_B \right) U \right] \right\| \right), \quad (\text{A26})$$

where we have defined $\hat{B}(\lambda) := \lim_{n \rightarrow \infty} \hat{B}_n(\lambda)$, $\sigma_S := \lim_{n \rightarrow \infty} \sigma_{S,n}$. Before continuing, we will first note the validity of Eq. (A26). We start by showing that $\hat{B}(\lambda)$ is trace class for $\lambda \in [0, 1]$. From Lemma 5 it follows

$$\|\hat{B}_n(\lambda)\|_1 \leq 2\tilde{Z}_{SB}^{n,1} \beta \sqrt{\|\hat{I}_n\|}, \quad (\text{A27})$$

for all $\lambda \in [0, 1]$ with the r.h.s. λ independent. By definition of $\tilde{Z}_{SB}^{n,\alpha}$, it follows that it is the partition function of a tensor product of thermal states on $\mathcal{S}(\mathcal{H}_S \otimes \mathcal{H}_{B,n})$ at inverse temperatures $\alpha\beta_S, \alpha\beta$. Since the Hamiltonians $\hat{H}_{B,1}, \hat{H}_{B,2}, \hat{H}_{B,3}, \dots, \hat{H}_B$ by definition have well defined thermal states (finite partition functions) for all positive temperatures, it follows that $\lim_{n \rightarrow \infty} \tilde{Z}_{SB}^{n,\alpha} < \infty$ for all $\alpha > 0$. Thus noting that by definition, $\lim_{n \rightarrow \infty} \|\hat{I}_n\| = \|\hat{I}\|$ and that \hat{I} is a bounded operator, it follows that

$$\|\hat{B}(\lambda)\|_1 = \lim_{n \rightarrow \infty} \|\hat{B}_n(\lambda)\|_1 = 2\tilde{Z}_{SB}^{\infty,1} \beta \sqrt{\|\hat{I}\|} < \infty, \quad (\text{A28})$$

Thus since $\tau_S + \sqrt{\lambda} \text{Tr}_B[\hat{B}(\lambda)]$ is finite dimensional and Hermitian, and the eigenvalues of finite dimensional Hermitian matrices are continuous in their entries [51, 52], it follows, since τ_S has full support, that there exists $0 < \lambda^* \leq 1$ such that for all $\lambda \in [0, \lambda^*]$, $\tau_S + \sqrt{\lambda} \text{Tr}_B[\hat{B}(\lambda)]$ has full support. Thus for all $\lambda \in [0, \lambda^*]$, the r.h.s. of Eq. (A26) is upper bounded by a finite quantity uniformly in $n \rightarrow \infty$ and thus since relative entropies are non-negative by definition, Eq. (A26) is well defined for all $\lambda \in [0, \lambda^*]$.

We now set Δ appearing in U to $\Delta = t/\lambda^2$ followed by taking the limit $\lambda \rightarrow 0^+$ while keeping t fixed in Eq. (A26) thus achieving

$$D(\rho_S \| \tau_S) - D(T_t(\rho_S) \| \tau_S) \geq D\left(\rho_S \left\| \text{Tr}_B[U^\dagger T_t(\rho_S) \otimes \tau_B U]\right.\right), \quad (\text{A29})$$

where we have used that by definition, $T_t(\cdot) = \lim_{\lambda \rightarrow 0^+} \text{Tr}_B[U(\cdot) \otimes \tau_B U^\dagger]$.

We now proceed to calculate the Petz's recovery map for the map $T_t(\cdot)$. The adjoint map is $\text{Tr}_B[\tau_B^{1/2} U^\dagger((\cdot) \otimes \mathbb{1}_B) U \tau_B^{1/2}]$. Hence from the definition in Eq.(A50) it follows that the Petz recovery map for $T_t(\cdot)$ is

$$\tilde{T}_t(\cdot) := \tau_S^{1/2} \text{Tr}_B \left[\tau_B^{1/2} U^\dagger \left(\tau_S^{-1/2}(\cdot) \tau_S^{-1/2} \otimes \mathbb{1}_B \right) U \tau_B^{1/2} \right] \tau_S^{1/2}. \quad (\text{A30})$$

Similarly to before, we define a traceless, self adjoint operator $\tilde{B} = \tilde{B}(\lambda) := \left(U \tau_S^{1/2} \otimes \tau_B^{1/2} U^\dagger - \tau_S^{1/2} \otimes \tau_B^{1/2} \right) / \sqrt{\lambda}$. In analogy with the reasoning which led to Eq. (A28), it follows from Lemma 5 that $\|\tilde{B}(\lambda)\|_1 = \lim_{n \rightarrow \infty} \|\tilde{B}_n(\lambda)\|_1 = 2\tilde{Z}_{SB}^{\infty, 1/2} \beta \sqrt{\|\tilde{I}\|} < \infty$, for all $\lambda \in [0, 1]$. For general $U = \exp(-i\Delta \hat{H}_{SB})$, we can now write

$$\tau_S^{1/2} \text{Tr}_B \left[\tau_B^{1/2} U^\dagger \left(\tau_S^{-1/2}(\cdot) \tau_S^{-1/2} \otimes \mathbb{1}_B \right) U \tau_B^{1/2} \right] \tau_S^{1/2} \quad (\text{A31})$$

$$= \text{Tr}_B \left[\left(U^\dagger \tau_S^{1/2} \otimes \tau_B^{1/2} + \sqrt{\lambda} U^\dagger \tilde{B} \right) \left(\tau_S^{-1/2}(\cdot) \tau_S^{-1/2} \otimes \mathbb{1}_B \right) \left(\tau_S^{1/2} \otimes \tau_B^{1/2} U + \sqrt{\lambda} \tilde{B} U \right) \right] \quad (\text{A32})$$

$$= \text{Tr}_B[U^\dagger((\cdot) \otimes \tau_B) U] + \sqrt{\lambda} \hat{g}_1(\cdot) + \lambda \hat{g}_2(\cdot) \in \mathcal{S}(\mathcal{H}_B), \quad (\text{A33})$$

where

$$\hat{g}_1(\cdot) = \text{Tr}_B \left[U^\dagger \tilde{B} (\tau_S^{-1/2}(\cdot) \otimes \tau_B^{1/2}) U \right] + \text{Tr}_B \left[U^\dagger ((\cdot) \tau_S^{-1/2} \otimes \tau_B^{1/2}) \tilde{B} U \right] \quad (\text{A34})$$

$$\hat{g}_2(\cdot) = \text{Tr}_B \left[U^\dagger \tilde{B} \left(\tau_S^{-1/2}(\cdot) \tau_S^{-1/2} \otimes \mathbb{1}_B \right) \tilde{B} U \right], \quad (\text{A35})$$

which are well defined since they are comprised of products of bounded operators. Similarly to before, in Eq. (A33) we now set Δ appearing in U to $\Delta = t/\lambda^2$ followed by taking the limit $\lambda \rightarrow 0^+$ while keeping t fixed achieving

$$\tilde{T}_t(\cdot) = \text{Tr}_B[U^\dagger((\cdot) \otimes \tau_B) U] \quad (\text{A36})$$

where we have used Eq. (A30). Hence substituting Eq. (A34) in to Eq. (A29) and noting the equations holds for all states $\rho_S \in \mathcal{S}(\mathcal{H}_B)$, we conclude the proof. \square

Remark 7. In the above proof, we have taken two independent limits, namely 1st the infinite bath volume limit ($n \rightarrow \infty$) followed by the Van Hove limit ($\lambda \rightarrow 0^+$ while keeping t fixed). This is the order in which Davies performed the limits [2, 47] when defining the Davies map. From physical reasoning, one would expect the Davies map to be equally valid if the order of the limits is reversed. We note that the proof of Theorem 2 follows also if the order of the these two limits is reversed but now with the new definitions

$$T_t(\cdot) = \lim_{n \rightarrow \infty} \lim_{\lambda \rightarrow 0^+} \text{Tr}_{B,n} [U_n(\tilde{t})(\cdot) \otimes \tau_{B,n} U_n^\dagger(\tilde{t})] \in \mathcal{S}(\mathcal{H}_S) \quad \text{subject to } \tilde{t}\lambda^2 = t \text{ fixed.} \quad (\text{A37})$$

$$\tilde{T}_t(\cdot) = \lim_{n \rightarrow \infty} \lim_{\lambda \rightarrow 0^+} \tau_S^{1/2} \text{Tr}_{B,n} \left[\tau_{B,n}^{1/2} U_n^\dagger(\tilde{t}) \left(\tau_S^{-1/2}(\cdot) \tau_S^{-1/2} \otimes \mathbb{1}_{B,n} \right) U_n(\tilde{t}) \tau_{B,n}^{1/2} \right] \tau_S^{1/2} \in \mathcal{S}(\mathcal{H}_S) \quad \text{subject to } \tilde{t}\lambda^2 = t \text{ fixed.} \quad (\text{A38})$$

An interesting technical question is whether the above limits commute i.e. whether Eqs. (A37), (A38) are identical to Eqs. (A4), (A30).

3. Quantum detailed balance and Petz recovery map

Now we show that all Davies maps have the peculiar property that they are the same as their Petz recovery map. This is because of a crucial property satisfied by their generators: *quantum detailed balance*. For Theorem 6 in the main text to hold, we require both the conditions of Section A 1 and the following Lemma to hold. For the sake of generality, we show that the results is true for any fixed point Ω with full support. We remind the reader, that a dynamical semigroup $M_t(\cdot)$ is a one parametre family of CPTP maps with a generator consisting of a unitary part $\theta(\cdot) = -[\tilde{H}_{\text{eff}}, \cdot]$, and a dissipative part called a Lindbladian, $\mathcal{L}(\cdot)$, such that all together we have

$$M_t(\cdot) = e^{ti\theta + t\mathcal{L}}(\cdot) \quad (\text{A39})$$

Theorem 8 (Dissipative Recovery map). A quantum dynamical semigroup $M_t(\cdot)$ with no unitary part, $\theta = 0$, and Lindbladian \mathcal{L} satisfying quantum detailed balance (Eq. (3)) for the state Ω with full rank, is equal to its corresponding Petz recovery map, namely,

$$M_t(\cdot) = \tilde{M}_t(\cdot), \quad (\text{A40})$$

where

$$\tilde{M}_t(\cdot) = \Omega^{1/2} M_t^\dagger (M_t(\Omega)^{-1/2} \cdot M_t(\Omega)^{-1/2}) \Omega^{1/2}. \quad (\text{A41})$$

Proof. The property of quantum detailed balance (also sometimes referred to as the reversibility, or KMS condition) reads

$$\langle A, \mathcal{L}^\dagger(B) \rangle_\Omega = \langle \mathcal{L}^\dagger(A), B \rangle_\Omega \quad (\text{A42})$$

for all $A, B \in \mathbb{C}^{d_S \times d_S}$, where \mathcal{L}^\dagger is the adjoint Lindbladian, and we define the scalar product

$$\langle A, B \rangle_\Omega := \text{Tr}[\Omega^{1/2} A^\dagger \Omega^{1/2} B]. \quad (\text{A43})$$

Because Eq. (A42) holds for all $A, B \in \mathbb{C}^{d_S \times d_S}$, Eq. (A42) implies that [53]

$$\mathcal{L}(\cdot) = \Omega^{1/2} \mathcal{L}^\dagger(\Omega^{-1/2} \cdot \Omega^{-1/2}) \Omega^{1/2}. \quad (\text{A44})$$

Eq. (A42) automatically implies that any power of the generator also obeys the same relation, that is, $\forall n \in \mathbb{N}^+$

$$\langle A, \mathcal{L}^{\dagger n}(B) \rangle_\Omega = \langle A, \Omega^{-1/2} \mathcal{L}(\Omega^{1/2} \dots \Omega^{-1/2} \mathcal{L}(\Omega^{1/2} B \Omega^{1/2}) \Omega^{-1/2} \dots \Omega^{1/2}) \Omega^{-1/2} \rangle_\Omega \quad (\text{A45})$$

$$= \langle A, \Omega^{-1/2} \mathcal{L}^n(\Omega^{1/2} B \Omega^{1/2}) \Omega^{-1/2} \rangle_\Omega \quad (\text{A46})$$

$$= \langle \mathcal{L}^{\dagger n}(A), B \rangle_\Omega, \quad (\text{A47})$$

where in the first line we use Eq. (A44) n times and the 2nd line follows from the definition of the adjoint map. Hence we can also write

$$\mathcal{L}^n(\cdot) = \Omega^{1/2} \mathcal{L}^{\dagger n}(\Omega^{-1/2} \cdot \Omega^{-1/2}) \Omega^{1/2}. \quad (\text{A48})$$

The semigroup can be written as $M_t(\cdot) = e^{\mathcal{L}t}(\cdot)$. Its adjoint semigroup is given by $e^{\mathcal{L}^\dagger t}$ and hence the Petz recovery map is (see Eq.(A50))

$$\tilde{M}_t(\cdot) = \Omega^{1/2} e^{\mathcal{L}^\dagger t}(\Omega^{-1/2} \cdot \Omega^{-1/2}) \Omega^{1/2}. \quad (\text{A49})$$

Since, $\tilde{M}_t(\cdot) = \Omega^{1/2} (\sum_{n=0}^{\infty} (t\mathcal{L})^{\dagger n}(\Omega^{-1/2} \cdot \Omega^{-1/2}) / (n!)) \Omega^{1/2}$, Eq. (A49) together with Eq. (A48), means that $\tilde{M}_t(\cdot) = M_t(\cdot)$.

We note that the Petz recovery map is defined in terms of a map $\Gamma(\cdot)$ and a state σ_S as the unique solution to

$$\langle A, \Gamma^\dagger(B) \rangle_{\sigma_S} = \langle \tilde{\Gamma}^\dagger(A), B \rangle_{\Gamma(\sigma_S)} \quad (\text{A50})$$

for all $A, B \in \mathbb{C}^{d_S \times d_S}$ and the scalar product is given by Eq. (A43). The solution takes the form [53]

$$\tilde{T}(\cdot) = \sigma_S^{1/2} T^\dagger(T(\sigma_S)^{-1/2} \cdot T(\sigma_S)^{-1/2}) \sigma_S^{1/2}, \quad (\text{A51})$$

such that we always have that $\tilde{T}(T(\sigma_S)) = \sigma_S$. Here this simplifies by choosing $\sigma_S = \Omega$ a fixed point of $M_t(\cdot)$. \square

When the generator is time-independent and $\theta = 0$, we thus have from Theorem 8 that the combination of a map for a time t and its recovery map is equivalent to applying the map for a time $2t$. That is $\tilde{M}_t(M_t(\cdot)) = M_{2t}(\cdot)$. This means we can write Eq. (A18) in a particularly simple form.

The following Lemma builds on Theorem 8 to extend it to the case in which the dynamical semigroup also includes a unitary part.

Lemma 9 (Dissipative and unitary Recovery map). Let $M_t(\cdot)$ be a quantum dynamical semigroup with unitary part θ and Lindbladian \mathcal{L} which: 1) satisfying quantum detailed balance (Eq. (3)) for the state Ω with full rank and 2) commute, $\theta(\mathcal{L}(\cdot)) = \mathcal{L}(\theta(\cdot))$. Then, $M_t(\cdot)$ has a Petz recovery map $\tilde{M}(\cdot)$ which is a dynamical semigroup with unitary part $-\theta$ and Lindbladian \mathcal{L} . Namely, if

$$M_t(\cdot) = e^{ti\theta + t\mathcal{L}}(\cdot), \quad (\text{A52})$$

satisfying 1) and 2), then

$$\tilde{M}_t(\cdot) = e^{-ti\theta + t\mathcal{L}}(\cdot). \quad (\text{A53})$$

Proof. We just need to note two facts:

- The Petz recovery map of a unitary map $U(\cdot)U^\dagger$ that had fixed point Ω is $U^\dagger(\cdot)U$.
- The Petz recovery map of a composition of two maps with the same fixed point, is equal to the composition of the Petz recovery maps of the individual maps, i.e. $\widetilde{\Gamma_1 \circ \Gamma_2} = \tilde{\Gamma}_2 \circ \tilde{\Gamma}_1$ (this is one of the key properties listed in [4]).

We hence can write the recovery map of $M_t(\cdot)$ as

$$\tilde{M}_t(\rho_S) = e^{t\mathcal{L}}(e^{iH_{\text{eff}}t}\rho_S e^{-iH_{\text{eff}}t}) = e^{iH_{\text{eff}}t}e^{t\mathcal{L}}(\rho_S)e^{-iH_{\text{eff}}t}. \quad (\text{A54})$$

The only difference between M_t and \tilde{M}_t is the change of sign in the time of the unitary evolution. The recovery map is then made up of the dissipative part evolving forwards, and the unitary part evolving backwards in time. \square

Theorem 10. (Theorem 2 of main text) Assume conditions in Section A 1 hold and $T_t(\cdot)$ satisfies quantum detailed balance (Eq. (3)) and has zero unitary part, $\theta = 0$. Then $T_t(\cdot)$ satisfies the inequality

$$D((\cdot)\|\tau_S) - D(T_t(\cdot)\|\tau_S) \geq D((\cdot)\|T_{2t}(\cdot)), \quad t \geq 0. \quad (\text{A55})$$

Proof. Direct consequence of Theorems 8, 6. \square

Remark 11 (When $\theta \neq 0$). Due to properties 3), 5) of the main text satisfied by Davies maps, and the unitary invariance of the Relative entropy (i.e. $D(U \cdot U^\dagger \| U \cdot U^\dagger) = D(\cdot \| \cdot)$), it follows

$$D((\cdot)\|\tau_S) - D(T_t(\cdot)\|\tau_S) = D((\cdot)\|\tau_S) - D(e^{t\mathcal{L}}(\cdot)\|\tau_S), \quad (\text{A56})$$

and thus the l.h.s. of Eq. (A55) is the same even when a non zero unitary part is included. Furthermore, we note that the canonical form of Davies maps have $\theta(\mathcal{L}(\cdot)) = \mathcal{L}(\theta(\cdot))$ by definition [see property 3) in main text] and thus, due to Lemma 9, even when $\theta \neq 0$, we have that

$$D((\cdot)\|\tilde{T}_t(T_t(\cdot))) = D((\cdot)\|e^{2t\mathcal{L}}(\cdot)), \quad (\text{A57})$$

which is the r.h.s. of Eq. (A55). Thus applying Theorem 2, we have

$$D((\cdot)\|\tau_S) - D(T_t(\cdot)\|\tau_S) = D((\cdot)\|\tau_S) - D(e^{t\mathcal{L}}(\cdot)\|\tau_S) \geq D((\cdot)\|e^{2t\mathcal{L}}(\cdot)), \quad (\text{A58})$$

for any θ .

4. Spohn's inequality: rate of entropy production

We give an alternative proof of a well-known result which was first shown in [3] that gives the expression for the infinitesimal rate of entropy production of a Davies map. This is stated without a proof in many standard references such as [25, 54]. Then we show in a similar way how in the infinitesimal time limit our lower bound becomes trivial.

First we need the following lemma, which proof can be found in, for instance, [55].

Lemma 12. Let $\mathbb{1} \in \mathbb{C}^{n \times n}$ be the identity matrix, and $A, B \in \mathbb{C}^{n \times n}$ be matrices such that both A and $A + tB$ are positive with $t \in \mathbb{R}$, we have that

$$\log(A + tB) - \log A = t \int_0^1 \frac{1}{(1-x)A + x\mathbb{1}} B \frac{1}{(1-x)A + x\mathbb{1}} dx + \mathcal{O}(t^2) \quad (\text{A59})$$

With this, we can show the following:

Theorem 13. Let $\mathcal{L}(\rho_S(t))$ be the generator of a dynamical semigroup, with a fixed point τ_S such that $\mathcal{L}(\tau_S) = 0$. We have that the entropy production rate $\sigma(\rho_S(t))$ is given by

$$\sigma(\rho_S(t)) := -\frac{dD(\rho_S(t)||\tau_S)}{dt} = \text{Tr}[\mathcal{L}(\rho_S(t))(\log \tau_S - \log \rho_S(t))] + \text{Tr}[\mathcal{L}(\rho_S(t))\Pi_{\rho_S(t)}] \geq 0, \quad (\text{A60})$$

where $\Pi_{\rho_S(t)}$ is the projector onto the support of $\rho_S(t)$. The second term of the sum vanishes at all times for which the rate is finite.

Proof. The last inequality (positivity) follows from the data processing inequality for the relative entropy, so we only need to prove the equality. The proof only requires Lemma 12 and some algebraic manipulations. We have that

$$\frac{dD(\rho_S(t)||\tau_S)}{dt} = \lim_{h \rightarrow 0} \frac{D(\rho_{t+h}) - D(\rho_S(t))}{h} \quad (\text{A61})$$

$$= \lim_{h \rightarrow 0} \frac{\text{Tr}[(\rho_S(t) + \mathcal{L}(\rho_S(t))h)(\log \{\rho_S(t) + \mathcal{L}(\rho_S(t))h\} - \log \tau_S)] - \text{Tr}[\rho_S(t)(\log \rho_S(t) - \log \tau_S)]}{h} \quad (\text{A62})$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \left[\text{Tr}[(\rho_S(t) + \mathcal{L}(\rho_S(t))h)\{\log(\rho_S(t)) + h \int_0^1 \frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} \mathcal{L}(\rho_S(t)) \frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} dx\} - \log \tau_S] \right. \\ \left. - \text{Tr}[\rho_S(t)(\log \rho_S(t) - \log \tau_S)] \right] \quad (\text{A63})$$

$$= \text{Tr}[\mathcal{L}(\rho_S(t))(\log \rho_S(t) - \log \tau_S)] + \text{Tr}[\rho_S(t) \int_0^1 \frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} \mathcal{L}(\rho_S(t)) \frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} dx] \quad (\text{A64})$$

$$= \text{Tr}[\mathcal{L}(\rho_S(t))(\log \rho_S(t) - \log \tau_S)] + \text{Tr}[\rho_S(t) \mathcal{L}(\rho_S(t)) \int_0^1 \left(\frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} \right)^2 dx]. \quad (\text{A65})$$

Where to go from the 2nd to the third line we used Lemma 12, and from the 4th to the 5th we use the cyclicity and linearity of the trace. Now note the following integral

$$\int_0^1 \left(\frac{1}{(1-x)p + x} \right)^2 dx = \frac{1}{p} \quad \forall p \neq 0. \quad (\text{A66})$$

This means that, on the support of $\rho_S(t)$,

$$\int_0^1 \left(\frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} \right)^2 dx = \frac{1}{\rho_S(t)}. \quad (\text{A67})$$

Note that outside the support of $\rho_S(t)$ this integral is not well defined. Given this, we can write

$$\frac{dD(\rho_S(t)||\tau_S)}{dt} = \text{Tr}[\mathcal{L}(\rho_S(t))(\log \rho_S(t) - \log \tau_S)] + \text{Tr}[\mathcal{L}(\rho_S(t))\Pi_{\rho_S(t)}], \quad (\text{A68})$$

where $\Pi_{\rho_S(t)}$ is the projector onto the support of $\rho_S(t)$. The Lindbladian is traceless $\text{Tr}[\mathcal{L}(\rho_S(t))] = 0$ and hence second term of this Equation vanishes as long as $\text{supp}(\mathcal{L}(\rho_S(t))) \subseteq \text{supp}(\rho_S(t))$, which we can expect for most times. At instants in time when this is not the case and this term may give a finite contribution (that is, when the rank increases), the first term in Eq. (A68) diverges logarithmically [3], and hence that finite contribution is negligible. \square

A similar reasoning can be used to show that the instantaneous lower bound on entropy production rate that we can get from our main result in Eq. (11) is trivial for most times. In particular, we can show

Lemma 14. The lower bound of Eq. (11) vanishes in the limit of infinitesimal time transformations. More precisely, we have that

$$\lim_{h \rightarrow 0} \frac{D(\rho_S(t) || \rho_S(t+2h))}{h} = -2\text{Tr}[\mathcal{L}(\rho_S(t))\Pi_{\rho_S(t)}], \quad (\text{A69})$$

where $\Pi_{\rho_S(t)}$ is the projector onto the support of $\rho_S(t)$. This vanishes as long as $\text{supp}(\mathcal{L}(\rho_S(t))) \subseteq \text{supp}(\rho_S(t))$.

Proof. The proof is similar to the one for Theorem 13 above.

$$\lim_{h \rightarrow 0} \frac{D(\rho_S(t) || \rho(t+2h))}{h} = \lim_{h \rightarrow 0} \frac{1}{h} \text{Tr}[\rho_S(t)(\log \rho_S(t) - \log(\rho_S(t) + 2h\mathcal{L}(\rho_S(t))))] \quad (\text{A70})$$

$$= \text{Tr}[-2\rho_S(t) \int_0^1 \frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} \mathcal{L}(\rho_S(t)) \frac{1}{(1-x)\rho_S(t) + x\mathbb{1}} dx] \quad (\text{A71})$$

$$= -2\text{Tr}[\mathcal{L}(\rho_S(t))\Pi_{\rho_S(t)}], \quad (\text{A72})$$

where in the second line we applied Lemma 12, and in the third we used Eq. (A67). \square

Hence for infinitesimal times, the lower bound gives the same condition as the positivity condition in Eq. (A60). It will be nonzero only when $\text{supp}(\mathcal{L}(\rho_S(t))) \not\subseteq \text{supp}(\rho_S(t))$, in which case the rate of entropy production diverges (at points in time when the rank of the system increases).

Appendix B: Maps Beyond Davies

Given that the inequality in Eq. (8) is saturated in some limits, such as when the evolution approaches the fixed point, it is unlikely that a stronger inequality of a similar kind can be derived even in particular cases. However, general results are known for CPTP maps, leading to weaker forms of such bound. In this Section we state the best known general result from [10] and show how they simplify in particular cases of maps with properties similar to Davies maps. This means that we can also bound the entropy production of maps that may not be Davies maps.

The result, which proof involves techniques from complex interpolation theory, is the following:

Theorem 15. (Main result of [10]) Let $\Gamma(\cdot)$ be a CPTP map, and ρ, σ any two quantum states. We have that

$$D(\rho || \sigma) - D(\Gamma(\rho) || \Gamma(\sigma)) \geq -2 \int_{\mathbb{R}} dt \, p(t) \log F(\rho, \tilde{\Gamma}_t(\Gamma(\rho))), \quad (\text{B1})$$

where $F(\rho, \sigma) = \text{Tr}[\sqrt{\sqrt{\sigma}\rho\sqrt{\sigma}}]$ is the quantum fidelity, the map $\tilde{\Gamma}_t$ is the *rotated* recovery map

$$\tilde{\Gamma}_t(\cdot) = \sigma^{it} \tilde{\Gamma}(\Gamma(\sigma)^{-it} \cdot \Gamma(\sigma)^{it}) \sigma^{-it} \quad (\text{B2})$$

and $p(t)$ is the probability density function $p(t) = \frac{\pi}{2} (\cosh(\pi t) + 1)^{-1}$.

Proof. See [10]. \square

We now observe that the rotated map can be simplified given the following conditions:

- If the map has a fixed point $\Gamma(\Omega) = \Omega$, the Petz recovery map simplifies to become

$$\tilde{\Gamma}_t(\cdot) = \Omega^{it} \tilde{\Gamma}(\Omega^{-it} \cdot \Omega^{it}) \Omega^{-it} \quad \forall t \in \mathbb{R} \quad (\text{B3})$$

This by itself implies that $\tilde{\Gamma}_t(\Omega) = \Omega$.

- The map may also obey the property of *time-translation symmetry*, where this is given by

$$\Gamma(\cdot) = \Omega^{it} \Gamma(\Omega^{-it} \cdot \Omega^{it}) \Omega^{-it}. \quad (\text{B4})$$

If a map obeys this symmetry, the adjoint map $\Gamma^\dagger(\cdot)$ also will. This can be seen through the definition of the adjoint, which is that for any two matrices A, B ,

$$\text{Tr}[A\Gamma(B)] = \text{Tr}[\Gamma^\dagger(A)B], \quad (\text{B5})$$

and in particular, it holds for the matrices $A' = \Omega^{it} A \Omega^{-it}$, $B' = \Omega^{-it} B \Omega^{it}$. This, together with Eq. (B4), means that

$$\text{Tr}[\Gamma^\dagger(A)B] = \text{Tr}[\Omega^{it}\Gamma^\dagger(\Omega^{-it} \cdot \Omega^{it})\Omega^{-it}(A)B] \quad (\text{B6})$$

Hence this property, together with the fixed-point property, means that the rotated recovery map becomes equal to the Petz map, and the integral in Eq. (B1) gets averaged out.

It may be the case, however, that the symmetry exists, but that the fixed point is not the thermal state, and hence the simplification does not occur. This may be the case for instance when there is weak coupling to a non-thermal environment.

- If on top of these two conditions the map has the property of quantum detailed balance, namely

$$\langle A, \Gamma^\dagger(B) \rangle_\Omega = \langle \Gamma^\dagger(A), B \rangle_\Omega, \text{ for all } A, B \in \mathbb{C}^{d_S \times d_S}, \quad (\text{B7})$$

the Petz recovery map and the original one are the same $\tilde{\Gamma}(\cdot) = \Gamma(\cdot)$. Examples of maps which satisfy detailed balance which are not Davies maps do exist. See [42] for general characterization of quantum dynamical semigroups.

When all these hold we have that Eq. (B1) becomes

$$D(\rho||\Omega) - D(\Gamma(\rho)||\Omega) \geq -2 \log F(\rho, \Gamma(\Gamma(\rho))). \quad (\text{B8})$$

This bound could be tightened by replacing the $-2 \log F(\rho, \Gamma(\Gamma(\rho)))$ with the measured relative entropy, $D_{\mathbb{M}}(\rho||\Gamma(\Gamma(\rho)))$ [12]. This would achieve a tighter bound, although at the expense of it being less explicit, unless one could solve the maximization problem in the definition of the measured relative entropy. If the map is a dynamical semigroup with a time-independent generator $\Gamma = M_t$, we may also write $M_t(M_t(\cdot)) = M_{2t}(\cdot)$.

Davies maps have all these properties. Further examples where all these properties appear are semigroups derived from the low-density limit (which models a system immersed in an ideal gas at low density, see [25] for details), or the so-called *heat bath generators* [56].

We note however that $D(\rho||\sigma) \geq -2 \log F(\rho, \sigma)$, and hence Eq. (B8) is a weaker bound than Eq. (8), and in particular is not tight as the fixed point is approached.

Appendix C: Equivalence of definitions of quantum detailed balance

In the literature, different nonequivalent definitions of the property of quantum detailed balance have been given. While in many places the one given is that of Eq. (3), an alternative definition, which can be found for instance in [25, 27] is that the Lindbladian is self-adjoint with respect to the inner product

$$\langle A, \mathcal{L}^\dagger(B) \rangle'_\Omega = \langle \mathcal{L}^\dagger(A), B \rangle'_\Omega, \quad (\text{C1})$$

for all $A, B \in \mathbb{C}^{d_S \times d_S}$, where the inner product is defined as

$$\langle A, B \rangle'_\Omega = \text{Tr}[\Omega A^\dagger B]. \quad (\text{C2})$$

Eq. C2 is different from that of Eq. (A43) due to the noncommutativity of the operators. The solution to Eq (C1) is [57]

$$\mathcal{L}(\cdot) = \Omega \mathcal{L}^\dagger(\Omega^{-1} \cdot), \quad (\text{C3})$$

while the solution to Eq. (3) is [53]

$$\mathcal{L}(\cdot) = \Omega^{1/2} \mathcal{L}^\dagger(\Omega^{-1/2} \cdot \Omega^{-1/2}) \Omega^{1/2}. \quad (\text{C4})$$

We now give a simple proof of the fact that, under the condition that the map is time-translation invariant w.r.t. fixed point, the two conditions are the same.

Theorem 16. For a Lindbladian operator $\mathcal{L}(\cdot)$ which obeys the property of time-translation symmetry w.r.t. fixed point Ω of full Rank (Eq. (14)), the quantum detailed balance conditions of Eq. (C3) and Eq. (C4) are equivalent.

Proof. We rewrite both Eq. (C3) and Eq. (C4) in terms of their individual matrix elements in the orthonormal basis $\{|i\rangle\}$ in which $\Omega = \sum_i p_i |i\rangle\langle i|$ is diagonal. Eq. (C3) can be written in the form

$$\langle i|\mathcal{L}(|n\rangle\langle m|)|j\rangle = \frac{p_i}{p_n} \langle i|\mathcal{L}^\dagger(|n\rangle\langle m|)|j\rangle \quad (\text{C5})$$

and Eq. (C4) is

$$\langle i|\mathcal{L}(|n\rangle\langle m|)|j\rangle = \sqrt{\frac{p_i p_j}{p_n p_m}} \langle i|\mathcal{L}^\dagger(|n\rangle\langle m|)|j\rangle. \quad (\text{C6})$$

We can see that for each matrix element the conditions only change by the factors multiplying in front, which are different unless $\frac{p_n}{p_m} = \frac{p_i}{p_j}$.

Let us now introduce the following decomposition of operators in $\mathbb{C}^{d_S \times d_S}$ in terms of their *modes of rotation*

$$A = \sum_{\omega} A_{\omega}, \quad (\text{C7})$$

where A_{ω} is defined as

$$A_{\omega} = \sum_{\substack{k,l \\ \text{s.t. } \omega = \log \frac{p_k}{p_l}}} |k\rangle\langle k| A |l\rangle\langle l|. \quad (\text{C8})$$

The name of *modes of rotation* is due to the fact that under the action of the unitary $\Omega^{-it} \cdot \Omega^{it}$ they rotate with a different Bohr frequency, that is

$$\Omega^{-it} A_{\omega} \Omega^{it} = A_{\omega} e^{-i\omega t}. \quad (\text{C9})$$

If the Lindbladian has the property of time-translational invariance w.r.t. the fixed point (Eq. (14)), it can be shown [58, 59] that each input mode is mapped to its corresponding output mode of the same Bohr frequency ω . We can write this fact as

$$\mathcal{L}(A_{\omega}) = \mathcal{L}(A)_{\omega}. \quad (\text{C10})$$

This means that in Eq. (C5) and (C6), $\langle i|\mathcal{L}(|n\rangle\langle m|)|j\rangle = 0$ unless the Bohr frequencies coincide at the input and the output, that is, when $\log \frac{p_n}{p_m} = \log \frac{p_i}{p_j}$. That is, the two conditions are nontrivial only in those particular matrix elements in which both are equivalent. \square