

**Conservation laws of the generalized Riemann equations at  $N = 2, 3, 4$** Binfang Gao,<sup>1, a)</sup> Kai Tian,<sup>1, b)</sup> Q. P. Liu,<sup>1, c)</sup> and Lujuan Feng<sup>1</sup>*Department of Mathematics, China University of Mining and Technology,  
Beijing 100083, People's Republic of China*

(Dated: 27 November 2024)

In this paper, we present infinitely many conserved densities satisfying particular conservation law  $F_t = (2uF)_x$  for the generalized Riemann equations at  $N = 2, 3, 4$ . In the  $N = 2$  case, we also construct conserved densities corresponding to new conservation laws containing an arbitrary smooth function. In virtue of reductions and/or changes of variables, related conserved densities are obtained for two component Hunter-Saxton equation, Hunter-Saxton equation, Gurevich-Zybin equation and Monge-Ampere equation.

Keywords: generalized Riemann equations, reciprocal transformation, conserved density, reduction

---

<sup>a)</sup>Electronic mail: ycugss2008@163.com

<sup>b)</sup>Electronic mail: tiankai@lsec.cc.ac.cn

<sup>c)</sup>Author to whom correspondence should be addressed. Electronic mail: qpl@cumtb.edu.cn

# I. INTRODUCTION

The Riemann equation

$$u_t + uu_x = 0, \tag{1}$$

where  $u = u(x, t)$  and subscripts denote specified partial derivatives, also known as inviscid Burgers equation or Hopf equation in literatures, has been extensively studied as a prototype for many phenomena related to hyperbolic systems.<sup>1</sup> Due to an interesting observation of Holm and Pavlov, the Riemann equation (1) was recently generalized to<sup>2-4</sup>

$$(\partial_t + u\partial_x)^N u = 0, \quad (N \in \mathbb{Z}_+) \tag{2}$$

which could be reformulated as a  $N$ -component system, namely

$$u_{j,t} = (1 - \delta_{jN})u_{j+1} - u_1u_{j,x}, \quad (j = 1, 2, \dots, N) \tag{3}$$

where  $u_1 = u$ ,  $u_{j+1} = (\partial_t + u\partial_x)u_j$  ( $1 \leq j < N$ ) and  $\delta_{jN}$  is the Kronecker delta. The equation (2), or equivalently the system (3), serves as a multi-component generalization of the Riemann equation (1) and will be referred to as the generalized Riemann equation, which has been shown to be closely related to some physically important models. In fact, the generalized Riemann equation at  $N = 2$  could be viewed as a regular Whitham type system,<sup>3</sup> and reduced to the Gurevich-Zybin system<sup>5</sup> or an equation describing non-local gas dynamics<sup>6</sup> as well. Through a Miura-type transformation, a  $N$ -component extension of the Hunter-Saxton equation<sup>7,8</sup> was deduced from the system (3).<sup>9</sup>

The generalized Riemann equation (3) is integrable, as certified by a universal matrix Lax representation at arbitrary  $N$ .<sup>9</sup> Furthermore, the bi-Hamiltonian structures for the generalized Riemann equations at  $N = 2, 3$  and  $4$  were constructed.<sup>2,4</sup> An extraordinary feature of the generalized Riemann equation (3) is attributed to abundant conserved densities of various types, which were constructed by different approaches, for instance, the recursion operator<sup>5,6</sup>, an iterative procedure<sup>2</sup> or the Lax representation.<sup>9</sup> It was already pointed out by Popowicz<sup>9</sup> that most conserved densities surprisingly satisfy a common conservation law, namely

$$\partial_t(F) = \partial_x(-u_1F), \tag{4}$$

where  $F$  is a certain smooth function of  $u_j$  ( $1 \leq j \leq N$ ) and their derivatives with respect to  $x$ . However, to the best of our knowledge there is still a lack of explanations to this richness of conserved densities.

Some clues from reduced equations of the generalized Riemann equation (3) may lead us to a better understanding about those conserved densities. On the one hand, the Riemann equation (1) was shown by Olver and Nutku to have a class of conservation law,<sup>10</sup> given by

$$\partial_t \left( u_x G(z_2, z_3, \dots, z_n) \right) = \partial_x \left( -u u_x G(z_2, z_3, \dots, z_n) \right),$$

where  $z_2 = u_x^{-3} u_{xx}$ ,  $z_{j+1} = u_x^{-1} (\partial_x z_j)$  ( $j \geq 2$ ), and  $G$  is an arbitrary smooth function of its arguments. On the other hand, similar results have recently been achieved for the Hunter-Saxton equation,<sup>11</sup> which also admits conserved densities involving arbitrary smooth functions. Supported by these facts, conserved densities of such kind are expected for the generalized Riemann equation (3).

In this paper, we will construct conserved densities for the generalized Riemann equations at  $N = 2, 3$  and  $4$ . With the help of appropriate changes of variables, we will manage to figure out the most general conserved densities for which equation (4) holds. It turns out that these conserved densities involve arbitrary functions. Similar discussions will be given to few systems related to the generalized Riemann equation (3), such as the Gurevich-Zybin system, the Monge-Ampere equation and the two-component Hunter-Saxton equation. The paper is organised as follows. In section II and III, we consider the generalized Riemann equation at  $N = 2$  and the related systems, respectively. In section IV, the generalized Riemann equations at  $N = 3, 4$  are studied. Last section presents some discussions.

## II. THE GENERALIZED RIEMANN EQUATION AT $N = 2$

We rescale dependent variables as  $u_1 = -2u$ ,  $u_2 = 4v$ , and write the generalized Riemann equation at  $N = 2$  as

$$u_t = 2uu_x - 2v, \tag{5a}$$

$$v_t = 2uv_x, \tag{5b}$$

whose general solutions in implicit form have been obtained by quadratures.<sup>12</sup>

Differentiating both equations (5a) and (5b) with respect to  $x$  yields

$$u_{xt} = 2uu_{xx} + 2u_x^2 - 2v_x, \tag{6a}$$

$$v_{xt} = \partial_x(2uv_x). \tag{6b}$$

Based on the trivial conservation law (6b), we introduce a reciprocal transformation

$$\mathcal{T}_1 : (x, t, u, v) \mapsto (y, \tau, \bar{u}, \bar{v}),$$

where new variables are defined by

$$dy = v_x dx + (2uv_x)dt, \quad d\tau = dt, \quad \bar{u}(y, \tau) = u_x, \quad \bar{v}(y, \tau) = v_x.$$

Applying  $\mathcal{T}_1$  to equations (6a) and (6b) respectively, we obtain a system of ODEs

$$\bar{u}_\tau = 2\bar{u}^2 - 2\bar{v}, \quad (7a)$$

$$\bar{v}_\tau = 2\bar{u}\bar{v}. \quad (7b)$$

If we change dependent variables as

$$\mathcal{T}_2 : (\bar{u}, \bar{v}) \mapsto (p, q) = \left( \frac{\bar{u}^2 - 2\bar{v}}{\bar{v}^2}, \frac{\bar{u}}{\bar{v}} \right),$$

then the system (7a)-(7b) is linearized to

$$p_\tau = 0, \quad (8a)$$

$$q_\tau = -2. \quad (8b)$$

Now suppose that  $F(u_x, v_x, \dots, u_{nx}, v_{nx})$  is a conserved density of the system (5a)-(5b) such that

$$\partial_t \left( F(u_x, v_x, \dots, u_{nx}, v_{nx}) \right) = \partial_x \left( 2uF(u_x, v_x, \dots, u_{nx}, v_{nx}) \right), \quad (9)$$

where  $u_{nx} = (\partial_x^n u)$  and  $v_{nx} = (\partial_x^n v)$ . Denote the conserved density  $F$  in the coordinates  $(y, \tau, p, q)$  as  $\widehat{F}$ , namely

$$\widehat{F}(p, q, p_y, q_y, \dots, p_{(n-1)y}, q_{(n-1)y}) = (\mathcal{T}_2 \circ \mathcal{T}_1)F(u_x, v_x, \dots, u_{nx}, v_{nx}),$$

then the equation (9) may be rewritten as

$$\partial_\tau \left( \widehat{F}(p, q, p_y, q_y, \dots, p_{(n-1)y}, q_{(n-1)y}) \right) = \frac{4q}{q^2 - p} \widehat{F}(p, q, p_y, q_y, \dots, p_{(n-1)y}, q_{(n-1)y}), \quad (10)$$

which identically holds provided that  $p$  and  $q$  solve (8a)-(8b).

It is easy to see that on the solutions of the system (8a)-(8b) the equation (10) reduces to

$$0 = \partial_\tau(\widehat{F}) - \frac{4q}{q^2 - p} \widehat{F} = -2 \frac{\partial \widehat{F}}{\partial q} - \frac{4q}{q^2 - p} \widehat{F},$$

which may be solved by the standard method, and yields

$$\widehat{F} = \frac{1}{q^2 - p} G(p, p_y, q_y, \dots, p_{(n-1)y}, q_{(n-1)y}),$$

where  $G$  is an arbitrary smooth function of its arguments. Switching to the coordinates  $(x, t, u, v)$  and taking account of the relations

$$p = u_x^2 v_x^{-2} - 2v_x^{-1}, \quad q = u_x v_x^{-1}, \quad \partial_y = v_x^{-1} \partial_x,$$

we obtain conserved densities with arbitrary function. Indeed,

$$\frac{v_x}{2} G(\alpha_1, \alpha_2, \beta_2, \dots, \alpha_n, \beta_n) \quad (11)$$

solves the conservation law (9), where new variables are defined as

$$\alpha_1 \equiv u_x^2 v_x^{-2} - 2v_x^{-1}, \quad \beta_1 \equiv u_x v_x^{-1}, \quad (12)$$

and

$$\alpha_{k+1} \equiv (v_x^{-1} \partial_x) \alpha_k, \quad \beta_{k+1} \equiv (v_x^{-1} \partial_x) \beta_k \quad (k \geq 1). \quad (13)$$

The conservation law (9) with the conserved density (11) could be directly proved with the following lemmas.

**Lemma 1.** *When the system(5a)-(5b) holds,  $\partial_t \alpha_k = 2u(\partial_x \alpha_k)$  ( $k \geq 1$ ),  $\partial_t \beta_1 = 2u(\partial_x \beta_1) - 2$  and  $\partial_t \beta_k = 2u(\partial_x \beta_k)$  ( $k \geq 2$ ).*

*Proof.* The evolution of  $\alpha_1$  and  $\beta_1$  is shown by direct calculation.

Both  $\alpha_k$  and  $\beta_k$  are defined by the same recursive relation  $\gamma_{k+1} = (v_x^{-1} \partial_x) \gamma_k$  ( $k \geq 1$ ). If we assume  $\partial_t \gamma_k = 2u(\partial_x \gamma_k) + c_k$ , where  $c_k$  is a certain constant, then based on the recursive relation, we have

$$\partial_t \gamma_{k+1} = -v_x^{-2} v_{xt} (\partial_x \gamma_k) + v_x^{-1} (\partial_x \partial_t \gamma_k) = 2u(-v_x^{-2} v_{2x} (\partial_x \gamma_k) + v_x^{-1} (\partial_x^2 \gamma_k)) = 2u(\partial_x \gamma_{k+1}).$$

Hence, the conclusion holds for all  $\alpha_k$ 's and  $\beta_k$ 's.  $\square$

For any  $n \in \mathbb{Z}_+$ , we introduce a transformation

$$\Gamma : (u_x, v_x, \dots, u_{nx}, v_{nx}) \mapsto (\alpha_1, \beta_1, \dots, \alpha_n, \beta_n),$$

where  $\alpha_k$ 's and  $\beta_k$ 's are defined by equations (12) and (13).

**Lemma 2.**  $\Gamma$  is invertible.

*Proof.* It is sufficient to formulate  $(u_x, v_x, \dots, u_{nx}, v_{nx})$  in terms of  $(\alpha_1, \beta_1, \dots, \alpha_n, \beta_n)$ . From equation (12), we immediately obtain

$$u_x = \frac{2\beta_1}{\beta_1^2 - \alpha_1}, \quad v_x = \frac{2}{\beta_1^2 - \alpha_1}.$$

From the recursive relation (13), we have

$$\begin{pmatrix} \alpha_k \\ \beta_k \end{pmatrix} = \begin{pmatrix} 2u_x v_x^{-k-1} & 2(v_x - u_x^2) v_x^{-k-2} \\ v_x^{-k} & -u_x v_x^{-k-1} \end{pmatrix} \begin{pmatrix} u_{kx} \\ v_{kx} \end{pmatrix} + \mathbb{G}_1(u_x, v_x, \dots, u_{(k-1)x}, v_{(k-1)x}),$$

where  $\mathbb{G}_1$  is a 2-dimensional vector function. The  $2 \times 2$  matrix in the right hand side is non-singular, so we could solve  $u_{kx}$  and  $v_{kx}$  step by step, and formulate them in terms of  $(\alpha_1, \beta_1, \dots, \alpha_k, \beta_k)$ .  $\square$

Lemmas 1 and 2 enable us to construct conserved densities explicitly depending on  $u$  and  $v$  for the system (5a)-(5b). Suppose that  $F(u, v, u_x, v_x, \dots, u_{nx}, v_{nx})$  is a conserved density such that

$$\partial_t \left( F(u, v, u_x, v_x, \dots, u_{nx}, v_{nx}) \right) = \partial_x \left( 2uF(u, v, u_x, v_x, \dots, u_{nx}, v_{nx}) \right). \quad (14)$$

According to lemma 2, any  $F(u, v, u_x, v_x, \dots, u_{nx}, v_{nx})$  could be reformulated in terms of  $(u, v, \alpha_1, \beta_1, \dots, \alpha_n, \beta_n)$  as

$$\hat{F}(u, v, \alpha_1, \beta_1, \dots, \alpha_n, \beta_n) = \Gamma F(u, v, u_x, v_x, \dots, u_{nx}, v_{nx}),$$

and with the help of lemma 1, the conservation law (14) could be calculated as

$$0 = \partial_t \hat{F} - \partial_x (2u\hat{F}) = -2v \frac{\partial \hat{F}}{\partial u} - 2 \frac{\partial \hat{F}}{\partial \beta_1} - \frac{4\beta_1}{\beta_1^2 - \alpha_1} \hat{F}.$$

Solving it by the standard method gives us conserved densities

$$\hat{F} = \frac{v_x}{2} G(v, \alpha_1, \beta_1 - uv^{-1}, \alpha_2, \beta_2, \dots, \alpha_n, \beta_n),$$

which takes conserved densities of the form (11) as special cases. The result is summarized as

**Proposition 1.** *On solutions of the generalized Riemann equation at  $N = 2$  (5a)-(5b), the conservation law (14) holds if and only if*

$$F(u, v, u_x, v_x, \dots, u_{nx}, v_{nx}) = \frac{v_x}{2} G(v, \alpha_1, \beta_1 - uv^{-1}, \alpha_2, \beta_2, \dots, \alpha_n, \beta_n),$$

where  $\alpha_k$ 's and  $\beta_k$ 's are defined by equations (12) and (13), and  $G$  is an arbitrary smooth function of its arguments.

In the rest part of the section, we manage to find new conservation laws other than (14) for the system (5a)-(5b). Let us consider a smooth function

$$H(u_x, v_x, \dots, u_{nx}, v_{nx}) = v_x P(\alpha_1, \beta_1, \dots, \alpha_n, \beta_n).$$

With the aid of lemma 1, it is straightforward to have

$$\partial_t H = \partial_x(2uv_x P) - 2v_x \frac{\partial P}{\partial \beta_1}.$$

If there exists  $Q(\alpha_1, \beta_1, \dots, \alpha_{n-1}, \beta_{n-1})$  such that

$$\partial_x Q = -2v_x \frac{\partial P}{\partial \beta_1}, \quad (15)$$

then  $H = v_x P(\alpha_1, \beta_1, \dots, \alpha_n, \beta_n)$  would be a conserved density such that

$$\partial_t H = \partial_x(2uH + Q).$$

Taking account of the recursive relation (13), we have

$$\partial_x Q = \sum_{k=1}^{n-1} \left( \frac{\partial Q}{\partial \alpha_k} (\partial_x \alpha_k) + \frac{\partial Q}{\partial \beta_k} (\partial_x \beta_k) \right) = v_x \sum_{k=1}^{n-1} \left( \frac{\partial Q}{\partial \alpha_k} \alpha_{k+1} + \frac{\partial Q}{\partial \beta_k} \beta_{k+1} \right).$$

Then equation (15) is rewritten as

$$\frac{\partial P}{\partial \beta_1} = -\frac{1}{2} \sum_{k=1}^{n-1} \left( \frac{\partial Q}{\partial \alpha_k} \alpha_{k+1} + \frac{\partial Q}{\partial \beta_k} \beta_{k+1} \right).$$

We summarize above discussions as

**Proposition 2.** *When the generalized Riemann equation at  $N = 2$  (5a)-(5b) holds, given an arbitrary smooth function  $Q(\alpha_1, \beta_1, \dots, \alpha_{n-1}, \beta_{n-1})$ , let*

$$P = -\frac{1}{2} \int \sum_{k=1}^{n-1} \left( \frac{\partial Q}{\partial \alpha_k} \alpha_{k+1} + \frac{\partial Q}{\partial \beta_k} \beta_{k+1} \right) d\beta_1, \quad (16)$$

then  $H = v_x P$  is conserved and satisfies the conservation law  $\partial_t H = \partial_x(2uH + Q)$ .

As an implementation of proposition 2, we construct a conserved density from the smooth function  $Q(\alpha_1)\beta_1$ . Following the formula (16), we have

$$P = -\frac{1}{2} \int \left( Q'(\alpha_1)\alpha_2\beta_1 + Q(\alpha_1)\beta_2 \right) d\beta_1 = -\frac{1}{4}Q'(\alpha_1)\alpha_2\beta_1^2 - \frac{1}{2}Q(\alpha_1)\beta_1\beta_2 + T(\alpha_1, \alpha_2, \beta_2),$$

where  $Q'(\alpha_1) = \partial Q/\partial\alpha_1$  and  $T(\alpha_1, \alpha_2, \beta_2)$  is the ‘‘constant’’ of integration. Then, we obtain a conserved density

$$H(u_x, v_x, u_{xx}, v_{xx}) = -\frac{1}{4}v_x Q'(\alpha_1)\alpha_2\beta_1^2 - \frac{1}{2}v_x Q(\alpha_1)\beta_1\beta_2 + v_x T(\alpha_1, \alpha_2, \beta_2)$$

such that

$$\partial_t H(u_x, v_x, u_{xx}, v_{xx}) = \partial_x (2uH(u_x, v_x, u_{xx}, v_{xx}) + Q(\alpha_1)\beta_1). \quad (17)$$

We notice that  $v_x T(\alpha_1, \alpha_2, \beta_2)$  is also a conserved density as implied by proposition 1, and satisfies

$$\partial_t (v_x T(\alpha_1, \alpha_2, \beta_2)) = \partial_x (2uv_x T(\alpha_1, \alpha_2, \beta_2)).$$

So the conservation law (17) could be simplified by replacing  $H(u_x, v_x, u_{xx}, v_{xx})$  by

$$-\frac{1}{4}v_x Q'(\alpha_1)\alpha_2\beta_1^2 - \frac{1}{2}v_x Q(\alpha_1)\beta_1\beta_2.$$

To conclude this section, we comment on that regarding the generalized Riemann equation at  $N = 2$  (5a)-(5b), while above two propositions provide most conserved densities, they by no means exhaust all possibilities since there are extra conservation laws explicitly depending on time,<sup>9</sup> such as

$$\begin{aligned} \partial_t (2tu_x v + v) &= \partial_x (t(4uu_x v - 2v^2) + 2uv), \\ \partial_t (u + 2tv + 2t^2 u_x v) &= \partial_x (t^2(4uu_x v - 2v^2) + 4tuv + u^2). \end{aligned}$$

### III. CONSERVATION LAWS OF RELATED SYSTEMS

Under appropriate changes of variables and/or reductions, the generalized Riemann equation at  $N = 2$  (5a)-(5b) is converted/reduced to the Gurevich-Zybin system, Monge-Ampere equation, two-component Hunter-Saxton equation and Hunter-Saxton equation. In this section, we will show that conserved densities of these systems could be easily deduced from those presented in section II.



## A. The Gurevich-Zybin system and Monge-Ampere equation

For the generalized Riemann equation at  $N = 2$  (5a)-(5b), we introduce  $u = -\hat{u}/2$  and  $v = -\Phi_x/4$ . Then equation (5a) is changed to

$$\hat{u}_t + \hat{u}\hat{u}_x + \Phi_x = 0, \quad (18)$$

while equation (5b) to

$$\Phi_{xt} + \hat{u}\Phi_{xx} = 0. \quad (19)$$

Differentiating equation (19) with respect to  $x$  and letting  $\rho = \Phi_{xx}$ , we obtain

$$\rho_t + \partial_x(\hat{u}\rho) = 0. \quad (20)$$

The coupled system (18) and (20) (N.B.  $\rho = \Phi_{xx}$ ) is nothing but the Gurevich-Zybin system in 1-dimensional space,<sup>13,14</sup> which has been shown to be linearisable and possesses infinitely many local Hamiltonian structures, local Lagrangian representations and local conservation laws (symmetries).<sup>5</sup> As observed by Pavlov, solving  $\hat{u}$  from equation (19) and substituting it into a conservation law of the Gurevich-Zybin system, i.e.

$$(\rho\hat{u})_t + (\rho\hat{u}^2 + \Phi_x^2/2)_x = 0,$$

we obtain the Monge-Ampere equation

$$\Phi_{tt} = \frac{\Phi_{xt}^2}{\Phi_{xx}} + \frac{1}{2}\Phi_x^2. \quad (21)$$

Taken the above connection into consideration, their conserved densities are easily deduced from those of the system (5a)-(5b). For instance, inferred from the conserved density (11), the Gurevich-Zybin system (18) and (20) has conserved densities of the form

$$\rho G(\bar{\alpha}_1, \bar{\alpha}_2, \bar{\beta}_2, \dots, \bar{\alpha}_n, \bar{\beta}_n),$$

such that the conservation law

$$\partial_t(\rho G) = \partial_x(-\hat{u}\rho G),$$

where  $\bar{\alpha}_1 = (\hat{u}_x^2 + 2\rho)\rho^{-2}$ ,  $\bar{\beta}_1 = \hat{u}_x\rho^{-1}$  and

$$\bar{\alpha}_{k+1} = (\rho^{-1}\partial_x)\bar{\alpha}_k, \quad \bar{\beta}_{k+1} = (\rho^{-1}\partial_x)\bar{\beta}_k \quad (k \geq 1).$$

Similarly, replacing  $u$  by  $\Phi_{xt}/(2\Phi_{xx})$ , while  $v$  by  $-\Phi_x/4$  in the conserved density (11) yields conserved densities of the Monge-Ampere equation (21).

## B. Two-component Hunter-Saxton equation

Let  $v_x = (u_x^2 + \eta^2)/2$ , then the system (6a)-(6b) is converted to

$$u_{xt} = 2uu_{xx} + u_x^2 - \eta^2, \quad (22a)$$

$$\eta_t = 2(u\eta)_x, \quad (22b)$$

which obviously reduces to the Hunter-Saxton equation<sup>7,8</sup> when  $\eta = 0$ , and is referred to as two-component Hunter-Saxton equation.<sup>2,5</sup>

Replacing  $v_x$  by  $(u_x^2 + \eta^2)/2$  in (11) immediately gives us conserved densities

$$(u_x^2 + \eta^2)G_1(\tilde{\alpha}_1, \tilde{\alpha}_2, \tilde{\beta}_2, \dots, \tilde{\alpha}_n, \tilde{\beta}_n) \quad (23)$$

for the two-component Hunter-Saxton equation (22a)-(22b), where  $\tilde{\alpha}_1 = \eta^2(u_x^2 + \eta^2)^{-2}$ ,  $\tilde{\beta}_1 = u_x(u_x^2 + \eta^2)^{-1}$  and

$$\tilde{\alpha}_{k+1} = (u_x^2 + \eta^2)^{-1} \partial_x \tilde{\alpha}_k, \quad \tilde{\beta}_{k+1} = (u_x^2 + \eta^2)^{-1} \partial_x \tilde{\beta}_k \quad (k \geq 1)$$

and the corresponding conservation law is given by

$$\partial_t \left( (u_x^2 + \eta^2)G_1 \right) = \partial_x \left( 2u(u_x^2 + \eta^2)G_1 \right)$$

When  $\eta = 0$ , all  $\tilde{\alpha}_k$ 's vanish and the quantity (23) reduces to

$$u_x^2 G_2 \left( (u_x^{-2} \partial_x) u_x^{-1}, (u_x^{-2} \partial_x)^2 u_x^{-1}, \dots, (u_x^{-2} \partial_x)^{n-1} u_x^{-1} \right),$$

where  $G_2 = G_1|_{\eta=0}$ . Thus we recover the conserved density with arbitrary function of the Hunter-Saxton equation recently reported by Tian and Liu.<sup>11</sup>

## IV. THE GENERALIZED RIEMANN EQUATIONS AT $N = 3, 4$

In this section, we will construct conserved densities with arbitrary smooth functions for the generalized Riemann equations at  $N = 3, 4$ . To this end, we will introduce new dynamical variables in such a way that their evolutions take simple forms. All results could be verified as we did in section II, so detailed proofs are omitted.

### A. $N = 3$

By rescaling dependent variables as  $u_1 = -2u, u_2 = 4v$  and  $u_3 = -8w$ , the generalized Riemann system at  $N = 3$  is rewritten as

$$u_t = 2uu_x - 2v, \quad (24a)$$

$$v_t = 2vv_x - 2w, \quad (24b)$$

$$w_t = 2ww_x. \quad (24c)$$

To get conserved densities with concise expressions, let

$$\alpha_1 \equiv (v_x^2 - 2u_x w_x)w_x^{-2}, \quad \beta_1 \equiv (v_x^3 + 3w_x^2 - 3u_x v_x w_x)w_x^{-3}, \quad \gamma_1 \equiv v_x w_x^{-1},$$

and

$$\alpha_{k+1} \equiv (w_x^{-1} \partial_x)^k \alpha_1, \quad \beta_{k+1} \equiv (w_x^{-1} \partial_x)^k \beta_1, \quad \gamma_{k+1} \equiv (w_x^{-1} \partial_x)^k \gamma_1 \quad (k \geq 1)$$

then in the same way as we proved lemmas 1 and 2, we obtain

**Lemma 3.** *when the system (24a)-(24c) holds,  $\partial_t \alpha_k = 2u(\partial_x \alpha_k)$ ,  $\partial_t \beta_k = 2u(\partial_x \beta_k)$  ( $k \geq 1$ ),  $\partial_t \gamma_1 = 2u(\partial_x \gamma_1) - 2$  and  $\partial_t \gamma_k = 2u(\partial_x \gamma_k)$  ( $k \geq 2$ ).*

**Lemma 4.** *For any  $n \in \mathbb{Z}_+$ , the transformation*

$$\Gamma_2 : (u_x, v_x, w_x, \dots, u_{nx}, v_{nx}, w_{nx}) \mapsto (\alpha_1, \beta_1, \gamma_1, \dots, \alpha_n, \beta_n, \gamma_n)$$

*is invertible.*

Regarding the generalized Riemann equation at  $N = 3$  (24a)-(24c), suppose that

$$F = F(u, v, w, u_x, v_x, w_x, \dots, u_{nx}, v_{nx}, w_{nx})$$

is a conserved density such that

$$\partial_t F = \partial_x (2uF). \quad (25)$$

According to lemmas 3 and 4, the conservation law (25) could be solved in coordinates  $(u, v, w, \alpha_1, \beta_1, \gamma_1, \dots, \alpha_k, \beta_k, \gamma_k)$ .

**Proposition 3.** *Equation (25) is qualified as a conservation law of the system (24a)-(24c) if and only if*

$$F = w_x G \left( w, v^2 - 2wu, \alpha_1, \beta_1, \gamma_1 - w^{-1}v, \alpha_2, \beta_2, \gamma_2, \dots, \alpha_n, \beta_n, \gamma_n \right).$$

In the spirit of proposition 2, we also find conservation laws of the form

$$\partial_t \left( w_x P \right) = \partial_x \left( 2u w_x P + Q \right)_x,$$

where  $Q = Q(\alpha_1, \beta_1, \gamma_1, \dots, \alpha_{n-1}, \beta_{n-1}, \gamma_{n-1})$  is an arbitrary smooth function and

$$P = -\frac{1}{2} \int \sum_{k=1}^{n-1} \left( \frac{\partial Q}{\partial \alpha_k} \alpha_{k+1} + \frac{\partial Q}{\partial \beta_k} \beta_{k+1} + \frac{\partial Q}{\partial \gamma_k} \gamma_{k+1} \right) d\gamma_1.$$

## B. $N = 4$

The generalized Riemann equation at  $N = 4$  is given by

$$u_t = 2u u_x - 2v, \tag{26a}$$

$$v_t = 2v v_x - 2w, \tag{26b}$$

$$w_t = 2w w_x - 2z, \tag{26c}$$

$$z_t = 2u z_x, \tag{26d}$$

where  $u = -u_1/2$ ,  $v = u_2/4$ ,  $w = -u_3/8$  and  $z = u_4/16$ . Let

$$\begin{aligned} \alpha_1 &\equiv (w_x^2 - 2v_x z_x) z_x^{-2}, & \beta_1 &\equiv (v_x^2 - 2u_x w_x + 2z_x) z_x^{-2}, \\ \gamma_1 &\equiv (w_x^3 - 3z_x v_x w_x + 3z_x^2 u_x) z_x^{-3}, & \delta_1 &\equiv w_x z_x^{-1}, \end{aligned}$$

and

$$\begin{aligned} \alpha_{k+1} &\equiv (z_x^{-1} \partial_x)^k \alpha_1, & \beta_{k+1} &\equiv (z_x^{-1} \partial_x)^k \beta_1, \\ \gamma_{k+1} &\equiv (z_x^{-1} \partial_x)^k \gamma_1, & \delta_{k+1} &\equiv (z_x^{-1} \partial_x)^k \delta_1 \quad (k \geq 1). \end{aligned}$$

then in terms of these new dynamical variables, we have

**Lemma 5.** *When the system (26a)-(26d) holds,  $\partial_t \alpha_k = 2u(\partial_x \alpha_k)$ ,  $\partial_t \beta_k = 2u(\partial_x \beta_k)$ ,  $\partial_t \gamma_k = 2u(\partial_x \gamma_k)$  ( $k \geq 1$ ),  $\partial_t \delta_1 = 2u(\partial_x \delta_1) - 2$  and  $\partial_t \delta_k = 2u(\partial_x \delta_k)$  ( $k \geq 2$ ).*

**Lemma 6.** *For any  $n \in \mathbb{Z}_+$ , the transformation*

$$\Gamma_3 : (u_x, v_x, w_x, z_x, \dots, u_{nx}, v_{nx}, w_{nx}, z_{nx}) \mapsto (\alpha_1, \beta_1, \gamma_1, \delta_1, \dots, \alpha_n, \beta_n, \gamma_n, \delta_n)$$

*is invertible.*

On the basis of above lemmas, the following results are proved.

**Proposition 4.**  $F = F(u, v, w, z, \dots, u_{nx}, v_{nx}, w_{nx}, z_{nx})$  is a conserved density such that  $\partial_t F = \partial_x(2uF)$  if and only if

$$F = z_x G \left( z, w^2 - 2vz, \frac{3uz^2 - 3vwz + w^3}{z^2}, \alpha_1, \beta_1, \gamma_1, \delta_1 - \frac{w}{z}, \alpha_2, \beta_2, \gamma_2, \delta_2, \dots, \alpha_n, \beta_n, \gamma_n, \delta_n \right).$$

**Proposition 5.** For any smooth function  $Q = Q(\alpha_1, \beta_1, \gamma_1, \delta_1, \dots, \alpha_{n-1}, \beta_{n-1}, \gamma_{n-1}, \delta_{n-1})$ ,

$$\partial_t (z_x P) = \partial_x (2uz_x P + Q),$$

is a conservation law of the system (26a)-(26d) if

$$P = -\frac{1}{2} \int \sum_{k=1}^{n-1} \left( \frac{\partial Q}{\partial \alpha_k} \alpha_{k+1} + \frac{\partial Q}{\partial \beta_k} \beta_{k+1} + \frac{\partial Q}{\partial \gamma_k} \gamma_{k+1} + \frac{\partial Q}{\partial \delta_k} \delta_{k+1} \right) d\delta_1.$$

## V. CONCLUSION

It is well known that the existence of infinitely many conserved densities dependent on arbitrary function implies strong integrability for a system. In this paper, by introducing special variables we obtain infinitely many conserved densities (two categories) for the generalized Riemann equations at  $N = 2, 3$  and  $4$ . Through the changes of variables or reductions, conserved densities for related systems are obtained. It is not difficult to find that the introduced variables such as  $\alpha_k, \beta_k, \gamma_k, \delta_k$  whose evolutions are simple all take rational form and could be endowed with degree. Thus it may be possible to consider the generalized Riemann equation for the general  $N$ . However in such case we do not have ready results.

## ACKNOWLEDGMENTS

This paper is supported by the National Natural Science Foundation of China (grant numbers: 11271366, 11331008 and 11505284) and the Fundamental Research Funds for Central Universities.

## REFERENCES

- <sup>1</sup>Whitham, G. B., "Linear and nonlinear waves" (Wiley, Canada, 1999).
- <sup>2</sup>Popowicz, Z. and Prykarpatsky, A. K., "The non-polynomial conservation laws and integrability analysis of generalized Riemann type hydrodynamical equations," Nonlinearity. **23**(10), 2517–2537 (2010).

- <sup>3</sup>Golenia, J., Pavlov, M. V., Popowicz, Z., and Prykarpatsky, A. K., “On a nonlocal Ostrovsky-Whitham type dynamical system, its Riemann type inhomogeneous regularizations and their integrability,” *SIGMA*. **6**, 002 (2010).
- <sup>4</sup>Prykarpatsky, A. K., Artemovych, O. D., Popowicz, Z., and Pavlov, M. V., “Differential-algebraic integrability analysis of the generalized Riemann type and Korteweg-de Vries hydrodynamical equations,” *J. Phys. A: Math. Theor.* **43**(29), (2010)295205.
- <sup>5</sup>Pavlov, M. V., “The Gurevich-Zybin system,” *J. Phys. A: Math. Gen.* **38**(17), 3823–3840 (2005).
- <sup>6</sup>Brunelli, J. C. and Das, A., “On an integrable hierarchy derived from the isentropic gas dynamics,” *J. Math. Phys.* **45**(7), 2633–2645 (2004).
- <sup>7</sup>Hunter, J. K. and Saxton, R., “Dynamics of director fields,” *SIAM J. Appl. Math.* **51**(6), 1498–1521 (1991).
- <sup>8</sup>Hunter, J. K. and Zheng, Y., “On a completely integrable nonlinear hyperbolic variational equation,” *Physica. D.* **79**(2-4), 361–384 (1994).
- <sup>9</sup>Popowicz, Z., “The matrix Lax representation of the generalized Riemann equations and its conservation laws,” *Phys. Lett. A.* **375**(37), 3268–3272 (2011).
- <sup>10</sup>Olver, P. J. and Nutku, Y., “Hamiltonian structures for systems of hyperbolic conservation laws,” *J. Math. Phys.* **29**(7), 1610–1619 (1988).
- <sup>11</sup>Tian, K. and Liu, Q. P., “Conservation laws and symmetries of Hunter-Saxton equation: revisited,” *Nonlinearity.* **29**(3), 737–755 (2016).
- <sup>12</sup>Sakovich, S., “On a Whitham-type equation,” *SIGMA*. **5**, 101 (2009).
- <sup>13</sup>Gurevich, A. V. and Zybin, K. P., “Nondissipative gravitational turbulence,” *Sov. Phys. JETP.* **67**(1), 1–12 (1988).
- <sup>14</sup>Gurevich, A. V. and Zybin, K. P., “Large-scale structure of the Universe. Analytic theory,” *Sov. Phys. Usp.* **38**(7), 687–722 (1995).