Level rearrangement in exotic-atom-like three-body systems

Jean-Marc Richard^{1,*} and Claude Fayard^{1,†}

¹Institut de Physique Nucléaire de Lyon, Université de Lyon, CNRS-IN2P3-UCBL, 4, rue Enrico Fermi, Villeurbanne, France (Dated: July 7, 2021)

We study systems of three bosons bound by a long-range interaction supplemented by a short-range potential of variable strength. This generalizes the usual two-body exotic atoms where the Coulomb interaction is modified by nuclear forces at short distances. The energy shift due to the short-range part of the interaction combines two-body terms similar to the ones entering the Trueman-Deser formula, and three-body contributions. A sudden variation of the energy levels is observed near the coupling thresholds of the short-range potential. But the patterns of rearrangement are significantly modified as compared to the two-body case.

I. INTRODUCTION

Exotic atoms have a long history, and have stimulated interesting developments in the quantum dynamics of systems involving both long-range and short-range forces. For the exploratory studies presented here, the refinements of effective theories [1] are not required, and we shall restrict ourselves to the Schrödinger framework, as reviewed, e.g., in [2] for the three-dimensional case, and extended in [3] for the two-dimensional one.

In units simplifying the treatment of the pure Coulomb case, where the strength becomes 1 and the reduced mass 1/2, an exotic atom can be modeled as

$$-\, \Delta \Psi + V\, \Psi = E\, \Psi \; , \quad V = -\frac{1}{r} + \lambda\, v(r) \; , \qquad \mbox{(1)} \label{eq:partial}$$

where r is the inter-particle distance and $\lambda v(r)$ the short-range correction, with a variable strength for the ease of discussion.

In most actual exotic atoms, there is a strong absorptive component in the short range interaction, so one has to use either a complex (optical) potential v or a coupled-channel formalism. Probably, the (\bar{D}_s, p) atom, with a proton and an anticharmed meson $\bar{D}_s = (\bar{c}s)$ of charge -1 and strangeness -1, escapes any absorption, since it lies below any threshold such as $(\bar{D}^0(\bar{c}u) \Lambda(sud))$, but it is not yet accessible experimentally. In this study, we make the somewhat drastic simplification of a purely real short-range term v.

When the above spectral problem is solved, the most striking observations are:

1. The energy shift $\delta E = E - E_n$, as compared to the pure-Coulomb energies $E_n = -1/(4\,n^2)$, is often rather small, but is usually not given by ordinary perturbation theory: for instance, an infinite hard-core of small radius corresponds to a small energy shift but to an infinite first order correction.

2. Each energy $E(\lambda)$, as a function of the strength parameter λ , is almost flat in a wide interval of λ , with a value close to some E_n , one of the pure Coulomb energies. For S-wave states δE is well approached by a formula by Deser et al., and Trueman [4],

$$\delta E \simeq \frac{a}{2 \, n^3} \,, \tag{2}$$

where a is the scattering length in the short-range interaction $\lambda v(r)$ alone.

- 3. If v(r) is attractive, when the strength λ approaches one of the positive critical values at which a first or a new bound state appears in the spectrum of λv alone, the energy $E(\lambda)$ quits its plateau and drops dramatically from the region of atomic energies to the one of deep nuclear binding. It is rapidly replaced in the plateau by the next level. This is known as *level rearrangement* [5]. Note that level rearrangement disappears if absorption becomes too strong [2, 6].
- 4. The above patterns are more general, and hold for any combination of a long-range and a short-range interaction, say $V=V_0+\lambda\,v(r)$, as encountered, e.g., in the physics of cold atoms where a long-range confining interaction is supplemented by the direct interaction among the atoms [7]. The Deser-Trueman formula of S-states is generalized as

$$\delta E \simeq 4\pi \, |\Psi_0(0)|^2 \, a \,,$$
 (3)

where Ψ_0 is the normalized wave-function in the external potential V_0 . If V_0 supports only one bound state, then the rearrangement "extracts" states from the continuum, instead of shifting radial excitations that are already bound. An illustration is given in Fig. 1, with a superposition of two exponential potentials of range parameters $\mu=1$ and $\mu=100$, namely

$$V(r) = 2 v_e(1, r) + \lambda v_e(100, r) ,$$

$$v_e(\mu, r) = -1.4458 \mu^2 \exp(-\mu r) ,$$
(4)

where $v_e(\mu, r)$ is tuned to start binding at unit strength.

^{*} j-m.richard@ipnl.in2p3.fr

[†] c.fayard@ipnl.in2p3.fr




FIG. 1. (Color online) Level rearrangement for a two-body system bound by the potential (4). The dashed line indicates the approximation corresponding to the Deser-Trueman formula (3). The dotted vertical lines show the coupling thresholds for the short-range part of the interaction.

5. There are several possible improvements and alternative formulations of (3). For instance, a, the bare scattering length in $\lambda v(r)$ can be replaced by the "longrange corrected" scattering length where the solutions of the radial equation for $\lambda v(r)$ are matched to the eigenfunction of the external potential. See, e.g., [8], and refs. there. In the physics of cold atoms, one is more familiar with the approach by Busch et al. [7]. It deals with the case of an harmonic oscillator modified at short distances, but the derivation can be generalized as follows. Let $u(E,r) = \sqrt{4\pi} r \Psi$ the s-wave reduced radial wavefunction for $V_0(r)$ that is regular at large r, at energy E < 0, with some normalization, e.g., $u(E,r) \exp(r\sqrt{-E}) \to 1$ for $r \to \infty$. The levels in V_0 correspond to the quantization condition $u(E_0,0) = 0$, where E_0 is an eigenenergy of V_0 , for instance the ground state. When a point-like interaction of scattering length a is added, then the boundary condition is modified into

$$u(E,0) + a \partial_r u(E,0) = 0, \qquad (5)$$

which can be expanded near E_0 to give

$$\delta E = E - E_0 \simeq -a \frac{\partial_r u(E_0, 0)}{\partial_E u(E_0, 0)}. \tag{6}$$

Now, the equivalence of (6) and (3), rewritten as $\delta E \simeq a \left[\partial_r u_N(E_0,0)\right]^2$, where u_N is the normalized version of u, comes from the relation

$$\partial_r u(E_0, 0) \, \partial_E u(E_0, 0) = -\int_0^\infty u(E_0, r)^2 \, \mathrm{d}r \,,$$
 (7)

which is easily derived from the Wronskian identity, widely used in some textbooks [9], here applied to energies E and E_0 .

6. The generalization to a number of dimension $d \neq 3$ is straightforward for d > 3. For d = 1, the first plateau is avoided, as the short-range potential, if attractive, develops its own discrete spectrum for any $\lambda > 0$. The case of d = 2 is more delicate: see, e.g., [6, 10].

Our aim here is to present a first investigation of the three-body analog of exotic atoms. There are already studies of systems such as (K^-,d) , where d=(p,n) is the deuteron, in which the neutron feels only the shortrange part of the interaction. We will study systems in which the three constituents are already bound by the long-range component of the potential. this is the first attempt, at least to our knowledge. We consider three identical bosons, relevant for three atoms in a confining trap¹. We have in mind less symmetric systems for future work. We address the following questions: Is there a pattern similar to the level rearrangement? Is there a generalization of the Deser-Trueman formula? What are the similarities with the case where the long-range interaction is replaced by an overall harmonic confinement?

Note that the occurrence of plateaus and sudden drops of the energies as a function of the coupling strength is not very usual, as these energies are monotonic and concave functions of any parameter entering linearly the Hamiltonian [11]. For instance, in a pioneering study of three-boson energies, Osborn [12] fund some type of rearrangements in the three-body spectrum corresponding to a Yukawa interaction, but it was later acknowledged that this calculation suffers from some numerical instability, as the computed Faddeev energies violate a rigorous lower bound [12, 13]. Unfortunately, the erroneous plot was reproduced in a seminal textbook on the three-body problem [14],

The paper is organized as follows. In Sec. II, we give some basic reminders about the spectrum of a three-boson systems from the Borromean limit of a single bound state to the regime of stronger binding, with a word about the numerical techniques. The results corresponding to a superposition $V_0(r) + \lambda v(r)$ are displayed in Sec. III. An interpretation is attempted in Sec. IV, with a three-body version of the Deser-Trueman formula. Section V is devoted to our conclusions.

II. THE THREE-BOSON SPECTRUM WITH A SIMPLE POTENTIAL

If two bosons interact through an attractive potential, or a potential with attractive parts, $\lambda \, v_0(r)$, a minimal strength is required to achieve binding, say $\lambda > \lambda_2^{\rm cr}$. A collection of values of $\lambda_2^{\rm cr}$ can be found, e.g., in the classic paper by Blatt and Jackson [15]. In the following, we shall normalize v_0 so that $\lambda_2^{\rm cr} = 1$.

If one assumes that $v_0(r)$ is attractive everywhere, once two bosons are bound, the 3-boson system is also bound. (The case of potentials with a strong inner repulsion would require a more detailed analysis which is beyond the scope of this preliminary investigation.) This

At least for harmonic confinement, an external potential can be rewritten as a sum of pair interactions

means that for a single monotonic potential

$$V_0 = \lambda \left[v_0(r_{12}) + v_0(r_{23}) + v_0(r_{31}) \right] , \tag{8}$$

where $r_{ij} = |\boldsymbol{r}_j - \boldsymbol{r}_i|$, the minimal coupling to achieve three-body binding, $\lambda = \lambda_3^{\rm cr}$, is less than 1, in our units. This is implicit in the seminal paper by Thomas [16]. The inequality $\lambda_3^{\rm cr} < \lambda_2^{\rm cr}$ is now referred to as "Borromean binding", after the study of neutron halos in nuclear physics [17]. One gets typically $\lambda_3^{\rm cr} \simeq 0.8$, i.e., about 20% of Borromean window [18].

In short, the three-boson spectrum has the following patterns:

- For $\lambda < \lambda_3^{\rm cr}$, no binding
- For $\lambda_3^{\rm cr} < \lambda < 1$, a single Borromean bound state, and, for λ close to 1, a second three-body bound state just below the two-body energy,
- Very near $\lambda = 1$, the very weakly bound Efimov states.
- For $1 < \lambda$, two bound states below the 2+1 breakup, and further bound states when λ becomes very large.

These patterns are independent of the detailed shape (exponential, Gaussian, ...) of the potential $v_0(r)$, if monotonic. For potentials with an internal or external repulsive barrier, one expects some changes, as already the width of the Borromean window is modified [19]. This is illustrated in Fig. 2 for the case of an exponential potential.

The first excited state can be considered as the first member of the sequence of the Efimov states occurring near $\lambda=1$. However, it differs from the other Efimov states in the sense that when the coupling is increased, it remains below the two-body break-up threshold, at least for the simple monotonic potentials we consider here. ²

The two-body energy is know analytically for a single exponential potential. For the three-body energies, we first made some investigations based on the hyperspherical expansion [20]. The results were finalized with a variational method based on either exponential of Gaussian wavefunctions, say

$$\Psi = \sum_{i} \gamma_{i} \left[\exp(-a_{i} r_{23}^{n} - b_{i} r_{31}^{n} - c_{i} r_{12}^{n}) + \cdots \right] , \quad (9)$$

where the dots stand for terms deduced by permuting the particles. For a given choice of range parameters a_i , b_i and c_i , the Schrödinger equation results in a generalized eigenvalue equation, whose eigenvectors are the




FIG. 2. (Color online) Two-boson (red) and three-boson spectrum (blue) in the potential (8), with $v_0(r) \propto -\exp(-r)$ at small coupling (top) and larger coupling (bottom). The first excited 3-body state (dashed) is always found below the 2-body energy. The second-excited state (dotted line in the bottom plot) and higher states (not shown) require a minimal coupling to be bound below the dissociation threshold.

coefficients $\{\gamma_1, \gamma_2, \ldots\}$ and eigenvalues upper (variational) bounds on the exacts energies. The range parameters are thus tuned to minimize the energy levels. For n = 1, we wrote our own code: the range parameters a_i , b_i and c_i , if restricted to be real, are chosen to belong to a geometric progression $\{\alpha, \alpha r, \alpha r^2, \ldots\}$ of common ration r > 1 and scale factor α . The lowest term α can be linked to the energy E by the relation $\alpha^2 = 14 E/15$ suggested by the Feshbach-Rubinow equation [21]. For excited states, we found it more efficient to introduce complex range parameters a_i , b_i and c_i as done, e.g., by Korobov [22] for three charge ions in atomic physics, and to adjust the a_i , b_i and c_i by random trials. For n = 2, we used the code made available by Suzuki and Varga [23], with minor changes. Anyhow, our aim was not to produce very accurate benchmark energies, but to identify the main patterns.

III. THE THREE-BODY ENERGIES WITH LONG- AND SHORT-RANGE FORCES

We now replace (8) by a superposition

$$\sum_{i < j} [\lambda_0 \, v_e(1, r_{ij}) + \lambda \, v_e(\mu, r_{ij})] , \qquad (10)$$

where $v_e(\mu, r)$ is significantly shorter ranged than the external potential $v_e(1, r)$. In practice, we will choose μ ranging from 10 to 30. Note that the computations be-

² We thank Pascal Naidon (RIKEN) for a correspondence on this point.

come rather delicate for larger values of μ , and would require dedicated techniques.

In Fig. 3 are shown the spectra for $\lambda_0=2$, i.e., twice the two-body critical coupling and varying λ , for $\mu=10,20$ and 30. The unit of energy is irrelevant, as it




FIG. 3. (Color online) Spectrum in the potential (10) for (top to bottom) a ratio of ranges $\mu=10,\,20$ and 30, as a function of the strength of the short-range potential. Red: Two-body energy, Blue: three-body levels, solid line for the ground state, dashed for the first excitation, dotted for the second excitation, visible only for $\mu=30$ (for smaller μ , the second excited state requires a coupling λ outside of the range of the plot)

can be modified by an overall rescaling of the distances. Comments are in order:

- As in Fig. 1, a convex behavior as a function of λ is observed for the excited energy-levels, i.e., $E_n(\lambda)$ with n > 1. This is permitted, provided that the sum of the first energies remains concave [11].
- As in the two-body case, the transition is sharper when the range of the additional potential becomes shorter.
- There is clearly a beginning of rearrangement, in the sense that for $\lambda \to \lambda_3^{\rm cr} \simeq 0.8$, the excited state falls suddenly near the unperturbed ground-state energy.
- However, there is no second plateau for the excited state, just somewhat a smoothing of the fall-off for $\lambda \gtrsim \lambda_3^{\rm cr}$, together with a tendency to smooth or to invert the concavity.

• The second excited state, of energy E_3^{**} , becomes bound (i.e., below the energy E_2 of its dissociation threshold) for smaller and smaller values of the coupling λ . We get typically a coupling threshold of about 5 for a single exponential, about 1 for $\mu = 10$, and about $\lambda = 0.5$ for $\mu = 30$. The binding energy $E_2 - E_3^{**}$ has seemingly some delicate pattern, as it is not monotonic as a function of λ , at least in our calculations. We cannot exclude that for another choice of potentials, the second excited state becomes bound, then is reabsorbed by the two-body threshold, and eventually reappears for larger λ . In this scenario, the number of normalizable three-body bound states below the two-body threshold would not be a monotonic function of the strength λ , as in the Efimov effect.

IV. INTERPRETATION

Let us first concentrate on the region of small λ . One can estimate the energy shifts $(\delta E)_{ij}$ corresponding to several external potentials $V_{0,i}$ with $i=1,\ldots N$ and short-range potentials V_j , with $j=1,\ldots N'$ and N,N'>3, and study empirically the properties of the matrix $\{(\delta E)_{ij}\}$.

In the two-body case, one finds that the 2×2 sub-determinants vanish almost exactly. This is compatible with a factorization

$$\delta E^{(2)} \simeq A_{\rm LR} B_{\rm SR} \,, \tag{11}$$

as a product of a long-range term depending only on v_0 and a short-range term depending only on λv . This factorization is achieved by the Deser-Trueman formula, with $A_{\rm LR}$ being the square of the wave function at r=0 (times 4π) and $B_{\rm SR}$ the scattering length.

In the three-body case, it is observed that the 2×2 sub-determinants still nearly vanish, especially for the smaller values of the short-range strength λ , but that the 3×3 sub-determinants vanish even better (of course we compared the determinants divided by the typical values of a product of 2 or $3 \delta E$). This is compatible with δE being a sum of two factorized contributions,

$$\delta E^{(3)} = A_{LR} B_{SR} + A'_{LR} B'_{SR} . \tag{12}$$

As explained, e.g., in the textbook by Ericson and Weise [24], the Deser-Trueman formula (3) can be understood as the perturbative correction due to a Fermi point-potential that includes non perturbatively the effect of the short-range interaction. Thus for a symmetric three-body system, the same prescription leads to a simple extension of (3) that reads

$$A_{LR} B_{SR} = 12 \pi |\Psi_{12}(0)|^2 a$$
, (13)

where $|\Psi_{12}(0)|^2$ is a short-hand notation for the two-body correlation factor $\langle \Psi | \delta^{(3)}(\boldsymbol{r}_2 - \boldsymbol{r}_1) | \Psi \rangle / \langle \Psi | \Psi \rangle$. It is

checked that this term dominates for small shifts, i.e., for small λ , see Fig. 4. However, this term alone would induce a sharp decrease of the atomic energies only for $|a| \to \infty$, i.e., for $\lambda \to 1$, the coupling threshold for two-binding, and not near $\lambda = 0.8$, as actually observed. For an asymmetric three-body system, the extension is, in an obvious notation,

$$3\pi \sum_{i < j} |\Psi_{ij}(0)|^2 a_{ij} . {14}$$




FIG. 4. (Color online) Calculated energy (thick blue line, dashed for the excited state) vs. the estimate using the first term in Eq. (12) (thin black line). The calculation is done for a superposition of two exponentials, one with unit range and strength $\lambda_0=2$, and another of range parameter $\mu=50$ and strength λ .

The second contribution in Eq. (13) should thus account for the genuine three-body effects, besides some higher-order two-body terms such as a contribution from effective range. The three-body part of $B'_{\rm SR}$ is a kind of generalized scattering length that blows up when λ approaches the coupling threshold $\lambda^{\rm cr}_3$ for three-body binding. As the theory of three-body scattering is a little intricate, we postpone the precise definition of $B'_{\rm SR}$ to some further study. As for the long-range factor $A'_{\rm LR}$ of this second term, the simplest guess is to assume that it is proportional to the square of the wavefunction at $r_1 = r_2 = r_3$, or in terms of the Jacobi variables x and y describing the relative motion, $A'_{\rm LR} \propto \langle \Psi | \delta^{(3)}(x) \, \delta^{(3)}(y) | \Psi \rangle$, but this is seemingly not the case.

Our study of generalized exotic atoms is related to the Efimov physics. In particular, the authors of refs. [25, 26], and probably some others, have studied how the Efimov effect is modified if each atom is submitted to an individual harmonic confinement. They also found that near a point where the two-body scattering length becomes infinite, there is a finite number of three-body bound states, instead of an infinite number in absence of confinement. The second-excited three-body bound state in Fig. 3 is slightly reminiscent of an Efimov state of the short-range potential, modified by the long-range potential.

V. CONCLUSIONS

The lowest states of three-bosons have been calculated with a superposition of long-range and shortrange attractive potentials. When the strength λ of the latter is increased, starting from $\lambda = 0$, the three-body energies decreases very slowly, and can be well approximated by a straightforward generalization of the Deser-Trueman formula involving only the two-body scattering length. However, when λ approaches 0.8 (in units where $\lambda = 1$ is the coupling threshold for binding in the short-range potential alone), there is a departure for the Deser-Trueman formula, which can be empirically accounted for by the product of a short-range and a longrange factor. The short-range factor is the three-body analog of the scattering length and becomes very large when $\lambda \simeq 0.8$ which corresponds to the occurrence of a Borromean bound state in the short-range potential

This exploratory investigation has been done with a simple variational method based on a few exponential or Gaussians functions, which is sufficient to show the main trends. More powerful minimization methods are probably required for the states at the edge of stability with respect to spontaneous dissociation. We also studied some modeling with separable potentials. The qualitative patterns of rearrangement are observed for separable potentials of rank 2 or higher, with some slight differences with respect to the case of local potentials. This will be presented in a forthcoming article.

Many other developments are required. What is the precise definition of the three-body short-range factor? What is the corresponding long-range factor? What is the minimal ratio of range parameters required for the occurrence of the third stable three-body state? When does a fourth state show up? What are the analogs for $N \ge 4$ bosons? We also aim at studying some asymmetric systems. For instance, a prototype of (K^-pp) could be built, with a Coulomb interaction, that is known to produce a stable ion, below the threshold for breakup into a (K^-p) atom and an isolated proton [27]. Then the strong interaction between the two protons and the strange meson K^- could be mimicked by a simple potential of range about 1 fm, first real, and then, more realistically, complex to include absorptive effects, to study how the existence of a nuclear bound state (K^-pp) modifies the atomic spectrum.

ACKNOWLEDGMENTS

One of us (JMR) would like to thank ECT*, Trento, for the hospitality extended to him.

- [1] B. R. Holstein. Hadronic atoms and effective interactions. *Phys. Rev. D*, 60(11):114030–+, December 1999.
- [2] A. Gal, E. Friedman, and C. J. Batty. On the interplay between Coulomb and nuclear states in exotic atoms. *Nuclear Physics A*, 606:283–291, February 1996.
- [3] M. Combescure, C. Fayard, A. Khare, and J.-M. Richard. Exotic atoms in two dimensions. *Journal of Physics A Mathematical General*, 44(27):275302, July 2011.
- [4] S. Deser, M. L. Goldberger, K. Baumann, and W. Thirring. Energy Level Displacements in π-Mesonic Atoms. *Physical Review*, 96:774–776, November 1954.
 T. Trueman. Energy level shifts in atomic states of strongly-interacting particles. *Nucl. Phys. A*, 26:57–67,
- [5] Ya.B. Zel'dovich, "Energy levels in a distorted Coulomb field", Sov. J. Solid State, 1 (1960) 1497.

July 1961.

- [6] M. Combescure, A. Khare, A. K. Raina, J. M. Richard, and C. Weydert. Level Rearrangement in Exotic Atoms and Quantum Dots. *International Journal of Modern Physics B*, 21:3765–3781, 2007.
- [7] Thomas Busch, Berthold-Georg Englert, Kazimierz Rzażewski, and Martin Wilkens. Two cold atoms in a harmonic trap. Foundations of Physics, 28(4):549–559, 1998.
- [8] J. Thaler. Coulomb-nuclear interference in hadronic atoms. *Journal of Physics G Nuclear Physics*, 9:1009–1024, September 1983.
- [9] Albert Messiah. Quantum mechanics. Dover, Mineola, NY, 2014.
- [10] N. T. Zinner. Universal Two-Body Spectra of Ultracold Harmonically Trapped Atoms in Two and Three Dimensions. J. Phys., A45:205302, 2012.
- [11] Walter Thirring. A course in mathematical physics 3: quantum mechanics of atoms and molecules. Springer, Vienna, 1981.
- [12] Thomas Arthur Osborn. *Faddeev's equations for local potentials*. PhD thesis, Stanford University, 1967.
- [13] J. W. Humberston, R. L. Hall, and T. A. Osborn. Bound states of three identical bosons. *Physics Letters B*, 27:195–198, July 1968.
- [14] Erich W Schmid and Horst Ziegelmann. *The quantum mechanical three-body problem*. Vieweg Tracts Pure Appl. Phys. Vieweg, Braunschweig, 1974.
- [15] J. M. Blatt and J. D. Jackson. On the Interpretation of

- Neutron-Proton Scattering Data by the Schwinger Variational Method. *Physical Review*, 76:18–37, July 1949.
- [16] L. H. Thomas. The Interaction Between a Neutron and a Proton and the Structure of H³. *Physical Review*, 47:903– 909, June 1935.
- [17] M. V. Zhukov, B. V. Danilin, D. V. Fedorov, J. M. Bang, I. J. Thompson, and J. S. Vaagen. Bound state properties of Borromean halo nuclei: ⁶He and ¹¹Li. *Phys. Rep.*, 231:151– 199, August 1993.
- [18] J.-M. Richard and S. Fleck. Limits on the domain of coupling constants for binding N-body systems with no bound subsystems. *Physical Review Letters*, 73:1464–1467, September 1994.
- [19] Steven Moszkowski, Sonia Fleck, Ali Krikeb, Lukas Theußl, Jean-Marc Richard, and Kálmán Varga. Binding three or four bosons without bound subsystems. *Phys. Rev. A*, 62:032504, Aug 2000.
- [20] R. Krivec. FBS Colloquium Hyperspherical-Harmonics Methods for Few-Body Problems. Few-Body Systems, 25:199–238, 1998.
- [21] H. Feshbach and S. I. Rubinow. Equivalent Two-Body Method for the Triton. *Physical Review*, 98:188–193, April 1955.
- [22] V. I. Korobov. Coulomb three-body bound-state problem: Variational calculations of nonrelativistic energies. *Phys. Rev. A* , 61(6):064503, June 2000.
- [23] K. Varga and Y. Suzuki. Solution of few-body problems with the stochastic variational method I. Central forces with zero orbital momentum. *Computer Physics Communications*, 106:157–168, October 1997.
- [24] Torleif Eric Oskar Ericson, Wolfram Weise, and W Weise. Pions and nuclei. Internat. Ser. Mono. Phys. Clarendon Press, Oxford, 1988.
- [25] M. Thøgersen, D. V. Fedorov, and A. S. Jensen. Universal properties of Efimov physics beyond the scattering length approximation. *Phys. Rev. A*, 78(2):020501, August 2008.
- [26] J. Portegies and S. Kokkelmans. Efimov Trimers in a Harmonic Potential. Few-Body Systems, 51:219–234, November 2011.
- [27] R. N. Hill. Proof that the H⁻ ion has only one bound state. Details and extension to finite nuclear mass. *Journal of Mathematical Physics*, 18:2316–2330, December 1977.