

Positron production in collision of heavy nuclei

I.B. Khriplovich¹

Department of Physics, St. Petersburg State University, 7/9 Universitetskaya Nab.,
St. Petersburg 199034, Russia

Abstract

We consider the electromagnetic production of positron in collision of slow heavy nuclei, with the simultaneously produced electron captured by one of the nuclei. The cross-section of the discussed process exceeds essentially the cross-section of e^+e^- production.

Keywords: positron production, heavy nuclei

The positron production in collision of slow heavy nuclei (as well as the production of e^+e^- pair), was addressed in numerous papers (see, for instance, [1 - 3]). In the present note we reconsider the problem of the positron production in collision of heavy nuclei, with the electron captured by another nucleus. We present a simple direct solution of this problem.

The process under discussion is presented in Fig. 1




Figure 1: Feynman diagram of positron production.

The heavy lines in the Figure refer to the propagation of nuclei. The wavy line refers to the virtual photon. In the produced pair e^+e^- , the electron is captured immediately by one of the nuclei. As to the positron, we assume that its interaction with the nuclei can be neglected. Simple symmetry arguments demonstrate that this repulsive interaction cannot be strong.

Then, it can be easily demonstrated that diagram Fig. 1 is the only one which survives in the limit of heavy mass $M \rightarrow \infty$ of colliding nuclei.

Let us come back now to Figure 1. The electromagnetic vertex on the upper heavy line reduces to $(p' + p)/2M \approx v$ (we neglect here the difference between p' and p). With the amplitude proportional to v , the discussed total cross-section is proportional to v also:

$$v^2/v = v; \tag{1}$$

¹khios231@mail.ru

v in the denominator of (1) originates from the particle flux. For the velocity of nuclei, we assume $v \simeq 0.1$.

Thus, we can confine to diagram 1, with the overall factor $Z^2\alpha^2$ in the cross-section. One factor 2 in this expression arises since the electron can be captured by any of the two nuclei. One more factor 2 is due to two polarizations of the produced positron.

The total cross-section is obviously a scalar, and its essential part is $1/m^2$.

In this way, we arrive at the following result for the total cross-section of the positron production:

$$\sigma = 4 Z^2 \alpha^2 \frac{v}{m^2}. \quad (2)$$

Numerically, this cross-section is

$$\sigma \simeq 10^{-22} \text{ cm}^2. \quad (3)$$

This cross-section exceeds essentially the cross-section of e^+e^- production in collision of the uranium nuclei, which is close to 10^{-25} cm^2 .

Acknowledgements

I am grateful to V.M. Shabaev for useful discussions.

References

- [1] U. Mueller, T.D. Reus, J. Reinhardt, B. Mueller, W. Greiner, G. Soff, Phys. Rev. A 37 (1988) 1449.
- [2] I.B. Khriplovich, JETP Letters 100 (2014) 552.
- [3] I.A. Maltsev, V.M. Shabaev, I.I. Tupitsin, A.I. Bondarev, Y.S. Kozhedub, G. Plunien, Th. Stolker, Phys. Rev. A 91 (2015) 032708.