## Once more on parastatistics

V.N. Tolstoy

Lomonosov Moscow State University, Skobeltsyn Institute of Nuclear Physics, Moscow 119991, Russian Federation

#### Abstract

Equivalence between algebraic structures generated by parastatistics triple relations of Green (1953) and Greenberg – Messiah (1965), and certain orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie superalgebras is found explicitly. Moreover, it is shown that such superalgebras give more complex para-Fermi and para-Bose systems then ones of Green – Greenberg – Messiah.

## 1 Introduction

The usual creation and annihilation operators of identical particles, fermions  $a_i^{\pm}$  (i = 1, ..., m) and bosons  $b_i^{\pm}$  (j = 1, ..., n), satisfy the canonical commutation relations:

$$\{a_i^{\zeta}, a_j^{\eta}\} = \frac{1}{2} |\eta - \zeta| \delta_{ij}, \qquad [b_i^{\zeta}, b_j^{\eta}] = \frac{1}{2} (\eta - \zeta) \delta_{ij}. \tag{1.1}$$

Here and elsewhere the Greek letters  $\zeta, \eta \in \{+, -\}$  if they are upper indexes, and they are interpreted as +1 and -1 in the algebraic expressions of the type  $\eta - \zeta$ .

From the relations (1.1) it follows the so-called "symmetrization postulate" (SP): States of more than one identical particle must be antisymmetric (fermions) or symmetric (bosons) under permutations.

In 1953 Green [1] proposed to refuse SP and he introduced algebras with the triple relations:

$$[[a_i^{\zeta}, a_j^{\eta}], a_k^{\xi}] = |\xi - \eta| \delta_{jk} a_i^{\zeta} - |\xi - \zeta| \delta_{ik} a_j^{\eta} \quad \text{(parafermions)}, \tag{1.2}$$

$$[\{b_i^{\zeta}, b_j^{\eta}\}, b_k^{\xi}] = (\xi - \eta)\delta_{jk}b_i^{\zeta} + (\xi - \zeta)\delta_{ik}b_j^{\eta} \quad \text{(parabosons)}.$$
(1.3)

The usual fermions and bosons satisfy these relations but also another solutions exist.

In 1962 Kamefuchi and Takahashi [2] (also see [3]) shown that the parafermionic algebra is isomorphic to the orthogonal Lie algebra  $\mathfrak{o}(2m+1) := \mathfrak{o}(2m+1, \mathbb{C})$ . Later in 1980 Ganchev and Palev [4] proved that the parabosonic algebra is isomorphic to the orthosymplectic  $\mathbb{Z}_2$ graded Lie superalgebra  $\mathfrak{osp}(1|2n)$ . In 1965 Greenberg and Messiah [5] considered parasystem consisting simultaneously from parafermions and parabosons and they defined the relative commutation rules between parafermions and parabosons. There are two types of such relations:

$$\begin{split} & [[a_i^{\zeta}, a_j^{\eta}], b_k^{\xi}] = 0, \\ & [[a_i^{\zeta}, b_j^{\eta}], a_k^{\xi}] = -|\xi - \zeta| \delta_{ik} b_j^{\eta}, \\ & \{[a_i^{\zeta}, b_j^{\eta}], b_k^{\xi}\} = (\xi - \eta) \delta_{jk} a_i^{\zeta}, \end{split}$$
(1.4)

$$[[a_i^{\zeta}, a_j^{\eta}], b_k^{\xi}] = 0, \qquad [\{b_i^{\zeta}, b_j^{\eta}\}, a_k^{\xi}] = 0, \\ \{\{a_i^{\zeta}, b_j^{\eta}\}, a_k^{\xi}\} = |\xi - \zeta| \delta_{ik} b_j^{\eta}, \qquad [\{a_i^{\zeta}, b_j^{\eta}\}, b_k^{\xi}] = (\xi - \eta) \delta_{jk} a_i^{\zeta},$$

$$(1.5)$$

where i, j, k = 1, 2, ..., m for the symbols *a*'s and i, j, k = 1, 2, ..., n for the symbols *b*'s. The first case (1.4) was called as *the relative para-Fermi set* and the second case (1.5) was called as *the relative para-Boson set*<sup>1</sup>.

In 1982 Palev [6] shown that the case (1.4) with (1.2) and (1.3) is isomorphic to the orthosymplectic  $\mathbb{Z}_2$ -graded Lie superalgebra  $\mathfrak{osp}(2m+1|2n)$ . No any similar solution for the second case (1.5) was known up to now.

Here we show that the case (1.5) is isomorphic to the orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebra  $\mathfrak{osp}(1, 2m|2n, 0)$ . Moreover it will demonstrate that the more general mixed parasystem, which simultaneously involves the relative para-Fermi and relative para-Bose sets, contains two sorts of parafermions and one sort of parabosons and it is isomorphic to the orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebra  $\mathfrak{osp}(2m_1+1, 2m_2|2n, 0)$ . All previous cases are particular (degenerated) variants of this general case.

The paper is organized as follows. Section 2 provides a definition and general structure of  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebras and also a matrix realization and a Cartan-Weyl basis of the general linear  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebra  $\mathfrak{gl}(m_1, m_2|n_1, n_2)$ . In Section 3 we describe the orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebra  $\mathfrak{osp}(2m_1 + 1, 2m_2|2n, 0)$  and show that a part of its defining triple relations in the terms of short root vectors coincides with the the relative para-Bose set.

# 2 Superalgebra $\mathfrak{gl}(m_1, m_2 | n_1, n_2)$

At first we remind a general definition of the  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebra [7], [8].

The  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded LSA  $\mathfrak{g}$ , as a linear space, is a direct sum of four graded components

$$\mathfrak{g} = \bigoplus_{\mathbf{a}=(a_1,a_2)} \mathfrak{g}_{\mathbf{a}} = \mathfrak{g}_{(0,0)} \oplus \mathfrak{g}_{(1,1)} \oplus \mathfrak{g}_{(1,0)} \oplus \mathfrak{g}_{(0,1)}$$
(2.1)

with a bilinear operation  $[\cdot, \cdot]$  satisfying the identities (grading, symmetry, Jacobi):

$$\deg(\llbracket x_{\mathbf{a}}, y_{\mathbf{b}} \rrbracket) = \deg(x_{\mathbf{a}}) + \deg(x_{\mathbf{b}}) = \mathbf{a} + \mathbf{b} = (a_1 + b_1, a_2 + b_2),$$
(2.2)

$$\llbracket x_{\mathbf{a}}, y_{\mathbf{b}} \rrbracket = -(-1)^{\mathbf{ab}} \llbracket y_{\mathbf{b}}, x_{\mathbf{a}} \rrbracket,$$
(2.3)

$$[\![x_{\mathbf{a}}, [\![y_{\mathbf{b}}, z]\!]]] = [\![[\![x_{\mathbf{a}}, y_{\mathbf{b}}]\!], z]\!] + (-1)^{\mathbf{ab}} [\![y_{\mathbf{b}}, [\![x_{\mathbf{a}}, z]\!]],$$
(2.4)

where the vector  $(a_1 + b_1, a_2 + b_2)$  is defined mod (2, 2) and  $\mathbf{ab} = a_1b_1 + a_2b_2$ . Here in (2.2)-(2.4)  $x_{\mathbf{a}} \in \mathfrak{g}_{\mathbf{a}}, x_{\mathbf{b}} \in \mathfrak{g}_{\mathbf{b}}$ , and the element  $z \in \mathfrak{g}$  is not necessarily homogeneous. From (2.2)

<sup>&</sup>lt;sup>1</sup>The names the relative para-Fermi and para-Boson set are directly related to type of the Lie bracket (commutator or anticommutator) given between parafermion and paraboson elements.

it is follows that  $\mathfrak{g}_{(0,0)}$  is a Lie subalgebra in  $\mathfrak{g}$ , and the subspaces  $\mathfrak{g}_{(1,1)}$ ,  $\mathfrak{g}_{(1,0)}$  and  $\mathfrak{g}_{(0,1)}$  are  $\mathfrak{g}_{(0,0)}$ -modules. It should be noted that  $\mathfrak{g}_{(0,0)} \oplus \mathfrak{g}_{(1,1)}$  is a Lie subalgebra in  $\mathfrak{g}$  and the subspace  $\mathfrak{g}_{(1,0)} \oplus \mathfrak{g}_{(0,1)}$  is a  $\mathfrak{g}_{(0,0)} \oplus \mathfrak{g}_{(1,1)}$ -module, and moreover  $\{\mathfrak{g}_{(1,1)}, \mathfrak{g}_{(1,0)}\} \subset \mathfrak{g}_{(0,1)}$  and vice versa  $\{\mathfrak{g}_{(1,1)}, \mathfrak{g}_{(0,1)}\} \subset \mathfrak{g}_{(1,0)}$ . From (2.2) and (2.3) it is follows that the general Lie bracket  $[\![\cdot, \cdot]\!]$  for homogeneous elements posses two value: commutator  $[\cdot, \cdot]$  and anticommutator  $\{\cdot, \cdot\}$  as well as in a case of usual  $\mathbb{Z}_2$ -graded Lie superalgebras [9].

Now we construct a  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded matrix superalgebras  $\mathfrak{gl}(m_1, m_2 | n_1, n_2)$ .

Let an arbitrary  $(m_1 + m_2 + n_1 + n_2) \times (m_1 + m_2 + n_1 + n_2)$ -matrix M be presented in the following block form<sup>2</sup>:

$$M = \begin{pmatrix} A_{(0,0)} & A_{(1,1)} & A_{(1,0)} & A_{(0,1)} \\ B_{(1,1)} & B_{(0,0)} & B_{(0,1)} & B_{(1,0)} \\ C_{(1,0)} & C_{(0,1)} & C_{(0,0)} & C_{(1,1)} \\ D_{(0,1)} & D_{(1,0)} & D_{(1,1)} & D_{(0,0)} \end{pmatrix},$$
(2.5)

where the diagonal block matrices  $A_{(0,0)}, B_{(0,0)}, C_{(0,0)}, D_{(0,0)}$  have the dimensions  $m_1 \times m_1$ ,  $m_2 \times m_2, n_1 \times n_1$  and  $n_2 \times n_2$  correspondingly, the dimensions of the non-diagonal block matrices  $A_{(1,1)}, A_{(1,0)}, A_{(0,1)}$ , etc. are easy determined by the dimensions of these diagonal block matrices. The matrix M can be split into the sum of four matrices:

$$M = M_{(0,0)} + M_{(1,1)} + M_{(1,0)} + M_{(0,1)} =$$

$$= \begin{pmatrix} A_{(0,0)} & 0 & 0 & 0 \\ 0 & B_{(0,0)} & 0 & 0 \\ 0 & 0 & C_{(0,0)} & 0 \\ 0 & 0 & 0 & D_{(0,0)} \end{pmatrix} + \begin{pmatrix} 0 & A_{(1,1)} & 0 & 0 \\ B_{(1,1)} & 0 & 0 & 0 \\ 0 & 0 & 0 & C_{(1,1)} \\ 0 & 0 & D_{(1,1)} & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 & A_{(0,1)} \\ 0 & 0 & B_{(0,1)} & 0 \\ 0 & 0 & D_{(1,0)} & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 & A_{(0,1)} \\ 0 & 0 & B_{(0,1)} & 0 \\ 0 & C_{(0,1)} & 0 & 0 \end{pmatrix}.$$

$$(2.6)$$

Let us define the general commutator  $[\cdot, \cdot]$  on a space of all such matrices by the following way:

$$\llbracket M_{(a_1,a_2)}, M'_{(b_1,b_2)} \rrbracket := M_{(a_1,a_2)} M'_{(b_1,b_2)} - (-1)^{a_1 b_1 + a_2 b_2} M'_{(b_1,b_2)} M_{(a_1,a_2)}.$$
(2.7)

for the homogeneous components  $M_{(a_1,a_2)}$  and  $M_{(b,b_2)}$ . For arbitrary matrices M and M' the commutator  $[\![\cdot, \cdot]\!]$  is extended by linearity. It is easy to check that

$$\llbracket M_{(a_1,a_2)}, M'_{(b_1,b_2)} \rrbracket = M''_{(a_1+a_2,b_1+b_2)},$$
(2.8)

where the sum  $(a_1 + a_2, b_1 + b_2)$  is defined mod (2, 2). Thus the grading condition (2.2) is available. The symmetry and Jacobi identities (2.3) and (2.4) are available too. Hence we obtain a Lie superalgebra which is called  $\mathfrak{gl}(m_1, m_2|n_1, n_2)$ . It should be noted that

$$\begin{bmatrix} M_{\mathbf{a}}, M_{\mathbf{b}}' \end{bmatrix} = \begin{bmatrix} M_{\mathbf{a}}, M_{\mathbf{b}}' \end{bmatrix} \quad \text{if} \quad \mathbf{ab} = 0, 2,$$
  
$$\begin{bmatrix} M_{\mathbf{a}}, M_{\mathbf{b}}' \end{bmatrix} = \{ M_{\mathbf{a}}, M_{\mathbf{b}}' \} \quad \text{if} \quad \mathbf{ab} = 1.$$
(2.9)

 $<sup>^{2}</sup>$ It is evidently supposed that all such matrices in each block-row or in each block-column have the same number of rows or columns

Now we consider the Cartan-Weyl basis of  $\mathfrak{gl}(m_1, m_2 | n_1, n_2)$  and its supercommutation  $(\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded) relations. In accordance with the block structure of the  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded matrix (2.5) we introduce a  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded function (grading)  $\mathbf{d}(\cdot)$  defined on the integer segment  $[1, 2, \ldots, m_1, m_1 + 1, \ldots, m_1 + m_2, m_1 + m_2 + 1, \ldots, m_1 + m_2 + n_1, m_1 + m_2 + n_1 + 1, \ldots, m_1 + m_2 + n_1 + n_2]$  as follows:

$$\mathbf{d}_{i} := \mathbf{d}(i) = \begin{cases} (0,0) & \text{for } i = 1, 2, \dots, m_{1}, \\ (1,1) & \text{for } i = m_{1} + 1, \dots, m_{1} + m_{2}, \\ (1,0) & \text{for } i = m_{1} + m_{2} + 1, \dots, m_{1} + m_{2} + n_{1}, \\ (0,1) & \text{for } i = m_{1} + m_{2} + n_{1} + 1, \dots, m_{1} + m_{2} + n_{1} + n_{2}. \end{cases}$$
(2.10)

Let  $e_{ij}$  be the  $(m_1 + m_2 + n_1 + n_2) \times (m_1 + m_2 + n_1 + n_2)$  matrix (2.5) with 1 is in the (i, j)-th place and other entries 0. The matrices  $e_{ij}$   $(i, j = 1, 2, ..., m_1 + m_2 + n_1 + n_2)$  are homogeneous, moreover, the grading deg $(e_{ij})$  is determined by

$$\deg(e_{ij}) = \mathbf{d}_{ij} := \mathbf{d}_i + \mathbf{d}_j \pmod{(2,2)}, \tag{2.11}$$

and the supercommutator for such matrices is given as follows

$$[\![e_{ij}, e_{kl}]\!] := e_{ij}e_{kl} - (-1)^{\mathbf{d}_{ij}\mathbf{d}_{kl}}e_{kl}e_{ij}.$$
(2.12)

It is easy to check that

$$[\![e_{ij}, e_{kl}]\!] = \delta_{jk} e_{il} - (-1)^{\mathbf{d}_{ij} \mathbf{d}_{kl}} \delta_{il} e_{kj}.$$
(2.13)

The elements  $e_{ij}$   $(i, j = 1, 2, ..., m_1 + m_2 + n_1 + n_2)$  with the relations (2.13) generates the Lie superalgebra  $\mathfrak{gl}(m_1, m_2|n_1, n_2)$ . The elements  $h_i := e_{ii}$   $(i, j = 1, 2, ..., m_1 + m_2 + n_1 + n_2)$  compose a basis in the Cartan subalgebra  $\mathfrak{h}(m_1 + m_2|n_1 + n_2) \subset \mathfrak{gl}(m_1, m_2|n_1, n_2)$ .

The Lie superalgebra  $\mathfrak{gl}(m_1, m_2 | n_1, n_2)$  play a special role among all finite dimensional  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie superalgebras. Namely, a general Ado's theorem is valid. It states: Any finite dimensional Lie  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebra can be realized in terms of a subalgebra of  $\mathfrak{gl}(m_1, m_2 | n_1, n_2)$ . This theorem was proved by Scheunert [10] for all finite dimensional graded generalized Lie algebras including our cases.

As an illustration of the Ado's theorem, in the next section we give realization of the orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded superalgebra  $\mathfrak{osp}(2m_1 + 1, 2m_2|2n, 0)$  in terms of the superalgebra  $\mathfrak{gl}(2m_1 + 1, 2m_2|2n, 0)$  and, moreover, we present a Cartan-Weyl basis of the orthosymplectic superalgebra and its explicit commutation relations and we also show that a subset of the short root vectors of the Cartan-Weyl basis generates this superalgebra and describe the parastatistics with the relative para-Fermi and para-Bose sets simultaneously.

# **3** Orthosymplectic superalgebra $\mathfrak{osp}(2m_1 + 1, 2m_2|2n, 0)$ and its relation with parastatistics

We start with an explicit description of embedding of the orthosymplectic Lie superalgebra  $\mathfrak{osp}(2m_1+1, 2m_2|2n, 0)$  in the general linear Lie superalgebra  $\mathfrak{gl}(2m_1+1, 2m_2|2n, 0)$ . For this propose the  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded integer segment  $\mathbb{S}_N^{(\mathbf{d})} := [1, 2, \ldots, 2N+1]$ , where  $N = m_1+m_2+n$ ,

with the grading  $d(\cdot)$  given by

$$\mathbf{d}_{i} := \mathbf{d}(i) = \begin{cases} (0,0) & \text{for } i = 1, 2, \dots, 2m_{1}, \\ (1,1) & \text{for } i = 2m_{1} + 1, \dots, 2m_{1} + 2m_{2}, \\ (1,0) & \text{for } i = 2m_{1} + 2m_{2} + 1, \dots, 2m_{1} + 2m_{2} + 2n. \end{cases}$$
(3.1)

is reindexed by the following way  $\tilde{\mathbb{S}}_N^{(\mathbf{d})} := [0, \pm 1, \pm 2, \dots, \pm N]$  with the grading  $\mathbf{d}(\cdot)$  given by

$$\mathbf{d}_{i} := \mathbf{d}(i) = \begin{cases} (0,0) & \text{for } i = 0, \pm 1, \pm 2, \dots, \pm m_{1}, \\ (1,1) & \text{for } i = \pm (m_{1}+1), \dots, \pm (m_{1}+m_{2}), \\ (1,0) & \text{for } i = \pm (m_{1}+m_{2}+1), \dots, \pm (m_{1}+m_{2}+n), \end{cases}$$
(3.2)

Rows and columns of  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded  $(2N+1) \times (2N+1)$ -matrices are enumerated by the indices  $0, 1, -1, 2, -2, \ldots, N, -N$   $(N = m_1 + m_2 + n)$ . Let  $e_{ij}(i, j \in \tilde{\mathbb{S}}_N^{(\mathbf{d})})$  be the standard (unit) basis of  $\mathfrak{gl}(2m_1 + 1, 2m_2|2n, 0)$  with the given indexing and the canonical supercommutation relations:

$$\llbracket e_{ij}, e_{kl} \rrbracket = \delta_{jk} e_{il} - (-1)^{\mathbf{d}_{ij} \mathbf{d}_{kl}} \delta_{il} e_{kj}, \qquad (3.3)$$

where  $\mathbf{d}_{ij} = \mathbf{d}_i + \mathbf{d}_j$  and the grading  $\mathbf{d}(\cdot)$  is given by (3.2).

The orthosymplectic ( $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded) Lie superalgebra  $\mathfrak{osp}(2m_1 + 1, 2m_2|2n, 0)$  is embedded in  $\mathfrak{gl}(2m_1 + 1, 2m_2|2n, 0)$  as a linear span of the elements

$$x_{ij} := e_{i,-j} - (-1)^{\mathbf{d}_i \mathbf{d}_j + \mathbf{d}_{ij}^2} \phi_i \phi_j e_{j,-i} \qquad (i, j \in \tilde{\mathbb{S}}_N^{(\mathbf{d})}),$$
(3.4)

where the index function  $\phi_i$  is given as follows

$$\phi_i := \begin{cases} 1 & \text{if } i = 0, \pm 1, \pm 2, \dots, \pm (m_1 + m_2), \\ 1 & \text{if } i = m_1 + m_2 + 1, \dots, m_1 + m_2 + n, \\ -1 & \text{if } i = -m_1 - m_2 - 1, \dots, -m_1 - m_2 - n. \end{cases}$$
(3.5)

It is easy to verify that the the elements (3.4) satisfy the following supercommutation relations

$$\begin{bmatrix} x_{ij}, x_{kl} \end{bmatrix} = \delta_{j,-k} x_{il} - \delta_{j,-l} (-1)^{\mathbf{d}_k \mathbf{d}_l + \mathbf{d}_{kl}^2} \phi_k \phi_l x_{ik} - \delta_{i,-k} (-1)^{\mathbf{d}_i \mathbf{d}_j + \mathbf{d}_{ij}^2} \phi_i \phi_j x_{jl} - \delta_{i,-l} (-1)^{\mathbf{d}_{ij} \mathbf{d}_{ik}} x_{kj}.$$

$$(3.6)$$

Not all elements (3.4) are linearly independent because they satisfy the relations

$$x_{ij} = -(-1)^{\mathbf{d}_i \mathbf{d}_j + \mathbf{d}_{ij}^2} \phi_i \phi_j x_{ji} \qquad (i, j \in \tilde{\mathbb{S}}_N^{(\mathbf{d})}),$$
(3.7)

and what is more

 $x_{ii} = 0$  for  $i = 0, \pm 1, \pm 2, \dots, \pm (m_1 + m_2).$  (3.8)

From the general supercommutation relations (3.6) it follows at once that the short root vectors  $x_{0i}$   $(i = \pm 1, \pm 2, \ldots, \pm (m_1 + m_2 + n))$  satisfy the following triple relations:

$$\llbracket \llbracket x_{0i}, x_{0j} \rrbracket, x_{0k} \rrbracket = -\delta_{j,-k} \phi_j x_{0i} + \delta_{i,-k} (-1)^{\mathbf{d}_i \mathbf{d}_j} \phi_i x_{0j}.$$
(3.9)

Conversely, let the abstract  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded generators  $x_{0i}$   $(i = \pm 1, \pm 2, \ldots, \pm (m_1 + m_2 + n))$  with the grading deg $(x_{0i}) = \mathbf{d}_{0i} \equiv \mathbf{d}_0 + \mathbf{d}_i = \mathbf{d}_i$ , where  $\mathbf{d}_i$  is given by (3.2), satisfy the relations (3.9), where the index function  $\phi_i$  is determined by (3.5), then it is not difficult to check that these relations generate for the superalgebra  $\mathfrak{osp}(2m_1 + 1, 2m_2|2n, 0)$ .

The defining relations (3.9) can be rewritten in detailed in accordance with the explicit grading of their generators using the following notations<sup>3</sup>:

$$\begin{aligned}
a_i^{-\zeta} &:= \zeta \sqrt{2} \, x_{0,\zeta i} & (\deg(a_i^{\zeta}) = (0,0)) & \text{for } i = 1, 2, \dots, m_1, \\
\tilde{a}_i^{-\zeta} &:= \zeta \sqrt{2} \, x_{0,\zeta(m_1+i)} & (\deg(\tilde{a}_i^{\zeta}) = (1,1)) & \text{for } i = 1, 2, \dots, m_2, \\
b_i^{-\zeta} &:= \zeta \sqrt{2} \, x_{0,\zeta(m+i)} & (\deg(b_i^{\zeta}) = (1,0)) & \text{for } i = 1, 2, \dots, n,
\end{aligned}$$
(3.10)

where  $m := m_1 + m_2$ . Substituting (3.10) in (3.9) we obtain the different types of defining triple relations.

#### 1. Parafermion relations:

(a1) the defining relations of  $\mathfrak{o}(2m_1+1)$ :

$$[[a_i^{\zeta}, a_j^{\eta}], a_k^{\xi}] = |\xi - \eta| \delta_{jk} a_i^{\zeta} - |\xi - \zeta| \delta_{ik} a_j^{\eta} \quad \text{for } i, j, k = 1, 2, \dots, m_1; \quad (3.11)$$

(a2) the defining relations of  $\mathfrak{o}(2m_2+1)$ :

$$[[\tilde{a}_{i}^{\zeta}, \tilde{a}_{j}^{\eta}], \tilde{a}_{k}^{\xi}] = |\xi - \eta| \delta_{jk} \tilde{a}_{i}^{\zeta} - |\xi - \zeta| \delta_{ik} \tilde{a}_{j}^{\eta} \quad \text{for } i, j, k = 1, 2, \dots, m_{2}; \quad (3.12)$$

(a3) the mixed parafermion relations :

$$[[a_i^{\zeta}, a_j^{\eta}], \tilde{a}_k^{\xi}] = 0, \qquad [[\tilde{a}_i^{\zeta}, \tilde{a}_j^{\eta}], a_k^{\xi}] = 0, \qquad (3.13)$$

$$[[a_i^{\zeta}, \tilde{a}_j^{\eta}], a_k^{\xi}] = -|\xi - \zeta| \delta_{ik} \tilde{a}_j^{\eta}, \qquad [[a_i^{\zeta}, \tilde{a}_j^{\eta}], \tilde{a}_k^{\xi}] = |\xi - \eta| \delta_{jk} a_i^{\zeta}, \qquad (3.14)$$

where  $i, j, k = 1, ..., m_1$  for the symbols a's and  $i, j, k = 1, ..., m_2$  for the symbols b's.

#### 2. Paraboson relations:

(b1) the defining relations of  $\mathfrak{osp}(1|2n)$ :

$$[\{b_i^{\zeta}, b_j^{\eta}\}, b_k^{\xi}] = (\xi - \eta)\delta_{jk}b_i^{\zeta} + (\xi - \zeta)\delta_{ik}b_j^{\eta} \quad \text{for } i, j, k = 1, 2, \dots, n.$$
(3.15)

2. Mixed parafermion and paraboson relations:

(ab1) the relative para-Fermi set:

$$[[a_i^{\zeta}, a_j^{\eta}], b_k^{\xi}] = 0, \qquad [\{b_i^{\zeta}, b_j^{\eta}\}, a_k^{\xi}] = 0, [[a_i^{\zeta}, b_j^{\eta}], a_k^{\xi}] = -|\xi - \zeta| \delta_{ik} b_j^{\eta}, \qquad \{[a_i^{\zeta}, b_j^{\eta}], b_k^{\xi}\} = (\xi - \eta) \delta_{jk} a_i^{\zeta},$$

$$(3.16)$$

where  $i, j, k = 1, 2, ..., m_1$  for the symbols a's and i, j, k = 1, ..., n for the symbols b's;

<sup>&</sup>lt;sup>3</sup>Here anywhere  $\zeta, \eta, \xi \in \{+, -\}$ .

(ab2) the relative para-Bose set:

$$\begin{bmatrix} [\tilde{a}_{i}^{\zeta}, \tilde{a}_{j}^{\eta}], b_{k}^{\xi}] = 0, \qquad [\{b_{i}^{\zeta}, b_{j}^{\eta}\}, \tilde{a}_{k}^{\xi}] = 0, \\ \{\{\tilde{a}_{i}^{\zeta}, b_{j}^{\eta}\}, \tilde{a}_{k}^{\xi}\} = |\xi - \zeta| \delta_{ik} b_{j}^{\eta}, \qquad [\{\tilde{a}_{i}^{\zeta}, b_{j}^{\eta}\}, b_{k}^{\xi}] = (\xi - \eta) \delta_{jk} \tilde{a}_{i}^{\zeta},$$

$$(3.17)$$

where  $i, j, k = 1, 2, ..., m_2$  for the symbols  $\tilde{a}$ 's and i, j, k = 1, ..., n for the symbols b's;

(ab3) the relations with distinct grading elements:

$$\{[a_i^{\zeta}, \tilde{a}_j^{\eta}], b_k^{\xi}\} = [\{\tilde{a}_j^{\eta}, b_k^{\xi}\}, a_i^{\zeta}] = \{[b_k^{\xi}, a_i^{\zeta}], \tilde{a}_j^{\eta}\} = 0.$$
(3.18)

The result connected with the relation (3.9) can be reformulated the following way. If we have two sorts of the parafermions  $a_i^{\zeta}$   $(i = 1, 2, ..., m_1)$  and  $\tilde{a}_i^{\zeta}$   $(i = 1, 2, ..., m_2)$  with the triple relations (3.11)–(3.14) and one sort of the parabosons  $b_i^{\zeta}$  (i = 1, 2, ..., n) with the triple relations (3.15) which together satisfy the relative para-Fermi set (3.16) and relative para-Bose set (3.17), and they obey also the triple relations of the form (3.18) then this parasystem generate the orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie superalgebra  $\mathfrak{osp}(2m_1 + 1, 2m_2|2n, 0)$ .

We consider two particular cases which are degenerations of  $\mathfrak{osp}(2m_1+1, 2m_2|2n, 0)$ .

- If the parasystem consist of only one sort of the parafermions  $a_i^{\zeta}$   $(i = 1, 2, ..., m_1)$ and one sort of the parabosons  $b_i^{\zeta}$  (i = 1, 2, ..., n) then we have the parasystem with the relative Fermi set and it generates the orthosymplectic  $\mathbb{Z}_2$ -graded Lie superalgebra  $\mathfrak{osp}(2m_1 + 1|2n) = \mathfrak{osp}(2m_1 + 1, 0|2n, 0).$
- If the parasystem contains one sort of the parafermions  $\tilde{a}_i^{\zeta}$   $(i = 1, 2, ..., m_2)$  and one sort of the parabosons  $b_i^{\zeta}$  (i = 1, 2, ..., n) then we have the case of a parasystem with the relative Bose set (see the relations (1.2), (1.3) and (1.5)) and it generates the orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie superalgebra  $\mathfrak{osp}(1, 2m_2|2n, 0)$ .

Thus we shown that the para-Fermi and para-Boose triple relations (1.2), (1.3) together with the relative para-Bose set (1.5) generate the orthosymplectic  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded Lie superalgebra  $\mathfrak{osp}(1, 2m|2n, 0)$ . Moreover, it was shown that the superalgebras  $\mathfrak{osp}(m_1, 2m_2|2n, 0)$  give more complex para-Fermi and para-Bose system which contain the relative para-Fermi and para-Bose sets simultaneously.

It should be noted that, probably, for the first time the  $\mathbb{Z}_2 \times \mathbb{Z}_2$ -graded structure of the relative para-Bose set (1.5) was observable in [11], [12] (also see [13]).

It should be also noted that the obtained relation between the parastatistics and the orthosymplectic superalgebras allows to apply all mathematical power of the representation theory of the superalgebras for a detailed description of the parastatistics, e.g. their Fock spaces etc. (for example, see [14] - [16]).

### Acknowledgments

The author would like to thank the Organizers for the kind invitation to speak at the International Workshop Supersymmetries and Quantum Symmetries, Dubna, July 29 - August 3, 2013. The paper was supported by the RFBR grant No.11-01-00980-a and the grant No.12-09-0064 of the Academic Fund Program of the National Research University Higher School of Economics.

### References

- Green H.S. A Generalized Method of Field Quantization // Phys. Rev. 1953. V. 9. P. 270-273.
- [2] Kamefuchi S., Takahashi Y. A generalization of field quantization and statistics // Nucl. Phys. 1962. V. 36. P. 177-206.
- [3] Ryan C., Sudarshan E.C.G. Representations of parafermi rings // Nucl. Phys. 1963. V. 47. P. 207-211.
- [4] Ganchev A.C., Palev T.D. A Lie superalgebraic interpretation of the para-Bose statistics // J. Math. Phys. 1980. V. 21. P. 797-799.
- [5] Greenberg O.W., Messiah A.M.L. Selection rules for parafields and the absence of paraparticles in nature // Phys. Rev. B. 1965. V. 138. P. 1155-1167.
- [6] Palev T.D. Para-Bose and para-Fermi operators as generators of orthosymplectic Lie superalgebras // J. Math. Phys. 1982. V. 23. P. 1100-1102.
- [7] Rittenberg V., Wyler D. Sequences of Z<sub>2</sub> × Z<sub>2</sub>-graded Lie algebras and superalgebras // J. Math. Phys. 1978. V. 19, No. 10. P. 2193-2200.
- [8] Rittenberg V., Wyler D. Generalized superalgebras // Nucl. Phys. B. 1978. V. 139, No. 10. P. 189-202.
- [9] Kac V.G. Lie superalgebras // Adv. Math. 1977. V. 26. P. 8-96.
- [10] Scheunert M. Generalized Lie algebras // J. Math. Phys. 1979. V. 20, No. 4. P. 712-720.
- [11] Yang W., Jing S. A new kind of graded Lie algebra and parastatistical supersymmetry // Science in China, Series A. 2001. V. 44, No. 9. P.1167-1173; arXiv:math-ph/0212004v1.
- [12] Yang W., Jing S. Graded Lie algebra generating of parastatistical algebraic structure // Commun. Theor. Phys. 2001. V. 36. P. 647-65; arXiv:math-ph/0212009v1.
- [13] Kanakoglou K., Daskaloyannis C., Herrera-Aguilar A. Mixed paraparticles, colors, braidings and a new class of realizations for Lie superalgebras // arXiv:0912.1070v1 [math-ph]. 2009. P. 1-19.
- [14] Lievens S. Stoilova N.I. Van der Jeugt J. The paraboson Fock space and unitary irreducible representations of the Lie superalgebra osp(1|2n) // Commun. Math. Phys. 2008. V. 281. P. 805-826; arXiv:0706.4196v2 [hep-th].
- [15] Stoilova N.I., Van der Jeugt J. The parafermion Fock space and explicit \$0(2n+1) representations // J. Phys. A. 2008. V. 41, 075202. P. 1-13; arXiv:0712.1485v1 [hep-th].
- [16] Stoilova N.I. The parastatistics Fock space and explicit Lie superalgebra representations // arXiv:1311.4042v1 [math-ph]. 2013. P. 1-14.