# **RULED AND QUADRIC SURFACES IN THE 3-DIMENSIONAL** EUCLIDEAN SPACE SATISFYING $\Delta^{III} x = \Lambda x$

#### HASSAN AL-ZOUBI AND STYLIANOS STAMATAKIS

ABSTRACT. We consider ruled and quadric surfaces in the 3-dimensional Euclidean space which are of coordinate finite type with respect to the third fundamental form III, i.e., their position vector  $\mathbf{x}$  satisfies the relation  $\Delta^{III}\mathbf{x} = \Lambda \mathbf{x}$  where  $\Lambda$  is a square matrix of order 3. We show that helicoids and spheres are the only surfaces in  $E^3$  satisfying the preceding relation.

## **1. INTRODUCTION**

Let S be a (connected) surface in a Euclidean 3-space  $E^3$  referred to any system of coordinates  $u^1$ ,  $u^2$ , which does not contain parabolic points, we denote by  $b_{ij}$  the components of the second fundamental form  $II = b_{ij} du^i du^j$  of S. Let  $\varphi(u^1, u^2)$  and  $\psi(u^1, u^2)$  be two sufficient differentiable functions on S. Then the first differential parameter of Beltrami with respect to the second fundamental form of S is defined by

$$abla^{II}(arphi, \psi) := b^{ij} arphi_{Ii} \psi_{Ii}$$

where  $\varphi_{i} := \frac{\partial \varphi}{\partial u^{i}}$  and  $(b^{ij})$  denotes the inverse tensor of  $(b_{ij})$ . Let  $e_{ij}$  be the components of the third fundamental form *III* of *S*. Then the second differential parameter of Beltrami with respect to the third fundamental form of S is defined by [9]

$$\Delta^{III}\varphi := -\frac{1}{\sqrt{e}}(\sqrt{e}e^{ij}\varphi_{Ii})_{Ij}^{1},$$

where  $(e^{ij})$  denotes the inverse tensor of  $(e_{ij})$  and  $e := \det(e_{ij})$ .

In [9], S. Stamatakis and H. Al-Zoubi showed, for the position vector  $\mathbf{x} = \mathbf{x}(u^1, u^2)$  of S, the relation

$$\Delta^{III} \boldsymbol{x} = \nabla^{III} (\frac{2H}{K}, \boldsymbol{n}) - \frac{2H}{K} \boldsymbol{n}, \qquad (1.1)$$

where *n* is the Gauss map, *K* the Gauss curvature and *H* the mean curvature of *S*. Moreover, in this context, the same authors proved that the surfaces  $S: \mathbf{x} = \mathbf{x}(u^1, u^2)$  satisfying the condition

$$\Delta^{III} \boldsymbol{x} = \lambda \boldsymbol{x}, \quad \lambda \in \mathbb{R}$$

i.e., for which all coordinate functions are eigenfunctions of  $\Delta^{III}$  with the same eigenvalue  $\lambda$ , are precisely either the minimal surfaces ( $\lambda = 0$ ), or the part of spheres ( $\lambda = 2$ ).

In [2] B.-Y. Chen introduced the notion of Euclidean immersions of finite type. In terms of B.-Y. Chen theory, a surface S is said to be of finite type, if its coordinate functions are a finite sum of eigenfunctions of the Beltrami operator  $\Delta^{III}$ . Therefore the two facts mentioned above can be stated as follows

<sup>2010</sup> Mathematics Subject Classification. 53A05, 47A75.

Key words and phrases. Surfaces in Euclidean space, Surfaces of coordinate finite type, Beltrami operator. <sup>1</sup> with sign convention such that  $\Delta = -\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2}$  for the metric  $ds^2 = dx^2 + dy^2$ .

- *S* is minimal if and only if *S* is of null type1.
- *S* lies in an ordinary sphere  $S^2$  if and only if *S* is of type1.

Following [2] we say that a surface *S* is of finite type with respect to the fundamental form *III*, or briefly of finite *III*-type, if the position vector  $\mathbf{x}$  of *S* can be written as a finite sum of nonconstant eigenvectors of the operator  $\Delta^{III}$ , i.e., if

$$\boldsymbol{x} = \boldsymbol{x}_0 + \sum_{i=1}^m \boldsymbol{x}_i, \quad \Delta^{III} \boldsymbol{x}_i = \lambda_i \boldsymbol{x}_i, \quad i = 1, ..., m,$$
(1.2)

where  $\mathbf{x}_0$  is a fixed vector and  $\lambda_1$ ,  $\lambda_2$ , ...,  $\lambda_m$  are eigenvalues of  $\Delta^{III}$ ; when there are exactly k nonconstant eigenvectors  $\mathbf{x}_1$ ,  $\mathbf{x}_2$ , ...,  $\mathbf{x}_k$  appearing in (1.2) which all belong to different eigenvalues  $\lambda_1$ ,  $\lambda_2$ , ...,  $\lambda_k$ , then S is said to be of *III*-type k, otherwise S is said to be of infinite type. When  $\lambda_i = 0$  for some i = 1, 2, ..., k, then S is said to be of null *III*-type k.

Up to now, very little is known about surfaces of finite *III*-type. Concerning this problem, the only known surfaces of finite *III*-type in  $E^3$  are parts of spheres, the minimal surfaces and the parallel of the minimal surfaces which are of null *III*-type 2, (see [9]).

In this paper we shall be concerned with the ruled and quadric surfaces in  $E^3$  which are connected, complete and which are of coordinate finite *III*-type, i.e., their position vector  $\mathbf{x} = \mathbf{x}(u^1, u^2)$  satisfies the relation

$$\Delta^{III} \boldsymbol{x} = \boldsymbol{\Lambda} \boldsymbol{x},\tag{1.3}$$

where  $\Lambda$  is a square matrix of order 3.

In [6] F. Dillen, J. Pas and L. Verstraelen studied coordinate finite type with respect to the first fundamental form  $I = g_{ij} du^i du^j$  and they proved that

**Theorem 1.** The only surfaces in  $\mathbb{R}^3$  satisfying

 $\Delta^{I} \boldsymbol{x} = A \boldsymbol{x} + B, \quad A \in M(3,3), \quad B \in M(3,1)$ 

are the minimal surfaces, the spheres and the circular cylinders.

(M(m,n) denotes the set of all matrices of the type (m,n)). While in [7] O. Garay showed that

**Theorem 2.** The only complete surfaces of revolutions in  $\mathbb{R}^3$ , whose component functions are eigenfunctions of their Laplacian, then the surface must be a catenoid, a sphere or a right circular cylinder.

Recently, H. Al-Zoubi and S. Stamatakis studied coordinate finite type with respect to the third fundamental form, more precisely, in [10] they proved the following

**Theorem 3.** A surface of revolution S in  $\mathbb{R}^3$ , satisfies (1.3), if and only if S is a catenoid or a part of a sphere.

## 2. MAIN RESULTS

Our main results are the following

**Proposition 1.** *The only ruled surfaces in the 3-dimensional Euclidean space that satisfies (1.3), are the helicoids.* 

**Proposition 2.** The only quadric surfaces in the 3-dimensional Euclidean space that satisfies (1.3), are the spheres.

Our discussion is local, which means that we show in fact that any open part of a ruled or a quadric satisfies (1.3), if it is an open part of a helicoid or an open part of a sphere respectively.

Before starting the proof of our main results we first show that the surfaces mentioned in the above propositions indeed satisfy the condition (1.3). On a helicoid the mean curvature vanishes, so, by virtue of (1.1),  $\Delta^{III} \mathbf{x} = \mathbf{0}$ . Therefore a helicoid satisfies (1.3), where  $\Lambda$  is the null matrix in M(3,3).

Let  $S^2(r)$  be a sphere of radius r centered at the origin. If x denotes the position vector field of  $S^2(r)$ , then the Gauss map n is given by  $-\frac{x}{r}$ . For the Gauss curvature K and the mean curvature H of  $S^2(r)$  we have  $K = \frac{1}{r^2}$  and  $H = \frac{1}{r}$ , so, by virtue of (1.1), we obtain

$$\Delta^{III} \boldsymbol{x} = 2\boldsymbol{x}$$

and we find that  $S^2(r)$  satisfies (1.3) with

$$\Lambda = \left[ \begin{array}{rrr} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{array} \right].$$

## 3. PROOF OF PROPOSITION 1

Let *S* be a ruled surface in  $E^3$ . We suppose that *S* is a non-cylindrical ruled surface. This surface can be expressed in terms of a directrix curve  $\alpha(s)$  and a unit vectorfield  $\beta(s)$  pointing along the rulings as

$$S: \mathbf{x}(s, t) = \mathbf{\alpha}(s) + t \mathbf{\beta}(s), \quad s \in J, \quad -\infty < t < \infty.$$

Moreover, we can take the parameter *s* to be the arc length along the spherical curve  $\beta(s)$ . Then we have

$$\langle \boldsymbol{\alpha}', \boldsymbol{\beta} \rangle = 0, \quad \langle \boldsymbol{\beta}, \boldsymbol{\beta} \rangle = 1, \quad \langle \boldsymbol{\beta}', \boldsymbol{\beta}' \rangle = 1,$$
(3.1)

where the prime denotes the derivative in s. The first fundamental form of S is

$$I = q ds^2 + dt^2,$$

while the second fundamental form is

$$II = \frac{p}{\sqrt{q}}ds^2 + \frac{2A}{\sqrt{q}}dsdt,$$

where

$$q := \langle \boldsymbol{\alpha}', \boldsymbol{\alpha}' \rangle + 2 \langle \boldsymbol{\alpha}', \boldsymbol{\beta}' \rangle t + t^{2},$$
  

$$p := (\boldsymbol{\alpha}', \boldsymbol{\beta}, \boldsymbol{\alpha}'') + [(\boldsymbol{\alpha}', \boldsymbol{\beta}, \boldsymbol{\beta}'') + (\boldsymbol{\beta}', \boldsymbol{\beta}, \boldsymbol{\alpha}'')] t + (\boldsymbol{\beta}', \boldsymbol{\beta}, \boldsymbol{\beta}'') t^{2},$$
  

$$A := (\boldsymbol{\alpha}', \boldsymbol{\beta}, \boldsymbol{\beta}').$$

For convenience, we put

$$\begin{aligned} \kappa &:= \langle \boldsymbol{\alpha}', \boldsymbol{\alpha}' \rangle, \qquad \lambda := \langle \boldsymbol{\alpha}', \boldsymbol{\beta}' \rangle, \\ \mu &:= \left( \boldsymbol{\beta}', \boldsymbol{\beta}, \boldsymbol{\beta}'' \right), \qquad \nu := \left( \boldsymbol{\alpha}', \boldsymbol{\beta}, \boldsymbol{\beta}'' \right) + \left( \boldsymbol{\beta}', \boldsymbol{\beta}, \boldsymbol{\alpha}'' \right), \\ \rho &:= \left( \boldsymbol{\alpha}', \boldsymbol{\beta}, \boldsymbol{\alpha}'' \right), \end{aligned}$$

and thus we have

$$q = t^2 + 2\lambda t + \kappa, \qquad p = \mu t^2 + \nu t + \rho.$$

For the Gauss curvature *K* of *S* we find

$$K = -\frac{A^2}{q^2}.$$
(3.2)

The Beltrami operator with respect to the third fundamental form, after a lengthy computation, can be expressed as follows

$$\Delta^{III} = -\frac{q}{A^2} \frac{\partial^2}{\partial s^2} + \frac{2qp}{A^3} \frac{\partial^2}{\partial s \partial t} - \left(\frac{q^2}{A^2} + \frac{qp^2}{A^4}\right) \frac{\partial^2}{\partial t^2} + \left(\frac{q_s}{2A^2} + \frac{qp_t}{A^3} - \frac{pq_t}{2A^3}\right) \frac{\partial}{\partial s} + \left(\frac{qp_s}{A^3} - \frac{pq_s}{2A^3} - \frac{pqA'}{A^4} - \frac{qq_t}{2A^2} + \frac{p^2q_t}{2A^4} - \frac{2qpp_t}{A^4}\right) \frac{\partial}{\partial t} = Q_1 \frac{\partial^2}{\partial s^2} + Q_2 \frac{\partial^2}{\partial s \partial t} + Q_3 \frac{\partial}{\partial s} + Q_4 \frac{\partial}{\partial t} + Q_5 \frac{\partial^2}{\partial t^2}$$
(3.3)

where

$$q_t := \frac{\partial q}{\partial t}, \quad q_s := \frac{\partial q}{\partial s}, \quad p_t := \frac{\partial p}{\partial t}, \quad p_s := \frac{\partial p}{\partial s}$$

and  $Q_1, Q_2, ..., Q_5$  are polynomials in *t* with functions in *s* as coefficients, and  $deg(Q_i) \le 6$ . More precisely we have

$$\begin{aligned} Q_{1} &= -\frac{1}{A^{2}} [t^{2} + 2\lambda t + \kappa], \\ Q_{2} &= \frac{2}{A^{3}} [\mu t^{4} + (2\lambda\mu + \nu) t^{3} + (2\lambda\nu + \rho + \kappa\mu) t^{2} \\ &+ (2\lambda\rho + \kappa\nu) t + \kappa\rho], \end{aligned}$$

$$\begin{aligned} Q_{3} &= \frac{1}{A^{3}} [\mu t^{3} + 3\lambda\mu t^{2} + (\lambda\nu - \rho + 2\kappa\mu + \lambda'A) t \\ &+ \frac{1}{2}\kappa'A - \lambda\rho + \kappa\nu], \end{aligned}$$

$$\begin{aligned} Q_{4} &= \frac{1}{A^{4}} [-3\mu^{2}t^{5} + (\mu'A - \mu A' - 4\mu\nu - 7\lambda\mu^{2}) t^{4} \\ &+ (\nu'A - \nu A' + 2\lambda\mu'A - 2\lambda\mu A' - \lambda'\mu A - A^{2} \\ &- 10\lambda\mu\nu - 2\mu\rho - \nu^{2} - 4\kappa\mu^{2}) t^{3} \\ &+ (\kappa\mu'A - \kappa\mu A' - \frac{1}{2}\kappa'\mu A + 2\lambda\nu'A - 2\lambda\nu A' - \lambda'\nu A \\ &- \rho A' + \rho'A - 3\lambda A^{2} - 3\lambda\nu^{2} - 6\lambda\mu\rho - 6\kappa\mu\nu) t^{2} \\ &+ (\kappa\nu'A - \kappa\nu A' - \frac{1}{2}\kappa'\nu A + 2\lambda\rho'A - 2\lambda\rho A' - \lambda'\rho A \\ &- \kappa A^{2} - 2\lambda^{2}A^{2} - 2\kappa\nu^{2} + \rho^{2} - 2\lambda\nu\rho - 4\kappa\mu\rho) t \\ &+ (\kappa\rho'A - \kappa\rho A' - \frac{1}{2}\kappa'\rho A + \lambda\rho^{2} - \kappa\lambda A^{2} - 2\kappa\nu\rho)], \end{aligned}$$

$$Q_{5} = -\frac{1}{A^{4}} [\mu^{2} t^{6} + (2\mu\nu + 2\lambda\mu^{2}) t^{5} + (2\mu\rho + \nu^{2} + 4\lambda\mu\nu + \kappa\mu^{2} + A^{2}) t^{4} + (2\nu\rho + 4\lambda\mu\rho + 2\lambda\nu^{2} + 2\kappa\mu\nu + 4\lambda A^{2}) t^{3} + (\rho^{2} + 4\lambda\nu\rho + 2\kappa\mu\rho + \kappa\nu^{2} + 4\lambda^{2}A^{2} + 2\kappa A^{2}) t^{2} + (2\lambda\rho^{2} + 2\kappa\nu\rho + 4\lambda\kappa A^{2}) t + (\kappa\rho^{2} + \kappa^{2}A^{2})].$$

Applying (3.3) for the position vector  $\boldsymbol{x}$  one finds

$$\Delta^{III} \boldsymbol{x} = Q_1 \boldsymbol{\alpha}^{\prime\prime} + Q_2 \boldsymbol{\beta}^{\prime} + Q_3 \boldsymbol{\alpha}^{\prime} + Q_4 \boldsymbol{\beta} + (Q_1 \boldsymbol{\beta}^{\prime\prime} + Q_3 \boldsymbol{\beta}^{\prime}) t.$$
(3.4)

Let  $\mathbf{x} = (x_1, x_2, x_3)$ ,  $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \alpha_3)$  and  $\boldsymbol{\beta} = (\beta_1, \beta_2, \beta_3)$  be the coordinate functions of  $\mathbf{x}, \boldsymbol{\alpha}$  and  $\boldsymbol{\beta}$ . By virtue of (3.4) we obtain

$$\Delta^{III} x_i = Q_1 \alpha''_i + Q_2 \beta'_i + Q_3 \alpha'_i + Q_4 \beta_i + (Q_1 \beta''_i + Q_3 \beta'_i) t, \qquad (3.5)$$
  

$$i = 1, 2, 3.$$

We denote by  $\lambda_{ij}$ , i, j = 1, 2, 3 the entries of the matrix  $\Lambda$ . Using (3.5) and condition (1.3) we have

$$Q_{1}\alpha_{i}'' + Q_{2}\beta_{i}' + Q_{3}\alpha_{i}' + Q_{4}\beta_{i} + (Q_{1}\beta_{i}'' + Q_{3}\beta_{i}') t$$
  
=  $\lambda_{i1}\alpha_{1} + \lambda_{i2}\alpha_{2} + \lambda_{i3}\alpha_{3} + (\lambda_{i1}\beta_{1} + \lambda_{i2}\beta_{2} + \lambda_{i3}\beta_{3}) t$ ,  
= 1,2,3.

Consequently

i

$$-3\mu^{2}\beta_{i}t^{5} + [(\mu'A - \mu A' - 4\mu v - 7\lambda\mu^{2})\beta_{i} + 3\mu A\beta'_{i}]t^{4} + [\mu A\alpha'_{i} - A^{2}\beta''_{i} + (2vA + 7\lambda\mu A)\beta'_{i} + (v'A - vA' + 2\lambda\mu'A - 2\lambda\mu A' - \lambda'\mu A - A^{2} - 10\lambda\mu v - 2\mu\rho - v^{2} - 4\kappa\mu^{2})\beta_{i}]t^{3} + [(\kappa\mu'A - \kappa\mu A' - \frac{1}{2}\kappa'\mu A + 2\lambda v'A - 2\lambda vA' - \lambda'vA - \rho A' + \rho'A - 3\lambda A^{2} - 3\lambda v^{2} - 6\lambda\mu\rho - 6\kappa\mu v)\beta_{i} + 3\lambda\mu A\alpha'_{i} - 2\lambda A^{2}\beta''_{i} - A^{2}\alpha''_{i} + (\lambda'A + 5\lambda v + 4\kappa\mu + \rho)A\beta'_{i}]t^{2} + [(\frac{1}{2}\kappa'A + 3\kappa v + 3\lambda\rho)A\beta'_{i} + (\kappa v'A - \kappa vA' - \frac{1}{2}\kappa'vA + 2\lambda\rho'A - 2\lambda\rho A' - \lambda'\rhoA - \kappa A^{2} - 2\lambda^{2}A^{2} - 2\kappa v^{2} + \rho^{2} - 2\lambda v\rho - 4\kappa\mu\rho)\beta_{i} - 2\lambda A^{2}\alpha''_{i} - \kappa A^{2}\beta''_{i} + (\lambda v - \rho + 2\kappa\mu + \lambda'A)A\alpha'_{i}]t - A^{4}(\lambda_{i1}\beta_{1} + \lambda_{i2}\beta_{2} + \lambda_{i3}\beta_{3})t + (\kappa\rho'A - \kappa\rho A' - \frac{1}{2}\kappa'\rho A + \lambda\rho^{2} - \kappa\lambda A^{2} - 2\kappa v\rho)\beta_{i} - \kappa A^{2}\alpha''_{i} + (\frac{1}{2}\kappa'A - \lambda\rho + \kappa v)A\alpha'_{i} - A^{4}(\lambda_{i1}\alpha_{1} + \lambda_{i2}\alpha_{2} + \lambda_{i3}\alpha_{3}) = 0.$$
 (3.6)

For i = 1, 2, 3, (3.6) is a polynomial in t with functions in s as coefficients. This implies that the coefficients of the powers of t in (3.6) must be zeros, so we obtain, for i = 1, 2, 3, the following equations

$$3\mu^2\beta_i = 0, \tag{3.7}$$

$$(\mu' A - \mu A' - 4\mu \nu - 7\lambda \mu^2) \beta_i + 3\mu A \beta'_i = 0, \mu A \alpha'_i - A^2 \beta''_i + (2\nu A + 7\lambda \mu A) \beta'_i + (\nu' A - \nu A' + 2\lambda \mu' A - 2\lambda \mu A' - \lambda' \mu A - A^2 - 10\lambda \mu \nu - 2\mu \rho - \nu^2 - 4\kappa \mu^2) \beta_i = 0,$$
 (3.8)

$$(\kappa\mu'A - \kappa\mu A' - \frac{1}{2}\kappa'\mu A + 2\lambda\nu'A - 2\lambda\nu A' - \lambda'\nu A - \rho A' + \rho'A - 3\lambda A^2 - 3\lambda\nu^2 - 6\lambda\mu\rho - 6\kappa\mu\nu)\beta_i + 3\lambda\mu A\alpha'_i - 2\lambda A^2\beta''_i - A^2\alpha''_i + (\lambda'A + 5\lambda\nu + 4\kappa\mu + \rho)A\beta'_i = 0,$$
(3.9)

$$(\kappa\nu'A - \kappa\nu A' - \frac{1}{2}\kappa'\nu A + 2\lambda\rho'A - 2\lambda\rho A' - \lambda'\rho A - \kappa A^{2} -2\lambda^{2}A^{2} - 2\kappa\nu^{2} + \rho^{2} - 2\lambda\nu\rho - 4\kappa\mu\rho)\beta_{i} - 2\lambda A^{2}\alpha_{i}'' - \kappa A^{2}\beta_{i}'' + (\frac{1}{2}\kappa'A + 2\kappa\nu + 4\lambda\rho)A\beta_{i}' + (\lambda\nu - \rho + 2\kappa\mu + \lambda'A)A\alpha_{i}' = A^{4}(\lambda_{i1}\beta_{1} + \lambda_{i2}\beta_{2} + \lambda_{i3}\beta_{3}), \qquad (3.10)$$

$$(\kappa \rho' A - \kappa \rho A' - \frac{1}{2} \kappa' \rho A + \lambda \rho^{2} - \kappa \lambda A^{2} - 2\kappa \nu \rho) \beta_{i} + 2\kappa \rho A \beta'_{i}$$
$$+ (\frac{1}{2} \kappa' A - \lambda \rho + \kappa \nu) A \alpha'_{i} - \kappa A^{2} \alpha''_{i}$$
$$= A^{4} (\lambda_{i1} \alpha_{1} + \lambda_{i2} \alpha_{2} + \lambda_{i3} \alpha_{3}).$$
(3.11)

From (3.7) one finds

$$\mu = \left(\boldsymbol{\beta}', \boldsymbol{\beta}, \boldsymbol{\beta}''\right) = 0, \qquad (3.12)$$

which implies that the vectors  $\beta'$ ,  $\beta$ ,  $\beta''$  are linearly dependent, and hence there exist two functions  $\sigma_1 = \sigma_1(s)$  and  $\sigma_2 = \sigma_2(s)$  such that

$$\boldsymbol{\beta}^{\prime\prime} = \sigma_1 \, \boldsymbol{\beta} + \sigma_2 \, \boldsymbol{\beta}^{\prime}. \tag{3.13}$$

On differentiating  $\langle \boldsymbol{\beta}', \boldsymbol{\beta}' \rangle = 1$ , we obtain  $\langle \boldsymbol{\beta}', \boldsymbol{\beta}'' \rangle = 0$ . So from (3.13) we have

$$\boldsymbol{\beta}^{\prime\prime} = \sigma_1 \boldsymbol{\beta}. \tag{3.14}$$

By taking the derivative of  $\langle \boldsymbol{\beta}, \boldsymbol{\beta} \rangle = 1$  twice, we find that

$$\langle \boldsymbol{\beta}', \boldsymbol{\beta}' \rangle + \langle \boldsymbol{\beta}, \boldsymbol{\beta}'' \rangle = 0.$$

But  $\langle \boldsymbol{\beta}', \boldsymbol{\beta}' \rangle = 1$ , and taking into account (3.14) we find that  $\sigma_1(s) = -1$ . Thus (3.14) becomes  $\boldsymbol{\beta}'' = -\boldsymbol{\beta}$  which implies that

$$\beta_i'' = -\beta_i, \quad i = 1, 2, 3. \tag{3.15}$$

Using (3.12) and (3.15) equation (3.8) reduces to

$$2\nu A\beta'_i + (\nu'A - \nu A' - \nu^2)\beta_i = 0, \quad i = 1, 2, 3$$

or, in vector notation

$$2\nu A\boldsymbol{\beta}' + (\nu'A - \nu A' - \nu^2)\boldsymbol{\beta} = \mathbf{0}.$$
(3.16)

By taking the derivative of  $\langle \boldsymbol{\beta}, \boldsymbol{\beta} \rangle = 1$ , we find that the vectors  $\boldsymbol{\beta}, \boldsymbol{\beta}'$  are linearly independent, and so from (3.16) we obtain that vA = 0. We note that  $A \neq 0$ , since from (3.2) the Gauss curvature vanishes, so we are left with v = 0. Then equation (3.9) becomes

$$-A^{2}\alpha_{i}^{\prime\prime} + (\lambda'A + \rho)A\beta_{i}^{\prime} + (\rho'A - \rho A' - \lambda A^{2})\beta_{i} = 0, \quad i = 1, 2, 3$$

or, in vector notation

$$-A^{2}\boldsymbol{\alpha}'' + (\lambda'A + \rho)A\boldsymbol{\beta}' + (\rho'A - \rho A' - \lambda A^{2})\boldsymbol{\beta} = \mathbf{0}.$$
(3.17)

Taking the inner product of both sides of the above equation with  $\beta'$  we find in view of (3.1) that

$$-A^{2}\langle \boldsymbol{\alpha}^{\prime\prime}, \boldsymbol{\beta}^{\prime} \rangle + \rho A + \lambda^{\prime} A^{2} = 0.$$
(3.18)

On differentiating  $\lambda = \langle \boldsymbol{\alpha}', \boldsymbol{\beta}' \rangle$  with respect to *s*, by virtue of (3.15) and (3.1), we get

$$\lambda' = \langle \boldsymbol{\alpha}'', \boldsymbol{\beta}' \rangle + \langle \boldsymbol{\alpha}', \boldsymbol{\beta}'' \rangle = \langle \boldsymbol{\alpha}'', \boldsymbol{\beta}' \rangle - \langle \boldsymbol{\alpha}', \boldsymbol{\beta} \rangle = \langle \boldsymbol{\alpha}'', \boldsymbol{\beta}' \rangle.$$
(3.19)

Hence, (3.18) reduces to  $\rho A = 0$ , which implies that  $\rho = 0$ . Thus the vectors  $\boldsymbol{\alpha}', \boldsymbol{\beta}, \boldsymbol{\alpha}''$  are linearly dependent, and so there exist two functions  $\sigma_3 = \sigma_3(s)$  and  $\sigma_4 = \sigma_4(s)$  such that

$$\boldsymbol{\alpha}^{\prime\prime} = \sigma_3 \,\boldsymbol{\beta} + \sigma_4 \,\boldsymbol{\alpha}^\prime. \tag{3.20}$$

Taking the inner product of both sides of the last equation with  $\boldsymbol{\beta}$  we find in view of (3.1) that  $\sigma_3 = \langle \boldsymbol{\alpha}'', \boldsymbol{\beta} \rangle$ .

Now, by taking the derivative of  $\langle \boldsymbol{\alpha}', \boldsymbol{\beta} \rangle = 0$ , we find  $\langle \boldsymbol{\alpha}'', \boldsymbol{\beta} \rangle + \langle \boldsymbol{\alpha}', \boldsymbol{\beta}' \rangle = 0$ , that is

$$\langle \boldsymbol{\alpha}^{\prime\prime}, \boldsymbol{\beta} \rangle + \lambda = 0, \tag{3.21}$$

and hence  $\sigma_3 = -\lambda$ .

Taking again the inner product of both sides of equation (3.20) with  $\beta'$  we find in view of (3.1) that

$$\langle \boldsymbol{\alpha}^{\prime\prime}, \boldsymbol{\beta}^{\prime} \rangle = \sigma_4 \lambda. \tag{3.22}$$

Using (3.19) we find  $\lambda' = \sigma_4 \lambda$ . Thus  $\sigma_4 = \frac{\lambda'}{\lambda}$ . Therefore

$$\boldsymbol{\alpha}^{\prime\prime} = -\lambda \,\boldsymbol{\beta} + \frac{\lambda^{\prime}}{\lambda} \boldsymbol{\alpha}^{\prime}. \tag{3.23}$$

We distinguish two cases

**Case 1**:  $\lambda = 0$ . Because of  $\rho = 0$  equation (3.17) would yield A = 0, which is clearly impossible for the surfaces under consideration.

**Case 2**:  $\lambda \neq 0$ . From (3.17), (3.23) and  $\rho = 0$  we find that

$$-\frac{\lambda'}{\lambda}A^2\boldsymbol{\alpha}' + \lambda'A^2\boldsymbol{\beta}' = \mathbf{0}$$

which implies that  $\lambda'(\alpha' - \lambda \beta') = 0$ .

If  $\lambda' \neq 0$ , then  $\alpha' = \lambda \beta'$ . Hence  $\alpha', \beta'$  are linearly dependent, and so A = 0 which contradicts our previous assumption. Thus  $\lambda' = 0$ . From (3.23) we have

$$\boldsymbol{\alpha}^{\prime\prime} = -\lambda \boldsymbol{\beta}. \tag{3.24}$$

On the other hand, by taking the derivative of  $\kappa$  and using the last equation we obtain that  $\kappa$  is constant. Hence equations (3.10) and (3.11) reduce to

$$\lambda_{i1}\beta_1 + \lambda_{i2}\beta_2 + \lambda_{i3}\beta_3 = 0,$$
  
$$\lambda_{i1}\alpha_1 + \lambda_{i2}\alpha_2 + \lambda_{i3}\alpha_3 = 0, i = 1, 2, 3$$

and so  $\lambda_{ij} = 0, i, j = 1, 2, 3$ .

Since the parameter *s* is the arc length of the spherical curve  $\beta(s)$ , and because of (3.12) we suppose, without loss of generality, that the parametrization of  $\beta(s)$  is

$$\boldsymbol{\beta}(s) = (\cos s, \sin s, 0).$$

Integrating (3.24) twice we get

$$\alpha(s) = (c_1 s + c_2 + \lambda \cos s, c_3 s + c_4 + \lambda \sin s, c_5 s + c_6)$$

where  $c_i$ , i = 1, 2, ..., 6 are constants.

Since  $\kappa = \langle \boldsymbol{\alpha}', \boldsymbol{\alpha}' \rangle$  is constant, it's easy to show that  $c_1 = c_3 = 0$ . Hence  $\boldsymbol{\alpha}(s)$  reduces to

$$\boldsymbol{\alpha}(s) = (c_2 + \lambda \cos s, c_4 + \lambda \sin s, c_5 s + c_6).$$

Thus we have

 $S: \mathbf{x}(s, t) = (c_2 + (\lambda + t)\cos s, c_4 + (\lambda + t)\sin s, c_5 s + c_6)$ 

which is a helicoid.

#### 4. PROOF OF PROPOSITION 2

Let now *S* be a quadric surface in the Euclidean 3-space  $E^3$ . Then *S* is either ruled, or of one of the following two kinds

$$z^2 - ax^2 - by^2 = c, \quad abc \neq 0$$
 (4.1)

or

$$z = \frac{a}{2}x^2 + \frac{b}{2}y^2, \quad a > 0, \ b > 0.$$
(4.2)

If *S* is ruled and satisfies (1.3), then by Proposition 1 *S* is a helicoid. We first show that a quadric of the kind (4.1) satisfies (1.3) if and only if a = -1 and b = -1, which means that *S* is a sphere. Next we show that a quadric of the kind (4.2) is never satisfying (1.3).

# 4.1. Quadrics of the first kind. This kind of quadric surfaces can be parametrized as follows

$$\boldsymbol{x}(u,v) = \left(u, v, \sqrt{c + au^2 + bv^2}\right).$$

Let's denote the function  $c + au^2 + bv^2$  by  $\omega$  and the function  $c + a(a+1)u^2 + b(b+1)v^2$  by *T*. Then the components  $g_{ij}$ ,  $b_{ij}$  and  $e_{ij}$  of the first, second and third fundamental tensors in (local) coordinates are the following

$$g_{11} = 1 + \frac{(au)^2}{\omega}, \quad g_{12} = \frac{abuv}{\omega}, \quad g_{22} = 1 + \frac{(bv)^2}{\omega},$$
$$b_{11} = \frac{a(c+bv^2)}{\omega\sqrt{T}}, \quad b_{12} = -\frac{abuv}{\omega\sqrt{T}}, \quad b_{22} = \frac{b(c+au^2)}{\omega\sqrt{T}}$$

and

$$e_{11} = \frac{a^2}{\omega T^2} [(buv)^2 + (bv^2 + c)^2 + b^2 v^2 \omega],$$
  

$$e_{12} = \frac{ab}{\omega T^2} [c(a+b)uv + abuv(u^2 + v^2 + \omega)],$$
  

$$e_{22} = \frac{b^2}{\omega T^2} [(auv)^2 + (au^2 + c)^2 + a^2 u^2 \omega].$$

Notice that  $\omega$  and T are polynomials in u and v. If for simplicity we put

 $C(u, v) = (buv)^{2} + (bv^{2} + c)^{2} + b^{2}v^{2}\omega,$   $B(u, v) = uv [c(a+b) + ab(u^{2} + v^{2} + \omega)],$  $A(u, v) = (auv)^{2} + (au^{2} + c)^{2} + a^{2}u^{2}\omega,$ 

then the third fundamental tensors  $e_{ij}$  turns into

$$e_{11} = \frac{a^2}{\omega T^2} C(u, v), \quad e_{12} = -\frac{ab}{\omega T^2} B(u, v), \quad e_{22} = \frac{b^2}{\omega T^2} A(u, v).$$

Hence the Beltrami operator  $\Delta^{III}$  of *S* can be expressed as follows

$$\Delta^{III} = -\frac{T}{(abc)^2} \left[ b^2 A \frac{\partial^2}{\partial u^2} + 2abB \frac{\partial^2}{\partial u \partial v} + a^2 C \frac{\partial^2}{\partial v^2} \right] -\frac{T}{(abc)^2} \left[ b \left( b \frac{\partial A}{\partial u} + a \frac{\partial B}{\partial v} \right) \frac{\partial}{\partial u} + a \left( a \frac{\partial C}{\partial v} + b \frac{\partial B}{\partial u} \right) \frac{\partial}{\partial v} \right] +\frac{T}{(abc)^2} \left[ \frac{ab^2}{\omega} (uA + vB) \frac{\partial}{\partial u} + \frac{a^2 b}{\omega} (uB + vC) \frac{\partial}{\partial v} \right] +\frac{1}{(abc)^2} [ab^2 ((a + 1) uA + (b + 1) vB) \frac{\partial}{\partial u} +a^2 b ((b + 1) vC + (a + 1) uB) \frac{\partial}{\partial v}].$$
(4.3)

We remark that

$$b\frac{\partial A}{\partial u} + a\frac{\partial B}{\partial v} = au \left[ 5ab(a+1)u^2 + 5ab(b+1)v^2 + c(3ab+5b+a) \right],$$
  

$$a\frac{\partial C}{\partial v} + b\frac{\partial B}{\partial u} = av \left[ 5ab(a+1)u^2 + 5ab(b+1)v^2 + c(3ab+5a+b) \right],$$
  

$$uA + vB = \left[ c + a(a+1)u^2 + a(b+1)v^2 \right] u\omega,$$
  

$$uB + vC = \left[ c + b(a+1)u^2 + b(b+1)v^2 \right] v\omega,$$
  

$$(a+1)uA + (b+1)vB = u \left[ c(a+1) + a(a+1)u^2 + a(b+1)v^2 \right] T,$$

 $(b+1) vC + (a+1) uB = v [c(b+1) + b(a+1)u^2 + b(b+1)v^2] T.$ We denote by  $\lambda_{ij}$ , i, j = 1, 2, 3 the entries of the matrix  $\Lambda$ . On account of (1.3) we get

$$\Delta^{III} x_1 = \Delta^{III} u = \lambda_{11} u + \lambda_{12} v + \lambda_{13} \sqrt{\omega}, \tag{4.4}$$

$$\Delta^{III} x_2 = \Delta^{III} v = \lambda_{21} u + \lambda_{22} v + \lambda_{23} \sqrt{\omega}, \qquad (4.5)$$

$$\Delta^{III} x_3 = \Delta^{III} \sqrt{\omega} = \lambda_{31} u + \lambda_{32} v + \lambda_{33} \sqrt{\omega}.$$

Applying (4.3) on the coordinate functions  $x_i$ , i = 1, 2 of the position vector  $\mathbf{x}$  and by virtue of (4.4) and (4.5), we find respectively

$$\Delta^{III} u = -\frac{uT}{c^2} \left[ 3(a+1)u^2 + 3(b+1)v^2 + \frac{c(3b+a+2ab)}{ab} \right]$$
  
=  $\lambda_{11}u + \lambda_{12}v + \lambda_{13}\sqrt{\omega},$  (4.6)

$$\Delta^{III} v = -\frac{vT}{c^2} \left[ 3(a+1)u^2 + 3(b+1)v^2 + \frac{c(b+3a+2ab)}{ab} \right]$$
  
=  $\lambda_{21}u + \lambda_{22}v + \lambda_{23}\sqrt{\omega},$  (4.7)

Putting v = 0 in (4.6), we obtain that

$$-\frac{3a(a+1)^2}{c^2}u^5 - \frac{(a+1)(6b+a+2ab)}{bc}u^3 - \frac{(3b+a+2ab)}{ab}u$$
$$= \lambda_{11}u + \lambda_{13}\sqrt{c+au^2}.$$

Since  $a \neq 0$  and  $c \neq 0$  this implies that a = -1. Similarly, if we put u = 0 in (4.7) we obtain that

$$-\frac{3b(b+1)^2}{c^2}v^5 - \frac{(b+1)(b+6a+2ab)}{ac}v^3 - \frac{(b+3a+2ab)}{ab}v$$
  
=  $\lambda_{22}v + \lambda_{23}\sqrt{c+bv^2}$ .

This implies that b = -1. Hence *S* must be a sphere.

4.2. **Quadrics of the second kind.** For this kind of surfaces we can consider a parametrization

$$\mathbf{x}(u,v) = \left(u,v,\frac{a}{2}u^2 + \frac{b}{2}v^2\right).$$

Then the components  $g_{ij}$ ,  $b_{ij}$  and  $e_{ij}$  of the first, second and third fundamental tensors are the following

$$g_{11} = 1 + (au)^2$$
,  $g_{12} = abuv$ ,  $g_{22} = 1 + (bv)^2$ ,

$$b_{11} = \frac{a}{\sqrt{g}}, \quad b_{12} = 0, \quad b_{22} = \frac{b}{\sqrt{g}},$$
$$e_{11} = \frac{a^2}{g^2}(1 + b^2v^2), \quad e_{12} = -\frac{a^2b^2}{g^2}uv, \quad e_{22} = \frac{b^2}{g^2}(1 + a^2u^2),$$

where  $g := \det(g_{ij}) = 1 + (au)^2 + (bv)^2$ .

A straightforward computation shows that the Beltrami operator  $\Delta^{III}$  of *S* takes the following form

$$\Delta^{III} = -\frac{g(1+a^2u^2)}{a^2}\frac{\partial^2}{\partial u^2} - \frac{g(1+b^2v^2)}{b^2}\frac{\partial^2}{\partial v^2} -2uvg\frac{\partial^2}{\partial u\partial v} - 2ug\frac{\partial}{\partial u} - 2vg\frac{\partial}{\partial v}.$$
(4.8)

On account of (1.3) we get

$$\Delta^{III} x_1 = \Delta^{III} u = \lambda_{11} u + \lambda_{12} v + \lambda_{13} \left( \frac{a}{2} u^2 + \frac{b}{2} v^2 \right), \tag{4.9}$$

$$\Delta^{III} x_2 = \Delta^{III} v = \lambda_{21} u + \lambda_{22} v + \lambda_{23} \left( \frac{a}{2} u^2 + \frac{b}{2} v^2 \right),$$

$$\Delta^{III} x_3 = \Delta^{III} \sqrt{\omega} = \lambda_{31} u + \lambda_{32} v + \lambda_{33} \left( \frac{a}{2} u^2 + \frac{b}{2} v^2 \right).$$
(4.10)

Applying (4.8) on the coordinate functions  $x_i$ , i = 1, 2 of the position vector  $\mathbf{x}$  and by virtue of (4.9) and (4.10) we find respectively

$$\Delta^{III} u = -2ug = \lambda_{11}u + \lambda_{12}v + \lambda_{13} \left(\frac{a}{2}u^2 + \frac{b}{2}v^2\right), \tag{4.11}$$

$$\Delta^{III} v = -2vg = \lambda_{21}u + \lambda_{22}v + \lambda_{23}\left(\frac{a}{2}u^2 + \frac{b}{2}v^2\right).$$
(4.12)

Putting v = 0 in (4.11), we obtain that

$$-2a^2u^3 - 2u = \lambda_{11}u + \lambda_{13}\frac{a}{2}u^2$$

This implies that *a* must be zero. Putting u = 0 in (4.12), we obtain that

$$-2b^2v^3 - 2v = \lambda_{22}v + \lambda_{23}\frac{b}{2}v^2$$

This implies that *b* must be zero, which is clearly impossible, since a > 0 and b > 0.

## References

- [1] C. BAIKOUSIS, D. BLAIR: On the Gauss map of ruled surfaces. Glasg. Math. J. 34, 355–359 (1992).
- [2] B.-Y. CHEN: Total mean curvature and submanifolds of finite type. World Scientific Publisher, 1984.
- [3] B.-Y. CHEN: Some open problems and conjectures on submanifolds of finite type. Soochow J. Math. 17, 169–188 (1991).
- [4] B.-Y. CHEN, F. DILLEN, L. VERSTRAELEN, L. VRANCKEN: *Ruled surfaces of finite type. Bull. Austral. Math. Soc. J.* **42**, 447–453 (1990).
- [5] B.-Y. CHEN, F. DILLEN: Quadrics of finite type. J. Geom. 38, 16–22 (1990).
- [6] F. DILLEN, J. PAS, L. VERSTRAELEN: On surfaces of finite type in Euclidean 3-space. Kodai Math. J. 13, 10–21 (1990).
- [7] O. GARAY: On a certain class of finite type of surfaces of revolution. Kodai Math. J. 11, 25–31 (1988).
- [8] S. STAMATAKIS: Der 2. Beltramishe operator der dritten Grundform einer Fläche des E<sup>3</sup>. Proceedings of the 4<sup>th</sup> international Congress of Geometry, Thessaloniki, 392–396 (1996).
- [9] S. STAMATAKIS, H. AL-ZOUBI: On surfaces of finite Chen-type. Results Math. 43, 181–190 (2003).
- [10] S. STAMATAKIS, H. AL-ZOUBI: Surfaces of revolution satisfying  $\Delta^{III} \mathbf{x} = A\mathbf{x}$ . J. Geom. Graph. 14, 181–186 (2010).
- [11] T. TAKAHASHI: Minimal immersions of Riemannian manifolds. J. Math. Soc. Japan 18, 380–385 (1966).

DEPARTMENT OF BASIC SCIENCES, AL-ZAYTOONAH UNIVERSITY OF JORDAN, AMMAN *E-mail address*: dr.hassanz@zuj.edu.jo

DEPARTMENT OF MATHEMATICS, ARISTOTLE UNIVERSITY OF THESSALONIKI *E-mail address*: stamata@math.auth.gr