

# On the values of Dedekind sums

Kurt Girstmair

## Abstract

Let  $s(a, b)$  denote the classical Dedekind sum and  $S(a, b) = 12s(a, b)$ . For a given denominator  $q \in \mathbb{N}$ , we study the numerators  $k \in \mathbb{Z}$  of the values  $k/q$ ,  $(k, q) = 1$ , of Dedekind sums  $S(a, b)$ . Our main result says that if  $k$  is such a numerator, then the whole residue class of  $k$  modulo  $(q^2 - 1)q$  consists of numerators of this kind. This fact reduces the task of finding all possible numerators  $k$  to that of finding representatives for finitely many residue classes modulo  $(q^2 - 1)q$ . By means of the proof of this result we have determined all possible numerators  $k$  for  $2 \leq q \leq 60$ , the case  $q = 1$  being trivial. The result of this search suggests a conjecture about all possible values  $k/q$ ,  $(k, q) = 1$ , of Dedekind sums  $S(a, b)$  for an arbitrary  $q \in \mathbb{N}$ .

## 1. Introduction and main results

Let  $a$  be an integer,  $b$  a natural number, and  $(a, b) = 1$ . The classical Dedekind sum  $s(a, b)$  is defined by

$$s(a, b) = \sum_{k=1}^b ((k/b))((ak/b)).$$

Here

$$((x)) = \begin{cases} x - [x] - 1/2 & \text{if } x \in \mathbb{R} \setminus \mathbb{Z}; \\ 0 & \text{if } x \in \mathbb{Z} \end{cases}$$

(see [5, p. 1]). It is often more convenient to work with

$$S(a, b) = 12s(a, b)$$

instead. We call  $S(a, b)$  a *normalized* Dedekind sum.

The values of normalized Dedekind sums are rational numbers, whose properties have been studied by many authors. For instance, it is known that they are dense in the field  $\mathbb{R}$  (see [4]); and zones of large and small values of normalized Dedekind sums have been described by the present author (see [2]). It is unknown, however, which rational numbers occur as the values of normalized Dedekind sums. It seems that there is not even a conjecture in this direction. In this paper we present a conjecture of this kind that is based on empirical data.

Our first main result says that the numerators of normalized Dedekind sums with a given denominator  $q$  do not appear isolated, but form certain residue classes.

**Theorem 1** *Let  $q$  be a natural number and  $k$  an integer with  $(k, q) = 1$ . If  $k/q$  is the value of a normalized Dedekind sum, then all numbers*

$$k'/q \text{ with } k' \in \mathbb{Z}, k' \equiv k \pmod{(q^2 - 1)q},$$

*are values of normalized Dedekind sums.*

The case  $q = 1$  of Theorem 1 is trivial, since  $(q^2 - 1)q = 0$  and 0 is the only integer value of a normalized Dedekind sum (see the first remark following the proof of Theorem 3 below).

Theorem 1 reduces the task of finding all values  $k/q$ ,  $(k, q) = 1$ , of normalized Dedekind sums to finitely many cases for each  $q$ . One just has to find a representative  $k$  of each residue class modulo  $(q^2 - 1)q$  in question.

The *proof* of this theorem provides the tools for finding these representatives in particular cases. In this way we have determined, for  $2 \leq q \leq 60$ , all possible numerators  $k$  of normalized Dedekind sums  $S(a, b) = k/q$ ,  $(k, q) = 1$ . The result is as follows.

**Theorem 2** *Let  $q$  be a natural number,  $2 \leq q \leq 60$ , and  $k$  an integer with  $(k, q) = 1$ . Then  $k/q$  is the value of a normalized Dedekind sum, if, and only if,  $k$  satisfies the following conditions.*

(a) *If  $3 \nmid q$ , then  $k \equiv 0 \pmod{3}$ .*

(b) *If  $2 \nmid q$ , then*

$$k \equiv \begin{cases} 2 \pmod{4}, & \text{if } q \equiv 3 \pmod{4}; \\ 0 \pmod{8}, & \text{if } q \text{ is a square}; \\ 0 \pmod{4}, & \text{otherwise.} \end{cases}$$

We will show that the conditions (a) and (b) necessarily hold if an arbitrary rational number  $k/q$ ,  $(k, q) = 1$ , is the value of a normalized Dedekind sum (see Propositions 1 and 2 below). By Theorem 2, these conditions are sufficient for  $2 \leq q \leq 60$ . This fact leads to the following conjecture.

*Conjecture.* Let  $q \geq 2$  be a natural number,  $k \in \mathbb{Z}$ ,  $(k, q) = 1$ . Then  $k/q$  is the value of a normalized Dedekind sum if, and only if,  $k$  satisfies the conditions (a) and (b) of Theorem 2.

## 2. The details

The basis of the present work is the following theorem.

**Theorem 3** *Let  $b, q$  be natural numbers,  $a$  an integer,  $(a, b) = 1$ . Then  $S(a, b)$  takes the form  $k/q$  for some  $k \in \mathbb{Z}$ ,  $(k, q) = 1$  if, and only if,  $b$  has the form*

$$b = \frac{q(a^2 + 1)}{t}, \tag{1}$$

*where  $t$  is a natural number,  $(t, q) = 1$ . (Note that the condition  $(t, q) = 1$  implies  $t \mid a^2 + 1$ .)*

*Proof.* It suffices to assume  $a \in \mathbb{N}$  since  $S(-a, b) = -S(a, b)$  and  $a^2 + 1 = (-a)^2 + 1$ . The reciprocity law for normalized Dedekind sums (see [5, p. 5]) gives

$$abS(a, b) = -abS(b, a) + a^2 + b^2 + 1 - 3ab.$$

By [5, p. 27, Th. 2],  $aS(b, a)$  is an integer, so  $abS(a, b) \equiv a^2 + 1 \pmod{b}$  or

$$aS(a, b) \equiv \frac{a^2 + 1}{b} \pmod{\mathbb{Z}}.$$

This means that  $q = b/(b, a^2 + 1)$  is the smallest possible denominator of  $aS(a, b)$ . Since  $bS(a, b) \in \mathbb{Z}$  and  $(a, b) = 1$ , it is also the smallest possible denominator of  $S(a, b)$ . Hence we may write  $b = q(a^2 + 1)/t$  with some natural number  $t$  dividing  $a^2 + 1$ . It is easy to see that  $(q, t) = 1$ , since, otherwise, we would obtain a smaller denominator than  $q$ . Conversely, if  $b$  has the form (1) with  $(t, q) = 1$ , one sees that  $(a^2 + 1)/t = (b, a^2 + 1)$ . Indeed, if  $(a^2 + 1)/t \cdot r$  divides both  $b$  and  $a^2 + 1$ , then  $r$  divides both  $q$  and  $t$ , and so  $r = 1$ . Accordingly,  $S(a, b)$  has the smallest possible denominator  $q = bt/(a^2 + 1)$ .  $\square$

*Remarks.* 1. For  $q = 1$ , Theorem 3 shows that  $b$  takes the form  $b = (a^2 + 1)/t$ . Hence  $a^2 \equiv -1 \pmod{b}$ . This, however, implies that  $S(a, b) = 0$ , since  $-a$  is a multiplicative inverse of  $a$  modulo  $b$ , and so  $S(a, b) = S(-a, b) = -S(a, b)$ . In other words, 0 is the only integer that is the value of a normalized Dedekind sum. This fact is well-known (see [5, p. 36]).

2. A weaker form of Theorem 3 can be found in the author's paper [3].

In what follows we assume that  $S(a, b) = k/q$ ,  $k \in \mathbb{Z}$ ,  $q \in \mathbb{N}$ ,  $(k, q) = 1$ . By Theorem 3 this is the same as saying  $b = q(a^2 + 1)/t$  for some  $t \in \mathbb{N}$  with  $(t, q) = 1$ . The following theorem expresses  $k/q$  in terms of two normalized Dedekind sums.

**Theorem 4** *Let  $q, t$  be natural numbers  $(t, q) = 1$ ,  $a$  an integer,  $(a, q) = 1$ , and  $b = q(a^2 + 1)/t$ . Then*

$$S(a, b) = \frac{(q^2 - 1)a}{tq} - S(aq, t) + S(at^*, q), \quad (2)$$

where  $t^*$  is an integer satisfying  $tt^* \equiv 1 \pmod{q}$ .

*Proof.* First we assume  $a \in \mathbb{N}$ . Then the reciprocity law gives

$$S(a, b) = -S(b, a) + \frac{a}{b} + \frac{b}{a} + \frac{1}{ab} - 3. \quad (3)$$

We apply the three-term relation for normalized Dedekind sums to  $S(b, a)$  (see [1, Lemma 1]). It gives

$$S(b, a) = S(aq, t) + S(r, q) + \frac{a}{tq} + \frac{t}{aq} + \frac{q}{at} - 3, \quad (4)$$

where  $r$  is defined as follows: Let  $j, l$  be integers such that  $-aqj + tl = 1$ . Then  $r = -al + bj$ . However, we only need the residue class of  $r \pmod{q}$ . Obviously, we have  $l \equiv t^* \pmod{q}$  and  $r \equiv -al \pmod{q}$ . So we may choose  $r = -at^*$ . Now equations (3) and (4), in combination with  $b = q(a^2 + 1)/t$ , yield (2).

In the case  $a < 0$  we write  $S(a, b) = -S(|a|, b)$  and expand  $S(|a|, b)$  as above. In the end, we obtain (2) again.  $\square$

*Proof of Theorem 1.* Let  $k \in \mathbb{Z}$ ,  $q \in \mathbb{N}$ ,  $(k, q) = 1$ , and suppose  $k/q = S(a, b)$ . By Theorem 3,  $b = q(a^2 + 1)/t$  with  $t \in \mathbb{N}$ ,  $(t, q) = 1$ . By Theorem 4,  $S(a, b)$  is given by (2). Let  $m \in \mathbb{Z}$  and  $a' = a + mtq$ ,  $b' = q(a'^2 + 1)/t$  (observe that  $a'^2 + 1 \equiv a^2 + 1 \equiv 0 \pmod{t}$ ). Then (2) gives

$$S(a', b') = \frac{(q^2 - 1)a'}{tq} - S(a'q, t) + S(a't^*, q).$$

Here, however,  $a'/(tq) = a/(tq) + m$ ; moreover,  $S(a'q, t) = S(aq, t)$ , since  $a' \equiv a \pmod{t}$ , and  $S(a't^*, q) = S(at^*, q)$ , since  $a' \equiv a \pmod{q}$ . Altogether, we have

$$S(a', b') = (q^2 - 1)m + S(a, b) = (q^2 - 1)m + \frac{k}{q}.$$

If we write  $(q^2 - 1)m + k/q$  in the form  $((q^2 - 1)qm + k)/q$ , we see that  $k'/q$  is the value of a normalized Dedekind sum for each number  $k' \equiv k \pmod{(q^2 - 1)q}$ .  $\square$

The next two propositions show that the conditions (a) and (b) of Theorem 2 are necessary for  $k/q$ ,  $(k, q) = 1$ , being the value of a normalized Dedekind sum.

**Proposition 1** *Let  $q \in \mathbb{N}$  and  $k \in \mathbb{Z}$ ,  $(k, q) = 1$  be such that  $k/q$  is the value of a normalized Dedekind sum. If  $3 \nmid q$ , then  $k \equiv 0 \pmod{3}$ . If  $q \equiv 3 \pmod{4}$ , then  $k \equiv 2 \pmod{4}$ . If  $q \equiv 1 \pmod{4}$ , then  $k \equiv 0 \pmod{4}$ .*

*Proof.* Suppose that  $k/q = S(a, b)$  with  $b = q(a^2 + 1)/t$  and  $(t, q) = 1$ . If  $3 \nmid q$ , we have  $3 \nmid b$ , since  $a^2 + 1/t \not\equiv 0 \pmod{3}$  (note that the prime divisors  $p$  of  $a^2 + 1$  are either  $p = 2$  or satisfy  $p \equiv 1 \pmod{4}$ ,  $-1$  being a quadratic residue mod  $p$ ). In this case  $bS(a, b) \equiv 0 \pmod{3}$  (see [5, p. 27, Th. 2]) and so  $k \equiv 0 \pmod{3}$ .

Now suppose that  $q$  is odd. We first consider the case that  $b$  is odd. A classical result of Dedekind says

$$bS(a, b) \equiv b + 1 - 2 \left( \frac{a}{b} \right) \pmod{8}, \quad (5)$$

where  $\left( \frac{a}{b} \right)$  is the Jacobi symbol (see [5, p. 34, formula (42)]). Since  $(a^2 + 1)/t$  is odd, it must be  $\equiv 1 \pmod{4}$ , by the above argument. Accordingly,  $b \equiv q \pmod{4}$ . If  $q$  (and, thus,  $b$ ) is  $\equiv 3 \pmod{4}$ , then (5) gives  $bS(a, b) \equiv 2 \pmod{4}$  and so  $k \equiv 2 \pmod{4}$ . In the case  $q \equiv 1 \pmod{4}$  we obtain, in the same way,  $k \equiv 0 \pmod{4}$ .

Let  $q$  be odd and  $b$  even. Then  $(a^2 + 1)/t$  is even. Further,  $t$  must be odd, since otherwise the common divisor 2 of  $a^2 + 1$  and  $t$  disappears from  $b$ . Put  $a' = a + tq$  and  $b' = q(a'^2 + 1)/t$ . From  $a'^2 + 1 = a^2 + 1 + 2atq + t^2q^2$  we see that  $(a'^2 + 1)/t$  is odd, and so  $b'$  is odd. Theorem 4 shows that  $S(a', b') = k'/q$  with  $k' = k + (q^2 - 1)q$ . However,  $q^2 - 1 \equiv 0 \pmod{8}$ , and so  $k' \equiv k \pmod{8}$ . From the above we have  $k' \equiv 2 \pmod{4}$  if  $q \equiv 3 \pmod{4}$ , and  $k' \equiv 0 \pmod{4}$  if  $q \equiv 1 \pmod{4}$ . The respective congruences also hold for  $k$  instead of  $k'$ .  $\square$

**Proposition 2** *Let  $q$  be an odd square and  $k \in \mathbb{Z}$ ,  $(k, q) = 1$ , be such that  $k/q$  is the value of a normalized Dedekind sum. Then  $k \equiv 0 \pmod{8}$ .*

*Proof.* Let  $k/q = S(a, b)$  with  $b = q(a^2 + 1)/t$ ,  $(t, q) = 1$ .

*Case 1:* Suppose that  $b$  is odd and  $a$  is even. Then  $t$  is odd. We write  $a = 2^j a'$  with an odd number  $a'$ . In view of (5), we have to evaluate  $\left( \frac{a}{b} \right)$ . Obviously,

$$\left( \frac{a}{b} \right) = \left( \frac{2}{b} \right)^j \left( \frac{a'}{b} \right). \quad (6)$$

Since  $b \equiv 1 \pmod{4}$ ,

$$\left(\frac{a'}{b}\right) = \left(\frac{b}{a'}\right) = \left(\frac{q(a^2+1)/t}{a'}\right) = \left(\frac{q}{a'}\right) \left(\frac{t}{a'}\right),$$

because

$$\left(\frac{(a^2+1)/t}{a'}\right) \left(\frac{t}{a'}\right) = \left(\frac{a^2+1}{a'}\right) = 1.$$

However,  $\left(\frac{q}{a'}\right) = 1$ , and so

$$\left(\frac{a'}{b}\right) = \left(\frac{t}{a'}\right) = \left(\frac{a'}{t}\right). \quad (7)$$

On the other hand,

$$\left(\frac{2}{b}\right) = \left(\frac{2}{q}\right) \left(\frac{2}{a^2+1}\right) \left(\frac{2}{t}\right).$$

Here  $\left(\frac{2}{q}\right) = 1$ . It is easy to see that

$$\left(\frac{2}{a^2+1}\right) = \begin{cases} -1, & \text{if } j = 1; \\ 1, & \text{if } j \geq 2. \end{cases} \quad (8)$$

From (6), (7) and (8) we conclude

$$\left(\frac{a}{b}\right) = \begin{cases} -\left(\frac{a}{t}\right), & \text{if } j = 1; \\ \left(\frac{a}{t}\right), & \text{if } j \geq 2. \end{cases} \quad (9)$$

In order to evaluate  $\left(\frac{a}{t}\right)$ , we consider an arbitrary prime divisor  $p$  of  $t$ . Since  $a^2 \equiv -1 \pmod{p}$ ,  $\bar{a}$  is an element of order 4 in the prime residue group  $(\mathbb{Z}/\mathbb{Z}p)^\times$ . Accordingly,  $\bar{a}$  is a square if, and only if, 8 divides the order of this group, i.e.,  $p \equiv 1 \pmod{8}$ . This gives

$$\left(\frac{a}{p}\right) = \begin{cases} 1, & \text{if } p \equiv 1 \pmod{8}; \\ -1, & \text{if } p \equiv 5 \pmod{8}. \end{cases} \quad (10)$$

If we write  $t = p_1 \cdots p_r$  with (not necessarily distinct) prime factors  $p_i$  and apply (10) to each of these factors, we see that  $\left(\frac{a}{t}\right) = 1$  if, and only if,  $t \equiv 1 \pmod{8}$ . In view of (9), we obtain that  $\left(\frac{a}{b}\right) = 1$  if, and only if,  $j = 1$  and  $t \equiv 5 \pmod{8}$  or  $j \geq 2$  and  $t \equiv 1 \pmod{8}$ . This is equivalent to  $(a^2+1)/t \equiv 1 \pmod{8}$ . Since  $q \equiv 1 \pmod{8}$ , we have

$$\left(\frac{a}{b}\right) = \begin{cases} 1, & \text{if } b \equiv 1 \pmod{8}; \\ -1, & \text{if } b \equiv 5 \pmod{8}. \end{cases}$$

Accordingly, (5) gives  $bS(a, b) \equiv 0 \pmod{8}$  and  $k \equiv 0 \pmod{8}$ .

*Case 2:*  $b$  and  $a$  are both odd. Then  $t$  is even, more precisely,  $t = 2t'$  with  $t' \equiv 1 \pmod{4}$ . We have

$$\left(\frac{a}{b}\right) = \left(\frac{b}{a}\right) = \left(\frac{q}{a}\right) \left(\frac{a^2+1}{a}\right) \left(\frac{t}{a}\right) = \left(\frac{t}{a}\right) = \left(\frac{2}{a}\right) \left(\frac{t'}{a}\right) = \left(\frac{2}{a}\right) \left(\frac{a}{t'}\right). \quad (11)$$

Now

$$\left(\frac{2}{a}\right) = \begin{cases} 1, & \text{if } a^2 \equiv 1 \pmod{16}; \\ -1, & \text{if } a^2 \equiv 9 \pmod{16}. \end{cases}$$

Moreover,  $(a^2 + 1)/2 \equiv 1 \pmod{8}$  if, and only if,  $a^2 \equiv 1 \pmod{16}$ , i.e.,  $\left(\frac{2}{a}\right) = 1$ . As in case 1, one shows that  $\left(\frac{a}{t'}\right) = 1$  if, and only if,  $t' \equiv 1 \pmod{8}$ . On observing

$$b = q \frac{a^2 + 1}{2} / t'$$

and  $q \equiv 1 \pmod{8}$ , we have  $b \equiv 1 \pmod{8}$  if, and only if,  $(a^2 + 1)/2 \equiv 1 \pmod{8}$  and  $t' \equiv 1 \pmod{8}$  or  $(a^2 + 1)/2 \equiv 5 \pmod{8}$  and  $t' \equiv 5 \pmod{8}$ . By the above, however, this is equivalent to  $\left(\frac{2}{a}\right) = 1$  and  $\left(\frac{a}{t'}\right) = 1$  or  $\left(\frac{2}{a}\right) = -1$  and  $\left(\frac{a}{t'}\right) = -1$ . In view of (11), we obtain that  $b \equiv 1 \pmod{8}$  if, and only if,  $\left(\frac{a}{b}\right) = 1$ . Again, (5) shows  $k \equiv 0 \pmod{8}$ .

*Case 3:*  $b$  is even. Then  $t$  must be odd. We reduce this case to the case “ $b$  odd” just as we did in the proof of Proposition 1: Put  $a' = a + tq$ ,  $b' = q(a^2 + 1)/t$ . Then  $b'$  is odd and  $S(a', b') = k'/q$  with  $k' \equiv k \pmod{8}$ . By the cases 1 and 2,  $k' \equiv 0 \pmod{8}$ , hence  $k \equiv 0 \pmod{8}$ .  $\square$

The *proof of Theorem 2* is delivered by the computer, of course. We describe the search procedure that gave us this theorem. In view of Theorem 4, we worked with a list of pairs  $(t, a_1)$ ,  $t \geq 1$ ,  $a_1 \geq 0$ , such that  $(t, q) = 1$ ,  $0 \leq a_1 \leq t/2$  and  $a_1^2 + 1 \equiv 0 \pmod{t}$ . Note that  $t$  may occur in this list more than once if it is the product of two or more distinct primes  $\equiv 1 \pmod{4}$ . To any such pair we determined the numbers  $a = a_1 + tj$ ,  $j = 0, \dots, q-1$ , such that  $(a, q) = 1$ . In this way  $a$  yields all residues  $a_2$ ,  $0 \leq a_2 \leq q-1$ ,  $(a_2, q) = 1$  modulo  $q$  (and the corresponding Dedekind sums  $S(a_2 t^*, q)$  in the sense of Theorem 4). Since  $a \equiv a_1 \pmod{t}$ ,  $t$  divides  $a^2 + 1$ . Therefore,  $b = q(a^2 + 1)/t \in \mathbb{N}$ . Of course,  $(a, b) = 1$ .

We computed  $S(a, b)$  and determined the residue class  $\overline{k}$  modulo  $(q^2 - 1)q$  of the numerator of  $S(a, b)$  by a representative  $k$ . In the case  $q = 2$  there is only the residue class  $\overline{3}$  modulo  $(q^2 - 1)q$  satisfying the conditions (a) and (b) of Theorem 2. If  $q \geq 3$ , these residue classes take the form  $\overline{\pm k}$ ,  $1 \leq k < (q^2 - 1)q/2$ . Observe that it suffices to find one of the numbers  $\pm k$ , since the other one arises from  $S(-a, b) = -S(a, b)$ .

It is easy to compute the number of those  $k$ ,  $1 \leq k < (q^2 - 1)q/2$ ,  $(k, q) = 1$ , for which  $\pm k$  satisfies the conditions (a) and (b) of Theorem 2. For instance, if  $q = 17$  this number is 192, and for  $q = 48$  it is 18424. If we compute the said normalized Dedekind sums  $S(a, b)$  successively, we see that the residue classes  $\overline{k}$  modulo  $(q^2 - 1)q$  appear with different frequencies. Some of them occur at an early stage of the computation and then repeatedly, others only after many attempts. This has the effect that a long list of pairs  $(t, a_1)$  may be necessary for verifying the occurrence of all possible residue classes. In the case  $q = 24$ , we needed about 65000 normalized Dedekind sums to verify the occurrence of all 2300 classes, in the case  $q = 60$ , however, about 2.5 million normalized Dedekind sums for all 28792 classes (in all cases up to the factor  $\pm 1$ ).

*Example.* Let  $q = 7$ , hence  $(q^2 - 1)q = 336$ . In this case there are 12 numbers  $k$ ,  $1 \leq k < 336/2$ ,  $(k, 7) = 1$ , such that  $\pm k$  satisfies conditions (a) and (b) of Theorem 2, namely  $k = 6, 18, 30, 54, 66, 78, 90, 102, 114, 138, 150, 162$ . In the following table we list the smallest numbers  $t$  together with  $a$  and  $b$ , for which the normalized Dedekind sum  $S(a, b) = k'/7$  is such that  $k' \equiv \pm k \pmod{336}$ . We also note which of the signs  $\pm$  holds

here. Recall that  $a$  has the form  $a = a_1 + tj$ ,  $j = 0, \dots, 6$ ,  $(a, 7) = 1$ , where  $a_1^2 + 1 \equiv 0 \pmod t$ . Further,  $b = 7(a^2 + 1)/t$ . The table shows that all numbers  $k$  appear with  $t = 1, 2, 5$ .

$t$	$a$	$b$	$S(a, b)$	$k$
1	1	14	78/7	78
1	2	35	102/7	102
1	3	70	138/7	138
2	1	7	30/7	30
2	3	35	66/7	66
2	5	91	90/7	90
5	2	7	6/7	6
5	12	203	162/7	162
5	17	406	186/7	-150
5	22	679	222/7	-114
5	27	1022	282/7	-54
5	32	1435	318/7	-18

## References

- [1] K. Girstmair, Dedekind sums with predictable signs, *Acta Arith.* 83 (1998), 283–294.
- [2] K. Girstmair, Zones of large and small values for Dedekind sums, *Acta Arith.* 109 (2003), 299–308.
- [3] K. Girstmair, Dedekind sums with small denominators, *Int. J. Number Th.* 8 (2012), 1965–1970.
- [4] D. Hickerson, Continued fractions and density results for Dedekind sums, *J. Reine Angew. Math.* 290 (1977), 113–116.
- [5] H. Rademacher, E. Grosswald, *Dedekind sums*, Mathematical Association of America, 1972.

Kurt Girstmair  
 Institut für Mathematik  
 Universität Innsbruck  
 Technikerstr. 13/7  
 A-6020 Innsbruck, Austria  
 Kurt.Girstmair@uibk.ac.at