

# Recursive constructions of $k$ -normal polynomials over finite fields

Mahmood Alizadeh

Department of Mathematics, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran  
alizadeh@iauahvaz.ac.ir

Saeid Mehrabi

Department of Mathematics, Farhangian University, Tehran, Iran  
saeid\_mehrabi@yahoo.com

## Abstract

The paper is devoted to produce infinite sequences of  $k$ -normal polynomials  $F_u(x) \in \mathbb{F}_q[x]$  of degrees  $np^u$  ( $u \geq 0$ ), for a suitably chosen initial  $k$ -normal polynomial  $F_0(x) \in \mathbb{F}_q[x]$  of degree  $n$  over  $\mathbb{F}_q$  by iteratively applying the transformation  $x \rightarrow \frac{x^p - x}{x^p - x + \delta}$ , where  $\delta \in \mathbb{F}_q$  and  $0 \leq k \leq n - 1$ .

Keywords: Finite fields; normal basis;  $k$ -normal element;  $k$ -normal polynomial.  
Mathematics Subject Classification: 12A20

## 1 Introduction

Let  $\mathbb{F}_q$  be the Galois field of order  $q = p^m$ , where  $p$  is a prime and  $m$  is a natural number, and  $\mathbb{F}_q^*$  be its multiplicative group. A *normal basis* of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$  is a basis of the form  $N = \{\alpha, \alpha^q, \dots, \alpha^{q^{n-1}}\}$ , i.e. a basis that consists of the algebraic conjugates of a fixed element  $\alpha \in \mathbb{F}_{q^n}^*$ . Such an element  $\alpha \in \mathbb{F}_{q^n}$  is said to generate a normal basis for  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$ , and for convenience, is called a *normal element*.

A monic irreducible polynomial  $F(x) \in \mathbb{F}_q[x]$  is called *normal polynomial* or *N-polynomial* if its roots are linearly independent over  $\mathbb{F}_q$ . Since the elements in a normal basis are exactly the roots of some N-polynomials, there is a canonical one-to-one correspondence between N-polynomials and normal elements. Normal bases have many applications, including coding theory, cryptography and computer algebra systems. For further details, see [9].

Recently, the  $k$ -normal elements over finite fields are defined and characterized by Huczynska et al [8]. For each  $0 \leq k \leq n - 1$ , the element  $\alpha \in \mathbb{F}_{q^n}$  is called a  $k$ -normal element if  $\deg(\gcd(x^n - 1, \sum_{i=0}^{n-1} \alpha^{q^i} x^{n-1-i})) = k$ .

By analogy with the case of normal polynomials, a monic irreducible polynomial  $P(x) \in \mathbb{F}_q[x]$  of degree  $n$  is called a  $k$ -normal polynomial (or  $N_k$ -polynomial) over  $\mathbb{F}_q$  if its roots are  $k$ -normal elements of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$ . Here,  $P(x)$  has  $n$  distinct conjugate roots, of which  $(n - k)$  are linearly independent. Recall that an element  $\alpha \in \mathbb{F}_{q^n}$  is called a proper element of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$  if  $\alpha \notin \mathbb{F}_{q^v}$  for any proper divisor  $v$  of  $n$ . So, the element  $\alpha \in \mathbb{F}_{q^n}$  is a proper  $k$ -normal element of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$  if  $\alpha$  is a  $k$ -normal and proper element of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$ .

Using the above mention, a normal polynomial (or element) is a 0-normal polynomial (or element). Since the proper  $k$ -normal elements of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$  are the roots of a  $k$ -normal polynomial of degree  $n$  over  $\mathbb{F}_q$ , hence the  $k$ -normal polynomials of degree  $n$  over  $\mathbb{F}_q$  is just another way of describing the proper  $k$ -normal elements of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$ . Some results regarding constructions of spacial sequences of  $k$ -normal polynomials over  $\mathbb{F}_q$ , in the cases  $k = 0$  and 1 can be found in [2, 4, 5, 10, 11] and [6], respectively.

In this paper, a recursive method for constructing the  $k$ -normal polynomials of higher degree from a given  $k$ -normal polynomial of degree  $n$  over  $\mathbb{F}_q$  for each  $0 \leq k \leq n - 1$  is given.

## 2 Preliminary notes

We use the definitions, notations and results given by Huczynska [8], Gao [7] and Kyuregyan [10, 11], where similar problems are considered. We need the following results for our further study.

The trace of  $\alpha$  in  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$ , is given by  $Tr_{\mathbb{F}_{q^n}|\mathbb{F}_q}(\alpha) = \sum_{i=0}^{n-1} \alpha^{q^i}$ . For convenience,  $Tr_{\mathbb{F}_{q^n}|\mathbb{F}_q}$  is denoted by  $Tr_{q^n|q}$ .

**Proposition 2.1.** [6] *Let  $n = n_1 p^e$ , for some  $e \geq 1$  and  $a, b \in \mathbb{F}_q^*$ . Then the element  $\alpha$  is a proper  $k$ -normal element of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$  if and only if  $a + b\alpha$  is a proper  $k$ -normal element of  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$ .*

Let  $p$  denote the characteristic of  $\mathbb{F}_q$  and let  $n = n_1 p^e = n_1 t$ , with  $\gcd(p, n_1) = 1$  and suppose that  $x^n - 1$  has the following factorization in  $\mathbb{F}_q[x]$  :

$$x^n - 1 = (\varphi_1(x)\varphi_2(x) \cdots \varphi_r(x))^t, \quad (1)$$

where  $\varphi_i(x) \in \mathbb{F}_q[x]$  are the distinct irreducible factors of  $x^n - 1$ . For each  $s$ ,  $0 \leq s < n$ , let there is a  $u_s > 0$  such that  $R_{s,1}(x), R_{s,2}(x), \dots, R_{s,u_s}(x)$  are all of the  $s$  degree polynomials dividing  $x^n - 1$ . So, from (1) we can write  $R_{s,i}(x) = \prod_{j=1}^r \varphi_j^{t_{ij}}(x)$ , for each  $1 \leq i \leq u_s$ ,  $0 \leq t_{ij} \leq t$ . Let

$$\phi_{s,i}(x) = \frac{x^n - 1}{R_{s,i}(x)}, \quad (2)$$

for  $1 \leq i \leq u_s$ . Then, there is a useful characterization of the  $k$ -normal polynomials of degree  $n$  over  $\mathbb{F}_q$  as follows.

**Proposition 2.2.** [6] *Let  $F(x)$  be an irreducible polynomial of degree  $n$  over  $\mathbb{F}_q$  and  $\alpha$  be a root of it. Let  $x^n - 1$  factor as (1) and let  $\phi_{s,i}(x)$  be as in (2). Then  $F(x)$  is a  $N_k$ -polynomial over  $\mathbb{F}_q$  if and only if, there is  $j$ ,  $1 \leq j \leq u_k$ , such that*

$$L_{\phi_{k,j}}(\alpha) = 0,$$

and also

$$L_{\phi_{s,i}}(\alpha) \neq 0,$$

for each  $s$ ,  $k < s < n$ , and  $1 \leq i \leq u_s$ , where  $u_s$  is the number of all  $s$  degree polynomials dividing  $x^n - 1$  and  $L_{\phi_{s,i}}(x)$  is the linearized polynomial defined by

$$L_{\phi_{s,i}}(x) = \sum_{v=0}^{n-s} t_{iv} x^{qv} \text{ if } \phi_{s,i}(x) = \sum_{v=0}^{n-s} t_{iv} x^v.$$

The following propositions are useful for constructing  $N_k$ -polynomials over  $\mathbb{F}_q$ .

**Proposition 2.3.** [3] *Let  $x^p - \delta_2 x + \delta_0$  and  $x^p - \delta_2 x + \delta_1$  be relatively prime polynomials in  $\mathbb{F}_q[x]$  and  $P(x) = \sum_{i=0}^n c_i x^i$  be an irreducible polynomial of degree  $n \geq 2$  over  $\mathbb{F}_q$ , and let  $\delta_0, \delta_1 \in \mathbb{F}_q$ ,  $\delta_2 \in \mathbb{F}_q^*$ ,  $(\delta_0, \delta_1) \neq (0, 0)$ . Then*

$$F(x) = (x^p - \delta_2 x + \delta_1)^n P\left(\frac{x^p - \delta_2 x + \delta_0}{x^p - \delta_2 x + \delta_1}\right)$$

is an irreducible polynomial of degree  $np$  over  $\mathbb{F}_q$  if and only if  $\delta_2^{\frac{q-1}{p-1}} = 1$  and

$$Tr_{q|p}\left(\frac{1}{A^p}\left((\delta_1 - \delta_0)\frac{P'(1)}{P(1)} - n\delta_1\right)\right) \neq 0,$$

where  $A^{p-1} = \delta_2$ , for some  $A \in \mathbb{F}_q^*$ .

**Proposition 2.4.** [1] Let  $x^p - x + \delta_0$  and  $x^p - x + \delta_1$  be relatively prime polynomials in  $\mathbb{F}_q[x]$  and let  $P(x)$  be an irreducible polynomial of degree  $n \geq 2$  over  $\mathbb{F}_q$ , and  $0 \neq \delta_1, \delta_0 \in \mathbb{F}_p$ , such that  $\delta_0 \neq \delta_1$ . Define

$$F_0(x) = P(x)$$

$$F_k(x) = (x^p - x + \delta_1)^{t_k-1} F_{k-1} \left( \frac{x^p - x + \delta_0}{x^p - x + \delta_1} \right), \quad k \geq 1$$

where  $t_k = np^k$  denotes the degree of  $F_k(x)$ . Suppose that

$$\text{Tr}_{q|p} \left( \frac{(\delta_1 - \delta_0)F'_0(1) - n\delta_1 F_0(1)}{F_0(1)} \right) \cdot \text{Tr}_{q|p} \left( \frac{(\delta_1 - \delta_0)F'_0\left(\frac{\delta_0}{\delta_1}\right) + n\delta_1 F_0\left(\frac{\delta_0}{\delta_1}\right)}{F_0\left(\frac{\delta_0}{\delta_1}\right)} \right) \neq 0.$$

Then  $(F_k(x))_{k \geq 0}$  is a sequence of irreducible polynomials over  $\mathbb{F}_q$  of degree  $t_k = np^k$ , for every  $k \geq 0$ .

**Lemma 2.5.** Let  $\gamma$  be a proper element of  $\mathbb{F}_{q^n}$  and  $\theta \in \mathbb{F}_p^*$ , where  $q = p^m$ , ( $m \in \mathbb{N}$ ). Then we have

$$\sum_{j=0}^{p-1} \frac{1}{\gamma + j\theta} = -\frac{1}{\gamma^p - \gamma}. \quad (3)$$

*Proof.* By observing that

$$\sum_{j=0}^{p-1} \frac{1}{\gamma + j\theta} = \frac{1}{\gamma^p - \gamma} \left( \sum_{j=0}^{p-1} \frac{\gamma^p - \gamma}{\gamma + j\theta} \right), \quad (4)$$

it is enough to show that

$$\sum_{j=0}^{p-1} \frac{\gamma^p - \gamma}{\gamma + j\theta} = -1. \quad (5)$$

We note that

$$\begin{aligned} \sum_{j=0}^{p-1} \frac{\gamma^p - \gamma}{\gamma + j\theta} &= \sum_{j=0}^{p-1} \left( (\gamma + j\theta)^{p-1} - 1 \right) \\ &= \sum_{j=0}^{p-1} (\gamma + j\theta)^{p-1} \\ &= \sum_{j=1}^{p-1} \theta^j \binom{p-1}{j} \left( \sum_{i=1}^{p-1} i^j \right), \end{aligned} \quad (6)$$

where

$$\binom{p-1}{j} = \frac{(p-1)!}{(p-1-j)!j!}, \quad j \in \mathbb{F}_p^*.$$

On the other side, we know that

$$\sum_{i=1}^{p-1} i^j = \begin{cases} 0 \pmod{p}, & \text{if } p-1 \nmid j \\ -1 \pmod{p}, & \text{if } p-1 \mid j \end{cases} \quad (7)$$

and also  $\theta^{p-1} = 1$ . Thus by (6) and (7), the proof is completed.  $\square$

### 3 Recursive construction $N_k$ -polynomials

In this section we establish theorems which will show how propositions 2.2, 2.3 and 2.4 can be applied to produce infinite sequences of  $N_k$ -polynomials over  $\mathbb{F}_q$ . Recall that, the polynomial  $P^*(x) = x^n P\left(\frac{1}{x}\right)$  is called the reciprocal polynomial of  $P(x)$ , where  $n$  is the degree of  $P(x)$ . In the case  $k = 0$ , some similar results of the following theorems has been obtained in [2], [4] and ([5], Theorems 3.3.1 and 3.4.1). We use of an analogous technique to that used in the above results, where similar problems are considered.

**Theorem 3.1.** *Let  $P(x) = \sum_{i=0}^n c_i x^i$  be an  $N_k$ -polynomial of degree  $n$  over  $\mathbb{F}_q$ , for each  $n = rp^e$ , where  $e \in \mathbb{N}$  and  $r$  equals 1 or is a prime different from  $p$  and  $q$  a primitive element modulo  $r$ . Suppose that  $\delta \in \mathbb{F}_q^*$  and*

$$F(x) = (x^p - x + \delta)^n P^* \left( \frac{x^p - x}{x^p - x + \delta} \right). \quad (8)$$

Then  $F^*(x)$  is an  $N_k$ -polynomial of degree  $np$  over  $\mathbb{F}_q$  if  $k < p^e$  and

$$\text{Tr}_{q|p} \left( \delta \frac{P^{*'}(1)}{P^*(1)} \right) \neq 0.$$

*Proof.* Since  $P^*(x)$  is an irreducible polynomial over  $\mathbb{F}_q$ , so Proposition 2.3 and theorem's hypothesis imply that  $F(x)$  is irreducible over  $\mathbb{F}_q$ .

Let  $\alpha \in \mathbb{F}_{q^n}$  be a root of  $P(x)$ . Since  $P(x)$  is an  $N_k$ -polynomial of degree  $n$  over  $\mathbb{F}_q$  by theorem's hypothesis, then  $\alpha \in \mathbb{F}_{q^n}$  is a proper  $k$ -normal element over  $\mathbb{F}_q$ .

Since  $q$  is a primitive modulo  $r$ , so in the case  $r > 1$  the polynomial  $x^{r-1} + \dots + x + 1$  is irreducible over  $\mathbb{F}_q$ . Thus  $x^n - 1$  has the following factorization in  $\mathbb{F}_q[x]$ :

$$x^n - 1 = (\varphi_1(x) \cdot \varphi_2(x))^t, \quad (9)$$

where  $\varphi_1(x) = x - 1$ ,  $\varphi_2(x) = x^{r-1} + \dots + x + 1$  and  $t = p^e$ .

Letting that for each  $0 \leq s < n$  and  $1 \leq i \leq u_s$ ,  $R_{s,i}(x)$  is the  $s$  degree polynomial dividing  $x^n - 1$ , where  $u_s$  is the number of all  $s$  degree polynomials dividing  $x^n - 1$ . So, from (9), we can write  $R_{s,i}(x) = (x - 1)^{s_{1,i}} \cdot (x^{r-1} + \dots + x + 1)^{s_{2,i}}$ , where  $s = s_{1,i} + s_{2,i} \cdot (r - 1)$  for each  $0 \leq s_{1,i}, s_{2,i} \leq t$ , except when  $s_{1,i} = s_{2,i} = t$ . So, we have

$$\phi_{s,i}(x) = \frac{x^n - 1}{R_{s,i}(x)} = \frac{x^n - 1}{(x - 1)^{s_{1,i}} \cdot (x^{r-1} + \dots + x + 1)^{s_{2,i}}} = \sum_{v=0}^{n-s} t_{s,i,v} x^v. \quad (10)$$

Since  $P(x)$  is an  $N_k$ -polynomial of degree  $n$  over  $\mathbb{F}_q$ , so by Proposition 2.2, there is a  $j$ ,  $1 \leq j \leq u_k$ , such that

$$L_{\phi_{k,j}}(\alpha) = 0, \quad (11)$$

and also

$$L_{\phi_{s,i}}(\alpha) \neq 0, \quad (12)$$

for each  $k < s < n$  and  $1 \leq i \leq u_s$ . Further, we proceed by proving that  $F^*(x)$  is a  $k$ -normal polynomial. Let  $\alpha_1$  be a root of  $F(x)$ . Then  $\beta_1 = \frac{1}{\alpha_1}$  is a root of its reciprocal polynomial  $F^*(x)$ . Note that by (9), the polynomial  $x^{np} - 1$  has the following factorization in  $\mathbb{F}_q[x]$ :

$$x^{np} - 1 = (\varphi_1(x) \cdot \varphi_2(x))^{pt}, \quad (13)$$

where  $\varphi_1(x) = x - 1$ ,  $\varphi_2(x) = x^{r-1} + \dots + x + 1$  and  $t = p^e$ .

Letting that for each  $0 \leq s' < np$  and  $1 \leq i' \leq u'_{s'}$ ,  $R'_{s',i'}(x)$  is the  $s'$  degree polynomial dividing  $x^{np} - 1$ , where  $u'_{s'}$  is the number of all  $s'$  degree polynomials dividing  $x^{np} - 1$ . So, from

(13) we can write  $R'_{s',i'}(x) = (x-1)^{s'_{1,i'}} \cdot (x^{r-1} + \dots + x + 1)^{s'_{2,i'}}$ , where  $s' = s'_{1,i'} + s'_{2,i'} \cdot (r-1)$  for each  $0 \leq s'_{1,i'}, s'_{2,i'} \leq pt$ , except when  $s'_{1,i'} = s'_{2,i'} = pt$ . Therefore by considering

$$H'_{s',i'}(x) = \frac{x^{np} - 1}{R'_{s',i'}(x)}, \quad (14)$$

and Proposition 2.2,  $F^*(x)$  is an  $N_k$ -polynomial of degree  $np$  over  $\mathbb{F}_q$  if and only if there is a  $j'$ ,  $1 \leq j' \leq u'_k$ , such that

$$L_{H'_{k,j'}}(\beta_1) = 0,$$

and also

$$L_{H'_{s',i'}}(\beta_1) \neq 0,$$

for each  $k < s' < np$  and  $1 \leq i' \leq u'_{s'}$ . Consider

$$\begin{aligned} H_{s,i}(x) &= \frac{x^{np} - 1}{R_{s,i}(x)} \\ &= \frac{x^n - 1}{R_{s,i}(x)} \left( \sum_{j=0}^{p-1} x^{jn} \right), \end{aligned} \quad (15)$$

for each  $0 \leq s < n$  and  $1 \leq i \leq u_s$ . By (10) we obtain

$$H_{s,i}(x) = \phi_{s,i}(x) \left( \sum_{j=0}^{p-1} x^{jn} \right) = \sum_{v=0}^{n-s} t_{s,i,v} \left( \sum_{j=0}^{p-1} x^{jn+v} \right).$$

It follows that

$$L_{H_{s,i}}(\beta_1) = \sum_{v=0}^{n-s} t_{s,i,v} \left( \sum_{j=0}^{p-1} (\beta_1)^{p^{jmn}} \right)^{p^{mv}},$$

or

$$L_{H_{s,i}}(\beta_1) = L_{H_{s,i}} \left( \frac{1}{\alpha_1} \right) = \sum_{v=0}^{n-s} t_{s,i,v} \left( \sum_{j=0}^{p-1} \left( \frac{1}{\alpha_1} \right)^{p^{jmn}} \right)^{p^{mv}}. \quad (16)$$

From (8), if  $\alpha_1$  is a zero of  $F(x)$ , then  $\frac{\alpha_1^p - \alpha_1 + \delta}{\alpha_1^p - \alpha_1}$  is a zero of  $P(x)$ , and therefore it may assume that

$$\alpha = \frac{\alpha_1^p - \alpha_1 + \delta}{\alpha_1^p - \alpha_1},$$

or

$$\frac{\alpha - 1}{\delta} = (\alpha_1^p - \alpha_1)^{-1}. \quad (17)$$

Now, by (17) and observing that  $P(x)$  is an irreducible polynomial of degree  $n$  over  $\mathbb{F}_q$ , we obtain

$$\frac{\alpha - 1}{\delta} = \left( \frac{\alpha - 1}{\delta} \right)^{p^{mn}} = (\alpha_1^{p^{mn+1}} - \alpha_1^{p^{mn}})^{-1}. \quad (18)$$

It follows from (17) and (18) that

$$(\alpha_1^{p^{mn+1}} - \alpha_1^{p^{mn}})^{-1} = (\alpha_1^p - \alpha_1)^{-1}. \quad (19)$$

Also observing that  $F(x)$  is an irreducible polynomial of degree  $np$  over  $\mathbb{F}_q$ , we have  $(\alpha_1^p - \alpha_1) \neq 0$  and  $(\alpha_1^{p^{mn+1}} - \alpha_1^{p^{mn}}) \neq 0$ . Hence by (19)

$$(\alpha_1^{p^{mn}} - \alpha_1)^p = (\alpha_1^{p^{mn}} - \alpha_1). \quad (20)$$

It follows from (20) that  $\alpha_1 p^{mn} - \alpha_1 = \theta \in \mathbb{F}_p^*$ . Hence  $\alpha_1 p^{mn} = \alpha_1 + \theta$  and

$$\alpha_1 p^{2mn} = (\alpha_1 + \theta) p^{mn} = \alpha_1 p^{mn} + \theta p^{mn} = \alpha_1 + \theta + \theta = \alpha_1 + 2\theta.$$

It is easy to show that

$$\alpha_1 p^{jmn} = \alpha_1 + j\theta, \quad \text{for } 1 \leq j \leq p-1,$$

or

$$\left(\frac{1}{\alpha_1}\right)^{p^{jmn}} = \frac{1}{\alpha_1 + j\theta}, \quad \text{for } 1 \leq j \leq p-1. \quad (21)$$

From (16) and (21), we immediately obtain

$$L_{H_{s,i}}(\beta_1) = \sum_{v=0}^{n-s} t_{s,i,v} \left( \sum_{j=0}^{p-1} \frac{1}{\alpha_1 + j\theta} \right)^{p^{mv}}. \quad (22)$$

Thus, by (17), (22) and Lemma 2.5 we have

$$\begin{aligned} L_{H_{s,i}}(\beta_1) &= \sum_{v=0}^{n-s} t_{s,i,v} \left( -\frac{1}{\alpha_1^p - \alpha_1} \right)^{p^{mv}} \\ &= \frac{1}{\delta} \sum_{v=0}^{n-s} t_{s,i,v} (1 - \alpha)^{p^{mv}} \\ &= L_{\phi_{s,i}} \left( \frac{1 - \alpha}{\delta} \right). \end{aligned} \quad (23)$$

Since  $\alpha$  is a zero of  $P(x)$ , then  $\alpha$  will be a  $k$ -normal element in  $\mathbb{F}_{q^n}$  over  $\mathbb{F}_q$ . Thus according to Proposition 2.1, the element  $\frac{1-\alpha}{\delta}$  will also be a  $k$ -normal element.

since  $\frac{1-\alpha}{\delta}$  is a root of  $P(-\delta x + 1)$ , so by (23) and Proposition 2.2, there is a  $j$ ,  $1 \leq j \leq u_k$ , such that

$$L_{H_{k,j}}(\beta_1) = 0,$$

and also

$$L_{H_{s,i}}(\beta_1) \neq 0,$$

for each  $s$ ,  $k < s < n$  and  $1 \leq i \leq u_s$ . So, there is a  $j'$ ,  $1 \leq j' \leq u'_k$ , such that,  $L_{H'_{k,j'}}(\beta_1) = L_{H_{k,j}}(\beta_1) = 0$ . On the other side, by (14) and (15), for each  $s'$ ,  $k < s' < np$  and  $1 \leq i' \leq u'_{s'}$ , there is  $s$ ,  $k < s < n$  and  $1 \leq i \leq u_s$  such that  $H'_{s',i'}(x)$  divide  $H_{s,i}(x)$ . It follows that

$$L_{H'_{s',i'}}(\beta_1) \neq 0,$$

for each  $s'$ ,  $k < s' < np$  and  $1 \leq i' \leq u'_{s'}$ . The proof is completed.  $\square$

In the following theorem, a computationally simple and explicit recurrent method for constructing higher degree  $N_k$ -polynomials over  $\mathbb{F}_q$  starting from an  $N_k$ -polynomial is described.

**Theorem 3.2.** *Let  $P(x)$  be an  $N_k$ -polynomial of degree  $n$  over  $\mathbb{F}_q$ , for each  $n = rp^e$ , where  $e \in \mathbb{N}$  and  $r$  equals 1 or is a prime different from  $p$  and  $q$  a primitive element modulo  $r$ . Define*

$$\begin{aligned} F_0(x) &= P^*(x) \\ F_u(x) &= (x^p - x + \delta)^{np^{u-1}} F_{u-1} \left( \frac{x^p - x}{x^p - x + \delta} \right), \end{aligned} \quad (24)$$

where  $\delta \in \mathbb{F}_p^*$ . Then  $(F_u^*(x))_{u \geq 0}$  is a sequence of  $N_k$ -polynomials of degree  $np^u$  over  $\mathbb{F}_q$  if  $k < p^e$  and

$$\text{Tr}_{q|p} \left( \frac{P^{*'}(0)}{P^*(0)} \right) \cdot \text{Tr}_{q|p} \left( \frac{P^{*'}(1)}{P^*(1)} \right) \neq 0,$$

where  $P^{*'}(0)$  and  $P^{*'}(1)$  are the formal derivative of  $P^*(x)$  at the points  $x = 0$  and  $x = 1$ , respectively.

*Proof.* By Proposition 2.4 and hypotheses of theorem for each  $u \geq 1$ ,  $F_u(x)$  is an irreducible polynomial over  $\mathbb{F}_q$ . Consequently,  $(F_u^*(x))_{u \geq 0}$  is a sequence of irreducible polynomials over  $\mathbb{F}_q$ . The proof of  $k$ -normality of the irreducible polynomials  $F_u^*(x)$ , for each  $u \geq 1$  is implemented by mathematical induction on  $u$ . In the case  $u = 1$ , by Theorem 3.1  $F_1^*(x)$  is a  $k$ -normal polynomial.

For  $u = 2$  we show that  $F_2^*(x)$  is also a  $k$ -normal polynomial. To this end we need to show that the hypothesis of Theorem 3.1 are satisfied. By Theorem 3.1,  $F_2^*(x)$  is a  $k$ -normal polynomial if

$$\text{Tr}_{q|p} \left( \frac{F_1'(1)}{F_1(1)} \right) \neq 0,$$

since  $\delta \in \mathbb{F}_p^*$ . We apply (24) to compute

$$F_u(0) = F_u(1) = \delta^{un} P^*(0), \quad u = 1, 2, \dots \quad (25)$$

We calculate the formal derivative of  $F_1^*(x)$  at the points  $x = 0$  and  $x = 1$ . According to (24) the first derivative of  $F_1(x)$  is

$$\begin{aligned} F_1'(x) &= -n(x^p - x + \delta)^{n-1} F_0' \left( \frac{x^p - x}{x^p - x + \delta} \right) \\ &+ (x^p - x + \delta)^n \cdot \left( \frac{(px^{p-1} - 1)(x^p - x + \delta) - (px^{p-1} - 1)(x^p - x)}{(x^p - x + \delta)^2} \right) \\ &\cdot F_0' \left( \frac{x^p - x}{x^p - x + \delta} \right) \\ &= -\delta(x^p - x + \delta)^{n-2} \cdot P^{*'} \left( \frac{x^p - x}{x^p - x + \delta} \right), \end{aligned}$$

and at the points  $x = 0$  and  $x = 1$

$$F_1'(0) = -F_1'(1) = -\delta^{n-1} P^{*'}(0) \quad (26)$$

which is not equal to zero by the condition  $\text{Tr}_{q|p} \left( \frac{P^{*'}(0)}{P^*(0)} \right) \neq 0$  in the hypothesis of theorem, since  $\delta \in \mathbb{F}_p^*$ . From (26) and (25)

$$\begin{aligned} \text{Tr}_{q|p} \left( \frac{F_1'(1)}{F_1(1)} \right) &= \text{Tr}_{q|p} \left( \frac{-\delta^{n-1} P^{*'}(0)}{\delta^n P^*(0)} \right) \\ &= -\frac{1}{\delta} \text{Tr}_{q|p} \left( \frac{P^{*'}(0)}{P^*(0)} \right), \end{aligned} \quad (27)$$

which is not equal to zero by hypothesis of theorem. Hence the polynomial  $F_2^*(x)$  is a  $k$ -normal polynomial. If induction holds for  $u - 1$ , then it must hold also for  $u$ , that is by assuming that  $F_{u-1}^*(x)$  is a  $k$ -normal polynomial, we show that  $F_u^*(x)$  is also a  $k$ -normal polynomial. Let  $u \geq 3$ . By Theorem 3.1,  $F_u^*(x)$  is a  $k$ -normal polynomial if

$$\text{Tr}_{q|p} \left( \frac{F_{u-1}'(1)}{F_{u-1}(1)} \right) \neq 0,$$

since  $\delta \in \mathbb{F}_p^*$ . We calculate the formal derivative of  $F'_{u-1}(x)$  at the points 1 and 0. By (24) the first derivative of  $F_{u-1}(x)$  is

$$\begin{aligned} F'_{u-1}(x) &= (x^p - x + \delta)^{np^{u-2}} \left( \frac{(px^{p-1} - 1)(x^p - x + \delta) - (px^{p-1} - 1)(x^p - x)}{(x^p - x + \delta)^2} \right) \\ &\quad \cdot F'_{u-2} \left( \frac{x^p - x}{x^p - x + \delta} \right) \\ &= -\delta(x^p - x + \delta)^{np^{u-2}-2} F'_{u-2} \left( \frac{x^p - x}{x^p - x + \delta} \right), \end{aligned}$$

and at the point  $x = 0$  and  $x = 1$

$$F'_{u-1}(0) = F'_{u-1}(1) = -\delta^{np^{u-2}-1} F'_{u-2}(0) = -\delta^{n-1} F'_{u-2}(0). \quad (28)$$

So we have

$$F'_{u-1}(0) = F'_{u-1}(1) = (-1)^{u-2} \delta^{(n-1)(u-2)} F'_1(0), \quad (29)$$

which is not equal to zero by (26) and the condition  $Tr_{q|p} \left( \frac{P^{*'}(0)}{P^*(0)} \right) \neq 0$  in the hypothesis of theorem, since  $\delta \in \mathbb{F}_p^*$ . Also

$$Tr_{q|p} \left( \frac{F'_{u-1}(1)}{F_{u-1}(1)} \right) = (-1)^{u-2} \frac{1}{\delta^{(u-2)}} Tr_{q|p} \left( \frac{F'_1(1)}{F_1(1)} \right),$$

which is not equal to zero by (27) and hypothesis of theorem. The theorem is proved.  $\square$

## References

- [1] S. Abrahamyan, M. K. Kyureghyan, A recurrent method for constructing irreducible polynomials over finite fields, proceeding of the 13th international conference on computer algebra in scientific computing, 2011, 1-9.
- [2] S. Abrahamyan, M. K. Kyureghyan, New recursive construction of normal polynomials over finite fields, proceeding of the 11th international conference on finite fields and their applications, 2013, 1-10.
- [3] S. Abrahamyan, M. Alizadeh, M. K. Kyureghyan, Recursive constructions of irreducible polynomials over finite fields, Finite Fields Appl., 18 (2012), 738-745.
- [4] M. Alizadeh, S. Abrahamyan, S. Mehrabi, M. K. Kyureghyan, Constructing of  $N$ -polynomials over finite fields, International Journal of Algebra, Vol. 5, No. 29, 2011, 1437-1442.
- [5] M. Alizadeh, Construction of Irreducible and Normal Polynomials over Finite Fields, Ph.D. Thesis, National Academy of Sciences of Armenia, Yerevan, 2013.
- [6] M. Alizadeh, Some notes on the  $k$ -normal elements and  $k$ -normal polynomials over finite fields, J. Algebra Appl., DOI:10.1142/S0219498817500062, 2016.
- [7] S. Gao, Normal bases over finite fields, Ph.D. Thesis, Waterloo, 1993.
- [8] S. Huczynska, G. L. Mullen, D. Panario and D. Thomson, Existence and properties of  $k$ -normal elements over finite fields, Finite Fields Appl., 24 (2013), 170-183.
- [9] D. Jungnickel, Finite Fields: Structure and Arithmetics, Wissenschaftsverlag, (Mannheim) 1993.



- [10] M. K. Kyuregyan, Iterated construction of irreducible polynomials over finite fields with linearly independent roots, *Finite Fields Appl.*, 10 (2004) 323-341.
- [11] M. K. Kyuregyan, Recursive construction of N-polynomials over  $GF(2^s)$ , *Discrete Applied Mathematics*, No. 156, 2008, 1554-1559.
- [12] H. W. Lenstra and R. Schoof, Primitive normal bases for finite fields, *Mathematics of Computation*, 48 (1987), 217-231.
- [13] S. Schwartz, Irreducible polynomials over finite fields with linearly independent roots, *Math. Slovaca*, Vol. 38, 1988, 147-158.