

The average size of the 3-isogeny Selmer groups of elliptic curves

$$y^2 = x^3 + k$$

Manjul Bhargava, Noam Elkies, and Ari Shnidman

October 19, 2016

Abstract

The elliptic curve $E_k: y^2 = x^3 + k$ admits a natural 3-isogeny $\phi_k: E_k \rightarrow E_{-27k}$. We compute the average size of the ϕ_k -Selmer group as k varies over the integers. Unlike previous results of Bhargava and Shankar on n -Selmer groups of elliptic curves, we show that this average can be very sensitive to congruence conditions on k ; this sensitivity can be precisely controlled by the Tamagawa numbers of E_k and E_{-27k} . As consequences, we prove that the average rank of the curves E_k , $k \in \mathbb{Z}$, is less than 1.21 and over 23% (resp. 41%) of the curves in this family have rank 0 (resp. 3-Selmer rank 1).

1 Introduction

Let F be a field of characteristic not 2 or 3, and let $k \in F$ be non-zero. The elliptic curve

$$E_k: y^2 = x^3 + k \tag{1}$$

has j -invariant 0, and every elliptic curve E/F with $j(E) = 0$ is isomorphic to E_k for some $k \in F$. The curves E_k and $E_{k'}$ are isomorphic if and only if $k' = km^6$ for some $m \in F^\times$.

Over a separable closure \bar{F} , each E_k has complex multiplication by the Eisenstein integers $\mathbb{Z}[\zeta_3]$, and hence admits an endomorphism $\sqrt{-3}$ of degree 3. This endomorphism is not generally defined over F , but its kernel is, and consequently there is a 3-isogeny $\phi = \phi_k: E_k \rightarrow E_{k'}$ for some $k' \in F$. By duality, there is a 3-isogeny $\hat{\phi} = \hat{\phi}_k: E_{k'} \rightarrow E_k$ in the reverse direction. In fact, we may take $k' = -27k$, and then these isogenies are given explicitly as:

$$\phi_k: (x, y) \mapsto \left(\frac{x^3 + 4k}{x^2}, \frac{y(x^3 - 8k)}{x^3} \right), \tag{2}$$

$$\hat{\phi}_k: (x, y) \mapsto \left(\frac{x^3 - 108k}{9x^2}, \frac{y(x^3 + 216k)}{27x^3} \right). \tag{3}$$

After identifying E_{3^6k} and E_k , we have $\hat{\phi}_k = \phi_{-27k}$.

If F is a number field, then the ϕ -Selmer group

$$\text{Sel}_\phi(E_k) \subset H^1(G_F, E_k[\phi])$$

associated to the isogeny ϕ consists of all “locally soluble” cohomology classes, i.e., those that are locally in the image of the connecting map

$$\partial_v: E_{-27k}(F_v) \longrightarrow H^1(G_{F_v}, E_k[\phi])$$

for every place v of F ; here, $G_F = \text{Gal}(\bar{F}/F)$.

In this paper, we take $F = \mathbb{Q}$ and study the average size of the ϕ -Selmer group for the elliptic curves E_k as k varies. Let

$$r = \frac{103 \cdot 229}{2 \cdot 3^2 \cdot 7^2 \cdot 13} \prod_{p \equiv 5 \pmod{6}} \frac{(1-p^{-1})(1+p^{-1} + \frac{5}{3}p^{-2} + p^{-3} + \frac{5}{3}p^{-4} + p^{-5})}{1-p^{-6}}. \quad (4)$$

Then we prove:

Theorem 1. *When the elliptic curves $E_k: y^2 = x^3 + k$, $k \in \mathbb{Z}$, are ordered by the absolute value of k , the average size of the ϕ_k -Selmer group associated to the 3-isogeny $\phi_k: E_k \rightarrow E_{-27k}$ is $1+r$ if k is negative, and $1+r/3$ if k is positive.*

We note that, in this theorem, we may take k to range over all integers or, if desired, only the sixth-power-free ones (so that we obtain each isomorphism class of elliptic curve of j -invariant 0 exactly once). We can calculate the product in (4) efficiently by approximating it by a product of powers of the values at $s = 2, 3, 4, \dots$ of $(1-2^{-s})(1-3^{-s})\zeta(s)$ and $(1+2^{-s})L(\chi, s)$ where χ is the Legendre symbol mod 3; we find that the product's numerical value is $1.033735512017364858\dots$, so $r = 2.1265\dots$, making $1+r = 3.1265\dots$ and $1+r/3 = 1.7088\dots$

In fact, we are able to determine the average size of the ϕ_k -Selmer group of $E_k: y^2 = x^3 + k$ where k varies in any subset S of \mathbb{Z} defined by finitely many—or, in suitable cases, infinitely many—congruence conditions. Let $S \subset \mathbb{Z}$ be any subset of integers defined by sign conditions and congruence conditions modulo a power of each prime, such that for all sufficiently large primes p , the closure S_p of S in \mathbb{Z}_p contains all elements of \mathbb{Z}_p not divisible by p^2 . We call such a set *acceptable*. For example, the set of all sixth-power-free integers is acceptable. The set of all positive squarefree integers congruent to 1 (mod 3) is also acceptable.

For any isogeny $\phi: A \rightarrow A'$ of abelian varieties over \mathbb{Q} and for any prime $p \leq \infty$, define the *local Selmer ratio* of ϕ at p to be

$$c_p(\phi) := \frac{|A'(\mathbb{Q}_p)/\phi(A(\mathbb{Q}_p))|}{|A[\phi](\mathbb{Q}_p)|} \quad (5)$$

where $\mathbb{Q}_p = \mathbb{R}$ when $p = \infty$. Then the following theorem gives the average size of the ϕ_k -Selmer group as k varies over the integers in S :

Theorem 2. *Let S be any acceptable set of integers. When the elliptic curves $E_k: y^2 = x^3 + k$, $k \in S$, are ordered by the absolute value of k , the average size of the ϕ -Selmer group associated to the 3-isogeny $\phi: E_k \rightarrow E_{-27k}$ is*

$$1 + \prod_{p \leq \infty} \frac{\int_{k \in S_p} c_p(\phi_k) dk}{\int_{k \in S_p} dk}, \quad (6)$$

where for $p < \infty$, S_p denotes the p -adic closure of S in \mathbb{Z}_p and dk denotes the usual additive Haar measure on \mathbb{Z}_p , and for $p = \infty$, S_p denotes the smallest interval in \mathbb{R} containing S and dk denotes the usual additive measure on \mathbb{R} .

Remark 3. The factor at $p = \infty$ is equal to $\frac{1}{3}$, 1, or $\frac{2}{3}$, depending on whether S contains only positive, only negative, or both positive and negative integers, respectively.

Remark 4. The group $H^1(G_F, E_k[\phi])$ parameterizes isomorphism classes of ϕ -coverings, i.e., maps $f : C \rightarrow E_{-27k}$ of curves that become isomorphic to ϕ over \bar{F} . Therefore, we can interpret the local Selmer ratio $c_p(\phi)$ as the number of soluble ϕ -coverings f over \mathbb{Q}_p , where each f is weighted by the inverse of the order of its automorphism group. Theorem 2 can thus be interpreted as saying that the expected (weighted) number of nontrivial locally soluble ϕ -coverings over \mathbb{Q} is simply the product of the (weighted) number of soluble ϕ -coverings over \mathbb{Q}_p . This suggests a Selmer group analogue of Bhargava's conjectures for number fields [2], which would include as a special case the conjecture of Bhargava and Shankar that the average size of $\text{Sel}_n(E)$ over any congruence family of elliptic curves over \mathbb{Q} is $\sum_{d|n} d$.

For any given set S , the p -adic integrals in Theorem 2 can be evaluated explicitly using Proposition 28 in §4.1. In particular, when $S = \mathbb{Z}$, we recover Theorem 1.

We note that unlike the results of [6, 7, 8, 9] for the 2-, 3-, 4-, and 5-Selmer groups, respectively, the average size of the ϕ -Selmer group in Theorem 2 can depend very much on the congruences defining the set S . However, we show that we may partition the set of non-zero integers into a countable union $\cup_{m=-\infty}^{\infty} T_m$ of sets T_m , where each T_m is itself the union of countably many sets defined by congruence conditions, such that if $S \subset T_m$ is an acceptable set, then the average size of the ϕ -Selmer group of $E_k : y^2 = x^3 + k$, as k varies in S , depends only on m .

More precisely, we define the *global Selmer ratio*

$$c(\phi_k) := \prod_p c_p(\phi_k) \tag{7}$$

to be the product of the local Selmer ratios. If ϕ is an ℓ -isogeny, for some prime ℓ , then the global Selmer ratio $c(\phi)$ is evidently a power of ℓ . The importance of the global Selmer ratio in the study of the Selmer groups $\text{Sel}_\phi(E_k)$ is clearly seen in the following theorem.

Theorem 5. *For each $m \in \mathbb{Z}$, let $T_m := \{k \in \mathbb{Z} : c(\phi_k) = 3^m\}$. Then:*

- (i) $\mathbb{Z} \setminus \{0\}$ is the disjoint union of the T_m ;
- (ii) Each T_m has positive density;
- (iii) If $k \in T_m$, and $E_k(\mathbb{Q})$ and $E_{-27k}(\mathbb{Q})$ have trivial 3-torsion, then $|\text{Sel}_\phi(E_k)| = 3^m |\text{Sel}_{\hat{\phi}}(E_{-27k})|$;
- (iv) If $S = T_m$, or if S is any acceptable set contained in T_m , then the average size of $\text{Sel}_\phi(E_k)$, $k \in S$, is equal to $1 + 3^m$, and the average size of $\text{Sel}_{\hat{\phi}}(E_{-27k})$, $k \in S$, is equal to $1 + 3^{-m}$.

Thus the average size of $\text{Sel}_\phi(E_k)$, $k \in S$, is independent of congruence conditions if we also fix the global Selmer ratio. The densities of the sets T_m , to three decimal places, are tabulated in Table 1, along with the densities of the subsets T_m^+ (respectively, T_m^-) of T_m consisting of its positive (respectively, negative) elements. We will describe the sets T_m via explicit congruence conditions in §7.4.

Using Theorem 5 and the rigorous computation of the densities $\mu(T_m)$, we may obtain bounds on the limsup of the average 3-ranks of the ϕ - and $\hat{\phi}$ -Selmer groups. These immediately imply a bound on the limsup of the average 3-rank of the 3-Selmer group $\text{Sel}_3(E_k)$, and hence a bound on the limsup of the average rank of $E_k(\mathbb{Q})$ as k varies in \mathbb{Z} ordered by absolute value. We prove the following theorem:

m	$\mu(T_m)$	$\mu(T_m^+)$	$\mu(T_m^-)$
-4	.000	.000	.000
-3	.004	.004	.000
-2	.067	.063	.004
-1	.295	.231	.063
0	.399	.167	.231
1	.199	.031	.167
2	.032	.000	.031
3	.000	.000	.000
4	.000	.000	.000

Table 1: Densities of the sets T_m , T_m^+ , and T_m^- for $|m| \leq 4$.

Theorem 6. *The (limsup of the) average rank of the elliptic curves $E_k: y^2 = x^3 + k$, $k \in \mathbb{Z}$, is less than 1.29.*

Theorem 6 immediately implies that a positive proportion of elliptic curves $E_k: y^2 = x^3 + k$ must have rank 0 or 1. To produce positive proportions of curves having the individual ranks 0 or 1, we may make use of the fact in Theorem 5(iii) that for 100% of $k \in T_m$, the 3-rank of the ϕ -Selmer group of E_k is m more than that of its $\hat{\phi}$ -Selmer group. This means, in particular, that 100% of curves in T_0 have even 3-Selmer rank (Proposition 42(ii)). Since the average sizes of both the ϕ and $\hat{\phi}$ -Selmer groups are equal to 2, we conclude that at least 1/2 of all curves in T_0 must in fact have 3-Selmer rank 0. Hence we obtain the following theorem.

Theorem 7. *At least 19.9% of all elliptic curves $E_k: y^2 = x^3 + k$, $k \in \mathbb{Z}$, have rank 0.*

Conditionally on the finiteness of the 3-primary part of the Shafarevich–Tate group, we may obtain a similar result for rank 1 elliptic curves by using instead the set $T_1 \cup T_{-1}$, for which the 3-Selmer parity is always odd.

Theorem 8. *Assume for all $k \in T_1 \cup T_{-1}$ that the 3-primary part of $\text{III}(E_k)$ is finite. Then at least 41.1% of all elliptic curves $E_k: y^2 = x^3 + k$, $k \in \mathbb{Z}$, have rank 1.*

Using forthcoming results of Skinner (private communication), we may be able to deduce this last result unconditionally.

Combining Theorems 7 and 8, we obtain:

Corollary 9. *At least 61% of all elliptic curves $E_k: y^2 = x^3 + k$, $k \in \mathbb{Z}$, have rank 0 or 1.*

Thus the majority of curves $E_k: y^2 = x^3 + k$ have rank 0 or 1.

Remark 10. Theorems 6, 7, and 8 and Corollary 9 can be improved slightly if we combine our results with those in Ruth’s thesis [30], where it is shown that the average size of $\text{Sel}_2(E_k)$, $k \in S$, is at most 3. Indeed, this gives a rank bound of 4/3 on any T_m , which turns out to be a better bound than ours for $m > 1$. The improvements are small since even T_2 has very small density: incorporating Ruth’s result, we can show that the average rank of E_k is at most 1.21, and the proportion of rank 0 curves is at least 23.2%.

Finally, we note that Theorem 5 implies:

Theorem 11. *For each $m \geq 0$, a positive proportion of elliptic curves $E_k: y^2 = x^3 + k$, $k \in \mathbb{Z}$, have 3-Selmer rank equal to m .*

Indeed, the average size of $\text{Sel}_{\hat{\phi}}(E_k)$ is $1 + 3^{-m}$ for $k \in T_m$ and, therefore, a positive proportion of $k \in T_m$ satisfy $|\text{Sel}_{\hat{\phi}}(E_k)| = 1$. By Theorem 5(iii), we then have $|\text{Sel}_{\phi}(E_k)| = 3^m$ and thus $|\text{Sel}_3(E_k)| = 3^m$ for most of these k . Taking into account Remark 10 and Selmer parity [17] we see that, for even m , Theorem 11 holds not just for $\text{Sel}_3(E_k)$ but for $\text{III}(E_k)[3]$ as well.

Our methods involve the connection between ϕ -Selmer groups and binary cubic forms. This connection was first studied by Selmer himself [34], and later by Cassels [13]. The rational theory was thoroughly treated by Satgé [31], where he studied the ϕ -Selmer group from the point of view of cubic fields. Later, Liverance [24] studied these Selmer groups using the classical invariant theory of binary cubics.

The boundedness of the average size of the ϕ - and $\hat{\phi}$ -Selmer groups—and thus the rank—of the elliptic curves $E_k: y^2 = x^3 + k$, $k \in \mathbb{Z}$, was first demonstrated by Fouvry [20], who used Satgé’s results to reduce the boundedness of the average rank to the theorem of Davenport and Heilbronn [16] on the boundedness of the average size of the 3-torsion subgroups of the class groups of quadratic fields. Fouvry’s method thus implicitly used binary cubic forms, as Davenport and Heilbronn’s proof on the mean size of the 3-torsion of the class groups of quadratic fields used a count of integral binary cubic forms to count cubic fields of bounded discriminant, together with class field theory to transform this count to one about 3-torsion elements in class groups of quadratic fields.

Recently, in [11] a more direct proof of Davenport and Heilbronn’s theorem on 3-torsion elements in class groups in quadratic fields was given. This proof used a count of *integer-matrix* binary cubic forms $ax^3 + 3bx^2y + 3cxy^2 + dy^3$ ($a, b, c, d \in \mathbb{Z}$), together with a direct correspondence between 3-torsion ideal classes in quadratic fields and integer-matrix binary cubic forms as studied in [19, 1] (see Section 2 for a description of this correspondence over any Dedekind domain). This suggested to us that perhaps a natural discriminant-preserving map from ϕ -Selmer groups $\text{Sel}_{\phi}(E_k)$ to orbits of integer-matrix binary cubic forms could be constructed. This construction is the key to our method, and is the subject of Sections 3–6.

With this direct map in hand, the problem of determining the average size of the ϕ -Selmer group is reduced to counting the appropriate set of integer-matrix binary cubic forms of bounded discriminant. For this, the counting method of Davenport [15] and a suitable adaptation of the sieve methods of [16] and [6] may be applied to obtain the optimal upper and lower bounds, and this is carried out in Section 7. This sieve renders the final answer as a product of local densities, and we prove that the density at the p -adic place for a given elliptic curve $E_k: y^2 = x^3 + k$ is given precisely by the local Selmer ratio $c_p(\phi)$ as defined by (5). For k varying over an acceptable set S , this yields Theorem 2 (see §7.3).

We may evaluate the local Selmer ratios by relating them to Tamagawa numbers and using Tate’s algorithm (Proposition 28). Setting $S = \mathbb{Z}$ then yields Theorem 1, while setting $S = T_m$, and applying a formula of Cassels [12] on the global Selmer ratio, yields Theorem 5 (see §§7.3–7.4). Finally, the results on average rank and on positive proportions of rank 0 and 3-Selmer rank 1 curves, as in Theorems 6–8 and Corollary 9, are deduced in Section 8.

We note that the connection between binary cubic forms and Selmer groups also has an interpretation in terms of Lie groups. In the language of Vinberg theory, the representation of SL_2 on binary

cubic forms arises from a $\mathbb{Z}/3\mathbb{Z}$ -grading of the Lie algebra of G_2 . This paper is one example of recent work connecting representations arising from Vinberg's theory to Selmer groups of Jacobians of algebraic curves. In this context, we believe that our theorems above give the first results for the exact and finite average size of a Selmer group associated to an isogeny that is not the multiplication-by- n isogeny. For results and conjectures about the latter see, e.g., [3, 37, 38, 6, 22, 25, 27, 29, 35]. A beautiful treatment of the connection between algebraic curves and Vinberg theory, using versal deformations, has been given by Thorne [37].

One may ask about the ϕ -Selmer group in families of quadratic twists $E_{km^3}: y^2 = x^3 + km^3$, for fixed k and m varying, as well as in families of cubic twists $E_{km^2}: y^2 = x^3 + km^2$. These are very sparse subsets of all curves E_k , and so our results in Theorem 2 do not apply. Nevertheless, in a forthcoming paper [5], it is shown that the analogue of Theorem 2 continues to hold even in these families of quadratic twists; in fact, the formula (6) for the average size of the ϕ -Selmer group holds more generally in any family of quadratic twists of an elliptic curve (not necessarily of j -invariant 0) having a rational 3-isogeny, leading to a proof of the boundedness of the average rank and a positive proportion having rank 0 also in such quadratic twist families. We note that the analogue of the formula in Theorem 2 holds also for the above cubic twist families [4], although in most such cases the product (6) does not converge and we find that the average ϕ -Selmer group sizes for curves in these families are unbounded, *except* in the case where $k = 1$.

Finally, in another forthcoming work, Kriz and Li (private communication) give a very different proof that a positive proportion of the curves $y^2 = x^3 + k$ have rank 0 (resp. 1); they show that around 10% of these curves have rank 0, and a similar proportion have rank 1. Their p -adic methods are completely different from ours, and in particular, their rank 1 results are unconditional and do not rely on the recent work of Skinner, W. Zhang, and others on converse theorems. It will be an interesting future question to see if there is a way to increase the rank proportions by combining our method with theirs.

2 Integer-matrix binary cubic forms over a Dedekind domain

Let $V^*(\mathbb{Z}) = \text{Sym}^3\mathbb{Z}^2$ be the lattice of integer-coefficient binary cubic forms, i.e., forms $f(u, v) = au^3 + bv^2y + cuv^2 + dv^3$ with $a, b, c, d \in \mathbb{Z}$. To ease notation, we write $f(u, v) = [a, b, c, d]$. The group $\text{GL}_2(\mathbb{Z})$ acts naturally on $V^*(\mathbb{Z})$ by linear change of variable, and the *discriminant*

$$\text{Disc}(f) = b^2c^2 - 4ac^3 - 4b^3d - 27a^2d^2 + 18abcd$$

is $\text{SL}_2(\mathbb{Z})$ -invariant.

In this paper, the more fundamental object will be the dual lattice $V(\mathbb{Z}) = \text{Sym}_3\mathbb{Z}^2$ consisting of *integer-matrix* binary cubic forms, i.e., forms $f(u, v) = [a, 3b, 3c, d]$ with $a, b, c, d \in \mathbb{Z}$. The lattice $V(\mathbb{Z})$ has its own (*reduced*) *discriminant*

$$\text{disc}(f) = -\frac{1}{27}\text{Disc}(f).$$

We note that the action of $\text{GL}_2(F)$ on the space $V(F) := V(\mathbb{Z}) \otimes F$ of binary cubic forms with coefficients in F satisfies

$$\text{disc}(g \cdot f) = \det(g)^6 \text{disc}(f)$$

for all $g \in \text{GL}_2(F)$ and $f \in V(F)$. For any ring R and $d \in R$, we write $V(R)_d$ for the set of $f \in V(R) := V(\mathbb{Z}) \otimes R$ with $\text{disc}(f) = d$.

In this section, we classify the orbits of $V(D)$, under the action of $\mathrm{SL}_2(D)$, for an arbitrary Dedekind domain D . We assume that D does not have characteristic 2 or 3. In later sections we will apply our classification to the case where D is a field, \mathbb{Z}_p , or \mathbb{Z} . Our result is a generalization of [1, Theorem 13], and is proved in the same way.

Theorem 12. *Let D be a Dedekind domain of characteristic not 2 or 3, and let $k \in D$ be any nonzero element. Let F be the fraction field of D , and let $S := D[z]/(z^2 - k)$ and $K := F[z]/(z^2 - k)$. Then there is a bijection between the orbits of $\mathrm{SL}_2(D)$ on $V(D)_{4k}$ and equivalence classes of triples (I, δ, s) , where I is a fractional S -ideal, $\delta \in K^*$, and $s \in F^*$, satisfying the relations $I^3 \subset \delta S$, $N(I)$ is the principal fractional ideal sD in F , and $N(\delta) = s^3$ in F^* . Two triples (I, δ, s) and (I', δ', s') are equivalent if there exists $\kappa \in K^*$ such that $I' = \kappa I$, $\delta' = \kappa^3 \delta$, and $s' = N(\kappa)s$. Under this correspondence, the stabilizer in $\mathrm{SL}_2(D)$ of $f \in V(D)_{4k}$ is isomorphic to $S(I)^*[3]_{N=1}$, where $S(I)$ is the ring of endomorphisms of I .*

Remark 13. We call triples (I, δ, s) satisfying the relations above *valid triples*.

Proof. We describe here the explicit bijection, but refer readers to [1] for the details of the proof. First, note that $S = D + D\tau$, with τ the image of x in S . Given a valid triple (I, δ, s) , since $N(I)$ is the principal D -ideal sD , the projective D -module I of rank 2 is free, and so we may write $I = D\alpha + D\beta$ for some $\alpha, \beta \in I$. Because $I^3 \subset \delta S$, we have

$$\begin{aligned} \alpha^3 &= \delta(e_0 + \tau a) \\ \alpha^2\beta &= \delta(e_1 + \tau b) \\ \alpha\beta^2 &= \delta(e_2 + \tau c) \\ \beta^3 &= \delta(e_3 + \tau d), \end{aligned} \tag{8}$$

for some integers $a, b, c, d, e_i \in \mathbb{Z}$. Then corresponding to (I, δ, s) is the binary cubic form $f = [a, 3b, 3c, d]$, which has discriminant $\mathrm{disc}(f) = 4k$. In more coordinate-free terms, f is the symmetric trilinear form

$$\frac{1}{\delta} \times : I \times I \times I \rightarrow S/D \cong D\tau. \tag{9}$$

We obtain an $\mathrm{SL}_2(D)$ -orbit of symmetric trilinear forms (integer-matrix binary cubic forms) over D by taking the symmetric $2 \times 2 \times 2$ matrix representation of this form with respect to any ordered basis $\langle \alpha, \beta \rangle$ of I that gives rise to the basis element $s(1 \wedge \tau)$ of the top exterior power of I over D . This normalization deals with the difference between $\mathrm{SL}_2(D)$ - and $\mathrm{GL}_2(D)$ -orbits. The stabilizer statement follows because elements in $S(I)^*[3]_{N=1}$ are precisely the elements of $K_{N=1}^*$ that preserve the map (9). \square

When D is a field, so that $D = F$, the previous result simplifies quite a bit. Let us write $(K^*/K^{*3})_{N=1}$ to denote the kernel of the norm map $K^*/K^{*3} \rightarrow F^*/F^{*3}$, and $(\mathrm{Res}_F^K \mu_3)_{N=1}$ for the kernel of the norm map $\mathrm{Res}_F^K \mu_3 \rightarrow \mu_3$.

Corollary 14. *There is a bijection between the set of $\mathrm{SL}_2(F)$ -orbits on $V(F)_{4k}$ and $(K^*/K^{*3})_{N=1}$. Moreover, the stabilizer of any $f \in V(F)_{4k}$ in $\mathrm{SL}_2(F)$ is isomorphic to $(\mathrm{Res}_F^K \mu_3)_{N=1}$.*

Proof. Both statements follow from taking $D = F$ in the previous theorem. The bijection sends $\delta \in (K^*/K^{*3})_{N=1}$ to the orbit of binary cubic forms corresponding to the triple (K, δ, s) , where s is any choice of cube root of $N(\delta)$. \square

Remark 15. Explicitly, if $\delta = a + b\tau \in K^*$, then the corresponding cubic form is $f = [ak, 3bk, 3a, b]$.

Another case of interest is when $D = \mathbb{Z}_p$ and S is the maximal order in K :

Proposition 16. *Let p be a prime and assume $S = \mathbb{Z}_p[z]/(z^2 - k)$ is the maximal order in $K = \mathbb{Q}_p[z]/(z^2 - k)$. Then the set of $\mathrm{SL}_2(\mathbb{Z}_p)$ -orbits on $V(\mathbb{Z}_p)_{4k}$ is in bijection with the unit subgroup $(S^*/S^{*3})_{N=1} \subset (K^*/K^{*3})_{N=1}$. Every rational $\mathrm{SL}_2(\mathbb{Q}_p)$ -orbit of discriminant $4k$ whose class lies in this unit subgroup contains a unique integral $\mathrm{SL}_2(\mathbb{Z}_p)$ -orbit. The stabilizer in $\mathrm{SL}_2(\mathbb{Q}_p)$ of an element in $V(\mathbb{Z}_p)_{4k}$ is equal to its stabilizer in $\mathrm{SL}_2(\mathbb{Z}_p)$.*

Proof. Indeed, every S -ideal I is principal in this case. From the \mathbb{Z}_p -version of Theorem 12, it follows that any integral orbit corresponds to a triple (S, δ, s) with $\delta \in S^*$. The stabilizer statement follows because $K^*[3]_{N=1} = S^*[3]_{N=1}$ when S is the maximal order in K . \square

Finally, we will need the following result on stabilizers of integer-matrix binary cubic forms over \mathbb{Z} . Let $G = \mathrm{SL}_2$ and for any ring R and $f \in V(R)$, write $\mathrm{Aut}_R(f)$ for $\mathrm{Stab}_{G(R)}(f)$.

Proposition 17. *Suppose $f \in V(\mathbb{Z})_{4k}$ is an integer-matrix binary cubic form of discriminant $4k$. Then*

$$|\mathrm{Aut}_{\mathbb{Q}}(f)| \sum_{f' \in O(f)} |\mathrm{Aut}_{\mathbb{Z}}(f')|^{-1} = \prod_p |\mathrm{Aut}_{\mathbb{Q}_p}(f)| \sum_{f'_p \in O_p(f)} |\mathrm{Aut}_{\mathbb{Z}_p}(f'_p)|^{-1},$$

where $O(f)$ is a set of representatives for the $G(\mathbb{Z})$ -orbits on $V(\mathbb{Z})_{4k}$ contained in the $G(\mathbb{Q})$ -orbit of f , and similarly $O_p(f)$ is a set of representatives for the $G(\mathbb{Z}_p)$ -orbits on $V(\mathbb{Z}_p)_{4k}$ contained in the $G(\mathbb{Q}_p)$ -orbit of f .

Proof. We follow the proof of [3, Proposition 8.9]. Let $K = \mathbb{Q}[z]/(z^2 - k)$, and fix a representative $\delta \in K^*$ of the class in $(K^*/K^{*3})_{N=1}$ corresponding to f . Let $m(\delta)$ be the number of ideals I of $S = \mathbb{Z}[z]/(z^2 - k)$ satisfying $I^3 \subset \delta S$ and the ideal equality $N(I)^3 = N(\delta)$. Similarly, let $m_p(\delta)$ be the number of ideals I_p of $S_p := S \otimes \mathbb{Z}_p$ with $I_p^3 \subset \delta S_p$ and $N(I_p)^3 = N(\delta)$. Note that for all but finitely many p , S_p is the maximal order and δ is a unit, so $m_p(\delta) = 1$ for all but finitely many p .

Since a lattice is determined by its local completions, and since a collection of local ideals produces a global ideal with the desired properties, we have

$$m(\delta) = \prod_p m_p(\delta). \tag{10}$$

Let s be any cube root of $N(\delta)$. Then the triple (I, δ, s) is valid and hence corresponds to an integer-matrix binary cubic form $f_I \in V(\mathbb{Z})_{4k}$ mapping to the rational orbit of δ . If s' is another choice of cube root, then $s' = s\zeta_3$ and (I, δ, s) is equivalent to (I, δ, s') . A triple (I, α, s) is in the same integral orbit as (cI, α, s) , for some $c \in K^\times$, exactly when $c^3 = 1$ and $N(c) = 1$. On the other hand, the ideals I and cI are equal if c is a unit in the ring of endomorphisms $S(I)$ of I . Thus, the number of distinct ideals giving the same integral orbit is the size of the group $K^*[3]_{N=1}/S(I)^*[3]_{N=1}$.

We have $K^*[3]_{N=1} \cong \mathrm{Aut}_{\mathbb{Q}}(f)$ and $S(I)^*[3]_{N=1} \cong \mathrm{Aut}_{\mathbb{Z}}(f_I)$ by Theorem 12. Thus, the number of distinct ideals associated to the integral orbit of f_I is $|\mathrm{Aut}_{\mathbb{Q}}(f)|/|\mathrm{Aut}_{\mathbb{Z}}(f_I)|$. We conclude that

$$m(\delta) = \sum_{f_I \in O(f)} |\mathrm{Aut}_{\mathbb{Q}}(f)|/|\mathrm{Aut}_{\mathbb{Z}}(f_I)|.$$

The same reasoning implies an analogous formula for $m_p(\delta)$, with global stabilizers replaced by local stabilizers. The proposition now follows from (10). \square

3 The elliptic curves E_k and orbits of binary cubic forms over a field

Let F be a field of characteristic not 2 or 3, and let $k \in F$ be non-zero. Recall that the elliptic curve

$$E_k: y^2 = x^3 + k \tag{11}$$

admits a 3-isogeny $\phi: E_k \rightarrow E_{-27k}$ defined over F , and a dual 3-isogeny $\hat{\phi}: E_{-27k} \rightarrow E_k$.

In this section, we describe the connection between the isogenies ϕ and $\hat{\phi}$ and binary cubic forms (i.e., symmetric trilinear forms) over F .

3.1 Galois cohomology of the 3-isogeny kernel and field arithmetic

Important in the study of ϕ - and $\hat{\phi}$ -descents on the curves E_k and E_{-27k} is the pair of “mirror” quadratic étale F -algebras

$$K = F[z]/(z^2 - k) \quad \text{and} \quad \hat{K} = F[z]/(x^2 + 27k).$$

The following result connects the arithmetic of the elliptic curves E_{-27k} and E_k to the arithmetic of K and \hat{K} :

Proposition 18. *There is an isomorphism of group schemes*

$$E_{-27k}[\hat{\phi}] \cong \ker(\text{Res}_F^K \mu_3 \rightarrow \mu_3),$$

and an induced isomorphism

$$H^1(G_F, E_{-27k}[\hat{\phi}]) \cong (K^*/K^{*3})_{N=1},$$

where $(K^*/K^{*3})_{N=1}$ denotes the kernel of the norm $N: K^*/K^{*3} \rightarrow F^*/F^{*3}$.

Proof. Duality gives a non-degenerate pairing

$$\langle \cdot, \cdot \rangle: E_{-27k}[\hat{\phi}] \times E_k[\phi] \rightarrow \mu_3.$$

Since $E_k[\phi](\bar{F})$ becomes a trivial Galois module when restricted to G_K , this induces an injective homomorphism of group schemes $\iota: E_{-27k}[\hat{\phi}] \rightarrow \text{Res}_F^K \mu_3$, given on points by

$$P \mapsto (\langle P, Q_1 \rangle, \langle P, Q_2 \rangle),$$

where Q_1 and Q_2 are the non-trivial points of $E_k[\phi]$. The image of ι is precisely the kernel of the norm map $\text{Res}_F^K \mu_3 \rightarrow \mu_3$, giving the desired isomorphism.

From Kummer theory and the long exact sequence attached to

$$0 \rightarrow E_{-27k}[\hat{\phi}] \rightarrow \text{Res}_F^K \mu_3 \rightarrow \mu_3 \rightarrow 0,$$

we obtain the isomorphism $H^1(G_F, E_{-27k}[\hat{\phi}]) \cong (K^*/K^{*3})_{N=1}$. □

Remark 19. The sequence $0 \rightarrow E_{-27k}[\hat{\phi}] \rightarrow E_{-27k} \rightarrow E_k \rightarrow 0$ induces a Kummer map

$$\partial: E_k(F) \rightarrow H^1(G_F, E_{-27k}[\hat{\phi}]) \cong (K^*/K^{*3})_{N=1},$$

which can be described explicitly as follows. If $(x, y) \notin E_k[\phi](F)$, then $\partial((x, y)) = y - \tau$, where τ is the image of x in $K = F[z]/(z^2 - k)$. If $P = (0, \pm\sqrt{k}) \in E_k[\phi]$, then $\partial(P) = \pm 1/2\tau$. See [12, §15].

Remark 20. Of course, the analogues of all these results hold also when $\hat{\phi}$ is replaced with ϕ and K is replaced with \hat{K} ; these analogues are obtained simply via the change of variable $k \mapsto -27k$.

3.2 Connection to binary cubic forms

We may now compare Corollary 14 and Proposition 18. This immediately yields the following canonical bijection:

Theorem 21. *There is a bijection between $H^1(G_F, E_{-27k}[\hat{\phi}])$ and the $\mathrm{SL}_2(F)$ -orbits on $V(F)_{4k}$. Moreover, the stabilizer in $\mathrm{SL}_2(F)$ of any $f \in V(F)_{4k}$ is isomorphic to $E_{-27k}[\hat{\phi}](F)$.*

Let $V(F)^{\mathrm{sol}}$ denote the set of binary cubic forms $f(x, y) \in V(F)$ that correspond under the bijection of Theorem 21 to classes in the image of the Kummer map $\partial: E_k(F) \rightarrow H^1(G_F, E_{-27k}[\hat{\phi}])$ for some $k \in F$. If $f \in V(F)^{\mathrm{sol}}$, then we also say that f is *soluble*. We write $V(F)_D^{\mathrm{sol}}$ for the elements of $V(F)^{\mathrm{sol}}$ having discriminant D .

Corollary 22. *There is a natural bijection between the $\mathrm{SL}_2(F)$ -orbits on $V(F)_{4k}^{\mathrm{sol}}$ and the elements of the group $E_k(F)/\hat{\phi}(E_{-27k}(F))$. Under this bijection, the identity element of $E_k(F)/\hat{\phi}(E_{-27k}(F))$ corresponds to the unique $\mathrm{SL}_2(F)$ -orbit of reducible binary cubic forms in $V(F)_{4k}^{\mathrm{sol}}$, namely the orbit of $f = [k, 0, 3, 0]$.*

Proof. The second part of the corollary follows from Remark 15, by taking $\delta = a = 1$ and $b = 0$. \square

The use of the term ‘soluble’ comes from the following fact:

Proposition 23. *A binary cubic form $f(x, y) \in V(F)_{4k}$ corresponds to an element δ in the image of $\partial: E_k(F) \rightarrow H^1(G_F, E_{-27k}[\hat{\phi}]) \cong (K/K^{*3})_{N=1}$ if and only if the curve $C_f: z^3 = f(x, y)$ in \mathbb{P}^2 has an F -rational point.*

Proof. Suppose $f \in V(F)_{4k}$, and let $\delta \in K^*$ be a representative element in $(K/K^{*3})_{N=1}$ corresponding to the $\mathrm{SL}_2(F)$ -orbit of f under the bijection of Corollary 14.

If δ represents the image of a point $(x, y) \in E_k(F)$ under ∂ , then, by Remarks 15 and 19, the form f is $\mathrm{SL}_2(F)$ -equivalent to $[1, 3y, 3k, yk]$, which represents a cube (namely, 1).

Conversely, if there exist $u, v, z \in F$ such that $z^3 = f(u, v)$, then by scaling u and v if necessary we may assume that $z = 1$. Therefore, we have by (8) or (9) that

$$(u + v\tau)^3 = \delta(g(u, v) + f(u, v)\tau) = \delta(g(u, v) + \tau) \quad (12)$$

where g is a binary cubic form over F , and so $g(u, v) \in F$. Writing $N(\delta) = s^3$, and then taking norms of both sides of (12), we obtain $N(u + v\tau)^3 = s^3(g(u, v)^2 - k)$. Thus the point $(x, y) = (N(u + v\tau)/s, g(u, v))$ lies on E_k , and this point (x, y) then maps to the class of δ in $(K/K^{*3})_{N=1}$ under ∂ . \square

Remark 24. Recall that a $\hat{\phi}$ -covering is a map of curves $C \rightarrow E_k$ over F which is a twist of $\hat{\phi}$. By descent, the group $H^1(G_F, E_{-27k}[\hat{\phi}])$ is in bijection with isomorphism classes of $\hat{\phi}$ -coverings. To construct the $\hat{\phi}$ -covering corresponding to $\delta \in (K/K^{*3})_{N=1}$, take any $s \in F^*$ such that $N(\delta) = s^3$, and let f be the corresponding binary cubic form over F under the bijection of Corollary 14. Then we take C_f to be the curve $z^3 = f(x, y)$ in \mathbb{P}^2 (whose Jacobian is easily computed to be E_k), and the $\hat{\phi}$ -covering map $C_f \rightarrow E_k$ corresponding to the class δ is given explicitly by

$$(u: v: z) \mapsto ((u^2 - kv^2)/s, g(u, v)),$$

where $g(u, v)$ is the cubic polynomial in the preceding proof.

Remark 25. One can also construct the $\hat{\phi}$ -covering using the invariant theory of binary cubics. If $f = [a, 3b, 3c, d] \in V(F)$, then the covariants of f are generated by the discriminant disc, the scaled Hessian

$$h(x, y) = \frac{1}{36}(f_{xx}f_{yy} - f_{xy}^2) = (ac - b^2)x^2 + (ad - bc)xy + (bd - c^2), \quad (13)$$

and the Jacobian derivative of f and h ,

$$g(x, y) = \frac{\partial(f, h)}{\partial(x, y)} = f_x h_y - f_y h_x, \quad (14)$$

which is a cubic polynomial in x, y whose coefficients are cubic polynomials in a, b, c, d . The cubic f and its covariants g, h are related by the syzygy¹

$$(g/3)^2 - \text{disc}(f)f^2 + 4h^3 = 0. \quad (15)$$

This gives us a degree-3 map from the genus-1 curve

$$C_f: z^3 = f(x, y)$$

to E_k with $k = \text{disc}/4$: divide both sides of (15) by $4z^6$ and solve for $(g/2z^3)^2$ to obtain

$$\left(\frac{1}{6} \frac{g}{z^3}\right)^2 = \left(-\frac{h}{z^2}\right)^3 + \frac{\text{disc}(f)}{4}. \quad (16)$$

Our curve E_{-27k} is the special case of C_f where $f(x, y) = kx^3 + 3xy^2$; we then see that $\text{disc}(f) = 4k$, and the map $(z, y/x) \mapsto (-h/z^2, g/6z^3)$ to E_k recovers our formula (3) for the 3-isogeny $\hat{\phi}$.

4 Soluble orbits over local fields

When F is a local field, we can give explicit formulas for the number of soluble $\text{SL}_2(F)$ -orbits of binary cubic forms of discriminant $4k$, i.e., the size of the group $E_k(F)/\hat{\phi}(E_{-27k}(F))$. Since $|E_{-27k}[\hat{\phi}](F)|$ is 3 or 1 depending on whether $-3k$ is a square in F , it is equivalent to give formulas for the (local) Selmer ratios

$$c(\hat{\phi}_k) = \frac{|\text{coker}(E_{-27k}(F) \rightarrow E_k(F))|}{|\text{ker}(E_{-27k}(F) \rightarrow E_k(F))|}.$$

We do this below for the local fields \mathbb{Q}_p, \mathbb{R} , and \mathbb{C} , though there are similar formulas for any finite extension of \mathbb{Q}_p and for equicharacteristic local fields such as $\mathbb{F}_p((t))$. If $F = \mathbb{Q}_p, \mathbb{R}$, or \mathbb{C} , we use the notation $c_p(\hat{\phi}_k)$, with $p \leq \infty$, to match with the introduction. We state the result for $c_p(\phi_k)$, the Selmer ratio of the original isogeny $\phi_k: E_k \rightarrow E_{-27k}$.

¹ These results are classical, at least for $F = \mathbb{C}$, and they go back at least to Hilbert [23, pp.68–69]; see also Schur's treatment [33, II, §8, Satz 2.24 on p. 77]. The syzygy (15) can be verified by direct computation, though it is easier to check it for one choice of f without repeated factors and then use the fact that $\text{SL}_2(\bar{F})$ acts transitively on such f . For example, $f = x^3 - y^3$ gives $\text{disc} = 1$, $h = -xy$ and $g = -3(x^3 + y^3)$, reducing the syzygy (15) to the identity $(x^3 + y^3)^2 - (x^3 - y^3)^2 = 4(xy)^3$.

4.1 Orbits over \mathbb{Q}_p

Suppose $F = \mathbb{Q}_p$. We first determine when E_k/\mathbb{Q}_p has good reduction.

Lemma 26. *Assume $k \in \mathbb{Z}_p$ is sixth-power-free. If $p > 3$, then E_k/\mathbb{Q}_p has good reduction if and only if $p \nmid k$. If $p = 3$, then E_k/\mathbb{Q}_3 has bad reduction. If $p = 2$, then E_k/\mathbb{Q}_2 has good reduction if and only if $k \equiv 16 \pmod{64}$.*

Proof. This follows from Tate's algorithm. If $k \equiv 16 \pmod{64}$, a model with good reduction at 2 is $y'^2 + y' = x'^3 + a_6$ where $k = 64a_6 + 16$ and $(x, y) = (4x', 8y' + 4)$. \square

Next, we express $c_p(\phi_k)$ in terms of the Tamagawa numbers of E_k and E_{-27k} (cf. [32, Lemma 3.8]).

Proposition 27. *If $k \in \mathbb{Z}_p$ is sixth-power-free, then*

$$c_p(\phi_k) = \frac{c_p(E_{-27k})}{c_p(E_k)} \times \begin{cases} 3 & \text{if } p = 3 \text{ and } 27 \mid k; \text{ and} \\ 1 & \text{otherwise,} \end{cases}$$

where $c_p(E) = |E(\mathbb{Q}_p)/E_0(\mathbb{Q}_p)|$ denotes the Tamagawa number of E .

Proof. Let ω_k and ω_{-27k} denote Néron differentials on E_k and E_{-27k} , respectively. Then for some $b, b' \in \mathbb{Q}$, we have $\omega = b \cdot \frac{dx}{y}$ and $\omega_{-27k} = b' \cdot \frac{dX}{Y}$ on $E_k: y^2 = x^3 + k$ and $E_{-27k}: Y^2 = X^3 - 27k$, respectively. The model $y^2 = x^3 + k$ is minimal except if $k \equiv 16 \pmod{64}$, in which case the model given in the proof of Lemma 26 is minimal. In either case, we compute that $b = b'$ when $27 \nmid k$ and $b' = 3b$ when $27 \mid k$. Since

$$\phi^* \left(b \cdot \frac{dX}{Y} \right) = \frac{b \cdot d \left(\frac{x^3 + 4k}{x^2} \right)}{\frac{y(x^3 - 8k)}{x^3}} = \frac{b \cdot \left(1 - \frac{8k}{x^3} \right) dx}{\frac{y(x^3 - 8k)}{x^3}} = b \cdot \frac{dx}{y},$$

we have $\phi^* \omega_{-27k} = a \omega_k$, where $a = 1$ if $27 \nmid k$ and $a = 3$ if $27 \mid k$.

We then compute

$$\int_{\hat{\phi}(E_k(\mathbb{Q}_p))} |\omega_{-27k}|_p = \frac{1}{|E_k[\phi](\mathbb{Q}_p)|} \cdot \int_{E_k(\mathbb{Q}_p)} |\phi^* \omega_{-27k}|_p = \frac{1}{|E_k[\phi](\mathbb{Q}_p)|} \cdot |a|_p \cdot \int_{E_k(\mathbb{Q}_p)} |\omega_k|_p.$$

Therefore,

$$c_p(\phi_k) = \frac{|(E_{-27k}(\mathbb{Q}_p)/\phi(E_k(\mathbb{Q}_p)))|}{|E_k[\phi](\mathbb{Q}_p)|} = \frac{\int_{E_{-27k}(\mathbb{Q}_p)} |\omega_{-27k}|_p}{|a|_p \int_{E_k(\mathbb{Q}_p)} |\omega_k|_p} = \frac{1}{|a|_p} \cdot \frac{c_p(E_{-27k})}{c_p(E_k)}$$

as desired. \square

We use Tate's algorithm [36] to compute the ratios of Tamagawa numbers in Proposition 27, and hence the local Selmer ratios $c_p(\phi_k)$. In particular, we find that the $c_p(\phi_k)$ are determined by congruence conditions on k :

Proposition 28. *Let $k \in \mathbb{Z}$ be sixth-power-free, and let χ_K denote the quadratic character attached to $K = \mathbb{Q}(\sqrt{k})$. If $p \neq 3$, then*

$$c_p(\phi_k) = \begin{cases} 3^{-\chi_K(p)} & \text{if } p \equiv 2 \pmod{3} \text{ and } v_p(4k) \in \{2, 4\}; \\ 1 & \text{otherwise.} \end{cases}$$

If $p = 3$, write $k = 3^{v_3(k)}k_3$. Then

$$c_p(\phi_k) = \begin{cases} 9 & \text{if } v_3(k) = 5, \text{ and } k_3 \equiv 2 \pmod{3}; \\ 3 & \text{if } v_3(k) = 0, \text{ and } k_3 \equiv 5 \text{ or } 7 \pmod{9}; \text{ or} \\ & v_3(k) \in \{1, 4\}, \text{ and } k_3 \equiv 2 \pmod{3}; \text{ or} \\ & v_3(k) = 3, \text{ and } k_3 \not\equiv 2 \text{ or } 4 \pmod{9}; \text{ or} \\ & v_3(k) = 5, \text{ and } k_3 \equiv 1 \pmod{3}; \\ \frac{1}{3} & \text{if } v_3(k) = 2, \text{ and } k_3 \equiv 1 \pmod{3}; \\ 1 & \text{otherwise.} \end{cases} \quad (17)$$

From Proposition 28, we may easily deduce explicit formulas for the number of soluble $\mathrm{SL}_2(\mathbb{Q}_p)$ -orbits on $V(\mathbb{Q}_p)_{4k}$ (see also Corollary 31 below). One may go further and describe the soluble classes as a subgroup of $(K^*/K^{*3})_{N=1}$. We describe this below in the cases where $p \neq 3$.

Proposition 29. *If $k \in \mathbb{Z}_p$ and E_k has good reduction, then the Kummer map induces an isomorphism*

$$E_k(\mathbb{Q}_p)/\hat{\phi}(E_{-27k}(\mathbb{Q}_p)) \cong (S_0^*/S_0^{*3})_{N=1},$$

where S_0 is the ring of integers of $K = \mathbb{Q}_p[x]/(x^2 - k)$.

Proof. See [14, Lemma 4.1] or [21, Lemma 6]. □

Proposition 30. *Assume $p \neq 3$ and that E_k/\mathbb{Q}_p has bad reduction. Then the natural map*

$$j: E_k[\phi](\mathbb{Q}_p) \rightarrow E_k(\mathbb{Q}_p)/\hat{\phi}(E_{-27k}(\mathbb{Q}_p))$$

is an isomorphism.

Proof. We first claim that $E_k[3](\mathbb{Q}_p) = E_k[\phi](\mathbb{Q}_p)$. Indeed, the six other 3-torsion points on $E_k(\bar{\mathbb{Q}}_p)$ are $(\sqrt[3]{-4k}, \sqrt{-3k})$, for the six possible choices of roots in this expression. It follows from the bad reduction of E_k and Lemma 26 that these points are defined over ramified extensions of \mathbb{Q}_p , so they do not lie in $E_k(\mathbb{Q}_p)$, proving the claim.

Next we show that j is injective. Indeed, if $P \in E_k[\phi](\mathbb{Q}_p)$ and $Q \in E_{-27k}(\mathbb{Q}_p)$ satisfy $\hat{\phi}(Q) = P$, then $Q \in E_{-27k}[3](\mathbb{Q}_p) = E_{-27k}[\hat{\phi}](\mathbb{Q}_p)$, where the equality follows from the above claim applied to E_{-27k} . Thus, $P = 0$ and j is injective.

Finally, it suffices to show that $|E_k(\mathbb{Q}_p)/\hat{\phi}(E_{-27k}(\mathbb{Q}_p))| \leq |E_k[\phi](\mathbb{Q}_p)|$. Note that the group $E_k(\mathbb{Q}_p)/3E_k(\mathbb{Q}_p)$ surjects onto $E_k(\mathbb{Q}_p)/\hat{\phi}(E_{-27k}(\mathbb{Q}_p))$. Since $p \neq 3$, an argument using the formal group [3, Lemma 12.3] shows that $E_k(\mathbb{Q}_p)/3E_k(\mathbb{Q}_p)$ has size $|E_k[3](\mathbb{Q}_p)|$, which is equal to $|E_k[\phi](\mathbb{Q}_p)|$, again by our claim above. Therefore, we indeed have

$$|E_k(\mathbb{Q}_p)/\hat{\phi}(E_{-27k}(\mathbb{Q}_p))| \leq |E_k[3](\mathbb{Q}_p)| = |E_k[\phi](\mathbb{Q}_p)|,$$

as desired. □

Corollary 31. *If $p \neq 3$, then the number of soluble orbits in $V(\mathbb{Q}_p)$ of discriminant $4k$ is equal to*

$$\begin{cases} |E_{-27k}[\hat{\phi}](\mathbb{Q}_p)| & \text{if } E_k/\mathbb{Q}_p \text{ has good reduction,} \\ |E_k[\phi](\mathbb{Q}_p)| & \text{if } E_k/\mathbb{Q}_p \text{ has bad reduction.} \end{cases}$$

Remark 32. At the start of the proof of Proposition 30, we noted that $E_k[3]$ consists of the three points of $E_k[\phi]$ together with the six points (x, y) with $x^3 = -4k$ and $y^2 = -3k$. If such a point is rational then $(k, x, y) = (-432m^6, 12m^2, 36m^3)$ for some nonzero m . In this case our elliptic curve $E_k \cong E_{-432}$ is isomorphic with the Fermat cubic $X^3 + Y^3 = Z^3$ (which is also isomorphic with the modular curve $X_0(27)$), and the isogenous curve $E_{27k} \cong E_{16}$ is isomorphic with $XY(X + Y) = Z^3$. These curves have good reduction at primes other than 3; in particular, they satisfy the condition of Lemma 26 for good reduction at 2, with minimal models $y'^2 + y' = x'^3 - 7$ and $y'^2 + y' = x'^3$ respectively.

4.2 Orbits over \mathbb{R} or \mathbb{C}

If $F = \mathbb{R}$ or \mathbb{C} , then every binary cubic form $f \in V(F)$ is reducible, so by Corollary 22, there is a unique $\mathrm{SL}_2(F)$ -orbit of binary cubic forms of discriminant $4k$. This agrees with the fact that $E_k(F)/\hat{\phi}(E_{-27k}(F))$ is always trivial. We conclude that $c_\infty(\phi_k) = 1/3$ if $F = \mathbb{C}$, and if $F = \mathbb{R}$, we have:

$$c_\infty(\phi_k) = \begin{cases} \frac{1}{3} & \text{if } k > 0; \\ 1 & \text{if } k < 0. \end{cases}$$

5 ϕ -Selmer groups and locally soluble orbits over a global field

We let F now be a global field of characteristic not 2 or 3. If $\varphi: A \rightarrow A'$ is an isogeny of abelian varieties over F , then the φ -Selmer group

$$\mathrm{Sel}_\varphi(A) \subset H^1(G_F, A[\varphi])$$

is the subgroup consisting of classes that are locally in the image of the Kummer map

$$\partial_v: A'(F_v) \longrightarrow H^1(G_{F_v}, A[\varphi])$$

for every place v of F . Equivalently, these are the classes locally in the kernel of the map

$$H^1(G_{F_v}, A[\varphi]) \rightarrow H^1(G_{F_v}, A)[\varphi],$$

for every place v of F , i.e., the classes corresponding to ϕ -coverings having a rational point over F_v for every place v .

Now let $V(F)^{\mathrm{loc. sol.}} \subset V(F)$ denote the set of *locally soluble* binary cubic forms, i.e. those $f(u, v) \in V(F)$ such that the equation $z^3 = f(u, v)$ has a non-zero solution over F_v for every place v of F . Then the following result follows immediately from Corollary 22.

Theorem 33. *Let $k \in F^*$. Then there is a bijection between the $\mathrm{SL}_2(F)$ -orbits on $V(F)^{\mathrm{loc. sol.}}$ having discriminant $4k$ and the elements of $\mathrm{Sel}_\phi(E_{-27k})$ corresponding to the isogeny $\hat{\phi}_k: E_{-27k} \rightarrow E_k$. Under this bijection, the identity element of $\mathrm{Sel}_\phi(E_{-27k})$ corresponds to the unique $\mathrm{SL}_2(F)$ -orbit of reducible binary cubic forms of discriminant $4k$, namely the orbit of $f(x, y) = kx^3 + 3xy^2$. Moreover, the stabilizer in $\mathrm{SL}_2(F)$ of any $f \in V(F)_{4k}^{\mathrm{loc. sol.}}$ is isomorphic to $E_{-27k}[\hat{\phi}](F)$.*

6 Existence of integral orbits

We now specialize to the case where $F = \mathbb{Q}$. The main goal of this section is to prove the following theorem.

Theorem 34. *Let $k \in 3\mathbb{Z}$. Then every locally soluble orbit for the action of $\mathrm{SL}_2(\mathbb{Q})$ on $V(\mathbb{Q})_{4k}$ has an integral representative, i.e., contains an element of $V(\mathbb{Z})_{4k}$.*

Proof. By Theorem 12 applied to the cases $D = \mathbb{Z}$ and $D = \mathbb{Z}_p$, it suffices to find a representative in $V(\mathbb{Z}_p)_{4k}$ for every soluble orbit of cubic forms $f \in V(\mathbb{Q}_p)_{4k}$, for each prime p . Indeed, such integral representatives would correspond to valid triples over \mathbb{Z}_p , which together determine a valid triple over \mathbb{Z} .

Let $S = \mathbb{Z}_p[z]/(z^2 - k)$ and $K = \mathbb{Q}_p[z]/(z^2 - k)$ as before, and assume for now that $p \neq 3$. The binary cubic form $f \in V(\mathbb{Q}_p)_{4k}^{\mathrm{sol}}$ corresponds to an element $\delta \in (K^*/K^{*3})_{N=1}$ in the image of the Kummer map. Let $P = (x, y) \in E_k(\mathbb{Q}_p)$ be a rational point mapping to δ . By Remark 19, we have $\delta = y - \tau$, where τ is the image of z in $K = \mathbb{Q}_p[z]/(z^2 - k)$. If x and y have negative valuation, then P lies in the subgroup $E_{k,1}(\mathbb{Q}_p) \subset E_k(\mathbb{Q}_p)$ isomorphic to the formal group of E_k . Since the formal group is pro- p and $p \neq 3$, it follows that there is a point $Q \in E_{k,1}(\mathbb{Q}_p)$ such that $3Q = P$; hence P is in $\hat{\phi}(E_{-27k}(\mathbb{Q}_p))$ and $\delta = \partial(P) = 1$. Thus, the valid triple $(S, 1, 1)$ corresponds to an integral representative in the orbit of f under the bijection of Theorem 12.

We may thus assume that x and y are in \mathbb{Z}_p . Define $I = \mathbb{Z}_p x + \mathbb{Z}_p \delta \subset K$. Then

$$\tau I = \mathbb{Z}_p x \tau + \mathbb{Z}_p \delta \tau = \mathbb{Z}_p (xy - x\delta) + \mathbb{Z}_p (x^3 - y\delta) \subset I,$$

and therefore I is an S -ideal. Furthermore, we have $N(I) = x\mathbb{Z}_p$, and $N(\delta) = N(y - \tau) = y^2 - k = x^3$. Finally, we note that the elements

$$\begin{aligned} \delta^{-1}(x^3) &= y + \tau \\ \delta^{-1}(x^2\delta) &= x^2 \\ \delta^{-1}(x\delta^2) &= xy - x\tau \\ \delta^{-1}(\delta^3) &= y^2 + k - 2y\tau, \end{aligned}$$

are each contained in $S = \mathbb{Z}_p + \mathbb{Z}_p \tau$, implying that $I^3 \subset \delta S$. Thus (I, δ, x) is a valid triple for S in the sense of Theorem 12 and Remark 13, and yields the desired integral representative in $V(\mathbb{Z}_p)_{4k}$.

Remark 35. One can even *characterize* the integral classes in terms of solubility conditions, at least if $p \neq 3$. Specifically, if $k \in \mathbb{Z}_p$ is sixth-power-free, then an $\mathrm{SL}_2(\mathbb{Q}_p)$ -orbit of forms $f \in V(\mathbb{Q}_p)_{4k}$ contains an integral form $f_0 \in V(\mathbb{Z}_p)_{4k}$ if and only if f is soluble over $\mathbb{Q}_p^{\mathrm{ur}}$, the maximal unramified extension of \mathbb{Q}_p .

Before considering the case $p = 3$, we state a general lemma. For any prime p , and for any $i \geq 0$, let S_i be the order of index p^i in the maximal order S_0 of $K = \mathbb{Q}_p[z]/(z^2 - k)$.

Lemma 36. *If (I, δ, s) is a valid triple for the ring S_i , with $i \geq 1$, then there exists an S_{i+1} -ideal J such that (J, δ, s) is a valid triple for the ring S_{i+1} . If $p = 3$, then this is true even if $i = 0$.*

Proof. This follows from the explicit bijection given in the proof of Theorem 12. If $f = [a, 3b, 3c, d]$ is the binary cubic form corresponding to the valid triple (I, δ, s) , then f has a root over \mathbb{F}_p , since

$$\mathrm{Disc}(f) = -27\mathrm{disc}(f) = -27 \cdot \mathrm{Disc}(S_i)$$

is divisible by p . So via a change of variables we may assume that $p \mid d$. If $\alpha\mathbb{Z}_p + \beta\mathbb{Z}_p$ is the corresponding basis of I , then $[p^2a, 3pb, 3c, d/p]$ is an integer-matrix binary cubic form corresponding to the triple $(p\alpha\mathbb{Z}_p + \beta\mathbb{Z}_p, \delta, s)$ for the ring S_{i+1} . If $p = 3$, then $\text{Disc}(f) = -27\text{disc}(f)$ is divisible by p , even if $i = 0$. \square

Now let $p = 3$ and let $f \in V(\mathbb{Q}_3)_{4k}^{\text{sol}}$ correspond to $\delta \in (K^*/K^{*3})_{N=1}$, with K and S as before. If the class of δ lies in the unit subgroup $(S_0^*/S_0^{*3})_{N=1}$, then (S_0, δ, s) is a valid triple for the ring S_0 , and so by Lemma 36, there exists J such that (J, δ, s) is valid for the ring S . This triple corresponds to the desired $f \in V(\mathbb{Z}_3)_{4k}$ in the orbit corresponding to δ .

If δ is not represented by a unit, then we must be in the case $K = \mathbb{Q}_3 \times \mathbb{Q}_3$, as this is the only étale \mathbb{Q}_3 -algebra where there exist δ 's not represented by units. Since $k \in 3\mathbb{Z}$, we must have $S = S_i$ for some $i \geq 1$. We may choose δ to be of the form $(3\pi, 3u)$, for some uniformizer $\pi \in 3\mathbb{Z}_3$ and unit $u \in \mathbb{Z}_3^*$. Then the triple $(3S_0, \delta, s)$ is valid for the ring S_1 , so by Lemma 36, there exist valid triples (J, δ, s) for S_i , for all $i \geq 1$, and in particular for S . This gives the desired $f \in V(\mathbb{Z}_3)_{4k}$ in the orbit corresponding to δ , and completes the proof of Theorem 34. \square

7 The average size of $\text{Sel}_\phi(E_k)$

In this section, for acceptable subsets $S \subset \mathbb{Z}$, we asymptotically count the number of $\text{SL}_2(\mathbb{Q})$ -classes of locally soluble integer-matrix binary cubic forms $f \in V(\mathbb{Z})$ of discriminant $3^6 4k$ for $|k| < X$, $k \in S$, as $X \rightarrow \infty$. By the work of Section 6, this will then allow us to deduce Theorems 1, 2, and 5.

7.1 The asymptotic number of binary cubic forms of bounded discriminant with weighted congruence conditions

Let $V(\mathbb{R})$ denote the vector space of binary cubic forms over \mathbb{R} . Let $V^{(0)}(\mathbb{R})$ denote the subset of elements in $V(\mathbb{R})$ having positive discriminant, and $V^{(1)}(\mathbb{R})$ the subset of elements having negative discriminant. We use $V^{(i)}(\mathbb{Z})$ to denote $V(\mathbb{Z}) \cap V^{(i)}(\mathbb{R})$.

Now let T be any set of integral binary cubic forms that is invariant under the action of $\text{SL}_2(\mathbb{Z})$. Let $N(T; X)$ denote the number of *irreducible* binary cubic forms contained in T , up to $\text{SL}_2(\mathbb{Z})$ -equivalence, having absolute discriminant at most X . Then we have the following theorem counting $\text{SL}_2(\mathbb{Z})$ -classes of integer-matrix binary cubic forms of bounded reduced discriminant, which easily follows from the work of Davenport [15] and Davenport–Heilbronn [16] (see [11, Theorem 19] for this deduction):

Theorem 37. *Let T be any set of integer-matrix binary cubic forms that is defined by finitely many congruence conditions modulo prime powers and that is invariant under the action of $\text{SL}_2(\mathbb{Z})$. For each prime p , let $\mu_p(T)$ denote the p -adic density of the p -adic closure of T in $V(\mathbb{Z}_p)$, where $V(\mathbb{Z}_p)$ is equipped with the usual additive p -adic measure normalized so that $\mu_p(V(\mathbb{Z}_p)) = 1$. Then*

$$(a) \quad N(T \cap V^{(0)}(\mathbb{Z}); X) = \frac{\pi^2}{12} \cdot \prod_p \mu_p(T) \cdot X + o(X);$$

$$(b) \quad N(T \cap V^{(1)}(\mathbb{Z}); X) = \frac{\pi^2}{4} \cdot \prod_p \mu_p(T) \cdot X + o(X).$$

For our particular application, we require a more general congruence version of our counting theorem for binary cubic forms, namely, one which not only allows appropriate infinite sets of congruence conditions to be imposed but which also permits weighted counts of lattice points (where weights are also assigned by congruence conditions). More precisely, we say that a function $\phi: V(\mathbb{Z}) \rightarrow [0, 1] \subset \mathbb{R}$ is *defined by congruence conditions* if, for all primes p , there exist functions $\phi_p: V(\mathbb{Z}_p) \rightarrow [0, 1]$ such that:

- (1) For all $f \in V(\mathbb{Z})$, we have $\phi(f) = \prod_p \phi_p(f)$.
- (2) For each prime p , the function ϕ_p is locally constant outside some set $Z_p \subset V(\mathbb{Z}_p)$ of measure zero.

Such a function ϕ is called *acceptable* if it is $\mathrm{SL}_2(\mathbb{Z})$ -invariant and, for sufficiently large primes p , we have $\phi_p(f) = 1$ whenever $p^2 \nmid \mathrm{disc}(f)$. For such an acceptable function ϕ , we let $N_\phi(V^{(i)}(\mathbb{Z}); X)$ denote the number of $\mathrm{SL}_2(\mathbb{Z})$ -equivalence classes of elements in $V^{(i)}(\mathbb{Z})$ having absolute discriminant at most X , where the equivalence class of $f \in V^{(i)}(\mathbb{Z})$ is weighted by $\phi(f)$.

We then have the following generalization of Theorem 37:

Theorem 38. *Let $\phi: V(\mathbb{Z}) \rightarrow [0, 1]$ be an acceptable function that is defined by congruence conditions via the local functions $\phi_p: V(\mathbb{Z}_p) \rightarrow [0, 1]$. Then*

$$N_\phi(V^{(i)}(\mathbb{Z}); X) = N(V^{(i)}(\mathbb{Z}); X) \prod_p \int_{f \in V_{\mathbb{Z}_p}} \phi_p(f) df + o(X). \quad (18)$$

Proof. The proof is exactly as in [6, Theorem 2.21], though we use [10, Proposition 29] in place of the uniformity estimate [6, Theorem 2.13]. \square

7.2 Weighted counts of binary cubic forms corresponding to Selmer elements

We wish to apply Theorem 38 to the set T of all integral binary cubic forms that lie in the correspondence of Theorems 33 and 34, with appropriately assigned weights. Namely, we need to count each $\mathrm{SL}_2(\mathbb{Z})$ orbit, $\mathrm{SL}_2(\mathbb{Z}) \cdot f$, weighted by $1/n(f)$, where $n(f)$ is equal to the number of $\mathrm{SL}_2(\mathbb{Z})$ -orbits inside the $\mathrm{SL}_2(\mathbb{Q})$ -equivalence class of f in $V(\mathbb{Z})$. For this purpose, it suffices to count the number of $\mathrm{SL}_2(\mathbb{Z})$ -orbits of locally soluble integral binary cubic forms having bounded discriminant and no rational linear factor with each orbit $\mathrm{SL}_2(\mathbb{Z}) \cdot f$ weighted by $1/m(f)$, where

$$m(f) := \sum_{f' \in O(f)} \frac{|\mathrm{Aut}_{\mathbb{Q}}(f')|}{|\mathrm{Aut}_{\mathbb{Z}}(f')|} = \sum_{f' \in O(f)} \frac{|\mathrm{Aut}_{\mathbb{Q}}(f)|}{|\mathrm{Aut}_{\mathbb{Z}}(f')|},$$

here $O(f)$ is a set of representatives for the action of $\mathrm{SL}_2(\mathbb{Z})$ on the $\mathrm{SL}_2(\mathbb{Q})$ -equivalence class of f in $V(\mathbb{Z})$ and $\mathrm{Aut}_{\mathbb{Q}}(f)$ (resp. $\mathrm{Aut}_{\mathbb{Z}}(f)$) denotes the stabilizer of f in $\mathrm{SL}_2(\mathbb{Q})$ (resp. $\mathrm{SL}_2(\mathbb{Z})$). The reason it suffices to weight by $1/m(f)$ instead of $1/n(f)$ is that, by the proof of [10, Lemma 22], all but a negligible number of $\mathrm{SL}_2(\mathbb{Z})$ -orbits of integral irreducible binary cubic forms with bounded discriminant have trivial stabilizer in $\mathrm{SL}_2(\mathbb{Q})$; thus all but a negligible number of $\mathrm{SL}_2(\mathbb{Z})$ -equivalence classes of integral binary cubic forms of bounded discriminant satisfy $m(f) = n(f)$. In the remainder of this subsection, we compute this weighted p -adic density of locally soluble integral binary cubic forms.

For a prime p and a binary cubic form $f \in V(\mathbb{Z}_p)$, define $m_p(f)$ by

$$m_p(f) := \sum_{f' \in O_p(f)} \frac{|\text{Aut}_{\mathbb{Q}_p}(f')|}{|\text{Aut}_{\mathbb{Z}_p}(f')|} = \sum_{f' \in O_p(f)} \frac{|\text{Aut}_{\mathbb{Q}_p}(f)|}{|\text{Aut}_{\mathbb{Z}_p}(f')|},$$

where $O_p(f)$ is a set of representatives for the action of $\text{SL}_2(\mathbb{Z}_p)$ on the $\text{SL}_2(\mathbb{Q}_p)$ -equivalence class of f in $V(\mathbb{Z}_p)$ and $\text{Aut}_{\mathbb{Q}_p}(f)$ (resp. $\text{Aut}_{\mathbb{Z}_p}(f)$) denotes the stabilizer of f in $\text{SL}_2(\mathbb{Q}_p)$ (resp. $\text{SL}_2(\mathbb{Z}_p)$). By Proposition 17, we have the factorization

$$m(f) = \prod_p m_p(f),$$

and so we have put ourselves in position to use Theorem 38.

Let S be an acceptable subset of \mathbb{Z} , and for each prime p , let S_p denote the closure of S in \mathbb{Z}_p . Let $B(S)$ denote the set of all locally soluble integral binary cubic forms having discriminant $3^6 4k$ for $k \in S$, and let $B_p(S)$ denote the p -adic closure of $B(S)$ in $V(\mathbb{Z}_p)$.

We now determine the p -adic density of $B_p(S)$, where each element $f \in B_p(S)$ is weighted by $1/m_p(f)$, in terms of the p -adic integral over S_p of the local Selmer ratio and the volume $\text{Vol}(\text{SL}_2(\mathbb{Z}_p))$ of the group $\text{SL}_2(\mathbb{Z}_p)$, which is computed with respect to a fixed generator ω of the rank 1 module of top-degree differentials of SL_2 over \mathbb{Z} , so that ω is well-determined up to sign.

Proposition 39. *There is a rational number $\mathcal{J} \in \mathbb{Q}^\times$, independent of S and p , such that*

$$\int_{B_p(S)} \frac{1}{m_p(f)} df = |\mathcal{J}|_p \cdot \text{Vol}(\text{SL}_2(\mathbb{Z}_p)) \cdot \int_{k \in S} \frac{|E_k(\mathbb{Q}_p)/\hat{\phi}(E_{-27k}(\mathbb{Q}_p))|}{|E_{-27k}(\mathbb{Q}_p)[\hat{\phi}]|} dk.$$

Proof. We make use of the facts that the number of $\text{SL}_2(\mathbb{Q}_p)$ -orbits of soluble binary cubic forms of discriminant $4k$ is equal to the cardinality of $E_{-27k}(\mathbb{Q}_p)/\hat{\phi}(E_k(\mathbb{Q}_p))$ (by Theorem 33), and the cardinality of $\text{Aut}_{\mathbb{Q}_p}(f)$ is equal to the cardinality of $E_k(\mathbb{Q}_p)[\hat{\phi}]$ (by Corollary 21); otherwise, the proof is identical to [6, Proposition 3.9]. \square

We note that the rational number \mathcal{J} also shows up in the archimedean factor of Theorem 38, since

$$N(V_{\mathbb{Z}}^{(i)}; X) = \frac{\text{Vol}(\text{SL}_2(\mathbb{Z}) \backslash \text{SL}_2(\mathbb{R})) |\mathcal{J}|_\infty}{n_i} X + o(X), \quad (19)$$

where $n_0 = 3$ and $n_1 = 1$; see [6, Proposition 3.11, Remark 3.14]. (In fact, it turns out that $\mathcal{J} = 3/2$, although we shall not need this fact.)

7.3 Proof of Theorems 1 and 2

Theorem 40. *Let $S \subset \mathbb{Z}$ be any acceptable subset of integers. Then when all elliptic curves $E_k: y^2 = x^3 + k$, $k \in S$, are ordered by the absolute value of k , the average size of the $\hat{\phi}$ -Selmer group associated to the 3-isogeny $\hat{\phi}: E_{-27k} \rightarrow E_k$ is*

$$1 + \prod_{p \leq \infty} \frac{\int_{k \in S_p} c_p(\hat{\phi}_k) dk}{\int_{k \in S_p} dk}, \quad (20)$$

where S_p denotes the p -adic closure of S in \mathbb{Z}_p for $p < \infty$, and S_∞ is the smallest interval in \mathbb{R} containing S .

Proof. Let $X = 3^6 4Y$, and $G = \mathrm{SL}_2$ and define $E'_k = E_{-27k}$. By the discussion in §7.2, to count the total number of non-identity elements in the groups $\mathrm{Sel}_{\hat{\phi}_k}(E_{-27k})$ with $k < Y$, it suffices to count the number of locally soluble irreducible $\mathrm{SL}_2(\mathbb{Z})$ -orbits of binary cubic forms $f \in V(\mathbb{Z})$ having discriminant in the set $3^6 4S$ and also bounded by X , and where each orbit is weighed by $1/m(f)$. This weighting function is acceptable by Proposition 16. Thus, by Theorem 38, Proposition 39, §4.2, and (19), the number of such weighted orbits divided by the total number of $k \in S$ with $k < Y$ approaches

$$\begin{aligned} & \mathrm{Vol}(G(\mathbb{Z}) \backslash G(\mathbb{R})) |\mathcal{J}|_\infty \frac{\sum_{\substack{k \in S \\ |k| < Y}} \frac{1}{|E'_k[\hat{\phi}](\mathbb{R})|}}{\sum_{\substack{k \in S \\ |k| < Y}} 1} \prod_p |\mathcal{J}|_p \mathrm{Vol}(G(\mathbb{Z}_p)) \frac{\prod_p \int_{k \in S_p} \frac{|E_k(\mathbb{Q}_p)/\hat{\phi}(E'_k(\mathbb{Q}_p))|}{|E'_k[\hat{\phi}](\mathbb{Q}_p)|} dk}{\prod_p \int_{k \in S_p} dk} \\ &= \frac{\sum_{\substack{k \in S \\ |k| < Y}} \frac{1}{|E'_k[\hat{\phi}](\mathbb{R})|}}{\sum_{\substack{k \in S \\ |k| < Y}} 1} \prod_p \frac{\int_{k \in S_p} \frac{|E_k(\mathbb{Q}_p)/\hat{\phi}(E'_k(\mathbb{Q}_p))|}{|E'_k[\hat{\phi}](\mathbb{Q}_p)|} dk}{\int_{k \in S_p} dk} \\ &= \prod_{p \leq \infty} \frac{\int_{k \in S_p} c_p(\hat{\phi}_k) dk}{\int_{k \in S_p} dk} \end{aligned}$$

as $Y \rightarrow \infty$. The first equality is due to the product formula $\prod_{p \leq \infty} |\mathcal{J}|_p = 1$, and the fact that the Tamagawa number $\mathrm{Vol}(G(\mathbb{Z}) \backslash G(\mathbb{R})) \prod_p \mathrm{Vol}(G(\mathbb{Z}_p))$ of G is equal to 1. This proves formula (20) of Theorem 40, after taking into account the identity element of each $\hat{\phi}$ -Selmer group. \square

Proof of Theorem 2. The formula comes from the change of variables $k \mapsto -27k$ in the result of Theorem 40, and the fact that $\hat{\phi}_k = \hat{\phi}_{-27k}$. \square

Proof of Theorem 1. We use Proposition 28 to compute the p -adic integrals in Theorem 2 for the set $S = \mathbb{Z}$. The result is that the product of the local densities at finite primes is equal to

$$\begin{aligned} r &= \left[\frac{(1 - 2^{-1}) \left(\frac{4}{3} + 2^{-1} + \frac{4}{3} 2^{-2} + 2^{-3} + 2^{-4} + 2^{-5} \right)}{1 - 2^{-6}} \right] \\ &\cdot \left[\frac{(1 - 3^{-1}) \left(\frac{5}{3} + 2 \cdot 3^{-1} + \frac{2}{3} \cdot 3^{-2} + \frac{7}{3} \cdot 3^{-3} + 2 \cdot 3^{-4} + 6 \cdot 3^{-5} \right)}{1 - 3^{-6}} \right] \\ &\cdot \prod_{p \equiv 5 \pmod{6}} \left[\frac{(1 - p^{-1}) \left(1 + p^{-1} + \frac{5}{3} p^{-2} + p^{-3} + \frac{5}{3} p^{-4} + p^{-5} \right)}{1 - p^{-6}} \right]. \end{aligned}$$

In each Euler factor, the six-term sum records the weighted average value of $c_p(\hat{\phi})$ on $p^i \mathbb{Z}_p - p^{i+1} \mathbb{Z}_p$ for $i = 0, \dots, 5$. Since $E_{km^6} \cong E_k$, these averages cyclically repeat themselves for $i \geq 6$, which explains the factor $1 - p^{-6}$ in the denominator. \square

7.4 Proof of Theorem 5

Claim (i) is clear, and (ii) follows immediately from Proposition 28. Claim (iii) follows from Cassels' formula [14],

$$c(\phi) = \frac{|E_{-27k}(\mathbb{Q})[\hat{\phi}]| \cdot |\text{Sel}_\phi(E_k)|}{|E_k(\mathbb{Q})[\phi]| \cdot |\text{Sel}_{\hat{\phi}}(E_{-27k})|}. \quad (21)$$

To prove (iv), we use Proposition 28 to express T_m as a disjoint union

$$T_m = \bigcup_{n \in \mathbb{Z}} T_{m,n},$$

where $T_{m,n}$ is the set of all $k \in T_m$ for which there are exactly $m+n$ primes p (possibly including $p = \infty$) satisfying $c_p(\phi_k) \geq 3$, with the caveat that if $c_3(\phi_k) = 9$ then $p = 3$ is counted twice. Each $T_{m,n}$ can itself be expressed as a disjoint union

$$T_{m,n} = \bigcup_{(\Pi_1, \Pi_2)} T_{\Pi_1, \Pi_2},$$

where the union is over all pairs (Π_1, Π_2) , where Π_1 is a multiset of $m+n$ primes, with each prime having multiplicity at most 1, except for $p = 3$ which may have multiplicity at most 2, and Π_2 is a set of n primes, disjoint from Π_1 . The set T_{Π_1, Π_2} consists of those integers $k \in T_m$ such that $c_p(\phi_k) \geq 3$ if and only if $p \in \Pi_1$ (and having multiplicity 2 if and only if $p = 3$ and $c_3(\phi_k) = 9$), and such that $c_p(\phi_k) = 1/3$ if and only if $p \in \Pi_2$.

Note that each set $T_{\Pi_1, \Pi_2} \subset \mathbb{Z}$ is acceptable, by the explicit congruence conditions of Proposition 28. Moreover, for each prime p , the function $c_p(\phi_k)$ is constant on T_{Π_1, Π_2} . Thus by Theorem 2, for any acceptable set $S \subset T_{\Pi_1, \Pi_2}$, the average size of $\text{Sel}_\phi(E_k)$ for $k \in S$ is

$$1 + \prod_p c_p(\phi_k) = 1 + c(\phi_k) = 1 + 3^m.$$

Since $c(\hat{\phi}) = c(\phi)^{-1}$, a similar argument shows that the average size of $\text{Sel}_{\hat{\phi}}(E_{-27k})$ for $k \in S$ is $1 + 3^{-m}$. Note that any acceptable set S contained in T_m is necessarily contained in some T_{Π_1, Π_2} , so this proves Theorem 5(iv) in the case that $S \subset T_m$ is an acceptable set.

If $S = T_m$, then since

$$T_m = \bigcup_n \bigcup_{(\Pi_1, \Pi_2)} T_{\Pi_1, \Pi_2},$$

we may write T_m as an ascending union $T_m = \bigcup_{i \geq 1} S_i$, where each S_i is the union of all T_{Π_1, Π_2} such that each prime $p \in \Pi_1 \cup \Pi_2$ is less than the i -th prime number p_i . Thus, each S_i is a finite union of acceptable sets, and so the average size of $\text{Sel}_\phi(E_k)$ for $k \in S_i$ is $1 + 3^m$. Moreover, by Proposition 28, the binary cubic forms $f \in V(\mathbb{Z})$ having discriminant in the complement $T_m - S_i$ have discriminant divisible by p_j^2 for some $j \geq i$. By the uniformity estimate in [10, Proposition 29], we have $\sum_{k \in T_m \setminus S_i; |k| < X} \text{Sel}_\phi(E_k)$ is $\sum_{j \geq i} O(X/p_j^2) = O(X/p_i)$, where the implied constant is independent of i . That is, the total number of Selmer elements of E_k over all $k \in T_m \setminus S_i$, $|k| < X$, is $O(X/p_i)$, while the total number of Selmer elements of E_k over all $|k| < X$ is of course $\gg X$. Letting i tend to infinity, we conclude that the average size of $\text{Sel}_\phi(E_k)$ for $k \in T_m$ is also equal to $1 + 3^m$. The same argument shows that the average size of $\text{Sel}_{\hat{\phi}}(E_k)$ for $k \in T_m$ is again equal to $1 + 3^{-m}$. This completes the proof of Theorem 5.

Remark 41. If $S \subset \mathbb{Z}$ is any acceptable set, then the average size of $\text{Sel}_\phi(E_k)$ for $k \in S$ is equal to $\sum_{m \in \mathbb{Z}} \mu(S \cap T_m)(1 + 3^m)$ and the average size of $\text{Sel}_{\hat{\phi}}(E_{-27k})$ is equal to $\sum_{m \in \mathbb{Z}} \mu(S \cap T_m)(1 + 3^{-m})$. Indeed, even though the sets $S \cap T_m$ are not in general acceptable, the sets of the form $S \cap T_{\Pi_1, \Pi_2}$ are, so we can argue as above.

8 Ranks of elliptic curves

In the previous section, we computed the average size of the ϕ -Selmer and $\hat{\phi}$ -Selmer groups in any acceptable family of elliptic curves with j -invariant 0. In this section, we deduce an upper bound on the average rank of $E(\mathbb{Q})$ in such families. We also deduce a lower bound on the density of curves E_k having rank 0 or (3-Selmer) rank 1, respectively.

8.1 An upper bound on the average rank of elliptic curves with vanishing j -invariant

For any elliptic curve E/\mathbb{Q} with j -invariant 0, we write $r(E)$, $r_3(E)$, $r_\phi(E)$, and $r_{\hat{\phi}}(E)$ for the ranks of the groups $E(\mathbb{Q})$, $\text{Sel}_3(E)$, $\text{Sel}_\phi(E)$, and $\text{Sel}_{\hat{\phi}}(E')$ respectively, where $E' = E/\ker \phi$.

Proposition 42. *For $\phi: E \rightarrow E'$ as above, we have*

$$(i) \quad r(E) \leq r_3(E) \leq r_\phi(E) + r_{\hat{\phi}}(E).$$

$$(ii) \quad \text{If } c(\phi) = 3^m, \text{ then } m \equiv r_3(E) - \dim_{\mathbb{F}_3} E[3](\mathbb{Q}) \pmod{2}.$$

Proof. Claim (i) follows from the exact sequence [28, Corollary 1],

$$0 \rightarrow \frac{E'(\mathbb{Q})[\hat{\phi}]}{\phi(E(\mathbb{Q})[3])} \rightarrow \text{Sel}_\phi(E) \rightarrow \text{Sel}_3(E) \rightarrow \text{Sel}_{\hat{\phi}}(E') \rightarrow \frac{\text{III}(E')[\hat{\phi}]}{\phi(\text{III}(E)[3])} \rightarrow 0. \quad (22)$$

As ϕ is adjoint with respect to the Cassels–Tate pairing [14, Theorem 1.2], there is an induced non-degenerate alternating pairing on the \mathbb{F}_3 -vector space $\frac{\text{III}(E')[\hat{\phi}]}{\phi(\text{III}(E)[3])}$. The latter therefore has even \mathbb{F}_3 -dimension, and (ii) now follows from (21) and (22). \square

Theorem 43. *Let $m \in \mathbb{Z}$, and suppose $S \subset T_m$ is an acceptable set. Then the (limsup of the) average rank of E_k , for $k \in S$ ordered by absolute value, is at most $|m| + 3^{-|m|}$.*

Proof. First assume that $m \geq 0$. Then one has the general inequality

$$2r_\phi(E) + 1 - 2m \leq 3^{r_\phi(E) - m}, \quad (23)$$

so that

$$r_\phi(E) \leq m - \frac{1}{2} + \frac{|\text{Sel}_\phi(E)|}{2 \cdot 3^m}.$$

By Theorem 5, the average size of $\text{Sel}_\phi(E_k)$ for $k \in S \subset T_m$ is $1 + 3^m$, so we conclude that the limsup of the average of $r_\phi(E_k)$ is at most $m + \frac{1}{2}3^{-m}$. Similarly, the average size of $\text{Sel}_{\hat{\phi}}(E_{-27k})$ is $1 + 3^{-m}$, so we conclude from the case $m = 0$ of inequality (23) that the limsup of the average of $r_{\hat{\phi}}(E_k)$ is at most $\frac{1}{2}3^{-m}$. Combining these bounds with Proposition 42(i), the limsup of the average rank of E_k for $k \in S$ is seen to be at most $m + 3^{-m}$. For $m < 0$, the roles of ϕ and $\hat{\phi}$ are reversed, so the average rank bounds that we obtain for m and $-m$ are the same. \square

Proof of Theorem 6. We observe that, by the explicit congruence description of T_m in §7.4, any $k \in T_m$ (for $|m| > 1$) must satisfy $p_j^2 \mid k$ for some $j \geq |m| - 1$, where we again use p_i to denote the i -th prime. Hence, by the uniformity estimate [10, Proposition 29], we have that

$$\sum_{|m| > M} \sum_{\substack{k \in T_m \\ k < X}} \text{Sel}_\phi(E_k) = \sum_{j \geq M} O(X/p_j^2) = O(X/p_M),$$

where the implied constant is independent of M . Theorem 43 therefore gives the following bound on the (limsup of the) average rank of the elliptic curves E_k for non-zero $k \in \mathbb{Z}$:

$$\limsup \text{avg}_{k \in \mathbb{Z}} r(E_k) \leq \lim_{M \rightarrow \infty} \sum_{m=-M}^M \mu(T_m)(|m| + 3^{-|m|}) + O(1/p_M) = \sum_{m=-\infty}^{\infty} \mu(T_m)(|m| + 3^{-|m|}),$$

where $\mu(T_m)$ denotes the density of integers in the set T_m . Using Proposition 28 and the explicit description given in §7.4 of T_m as the disjoint union

$$T_m = \bigcup_n T_{\Pi_1, \Pi_2},$$

we can estimate the densities $\mu(T_m)$ to arbitrary precision. One may also prove, e.g., that

$$\sum_{|m| > 5} \mu(T_m)(|m| + 3^{-|m|}) < 0.001.$$

A computation in Mathematica produces the table of densities in the introduction, and it follows that the average rank of E_k is less than 1.29. We can obtain a better bound on the average rank if we use the the work of [30]. Ruth's main result—that the average size of the 2-Selmer group of E_k , $k \in S$, is 3—is stated for the set $S = \mathbb{Z}$, but the proof appears to hold for any acceptable set S . Since the 3-Selmer—and therefore (due to the work of [17]) the 2-Selmer parity—is constant for 100% of E_k for $k \in T_m$ by Theorems 5(iii) and 42(ii), this implies that the average rank of E_k , $k \in S$, is less than $4/3$ on any T_m , giving the bound:

$$\text{avg}_{k \in \mathbb{Z}} r(E) \leq \sum_{m=-1}^1 \mu(T_m)(|m| + 3^{-|m|}) + \frac{4}{3}(1 - \mu(S_1 \cup S_0 \cup S_{-1})) < 1.21.$$

□

8.2 A lower bound on the proportion of elliptic curves with vanishing j -invariant having 3-Selmer rank 0 or 1

Proof of Theorem 7. Let s_0 denote the liminf of the natural density of $k \in T_0$ for which $r_\phi(E_k) = 0$. Then by Theorem 5(iv), s_0 is also the liminf of the natural density of $k \in T_0$ for which $r_\phi(E_k) = 0$. Since the average size of $|\text{Sel}_\phi(E_k)|$, $k \in \mathbb{Z}$, is 2, we must have

$$s_0 \cdot 1 + (1 - s_0) \cdot 3 \leq 2,$$

implying $s_0 \geq 1/2$. It follows that at least $1/2$ of the curves E_k , $k \in T_0$, satisfy $r_\phi(E_k) = r_\phi(E_k) = r_3(E_k) = 0$, and thus $r(E_k) = 0$. Theorem 7 follows, since the density of T_0 is at least .399, and $(1/2)(.399) > .199$. □

Proof of Theorem 8. Let s_1 denote the liminf of the natural density of $k \in T_1$ for which $r_\phi(E_k) = 1$. Then by Theorem 5(iv), s_1 is also the liminf of the natural density of $k \in T_1$ for which $r_{\hat{\phi}}(E_k) = 0$. Since the average size of $|\text{Sel}_\phi(E_k)|$, $k \in T_1$, is 4, we see that

$$s_1 \cdot 3 + (1 - s_1) \cdot 9 \leq 4,$$

implying $s_1 \geq 5/6$. In conjunction with the exact sequence (22), this implies that a lower density of at least $5/6$ of the curves E_k , $k \in T_1$, satisfy $r_{\hat{\phi}}(E_k) = 0$ and $r_\phi(E_k) = r_3(E_k) = 1$ (and under the assumption that $\text{III}(E_k)[3^\infty]$ is finite, that $r(E_k) = 1$ as well). The identical argument with T_{-1} in place of T_1 shows that a lower density of at least $5/6$ of the curves E_k , $k \in T_{-1}$, satisfy $r_\phi(E_k) = 0$ and $r_{\hat{\phi}}(E_k) = 1$. By (22) and the fact that $\frac{\text{III}(E')[\hat{\phi}]}{\phi(\text{III}(E)[3])}$ has even rank, these curves also satisfy $r_3(E_k) = 1$ (and again $r(E_k) = 1$ under the assumption that $\text{III}(E_{-27k})[3]$ is trivial). Theorem 8 now follows, because the density of $T_1 \cup T_{-1}$ is at least .494, and $(5/6)(.494) > .411$. \square

Since $.199 + .411 = .61$, we conclude that at least 61% of all E_k have rank 0 or 1, which is Corollary 9.

Acknowledgments

We are grateful to Benedict Gross and Ila Varma for helpful conversations. The first author was supported by a Simons Investigator Grant and NSF grant DMS-1001828. The second author was supported by NSF grants DMS-1100511 and DMS-1502161. The third author was partially supported by NSF grant DMS-0943832.

References

- [1] M. Bhargava, Higher composition laws. I. A new view on Gauss composition, and quadratic generalizations, *Ann. of Math.* **159** (2004), no. 1, 217–250.
- [2] M. Bhargava, Mass formulae for local fields, and conjectures on the density of number fields discriminants, *Internat. Math. Research Notices* (2007), Article ID rnm052, 20 pages.
- [3] M. Bhargava and B. Gross, The average size of the 2-Selmer group of Jacobians of hyperelliptic curves having a rational Weierstrass point. *Automorphic representations and L-functions*, 23–91, *Tata Inst. Fundam. Res. Stud. Math.* **22**, Tata Inst. Fund. Res., Mumbai, 2013.
- [4] M. Bhargava, N. D. Elkies, D. Kane, and A. Shnidman, forthcoming.
- [5] M. Bhargava, Z. Klagsbrun, R. Lemke Oliver, and A. Shnidman, Average sizes of Selmer groups in families of quadratic twists with a 3-isogeny, preprint.
- [6] M. Bhargava and A. Shankar, Binary quartic forms having bounded invariants, and the boundedness of the average rank of elliptic curves, *Ann. of Math. (2)* **181** (2015), no. 1, 191–242.
- [7] M. Bhargava and A. Shankar, Ternary cubic forms having bounded invariants, and the existence of a positive proportion of elliptic curves having rank 0, *Ann. of Math. (2)* **181** (2015), no. 2, 587–621.

- [8] M. Bhargava and A. Shankar, The average number of elements in the 4-Selmer groups of elliptic curves is 7, preprint.
- [9] M. Bhargava and A. Shankar, The average size of the 5-Selmer group of elliptic curves is 6, and the average rank is less than 1, preprint.
- [10] M. Bhargava, A. Shankar, and J. Tsimerman. On the Davenport-Heilbronn theorems and second order terms. *Invent. Math.* **193** (2013), no. 2, 439–499.
- [11] M. Bhargava and I. Varma, The mean number of 3-torsion elements in the class groups and ideal groups of quadratic orders, *Proc. Lond. Math. Soc.* (3) **112** (2016), no. 2, 235–266.
- [12] J. W. S. Cassels, *Lectures on elliptic curves*, London Mathematical Society Student Texts **24**, Cambridge University Press, Cambridge, 1991.
- [13] J. W. S. Cassels, Diophantine equations with special reference to elliptic curves, *J. London Math. Soc.* **41** 1966, 193–291.
- [14] J. W. S. Cassels, Arithmetic on curves of genus 1, VIII: On conjectures of Birch and Swinnerton-Dyer, *J. Reine Angew. Math.* **217** (1965), 180–199.
- [15] H. Davenport, On the class-number of binary cubic forms I and II, *J. London Math. Soc.* **26** (1951), 183–198.
- [16] H. Davenport and H. Heilbronn, On the density of discriminants of cubic fields II, *Proc. Roy. Soc. London Ser. A* **322** (1971), no. 1551, 405–420.
- [17] T. Dokchitser and V. Dokchitser, On the Birch–Swinnerton-Dyer quotient modulo squares, *Annals of Math.* **172** (2010), no.1, 567–596.
- [18] V. Dokchitser, Root numbers of non-abelian twists of elliptic curves (with an appendix by Tom Fisher), *Proc. London Math. Soc.* (3) **91** (2005), no. 2, 300–324.
- [19] G. Eisenstein, Théorèmes sur les formes cubiques et solution d’une équation du quatrième degré à quatre indéterminées, *J. Reine Angew. Math.* **27** (1844), 75–79.
- [20] É. Fouvry, Sur le comportement en moyenne du rang des courbes $y^2 = x^3 + k$, *Séminaire de Théorie des Nombres, Paris, 1990-91, Progr. Math.* **108**, Birkhäuser Boston, MA, 1993.
- [21] B. Gross, J. Parson, On the local divisibility of Heegner points. *Number theory, analysis and geometry*, 215–241, Springer, New York, 2012.
- [22] D. R. Heath-Brown, The size of Selmer groups for the congruent number problem, *Invent. Math.* **111** (1993), no. 1, 171–195.
- [23] D. Hilbert, *Theory of Algebraic Invariants* (1897; trans. R. C. Laubenchacher, ed. B. Sturmfels), Cambridge University Press 1993.
- [24] E. Liverance, Binary cubic forms with many integral points,
<http://www.kurims.kyoto-u.ac.jp/~kyodo/kokyuroku/contents/pdf/0998-8.pdf>

- [25] D. Kane, On the Ranks of the 2-Selmer Groups of Twists of a Given Elliptic Curve, preprint (arXiv:1111.2321).
- [26] D. Kane, J. Thorne, On the ϕ -Selmer groups of the elliptic curves $y^2 = x^3 - Dx$, *Mathematical Proceedings of the Cambridge Philosophical Society* (2016), <http://dx.doi.org/10.17863/CAM.1298>.
- [27] Z. Klagsbrun, B. Mazur, and K. Rubin, Selmer ranks of quadratic twists of elliptic curves, preprint (arXiv:1111.2321v1).
- [28] R. Kloosterman and E. Schaefer, Selmer groups of elliptic curves that can be arbitrarily large, *J. Number Theory* **99** (2003), no. 1, 148–163.
- [29] B. Poonen and E. Rains, Random maximal isotropic subspaces and Selmer groups, *J. Amer. Math. Soc.*, **25** (2012), no. 1, 245–269. arXiv: 1009.0287
- [30] S. Ruth, A bound on the average rank of j -invariant zero elliptic curves, Princeton Ph.D. Thesis (2014).
- [31] P. Satgé, Groupes de Selmer et corps cubiques, *J. Number Theory* **23** (1986), no. 3, 294–317.
- [32] E. Schaefer, Class groups and Selmer groups, *J. Number Theory* **56** (1996), no. 1, 79–114.
- [33] I. Schur, *Vorlesungen über Invariantentheorie*, Grundlehren der mathematischen Wissenschaften **143**, Springer: Berlin 1968.
- [34] E. Selmer, The Diophantine equation $ax^3 + by^3 + cz^3 = 0$, *Acta Math.* **85** (1951), 203–362.
- [35] H. P. F. Swinnerton-Dyer, The effect of twisting on the 2-Selmer group, *Math. Proc. Cambridge Philos. Soc.* **145** (2008), no. 3, 513–526.
- [36] J. Tate, Algorithm for determining the type of a singular fiber in an elliptic pencil, *Lecture Notes in Mathematics* **476** (Modular Forms in One Variable IV, Antwerp 1972; B. J. Birch and W. Kuyk, eds.), 33–52.
- [37] J. Thorne, Vinberg’s representations and arithmetic invariant theory, *Algebra Number Theory* **7** (2013), no. 9, 2331–2368.
- [38] J. Thorne, E_6 and the arithmetic of a family of non-hyperelliptic curves of genus 3, *Forum Math. Pi* **3** (2015), e1, 41 pp.