# $(\varphi_1, \varphi_2)$ –Variational principle

Abdelhakim MAADEN and Abdelkader STOUTI

Université Sultan Moulay Slimane Faculté des Sciences et Techniques. Laboratoire de Mathématiques et Applications. B.P. 523. BENI - MELLAL, MAROC. Corresponding author: E-mail address: hmaaden2002@yahoo.fr

#### Abstract

In this paper we prove that if X is a Banach space, then for every lower semicontinuous bounded below function f, there exists a  $(\varphi_1, \varphi_2)$ -convex function g, with arbitrarily small norm, such that f + g attains its strong minimum on X. This result extends some of the well-known variational principles as that of Ekeland [18], that of Borwein-Preiss [6] and that of Deville-Godefroy-Zizler [14, 15].

Keywords:  $(\varphi_1, \varphi_2)$  -convex function,  $(\varphi_1, \varphi_2)$  -variational principle, Ekeland's variational principle, smooth variational principle.

Mathematics subject classification (2010): 26A51, 52A30, 46B20.

### 1 Introduction

Let  $(X, \|.\|)$  be a Banach space. Let f be a real-valued function defined on X, lower semicontinuous and bounded below. Let P be a class of functions in X which serves as a source of possible perturbations for f. By a variational principle we mean an assertion ensuring the existence of at least one perturbation g from P such that f + g attains its minimum on X.

The first variational principle, based on the Bishop-Phelps lemma [3, 27], was established by Ekeland [18]. It says that P is a family of suitable translations of the norm.

If g is required to be smooth, then we speak about the smooth variational principle. The first result of this type was shown by Stegall [31, 27], where P is the elements of the dual space  $X^*$ . He proved that if X has the Radon-Nikodym property in particular, if X is reflexive, then one can take for g even a linear functional, with arbitrarily small norm. In [17], Deville-Maaden shown under the same hypotheses as Stegall that X has the Radon-Nikodym property and the function f is lower semicontinuous and superlinear with the set P consisting of radial smooth functions. However, this principle does not cover some important Banach spaces. For example the space  $c_0$  does not have the Radon-Nikodym property while it, in fact, admits a smooth norm [5]. In this direction Borwein-Preiss [6] proved a smooth variational principle imposing no additional conditions on the space, except, the presence of some equivalent smooth norm with the set P being the family of smooth combinations of the norm in X. Haydon's work [23], shows that there exists a Banach spaces with smooth bump function without an equivalent smooth norm (a function b is bump if it has a non empty and bounded support). So, the variational principle of Borwein-Preiss is not applicable in these spaces. So that, Deville et al [14, 15] extended the Borwein-Preiss variational principle to spaces with smooth bump function, with P equal to the family of Lipschitz smooth functions.

In an analytical approach we can often associate a geometrical approach to complete study of which or stimulates the analytical approche. From this perspective Browder [8] gaves a geometrical result which bears at present the name of the Drop Theorem (see also [10]). Penot in [26, 21] showed that the drop theorem is a geometrical version of the Ekeland's variational principle. After this, Maaden in [25, 22] introduced and studied the notion of the smooth drop which can be seen as a geometrical version of the smooth variational principle of Borwein-Preiss.

Those variational principles are a tools that have been very important in nonlinear analysis, in that they enjoyed a big deal of applications from the geometry of Banach spaces [3, 4, 7] to the optimization theory [18, 19, 30] and of generalized differential and sub-differential calculus [1, 2, 6, 11, 12, 13, 26], calculus of variations [9, 18] up to the nonlinear semi-groups theory [7, 18] and the viscosity solutions of Hamilton-Jacobi equations [12, 13, 15].

In [28, 29], Pini et all defined the notion of  $(\varphi_1, \varphi_2)$ -convex functions. They say that a real valued function f defined on a non empty subset D of  $\mathbb{R}^n$  is  $(\varphi_1, \varphi_2)$ -convex if  $f(\varphi_1(x, y, \lambda)) \leq \varphi_2(x, y, \lambda, f)$  for all  $x, y \in D$  and for all  $\lambda \in [0, 1]$ . Where  $\varphi_1$  is a function from  $D \times D \times [0,1]$  in  $\mathbb{R}^n$  and  $\varphi_2$  is a function from  $D \times D \times [0,1] \times F$  in  $\mathbb{R}$ , with F is a given vector space of real valued functions defined on the set D. In this paper we shall use the same definition of  $(\varphi_1, \varphi_2)$ -convex functions as above with using any Banach spaces instead of  $\mathbb{R}^n$ . In this way, we prove that under suitable choices of the functions  $\varphi_1$  and  $\varphi_2$  a new variational principle for the set of  $(\varphi_1, \varphi_2)$ -convex functions (see Theorem 3.1). This  $(\varphi_1, \varphi_2)$  – variational principle is providing a unified framework to deduce Ekeland's, Borwein-Preiss's and Deville's variational principles.

### 2 Auxiliaries results

(

In this section we shall give some definitions and establish some auxiliaries results which we shall use to prove our main result in this paper.

Let  $(X, \|.\|)$  be a Banach space. For a continuous function  $f: X \longrightarrow \mathbb{R}$  we define

$$\mu\left(f\right) = \sum_{n=1}^{\infty} \frac{\|f\|_n}{2^n},$$

where

$$||f||_n = \sup \{ |f(x)|; x \in X, ||x|| \le n \}.$$

Let M be the set of all continuous functions f such that  $\mu(f) < \infty$ . It is routine to check that  $(M, \mu)$  is a Banach space.

Let  $\varphi_1 : X \times X \times [0,1] \longrightarrow X$  and  $\varphi_2 : X \times X \times [0,1] \times F \longrightarrow \mathbb{R}$ , two functions where F is a given set of real functions on X. Define,

**Definition 2.1** A function  $f: X \longrightarrow \mathbb{R}$  is said to be  $(\varphi_1, \varphi_2)$  -convex if

$$f\left(\varphi_{1}\left(x, y, \lambda\right)\right) \leq \varphi_{2}\left(x, y, \lambda, f\right), \forall x, y \in X, \forall \lambda \in [0, 1].$$

Remarking that under suitable assumptions on  $\varphi_1$  and/or  $\varphi_2$ , the class of  $(\varphi_1, \varphi_2)$  – convex functions is a convex cone. For example:

1) If  $\varphi_2$  is super-linear with respect to  $f \in F$  (that  $\varphi_2$  is super-additive and positively homogeneous), then the class of  $(\varphi_1, \varphi_2)$  -convex functions is a convex cone.

Indeed, let f, g are two  $(\varphi_1, \varphi_2)$  -convex functions and  $\alpha > 0$ . Then, for  $x, y \in X$ and  $\lambda \in [0, 1]$  we have

$$\begin{aligned} \left(f+g\right)\left(\varphi_{1}\left(x,y,\lambda\right)\right) &\leq \varphi_{2}\left(x,y,\lambda,f\right) + \varphi_{2}\left(x,y,\lambda,g\right) \\ &\leq \varphi_{2}\left(x,y,\lambda,f+g\right) \end{aligned}$$

and

$$(\alpha f) (\varphi_1 (x, y, \lambda)) = \alpha (f (\varphi_1 (x, y, \lambda)))$$
  
$$\leq \alpha \varphi_2 (x, y, \lambda, f)$$
  
$$= \varphi_2 (x, y, \lambda, \alpha f).$$

2) If  $\varphi_2(x, y, \lambda, f) = C((1 - \lambda) f(x) + \lambda f(y))$  for some C > 0, the set of  $(\varphi_1, \varphi_2)$  -convex functions is a convex cone.

In all the sequel, we define the following sets :

$$\Phi = \{ f \in M : f \text{ is } (\varphi_1, \varphi_2) - \text{convex and } f \ge 0 \},\$$
$$F = \{ f \in \Phi : f(x) \longrightarrow +\infty \text{ as } ||x|| \longrightarrow +\infty \}.$$

The metric  $\rho$  on  $\Phi$  is defined as:

$$\rho(f,g) = \mu(f-g) = \sum_{n \ge 1} \frac{\|f-g\|_n}{2^n} \text{ for all } f,g \in \Phi,$$

and it is easy to show that  $(\Phi, \rho)$  is a complete metric space.

- Throughout this paper, the functions  $\varphi_1$  and  $\varphi_2$  satisfies the following assumptions:  $(P_1) \ \varphi_1(x, x, 0) = x; \forall x \in X;$   $(P_2) \ \varphi_1(x, y, \lambda) + \varphi_1(z, z, 0) = \varphi_1(x + z, y + z, \lambda); \forall x, y, z \in X, \forall \lambda \in [0, 1];$   $(P_3) \ \exists C \ge 1$ , such that  $\varphi_2(\lambda x, \lambda x, 0, h) \le C[(1 - \lambda) h(0) + \lambda h(x)]; \forall x \in X, \forall \lambda \in [0, 1], \forall h \in \Phi;$   $(P_4) \ \text{For } x_0 \in X, \ \varphi_2(x - x_0, y - x_0, \lambda, h) \le \varphi_2(x, y, \lambda, h(. - x_0)); \forall x, y \in X, \forall \lambda \in [0, 1]; \forall h \in \Phi;$  $(P_4) \ \text{The class of } (x_0, x_0) = convex functions is a convex cone.$
- $(P_5)$  The class of  $(\varphi_1, \varphi_2)$  convex functions is a convex cone.

We will also assume that  $\varphi_1$  is continuous with respect to  $\lambda$ .

We present now three preliminaries lemmas, which are useful for the proof of our principal result of this paper. In the first, we use  $(P_1)$  and  $(P_3)$  to prove the following:

**Lemma 2.2** Let  $h \in \Phi$  and let  $y = \lambda x$ ,  $\lambda > 1$ . Then,  $h(y) - h(0) \ge \frac{\lambda}{C} (h(x) - Ch(0))$ .

Proof:

Let  $\mu = 1/\lambda$ . Then  $x = \mu y$ . By using  $(P_1)$  and  $(P_3)$  we obtain that:

$$h(x) = h(\mu y)$$
  
=  $h(\varphi_1(\mu y, \mu y, 0))$   
 $\leq \varphi_2(\mu y, \mu y, 0, h)$   
 $\leq C((1 - \mu) h(0) + \mu h(y)).$ 

Consequently, we get

$$h(x) - Ch(0) \le C\mu(h(y) - h(0)).$$

Since c > 0 and  $\mu > 0$ , we deduce that

$$h(y) - h(0) \ge \frac{1}{C\mu} (h(x) - Ch(0)) = \frac{\lambda}{C} (h(x) - Ch(0))$$

and the proof is complete.  $\blacksquare$ 

We have now all tools to confirm that  $(F, \rho)$  is a Baire space. For this, it suffices to show that

**Lemma 2.3**  $(F, \rho)$  is open in  $\Phi$ .

Proof:

Let f in F. Let N > Cf(0) (where C is given by  $(P_3)$ ) and let  $\varepsilon > 0$  be such that

$$0 < \varepsilon < \frac{N - Cf(0)}{2C + 1}.$$
(1)

Since  $f \in F; f \longrightarrow +\infty$  as  $||x|| \longrightarrow +\infty$ ; there exists  $n \in \mathbb{N}$  such that

$$f(x) > N$$
 whenever  $||x|| \ge n.$  (2)

Let  $g \in \Phi$  such that  $\rho(f,g) < \frac{\varepsilon}{2^n}$ . Then

$$\frac{\|f-g\|_n}{2^n} < \frac{\varepsilon}{2^n}$$

Hence,

$$|f(x) - g(x)| < \varepsilon$$
 whenever  $||x|| \le n.$  (3)

In particular, we have

$$|f(0) - g(0)| < \varepsilon. \tag{4}$$

Combining (2) and (3) we obtain for ||x|| = n,

$$g\left(x\right) > N - \varepsilon. \tag{5}$$

On the first, for  $y \in X$  such that  $||y|| \to +\infty$ , there exist  $x \in X$  with ||x|| = n and  $\lambda > 1$  such that  $y = \lambda x$ , and we have  $\lambda \to +\infty$ . Therefore, combining Lemma 2.2, (5), (4) and (1) we obtain

$$g(y) - g(0) \geq \frac{\lambda}{C} (g(x) - Cg(0))$$
  
>  $\frac{\lambda}{C} (N - \varepsilon - Cg(0))$   
$$\geq \frac{\lambda}{C} (N - \varepsilon - Cf(0) - C\varepsilon)$$
  
>  $\lambda \varepsilon,$ 

which implies that

$$g(y) > \lambda \varepsilon + g(0) \ge \lambda \varepsilon > 0.$$

On the other hand, we know that  $\lambda \longrightarrow +\infty$  as  $||y|| \longrightarrow +\infty$ . So, we get that  $g(y) \longrightarrow +\infty$  as  $||y|| \longrightarrow +\infty$ . So that,  $g \in F$  and F is open.

Next, by using  $(P_1), (P_2)$  and  $(P_4)$  we obtain the following:

**Lemma 2.4** Let  $\theta$  be a  $(\varphi_1, \varphi_2)$  - convex function and let  $h(x) = \theta(x - x_0)$ . Then, h is a  $(\varphi_1, \varphi_2)$  - convex function.

Proof:

Let  $x, y \in X$  and  $\lambda \in [0, 1]$ . By using  $(P_1), (P_2)$  and  $(P_4)$  we get

$$h (\varphi_1 (x, y, \lambda)) = \theta (\varphi_1 (x, y, \lambda) - x_0)$$
  
=  $\theta (\varphi_1 (x, y, \lambda) + \varphi_1 (-x_0, -x_0, 0))$   
=  $\theta (\varphi_1 (x - x_0, y - x_0, \lambda))$   
 $\leq \varphi_2 (x - x_0, y - x_0, \lambda, \theta)$   
 $\leq \varphi_2 (x, y, \lambda, \theta (. - x_0))$   
=  $\varphi_2 (x, y, \lambda, h)$ ,

which shows that h is a  $(\varphi_1, \varphi_2)$  -convex function.

**Corollary 2.5** Let  $\theta$  be a  $(\varphi_1, \varphi_2)$  - convex function in F then the function  $h(x) = \theta(x - x_0)$  is in F.

#### 3 The main result

In this section we shall establish a  $(\varphi_1, \varphi_2)$ -variational principle. We show that the set P; which is a source of perturbation for f; is a class of  $(\varphi_1, \varphi_2)$ -convex functions. Furthermore we can take them of  $C^{\infty}$  in smooth Banach spaces.

In the mathematical field of topology, a  $G_{\delta}$  set is a subset of a topological space that is a countable intersection of open sets. In a complete metric space, a countable union of nowhere dense sets is said to be meagre; the complement of such a set is a residual set.

An element y of a Banach space X is said a strong minimum for a real function f defined on the space X, if f(y) is the infimum of f and any minimizing sequence for f converges to y.

The aim result in this paper is the following variational principle:

**Theorem 3.1** Let X be a Banach space. Let  $f : X \longrightarrow \mathbb{R} \cup \{+\infty\}$  be a lower semicontinuous function bounded from below. Let Y be a subset of F such that:

i) the metric  $\rho_Y$  in Y is such that  $\rho_Y(f,g) = \mu_Y(f-g) \ge \mu(f-g)$ , for all  $f,g \in Y$ .

ii)  $(Y, \rho_Y)$  is a Baire space.

*iii)* there exists  $\theta \in Y$  such that  $\mu_Y(\theta) < +\infty, \theta(0) = 0$ , there is  $k \in ]0, 1[$  such that for every  $||x|| \ge k$  we have  $\theta(x) \ge k^2$  and  $\mu_Y(\theta(.-x_0)) \le \mu_Y(\theta) + ||\theta||_{||x_0||}$ .

Then the set

$$\{g \in Y : f + g \text{ attains its strong minimum on } X\}$$

is residual in Y.

Next, we shall show that Theorem 3.1 is providing a unified framework to deduce Ekeland's variational principle [18], Borwein-Preiss's [6] variational principle and Deville-Godefroy-Zizler's Variational principle [15].

**Application 1.** As a first application we get the Ekeland's variational principle [18].

Let  $(X, \|.\|)$  be a Banach space. Assume that  $\varphi_1(x, y, \lambda) = \lambda x + (1 - \lambda) y$  and  $\varphi_2(x, y, \lambda, f) = \lambda f(x) + (1 - \lambda) f(y)$ . Then  $\varphi_1$  and  $\varphi_2$  satisfies  $(P_1), (P_2), (P_3)$  and  $(P_4)$ . Let

 $Y = \{f: X \longrightarrow \mathbb{R} : f \text{ convex, Lipschitz, } \ge 0, f \longrightarrow +\infty \text{ as } ||x|| \longrightarrow +\infty \}.$ 

We define on Y the metric  $\rho_Y$  such that for  $f, g \in Y$ ,

$$\rho_Y(f,g) = \mu_Y(f-g) = \sum_{n \ge 1} \frac{||f-g||_n}{2^n} + \sup\left\{\frac{|(f-g)(x) - (f-g)(y)|}{||x-y||}; x \ne y\right\}.$$

It is clear that  $(Y, \rho_Y)$  satisfies  $(P_5)$  and the conditions (i) and (ii) of Theorem 3.1. Also, the function  $\theta = ||x||$  satisfies the assertion (iii) of Theorem 3.1. Consequently we have the following:

**Corollary 3.2** Let (X, ||.||) be a Banach space, consider a lower semi-continuous bounded below function  $f : X \longrightarrow \mathbb{R} \cup \{+\infty\}$ . Then for each  $\varepsilon > 0$ , there exists  $x_0 \in X$  such that

$$f(x) + \varepsilon ||x - x_0|| \ge f(x_0).$$

Proof:

From Theorem 3.1, for each  $\varepsilon > 0$ , there exits  $g \in Y$  such that  $\mu_Y(g) < \varepsilon$  and f + g attains a strong minimum at  $x_0$ . Therefore, for all  $x \in X$ ,

$$f(x) + g(x) \ge f(x_0) + g(x_0)$$
 and  $\sum_{n\ge 1} \frac{||g||_n}{2^n} + \sup\left\{\frac{|g(x) - g(y)|}{||x - y||}; x \ne y\right\} < \varepsilon$ ,

which implies that

$$f(x) \geq f(x_0) + g(x_0) - g(x)$$
  
$$\geq f(x_0) - \varepsilon ||x - x_0||.$$

Application 2. Let  $(X, \|.\|)$  be a Banach space with smooth norm. Assume that  $\varphi_1(x, y, \lambda) = \lambda x + (1 - \lambda) y$  and  $\varphi_2(x, y, \lambda, f) = \lambda f(x) + (1 - \lambda) f(y)$ . Then  $\varphi_1$  and  $\varphi_2$  satisfies  $(P_1), (P_2), (P_3)$ , and  $(P_4)$ . Let

 $Y = \left\{ f \text{ is } C^1 \text{-smooth, convex,} \ge 0 \text{ and } f \longrightarrow +\infty \text{ as } ||x|| \longrightarrow +\infty \right\}.$ 

We define the metric  $\rho_Y$  in Y by:

$$\rho_Y(f,g) = \mu_Y(f-g) = \sum_{n \ge 1} \frac{||f-g||_n}{2^n} + ||(f-g)'||_\infty \text{ for all } f,g \in Y$$

where  $||f'||_{\infty} := \sup_{\|x\| \le 1} ||f'(x)||_{X^*}$  and the space  $(Y, \rho_Y)$  satisfies (*i*) and (*ii*) of Theorem 3.1 and so also  $(P_5)$ .

Let

$$\begin{aligned} h: & [0, +\infty[ & \longrightarrow [0, +\infty[ \\ t & \longmapsto \begin{cases} t^2 & \text{if } 0 \leq t \leq 1\\ 2t-1 & \text{if } t > 1. \end{cases} \end{aligned}$$

The function  $\theta(x) = h(||x||) \in Y$  satisfies the assertion *(iii)* of Theorem 3.1.

Therefore we have the Borwein-Preiss's variational principle [6, 27]:

**Corollary 3.3** Let (X, ||.||) be a Banach space with a smooth norm and consider a lower semi-continuous function  $f : X \longrightarrow \mathbb{R} \cup \{+\infty\}$  bounded from below. Then the set

 $\{g \in Y : f + g \text{ attains its strong minimum on } X\}$ 

is residual in Y.

Application 3. Let X be a Banach space admitting Lipschitz  $C^1$ -smooth bump function. According to a construction of Leduc [24], there exists a Lipschitz function  $d: X \longrightarrow \mathbb{R}$  which is  $C^1$ -smooth on  $X \setminus \{0\}$  and satisfies:

- i)  $d(\lambda x) = \lambda d(x)$  for all  $\lambda > 0$  and for all  $x \in X$ ,
- ii) there exists  $C \ge 1$  such that  $||x|| \le d(x) \le C ||x||$  for all  $x \in X$ .

Moreover the function  $d^2$  is  $C^1$ -smooth on all the space X.

Let  $\varphi_1(x, y, \lambda) = \lambda x + (1 - \lambda) y$  and  $\varphi_2(x, y, \lambda, f) = C^2[\lambda f(x) + (1 - \lambda) f(y)]$ . Then  $\varphi_1$  and  $\varphi_2$  satisfies  $(P_1), (P_2), (P_3)$  and  $(P_4)$ . Let  $\theta(x) = d^2(x)$ . We have

$$d^2 \left(\lambda x + (1 - \lambda) y\right) \le C^2 ||\lambda x + (1 - \lambda) y||^2.$$

Since the function  $||.||^2$  is convex we deduce that

$$d^{2}\left(\lambda x + (1-\lambda)y\right) \leq C^{2}\left(\lambda d^{2}(x) + (1-\lambda)d^{2}(y)\right).$$

That is the function  $d^2$  is a  $(\varphi_1, \varphi_2)$  -convex function.

Let

$$Y = \left\{ f \text{ a } (\varphi_1, \varphi_2) - \text{convex}, C^1 - \text{Lipschitz}, \ge 0 \text{ and } f \longrightarrow +\infty \text{ as } ||x|| \longrightarrow +\infty \right\}$$

and so the set Y satisfies  $(P_5)$ .

The metric  $\rho_Y$  on Y is such that, for  $f, g \in Y$ 

$$\rho_Y(f,g) = \mu_Y(f-g) = \sum_{n \ge 1} \frac{||f-g||_n}{2^n} + \sum_{n \ge 1} \frac{||(f-g)'||_n}{2^n}$$

where  $\left\|f'\right\|_n = \sup_{||x|| \leq n} \left\|f'(x)\right\|_{X^*}.$ 

In the other hand, let  $\theta(x) = d^2(x)$ . So that,

- i)  $\theta(0) = 0$
- ii)  $\mu_Y(\theta) < \infty$

iii) let 0 < k < 1. Hence, for all  $x \in X$  such that  $||x|| \ge k$  we have  $d^2(x) \ge ||x||^2 \ge k^2$ .

Therefore the function  $\theta \in Y$  and satisfies *(iii)* of Theorem 3.1.

Thus we have the following variational principle (for unbounded functions) of Deville-Godefroy-Zizler [14, 15, 16, 20]:

**Corollary 3.4** Let (X, ||.||) be a Banach space admitting a  $C^1$ -Lipschitz bump function and consider a lower semi-continuous bounded below function  $f : X \longrightarrow \mathbb{R} \cup \{+\infty\}$ . Then the set

 $\{g \in Y : f + g \text{ attains its strong minimum on } X\}$ 

is residual in Y.

Now, we are ready to give the proof of Theorem 3.1.

#### Proof of Theorem 3.1

Following the method of [15, 20], for  $n \in \mathbb{N} \setminus \{0\}$ , we let

$$G_n = \{g \in Y : \exists x_0 \in X, (f+g)(x_0) < \inf\{(f+g)(x) : ||x-x_0|| \ge 1/n\}\}.$$

Claim 1. We claim that  $G_n$  is open for each n. Indeed, let  $n \in \mathbb{N}$  and  $g \in G_n$ . So that there is  $x_0$  in X such that

$$(f+g)(x_0) < \inf \{(f+g)(x) : ||x-x_0|| \ge 1/n\}.$$

Let  $0 < \varepsilon < 1$  such that

$$(f+g)(x_0) + 2\varepsilon < \inf \{ (f+g)(x) : ||x-x_0|| \ge 1/n \}.$$
(1)

Let  $A = C(f+g)(x_0) + C(g(0) - \inf(f)) + (2C+3)\varepsilon$ , where C is given by  $(P_3)$ . Since  $g \in Y$ , g goes to  $+\infty$  as ||x|| goes to  $+\infty$ . This means that, there is k in  $\mathbb{N}$  such that  $k > ||x_0||$  and g(x) > A whenever  $||x|| \ge k$ . This is equivalent to say that

$$g(x) > C(f+g)(x_0) + C(g(0) - \inf(f)) + (2C+3)\varepsilon$$
 whenever  $||x|| \ge k$ . (2)

Let  $h \in Y$  such that  $\rho_Y(h,g) < \frac{\varepsilon}{2^k}$ . We have

$$\sum_{n\geq 1} \frac{\|h-g\|_n}{2^n} \leq \rho_Y(h,g) = \mu_Y(h-g) < \frac{\varepsilon}{2^k}.$$

Thus

$$\frac{\|h-g\|_k}{2^k} < \frac{\varepsilon}{2^k}.$$

So that

$$|h(x) - g(x)| < \varepsilon$$
 whenever  $||x|| \le k$ , (3)

in particular

$$|h(x_0) - g(x_0)| < \varepsilon.$$
(4)

Combining (2) with (3) we obtain that

$$h(x) > C(f+g)(x_0) + C(g(0) - \inf(f)) + (2C+2)\varepsilon > 0$$
 whenever  $||x|| = k$ .

Since  $C \ge 1$  and  $h \ge 0$ , we deduce that for ||x|| = k

$$h(x) \ge \frac{h(x)}{C} > (f+g)(x_0) + (g(0) - \inf(f)) + (2 + (2/C))\varepsilon.$$
(5)

In the first hand, let  $y \in X$  such that ||y|| > k. Then, there exist  $\lambda > 1$  and  $x \in X$  with ||x|| = k, such that  $y = \lambda x$ . By using Lemma 2.2, we deduce that

$$h(y) - h(0) \ge \frac{\lambda}{C} (h(x) - Ch(0)) \ge \frac{1}{C} (h(x) - Ch(0)) = \frac{h(x)}{C} - h(0).$$

Combining this with (5) we show for  $||y|| \ge k$  that,

$$h(y) - h(0) > (f + g)(x_0) + g(0) - \inf f + \left(2 + \frac{2}{C}\right)\varepsilon - h(0).$$
(6)

Combining the fact that  $h \ge 0, (6), (3)$  and (4) we obtain for all  $x \in X$  such that  $||x|| \ge k$ :

$$(f+h)(x) \geq \inf (f) + h (x) \geq \inf (f) + h (x) - h (0) > \inf (f) + (f+g) (x_0) + g (0) - \inf (f) + \left(2 + \frac{2}{C}\right) \varepsilon - h (0) > (f+g) (x_0) + \left(1 + \frac{2}{C}\right) \varepsilon > (f+h) (x_0) + \frac{2}{C} \varepsilon > (f+h) (x_0).$$

Therefore for all  $x \in X$  such that  $||x|| \ge k$ , we have

$$(f+h)(x) > (f+h)(x_0).$$

In other hand, if  $||x|| \leq k$  and  $||x - x_0|| \geq 1/n$ , and combining (4), (1) and (3) we obtain that

$$(f+h)(x_0) < (f+g)(x_0) + \varepsilon$$
  

$$\leq \inf \{(f+g)(x) : ||x-x_0|| \ge 1/n\} - 2\varepsilon + \varepsilon$$
  

$$\leq (f+g)(x) - \varepsilon$$
  

$$< (f+h)(x).$$

Then for all x such that  $||x - x_0|| \ge 1/n$  we have

$$(f+h)(x_0) < (f+h)(x).$$

Hence  $h \in G_n$  and  $G_n$  is open.

Claim 2. We confirm that  $G_n$  is dense in Y. Indeed, let  $g \in Y$  and  $0 < \varepsilon < 1$ . Let c > 0 be such that

$$(f+g)(x) > \inf (f+g) + 1$$
 whenever  $||x|| > c$ .

Let  $1 > \delta > 0$  be such that  $\delta(\mu_Y(\theta) + \|\theta\|_c) < \varepsilon$ . Let  $x_0 \in X$  be such that

$$(f+g)(x_0) < \inf(f+g) + \frac{\delta}{n^2}.$$
 (6)

Since  $\frac{\delta}{n^2} < 1$ , we deduce that

 $\|x_0\| \le c. \tag{7}$ 

Let  $h(x) = \delta \theta(x - x_0)$ . Now Corollary 2.5 ensure that h is a  $(\varphi_1, \varphi_2)$  -convex function in F.

From the hypothesis (iii) of Theorem 3.1 and (7), we get

$$\rho_Y(h,0) = \mu_Y(h) = \delta\mu_Y(\theta(.-x_0)) \le \delta\mu_Y(\theta) + \delta \|\theta\|_{\|x_0\|} \le \delta (\mu_Y(\theta) + \|\theta\|_c) < \varepsilon.$$

Now if  $||x - x_0|| \ge 1/n$ , and by (*iii*) of Theorem 3.1 we deduce that,

$$h(x) = \delta \theta(x - x_0) \ge \frac{\delta}{n^2}.$$

By using (6), we get

$$\inf \{f + g + h : \|x - x_0\| \ge 1/n\} \ge \inf \{f + g : \|x - x_0\| \ge 1/n\} + \frac{\delta}{n^2}$$
$$\ge \inf \{f + g\} + \frac{\delta}{n^2}$$
$$> (f + g)(x_0) - \frac{\delta}{n^2} + \frac{\delta}{n^2}.$$

Moreover  $h(x_0) = \delta \theta(0) = 0$ , then,

$$\inf \{f + g + h : \|x - x_0\| \ge 1/n\} > (f + g)(x_0) = (f + g + h)(x_0).$$

Thus  $(g+h) \in G_n$  and  $G_n$  is a dense subset in Y.

Therefore the set  $\bigcap_{n\geq 1} G_n$  is residual in Y. Following the proof of [15], we can show f + g attains its strong minimum for each  $g \in \bigcap_{n\geq 1} G_n$ .

## References

 J.P. Aubin, Lipshits behavior of solutions to convex minimization problems, Oper. Res. 9 (1) (1984) 87 - 111.

- [2] J.P. Aubin, I. Ekeland, Applied nonlinear analysis, Wiley-Interscience, New York, 1984.
- [3] E. Bishop, R.R. Phelps, The support functional of convex set. Convexity, Klee ed. Proc. Sympos. Pure Math., 9, A.M.S., Providence, R.I, 1963
- [4] E. Bishop, R.R. Phelps, The support cones in Banach spaces and their applications, Adv. In Math. 13 (1974) 1 - 19.
- [5] N. Bonic, J. Frampton, Smooth functions on Banach manifolds, J. Math. Mech. 15 (1966) 877 - 898.
- [6] J. Borwein, D. Preiss, A smooth variational principle with applications to subdifferentiability and to differentiability of convex functions, Trans. Amer. Math. Soc. 303 (1987) 517 - 527.
- H. Brezis, F. Browder, A general ordering principle in nonlinear functional analysis, Adv. In Math. 21 (1976) 355 - 364.
- [8] F.E. Browder, Normal solvability and the Fredholm altertive for mapping into infinite dimensional manifolds, J. Funct. Anal. 8 (1971) 250 274.
- [9] F.H. Clarke, Optimization and nonsmooth analysis, Wiley-Interscience, New York, 1983.
- [10] J. Danes, A geometric theorem useful in non-linear functional analysis, Boll. Un. Math. Ital. 6 (1972) 369 - 375.
- [11] R. Deville, Stability of subdifferentials of nonconvex functions in Banach spaces, Set-Valued Anal. 2 (1994) 141 - 157.
- [12] R. Deville, E.M. El Haddad, The subdifferential of the sum of two functions in Banach spaces. II. Second order case, J. Convex Anal. 3 (2) (1995) 295 - 308.
- [13] R. Deville, E.M. El Haddad, The subdifferential of the sum of two functions in Banach spaces. I. First order case, Bull. Austral. Math. Soc. 51 (1996) 235 -248.

- [14] R. Deville, G. Godefroy, V. Zizler, Un principe variationel utilisant des fonctions bosses, C. R. Acad. Sci. (Paris). Ser.I, 312 (1991) 281 - 286.
- [15] R. Deville, G. Godefroy, V. Zizler, A smooth variational principle with applications to Hamilton-Jacobi equations in infinite dimensions, J. Funct. Anal. 111 (1993) 197 - 212.
- [16] R. Deville, G. Godefroy, V. Zizler, Smoothness and renormings in Banach spaces, Longman Scientific and Technical, Pitman Monographs and surveys in Pure and Applied Math., 64, 1993
- [17] R. Deville, A. Maaden, Smooth variational principles in Radon-Nikodym spaces, Bull. Austral. Math. Soc. 60, (1999) 109 - 118.
- [18] I. Ekeland, On the variational principle, J. Math. Anal. Appl. 47 (1974) 323 -353.
- [19] I. Ekeland, Nonconvex minimization problems, Bull. Amer. Math. Soc. (New Series)I (1979) 443 - 474.
- [20] M. Fabian, P. Hajeik, J. Vanderwerff, On smooth variational principles in Banach spaces, J. Math. Anal. Appl. 197 (1) (1996) 153 - 172.
- [21] P. Georgiev, The strong Ekeland variational principle, the strong drop theorem and applications, J. Math. Anal. Appl. 131 (1988) 1 - 21.
- [22] P. Georgiev, D. Kutzarova, A. Maaden, On the smooth drop property, Nonlinear Anal. Theo. Meth. Appl. 26 (3) (1996) 595 - 602.
- [23] R. Haydon, A counterexample in several questions about scattered compact spaces, Bull. London. Math. Soc. 22 (1990) 261 - 268.
- [24] M. Leduc, Densité de certaines familles d'hyperplans tangents, C. R. Acd. Sci (Paris). Ser. A. 270 (1970) 326 - 328.
- [25] A. Maaden, Théorème de la goutte lisse, Rocky Mountain J. Math. 25 (3) (1995) 1093 - 1101.

- [26] J.P. Penot, The drop theorem, the petal theorem and the Ekeland's variational principle, J. Nonlin. Anal. Appl. 10 (9) (1986) 813 - 822.
- [27] R.R. Phelps, Convex functions, monotone operators and differentiability, Lecture notes in Math., 1364, 1991.
- [28] R. Pini, C. Singh,  $(\varphi_1, \varphi_2)$  -convexity, Optimization, 40 (2) (1997) 103 120.
- [29] R. Pini, C. Singh, (φ<sub>1</sub>, φ<sub>2</sub>) –optimality and duality under differentiability, Optimization. 41 (2) (1997) 101 - 116.
- [30] H. Riahi, Application du principe variationnel d'Ekeland à l'existence d'optima de Pareto, Extr. Math. 7 (1) (1992) 42 - 45.
- [31] C. Stegall, Optimization of functions on certain subset of Banach spaces, Math. Ann. 236 (1978) 171 - 176.