

Asymptotic formulas for general colored partition functions

Yong-Gao Chen*, Ya-Li Li

School of Mathematical Sciences and Institute of Mathematics,

Nanjing Normal University, Nanjing 210023, P. R. China

Abstract

In 1917, Hardy and Ramanujan obtained the asymptotic formula for the classical partition function $p(n)$. The classical partition function $p(n)$ has been extensively studied. Recently, Luca and Ralaivaosaona obtained the asymptotic formula for the square-root function. Many mathematicians have paid much attention to congruences on some special colored partition functions. In this paper, we investigate the general colored partition functions. Given positive integers $1 = s_1 < s_2 < \cdots < s_k$ and l_1, l_2, \dots, l_k . Let $g(\mathbf{s}, \mathbf{l}, n)$ be the number of l -colored partitions of n with l_i of the colors appearing only in multiplies of s_i ($1 \leq i \leq k$), where $l = l_1 + \cdots + l_k$. By using the elementary method we obtain an asymptotic formula for the partition function $g(\mathbf{s}, \mathbf{l}, n)$ with an explicit error term.

Keyword: colored partition; partition function; asymptotic formula; Gaussian integral

2010 Mathematics Subject Classifications: 11P82;11N37;05A17

*Corresponding author, ygchen@njnu.edu.cn(Y.-G. Chen)

1 Introduction

Let $p(n)$ denote the number of partitions of n , i.e.

$$n = a_1 + \cdots + a_k$$

with integers $1 \leq a_1 \leq \cdots \leq a_k$. The generating function of $p(n)$ is

$$f(z) = 1 + \sum_{n=1}^{\infty} p(n)z^n = \prod_{n=1}^{\infty} \frac{1}{1 - z^n}. \quad (1.1)$$

Ramanujan [16] obtained many congruent identity for $p(n)$. Hardy and Ramanujan [9] and Uspensky [18] independently proved that

$$p(n) \sim \frac{1}{4n\sqrt{3}} \exp\left(\pi\sqrt{\frac{2n}{3}}\right). \quad (1.2)$$

An elementary proof for this formula is given by Erdős [8] with no explicit constant $(4\sqrt{3})^{-1}$. Lehmer [12] gave the series for the partition function $p(n)$. Odlyzko [15, (1.6)] gave an asymptotic formula for $p(n)$ with an explicit error term. That is,

$$p(n) = \frac{1 + O(n^{-1/2})}{4n\sqrt{3}} \exp\left(\pi\sqrt{\frac{2n}{3}}\right). \quad (1.3)$$

The partition function $p(n)$ has a long history and generates varieties.

Recently, Luca and Ralaivaosaona [14] obtained the asymptotic formula for the square-root function $q(n)$, which is defined to be the number of solutions of

$$n = [\sqrt{a_1}] + [\sqrt{a_2}] + \cdots + [\sqrt{a_k}]$$

with integers $1 \leq a_1 \leq a_2 \leq \cdots \leq a_k$. For related results, one may refer to Balasubramanian and Luca [2] and Chen and Li [7] and [13].

Let $p_k(n)$ be the number of 2-color partitions of n where one of the colors appears only in parts that are multiples of k . The generating function of $p_k(n)$ is

$$1 + \sum_{n=1}^{\infty} p_k(n)z^n = \prod_{n=1}^{\infty} \frac{1}{(1 - z^n)(1 - z^{kn})}.$$

Chan [3], Kim [11] and Sinick [17] studied some results of the case $k = 2$. Recently, Ahmed, Baruah and Dastidar [1] and Chern [6] obtained many congruences of $p_k(n)$ for some k .

By further analogy, Chan and Cooper [4] and Chen [5] considered a special partition function $c(n)$ which is the number of 4-colored partitions of n with two of the colors appearing only in multiplies of 3. The generating function of $c(n)$ is

$$1 + \sum_{n=1}^{\infty} c(n)z^n = \prod_{n=1}^{\infty} \frac{1}{(1-z^n)^2(1-z^{3n})^2}.$$

In this paper, we focus on the asymptotic formula for the general colored partition functions.

Given integers $1 = s_1 < s_2 < \dots < s_k$ and $\ell_1, \ell_2, \dots, \ell_k$. Let $g(\mathbf{s}, \mathbf{l}, n)$ be the number of ℓ -colored partitions of n with ℓ_i of the colors appearing only in multiplies of s_i ($1 \leq i \leq k$), where $\ell = \ell_1 + \dots + \ell_k$. Write

$$\mathbf{s} = (s_1, s_2, \dots, s_k)$$

and

$$\mathbf{l} = (\ell_1, \ell_2, \dots, \ell_k).$$

We call $g(\mathbf{s}, \mathbf{l}, n)$ the (\mathbf{s}, \mathbf{l}) -colored partition function. For convenience, we define $g(\mathbf{s}, \mathbf{l}, 0) = 1$ and $g(\mathbf{s}, \mathbf{l}, n) = 0$ for all $n < 0$. The generating function of $g(\mathbf{s}, \mathbf{l}, n)$ is

$$\sum_{n=0}^{\infty} g(\mathbf{s}, \mathbf{l}, n)x^n = \prod_{n=1}^{\infty} \frac{1}{(1-z^{s_1 n})^{\ell_1} \dots (1-z^{s_k n})^{\ell_k}}. \quad (1.4)$$

In this paper, the following result is proved.

Theorem 1.1. *For any given $\varepsilon > 0$ (small), we have*

$$g(\mathbf{s}, \mathbf{l}, n) = c(\mathbf{s}, \mathbf{l})n^{d(\mathbf{l})} \exp\left(\pi\sqrt{\frac{2a(\mathbf{s}, \mathbf{l})n}{3}}\right) + O\left(n^{d(\mathbf{l})-\frac{1}{4}+\varepsilon} \exp\left(\pi\sqrt{\frac{2a(\mathbf{s}, \mathbf{l})n}{3}}\right)\right),$$

where

$$a(\mathbf{s}, \mathbf{l}) = \sum_{i=1}^k \frac{\ell_i}{s_i}, \quad d(\mathbf{l}) = -\frac{3}{4} - \frac{1}{4}(\ell_1 + \dots + \ell_k),$$

$$c(\mathbf{s}, \mathbf{l}) = 2^{-(3\ell_1 + \dots + 3\ell_k + 5)/4} 3^{-(\ell_1 + \dots + \ell_k + 1)/4} a(\mathbf{s}, \mathbf{l})^{(\ell_1 + \dots + \ell_k + 1)/4} s_1^{\ell_1/2} \dots s_k^{\ell_k/2}.$$

Remark 1.2. *By employing the Tauberian theorem of Ingham [10], under its form in the special case, it is possible to get an asymptotic formula for $g(\mathbf{s}, \mathbf{l}, n)$ without the error term. We do not intend to give the details here.*

Let $k = 2, s_1 = 1, s_2 = 3$ and $\ell_1 = \ell_2 = 2$. By Theorem 1.1, we can give an asymptotic formula for $c(n)$, that is,

$$c(n) = \frac{1}{3\sqrt{6}n^{7/4}} \exp\left(\frac{4}{3}\pi\sqrt{n}\right) + O\left(n^{-2+\varepsilon} \exp\left(\frac{4}{3}\pi\sqrt{n}\right)\right).$$

We believe that there are many congruences for the (\mathbf{s}, \mathbf{l}) -colored partition functions as many known various partition functions.

2 The main ingredients

For convenience, let

$$a = a(\mathbf{s}, \mathbf{l}) = \sum_{i=1}^k \frac{\ell_i}{s_i}, \quad g(n) = g(\mathbf{s}, \mathbf{l}, n), \quad c_1 = \pi\sqrt{\frac{2}{3}}, \quad c_2 = \frac{1}{4\sqrt{3}}. \quad (2.1)$$

Then (1.3) becomes

$$p(n) = \frac{c_2 + O(n^{-1/2})}{n} \exp(c_1\sqrt{n}). \quad (2.2)$$

If $k = 1$ and $\ell_1 = 1$, then $g(n) = p(n)$. In this case, Theorem 1.1 follows from (2.2). Now we assume that $k + \ell_1 \geq 3$. So $as_k > 1$.

Lemma 2.1. *We have*

$$g(n) = \sum_{\mathbf{u} \in U_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}),$$

where U_n is the set of all $\ell_1 + \dots + \ell_k$ tuples

$$\mathbf{u} = (u_{1,1}, \dots, u_{1,\ell_1}, \dots, u_{k,1}, \dots, u_{k,\ell_k})$$

of nonnegative integers with

$$s_1 u_{1,1} + \dots + s_1 u_{1,\ell_1} + \dots + s_k u_{k,1} + \dots + s_k u_{k,\ell_k} = n.$$

Proof. By (1.1) and (1.4), we have

$$\begin{aligned}
\sum_{n=0}^{\infty} g(n)z^n &= \left(\sum_{n=0}^{\infty} p(n)z^{s_1 n} \right)^{\ell_1} \cdots \left(\sum_{n=0}^{\infty} p(n)z^{s_k n} \right)^{\ell_k} \\
&= \sum_{u_{1,1}=0}^{\infty} p(u_{1,1})z^{s_1 u_{1,1}} \cdots \sum_{u_{1,\ell_1}=0}^{\infty} p(u_{1,\ell_1})z^{s_1 u_{1,\ell_1}} \cdots \\
&\quad \sum_{u_{k,1}=0}^{\infty} p(u_{k,1})z^{s_k u_{k,1}} \cdots \sum_{u_{k,\ell_k}=0}^{\infty} p(u_{k,\ell_k})z^{s_k u_{k,\ell_k}} \\
&= \sum_{n=0}^{\infty} \left(\sum_{\mathbf{u} \in U_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) \right) z^n.
\end{aligned}$$

Now Lemma 2.1 follows immediately. \square

We will divide U_n into two parts U'_n and U''_n which will be given later such that $u_{i,j} \rightarrow +\infty$ as $n \rightarrow +\infty$ for any $\mathbf{u} = (u_{1,1}, \dots, u_{1,\ell_1}, \dots, u_{k,1}, \dots, u_{k,\ell_k}) \in U'_n$. By Lemma 2.1, we have

$$g(n) = \sum_{\mathbf{u} \in U'_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) + \sum_{\mathbf{u} \in U''_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}). \quad (2.3)$$

We expect that U'_n contributes to $g(n)$ the main term and U''_n contributes to $g(n)$ the remainder term.

By (2.2), we have

$$\begin{aligned}
&\sum_{\mathbf{u} \in U'_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) \\
&= \sum_{\mathbf{u} \in U'_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \frac{c_2 + O(u_{i,j}^{-1/2})}{u_{i,j}} \exp(c_1 \sqrt{u_{i,j}}) \\
&= \sum_{\mathbf{u} \in U'_n} \frac{c_2^{\ell_1 + \dots + \ell_k} + O(\sum u_{i,j}^{-1/2})}{\prod u_{i,j}} \exp\left(c_1 \sum_{i,j} \sqrt{u_{i,j}}\right) \quad (2.4)
\end{aligned}$$

and

$$\sum_{\mathbf{u} \in U''_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) \ll \sum_{\mathbf{u} \in U''_n} \exp\left(c_1 \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \sqrt{u_{i,j}}\right). \quad (2.5)$$

Since the function $\exp(x)$ increases rapidly, it infers from (2.4) and (2.5) that the maximal value of $\sum_{i,j} \sqrt{u_{i,j}}$ with $\mathbf{u} \in U_n$ gives the main contribution to $g(n)$. Now we find the maximal value of $\sum_{i,j} \sqrt{u_{i,j}}$ with $\mathbf{u} \in U_n$. By the Cauchy-Schwarz inequality, for any $\mathbf{u} \in U_n$, we have

$$\begin{aligned} \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \sqrt{u_{i,j}} &= \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \frac{1}{\sqrt{s_i}} \sqrt{s_i u_{i,j}} \\ &\leq \left(\sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \frac{1}{s_i} \right)^{1/2} \left(\sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} s_i u_{i,j} \right)^{1/2} \\ &= \left(\sum_{i=1}^k \frac{\ell_i}{s_i} \right)^{1/2} \sqrt{n}, \end{aligned}$$

where the equality holds if and only if

$$u_{i,j} = \frac{n}{s_i^2 a}, \quad 1 \leq i \leq k, 1 \leq j \leq \ell_i,$$

where

$$a = \sum_{i=1}^k \frac{\ell_i}{s_i}.$$

Let

$$v_{i,j} = \frac{n}{s_i^2 a}, \quad 1 \leq i \leq k, 1 \leq j \leq \ell_i.$$

It is clear that

$$\sum_{i,j} s_i v_{i,j} = \sum_{i=1}^k \frac{\ell_i n}{s_i a} = n.$$

Now we have proved that the maximal value of $\sum_{i,j} \sqrt{u_{i,j}}$ is obtained if and only if

$$\mathbf{u} = (u_{1,1}, \dots, u_{1,\ell_1}, \dots, u_{k,1}, \dots, u_{k,\ell_k}) = (v_{1,1}, \dots, v_{1,\ell_1}, \dots, v_{k,1}, \dots, v_{k,\ell_k}) := \mathbf{v}.$$

Basing on the above intuition that the maximal value of $\sum_{i,j} \sqrt{u_{i,j}}$ with $\mathbf{u} \in U_n$ gives the main contribution to $g(n)$, we take U'_n to be the set of $\mathbf{u} \in U_n$ for which every $u_{i,j}$ is near to $v_{i,j}$ and U''_n to be the set of $\mathbf{u} \in U_n$ for which $u_{i,j}$ is far from $v_{i,j}$ for some pair i, j . Since

$$s_1 u_{1,1} + \dots + s_1 u_{1,\ell_1} + \dots + s_k u_{k,1} + \dots + s_k u_{k,\ell_k} = n,$$

it is enough to take U'_n to be those $\mathbf{u} \in U_n$ for which $u_{i,j}$ is near to $v_{i,j}$ for all i, j with $i + j > 2$ and U''_n to be those $\mathbf{u} \in U_n$ for which $u_{i,j}$ is far from $v_{i,j}$ for some pair i, j with $i + j > 2$. Now we give explicit U'_n and U''_n .

We appoint a real number η such that

$$\frac{3}{4} < \eta < \min \left\{ \frac{5}{6}, \frac{3}{4} \frac{\ell_1 + \cdots + \ell_k - 1}{\ell_1 + \cdots + \ell_k - 2} \right\}.$$

Since the right hand side is more than $3/4$, it follows that such η exists. Since $3/4 < \eta < 5/6$, we have $2\eta - 3/2 > 0$ and $3\eta - 5/2 < 0$.

Let

$$U'_n = \{\mathbf{u} \in U_n : |u_{i,j} - v_{i,j}| < v_{i,j}^\eta \text{ for all pairs } i, j \text{ with } i + j > 2\}$$

and

$$U''_n = \{\mathbf{u} \in U_n : |u_{i,j} - v_{i,j}| \geq v_{i,j}^\eta \text{ for some pair } i, j \text{ with } i + j > 2\}.$$

We hope that U'_n contributes to $g(n)$ the main term and U''_n contributes to $g(n)$ the remainder term. These will be proved in the next section.

3 Preliminary Lemmas

Firstly we prove that U''_n contributes to $g(n)$ the remainder term.

Lemma 3.1. *We have*

$$\sum_{\mathbf{u} \in U''_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) = O\left(\exp\left(c_1 \sqrt{an} - c_3 n^{2\eta-3/2}\right)\right),$$

where c_1 is given by (2.1) and c_3 is a positive constant.

Proof. Let $\mathbf{u} \in U''_n$. Without loss of generality, we assume that

$$|u_{k,\ell_k} - v_{k,\ell_k}| \geq v_{k,\ell_k}^\eta. \tag{3.1}$$

Let

$$u_{k,\ell_k} = v_{k,\ell_k} + \alpha v_{k,\ell_k}.$$

Then $|\alpha| \geq v_{k,\ell_k}^{-(1-\eta)}$. Since $\mathbf{u} \in U_n''$, it follows that $n \geq s_k u_{k,\ell_k}$. Noting that $n = s_k^2 a v_{k,\ell_k}$, we have

$$v_{k,\ell_k} + \alpha v_{k,\ell_k} = u_{k,\ell_k} \leq \frac{n}{s_k} = s_k a v_{k,\ell_k}.$$

It follows that $1 + \alpha \leq s_k a$. So $(s_k a - 1)^{-1} \alpha \leq 1$.

By the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \sqrt{u_{i,j}} &= \sqrt{u_{k,\ell_k}} + \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i, (i,j) \neq (k,\ell_k)}} \frac{1}{\sqrt{s_i}} \sqrt{s_i u_{i,j}} \\ &\leq \sqrt{u_{k,\ell_k}} + \left(\sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i, (i,j) \neq (k,\ell_k)}} \frac{1}{s_i} \right)^{1/2} \left(\sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i, (i,j) \neq (k,\ell_k)}} s_i u_{i,j} \right)^{1/2} \\ &= \sqrt{u_{k,\ell_k}} + \left(a - \frac{1}{s_k} \right)^{1/2} \sqrt{n - s_k u_{k,\ell_k}} \\ &= (v_{k,\ell_k} + \alpha v_{k,\ell_k})^{1/2} \\ &\quad + \left(a - \frac{1}{s_k} \right)^{1/2} (s_k^2 a v_{k,\ell_k} - s_k v_{k,\ell_k} - \alpha s_k v_{k,\ell_k})^{1/2} \\ &= f(\alpha) \sqrt{v_{k,\ell_k}}, \end{aligned}$$

where

$$f(x) = (1+x)^{1/2} + (s_k a - 1) (1 - (s_k a - 1)^{-1} x)^{1/2}.$$

Since

$$f'(x) = \frac{1}{2}(1+x)^{-1/2} - \frac{1}{2}(1 - (s_k a - 1)^{-1} x)^{-1/2},$$

it follows that $f'(x) > 0$ for $x < 0$ and $f'(x) < 0$ for $x > 0$. By $|\alpha| \geq v_{k,\ell_k}^{-(1-\eta)}$, we have

$$f(\alpha) \leq \max\{f(-v_{k,\ell_k}^{-(1-\eta)}), f(v_{k,\ell_k}^{-(1-\eta)})\}.$$

For $\beta \in \{-v_{k,\ell_k}^{-(1-\eta)}, v_{k,\ell_k}^{-(1-\eta)}\}$, we have

$$\begin{aligned} f(\beta) &= 1 + \frac{1}{2}\beta - \frac{1}{8}\beta^2 + O(\beta^3) \\ &\quad + (s_k a - 1) \left(1 - \frac{1}{2}(s_k a - 1)^{-1}\beta - \frac{1}{8}(s_k a - 1)^{-2}\beta^2 + O(\beta^3) \right) \\ &= s_k a - \frac{s_k a}{8(s_k a - 1)}\beta^2 + O(\beta^3) \\ &= s_k a - \frac{s_k a}{8(s_k a - 1)} v_{k,\ell_k}^{-2(1-\eta)} + O(v_{k,\ell_k}^{-3(1-\eta)}). \end{aligned}$$

Hence

$$\begin{aligned} \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \sqrt{u_{i,j}} &\leq s_k a \sqrt{v_{k,\ell_k}} - \frac{s_k a}{8(s_k a - 1)} v_{k,\ell_k}^{-2(1-\eta)+1/2} + O(v_{k,\ell_k}^{-3(1-\eta)+1/2}) \\ &\leq \sqrt{an} - \delta n^{2\eta-3/2} \end{aligned}$$

for all sufficiently large integers n , where δ is a positive constant. Thus, noting that $s_k^2 a v_{k,\ell_k} = n$, for any $\mathbf{u} \in U_n''$, and by (2.5), we have

$$\begin{aligned} &\sum_{\mathbf{u} \in U_n''} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) \\ &\ll \sum_{\mathbf{u} \in U_n''} \exp\left(c_1 \sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} \sqrt{u_{i,j}}\right) \\ &\ll \sum_{\mathbf{u} \in U_n''} \exp\left(c_1 \sqrt{an} - c_1 \delta n^{2\eta-3/2}\right) \\ &\ll (n+1)^{\ell_1+\dots+\ell_k-1} \exp\left(c_1 \sqrt{an} - c_1 \delta n^{2\eta-3/2}\right) \\ &\ll \exp\left(c_1 \sqrt{an} - c_3 n^{2\eta-3/2}\right), \end{aligned}$$

for all sufficiently large integers n , where c_3 is a positive constant. This completes the proof of Lemma 3.1. \square

Lemma 3.1 deals with those \mathbf{u} for which $u_{i,j}$ is far from $v_{i,j}$ for some pair i, j with $i+j > 2$. These \mathbf{u} contribute to $g(n)$ with the remainder. Now we deal with all \mathbf{u} for which $u_{i,j}$ is near to $v_{i,j}$ for every pair i, j with $i+j > 2$.

Lemma 3.2. *We have*

$$\sum_{\mathbf{u} \in U_n'} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) = \frac{c_2^{\ell_1+\dots+\ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \sum_{\mathbf{u} \in U_n'} \exp\left(c_1 \sum_{i,j} \sqrt{u_{i,j}}\right),$$

where c_1 and c_2 are given by (2.1).

Proof. Recall that

$$U_n' = \{\mathbf{u} \in U_n : |u_{i,j} - v_{i,j}| < v_{i,j}^\eta \text{ for all } i+j > 2\}.$$

Let $\mathbf{u} \in U_n'$. By $\sum_{i,j} s_i v_{i,j} = n$ and $|u_{i,j} - v_{i,j}| < v_{i,j}^\eta$, we have

$$|u_{1,1} - v_{1,1}| = \left| \sum_{i+j>2} s_i (u_{i,j} - v_{i,j}) \right| < \sum_{i+j>2} s_i v_{i,j}^\eta < \sum_{i+j>2} s_i n^\eta.$$

Thus $u_{i,j} \rightarrow +\infty$ as $n \rightarrow +\infty$ for $i + j \geq 2$. Moreover, by $v_{i,j} = O(n)$, we have $|u_{i,j} - v_{i,j}| \ll n^\eta$ for all i, j with $i + j \geq 2$. Thus $u_{i,j} = O(n)$ all i, j with $i + j \geq 2$. Hence

$$\frac{1}{u_{i,j}} = \frac{1}{v_{i,j} + O(n^\eta)} = \frac{1}{v_{i,j}(1 + O(n^{\eta-1}))} = \frac{1}{v_{i,j}}(1 + O(n^{\eta-1})), \quad i + j \geq 2$$

and

$$\sum_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} u_{i,j}^{-1/2} = O(n^{-1/2}).$$

Noting that

$$\prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} (1 + O(n^{\eta-1})) = 1 + O(n^{\eta-1}),$$

from the above arguments, (2.4) and $\frac{3}{4} < \eta < \frac{5}{6}$, we have

$$\begin{aligned} & \sum_{\mathbf{u} \in U'_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) \\ &= \frac{c_2^{\ell_1 + \dots + \ell_k} + O(n^{-1/2})}{\prod v_{i,j}} (1 + O(n^{\eta-1})) \sum_{\mathbf{u} \in U'_n} \exp\left(c_1 \sum_{i,j} \sqrt{u_{i,j}}\right) \\ &= \frac{c_2^{\ell_1 + \dots + \ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \sum_{\mathbf{u} \in U'_n} \exp\left(c_1 \sum_{i,j} \sqrt{u_{i,j}}\right). \end{aligned}$$

□

The following two lemmas devote to convert summations on integral variables into integrals.

Lemma 3.3. *Suppose that f is a function on $[a, b]$ such that f' exists with m zero points on (a, b) . Then*

$$\sum_{a \leq n \leq b} f(n) = \int_a^b f(x) dx + O\left((m+1) \max_{a \leq x \leq b} |f(x)|\right).$$

Proof. We divide $[a, b]$ into $m + 1$ intervals $[a_0, a_1], [a_1, a_2], \dots, [a_m, a_{m+1}]$ such that $f' \neq 0$ on (a_i, a_{i+1}) . Given $0 \leq i \leq m$. Without loss of generality, we assume that $f' > 0$ on (a_i, a_{i+1}) . For any integer $n \in (a_i + 1, a_{i+1} - 1)$, we have

$$\int_{n-1}^n f(x) dx \leq f(n) \leq \int_n^{n+1} f(x) dx.$$

Summing on n , we have

$$\left| \sum_{a_i \leq n \leq a_{i+1}} f(n) - \int_{a_i}^{a_{i+1}} f(x) dx \right| = O\left(\max_{a_i \leq x \leq a_{i+1}} |f(x)|\right) = O\left(\max_{a \leq x \leq b} |f(x)|\right).$$

Then

$$\begin{aligned} \left| \sum_{a \leq n \leq b} f(n) - \int_a^b f(x) dx \right| &\leq \sum_{i=0}^m \left| \sum_{a_i \leq n \leq a_{i+1}} f(n) - \int_{a_i}^{a_{i+1}} f(x) dx \right| \\ &= O\left((m+1) \max_{a \leq x \leq b} |f(x)|\right). \end{aligned}$$

This completes the proof. □

Lemma 3.4. *Suppose that f is a function on*

$$D = \{(x_1, \dots, x_s) : a_i \leq x_i \leq a_i + b_i, 1 \leq i \leq s\}$$

such that, for each i and fixed $\{x_1, \dots, x_s\} \setminus \{x_i\}$, the partial derivative $\frac{\partial}{\partial x_i} f$ exists with at most m_i zero points on $[a_i, a_i + b_i]$. Then

$$\begin{aligned} &\sum_{(n_1, \dots, n_s) \in D \cap \mathbb{Z}^s} f(n_1, \dots, n_s) \\ &= \int_D \cdots \int f(x_1, \dots, x_s) dx_1 \cdots dx_s + O\left(\sum_{1 \leq i \leq s} \frac{m_i + 1}{b_i} BM\right), \end{aligned}$$

where $B = b_1 b_2 \cdots b_s$ and $M = \max_D |f(x_1, \dots, x_s)|$.

Proof. Let $c_i = a_i + b_i (1 \leq i \leq s)$. By Lemma 3.3, we have

$$\begin{aligned}
& \sum_{(n_1, \dots, n_s) \in D \cap \mathbb{Z}^s} f(n_1, \dots, n_s) \\
&= \sum_{\substack{a_2 \leq n_2 \leq c_2 \\ \dots \\ a_s \leq n_s \leq c_s}} \left(\int_{a_1}^{c_1} f(x_1, n_2, \dots, n_s) dx_1 + O((m_1 + 1)M) \right) \\
&= \sum_{\substack{a_2 \leq n_2 \leq c_2 \\ \dots \\ a_s \leq n_s \leq c_s}} \int_{a_1}^{c_1} f(x_1, n_2, \dots, n_s) dx_1 + O\left(\frac{m_1 + 1}{b_1} BM\right) \\
&= \sum_{\substack{a_3 \leq n_3 \leq c_3 \\ \dots \\ a_s \leq n_s \leq c_s}} \left(\int_{a_2}^{c_2} \int_{a_1}^{c_1} f(x_1, x_2, n_3, \dots, n_s) dx_1 dx_2 + O((m_2 + 1)b_1 M) \right) \\
&\quad + O\left(\frac{m_1 + 1}{b_1} BM\right) \\
&= \sum_{\substack{a_3 \leq n_3 \leq c_3 \\ \dots \\ a_s \leq n_s \leq c_s}} \int_{a_2}^{c_2} \int_{a_1}^{c_1} f(x_1, x_2, n_3, \dots, n_s) dx_1 dx_2 + O\left(\left(\frac{m_1 + 1}{b_1} + \frac{m_2 + 1}{b_2}\right) BM\right) \\
&\quad \vdots \\
&= \int_D \cdots \int f(x_1, \dots, x_s) dx_1 \cdots dx_s + O\left(\sum_{1 \leq i \leq s} \frac{m_i + 1}{b_i} BM\right).
\end{aligned}$$

This completes the proof of Lemma 3.4. \square

Lemma 3.5. *We have*

$$\begin{aligned}
\sum_{\mathbf{u} \in U'_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) &= \frac{c_2^{\ell_1 + \dots + \ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \int_{D_1} \cdots \int \exp\left(c_1 \sum_{i,j} \sqrt{x_{i,j}}\right) dx_{1,2} \cdots dx_{k,\ell_k} \\
&\quad + O\left(n^{-(1-\eta)(\ell_1 + \dots + \ell_k) - 2\eta} \exp(c_1 \sqrt{an})\right),
\end{aligned}$$

where c_1 and c_2 are given by (2.1), and D_1 is the set of all $\ell_1 + \dots + \ell_k - 1$ tuples $(x_{1,2}, \dots, x_{k,\ell_k})$ of real numbers with

$$|x_{i,j} - v_{i,j}| \leq v_{i,j}^\eta \quad (3.2)$$

for all $i + j > 2$, and

$$x_{1,1} = n - \sum_{i+j>2} s_i x_{i,j}.$$

Proof. By Lemma 3.2, we have

$$\sum_{\mathbf{u} \in U'_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) = \frac{c_2^{\ell_1 + \dots + \ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \sum_{\mathbf{u} \in U'_n} \exp\left(c_1 \sum_{i,j} \sqrt{u_{i,j}}\right).$$

For the function

$$f(x_{1,2}, \dots, x_{1,\ell_1}, \dots, x_{k,1}, \dots, x_{k,\ell_k}) = \exp\left(\sqrt{n - \sum_{i+j>2} s_i x_{i,j}} + \sum_{i+j>2} \sqrt{x_{i,j}}\right),$$

by Lemma 3.4, we have

$$\begin{aligned} & \sum_{\mathbf{u} \in U'_n} \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell_i}} p(u_{i,j}) \\ &= \frac{c_2^{\ell_1 + \dots + \ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \int_{D_1} \dots \int \exp\left(c_1 \sum_{i,j} \sqrt{x_{i,j}}\right) dx_{1,2} \dots dx_{k,\ell_k} \\ & \quad + O\left(n^{-(1-\eta)(\ell_1 + \dots + \ell_k) - 2\eta} \max_{D_1} \exp\left(c_1 \sum_{i,j} \sqrt{x_{i,j}}\right)\right). \end{aligned}$$

By the Cauchy-Schwarz inequality, we have

$$\begin{aligned} \sum_{i,j} \sqrt{x_{i,j}} &= \sum_{i,j} \frac{1}{\sqrt{s_i}} \sqrt{s_i x_{i,j}} \\ &\leq \left(\sum_{i,j} \frac{1}{s_i}\right)^{1/2} \left(\sum_{i,j} s_i x_{i,j}\right)^{1/2} \\ &= \sqrt{an}. \end{aligned}$$

Thus, we have

$$\begin{aligned} & O\left(n^{-(1-\eta)(\ell_1 + \dots + \ell_k) - 2\eta} \max_{D_1} \exp\left(c_1 \sum_{i,j} \sqrt{x_{i,j}}\right)\right) \\ &= O\left(n^{-(1-\eta)(\ell_1 + \dots + \ell_k) - 2\eta} \exp\left(c_1 \sqrt{an}\right)\right). \end{aligned}$$

□

Lemma 3.6. *We have*

$$\begin{aligned} g(n) &= \frac{c_2^{\ell_1 + \dots + \ell_k} + O(n^{3\eta-5/2})}{v_{1,1}} \left(\frac{8}{c_1}\right)^{(\ell_1 + \dots + \ell_k - 1)/2} \left(\frac{v_{1,1}}{\prod v_{i,j}}\right)^{1/4} \exp(c_1 \sqrt{an}) \\ & \quad \sqrt{s_1}^{\ell_1-1} \sqrt{s_2}^{\ell_2} \dots \sqrt{s_k}^{\ell_k} \cdot \int_{\Omega} \dots \int \exp\left(-\sum_{i,j} s_i w_{i,j}^2\right) dw_{1,2} \dots dw_{k,\ell_k} \\ & \quad + O\left(n^{-(1-\eta)(\ell_1 + \dots + \ell_k) - 2\eta} \exp\left(c_1 \sqrt{an}\right)\right), \end{aligned}$$

where Ω is the set of all $\ell_1 + \dots + \ell_k - 1$ tuples $(w_{1,2}, \dots, w_{k,\ell_k})$ of real numbers with $|w_{i,j}| < \sqrt{\frac{c_1}{8s_i} v_{i,j}^{\eta-3/4}}$ for all $i + j > 2$, and $w_{1,1} = -\sum_{i+j>2} w_{i,j}$.

Proof. Noting that

$$\exp(c_1\sqrt{an} - c_2n^{2\eta-3/2}) = O(n^{-(1-\eta)(\ell_1+\dots+\ell_k)-2\eta} \exp(c_1\sqrt{an})),$$

by (2.3), Lemmas 3.1 and 3.5, we have

$$\begin{aligned} g(n) &= \frac{c_2^{\ell_1+\dots+\ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \int \cdots \int_{D_1} \exp\left(c_1 \sum_{i,j} \sqrt{x_{i,j}}\right) dx_{1,2} \cdots dx_{k,\ell_k} \\ &\quad + O(n^{-(1-\eta)(\ell_1+\dots+\ell_k)-2\eta} \exp(c_1\sqrt{an})). \end{aligned} \quad (3.3)$$

Recall that D_1 is the set of all $\ell_1 + \dots + \ell_k - 1$ tuples $(x_{1,2}, \dots, x_{k,\ell_k})$ of real numbers with

$$|x_{i,j} - v_{i,j}| \leq v_{i,j}^\eta$$

for all $i + j > 2$, and

$$x_{1,1} = n - \sum_{i+j>2} s_i x_{i,j}.$$

Let $x_{i,j} = v_{i,j} + v_{i,j} y_{i,j}$ for all i, j . Then

$$\begin{aligned} &\int \cdots \int_{D_1} \exp\left(c_1 \sum_{i,j} \sqrt{x_{i,j}}\right) dx_{1,2} \cdots dx_{k,\ell_k} \\ &= \frac{\prod v_{i,j}}{v_{1,1}} \int \cdots \int_{D_2} \exp\left(c_1 \sum_{i,j} \sqrt{v_{i,j} + v_{i,j} y_{i,j}}\right) dy_{1,2} \cdots dy_{k,\ell_k}, \end{aligned}$$

where D_2 is the set of all $\ell_1 + \dots + \ell_k - 1$ tuples $(y_{1,2}, \dots, y_{k,\ell_k})$ of real numbers with

$$|y_{i,j}| \leq v_{i,j}^{-(1-\eta)} \quad (3.4)$$

for all $i + j > 2$, and

$$s_1 v_{1,1} y_{1,1} = - \sum_{i+j>2} s_i v_{i,j} y_{i,j}. \quad (3.5)$$

The last equality comes from $s_1 = 1$ and

$$\sum_{i,j} s_i x_{i,j} = n = \sum_{i,j} s_i v_{i,j}.$$

By (3.4) and (3.5), we have

$$|y_{i,j}| \ll n^{-(1-\eta)} \quad (3.6)$$

for all i, j . By the definition of $v_{i,j}$, we have

$$s_i v_{i,j} = \frac{n}{s_i a}.$$

Thus, by (3.5), we have

$$\sum_{i,j} \frac{n}{s_i a} y_{i,j} = 0.$$

That is,

$$\sum_{i,j} \frac{y_{i,j}}{s_i} = 0.$$

Hence

$$\sum_{i,j} \sqrt{v_{i,j}} y_{i,j} = \sum_{i,j} \sqrt{\frac{n}{s_i^2 a}} y_{i,j} = \sqrt{\frac{n}{a}} \sum_{i,j} \frac{y_{i,j}}{s_i} = 0.$$

Thus

$$\begin{aligned} & \sum_{i,j} \sqrt{v_{i,j} + v_{i,j} y_{i,j}} \\ &= \sum_{i,j} \sqrt{v_{i,j}} \left(1 + \frac{1}{2} y_{i,j} - \frac{1}{8} y_{i,j}^2 + O(y_{i,j}^3) \right) \\ &= \sum_{i,j} \sqrt{v_{i,j}} - \frac{1}{8} \sum_{i,j} \sqrt{v_{i,j}} y_{i,j}^2 + O(n^{3\eta-5/2}) \\ &= \sum_{i,j} \sqrt{\frac{n}{s_i^2 a}} - \frac{1}{8} \sum_{i,j} \sqrt{v_{i,j}} y_{i,j}^2 + O(n^{3\eta-5/2}) \\ &= \sqrt{an} - \frac{1}{8} \sum_{i,j} \sqrt{v_{i,j}} y_{i,j}^2 + O(n^{3\eta-5/2}). \end{aligned}$$

Therefore

$$\begin{aligned} & \int \cdots \int_{D_2} \exp \left(c_1 \sum_{i,j} \sqrt{v_{i,j} + v_{i,j} y_{i,j}} \right) dy_{1,2} \cdots dy_{k,\ell_k} \\ &= \exp(O(n^{3\eta-5/2})) \exp(c_1 \sqrt{an}) \int \cdots \int_{D_2} \exp \left(-\frac{c_1}{8} \sum_{i,j} \sqrt{v_{i,j}} y_{i,j}^2 \right) dy_{1,2} \cdots dy_{k,\ell_k} \\ &= (1 + O(n^{3\eta-5/2})) \exp(c_1 \sqrt{an}) \int \cdots \int_{D_2} \exp \left(-\frac{c_1}{8} \sum_{i,j} \sqrt{v_{i,j}} y_{i,j}^2 \right) dy_{1,2} \cdots dy_{k,\ell_k}. \end{aligned}$$

Let

$$v_{i,j}^{1/4} y_{i,j} = z_{i,j}, \quad \sqrt{\frac{c_1}{8}} z_{i,j} = \sqrt{s_i} w_{i,j}$$

for all i, j . Then

$$\begin{aligned} & \int \cdots \int_{D_2} \exp\left(-\frac{c_1}{8} \sum_{i,j} \sqrt{v_{i,j}} y_{i,j}^2\right) dy_{1,2} \cdots dy_{k,\ell_k} \\ &= \left(\frac{v_{1,1}}{\prod v_{i,j}}\right)^{1/4} \int \cdots \int_{D_3} \exp\left(-\frac{c_1}{8} \sum_{i,j} z_{i,j}^2\right) dz_{1,2} \cdots dz_{k,\ell_k} \\ &= \left(\frac{v_{1,1}}{\prod v_{i,j}}\right)^{1/4} \left(\frac{8}{c_1}\right)^{(\ell_1+\cdots+\ell_k-1)/2} \sqrt{s_1}^{\ell_1-1} \sqrt{s_2}^{\ell_2} \cdots \sqrt{s_k}^{\ell_k} \\ & \quad \cdot \int \cdots \int_{\Omega} \exp\left(-\sum_{i,j} s_i w_{i,j}^2\right) dw_{1,2} \cdots dw_{k,\ell_k}, \end{aligned}$$

where D_3 is the set of all $\ell_1 + \cdots + \ell_k - 1$ tuples $(z_{1,2}, \dots, z_{k,\ell_k})$ of real numbers with

$$|z_{i,j}| \leq v_{i,j}^{\eta-3/4}$$

for all $i + j > 2$, and

$$s_1 v_{1,1}^{3/4} z_{1,1} = - \sum_{i+j>2} s_i v_{i,j}^{3/4} z_{i,j},$$

Ω is the set of all $\ell_1 + \cdots + \ell_k - 1$ tuples $(w_{1,2}, \dots, w_{k,\ell_k})$ of real numbers with

$$|w_{i,j}| < \sqrt{\frac{c_1}{8s_i}} v_{i,j}^{\eta-3/4}$$

for all $i + j > 2$, and

$$s_1^{3/2} v_{1,1}^{3/4} w_{1,1} = - \sum_{i+j>2} s_i^{3/2} v_{i,j}^{3/4} w_{i,j}. \quad (3.7)$$

Noting that $s_i^{3/2} v_{i,j}^{3/4} = (n/a)^{3/4}$, (3.7) is equivalent to

$$w_{1,1} = - \sum_{i+j>2} w_{i,j}. \quad (3.8)$$

Hence

$$\begin{aligned}
& \int \cdots \int_{D_1} \exp \left(c_1 \sum_{i,j} \sqrt{x_{i,j}} \right) dx_{1,2} \cdots dx_{k,\ell_k} \\
&= \frac{\prod v_{i,j}}{v_{1,1}} \int \cdots \int_{D_2} \exp \left(c_1 \sum_{i,j} \sqrt{v_{i,j} + v_{i,j} y_{i,j}} \right) dy_{1,2} \cdots dy_{k,\ell_k} \\
&= \frac{\prod v_{i,j}}{v_{1,1}} (1 + O(n^{3\eta-5/2})) \exp(c_1 \sqrt{an}) \int \cdots \int_{D_2} \exp \left(-\frac{c_1}{8} \sum_{i,j} \sqrt{v_{i,j}} y_{i,j}^2 \right) dy_{1,2} \cdots dy_{k,\ell_k} \\
&= \frac{\prod v_{i,j}}{v_{1,1}} (1 + O(n^{3\eta-5/2})) \exp(c_1 \sqrt{an}) \left(\frac{v_{1,1}}{\prod v_{i,j}} \right)^{1/4} \left(\frac{8}{c_1} \right)^{(\ell_1 + \cdots + \ell_k - 1)/2} \\
&\quad \sqrt{s_1}^{\ell_1 - 1} \sqrt{s_2}^{\ell_2} \cdots \sqrt{s_k}^{\ell_k} \cdot \int \cdots \int_{\Omega} \exp \left(-\sum_{i,j} s_i w_{i,j}^2 \right) dw_{1,2} \cdots dw_{k,\ell_k}.
\end{aligned}$$

It follows from (3.3) and $\frac{3}{4} < \eta < \frac{5}{6}$ that

$$\begin{aligned}
g(n) &= \frac{c_2^{\ell_1 + \cdots + \ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \int \cdots \int_{D_1} \exp \left(c_1 \sum_{i,j} \sqrt{x_{i,j}} \right) dx_{1,2} \cdots dx_{k,\ell_k} \\
&\quad + O(n^{-(1-\eta)(\ell_1 + \cdots + \ell_k) - 2\eta} \exp(c_1 \sqrt{an})) \\
&= \frac{c_2^{\ell_1 + \cdots + \ell_k} + O(n^{\eta-1})}{\prod v_{i,j}} \frac{\prod v_{i,j}}{v_{1,1}} (1 + O(n^{3\eta-5/2})) \exp(c_1 \sqrt{an}) \left(\frac{v_{1,1}}{\prod v_{i,j}} \right)^{1/4} \\
&\quad \left(\frac{8}{c_1} \right)^{(\ell_1 + \cdots + \ell_k - 1)/2} \sqrt{s_1}^{\ell_1 - 1} \sqrt{s_2}^{\ell_2} \cdots \sqrt{s_k}^{\ell_k} \\
&\quad \cdot \int \cdots \int_{\Omega} \exp \left(-\sum_{i,j} s_i w_{i,j}^2 \right) dw_{1,2} \cdots dw_{k,\ell_k} \\
&\quad + O(n^{-(1-\eta)(\ell_1 + \cdots + \ell_k) - 2\eta} \exp(c_1 \sqrt{an})) \\
&= \frac{c_2^{\ell_1 + \cdots + \ell_k} + O(n^{3\eta-5/2})}{v_{1,1}} \left(\frac{8}{c_1} \right)^{(\ell_1 + \cdots + \ell_k - 1)/2} \left(\frac{v_{1,1}}{\prod v_{i,j}} \right)^{1/4} \exp(c_1 \sqrt{an}) \\
&\quad \sqrt{s_1}^{\ell_1 - 1} \sqrt{s_2}^{\ell_2} \cdots \sqrt{s_k}^{\ell_k} \cdot \int \cdots \int_{\Omega} \exp \left(-\sum_{i,j} s_i w_{i,j}^2 \right) dw_{1,2} \cdots dw_{k,\ell_k} \\
&\quad + O(n^{-(1-\eta)(\ell_1 + \cdots + \ell_k) - 2\eta} \exp(c_1 \sqrt{an})).
\end{aligned}$$

□

Now we determine the value of integral in Lemma 3.6

$$\int_{\Omega} \cdots \int \exp\left(-\sum_{i,j} s_i w_{i,j}^2\right) dw_{1,2} \cdots dw_{k,\ell_k}.$$

To do this, we need the following general lemma. We believe it should appear in somewhere.

Lemma 3.7. *If $\mathbf{x}^T A \mathbf{x}$ is a positive definite quadratic form in $\mathbf{x}^T = (x_1, \dots, x_k)$ and U is a region in \mathbb{R}^k , then there is a linear transformation $\mathbf{x} = C\mathbf{y}$ such that*

$$\int_U \cdots \int e^{-\mathbf{x}^T A \mathbf{x}} d\mathbf{x} = \frac{1}{\sqrt{\det(A)}} \int_{U'} \cdots \int e^{-y_1^2 - \cdots - y_k^2} d\mathbf{y},$$

where

$$U' = \{\mathbf{y} : \mathbf{y} = C^{-1}\mathbf{x}, \mathbf{x} \in U\}.$$

Proof. Let $\mathbf{x} = C\mathbf{y}$ be a linear transformation such that $C^T A C = I$ is the unit matrix. Then

$$|\det(C)| = \frac{1}{\sqrt{\det(A)}}.$$

Therefore

$$\begin{aligned} \int_U \cdots \int e^{-\mathbf{x}^T A \mathbf{x}} d\mathbf{x} &= |\det(C)| \int_{U'} \cdots \int e^{-y_1^2 - \cdots - y_k^2} d\mathbf{y} \\ &= \frac{1}{\sqrt{\det(A)}} \int_{U'} \cdots \int e^{-y_1^2 - \cdots - y_k^2} d\mathbf{y}. \end{aligned}$$

□

Lemma 3.8. *Let a_0, a_1, \dots, a_k be $k+1$ positive real numbers. Then there is a linear transformation $\mathbf{x} = C\mathbf{y}$ such that*

$$\begin{aligned} &\int_U \cdots \int \exp\left(-a_0(x_1 + \cdots + x_k)^2 - a_1 x_1^2 - \cdots - a_k x_k^2\right) d\mathbf{x} \\ &= \left(a_0 a_1 \cdots a_k \sum_{i=0}^k \frac{1}{a_i}\right)^{-1/2} \int_{U'} \cdots \int e^{-y_1^2 - \cdots - y_k^2} d\mathbf{y}, \end{aligned}$$

where

$$U' = \{\mathbf{y} : \mathbf{y} = C^{-1}\mathbf{x}, \mathbf{x} \in U\}.$$

Proof. It is clear that the quadratic form

$$a_0(x_1 + \cdots + x_k)^2 + a_1x_1^2 + \cdots + a_kx_k^2$$

is positive definite. Its matrix is

$$A_k = \begin{pmatrix} a_0 + a_1 & a_0 & a_0 & \cdots & a_0 \\ a_0 & a_0 + a_2 & a_0 & \cdots & a_0 \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ a_0 & a_0 & a_0 & \cdots & a_0 + a_k \end{pmatrix}.$$

It is not difficult to see that

$$\det(A_k) = a_k \det(A_{k-1}) + a_0 a_1 \cdots a_{k-1}.$$

By induction on k , we have

$$\det(A_k) = a_0 a_1 \cdots a_k \sum_{i=0}^k \frac{1}{a_i}.$$

Now Lemma 3.8 follows from Lemma 3.7. □

4 Proof of Theorem 1.1

It follows from Lemma 3.6 that

$$\begin{aligned} g(n) &= \frac{c_2^{\ell_1 + \cdots + \ell_k} + O(n^{3\eta - 5/2})}{v_{1,1}} \left(\frac{8}{c_1} \right)^{(\ell_1 + \cdots + \ell_k - 1)/2} \left(\frac{v_{1,1}}{\prod v_{i,j}} \right)^{1/4} \exp(c_1 \sqrt{an}) \\ &\quad \sqrt{s_1}^{\ell_1 - 1} \sqrt{s_2}^{\ell_2} \cdots \sqrt{s_k}^{\ell_k} \cdot \int \cdots \int_{\Omega} \exp \left(- \sum_{i,j} s_i w_{i,j}^2 \right) dw_{1,2} \cdots dw_{k,\ell_k} \\ &\quad + O \left(n^{-(1-\eta)(\ell_1 + \cdots + \ell_k) - 2\eta} \exp(c_1 \sqrt{an}) \right) \end{aligned}$$

Thus, we only need estimate $\int \cdots \int_{\Omega} \exp\left(-\sum_{i,j} s_i w_{i,j}^2\right) dw_{1,2} \cdots dw_{k,\ell_k}$. By Lemma 3.8, we have

$$\begin{aligned} & \int \cdots \int_{\Omega} \exp\left(-\sum_{i,j} s_i w_{i,j}^2\right) dw_{1,2} \cdots dw_{k,\ell_k} \\ &= \left(s_1^{\ell_1} s_2^{\ell_2} \cdots s_k^{\ell_k} \sum_{i=1}^k \frac{\ell_i}{s_i}\right)^{-1/2} \int \cdots \int_{\Omega'} \exp\left(-\sum_{i+j>2} w_{i,j}^2\right) dw'_{1,2} \cdots dw'_{k,\ell_k}, \end{aligned}$$

where Ω' is the set of all $\ell_1 + \cdots + \ell_k - 1$ tuples $(w'_{1,2}, \dots, w'_{k,\ell_k})$ of real numbers with

$$|w'_{i,j}| < c'_{i,j} v_{i,j}^{\eta-3/4}$$

for all $i + j > 2$ and some positive constants $c'_{i,j}$. Noting that, for $x > 1$,

$$\int_x^{\infty} e^{-t^2} dt = \int_x^{\infty} e^{-t^2+t-t} dt \leq e^{-x^2+x} \int_x^{\infty} e^{-t} dt = e^{-x^2}$$

and the well known Gaussian integral (also known as the Euler Poisson integral)

$$\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi},$$

for $\eta > \frac{3}{4}$, we have

$$\begin{aligned} \int_{-c'_{i,j} v_{i,j}^{\eta-3/4}}^{c'_{i,j} v_{i,j}^{\eta-3/4}} e^{-x^2} dx &= \int_{-c''_{i,j} n^{\eta-3/4}}^{c''_{i,j} n^{\eta-3/4}} e^{-x^2} dx \\ &= \int_{-\infty}^{\infty} e^{-x^2} dx + O(e^{-(c''_{i,j} n^{\eta-3/4})^2}) \\ &= \sqrt{\pi} + O(e^{-(c''_{i,j} n^{\eta-3/4})^2}), \end{aligned}$$

where $c''_{i,j}$ are some positive constants. Thus

$$\begin{aligned} & \int \cdots \int_{\Omega'} \exp\left(-\sum_{i+j>2} w_{i,j}^2\right) dw'_{1,2} \cdots dw'_{k,\ell_k} \\ &= \left(\sqrt{\pi} + O\left(e^{-(c''_{1,2} n^{\eta-3/4})^2}\right)\right) \cdots \left(\sqrt{\pi} + O\left(e^{-(c''_{k,\ell_k} n^{\eta-3/4})^2}\right)\right) \\ &= \pi^{(\ell_1 + \cdots + \ell_k - 1)/2} + O\left(e^{-cn^{2\eta-3/2}}\right), \end{aligned}$$

where c is a positive constant. Therefore,

$$\begin{aligned}
g(n) &= \frac{c_2^{\ell_1+\dots+\ell_k} + O(n^{3\eta-5/2})}{v_{1,1}} \left(\frac{8}{c_1}\right)^{(\ell_1+\dots+\ell_k-1)/2} \left(\frac{v_{1,1}}{\prod v_{i,j}}\right)^{1/4} \exp(c_1\sqrt{an}) \\
&\quad \cdot \sqrt{s_1}^{\ell_1-1} \sqrt{s_2}^{\ell_2} \cdots \sqrt{s_k}^{\ell_k} \cdot \left(s_1^{\ell_1} s_2^{\ell_2} \cdots s_k^{\ell_k} \sum_{i=1}^k \frac{\ell_i}{s_i}\right)^{-1/2} \\
&\quad \cdot (\pi^{(\ell_1+\dots+\ell_k-1)/2} + O(e^{-(cn^{2\eta-3/2})})) \\
&\quad + O\left(n^{-(1-\eta)(\ell_1+\dots+\ell_k)-2\eta} \exp(c_1\sqrt{an})\right) \\
&= \frac{c_2^{\ell_1+\dots+\ell_k} \pi^{(\ell_1+\dots+\ell_k-1)/2} + O(n^{3\eta-5/2})}{v_{1,1}} \left(\frac{8}{c_1}\right)^{(\ell_1+\dots+\ell_k-1)/2} \left(\frac{v_{1,1}}{\prod v_{i,j}}\right)^{1/4} \\
&\quad \exp(c_1\sqrt{an}) \sqrt{s_1}^{\ell_1-1} \sqrt{s_2}^{\ell_2} \cdots \sqrt{s_k}^{\ell_k} \cdot \left(s_1^{\ell_1} s_2^{\ell_2} \cdots s_k^{\ell_k} \sum_{i=1}^k \frac{\ell_i}{s_i}\right)^{-1/2} \\
&\quad + O\left(n^{-(1-\eta)(\ell_1+\dots+\ell_k)-2\eta} \exp(c_1\sqrt{an})\right) \\
&= 2^{-(3\ell_1+\dots+3\ell_k+5)/4} 3^{-(\ell_1+\dots+\ell_k+1)/4} a^{(\ell_1+\dots+\ell_k+1)/4} s_1^{\ell_1/2} \cdots s_k^{\ell_k/2} \\
&\quad \cdot n^{-3/4-(\ell_1+\dots+\ell_k)/4} \exp(c_1\sqrt{an}) \\
&\quad + O\left(n^{3\eta-\frac{5}{2}-1-(\ell_1+\dots+\ell_k-1)/4} \exp(c_1\sqrt{an})\right) \\
&\quad + O\left(n^{-(1-\eta)(\ell_1+\dots+\ell_k)-2\eta} \exp(c_1\sqrt{an})\right) \\
&= c(\mathbf{s}, \mathbf{l}) n^{d(\mathbf{l})} \exp(c_1\sqrt{an}) \\
&\quad + O\left(n^{3\eta-\frac{7}{2}-(\ell_1+\dots+\ell_k-1)/4} \exp(c_1\sqrt{an})\right) \\
&\quad + O\left(n^{-(1-\eta)(\ell_1+\dots+\ell_k)-2\eta} \exp(c_1\sqrt{an})\right).
\end{aligned}$$

Noting that η is a real number that subjects to

$$\frac{3}{4} < \eta < \min\left\{\frac{5}{6}, \frac{3}{4} \frac{\ell_1 + \dots + \ell_k - 1}{\ell_1 + \dots + \ell_k - 2}\right\},$$

we choose $\eta = \frac{3}{4} + \varepsilon'$, where ε' is sufficiently small positive real number. Thus

$$\begin{aligned}
&O(n^{3\eta-\frac{7}{2}-(\ell_1+\dots+\ell_k-1)/4} \exp(c_1\sqrt{an})) + O(n^{-(1-\eta)(\ell_1+\dots+\ell_k)-2\eta} \exp(c_1\sqrt{an})) \\
&= O(n^{-1+\varepsilon-(\ell_1+\dots+\ell_k)/4} \exp(c_1\sqrt{an})) \\
&= O(n^{d(\mathbf{l})-\frac{1}{4}+\varepsilon} \exp(c_1\sqrt{an})).
\end{aligned}$$

Noting that

$$\exp(c_1\sqrt{an}) = \exp\left(\pi\sqrt{\frac{2a(\mathbf{s}, \mathbf{l})n}{3}}\right),$$

we obtain a proof of Theorem 1.1.

Acknowledgments

This work was supported by the National Natural Science Foundation of China, Grant No. 11371195 and a project funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions.

References

- [1] Z. Ahmed, N. Baruah, M.G. Dastidar, *New congruences modulo 5 for the number of 2-color partitions*, J. Number Theory 157 (2015), 184–98, <http://dx.doi.org/10.1016/j.jnt.2015.05.002>.
- [2] R. Balasubramanian, F. Luca, *On the number of factorizations of an integer*, Integers 11 (2011), 139–143, <http://dx.doi.org/10.1515/integ.2011.012>, A12, 5 pp.
- [3] H.C. Chan, *Ramanujan’s cubic continued fraction and an analog of his “most beautiful identity”*, Int. J. Number Theory 6 (2010), no. 3, 673–680, <http://dx.doi.org/10.1142/S1793042110003150>.
- [4] H.C. Chan, S. Cooper, *Congruences modulo powers of 2 for a certain partition function*, Ramanujan J. 22 (2010), 101–117. <http://dx.doi.org/10.1007/s11139-009-9197-6>.
- [5] S.C. Chen, *Congruences for a certain partition function*, Ann. Comb. 18 (2014), 607–615, <http://dx.doi.org/10.1007/s00026-014-0240-y>.
- [6] S. Chern, *New congruences for 2-color partitions*, J. Number Theory 163 (2016), 474–481, <http://dx.doi.org/10.1016/j.jnt.2015.12.020>.
- [7] Y.-G. Chen, Y.-L. Li, *On the square-root partition function*, C. R. Math. Acad. Sci. Paris 353 (4) (2015), 287–290, <http://dx.doi.org/10.1016/j.crma.2015.01.013>.
- [8] P. Erdős, *On an elementary proof of some asymptotic formulas in the theory of partitions*, Ann. Math. 43 (1942), 437–450, <http://dx.doi.org/10.2307/1968802>.

- [9] G.H. Hardy, S. Ramanujan, *Asymptotic formula for the distribution of integers of various types*, Proc. London Math. Soc. (2) 16 (1917), 112–132.
- [10] A.E. Ingham, *A Tauberian theorem for partitions*, Ann. of Math. (2) 42 (1941), 1075–1090, <http://dx.doi.org/10.2307/1970462>.
- [11] B. Kim, *An analog of crank for a certain kind of partition function arising from the cubic continued fraction*, Acta Arith. 148 (2011), no. 1, 1–19, <http://dx.doi.org/10.4064/aa148-1-1>.
- [12] D.H. Lehmer, *On the remainders and convergence of the series for the partition function*, Trans. Amer. Math. Soc. 46 (1939), 362–373, <http://dx.doi.org/10.2307/1989927>.
- [13] Y. L. Li and Y. G. Chen, *On the r -th Root Partition Function*, Taiwanese J. Math. 20 (2016), 545–551, <http://dx.doi.org/10.11650/tjm.20.2016.6812>.
- [14] F. Luca and D. Ralaivaosaona, *An explicit bound for the number of partitions into roots*, J. Number Theory 169 (2016), 250–264, <http://dx.doi.org/10.1016/j.jnt.2016.05.017>.
- [15] A.M. Odlyzko, *Asymptotic enumeration methods*, Handbook of combinatorics. 1, 2 (1995), 1063–1229.
- [16] S. Ramanujan, *Congruence properties of partitions*, Math. Z. 9 (1921), no. 1-2, 147–153, <http://dx.doi.org/10.1007/BF01378341>.
- [17] J. Sinick, *Ramanujan congruences for a class of eta quotients*, Int. J. Number Theory 6 (2010), 835–847, <http://dx.doi.org/10.1142/S1793042110003253>.
- [18] Y.V. Uspensky *Asymptotic expressions of numerical functions occurring in problems concerning the partition of numbers into summands*, Bull Acad Sci de Russie 14 (1920), 199–218.