# COVERINGS, MATCHINGS AND THE NUMBER OF MAXIMAL INDEPENDENT SETS OF GRAPHS

### DO TRONG HOANG AND TRAN NAM TRUNG

ABSTRACT. We determine the maximum number of maximal independent sets of arbitrary graphs in terms of their covering numbers and we completely characterize the extremal graphs. As an application, we give a similar result for König-Egerváry graphs in terms of their matching numbers.

## 1. Introduction

Let m(G) be the number of maximal independent sets of a simple graph G. Around 1960, Erdös and Moser raised the problem of determining the largest number of m(G) in terms of order of G, which we shall denote by n in this paper, and determining the extremal graphs. In 1965, Moon and Moser [14] solved this problem for any simple graph.

This problem now has been focused in investigating various classes of graphs such as: for connected graphs by Füredi [5]; and independently Griggs et al. [8]; for triangle-free graphs by Hujter and Tuza [10] and for connected triangle-free graphs by Chang and Jou [3]; Sagan and Vatter [16] and Goh et al. [6] solved the problem for graphs with at most r cycles; for connected unicyclic graphs by Koh et al. [11]; for trees independently by Cohen [4], Griggs and Grinstead [7], Sagan [15], Wilf [17]; for bipartite graphs by Liu [13] and bipartite graphs with at least one cycle by Li et al. [12].

The goal of this paper is to determine the maximum number of m(G) of arbitrary simple graph G in terms of its covering number, denoted by  $\beta(G)$ ; and to characterize the extremal graphs. On that basis we will consequently improve some certain results among those mentioned above. Before stating our results, recall that a matching in G is a set of edges, no two of which meet a common vertex. The matching number  $\nu(G)$  of G is the maximum size of matchings of G. An induced matching M in a graph G is a matching where no two edges of G are joined by an edge of G. The induced matching number  $\nu_0(G)$  of G is the maximum size of induced matchings of G. We always have  $\nu_0(G) \leq \nu(G)$ ; and if  $\nu_0(G) = \nu(G)$  then G is called a Cameron-Walker graph according to Hibi et al. [9]. The main result of the paper is as follows:

**Theorem** (Theorem 2.7 and Theorem 3.3). Let G be a graph. Then  $m(G) \leq 2^{\beta(G)}$ , and the equality holds if and only if G is a Cameron-Walker bipartite graph.

<sup>2010</sup> Mathematics Subject Classification. 05C30, 05D15.

Key words and phrases. Independent set, Matching number, Covering number.

A graph G is called a  $K\"{o}nig$ - $Egerv\'{a}ry$  graph if the matching number is equal to the covering number that is  $\beta(G) = \nu(G)$ . As an application, we determine the maximum number of m(G) for  $K\"{o}nig$ - $Egerv\'{a}ry$  graphs G, and characterize the extremal graphs.

Corollary 3.4. Let G be a König-Egerváry graph. Then

$$m(G) \le 2^{\nu(G)},$$

and the equality holds if and only if G is a Cameron-Walker bipartite graph.

It is well-known that all bipartite graphs are König-Egerváry (see [1, Theorem 8.32]). In general,  $\nu(G) \leq \lfloor \frac{n}{2} \rfloor$ , where n is the order of G. Thus Corollary 3.4 improves the main result of Liu (see [13, Theorem 2.1]) for bipartite graphs.

# 2. Bounds for m(G)

We now recall some basic concepts and terminology from graph theory (see [1]). Let G be a simple graph with vertex set V(G) and edge set E(G). An edge  $e \in E(G)$  connecting two vertices x and y will be also written as xy (or yx). For a subset S of V(G), we denote by G[S] the induced subgraph of G on the vertex set S; and denote  $G \setminus S$  by  $G[V(G) \setminus S]$ . The neighborhood of S in G is the set

$$N_G(S) := \{ y \in V(G) \setminus S \mid xy \in E(G) \text{ for some } x \in S \},$$

the close neighborhood of S is  $N_G[S] := S \cup N_G(S)$  and the localization of G with respect to S is  $G_S := G \setminus N_G[S]$ . If  $S = \{x\}$ , we write  $N_G(x)$  (resp.  $N_G[x]$ ,  $G_x$ ,  $G \setminus x$ ) instead of  $N_G(\{x\})$  (resp.  $N_G[\{x\}]$ ,  $G_{\{x\}}$ ,  $G \setminus \{x\}$ ). The number  $\deg_G(x) := |N_G(x)|$  is called the degree of x in G. A vertex in G of degree zero is called an isolated vertex of G. A vertex x of G is called leaf adjacent to g if  $\deg_G(x) = 1$  and g is an edge of g. A complete graph with g vertices is denoted by g. A graph g is called triangle. The union of two disjoint graphs g and g is the graph g with vertex set g and g is denoted by g where g is a positive integer.

A graph is called *totally disconnected* if it is either a null graph or containing no edge. Thus, m(G) = 1 whenever G is totally disconnected. The following basic lemmas on determining m(G) for arbitrary graph G will be frequently used later.

**Lemma 2.1.** [10, Lemma 1] Let G be a graph. Then

- (1)  $\operatorname{m}(G) \leq \operatorname{m}(G_x) + \operatorname{m}(G \setminus x)$ , for any vertex x of G.
- (2) If x is a leaf adjacent to y of G, then  $m(G) = m(G_x) + m(G_y)$ .
- (3) If  $G_1, \ldots, G_s$  are connected components of G, then

$$m(G) = \prod_{i=1}^{s} m(G_i).$$

**Lemma 2.2.** If H is an induced subgraph of G, then  $m(H) \leq m(G)$ .

We first give an upper bound for m(G) in terms of  $\nu(G)$ , and the extremal graphs. However, that upper bound does not cover the result of Moon and Moser [14].

**Proposition 2.3.** Let G be a graph. Then,  $m(G) \leq 3^{\nu(G)}$  and the equality holds if and only if  $G \cong sK_3 \cup tK_1$ , where  $s = \nu(G)$  and t = |V(G)| - 3s.

*Proof.* We prove the proposition by induction on  $\nu(G)$ . If  $\nu(G) = 0$ , then G is totally disconnected, and then the assertion is trivial.

If  $\nu(G) = 1$ , let xy be an edge of G and let  $S := V(G) \setminus \{x,y\}$ . Then G[S] is totally disconnected and if we have two vertices in S, say u and v, such that xu and yv are edges of G, then  $u \equiv v$ . In particular, there is at most one vertex in S that is adjacent to both x and y. We now consider three cases:

Case 1: x and y are not adjacent to any vertex in S. In this case, we have m(G) = 2, and the proposition holds.

Case 2: x is not adjacent to any vertex in S and y is adjacent to some vertices in S. Then, we have m(G) = 2, and the proposition holds.

Case 3: There is a vertex in S, say z, that is adjacent to both x and y. In this case, every other vertex of S is not adjacent to either x or y. Thus,  $G = K_3 \cup tK_1$ , where t = |V(G)| - 3 and  $m(G) = 3 = 3^{\nu(G)}$ . Therefore, the proposition is proved in this case.

Assume that  $\nu(G) \ge 2$ . Let xy be an edge of G. Since both x and y are not vertices of the following graphs:  $G_x$ ,  $G_y$  and  $G \setminus \{x,y\}$ , we deduce that

$$\nu(G_x) \le \nu(G) - 1, \ \nu(G_y) \le \nu(G) - 1 \ \text{ and } \nu(G \setminus \{x, y\}) \le \nu(G) - 1.$$

Thus, by the induction hypothesis, we obtain

$$m(G_x) \leqslant 3^{\nu(G)-1}, \ m(G_y) \leqslant 3^{\nu(G)-1} \ \text{and} \ \ m(G \setminus \{x,y\}) \leqslant 3^{\nu(G)-1}.$$

Note that  $(G \setminus x)_y = G_y$ . Combining with Lemma 2.1, we obtain

$$m(G) \leq m(G_x) + m(G \setminus x)$$

$$\leq m(G_x) + m(G_y) + m(G \setminus \{x, y\})$$

$$\leq 3^{\nu(G)-1} + 3^{\nu(G)-1} + 3^{\nu(G)-1} = 3^{\nu(G)}.$$

This proves the first conclusion of the proposition. The equality  $m(G) = 3^{\nu(G)}$  occurs if and only if

$$m(G) = m(G_x) + m(G \setminus x), \ m(G_x) = m(G_y) = m(G \setminus \{x, y\}) = 3^{\nu(G)-1}$$

and

$$\nu(G_x) = \nu(G_y) = \nu(G \setminus \{x, y\}) = \nu(G) - 1.$$

If  $G = sK_3 \cup tK_1$ , then  $s = \nu(G)$  and  $m(G) = 3^{\nu(G)}$ . Therefore, the necessary condition of the second conclusion of the proposition is followed. Now, it remains to prove that if  $m(G) = 3^{\nu(G)}$  then  $G \cong sK_3 \cup tK_1$ .

Indeed, by the induction hypothesis, it follows that  $G_x, G_y$  and  $G \setminus \{x, y\}$  have the same component without isolated vertices, that is  $(s-1)K_3$ , where  $s = \nu(G)$ . In

particular, x and y are not adjacent to any vertex of  $(s-1)K_3$ . Let H be an induced subgraph of G on the vertex set  $V(G)\backslash V((s-1)K_3)$ . Then, H and  $(s-1)K_3$  are disjoint subgraphs of G. By Lemma 2.1, we imply  $\mathrm{m}(G)=\mathrm{m}(H)\,\mathrm{m}((s-1)K_3)=\mathrm{m}(H)3^{s-1}$ . Since  $\mathrm{m}(G)=3^s$ ,  $\mathrm{m}(H)=3$ . Note that  $\nu(H)=1$ , so the induction hypothesis again yields  $H=K_3\cup tK_1$ . Thus,  $G=sK_3\cup tK_1$ . The proof is complete.

The following lemma gives a lower bound for m(G) in terms of the induced matching number  $\nu_0(G)$ .

**Lemma 2.4.** Let G be a graph. Then,  $m(G) \ge 2^{\nu_0(G)}$ .

*Proof.* Let 
$$\{x_1y_1, \ldots, x_ry_r\}$$
 be an induced matching of  $G$ , where  $r = \nu_0(G)$ . Set  $H := G[\{x_1, \ldots, x_r, y_1, \ldots, y_r\}]$ . By Lemma 2.2,  $m(G) \ge m(H) = 2^{\nu_0(G)}$ .

Recall that a vertex cover of G is a subset C of V(G) such that for each  $xy \in E(G)$ , we have either  $x \in C$  or  $y \in C$ . The covering number of G, denoted by  $\beta(G)$ , is the minimum size of vertex covers of G. From this definition, the following two lemmas are obvious.

**Lemma 2.5.** Let H be an induced subgraph of G. Then,

- (1) If S is a vertex cover of G, then  $S \cap V(H)$  is a vertex cover of H.
- (2)  $\beta(H) \leq \beta(G)$ .

**Lemma 2.6.** Assume S is a vertex cover of G. If  $U \subseteq S$ , then

- (1)  $S \setminus U$  is a vertex cover of  $G \setminus U$ ; and
- $(2) \ \beta(G \setminus U) \le \beta(G) |U|.$

We conclude this section by giving a upper bound for m(G) in terms of  $\beta(G)$ .

**Theorem 2.7.** Let G be a graph. Then,  $m(G) \leq 2^{\beta(G)}$ .

*Proof.* We prove the lemma by induction on  $\beta(G)$ . If  $\beta(G) = 0$ , then G is totally disconnected, and so the assertion is trivial.

Assume that  $\beta(G) \ge 1$ . Let S be a vertex cover of G such that  $|S| = \beta(G)$ . Let  $x \in S$ . By Lemma 2.6, we have  $\beta(G \setminus x) \le \beta(G) - 1$ . Hence,  $m(G \setminus x) \le 2^{\beta(G \setminus x)}$  by the induction hypothesis.

Since  $G_x$  is an induced subgraph of  $G \setminus x$ ,  $\mathrm{m}(G_x) \leq \mathrm{m}(G \setminus x)$  by Lemma 2.2. Together with Lemma 2.1, we obtain

$$\operatorname{m}(G) \leq \operatorname{m}(G \setminus x) + \operatorname{m}(G_x)$$
  
  $\leq 2 \cdot \operatorname{m}(G \setminus x) \leq 2^{\beta(G \setminus x) + 1} \leq 2^{\beta(G)},$ 

as required.

### 3. Extremal graphs

A graph G is called *bipartite* if its vertex set can be partitioned into two subsets A and B so that every edge has one end in A and one end in B; such a partition is called a *bipartition* of the graph, and denoted by (A, B). If every vertex in A is joined

to every vertex in B then G is called a complete bipartite graph, which is denoted by  $K_{|A|,|B|}$ . A star is the complete bipartite graph  $K_{1,m}$   $(m \ge 0)$  consisting of m+1 vertices. A star triangle is a graph joining some triangles at one common vertex.

Cameron and Walker [2] gave a classification of the simple graphs G with  $\nu(G) = \nu_0(G)$ ; such graphs now are the so-called Cameron-Walker graphs (see [9]).


**Lemma 3.1.** ([2, Theorem 1] or [9, p.258]) A graph G is Cameron-Walker if and only if it is one of the following graphs:

- (1) a star;
- (2) a star triangle;
- (3) a finite graph consisting of a connected bipartite graph with bipartition (A, B) such that there is at least one leaf edge attached to each vertex  $i \in A$  and that there may be possibly some pendant triangles attached to each vertex  $j \in B$ .

**Example 3.2.** Let G be Cameron-Walker graph with 8 vertices in Figure 1. Then  $\nu(G) = 2$  and the maximal independent sets of G are

$$\{1, 2, 5, 6, 7, 8\}; \{3, 4\}; \{3, 5, 6\}; \{4, 7, 8\}$$

Hence, m(G) = 4.



**Theorem 3.3.** Let G be a graph. Then  $m(G) = 2^{\beta(G)}$  if and only if G is a Cameron-Walker bipartite graph.

*Proof.* If G is a Cameron-Walker bipartite graph, then  $\nu_0(G) = \nu(G) = \beta(G)$ . Together with Lemma 2.4 and Theorem 2.7, this fact yields  $m(G) = 2^{\beta(G)}$ .

Conversely, assume that  $m(G) = 2^{\beta(G)}$ . We will prove that G is Cameron-Walker bipartite by induction on  $\beta(G)$ .

If  $\beta(G) = 0$ , then G is totally disconnected and so the assertion is trivial. If  $\beta(G) = 1$ , then G is a union of a star and isolated vertices. In this case, G is a Cameron-Walker bipartite graph by Lemma 3.1.

Assume that  $\beta(G) \geq 2$ . Let S be a minimal vertex cover of G such that  $|S| = \beta(G)$ . We first prove two following claims.

Claim 1: S is an independent set of G.

Indeed, assume on the contrary that there would be an edge, say xy, with  $x, y \in S$ . By Lemma 2.5,  $S \cap V(G_x)$  is a vertex cover of  $G_x$ . Since  $S \cap V(G_x) \subseteq S \setminus \{x, y\}$ , we deduce that

$$\beta(G_x) \leqslant |S| - 2 = \beta(G) - 2.$$

Similarly,  $S \setminus \{x\}$  is a vertex cover of  $G \setminus x$ . Thus  $\beta(G \setminus x) \leq \beta(G) - 1$ . Together those inequalities with Lemma 2.1 and Theorem 2.7, we have

$$m(G) \le m(G_x) + m(G \setminus x) \le 2^{\beta(G)-2} + 2^{\beta(G)-1} < 2^{\beta(G)}.$$

This inequality contradicts our assumption. Therefore, S is an independent set of G.

Claim 2: 
$$m(G_U) = 2^{\beta(G_U)}$$
 and  $\beta(G_U) = \beta(G) - |U|$  for any  $U \subseteq S$ .

Indeed, we prove the claim by the induction on |U|. If |U| = 0, i.e., U is empty, then there is nothing to prove.

If |U|=1, then  $U=\{x\}$  for some vertex x. Since  $x\in S$ , by Lemmas 2.5 and 2.6, we have  $\beta(G_x)\leq \beta(G\setminus x)\leq \beta(G)-1$ . By Theorem 2.7,  $\mathrm{m}(G\setminus x)\leq 2^{\beta(G\setminus x)}$  and  $\mathrm{m}(G_x)\leq 2^{\beta(G_x)}$ . Together these inequalities with equality  $\mathrm{m}(G)=2^{\beta(G)}$ , Lemma 2.1 gives

$$2^{\beta(G)} = m(G) \le m(G \setminus x) + m(G_x) \le 2^{\beta(G \setminus x)} + 2^{\beta(G_x)} \le 2^{\beta(G)-1} + 2^{\beta(G)-1} \le 2^{\beta(G)}.$$

Hence,  $m(G_x) = 2^{\beta(G_x)}$  and  $\beta(G_x) = \beta(G) - 1$ , and the claim holds in this case.

We now assume  $|U| \geq 2$ . Let  $x \in U$  and let  $T := U \setminus \{x\}$ . Note that T is a nonempty independent set of S and |T| = |U| - 1. By the induction hypothesis of our claim,  $m(G_T) = 2^{\beta(G_T)}$  and  $\beta(G_T) = \beta(G) - |T|$ .

Note that, by Claim 1, S is an independent set of G. Thus  $S \setminus T = S \setminus N_G[T]$ . By Lemma 2.5,  $S \setminus T$  is a vertex cover of  $G_T$ . Since  $x \in S \setminus T$ , by the same argument in the inductive step of our claim with  $G_T$  replacing by G, we have  $\mathrm{m}((G_T)_x) = 2^{\beta((G_T)_x)}$  and  $\beta((G_T)_x) = \beta(G_T) - 1$ .

Since  $G_U = (G_T)_x$ , we obtain  $m(G_U) = 2^{\beta(G_U)}$  and

$$\beta(G_U) = \beta(G_T) - 1 = \beta(G) - (|T| + 1) = \beta(G) - |U|,$$

as claimed.

We turn back to the proof of the theorem. By Claim 1, S is both a vertex cover and an independent set of G. Therefore G is a bipartite graph with bipartition  $(S, V(G) \setminus S)$ . It remains to prove G is a Cameron-Walker graph.

For each  $x \in S$ , let  $U := S \setminus \{x\}$ . By Claim 2,  $\beta(G_U) = \beta(G) - |U| = 1$ . Hence,  $G_U$  is a union of a star with bipartition  $(\{x\}, Y)$ , where  $\emptyset \neq Y \subseteq V(G) \setminus S$  and isolated vertices. Thus, there is a vertex  $y \in Y$  such that  $\deg_{G_U}(y) = 1$  and  $xy \in E(G)$ . Since  $V(G) \setminus S$  is an independent set, the equality  $\deg_{G_U}(y) = 1$  forces  $\deg_G(y) = 1$ . By using Lemma 3.1, we conclude that G is a Cameron-Walker graph, and the proof is complete.

If G is a König-Egerváry graph, then  $\beta(G) = \nu(G)$ . Together Theorems 2.7 and 3.3, this fact yields.

Corollary 3.4. Let G be a König-Egerváry graph. Then

$$m(G) < 2^{\nu(G)}$$
,

and the equality holds if and only if G is a Cameron-Walker bipartite graph.

**Acknowledgment.** This work is partially supported by NAFOSTED (Vietnam), Project 101.04-2015.02.

## References

- J. A. Bondy, U. S. R. Murty, Graph Theory, no. 244 in Graduate Texts in Mathematics, Springer, 2008.
- [2] K. Cameron and T. Walker, The graphs with maximum induced matching and maximum matching the same size, Discrete Math. **299** (2005) 49-55.
- [3] G. J. Chang and M. J. Jou, The number of maximal independent sets in connected triangle-free graphs, Discrete Math. 197-198 (1999), 169-178.
- [4] D. Cohen, Counting stable sets in trees, in Seminaire Lotharingien de combinatoire, 10 session, R. König, ed., Institute de Recherche Mathámatique Avancée pub., Strasbourg, France, 1984, 48-52.
- [5] Z. Füredi, The number of maximal independent sets in connected graphs, J. Graph Theory, 11 (1987), 463-470.
- [6] C. Y. Goh, K. M. Koh, B. E. Sagan, and V. R. Vatter, Maximal independent sets in graphs with at most r cycles, J. Graph Theory, 53 (2006) 207-282.
- [7] J. R. Griggs and C. M. Grinstead, unpublished result, 1986.
- [8] J. R. Griggs, C. M. Grinstead and D. R. Guichard, The number of maximal independent sets in a connected graph, Discrete Math. 68 (1988), 211-220.
- [9] T. Hibi, A. Higashitani, K. Kimura and A. B. O'Keefe, Algebraic study on CameronWalker graphs, J. Algebra 422 (2015), 257-269.
- [10] M. Hujter, Z. Tuza, The number of maximal independent sets in triangle-free graphs, SIAM J. Discrete Math. 6 (1993) 284-288.
- [11] K. M. Koh, C. Y. Goh and F. M. Dong, The maximum number of maximal independent sets in unicyclic connected graphs, Discrete Math. 380 (2008), 3761-3769.
- [12] S. Li, H. Zhang and X. Zhang, Maximal independent sets in bipartite graphs with at least one cycle, Discrete Math. and Theor. Comput. Sci. 15 (2013), 243-248.
- [13] J. Liu, Maximal independent sets in bipartite graphs, J. Graph Theory, 17 (1993), 495-507.
- [14] J. W. Moon and L. Moser, On cliques in graphs, Israel J. Math., 3 (1965), pp. 23-28.
- [15] B. E. Sagan, A note on independent sets in trees, SIAM J. Discrete Math. 1 (1988), 105-108.
- [16] B. E. Sagan, V. R. Vatter, Maximal and maximum independent sets in graphs with at most r cycles, J. Graph Theory 53 (2006) 283-314.
- [17] H. S. Wilf, The number of maximal independent sets in a tree, SIAM J. Algebraic Discrete Methods 7 (1986), 125-130.

E-mail address: dthoang@math.ac.vn

Institute of Mathematics, VAST, 18 Hoang Quoc Viet, 10307 Hanoi, Vietnam

E-mail address: tntrung@math.ac.vn

Institute of Mathematics, VAST, 18 Hoang Quoc Viet, 10307 Hanoi, Vietnam