

DEGREE OF W-OPERATOR AND NONCROSSING PARTITION

HAO SUN

ABSTRACT. Goulden and Jackson first introduced the cut-and-join operator. The cut-and-join is widely used in studying the Hurwitz number and many other topological recursion problems. Mironov, Morosov and Natanzon give a more general construction and call it W -operator $W([n])$. As a special case, the cut-and-join operator is $W([2])$. In this paper, we study the structure of $W([n])$. We prove that $W([n])$ can be written as the sum of $n!$ terms and each term corresponds uniquely to a permutation in S_n . We also define the degree of each term. We prove that there is a correspondence between the terms of $W([n])$ with highest degree and the noncrossing partitions.

AMS Subject Classification Number: 05E15, 05A18

Key Words: W -operator, cut-and-join operator, noncrossing partition

1. INTRODUCTION

The cut-and-join operator Δ

$$\Delta = \frac{1}{2} \sum_{i,j \geq 1} \left((i+j)p_i p_j \frac{\partial}{\partial p_{i+j}} + ij p_{i+j} \frac{\partial^2}{\partial p_i \partial p_j} \right)$$

is introduced by Goulden and Jackson [1, 2]. It is an infinite sum of differential operators in variables p_i , $i \geq 1$. This operator plays an important role in calculating the simple Hurwitz number [3, 4, 6, 7] and many other enumerative geometry problems [5, 9, 13].

Mironov, Morosov and Natanzon [10, 11] construct the W -operators $W(\lambda)$, where $\lambda = (1^{i_1} 2^{i_2} \dots n^{i_n})$ is a partition of a positive integer n . The W -operators are differential operators acting on the formal power series $\mathbb{C}[[X_{ij}]]_{i,j \geq 1}$, where X_{ij} are coordinate functions on the positive-half-infinite matrix. A subring of $\mathbb{C}[[X_{ij}]]_{i,j \geq 1}$ is $\mathbb{C}[p_1, p_2, \dots]$, where $p_k = \text{Tr}(X^k)$ and $X = (X_{ij})_{i,j \geq 1}$. Let $[n] = (n^1)$ be the partition of n . A direct calculation shows that $W([2])$ is the cut-and-join operator Δ on the ring $\mathbb{C}[p_1, p_2, \dots]$. In this paper, we study the structure of the operators $W([n])$, $n \geq 1$. We consider $W([n])$ as operators on the ring $\mathbb{C}[p_1, p_2, \dots]$.

In §2, we review the natural quiver structure of permutations. Based on the quiver structure, we give a construction (Construction 2.2) that all permutations in S_{n+1} can be constructed from permutations in S_n . Generally speaking, given any permutation in S_n , we construct $n+1$ permutations in S_{n+1} . The key part in §2 is Construction 2.2. Construction 2.2 plays an important role in proving the structure theorem of $W([n])$ (Theorem 3.7).

In §3, we review the definition and some properties of $W([n])$. We prove the following theorem about the structure of $W([n])$.

Theorem. 3.7 *$W([n])$ is a well-defined operator on $\mathbb{C}[p_1, p_2, \dots]$. It can be written as the sum of $n!$ summations FS_α , each of which corresponds to a unique quiver \hat{Q}_α , or equivalently a unique permutation $\alpha \in S_n$.*

In §4, we define the degree of each summation (term) of $W([n])$. We show that the maximal degree is $n + 1$. The maximal degree summations are used to study the generating function of minimal d -Hurwitz number [8], where d stands for d -cycles.

In §5, we study the number of summations of $W([n])$ with maximal degree. We show that there is an one-to-one correspondence between summations with maximal degree and noncrossing permutations of $[n] = \{1, \dots, n\}$ (Theorem 5.4). By Simion's study of noncrossing partition [12], we show that the number of summations with maximal degree is exactly the Catalan number (Corollary 5.5).

2. PERMUTATION GROUP AND QUIVER

In this section, we first review the quiver structure of permutations. Then, we give a new construction about all permutations in S_n by induction. Roughly speaking, we construct n distinct permutations in S_n from a given permutation in S_{n-1} . This construction comes from some operations on the quiver, and the idea of this construction comes from the calculation of W -operator (Remark 3.6). We will discuss the idea in the next section.

We begin with the definition of the quiver. A *quiver* is a directed graph. So, as usual, a *quiver* $Q = (V, A, s, t)$ is a quadruple, where V is the set of vertices, A is the set of arrows, s and t are two maps $A \rightarrow V$. If $a \in A$, $s(a)$ is the source of this arrow and $t(a)$ is the target. We assume V and A to be finite sets. If B is a subset of A , $V_B = \{s(a), t(a), a \in B\}$, then we call (V_B, B, s', t') the *subquiver* of Q , where $s' = s|_B, t' = t|_B$. A quiver $Q = (V, A, s, t)$ is *connected* if the underlying undirected graph of Q is connected. A connected quiver $Q = (V, A, s, t)$ is a *loop*, if for any vertex $v \in V$, there is a unique arrow $a \in A$ such that $s(a) = v$ and a unique arrow $b \in A$ such that $t(b) = v$. A *chain* is obtained by omitting a single arrow in a loop.

Any permutation $\alpha \in S_n$ has a natural quiver structure $Q_\alpha = (V_\alpha, A_\alpha, s, t)$, where $V_\alpha = \{1, \dots, n\}$ and $A_\alpha = \{i \rightarrow \alpha(i), 1 \leq i \leq n\}$. Notice that the quiver Q_α only contains loops. Since we want to use induction, we construct another quiver \hat{Q}_α from Q_α .

Construction 2.1. Given $\alpha \in S_n$, let Q_α be the corresponding quiver. There is a unique arrow a in Q_α such that $s(a) = 1$. We substitute this arrow by a new one \hat{a} , where $s(\hat{a}) = n + 1$ and $t(\hat{a}) = t(a)$. Denote by \hat{A}_α the new set of arrows. The new vertex set is $\hat{V}_\alpha = \{1, \dots, n, n + 1\}$. Denote by \hat{Q}_α the new quiver, $\hat{Q}_\alpha = (\hat{V}_\alpha, \hat{A}_\alpha, s, t)$.

We give an example about this construction. Take $\alpha = (321) \in S_3$, then Q_α is

$$Q_\alpha : \quad 3 \longrightarrow 2 \longrightarrow 1 .$$

\hat{Q}_α is

$$\hat{Q}_\alpha : \quad 4 \longrightarrow 3 \longrightarrow 2 \longrightarrow 1 .$$

Clearly, Q_α is a loop and \hat{Q}_α is a chain. In general, \hat{Q}_α consists of one chain and possibly a number of loops. The chain in \hat{Q}_α always start from $n + 1$ and stop at 1. Clearly, we can construct Q_α uniquely from \hat{Q}_α . Thus we have an one-to-one correspondence between permutations α and quivers \hat{Q}_α .

Now we will work on the quiver \hat{Q}_α and construct $n + 1$ quivers corresponding to permutations in S_{n+1} . Generally speaking, if we want to construct a new quiver \hat{Q}_β representing an element $\beta \in S_{n+1}$, we should add one more vertex $n + 2$ into \hat{V}_α and add arrows a_1, a_2 in \hat{A}_α

such that

$$s(a_1) = n + 2, \quad t(a_2) = n + 1,$$

where a_1, a_2 can be the same arrow. Here is the construction.

Construction 2.2. Given any $\alpha \in S_n$, we write α as the product of disjoint cycles $\alpha = \alpha_1 \alpha_2 \dots \alpha_k$. We assume $1 \in \alpha_1$. So, the corresponding subquiver for α_1 in \hat{Q}_α is the chain as following

$$\hat{Q}_{\alpha_1} : \quad n + 1 \longrightarrow \dots \longrightarrow 1 .$$

• **Case 0**

We extend the quiver for α_1 directly

$$\hat{Q}_{\beta_1} : \quad n + 2 \longrightarrow n + 1 \longrightarrow \dots \longrightarrow 1 .$$

Clearly, this subquiver represents a well-defined cycle β_1 (by replacing $n + 2$ by 1). In this way, we construct a permutation $\beta \in S_{n+1}$, where $\beta = \beta_1 \alpha_2 \dots \alpha_k$. In this case, a_1, a_2 are the same arrow

$$a_1 = a_2 : \quad n + 2 \longrightarrow n + 1 .$$

Next we consider the general case. The idea is cutting an arrow in \hat{Q}_α and reconnect the chain and loops in \hat{Q}_α . Since there are n choices of arrows in \hat{Q}_α , we can construct n permutations. We choose an arbitrary arrow $a : i \rightarrow j$ in \hat{Q}_α .

• **Case 1 (Cut Case), $a \in \hat{Q}_{\alpha_1}$**

In this case, \hat{Q}_{α_1} is

$$\hat{Q}_{\alpha_1} : \quad n + 1 \longrightarrow \dots \longrightarrow i \longrightarrow j \longrightarrow \dots \longrightarrow 1 .$$

First, cut the arrow $i \rightarrow j$. We get

$$n + 1 \longrightarrow \dots \longrightarrow i, \quad j \longrightarrow \dots \longrightarrow 1 .$$

Then, we add the following two arrows

$$a_1 : n + 2 \longrightarrow j \quad a_2 : i \longrightarrow n + 1 .$$

Finally, we get the following quiver,

$$\hat{Q}_{\beta_1 \beta_2} : \quad n + 2 \longrightarrow j \longrightarrow \dots \longrightarrow 1, \quad i \xrightarrow{\quad \curvearrowright \quad} n + 1 \longrightarrow \dots \longrightarrow 1 .$$

They represent two disjoint cycles in S_{n+1} by replacing $n + 2$ by 1. Denote by β_1 and β_2 , $1 \in \beta_1$. So, $\beta = \beta_1 \beta_2 \alpha_2 \dots \alpha_k$ is the permutation in S_{n+1} constructed by cutting the arrow a .

• **Case 2 (Join Case), $a \notin \hat{Q}_{\alpha_1}$**

Without loss of generality, we assume that $a \in \hat{Q}_{\alpha_2}$. The corresponding quiver for α_1 and α_2 are

$$\hat{Q}_{\alpha_1 \alpha_2} : \quad n + 1 \longrightarrow \dots \longrightarrow 1, \quad i \xrightarrow{\quad \curvearrowright \quad} j \longrightarrow \dots \longrightarrow 1 .$$

Similar to **Case 1**, we cut the arrow $i \rightarrow j$ and we get

$$n+1 \longrightarrow \cdots \longrightarrow 1, \quad j \longrightarrow \cdots \longrightarrow i.$$

Then, we add the following two arrows

$$a_1 : n+2 \longrightarrow j \quad a_2 : i \longrightarrow n+1.$$

Finally, we get the chain

$$\hat{Q}_{\beta_1} : n+2 \longrightarrow j \longrightarrow \cdots \longrightarrow i \longrightarrow n+1 \longrightarrow \cdots \longrightarrow 1.$$

It represents a cycle in S_{n+1} by replacing $n+2$ by 1. Denote by β_1 . So, $\beta = \beta_1 \alpha_3 \dots \alpha_k$ is the permutation in S_{n+1} constructed in this case.

The following theorem is a direct result from Construction 2.2.

Theorem 2.3. *For any $\alpha \in S_n$, Construction 2.2 gives $n+1$ distinct permutations in S_{n+1} . In fact, if we do it for all $\alpha \in S_n$, we will get $(n+1)!$ elements, which are exactly all permutations in the group S_{n+1} .*

We define the following notation $[\alpha, j]$, which will be used in the next section.

Definition 2.4. Let α be a permutation in S_n . Denote by $[\alpha, j]$ the permutation in S_{n+1} constructed from α , where j is an integer, $0 \leq j \leq n$.

- (1) $j = 0$ $[\alpha, 0]$ corresponds to the **Case 0** in Construction 2.2.
- (2) $j > 0$ $[\alpha, j]$ corresponds to **Case 1,2** by cutting the arrow a such that $t(a) = j$.

3. W -OPERATOR

We first review the definition and some properties about the W -operator $W([n])$. Details can be found in [10, 11]. Then, we will show how to calculate $W([n+1])$ from $W([n])$ by induction, which relates to Construction 2.2 in the last section. In other words, the structure of the W -operator $W([n])$ is closely related to the permutation group S_n and the quivers we studied in §2.

Let X be an infinite matrix with variable X_{ab} in the (a, b) -entry, i.e. $X := (X_{ab})_{a \geq 1, b \geq 1}$. Given a positive integer k , denote by p_k the trace of X^k ,

$$p_k = \sum_{a_1, \dots, a_k \geq 1} X_{a_1 a_k} X_{a_k a_{k-1}} \cdots X_{a_2 a_1}.$$

Clearly, p_k is a power series in $\mathbb{C}[[X_{ab}]]_{a, b \geq 1}$.

The operator matrix D is the infinite matrix with D_{ab} in the (a, b) -entry, where

$$D_{ab} = \sum_{c=1}^{\infty} X_{ac} \frac{\partial}{\partial X_{bc}}.$$

In the rest of the paper, we prefer to write $D_{ab} = X_{ac} \frac{\partial}{\partial X_{bc}}$ with the sum over c implied. As differential operators, the normal ordered product of D_{ab} and D_{cd} is defined as

$$: D_{ab} D_{cd} := X_{ae_1} X_{ce_2} \frac{\partial}{\partial X_{be_1}} \frac{\partial}{\partial X_{de_2}}.$$

It means that we always calculate the differentiation first. The normal ordered product $: D_{a_{n+1} a_n} D_{a_n a_{n-1}} \cdots D_{a_2 a_1} :$ is defined similarly.

Definition 3.1. For any positive integer n , the W -operator $W([n])$ is defined as follows

$$W([n]) := \frac{1}{n} : tr(D^n) := \frac{1}{n} \sum_{a_1, \dots, a_n \geq 1} : D_{a_1 a_n} D_{a_n a_{n-1}} \dots D_{a_2 a_1} : .$$

Before we calculate $W([n+1])$ by induction, we first review some important formulas.

Lemma 3.2 ([11]). *Let $F(p)$ be an element in $\mathbb{C}[p_1, p_2, \dots]$. We have*

$$D_{ab}F(p) = \sum_{k=1}^{\infty} k(X^k)_{ab} \frac{\partial F(p)}{\partial p_k}.$$

As a differential operator, the above formula of D_{ab} can be written as

$$D_{ab} = \sum_{k=1}^{\infty} k(X^k)_{ab} \frac{\partial}{\partial p_k}.$$

Lemma 3.3 ([11]).

$$D_{cd}(X^k)_{ab} = \sum_{j=0}^{k-1} (X^j)_{ad} (X^{k-j})_{cb}.$$

In particular, Lemma 3.3 implies the following formula

$$\begin{aligned} \sum_{k_j=1}^{\infty} D_{a_{n+1} a_n} (X^{k_j})_{a_i a_j} &= \sum_{k_j=1}^{\infty} \sum_{k_n=0}^{k_j-1} (X^{k_n})_{a_i a_n} (X^{k_j-k_n})_{a_{n+1} a_j} \\ &= \sum_{k_j=1}^{\infty} \sum_{k_n=1}^{\infty} (X^{k_n})_{a_i a_n} (X^{k_j})_{a_{n+1} a_j} \end{aligned}$$

by setting $a = a_i, b = a_j, c = a_{n+1}, d = a_n$.

Lemma 3.4 ([10, 11]).

$$\begin{aligned} D_{a_{n+2} a_{n+1}} D_{a_i a_j} &= \sum_{k \geq 1, j \geq 0} ((k+j)(X^j)_{a_i a_{n+1}} (X^k)_{a_{n+2} a_j} \frac{\partial}{\partial p_{k+j}}) \\ &\quad + \sum_{k, j \geq 1} (kj(X^k)_{a_{n+2} a_{n+1}} (X^j)_{a_i a_j} \frac{\partial^2}{\partial p_k \partial p_j}), \\ : D_{a_{n+2} a_{n+1}} D_{a_i a_j} : &= \sum_{k, j \geq 1} ((k+j)(X^j)_{a_i a_{n+1}} (X^k)_{a_{n+2} a_j} \frac{\partial}{\partial p_{k+j}}) \\ &\quad + \sum_{k, j \geq 1} (kj(X^k)_{a_{n+2} a_{n+1}} (X^j)_{a_i a_j} \frac{\partial^2}{\partial p_k \partial p_j}). \end{aligned}$$

Remark 3.5. The calculation of $D_{a_{n+2} a_{n+1}} D_{a_i a_j}$ in Lemma 3.4 is a result of Lemma 3.2 and 3.3. The operator $D_{a_{n+2} a_{n+1}}$ acts on the formula in Lemma 3.2. Since $D_{a_{n+2} a_{n+1}} D_{a_i a_j}$ is a differential operator, the chain rule gives us the result, where the first line comes from the action on the polynomial part (Lemma 3.3) and the action on the differential part gives us the second line (Lemma 3.2).

Notice that the only difference between the above two formulas in Lemma 3.4 is that the subscript j in the first sum starts from 0 in the first formula, while it goes from 1 in the second

one. In fact, the formula of normal ordered product $: D_{a_{n+2}a_{n+1}} D_{a_i a_j} :$ in Lemma 3.4 comes from $D_{a_{n+2}a_{n+1}} D_{a_i a_j}$ by subtracting an extra term. More precisely, by definition of normal ordered product, we have the following equation

$$D_{a_{n+2}a_{n+1}} D_{a_{n+1}a_{n-1}} =: D_{a_{n+2}a_{n+1}} D_{a_{n+1}a_{n-1}} : + X_{a_{n+2}e_1} \left[\frac{\partial}{\partial X_{a_{n+1}e_1}}, X_{a_{n+1}e_2} \right] \frac{\partial}{\partial X_{a_n e_2}}.$$

The last term is exactly the term missing in the normal ordered product.

As for the normal ordered product $: D_{a_{n+2}a_{n+1}} \cdots D_{a_2 a_1} :$, the reader can use the same approach to calculate $: D_{a_{n+2}a_{n+1}} \cdots D_{a_2 a_1} :$ from the product $D_{a_{n+2}a_{n+1}} \cdots D_{a_2 a_1}$. In the formula of normal ordered product $: D_{a_{n+2}a_{n+1}} \cdots D_{a_2 a_1} :$, the sum always goes from 1 to infinity, while some subscripts start from 0 in the formula of $D_{a_{n+2}a_{n+1}} \cdots D_{a_2 a_1}$.

Remark 3.6. In this remark, we will briefly explain how the above formulas relates to Construction 2.2. We fix a permutation $\alpha \in S_n$, positive integers k, k_j , and a_j $1 \leq j \leq n$. Let $\hat{Q}_\alpha = (\hat{V}_\alpha, \hat{A}_\alpha)$ be the quiver we defined in Construction 2.1. We consider a special differential operator

$$\prod_{b \in \hat{A}_\alpha} (X^{k_j})_{a_s(b) a_t(b)} \frac{\partial}{\partial p_k},$$

where the polynomial part $\prod_{b \in \hat{A}_\alpha} (X^{k_j})_{a_s(b) a_t(b)}$ corresponds to the quiver \hat{Q}_α .

We want to calculate $D_{a_{n+2}a_{n+1}} \left(\prod_{b \in \hat{A}_\alpha} (X^{k_j})_{a_s(b) a_t(b)} \frac{\partial}{\partial p_k} \right)$. By the chain rule, the operator $D_{a_{n+2}a_{n+1}}$ will act on the polynomial part $\prod_{b \in \hat{A}_\alpha} (X^{k_j})_{a_s(b) a_t(b)}$ and the differential part $\frac{\partial}{\partial p_k}$ separately.

- (1) When $D_{a_{n+2}a_{n+1}}$ acts on the differential part, the formula in Lemma 3.2 gives us the result. In the language of quivers, we add one more arrow $a_{n+2} \rightarrow a_{n+1}$ to the quiver \hat{Q}_α , which corresponds to **Case 0** in Construction 2.2.
- (2) When $D_{a_{n+2}a_{n+1}}$ acts on the polynomial part, assume that it acts on $(X^{k_j})_{a_i a_j}$. Lemma 3.3 gives us the calculation. In the language of quivers, it means that we cut the arrow $i \rightarrow j$ and add two arrows $i \rightarrow n$ and $n+1 \rightarrow j$, which corresponds to **Case 1, 2** in Construction 2.2.

This is the reason why we do Construction 2.2 in §2.

Now we are ready to calculate the W -operator $W([n])$ by induction.

Theorem 3.7. $W([n])$ is a well-defined operator on $\mathbb{C}[p_1, p_2, \dots]$. It can be written as the sum of $n!$ summations FS_α , each of which corresponds to a unique quiver \hat{Q}_α , or equivalently a unique permutation $\alpha \in S_n$.

Proof. To calculate $W([n])$, we have to figure out the formula of $: D_{a_1 a_n} D_{a_n a_{n-1}} \cdots D_{a_2 a_1} :$ for any positive integer $a_i \geq 1$, $1 \leq i \leq n$. By Lemma 3.4 and Remark 3.5, it is equivalent for us to calculate the product $D_{a_1 a_n} D_{a_n a_{n-1}} \cdots D_{a_2 a_1}$. Since we want to use induction to calculate this product, we replace $D_{a_1 a_n}$ by $D_{a_{n+1} a_n}$.

Let's start from the base step. When $n = 1$, by Lemma 3.2, we have

$$D_{a_2 a_1} = \sum_{k_1=1}^{\infty} k_1 (X^{k_1})_{a_2 a_1} \frac{\partial}{\partial p_{k_1}}.$$

We associate this summation to the quiver

$$\hat{Q}_{(1)} : 2 \longrightarrow 1 ,$$

which corresponds to the subscript of $(X^{k_1})_{a_2 a_1}$. Note that there is only one summation. Thus we define

$$FS'_{(1)} = \sum_{k_1=1}^{\infty} k_1 (X^{k_1})_{a_2 a_1} \frac{\partial}{\partial p_{k_1}} .$$

Replacing a_2 by a_1 and taking the sum over a_1 , we have

$$W([1]) = \underbrace{\sum_{k_1 \geq 1} k_1 p_{k_1} \frac{\partial}{\partial p_{k_1}}}_{FS_{(1)}} .$$

Denote by $FS_{(1)}$ the summation in $W([1])$ corresponding to $FS'_{(1)}$ in $D_{a_2 a_1}$. Note that $FS'_{(1)} = D_{a_2 a_1}$ in the base case. This is for the base case $n = 1$.

We do the next step $n = 2$ in fully details. The induction step follows directly from the discussion for $n = 2$. When $n = 2$, we have to calculate $D_{a_3 a_2} D_{a_2 a_1}$,

$$D_{a_3 a_2} D_{a_2 a_1} = \sum_{k_1=1}^{\infty} \left(D_{a_3 a_2} (k_1 (X^{k_1})_{a_2 a_1}) \right) \frac{\partial}{\partial p_{k_1}} + \sum_{k_1=1}^{\infty} k_1 (X^{k_1})_{a_2 a_1} \left(D_{a_3 a_2} \circ \frac{\partial}{\partial p_{k_1}} \right) .$$

By Lemma 3.4, we have

$$\begin{aligned} D_{a_3 a_2} D_{a_2 a_1} &= \sum_{k_1 \geq 1, k_2 \geq 0} ((k_1 + k_2) (X^{k_2})_{a_2 a_2} (X^{k_1})_{a_3 a_1} \frac{\partial}{\partial p_{k_1 + k_2}}) \\ &+ \sum_{k_1, k_2 \geq 1} (k_1 k_2 (X^{k_2})_{a_3 a_2} (X^{k_1})_{a_2 a_1}) \frac{\partial^2}{\partial p_{k_1} \partial p_{k_2}} . \end{aligned}$$

We associate the first summation to the quiver $\hat{Q}_{(1)(2)}$

$$\hat{Q}_{(1)(2)} : 2 \overset{\curvearrowright}{\longrightarrow} 2 , \quad 3 \longrightarrow 1 ,$$

which comes from the subscripts of the polynomial part $(X^{k_2})_{a_2 a_2} (X^{k_1})_{a_3 a_1}$. Similarly, the second summation corresponds to the quiver $\hat{Q}_{(12)}$

$$\hat{Q}_{(12)} : 3 \longrightarrow 2 \longrightarrow 1 .$$

We know that $D_{a_3 a_2}$ acting on $(X^{k_1})_{a_2 a_1}$ gives the first summation, which corresponds to **Case 1** of cutting the arrow $2 \rightarrow 1$ in $\hat{Q}_{(1)}$ in Construction 2.2. The same argument holds for the second summation, where $D_{a_3 a_2}$ acts on $\frac{\partial}{\partial p_{k_1}}$ and it corresponds to the **Case 0** in Construction 2.2.

By Lemma 3.4 and Remark 3.5, we know that : $D_{a_3 a_2} D_{a_2 a_1}$: and $D_{a_3 a_2} D_{a_2 a_1}$ are almost the same and the only difference comes from the term with subscript $j = 0$ in the first summation.

Hence, we can use quivers to describe the summations of $: D_{a_3 a_2} D_{a_2 a_1} :$ in the same way as $D_{a_3 a_2} D_{a_2 a_1}$. We use the notation FS'_α for the summation corresponding to $\alpha \in S_2$. We have

$$: D_{a_3 a_2} D_{a_2 a_1} := \sum_{\alpha \in S_2} FS'_\alpha.$$

In conclusion, we find that $: D_{a_3 a_2} D_{a_2 a_1} :$ can be written as the sum of two summations, which correspond to quivers \hat{Q}_α , $\alpha \in S_2$,

$$\begin{aligned} : D_{a_3 a_2} D_{a_2 a_1} := & \underbrace{\sum_{k_1, k_2 \geq 1} ((k_1 + k_2)(X^{k_2})_{a_2 a_2} (X^{k_1})_{a_3 a_1} \frac{\partial}{\partial p_{k+j}})}_{FS'_{(1)(2)}} \\ & + \underbrace{\sum_{k_1, k_2 \geq 1} (k_1 k_2 (X^{k_2})_{a_3 a_2} (X^{k_1})_{a_2 a_1}) \frac{\partial^2}{\partial p_{k_1} \partial p_{k_2}}}_{FS'_{(12)}}. \end{aligned}$$

Replacing a_3 by a_1 and taking the sum over a_1, a_2 , we have

$$W([2]) = \underbrace{\frac{1}{2} \sum_{k_1, k_2 \geq 1} (k_1 + k_2) p_{k_1} p_{k_2} \frac{\partial}{\partial p_{k_1 + k_2}}}_{FS_{(1)(2)}} + \underbrace{\frac{1}{2} \sum_{k_1, k_2 \geq 1} k_1 k_2 p_{k_1 + k_2} \frac{\partial^2}{\partial p_{k_1} \partial p_{k_2}}}_{FS_{(12)}}.$$

We define FS_α to be summation of $W([2])$ which corresponds to FS'_α in the formula of $: D_{a_3 a_2} D_{a_2 a_1} :$.

When $n = 3$, we can calculate the normal ordered product $: D_{a_4 a_3} D_{a_3 a_2} D_{a_2 a_1} :$ from the product $D_{a_4 a_3} D_{a_3 a_2} D_{a_2 a_1}$ similarly. We use the operator $D_{a_4 a_3}$ acting on $D_{a_3 a_2} D_{a_2 a_1}$. Since the polynomial part of each summation is a product of two terms (or the corresponding quiver has two arrows), we will get three new summations by the chain rule where two comes from the polynomial part and one from the differential part. By Theorem 2.3 and Remark 3.6, each of the new summation corresponds to a unique permutation in S_3 and this correspondence is a one-to-one correspondence. Thus the normal ordered product $: D_{a_4 a_3} D_{a_3 a_2} D_{a_2 a_1} :$ can be written as the sum of summations FS'_α labelled by permutations α in S_3 . Replacing a_4 by a_1 and taking the sum over a_1, a_2, a_3 , we get the formula for $W([3])$. We define FS_α to be the summation of $W([3])$, which corresponds to the summation FS'_α . In this way, the operator $W([n])$ can be written as the sum of $n!$ summations by induction, and each summation corresponds to a unique permutation in S_n .

At the end of the proof, we will show that $W([n])$ is a well defined operator on $\mathbb{C}[p_1, p_2, \dots]$ by giving two examples. When $n = 1$, consider $D_{a_2 a_1}$. Let $a_2 = a_1$ and take the sum over a_1 . We have

$$W([1]) = \sum_{k_1} k_1 p_{k_1} \frac{\partial}{\partial p_{k_1}}.$$

Now we consider $: D_{a_3 a_2} D_{a_2 a_1} :$. Let $a_3 = a_1$ and sum over a_1, a_2 . We have

$$W([2]) = \frac{1}{2} \sum_{a_1, a_2 \geq 1} : D_{a_1 a_2} D_{a_2 a_1} := \frac{1}{2} \sum_{k_1, k_2 \geq 1} ((k_1 + k_2) p_{k_1} p_{k_2} \frac{\partial}{\partial p_{k_1 + k_2}} + k_1 k_2 p_{k_1 + k_2} \frac{\partial^2}{\partial p_{k_1} \partial p_{k_2}}).$$

Clearly, $W([1])$ and $W([2])$ are well-defined operator on $\mathbb{C}[p_1, p_2, \dots]$. The operator $W([n])$ can be proved to be a well-defined operator on $\mathbb{C}[p_1, p_2, \dots]$ by induction. \square

4. DEGREE OF SUMMATION FS_α

Definition 4.1. Given any summation FS_α of $W([n])$, define $dP(FS_\alpha)$ to be the degree of its polynomial part and $dD(FS_\alpha)$ the order of its differential part. The degree $d(FS_\alpha)$ of the summation FS_α is $d(FS_\alpha) = dP(FS_\alpha) + dD(FS_\alpha)$.

We give an easy example to explain this definition. Consider the summation

$$FS_{(1)(2)} = \frac{1}{2} \sum_{k_1, k_2 \geq 1} (k_1 + k_2) p_{k_1} p_{k_2} \frac{\partial}{\partial p_{k_1+k_2}}.$$

We have

$$dP(FS_\alpha) = 2, \quad dD(FS_\alpha) = 1, \quad d(FS_\alpha) = 3.$$

Similarly, the degree data of $FS_{(12)}$ is

$$dP(FS_\alpha) = 1, \quad dD(FS_\alpha) = 2, \quad d(FS_\alpha) = 3.$$

The following lemma describes the relation between the degree of FS_β and FS_α , where $\beta = [\alpha, j]$ (see Definition 2.4).

Lemma 4.2. For any $\alpha \in S_n$,

(1) If $[\beta] = [\alpha, 0]$, we have

$$dP(FS_\beta) = dP(FS_\alpha), \quad dD(FS_\beta) = dD(FS_\alpha) + 1, \quad d(FS_\beta) = d(FS_\alpha) + 1.$$

(2) If $[\beta] = [\alpha, j]$ and j is a vertex in the chain of \hat{Q}_α , then, we have

$$dP(FS_\beta) = dP(FS_\alpha) + 1, \quad dD(FS_\beta) = dD(FS_\alpha), \quad d(FS_\beta) = d(FS_\alpha) + 1.$$

(3) If $[\beta] = [\alpha, j]$ and j is not a vertex in the chain of \hat{Q}_α , we have

$$dP(FS_\beta) = dP(FS_\alpha) - 1, \quad dD(FS_\beta) = dD(FS_\alpha), \quad d(FS_\beta) = d(FS_\alpha) - 1.$$

Proof. Notice that the $dP(FS_\alpha)$ is exactly the number of disjoint cycles of α .

When $j = 0$, the differential degree of FS'_β increases by one by Lemma 3.2. The disjoint cycle of $\beta = [\alpha, 0]$ is the same as α . Hence, $dP(FS_\beta) = dP(FS_\alpha)$.

When $j \geq 1$, Lemma 3.3 and Remark 3.6 implies that the operator $D_{a_n+n a_{n+1}}$ fixes the differential degree. Now we consider the polynomial degree. If j is in the chain of \hat{Q}_α , **Case 1** in Construction 2.2 tells us that β has one more disjoint cycle than α . When j is not in the chain of \hat{Q}_α , β corresponds to **Case 2** and $dP(FS_\beta) = dP(FS_\alpha) - 1$. This finishes the proof of the lemma. \square

From the above lemma, it is easy to find that the highest degree of summations in $W([n])$ is $n + 1$ and the other possible degrees are $n - 1, n - 3, \dots$.

Definition 4.3 (Ordinary Summation). Let α be a permutation in S_n . We say that FS_α is an *ordinary summation* (OS), if $d(FS_\alpha) = n + 1$.

An ordinary summation FS_α is of *type* $(r, n - r + 1)$ if $dP(FS_\alpha) = r$ and $dD(FS_\alpha) = n - r + 1$.

5. NONCROSSING PERMUTATION

In this section, we prove that if the permutation α is a noncrossing permutation if and only if FS_α is of maximal degree. As a corollary of this correspondence, we show that the number of ordinary summations with maximal degree is the Catalan number.

The noncrossing permutation comes from noncrossing partition with respect to a fixed order of objects. Recall that a partition of $[n] = \{1, \dots, n\}$ is *noncrossing* if whenever four elements, $1 \leq a < b < c < d \leq n$, are such that a, c are in the same block and b, d are in the same block, then the two blocks coincide. With respect to the natural order of integers, each noncrossing partition corresponds to a unique permutation, where each block corresponds to a disjoint cycle and the order $i > j$ implies an arrow $i \rightarrow j$ in the disjoint cycle. Here is the definition of noncrossing permutation.

Definition 5.1 (Noncrossing permutation). Let α be a permutation in S_n . Let $\alpha = \alpha_1 \dots \alpha_r$ be the decomposition of α into disjoint cycles. The permutation α is a *noncrossing permutation* if it satisfies the following conditions

- ($*_1$) For each arrow a in the unique chain of \hat{Q}_α , we have $t(a) < s(a)$, and there is only one arrow b in each loop of \hat{Q}_α such that $s(b) < t(b)$.
- ($*_2$) Given any two distinct cycles α_i and α_j , they satisfy at least one of the following conditions:
 - (a) pick an arbitrary element m in α_i , either $m > n$ for any n in α_j or $m < n$ for any n in α_j ;
 - (b) pick an arbitrary element m in α_j , either $m > n$ for any n in α_i or $m < n$ for any n in α_i .

Condition ($*_1$) means that we have an order on the finite set which determines the permutation. Condition ($*_2$) corresponds to the *noncrossing* condition. If α_i and α_j satisfy only one of the condition (a), (b), then then one is contained in the other one. If they satisfy both these two conditions, they are disjoint. For instance, consider the following examples,

$$\tau_1 = (123)(45), \quad \tau_2 = (125)(34), \quad \tau_3 = (124)(35).$$

The two disjoint cycles in τ_1 satisfies both these two conditions. If we consider the integers are in a straight line

$$(\ 5 \ 4 \) \ (\ 3 \ 2 \ 1 \),$$

the disjoint cycles (123) and (45) are disjoint.

The permutation τ_2 is an example that (125) and (34) only satisfy one of the conditions (a), (b). It is obvious that the disjoint cycle (34) is contained in (125),

$$(\ 5 \ (\ 4 \ 3 \) \ 2 \ 1 \).$$

Now we will prove that FS_α is an ordinary summation if and only if α is a noncrossing permutation in several steps.

Lemma 5.2. *Given $\alpha \in S_n$, if FS_α is an OS, then α satisfies the condition ($*_1$).*

Proof. We prove this lemma by induction on n . For the base step $n = 1$, $\hat{Q}_{(1)}$ is the only quiver and $FS_{(1)}$ is an OS. There is only one arrow $2 \rightarrow 1$ in the quiver $\hat{Q}_{(1)}$. Clearly, (1) satisfies the condition ($*_1$).

By induction, we assume that, for all $\alpha \in S_{k-1}$, if FS_α is an OS, then α satisfies ($*_1$). Let $\beta = [\alpha, j]$ be a permutation in S_k . The condition FS_β is an OS implies that FS_α is also an


OS. Indeed, if FS_α is not an OS, then $d(FS_\alpha) < k$. By Lemma 4.2, $d(FS_\beta) < k + 1$, violating the fact that β is an OS.

Let $\alpha = \alpha_1 \dots \alpha_r$ be the decomposition of α into disjoint cycles with $1 \in \alpha_1$. By Lemma 4.2, the integer j is either zero or the target of some arrow in the chain of \hat{Q}_α . Now we discuss these two cases.

- (1) If $j = 0$, then $\beta = \beta_1 \alpha_2 \dots \alpha_r$, where \hat{Q}_{β_1} is constructed from \hat{Q}_{α_1} by adding another arrow $k + 1 \rightarrow k$. By induction, the statement is true.
- (2) If $j \neq 1$, then β is constructed from α by cutting the arrow $a : i \rightarrow j$, which is an arrow a in the chain of \hat{Q}_α . We use the same notation as **Case 1** in Construction 2.2. Let $\beta = \beta_1 \beta_2 \alpha_2 \dots \alpha_r$. The quiver \hat{Q}_{β_1} of the cycle β_1 is

$$\hat{Q}_{\beta_1} : \quad k + 2 \longrightarrow j \longrightarrow \cdots \longrightarrow 1 ,$$

where $j \rightarrow \cdots \rightarrow 1$ is a subquiver of α_1 . Hence, all arrows in this chain satisfy that the source is larger than the target. The quiver \hat{Q}_{β_2} is

$$\hat{Q}_{\beta_2} : \quad i \longrightarrow k + 1 \longrightarrow \cdots \longrightarrow ,$$


where $k + 1 \rightarrow \cdots \rightarrow i$ is a subquiver of α_1 by construction. So the only arrow a in the cycle \hat{Q}_{β_2} satisfying $s(a) < t(a)$ is $i \rightarrow k + 1$. Hence, the statement is true for $n = k$. \square

Lemma 5.3. *If FS_α is an OS, then α satisfies the condition $(*_2)$.*

Proof. Similar to the proof of Lemma 5.2, we prove this lemma by induction on the permutation group S_n . When $n = 1$, it is clear that the unique permutation (1) in S_1 satisfies the condition $(*_2)$.

By induction, we assume that for all $\alpha \in S_{k-1}$ if FS_α is an OS, then α satisfies $(*_2)$. Let $\beta = [\alpha, j] \in S_k$. Let $\alpha = \alpha_1 \dots \alpha_r$ be the decomposition of α into disjoint cycles. We will prove that if FS_β is an OS, then β satisfies the condition $(*_2)$. Before we give the proof, we remind the reader that if $[\beta] = [\alpha, j]$ and FS_β is an OS, then FS_α is also an OS. This property comes from the proof of Lemma 5.2.

If $j = 0$, then $\beta = \beta_1 \alpha_2 \dots \alpha_r$, where \hat{Q}_{β_1} is constructed from \hat{Q}_{α_1} by adding another arrow $k + 1 \rightarrow k$. In other words, we put another element k into the cycle α_1 (see Construction 2.2). By assumption that any two disjoint cycles of $\alpha \in S_{k-1}$ satisfy at least one of the conditions, we only have to check whether the pair (β_1, α_i) satisfies the condition $(*_2)$, $2 \leq i \leq r$. Since α_1 contains the smallest element 1, so if α_1 and α_i are disjoint, then any element in α_1 is smaller than any element in α_i . Since k is the largest element, so the statement is true for β_1 and α_i . Now we consider that α_1 and α_i are not disjoint. Since 1 is contained in α_1 , so α_i is contained in α_1 . Clearly, it still holds for β_1 and α_i . So, (β_1, α_i) satisfies the condition $(*_2)$.

Now let's consider the case that β is constructed from α by cutting the arrow $a : i \rightarrow j$ lying in the chain of α . We use the same notation as **Case 1** in Construction 2.2. Let $\beta = \beta_1 \beta_2 \alpha_2 \dots \alpha_r$. So, we have to check whether the following three types of pairs satisfy the condition:

$$(\beta_1, \beta_2), \quad (\beta_1, \alpha_i), \quad (\beta_2, \alpha_i),$$

where $2 \leq i \leq r$.

- (β_1, β_2)

Since FS_α is OS, all arrows a in \hat{Q}_{α_1} satisfy $t(a) < s(a)$. Hence, when cutting the arrow $i \rightarrow j$, any elements in β_2 is larger than any elements in β_1 . It is true in this case.

- (β_1, α_i)

By induction, we know that the lemma is true for (α_1, α_i) , $2 \leq i \leq r$. Since the elements of β_1 is a subset of the elements of α_1 , so it is true for (β_1, α_i) , $2 \leq i \leq r$.

- (β_2, α_i)

If β_2 is a single disjoint "one cycle" (k) , the statement is true. If $\beta_2 \neq (k)$, assume the largest element in β_2 except k is ϕ . If ϕ is smaller than the smallest element in α_i , then any element u except k in β_2 is smaller than any element in α_i . Also, k is larger than any element in α_i . Hence, the statement is true in this case. Now let's consider the case that ϕ is larger than the smallest element in α_i . By construction, ϕ is an element in α_1 , which contains 1. Hence, ϕ is larger than any elements in α_i by induction. Similarly, any other elements in β_2 is larger or smaller to all elements in α_i by induction. So, the statement is true.

This finishes the proof of this lemma. \square

Theorem 5.4. *The summation FS_α is an OS if and only if α is a noncrossing permutation.*

Proof. The "only if" part is exactly Lemma 5.2 and 5.3. So, we only have to prove the "if" part. We prove this theorem by induction on n .

When $n = 1$, it is easy to prove, since (1) is the only permutation.

By induction, we assume that if $\alpha \in S_{k-1}$ satisfies the condition (*), then FS_α is an OS. We will prove that if $\beta \in S_k$ satisfies the condition (*), then FS_β is an OS. Assume $[\beta] = [\alpha, j]$ for some α in S_{n-1} and some nonnegative integer j . We claim that j is 0 or in the chain of \hat{Q}_α (**Claim 1**). Also, we claim that α is a noncrossing permutation (**Claim 2**).

By **Claim 1**, j is 0 or in the chain of \hat{Q}_α . By **Claim 2**, FS_α is an ordinary summation. By Construction 2.2 and Lemma 4.2, we know FS_β is also an OS. Now we are going to prove these two claims.

Proof of Claim 1. If not, β is constructed from α by cutting arrow $a : i \rightarrow j$ which is not in the chain of \hat{Q}_α . Hence, by **Case 2** in Construction 2.2, we will get a long chain

$$k + 1 \longrightarrow j \longrightarrow \cdots \longrightarrow i \longrightarrow k \longrightarrow \cdots \longrightarrow 1 .$$

In this chain, we have $i < k$, which contradicts with our assumptions that β satisfies the condition $(*_1)$. So, j must be in the chain of \hat{Q}_α or $j = 0$. \square

Proof of Claim 2. By **Claim 1**, we know that $j = 0$ or j is in the chain of \hat{Q}_α . If $j = 0$, it is easy to prove α is a noncrossing permutation. We leave it for the reader. Now we assume that j is in the chain of \hat{Q}_α . With the same notation as in Construction 2.2, let $\beta = \beta_1 \beta_2 \alpha_2 \dots \alpha_r$ with $1 \in \beta_1$.

First, we have to check α satisfies the condition $(*_1)$. By the assumption of β , there is exactly one arrow a in the quiver of α_i such that $t(a) > s(a)$, where $2 \leq i \leq r$. So, we have to show all arrows a in the chain of \hat{Q}_α satisfying $t(a) < s(a)$. We assume that there is an arrow a in the chain of \hat{Q}_α such that $s(a) < t(a)$. If $t(a) \neq j$, then this arrow will be in either β_1 or β_2 , which contradicts with the assumption of β . If $t(a) = j$, then we get β_1

$$\hat{Q}_{\beta_1} : k + 1 \longrightarrow j \longrightarrow \cdots \longrightarrow 1 .$$

and β_2

$$\hat{Q}_{\beta_2} : \quad i \longrightarrow k \longrightarrow \cdots \longrightarrow .$$

Since $k > j > i$, so (β_1, β_2) does not satisfy the condition $(*_2)$. Hence, we have $t(a) < s(a)$ for each arrow a in the chain of \hat{Q}_α and there is exactly one arrow b in each loop of \hat{Q}_α such that $s(b) < t(b)$.

Now, we are going to prove that α satisfies the condition $(*_2)$. The problem pair is (α_1, α_i) , $2 \leq i \leq r$. By assumption, β_1 contains the smallest element 1 and β_2 contains the element k . Hence, by Construction 2.2 and Lemma 5.3, we know that any element in β_1 is smaller than any element in β_2 . Since β is a noncrossing permutation, so for any cycle α_i , $2 \leq i \leq r$, there are three possible cases

- α_i is contained in β_1 , i.e. if we pick an arbitrary element m in β_1 , then we have $m > n$ for any n in α_i or $m < n$ for any n in α_i ;
- α_i is contained in β_2 , i.e. if we pick an arbitrary element m in β_2 , then we have $m > n$ for any n in α_i or $m < n$ for any n in α_i ;
- α_i is disjoint with β_1 and β_2 , i.e. any element in α_i is larger than any element in β_1 and smaller than any element in β_2 .

In the first case, if α_i is "contained" in β_1 , then any element in β_2 is larger than any element in α_i , because the element in β_2 is always larger than the element in β_1 . By the construction of α_1 , the condition is true for (α_1, α_i) . The same argument holds for the second case. For the third case, β_1 and β_2 are constructed from α_1 by cutting the arrow with target j and add another element k . Hence, α_i is "contained" in α_1 . Hence, α satisfies the condition $(*_2)$. \square

\square

Corollary 5.5. *The number of $(r, n - r + 1)$ -type OS in $W([n])$ is the Narayana number:*

$$\frac{1}{n+1} \binom{n+1}{r} \binom{n-1}{r-1}.$$

The number of all ordinary summations in $W([n])$ is the Catalan number

$$\sum_{r \geq 1}^n \frac{1}{n+1} \binom{n+1}{r} \binom{n-1}{r-1} = \frac{1}{n+1} \binom{2n}{n}.$$

Proof. By Theorem 5.4, there is a one-to-one correspondence between the ordinary summation and noncrossing permutation (also noncrossing partition). The number of $(r, n - r + 1)$ -type OS is exactly the number of noncrossing partition with r blocks, which is the Narayama number $\frac{1}{n+1} \binom{n+1}{r} \binom{n-1}{r-1}$ [12]. The number of all ordinary summations in $W([n])$ is the Catalan number, which is the sum of Narayama number. \square

Acknowledgements

This project is based on part of the author's PhD thesis. The author want to thank his advisor Maarten Bergvelt for helpful discussions and invaluable suggestions.

REFERENCES

- [1] Goulden, I. P. (1994). A differential operator for symmetric functions and the combinatorics of multiplying transpositions. *Transactions of the American Mathematical Society*, 344(1), 421-440.
- [2] Goulden, I., Jackson, D. (1997). Transitive factorisations into transpositions and holomorphic mappings on the sphere. *Proceedings of the American Mathematical Society*, 125(1), 51-60.
- [3] Goulden, I. P., Jackson, D. M. (1999). A proof of a conjecture for the number of ramified coverings of the sphere by the torus. *Journal of Combinatorial Theory, Series A*, 88(2), 246-258.
- [4] Goulden, I. P., Jackson, D. M. (1999). The number of ramified coverings of the sphere by the double torus, and a general form for higher genera. *Journal of Combinatorial Theory, Series A*, 88(2), 259-275.
- [5] Goulden, I. P., Jackson, D. M., Vakil, R. (2005). Towards the geometry of double Hurwitz numbers. *Advances in Mathematics*, 198(1), 43-92.
- [6] Goulden, I. P., Jackson, D. M., Vainshtein, A. (2000). The number of ramified coverings of the sphere by the torus and surfaces of higher genera. *Annals of Combinatorics*, 4(1), 27-46.
- [7] Goulden, I., Jackson, D. (1997). Transitive factorisations into transpositions and holomorphic mappings on the sphere. *Proceedings of the American Mathematical Society*, 125(1), 51-60.
- [8] Goulden, I. P., Jackson, D. M. (2000). Transitive factorizations in the symmetric group, and combinatorial aspects of singularity theory. *European Journal of Combinatorics*, 21(8), 1001-1016.
- [9] Lando, S. K., Zvonkin, A. K. (2013). *Graphs on surfaces and their applications* (Vol. 141). Springer Science Business Media.
- [10] Mironov, A., Morozov, A., Natanzon, S. (2012). Algebra of differential operators associated with Young diagrams. *Journal of Geometry and Physics*, 62(2), 148-155.
- [11] Mironov, A. D., Morozov, A. Y., Natanzon, S. M. (2011). Complete set of cut-and-join operators in the Hurwitz-Kontsevich theory. *Theoretical and Mathematical Physics*, 166(1), 1-22.
- [12] Simion, R. (2000). Noncrossing partitions. *Discrete Mathematics*, 217(1-3), 367-409.
- [13] Sun, H. (2019). A formula about W -operator and its application to Hurwitz number. *Discrete Mathematics*, 342(3), 715-722.

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY
135 Xingang W Rd, BinJiang Lu, Haizhu Qu, Guangzhou Shi, Guangdong Sheng, China
E-mail address: sunh66@mail.sysu.edu.cn