

SOLVABLE NORMAL SUBGROUPS OF 2-KNOT GROUPS

J.A.HILLMAN

ABSTRACT. If X is an orientable, strongly minimal PD_4 -complex and $\pi_1(X)$ has one end then it has no nontrivial locally-finite normal subgroup. Hence if π is a 2-knot group then (a) if π is virtually solvable then either π has two ends or $\pi \cong \Phi$, with presentation $\langle a, t | ta = a^2t \rangle$, or π is torsion-free and polycyclic of Hirsch length 4; (b) either π has two ends, or π has one end and the centre $\zeta\pi$ is torsion-free, or π has infinitely many ends and $\zeta\pi$ is finite; and (c) the Hirsch-Plotkin radical $\sqrt{\pi}$ is nilpotent.

A 2-knot is a topologically locally flat embedding of S^2 in S^4 . The main result of this note is that if the group of a 2-knot is virtually solvable then it either has finite commutator subgroup or is torsion-free. All such groups are known, and in the torsion-free case the knot exteriors are determined up to homeomorphism by the knot group π together with the $Aut(\pi)$ -orbit of a meridian. With the exception of one group, such knots may be constructed by elementary surgery on a section of the mapping torus of a diffeomorphism of a 3-manifold; the cocore of the surgery is then a smooth 2-knot in a smooth homotopy 4-sphere. (The exceptional group is the group Φ of Examples 10 and 11 in [5], which are ribbon knots.) In many cases the homotopy 4-sphere is standard, but the question remains open in general. (This construction is perhaps the simplest source for possible counter-examples to the smooth 4-dimensional Poincaré conjecture [8].)

The first section establishes some notation. In §2 we show that if X is an orientable PD_4 -complex such that $\pi_1(X)$ has one end and the equivariant intersection pairing on $\pi_2(X)$ is 0 then $\pi_1(X)$ has no nontrivial locally-finite normal subgroup. This is used in §3 to show that if a 2-knot group π is virtually solvable then either π' is finite or $\pi \cong \Phi = Z*_2$, with presentation $\langle a, t | ta = a^2t \rangle$, or π is torsion-free polycyclic and of Hirsch length 4. (The final family of such groups was found in [14].) More generally, if S is an infinite solvable normal

1991 *Mathematics Subject Classification.* 57Q45.

Key words and phrases. centre, coherent, Hirsch-Plotkin radical, 2-knot, torsion.

subgroup and π is not itself solvable then $S \cong \mathbb{Z}^2$ or is virtually torsion-free abelian of rank 1. We show also that the Hirsch-Plotkin radical $\sqrt{\pi}$ of every 2-knot group is nilpotent. Finally we consider the centre $\zeta\pi$. If π has one end then $\zeta\pi \cong \mathbb{Z}^2$ or is torsion-free, of rank ≤ 1 . In particular, this is so if the commutator subgroup π' is infinite and $\zeta\pi$ has rank > 0 . If π has two ends $\zeta\pi$ has rank 1, and may be either \mathbb{Z} or $\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$, while if π has infinitely many ends $\zeta\pi$ is finite. We extend a construction of [27] to give examples with $\sqrt{\pi}$ cyclic of order q or $2q$, with q odd.

The group Φ is the only nontrivial higher dimensional knot group which is solvable and has deficiency 1. In §4 we sketch briefly how the determination of 2-knots with this group might extend to other groups with geometric dimension 2, and perhaps to all with deficiency 1. In view of the present limitations on 4-dimensional surgery techniques, we can expect only a classification up to s -concordance (relative s -cobordism of knots). With this proviso, we can show that a 2-knot whose group π is a 1-knot group is determined up to s -concordance, reflections and the Gluck ambiguity (whether a knot is determined by its exterior [7]) by π and the $Aut(\pi)$ -orbit of a meridian (Theorem 5). In particular, a 2-knot with group π and meridian μ is s -concordant to the Artin spin of a fibred 1-knot if and only if π' is finitely generated, $C_\pi(\mu) \cong \mathbb{Z}^2$ and $(\pi, C_\pi(\mu))$ is a PD_3 -pair (Corollary 6.1).

1. NOTATION

If G is a group let ζG and G' be the centre and commutator subgroup of G , respectively. Let $C_G(H)$ be the centralizer of a subgroup $H \leq G$. The product of locally-nilpotent normal subgroups of G is again a locally-nilpotent normal subgroup, by the Hirsch-Plotkin Theorem, and the Hirsch-Plotkin radical \sqrt{G} is the (unique) maximal such subgroup. (See Proposition 12.1.2 of [23].) In general, \sqrt{G} is elementary amenable, but need not be nilpotent.

If E is an elementary amenable group then it has a well-defined Hirsch length $h(E) \in \mathbb{N} \cup \{\infty\}$. (See [15] or Chapter 1 of [11].)

If X is a space let $\beta_i^{(2)}(X)$ be the i th L^2 Betti number of X , and let $\beta_i^{(2)}(G) = \beta_i^{(2)}(K(G, 1))$. (See [21] for a comprehensive exposition of L^2 -theory.)

2. PD_4 -COMPLEXES WITH $\chi = 0$

A PD_4 -complex X with fundamental group π is *strongly minimal* if the equivariant intersection pairing on $\pi_2(X)$ is 0, equivalently, if the homomorphism from $H^2(\pi; \mathbb{Z}[\pi])$ to $H^2(X; \mathbb{Z}[\pi])$ induced by the

classifying map $c_X : X \rightarrow K(\pi, 1)$ is an isomorphism [12]. Note that X is aspherical if and only if it is strongly minimal and $H^i(\pi; \mathbb{Z}[\pi]) = 0$ for $i \leq 2$.

Lemma 1. *Let X be a finite PD_4 -complex with fundamental group π . If $\chi(X) = 0$ and $\beta_1^{(2)}(\pi) = 0$ then X is strongly minimal.*

Proof. Since X is a finite complex the L^2 -Euler characteristic formula holds, and so $\chi(X) = \beta_2^{(2)}(X) - 2\beta_1^{(2)}(X)$. Hence $\beta_2^{(2)}(X) = 0$ also. Since $\beta_1^{(2)}(X) = \beta_1^{(2)}(\pi) = 0$, $\beta_2^{(2)}(X) \geq \beta_2^{(2)}(\pi) \geq 0$ and $\chi(X) = 0$, it follows that $\beta_2^{(2)}(X) = \beta_2^{(2)}(\pi)$. Hence $H^2(c_X; \mathbb{Z}[\pi])$ is an isomorphism, by part (3) of Theorem 3.4 of [11]. \square

The condition $\beta_1^{(2)}(\pi) = 0$ holds if π has an infinite normal subgroup which is amenable, or is finitely generated and of infinite index. (See Chapter 7 of [21].)

Lemma 2. *Let G be a group. If T is a locally finite normal subgroup of G then T acts trivially on $H^j(G; \mathbb{Z}[G])$, for all $j \geq 0$.*

Proof. If T is finite then $H^j(G; \mathbb{Z}[G]) \cong H^j(G/T; \mathbb{Z}[G/T])$, for all j , and the result is clear. Thus we may assume that T and G are infinite. Hence $H^0(G; \mathbb{Z}[G]) = 0$, and T acts trivially. We may write $T = \cup_{n \geq 1} T_n$ as a strictly increasing union of finite subgroups. Then there are short exact sequences [16]

$$0 \rightarrow \varprojlim^1 H^{s-1}(T_n; \mathbb{Z}[\pi]) \rightarrow H^s(T; \mathbb{Z}[\pi]) \rightarrow \varprojlim H^s(T_n; \mathbb{Z}[\pi]) \rightarrow 0.$$

Hence $H^s(T; \mathbb{Z}[\pi]) = 0$ if $s \neq 1$ and $H^1(T; \mathbb{Z}[\pi]) = \varprojlim^1 H^0(T_n; \mathbb{Z}[\pi])$, and so the LHS spectral sequence collapses to give $H^j(G; \mathbb{Z}[G]) \cong H^{j-1}(G/T; H^1(T; \mathbb{Z}[G]))$, for all $j \geq 1$. Let $g \in T$. We may assume that $g \in T_n$ for all n , and so g acts trivially on $H^0(T_n; \mathbb{Z}G)$, for all j and n . But then g acts trivially on $\varprojlim^1 H^0(T_n; \mathbb{Z}[\pi])$, by the functoriality of the construction. Hence every element of T acts trivially on $H^{j-1}(G/T; H^1(T; \mathbb{Z}[G]))$, for all $j \geq 1$. \square

If π is a group and M is a right $\mathbb{Z}[\pi]$ -module let \overline{M} be the conjugate left module, with π -action defined by $g.m = mg^{-1}$ for all $m \in M$ and $g \in \pi$.

Theorem 3. *Let X be an orientable PD_4 -complex with fundamental group π . If X is strongly minimal and π has one end then π has no non-trivial locally-finite normal subgroup.*

Proof. Let \tilde{X} be the universal cover of X , and let $C_* = C_*(X; \mathbb{Z}[\pi]) = C_*(\tilde{X}; \mathbb{Z})$ be the cellular chain complex of \tilde{X} . We assume that π acts

on the left on \tilde{X} , and so C_* is a complex of free left $\mathbb{Z}[\pi]$ -modules. Since π has one end, $H_s(\tilde{X}; \mathbb{Z}) = H_s(X; \mathbb{Z}[\pi]) = 0$ for $s \neq 0$ or 2 . Let $\Pi = \pi_2(X) \cong H_2(\tilde{X}; \mathbb{Z}) = H_2(X; \mathbb{Z}[\pi])$. Poincaré duality and c_X give an isomorphism $\Pi \cong \overline{H^2(X; \mathbb{Z}[\pi])}$. Since X is strongly minimal, this in turn is isomorphic to $H^2(\pi; \mathbb{Z}[\pi])$.

Suppose that π has a nontrivial locally-finite normal subgroup T . Let $g \in T$ have prime order p , and let $C = \langle g \rangle \cong Z/pZ$. Then C acts freely on \tilde{X} , which has homology only in degrees 0 and 2. On considering the homology spectral sequence for the classifying map $c_{\tilde{X}/C} : \tilde{X}/C \rightarrow K(C, 1)$, we see that $H_{i+3}(C; \mathbb{Z}) \cong H_i(C; \Pi)$, for all $i \geq 2$. (See Lemma 2.10 of [11].) Since C has cohomological period 2 and acts trivially on Π , by Lemma 1, there is an exact sequence

$$0 \rightarrow Z/pZ \rightarrow \Pi \rightarrow \Pi \rightarrow 0.$$

On the other hand, since π is finitely presentable, $\Pi \cong H^2(\pi; \mathbb{Z}[\pi])$ is torsion-free, by Proposition 13.7.1 of [6]. Hence T has no such element g and so π has no such finite normal subgroup. \square

Corollary 3.1. *Every ascending locally finite subgroup of π is trivial.*

Proof. If T is an ascending locally finite subgroup of π then a transfinite induction shows that the normal closure of T in π is locally finite. \square

Theorems 8.1 and 9.1 of [11] are the bases for characterizations of the homotopy types of 4-dimensional infrasolvmanifolds and Seifert fibred 4-manifolds, respectively. Each assumes at some point that $H^2(\pi; \mathbb{Z}[\pi]) = 0$. We have long felt that this condition might be redundant. The results above justify this expectation for Theorem 9.1, which may now be stated as follows.

Theorem. *Let M be a closed 4-manifold with fundamental group π . If $\chi(M) = 0$ and π has an elementary amenable normal subgroup ρ with $h(\rho) = 2$ and $[\pi : \rho] = \infty$ then M is aspherical and ρ is virtually abelian. \square*

The key points are that π has one end, since $h(\rho) > 1$, and the quotient of ρ by its maximal locally-finite normal subgroup τ is virtually solvable, since $h(\rho) < \infty$. Hence $\tau = 1$, so ρ is virtually solvable, and then π has a nontrivial torsion-free abelian normal subgroup. Theorem 1.17 of [11] may then be used to show that $H^2(\pi; \mathbb{Z}[\pi]) = 0$, and the rest of the argument is as in [11].

We do not yet have a corresponding improvement of Theorem 8.1, as we do not know how to show that $H^2(\rho; W) = 0$ if ρ is elementary

amenable, has no nontrivial locally-finite normal subgroup and $h(\rho) = \infty$, and W is a free $\mathbb{Z}[\rho]$ -module. (This holds if ρ is torsion-free [20].)

3. CENTRES, HIRSCH-PLOTKIN RADICALS AND VIRTUALLY SOLVABLE 2-KNOT GROUPS

We work with topological manifolds and locally flat embeddings. Thus a 2-knot is an embedding $K : S^2 \rightarrow S^4$ which extends to a product neighbourhood $N(K) \cong S^2 \times D^2$. The knot exterior is $X(K) = \overline{S^4 \setminus N(K)}$, and the knot group is $\pi = \pi K = \pi_1(X(K))$. Let $M(K) = X(K) \cup D^3 \times S^1$ be the closed 4-manifold obtained by elementary surgery on K . Then $\pi_1(M(K)) \cong \pi$ and $\chi(M(K)) = 0$.

We can recover the exterior from the knot manifold $M(K)$ by specifying the isotopy class of the core $\{0\} \times S^1$. By general position, this is determined by the conjugacy class of a meridian. A *weight orbit* in a knot group π is the orbit of a normal generator under the action of $\text{Aut}(\pi)$. If each automorphism of π is realizable by a self-homeomorphism of $M(K)$ then $X(K)$ is determined by $M(K)$ together with the weight orbit of a meridian for K . (In general, a 2-knot group may have many distinct weight orbits, corresponding to distinct knots giving rise to the same knot manifold. See §7 of Chapter 14 of [11].)

If we reattach $S^2 \times D^2$ to $X(K)$ via an identification $\partial X(K) \cong S^2 \times S^1$ we may recover a 2-knot, represented by the core $S^2 \times \{0\}$. However there is a potential ambiguity of order 2 here, since $S^n \times S^1$ has an involution which does not extend across $S^n \times D^2$, when $n > 1$ [7]. If this involution extends across $X(K)$ then K is determined by its exterior; we say that K is *reflexive*.

Theorem 4. *Let K be a 2-knot with group $\pi = \pi K$. If π has normal subgroups $A \leq E$ with A a nontrivial abelian group and E an infinite elementary amenable group then either π' is finite or E is virtually torsion-free solvable. If $h(E) = 1$ then E is abelian or virtually \mathbb{Z} ; if $h(E) = 2$ then $E \cong \mathbb{Z}^2$, and if $h(E) > 2$ then E is torsion-free polycyclic, and $h(E) = 3$ or 4 .*

Proof. We may assume that π' is infinite. Then π has one end, and $\beta_1^{(2)}(\pi) = 0$, since π has an infinite amenable normal subgroup. Since $M(K)$ is a closed 4-manifold, it is homotopy equivalent to a finite PD_4 -complex, and so is strongly minimal, by Lemma 1. The torsion subgroup of A is characteristic, and so is normal in π . Hence A is torsion-free, by Theorem 3. Therefore either $\pi \cong \Phi$ or $M(K)$ is aspherical or $A \cong \mathbb{Z}$ and π/A has infinitely many ends, by Theorem 15.7

of [11]. If $M(K)$ is aspherical then E has finite cohomological dimension, and so is virtually solvable [15]. (See Corollary 1.9.2 of [11].) If π/A has infinitely many ends then E/A is finite, since it is an elementary amenable normal subgroup in π/A . In all cases E is in fact virtually torsion-free solvable.

Since π has one end, E has no non-trivial finite normal subgroup. Hence if $h(E) = 1$ then \sqrt{E} is torsion-free, abelian and of index ≤ 2 in E . If E is not finitely generated, neither is \sqrt{E} . In this case $\pi \cong \Phi$ or $M(K)$ is aspherical (by Theorem 15.7 of [11], as above), so π is torsion-free, and E must be abelian.

If $h(E) = 2$ then $M(K)$ is aspherical, E is torsion-free and π/E is virtually a PD_2 -group, by Theorems 9.1 and 16.2 of [11]. Since $M(K)$ is orientable, E cannot be the Klein bottle group, and so $E \cong \mathbb{Z}^2$. If $h(E) > 2$ then π is torsion-free polycyclic and $h(\pi) = 4$, by Theorem 8.1 of [11], so $h(E) = 3$ or 4 . \square

Corollary 4.1. *The following are equivalent:*

- (1) π is elementary amenable and $h(\pi) < \infty$;
- (2) π is virtually solvable;
- (3) either π' is finite or $\pi \cong \Phi$ or π is torsion-free polycyclic and $h(\pi) = 4$.

Proof. If π is elementary amenable and $h(\pi) < \infty$ then either π' is finite or π has one end. In the latter case π has no nontrivial locally-finite normal subgroup, by Lemma 1 and Theorem 3, and so π is virtually solvable, by Corollary 1.9.2 of [11].

If π has a solvable normal subgroup S of finite index then the lowest nontrivial term of the derived series for S is abelian, and is characteristic in S . Hence it is normal in π , and Theorem 4 applies.

Finally, (3) clearly implies the other conditions. \square

It is enough to assume that π is elementary amenable and has a nontrivial abelian normal subgroup. Can we relax “virtually solvable” further to “elementary amenable”, or even just “amenable”?

If $\pi \cong \Phi$ then K is TOP isotopic to Fox’s Example 10 or its reflection Example 11 [12], while if π is torsion-free polycyclic then it is one of the groups described in §4 of Chapter 16 of [11], and π determines $M(K)$ up to homeomorphism, by Theorem 17.4 of [11].

Corollary 4.2. *The Hirsch-Plotkin radical $\sqrt{\pi}$ is nilpotent.*

Proof. Since $\sqrt{\pi}$ is locally nilpotent it has a maximal locally-finite normal subgroup T with torsion-free quotient. If $\sqrt{\pi}$ is finitely generated there is nothing to prove. Otherwise, π has one end and so T is trivial, by Theorem 3. If $T = 1$ and $h(\sqrt{\pi}) \leq 2$ then $\sqrt{\pi}$ is abelian; if

$h(\sqrt{\pi}) > 2$ then π is virtually polycyclic and $h(\pi) = 4$, by Theorem 8.1 of [11]. In each case $\sqrt{\pi}$ is nilpotent. \square

Every finitely generated abelian group is the centre of some high-dimensional knot group [10]. On the other hand, the only 1-knots whose groups have nontrivial abelian normal subgroups are the torus knots, for which $\sqrt{\pi} = \zeta\pi \cong \mathbb{Z}$ and $\zeta\pi \cap \pi' = 1$. The intermediate case of 2-knots is less clear. If $\zeta\pi$ has rank > 1 then it is \mathbb{Z}^2 ; most twist spins of torus knots have such groups. There are examples with centre 1, $Z/2Z$, \mathbb{Z} or $\mathbb{Z} \oplus Z/2Z$. (See Chapters 15–17 of [11].)

Corollary 4.3. *Let K be a 2-knot with group $\pi = \pi K$. Then*

- (1) *if π has two ends then $\zeta\pi \cong \mathbb{Z} \oplus Z/2Z$ if π' has even order; otherwise $\zeta\pi \cong \mathbb{Z}$;*
- (2) *if π has one end then $\zeta\pi \cong \mathbb{Z}^2$, or is torsion-free of rank ≤ 1 ;*
- (3) *if π has infinitely many ends then $\zeta\pi$ is finite.*

Proof. If π has two ends then π' is finite, and so $\zeta\pi$ is finitely generated and of rank 1. It follows from the classification of such 2-knot groups (see Chapter 15 of [11]) that $\zeta\pi \cong \mathbb{Z} \oplus Z/2Z$ if π' has even order, and otherwise $\zeta\pi \cong \mathbb{Z}$.

Part (2) follows from Theorem 4, while part (3) is clear. \square

Note that π has finitely many ends if π' is finitely generated, or if $\zeta\pi$ is infinite. When π has more than one end Lemma 2.10 of [11] either does not lead to a contradiction or does not apply.

If $\zeta\pi$ is a nontrivial torsion group then it is finite. Yoshikawa constructed an example of a 2-knot whose group π has centre of order 2 [28]. It is easy to see that $\sqrt{\pi} = \zeta\pi$ in this case. The construction may be extended as follows. Let $q > 0$ be odd and let k_q be a 2-bridge knot such that the 2-fold branched cyclic cover of S^3 , branched over k_q is a lens space $L(3q, r)$, for some r relatively prime to q . Let $K_1 = \tau_2 k_q$ be the 2-twist spin of k_q , and let $K_2 = \tau_3 k$ be the 3-twist spin of a nontrivial knot k . Let γ be a simple closed curve in $X(K_1)$ with image $[\gamma] \in \pi K_1$ of order $3q$, and let w be a meridian for K_2 . Then w^3 is central in πK_2 . The group of the satellite of K_1 about K_2 relative to γ is the generalized free product

$$\pi = \pi K_2 / \langle\langle w^{3q} \rangle\rangle *_{w=[\gamma]} \pi K_1.$$

(See [17].) Hence $\sqrt{\pi} = \langle w^3 \rangle \cong Z/qZ$, while $\zeta\pi = 1$.

If we use a 2-knot K_1 with group $(Q(8) \times Z/3qZ) \rtimes_{\theta} \mathbb{Z}$ instead and choose γ so that $[\gamma]$ has order $6q$ then we obtain examples with $\sqrt{\pi} \cong Z/2qZ$ and $\zeta\pi = Z/2Z$. (Knots K_1 with such groups may be

constructed by surgery on sections of mapping tori of homeomorphisms of 3-manifolds with fundamental group $Q(8) \times Z/3qZ$ [27].)

If $\zeta\pi$ has rank 1 and nontrivial torsion then π' is finite, and $\zeta\pi$ is finitely generated.

If $\zeta\pi$ has rank 1 but is not finitely generated then $M(K)$ is aspherical. It is not known whether there are such 2-knots (nor, more generally, whether abelian normal subgroups of rank 1 in PD_n -groups with $n > 3$ must be finitely generated). What little we know about this case is as follows. Since $\zeta\pi < \pi'$ and $\pi/\pi' \cong \mathbb{Z}$, we must have $\zeta\pi \leq \pi''$. Since $\zeta\pi$ is torsion-free of rank 1 but is not finitely generated, *c.d.* $\zeta\pi = 2$. Hence if G is a nonabelian subgroup which contains $\zeta\pi$ then *c.d.* $G \geq 3$, by Theorem 8.6 of [2]. If H is a subgroup of π such that $H \cap \zeta\pi = 1$ then $H \cdot \zeta\pi \cong H \times \zeta\pi$ is not finitely generated, and so has infinite index in π . Hence *c.d.* $H \times \mathbb{Z} \leq c.d.H \times \zeta\pi \leq 3$ [24]. Theorem 5.5 of [2] gives, firstly, that *c.d.* $H \leq 2$, and then, that if H is FP_2 then *c.d.* $H \leq 1$, and so H is free. Thus if π is almost coherent every subgroup either meets $\zeta\pi$ nontrivially or is locally free.

If $\zeta\pi$ has rank > 1 then $M(K)$ is aspherical and $\zeta\pi \cong \mathbb{Z}^2$, by Theorem 16.3 of [11].

The following questions remain open:

- (i) if $\zeta\pi$ has rank 1, must it be finitely generated?
- (ii) if $\zeta\pi$ is finite, must it be $Z/2Z$ or 1?
- (iii) is there a 2-knot group π with $\sqrt{\pi}$ a non-cyclic finite group?
- (iv) if π is elementary amenable is it virtually solvable?

In each case the answer is “yes” if π' is finitely presentable, for then the infinite cyclic cover $M(K)'$ is homotopy equivalent to a PD_3 -complex, by Theorem 4.5 of [11].

4. GEOMETRIC DIMENSION 2

The only solvable 2-knot groups with deficiency 1 are \mathbb{Z} and Φ . One of the early triumphs of 4-dimensional surgery was Freedman’s unknotting criterion: 2-knots with group \mathbb{Z} are topologically trivial. In [12] we showed that any 2-knot K with group Φ is topologically equivalent to Fox’s Example 10 or its reflection (Example 11). The argument involved showing that

- (i) $\pi = \pi K$ determines the homotopy type of $M(K)$;
- (ii) $Wh(\pi) = 0$ and the surgery assembly homomorphisms $\mathcal{A}_*(\pi) : H_*(\pi; \mathbb{L}_0) \rightarrow L_*(\pi)$ are isomorphisms;
- (iii) 4-dimensional TOP surgery;
- (iv) π has an unique weight orbit, up to inversion; and
- (v) Fox’s examples are reflexive.

The group Φ has geometric dimension 2 ($g.d.\Phi = 2$): there is a finite 2-dimensional $K(\Phi, 1)$ complex. Our strategy for (i) uses the fact that $M(K)$ is strongly minimal if $g.d.\pi = 2$, since $\chi(M(K)) = 0 = 2\chi(\pi)$. It should extend to all knot groups π with $g.d.\pi = 2$, but at present is incomplete; there is a 2-torsion condition which holds for Φ , but is not otherwise easily verified. (See [13] for a much expanded exposition of [12] and earlier work.) When $g.d.\pi = 2$ and π' is finitely generated there is a quite different argument. For then π' is free of finite rank, and $M(K)$ is homotopy equivalent to the mapping torus of a self-homeomorphism of $\#^r(S^2 \times S^1)$, where r is the rank of π' . (See Corollary 4.5.3 of [11].)

Step (ii) holds if π is a 1-knot group, or is square root closed accessible, or is in the class \mathcal{X} of fundamental groups of finite graphs of groups with all vertex and edge groups \mathbb{Z} . (See Lemma 6.9 of [11].) Semidirect products $F(r) \rtimes \mathbb{Z}$ are square root closed accessible, while Φ is in \mathcal{X} .

At the time of writing, the largest known class of groups for which 4-dimensional topological surgery works is the class SG obtained from subexponential groups by taking increasing unions and extensions. To go beyond this class we must work modulo TOP s -cobordisms (or s -concordances of 2-knots) rather than expect homeomorphisms.

Step (iv) fails for other knot groups π with $g.d.\pi = 2$. In general, we must specify a weight orbit in order to recover $X(K)$ from $M(K)$.

The final step (v) is not satisfied by all 2-knots. However, those which have a Seifert hypersurface with free fundamental group are reflexive. Fibred 2-knots with π' free and ribbon 2-knots have such Seifert hypersurfaces [26].

The difficulty with (i) has been bypassed in [9], which shows directly that an orientable 4-manifold M with $\pi = \pi_1(M)$ such that $g.d.\pi = 2$ and π “satisfies properties W -AA” (i.e., (ii) holds) is determined up to s -cobordism by π , the second Wu class $w_2(M)$ and the equivariant intersection pairing on $\pi_2(M)$.

Theorem 5. *Let π be a 2-knot group with deficiency 1. If either*

- (1) π is the group of a 1-knot; or
- (2) π' is finitely generated; or
- (3) $\beta_1^{(2)}(\pi) = 0$ and π is square-root closed accessible,

then any 2-knot with group π is determined up to TOP s -concordance, reflections and the Gluck ambiguity by π and the weight orbit of a meridian.

Proof. We may assume that $\pi \not\cong \mathbb{Z}$.

If π is a 1-knot group then $g.d.\pi = 2$ and $\beta_1^{(2)}(\pi) = 0$, by Theorem 4.1 of [21], while $Wh(\pi) = 0$ and the surgery assembly homomorphisms $\mathcal{A}_*(\pi)$ are isomorphisms [1]. If π' is finitely generated then $\beta_1^{(2)}(\pi) = 0$, while if $\text{def}(\pi) = 1$ and $\beta_1^{(2)}(\pi) = 0$ then $g.d.\pi = 2$, by Theorem 2.5 of [11]. In particular, if π' is finitely generated and $\text{def}(\pi) = 1$ then π' is free and so $\pi \cong \pi' \rtimes \mathbb{Z}$ is square root closed accessible. Hence (2) \Rightarrow (3), which implies that $Wh(\pi) = 0$ and the $\mathcal{A}_*(\pi)$ are isomorphisms [3]. Thus in each case π satisfies properties *W-AA* of [9].

In each case π is torsion-free, and so has one end [19], and $\beta_1^{(2)}(\pi) = 0$. Hence $M(K)$ is strongly minimal, by Lemma 1. Since $M(K)$ is orientable, $w_2(M(K)) = 0$, the equivariant intersection pairing on $\pi_2(M(K))$ is trivial and π satisfies properties *W-AA*, π determines $M(K)$ up to *s-cobordism*, by the main result of [9]. More precisely, if K_0 and K_1 each have group π and their weight orbits correspond under an isomorphism $\theta : \pi K_0 \cong \pi K_1$ then there is a (5-dimensional) *s-cobordism* W from $M(K_0)$ to $M(K_1)$ such that $\pi_1(j_0) = \pi_1(j_1)\theta$, where $j_i : M(K_i) \rightarrow W$ is the natural inclusion, for $i = 0, 1$. The weight orbit of a meridian determines a properly embedded annulus in W with boundary components meridians for K_0 and K_1 . Excising an open product neighbourhood of this annulus gives a relative *s-cobordism* between the exteriors, and hence an *s-concordance* between the knots, modulo reflections and the Gluck ambiguity. \square

Beyond this, there is some reason to hope that all 2-knots with groups of deficiency 1 should be so determined. The exact state of affairs is contingent upon the outcome of several well-known conjectures. Let \mathcal{DW} be the class of groups with deficiency 1 and weight 1. Kervaire showed that every group in \mathcal{DW} is the group of a smooth knot in a smooth homotopy 4-sphere [18]. He first constructs a knot manifold as the boundary of a 5-manifold obtained from D^5 by attaching 1- and 2-handles, and so the groups are realized by homotopy ribbon knots.

If the *Andrews-Curtis Conjecture* is true then groups in \mathcal{DW} have Wirtinger presentations of deficiency 1, and so are groups of ribbon 2-knots [29]. (In particular, these are smooth knots in S^4 .) If the *Whitehead Conjecture* is true then $g.d.\pi \leq 2$, for all $\pi \in \mathcal{DW}$. (If $\pi \in \mathcal{DW}$ then $g.d.G = 2 \Leftrightarrow c.d.G = 2$, by Theorem 2.8 of [11].) If the *Fibred Isomorphism Conjecture* holds for groups of cohomological dimension 2 then (ii) holds for such groups.

A conjecture which has perhaps been less studied has implications for characterizing 1-knot groups, and thus for recognizing the Artin spins of 1-knots. If π is a nontrivial 1-knot group and $T \cong \mathbb{Z}^2$ is the

subgroup generated by a meridian-longitude pair (μ, λ) then (π, T) is a PD_3 -pair. (When the knot is prime the subgroup T is unique up to conjugacy.) If *finitely presentable PD_3 -groups are fundamental groups of aspherical closed 3-manifolds* then the converse holds: every PD_3 -pair (π, T) in which $T \cong \mathbb{Z}^2$ contains a normal generator is the group of a 1-knot.

This is known in the fibred case. The following result is essentially Theorem 5 of [25], which is in turn a reformulation of Theorem 9.2.3 of [22], taking into account first the determination of PD_2 -groups and pairs, and more recently Perelman's proof of the Poincaré Conjecture.

Theorem 6. *Let π be a group with deficiency 1 and weight 1, and suppose that $\pi \not\cong \mathbb{Z}$. Then the following are equivalent:*

- (1) π is the group of a fibred 1-knot ;
- (2) π has a subgroup $\Delta \cong \mathbb{Z}^2$ which contains a normal generator and such that (π, Δ) is a PD_3 -pair, and π' is finitely generated;
- (3) π has a subgroup $\Delta = \langle \mu, \lambda \rangle \cong \mathbb{Z}^2$, where μ is a normal generator and $\lambda \in \pi''$, and such that $(\pi', \langle \lambda \rangle)$ is a PD_2 -pair.

Proof. If $\pi \cong \pi k$, where k is a nontrivial 1-knot, then $X(k)$ is aspherical, $\partial X(k)$ is an incompressible torus, and π is normally generated by a meridian on $\partial X(k)$. Hence $\Delta = \pi_1(\partial X(k)) \cong \mathbb{Z}^2$ contains a normal generator and (π, Δ) is a PD_3 -pair. If $X(k)$ fibres over S^1 with fibre F then $\pi' \cong \pi_1(F)$ is finitely generated. Thus (1) implies (2).

If (2) holds then $\pi/\pi' \cong \mathbb{Z}$, and so T has a basis μ, λ , where μ is a normal generator and λ generates $T \cap \pi' \cong \mathbb{Z}$. In fact, λ is in π'' , since μ commutes with λ and $\pi/\pi' \cong \mathbb{Z}$. Since $\text{def}(\pi) = 1$ and π' is finitely generated, π' is free, and so is FP . Hence $(\pi', \langle \lambda \rangle)$ is a PD_2 -pair, by a Shapiro's Lemma argument, as in Theorem 1.19 of [11]. Thus (2) implies (3).

If (3) holds then $(\pi', \langle \lambda \rangle)$ is the fundamental group pair of a surface S with boundary $\partial S = S^1$ [4]. The group π is a semidirect product $\pi \cong \pi' \rtimes_{\theta} \mathbb{Z}$, where the monodromy θ fixes λ . Hence it is realized by a self-homeomorphism f of S , by Nielsen's Theorem. (See Theorem 5.7.1 of [30].) We may adjoin a solid torus to the mapping torus $M(f)$ to get a homotopy 3-sphere containing a knot with exterior $M(f)$. Thus π is the group of a fibred 1-knot, by the validity of the Poincaré Conjecture. \square

Corollary 6.1. *A nontrivial 2-knot with group π and meridian μ is s -concordant to the Artin spin of a fibred 1-knot if and only if π' is finitely generated, $C_{\pi}(\mu) \cong \mathbb{Z}^2$ and $(\pi, C_{\pi}(\mu))$ is a PD_3 -pair.*

Proof. This follows immediately from Theorems 5 and 6, since Artin spins of 1-knots are reflexive [7]. \square

REFERENCES

- [1] Aravinda, C.S., Farrell, F.T. and Roushon, S.K. Surgery groups of knot and link complements, *Bull. London Math. Soc.* 29 (1997), 400–406.
- [2] Bieri, R. *Homological Dimension of Discrete Groups*, Queen Mary College Lecture Notes in Mathematics, London (1976).
- [3] Cappell, S.E. Mayer-Vietoris sequences in Hermitean K -theory, in *Hermitean K -Theory and Geometric Applications* (edited by H.Bass), Lecture Notes in Mathematics 343, Springer-Verlag, Berlin - Heidelberg - New York (1973), 478–512.
- [4] Eckmann, B. and Müller, H. Poincaré duality groups of dimension two, *Comment. Math. Helvetici* 55 (1980), 510–520.
- [5] Fox, R.H. A quick trip through knot theory, in *Topology of 3-Manifolds and Related Topics* (edited by M.K.Fort, Jr), Prentice-Hall, Englewood Cliffs, N.J.(1962), 120–167.
- [6] Geoghegan, R. *Topological Methods in Group Theory*, Graduate Texts in Mathematics 243, Springer-Verlag, Berlin – Heidelberg – New York (2008).
- [7] Gluck, H. The embedding of two-spheres in the four-sphere, *Trans. Amer. Math. Soc.* 104 (1962), 308–333.
- [8] Gompf, R. More Cappell-Shaneson spheres are standard, *Alg. Geom. Top* 10 (2010), 1665–1681.
- [9] Hambleton, I., Kreck, M. and Teichner, P. Topological 4-manifolds with geometrically two-dimensional fundamental groups, *J. Topol. Anal.* 1(2009), 123–151.
- [10] Hausmann, J.-C. and Kervaire, M. Sur le centre des groupes de noeuds multidimensionnelles, *C.R. Acad. Sci. Paris* 287 (1978), 699–702.
- [11] Hillman, J. A. *Four-Manifolds, Geometries and Knots*, Geometry and Topology Monographs 5, Geometry and Topology Publications (2002). (Revisions 2007 and 2014).
- [12] Hillman, J.A. Strongly minimal PD_4 -complexes, *Top. Appl.* 156 (2009), 1565–1577.
- [13] Hillman, J.A. PD_4 -complexes and 2-dimensional duality groups, arXiv: 1303.5486 v7 [math.GT]
- [14] Hillman, J.A. and Howie, J. Seifert fibred knot manifolds, *J. Knot Theory Ramif.* 22 (2013), 1350082.
- [15] Hillman, J.A. and Linnell, P.A. Elementary amenable groups of finite Hirsch length are locally finite by virtually solvable, *J. Austral. Math. Soc.* 52 (1992), 237–241.
- [16] Jensen, C.U. *Les foncteurs dérivés de \varprojlim et ses applications a la théorie des modules*, Lecture Notes in Mathematics 254, Springer-Verlag, Berlin – Heidelberg – New York (1972).
- [17] Kanenobu, T. Groups of higher dimensional satellite knots, *J. Pure Appl. Alg.* 28 (1983), 179–188.
- [18] Kervaire, M.A. Les noeuds de dimensions supérieures, *Bull. Soc. Math. France* 93 (1965), 225–271.

- [19] Klyachko, A. Funny property of sphere and equations over groups, *Comm. Alg.* 21 (1993), 2555–2575.
- [20] Kropholler, P.H. Soluble groups of type $(FP)_{\infty}$ have finite torsion-free rank, *Bull. London Math. Soc.* 25 (1993), 558–566.
- [21] Lück, W. *L^2 -Invariants: Theory and Applications to Geometry and K -Theory*, *Ergebnisse der Mathematik und ihrer Grenzgebiete 3 Folge*, Bd. 44, Springer-Verlag, Berlin - Heidelberg - New York (2002).
- [22] Neuwirth, L.P. *Knot Groups*, *Ann. Math. Studies* 56, Princeton University Press, Princeton, N.J. (1965).
- [23] Robinson, D.J.S. *A Course in the Theory of Groups*, *Graduate Texts in Mathematics* 80, Springer-Verlag, Berlin – Heidelberg – New York (1982).
- [24] Strebel, R. A remark on subgroups of infinite index in Poincaré duality groups, *Comment. Math. Helvetici* 52 (1977), 317–324.
- [25] Thomas, C.B. Splitting theorems for certain PD^3 -groups, *Math. Z.* 186 (1984), 201–209.
- [26] Yanagawa, T. On ribbon 2-knots: the 3-manifold bounded by the 2-knot, *Osaka J. Math.* 6 (1969), 447-464.
- [27] Yoshikawa, K. On 2-knot groups with the finite commutator subgroups, *Math. Seminar Notes Kobe University* 8 (1980), 321–330.
- [28] Yoshikawa, K. On a 2-knot group with nontrivial centre, *Bull. Austral.Math. Soc.* 25 (1982), 321–326.
- [29] Yoshikawa, K. A note on Levine’s condition for knot groups, *Math. Sem. Notes Kobe University* 10 (1982), 633–636.
- [30] Zieschang, H., Vogt, E. and Coldewey, H.-D. *Surfaces and Planar Discontinuous Groups*, *Lecture Notes in Mathematics* 835, Springer-Verlag, Berlin – Heidelberg – New York (1980).

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF SYDNEY, NSW
2006, AUSTRALIA

E-mail address: jonathan.hillman@sydney.edu.au