

HECKE OPERATORS IN KK -THEORY AND THE K -HOMOLOGY OF BIANCHI GROUPS

BRAM MESLAND AND MEHMET HALUK ŞENGÜN

Abstract. Let Γ be a torsion-free arithmetic group acting on its associated global symmetric space X . Assume that X is of non-compact type and let Γ act on the geodesic boundary ∂X of X . Via general constructions in KK -theory, we endow the K -groups of the arithmetic manifold X/Γ , of the reduced group C^* -algebra $C_r^*(\Gamma)$ and of the boundary crossed product algebra $C(\partial X) \rtimes \Gamma$ with Hecke operators. The K -theory and K -homology groups of these C^* -algebras are related by a Gysin six-term exact sequence. In the case when Γ is a group of real hyperbolic isometries, we show that this Gysin sequence is Hecke equivariant. Finally, when Γ is a Bianchi group, we assign explicit unbounded Fredholm modules (i.e. spectral triples) to (co)homology classes, inducing Hecke-equivariant isomorphisms between the integral cohomology of Γ and each of these K -groups. Our methods apply to case $\Gamma \subset \mathbf{PSL}_2(\mathbf{Z})$ as well. As these results are achieved in the context of unbounded Fredholm modules, they shed light on noncommutative geometric aspects of the purely infinite boundary crossed product algebra.

Contents

	Page
Introduction	1
1. Prelude: Kasparov's spectral sequence	6
2. Hecke operators and KK -theory	8
3. Gysin sequence and Hecke operators	13
4. K -cycles for the reduced Bianchi group C^* -algebra	20
5. K -cycles for Bianchi manifolds	27
6. The case of $\mathbf{PSL}_2(\mathbf{Z})$	32
7. The extension class as a hypersingular integral operator	33
8. Embedded surfaces and K -cycles for the boundary crossed product	39
References	46

Introduction

Let \mathbf{G} be a semi-simple algebraic group over \mathbf{Q} and $\Gamma \subset \mathbf{G}(\mathbf{Q})$ be an arithmetic group. The cohomology of Γ can be equipped with a commuting family of endomorphisms called ‘‘Hecke operators’’ which arise from correspondences on the associated arithmetic manifold X/Γ . The cohomology of Γ as a Hecke module plays an important role in modern number theory. It is well-known that, as a Hecke module, the complex cohomology of Γ can be completely accounted for

Date: November 16, 2018.

2010 Mathematics Subject Classification. 11F75, 46L80, 19K35, 58B34, 53C23.

by various spaces of automorphic forms (see [25]). The Langlands programme predicts that classes in the complex (resp. mod p) cohomology of Γ that are simultaneous eigenvectors of the Hecke operators correspond to p -adic (resp. mod p) Galois representations. This is known in many cases (see, for example, [68]).

Let X denote the symmetric space associated to the real Lie group $\mathbf{G}(\mathbf{R})$. Assume that X is of non-compact type and let ∂X denote the geodesic boundary of X . We consider three C^* -algebras that are naturally associated to Γ , namely, the algebra $C_0(X/\Gamma)$ of functions on the arithmetic manifold X/Γ , the reduced group C^* -algebra $C_r^*(\Gamma)$ and the boundary crossed product algebra $C(\partial X) \rtimes \Gamma$, which we call the *arithmetic C^* -algebras* associated to Γ .

Via general constructions in KK -theory, we endow the K -groups of arithmetic C^* -algebras with Hecke operators. As in the case of the cohomology of arithmetic groups, we expect that these Hecke modules are related to the arithmetic of automorphic forms and to Galois representations. We show that this is the case when Γ is a Bianchi group by proving that the *integral* cohomology of Γ and the K -homology of the arithmetic C^* -algebras associated to Γ are isomorphic as Hecke modules. We realize these isomorphisms by constructing explicit unbounded Fredholm modules (i.e. spectral triples) for each of the arithmetic C^* -algebras. Our results also apply to the case of $\mathbf{PSL}_2(\mathbf{Z})$.

Bianchi groups are among the simplest generalizations of the $\mathbf{PSL}_2(\mathbf{Z})$, yet from the perspective of the Langlands programme, many fundamental questions are still wide open (see, for example, [70]) and moreover new phenomena ([9, 15, 68, 69]) that are not present in the setting of $\mathbf{PSL}_2(\mathbf{Z})$ (or, more generally, in the setting of Shimura varieties) arise. Our results allow the possibility of using ideas and tools from K -theory (both topological and analytic) and noncommutative geometry in investigating these problems and phenomena. For example, one of our results implies that, up to normalization, a period of a Bianchi (or classical) modular form can be captured as the index of a Fredholm operator.

Our work is inspired by the works of Manin and Marcolli [47, 48, 49] which pursue number theoretic questions around the ‘noncommutative modular curves’ via tools of Connes’ noncommutative geometry [18, 19]. The K -theory of C^* -algebras associated to Kleinian and Fuchsian groups have been studied by many authors e.g. [2, 10, 17, 23], with K -homology receiving attention only recently [24, 26, 62, 63]. The novelty of our work lies in the introduction of the Hecke module structure within KK -theory and the construction of new unbounded K -cycles using the unbounded Kasparov product [34, 53, 54]. This is of particular interest for the purely infinite boundary crossed product C^* -algebra $C(\partial X) \rtimes \Gamma$.

Description of results and plan of the paper. We first carry out a very general treatment of Hecke operators as they are crucial for the link with arithmetic that we are aiming to establish. This is done in Section 2. A novelty of our treatment is the introduction of Hecke operators via KK -theory. This proves to be a powerful, and natural, way of treating Hecke operators that allows us to establish various results related to Hecke operators in a robust and efficient manner.

Let G be a group acting on a locally compact Hausdorff space X . Given a subgroup Γ of G which acts freely and properly on X , put $M = X/\Gamma$. For any element g in the commensurator group of Γ in G , we define a bimodule T_g^M and let $[T_g^M] \in KK_0(C_0(M), C_0(M))$ be its class. Then we define the Hecke operator

$$T_g : K^*(C_0(M)) \rightarrow K^*(C_0(M)), \quad x \mapsto [T_g^M] \otimes x,$$

to be the Kasparov product with this class. Now let B an arbitrary $C_G(\Gamma)$ - C^* -algebra. Similarly we define a class $[T_g^\Gamma] \in KK_0(B \rtimes_r \Gamma, B \rtimes_r \Gamma)$ and define the Hecke operator

$$T_g : K^*(B \rtimes_r \Gamma) \rightarrow K^*(B \rtimes_r \Gamma), \quad x \mapsto [T_g^\Gamma] \otimes x,$$

as the Kasparov product with this class. Of course, in both cases, we obtain Hecke operators on K -theory as well. These two constructions allow us to define Hecke operators on the K -groups of arithmetic C^* -algebras.

Next, in Section 3, we study the Hecke equivariance of a certain Gysin exact sequence, which shall play an important role in our investigation. Similar exact sequences were studied by Emerson-Meyer [23] in K -theory and later by Emerson-Nica [24] in K -homology. Our treatment is again general here. Let Γ be a group acting freely and properly on hyperbolic n -space \mathbf{H}_n via isometries such that the hyperbolic manifold $M = \mathbf{H}_n/\Gamma$ has finite volume. Let $C_r^*(\Gamma)$ denote the reduced group C^* -algebra of Γ . Starting from the Γ -equivariant short exact sequence of C^* -algebras

$$0 \rightarrow C_0(\mathbf{H}_n) \rightarrow C(\mathbf{H}_n \cup \partial\mathbf{H}_n) \rightarrow C(\partial\mathbf{H}_n) \rightarrow 0,$$

we show the following in Theorem 3.10 below.

Theorem A. The K -homology Gysin exact sequence takes the form

$$\begin{array}{ccccccc} 0 & \longrightarrow & K^1(C_0(M)) & \longrightarrow & K^0(C(\partial\mathbf{H}_n) \rtimes \Gamma) & \longrightarrow & K^0(C_r^*(\Gamma)) \\ & & & & & & \downarrow \\ 0 & \longleftarrow & K^1(C_r^*(\Gamma)) & \longleftarrow & K^1(C(\partial\mathbf{H}_n) \rtimes \Gamma) & \longleftarrow & K^0(C_0(M)) \end{array}$$

and is *Hecke equivariant*.

This is done by representing the boundary extension as a Fredholm module constructed from the field of harmonic measures on the boundary. Since we work within KK -theory, the results of Section 2 and Section 3 have counterparts for K -theory groups as well.

At this point, we specialize to the case of Bianchi groups and study the above Hecke equivariant Gysin exact sequence in great detail. Let K be an imaginary quadratic field with ring of integers \mathbf{Z}_K . Let Γ be a torsion-free finite index subgroup of the Bianchi group $\mathbf{PSL}_2(\mathbf{Z}_K)$ acting on the hyperbolic 3-space \mathbf{H}_3 and its boundary $\partial\mathbf{H}_3$. In this case, the K -homology Gysin exact sequence splits into two exact sequences,

$$(\star) \quad 0 \rightarrow K^{i+1}(C_0(M)) \rightarrow K^i(C(\partial\mathbf{H}_3) \rtimes \Gamma) \rightarrow K^i(C_r^*(\Gamma)) \rightarrow 0,$$

with $i = 0, 1$, revealing that K -homology of $C(\partial\mathbf{H}_3) \rtimes \Gamma$ is ‘made of’ that of $C_r^*(\Gamma)$ and that of M . We then study the K -groups of these two parts in Section 4 and Section 5, relating them to the ordinary (co)homology of Γ and M . We do not consider K -theory groups anymore except in Section 4. To describe the K -homology isomorphism at the unbounded level, we give a construction of unbounded Fredholm modules from group cocycles. The fact that Kasparov’s γ -element is equal to the identity in $KK_0^\Gamma(\mathbf{C}, \mathbf{C})$ (see [38, 39]) is a vital ingredient in the construction.

Let $c : \Gamma \rightarrow \mathbf{Z}$ be 1-cocycle with kernel Γ_c and $(C_r^*(\Gamma), E, D_c)$ the associated unbounded $(C_r^*(\Gamma), C_r^*(\Gamma_c))$ Kasparov module (see Proposition 4.2). For $0 < s < 1$, the γ -element induces an unbounded Fredholm module

$$(C_r^*(\Gamma_c), L^2(\wedge^* \mathbf{H}_n), \mathcal{D}_s := \mathcal{D}_{HR} + \rho^s \hat{c}(d\rho)),$$

where \mathbb{D}_{HR} is the Hodge-deRham operator, $\rho(x) = d_{\mathbf{H}_n}(0, x)$ the distance function on the hyperbolic ball and \hat{c} the Clifford multiplication. We prove the following result in Theorem 4.10.

Theorem B. The operators D_c and \mathbb{D}_s assemble into an $(1 - s)$ unbounded Fredholm module

$$(C_r^*(\Gamma), E \otimes_{C_r^*(\Gamma_c)} L^2(\wedge^* \mathbf{H}_n), D_c \otimes \sigma + 1 \otimes_{\nabla_g} \mathbb{D}_s).$$

For $n = 3$, this induces Hecke equivariant isomorphisms

$$H^1(\Gamma, \mathbf{Z}) \simeq K^1(C_r^*(\Gamma)), \quad c \mapsto [D_{c,s}] \quad H_1(\Gamma, \mathbf{Z}) \simeq K_1(C_r^*(\Gamma)), \quad [\gamma] \mapsto u_\gamma.$$

Moreover under these isomorphisms, the homological pairing $H^* \times H_* \rightarrow \mathbf{Z}$ corresponds to the index pairing $K^* \times K_* \rightarrow \mathbf{Z}$.

The K -theory isomorphism comes from the results in [11, 51]. In Section 5 we exploit the equivalence between geometric and analytic K -homology ([7]) of the non-compact manifold M and in Theorem 5.5 establish the following:

Theorem C. There is an explicit Hecke equivariant isomorphism

$$H_2(\overline{M}, \partial \overline{M}, \mathbf{Z}) \simeq K^0(C_0(M)), \quad (N, \partial N) \mapsto (C_0(M), L^2(\overset{\circ}{N}, \mathcal{S}_{\overset{\circ}{N}}), \mathbb{D}_{\overset{\circ}{N}})$$

where \overline{M} denotes the Borel-Serre compactification of M and $(N, \partial N) \subset (\overline{M}, \partial \overline{M})$ is a properly embedded hypersurface. The interior $\overset{\circ}{N}$ inherits a complete Riemannian metric from M and hence the Dirac operators $\mathbb{D}_{\overset{\circ}{N}}$ are self-adjoint.

Note that $H^1(\Gamma, \mathbf{Z}) \simeq H_2(\overline{M}, \partial \overline{M}, \mathbf{Z})$. Our methods above also apply to the case of torsion-free finite index subgroups of $\mathbf{PSL}_2(\mathbf{Z})$ which we discuss in Section 6.

With these results in place, we proceed to describe the unbounded Fredholm modules that exhaust the K -homology of the purely infinite boundary crossed product $C(\partial \mathbf{H}_3) \rtimes \Gamma$. To this end we use the maps $K^1(C(\partial \mathbf{H}_3) \rtimes \Gamma) \rightarrow K^1(C_r^*(\Gamma))$, for which we construct an explicit section in Section 4, and $\partial : K^0(C_0(M)) \rightarrow K^1(C(\partial \mathbf{H}_3) \rtimes \Gamma)$ coming from (\star).

To compute the latter map at the unbounded level, we construct an unbounded representative for the extension class by means of a hypersingular integral operator built from the harmonic measures ν_x and associated metrics d_x on $\partial \mathbf{H}_3$ based at $x \in \mathbf{H}$. Let $T_1 \mathbf{H}_n = \mathbf{H}_n \times \partial \mathbf{H}_n$ be the unit tangent bundle of \mathbf{H}_n and $L^2(T_1 \mathbf{H}_n, \nu_x)$ the associated C^* -module completion. The integral operators

$$\Delta \Psi(x, \xi) = \int_{\partial \mathbf{H}_n} \frac{\Psi(x, \xi) - \Psi(x, \eta)}{d_x(\xi, \eta)^{n-1}} d\nu_x \eta, \quad p \Psi(x, \xi) = \int_{\partial \mathbf{H}_n} \Psi(x, \eta) d\nu_x \eta,$$

are G -invariant and the multiplication operator $\rho \Psi(x, \xi) = d_{\mathbf{H}_n}(0, x) \Psi(x, \xi)$ commutes with G boundedly in $L^2(T_1 \mathbf{H}, \nu_x)$. In Section 7 we prove:

Theorem D. The operators Δ, ρ and p assemble into a G -equivariant unbounded Kasparov module

$$(C(\partial \mathbf{H}), L^2(T_1 \mathbf{H}, \nu_x), -\Delta + (2p - 1)\rho),$$

representing the class of the G -equivariant extension

$$0 \rightarrow C_0(\mathbf{H}_n) \rightarrow C(\mathbf{H}_n \cup \partial \mathbf{H}_n) \rightarrow C(\partial \mathbf{H}_n) \rightarrow 0.$$

In Section 8 we then compute the unbounded Kasparov product of this operator with the self-adjoint Dirac operators on embedded hypersurfaces from Section 5. The main result here can be found in Theorem 8.8. The results in Sections 4 and 8 can be summarized to describe the structure of the K -homology of the purely infinite simple C^* -algebra $C(\partial\mathbf{H}_3) \rtimes \Gamma$ as in the following theorem.

Theorem E. The unbounded Fredholm modules in Theorem B extend to $C(\partial\mathbf{H}_n) \rtimes \Gamma$. A self-adjoint Dirac operator $\mathcal{D}_{\mathring{N}}$ associated to an embedded hypersurface and the integral operator $S := -\Delta + (2p - 1)\rho$ of Theorem D assemble into an unbounded Kasparov product

$$(C(\partial\mathbf{H}_n) \rtimes \Gamma, L^2(T_1\mathbf{H}_n) \otimes_{C_0(M)} L^2(\mathring{N}, \mathcal{S}_{\mathring{N}}), S \otimes \sigma + 1 \otimes_{\nabla} \mathcal{D}_{\mathring{N}}).$$

Consequently, for $n = 3$, there is an explicit Hecke equivariant isomorphism

$$H^1(\Gamma, \mathbf{Z}) \oplus H_2(\overline{M}, \partial\overline{M}) \simeq K^1(C(\partial\mathbf{H}_3) \rtimes \Gamma), \quad ([c], [(N, \partial N)]) \mapsto [D_{c,s}] + [S \otimes 1 + 1 \otimes_{\nabla} \mathcal{D}_{\mathring{N}}]$$

defined at the level of unbounded Fredholm modules.

The two pieces of $K^1(C(\partial\mathbf{H}_3) \rtimes \Gamma)$ give rise to very different unbounded Fredholm modules, which by virtue of the Gysin sequence pair with K -theory in distinct ways.

Questions.

- (1) Given an arithmetic group Γ , we construct Hecke operators on the K -groups of arithmetic C^* -algebras associated to Γ . In the case of Bianchi groups, we show that our Hecke operators correspond to the classical Hecke operators on the cohomology groups of Γ . While this convinces us that our construction is natural and correct, for general Γ such a comparison is still to be made. A natural question to ask here is, does the Chern character homomorphism

$$K^i(B\Gamma) \rightarrow \bigoplus_{n \geq 0} H^{2n+i}(B\Gamma, \mathbf{Q}),$$

where $i = 0, 1$ commute with the Hecke operators?

- (2) Torsion in the homology of arithmetic groups has gained a lot of interest in recent years. What can we say about the torsion in the K -theory of arithmetic C^* -algebras? We observe in Section 1 that for Bianchi groups, H_1 and K_1 hold the same torsion. However this is a coincidence of low dimensionality and in general the torsion on the two sides will not agree. Note that it is natural expect that p -torsion Hecke eigenclasses in the K -homology of arithmetic C^* -algebras have associated mod p Galois representations.
- (3) Can the K -homology of the arithmetic C^* -algebras as Hecke modules be accounted for by automorphic forms as is the case for cohomology? If so, what are these ‘ K -theoretic’ automorphic forms? Are they the same as cohomological ones? In the case of $\mathbf{PSL}_2(\mathbf{Z})$ and Bianchi groups, we show that they are the same. In these cases, can we directly associate a K -homology class associated to a Bianchi (or classical) modular form?
- (4) What can we say about the summability properties of the (un)bounded Fredholm modules that we construct? Do their spectral zeta functions relate to the arithmetic of Bianchi modular forms?

Acknowledgements. We gratefully acknowledge that we benefited from conversations with Michael Atiyah, Alain Connes, Gunther Cornelissen, Robin Deeley, Nathan Dunfield, Magnus Goffeng, Paul Gunnells, Joel Hass, Matilde Marcolli, Sergey Neshveyev, Ryszard Nest, Bogdan Nica, Elmar Schrohe, René Schulz, Richard Sharp, Aidan Sims and Alain Valette. We thank the Leibniz Universität Hannover and the MSRC of the University of Sheffield for their financial support in facilitating this collaboration. We are also grateful to the University of Warwick and the Nesin Mathematics Village for providing excellent working environments. The first author was partially supported by EPSRC grant EP/J006580/2, Simons Foundation grant 346300 and the Polish Government MNiSW 2015-2019 matching fund. The second author was a Marie Curie Intra-European Fellow during the preparation of a significant part of this work.

1. Prelude: Kasparov's spectral sequence

Let Γ be a torsion-free cofinite discrete subgroup of $\mathrm{PSL}_2(\mathbf{C})$ acting on the hyperbolic 3-space \mathbf{H}_3 and its boundary $\partial\mathbf{H}_3$. The limit set of Γ is all of $\partial\mathbf{H}_3$ on which it acts with dense orbits. We can identify $\partial\mathbf{H}_3$ with $S^2 \simeq \mathbb{P}^1(\mathbf{C})$ and the action of Γ with the usual Möbius action. Let M denote the hyperbolic 3-manifold \mathbf{H}_3/Γ . In this section, we employ a KK -theory spectral sequence and get a description of K -groups of $C(\partial\mathbf{H}_3) \rtimes \Gamma$, $C_r^*(\Gamma)$ and $C_0(M)$ in terms of the cohomology of Γ .

The abstract isomorphisms that will come out of the spectral sequence will motivate the main task we accomplish in the present paper: Can we equip the respective K -groups with a Hecke module structure and find *explicit Hecke equivariant* isomorphisms from (co)homology to K -groups?

We let D be a C^* -algebra with a Γ -action and $D \rtimes_r \Gamma$ be the reduced crossed product algebra of Γ and D . Let $\gamma \in KK^\Gamma(\mathbf{C}, \mathbf{C})$ denote the Kasparov idempotent. A spectral sequence of Kasparov (see [39, Section 6.10.], see also [67]) calculates the γ -parts of the K -groups of $D \rtimes_r \Gamma$ out of those of D . For discrete subgroups of $\mathrm{Isom}(\mathbf{H}_n)$, it holds that $\gamma = 1$, a fact that will be of importance in several places in the present paper (see for instance [73, Chapter 9] and [38, 39]). For our groups, the γ -part of a K -group of $D \rtimes_r \Gamma$ is itself.

Theorem 1.1. *There is a cohomological spectral sequence (E_r, d_r) with differentials $d_r^{p,q} : E_r^{p,q} \rightarrow E_r^{p+r,q-r+1}$ and the term $E_2^{p,q} = H^p(\Gamma, K^q(D))$ converging to the K -homology groups of $D \rtimes_r \Gamma$. There is an analogous homological spectral sequence converging to the K -theory groups of $D \rtimes_r \Gamma$.*

By setting D equal to $\mathbf{C} = C(\mathbf{pt})$, $C_0(\mathbf{H}_3)$ and $C(\partial\mathbf{H}_3)$, we shall use the above spectral sequence to obtain information on the K -groups of $C(\partial\mathbf{H}_3) \rtimes_r \Gamma$, $C_r^*(\Gamma)$ and $C_0(M)$ respectively. We first set $D = C(\partial\mathbf{H}_3)$, and note that the action of Γ on $\partial\mathbf{H}_3$ is amenable so the full and reduced crossed products coincide (see [46, Lemma 3.8]). The following well known lemma computes the K -homology groups of D as Γ -modules in this case.

Lemma 1.2. *We have*

$$K^0(C(\partial\mathbf{H}_3)) \simeq \mathbf{Z}^2, \quad K^1(C(\partial\mathbf{H}_3)) = \{0\},$$

where the action of Γ on $K^0(C(\partial\mathbf{H}_3))$ is trivial.

Proof. The K -homology of the two-sphere $S^2 \simeq \partial\mathbf{H}_3$ is well-known. The triviality of the action of Γ on $K^0(C(\partial\mathbf{H}_3))$ follows from the facts that it is the restriction of the action of $\mathrm{PSL}_2(\mathbf{C})$, which is a connected group, and K -homology is homotopy invariant. \square

Proposition 1.3. *There is a short exact sequence*

$$(1.1) \quad 0 \rightarrow H^0(\Gamma, \mathbf{Z}^2) \rightarrow K^0(C(\partial\mathbf{H}_3) \rtimes \Gamma) \rightarrow H^2(\Gamma, \mathbf{Z}^2) \rightarrow 0$$

and an isomorphism

$$(1.2) \quad H^1(\Gamma, \mathbf{Z}^2) \simeq K^1(C(\partial\mathbf{H}_3) \rtimes \Gamma)$$

where the action of Γ on \mathbf{Z}^2 is trivial.

Proof. We apply Theorem 1.1 with $D = C(\partial\mathbf{H}_3)$ and use Lemma 1.2. The cohomological dimension of Γ is two. As Γ is torsion-free, all its integral cohomology above degree two vanishes and we see that the E_2 page of the spectral sequence looks like this:

\vdots	\vdots	\vdots	\vdots	\vdots	\vdots	\dots
0	0	0	0	0	0	\dots
$H^0(\Gamma, \mathbf{Z}^2)$	$H^1(\Gamma, \mathbf{Z}^2)$	$H^2(\Gamma, \mathbf{Z}^2)$	0	0	0	\dots
0	0	0	0	0	0	\dots
$H^0(\Gamma, \mathbf{Z}^2)$	$H^1(\Gamma, \mathbf{Z}^2)$	$H^2(\Gamma, \mathbf{Z}^2)$	0	0	0	\dots

The E_2 page of the spectral sequence.

□

Note that since the action of Γ is trivial on \mathbf{Z}^2 , we have $H^*(\Gamma, \mathbf{Z}^2) \simeq H^*(\Gamma, \mathbf{Z}) \otimes \mathbf{Z}^2$. In particular, $H^1(\Gamma, \mathbf{Z}^2) \cong H^1(\Gamma, \mathbf{Z})^{\oplus 2}$ and thus $K^1(C(\partial\mathbf{H}_3) \rtimes \Gamma)$ holds two copies of $H^1(\Gamma, \mathbf{Z})$. Moreover, $H^0(\Gamma, \mathbf{Z}^2) \simeq \mathbf{Z}^2$. As $H^2(\Gamma, \mathbf{Z})$ typically has a lot of torsion (see [69, 9]), the sequence 1.1 does not split.

Next, we apply Theorem 1.1 to the case $D = \mathbf{C} = C(\mathbf{pt})$. Note that $C(\mathbf{pt}) \rtimes \Gamma \simeq C_r^*(\Gamma)$. Since $K^0(\mathbf{C}) \simeq \mathbf{Z}$, $K^1(\mathbf{C}) = \{0\}$, and the Γ -action is trivial, we find

Proposition 1.4. *There is a short exact sequence*

$$(1.3) \quad 0 \rightarrow H^0(\Gamma, \mathbf{Z}) \rightarrow K^0(C_r^*(\Gamma)) \rightarrow H^2(\Gamma, \mathbf{Z}) \rightarrow 0$$

and an isomorphism

$$(1.4) \quad H^1(\Gamma, \mathbf{Z}) \simeq K^1(C_r^*(\Gamma)),$$

where the action of Γ on \mathbf{Z} is trivial.

Lastly, we apply Theorem 1.1 to the case $D = C_0(\mathbf{H}_3)$. Note that $C_0(\mathbf{H}_3) \rtimes \Gamma$ is Morita equivalent to $C_0(M)$ as the action of Γ on \mathbf{H}_3 is free and properly discontinuous. In particular, they have the same K -groups. It is well known that $K^0(C_0(\mathbf{H}_3)) \simeq \{0\}$ and that $K^1(C_0(\mathbf{H}_3)) = \mathbf{Z}$ with trivial Γ -action.

Proposition 1.5. *There is a short exact sequence*

$$(1.5) \quad 0 \rightarrow H^0(\Gamma, \mathbf{Z}) \rightarrow K^1(C_0(M)) \rightarrow H^2(\Gamma, \mathbf{Z}) \rightarrow 0$$

and an isomorphism

$$(1.6) \quad H^1(\Gamma, \mathbf{Z}) \simeq K^0(C_0(M))$$

where the action of Γ on \mathbf{Z} is trivial.

We summarize the results of the homological spectral sequence and omit the details as they are the same as the above.

Proposition 1.6. *There are isomorphisms*

$$(1.7) \quad H_1(\Gamma, \mathbf{Z}^2) \simeq K_1(C(\partial\mathbf{H}_3) \rtimes \Gamma),$$

and

$$(1.8) \quad H_1(\Gamma, \mathbf{Z}) \simeq K_1(C_r^*(\Gamma)),$$

and

$$(1.9) \quad H_1(\Gamma, \mathbf{Z}) \simeq K_0(C_0(M)),$$

and short exact sequences

$$(1.10) \quad 0 \rightarrow H_0(\Gamma, \mathbf{Z}^2) \rightarrow K_0(C(\partial\mathbf{H}_3) \rtimes \Gamma) \rightarrow H_2(\Gamma, \mathbf{Z}^2) \rightarrow 0,$$

$$(1.11) \quad 0 \rightarrow H_0(\Gamma, \mathbf{Z}) \rightarrow K_0(C_r^*(\Gamma)) \rightarrow H_2(\Gamma, \mathbf{Z}) \rightarrow 0,$$

$$(1.12) \quad 0 \rightarrow H_0(\Gamma, \mathbf{Z}) \rightarrow K_1(C_0(M)) \rightarrow H_2(\Gamma, \mathbf{Z}) \rightarrow 0.$$

where the actions of Γ on \mathbf{Z}^2 and on \mathbf{Z} are trivial.

With these abstract isomorphisms in hand, we set ourselves two tasks. The first is to equip the K -groups of the above arithmetic C^* -algebras with natural Hecke module structures. Secondly, we would like to compare the K -groups and (co)homology groups as Hecke modules. For this purpose, the abstract isomorphism coming from Kasparov's spectral sequence above cannot help, so we need to construct explicit isomorphisms between the respective groups appearing in this section. In the rest of the paper, we accomplish both of our tasks. However it should be noted that while our treatment of the first task is general, our treatment of the second task is very specific to the case of Bianchi groups (see Question 2 in the Introduction).

2. Hecke operators and KK -theory

The various (co)homology groups associated with an arithmetic group Γ come equipped with so called *Hecke operators*. These arise from elements in the commensurator $C_G(\Gamma)$ of Γ in its ambient real Lie group G :

$$C_G(\Gamma) := \{g \in G : \Gamma \cap g\Gamma g^{-1} \text{ has finite index in both } \Gamma \text{ and } g\Gamma g^{-1}\}.$$

We start by quickly recalling the definition of Hecke operators on the (co)homology groups that we deal with in the paper. Afterwards, for each element in $C_G(\Gamma)$, we construct elements in KK -rings $KK_0(A, A)$ of the arithmetic C^* -algebras A associated to Γ . The elements that we

construct will give rise to endomorphisms which play the rôle of Hecke operators on K -groups of A .

2.1. Homological definitions. Let $\Gamma \subset \mathbf{PSL}_2(\mathbf{C}) =: G$ be a torsion-free finite-index subgroup of a Bianchi group $\mathbf{PSL}_2(\mathbf{Z}_K)$, acting on \mathbf{H}_3 freely and proper discontinuously. In this case, we have $C_G(\Gamma) = \mathbf{PGL}_2(K) \subset \mathbf{PGL}_2(\mathbf{C}) \cong G$. For our purposes, the main distinction to be made is that between algebraically defined Hecke operators on $H^*(\Gamma, \mathbf{Z})$ and topologically defined Hecke operators on $H_*(\overline{M}, \partial\overline{M}, \mathbf{Z})$ where M is the associated hyperbolic 3-manifold \mathbf{H}_3/Γ and \overline{M} is its Borel-Serre bordification.

2.1.1. On Group Homology. For a subgroup $\Delta \subset \Gamma$ of finite index d , any choice of coset representatives

$$\gamma_i \in \Gamma, \quad \Gamma = \bigsqcup_{i=1}^d \gamma_i \Delta,$$

gives a map $s : \Gamma \rightarrow \Delta^d$ determined by $\gamma\gamma_i = \gamma_{\gamma(i)}s_i(\gamma)$, where $s_i(\gamma) \in \Delta$ and $\gamma(i)$ is a permutation of $1, \dots, d$. This determines the *transfer* or *corestriction map*

$$\text{cores} : H^1(\Delta, \mathbf{Z}) \rightarrow H^1(\Gamma, \mathbf{Z}), \quad \text{cores } c(\gamma) = \sum_{i=1}^d c(s_i(\gamma)),$$

which is independent of the choice of coset representatives γ_i . For $g \in C_G(\Gamma)$, write $\Gamma_g := \Gamma \cap g\Gamma g^{-1}$ and the Hecke operator on group cohomology is given by

$$(2.1) \quad T_g : H^1(\Gamma, \mathbf{Z}) \xrightarrow{\text{res}} H^1(\Gamma_g, \mathbf{Z}) \xrightarrow{\text{Ad}_g} H^1(\Gamma_{g^{-1}}, \mathbf{Z}) \xrightarrow{\text{cores}} H^1(\Gamma, \mathbf{Z}).$$

Operators $T_g : H_1(\Gamma, \mathbf{Z}) \rightarrow H_1(\Gamma, \mathbf{Z})$ are defined analogously. To compute the operator T_g , one uses the disjoint union decomposition of the double coset

$$(2.2) \quad \Gamma g^{-1} \Gamma = \bigsqcup_{i=1}^d g_i \Gamma, \quad g_i = \delta_i g^{-1} \in G, \quad \delta_i \in \Gamma.$$

The elements δ_i form a complete set of coset representatives for $\Gamma/\Gamma_{g^{-1}}$. The group Γ acts on the double coset $\Gamma g^{-1} \Gamma$, and thus permutes the cosets $g_i \Gamma$. As above there are indices $\gamma(i)$ and group elements $t_i(\gamma) \in \Gamma$ such that $\gamma g_i = g_{\gamma(i)} t_i(\gamma)$, determining a map $\Gamma \rightarrow \Gamma_{g^{-1}}^d$. The Hecke operators $T_g : H^1(\Gamma, \mathbf{Z}) \rightarrow H^1(\Gamma, \mathbf{Z})$ and $T_g : H_1(\Gamma, \mathbf{Z}) \rightarrow H_1(\Gamma, \mathbf{Z})$ are then given explicitly by

$$(2.3) \quad (T_g c)(\gamma) := \sum_{i=1}^d c(t_i(\gamma)), \quad T_g([\gamma]) = \sum_i [t_i(\gamma)]$$

which is independent of the choice of coset representatives δ_i .

2.1.2. On Simplicial Homology. We start with the manifold $M_g := \mathbf{H}_3/\Gamma_g$ and the associated finite covering $\pi_g : M_g \rightarrow M$. This finite covering induces a corestriction map $\pi_g^* : H_*(M) \rightarrow H_*(M_g)$ by mapping a simplex to the sum of its inverse images. Similarly there is a covering $\pi_{g^{-1}} : M_{g^{-1}} \rightarrow M$, and the isometry $g : \mathbf{H}_3 \rightarrow \mathbf{H}_3$ induces a homeomorphism $g_* : M_g \rightarrow M_{g^{-1}}$ because $g^{-1}\Gamma_g g = \Gamma_{g^{-1}}$. Thus we obtain a second covering $\tau_g := \pi_{g^{-1}} \circ g_* : M_g \rightarrow M$.

For $g \in C_G(\Gamma)$, we define Hecke operators, both denoted T_g , on homology and on cohomology as the group homomorphisms

$$T_g := \tau_{g*} \circ \pi_g^* : H_*(M, \mathbf{Z}) \rightarrow H_*(M, \mathbf{Z}),$$

$$T_g := \tau_g^* \circ \pi_{g*} : H^*(M, \mathbf{Z}) \rightarrow H^*(M, \mathbf{Z}).$$

We shall need Hecke operators also on the homology of the Borel-Serre compactifications. In our low-dimensional cases, these compactifications can be described concretely as follows (see [13] and also [12, III.5.15], [8, §2.8]). We first construct a partial compactification $\widehat{\mathbf{H}}_3$ of \mathbf{H}_3 by adding a copy of the complex plane \mathbf{C} to every boundary point in $\mathbb{P}^1(K) \subset \mathbb{P}^1(\mathbf{C}) = \partial\mathbf{H}_3$, more precisely

$$\widehat{\mathbf{H}}_3 = \mathbf{H}_3 \bigsqcup_{z \in \mathbb{P}^1(K)} \mathbb{P}^1(\mathbf{C}) \setminus \{z\}.$$

The copy $\mathbb{P}^1(\mathbf{C}) \setminus \{z\} = \mathbf{C}$ is the parameter space of all geodesics in \mathbf{H}_3 converging to the boundary point $z \in \mathbb{P}^1(K)$. The action of $\mathbf{PGL}_2(K)$, but not of G , on \mathbf{H}_3 extends to an action on $\widehat{\mathbf{H}}_3$ by sending $\omega \in \mathbb{P}^1(\mathbf{C}) \setminus \{z\}$ to $\omega\gamma \in \mathbb{P}^1(\mathbf{C}) \setminus \{z\gamma\}$. One can topologize $\widehat{\mathbf{H}}_3$ in such a way that the action of $\mathbf{PGL}_2(\mathbf{K})$ is continuous. The action of Γ on $\widehat{\mathbf{H}}_3$, unlike its action on the geodesic completion $\overline{\mathbf{H}}_3$, is free and proper. The quotient $\widehat{\mathbf{H}}_3/\Gamma$ can be shown to be a compact 3-manifold with boundary which we call the Borel-Serre compactification of M and denote by \overline{M} . The connected components of its boundary are 2-tori, attached at ‘infinity’ to each cusp of M . Note that M is the interior of \overline{M} and thus they are homotopy equivalent.

Just as before, we obtain finite coverings $\overline{\pi}_g, \overline{\tau}_g : \overline{M}_g \rightarrow \overline{M}$, extending $\pi_g, \tau_g : M_g \rightarrow M$, and construct the Hecke operator

$$T_g := (\pi_{g^{-1}})_* \circ g_* \circ \pi_g^* : H_*(\overline{M}, \mathbf{Z}) \rightarrow H_*(\overline{M}, \mathbf{Z}).$$

As $\overline{\pi}_g, \overline{\tau}_g$ restrict to finite coverings on the boundaries, we also obtain Hecke operators on the relative homology groups

$$T_g : H_*(\overline{M}, \partial\overline{M}, \mathbf{Z}) \rightarrow H_*(\overline{M}, \partial\overline{M}, \mathbf{Z}).$$

These Hecke operators are compatible with the Lefschetz duality isomorphism

$$H_*(\overline{M}, \partial\overline{M}, \mathbf{Z}) \cong H^{n-*}(\overline{M}, \mathbf{Z}),$$

see [3, Lemma 1.4.3], and the isomorphisms

$$H^*(\overline{M}, \mathbf{Z}) \cong H^*(M, \mathbf{Z}) \cong H^*(\Gamma, \mathbf{Z}),$$

see, for example, [45, Section 6].

2.2. Hecke operators in KK -theory. Let X be a locally compact Hausdorff space and assume that G acts on X and that $\Gamma \subset G$ acts freely and properly on X . Suggestively, denote by $M := X/\Gamma$ the quotient space which is locally compact and Hausdorff.

The finite coverings $M \xleftarrow{\tau_g} M_g \xrightarrow{\pi_g} M$ form a *correspondence* in the sense of [20] and define a class $[T_g^M] \in KK_0(C_0(M), C_0(M))$. The conditional expectation and right module structure

$$\rho_g : C_0(M_g) \rightarrow C_0(M), \quad \rho(\psi)(m) = \sum_{x \in \pi_g^{-1}(m)} \psi(x), \quad \psi \cdot f(x) := \psi(x)f(\pi_g(x))$$

give a right $C_0(M)$ -module denoted by T_g^M . Because the map $\tau_g : M_g \rightarrow M$ is proper, there is a left action by compact operators

$$C_0(M) \rightarrow \mathbb{K}(T_g^M), \quad f \cdot \psi(x) = f(\tau_g(x))\psi(x).$$

The class $[T_g^M] \in KK_0(C_0(M), C_0(M))$ coincides with the class of this bimodule.

Definition 2.1. Let $M = X/\Gamma$ as above. For any separable C^* -algebra C , the Hecke operators $T_g : KK_*(C_0(M), C) \rightarrow KK_*(C_0(M), C)$, $T_g : KK_*(C, C_0(M)) \rightarrow KK_*(C, C_0(M))$, are defined to be the Kasparov product with the class $[T_g^M] \in KK_0(C_0(M), C_0(M))$.

For the moment, we denote by B an arbitrary $C_G(\Gamma)$ - C^* -algebra and by $g : b \mapsto g(b)$ the $C_G(\Gamma)$ -action. Let $C_c(\Gamma, B)$ denote the compactly supported B -valued functions on Γ . The Γ - C^* -module

$$\ell^2(\Gamma, B) := \left\{ \psi : \Gamma \rightarrow B : \sum_{\gamma \in \Gamma} \psi(\gamma)^* \psi(\gamma) < \infty \right\},$$

of ℓ^2 functions on Γ with values in B is constructed as a completion of $C_c(\Gamma, B)$. The convolution product and involution given by (see [40])

$$(2.4) \quad f * \psi(\gamma) = \sum_{\delta \in \Gamma} f(\delta) \delta(\psi(\delta^{-1}\gamma)), \quad f^*(\gamma) := \gamma f(\gamma^{-1})^*$$

make $C_c(\Gamma, B)$ into a $*$ -algebra and define a $*$ -representation $C_c(\Gamma, B) \rightarrow \text{End}_B^*(\ell^2(\Gamma, B))$. The reduced crossed product $B \rtimes_r \Gamma$ is defined as the closure of $C_c(\Gamma, B)$ in this representation.

For a subgroup $\Delta \subset \Gamma$, restriction of functions $C_c(\Gamma, B) \rightarrow C_c(\Delta, B) \subset C_c(\Gamma, B)$ defines a projection $p_\Delta \in \text{End}_B^*(\ell^2(\Gamma, B))$. This gives a contractive conditional expectation

$$\rho_\Delta : B \rtimes_r \Gamma \rightarrow B \rtimes_r \Delta, \quad a \mapsto p_\Delta a p_\Delta,$$

extending the restriction map $C_c(\Gamma, B) \rightarrow C_c(\Delta, B)$. Thus, for $g \in C_G(\Gamma)$ we obtain the expectation $\rho_{g^{-1}} : B \rtimes_r \Gamma \rightarrow B \rtimes_r \Gamma_{g^{-1}}$ and a $(B \rtimes_r \Gamma, B \rtimes_r \Gamma_{g^{-1}})$ bimodule $(B \rtimes_r \Gamma)_{\rho_{g^{-1}}}$. Using the $*$ -homomorphism

$$B \rtimes_r \Gamma_{g^{-1}} \xrightarrow{\text{Ad}_g} B \rtimes_r \Gamma_g \hookrightarrow B \rtimes_r \Gamma, \quad \text{Ad}_g(f)(\gamma) := g f(g^{-1}\gamma g),$$

we form the interior C^* -module tensor product

$$T_g^\Gamma := (B \rtimes_r \Gamma)_{\rho_{g^{-1}}} \otimes_{\text{Ad}_g} B \rtimes_r \Gamma,$$

which is a $B \rtimes_r \Gamma$ -bimodule.

Definition 2.2. Let B be a separable $C_G(\Gamma)$ - C^* -algebra and C a separable C^* -algebra. The Hecke operators

$$T_g : KK_*(B \rtimes_r \Gamma, C) \rightarrow KK_*(B \rtimes_r \Gamma, C), \quad T_g : KK_*(C, B \rtimes_r \Gamma) \rightarrow KK_*(C, B \rtimes_r \Gamma).$$

are defined to be the Kasparov product with the class $[T_g^\Gamma] \in KK_0(B \rtimes_r \Gamma, B \rtimes_r \Gamma)$.

Let A be any of the C^* -algebras and T_g any of the KK -theoretic Hecke operators discussed above. If $\langle x, y \rangle$ denotes the index pairing of elements $x \in K_*(A)$ and $y \in K^*(A)$, associativity of the Kasparov product gives $\langle T_g x, y \rangle = \langle x, T_g y \rangle$. That is, the Hecke action is self-adjoint with respect to the index pairing between K -theory and K -homology.

2.3. Explicit formulae for the reduced crossed product. To describe the $B \rtimes_r \Gamma$ -bimodule T_g^Γ , let δ_i be as in (2.2) and $\chi_i \in C_c(\Gamma, M(B))$ be the function that is 1 at δ_i and 0 elsewhere. It is straightforward to check that $\sum_{i=1}^d \chi_i * \rho(\chi_i^* * f) = f$, and $\rho(\chi_i^* * \chi_j) = \delta_{ij}$. This implements a unitary isomorphism of right modules

$$(2.5) \quad u : T_g^\Gamma = (B \rtimes_r \Gamma)_{\rho_{g^{-1}}} \otimes_{\text{Ad}_g} B \rtimes_r \Gamma \rightarrow (B \rtimes_r \Gamma)^d, \quad f \otimes k \mapsto (\rho(\chi_i^* * f) * k),$$

where $d = [\Gamma : \Gamma_{g^{-1}}]$. To describe the left $B \rtimes_r \Gamma$ action on $T_g^\Gamma \simeq (B \rtimes_r \Gamma)^d$ we consider the dense submodule $C_c(\Gamma, B^d)$, the elements of which we view as columns $\Psi := (\Psi_i)_{i=1}^d$ of maps $\Psi_i : \Gamma \rightarrow B$. First we collect some useful facts and relations for the elements $t_i(\gamma)$.

Lemma 2.3. *We have the relations:*

- 1.) $t_i(\gamma) = g\delta_{\gamma(i)}^{-1}\gamma\delta_i g^{-1} = g_{\gamma(i)}^{-1}\gamma g_i$;
- 2.) $t_i(\gamma_1\gamma_2) = t_{\gamma_2(i)}(\gamma_1)t_i(\gamma_2)$;
- 3.) $t_i(\gamma^{-1}) = t_{\gamma^{-1}(i)}(\gamma)^{-1}$.

Proof. All relations are checked by direct computation using the definitions in section 2.1 \square

For $\gamma \in \Gamma$ denote by $u_\gamma \in C_c(\Gamma, M(B))$ the function which is 1 at γ and 0 elsewhere. We identify $B \subset C_c(\Gamma, B)$ with the function that takes the value b at $e \in \Gamma$ and 0 elsewhere.

Proposition 2.4. *The left $B \rtimes_r \Gamma$ module structure on $T_g^\Gamma \simeq (B \rtimes_r \Gamma)^d$ is given by*

$$(2.6) \quad (t_g(f)\Psi)_i(\delta) = \sum_{\gamma} g_i^{-1} f(\gamma) t_i(\gamma^{-1})^{-1} \Psi_{\gamma^{-1}(i)}(t_i(\gamma^{-1})\delta).$$

Equivalently, we have the covariant representation

$$(2.7) \quad (t_g(b) \cdot \Psi)_i(\delta) := g_i^{-1}(b)\Psi_i(\delta), \quad (t_g(u_\gamma)\Psi)_i(\delta) := t_i(\gamma^{-1})^{-1}(\Psi_{\gamma^{-1}(i)}(t_i(\gamma^{-1})\delta)).$$

Moreover, for $\gamma \in \Gamma$ we have a factorisation $t_g(u_\gamma) = \tau(\gamma)\text{diag}(u_{t_k(\gamma)})$, where $\tau(\gamma) \in M_d(\mathbf{C})$ is a permutation matrix.

Proof. By right $B \rtimes_r \Gamma$ linearity, it suffices to prove (2.6) for elements Ψ with $\text{supp } \Psi \subset \Gamma_g$. Using u as in (2.5) and the relations in Lemma 2.3 one computes, for $h \in C_c(\Gamma, B)$ and $\delta \in \Gamma_g$:

$$\begin{aligned} (2.8) \quad (t_g(h)\Psi)_i(\delta) &= (u(h * u^*\Psi))_i(\delta) = \text{Ad}_g \rho(\chi_i^* * h * u^*\Psi)(\delta) \\ &= g\rho(\chi_i^* * h * u^*\Psi)(g^{-1}\delta g) = g\delta_i^{-1}(h * u^*\Psi)(\delta_i g^{-1}\delta g) \\ &= \sum_{j:\gamma} g\delta_i^{-1}(h(\gamma)\gamma\delta_j g^{-1}\Psi_j(g\delta_j^{-1}\gamma^{-1}\delta_i g^{-1}\delta)) \\ (2.9) \quad &= \sum_{\gamma} g\delta_i^{-1}h(\gamma)g\delta_i^{-1}\gamma\delta_{\gamma^{-1}(i)}g^{-1}\Psi_{\gamma^{-1}(i)}(g\delta_{\gamma^{-1}(i)}^{-1}\gamma^{-1}\delta_i g^{-1}\delta) \\ &= \sum_{\gamma} g_i^{-1}h(\gamma)t_i(\gamma^{-1})^{-1}\Psi_{\gamma^{-1}(i)}(t_i(\gamma^{-1})\delta). \end{aligned}$$

The step from (2.8) to (2.9) follows since $g\delta_j^{-1}\gamma^{-1}\delta_i g^{-1}\delta \in \Gamma_g \Leftrightarrow j = \gamma^{-1}(i)$. Thus we have established (2.6) and (2.7) follows. Let $\tau(\gamma) \in M_d(\mathbf{C})$ be the permutation matrix corresponding

to $(\tau(\gamma)\Psi)_i = \Psi_{\gamma^{-1}(i)}$. To prove the last statement we compute

$$\begin{aligned} (\tau(\gamma)\text{diag}(u_{t_k(\gamma)})\Psi)_i(\delta) &= (\text{diag}(u_{t_k(\gamma)})\Psi)_{\gamma^{-1}(i)}(\delta) = t_{\gamma^{-1}(i)}(\gamma)^{-1}(\Psi_{\gamma^{-1}(i)}(t_{\gamma^{-1}(i)}(\gamma)\delta)) \\ &= t_i(\gamma^{-1})^{-1}(\Psi_{\gamma^{-1}(i)}(t_i(\gamma^{-1})\delta)) = (t_g(u_\gamma)\Psi)_i(\delta), \end{aligned}$$

as required. \square

3. Gysin sequence and Hecke operators

The paper [23] is an extensive study of the Gysin sequence in K -theory arising from a group action on a space X and the associated boundary action on ∂X , e.g. the Furstenberg or Gromov boundary. In [24, Section 10], the K -homological version is described for hyperbolic groups Γ with cocompact classifying space for proper actions $\mathcal{E}\Gamma$. We will describe the Gysin sequence in the setting of hyperbolic $n + 1$ -space \mathbf{H} , the geodesic compactification $\overline{\mathbf{H}}$ and its boundary sphere $\partial\mathbf{H} = S^n$.

3.1. The K -homology exact sequence. Let $G = \text{Isom}(\mathbf{H})$ and $\overline{\mathbf{H}} := \mathbf{H} \cup \partial\mathbf{H}$ the geodesic compactification of \mathbf{H} on which G acts as well. We consider the G -equivariant extension

$$(3.1) \quad 0 \rightarrow C_0(\mathbf{H}) \rightarrow C(\overline{\mathbf{H}}) \rightarrow C(\partial\mathbf{H}) \rightarrow 0,$$

defining a class in $KK_1^G(C(\partial\mathbf{H}), C_0(\mathbf{H}))$. Thus, for any subgroup $\Gamma \subset G$ we obtain a class in $KK_1^\Gamma(C(\partial\mathbf{H}), C_0(\mathbf{H}))$ through restriction, and a long exact sequence in equivariant K -homology:

$$(3.2) \quad \cdots \rightarrow K_1^i(C_0(\mathbf{H})) \rightarrow K_1^{i+1}(C(\partial\mathbf{H})) \rightarrow K_1^{i+1}(C(\overline{\mathbf{H}})) \rightarrow \cdots$$

Lemma 3.1. *Suppose $\Gamma \subset G$ is discrete and torsion-free. Then the inclusion $i : \mathbf{C} \rightarrow C(\overline{\mathbf{H}})$ as constant functions induces an isomorphism $i^* : K_1^i(C(\overline{\mathbf{H}})) \rightarrow K_1^i(\mathbf{C})$.*

Proof. Because Γ is torsion-free and $\overline{\mathbf{H}}$ is contractible, H -equivariant contractibility [55] of $\overline{\mathbf{H}}$ for finite subgroups $H \subset \Gamma$ follows trivially. Then the argument of [24, Lemma 10.6] applies. \square

The extension (3.1) induces an extension of crossed products

$$(3.3) \quad 0 \rightarrow C_0(\mathbf{H}) \rtimes \Gamma \rightarrow C(\overline{\mathbf{H}}) \rtimes \Gamma \rightarrow C(\partial\mathbf{H}) \rtimes \Gamma \rightarrow 0,$$

as the Γ -action on either of the algebras in (3.1) is amenable, and the full and reduced crossed products coincide. Let $L^2(\wedge^*\mathbf{H})$ be the Hilbert space of L^2 -sections of the exterior algebra bundle of \mathbf{H} , and \not{D}_{HR} the Hodge-DeRham operator. The triple $[(\mathbf{C}, L^2(\wedge^*\mathbf{H}), \not{D}_{HR})]$ defines an element in $KK_0^G(\mathbf{C}, \mathbf{C})$ and thus in $KK_0^\Gamma(\mathbf{C}, \mathbf{C})$ for any subgroup $\Gamma \subset G$. We will refer to each of these elements as the *Euler class* (cf. [23, 24]). We obtain the following Proposition.

Proposition 3.2. *For a discrete torsion-free subgroup $\Gamma \subset G$ there is an exact hexagon*

$$(3.4) \quad \begin{array}{ccccc} K^1(C_0(M)) & \xrightarrow{\partial} & K^0(C(\partial\mathbf{H}) \rtimes \Gamma) & \xrightarrow{i^*} & K^0(C_r^*(\Gamma)) \\ \text{Eul}_1 \uparrow & & & & \downarrow \text{Eul}_0 \\ K^1(C_r^*(\Gamma)) & \xleftarrow{i^*} & K^1(C(\partial\mathbf{H}) \rtimes \Gamma) & \xleftarrow{\partial} & K^0(C_0(M)), \end{array}$$

where i^* is induced from the inclusion $i : \mathbf{C} \rightarrow C(\overline{\mathbf{H}})$ and the maps Eul_* are induced from the Kasparov product with the Euler class $[(\mathbf{C}, L^2(\wedge^*\mathbf{H}), \not{D}_{HR})]$.

Proof. This follows from the arguments in [23, 24]. Since $\gamma = 1 \in KK_0^\Gamma(\mathbf{C}, \mathbf{C})$, there is an isomorphism $K_\Gamma^i(\mathbf{C}) \cong K^i(C_r^*(\Gamma))$, by descent in the first variable. Thus, Lemma 3.1 gives isomorphisms

$$K^*(C(\overline{\mathbf{H}}) \rtimes \Gamma) \xrightarrow{\sim} K_\Gamma^*(\mathbf{C}) \xrightarrow{\sim} K^*(C_r^*(\Gamma)).$$

Because the action of Γ on \mathbf{H} is free and proper, we have isomorphisms

$$K_\Gamma^*(C_0(\mathbf{H})) \xrightarrow{\sim} K^*(C_0(\mathbf{H}) \rtimes \Gamma) \xrightarrow{\sim} K^*(C_0(M)).$$

Lastly, [46, Lemma 3.8] and $\gamma = 1$ give

$$K_\Gamma^*(C(\partial\mathbf{H})) \xrightarrow{\sim} K^*(C(\partial\mathbf{H}) \rtimes \Gamma).$$

Via these isomorphisms, the sequence (3.2) can be identified with the six-term exact sequence associated to the extension (3.3). The identification of the maps $K^*(C_r^*(\Gamma)) \rightarrow K^*(C_0(M))$ as induced by taking the Kasparov product with the Euler class now follows by combining the argument in [23, Proposition 9] with [23, Theorem 38], yielding (3.4). \square

The exact sequence (3.4) simplifies further. We denote by $[\text{pt}] \in K^0(C_0(M))$ the class given by the homomorphism $C_0(M) \rightarrow \mathbf{C}, f \mapsto f(x)$, for some $x \in M$. Since M is connected this does not depend on the choice of x . Furthermore we denote by

$$\chi(M) := \sum_{k=0}^{\dim M} (-1)^k \text{rank } H^k(M, \mathbf{Z}),$$

the Euler characteristic of M . The following result uses the method of [24, Theorem 10.7].

Theorem 3.3. *For a discrete torsion-free subgroup $\Gamma \subset \text{Isom}(\mathbf{H})$, the homomorphism $\text{Eul}_1 : K^1(C_r^*(\Gamma)) \rightarrow K^1(C_0(M))$ vanishes and Eul_0 is given by*

$$\text{Eul}_0 : K^0(C_r^*(\Gamma)) \rightarrow K^0(C_0(M)), \quad [(C_r^*(\Gamma), \mathcal{H}, D)] \mapsto \chi(M) \text{Ind}(D^+) [\text{pt}].$$

In particular, if Γ is noncocompact or \mathbf{H} has odd dimension, there are short exact sequences

$$(3.5) \quad 0 \rightarrow K^1(C_0(M)) \xrightarrow{\partial} K^0(C(\partial\mathbf{H}) \rtimes \Gamma) \xrightarrow{i^*} K^0(C_r^*(\Gamma)) \rightarrow 0,$$

$$(3.6) \quad 0 \rightarrow K^0(C_0(M)) \xrightarrow{\partial} K^1(C(\partial\mathbf{H}) \rtimes \Gamma) \xrightarrow{i^*} K^1(C_r^*(\Gamma)) \rightarrow 0.$$

Proof. The exact sequences (3.5) and (3.6) are derived directly from Proposition 3.4. We now prove the statements about the maps Eul_* . Let $x \in M$, $\pi : \mathbf{H} \rightarrow M$ the quotient map and $\rho_x : C_0(\mathbf{H}) \rtimes \Gamma \rightarrow B(\ell^2(\pi^{-1}(x)))$ the induced representation. By [23, Example 24] and [23, Theorem 30] we find that

$$(3.7) \quad [(C_0(\mathbf{H}), L^2(\wedge^* \mathbf{H}), \mathcal{D}_{HR})] = \chi(M) [\pi_x] \in KK_0^\Gamma(C_0(\mathbf{H}), \mathbf{C}).$$

There is a factorisation $[\pi_x] = [\varphi_x] \otimes [1]$ where $[1] \in KK_0^\Gamma(C_0(\Gamma), \mathbf{C}) \simeq \mathbf{Z}$ is the class of the map $C_0(\Gamma) \rightarrow B(\ell^2(\Gamma))$ and $\varphi_x : C_0(\mathbf{H}) \rightarrow C_0(\Gamma)$ is defined through $\Gamma \rightarrow \mathbf{H}, \gamma \mapsto x\gamma$. This yields the explicit form of Eul_0 . Since $KK_1^\Gamma(C_0(\Gamma), \mathbf{C}) = KK_1(C_0(\Gamma) \rtimes \Gamma, \mathbf{C}) = KK_1(\mathbb{K}(\ell^2(\Gamma)), \mathbf{C}) = 0$, the statement $\text{Eul}_1 = 0$ follows. \square

In particular the Gysin sequence simplifies for all Bianchi groups and some Fuchsian groups.

3.2. The extension class. For a subgroup $H \subset G$, we denote the class defined through the exact sequence (3.1) by $[\text{Ext}] \in KK_1^H(C(\partial\mathbf{H}), C_0(\mathbf{H}))$. We now construct an equivariant Kasparov module representing $[\text{Ext}]$, and then employ Kasparov descent and Morita equivalence to obtain an explicit representative $[\partial] \in KK_1(C(\partial\mathbf{H}) \rtimes \Gamma, C_0(M))$ for torsion-free discrete subgroups $\Gamma \subset G$.

In the Poincaré ball model of hyperbolic $n + 1$ space \mathbf{H} , the boundary $\partial\mathbf{H}$ is the unit sphere in \mathbf{R}^{n+1} . For an element $g \in G$, write $|g'(\xi)| = |\det J_g(\xi)|$, the determinant of the Jacobian of the conformal transformation g . Consider $T_1\mathbf{H} := \mathbf{H} \times \partial\mathbf{H}$, which can be thought of as the unit tangent bundle of \mathbf{H} . The *Poisson kernel* is the map

$$(3.8) \quad P : T_1\mathbf{H} \rightarrow (0, \infty), \quad P(x, \xi) := \frac{1 - \|x\|^2}{\|x - \xi\|^2},$$

which for $g \in G$ satisfies the transformation rule

$$(3.9) \quad P(xg, \xi g) = |g'(\xi)|^{-1} P(x, \xi)$$

(see [56, Equation 5.1.2]). The *harmonic measure* ν_x on $\partial\mathbf{H}$ based at $x \in \mathbf{H}$ is defined to be unique probability measure on $\partial\mathbf{H}$ that is invariant under the action of the stabiliser G_x of x . Then ν_0 is normalised Lebesgue measure on $\partial\mathbf{H}$ and the measures ν_x satisfy

$$(3.10) \quad d\nu_x(\xi) = P(x, \xi)^n d\nu_0(\xi), \quad d\nu_{xg}(\xi g) = d\nu_x(\xi).$$

We consider the $G - C^*$ -algebra $C_0(\mathbf{H})$ as a G -equivariant C^* -module over itself. A second C^* -module is constructed using the harmonic measures on the boundary. The harmonic measures give an expectation

$$C_c(T_1\mathbf{H}) \rightarrow C_c(\mathbf{H}), \quad \rho(\Psi)(x, \xi) := \int \Psi(\xi, x) d\nu_x(\xi),$$

and hence a $C_0(\mathbf{H})$ -module $L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}$. This module carries a representation of the boundary algebra $C(\partial\mathbf{H})$ by pointwise multiplication.

Theorem 3.4. *Let $p = ww^*$ be the projection defined from the adjointable isometry*

$$w : C_0(\mathbf{H}) \rightarrow L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}, \quad w\Psi(x, \xi) := \Psi(x).$$

Then p commutes with G and has compact commutators with the $C(\partial\mathbf{H})$ -representation. The triple $(C(\partial\mathbf{H}), L^2(T_1\mathbf{H})_{C_0(\mathbf{H})}, F_p)$, with $F_p := 2p - 1$ is a G -equivariant KK -cycle for $(C(\partial\mathbf{H}), C_0(\mathbf{H}))$ representing the class of the boundary extension (3.1)

Proof. To see that w is adjointable define $w^*f(x) := \int_{\partial\mathbf{H}} f(\xi, x) d\nu_x\xi$. A quick computation shows that w and w^* are mutually adjoint:

$$\langle w\Psi, \Phi \rangle = \int \overline{w\Psi(\xi, x)} \Phi(\xi, x) d\nu_x(\xi) = \int \overline{\Psi(x)} \Phi(\xi, x) d\nu_x(\xi) = \langle \Psi, w^*\Phi \rangle.$$

Then computing the composition $w^*wf(x) = \int wf(\xi, p) d\nu_x(\xi) = \int f(x) d\nu_x(\xi) = f(x)$, that is $w^*w = 1$ and w is an isometry. Consider

$$(3.11) \quad w^*fw\Psi(x) = \left(\int f(\xi) d\nu_x\xi \right) \Psi(x).$$

The function $x \mapsto \int f(\xi) d\nu_x\xi$ is hyperbolically harmonic and continuous up to the boundary with limit f by [1, page 69] (see also [56, Theorem 5.1.5]). Hence $f \rightarrow w^*fw$ defines a G -invariant completely positive linear splitting $C(\partial\mathbf{H}) \rightarrow C_0(\mathbf{H})$.

Thus $(C(\partial\mathbf{H}), L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}, F_p)$ is a G -equivariant Kasparov module (see for example [28, Lemma 3.7]), and the usual Stinespring dilation argument shows that it represents the extension (3.1). \square

3.3. Kasparov descent and Morita equivalence. Consider the universal cover $\pi : \mathbf{H} \rightarrow M$, and the associated expectation

$$(3.12) \quad \rho_M : C_c(\mathbf{H}) \rightarrow C_c(M), \quad \rho_M(\Psi)(m) := \sum_{h \in \pi^{-1}(m)} \Psi(h),$$

defining a $C_c(M)$ -valued inner product on $C_c(\mathbf{H})$ by $\langle \Phi, \Psi \rangle := \rho_M(\overline{\Phi}\Psi)$. Denote its completion by $L^2_\pi(\mathbf{H})_{C_0(M)}$. The following result is a special case of the well known Morita equivalence for free and proper actions.

Lemma 3.5. *The C^* -algebra $\mathbb{K}(L^2_\pi(\mathbf{H})_{C_0(M)})$ is isomorphic to $C_0(\mathbf{H}) \rtimes \Gamma$, implementing the Morita equivalence with $C_0(M)$. The C^* -algebra $C(\overline{\mathbf{H}}) \rtimes \Gamma$ acts faithfully on $L^2_\pi(\mathbf{H})_{C_0(M)}$.*

The well known descent homomorphism [40, Theorem 6.1] is a map

$$(3.13) \quad j_\Gamma : KK_*^\Gamma(A, B) \rightarrow KK_*(A \rtimes \Gamma, B \rtimes \Gamma),$$

which can be explicitly defined on the level of cycles. We will describe the image of the cycle $(C(\partial\mathbf{H}), L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}, F_p)$ from Theorem 3.4 under the map j_Γ as well as its composition with the Morita equivalence from Lemma 3.5. This will furnish us with a representative of the mapping $K^*(C_0(M)) \rightarrow K^{*+1}(C(\partial\mathbf{H}) \rtimes \Gamma)$ appearing in the exact sequences (3.5) and (3.6).

Following [40, Section 6.1], the underlying $C_0(\mathbf{H}) \rtimes \Gamma$ module for the class $j_\Gamma([\text{Ext}])$ is given as the completion of $C_c(T_1\mathbf{H} \times \Gamma)$ in the $C_c(\mathbf{H}) \rtimes \Gamma$ -valued inner product

$$\langle \Phi, \Psi \rangle(x, \gamma) := \sum_{\delta \in \Gamma} \int_{\partial\mathbf{H}} \overline{\Phi(\xi, x, \delta)} \Psi(\xi\delta^{-1}, x\delta^{-1}, \delta\gamma) d\nu_x(\xi),$$

and left and right module structures

$$(f \cdot \Psi)(\xi, x, \gamma) := \sum_{\delta \in \Gamma} f(\xi, \delta) \Psi(\xi\delta, x\delta, \delta^{-1}\gamma), \quad (\Psi \cdot g)(\xi, x, \gamma) = \sum_{\delta \in \Gamma} \Psi(\xi, x, \delta) g(x\delta, \delta^{-1}\gamma).$$

The operator F_p is defined by viewing $\Psi_\gamma(\xi, x) := \Psi(\xi, x, \gamma)$ as an element of $C_c(T_1\mathbf{H})$ for each $\gamma \in \Gamma$. The product with the Morita equivalence $L^2_\pi(\mathbf{H})_{C_0(M)}$ is now easily described. The map

$$m : C_c(T_1\mathbf{H} \times \Gamma) \otimes_{C_c(\mathbf{H} \times \Gamma)} C_c(\mathbf{H}) \rightarrow C_c(T_1\mathbf{H}), \quad m(\Psi \otimes \Phi)(\xi, x) = \sum_{\delta \in \Gamma} \Psi(\xi, x, \delta) \Phi(x\delta),$$

is surjective and compatible with the balancing relation. The resulting $C_0(M)$ -valued inner product on $C_c(T_1(\mathbf{H}))$ is given by

$$\langle \Psi, \Phi \rangle(m) := \sum_{x \in \pi^{-1}(m)} \int_{\partial\mathbf{H}} \overline{\Psi(\xi, x)} \Phi(\xi, x) d\nu_x \xi.$$

The left representation and the operator $F_p = 2p - 1$ are induced from tensoring with the identity operator. We summarize the above findings:

Corollary 3.6 (of Theorem 3.4). *The class*

$$[\partial] := j_\Gamma([\text{Ext}]) \otimes_{C_0(\mathbf{H}) \rtimes \Gamma} [L_\pi^2(\mathbf{H})_{C_0(M)}] \in KK_1(C(\partial\mathbf{H}) \rtimes \Gamma, C_0(M)),$$

is represented by the bounded Kasparov module $(C(\partial\mathbf{H}) \rtimes \Gamma, L_\pi^2(T_1\mathbf{H})_{C_0(M)}, F_p)$.

Consequently the boundary map $\partial : K^0(C_0(M)) \rightarrow K^1(C(\partial\mathbf{H}) \rtimes \Gamma)$ is implemented by the Kasparov product with $(C(\partial\mathbf{H}) \rtimes \Gamma, L_\pi^2(T_1\mathbf{H}, \nu_x)_{C_0(M)}, F_p)$.

3.4. Hecke equivariance. Our purpose is now to show that the exact hexagon (3.4) is equivariant for the action of the Hecke operator T_g on the various algebras appearing in (3.4). We first consider compatibility of the Hecke operators with Morita equivalences arising from free and proper actions.

Let X be a G -space such that Γ acts freely and properly on X , $\pi : X \rightarrow M := X/\Gamma$ the covering map and $L_\pi^2(X)_{C_0(M)}$ the associated $C_0(X) \rtimes \Gamma$ - $C_0(M)$ Morita equivalence bimodule. There is a well-known unitary isomorphism

$$(3.14) \quad T_g^\Gamma \otimes_{C_0(X) \rtimes \Gamma} L_\pi^2(X)_{C_0(M)} \rightarrow L_\pi^2(X)_{C_0(M)}^d, \quad (\Phi_i) \otimes \Psi \mapsto (\Phi_i \Psi),$$

of right $C_0(M)$ -modules. We show that the same is true for $L_\pi^2(X) \otimes_{C_0(M)} T_g^M$ and then compare the left actions. Consider the *fiber product* with its natural covering maps

$$X \times_{\tau_g} M_g := \{(x, m) \in X \times M_g : \pi(x) = \tau_g(m)\}, \quad M \xleftarrow{\pi} X \times_{\tau_g} M_g \xrightarrow{\pi_g} M,$$

making $C_c(X \times_{\tau_g} M_g)$ into a $C_c(M)$ inner product bimodule. There is a well-defined map

$$w : C_c(X) \otimes_{C_c(M)} C_c(M_g) \rightarrow C_c(X \times_{\tau_g} M_g), \quad w(\Psi \otimes \Phi)(x, m) = \Psi(x)\Phi(m),$$

of right $C_c(M)$ -modules preserving the inner product. By standard support arguments, w is shown to be surjective, and $L_\pi^2(X) \otimes_{C_0(M)} T_g^M$ is obtained as a completion of $C_c(X \times_{\tau_g} M_g)$.

Lemma 3.7. *Let δ_i be a set of right coset representatives for $\Gamma_{g^{-1}}$, $\Gamma = \bigsqcup_{i=1}^d \delta_i \Gamma_{g^{-1}}$. The continuous open maps*

$$\varphi_i : X \rightarrow X \times_{\tau_g} M_g, \quad x \mapsto (xg\delta_i^{-1}, [x]), \quad i = 1, \dots, d,$$

assemble to a homeomorphism $\varphi : \bigsqcup_{i=1}^d X \rightarrow X \times_{\tau_g} M_g$.

Proof. Since $\tau_g([x]) = \pi(xg) = \pi(xg\delta_i^{-1})$, each φ_i is well-defined and injective. Moreover, the φ_i assemble to an injective map on the disjoint union

$$\varphi : \bigsqcup_{i=1}^d X \rightarrow X \times_{\tau_g} M_g.$$

This can be seen by assuming that $\varphi_i(x) = \varphi_j(x')$ for some $x, x' \in X$. Then

$$xg\delta_i^{-1} = x'g\delta_j^{-1}, \quad [x] = [x'] \Leftrightarrow \exists \gamma \in \Gamma_g \quad x\gamma = x',$$

which gives $xg\delta_i^{-1} = x\gamma g\delta_j^{-1} = xg\delta_j^{-1}\gamma g\delta_j^{-1}$.

Now $g^{-1}\gamma g\delta_j^{-1} \in \Gamma$ because $\gamma \in \Gamma_g$ and freeness of the Γ action on X gives $g^{-1}\gamma g\delta_j^{-1} = \delta_i^{-1}$. Hence we find $\delta_i^{-1}\delta_j = g^{-1}\gamma g \in \Gamma_{g^{-1}}$, which gives $i = j$ and hence $x = x'$ as well, as desired.

It remains to show that φ is surjective, so let $(x', [x]) \in X \times_{\tau_g} M_g$. Since $\pi(x) = \pi(x'g)$, there is $\gamma' \in \Gamma$ such that $xg = x'\gamma' = x'\delta_i\gamma'^{-1}$ for some unique i and $\gamma' \in \Gamma_{g^{-1}}$. Hence, we find

$x' = xg\gamma\delta_i^{-1} = xg\gamma g^{-1}g\delta_i^{-1}$. Now in M_g it holds that $[x] = [xg\gamma g^{-1}]$, because $g\gamma g^{-1} \in \Gamma_g$. Therefore

$$(x, [x']) = (xg\gamma g^{-1}g\delta_i^{-1}, [xg\gamma g^{-1}]) = \varphi_i(xg\gamma g^{-1}),$$

proving that φ is surjective. \square

Proposition 3.8. *Let X be a G -space such that Γ acts freely and properly on X , $M := X/\Gamma$ and $L_\pi^2(X)_{C_0(M)}$ the associated $C_0(X) \rtimes \Gamma$ - $C_0(M)$ Morita equivalence bimodule. For each $g \in C_G(\Gamma)$ there is a unitary isomorphism of $C_0(X) \rtimes \Gamma$ - $C_0(M)$ -bimodules*

$$T_g^\Gamma \otimes_{C_0(X) \rtimes \Gamma} L_\pi^2(X) \xrightarrow{\sim} L_\pi^2(X) \otimes_{C_0(M)} T_g^M.$$

In particular we have the identity

$$[T_g^\Gamma] \otimes_{C_0(\mathbf{H}) \rtimes \Gamma} [L_\pi^2(X)] = [L_\pi^2(X)] \otimes_{C_0(M)} [T_g^M] \in KK_0(C_0(X) \rtimes \Gamma, C_0(M)).$$

Proof. The homeomorphism φ from Lemma 3.7 induces a unitary isomorphism of right $C_0(M)$ -modules

$$L_\pi^2(X) \otimes_{C_0(M)} T_g^M \cong L_\pi^2(X)^d \cong T_g^\Gamma \otimes_{C_0(X) \rtimes \Gamma} L_\pi^2(X)_{C_0(M)}.$$

As before we write elements of the module $L_\pi^2(\bigsqcup_{i=1}^d X)_{C_0(M)}$ as columns $\Psi = (\Psi_i)_{i=1}^d$. The action of a function $f \in C_0(X)$ is given by

$$\varphi^* f \varphi^{*-1}(\Psi)_i(x) = f \varphi^{*-1}(\Psi)((x\delta_i g^{-1}, [x])) = (f(xg\delta_i^{-1})\Psi_i(x)),$$

which equals the action coming from the identification $L_\pi^2(X)^d \cong T_g^\Gamma \otimes_{C_0(X) \rtimes \Gamma} L_\pi^2(X)$. The action of a group element γ is given by

$$\varphi^* u_\gamma \varphi^{*-1}(\Psi)_i(x) = u_\gamma \varphi^{*-1}(\Psi)(xg\delta_i^{-1}, [x]) = \varphi^{*-1}(\Psi)(xg\delta_i^{-1}\gamma, [x]).$$

We compute further by using Lemma 2.3 and observing that

$$xg\delta_i^{-1}\gamma = xt_i(\gamma^{-1})^{-1}g\delta_{\gamma^{-1}(i)}^{-1},$$

and since $t_i(\gamma^{-1})^{-1} \in \Gamma_g$ we find $[x] = [xt_i(\gamma^{-1})^{-1}]$ so

$$\begin{aligned} \varphi^{*-1}(\Psi)(xg\delta_i^{-1}\gamma, [x]) &= \varphi^{*-1}(\Psi)(xt_i(\gamma^{-1})^{-1}g\delta_{\gamma^{-1}(i)}^{-1}, [xt_i(\gamma^{-1})^{-1}]) \\ &= \Psi_{\gamma^{-1}(i)}(xt_i(\gamma^{-1})^{-1}) = \Psi_{\gamma^{-1}(i)}(xt_{\gamma^{-1}(i)}(\gamma)). \end{aligned}$$

As above, this equals the action coming from the identification

$$L_\pi^2(X)^d \cong T_g^\Gamma \otimes_{C_0(X) \rtimes \Gamma} L_\pi^2(X).$$

This completes the proof. \square

As a right $C(\partial\mathbf{H}) \rtimes \Gamma$ -module, T_g^Γ is free of rank d . Therefore, as in (3.14) the map

$$(3.15) \quad u : T_g^\Gamma \otimes_{C(\partial\mathbf{H}) \rtimes \Gamma} L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x) \rightarrow L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)^d, \quad (\Psi_i) \otimes f \mapsto (\Psi_i f),$$

defined through coordinatewise product, is a unitary right module map. Using this map we can define the operator $u^* \text{diag}(p)u$ on $T_g^\Gamma \otimes_{C(\partial\mathbf{H}) \rtimes \Gamma} L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)$. By a slight abuse of notation, we denote this operator by $1 \otimes p$.

Theorem 3.9. *There is a unitary isomorphism of $(C(\partial\mathbf{H}) \rtimes \Gamma, C_0(\mathbf{H}) \rtimes \Gamma)$ -bimodules*

$$L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x) \otimes_{C_0(\mathbf{H}) \rtimes \Gamma} T_g^\Gamma \xrightarrow{\sim} T_g^\Gamma \otimes_{C(\partial\mathbf{H}) \rtimes \Gamma} L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x),$$

intertwining the operators $p \otimes 1$ and $1 \otimes p$. We have the identity

$$[T_g^\Gamma] \otimes [\partial] = [\partial] \otimes [T_g^M] \in KK_1(C(\partial\mathbf{H}) \rtimes \Gamma, C_0(M)).$$

In particular, the boundary map $\partial : K^0(C_0(M)) \rightarrow K^1(C(\partial\mathbf{H}) \rtimes \Gamma)$ is Hecke equivariant.

Proof. By Corollary 3.6 and Lemma 3.8, the second statement follows from the first because it implies that

$$j_\Gamma([\text{Ext}]) \otimes [T_g^\Gamma] = [T_g^\Gamma] \otimes j_\Gamma([\text{Ext}]),$$

and the class $j_\Gamma([\text{Ext}])$ is represented by the Kasparov module $(L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)_{C_0(\mathbf{H}) \rtimes \Gamma}, F_p)$. First we compare the bimodules $L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x) \otimes_{C_0(\mathbf{H}) \rtimes \Gamma} T_g^\Gamma$ and $T_g^\Gamma \otimes_{C(\partial\mathbf{H}) \rtimes \Gamma} L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)$. The right $C_0(\mathbf{H}) \rtimes \Gamma$ -module $L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)_{C_0(\mathbf{H}) \rtimes \Gamma}^d$ is a left $B \rtimes_r \Gamma$ module for either of the Γ - C^* -algebras $B = C(\partial\mathbf{H}), C_0(\mathbf{H}), C_0(T_1\mathbf{H})$ via Equation (2.7). The map u in Equation (3.15) is readily seen to be a left $C(\partial\mathbf{H}) \rtimes \Gamma$ module map.

We now construct a unitary isomorphism of $(C(\partial\mathbf{H}) \rtimes \Gamma, C_0(\mathbf{H}) \rtimes \Gamma)$ -bimodules

$$\alpha : L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x) \otimes_{C_0(\mathbf{H}) \rtimes \Gamma} T_g^\Gamma \xrightarrow{\sim} L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)_{C_0(\mathbf{H}) \rtimes \Gamma}^d.$$

To achieve this we view $L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)$ as a completion of $C_c(T_1\mathbf{H} \rtimes \Gamma)$ and we consider the embedding of right $C_0(\mathbf{H}) \rtimes \Gamma$ bimodules

$$\beta : T_g^\Gamma = (C_0(\mathbf{H}) \rtimes \Gamma)^d \rightarrow L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)^d, \quad \beta(\Psi_i)(x, \xi, \gamma) := (\Psi_i(x, \gamma)).$$

We view $L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)^d$ as a left $C_c(X \rtimes \Gamma)$ -module, where $X = T_1\mathbf{H}$ or $X = \partial\mathbf{H}$, via

$$C_c(X \rtimes \Gamma) \times L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)^d \rightarrow L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)^d, \quad f \cdot (\Psi_i) := t_g(f)(\Psi_i),$$

using Equation (2.7) in Proposition 2.4. Now define a map

$$\alpha : C_c(T_1\mathbf{H} \rtimes \Gamma) \otimes_{C_c(\mathbf{H} \rtimes \Gamma)} T_g^\Gamma \rightarrow L^2(T_1\mathbf{H} \rtimes \Gamma, \nu_x)^d, \quad \alpha(f \otimes (\Psi_i)) := f \cdot \beta(\Psi_i),$$

which respects the $(C_c(\partial\mathbf{H} \rtimes \Gamma), C_c(\mathbf{H} \rtimes \Gamma))$ bimodule structures because

$$\alpha(f * h \otimes (\Psi_i)) = t_g(f * h)(\beta(\Psi_i)) = t_g(f)t_g(h)\beta(\Psi_i) = t_g(f)\alpha(h \otimes (\Psi_i)),$$

and the right module structure is respected because β is a right module map. We find

$$\begin{aligned} \alpha(pf \otimes (\Psi_i))(\delta) &= t_g(pf)\beta(\Psi_i)(\delta) = \sum_{\gamma} g_i^{-1} pf(\gamma) t_i(\gamma^{-1})^{-1} \beta\Psi_{\gamma^{-1}(i)}(t_i(\gamma^{-1})\delta) \\ &= p \left(\sum_{\gamma} g_i^{-1} f(\gamma) t_i(\gamma^{-1})^{-1} \beta\Psi_{\gamma^{-1}(i)}(t_i(\gamma^{-1})\delta) \right) = p(t_g(f)\beta(\Psi_i)), \end{aligned}$$

by Proposition 2.4 using that $p(\Phi\beta(\Psi_i)) = (p\Phi)(\beta(\Psi))_i$ and p is G -invariant. Thus,

$$\text{diag}(p)\alpha(f \otimes (\Psi_i)) = \alpha(pf \otimes (\Psi_i)), \quad (\text{diag}(p)) \circ \alpha = \alpha \circ (p \otimes 1).$$

The fact that α is unitary follows from a lengthy but straightforward calculation which we omit. \square

We arrive at the main general result of this section, expressing the compatibility of the various Hecke operators we construct.

Theorem 3.10. *The Gysin-sequences in K -homology*

$$\begin{array}{ccccccc} 0 & \longrightarrow & K^1(C_0(M)) & \xrightarrow{\partial} & K^0(C(\partial\mathbf{H}) \rtimes \Gamma) & \longrightarrow & K^0(C_r^*(\Gamma)) \\ & & & & & & \downarrow \\ 0 & \longleftarrow & K^1(C_r^*(\Gamma)) & \longleftarrow & K^1(C(\partial\mathbf{H}) \rtimes \Gamma) & \xleftarrow{\partial} & K^0(C_0(M)) \end{array}$$

and K -theory

$$\begin{array}{ccccccc} 0 & \longrightarrow & K_1(C_r^*(\Gamma)) & \longrightarrow & K_1(C(\partial\mathbf{H}) \rtimes \Gamma) & \xrightarrow{\partial} & K_0(C_0(M)) \\ & & & & & & \downarrow \\ 0 & \longleftarrow & K_1(C_0(M)) & \xleftarrow{\partial} & K_0(C(\partial\mathbf{H}) \rtimes \Gamma) & \longleftarrow & K_0(C_r^*(\Gamma)) \end{array}$$

are Hecke-equivariant.

Proof. This follows by combining Proposition 3.8, 3.9 and the observation that the inclusion $K_*(C_r^*(\Gamma)) \rightarrow K_*(C(\partial\mathbf{H}) \rtimes \Gamma)$ and restriction $K_*(C(\partial\mathbf{H}) \rtimes \Gamma) \rightarrow K_*(C_r^*(\Gamma))$ are Hecke equivariant by construction. Hecke equivariance of the Euler maps $K^*(C_0(M)) \rightarrow K^*(C_r^*(\Gamma))$ follows from the commutative diagram

$$\begin{array}{ccc} K^*(C_r^*(\Gamma)) & \xrightarrow{(\iota^*)^{-1}} & K^*(C(\overline{\mathbf{H}}) \rtimes \Gamma) \\ \text{Eul} \downarrow & & \downarrow \\ K^*(C_0(M)) & \longleftarrow & K^*(C_0(\mathbf{H}) \rtimes \Gamma) \end{array}$$

and since $T_g \iota^* = \iota^* T_g$ implies $(\iota^*)^{-1} T_g = T_g (\iota_*)^{-1}$, the map Eul is a composition of Hecke equivariant maps, whence Hecke equivariant. The argument for the K -theory sequence is identical. \square

4. K -cycles for the reduced Bianchi group C^* -algebra

We will describe a naturally defined map $s : H^1(\Gamma, \mathbf{Z}) \rightarrow K^1(C_r^*(\Gamma))$ for a discrete group Γ of hyperbolic isometries and show that in the special case when Γ is a torsion-free finite index subgroup of a Bianchi group, our explicit map s is a Hecke equivariant isomorphism.

It is well known that for a general discrete group Γ , there is a homomorphism $t : H_1(\Gamma, \mathbf{Z}) \rightarrow K_1(C_r^*(\Gamma))$. This homomorphism has been studied for instance by Matthey [51] in the context of the Baum-Connes conjecture. We show that when Γ is a torsion-free finite index subgroup of a Bianchi group, t is an isomorphism and that the homological pairing $H^1 \times H_1 \rightarrow \mathbf{Z}$ and the index pairing $K^1 \times K_1 \rightarrow \mathbf{Z}$ commute with the isomorphisms s and t .

4.1. Group cocycles and index theory. In this subsection, Γ is an arbitrary countable discrete group. Let $c : \Gamma \rightarrow \mathbf{Z}$ be an integral group cocycle, simply a group homomorphism, and denote by Γ_c its kernel. The multiplication operator

$$D_c : C_c(\Gamma) \rightarrow C_c(\Gamma),$$

defined through $(D_c f)(\gamma) := -c(\gamma)f(\gamma)$ extends to a selfadjoint regular operator in the C^* -completion E_c of $C_c(\Gamma)$ over Γ_c . This gives an unbounded Kasparov module (E_c, D_c) and

an element in the group $KK_1(C_r^*(\Gamma), C_r^*(\Gamma_c))$, as a special case of the construction in [52, **Theorem 3.2.2, Lemma 3.4.1**],

To describe the pairing of this cycle with $K_1(C_r^*(\Gamma))$, we need a concrete description of the latter group. First note that here we use the surjective (Hurewicz) map $\Gamma \rightarrow \Gamma^{ab} \simeq H_1(\Gamma, \mathbf{Z})$, we can represent homology classes by elements $\delta \in \Gamma$. A group element $\delta \in \Gamma$ defines a unitary u_δ in the reduced C^* -algebra $C_r^*(\Gamma)$, and thus a class $[u_\delta] \in K_1(C_r^*(\Gamma))$ via the standard picture of K_1 . This gives us a homomorphism $H_1(\Gamma, \mathbf{Z}) \rightarrow K_1(C_r^*(\Gamma))$, for any discrete group Γ .

Definition 4.1. *We define the norm of a cocycle $c : \Gamma \rightarrow \mathbf{Z}$ to be the nonnegative integer*

$$|c| := \min\{|\langle c(\gamma) \rangle| : \gamma \in \Gamma, \gamma \notin \Gamma_c\}.$$

A cocycle c is normalised if $1 \in c(\Gamma) \subset \mathbf{Z}$. The norm of the 0-cocycle is defined to be ∞ .

Since $c(\Gamma) = |c|\mathbf{Z}$, the statement that c is normalised is equivalent to saying that $c(\Gamma) = \mathbf{Z}$. Any cocycle is an integral multiple of a normalised cocycle, and thus $H^1(\Gamma, \mathbf{Z})$ is generated by normalised cocycles. If c is normalised, the short exact sequence of groups

$$0 \rightarrow \Gamma_c \rightarrow \Gamma \rightarrow \mathbf{Z} \rightarrow 0,$$

admits a non canonical splitting by choosing $g \in c^{-1}(1)$ and define $s : \mathbf{Z} \rightarrow \Gamma$, $n \mapsto g^n$. Any such splitting determines a group isomorphism $\Gamma \cong \Gamma_c \rtimes_s \mathbf{Z}$ (semidirect product) and a C^* -algebra isomorphism $C_r^*(\Gamma) \cong C_r^*(\Gamma_c) \rtimes \mathbf{Z}$.

Proposition 4.2. *Let $c : \Gamma \rightarrow \mathbf{Z}$ be a normalised cocycle. The Kasparov product*

$$K_1(C_r^*(\Gamma)) \times KK_1(C_r^*(\Gamma), C_r^*(\Gamma_c)) \rightarrow K_0(C_r^*(\Gamma_c)),$$

maps the pair $([u_\delta], [D_c])$ to the class

$$\text{sgn}(c(\delta))[C_r^*(\Gamma_c)^{|c(\delta)|}] = c(\delta)[1_{C_r^*(\Gamma_c)}].$$

Proof. Choosing $g \in c^{-1}(1)$ gives a generating set $\{e_{g^n}\}_{n \in \mathbf{Z}}$ for the module E and a decomposition

$$(4.1) \quad E \simeq \bigoplus_{n \in \mathbf{Z}} C_r^*(\Gamma_c), \quad e_\gamma \mapsto e_{g^{c(\gamma)}} u_{g^{-c(\gamma)}\gamma},$$

under which the operator D_c becomes multiplication by $-n \in \mathbf{Z}$. Denote by $p_c : E_c \rightarrow E_c$ the projection onto the positive spectrum of D_c , which is adjointable by the decomposition (4.1). The Fredholm operator given by $F_c := D_c(1 + D_c^2)^{-\frac{1}{2}}$ is a compact perturbation of the adjointable operator $S_c = 2p_c - 1$ and defines the same class as $[D_c] = [F_c] \in KK_1(C_r^*(\Gamma), C_r^*(\Gamma_c))$. For $\delta \in \Gamma$, since $u_\delta e_{g^n} = e_{g^{n+c(\delta)}} u_{g^{-n-c(\delta)}\delta} g^n$ it follows that

$$\text{imp}_c u_\delta p_c = \text{span}\{e_{g^n} : n \leq \min\{0, -c(\delta)\}\},$$

which is a complemented submodule, and hence so is $\text{imp}_c u_\delta^* p_c$. Thus by [44, **Ch. 3**] the operator $p_c u_\delta p_c + 1 - p_c$ admits a polar decomposition and by [36, **Theorem 7.8**] and the argument in [36, **Lemma 2.1**], the Kasparov product may be computed as a higher index, that is

$$[u_\delta] \otimes [(E_c, D_c)] = [\ker p_c u_\delta p_c] - [\text{coker } p_c u_\delta p_c] \in K_0(C_r^*(\Gamma_c)).$$

As above we have

$$\ker p_c u_\delta p_c = \text{span}\{e_{g^n} : -c(\delta) < n \leq 0\},$$

and since c is normalised, this module is isomorphic to the free module of rank $|c(\delta)|$ over $C_r^*(\Gamma_c)$ if $c(\delta) > 0$ and 0 otherwise. Since

$$\operatorname{coker} p_c u_\delta p_c = \ker p_c u_{\delta^{-1}} p_c,$$

the statement follows. \square

In order to obtain a genuine unbounded Fredholm module from a cocycle, we need to get rid of the algebra $C_r^*(\Gamma_c)$ in Proposition 4.2. It is not clear how to do this without making more assumptions on Γ . In the next subsection, we achieve this when Γ is a discrete group of hyperbolic isometries.

4.2. The unbounded γ -element. In case Γ is a group of isometries of a simply connected, complete Riemannian manifold X with nonpositive sectional curvature, Kasparov's Dirac/dual-Dirac construction [39] gives a canonical element $[\gamma_X] \in K^0(C_r^*(\Gamma_c))$. In this section we work with the real hyperbolic $n+1$ -space $X = \mathbf{H}$, but this is not necessary. Let ρ denote the function $\rho(x) := d_{\mathbf{H}}(0, x)$, $L^2(\wedge^* \mathbf{H})$ the Hilbert space of L^2 -sections of the exterior algebra bundle of \mathbf{H} , \mathcal{D}_{HR} the Hodge-DeRham operator, \hat{c} the Clifford multiplication and d the exterior derivative.

The following lemma is well-known in the case $s = 1$ and we state it for convenience.

Lemma 4.3 ([39]). *For $0 < s \leq 1$ the triple $(\mathbf{C}, L^2(\wedge^* \mathbf{H}), \mathcal{D}_{\rho,s} := \mathcal{D}_{HR} + \rho^s \hat{c}(d\rho))$ is a G -equivariant unbounded Fredholm module representing the class $[\gamma_{\mathbf{H}}] = 1 \in KK^G(\mathbf{C}, \mathbf{C})$. In particular, for any discrete subgroup $\Gamma \subset G$ the triple*

$$(C_r^*(\Gamma), L^2(\wedge^* \mathbf{H}), \mathcal{D}_{\rho,s}),$$

is an unbounded Fredholm module and $\operatorname{Ind}(\mathcal{D}_{\rho,s}^+) = 1$.

Proof. The statement follows from the observation that $(1 + \rho)^s \hat{c}(d\rho)$ a bounded perturbation of $\rho^s \hat{c}(d\rho)$ and is a representative for Kasparov's dual Dirac element. The graded commutator

$$[\mathcal{D}_{HR}, (1 + \rho)^s \hat{c}(d\rho)] = [\mathcal{D}_{HR}, (1 + \rho)^s] \hat{c}(d\rho) + (1 + \rho)^s [\mathcal{D}_{HR}, \hat{c}(d\rho)],$$

is relatively bounded to $(1 + \rho)^s$, so $\mathcal{D}_{\rho,s}$ represents the Kasparov product of the Dirac and dual-Dirac element and is hence in the class of $[\gamma_{\mathbf{H}}]$. For a discrete subgroup $\Gamma \subset G$ its action on \mathbf{H} defines a representation of $C_r^*(\Gamma)$ on $L^2(\wedge^* \mathbf{H})$. The statement about the index is immediate. \square

Lemma 4.4. *For $0 < s \leq 1$ and any element $g \in G$ we have the estimate*

$$\|\rho^s [\hat{c}(d\rho), u_g]\| \leq 2d_{\mathbf{H}}^s(0, 0g).$$

Proof. By the proof of [39, Lemma 5.3], for $x \neq 0$ it holds that

$$\|[\hat{c}(d_x \rho), u_g]\| \leq 2d_{\mathbf{H}}(0, 0g)(d_{\mathbf{H}}(0, x) + d_{\mathbf{H}}(0, xg))^{-1},$$

so we assume $0g \neq 0$ as well. This yields the estimate

$$\begin{aligned} \|d_{\mathbf{H}}^s(0, x)[\hat{c}(d_x \rho), u_g]\| &\leq 2d_{\mathbf{H}}(0, 0g)d_{\mathbf{H}}^s(0, x)(d_{\mathbf{H}}(0, x) + d_{\mathbf{H}}(0, xg))^{-1} \\ &\leq 2d_{\mathbf{H}}(0, 0g)(d_{\mathbf{H}}(0, x) + d_{\mathbf{H}}(0, xg))^{s-1} \\ &\leq 2d_{\mathbf{H}}(0, 0g)d_{\mathbf{H}}(0, 0g)^{s-1} = 2d_{\mathbf{H}}(0, 0g)^s, \end{aligned}$$

which produces the claimed norm estimate. \square

Lemma 4.5. *Let $0 < s \leq 1$, $\gamma_1, \dots, \gamma_k \in \Gamma$ and $x \in \mathbf{H}$. Then*

$$d_{\mathbf{H}}^s(x, x\gamma_k \cdots \gamma_1) \leq \left(\sum_{i=1}^k d_{\mathbf{H}}(x, x\gamma_i) \right)^s \leq \sum_{i=1}^k d_{\mathbf{H}}^s(x, x\gamma_i).$$

Proof. This is a straightforward induction. For $k = 1$ there is nothing to prove. Then for $k > 1$ we write

$$\begin{aligned} d_{\mathbf{H}}^s(x, x\gamma_k \cdots \gamma_1) &\leq (d_{\mathbf{H}}(x, x\gamma_{k-1} \cdots \gamma_1) + d_{\mathbf{H}}(x\gamma_{k-1} \cdots \gamma_1, x\gamma_k \cdots \gamma_1))^s \\ &= (d_{\mathbf{H}}(x, x\gamma_{k-1} \cdots \gamma_1) + d_{\mathbf{H}}(x, x\gamma_k))^s \leq \left(\sum_{i=1}^k d_{\mathbf{H}}(x, x\gamma_i) \right)^s \leq \sum_{i=1}^k d_{\mathbf{H}}^s(x, x\gamma_i), \end{aligned}$$

which are the desired inequalities. \square

We wish to construct the Kasparov product of the element $[D_c] \in KK_1(C_r^*(\Gamma), C_r^*(\Gamma_c))$ and $[\gamma_{\mathbf{H}}] \in K^1(C_r^*(\Gamma_c))$ in order to obtain an unbounded Fredholm module and a class in $K^1(C_r^*(\Gamma))$. In order to do this we define, for $g \in c^{-1}(|c|)$

$$(4.2) \quad 1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}(e_\gamma \otimes \psi) := e_{g^{c(\gamma)}} \otimes \mathcal{D}_{\rho,s}(u_{g^{-c(\gamma)}} \gamma \psi)$$

which is a densely defined symmetric operator with initial domain $C_c(\Gamma) \otimes_{C_c(\Gamma_c)} \text{Dom } \mathcal{D}_{\rho,s}$. We then consider the densely defined symmetric operator

$$(4.3) \quad D_c \otimes \sigma + 1 \otimes_{\nabla} \mathcal{D}_{\rho,s} = \begin{pmatrix} -c & 1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}^+ \\ 1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}^- & c \end{pmatrix},$$

on the Hilbert space $E \otimes_{C_r^*(\Gamma_c)} L^2(\wedge^* \mathbf{H})$ with grading operator σ , decomposed according to even and odd forms $L^2(\wedge^* \mathbf{H}) = L^2(\wedge^+ \mathbf{H}) \oplus L^2(\wedge^- \mathbf{H})$.

We recall from the appendix to [29], that the notion of unbounded Fredholm module can be loosened.

Definition 4.6. *An unbounded Fredholm module is a triple (\mathcal{A}, H, D) , where*

- (1) \mathcal{A} is a $*$ -algebra represented on the $\mathbf{Z}/2$ -graded Hilbert space H ;
- (2) D is a self-adjoint operator such that $a(D \pm i)^{-1} \in \mathbf{K}(H)$;
- (3) for all $a \in \mathcal{A}$, $a \text{Dom } D \subset \text{Dom } D$ and there exists $\varepsilon > 0$ such that $[D, a](1 + D^2)^{-\frac{1-\varepsilon}{2}}$ and $(1 + D^2)^{-\frac{1-\varepsilon}{2}}[D, a]$ extend to bounded operators.

If ε can be chosen independent of $a \in \mathcal{A}$ then (\mathcal{A}, H, D) is called an ε -unbounded Fredholm module.

Theorem 4.7. *Let $\Gamma \subset \text{Isom}(\mathbf{H})$ be a discrete group, $0 < s < 1$ and $c : \Gamma \rightarrow \mathbf{Z}$ be a normalised cocycle. The Kasparov product of the classes $[(E_c, D_c)]$ and $[\gamma_{\mathbf{H}}]$ is represented by the $(1 - s)$ unbounded Fredholm module*

$$(C_r^*(\Gamma), E \otimes_{C_r^*(\Gamma_c)} L^2(\wedge^* \mathbf{H}), \mathcal{D}_{c,s} := D_c \otimes \sigma + 1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}),$$

and in particular is independent of the choice of $g \in c^{-1}(1)$.

Proof. Essential self-adjointness and compact resolvent of the operator $\mathcal{D}_{c,s}$ in (4.3) follows from general considerations in [54]. It remains to show condition 3 of Definition 4.6 is satisfied

for the unitaries u_γ generating $C_r^*(\Gamma)$ and $\varepsilon = 1 - s$. For the operator $D_c \otimes \sigma$ this follows from the fact that $[D_c, u_\gamma]$ defines an adjointbale operator on E_c .

For $1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}$, Equation (4.2) shows that the commutator can be expressed as

$$[1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}, u_\gamma](e_{g^n} \otimes \psi) = e_{g^{n+c(\gamma)}} \otimes [\mathcal{D}_{\rho,s}, u_{g^{-n-c(\gamma)}\gamma g^n}]\psi.$$

The Hilbert space $E \otimes L^2(\wedge^* \mathbf{H})$ decomposes as a direct sum

$$E \otimes_{C_r^*(\Gamma_c)} L^2(\wedge^* \mathbf{H}) \cong \bigoplus_{n \in \mathbf{Z}} e_{g^n} \otimes L^2(\wedge^* \mathbf{H}),$$

and it suffices to control the supremum of the norms $\|\cdot\|_n$ of the operators

$$[1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}, u_\gamma](1 + D_c^2 + (1 \otimes_{\nabla} \mathcal{D}_{\rho,s})^2)^{-\frac{s}{2}} : e_{g^n} \otimes L^2(\wedge^* \mathbf{H}) \rightarrow e_{g^{n+c(\gamma)}} \otimes L^2(\wedge^* \mathbf{H}).$$

To compute the commutator $[\mathcal{D}_{\rho,s}, u_{g^{-n-c(\gamma)}\gamma g^n}]$, we observe that $\mathcal{D}_{\rho,s} = \mathcal{D}_{HR} + \rho^s \hat{c}(d\rho)$, and \mathcal{D}_{HR} commutes with u_δ for all δ . So we need only concern ourselves with the dual Dirac part. To this end we expand

$$[\rho^s \hat{c}(d\rho), u_{g^{-n-c(\gamma)}\gamma g^n}] = [\rho^s, u_{g^{-n-c(\gamma)}\gamma g^n}] \hat{c}(d\rho) + \rho^s [\hat{c}(d\rho), u_{g^{-n-c(\gamma)}\gamma g^n}].$$

Since $\|\hat{c}(d\rho)\| = 1$, the norm of the first term is controlled by

$$\sup_{x \in \mathbf{H}} |(d_{\mathbf{H}}^s(0, x) - d_{\mathbf{H}}^s(0, xg^{-n-c(\gamma)}\gamma g^n))| \leq d_{\mathbf{H}}^s(0, 0g^{-n-c(\gamma)}\gamma g^n),$$

whereas Lemma 4.4 takes care of the second term with the estimate

$$\|\rho^s [\hat{c}(d\rho), u_{g^{-n-c(\gamma)}\gamma g^n}]\| \leq 2d_{\mathbf{H}}^s(0, 0g^{-n-c(\gamma)}\gamma g^n).$$

Thus the size of the commutator is determined by the distance $d_{\mathbf{H}}^s(0, 0g^{-n-c(\gamma)}\gamma g^n)$. Using lemma 4.5 we can estimate

$$\begin{aligned} d_{\mathbf{H}}^s(0, 0g^{-n-c(\gamma)}\gamma g^n) &\leq 2d_{\mathbf{H}}^s(0, 0g^n) + d_{\mathbf{H}}^s(0, 0\gamma) + d_{\mathbf{H}}^s(0, 0g^{c(\gamma)}) \\ &\leq 2n^s d_{\mathbf{H}}^s(0, 0g) + d_{\mathbf{H}}^s(0, 0\gamma) + d_{\mathbf{H}}^s(0, 0g^{c(\gamma)}). \end{aligned}$$

Thus, the norm of the operator

$$[1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}, u_\gamma] : e_{g^n} \otimes L^2(\wedge^* \mathbf{H}) \rightarrow e_{g^{n+c(\gamma)}} \otimes L^2(\wedge^* \mathbf{H}),$$

satisfies $\|[1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}, u_\gamma]\|_n \leq C_\gamma + 2n^s$. Since we also have the estimate

$$\|(1 + D_c^2 \otimes 1 + (1 \otimes_{\nabla} \mathcal{D}_{\rho,s})^2)^{-\frac{s}{2}}\|_n \leq (1 + n^2)^{-\frac{s}{2}},$$

we find that

$$\sup_n \|[1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}, u_\gamma](1 + D_c^2 + (1 \otimes_{\nabla} \mathcal{D}_{\rho,s})^2)^{-\frac{s}{2}}\|_n \leq 2d_{\mathbf{H}}^s(0, 0g) + C_\gamma.$$

The operator $(1 + D_c^2 + (1 \otimes_{\nabla} \mathcal{D}_{\rho,s})^2)^{-\frac{s}{2}}[1 \otimes_{\nabla_g} \mathcal{D}_{\rho,s}, u_\gamma]$ is shown to be bounded by noting that $u_\gamma = u_{\gamma^{-1}}^*$. \square

Our next result says that the index pairing between K^1 and K_1 when applied to the K -cycles constructed from group cocycles in Theorem 4.7 and unitaries $[u_\delta] \in K_1(C_r^*(\Gamma))$ recovers the pairing between H_1 and H^1 . This result will play an important rôle in what follows.

Proposition 4.8. *Let $\Gamma \subset \text{Isom}(\mathbf{H})$ be a discrete group and $c : \Gamma \rightarrow \mathbf{Z}$ a normalised cocycle. The index pairing*

$$K_1(C_r^*(\Gamma)) \times K^1(C_r^*(\Gamma)) \rightarrow \mathbf{Z}$$

maps the pair $([u_\delta], [\mathcal{D}_{c,s}])$ to the integer $c(\delta)$, and thus recovers the (co)homology pairing

$$H_1(\Gamma, \mathbf{Z}) \times H^1(\Gamma, \mathbf{Z}) \rightarrow \mathbf{Z}.$$

Proof. We use that the Kasparov product is associative:

$$[u_\delta] \otimes [\mathcal{D}_{c,s}] = [u_\delta] \otimes [\mathcal{D}_{c,s}] \otimes [\gamma_{\mathbf{H}}],$$

and apply Proposition 4.2 and Lemma 4.3 to obtain that this equals

$$c(\delta)[1_{C_r^*(\Gamma_c)}] \otimes [\gamma_{\mathbf{H}}] = c(\delta)\text{Ind}(\mathcal{D}_{\rho,s}^+) = c(\delta).$$

This proves the proposition. \square

4.3. A Hecke equivariant isomorphism. We return to the specific setting of Bianchi groups in dimension 3. We saw in Propositions 1.4 and 1.6 that for a Bianchi group Γ there are isomorphisms

$$K^1(C_r^*(\Gamma)) \simeq H^1(\Gamma, \mathbf{Z}), \quad K_1(C_r^*(\Gamma)) \simeq H_1(\Gamma, \mathbf{Z}).$$

However for our purposes, these isomorphisms are not useful as they are only given abstractly. In this subsection we set out to show that the construction of the previous section gives explicit isomorphisms between the above. We prove their the Hecke equivariance and construct a section for the restriction map $K^1(C(\partial\mathbf{H}_3) \rtimes \Gamma) \rightarrow K^1(C_r^*(\Gamma))$ in the Gysin sequence.

Proposition 4.9. *Let $\Gamma \subset \text{PSL}_2(\mathbf{C})$ be a noncompact torsion-free discrete subgroup. The map*

$$H_1(\Gamma, \mathbf{Z}) \rightarrow K_1(C_r^*(\Gamma)), \quad [\delta] \mapsto [u_\delta],$$

is an isomorphism.

Proof. The quotient manifold $M = \mathbf{H}_3/\Gamma$ is a model for $B\Gamma$. Since $H_3(\Gamma, \mathbf{Z}) = 0$, by [51, Proposition 2.1.ii)] there is an isomorphism $\beta_1^M : H_1(\Gamma, \mathbf{Z}) \rightarrow K_1^{\text{geo}}(M)$, which we compose with the Novikov assembly map $\nu_1^\Gamma : K_1^{\text{geo}}(M) \rightarrow K_1(C_r^*(\Gamma))$. Since the Baum-Connes conjecture holds for Γ , ν_1^Γ is an isomorphism. The composition $\nu_1^\Gamma \circ \beta_1^M$ is shown to coincide with the map $[\delta] \mapsto [u_\delta]$ in [11, Theorem 10.4]. \square

Theorem 4.10. *Let $\Gamma \subset \text{PSL}_2(\mathbf{C})$ be a noncompact torsion-free discrete subgroup. The maps*

$$\begin{array}{ccc} H_1(\Gamma, \mathbf{Z}) & \longrightarrow & K_1(C_r^*(\Gamma)) & \text{and} & H^1(\Gamma, \mathbf{Z}) & \longrightarrow & K^1(C_r^*(\Gamma)) \\ [\delta] & \mapsto & [u_\delta] & & [c] & \mapsto & |c| \cdot [\mathcal{D}_{c,s}] \end{array}$$

are isomorphisms compatible with the pairings of the respective groups.

Proof. Proposition 4.9 gives the K -theory isomorphism. To show that the K -homology map is a homomorphism, we use that $C_r^*(\Gamma)$ is KK -equivalent to $C(\overline{\mathbf{H}}_3) \rtimes \Gamma$ which is in the bootstrap class. By the Universal Coefficient Theorem (UCT) [64, Theorem 1.17, Corollary 1.18] there is a short exact sequence

$$0 \rightarrow \text{Ext}_{\mathbf{Z}}^1(K_0(C_r^*(\Gamma)), \mathbf{Z}) \rightarrow K^1(C_r^*(\Gamma)) \xrightarrow{\otimes} \text{Hom}(K_1(C_r^*(\Gamma)), \mathbf{Z}) \rightarrow 0,$$

where \otimes denotes the map induced by the Kasparov product. By (1.11) $K_0(C_r^*(\Gamma))$ is finitely generated and torsion-free, so the Ext group vanishes and $K^1(C_r^*(\Gamma)) \cong \text{Hom}(K_1(C_r^*(\Gamma)), \mathbf{Z})$. That

is classes in the K -homology $K^1(C_r^*(\Gamma))$ are determined by the index pairing. For an arbitrary cocycle $c : \Gamma \rightarrow \mathbf{Z}$, $\frac{c}{|c|}$ is normalised and $D_{\frac{c}{|c|}} = |c|D_c$ is a scalar multiple of D_c . Thus

$$[D_c] = [D_{\frac{c}{|c|}}] \in KK_1(C_r^*(\Gamma), C_r^*(\Gamma_c)).$$

Theorem 4.8 and the K -theory isomorphism show that the classes $|c|[D_c] + |c'|[D_{c'}]$ and $|c + c'|[D_{c+c'}]$ have the same index pairing and hence are equal, proving that the map $c \mapsto |c|[\mathcal{D}_{c,s}]$ is a homomorphism. Injectivity follows in the same way. For surjectivity, let (H, F) be an odd Fredholm module and p_+ the positive spectral projection of F . Then $c : \gamma \mapsto \text{Ind} p_+ u_\gamma p_+$ is a 1-cocycle on Γ , and $|c|[\mathcal{D}_{c,s}]$ is an unbounded Fredholm module whose index pairing coincides with F . Therefore $[(H, F)] = |c|[\mathcal{D}_{c,s}]$ proving surjectivity. \square

We now show that the explicit isomorphism of abelian groups $H^1(\Gamma, \mathbf{Z}) \xrightarrow{\sim} K^1(C_r^*(\Gamma))$ is Hecke equivariant and construct an explicit section for the restriction map $K^i(C(\partial\mathbf{H}_3) \rtimes \Gamma) \rightarrow K^i(C_r^*(\Gamma))$ in the Gysin sequence.

Proposition 4.11. *Let $[T_g^\Gamma] \in KK_0(C_r^*(\Gamma), C_r^*(\Gamma))$ be the Hecke class from Definition 2.2, $c : \Gamma \rightarrow \mathbf{Z}$ a cocycle and $\delta \in \Gamma$. We have the identities*

$$[u_\delta] \otimes [T_g^\Gamma] = [u_{T_g(\delta)}] \in K_1(C_r^*(\Gamma)), \quad [T_g^\Gamma] \otimes |c|[\mathcal{D}_{c,s}] = |T_g(c)|[\mathcal{D}_{T_g(c),s}] \in K^1(C_r^*(\Gamma)).$$

In particular the isomorphisms $H_1(\Gamma, \mathbf{Z}) \rightarrow K_1(C_r^(\Gamma))$ and $H^1(\Gamma, \mathbf{Z}) \rightarrow K^1(C_r^*(\Gamma))$ are Hecke equivariant.*

Proof. By Proposition 2.4 we have $t_g(u_\delta) = \tau(\delta)\text{diag}(u_{\chi_k(\delta)})$ and since $\tau(\delta) \in M_n(\mathbf{C})$ we have $[\tau(\gamma)] = 0 \in K_1(C_r^*(\Gamma))$. So together with (2.3) we find

$$[T_g(u_\delta)] = [t_g(u_\delta)] = [\tau(\gamma)\text{diag}(\chi_i(\gamma))] = [\text{diag}(u_{\chi_i(\gamma)})] = \sum_{i=1}^d [u_{\chi_i(\gamma)}] = [u_{T_g([\delta])}].$$

Thus Hecke equivariance of the map $[\delta] \mapsto [u_\delta]$ is proved. For K -homology, by the UCT, it suffices to show that for all $\gamma \in \Gamma$ it holds that

$$(T_g^\Gamma \otimes |c|[D_c], u_\gamma) = (|T_g(c)|[D_{T_g(c)}], u_\gamma).$$

By Theorem 4.8, we can compute the right handside to equal $T_g(c)(\gamma)$. For the left handside, observe that the class $T_g^\Gamma \otimes [\mathcal{D}_{c,s}]$ is represented by $(\bigoplus_{i=1}^d E \otimes_{C_r^*(\Gamma_c)} L^2(\wedge^* \mathbf{H}_3), \text{diag}(\mathcal{D}_{c,s}))$. The representation of a unitary u_γ is given by $\alpha_g(u_\gamma)(h_i) = (\chi_i(\gamma)h_{\gamma(i)})$. The positive spectral projection of $\text{diag}(\mathcal{D}_{c,s})$ is $\widetilde{p}_+ = \text{diag}(p_+)$ and thus the index pairing becomes

$$(T_g^\Gamma \otimes |c|[D_c], u_\gamma) = |c| \text{Ind} \widetilde{p}_+ \alpha_g(u_\gamma) \widetilde{p}_+ = |c| \sum_{i=1}^d \text{Ind} p_+ u_{\chi_i(\gamma)} p_+ = \sum_{i=1}^d c(\chi_i(\gamma)) = T_g(c)(\gamma),$$

as required. \square

Unlike the previous results in this section, the following theorem is valid for discrete subgroups $\Gamma \subset \text{Isom} \mathbf{H}$ in any dimension.

Theorem 4.12. *The $(1-s)$ -unbounded Fredholm modules in Theorem 4.7 extend to $(1-s)$ -unbounded Fredholm modules for $C(\partial\mathbf{H}) \rtimes \Gamma$ such that $C(\partial\mathbf{H})$ commutes with $\mathcal{D}_{c,s}$. The extension is compatible with the restriction map $K^1(C(\partial\mathbf{H}) \rtimes \Gamma) \rightarrow K^1(C_r^*(\Gamma))$.*

Proof. Let $X = X_\Gamma \subset \mathbf{H}$ be an open connected fundamental domain for Γ . The disjoint union $\bigcup_{\gamma \in \Gamma} X\gamma$ is dense in \mathbf{H} and

$$\tau : \mathbf{H} \rightarrow \Gamma, \quad \tau(x) = \tau_X(x) = g \Leftrightarrow xg^{-1} \in X,$$

is an almost everywhere defined equivariant measurable map.

The tensor product $E \otimes L^2(\wedge^* \mathbf{H})$ can be identified with the Hilbert space $\bigoplus_{n \in \mathbf{Z}} L^2(\wedge^* \mathbf{H})$ by choosing $g \in g^{-1}(|c|)$ and using Equation (4.1). By choosing a point $\xi \in \partial \mathbf{H}$, representations of $C_r^*(\Gamma)$ and $C(\partial \mathbf{H})$ are defined, for $\psi = (\psi_n)_{n \in \mathbf{Z}}$, by

$$u_\delta(\psi)_n(h) = \psi_{n-c(\delta)}(hg^{-n}\delta g^{n-c(\delta)}), \quad (\pi_{X,\xi}(f)\psi)_n(h) := f(\xi\tau(h)g^{-n})\psi_n(h),$$

and form a covariant pair. Thus we obtain a representation of $C(\partial \mathbf{H}) \rtimes \Gamma$ on $E \otimes_{C_r^*(\Gamma_c)} L^2(\wedge^* \mathbf{H})$.

The representation $\pi_{X,\xi}$ clearly commutes with the multiplication operators ρ and c . Because $\pi_{X,\xi}(f)$ is constant on each $X\gamma$, it also commutes with the Dirac operator \not{D} . Therefore the $(1-s)$ spectral triples from Theorem 4.7 extend to $C(\partial \mathbf{H}) \rtimes \Gamma$. Since $\partial \mathbf{H}$ is connected, the choice of $\xi \in \partial \mathbf{H}$ does not affect the homotopy class of the spectral triple. If Y is another open connected fundamental domain for Γ , there exists $\delta \in \text{Isom}(\mathbf{H})$ such that $X = Y\delta$ and thus $\tau_X(x) = \delta\tau_Y(x)$. This implies that $\pi_{Y,\xi} = \pi_{X,\xi\delta}$. Therefore the representations $\pi_{X,\xi}$ and $\pi_{Y,\xi}$ are homotopic as well. \square

5. K -cycles for Bianchi manifolds

Let Γ be a torsion-free finite-index subgroup of a Bianchi group and M be the associated hyperbolic 3-manifold. We already know from Propositions 1.5 and 1.6 that there is an abstract isomorphism

$$K^0(C_0(M)) \simeq H^1(\Gamma, \mathbf{Z}).$$

In this section, we shall construct an explicit Hecke equivariant isomorphism

$$K^0(C_0(M)) \simeq H_2(\overline{M}, \partial \overline{M}, \mathbf{Z}),$$

where \overline{M} is the Borel-Serre compactification of M (see Section 2.1.2). Recall that

$$H_2(\overline{M}, \partial \overline{M}, \mathbf{Z}) \cong H^1(\overline{M}, \mathbf{Z}) \cong H^1(\Gamma, \mathbf{Z})$$

and these isomorphisms are Hecke equivariant. Our approach uses geometric K -homology and employ work of Matthey [51] on geometric K -homology of low-dimensional CW -complexes.

5.1. Complex spin structures. Spin structures on M are in bijection with lifts of the holonomy representation $\Gamma \hookrightarrow \text{PSL}_2(\mathbf{C})$ to $\text{SL}_2(\mathbf{C}) \simeq \text{Spin}(3, 1)$ (see, e.g. [58, Section 2.7]). It is known that such lifts exist and thus M admits a spin structure. Let us fix a lift of the holonomy map of M and denote the corresponding spin structure on M by σ . It is well known that any compact oriented 3-manifold admits a spin structure (see [41, Section IV]), in particular, the Borel-Serre compactification \overline{M} of M admits a spin structure. It turns out that, see [41, Proposition 1, Section IV], we can choose a spin structure on \overline{M} so that the induced spin structure on M agrees with our fixed σ . We fix such a spin structure δ on \overline{M} .

A spin structure induces a complex spin (or spin^c) structure, in a canonical way. We denote the corresponding spin^c -structures on M and \overline{M} with the same symbols σ and δ respectively. This will not cause confusion as we shall only consider spin^c structures. In the rest of the paper, we will endow all codimension 0 and codimension 1 submanifolds of \overline{M} with the canonical spin^c structure arising from δ .

5.2. Geometric K -homology. Let us describe $K^0(C_0(M))$ as a relative group in the Baum-Douglas model for K -homology of manifolds [6]. For a CW-pair (X, Y) , a *geometric cycle* is a triple (N, E, φ) consisting of a compact spin^c manifold N with boundary ∂N , a vector bundle $E \rightarrow N$ and a continuous map $\varphi : N \rightarrow X$ such that $\varphi(\partial N) \subset Y$. The parity $*$ = 0, 1 corresponds to the dimension of N being even or odd. Modulo a suitable equivalence relation, such cycles generate the *geometric K -homology* $K_*^{\text{geo}}(X, Y)$ of the pair (X, Y) . By taking $Y = \emptyset$, we obtain the geometric K -homology group $K_*^{\text{geo}}(X) := K_*^{\text{geo}}(X, \emptyset)$. For details see [7, 33].

The paper [51] describes explicit relationships between ordinary homology and geometric K -homology of low dimensional CW-complexes. Recall the Hurewicz homomorphism $h : \pi_1(M) \rightarrow H_1(M, \mathbf{Z})$, which sends the class of a map $\varphi : S^1 \rightarrow M$ to $\varphi_*([S^1])$, where $[S^1] \in H_1(S^1, \mathbf{Z}) \simeq \mathbf{Z}$ is the fundamental class. By a slight abuse of notation, we denote $h([\varphi]) \in H_1(M, \mathbf{Z})$ by $[\varphi]$. By surjectivity of h , the group $H_1(M, \mathbf{Z})$ is exhausted by the elements $[\varphi]$. Similarly, any nontrivial class $z \in H_2(M, \mathbf{Z})$ can be represented by an embedded surface, that is, there is a compact oriented surface N and an embedding $\varphi : N \rightarrow M$ such that $\varphi_*([N]) = z$ where $[N] \in H^2(N, \mathbf{Z}) \simeq \mathbf{Z}$ is the fundamental class, [72, Corollaire III.7.]. We shall denote $\varphi_*([N])$ by $[N, \varphi]$.

Proposition 5.1 ([51]). *Let X be a connected CW-complex such that $H_k(X, \mathbf{Z}) = 0$ for all $k \geq 3$. There are explicit natural isomorphisms*

$$\begin{aligned} \beta_{\text{odd}} : H_1(X, \mathbf{Z}) &\rightarrow K_1^{\text{geo}}(X), & [\varphi] &\mapsto [S^1, 1_{S^1}, \varphi] \\ \beta_{\text{ev}} : H_0(X, \mathbf{Z}) \oplus H_2(X, \mathbf{Z}) &\rightarrow K_0^{\text{geo}}(X), & ([\text{pt}], [N, \varphi]) &\mapsto [\text{pt}, 1_{\text{pt}}, i] + [N, 1_N, \varphi], \end{aligned}$$

where $[\varphi] \in H_1(X, \mathbf{Z})$ and $[N, \varphi] \in H_2(X, \mathbf{Z})$ are as above, and $i : \text{pt} \rightarrow X$ is any choice of inclusion.

Proof. This result follows by Theorem 2.1 and Propositions 3.2, 3.3, 3.4 and 3.6 in [51]. \square

Given a geometric cycle (N, E, φ) for a CW-pair (X, Y) , let $S_N \rightarrow N$ be the spinor bundle and D_E the associated symmetric Dirac operator on the bundle $E \otimes S$. The restriction of φ to $N \setminus \varphi^{-1}(Y)$ gives a continuous map $\varphi : N \setminus \varphi^{-1}(Y) \rightarrow X \setminus Y$, which by the Tietze extension theorem gives a *-homomorphism $C_0(X \setminus Y) \rightarrow C_0(\overset{\circ}{N})$. Here $\overset{\circ}{N} = N \setminus \partial N \subset N$ denotes the interior of N . We so obtain a representation $C_0(X \setminus Y) \rightarrow B(L^2(\overset{\circ}{N}, S))$. The symmetric operator D_E then defines a K -homology class by [31, Theorem 3.2]. The relation between geometric and analytic K -homology is given by the following result.

Lemma 5.2. *Let \overline{M} denote a topological compactification of M and $\partial \overline{M} := \overline{M} \setminus M$. If $(\overline{M}, \partial \overline{M})$ is a CW-pair, then the map*

$$K_0^{\text{geo}}(\overline{M}, \partial \overline{M}) \xrightarrow{\sim} K^0(C_0(M)) \quad (N, E, \varphi) \mapsto (C_0(M), L^2(\overset{\circ}{N}, E \otimes S), D_E)$$

is a natural isomorphism.

Proof. This is the statement of [7, Theorem 6.2]. \square

In view of the last lemma, we consider the Borel-Serre compactification \overline{M} of M , see Section 2.1.2. The pair $(\overline{M}, \partial \overline{M})$ form a CW-pair. In view of Proposition 5.1 and Lemma 5.2, we aim to construct a relative version of the map β_{ev} . We begin with a relative version of Steenrod representability for H_2 , which can be found in [50, Proposition 1.7.16] (see also [37, Lemma 2.9] and the remark after its proof).

Lemma 5.3. *Any nontrivial class $z \in H_2(\overline{M}, \partial\overline{M}, \mathbf{Z})$ can be represented by a properly embedded surface, that is, there is a compact oriented surface N and an embedding $\varphi : N \rightarrow \overline{M}$ such that $\varphi(\partial N) = \varphi(N) \cap \partial\overline{M}$ and $\varphi_*([N]) = z$ where $[N] \in H_2(N, \partial N, \mathbf{Z}) \simeq \mathbf{Z}$ is the fundamental class. Moreover N can be chosen so that all its components have negative Euler characteristic.*

As before, we denote $\varphi_*([N])$ by $[N, \varphi]$. For convenience we will write $(N, \partial N) \subset (\overline{M}, \partial\overline{M})$ to mean that N is a compact surface with boundary that is properly embedded into \overline{M} as in Lemma 5.3.

As $N \subset \overline{M}$ is an embedded hypersurface, the spin structure on \overline{M} descends to N and $(N, 1_N, \varphi)$ is a geometric K -cycle for $(\overline{M}, \partial\overline{M})$. We now show that these cycles exhaust the group $K_0^{\text{geo}}(\overline{M}, \partial\overline{M})$.

Proposition 5.4. *There is a natural isomorphism*

$$(5.1) \quad \beta_2^{\text{rel}} : H_2(\overline{M}, \partial\overline{M}, \mathbf{Z}) \xrightarrow{\simeq} K_0^{\text{geo}}(\overline{M}, \partial\overline{M}), \quad [N, \varphi] \mapsto [N, 1_N, \varphi],$$

where \overline{M} is the Borel-Serre compactification of M and $\varphi : (N, \partial N) \rightarrow (\overline{M}, \partial\overline{M})$ is an embedding.

Proof. The notation β_2^{rel} is in accordance with [51] and Proposition 5.1, as the maps $\beta_{\text{ev}} = \beta_0 \oplus \beta_2$ and $\beta_{\text{odd}} = \beta_1$.

To show that β_2^{rel} is well-defined, let $[N_1, \varphi_1], [N_2, \varphi_2]$ represent the same homology class. Consider the oriented bordism group $\Omega_2^{SO}(\overline{M}, \partial\overline{M})$ (see [16, Section 4]). Noting that $\Omega_0^{SO} \simeq \mathbf{Z}$ and $\Omega_1^{SO} = \Omega_2^{SO} = 0$, and that $H_2(\overline{M}, \partial\overline{M}, \mathbf{Z})$ is finitely generated and is torsion-free (as it is isomorphic to $H^1(\Gamma, \mathbf{Z}) = \text{Hom}(\Gamma, \mathbf{Z})$), we conclude by Theorem 15.2 of [16] that the representation map $\mu : \Omega_2(\overline{M}, \partial\overline{M}) \rightarrow H_2(\overline{M}, \partial\overline{M}, \mathbf{Z})$ is an isomorphism. This implies that $[N_1, \varphi_1]$ and $[N_2, \varphi_2]$ are bordant in $(\overline{M}, \partial\overline{M})$. As we consider codimension 0 and codimension 1 submanifolds of \overline{M} with the spin^c structure inherited from that of \overline{M} , it follows immediately that the cycles $[N_1, 1_{N_1}, \varphi_1]$ and $[N_2, 1_{N_2}, \varphi_2]$ are spin^c -bordant and thus represent the same geometric K -homology class. As the addition operation on both groups is given by disjoint union, it is now clear that we have a homomorphism $H_2(\overline{M}, \partial\overline{M}, \mathbf{Z}) \rightarrow K_0^{\text{geo}}(\overline{M}, \partial\overline{M})$.

As group operations on both sides amount to taking disjoint unions of manifolds representing classes, it is clear that β_2 is a homomorphism.

To show that the map β_2^{rel} is an isomorphism, recall the long exact sequence in homology associated to the pair $(\overline{M}, \partial\overline{M})$ which takes the form

$$\begin{array}{ccccccccccc} 0 & \longleftarrow & H_0(\overline{M}) & \longleftarrow & H_0(\partial\overline{M}) & \longleftarrow & H_1(\overline{M}, \partial\overline{M}) & \longleftarrow & H_1(\overline{M}) & \longleftarrow & H_1(\partial\overline{M}) \\ & & & & & & & & & & \uparrow \\ 0 & \longrightarrow & H_3(\overline{M}, \partial\overline{M}) & \longrightarrow & H_2(\partial\overline{M}) & \longrightarrow & H_2(\overline{M}) & \longrightarrow & H_2(\overline{M}, \partial\overline{M}) & & \end{array}$$

due to the facts that $H^0(\overline{M}, \partial\overline{M}) \simeq 0 \simeq H_3(\partial\overline{M})$. Next, consider the six-term exact sequence of geometric K -homology groups (see for instance [7, 33]):

$$\begin{array}{ccccc} K_0^{\text{geo}}(\partial\overline{M}) & \longrightarrow & K_0^{\text{geo}}(\overline{M}) & \longrightarrow & K_0^{\text{geo}}(\overline{M}, \partial\overline{M}) \\ & & \uparrow \partial & & \downarrow \partial \\ K_1^{\text{geo}}(\overline{M}, \partial\overline{M}) & \longleftarrow & K_1^{\text{geo}}(\overline{M}) & \longleftarrow & K_1^{\text{geo}}(\partial\overline{M}) \end{array}$$

Writing $H_{\text{ev}}(X) = H_0(X) \oplus H_2(X)$, and $\iota : (\overline{M}, \emptyset) \rightarrow (\overline{M}, \partial\overline{M})$ for the inclusion of CW-pairs, Proposition 5.1 yields a diagram with exact rows

$$\begin{array}{ccccccccc} H_{\text{ev}}(\partial\overline{M}) & \longrightarrow & H_{\text{ev}}(\overline{M}) & \xrightarrow{\iota_*} & H_2(\overline{M}, \partial\overline{M}) & \xrightarrow{\partial} & H_1(\partial\overline{M}) & \longrightarrow & H_1(\overline{M}) \\ \beta_{\text{ev}} \downarrow & & \beta_{\text{ev}} \downarrow & & \beta_2^{\text{rel}} \downarrow & & \beta_{\text{odd}} \downarrow & & \beta_{\text{odd}} \downarrow \\ K_0^{\text{geo}}(\partial\overline{M}) & \longrightarrow & K_0^{\text{geo}}(\overline{M}) & \xrightarrow{\iota_*} & K_0^{\text{geo}}(\overline{M}, \partial\overline{M}) & \xrightarrow{\partial} & K_1^{\text{geo}}(\partial\overline{M}) & \longrightarrow & K_1^{\text{geo}}(\overline{M}), \end{array}$$

whose outer squares commute. If we show that the inner squares commute as well, then the five Lemma and the fact that $\beta_{\text{ev}}, \beta_{\text{odd}}$ are isomorphisms, implies that β_2^{rel} is an isomorphism as well.

To show that $\beta_2^{\text{rel}} \circ \iota_* = \iota_* \circ \beta_{\text{ev}}$, observe that the $H_0(\overline{M})$ summand of $H_{\text{ev}}(\overline{M})$ is annihilated by ι_* , as is the class of a point in $K_0^{\text{geo}}(\overline{M})$. For a surface class $[(N, \varphi)] \in H_2(\overline{M})$ we find that

$$\beta_2^{\text{rel}} \circ \iota_* [(N, \varphi)] = \beta_2^{\text{rel}} [(N, \varphi)] = [(N, 1_N, \varphi)] = \iota_* [(N, 1_N, \varphi)] = \iota_* \circ \beta_{\text{ev}} [(N, \varphi)],$$

as desired. We now prove that $\beta_{\text{odd}} \circ \partial = \partial \circ \beta_2^{\text{rel}}$. By Lemma 5.3 all classes in $H_2(\overline{M}, \partial\overline{M})$ are of the form $[(N, \varphi)]$. The boundary ∂N is a compact 1-dimensional manifold, and therefore decomposes as a disjoint union $\partial N = \bigsqcup_{i=1}^k S^1$ of circles S^1 . Denote by φ_i the restriction of φ to the i -th circle in this decomposition. We compute the composition

$$\begin{aligned} \beta_{\text{odd}} \circ \partial [(N, \varphi)] &= \beta_{\text{odd}} [(\partial N, \varphi|_{\partial N})] = \sum_{i=1}^k \beta_{\text{odd}} [(S^1, \varphi_i)] \\ &= \sum_{i=1}^k [(S^1, 1_{S^1}, \varphi_i)] = [(\partial N, 1_N, \varphi)] = \partial [(N, 1_N, \varphi)] = \partial \circ \beta_2^{\text{rel}} [(N, \varphi)]. \end{aligned}$$

This completes the proof that β_2^{rel} is an isomorphism. \square

Note that Lemma 5.3 implies that $\mathring{N} = N \cap M \subset M$ is a closed embedded hypersurface. We equip \mathring{N} with the metric inherited from the hyperbolic metric on M as well as with the inherited spin^c structure. The Riemannian distances $d_{\mathring{N}}, d_M$ satisfy $d_M(x, y) \leq d_{\mathring{N}}(x, y)$ for $x, y \in \mathring{N}$. Since \mathring{N} carries the relative topology as a subset of M and M is complete, it follows that \mathring{N} is complete. The spinor bundle $\mathcal{S}_{\mathring{N}} \rightarrow \mathring{N}$ is the restriction of the spinor bundle $\mathcal{S}_M \rightarrow M$ to \mathring{N} (see [5, 32]). Thus \mathring{N} is a complete Riemannian spin^c manifold and we denote by $\mathcal{D}_{\mathring{N}}$ its Dirac operator, which is essentially self-adjoint on $C_c^1(\mathring{N}, \mathcal{S}_{\mathring{N}})$, the compactly supported C^1 -sections.

Theorem 5.5. *Let $\Gamma \subset \text{PSL}_2(\mathbb{C})$ be a noncocompact torsion-free discrete subgroup. There is a natural isomorphism*

$$(5.2) \quad H_2(\overline{M}, \partial\overline{M}) \xrightarrow{\sim} K^0(C_0(M)), \quad [(N, \varphi)] \mapsto (C_0(M), \varphi L^2(\mathring{N}, \mathcal{S}_{\mathring{N}}), \mathcal{D}_{\mathring{N}}),$$

where \hat{N} is viewed as a spin^c surface with associated Dirac operator $\hat{D}_{\hat{N}}$.

Proof. By Lemma 5.2, we obtain a map $(N, 1_N, \varphi) \rightarrow (C_0(M), L^2(\hat{N}, S), D_{\hat{N}})$ where $D_{\hat{N}}$ is the symmetric operator obtained from the manifold with boundary N . Since we have chosen the spin structure on \overline{M} to be compatible with that on M , the spin structure that N inherits from \overline{M} is compatible with the spin structure that \hat{N} inherits from M . By [30, Proposition 11.27] it follows that

$$[(C_0(M), L^2(\hat{N}, S), D_{\hat{N}})] = [(C_0(M), L^2(\hat{N}, \mathcal{S}|_{\hat{N}}), \hat{D}_{\hat{N}})] \in K^0(C_0(M)).$$

Combining Lemmas 5.2, 5.3 and Proposition 5.4, it thus follows that the map (5.2) is an isomorphism. \square

5.3. Hecke equivariance. Given a class $[N, \varphi] \in H_2(\overline{M}, \partial\overline{M}, \mathbf{Z})$ and a Hecke operator T_g , it can be seen that the class $T_g([N, \varphi])$ is represented by

$$(5.3) \quad T_g([N, \varphi]) = [(\pi_g^{-1}(N), \tau_g)] = [(N_g, \tau_g)]$$

where π_g, τ_g are as in Section 2.1.2 and N_g is a compact surface with boundary $\partial N_g \subset \partial\overline{M}_g$ given by the fiber product

$$(5.4) \quad N_g := (\overline{M}_g)_{\pi_g} \times_{\varphi} N \simeq \pi_g^{-1}(N) \subset \overline{M}_g.$$

The reader should compare this with the discussion in [22, Section 3].

Proposition 5.6. *The isomorphism $H_2(\overline{M}, \partial\overline{M}, \mathbf{Z}) \rightarrow K^0(C_0(M))$ (cf. (5.2)) is Hecke equivariant.*

Proof. Take a class $[N, \varphi] \in H_2(\overline{M}, \partial\overline{M}, \mathbf{Z})$. Comparing 5.3 and the isomorphism (5.2), we see that we need to show that

$$[(C_0(M), \tau_g, L^2(\hat{N}_g, \mathcal{S}_{\hat{N}_g}), \hat{D}_{N_g})] = [T_g^M] \otimes [(C_0(M), \varphi, L^2(\hat{N}, \mathcal{S}_{\hat{N}}), \hat{D}_{\hat{N}})],$$

in the group $KK_0(C_0(M), C_0(M))$. Viewing \hat{N}_g as the inverse image $\pi_g^{-1}(\hat{N}) \subset M_g$ using (5.4) it is straightforward to show that

$$w : T_g^M \otimes_{C_0(M)} L^2(\hat{N}, \mathcal{S}_{\hat{N}}) \rightarrow_{\tau_g} L^2(\hat{N}_g, \mathcal{S}_{\hat{N}_g}), \quad w(\chi \otimes \psi)(n) = \chi(n)\pi_g^*\psi(n),$$

is a unitary isomorphism intertwining the left $C_0(M)$ -representations. To prove that $\hat{D}_{\hat{N}_g}$ represents the Kasparov product we need to check conditions i-iii in [42, Theorem 13], of which ii and iii are trivial since the module T_g^M carries the 0 operator. Now suppose that $\chi \in C_c^1(\hat{M}_g)$ is such that $\text{supp } \chi \subset U$, with U an open set such that $\pi_g|_U$ is injective. Then we can choose $\zeta \in C_c^1(\hat{M})$ with $\chi = (\pi_g^*\zeta)|_U$. Then for $\psi \in L^2(\hat{N}, \mathcal{S}_{\hat{N}})$ we have $\hat{D}_{\hat{N}_g}\chi\pi_g^*\psi = \hat{D}_{\hat{N}_g}\pi_g^*(\zeta\psi)|_U = \pi_g^*(\hat{D}_{\hat{N}}\zeta\psi)|_U$. Thus we find

$$\hat{D}_{\hat{N}_g}\chi\pi_g^*\psi - \chi\pi_g^*\hat{D}_{\hat{N}}\psi = \pi_g^*(\hat{D}_{\hat{N}}\zeta\psi - \zeta\hat{D}_{\hat{N}}\psi)|_U = \pi_g^*(c(\mathbf{d}_{\hat{N}}(\zeta|_{\hat{N}}))\psi)|_U$$

where c denotes Clifford multiplication of forms. Since

$$\|\pi_g^*(c(\mathbf{d}_{\hat{N}}(\zeta|_{\hat{N}}))\psi)|_U\|_{L^2(\hat{N}_g, \mathcal{S}_{\hat{N}_g})} = \|c(\mathbf{d}_{\hat{N}}(\zeta|_{\hat{N}}))\psi\|_{L^2(\hat{N}, \mathcal{S})} \leq \|c(\mathbf{d}_{\hat{N}}(\zeta|_{\hat{N}}))\| \|\psi\|_{L^2(\hat{N}, \mathcal{S})}$$

it follows that $\psi \mapsto \hat{D}_{\hat{N}_g}\chi\pi_g^*\psi - \chi\pi_g^*\hat{D}_{\hat{N}}\psi$ extends to a bounded operator. The submodule of T_g^M generated by elements $\chi \in C_c^1(\hat{M}_g)$ of small support is dense in T_g^M . Hence condition i of [42, Theorem 13] is satisfied and we are done. \square

6. The case of $\mathrm{PSL}_2(\mathbf{Z})$

Let Γ be a torsion-free finite index subgroup of $\mathrm{PSL}_2(\mathbf{Z})$. Then it acts properly discontinuously on the hyperbolic plane \mathbf{H}_2 and the quotient $M = \mathbf{H}_2/\Gamma$ is a finite volume hyperbolic surface with cusps. The boundary of \mathbf{H}_2 can be identified with $\mathbb{P}^1(\mathbf{R})$.

The analogue of Proposition 1.3 in this case is the following (note that the cohomological dimension of Γ is one). For $i = 0, 1$, we have

$$K_i(C(\mathbb{P}^1(\mathbf{R})) \rtimes \Gamma) \simeq H_0(\Gamma, \mathbf{Z}) \oplus H_1(\Gamma, \mathbf{Z})$$

and

$$K^i(C(\mathbb{P}^1(\mathbf{R})) \rtimes \Gamma) \simeq H^0(\Gamma, \mathbf{Z}) \oplus H^1(\Gamma, \mathbf{Z}).$$

This is actually well known, it is a special case of the work of Anantharaman-Delaroche (see [2]) who treated cofinite discrete subgroups of $\mathrm{PSL}_2(\mathbf{R})$. Note that $H^0(\Gamma, \mathbf{Z}) \simeq \mathbf{Z}$ and $H^1(\Gamma, \mathbf{Z}) \simeq \mathbf{Z}^{2g+c-1}$ where g is the genus of Γ and $c \geq 1$ is the number of cusps of M .

In [48], Manin and Marcolli describe the above isomorphisms in terms of Manin symbols using Pimsner's 6-term exact sequence [60] of which Kasparov's spectral sequence can be viewed as a generalization.

Much of Section 3 carries through and we obtain the Hecke equivariant exact hexagon

$$(6.1) \quad \begin{array}{ccccc} K^1(C_0(M)) & \longrightarrow & K^0(C(\mathbb{P}^1(\mathbf{R})) \rtimes \Gamma) & \longrightarrow & K^0(C_r^*(\Gamma)) \\ & & \uparrow & & \downarrow \\ & & K^1(C_r^*(\Gamma)) & \longleftarrow & K^1(C(\mathbb{P}^1(\mathbf{R})) \rtimes \Gamma) & \longleftarrow & K^0(C_0(M)) \end{array}$$

As M is non-compact, this hexagon (6.1) breaks apart into two short exact sequences (see [23])

$$(6.2) \quad 0 \rightarrow K^1(C_0(M)) \rightarrow K^0(C(\mathbb{P}^1(\mathbf{R})) \rtimes \Gamma) \rightarrow K^0(C_r^*(\Gamma)) \rightarrow 0,$$

$$(6.3) \quad 0 \rightarrow K^0(C_0(M)) \rightarrow K^1(C(\mathbb{P}^1(\mathbf{R})) \rtimes \Gamma) \rightarrow K^1(C_r^*(\Gamma)) \rightarrow 0$$

as in the Bianchi case.

It is well-known that Γ is a free group on $2g + c - 1$ generators. It follows, for example, from work of Cuntz [21] and of Lance [43], that $K^0(C_r^*(\Gamma)) \simeq \mathbf{Z}$ and $K^1(C_r^*(\Gamma)) \simeq \mathbf{Z}^{2g+c-1}$. As a result, the sequences (6.2) and (6.3) split and also we get $K^1(C_0(M)) \simeq \mathbf{Z}^{2g+c-1}$ and $K^0(C_0(M)) \simeq \mathbf{Z}$.

The analogue of Proposition 1.4 reads as follows. For $i = 0, 1$, we have

$$K^i(C_r^*(\Gamma)) \simeq H^i(\Gamma, \mathbf{Z}),$$

$$K_i(C_r^*(\Gamma)) \simeq H_i(\Gamma, \mathbf{Z}).$$

The map we constructed in Section 4 is defined here as well and we get

Theorem 6.1. *The maps*

$$\begin{array}{ccc} H_1(\Gamma, \mathbf{Z}) & \longrightarrow & K_1(C_r^*(\Gamma)) & \text{and} & H^1(\Gamma, \mathbf{Z}) & \longrightarrow & K^1(C_r^*(\Gamma)) \\ [\delta] & \mapsto & [u_\delta] & & [c] & \mapsto & |c| \cdot [D_{\frac{c}{|c|}, \mathbf{H}}] \end{array}$$

are Hecke equivariant isomorphisms compatible with the pairings of the respective groups.

Our results in Section 5 adapt straightforwardly to the case of $\mathrm{PSL}_2(\mathbf{Z})$. The 1-dimensional analogue of Lemma 5.3 holds and we get the following (using the notation of Section 5):

Theorem 6.2. *There is a Hecke-equivariant isomorphism*

$$H_1(\overline{M}, \partial\overline{M}, \mathbf{Z}) \rightarrow K^1(C_0(M))$$

sending the homology class $[N, \varphi]$ to the class $[(C_0(M), L^2(\overset{\circ}{N}, E \otimes S), D_E)]$ where \overline{M} denotes the Borel-Serre compactification of M .

Note that $H_1(\overline{M}, \partial\overline{M}, \mathbf{Z}) \simeq H^1(\Gamma, \mathbf{Z})$ as Hecke modules.

7. The extension class as a hypersingular integral operator

In section 3.2 we used the harmonic measures ν_x on the boundary $\partial\mathbf{H}$ to represent the boundary extension (3.1) as a Kasparov module. In order to compute the K -homology boundary map $\partial : K^0(C_0(M)) \rightarrow K^1(C(\partial\mathbf{H}) \rtimes \Gamma)$, we now construct an unbounded Kasparov module [4] representing the extension class.

7.1. Harmonic calculus on $T_1\mathbf{H}$. As before, \mathbf{H} denotes the Poincaré ball model of hyperbolic $n + 1$ space. To construct an unbounded representative for the boundary extension we discuss hypersingular integral operators defined using the harmonic measures ν_x and a family of metrics d_x on $\partial\mathbf{H}$ which we now describe. For all $x, y \in \overline{\mathbf{H}}$ and $g \in G$ we have (cf. [56, Equation 1.3.2]):

$$(7.1) \quad \|xg - yg\| = |g'(x)|^{\frac{1}{2}} |g'(y)|^{\frac{1}{2}} \|x - y\|,$$

where $\|\cdot\|$ denotes the Euclidean norm on \mathbf{R}^{n+1} . Using the Poisson kernel (3.8) the function

$$(7.2) \quad d_x(\xi, \eta) := P(x, \xi)^{1/2} P(x, \eta)^{1/2} \|\xi - \eta\|,$$

satisfies $d_{xg}(\xi g, \eta g) = d_x(\xi, \eta)$ by (3.9) and so d_x is a metric on $\partial\mathbf{H}$, as this holds for d_0 and g acts transitively on \mathbf{H} (compare [56, Lemma 3.4.2] and [61, Section 3.3]).

For a pair $(x, \xi) \in T_1\mathbf{H}$ we denote by $r_{(x, \xi)} : \mathbf{R} \rightarrow \mathbf{H}$ the geodesic ray with $r_{(x, \xi)}(0) = x$ and $\lim_{t \rightarrow +\infty} r_{(x, \xi)}(t) = \xi$. Recall that for $r > 0$ and $x, y \in \mathbf{H}$, the *Sullivan shadow* is the set

$$\mathcal{O}_r(x, y) := \{\xi \in \partial\mathbf{H} : \inf\{d_{\mathbf{H}}(r_{x, \xi}(t), y) : t \in [0, \infty)\} \leq r\} \subset \partial\mathbf{H}.$$

A proof of the following result can be found in [56, 61].

Proposition 7.1 (Sullivan's shadow lemma [71]). *For all $\xi, \eta \in \mathcal{O}_r(x, y)$ it holds that*

$$e^{-r} e^{d(x, y)} \leq d_x(\xi, \eta) \leq e^{d(x, y)}.$$

Moreover there exists r_0 such that for all $r \geq r_0$ there exists $C_r > 0$ for which

$$C_r^{-1} e^{-nd(x, y)} \leq \nu_x(\mathcal{O}_r(x, y)) \leq C_r e^{-nd(x, y)},$$

for all $x, y \in \mathbf{H}$.

We start with the following observation, concerning the Riesz potentials commonly studied in metric measure theory (see for instance [74]).

Lemma 7.2. *For all $0 < s < n$, the integral*

$$I_s = I_s(x, \xi) := \int \frac{1}{d_x(\xi, \eta)^{n-s}} d\nu_x \eta,$$

is finite and independent of (x, ξ) .

Proof. To see that the integral is finite for fixed (x, ξ) , we only consider the case $0 < s < n$, as the case $s \geq n$ is immediate. Choose points x_k on the geodesic from x to ξ such that $d(x, x_k) = k$. We write $\partial\mathbf{H}$ as a disjoint union

$$\partial\mathbf{H} = \bigcup_{k=0}^{\infty} \mathcal{O}_r(x, x_k) \setminus \mathcal{O}_r(x, x_{k+1}),$$

and expand the integral and then estimate using Proposition 7.1:

$$\begin{aligned} \int \frac{1}{(d_x(\xi, \eta))^{n-s}} d\nu_x \eta &= \sum_{k=0}^{\infty} \int_{\mathcal{O}_r(x, x_k) \setminus \mathcal{O}_r(x, x_{k+1})} \frac{1}{d_x(\xi, \eta)^{n-s}} d\nu_x \eta \leq C_r \sum_{k=0}^{\infty} \frac{\nu_x(\mathcal{O}_r(x, x_k))}{(e^{-d(x, x_{k+1})})^{n-s}} \\ &\leq C_r \sum_{k=0}^{\infty} e^{(n-s)(k+1)} e^{-nk} = C_r e^{n-s} \sum_{k=0}^{\infty} e^{-sk} < \infty. \end{aligned}$$

Thus the integral $I_s \Psi(x, \xi)$ is finite for fixed (x, ξ) . For $g \in G_x$ we have

$$\int \frac{1}{(d_x(\xi, \eta))^{n-s}} d\nu_x(\eta) = \int \frac{1}{(d_x(\xi g, \eta g))^{n-s}} d\nu_x(\eta g) = \int \frac{1}{(d_x(\xi g, \eta))^{n-s}} d\nu_x(\eta),$$

and since G_x acts transitively on $\partial\mathbf{H}$, we see that $I_s(x, \xi)$ is constant in ξ . Using this fact, we can write for $g \in G$:

$$\int \frac{1}{(d_x(\xi, \eta))^{n-s}} d\nu_x(\eta) = \int \frac{1}{d_{xg}(\xi g, \eta g)^{n-s}} d\nu_{xg}(\eta g) = \int \frac{1}{(d_{xg}(\xi, \eta))^{n-s}} d\nu_{xg}(\eta),$$

and since G acts transitively on \mathbf{H} we see that $I_s(x, \xi)$ is independent of x as well. \square

Now we consider the spherical hypersingular operator $\Delta_0 : \text{Lip}(\partial\mathbf{H}) \rightarrow L^2(\partial\mathbf{H}, \nu_0)$ and the projection $p_0 \in \mathbb{B}(L^2(\partial\mathbf{H}, \nu_0))$ given by

$$(7.3) \quad \Delta_0 f(\xi) = \int \frac{f(\xi) - f(\eta)}{d_0(\xi, \eta)^n} d\nu_0 \eta, \quad p_0 f(\xi) := \int f(\eta) d\nu_0 \eta.$$

Operators of exponent $n+\varepsilon$ in the denominator have been extensively studied by Samko [65, 66].

Lemma 7.3. *The operator Δ_0 maps $\text{Lip}(\partial\mathbf{H}) \subset L^2(\partial\mathbf{H}, \nu_0)$ into bounded functions on $\partial\mathbf{H}$, is essentially self-adjoint on $C^1(\partial\mathbf{H}) \subset \text{Lip}(\partial\mathbf{H})$ and has compact resolvent. Moreover $\ker \Delta_0 = \text{Im } p_0$ and $\Delta_0 + p_0$ is strictly positive.*

Proof. Observe that for $f \in \text{Lip}(\partial\mathbf{H})$, by Hölder's inequality and Lemma 7.2 we have

$$\left| \int \frac{f(\xi) - f(\eta)}{d_0(\xi, \eta)^n} d\nu_0 \eta \right| \leq \int \frac{|f(\xi) - f(\eta)|}{d_0(\xi, \eta)^n} d\nu_0 \eta \leq \|f\|_{\text{Lip}} \int \frac{1}{d_0(\xi, \eta)^{n-1}} d\nu_0 \eta = I_1 \|f\|_{\text{Lip}},$$

so $\Delta_0 f$ is a bounded function and thus $\Delta_0 f \in L^2(\partial\mathbf{H}, \nu_0)$. In particular, any orthonormal family of spherical harmonics $Y_{m,k} \in C^1(\partial\mathbf{H})$ is in the domain of Δ_0 . Using the the method of Samko [66, Lemma 6.25] we see that the multiplier of Δ_0 on spherical harmonics is given by

$$Y_{m,k} \mapsto \lambda_m Y_{m,k}, \quad \lambda_m = \int_{-1}^1 (1+t)^{\frac{n-3}{2}} (1-t)^{-1} (1 - P_m(t)) dt.$$

Here $P_m(t) = \sum_{k=0}^m \left(\frac{-1}{2}\right)^k \binom{m}{k} \binom{m+k}{k} (1-t)^k$ is the m -th Legendre polynomial. In particular we have $\lambda_0 = 0$ since $P_0(t) = 1$. For $m > 0$ we find

$$(1-t)^{-1}(1-P_m(t)) = \sum_{k=1}^m \left(\frac{-1}{2}\right)^{k-1} \binom{m}{k} \binom{m+k}{k} (1-t)^{k-1},$$

so we are concerned with the integrals

$$\int_{-1}^1 (1-t)^{k-1} (1+t)^\alpha dt = \frac{(-2)^{k-1} 2^\alpha (k-1)!}{(\alpha+1) \cdots (\alpha+k-1)}, \quad \alpha = \frac{n-3}{2}.$$

The proof of this equality follows by induction on k . Thus we find, for $m \geq \alpha$

$$\begin{aligned} \lambda_m &= 2^\alpha \sum_{k=1}^m \left(\binom{m}{k} \binom{m+k}{k} \frac{(k-1)!}{(\alpha+1) \cdots (\alpha+k-1)} \right) \\ &= 2^\alpha \sum_{k=1}^m \left(\frac{(m+k)!}{k!(m-k)!} \frac{k}{(\alpha+1) \cdots (\alpha+k-1)} \right) \geq 2^\alpha \sum_{k=1}^m \frac{m!}{k!(m-k)!} = 2^\alpha (2^m - 1) \end{aligned}$$

This proves that $\lambda_m > 0$ for all $m > 0$ and $\lambda_m \rightarrow \infty$ for $m \rightarrow \infty$. Hence Δ_0 is essentially self-adjoint on $\text{Lip}(\partial\mathbf{H})$, and has compact resolvent in $L^2(\partial\mathbf{H}, \nu_0)$. Moreover Δ_0 is positive with kernel the constant functions, on which p_0 projects so $\Delta_0 + p_0$ is strictly positive. \square

We wish to extend the operator Δ_0 to the module $L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}$ in a way compatible with the action of $G = \text{Isom}\mathbf{H}$ on this module.

Lemma 7.4. *The map $\Psi \mapsto P^{n/2}\Psi$, where P is the Poisson kernel (3.8), extends to a unitary isomorphism*

$$(7.4) \quad v : L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})} \rightarrow C_0(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \simeq L^2(\partial\mathbf{H}, \nu_0) \otimes C_0(\mathbf{H}),$$

of right $C_0(\mathbf{H})$ -modules.

Proof. Since $d\nu_x(\xi) = P(x, \xi)^n d\nu_0(\xi)$ it is straightforward that v is innerproduct preserving. Since it maps $C_c(T_1\mathbf{H})$ into itself, it is surjective. \square

In $L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}$ we consider the operator

$$(7.5) \quad \Delta\Psi(x, \xi) := \int \frac{\Psi(x, \xi) - \Psi(x, \eta)}{d_x(\xi, \eta)^n} d\nu_x\eta,$$

which is initially defined on the dense subspace $C_c^1(T_1\mathbf{H}) \subset L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}$.

Lemma 7.5. *The function $H : T_1\mathbf{H} \rightarrow \mathbf{R}$ given by*

$$(7.6) \quad H(x, \xi) = \int \frac{\frac{P(x, \xi)^{n/2}}{P(x, \eta)^{n/2}} - 1}{d_0(\xi, \eta)^n} d\nu_0\eta = P(x, \xi)^{n/2} \int \frac{P(x, \eta)^{-n/2} - P(x, \xi)^{-n/2}}{d_0(\xi, \eta)^n} d\nu_0\eta,$$

defines an element of $C^1(\mathbf{H}, L^\infty(\partial\mathbf{H}, \nu_0))$ via $x \mapsto H_x$, $H_x(\xi) := H(x, \xi)$.

Proof. For fixed x , $P_x : \xi \mapsto P(x, \xi)$ is nonvanishing and Lipschitz in ξ . Therefore the function

$$H_x : \partial\mathbf{H} \rightarrow \partial\mathbf{H}, \quad H_x : \xi \mapsto H(x, \xi) = P_x(\xi)^{n/2} \Delta_0(P_x^{-n/2})(\xi)$$

is bounded by Lemma 7.3 and thus defines an element of $L^2(\partial\mathbf{H}, \nu_0)$. For $x, y \in \mathbf{H}$, the function $P_x^{-n/2} - P_y^{-n/2} : \partial\mathbf{H} \rightarrow \mathbf{R}$ is defined on an open neighbourhood of $\partial\mathbf{H} \subset \mathbf{R}^{n+1}$. Denote by

grad the Euclidean gradient by and $h(x, y) := \sup_{\xi} \|\text{grad}(P_x^{-n/2}(\xi) - P_y^{-n/2}(\xi))\|$, which is a continuous function of (x, y) that vanishes on the diagonal. The mean value theorem gives

$$\|P_x(\xi)^{-n/2} - P_y(\xi)^{-n/2} - P_x(\eta)^{-n/2} + P_y(\eta)^{-n/2}\| \leq h(x, y)\|\xi - \eta\|.$$

Thus, with I_1 as in Lemma 7.2 we can estimate

$$\begin{aligned} \|\Delta_0(P_x^{\frac{-n}{2}} - P_y^{\frac{-n}{2}})\|_{\infty} &\leq \left\| \int \frac{P_x(\eta)^{\frac{-n}{2}} - P_x(\xi)^{\frac{-n}{2}} - P_y(\eta)^{\frac{-n}{2}} + P_y(\xi)^{\frac{-n}{2}}}{d_0(\xi, \eta)^n} d\nu_0 \eta \right\|_{\infty} \\ &\leq h(x, y) \int_{\partial \mathbf{H}} \frac{1}{d_0(\xi, \eta)^{n-1}} d\nu_0 \eta = I_1 h(x, y). \end{aligned}$$

Then the estimate

$$\|P_x^{n/2} \Delta_0 P_x^{-n/2} - P_y^{n/2} \Delta_0 P_y^{-n/2}\|_{\infty} \leq \|P_x^{n/2} - P_y^{n/2}\|_{\infty} \|\Delta_0 P_x^{-n/2}\|_{\infty} + \|P_y^{n/2}\|_{\infty} I_1 h(x, y),$$

shows that $\lim_{x \rightarrow y} \|P_x^{n/2} \Delta_0 P_x^{-n/2} - P_y^{n/2} \Delta_0 P_y^{-n/2}\|_{\infty} = 0$, so $x \mapsto H_x$ is a continuous map $\mathbf{H} \rightarrow L^{\infty}(\partial \mathbf{H}, \nu_0)$. The partial derivatives $\partial_{x_i} H(x, \xi)$ are continuously differentiable in ξ , and so a similar argument shows that the maps $x \mapsto \partial_{x_i} H(x, \xi)$ are continuous maps $\mathbf{H} \rightarrow L^{\infty}(\mathbf{H}, \nu_0)$ as well. \square

For a Banach space E , we denote by $C_c^1(\mathbf{H}, E)$ the space of compactly supported continuously differentiable functions on \mathbf{H} with values in E . We will always consider $\text{Dom } \Delta_0$ as a Hilbert space in the graph norm. Using the injection $\text{Dom } \Delta_0 \rightarrow L^2(\partial \mathbf{H}, \nu_0)$ we view $C_c^1(\mathbf{H}, \text{Dom } \Delta_0)$ as a subspace of $C_0(\mathbf{H}, L^2(\partial \mathbf{H}, \nu_0))$, which both are $C_0(\mathbf{H})$ -bimodules

Proposition 7.6. *Let v be as in Lemma 7.4 and Δ as in (7.5). The operator Δ is essentially self-adjoint and regular on $v^* C_c^1(\mathbf{H}, C^1(\partial \mathbf{H})) \subset L^2(T_1 \mathbf{H}, \nu_x)_{C_0(\mathbf{H})}$ and satisfies*

$$\Delta : v^* C_c^1(\mathbf{H}, \text{Dom } \Delta_0) \rightarrow v^* C_c^1(\mathbf{H}, L^2(\partial \mathbf{H}, \nu_0)).$$

It has $C_0(\mathbf{H})$ locally compact resolvent in $L^2(T_1 \mathbf{H}, \nu_x)_{C_0(\mathbf{H})}$. In particular, for all $f \in C_0(\mathbf{H})$ we have $f(1 + \Delta)^{-1} \in \mathbb{K}(L^2(T_1 \mathbf{H}, \nu_x)_{C_0(\mathbf{H})})$.

Proof. Using the map (7.4), it suffices to show that the operator $v D v^*$ with domain

$$C_c^1(\mathbf{H}, \text{Dom } \Delta_0) \subset v(C_c^1(T_1 \mathbf{H})),$$

is essentially self-adjoint, regular and has $C_0(\mathbf{H})$ -compact resolvent in $C_0(\mathbf{H}, L^2(\partial \mathbf{H}, \nu_0))$. We see that it is given by

$$(7.7) \quad v \Delta v^* \Psi(x, \xi) = \int \frac{\Psi(x, \xi) \frac{P_x^{n/2}(x, \xi)}{P_x^{n/2}(x, \eta)} - \Psi(x, \eta)}{d_0(\xi, \eta)^n} d\nu_0 \eta = (\Delta_0 \otimes 1 + H(x, \xi)) \Psi(x, \xi),$$

where $H(x, \xi) \in C^1(\mathbf{H}, L^{\infty}(\partial \mathbf{H}, \nu_0))$ is the function from (7.6). In particular, $v \Delta v^*$ maps $C_c^1(\mathbf{H}, \text{Dom } \Delta_0)$ into $C_c^1(\mathbf{H}, L^2(\partial \mathbf{H}, \nu_0))$. It follows that Δ is a map

$$\Delta : v^* C_c^1(\mathbf{H}, \text{Dom } \Delta_0) \rightarrow v^* C_c^1(\mathbf{H}, L^2(\partial \mathbf{H}, \nu_0)) \subset L^2(T_1 \mathbf{H}, \nu_x)_{C_0(\mathbf{H})},$$

and thus Δ is a densely defined symmetric operator. By [59, Théorème 1.18], $v \Delta v^*$ is self-adjoint and regular if and only if for all $x \in \mathbf{H}$ the localisation $(v \Delta v^*)_x$ is self-adjoint in $L^2(\partial \mathbf{H}, \nu_0)$. But $(v \Delta v^*)_x$ is a bounded perturbation of Δ_0 by (7.7), and therefore self-adjoint.

Thus $v\Delta v^*$ is self-adjoint and regular. By the same argument, because Δ_0 has compact resolvent, $(v\Delta v^*)_x$ has compact resolvent and thus $v\Delta v^*$ has $C_0(\mathbf{H})$ -compact resolvent. Since v is unitary, it follows that Δ is self-adjoint and regular, with $C_0(\mathbf{H})$ -compact resolvent. \square

Proposition 7.7. *The operator Δ is positive and G -equivariant. Moreover, for p the projection of Theorem 3.4, $\Delta + p$ is strictly positive and $\Delta p = p\Delta = 0$.*

Proof. By Proposition 7.6 Δ_0 is positive and $\Delta_0 + p_0$ is strictly positive. Now consider g with $0 = xg$ and for $\Psi \in C_c(\mathbf{H}, C^1(\partial\mathbf{H}))$, which is a core for Δ , write $\Psi_x^g(\xi) = \Psi(x, \xi g^{-1})$, so $\Psi_x^g \in C^1(\partial\mathbf{H})$. Then compute

$$\begin{aligned} \langle \Delta \Psi, \Psi \rangle(x) &= \int \int \frac{|\Psi(x, \xi)|^2 - \overline{\Psi(x, \eta)} \Psi(x, \xi)}{d_x(\xi, \eta)^n} d\nu_x \xi d\nu_x \eta \\ &= \int \int \frac{|\Psi(x, \xi)|^2 - \overline{\Psi(x, \eta)} \Psi(x, \xi)}{d_0(\xi g, \eta g)^n} d\nu_0(\xi g) d\nu_0(\eta g) \\ &= \int \int \frac{|\Psi(x, \xi g^{-1})|^2 - \overline{\Psi(x, \eta g^{-1})} \Psi(x, \xi g^{-1})}{d_0(\xi, \eta)^n} d\nu_0 \xi d\nu_0 \eta = \langle \Delta_0 \Psi_x^g, \Psi_x^g \rangle_0, \end{aligned}$$

and similarly one shows that $\langle p\Psi, \Psi \rangle(x) = \langle p_0 \Psi_x^g, \Psi_x^g \rangle_0$. It thus follows that Δ is positive and $\Delta + p$ is strictly positive. A simple change of variables establishes G -equivariance. The equality $\Delta p = 0$ follows because $p\Psi(x, \xi)$ is constant in ξ , and $p\Delta = 0$ thus follows by taking adjoints. \square

7.2. An unbounded Kasparov module for the extension class. In Section 4.2 the function $\rho(x) = d_{\mathbf{H}}(0, x)$ was introduced, and on $C_c(\mathbf{H}, \text{Dom } \Delta_0)$ we consider the multiplication operator $\rho\Psi(x, \xi) = \rho(x)\Psi(x, \xi)$. It is straightforward to show that Δ and ρ commute on this domain and that $\Delta + \rho$ is essentially self-adjoint and regular.

Proposition 7.8. *The positive self-adjoint regular operator $\Delta + \rho$ has compact resolvent in the C^* -module $L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}$.*

Proof. Because both Δ and ρ are positive regular operators, $(1 + \Delta + \rho)^{-1}$, $(1 + \Delta)^{-1}$ define adjointable operators and the function $(1 + \rho)^{-1}$ is an element of $C_0(\mathbf{H})$. Hence if u_n is an increasing approximate unit in $C_0(\mathbf{H})$, for $n > m$ we have the operator inequalities

$$0 \leq (u_n - u_m)(1 + \Delta + \rho)^{-1}(u_n - u_m) \leq (u_n - u_m)(1 + \rho)^{-1}(u_n - u_m) \rightarrow 0.$$

Thus the sequence $u_n(1 + \Delta + \rho)^{-1}$ is Cauchy. On the other hand

$$u_n(1 + \Delta + \rho)^{-1}u_n \leq u_n(1 + \Delta)^{-1}u_n \in \mathbb{K}_{C_0(\mathbf{H})}(L^2(T_1\mathbf{H}, \nu_x)),$$

because $u_n(1 + \Delta)^{-1} \in \mathbb{K}$ by Proposition 7.6. Since \mathbb{K} is an ideal, it is a hereditary subalgebra, and thus the operator $u_n(1 + \Delta + \rho)^{-1}u_n \in \mathbb{K}$ from which it follows that $u_n(1 + \Delta + \rho)^{-1/2} \in \mathbb{K}$, and since this sequence is Cauchy, its limit is in \mathbb{K} as well. Because the sequence converges pointwise to $(1 + \Delta + \rho)^{-1/2}$, it follows that $(1 + \Delta + \rho)^{-1} \in \mathbb{K}$, as desired. \square

Next we address the commutator properties of Δ and ρ with functions $f \in \text{Lip}(\partial\mathbf{H})$.

Lemma 7.9. *For $f \in \text{Lip}(\partial\mathbf{H})$, the operator $[\Delta, f]$ extends to a bounded operator.*

Proof. This commutator can be computed using the explicit formula for $v\Delta v^*$ from Equation (7.7) to find:

$$[v\Delta v^*, f]\Psi(x, \xi) = \int \frac{(f(\xi) - f(\eta))\Psi(x, \eta)}{d_0(\xi, \eta)^n} d\mu_0(\eta).$$

Using Hölder's inequality, the fact that f is Lipschitz and Lemma 7.2 we estimate

$$\begin{aligned} | \langle [v\Delta v^*, f]\Psi, \Phi \rangle(x) | &= \left| \int \int \frac{(f(\xi) - f(\eta))\Psi(x, \eta)}{d_0(\xi, \eta)^n} d\mu_0(\eta) \Phi(x, \xi) d\mu_0(\xi) \right| \\ &\leq \|f\|_{\text{Lip}} \int \int \frac{|\Psi(x, \eta)| |\Phi(x, \xi)|}{d_0(\xi, \eta)^{n-1}} d\mu_0(\eta) d\mu_0(\xi) \\ &\leq \|f\|_{\text{Lip}} \left(\int \int \frac{|\Psi(x, \eta)|^2}{d_0(\xi, \eta)^{n-1}} d\mu_0(\eta) d\mu_0(\xi) \right)^{\frac{1}{2}} \left(\int \int \frac{|\Phi(x, \xi)|^2}{d_0(\xi, \eta)^{n-1}} d\mu_0(\eta) d\mu_0(\xi) \right)^{\frac{1}{2}} \\ &\leq \|f\|_{\text{Lip}} I_1 \|\Phi\|_2 \|\Psi\|_2, \end{aligned}$$

which is independent of x . Thus $[v\Delta v^*, f]$ extends to an adjointable operator. \square

Next we recall the function $\rho(x) = d_{\mathbf{H}}(0, x)$ and the projection p from Theorem 3.4.

Lemma 7.10. *For $f \in \text{Lip}(\partial\mathbf{H})$, the operator $[p, f]\rho$ extends to a bounded operator.*

Proof. The proof consists of pointwise estimates in $x \in \mathbf{H}$:

$$\begin{aligned} | \langle [p, f]\Psi, \Phi \rangle(x) | &\leq \int \int |\Psi(x, \eta) \Phi(x, \xi)| |f(\xi) - f(\eta)| d\nu_x(\xi) d\nu_x(\eta) \\ &\leq \|f\|_{\text{Lip}} \int \int |\Psi(x, \eta) \Phi(x, \xi)| \|\xi - \eta\| d\nu_x(\xi) d\nu_x(\eta) \\ &\leq \|f\|_{\text{Lip}} (1 - \|x\|^2) \int \int |\Psi(x, \eta) \Phi(x, \xi)| \frac{\|x - \xi\| + \|x - \eta\|}{1 - \|x\|^2} d\nu_x(\xi) d\nu_x(\eta) \\ (7.8) \quad &\leq \|f\|_{\text{Lip}} (1 - \|x\|^2) \int \int |\Psi(x, \eta) \Phi(x, \xi)| (P(x, \xi)^{-\frac{1}{2}} + P(x, \eta)^{-\frac{1}{2}}) d\nu_x(\xi) d\nu_x(\eta) \end{aligned}$$

Thus we estimate

$$\int \int |\Phi(x, \xi) \Psi(x, \eta)| P^{-1/2}(x, \xi) d\nu_x(\xi) d\nu_x(\eta) \leq \|\Phi\|_2 \|\Psi\|_2 \left(\int \int P^{-1}(x, \xi) d\nu_x(\xi) d\nu_x(\eta) \right)^{\frac{1}{2}}.$$

Using Hölder's inequality, the fact that ν_x is a probability measure and $d\nu_x = P^n d\nu_0$ we find

$$\int P^{-1}(x, \xi) d\nu_x(\xi) = \int P^{n-1}(x, \xi) d\nu_0(\xi) \leq \left(\int P^n(x, \xi) d\nu_0(\xi) \right)^{\frac{n-1}{n}} = 1.$$

Combining this with (7.8) we obtain the estimate

$$| [p, f]\Psi, \Phi \rangle(x) | \leq 2 \|f\|_{\text{Lip}} \|\Phi\|_2 \|\Psi\|_2 (1 - \|x\|^2).$$

An elementary computation using the explicit distance formula on the hyperbolic ball (see [14, Section I.6.7]) shows that $\rho(x) = \log \frac{(1+\|x\|)^2}{1-\|x\|^2}$. It thus follows that $\rho[p, f]$ is bounded. \square

This leads us to consider the operator $S := -\Delta + \rho F_p$, as a candidate for the unbounded representative of the Fredholm module constructed in Theorem 3.4. We arrive at the main result of this section.

Theorem 7.11. *The triple $(C(\partial\mathbf{H}), L^2(T_1\mathbf{H}, \nu_x)_{C_0(\mathbf{H})}, S)$ is an unbounded G -equivariant Kasparov module representing the G -equivariant extension*

$$0 \rightarrow C_0(\mathbf{H}) \rightarrow C(\overline{\mathbf{H}}) \rightarrow C(\partial\mathbf{H}) \rightarrow 0.$$

Proof. We have $S = -\Delta + \rho F_p = F_p(\Delta + \rho) = F_p(1 + \Delta + \rho) - F_p$ by Proposition 7.7. Now $1 + \Delta + \rho$ commutes with F_p and has compact inverse by Proposition 7.8, so S has compact resolvent in $L^2(T_1\mathbf{H})_{C_0(\mathbf{H})}$. We note that since

$$[S, f] = [\rho, f]F_p - [\Delta, f] + 2[p, f]\rho,$$

it follows by Lemmas 7.9 and 7.10 that $[S, f]$ extends to a bounded adjointable operator whenever $f \in \text{Lip}(\partial\mathbf{H})$. The operators Δ and F_p are G -invariant by Theorem 3.4 and Proposition 7.7, and the function ρ commutes boundedly with G . Thus S defines a G -equivariant cycle, and by construction, the bounded transform defines the same class as F_p , so we are done. \square

8. Embedded surfaces and K -cycles for the boundary crossed product

In this section we explicitly compute the boundary map $\partial : K^0(C_0(M)) \rightarrow K^1(C(\partial\mathbf{H}) \rtimes \Gamma)$ for classes $[\not{D}_N] \in K^0(C_0(M))$ attached to a surface $(N, \partial N) \subset (\overline{M}, \partial\overline{M})$. For Bianchi groups, this gives an exhaustive description of the map ∂ by Theorem 5.5.

8.1. Hyperbolic Dirac operator and Poisson kernel. We now prove several technical Lemmas concerning the commutation relations of the hypersingular integral operator Δ (7.5), the projection p from Theorem 3.4 and the hyperbolic Dirac operator $\not{D}_{\mathbf{H}}$. In the next section we use these results to compute the Kasparov product of the boundary extension with Dirac operators on embedded surfaces.

We set some conventions. Following Patterson [57, Page 294] we view \mathbf{H} as the unit ball in \mathbf{R}^{n+1} as before, with the Riemannian metric $ds^2 = \frac{dx^2}{(1-\|x\|^2)^2}$. Let $\mathcal{S} \rightarrow \mathbf{H}$ be the spinor bundle and consider the internal tensor product of $C_0(\mathbf{H})$ - C^* -modules

$$C_0(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\mathbf{H})} C_0(\mathbf{H}, \mathcal{S}) \simeq L^2(\partial\mathbf{H}, \nu_0) \otimes C_0(\mathbf{H}, \mathcal{S}),$$

and the dense subspace

$$(8.1) \quad W = W_{\mathbf{H}} = C_c^1(\mathbf{H}, \text{Dom } \Delta_0) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S}).$$

Let $\Psi \in C_c^1(\mathbf{H}, \mathcal{S})$ be a compactly supported C^1 -section and \mathbf{x} a tangent vector field. We denote the Clifford representation associated to the hyperbolic metric on \mathbf{H} by $\psi \mapsto c(\mathbf{x})\psi$. Let $e_i \in \mathbf{R}^{n+1}$ denote the i -th standard basis vector. The vector fields $e_i(x) = e_i(1 - \|x\|^2)$ define a global orthonormal frame for the tangent bundle on \mathbf{H} . The hyperbolic Dirac operator $\not{D}_{\mathbf{H}}$ can be computed by elementary methods (see for instance [27, Theorem 5.3.5]) and is given by

$$\not{D}_{\mathbf{H}} : C_c^1(\mathbf{H}, \mathcal{S}) \rightarrow C_c(\mathbf{H}, \mathcal{S}), \quad \psi \mapsto \sum_{i=0}^n (1 - \|x\|^2) c(\mathbf{e}_i) \partial_i \psi + L_i c(\mathbf{e}_i) \psi.$$

Here the L_i are bounded functions on \mathbf{H} . It induces an operator

$$T_{\mathbf{H}} : C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S}) \rightarrow C_0(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\mathbf{H})}^{\text{alg}} C_0(\mathbf{H}, \mathcal{S}),$$

$$\begin{aligned} T_{\mathbf{H}}(\Psi \otimes \psi) &= \Psi \otimes \mathcal{D}_{\mathbf{H}}\psi + \partial_i \Psi \otimes c(\mathbf{e}_i)\psi \\ &= (1 - \|x\|^2) \left(\sum_{i=0}^n \Psi \otimes c(\mathbf{e}_i)(\partial_i \psi) + (\partial_i \Psi) \otimes c(\mathbf{e}_i)\psi \right) + L_i \Psi \otimes c(\mathbf{e}_i)\psi, \end{aligned}$$

For $s \in \mathbf{R}$, the powers of the Poisson kernel define multiplication operators

$$P^s : C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \rightarrow C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)), \quad \Psi \mapsto P^s \Psi.$$

Lemma 8.1. *Let $s \in \mathbf{R}$ and $\Psi \otimes \psi \in W_{\mathbf{H}}$. The operators P^s and $T_{\mathbf{H}}$ satisfy*

$$\begin{aligned} T_{\mathbf{H}} : W_{\mathbf{H}} &\rightarrow C_c(\mathbf{H}, \text{Dom } \Delta_0) \otimes_{C_c(\mathbf{H})}^{\text{alg}} C_c(\mathbf{H}, \mathcal{S}), \\ P^s : W_{\mathbf{H}} &\rightarrow C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S}). \end{aligned}$$

There are functions $u_i \in C_b(T_1\mathbf{H})$ with $\sum_{i=0}^n u_i^2 = 1$ such that

$$[T_{\mathbf{H}}, P^s](\Psi \otimes \psi) = 2sP^s \sum_{i=0}^n c(\mathbf{e}_i)u_i(\Psi \otimes \psi).$$

Proof. The domain mapping properties are straightforward to check and guarantee that commutator $[T_{\mathbf{H}}, P^s]$ is well-defined. By the derivation property of $T_{\mathbf{H}}$, we need to compute

$$(1 - \|x\|^2)\partial_i(P(x, \xi)^s) = 2sP(x, \xi)^s \frac{(\xi_i - x_i)(1 - \|x\|^2) - x_i\|\xi - x\|^2}{\|x - \xi\|^2},$$

so we have

(8.2)

$$[T_{\mathbf{H}}, P^s](\Psi \otimes \psi) = 2sP^s \sum_i c(\mathbf{e}_i)u_i(\Psi \otimes \psi), \quad u_i(x, \xi) = \frac{(\xi_i - x_i)(1 - \|x\|^2) - x_i\|\xi - x\|^2}{\|x - \xi\|^2}$$

Now since

$$|\xi_i - x_i| \leq \left(\sum_{k=0}^n (x_k - \xi_k)^2 \right)^{\frac{1}{2}} \leq \|x - \xi\|, \quad 1 - \|x\|^2 \leq (1 + \|x\|)\|x - \xi\|$$

we find $|u_i(x, \xi)| \leq 1 + \|x\| - x_i$ which is a bounded function. The proof that $\sum u_i^2 = 1$ is an elementary computation which we omit. \square

The operator $v\Delta v^*$ from (7.7) induces an operator

$$(v\Delta v^*) \otimes 1 : W_{\mathbf{H}} \rightarrow C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S}),$$

which we will denote by $v\Delta v^*$ as well.

Proposition 8.2. *The operators $(v\Delta v^*) \otimes 1$ and $(vpv^*) \otimes 1$ satisfy*

$$(v\Delta v^*) \otimes 1, (vpv^*) \otimes 1 : W_{\mathbf{H}} \rightarrow C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S}).$$

For $\Psi \otimes \psi \in W_{\mathbf{H}}$ it holds that

$$[T_{\mathbf{H}}, vpv^*](\Psi \otimes \psi) = n \sum_i (v(u_i p + p u_i) v^*) \Psi \otimes c(\mathbf{e}_i)\psi,$$

$$[v\Delta v^*, T_{\mathbf{H}}](\Psi \otimes \psi) = n \sum_i g_i \Psi \otimes c(\mathbf{e}_i)\psi, \quad g_i(x, \xi) := \left(\int \frac{(u_i(x, \xi) - u_i(x, \eta))}{d_x(\xi, \eta)^n} d\nu_x(\eta) \right).$$

Moreover, the functions $g_i : T_1\mathbf{H} \rightarrow \mathbf{R}$ satisfy $\sup_{(x,\xi)} \sum g_i(x, \xi)^2 < \infty$.

Proof. The operator vpv^* can be written as

$$vpv^*\Psi(x, \xi) = P(x, \xi)^{n/2} \int \Psi(x, \eta) P^{n/2}(x, \eta) d\nu_0\eta,$$

from which the domain mapping property follows readily. The formula for commutator is a direct application of Equation 8.2. We turn to the operator $v\Delta v^*$. Recall from Equation (7.7) that the operator $v\Delta v^*$ can be written as $v\Delta v^* = \Delta_0 \otimes 1 + H$, with H as in Equation (7.6). By Lemma 7.5, $H \in C^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0))$ and thus H multiplies $C_c^1(\mathbf{H}, \text{Dom } \Delta_0)$ into $C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0))$. Clearly $\Delta_0 \otimes 1$ maps $C_c^1(\mathbf{H}, \text{Dom } \Delta_0)$ into $C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0))$ as well. Therefore

$$(v\Delta v^*) \otimes 1 : W_{\mathbf{H}} \rightarrow C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S}),$$

and the commutator $[T_{\mathbf{H}}, (v\Delta v^*) \otimes 1]$ is well defined and equals $[T_{\mathbf{H}}, H \otimes 1]$. By Lemma 8.1 and the Leibniz rule

$$[T_{\mathbf{H}}, P(x, \eta)^{n/2} P(x, \xi)^{-n/2}] = nP(x, \eta)^{n/2} P(x, \xi)^{-n/2} \sum_i c(\mathbf{e}_i)(u_i(x, \xi) - u_i(x, \eta)).$$

It now follows that on $W_{\mathbf{H}}$

$$\begin{aligned} [T_{\mathbf{H}}, H \otimes 1] &= n \sum_i c(\mathbf{e}_i) \int \frac{P(x, \eta)^{n/2} P(x, \xi)^{-n/2} (u_i(x, \xi) - u_i(x, \eta))}{d_0(\xi, \eta)} d\nu_0\eta \\ &= n \sum_i c(\mathbf{e}_i) \int \frac{(u_i(x, \xi) - u_i(x, \eta))}{P(x, \xi)^{n/2} P(x, \eta)^{n/2} d_0(\xi, \eta)} P(x, \eta)^n d\nu_0\eta \\ &= n \sum_i c(\mathbf{e}_i) \int \frac{(u_i(x, \xi) - u_i(x, \eta))}{d_x(\xi, \eta)} d\nu_x\eta, \end{aligned}$$

as claimed. To prove that $\sup_{(x,\xi)} \sum g_i(x, \xi)^2 < \infty$, it suffices to show that

$$(8.3) \quad \sup_{(x,\xi)} \sum_i \left(\int \frac{(u_i(x, \xi) - u_i(x, \eta))}{d_x(\xi, \eta)^n} d\mu_x(\eta) \right)^2 \leq \sup_{(x,\xi)} \sum_i \left(\int \frac{|u_i(x, \xi) - u_i(x, \eta)|}{d_x(\xi, \eta)^n} d\mu_x(\eta) \right)^2,$$

is finite. By Hölder's inequality we have, for $0 < s < 1$ that

$$(8.4) \quad \left(\int \frac{|u_i(x, \xi) - u_i(x, \eta)|}{d_x(\xi, \eta)^n} d\nu_x(\eta) \right)^2 \leq \int \frac{(u_i(x, \xi) - u_i(x, \eta))^2}{d_x(\xi, \eta)^{n+s}} d\nu_x(\eta) \int \frac{1}{d_x(\xi, \eta)^{n-s}} d\nu_x(\eta).$$

A lengthy but elementary calculation shows that

$$(8.5) \quad \sum_{i=0}^n (u_i(x, \xi) - u_i(x, \eta))^2 = \|\xi - \eta\|^2 P(x, \xi) P(x, \eta) = d_x(\xi, \eta)^2.$$

Combining (8.3), (8.4) and (8.5), we find

$$\begin{aligned} \sum_i \left(\int \frac{(u_i(x, \xi) - u_i(x, \eta))}{d_x(\xi, \eta)^n} d\nu_x(\eta) \right)^2 &\leq I_s \sum_i \int \frac{(u_i(x, \xi) - u_i(x, \eta))^2}{d_x(\xi, \eta)^{n+s}} d\nu_x(\eta) \\ &= I_s \int \frac{d_x(\xi, \eta)^2}{d_x(\xi, \eta)^{n+s}} d\nu_x(\eta) = I_s \int \frac{1}{d_x(\xi, \eta)^{n+s-2}} d\nu_x(\eta) = I_s I_{2-s}, \end{aligned}$$

which proves boundedness by Lemma 7.2. \square

8.2. Kasparov products with embedded surfaces. By Theorem 5.5, any element of the group $K^0(C_0(M))$ of a Bianchi manifold M can be represented by the self-adjoint Dirac operator on a closed embedded hypersurface $\dot{N} \rightarrow M$. Throughout we will use that the spinor bundle on a closed embedded hypersurface is the restriction of the spinor bundle of the ambient manifold (cf. [5, 32]).

We will consider the embedded hypersurface $\Sigma := \pi^{-1}(\dot{N}) \subset \mathbf{H}$ inside the universal cover \mathbf{H} of M and denote by \mathbf{n} the unit normal vector field of $\Sigma \subset \mathbf{H}$. Let $\mathcal{S} \rightarrow \mathbf{H}$ be the spinor bundle of \mathbf{H} , which is the pullback of the spinor bundle of M under the covering map $\pi : \mathbf{H} \rightarrow M$. The Clifford module structure on $\mathcal{S}|_\Sigma$ is given by

$$c_\Sigma(\mathbf{x})\psi := c(\mathbf{x})c(\mathbf{n})\psi,$$

with \mathbf{x} a vector field on Σ . We denote by $\not{D}_\Sigma : C_c^1(\Sigma, \mathcal{S}|_\Sigma) \rightarrow C_c(\Sigma, \mathcal{S}|_\Sigma)$ the Dirac operator on $\mathcal{S}|_\Sigma$ associated to this Clifford module structure. The map

$$(8.6) \quad \sigma : \mathcal{S}|_\Sigma \rightarrow \mathcal{S}|_\Sigma, \quad \psi \mapsto ic(\mathbf{n})\psi$$

is self-adjoint for the Riemannian inner product on \mathcal{S} and squares to 1. As such σ induces a grading on $\mathcal{S}|_\Sigma$, giving a decomposition $\mathcal{S}|_\Sigma^\pm \oplus \mathcal{S}|_\Sigma^\mp$. Moreover, since $\mathcal{S}|_{\mathbf{H}}$ is G -equivariant, $\mathcal{S}|_\Sigma$ is Γ -equivariant. Similar relations hold for the spinor bundles of M and \dot{N} .

Let $\langle \psi, \varphi \rangle_{\mathcal{S}}$ denote the inner product on the spinor bundle and $C_0(\Sigma, \mathcal{S})$ the associated $C_0(\Sigma)$ module of sections. Moreover, we write $L_\pi^2(\Sigma, \mathcal{S}|_\Sigma)$ for the $C_0(\dot{N})$ -module obtained as the completion of $C_c(\Sigma, \mathcal{S})$ in the inner product norm given by

$$\langle \psi, \varphi \rangle(n) := \sum_{x \in \pi^{-1}(n)} \langle \psi, \varphi \rangle_{\mathcal{S}}(x) \in C_0(\dot{N}).$$

Proposition 8.3. *The right $C_c(\dot{N})$ module map*

$$v : C_c(T_1(\mathbf{H})) \otimes_{C_c(M)}^{\text{alg}} C_c(\dot{N}, \mathcal{S}_{\dot{N}}) \rightarrow C_c(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c(\Sigma)}^{\text{alg}} C_c(\Sigma, \mathcal{S}|_\Sigma),$$

$$\Psi \otimes \psi \mapsto (P^{n/2} \cdot \Psi)|_\Sigma \otimes \pi^* \psi$$

is well-defined and extends to a unitary isomorphism of $C_0(\dot{N})$ modules

$$v : L_\pi^2(T_1\mathbf{H}, \nu_x) \otimes_{C_0(M)} C_0(\dot{N}, \mathcal{S}_{\dot{N}}) \rightarrow C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L_\pi^2(\Sigma, \mathcal{S}|_\Sigma)_{C_0(\dot{N})},$$

and to a unitary isomorphism of Hilbert spaces

$$v : L_\pi^2(T_1\mathbf{H}, \nu_x) \otimes_{C_0(M)} L^2(\dot{N}, \mathcal{S}_{\dot{N}}) \rightarrow C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S}|_\Sigma).$$

Here the latter Hilbert space is the completion of $C_c(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c(\Sigma)}^{\text{alg}} C_c(\Sigma, \mathcal{S}|_\Sigma)$ in the inner product

$$(8.7) \quad \langle \Psi \otimes \psi, \Phi \otimes \varphi \rangle_\mu = \int_\Sigma \langle \Psi \otimes \psi, \Phi \otimes \varphi \rangle_{C_0(\Sigma)}(x) d\mu(x),$$

where $\langle \Psi \otimes \psi, \Phi \otimes \varphi \rangle_{C_0(\Sigma)}$ denotes the inner product on $C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} C_0(\Sigma, \mathcal{S})$. The algebra $C(\partial\mathbf{H})$ acts by pointwise multiplication and the left Γ -representation is given by

$$(8.8) \quad v(u_\gamma \otimes 1)v^*(\Psi \otimes \psi) = |\gamma'|^{-\frac{n}{2}}(\gamma \circ \Psi) \otimes (\gamma \circ \psi).$$

Proof. First note that v is well defined for if $\chi \in C_c(\Sigma)$ is a function such that $\chi = 1$ on $\text{supp } \Psi|_\Sigma$, then

$$(P^{n/2} \cdot \Psi)|_\Sigma \otimes \pi^* \psi = (P^{n/2} \cdot \Psi)|_\Sigma \otimes \chi \pi^* \psi \in C_c(\Sigma, L^2(\partial \mathbf{H}, \nu_0)) \otimes_{C_c(\Sigma)} C_c(\Sigma, \mathcal{S}|_\Sigma).$$

The balancing relation is respected by v for if $g \in C_0(M)$ and χ is as above then

$$\begin{aligned} v(\Psi \cdot g \otimes \psi) &= v(\Psi \chi \pi^*(g) \otimes \psi) = P^{n/2} \cdot \Psi \chi \pi^*(g) \otimes \pi^*(\psi) \\ &= P^{n/2} \cdot \Psi \otimes \chi \pi^*(g)|_\Sigma \pi^* \psi = P^{n/2} \cdot \Psi \otimes \chi \pi^*(g|_N \psi) = v(\Psi \otimes g \cdot \psi). \end{aligned}$$

Compatibility of the inner products follows by

$$\begin{aligned} \langle v(\Psi \otimes \psi), v(\Phi \otimes \varphi) \rangle(n) &= \langle \pi^* \psi, \langle P^{n/2} \Psi, P^{n/2} \Phi \rangle_{C_0(\Sigma, L^2)} \pi^* \varphi \rangle(n) \\ &= \sum_{x \in \pi^{-1}(n)} \langle \pi^* \psi, \langle P^{n/2} \Psi, P^{n/2} \Phi \rangle_{C_0(\Sigma, L^2)} \pi^* \varphi \rangle_{\mathcal{S}_\Sigma}(x) \\ &= \sum_{x \in \pi^{-1}(n)} \langle \psi(n), \langle \Psi, \Phi \rangle_{L^2(T_1 \mathbf{H}, \nu_x)}(x) \varphi(n) \rangle_{\mathcal{S}_{\dot{N}}} \\ &= \langle \psi(n), \langle \Psi, \Phi \rangle_{L^2_\pi(T_1 \mathbf{H}, \nu_x)_{C_0(M)}}(n) \varphi(n) \rangle_{\mathcal{S}_{\dot{N}}} = \langle \Psi \otimes \psi, \Phi \otimes \varphi \rangle(n), \end{aligned}$$

so it remains to show that v has dense range. Choose a pre-compact open cover $\{U_i\}$ of Σ with the property that $U_i \gamma \cap U_i = \emptyset$ whenever $\gamma \neq e$ and let χ_i^2 be a partition of unity subordinate to $\{U_i\}$. Then for each χ_i , $\psi \in C_c(\Sigma, \mathcal{S}|_\Sigma)$ and $\Psi \in C_c(\Sigma, L^2(\partial \mathbf{H}, \nu_0))$ there is a section $\psi_i \in C_c(\dot{N}, \mathcal{S})$ such that $\chi_i \psi = \pi^* \psi_i|_{U_i}$, and a function $\Psi_i \in C_c(T_1 \mathbf{H})$ such that $\Psi \chi_i = P^{n/2} \Psi_i$. Now choose functions $f_i \in C_c(\Sigma)$ with $f_i = 1$ on U_i , so that

$$\Psi \otimes \psi = \sum \Psi \chi_i f_i \otimes \chi_i \psi = \sum \Psi \chi_i \otimes f_i \pi^* \psi_i = \sum \Psi \chi_i \otimes \pi^* \psi_i = \sum v(\Psi_i \otimes \varphi_i),$$

which shows that v is surjective and thus extends to unitary isomorphism of the C^* -completions. To see that v extends to the Hilbert space completions, we need only observe that $\pi : \Sigma \rightarrow \dot{N}$ is a local isometry, so $L^2(\Sigma, \mathcal{S}|_\Sigma, \mu) \simeq C_0(\Sigma, \mathcal{S}|_\Sigma) \otimes_{C_0(\dot{N})} L^2(\dot{N}, \mu)$. The statements about the algebra representations follow by straightforward calculation. \square

Similar to (8.1) we consider the subspace $W_\Sigma := C_c^1(\Sigma, \text{Dom } \Delta_0) \otimes_{C_c^1(\Sigma)}^{\text{alg}} C_c^1(\Sigma, \mathcal{S}|_\Sigma)$. The restriction maps

$$C_c^1(\mathbf{H}) \rightarrow C_c^1(\Sigma), \quad C_c^1(\mathbf{H}, \mathcal{S}) \rightarrow C_c^1(\Sigma, \mathcal{S}|_\Sigma), \quad C_c^1(\mathbf{H}, \text{Dom } \Delta_0) \rightarrow C_c^1(\Sigma, \text{Dom } \Delta_0),$$

are all surjective and we conclude that restriction gives a surjection $W_{\mathbf{H}} \rightarrow W_\Sigma$.

The closed embedded surface $\Sigma \subset \mathbf{H}$ admits unit normal vector field \mathbf{n} , which we can extend locally to a vector field on \mathbf{H} . For $x \in \Sigma$ let $\{\mathbf{x}_i\}_{i=0}^n$ be a local orthonormal frame at x for the tangent bundle of \mathbf{H} , with $\mathbf{x}_n = \mathbf{n}$. We define $T_\Sigma : W_\Sigma \rightarrow C_0(\Sigma, L^2) \otimes_{C_0(\Sigma)} C_0(\Sigma, \mathcal{S}|_\Sigma)$ by

$$T_\Sigma(\Psi \otimes \psi)(x) := (1 - \|x\|^2) \left(\sum_{k=0}^{n-1} \partial_{\mathbf{x}_k} \Psi(x) \otimes c_\Sigma(\mathbf{x}_k) \psi(x) \right) + \Psi(x) \otimes (\not{D}_\Sigma \psi)(x).$$

Lemma 8.4. *The operator T_Σ is essentially self-adjoint on $W_\Sigma \subset C_0(\Sigma, L^2(\partial \mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S})$. Moreover T_Σ commutes with functions $f \in C(\partial \mathbf{H})$ and the Γ -representation (8.8).*

Proof. The operator $\mathcal{D}_\Sigma : C_c^1(\Sigma, \mathcal{S}) \rightarrow C_c(\Sigma, \mathcal{S})$ is essentially self-adjoint because Σ is a complete manifold. The subspace $L^2(\partial\mathbf{H}, \nu_0) \otimes^{\text{alg}} C_c(\Sigma) \subset C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0))$ is dense and

$$L^2(\partial\mathbf{H}, \nu_0) \otimes^{\text{alg}} C_c(\Sigma) \otimes_{C_c(\Sigma)} C_c(\Sigma, \mathcal{S}) \rightarrow L^2(\partial\mathbf{H}, \nu_0) \otimes C_c(\Sigma, \mathcal{S}), \quad f \otimes g \otimes \psi \mapsto f \otimes g\psi,$$

extends to a unitary isomorphism

$$(8.9) \quad \alpha : C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S}) \rightarrow L^2(\partial\mathbf{H}, \nu_0) \otimes L^2(\Sigma, \mathcal{S}).$$

For $f \in L^2(\partial\mathbf{H}, \nu_0)$, $g \in C_c^1(\Sigma)$ and $\psi \in C_c^1(\Sigma, \mathcal{S})$ we have $f \otimes g \otimes \psi \in W_\Sigma$ and

$$(\alpha \circ T_\Sigma)(f \otimes g \otimes \psi) = f \otimes \mathcal{D}_\Sigma(g\psi), \quad \alpha T_\Sigma \alpha^{-1} = 1 \otimes \mathcal{D}_\Sigma.$$

While the elements $g\psi$ span the core $C_c^1(\Sigma, \mathcal{S})$ on which \mathcal{D}_Σ is essentially selfadjoint it follows that $\alpha \circ (T_\Sigma \pm i)$ has dense range, and hence so does $T_\Sigma \pm i$. Lastly, it is clear that T_Σ commutes with functions $f \in C(\partial\mathbf{H})$. For the Γ -representation (8.8), it is enough to observe that under the map α above we have

$$\alpha[T_\Sigma, vu_\gamma v^*] \alpha^{-1} f \otimes \psi = [1 \otimes \mathcal{D}_\Sigma, |\gamma'|^{-\frac{n}{2}} u_\gamma] f \otimes \psi = 0,$$

because \mathcal{S} is Γ -equivariant and γ' does not depend on x . \square

Let $\psi \in C_c^1(\mathbf{H}, \mathcal{S})$ and $x \in \Sigma$. By [5, Proposition 2.2] the hyperbolic Dirac operator $\mathcal{D}_\mathbf{H}$ and the surface Dirac operator \mathcal{D}_Σ are related by the formulae

$$c(\mathbf{n})\mathcal{D}_\mathbf{H}\psi(x) = \mathcal{D}_\Sigma\psi(x) - K\psi(x) + \nabla_\mathbf{n}^{\mathbf{H}}\psi(x), \quad (\mathcal{D}_\mathbf{H}c(\mathbf{n}) - c(\mathbf{n})\mathcal{D}_\mathbf{H})\psi(x) = 2\mathcal{D}_\Sigma\psi(x).$$

Here ψ is a section of the spinor bundle on \mathbf{H} defined in a neighbourhood of Σ , $\nabla^{\mathbf{H}}$ is the spin connection in the spinor bundle of \mathbf{H} and K is the *mean curvature* of the surface Σ . For $\Psi \otimes \psi \in W_\mathbf{H}$ and $x \in \Sigma$ we find the simple formula

$$(8.10) \quad (T_\mathbf{H}c(\mathbf{n}) - c(\mathbf{n})T_\mathbf{H})(\Psi \otimes \psi)(x) = 2T_\Sigma(\Psi \otimes \psi)(x).$$

Choose $\lambda_k^i \in C_c(\mathbf{H})$ such that $\mathbf{e}_i = \sum_k \lambda_k^i \mathbf{x}_k$ in a neighbourhood of x . The following Lemma allows us to exploit the commutator computations of the previous section.

Lemma 8.5. *Let $R : W_\mathbf{H} \rightarrow C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S})$ be an operator such that $[T_\mathbf{H}, R] = \sum_{i=0}^n c(\mathbf{e}_i)R_i$, where*

$$R_i : W_\mathbf{H} \rightarrow C_c^1(\mathbf{H}, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_c^1(\mathbf{H})}^{\text{alg}} C_c^1(\mathbf{H}, \mathcal{S})$$

are operators that commute with the Clifford action. Then for $\Psi \otimes \psi \in W_\mathbf{H}$

$$(8.11) \quad ([T_\Sigma, R])(\Psi \otimes \psi)(x) = \sum_{k=0}^{n-1} \sum_{i=0}^n \lambda_k^i R_i c_\Sigma(\mathbf{x}_k)(\Psi \otimes \psi)(x),$$

locally at x . If the R_i define bounded operators on $C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S}|_\Sigma)$ then $[T_\Sigma, R]$ extends to a bounded operator on $C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S}|_\Sigma)$.

Proof. Equation (8.11) is obtained by linear algebra using Equation (8.10) and the local relations $c(\mathbf{x}_k)^2 = 1$ for $0 \leq k \leq n$ and $c(\mathbf{x}_i)c(\mathbf{x}_j) = -c(\mathbf{x}_j)c(\mathbf{x}_i)$ for $i \neq j$. To see that $[T_\Sigma, R]$ defines

a bounded operator whenever the R_i are bounded we first compute the $C_0(\Sigma)$ -valued inner product $\langle \cdot, \cdot \rangle_{C_0(\Sigma)}$ from Equation 8.7, and do a pointwise estimate:

$$\begin{aligned} \langle [T_{C_0(\Sigma)}, R]w, [T_\Sigma, R]w \rangle_{C_0(\Sigma)}(x) &= \sum_{i=0}^n \left(\sum_{k=1}^{n-1} \lambda_k^i(x)^2 \right) \langle R_i w, R_i w \rangle_{C_0(\Sigma)}(x) \\ &\leq \sum_{i=0}^n \langle R_i w, R_i w \rangle_{C_0(\Sigma)}(x). \end{aligned}$$

Therefore, integration against the measure μ , for the inner product (8.7) we find

$$\langle [T_\Sigma, R]w, [T_\Sigma, R]w \rangle_\mu \leq \sum_{i=0}^n \langle R_i w, R_i w \rangle_\mu \leq \left(\sum_{i=0}^n \|R_i\|^2 \right) \langle w, w \rangle_\mu,$$

which proves boundedness of $[T_\Sigma, R]$ on $C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S}|_\Sigma)$. \square

We write $\sigma := ic(\mathbf{n})$, which satisfies $\sigma^* = \sigma$ and $\sigma^2 = 1$ and commutes with the operator S constructed in Theorem 7.11.

Proposition 8.6. *The operator $v\sigma S v^* + T_\Sigma : W_\Sigma \rightarrow C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S}|_\Sigma)$ is essentially self-adjoint.*

Proof. Write $t := \mathcal{D}_\Sigma$ and $s := v\sigma S v^*$. On $W_\Sigma \subset \text{Dom } s \cap \text{Dom } t$ we can write

$$\begin{aligned} st + ts &= T_\Sigma v\sigma S v^* + v\sigma S v^* T_\Sigma = \sigma(T_\Sigma v\sigma S v^* - v\sigma S v^* T_\Sigma) \\ &= \sigma[T_\Sigma, v\Delta v^*] + \sigma\rho[T_\Sigma, vpv^*] + \sigma[T_\Sigma, \rho]vpv^*. \end{aligned}$$

It is straightforward to check that formula (7.7) holds for the isometry v from Proposition 8.3. By Proposition 8.2 and Lemma 8.5, the operators $[T_\Sigma, v\Delta v^*]$, $[T_\Sigma, vpv^*]$ and $[T_\Sigma, \rho]$ extend to bounded operators. The unbounded multiplication operator ρ commutes with the other operators involved. Now since

$$S^2 = (-\Delta - \rho + 2p\rho)^2 = \Delta^2 + 2\Delta\rho + \rho^2 \geq \rho^2,$$

it follows that $\rho^2(1+S^2)^{-1} \leq \rho^2(1+\rho^2)^{-1} \leq 1$ and hence $\rho(vSv^* \pm i)^{-1}$ extends to a bounded operator. The operators $(s \pm i)^{-1}$ preserve the core W_Σ and we have shown that $(st+ts)(s \pm i)^{-1}$ extends to a bounded operator. Thus s and t satisfy [54, Definition A.1] (see also [34]) and by [54, Theorem A.4] the sum operator $s + t$ is self-adjoint on $\text{Dom } s \cap \text{Dom } t$. \square

For $\chi \in C_c(\mathbf{H})$ and $f \in L^2(\partial\mathbf{H}, \nu_0)$ we denote by $f \cdot \chi$ the function $f \cdot \chi(x, \xi) := \chi(x)f(\xi)$. Using that χ has compact support, it is straightforward to check that $f \cdot \chi \in L^2(T_1\mathbf{H}, \nu_x)_{C_0(M)}$.

Lemma 8.7. *For any function $\chi \in C_c^1(\mathbf{H})$ for which $\pi : \text{supp } \chi \rightarrow M$ is injective and $f \in L^2(\partial\mathbf{H}, \nu_0)$, the operator*

$$\psi \mapsto v^* T_\Sigma v(f \cdot \chi \otimes \psi) - f \cdot \chi \otimes \mathcal{D}_{\mathring{N}} \psi,$$

extends to a bounded operator $L^2(\mathring{N}, \mathcal{S}) \rightarrow L^2_\pi(T_1\mathbf{H}, \nu_x) \otimes_{C_0(M)} L^2(\mathring{N}, \mathcal{S})$.

Proof. As in the proof of Proposition 5.6, there exists $\zeta \in C_c^1(M)$ such that $\chi = (\pi^* \zeta)|_U$. Then

$$\begin{aligned} T_\Sigma v(f \cdot \chi \otimes \psi) - v(f \cdot \chi \otimes \mathcal{D}_{\mathring{N}} \psi) &= T_\Sigma(P^{n/2} f \cdot \chi \otimes \pi^* \psi) - P^{n/2} f \cdot \chi \otimes \pi^* \mathcal{D}_{\mathring{N}} \psi \\ &= [T_\Sigma, P^{n/2}] f \cdot \chi \otimes \pi^* \psi + P^{n/2} f \cdot (T_\Sigma \chi \otimes \pi^* \psi - \chi \otimes \pi^* \mathcal{D}_{\mathring{N}} \psi) \end{aligned}$$

and we consider both summands separately. Writing $\bar{\mathbf{x}}_k$ for the vector fields on \mathring{N} , satisfying $c(\mathbf{x}_k)(x) = \pi^*c(\bar{\mathbf{x}}_k)(x)$, Lemmas 8.2 and 8.5 give the local expression

$$\begin{aligned} [T_\Sigma, P^{n/2}]f \cdot \chi \otimes \pi^*\psi(x) &= nP^{n/2} \sum_{k=0}^{n-1} \sum_i \lambda_k^i u_i f \cdot \chi \otimes c_\Sigma(\mathbf{x}_k) \pi^*\psi(x) \\ &= nv \left(\sum_{k=0}^{n-1} \sum_i \lambda_k^i u_i f \cdot \chi \otimes c_N(\bar{\mathbf{x}}_k) \psi \right) (x). \end{aligned}$$

This is shown to be a bounded operator as in the proof of Lemma 8.5. For the second summand,

$$\begin{aligned} P^{n/2} f \cdot (T_\Sigma \pi^* \zeta \otimes \pi^* \psi - \pi^* \zeta \otimes \pi^* \mathcal{D}_N \psi)|_U &= \sum_{k=0}^{n-1} P^{n/2} f \cdot \pi^*(\partial_{\bar{\mathbf{x}}_k} \zeta)|_U \otimes (\pi^* c_N(\bar{\mathbf{x}}_k) \psi)|_U \\ &= v \left(\sum_{k=0}^{n-1} f \cdot \pi^*(\partial_{\bar{\mathbf{x}}_k} \zeta)|_U \otimes c_N(\bar{\mathbf{x}}_k) \psi \right), \end{aligned}$$

which defines a bounded operator as in Proposition 5.6. \square

Theorem 8.8. *The triple $(C(\partial\mathbf{H}) \rtimes \Gamma, L_\pi^2(T_1\mathbf{H}, \nu_x) \otimes_{C_0(M)} L^2(\mathring{N}, \mathcal{S}_N), \sigma S + v^* T_\Sigma v)$ is an unbounded Fredholm module representing the class $\partial[\mathcal{D}_N] \in K^1(C(\partial\mathbf{H}) \rtimes \Gamma)$.*

Proof. The operator $\sigma S + v^* T_\Sigma v$ is self-adjoint by Proposition 8.6. By Theorem 7.11 and Proposition 8.4 the operator $\sigma S + v^* T_\Sigma v$ has bounded commutators with functions $f \in C(\partial\mathbf{H})$ and with group elements u_γ . By Lemma 8.7 and combining arguments in [54, Lemma 4.3], [35, Theorem 6.7] and [54, Theorem 4.4], it follows that the triple $(C(\partial\mathbf{H}) \rtimes \Gamma, L_\pi^2(T_1\mathbf{H}, \nu_x) \otimes_{C_0(M)} L^2(\mathring{N}, \mathcal{S}_N), \sigma S + v^* T_\Sigma v)$ is a K -cycle representing the product $[\partial] \otimes [\mathcal{D}_N] = \partial[\mathcal{D}_N]$. \square

It should be noted that under the isomorphism (8.9),

$$\alpha : C_0(\Sigma, L^2(\partial\mathbf{H}, \nu_0)) \otimes_{C_0(\Sigma)} L^2(\Sigma, \mathcal{S}) \rightarrow L^2(\partial\mathbf{H}, \nu_0) \otimes L^2(\Sigma, \mathcal{S}),$$

the unbounded Fredholm module in Theorem 8.8 admits a simple description. It can be represented on the module $L^2(\partial\mathbf{H}, \nu_0) \otimes L^2(\Sigma, \mathcal{S})$ using the Γ -representation

$$u_\gamma(f \otimes \psi)(\xi, x) = |\gamma'(\xi)| f(\xi\gamma) \otimes \psi(x\gamma),$$

and the operator $\sigma(\Delta_0 \otimes 1 + H + \rho F_p) + 1 \otimes \mathcal{D}_\Sigma$. However, proving that this operator is self-adjoint with compact resolvent requires the analysis presented above.

References

- [1] L. Ahlfors, *Möbius transformations in several dimensions*, Lecture notes, School of Mathematics, University of Minnesota, Minneapolis 1981.
- [2] C. Anantharaman-Delaroche, *C^* -Algèbres de Cuntz-Krieger et groupes fuchsien*. Operator Theory, Operator Algebras and Related Topics (Timișoara, 1996) 17–35. Theta Found., Bucharest, 1997.
- [3] A. Ash and G. Stevens, *Cohomology of arithmetic groups and congruences between systems of Hecke eigenvalues*, J. Reine Angew. Math. 365 (1986), 192–220.
- [4] S. Baaj, P. Julg, *Théorie bivariante de Kasparov et opérateurs non bornées dans les C^* -modules hilbertiens*, C. R. Acad. Sci. Paris, 296 (1983), 875–878.
- [5] C. Bär, *Metrics with harmonic spinors*, Geom.Funct. Anal. 6 (1996), 899–942.

- [6] P. Baum and R. Douglas, *K-homology and index theory*, Operator Algebras and Applications, volume 38 of proceedings of Symposia in Pure Mathematics, 117–173, Providence, RI, 1982. American Mathematical Society.
- [7] P. Baum, N. Higson and T. Schick, *On the equivalence of geometric and analytic K-homology*, Pure and Appl. Math. Q. 3 (2007), 1–24.
- [8] T. Berger, *An Eisenstein ideal for imaginary quadratic fields*. Ph.D. Thesis, University of Michigan, 2005.
- [9] N. Bergeron and A. Venkatesh, *The asymptotic growth of torsion homology for arithmetic groups*. J. Inst. Math. Jussieu 12 (2013), 391–447.
- [10] E. Berkove and D. Juan-Pineda, *On the K-theory of Bianchi groups*. Bol. Soc. Mat. Mexicana (3) 2.1 (1996): 15–29.
- [11] H. Betttaieb, M. Matthey and A. Valette, *Unbounded symmetric operators in K-homology and the Baum-Connes conjecture*, J. Funct. Anal. 229 (2005) 184–237.
- [12] A. Borel and L. Ji, *Compactifications of Symmetric and Locally Symmetric Spaces*, Birkhäuser, Boston, 2006.
- [13] A. Borel and J.-P. Serre, *Corners and arithmetic groups*, Comment. Math. Helv. 48 (1973), 436–491.
- [14] M. R. Bridson and A. Haefliger, *Metric spaces of nonpositive curvature*, Springer-Verlag, Berlin, 1999.
- [15] F. Calegari and A. Venkatesh, *Towards a Torsion Jacquet-Langlands Correspondence*, [arXiv:1212.3847].
- [16] P.E. Conner and E. E. Floyd, *Differentiable Periodic Maps*, Academic Press, New York, 1964.
- [17] A. Connes, *Cyclic homology and the transverse fundamental class of a foliation*, Geometric methods in operator algebras (Kyoto, 1983). Pitman Res. Notes Math. Ser., 123, Longman Sci. Tech. Harlow, (1986), 52–144.
- [18] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego, 1994.
- [19] A. Connes, *Geometry from the spectral point of view*, Lett. Math. Phys. 34(1995), 203–238.
- [20] A. Connes and G. Skandalis, *The longitudinal index theorem for foliations*, Publ. Res. Inst. Math. Sci. 20(1984), 1139–1183.
- [21] J. Cuntz, *K-theoretic amenability for discrete groups*. J. Reine Angew. Math. 344 (1983): 180–195.
- [22] N.M. Dunfield and D. Ramakrishnan, *Increasing the number of fibered faces of arithmetic hyperbolic 3-manifolds*, Amer. J. Math. 132 (2010), 53–97.
- [23] H. Emerson and R. Meyer, *Euler characteristics and Gysin sequences for group actions on boundaries*, Math. Ann. 334 (2006), 853–904.
- [24] H. Emerson and B. Nica, *K-homological finiteness and hyperbolic groups*, J. Reine. Angew. Math, in press, [arXiv:1208.0856].
- [25] J. Franke, *Harmonic analysis in weighted L_2 -spaces*. Ann. Sci. École Norm. Sup. (4) 31.2 (1998): 181–279.
- [26] M. Fuchs, *Equivariant K-homology of Bianchi groups in the case of non-trivial class group*, Münster J. Math. 7 (2014), 589–607.
- [27] J. E. Gilbert and M. A. M. Murray, *Clifford algebras and Dirac operators in harmonic analysis*, Cambridge University Press, Cambridge, 1991.
- [28] M. Goffeng, *Equivariant extensions of $*$ -algebras*, New York J. Math. 16 (2010), 369–385.
- [29] M. Goffeng and B. Mesland, *Spectral triples and finite summability on Cuntz-Krieger algebras*, Documenta Math. 20 (2015), 89–170.
- [30] N. Higson and J. Roe, *Analytic K-Homology*. Oxford University Press, Oxford, 2001.
- [31] M. Hilsum, *Bordism invariance in KK-theory*, Math. Scand. 107 (2010), 73–89.
- [32] O. Hijazi, S. Montiel and X. Zhang, *Dirac operator on embedded hypersurfaces*, Math. Res. Lett. 8 (2001), 195–208.
- [33] M. Jakob, *A bordism-type description of homology*, Manuscripta Math. 96 (1998), 67–80 .
- [34] J. Kaad and M. Lesch, *A local-global principle for regular operators on Hilbert modules*, J. Funct. Anal. 262 (2012), 4540–4569.
- [35] J. Kaad and M. Lesch, *Spectral flow and the unbounded Kasparov product*, Adv. Math. 248 (2013), 495–530.
- [36] J. Kaad, R. Nest and A. Rennie, *KK-theory and spectral flow in von Neumann algebras*, J. K-Theory 10 (2012), 241–277.
- [37] M. Kapovich, *Hyperbolic Manifolds and Discrete Groups*, Birkhäuser, Boston, 2001.
- [38] G.G. Kasparov, *Lorentz groups: K-theory of unitary representations and crossed products*, Dokl. Akad. Nauk SSSR 275 (1984), 541–545.
- [39] G.G. Kasparov, *Equivariant KK-theory and the Novikov conjecture*, Invent. Math. 91, (1988) 147–201.

- [40] G.G. Kasparov, *K-theory, group C^* -algebras, and higher signatures*, Novikov conjectures, index theorems and rigidity 1, London Math. Soc. Lecture Notes 226, 101–146.
- [41] R. C. Kirby, *The Topology of 4-Manifolds*, Springer-Verlag, Berlin, 1989.
- [42] D. Kucerovsky, *The KK -product of unbounded modules*, K-Theory 11 (1997), 17–34.
- [43] E.C. Lance, *K-theory for certain group C^* -algebras*, Acta Math. 151 (1983), 209–230.
- [44] E.C. Lance, *Hilbert C^* -Modules: a toolkit for operator algebraists*, Cambridge University Press, Cambridge, 1995.
- [45] M.H. Lee, *Hecke operators on cohomology*, Rev. Un. Mat. Argentina 50 (2009), 99–144.
- [46] J. Lott, *Limit sets as examples in noncommutative geometry*, K-Theory 34 (2005), 283–326.
- [47] Y.I. Manin and M. Marcolli, *Holography principle and arithmetic of algebraic curves*, Adv. Theor. Math. Phys. 5 (2001), 617–650.
- [48] Y.I. Manin and M. Marcolli, *Continued fractions, modular symbols, and noncommutative geometry*, Selecta Math. (N.S.) 8 (2002), 475–521.
- [49] Y.I. Manin and M. Marcolli, *Modular shadows and the Lévy-Mellin ∞ -adic transform*, Modular Forms on Schiermonnikoog, Cambridge Univ. Press, 2008, 189–238.
- [50] B. Martelli, *An Introduction to Geometric Topology*, [arXiv1610.02592v1].
- [51] M. Matthey, *Mapping the homology of a group to the K -theory of its C^* -algebra*, Illinois J. Math. 46(2002), 953–977.
- [52] B. Mesland, *Groupoid cocycles and K -theory*, Munst. J. Math. 4 (2011), 227–250.
- [53] B. Mesland, *Unbounded bivariant K -theory and correspondences in noncommutative geometry*, J. Reine Angew. Math., 691 (2014), 101–172.
- [54] B. Mesland and A. Rennie, *Nonunital spectral triples and metric completeness in unbounded KK -theory*, J. Funct. Anal. 271 (2016), 2460–2538.
- [55] R.Meyer and R.Nest, *The Baum-Connes conjecture via localisation of categories*, Topology 45 (2006), 209–259.
- [56] P. Nicholls, *The Ergodic Theory of Discrete Groups*, Cambridge University Press, Cambridge, 1989.
- [57] S. J. Patterson, *Lectures on measures on limit sets of Kleinian groups*, Analytical and geometric aspects of hyperbolic space (Coventry/Durham, 1984) 111 (1987), Cambridge Univ. Press, 281–323.
- [58] J. Pfaff, *Selberg zeta functions on odd-dimensional hyperbolic manifolds of finite volume*, J. Reine Angew. Math. 703 (2015), 115–145.
- [59] F. Pierrot, *Opérateurs réguliers dans les C^* -modules et structure des C^* -algèbres de groupes de Lie semisimples complexes simplement connexes*, J. Lie Th. 16 (2006), 651–689.
- [60] M.V. Pimsner, *KK -groups of crossed products by groups acting on trees*, Invent. Math. 86 (1986), 603–634.
- [61] J.F. Quint, *An overview of Patterson-Sullivan theory*, online lecture notes, last accessed 14.10.2016.
- [62] A. Rahm, *Homology and K -theory of the Bianchi groups*, C. R. Math. Acad. Sci. Paris 349 (2011) 11–12, 615–619.
- [63] A. Rahm, *On the equivariant of K -homology of PSL_2 of the imaginary quadratic integers*, Ann. Inst. Fourier 66 (4) (2016) 1667–1689.
- [64] J. Rosenberg and C. Schochet, *The Künneth Theorem and the Universal Coefficient Theorem for Kasparov’s generalized K -functor*, Duke Math. J. 55 (1987), 431–474.
- [65] S. Samko, *On inversion of fractional spherical potentials by spherical hypersingular operators*, Proceedings of the Conference IWOTA-2000, Birkhäuser, 2002.
- [66] S. Samko, *Hypersingular integrals and their applications*, Taylor & Francis, London, 2002.
- [67] C. Schochet, *Topological methods for C^* -algebras I. Spectral sequences*, Pacific J. Math. 96 (1981), 193–211.
- [68] P. Scholze, *On torsion in the cohomology of locally symmetric varieties*, Ann. of Math. 182 (2015), 945–1066.
- [69] M.H. Şengün, *On the integral cohomology of Bianchi groups*, Exp. Math. 20 (2011), 487–505.
- [70] M. H. Şengün, *Arithmetic aspects of Bianchi groups*, Computing with Modular Forms, Springer-Verlag, 2011.
- [71] D. Sullivan, *The density at infinity of a discrete group of hyperbolic motions*, Publ. Math. IHES 50 (1979), 171–202.
- [72] R. Thom, *Quelques propriétés globales des variétés différentiables*, Comment. Math. Helv. 28 (1954), 17–86.
- [73] A. Valette, *Introduction to the Baum-Connes conjecture*, Lectures in Mathematics, ETH, Zürich (2002).
- [74] M. Zahle, *Harmonic calculus on fractals, a measure geometric approach. II*, Trans. Amer. Math. Soc. 357, 3407–3423.

Institut für Analysis, Leibniz Universität Hannover, Welfengarten 1, 30167, Hannover, Germany
E-mail address: mesland@math.uni-hannover.de

University of Sheffield, School of Mathematics and Statistics, Hicks Building, Sheffield, UK
E-mail address: m.sengun@sheffield.ac.uk