

Differential equations satisfied by modular forms of level 5

By Kazuhide Matsuda

Faculty of Fundamental Science, National Institute of Technology, Niihama College,
 7-1 Yagumo-chou, Niihama, Ehime 792-8580, Japan
 E-mail: matsuda@sci.niihama-nct.ac.jp
 Fax: 81-0897-37-7809

Abstract This paper describes the derivation of the level 5 versions of Ramanujan's system of ordinary differential equations satisfied by the Eisenstein series, $E_2(q)$, $E_4(q)$, and $E_6(q)$.

Key Words: theta function; theta derivatives; rational characteristics.

MSC(2010) 14K25; 11E25

1 Introduction

Let \mathbb{N}_0 and \mathbb{N} denote sets of nonnegative and positive integers, respectively. For k and $n \in \mathbb{N}$, $\sigma_k(n)$ is the sum of the k -th power of the positive divisors of n , and $\sigma_k(n) = 0$ for $n \in \mathbb{Q} \setminus \mathbb{N}_0$.

The *upper half plane* \mathbb{H}^2 is defined by $\mathbb{H}^2 = \{\tau \in \mathbb{C} \mid \Im\tau > 0\}$. Throughout this paper, we set $q = \exp(2\pi i\tau)$ and define the *Dedekind eta function* as $\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n)$.

In [4], Darboux studied a problem in mechanics and differential geometry, and encountered the system of ordinary differential equations (ODEs),

$$x \frac{d}{dx}(u_1 + u_2) = u_1 u_2, \quad x \frac{d}{dx}(u_1 + u_3) = u_1 u_3, \quad x \frac{d}{dx}(u_2 + u_3) = u_2 u_3, \quad x = \exp(\pi i\tau). \quad (1.1)$$

In [6, pp. 330], Halphen provided the following solution to the problem studied by Darboux,

$$u_1 = 1 + 8 \sum_{n=1}^{\infty} \frac{x^{2n}}{(1+x^{2n})^2}, \quad u_2 = -8 \sum_{n=1}^{\infty} \frac{x^{2n-1}}{(1-x^{2n-1})^2}, \quad u_3 = 8 \sum_{n=1}^{\infty} \frac{x^{2n-1}}{(1+x^{2n-1})^2}.$$

Equation (1.1) is called Halphen's system.

For studying Painlevé type equations of the third order, Chazy [2] considered the nonlinear differential equation for the complex function

$$y''' = 2yy'' - 3(y')^2. \quad (1.2)$$

Note that $y = u_1 + u_2 + u_3 = \pi i E_2(\tau)$ is a solution of Chazy's equation (1.2), where the Eisenstein series E_2 , E_4 , and E_6 are respectively defined by

$$E_2(q) = E_2(\tau) := 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^n, \quad E_4(q) = E_4(\tau) := 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) q^n,$$

$$E_6(q) = E_6(\tau) := 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^n.$$

In [11], Ramanujan derived the following system of ODEs,

$$q \frac{dE_2}{dq} = \frac{(E_2)^2 - E_4}{12}, \quad q \frac{dE_4}{dq} = \frac{E_2 E_4 - E_6}{3}, \quad q \frac{dE_6}{dq} = \frac{E_2 E_6 - (E_4)^2}{2}. \quad (1.3)$$

In particular, by eliminating E_4 and E_6 , $y(\tau) = \pi i E_2(\tau)$ is observed to be a solution to Chazy's equation (1.2). Further, Ohya [10] showed that Halphen's differential field is an extension of Ramanujan's differential field, whose Galois group is the symmetric group S_3 .

In [8] and [9], ODEs satisfied by the cubic theta functions were derived as

$$a(q) = \sum_{m,n \in \mathbb{Z}} q^{m^2+mn+n^2}, \quad b(q) = \sum_{m,n \in \mathbb{Z}} \omega^{n-m} q^{m^2+mn+n^2},$$

$$c(q) = \sum_{m,n \in \mathbb{Z}} q^{(n+\frac{1}{3})^2+(n+\frac{1}{3})(m+\frac{1}{3})+(m+\frac{1}{3})^2}, \quad \omega = e^{\frac{2\pi i}{3}}, \quad |q| < 1.$$

Therefore, Huber used Ramanujan's theory of theta functions. In addition, Cooper [3] also treated ODEs satisfied by modular forms by using Ramanujan's theory. In the present study, the theory of theta functions with rational characteristics has been adopted. Noted is that Hermite [7] introduced theta functions with characteristics, and Farkas and Kra [5] developed the theory of theta functions with rational characteristics.

The aim of the current research was to derive the ODEs satisfied by the modular forms of level 5, and to obtain an extension of Ramanujan's differential field by using the modular forms of level 5.

The main theorems of this study are as follows.

Theorem 1.1. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} P(q) &= \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \\ Q(q) &= \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \\ R(q) = E_2(q) &= 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^n. \end{aligned}$$

Then, we have

$$\begin{aligned} q \frac{d}{dq} P &= \frac{-13P^3 - 39P^2Q + 47PQ^2 - 9Q^3 + 2PR}{24}, \quad q \frac{d}{dq} Q = \frac{9P^3 + 47P^2Q + 39PQ^2 - 13Q^3 + 2QR}{24}, \\ q \frac{d}{dq} R &= \frac{5P^4 - 15P^3Q - 155P^2Q^2 + 15PQ^3 + 5Q^4 + R^2}{12}. \end{aligned}$$

Theorem 1.2. Further, for $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} P(q) &= 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \quad Q(q) = 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n, \\ R(q) = E_2(q^5) &= 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^{5n}. \end{aligned}$$

Then, we have

$$\begin{aligned} q \frac{d}{dq} P &= \frac{13P^3 + 39P^2Q - 47PQ^2 + 9Q^3 + 50PR}{120}, \quad q \frac{d}{dq} Q = \frac{-9P^3 - 47P^2Q - 39PQ^2 + 13Q^3 + 50QR}{120}, \\ q \frac{d}{dq} R &= \frac{P^4 - 3P^3Q - 31P^2Q^2 + 3PQ^3 + Q^4 + 125R^2}{300}. \end{aligned}$$

The remainder of this paper is organized as follows. Section 2 provides an overview of Farkas' and Kra's theory of theta functions with rational characteristics. Next, Section 3 discusses Weierstrass elliptic function theory, and Section 4 describes the preliminary results obtained for P , Q , and R . Further, Section 5 derives the identities of the theta derivatives. Sections 6 and 7 provide the proofs for Theorems 1.1 and 1.2, respectively. Section 8 presents the eta products, $\eta^5(\tau)/\eta(5\tau)$ and $\eta^5(5\tau)/\eta(\tau)$. Section 9 discusses the theorem of Farkas and Kra stating the theta derivatives. Finally, Section 10 derives the Riccati equations satisfied by the modular forms of level 5.

Acknowledgments

This work was supported by JSPS KAKENHI Grant Number JP17K14213.

2 Properties of the theta functions

2.1 Definitions

Following the work of Farkas and Kra [5], we introduce the *theta function with characteristics*, which is defined by

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(\zeta, \tau) = \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(\zeta) := \sum_{n \in \mathbb{Z}} \exp \left(2\pi i \left[\frac{1}{2} \left(n + \frac{\epsilon}{2} \right)^2 \tau + \left(n + \frac{\epsilon}{2} \right) \left(\zeta + \frac{\epsilon'}{2} \right) \right] \right),$$

where $\epsilon, \epsilon' \in \mathbb{R}$, $\zeta \in \mathbb{C}$, and $\tau \in \mathbb{H}^2$.

The relation between theta functions with rational characteristics and Jacobi theta functions is given by

$$\vartheta_1(\pi\zeta) = -\theta \begin{bmatrix} 1 \\ 1 \end{bmatrix}(\zeta), \quad \vartheta_2(\pi\zeta) = \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}(\zeta), \quad \vartheta_3(\pi\zeta) = \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix}(\zeta), \quad \vartheta_4(\pi\zeta) = \theta \begin{bmatrix} 0 \\ 1 \end{bmatrix}(\zeta).$$

The *theta constants* are given by

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(0, \tau).$$

Furthermore, we denote the *theta derivatives* by

$$\theta' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \frac{\partial}{\partial \zeta} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(\zeta, \tau) \Big|_{\zeta=0}, \quad \theta'' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \frac{\partial^2}{\partial \zeta^2} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(\zeta, \tau) \Big|_{\zeta=0},$$

and

$$\theta''' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \frac{\partial^3}{\partial \zeta^3} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(\zeta, \tau) \Big|_{\zeta=0}, \quad \theta^{(n)} \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \frac{\partial^n}{\partial \zeta^n} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}(\zeta, \tau) \Big|_{\zeta=0}, \quad (n = 1, 2, 3, 4, \dots).$$

In particular, Jacobi's derivative formula is given by

$$\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} = -\pi \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \tag{2.1}$$

2.2 Basic properties

First, we note that for $m, n \in \mathbb{Z}$,

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta + n + m\tau, \tau) = \exp(2\pi i) \left[\frac{n\epsilon - m\epsilon'}{2} - m\zeta - \frac{m^2\tau}{2} \right] \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau), \quad (2.2)$$

and

$$\theta \begin{bmatrix} \epsilon + 2m \\ \epsilon' + 2n \end{bmatrix} (\zeta, \tau) = \exp(\pi i \epsilon n) \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau). \quad (2.3)$$

Furthermore, it is easy to see that

$$\theta \begin{bmatrix} -\epsilon \\ -\epsilon' \end{bmatrix} (\zeta, \tau) = \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (-\zeta, \tau) \text{ and } \theta' \begin{bmatrix} -\epsilon \\ -\epsilon' \end{bmatrix} (\zeta, \tau) = -\theta' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (-\zeta, \tau).$$

For $m, n \in \mathbb{R}$, we get

$$\begin{aligned} & \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} \left(\zeta + \frac{n + m\tau}{2}, \tau \right) \\ &= \exp(2\pi i) \left[-\frac{m\zeta}{2} - \frac{m^2\tau}{8} - \frac{m(\epsilon' + n)}{4} \right] \theta \begin{bmatrix} \epsilon + m \\ \epsilon' + n \end{bmatrix} (\zeta, \tau). \end{aligned} \quad (2.4)$$

Note that $\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau)$ has only one zero in the fundamental parallelogram, and is given by

$$\zeta = \frac{1 - \epsilon}{2}\tau + \frac{1 - \epsilon'}{2}.$$

2.3 Jacobi's triple product identity

All the theta functions have infinite product expansions given by

$$\begin{aligned} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) &= \exp\left(\frac{\pi i \epsilon \epsilon'}{2}\right) x^{\frac{\epsilon^2}{4}} z^{\frac{\epsilon}{2}} \\ &\times \prod_{n=1}^{\infty} (1 - x^{2n})(1 + e^{\pi i \epsilon'} x^{2n-1+\epsilon} z)(1 + e^{-\pi i \epsilon'} x^{2n-1-\epsilon}/z), \end{aligned} \quad (2.5)$$

where $x = \exp(\pi i \tau)$ and $z = \exp(2\pi i \zeta)$. Therefore, according to Jacobi's derivative formula (2.1), it follows that

$$\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} (0, \tau) = -2\pi q^{\frac{1}{8}} \prod_{n=1}^{\infty} (1 - q^n)^3, \quad q = \exp(2\pi i \tau).$$

2.4 Spaces of N -th order θ -functions

Following the theory by Farkas and Kra [5], we define $\mathcal{F}_N \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right]$ as the set of all functions f satisfying the two functional equations,

$$f(\zeta + 1) = \exp(\pi i \epsilon) f(\zeta),$$

and

$$f(\zeta + \tau) = \exp(-\pi i)[\epsilon' + 2N\zeta + N\tau] f(\zeta), \quad \zeta \in \mathbb{C}, \tau \in \mathbb{H}^2,$$

where N is a positive integer and $\left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] \in \mathbb{R}^2$. This set of functions is called the space of N -th order θ -functions with characteristics $\left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right]$. Note that

$$\dim \mathcal{F}_N \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] = N.$$

For the proof, please refer to [5, pp.133].

2.5 The heat equation

The theta function satisfies the following heat equation:

$$\frac{\partial^2}{\partial \zeta^2} \theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] (\zeta, \tau) = 4\pi i \frac{\partial}{\partial \tau} \theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] (\zeta, \tau). \quad (2.6)$$

3 Weierstrass elliptic function theory

In this paper, we introduce Weierstrass \wp -function and σ -function as follows:

$$\begin{aligned} \wp(z; \omega_1, \omega_2) &= \wp(z) = \frac{1}{z^2} + \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m, n) \neq (0, 0)}} \left(\frac{1}{(z - m\omega_1 - n\omega_2)^2} - \frac{1}{(m\omega_1 + n\omega_2)^2} \right), \\ \sigma(z; \omega_1, \omega_2) &= \sigma(z) = z \prod_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m, n) \neq (0, 0)}} \left(1 - \frac{z}{m\omega_1 + n\omega_2} \right) \exp \left(\frac{z}{m\omega_1 + n\omega_2} + \frac{z^2}{2(m\omega_1 + n\omega_2)^2} \right), \end{aligned}$$

where z, ω_1, ω_2 are complex numbers with $\omega_2/\omega_1 \notin \mathbb{R}$.

The following formula is recalled from the study by Whittaker and Watson [12, pp. 437, 459]:

$$\begin{aligned}\wp'(z)^2 &= 4\wp(z)^3 - g_2\wp(z) - g_3, \\ g_2(\omega_1, \omega_2) &= 60G_4(\omega_1, \omega_2) = 60 \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m, n) \neq (0, 0)}} \frac{1}{(m\omega_1 + n\omega_2)^4}, \\ g_3(\omega_1, \omega_2) &= 140G_6(\omega_1, \omega_2) = 140 \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m, n) \neq (0, 0)}} \frac{1}{(m\omega_1 + n\omega_2)^6},\end{aligned}$$

which implies

$$\wp''(z) = 6\wp^2(z) - \frac{1}{2}g_2. \quad (3.1)$$

For $k \in \mathbb{N}$, the Eisenstein series is defined by

$$G_{2k}(\tau) = G_{2k}(1, \tau) = \sum_{\substack{(m, n) \in \mathbb{Z}^2 \\ (m, n) \neq (0, 0)}} \frac{1}{(m + n\tau)^{2k}}, \quad \tau \in \mathbb{H}^2.$$

Moreover, from [5, pp. 124], we recall

$$\begin{aligned}\wp(z; 1, \tau) &= \frac{1}{3} \frac{\theta''' \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}} - \frac{d^2}{dz^2} \log \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z, \tau), \\ \sigma(z; \omega_1, \omega_2) &= \exp \left(\frac{\eta_1 z^2}{2\omega_1} \right) \frac{\omega_1}{\theta' \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}} \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} \left(\frac{z}{\omega_1}, \tau \right),\end{aligned}$$

where $\omega_2/\omega_1 = \tau \in \mathbb{H}^2$.

4 Preliminary results

Proposition 4.1. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} = -\pi \left\{ \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n \right\},$$

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = -\pi \left\{ \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n \right\},$$

where $q = \exp(2\pi i\tau)$.

Proof. The proposition follows from Jacobi's triple product identity (2.5). \square

Proposition 4.2. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} = \frac{\pi i}{5} \left\{ 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))y^n \right\}, \quad \frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} = \frac{\pi i}{5} \left\{ 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))y^n \right\},$$

where $y = \exp(2\pi i\tau/5)$.

Proof. The proposition follows from Jacobi's triple product identity (2.5). \square

Proposition 4.3. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} = 4\pi i \frac{d}{d\tau} \log \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} = -\pi^2 E_2(q), \quad q = \exp(2\pi i\tau).$$

Proof. The proposition follows from Jacobi's triple product identity (2.5). \square

5 Applications of the residue theorem

Theorem 5.1. *The sum of all the residues of an elliptic function in the fundamental parallelogram is zero.*

5.1 Derivative formulas of level five

Theorem 5.2. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} = \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{\left(\theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} - 3\theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} \right)}{10\theta^3 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^3 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}, \quad (5.1)$$

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{\left(3\theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} + \theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} \right)}{10\theta^3 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^3 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}. \quad (5.2)$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau)}{\theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} (z, \tau)} \text{ and } \psi(z) = \frac{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau)}{\theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ -\frac{1}{5} \end{bmatrix} (z, \tau)}.$$

Note that, in the fundamental parallelogram, the poles of $\varphi(z)$ are $z = 2/5$ and $z = 1/5$. Thus, the direct calculation yields

$$\text{Res} \left(\varphi(z), \frac{2}{5} \right) = \frac{\theta^3 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^2 \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} \left\{ -3 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} + \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} \right\}$$

and

$$\text{Res} \left(\varphi(z), \frac{1}{5} \right) = - \frac{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}.$$

Because $\text{Res}(\varphi(z), 2/5) + \text{Res}(\varphi(z), 1/5) = 0$, it is understood that

$$3 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = - \frac{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta^3 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}.$$

From $\psi(z)$, we have

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} + 3 \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = \frac{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta^3 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}},$$

which proves the theorem. \square

Theorem 5.3. For every $\tau \in \mathbb{H}^2$, we have

$$\frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} = -\zeta_5^3 \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{\left(\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} + 3\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \right)}{10\theta^3 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^3 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}, \quad (5.3)$$

$$\frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} = -\zeta_5^3 \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{\left(3\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} - \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \right)}{10\theta^3 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^3 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}. \quad (5.4)$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z, \tau)}{\theta^2 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}(z, \tau) \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}(z, \tau)} \text{ and } \psi(z) = \frac{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z, \tau)}{\theta^2 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}(z, \tau) \theta \begin{bmatrix} -\frac{1}{5} \\ 1 \end{bmatrix}(z, \tau)}.$$

Then, from $\varphi(z)$ and $\psi(z)$, we have

$$3 \frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} - \frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} = -\zeta_5^3 \frac{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \theta^2 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta^3 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \quad (5.5)$$

and

$$\frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} + 3 \frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} = -\zeta_5^3 \frac{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \theta^2 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta^3 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}. \quad (5.6)$$

Thus, the theorem is proved. \square

5.2 Second-order theta derivatives

Proposition 5.4. *For every $\tau \in \mathbb{H}^2$, we have*

$$2 \frac{\theta'' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] + \theta'' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} - \frac{\theta''' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]}{\theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]} + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} \right\}^2 + 4 \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} \cdot \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]} = 0, \quad (5.7)$$

$$\frac{\theta'' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] + 2 \theta'' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} - \frac{\theta''' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]}{\theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]} - 4 \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} \cdot \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]} + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]} \right\}^2 = 0. \quad (5.8)$$

Proof. Consider the following elliptic functions:

$$\frac{\theta^2 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] (z, \tau) \theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] (z, \tau)}{\theta^3 \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] (z, \tau)}, \quad \frac{\theta \left[\begin{smallmatrix} -\frac{1}{5} \end{smallmatrix} \right] (z, \tau) \theta^2 \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] (z, \tau)}{\theta^3 \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] (z, \tau)}.$$

The proposition can be proved using Theorem 5.1. \square

Proposition 5.5. *For every $\tau \in \mathbb{H}^2$, we have*

$$2 \frac{\theta'' \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right] + \theta'' \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]} - \frac{\theta''' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]}{\theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]} + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]} \right\}^2 + 4 \frac{\theta' \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]} \cdot \frac{\theta' \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]} = 0, \quad (5.9)$$

$$\frac{\theta'' \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right] + 2 \theta'' \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]} - \frac{\theta''' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]}{\theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]} - 4 \frac{\theta' \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right]} \cdot \frac{\theta' \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]} + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right]} \right\}^2 = 0. \quad (5.10)$$

Proof. Consider the following elliptic functions:

$$\frac{\theta^2 \left[\begin{smallmatrix} \frac{1}{5} \\ 1 \end{smallmatrix} \right] (z, \tau) \theta \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right] (z, \tau)}{\theta^3 \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] (z, \tau)}, \quad \frac{\theta \left[\begin{smallmatrix} -\frac{1}{5} \\ 1 \end{smallmatrix} \right] (z, \tau) \theta^2 \left[\begin{smallmatrix} \frac{3}{5} \\ 1 \end{smallmatrix} \right] (z, \tau)}{\theta^3 \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] (z, \tau)}.$$

The proposition can be proved using Theorem 5.1. \square

5.3 Third-order theta derivatives

Proposition 5.6. *For every $\tau \in \mathbb{H}^2$, we have*

$$\begin{aligned} & \frac{\theta^{(3)}\left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array}\right] + 3\theta^{(3)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right] + 9\theta^{(2)}\left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array}\right]\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right] + 18\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array}\right]\left\{\frac{\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}{\theta\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}\right\}^2 + 9\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array}\right]\theta^{(2)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}{\theta\left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array}\right]\theta\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]} \\ & + 6\left\{\frac{\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}{\theta\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}\right\}^3 + 18\frac{\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]\theta^{(2)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}{\theta\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]\theta\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]} - 4\frac{\theta^{(3)}\left[\begin{array}{c} 1 \\ 1 \end{array}\right]\left(\frac{\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array}\right]}{\theta\left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array}\right]} + 3\frac{\theta^{(1)}\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}{\theta\left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array}\right]}\right)}{\theta^{(1)}\left[\begin{array}{c} 1 \\ 1 \end{array}\right]} = 0, \end{aligned}$$

and

$$3 \frac{\theta^{(3)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} - \frac{\theta^{(3)} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} + 9 \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^{(2)} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} - 18 \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} \left\{ \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} \right\}^2 - 9 \frac{\theta^{(2)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} \\ + 6 \left\{ \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} \right\}^3 + 18 \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^{(2)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} - 4 \frac{\theta^{(3)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \left(3 \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} - \frac{\theta^{(1)} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} \right) = 0.$$

Proof. Consider the following elliptic functions:

$$\frac{\theta \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] (z) \theta^3 \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right] (z)}{\theta^4 \left[\begin{array}{c} 1 \\ 1 \end{array} \right] (z)}, \quad \frac{\theta^3 \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] (z) \theta \left[\begin{array}{c} 1 \\ -\frac{3}{5} \end{array} \right] (z)}{\theta^4 \left[\begin{array}{c} 1 \\ 1 \end{array} \right] (z)}.$$

Proposition 5.7. *For every $\tau \in \mathbb{H}^2$, we have*

$$\begin{aligned} & \frac{\theta^{(3)}\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]} + 3 \frac{\theta^{(3)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} + 9 \frac{\theta^{(2)}\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]} \frac{\theta^{(1)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} + 18 \frac{\theta^{(1)}\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]} \left\{ \frac{\theta^{(1)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} \right\}^2 + 9 \frac{\theta^{(1)}\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]} \frac{\theta^{(2)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} \\ & + 6 \left\{ \frac{\theta^{(1)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} \right\}^3 + 18 \frac{\theta^{(1)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} \frac{\theta^{(2)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} - 4 \frac{\theta^{(3)}\left[\begin{array}{c} 1 \\ 1 \end{array}\right]}{\theta^{(1)}\left[\begin{array}{c} 1 \\ 1 \end{array}\right]} \left(\frac{\theta^{(1)}\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{1}{5} \\ 1 \end{array}\right]} + 3 \frac{\theta^{(1)}\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]}{\theta\left[\begin{array}{c} \frac{3}{5} \\ 1 \end{array}\right]} \right) = 0, \end{aligned}$$

and

$$3 \frac{\theta^{(3)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} - \frac{\theta^{(3)} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} + 9 \frac{\theta^{(1)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^{(2)} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} - 18 \frac{\theta^{(1)} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} \left\{ \frac{\theta^{(1)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \right\}^2 - 9 \frac{\theta^{(2)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \theta^{(1)} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \\ + 6 \left\{ \frac{\theta^{(1)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \right\}^3 + 18 \frac{\theta^{(1)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^{(2)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} - 4 \frac{\theta^{(3)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \left(3 \frac{\theta^{(1)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} - \frac{\theta^{(1)} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} \right) = 0.$$

Proof. Furthermore, consider the following elliptic functions:

$$\frac{\theta \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] (z) \theta^3 \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right] (z)}{\theta^4 \left[\begin{array}{c} 1 \\ 1 \end{array} \right] (z)}, \quad \frac{\theta^3 \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] (z) \theta \left[\begin{array}{c} -1 \\ \frac{3}{5} \end{array} \right] (z)}{\theta^4 \left[\begin{array}{c} 1 \\ 1 \end{array} \right] (z)}.$$

1

5.4 Fourth-order theta derivatives

Proposition 5.8. *For every $\tau \in \mathbb{H}^2$ and $j = 1, 3$, we have*

$$\begin{aligned} & \frac{\theta^{(4)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]} + 16 \frac{\theta^{(1)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right] \theta^{(3)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]^2} + 12 \left\{ \frac{\theta^{(2)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]} \right\}^2 + 72 \left\{ \frac{\theta^{(1)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]} \right\}^2 \frac{\theta^{(2)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]} \\ & + 24 \left\{ \frac{\theta^{(1)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]} \right\}^4 - 40 \frac{\theta^{(3)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]} \left\{ \frac{\theta^{(1)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]} \right\}^2 - 10 \frac{\theta^{(3)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right] \theta^{(2)} \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right] \theta \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right]} \\ & - \frac{1}{5} \frac{\theta^{(5)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]} + 2 \left\{ \frac{\theta^{(3)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]} \right\}^2 = 0. \end{aligned}$$

Proof. Consider the following elliptic functions: $f_j(z) = \theta^5 \left[\begin{matrix} 1 \\ \frac{j}{5} \end{matrix} \right] (z, \tau) / \theta^5 \left[\begin{matrix} 1 \\ 1 \end{matrix} \right] (z, \tau)$.

□

Proposition 5.9. *For every $\tau \in \mathbb{H}^2$ and $j = 1, 3$, we have*

$$\begin{aligned} & \frac{\theta^{(4)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]} + 16 \frac{\theta^{(1)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right] \theta^{(3)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]^2} + 12 \left\{ \frac{\theta^{(2)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]} \right\}^2 + 72 \left\{ \frac{\theta^{(1)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]} \right\}^2 \frac{\theta^{(2)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]} \\ & + 24 \left\{ \frac{\theta^{(1)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]} \right\}^4 - 40 \frac{\theta^{(3)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]} \left\{ \frac{\theta^{(1)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]} \right\}^2 - 10 \frac{\theta^{(3)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right] \theta^{(2)} \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right] \theta \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right]} \\ & - \frac{1}{5} \frac{\theta^{(5)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]} + 2 \left\{ \frac{\theta^{(3)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]}{\theta^{(1)} \left[\begin{matrix} 1 \\ 1 \end{matrix} \right]} \right\}^2 = 0. \end{aligned}$$

Proof. Consider the following elliptic functions: $f_j(z) = \theta^5 \left[\begin{matrix} \frac{j}{5} \\ 1 \end{matrix} \right] (z, \tau) / \theta^5 \left[\begin{matrix} 1 \\ 1 \end{matrix} \right] (z, \tau)$.

□

6 Proof of Theorem 1.1

6.1 Notations

For every $\tau \in \mathbb{H}^2$, set $q = \exp(2\pi i\tau)$ and

$$X = \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}, \quad Y = \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}, \quad Z = \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}.$$

6.2 ODEs for $P(q)$ and $Q(q)$

Proof. Based on Proposition 5.4, we have

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} = \frac{1}{3}Z - 4XY - \frac{4}{3}X^2 + \frac{2}{3}Y^2, \quad \frac{\theta'' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = \frac{1}{3}Z + 4XY + \frac{2}{3}X^2 - \frac{4}{3}Y^2. \quad (6.1)$$

Based on Proposition 5.6, we obtain

$$\frac{\theta''' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} = 3X^3 + 9X^2Y - 15XY^2 + 3Y^3 + XZ, \quad \frac{\theta''' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = -3X^3 - 15X^2Y - 9XY^2 + 3Y^3 + YZ. \quad (6.2)$$

Because

$$\frac{\theta''' \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}} = 4\pi i \frac{d}{d\tau} \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}} \right\} + \frac{\theta' \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}} \cdot \frac{\theta'' \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{j}{5} \end{bmatrix}}, \quad (j = 1, 3),$$

it follows that

$$\begin{aligned} 4\pi i X' &= \frac{13}{3}X^3 + 13X^2Y - \frac{47}{3}XY^2 + 3Y^3 + \frac{2}{3}XZ, \\ 4\pi i Y' &= -3X^3 - \frac{47}{3}X^2Y - 13XY^2 + \frac{13}{3}Y^3 + \frac{2}{3}YZ, \end{aligned}$$

where $' = d/d\tau$. By considering $d/d\tau = 2\pi iq d/dq$, we obtain the ODEs for $P(q)$ and $Q(q)$. \square

6.3 ODE for $R(q)$

Proof. First, note that

$$\begin{aligned} \frac{\theta^{(4)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} &= 4\pi i \frac{d}{d\tau} \left\{ \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} \right\} + \left\{ \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} \right\}^2, \\ &= \frac{4\pi i}{3} \frac{dZ}{d\tau} + \frac{20}{9} X^4 + \frac{52}{3} X^3 Y + \frac{316}{9} X^2 Y^2 + \frac{44}{3} X Y^3 - \frac{52}{9} Y^4 \\ &\quad - 8 X Y Z - \frac{8}{3} X^2 Z + \frac{4}{3} Y^2 Z + \frac{1}{9} Z^2, \end{aligned}$$

and

$$\frac{\theta^{(5)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} = 4\pi i \frac{d}{d\tau} \left\{ \frac{\theta^{(3)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \right\} + \left\{ \frac{\theta^{(3)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \right\}^2 = 4\pi i \frac{dZ}{d\tau} + Z^2.$$

By using Proposition 5.8, we have

$$4\pi i Z' = \frac{10}{3} X^4 - 10 X^3 Y - \frac{310}{3} X^2 Y^2 + 10 X Y^3 + \frac{10}{3} Y^4 + \frac{2}{3} Z^2,$$

and

$$\frac{\theta^{(4)} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}} = \frac{10}{3} X^4 + 14 X^3 Y + \frac{2}{3} X^2 Y^2 + 18 X Y^3 - \frac{14}{3} X Y^3 + \frac{1}{3} Z^2 - 8 X Y Z - \frac{8}{3} X^2 Z + \frac{4}{3} Y^2 Z,$$

where $' = d/d\tau$. By considering $d/d\tau = 2\pi i q d/dq$, we obtain the ODE for $R(q)$. \square

6.4 Note on E_4 and E_6

Theorem 6.1. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} P(q) &= \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \\ Q(q) &= \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n. \end{aligned}$$

Then, we have

$$E_4(q) = -5P^4 + 15P^3Q + 155P^2Q^2 - 15PQ^3 - 5Q^4,$$

$$E_6(q) = \frac{-395P^6 - 3690P^5Q - 8025P^4Q^2 - 8025P^2Q^4 + 3690PQ^5 - 395Q^6}{8}.$$

Proof. For the proof, we use

$$\wp = \wp(z; 1, \tau) = \frac{1}{3} \frac{\theta'''}{\theta'} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{d^2}{dz^2} \log \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z).$$

Note that

$$\wp'' = 6\wp^2 - g_2, \quad g_2(1, \tau) = \frac{4\pi^4}{3} E_4(q).$$

By substituting $z = -2/5$, we have

$$g_2 = -\frac{20}{3}X^4 + 20X^3Y + \frac{620}{3}X^2Y^2 - 20XY^3 - \frac{20}{3}Y^4, \quad (6.3)$$

which implies

$$E_4(q) = -5P^4 + 15P^3Q + 155P^2Q^2 - 15PQ^3 - 5Q^4.$$

Next, note that

$$(\wp')^2 = 4\wp^3 - g_2\wp - g_3, \quad g_3(1, \tau) = \frac{8\pi^6}{27} E_6(q).$$

By substituting $z = -2/5$, we obtain

$$g_3 = -\frac{395}{27}X^6 - \frac{410}{3}X^5Y - \frac{2675}{9}X^4Y^2 - \frac{2675}{9}X^2Y^4 + \frac{410}{3}XY^5 - \frac{395}{27}Y^6, \quad (6.4)$$

which implies

$$E_6(q) = \frac{-395P^6 - 3690P^5Q - 8025P^4Q^2 - 8025P^2Q^4 + 3690PQ^5 - 395Q^6}{8}.$$

□

Corollary 6.2. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} S(q) &= \frac{\sqrt{250 - 110\sqrt{5}}}{5} \prod_{n=1}^{\infty} \frac{(1 - q^n)^5}{(1 - q^{5n})^3} \left(1 + \frac{1 - \sqrt{5}}{2} q^n + q^{2n} \right)^5 \\ &= \frac{\sqrt{250 - 110\sqrt{5}}}{5} \left\{ 1 - \frac{5}{2}(1 + \sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7 + 3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n \right\}, \\ T(q) &= \frac{\sqrt{250 + 110\sqrt{5}}}{5} \prod_{n=1}^{\infty} \frac{(1 - q^n)^5}{(1 - q^{5n})^3} \left(1 + \frac{1 + \sqrt{5}}{2} q^n + q^{2n} \right)^5 \\ &= \frac{\sqrt{250 + 110\sqrt{5}}}{5} \left\{ 1 + \frac{5}{2}(-1 + \sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7 - 3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n \right\}. \end{aligned}$$

Then we have

$$\begin{aligned} E_4(q) &= \frac{S^4 + 12S^3T + 14S^2T^2 - 12ST^3 + T^4}{16}, \\ E_6(q) &= -\frac{1}{64} (S^2 + T^2) (S^4 + 18S^3T + 74S^2T^2 - 18ST^3 + T^4), \end{aligned}$$

and

$$\begin{aligned} q(q : q)_{\infty}^{24} &= \frac{(E_4(q))^3 - (E_6(q))^2}{1728} = -\frac{S^5T^5(S^2 + 11ST - T^2)}{4096}, \\ j(q) &= -\frac{(S^4 + 12S^3T + 14S^2T^2 - 12ST^3 + T^4)^3}{S^5T^5(S^2 + 11ST - T^2)} = -\frac{(f^4 - 12f^3 + 14f^2 + 12f + 1)^3}{f^5(-f^2 + 11f + 1)}, \quad f = T/S. \end{aligned}$$

Proof. The corollary can be obtained by considering

$$S = -3P + Q, \quad T = P + 3Q, \quad \text{and } f = T/S.$$

□

Corollary 6.3. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} V(q) &= \prod_{n=1}^{\infty} \frac{(1-q^n)^5}{(1-q^{5n})^3} \left(1 + \frac{1-\sqrt{5}}{2}q^n + q^{2n}\right)^5 \\ &= 1 - \frac{5}{2}(1+\sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7+3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \\ W(q) &= \prod_{n=1}^{\infty} \frac{(1-q^n)^5}{(1-q^{5n})^3} \left(1 + \frac{1+\sqrt{5}}{2}q^n + q^{2n}\right)^5 \\ &= 1 + \frac{5}{2}(-1+\sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7-3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n. \end{aligned}$$

Then we have

$$\begin{aligned} E_4(q) &= \frac{(123 - 55\sqrt{5})V^4 + 12(-11 + 5\sqrt{5})V^3W + 28V^2W^2 - 12(11 + 5\sqrt{5})VW^3 + (123 + 55\sqrt{5})W^4}{10}, \\ E_6(q) &= \frac{1}{25}(-1525 + 682\sqrt{5})V^6 - \frac{9}{25}(-275 + 123\sqrt{5})V^5W + \frac{3}{2}(-25 + 11\sqrt{5})V^4W^2 \\ &\quad - \frac{3}{2}(25 + 11\sqrt{5})V^2W^4 + \frac{9}{25}(275 + 123\sqrt{5})VW^5 - \frac{1}{25}(1525 + 682\sqrt{5})W^6, \end{aligned}$$

and

$$\begin{aligned} q(q; q)_\infty^{24} &= -\frac{1}{250}V^5W^5(V-W)\left(\left(-11 + 5\sqrt{5}\right)V + \left(11 + 5\sqrt{5}\right)W\right) \\ &= \frac{1}{250}(11 - 5\sqrt{5})V^7W^5 - \frac{11}{125}V^6W^6 + \frac{1}{250}(11 + 5\sqrt{5})V^5W^7. \end{aligned}$$

Theorem 6.4. For every $\tau \in \mathbb{H}^2$, we have

$$G_4(\tau) = \frac{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^4}{720 \cdot \theta^{12} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^{12} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} (A^4 - 12A^3B + 14A^2B^2 + 12AB^3 + B^4),$$

and

$$G_6(\tau) = \frac{-\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^6}{30240 \cdot \theta^{18} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^{18} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} (A^2 + B^2)(A^4 - 18A^3B + 74A^2B^2 + 18AB^3 + B^4),$$

where

$$A = \theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}, \quad B = \theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}.$$

Proof. The theorem is based on equations (6.3) and (6.4) and the derivative formulas of Theorem 5.2. \square

Theorem 6.5. For every $\tau \in \mathbb{H}^2$, we have

$$\Delta = g_2^3 - 27g_3^2 = \frac{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^{12}}{\theta^{11} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^{11} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} \left(\theta^{10} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} - 11\theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} - \theta^{10} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} \right),$$

and

$$j = \frac{12^3 g_2^3}{\Delta} = \frac{(f^4(\tau) - 12f^3(\tau) + 14f^2(\tau) + 12f(\tau) + 1)^3}{f^5(\tau)(f^2(\tau) - 11f(\tau) - 1)},$$

where

$$f(\tau) = \theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} / \theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} = \frac{11 + 5\sqrt{5}}{2} \prod_{n=1}^{\infty} \frac{\left(1 + \frac{1+\sqrt{5}}{2}q^n + q^{2n}\right)^5}{\left(1 + \frac{1-\sqrt{5}}{2}q^n + q^{2n}\right)^5}, \quad q = \exp(2\pi i\tau).$$

Proof. By using equations (6.3) and (6.4), we have

$$\begin{aligned} \Delta &= -25(3X - Y)^5(X + 3Y)^5(X^2 + 4XY - Y^2) = -\frac{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^{12}}{\theta^{36} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^{36} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} A^5 B^5 (A^2 - 11AB - B^2), \\ j &= \frac{320(X^4 - 3X^3Y - 31X^2Y^2 + 3XY^3 + Y^4)^3}{27(3X - Y)^5(X + 3Y)^5(X^2 + 4XY - Y^2)} = \frac{(A^4 - 12A^3B + 14A^2B^2 + 12AB^3 + B^4)^3}{A^5 B^5 (A^2 - 11AB - B^2)}, \end{aligned}$$

where

$$A = \theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}, \quad B = \theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}.$$

\square

Corollary 6.6. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} P(q) &= \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \\ Q(q) &= \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n. \end{aligned}$$

Then, we have

$$j(\tau) = j(q) = \frac{320(P^4 - 3P^3Q - 31P^2Q^2 + 3PQ^3 + Q^4)^3}{27(3P - Q)^5(P + 3Q)^5(P^2 + 4PQ - Q^2)},$$

and

$$f(\tau) = \theta^5 \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] / \theta^5 \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right] = -\frac{P + 3Q}{3P - Q}, \quad q = \exp(2\pi i\tau).$$

Proof. The expression of $f(\tau)$ follows from the derivative formulas (5.1) and (5.2), which implies that

$$f(\tau) = -\frac{X + 3Y}{3X - Y}.$$

□

Theorem 6.7. For every $\tau \in \mathbb{H}^2$, we have

$$\theta' \left[\begin{array}{c} 1 \\ 1 \end{array} \right]^4 = \frac{(2\pi)^4 \theta^{11} \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] \theta^{11} \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right]}{\theta^{10} \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] - 11\theta^5 \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right] \theta^5 \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right] - \theta^{10} \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right]}. \quad (6.5)$$

Proof. Recall the following formula from Apostol [1, pp. 51]:

$$\Delta = (2\pi)^{12} \eta^{24}(\tau) = (2\pi)^4 \theta' \left[\begin{array}{c} 1 \\ 1 \end{array} \right]^8. \quad (6.6)$$

The theorem follows from Theorem 6.5. □

Theorem 6.8.

- (1) For each positive integer k , the Eisenstein series $G_{4k}(\tau)$ can be expressed using a rational expression as follows:

$$\theta \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right](0, \tau), \text{ and } \theta \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right](0, \tau).$$

- (2) For each positive integer k , the Eisenstein series $G_{4k+2}(\tau)$ can be expressed using a rational expression as follows:

$$\theta \left[\begin{array}{c} 1 \\ \frac{1}{5} \end{array} \right](0, \tau), \theta \left[\begin{array}{c} 1 \\ \frac{3}{5} \end{array} \right](0, \tau), \text{ and } \theta' \left[\begin{array}{c} 1 \\ 1 \end{array} \right](0, \tau).$$

Proof. From the study by Apostol [1, pp. 118], we recall that for $k = 2, 3, 4 \dots$,

$$G_{2k}(\tau) = \sum_{a,b} c_{a,b} G_4^a(\tau) G_6^b(\tau),$$

where $c_{a,b}$ are complex numbers, and the sum is extended over all integers $a \geq 0, b \geq 0$ such that $4a + 6b = k$. The theorem follows from Theorems 6.4 and 6.7. \square

6.5 Proof of Ramanujan's ODEs (1.3)

Theorem 6.9. For $q \in \mathbb{C}$ with $|q| < 1$, we have

$$q \frac{dE_2}{dq} = \frac{(E_2)^2 - E_4}{12}, \quad q \frac{dE_4}{dq} = \frac{E_2 E_4 - E_6}{3}, \quad q \frac{dE_6}{dq} = \frac{E_2 E_6 - (E_4)^2}{2}.$$

Proof. According to Theorems 1.1 and 6.1,

$$q \frac{d}{dq} E_2 = q \frac{d}{dq} R = \frac{5P^4 - 15P^3Q - 155P^2Q^2 + 15PQ^3 + 5Q^4 + R^2}{12} = \frac{E_2^2 - E_4}{12}.$$

Next, it is observed that

$$\begin{aligned} q \frac{d}{dq} E_4 &= q \frac{d}{dq} (-5P^4 + 15P^3Q + 155P^2Q^2 - 15PQ^3 - 5Q^4) \\ &= \frac{395P^6}{24} + \frac{615P^5Q}{4} + \frac{2675P^4Q^2}{8} + \frac{2675P^2Q^4}{8} - \frac{615PQ^5}{4} + \frac{395Q^6}{24} \\ &\quad - \frac{5P^4R}{3} + 5P^3QR + \frac{155}{3}P^2Q^2R - 5PQ^3R - \frac{5Q^4R}{3} \\ &= \frac{E_2 E_4 - E_6}{3}. \end{aligned}$$

Finally, we obtain

$$\begin{aligned} q \frac{d}{dq} E_6 &= q \frac{d}{dq} \left\{ \frac{-395P^6 - 3690P^5Q - 8025P^4Q^2 - 8025P^2Q^4 + 3690PQ^5 - 395Q^6}{8} \right\} \\ &= -\frac{25P^8}{2} + 75P^7Q + \frac{1325P^6Q^2}{2} - 2400P^5Q^3 - \frac{23625P^4Q^4}{2} \\ &\quad + 2400P^3Q^5 + \frac{1325P^2Q^6}{2} - 75PQ^7 - \frac{25Q^8}{2} \\ &\quad - \frac{395P^6R}{16} - \frac{1845}{8}P^5QR - \frac{8025}{16}P^4Q^2R - \frac{8025}{16}P^2Q^4R + \frac{1845}{8}PQ^5R - \frac{395Q^6R}{16} \\ &= \frac{E_2 E_6 - (E_4)^2}{2}. \end{aligned}$$

\square

7 Proof of Theorem 1.2

7.1 Notations

For every $\tau \in \mathbb{H}^2$, set $y = \exp(2\pi i\tau/5)$ and

$$X = \frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}, \quad Y = \frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}, \quad Z = \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}.$$

7.2 ODEs for $P(q)$ and $Q(q)$

Proof. According to Proposition 5.5, we have

$$\frac{\theta'' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} = \frac{1}{3}Z - 4XY - \frac{4}{3}X^2 + \frac{2}{3}Y^2, \quad \frac{\theta'' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} = \frac{1}{3}Z + 4XY + \frac{2}{3}X^2 - \frac{4}{3}Y^2. \quad (7.1)$$

According to Proposition 5.7, we obtain

$$\frac{\theta''' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} = 3X^3 + 9X^2Y - 15XY^2 + 3Y^3 + XZ, \quad \frac{\theta''' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} = -3X^3 - 15X^2Y - 9XY^2 + 3Y^3 + YZ. \quad (7.2)$$

Because

$$\frac{\theta''' \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}} = 4\pi i \frac{d}{d\tau} \left\{ \frac{\theta' \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}} \right\} + \frac{\theta' \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}} \cdot \frac{\theta'' \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{j}{5} \\ 1 \end{bmatrix}}, \quad (j = 1, 3),$$

we get

$$4\pi i X' = \frac{13}{3}X^3 + 13X^2Y - \frac{47}{3}XY^2 + 3Y^3 + \frac{2}{3}XZ,$$

$$4\pi i Y' = -3X^3 - \frac{47}{3}X^2Y - 13XY^2 + \frac{13}{3}Y^3 + \frac{2}{3}YZ,$$

where $' = d/d\tau$. Thus, by changing $\tau \rightarrow 5\tau$, we obtain the ODEs for $P(q)$ and $Q(q)$. \square

7.3 ODE for $R(q)$

Proof. First, note that

$$\begin{aligned} \frac{\theta^{(4)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} &= 4\pi i \frac{d}{d\tau} \left\{ \frac{\theta'' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \right\} + \left\{ \frac{\theta'' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \right\}^2, \\ &= \frac{4\pi i}{3} \frac{dZ}{d\tau} + \frac{20}{9} X^4 + \frac{52}{3} X^3 Y + \frac{316}{9} X^2 Y^2 + \frac{44}{3} X Y^3 - \frac{52}{9} Y^4 \\ &\quad - 8 X Y Z - \frac{8}{3} X^2 Z + \frac{4}{3} Y^2 Z + \frac{1}{9} Z^2, \end{aligned}$$

and

$$\frac{\theta^{(5)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} = 4\pi i \frac{d}{d\tau} \left\{ \frac{\theta^{(3)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \right\} + \left\{ \frac{\theta^{(3)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta^{(1)} \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \right\}^2 = 4\pi i \frac{dZ}{d\tau} + Z^2.$$

Thus, according to Proposition 5.9, we have

$$4\pi i Z' = \frac{10}{3} X^4 - 10 X^3 Y - \frac{310}{3} X^2 Y^2 + 10 X Y^3 + \frac{10}{3} Y^4 + \frac{2}{3} Z^2,$$

and

$$\frac{\theta^{(4)} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} = \frac{10}{3} X^4 + 14 X^3 Y + \frac{2}{3} X^2 Y^2 + 18 X Y^3 - \frac{14}{3} X Y^3 + \frac{1}{3} Z^2 - 8 X Y Z - \frac{8}{3} X^2 Z + \frac{4}{3} Y^2 Z,$$

where $' = d/d\tau$. Thus, by changing $\tau \rightarrow 5\tau$, we obtain the ODE for $R(q)$. \square

Theorem 7.1. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$P(q) = 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \quad Q(q) = 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n.$$

Then, we have

$$\begin{aligned} E_4(q^5) &= \frac{-P^4 + 3P^3Q + 31P^2Q^2 - 3PQ^3 - Q^4}{125}, \\ E_6(q^5) &= \frac{79P^6 + 738P^5Q + 1605P^4Q^2 + 1605P^2Q^4 - 738PQ^5 + 79Q^6}{25000}. \end{aligned}$$

Proof. For the proof, we use

$$\wp = \wp(z; 1, \tau) = \frac{1}{3} \frac{\theta'''}{\theta'} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{d^2}{dz^2} \log \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z).$$

First, note that

$$\wp'' = 6\wp^2 - g_2, \quad g_2(1, \tau) = \frac{4\pi^4}{3} E_4(q).$$

Then, by substituting $z = -2\tau/5$, we have

$$g_2 = -\frac{20}{3}X^4 + 20X^3Y + \frac{620}{3}X^2Y^2 - 20XY^3 - \frac{20}{3}Y^4, \quad (7.3)$$

which implies

$$E_4(q^5) = \frac{-P^4 + 3P^3Q + 31P^2Q^2 - 3PQ^3 - Q^4}{125}.$$

Next, it is observed that

$$(\wp')^2 = 4\wp^3 - g_2\wp - g_3, \quad g_3(1, \tau) = \frac{8\pi^6}{27} E_6(q).$$

Thus, by substituting $z = -2\tau/5$, we obtain

$$g_3 = -\frac{395}{27}X^6 - \frac{410}{3}X^5Y - \frac{2675}{9}X^4Y^2 - \frac{2675}{9}X^2Y^4 + \frac{410}{3}XY^5 - \frac{395}{27}Y^6, \quad (7.4)$$

which implies

$$E_6(q^5) = \frac{79P^6 + 738P^5Q + 1605P^4Q^2 + 1605P^2Q^4 - 738PQ^5 + 79Q^6}{25000}.$$

□

Corollary 7.2. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$S(q) = q \prod_{n=1}^{\infty} \frac{(1-q^n)^2}{(1-q^{5n-2})^5(1-q^{5n-3})^5} = \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n - 3 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n,$$

and

$$T(q) = \prod_{n=1}^{\infty} \frac{(1-q^n)^2}{(1-q^{5n-1})^5(1-q^{5n-4})^5} = 1 + 3 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n.$$

Then, we have

$$E_4(q^5) = S^4 + 12S^3T + 14S^2T^2 - 12ST^3 + T^4,$$

$$E_6(q^5) = (S^2 + T^2)(S^4 + 18S^3T + 74S^2T^2 - 18ST^3 + T^4),$$

and

$$q^5(q^5; q^5)_\infty = -S^5T^5(S^2 + 11ST - T^2)$$

$$j(q^5) = -\frac{(S^4 + 12S^3T + 14S^2T^2 - 12ST^3 + T^4)^3}{S^5T^5(S^2 + 11ST - T^2)} = -\frac{(g^4 + 12g^3 + 14g^2 - 12g + 1)^3}{g^5(g^2 + 11g - 1)}, \quad g = S/T.$$

Proof. The corollary can be obtained by considering

$$S = -\frac{3}{10}P + \frac{1}{10}Q, \quad T = \frac{1}{10}P + \frac{3}{10}Q.$$

□

Theorem 7.3. For every $\tau \in \mathbb{H}^2$, we have

$$G_4(\tau) = \frac{\zeta_5^2 \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^4}{720 \cdot \theta^{12} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^{12} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} (A^4 + 12A^3B + 14A^2B^2 - 12AB^3 + B^4),$$

and

$$G_6(\tau) = \frac{-\zeta_5^3 \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^6}{30240 \cdot \theta^{18} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^{18} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} (A^2 + B^2)(A^4 + 18A^3B + 74A^2B^2 - 18AB^3 + B^4),$$

where $\zeta_5 = \exp(2\pi i/5)$, and

$$A = \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}, \quad B = \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}.$$

Proof. The theorem follows from equations (7.3) and (7.4) and the derivative formulas of Theorem 5.3. □

Theorem 7.4. For every $\tau \in \mathbb{H}^2$, we have

$$\Delta(\tau) = g_2^3 - 27g_3^2 = \frac{\zeta_5 \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^{12}}{\theta^{11} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^{11} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} \left(\theta^{10} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - 11\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} - \theta^{10} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \right),$$

and

$$j(5\tau) = \frac{12^3 g_2(5\tau)^3}{\Delta(5\tau)} = \frac{(g^4(\tau) - 12g^3(\tau) + 14g^2(\tau) + 12g(\tau) + 1)^3}{g^5(\tau)(g^2(\tau) - 11g(\tau) - 1)},$$

where $\zeta_5 = \exp(2\pi i/5)$,

$$g(\tau) = \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, 5\tau) / \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, 5\tau) = -q \prod_{n=1}^{\infty} \frac{(1 - q^{5n-1})^5 (1 - q^{5n-4})^5}{(1 - q^{5n-2})^5 (1 - q^{5n-3})^5}, \quad q = \exp(2\pi i\tau).$$

Proof. Based on equations (7.3) and (7.4), we have

$$\begin{aligned} \Delta &= -25(3X - Y)^5(X + 3Y)^5(X^2 + 4XY - Y^2) = \frac{-\zeta_5 \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^{12}}{\theta^{36} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^{36} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} A^5 B^5 (A^2 + 11AB - B^2), \\ j &= \frac{320(X^4 - 3X^3Y - 31X^2Y^2 + 3XY^3 + Y^4)^3}{27(3X - Y)^5(X + 3Y)^5(X^2 + 4XY - Y^2)} = -\frac{(A^4 + 12A^3B + 14A^2B^2 - 12AB^3 + B^4)^3}{A^5 B^5 (A^2 + 11AB - B^2)}, \end{aligned}$$

where

$$A = \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}, \quad B = \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}.$$

□

Corollary 7.5. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$P(q) = 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \quad Q(q) = 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n.$$

Then, we have

$$j(5\tau) = j(q^5) = \frac{320(P^4 - 3P^3Q - 31P^2Q^2 + 3PQ^3 + Q^4)^3}{27(3P - Q)^5(P + 3Q)^5(P^2 + 4PQ - Q^2)},$$

and

$$g(\tau) = \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, 5\tau) / \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, 5\tau) = \frac{3P - Q}{P + 3Q}, \quad q = \exp(2\pi i\tau).$$

Proof. The expression of $g(\tau)$ follows from the derivative formulas (5.3) and (5.4), which imply that

$$g(\tau/5) = \frac{3X - Y}{X + 3Y}.$$

□

Theorem 7.6. For every $\tau \in \mathbb{H}^2$, set

$$f(\tau) = \frac{\theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} (0, \tau)}{\theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} (0, \tau)} = \frac{11 + 5\sqrt{5}}{2} \prod_{n=1}^{\infty} \frac{\left(1 + \frac{1+\sqrt{5}}{2}q^n + q^{2n}\right)^5}{\left(1 + \frac{1-\sqrt{5}}{2}q^n + q^{2n}\right)^5},$$

and

$$g(\tau) = \frac{\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, 5\tau)}{\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, 5\tau)} = -q \prod_{n=1}^{\infty} \frac{(1 - q^{5n-1})^5 (1 - q^{5n-4})^5}{(1 - q^{5n-2})^5 (1 - q^{5n-3})^5}, \quad q = \exp(2\pi i\tau).$$

Then, we have

$$g(\tau) = \frac{\frac{1}{2}(-11 + 5\sqrt{5})f(\tau) - 1}{-f(\tau) - \frac{1}{2}(-11 + 5\sqrt{5})}. \quad (7.5)$$

Proof. We first set

$$P_2(q) := 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \quad Q_2(q) := 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n,$$

which implies that

$$\begin{aligned} P_1(q) &:= \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \\ &= -\frac{2}{5} \sin \frac{2\pi}{5} P_2(q) + \frac{2}{5} \sin \frac{4\pi}{5} Q_2(q), \\ Q_1(q) &:= \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \\ &= \frac{2}{5} \sin \frac{4\pi}{5} P_2(q) + \frac{2}{5} \sin \frac{2\pi}{5} Q_2(q). \end{aligned}$$

The theorem is derived from Corollaries 6.6 and 7.5. □

Theorem 7.7. For every $\tau \in \mathbb{H}^2$, we have

$$\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^4 = \frac{-(2\pi)^4 \zeta_5^4 \theta^{11} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^{11} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta^{10} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} + 11\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - \theta^{10} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}, \quad (7.6)$$

where $\zeta_5 = \exp(2\pi i/5)$.

Proof. The theorem is derived from formula (6.6) and Theorem 7.4. \square

Theorem 7.8.

- (1) For each positive integer k , the Eisenstein series $G_{4k}(\tau)$ can be expressed using a rational expression

$$\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, \tau), \text{ and } \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, \tau).$$

- (2) For each positive integer k , the Eisenstein series $G_{4k+2}(\tau)$ can be expressed using a rational expression

$$\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, \tau), \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, \tau), \text{ and } \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} (0, \tau).$$

Proof. The theorem can be proved in the same way as Theorem 6.8. \square

8 Some product-series identities

8.1 Some theta functional formulas

Proposition 8.1. For every $(z, \tau) \in \mathbb{C} \times \mathbb{H}^2$, we have

$$\begin{aligned} & \theta^2 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (z) \theta \begin{bmatrix} \frac{1}{9} \\ \frac{9}{5} \end{bmatrix} (z) - \theta^2 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{3} \\ \frac{3}{5} \end{bmatrix} (z) \theta \begin{bmatrix} \frac{1}{7} \\ \frac{5}{5} \end{bmatrix} (z) \\ & + \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{3} \\ \frac{3}{5} \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z) = 0, \end{aligned} \quad (8.1)$$

$$\begin{aligned} & -\zeta_5^2 \theta^2 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (z) \theta \begin{bmatrix} \frac{9}{5} \\ 1 \end{bmatrix} (z) + \zeta_5^3 \theta^2 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (z) \theta \begin{bmatrix} \frac{7}{5} \\ 1 \end{bmatrix} (z) \\ & + \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z) = 0, \end{aligned} \quad (8.2)$$

where $\zeta_5 = \exp(2\pi i/5)$.

Proof. Here, we provide the proof of equation (8.1). Equation (8.2) can be proved in the same manner. First, note that $\dim \mathcal{F}_2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} = 2$, and

$$\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{9}{5} \end{bmatrix} (z, \tau), \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{7}{5} \end{bmatrix} (z, \tau), \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau) \in \mathcal{F}_2 \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Therefore, there exist some complex numbers, x_1, x_2 , and x_3 , which are not all zeros, such that

$$x_1 \theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{9}{5} \end{bmatrix} (z, \tau) + x_2 \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{7}{5} \end{bmatrix} (z, \tau) + x_3 \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau) = 0.$$

Note that in the fundamental parallelogram, the zero of $\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} (z)$, $\theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} (z)$, or $\theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)$ is $z = 2/5, 1/5$ or 0 . By substituting $z = 2/5, 1/5$, and 0 , we have

$$\begin{aligned} x_2 \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} + x_3 \theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} &= 0, \\ -x_1 \theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} + x_3 \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} &= 0, \\ -x_1 \theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} - x_2 \theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} &= 0. \end{aligned}$$

By solving this system of equations, we get

$$(x_1, x_2, x_3) = \alpha \left(\theta^2 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}, -\theta^2 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}, \theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} \right) \text{ for some } \alpha \in \mathbb{C} \setminus \{0\},$$

which proves the proposition. \square

Lemma 8.2. *For every $(z, \tau) \in \mathbb{C} \times \mathbb{H}^2$, we have*

$$\left\{ \frac{\theta' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)}{\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)} \right\}^2 = \frac{\theta'' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)}{\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)} - \frac{d^2}{dz^2} \log \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z).$$

Proof. The lemma can be proved by direct calculation. \square

Theorem 8.3. For every $\tau \in \mathbb{H}^2$, we have

$$\frac{\eta^5(\tau)}{\eta(5\tau)} = \frac{(q; q)_\infty^5}{(q^5; q^5)_\infty} = 1 - 5 \sum_{n=1}^{\infty} \left(\sum_{d|n} d \left(\frac{d}{5} \right) \right) q^n, \quad (8.3)$$

$$\frac{\eta^5(5\tau)}{\eta(\tau)} = q \frac{(q^5; q^5)_\infty^5}{(q; q)_\infty} = \sum_{n=1}^{\infty} \left(\sum_{d|n} \frac{n}{d} \left(\frac{d}{5} \right) \right) q^n, \quad q = \exp(2\pi i\tau), \quad (8.4)$$

where for each $m \in \mathbb{N}$,

$$\left(\frac{m}{5} \right) = \begin{cases} 1 & \text{if } m \equiv \pm 1 \pmod{5}, \\ -1 & \text{if } m \equiv \pm 2 \pmod{5}, \\ 0 & \text{if } m \equiv 0 \pmod{5}. \end{cases}$$

Proof. By comparing the coefficients of term z^2 in equations (8.1) and (8.2), we have

$$\frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = \frac{d^2}{dz^2} \log \theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix}(z) \Big|_{z=0} - \frac{d^2}{dz^2} \log \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}(z) \Big|_{z=0}, \quad (8.5)$$

and

$$\zeta_5 \frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} = \frac{d^2}{dz^2} \log \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}(z) \Big|_{z=0} - \frac{d^2}{dz^2} \log \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}(z) \Big|_{z=0}. \quad (8.6)$$

Therefore, the theorem can be obtained using Jacobi's triple product identity (2.5). \square

Theorem 8.4. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$P(q) = \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n,$$

$$Q(q) = \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n.$$

Then, we have

$$\frac{(q; q)_\infty^5}{(q^5; q^5)_\infty} = -\frac{\sqrt{5}}{4} (3P - Q)(P + 3Q). \quad (8.7)$$

Proof. The theorem follows from equations (6.1) and (8.5). \square

Theorem 8.5. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$P(q) = 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \quad Q(q) = 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n.$$

Then, we have

$$q \frac{(q^5; q^5)_\infty^5}{(q; q)_\infty} = -\frac{1}{100} (3P - Q)(P + 3Q), \quad (8.8)$$

and

$$\frac{(q; q)_\infty^5}{(q^5; q^5)_\infty} = \frac{1}{4} (P^2 + 4PQ - Q^2). \quad (8.9)$$

Proof. The theorem can be proved in the same way as Theorem 8.4. \square

9 Some theorem of Farkas and Kra

Theorem 9.1. (Farkas and Kra [5, pp. 318]) For every $\tau \in \mathbb{H}^2$, we have

$$\frac{d}{d\tau} \log \left(\frac{\eta(5\tau)}{\eta(\tau)} \right) + \frac{1}{2\pi i \cdot 3} \left[\left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] (0, \tau)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] (0, \tau)} \right\}^2 + \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] (0, \tau)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] (0, \tau)} \right\}^2 \right] = 0.$$

Proof. Summing both sides of equations (5.7) and (5.8) yields

$$3 \frac{\theta'' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} + 3 \frac{\theta'' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]} - 2 \frac{\theta''' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]}{\theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right]} + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} \right\}^2 + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]} \right\}^2 = 0.$$

The heat equation, Eq. (2.6), implies that

$$4\pi i \frac{d}{d\tau} \log \frac{\theta^3 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta^3 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\left\{ \theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \right\}^2} + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]} \right\}^2 + 2 \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}{\theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]} \right\}^2 = 0.$$

Jacobi's triple product identity (2.5) yields

$$\frac{\theta^3 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta^3 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right]}{\left\{ \theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \right\}^2} = \frac{(\sqrt{5})^3}{4\pi^2} \frac{\eta^3(5\tau)}{\eta^3(\tau)},$$

which proves the theorem. \square

Corollary 9.2. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$P(q) = \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n,$$

$$Q(q) = \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n.$$

Then, we have

$$P^2 + Q^2 = \frac{1}{2} \{ -E_2(q) + 5E_2(q^5) \}. \quad (9.1)$$

Theorem 9.3. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$P(q) = \cot \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n - 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n,$$

$$Q(q) = \cot \frac{\pi}{5} + 4 \sin \frac{2\pi}{5} \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n + 4 \sin \frac{4\pi}{5} \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n.$$

Then, we have

$$E_4(q^5) = P^4 + 3P^3Q - P^2Q^2 - 3PQ^3 + Q^4,$$

$$E_6(q^5) = \frac{1}{40} (P^2 + Q^2) (41P^4 + 198P^3Q + 154P^2Q^2 - 198PQ^3 + 41Q^4).$$

Proof. Changing $q \rightarrow q^5$ in Ramanujan's system (1.3), we have

$$q \frac{d}{dq} E_2(q^5) = \frac{5}{12} \{ (E_2(q^5))^2 - E_4(q^5) \}, \quad q \frac{d}{dq} E_4(q^5) = \frac{5}{3} \{ E_2(q^5) E_4(q^5) - E_6(q^5) \}.$$

From Corollary 9.2, we have

$$E_2(q^5) = \frac{1}{5} (2P^2 + 2Q^2 + R), \quad R = E_2(q),$$

which implies that

$$q \frac{d}{dq} E_2(q^5) = -\frac{7P^4}{20} - \frac{5P^3Q}{4} + \frac{11P^2Q^2}{20} + \frac{5PQ^3}{4} - \frac{7Q^4}{20} + \frac{P^2R}{15} + \frac{Q^2R}{15} + \frac{R^2}{60}.$$

Therefore, it follows that

$$E_4(q^5) = (E_2(q^5))^2 - \frac{12}{5}q \frac{d}{dq} E_2(q^5) = P^4 + 3P^3Q - P^2Q^2 - 3PQ^3 + Q^4.$$

In the same way, we note that

$$\begin{aligned} q \frac{d}{dq} E_4(q^5) &= -\frac{25P^6}{24} - \frac{25P^5Q}{4} - \frac{65P^4Q^2}{8} - \frac{65P^2Q^4}{8} + \frac{25PQ^5}{4} - \frac{25Q^6}{24} \\ &\quad + P^3QR - \frac{1}{3}P^2Q^2R - PQ^3R + \frac{P^4R}{3} + \frac{Q^4R}{3}, \end{aligned}$$

which implies that

$$\begin{aligned} E_6(q^5) &= E_2(q^5)E_4(q^5) - \frac{3}{5}q \frac{d}{dq} E_4(q^5) \\ &= \frac{1}{40} (P^2 + Q^2) (41P^4 + 198P^3Q + 154P^2Q^2 - 198PQ^3 + 41Q^4). \end{aligned}$$

□

Corollary 9.4. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} S(q) &= \frac{\sqrt{250 - 110\sqrt{5}}}{5} \prod_{n=1}^{\infty} \frac{(1 - q^n)^5}{(1 - q^{5n})^3} \left(1 + \frac{1 - \sqrt{5}}{2}q^n + q^{2n} \right)^5 \\ &= \frac{\sqrt{250 - 110\sqrt{5}}}{5} \left\{ 1 - \frac{5}{2}(1 + \sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7 + 3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n \right\}, \\ T(q) &= \frac{\sqrt{250 + 110\sqrt{5}}}{5} \prod_{n=1}^{\infty} \frac{(1 - q^n)^5}{(1 - q^{5n})^3} \left(1 + \frac{1 + \sqrt{5}}{2}q^n + q^{2n} \right)^5 \\ &= \frac{\sqrt{250 + 110\sqrt{5}}}{5} \left\{ 1 + \frac{5}{2}(-1 + \sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7 - 3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n \right\}. \end{aligned}$$

Then we have

$$\begin{aligned} E_4(q^5) &= \frac{S^4 - 228S^3T + 494S^2T^2 + 228ST^3 + T^4}{10000} \\ E_6(q^5) &= - \frac{(S^2 + T^2)(S^4 + 522S^3T - 10006S^2T^2 - 522ST^3 + T^4)}{1000000}, \end{aligned}$$

and

$$\begin{aligned} q^5(q^5 : q^5)_\infty^{24} &= \frac{(E_4(q^5))^3 - (E_6(q^5))^2}{1728} = -\frac{ST(S^2 + 11ST - T^2)^5}{10000000000000} \\ j(q^5) &= -\frac{(S^4 - 228S^3T + 494S^2T^2 + 228ST^3 + T^4)^3}{ST(S^2 + 11ST - T^2)^5} \\ &= -\frac{(f^4 + 228f^3 + 494f^2 - 228f + 1)^3}{f(-f^2 + 11f + 1)^5}, \quad f = T/S. \end{aligned}$$

Proof. The corollary can be obtained by considering

$$S = -3P + Q, \quad T = P + 3Q, \quad \text{and} \quad f = T/S.$$

□

Corollary 9.5. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$\begin{aligned} V(q) &= \prod_{n=1}^{\infty} \frac{(1-q^n)^5}{(1-q^{5n})^3} \left(1 + \frac{1-\sqrt{5}}{2}q^n + q^{2n}\right)^5 \\ &= 1 - \frac{5}{2}(1+\sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7+3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \\ W(q) &= \prod_{n=1}^{\infty} \frac{(1-q^n)^5}{(1-q^{5n})^3} \left(1 + \frac{1+\sqrt{5}}{2}q^n + q^{2n}\right)^5 \\ &= 1 + \frac{5}{2}(-1+\sqrt{5}) \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \frac{5}{2}(7-3\sqrt{5}) \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n. \end{aligned}$$

Then we have

$$\begin{aligned} E_4(q^5) &= \frac{1}{6250} \times \\ &\times \left\{ \left(123 - 55\sqrt{5}\right)V^4 + 228\left(11 - 5\sqrt{5}\right)V^3W + 988V^2W^2 + 228\left(11 + 5\sqrt{5}\right)VW^3 + \left(123 + 55\sqrt{5}\right)W^4 \right\}, \\ E_6(q^5) &= -\frac{\left((11\sqrt{5} - 25)V^2 - (25 + 11\sqrt{5})W^2\right)}{1562500} \times \\ &\times \left\{ \left(55\sqrt{5} - 123\right)V^4 - 522\left(5\sqrt{5} - 11\right)V^3W + 20012V^2W^2 \right. \\ &\quad \left. + 522\left(11 + 5\sqrt{5}\right)VW^3 - \left(123 + 55\sqrt{5}\right)W^4 \right\} \end{aligned}$$

and

$$q^5(q^5; q^5)_\infty^{24} = -\frac{VW(V-W)^5}{61035156250} \times \\ \times \left\{ \left(75025\sqrt{5} - 167761 \right) V^5 + 10 \left(305\sqrt{5} - 682 \right) V^4 W + 10 \left(5\sqrt{5} - 11 \right) V^3 W^2 \right. \\ \left. + 10 \left(11 + 5\sqrt{5} \right) V^2 W^3 + 10 \left(682 + 305\sqrt{5} \right) V W^4 + \left(167761 + 75025\sqrt{5} \right) W^5 \right\}.$$

Theorem 9.6. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \left(\frac{\eta(\tau/5)}{\eta(\tau)} \right) + \frac{1}{2\pi i \cdot 3} \left[\left\{ \frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, \tau)} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, \tau)} \right\}^2 \right] = 0.$$

Proof. Summing both sides of equations (5.9) and (5.10) yields

$$3 \frac{\theta'' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} + 3 \frac{\theta'' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} - 2 \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \right\}^2 + 2 \left\{ \frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} \right\}^2 = 0.$$

The heat equation (2.6) implies that

$$4\pi i \frac{d}{d\tau} \log \frac{\theta^3 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^3 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2} + 2 \left\{ \frac{\theta' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \right\}^2 + 2 \left\{ \frac{\theta' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} \right\}^2 = 0.$$

Jacobi's triple product identity (2.5) yields

$$\frac{\theta^3 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^3 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}}{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2} = \frac{\zeta_5^3}{4\pi^2} \frac{\eta^3(\tau/5)}{\eta^3(\tau)},$$

which proves the theorem. \square

Corollary 9.7. *For $q \in \mathbb{C}$ with $|q| < 1$, set*

$$P(q) = 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n)) q^n, \quad Q(q) = 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n)) q^n.$$

Then, we have

$$P^2 + Q^2 = \frac{5}{2} \{-E_2(q) + 5E_2(q^5)\}. \quad (9.2)$$

Theorem 9.8. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$P(q) = 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \quad Q(q) = 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n.$$

Then, we have

$$\begin{aligned} E_4(q) &= P^4 + 3P^3Q - P^2Q^2 - 3PQ^3 + Q^4, \\ E_6(q) &= -\frac{1}{40} (P^2 + Q^2) (41P^4 + 198P^3Q + 154P^2Q^2 - 198PQ^3 + 41Q^4), \end{aligned}$$

and

$$\begin{aligned} 1728q(q;q)_\infty^{24} &= (E_4(q))^3 - (E_6(q))^2 = -\frac{27(3P - Q)(P + 3Q)(P^2 + 4PQ - Q^2)^5}{1600}, \\ j(q) &= \frac{1728(E_4(q))^3}{(E_4(q))^3 - (E_6(q))^2} = -\frac{102400 (P^4 + 3P^3Q - P^2Q^2 - 3PQ^3 + Q^4)^3}{(3P - Q)(P + 3Q)(P^2 + 4PQ - Q^2)^5} \\ &= \frac{(g^4 + 228g^3 + 494g^2 - 228g + 1)^3}{g(g^2 - 11g - 1)^5}, \end{aligned}$$

where

$$g(q) = \frac{3P - Q}{P + 3Q} = -q \prod_{n=1}^{\infty} \frac{(1 - q^{5n-1})^5(1 - q^{5n-4})^5}{(1 - q^{5n-2})^5(1 - q^{5n-3})^5}.$$

Proof. From Corollary 9.7, we have

$$E_2(q) = \frac{-2P^2 - 2Q^2 + 25R}{5}, \quad R = E_2(q^5),$$

which implies that

$$q \frac{d}{dq} E_2(q) = -\frac{7P^4}{100} - \frac{P^3Q}{4} + \frac{11P^2Q^2}{100} + \frac{PQ^3}{4} - \frac{7Q^4}{100} - \frac{P^2R}{3} - \frac{Q^2R}{3} + \frac{25R^2}{12}.$$

By Ramanujan's system (1.3), we obtain

$$E_4(q) = \{E_2(q)\}^2 - 12q \frac{d}{dq} E_2(q) = P^4 + 3P^3Q - P^2Q^2 - 3PQ^3 + Q^4.$$

In the same way, we note that

$$\begin{aligned} q \frac{d}{dq} E_4(q) &= \frac{5P^6}{24} + \frac{5P^5Q}{4} + \frac{13P^4Q^2}{8} + \frac{13P^2Q^4}{8} - \frac{5PQ^5}{4} + \frac{5Q^6}{24} \\ &\quad - \frac{5}{3}P^2Q^2R + 5P^3QR - 5PQ^3R + \frac{5P^4R}{3} + \frac{5Q^4R}{3}, \end{aligned}$$

which implies that

$$\begin{aligned} E_6(q) &= E_2(q)E_4(q) - 3q \frac{d}{dq} E_4(q) \\ &= -\frac{1}{40} (P^2 + Q^2) (41P^4 + 198P^3Q + 154P^2Q^2 - 198PQ^3 + 41Q^4). \end{aligned}$$

□

Corollary 9.9. For $q \in \mathbb{C}$ with $|q| < 1$, set

$$S(q) = q \prod_{n=1}^{\infty} \frac{(1-q^n)^2}{(1-q^{5n-2})^5(1-q^{5n-3})^5} = \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n - 3 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n,$$

and

$$T(q) = \prod_{n=1}^{\infty} \frac{(1-q^n)^2}{(1-q^{5n-1})^5(1-q^{5n-4})^5} = 1 + 3 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n + \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n.$$

Then, we have

$$\begin{aligned} E_4(q) &= S^4 - 228S^3T + 494S^2T^2 + 228ST^3 + T^4, \\ E_6(q) &= (S^2 + T^2) (S^4 + 522S^3T - 10006S^2T^2 - 522ST^3 + T^4), \end{aligned}$$

and

$$\begin{aligned} q(q;q)_{\infty} &= -ST (S^2 + 11ST - T^2)^5 \\ j(q) &= -\frac{(S^4 - 228S^3T + 494S^2T^2 + 228ST^3 + T^4)^3}{ST (S^2 + 11ST - T^2)^5} = -\frac{(g^4 - 228g^3 + 494g^2 + 228g + 1)^3}{g(g^2 + 11g - 1)^5}, \end{aligned}$$

where

$$g = S/T = q \prod_{n=1}^{\infty} \frac{(1-q^{5n-1})^5(1-q^{5n-4})^5}{(1-q^{5n-2})^5(1-q^{5n-3})^5}.$$

Remark

Corollary 9.7 and Theorem 9.8 enable us to consider the extension of the differential fields,

$$\mathbb{Q}(E_2(q), E_4(q), E_6(q)) \subset \mathbb{Q}(P(q), Q(q), E_2(q)),$$

where

$$P(q) = 1 + 10 \sum_{n=1}^{\infty} (d_{2,5}(n) - d_{3,5}(n))q^n, \quad Q(q) = 3 + 10 \sum_{n=1}^{\infty} (d_{1,5}(n) - d_{4,5}(n))q^n.$$

10 Riccati equations satisfied by level 5 modular forms

10.1 Derivation of Riccati equation (1)

Proposition 10.1. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} - \theta'' \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} - \theta \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} = \frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2}{100\theta^6 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^6 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix}} \left\{ -8\theta^{10} \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} + 88\theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} \theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} + 8\theta^{10} \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} \right\}.$$

Proof. The proposition can be obtained by considering the derivative formulas (5.1) and (5.2) and by comparing the coefficients of term z^2 in the equation (8.1). \square

Theorem 10.2. *For every $\tau \in \mathbb{H}^2$, set*

$$W = \frac{\theta^5 \begin{bmatrix} 1 \\ \frac{1}{5} \end{bmatrix} (0, \tau)}{\theta^5 \begin{bmatrix} 1 \\ \frac{3}{5} \end{bmatrix} (0, \tau)} = \frac{11 + 5\sqrt{5}}{2} \prod_{n=1}^{\infty} \frac{\left(1 + \frac{1+\sqrt{5}}{2}q^n + q^{2n}\right)^5}{\left(1 + \frac{1-\sqrt{5}}{2}q^n + q^{2n}\right)^5}, \quad q = \exp(2\pi i\tau).$$

W then satisfies the following Riccati equation:

$$q \frac{d}{dq} W = \frac{1}{(\sqrt{5})^3} \frac{(q; q)_\infty^5}{(q^5; q^5)_\infty} (W^2 - 11W - 1). \quad (10.1)$$

Proof. The heat equation (2.6) and Proposition 10.1 imply that

$$4\pi i \frac{d}{d\tau} \log \theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] - 4\pi i \frac{d}{d\tau} \log \theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] \\ = \frac{\left\{ \theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \right\}^2 \left\{ -8\theta^{10} \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] + 88\theta^5 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] + 8\theta^{10} \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] \right\}}{100\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]},$$

which shows that

$$4\pi i \frac{d}{d\tau} \log \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] - 4\pi i \frac{d}{d\tau} \log \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] \\ = \frac{\left\{ \theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \right\}^2 \left\{ -8\theta^{10} \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] + 88\theta^5 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] + 8\theta^{10} \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] \right\}}{20\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]}.$$

Jacobi's triple product identity (2.5) yields

$$\frac{\left\{ \theta' \left[\begin{smallmatrix} 1 \\ 1 \end{smallmatrix} \right] \right\}^2}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right] \theta \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right]} = \frac{4\pi^2}{\sqrt{5}} \prod_{n=1}^{\infty} \frac{(1-q^n)^5}{(1-q^{5n})} = \frac{4\pi^2}{\sqrt{5}} \frac{\eta(\tau)^5}{\eta(5\tau)}.$$

By setting

$$(X, Y) = \left(\theta^5 \left[\begin{smallmatrix} 1 \\ \frac{1}{5} \end{smallmatrix} \right], \theta^5 \left[\begin{smallmatrix} 1 \\ \frac{3}{5} \end{smallmatrix} \right] \right),$$

we obtain

$$\frac{dX}{d\tau} Y - X \frac{dY}{d\tau} = \frac{2\pi i}{(\sqrt{5})^3} \frac{\eta^5(\tau)}{\eta(5\tau)} (X^2 - 11XY - Y^2),$$

which implies that

$$\frac{d}{d\tau} W = \frac{d}{d\tau} \left(\frac{X}{Y} \right) = \frac{\frac{dX}{d\tau} Y - X \frac{dY}{d\tau}}{Y^2} = \frac{2\pi i}{(\sqrt{5})^3} \frac{\eta^5(\tau)}{\eta(5\tau)} (W^2 - 11W - 1).$$

The theorem can be obtained by considering that $\frac{d}{d\tau} = \frac{dq}{d\tau} \frac{d}{dq} = 2\pi iq \frac{d}{dq}$. \square

10.2 Derivation of a Riccati equation (2)

Proposition 10.3. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta'' \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - \theta'' \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} = \frac{2\zeta_5}{25} \frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2}{\theta^6 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^6 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix}} \left\{ \theta^{10} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} + 11\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - \theta^{10} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \right\}.$$

Proof. The proposition can be obtained by considering the derivative formulas (5.3) and (5.4) and by comparing the coefficients of the term z^2 in equation (8.2). \square

Theorem 10.4. *For every $\tau \in \mathbb{H}^2$, set*

$$W = \frac{\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, 5\tau)}{\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, 5\tau)} = -q \prod_{n=1}^{\infty} \frac{(1 - q^{5n-1})^5 (1 - q^{5n-4})^5}{(1 - q^{5n-2})^5 (1 - q^{5n-3})^5}, \quad q = \exp(2\pi i\tau).$$

W then satisfies the following Riccati equation:

$$q \frac{d}{dq} W = q \frac{(q^5; q^5)_\infty^5}{(q; q)_\infty} (W^2 - 11W - 1). \quad (10.2)$$

Proof. The heat equation (2.6) and Proposition 10.3 imply that

$$\begin{aligned} & 4\pi i \frac{d}{d\tau} \log \theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - 4\pi i \frac{d}{d\tau} \log \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \\ &= -\frac{2\zeta_5}{25} \frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \frac{\left\{ \theta^{10} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - 11\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} - \theta^{10} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \right\}}{\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}, \end{aligned}$$

which shows that

$$\begin{aligned} & 4\pi i \frac{d}{d\tau} \log \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - 4\pi i \frac{d}{d\tau} \log \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \\ &= -\frac{2\zeta_5}{5} \frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} \frac{\left\{ \theta^{10} \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} - 11\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} - \theta^{10} \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} \right\}}{\theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}}. \end{aligned}$$

Jacobi's triple product identity (2.5) yields

$$\frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2}{\theta \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} \theta \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix}} = \frac{4\pi^2}{\zeta_5} y \prod_{n=1}^{\infty} \frac{(1-y^{5n})^5}{(1-y^n)} = \frac{4\pi^2}{\zeta_5} \frac{\eta^5(\tau)}{\eta(\tau/5)}, \quad y = \exp(2\pi i \tau/5).$$

By setting

$$(X, Y) = \left(\theta^5 \begin{bmatrix} \frac{1}{5} \\ 1 \end{bmatrix} (0, \tau), \theta^5 \begin{bmatrix} \frac{3}{5} \\ 1 \end{bmatrix} (0, \tau) \right),$$

we obtain

$$y \frac{d}{dy} W = y \frac{d}{dy} \left(\frac{Y}{X} \right) = y \frac{(y^5; y^5)_\infty^5}{(y; y)_\infty} (W^2 - 11W - 1).$$

The theorem can be obtained by changing $\tau \rightarrow 5\tau$. □

References

- [1] T. M. Apostol, *Modular functions and Dirichlet series in number theory*, Second edition. Grad. Texts in Math. **41** Springer-Verlag, New York, 1990.
- [2] J. Chazy, *Sur les équations différentielles du troisième ordre et d'ordre supérieur dont l'integrale générale a ses points critiques fixes*, Acta Math. **34** (1911), 317-385.
- [3] S. Cooper, *Ramanujan's theta functions*. Springer, Cham, 2017.
- [4] G. Darboux, *Sur la théorie des coordonnées curvilignes et les systèmes orthogonaux*, Ann. Ecole Normale Supérieure **7** (1878) 101-150
- [5] H. M. Farkas and I. Kra, *Theta constants, Riemann surfaces and the modular group*, AMS Grad. Studies in Math. **37** (2001).
- [6] G. Halphen, *Sur une system d'équations différ entielles*, C. R. Acad. Sci., Paris **92** (1881), 1101-1103.
- [7] C. Hermite, Oeuvres, Vol. II, Gauthier-Villars, Paris, 1908. *Note sur la théorie des fonctions elliptiques*, pp125-238.
- [8] T. Huber, *Differential equations for cubic theta functions*, Int. J. Number Theory **71** (2011), 1945-1957.

- [9] K. Matsuda, *Differential equations satisfied by $a(q) = \sum_{m,n \in \mathbb{Z}} q^{m^2+mn+n^2}$* , arXiv:1609.07481
- [10] Y. Ohyama, *Differential relations of theta functions.* Osaka J. Math. 32 (1995), 431-450.
- [11] S. Ramanujan, *On certain arithmetical functions,* Trans. Cambridge Philos. Soc. **22**, no. 9, (1916) 159-184.
- [12] E. T. Whittaker, and G. N. Watson, *A course of modern analysis. An introduction to the general theory of infinite processes and of analytic functions; with an account of the principal transcendental functions,* Fourth edition. Reprinted Cambridge University Press, New York 1962