

Note on some theorem of Farkas and Kra

By Kazuhide Matsuda

Faculty of Fundamental Science, National Institute of Technology, Niihama College,
7-1 Yagumo-chou, Niihama, Ehime, Japan, 792-8580.

E-mail: matsuda@sci.niihama-nct.ac.jp

Fax: 81-0897-37-7809

Abstract In this paper, we apply high level versions of Jacobi's derivative formula to number theory such as quarternary quadratic forms and convolution sums of some arithmetical functions.

Key Words: theta function; theta constant; rational characteristics; Hecke group.

MSC(2010) 14K25; 11E25

1 Introduction

Throughout this paper, let \mathbb{N}_0, \mathbb{N} , denote the set of nonnegative integers, positive integers, and denote the set of squares by $\{x^2 : x \in \mathbb{Z}\}$. A *triangular number* is defined as $t_x = x(x+1)/2$, where $x \in \mathbb{Z}$.

For the positive integers j, k , and $n \in \mathbb{N}$, $d_{j,k}(n)$ denotes the number of positive divisors d of n such that $d \equiv j \pmod k$, and $d_{j,k}^*(n)$ denotes the number of positive divisors d of n such that $d \equiv j \pmod k$ and n/d is odd, which implies that $d_{j,k}^*(n) = d_{j,k}(n) - d_{j,k}(n/2)$.

For each $n \in \mathbb{N}$, $\sigma(n)$ is the sum of positive divisors of n , and $\sigma^*(n)$ is the sum of positive divisors d of n such that n/d is odd, which implies that $\sigma^*(n) = \sigma(n) - \sigma(n/2)$. Moreover, we set $d_{j,k}(n) = \sigma(n) = 0$ for $n \in \mathbb{Q} \setminus \mathbb{N}_0$.

The *upper half plane* \mathbb{H}^2 is defined by $\mathbb{H}^2 = \{\tau \in \mathbb{C} \mid \Im\tau > 0\}$, and the *Dedekind eta function* is defined by $\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n)$, $q = \exp(2\pi i\tau)$ for $\tau \in \mathbb{H}^2$.

Following Farkas and Kra [4], we introduce the *theta function with characteristics*, which is defined by

$$\theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] (\zeta, \tau) = \theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] (\zeta) := \sum_{n \in \mathbb{Z}} \exp \left(2\pi i \left[\frac{1}{2} \left(n + \frac{\epsilon}{2} \right)^2 \tau + \left(n + \frac{\epsilon}{2} \right) \left(\zeta + \frac{\epsilon'}{2} \right) \right] \right),$$

where $\epsilon, \epsilon' \in \mathbb{R}$, $\zeta \in \mathbb{C}$, and $\tau \in \mathbb{H}^2$. The *theta constants* are given by

$$\theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] := \theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] (0, \tau).$$

Furthermore, we denote the derivative coefficients of the theta function by

$$\theta' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \frac{\partial}{\partial \zeta} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) \Big|_{\zeta=0}, \quad \theta'' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} := \frac{\partial^2}{\partial \zeta^2} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) \Big|_{\zeta=0}.$$

Farkas and Kra [4] treated the theta constants with *rational characteristics*, that is, the case where ϵ and ϵ' are both rational numbers, and derived a number of interesting theta constant identities. For each positive integer k , define the *Hecke group* $\Gamma_o(k)$ by

$$\Gamma_o(k) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL(2, \mathbb{Z}); c \equiv 0 \pmod{k} \right\}.$$

In [4, pp. 318], they proved the following result:

Theorem 1.1. (Farkas and Kra) *For each odd prime k ,*

$$\frac{d}{d\tau} \log \left(\frac{\eta(k\tau)}{\eta(\tau)} \right) + \frac{1}{2\pi i(k-2)} \sum_{l=0}^{\frac{k-3}{2}} \left(\frac{\theta' \begin{bmatrix} 1 \\ \frac{1+2l}{k} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{1+2l}{k} \end{bmatrix} (0, \tau)} \right)^2, \quad \tau \in \mathbb{H}^2,$$

is a cusp form for $\Gamma_o(k)$. This form is identically zero provided $k \leq 13$, $k \neq 11$.

The aim of this paper is to consider analogues of Theorem 1.1 for $k = 4, 6, 8$ and their applications to number theory such as quarternary quadratic forms and convolutions sums of some arithmetical functions. For this purpose, we use the results of our papers [5, 6, 7], where we considered high level versions of *Jacobi's derivative formula*:

$$\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} = -\pi \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \quad (1.1)$$

The remainder of this paper is as follows. In Sections 2 and 3, we review the basic properties and the derivative formulas of the theta functions, respectively. In Section 4, we deal with $\Gamma_o(4)$ and prove 4 squares theorem and 4 triangular numbers theorem.

In Section 5, we treat $\Gamma_o(6)$ and apply the results to the quadratic forms,

$$x^2 + xy + y^2 + z^2 + zw + w^2, \quad x^2 + y^2 + 3z^2 + 3w^2, \quad x^2 + xy + y^2 + 2(z^2 + zw + w^2),$$

and convolution sums of the arithmetical functions,

$$\delta^*(n) = d_{1,3}^*(n) - d_{2,3}^*(n), \quad \epsilon^*(n) = d_{1,6}^*(n) + d_{2,6}^*(n) - d_{4,6}^*(n) - d_{5,6}^*(n) \text{ for each } n \in \mathbb{N}.$$

In Section 6, we deal with $\Gamma_o(8)$ and apply the result to the mixed sums of squares and triangular numbers,

$$x^2 + y^2 + 2z^2 + 2w^2, \quad x^2 + 2y^2 + 4t_z + 4t_w, \quad x^2 + y^2 + 4t_z + 4t_w, \quad x^2 + 2y^2 + 2z^2 + 4t_w,$$

and

$$x^2 + y^2 + z^2 + 2w^2, \quad x^2 + 2y^2 + 2z^2 + 2w^2, \quad x^2 + y^2 + z^2 + 4t_w, \quad x^2 + 4t_y + 4t_z + 4t_w.$$

2 The properties of the theta functions

2.1 Basic properties

We first note that for $m, n \in \mathbb{Z}$,

$$\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta + n + m\tau, \tau) = \exp(2\pi i) \left[\frac{n\epsilon - m\epsilon'}{2} - m\zeta - \frac{m^2\tau}{2} \right] \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau), \quad (2.1)$$

and

$$\theta \begin{bmatrix} \epsilon + 2m \\ \epsilon' + 2n \end{bmatrix} (\zeta, \tau) = \exp(\pi i \epsilon n) \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau). \quad (2.2)$$

Furthermore, it is easy to see that

$$\theta \begin{bmatrix} -\epsilon \\ -\epsilon' \end{bmatrix} (\zeta, \tau) = \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (-\zeta, \tau) \text{ and } \theta' \begin{bmatrix} -\epsilon \\ -\epsilon' \end{bmatrix} (\zeta, \tau) = -\theta' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (-\zeta, \tau).$$

For $m, n \in \mathbb{R}$, we see that

$$\begin{aligned} & \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} \left(\zeta + \frac{n + m\tau}{2}, \tau \right) \\ &= \exp(2\pi i) \left[-\frac{m\zeta}{2} - \frac{m^2\tau}{8} - \frac{m(\epsilon' + n)}{4} \right] \theta \begin{bmatrix} \epsilon + m \\ \epsilon' + n \end{bmatrix} (\zeta, \tau). \end{aligned} \quad (2.3)$$

We note that $\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau)$ has only one zero in the fundamental parallelogram, which is given by

$$\zeta = \frac{1 - \epsilon}{2}\tau + \frac{1 - \epsilon'}{2}.$$

2.2 Jacobi's triple product identity

All the theta functions have infinite product expansions, which are given by

$$\begin{aligned} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) &= \exp\left(\frac{\pi i \epsilon \epsilon'}{2}\right) x^{\frac{\epsilon}{4}} z^{\frac{\epsilon'}{2}} \\ &\times \prod_{n=1}^{\infty} (1 - x^{2n})(1 + e^{\pi i \epsilon'} x^{2n-1+\epsilon} z)(1 + e^{-\pi i \epsilon'} x^{2n-1-\epsilon}/z), \end{aligned} \quad (2.4)$$

where $x = \exp(\pi i \tau)$ and $z = \exp(2\pi i \zeta)$. Therefore, it follows from Jacobi's derivative formula (1.1) that

$$\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} (0, \tau) = -2\pi q^{\frac{1}{8}} \prod_{n=1}^{\infty} (1 - q^n)^3, \quad q = \exp(2\pi i \tau).$$

2.3 Spaces of N -th order θ -functions

Following Farkas and Kra [4], we define $\mathcal{F}_N \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}$ to be the set of entire functions f that satisfy the two functional equations,

$$f(\zeta + 1) = \exp(\pi i \epsilon) f(\zeta),$$

and

$$f(\zeta + \tau) = \exp(-\pi i)[\epsilon' + 2N\zeta + N\tau] f(\zeta), \quad \zeta \in \mathbb{C}, \tau \in \mathbb{H}^2,$$

where N is a positive integer and $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} \in \mathbb{R}^2$. This set of functions is called the space of N -th order θ -functions with characteristic $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}$. Note that

$$\dim \mathcal{F}_N \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} = N.$$

For its proof, see Farkas and Kra [4, pp.133].

2.4 The heat equation

The theta function satisfies the following heat equation:

$$\frac{\partial^2}{\partial \zeta^2} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) = 4\pi i \frac{\partial}{\partial \tau} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau). \quad (2.5)$$

2.5 Lemma of Farkas and Kra

We recall the lemma of Farkas and Kra [4, pp.78].

Lemma 2.1. *For all characteristics $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}$, $\begin{bmatrix} \delta \\ \delta' \end{bmatrix}$ and all $\tau \in \mathbb{H}^2$, we have*

$$\begin{aligned} & \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (0, \tau) \theta \begin{bmatrix} \delta \\ \delta' \end{bmatrix} (0, \tau) \\ = & \theta \begin{bmatrix} \frac{\epsilon+\delta}{2} \\ \epsilon' + \delta' \end{bmatrix} (0, 2\tau) \theta \begin{bmatrix} \frac{\epsilon-\delta}{2} \\ \epsilon' - \delta' \end{bmatrix} (0, 2\tau) + \theta \begin{bmatrix} \frac{\epsilon+\delta}{2} + 1 \\ \epsilon' + \delta' \end{bmatrix} (0, 2\tau) \theta \begin{bmatrix} \frac{\epsilon-\delta}{2} + 1 \\ \epsilon' - \delta' \end{bmatrix} (0, 2\tau). \end{aligned}$$

3 Derivative formulas

From Matsuda [5, 6, 7], recall the following derivative formulas:

Theorem 3.1. *For every $\tau \in \mathbb{H}^2$, we have*

$$\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} (0, \tau) = -\pi\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} (0, \tau), \quad (3.1)$$

$$\theta' \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix} (0, \tau) = -\pi\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix} (0, \tau), \quad (3.2)$$

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} = \frac{1}{6}\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{\left(\theta^4 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} - 3\theta^4 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} \right)}{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta^3 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}, \quad \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} = \frac{1}{3}\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \frac{\theta^4 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta^3 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}, \quad (3.3)$$

$$\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix} (0, \tau) = -\pi\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix} (0, \tau)\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \left\{ \sqrt{2}\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) - \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \right\}, \quad (3.4)$$

$$\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix} (0, \tau) = -\pi\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix} (0, \tau)\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \left\{ \sqrt{2}\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) + \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \right\}, \quad (3.5)$$

$$\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (0, \tau) = -\pi\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (0, \tau)\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) \left\{ \sqrt{2}\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) - \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) \right\}, \quad (3.6)$$

$$\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix} (0, \tau) = -\pi\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix} (0, \tau)\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) \left\{ \sqrt{2}\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) + \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) \right\}. \quad (3.7)$$

4 Case for $\Gamma_o(4)$

4.1 Case where $\epsilon = 1, \epsilon' = 1/2$

4.1.1 Theorem for $\epsilon = 1, \epsilon' = 1/2$

Theorem 4.1. (Farkas and Kra [4, pp. 321]) *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta(4\tau)}{\eta(\tau)} + \frac{1}{2\pi i \cdot 2} \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} (0, \tau)} \right\}^2 = 0.$$

Proof. Consider the following elliptic function:

$$\varphi(z) = \frac{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z) \theta^2 \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}.$$

In the fundamental parallelogram, the pole of $\varphi(z)$ is $z = 0$, which implies that $\text{Res}(\varphi(z), 0) = 0$. Therefore, it follows that

$$\frac{\theta'' \begin{bmatrix} 1 \\ 0 \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \right\}^2 = 0.$$

The heat equation (2.5) implies that

$$4\pi i \frac{d}{d\tau} \log \frac{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \right\}^2 = 0.$$

The theorem follows from Jacobi's triple product identity (2.4). □

4.1.2 4 squares theorem

Theorem 4.2. For each $n \in \mathbb{N}$,

$$S_4(n) =: \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + y^2 + z^2 + w^2 = n\}.$$

Then, we have

$$S_4(n) = 8\sigma(n) - 32\sigma(n/4).$$

Proof. Set $q = \exp(2\pi i\tau)$. The derivative formula (3.1) and Theorem 4.1 imply that

$$\frac{d}{d\tau} \log \frac{\eta(4\tau)}{\eta(\tau)} + \frac{1}{2\pi i \cdot 2} \left\{ -\pi \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \right\}^2 = 0,$$

which shows that

$$1 + \sum_{n=1}^{\infty} S_4(n) q^n = 1 + 8 \sum_{n=1}^{\infty} (\sigma(n) - 4\sigma(n/4)) q^n.$$

□

4.2 Case where $\epsilon = 0, \epsilon' = 1/2$

4.2.1 Theorem for $\epsilon = 0, \epsilon' = 1/2$

Theorem 4.3. (Farkas and Kra [4, pp. 321]) *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta^3(2\tau)}{\eta^2(\tau)\eta(4\tau)} + \frac{1}{2\pi i \cdot 2} \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix} (0, \tau)} \right\}^2 = 0.$$

Proof. Consider the following elliptic function:

$$\psi(z) = \frac{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z) \theta^2 \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}.$$

In the fundamental parallelogram, the pole of $\psi(z)$ is $z = 0$, which implies that $\text{Res}(\psi(z), 0) = 0$. Therefore, it follows that

$$\frac{\theta'' \begin{bmatrix} 1 \\ 0 \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix}} \right\}^2 = 0.$$

The heat equation (2.5) implies that

$$4\pi i \frac{d}{d\tau} \log \frac{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta^2 \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{2} \end{bmatrix}} \right\}^2 = 0.$$

The theorem follows from Jacobi's triple product identity (2.4). □

4.2.2 4 triangular numbers theorem

Theorem 4.4. *For each $n \in \mathbb{N}_0$, set*

$$T_4(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid t_x + t_y + t_z + t_w = n\}.$$

Then, we have

$$T_4(n) = 16\sigma(2n + 1).$$

Proof. Set $q = \exp(2\pi i\tau)$. The derivative formula (3.2) and Theorem 4.3 imply that

$$\frac{d}{d\tau} \log \frac{\eta^3(2\tau)}{\eta^2(\tau)\eta(4\tau)} + \frac{1}{2\pi i \cdot 2} \left\{ -\pi\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau) \right\}^2 = 0,$$

which shows that

$$\begin{aligned} \sum_{n=0}^{\infty} T_4(n)q^{2n+1} &= 16 \sum_{n=1}^{\infty} (\sigma(n) - 3\sigma(n/2) + 2\sigma(n/4))q^n \\ &= 16 \sum_{n=0}^{\infty} \sigma(2n+1)q^{2n+1}. \end{aligned}$$

□

5 Case for $\Gamma_o(6)$

In this section, we introduce the following formula:

$$a(q) =: \sum_{m,n \in \mathbb{Z}} q^{m^2+mn+n^2} = 1 + 6 \sum_{n=1}^{\infty} (d_{1,3}(n) - d_{2,3}(n))q^n \text{ for } q \in \mathbb{C} \text{ with } |q| < 1, \quad (5.1)$$

which is equivalent to the following formula: for each $n \in \mathbb{N}$,

$$\#\{(x, y) \in \mathbb{Z}^2 \mid x^2 + xy + y^2\} = 6(d_{1,3}(n) - d_{2,3}(n)). \quad (5.2)$$

For the proof of the formula (5.1), see Berndt [1, pp. 79]. For the elementary proof of the formula (5.2), see Dickson [2, pp. 68].

5.1 Preliminary results

Proposition 5.1. *For every $\tau \in \mathbb{H}^2$, set $q = \exp(2\pi i\tau)$. Then, for every $\tau \in \mathbb{H}^2$ and $j = 1, 2$, we have*

$$\begin{aligned} \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, \tau)} &= -\frac{\pi}{\sqrt{3}}a(q), \quad \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, \tau)} - \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (0, \tau)} = -2 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, 2\tau)}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, 2\tau)}, \\ \frac{\theta' \begin{bmatrix} 0 \\ \frac{j}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{j}{3} \end{bmatrix} (0, \tau)} &= \frac{\theta' \begin{bmatrix} 1 \\ \frac{j}{3} \end{bmatrix} (0, \tau/2)}{\theta \begin{bmatrix} 1 \\ \frac{j}{3} \end{bmatrix} (0, \tau/2)} - \frac{\theta' \begin{bmatrix} 1 \\ \frac{j}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{j}{3} \end{bmatrix} (0, \tau)}. \end{aligned}$$

Proof. Set $x = \exp(\pi i \tau)$. Jacobi's triple product identity (2.4) yields

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} = -\frac{\pi}{\sqrt{3}} \left\{ 1 + 6 \sum_{n=1}^{\infty} (d_{1,3}(n) - d_{2,3}(n)) q^n \right\},$$

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} = -\sqrt{3}\pi \left\{ 1 + 2 \sum_{n=1}^{\infty} (d_{1,6}(n) + d_{2,6}(n) - d_{4,6}(n) - d_{5,6}(n)) q^n \right\}.$$

$$\frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} = -2\sqrt{3}\pi \sum_{n=1}^{\infty} (d_{1,3}^*(n) - d_{2,3}^*(n)) x^n,$$

and

$$\frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} = -2\sqrt{3}\pi \sum_{n=1}^{\infty} (d_{1,6}^*(n) + d_{2,6}^*(n) - d_{4,6}^*(n) - d_{5,6}^*(n)) x^n,$$

which prove the proposition. □

5.2 Theorem for $\epsilon = 1, \epsilon' = 1/3$

5.2.1 Theorem for $\epsilon = 1, \epsilon' = 1/3$

Theorem 5.2. (Farkas and Kra [4, pp. 318]) *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta(3\tau)}{\eta(\tau)} + \frac{1}{2\pi i} \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (0, \tau)} \right\}^2 = 0.$$

Proof. Consider the following elliptic function:

$$\varphi(z) = \frac{\theta^3 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}.$$

In the fundamental parallelogram, the pole of $\varphi(z)$ is $z = 0$, which implies that $\text{Res}(\varphi(z), 0) = 0$. Therefore, it follows that

$$3 \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 6 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} \right\}^2 = 0. \quad (5.3)$$

The heat equation (2.5) implies that

$$4\pi i \frac{d}{d\tau} \log \frac{\theta^3 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 6 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} \right\}^2 = 0.$$

The theorem follows from Jacobi's triple product identity (2.4). \square

5.2.2 On $x^2 + xy + y^2 + z^2 + zw + w^2$

Theorem 5.3. For each $n \in \mathbb{N}$, set

$$s_2(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + xy + y^2 + z^2 + zw + w^2 = n\}.$$

Then, we have

$$s_2(n) = 12\sigma(n) - 36\sigma(n/3).$$

Proof. The theorem follows from Proposition 5.1 and Theorem 5.2. \square

5.2.3 On $x^2 + y^2 + 3z^2 + 3w^2$

Theorem 5.4. For each $n \in \mathbb{N}$, set

$$S_{1,1,3,3}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + y^2 + 3z^2 + 3w^2 = n\}.$$

Then,

$$S_{1,1,3,3}(n) = 4(-1)^{n-1}(\sigma(n) - 4\sigma(n/2) - 3\sigma(n/3) + 12\sigma(n/6)).$$

Proof. Consider the following elliptic function:

$$\psi(z) = \frac{\theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}(z) \theta \begin{bmatrix} 1 \\ -\frac{1}{3} \end{bmatrix}(z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix}(z)}.$$

In the fundamental parallelogram, the pole of $\psi(z)$ is $z = 0$, which implies that $\text{Res}(\psi(z), 0) = 0$. Therefore, it follows that

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} - 4 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} \right\}^2 = 0. \quad (5.4)$$

The heat equation (2.5) and the derivative formulas (3.3) imply that

$$4\pi i \frac{d}{d\tau} \log \frac{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + \frac{2}{3} \frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2 \theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} = 0. \quad (5.5)$$

Jacobi's triple product identity (2.4) yields

$$\frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2 \theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} = \left\{ -\sqrt{3}\pi \frac{\eta^2(\tau)\eta^2(3\tau)}{\eta(2\tau)\eta(6\tau)} \right\}^2,$$

$$\left\{ \frac{\eta^2(\tau)\eta^2(3\tau)}{\eta(2\tau)\eta(6\tau)} \right\}^2 = 1 - 4 \sum_{n=1}^{\infty} (\sigma(n) - 4\sigma(n/2) - 3\sigma(n/3) + 12\sigma(n/6)) q^n, \quad q = \exp(2\pi i\tau),$$

and

$$\theta^2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} (0, 6\tau) = \left\{ \left(\sum_{m \in \mathbb{Z}} (-1)^m q^{m^2} \right) \left(\sum_{n \in \mathbb{Z}} (-1)^n q^{3n^2} \right) \right\}^2 = \left\{ \frac{\eta^2(\tau)\eta^2(3\tau)}{\eta(2\tau)\eta(6\tau)} \right\}^2,$$

which imply that

$$\sum_{n=0}^{\infty} (-1)^n S_{1,1,3,3}(n) q^n = 1 - 4 \sum_{n=1}^{\infty} (\sigma(n) - 4\sigma(n/2) - 3\sigma(n/3) + 12\sigma(n/6)) q^n.$$

□

5.3 Theorem for $\epsilon = 1, \epsilon' = 2/3$

Proposition 5.5. *For every $(z, \tau) \in \mathbb{C} \times \mathbb{H}^2$, we have*

$$\begin{aligned} \theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z, \tau) + \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{4}{3} \end{bmatrix} (z, \tau) \\ - \theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau) = 0. \end{aligned} \quad (5.6)$$

Proof. We prove equation (5.6). We first note that $\dim \mathcal{F}_2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} = 2$, and

$$\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z, \tau), \theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{4}{3} \end{bmatrix} (z, \tau), \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau) \in \mathcal{F}_2 \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Therefore, there exist some complex numbers, x_1, x_2 and x_3 not all zero such that

$$x_1 \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z, \tau) + x_2 \theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (z, \tau) \theta \begin{bmatrix} 1 \\ \frac{4}{3} \end{bmatrix} (z, \tau) + x_3 \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau) = 0.$$

Note that in the fundamental parallelogram, the zero of $\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z)$, $\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (z)$, or $\theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)$ is $z = 1/2, 1/6$ or 0 . Substituting $z = 1/2, 1/6$, and 0 , we have

$$\begin{aligned} x_2 \theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} + x_3 \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} &= 0, \\ x_1 \theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} + x_3 \theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} &= 0, \\ x_1 \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} - x_2 \theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} &= 0. \end{aligned}$$

Solving this system of equations, we have

$$(x_1, x_2, x_3) = \alpha \left(\theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}, \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}, -\theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \right) \text{ for some } \alpha \in \mathbb{C} \setminus \{0\},$$

which proves the proposition. \square

Theorem 5.6. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta^4(6\tau)}{\eta^3(\tau)\eta(3\tau)} + \frac{1}{2\pi i} \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix} (0, \tau)} \right\}^2 = 0.$$

Proof. Comparing the coefficients of the term z^2 in equation (5.6) yields

$$\frac{\theta'' \begin{bmatrix} 1 \\ 0 \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}} - \frac{\theta'' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} - \frac{\left\{ \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}^2 \theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} \right\}^2 = 0.$$

The heat equation (2.5) and equation (5.5) imply that

$$2\pi i \frac{d}{d\tau} \log \left\{ \frac{\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta^3 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta^4 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^3} \right\} + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} \right\}^2 = 0.$$

Jacobi's triple product identity (2.4) yields

$$\frac{\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \theta^3 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta^4 \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^3} = -\frac{3\sqrt{3}}{2\pi^3} \frac{\eta^4(6\tau)}{\eta^3(\tau)\eta(3\tau)},$$

which proves the theorem. □

5.3.1 On $x^2 + xy + y^2 + 2(z^2 + zw + w^2)$

Theorem 5.7. For each $n \in \mathbb{N}$, set

$$s_{1,2}(n) = \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + xy + y^2 + 2(z^2 + zw + w^2) = n\}.$$

Then, we have

$$s_{1,2}(n) = 6\sigma(n) - 12\sigma(n/2) + 18\sigma(n/3) - 36\sigma(n/6).$$

Proof. Set $q = \exp(2\pi i\tau)$. Summing both sides of equations (5.3) and (5.4) yields

$$2 \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} \right\} + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{2}{3} \end{bmatrix}} \right\}^2 = 0.$$

The heat equation (2.5), Proposition 5.1 and Theorems 5.2 and 5.6 imply that

$$\frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)} \cdot \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, 2\tau)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, 2\tau)} = \frac{2\pi i}{2} \frac{d}{d\tau} \log \left\{ \frac{\eta(\tau)\eta(3\tau)}{\eta(2\tau)\eta(6\tau)} \right\}. \quad (5.7)$$

Proposition 5.1 shows that

$$\begin{aligned} a(q)a(q^2) &= 6q \frac{d}{dq} \log q^{\frac{1}{6}} \prod_{n=1}^{\infty} \frac{(1-q^{2n})(1-q^{6n})}{(1-q^n)(1-q^{3n})} \\ &= 1 + 6 \sum_{n=1}^{\infty} (\sigma(n) - 2\sigma(n/2) + 3\sigma(n/3) - 6\sigma(n/6))q^n. \end{aligned}$$

□

5.4 Theorem for $\epsilon = 0, \epsilon' = 1/3$

Theorem 5.8. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta^4(3\tau/2)}{\eta^3(\tau)\eta(3\tau)} + \frac{1}{2\pi i} \left\{ \frac{\theta' \left[\begin{smallmatrix} 0 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)}{\theta \left[\begin{smallmatrix} 0 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)} \right\}^2 = 0.$$

Proof. Proposition 5.1 implies that

$$\begin{aligned} & \left\{ \frac{\theta' \left[\begin{smallmatrix} 0 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)}{\theta \left[\begin{smallmatrix} 0 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)} \right\}^2 \\ &= \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau/2)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau/2)} \right\}^2 - 2 \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau/2)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau/2)} \cdot \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)} + \left\{ \frac{\theta' \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)}{\theta \left[\begin{smallmatrix} 1 \\ \frac{1}{3} \end{smallmatrix} \right] (0, \tau)} \right\}^2. \end{aligned}$$

Theorem 5.8 and equation (5.7) imply that

$$\begin{aligned} & \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix} (0, \tau)} \right\}^2 \\ &= -2 \cdot 2\pi i \frac{d}{d\tau} \log \frac{\eta(3\tau/2)}{\eta(\tau/2)} - 2 \cdot 2 \cdot \frac{2\pi i}{2} \frac{d}{d\tau} \log \frac{\eta(\tau/2)\eta(3\tau/2)}{\eta(\tau)\eta(3\tau)} - 2\pi i \frac{d}{d\tau} \log \frac{\eta(3\tau)}{\eta(\tau)}, \end{aligned}$$

which proves the theorem. \square

5.5 Theorem for $\epsilon = 0, \epsilon' = 2/3$

Theorem 5.9. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta^{11}(3\tau)}{\eta^3(\tau)\eta^4(3\tau/2)\eta^4(6\tau)} + \frac{1}{2\pi i} \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix} (0, \tau)} \right\}^2 = 0.$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^2 \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix} (z) \theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}, \quad \psi(z) = \frac{\theta^2 \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix} (z) \theta \begin{bmatrix} 1 \\ -\frac{1}{3} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}.$$

In the fundamental parallelogram, the pole of $\varphi(z)$ or $\psi(z)$ is $z = 0$, which implies that $\text{Res}(\varphi(z), 0) = \text{Res}(\psi(z), 0) = 0$. Therefore, it follows that

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 4 \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} \right\}^2 = 0,$$

and

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} - 4 \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} \right\}^2 = 0.$$

Summing both sides of these equations yields

$$\begin{aligned} & \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \\ & + 2 \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} \right\} \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} + \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} \right\}^2 = 0. \end{aligned}$$

The theorem follows from the heat equation (2.5), Proposition 5.1, equation (5.7) and Theorems 5.2, 5.8. \square

5.6 Convolution sums

Theorem 5.10. *For each $n \in \mathbb{N}$, set*

$$\delta^*(n) = d_{1,3}^*(n) - d_{2,3}^*(n), \text{ and } \epsilon^*(n) = d_{1,6}^*(n) + d_{2,6}^*(n) - d_{4,6}^*(n) - d_{5,6}^*(n).$$

Then, for $n \geq 2$,

$$\sum_{k=1}^{n-1} \delta^*(k) \delta^*(n-k) = \sigma(n/2) - 2\sigma(n/3) + \sigma(n/6), \quad (5.8)$$

and

$$\sum_{k=1}^{n-1} \epsilon^*(k) \epsilon^*(n-k) = \sigma(n/2) + 2\sigma(n/3) - 11\sigma(n/6) + 8\sigma(n/12). \quad (5.9)$$

Proof. For every $\tau \in \mathbb{H}^2$, set $x = \exp(\pi i \tau)$. Jacobi's triple product identity (2.4) yields

$$\frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} = -2\sqrt{3}\pi \sum_{n=1}^{\infty} (d_{1,3}^*(n) - d_{2,3}^*(n)) x^n,$$

and

$$\frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} = -2\sqrt{3}\pi \sum_{n=1}^{\infty} (d_{1,6}^*(n) + d_{2,6}^*(n) - d_{4,6}^*(n) - d_{5,6}^*(n)) x^n.$$

The theorem follows from Theorems 5.8 and 5.9. \square

Theorem 5.11. For each $n \in \mathbb{N}$, set

$$\delta^*(n) = d_{1,3}^*(n) - d_{2,3}^*(n), \text{ and } \epsilon^*(n) = d_{1,6}^*(n) + d_{2,6}^*(n) - d_{4,6}^*(n) - d_{5,6}^*(n).$$

Then, for $n \geq 2$,

$$\sum_{k=1}^{n-1} \delta^*(k)\epsilon^*(n-k) = \begin{cases} 0 & \text{if } n \text{ is odd,} \\ \sigma(n) - 2\sigma(n/2) - \sigma(n/3) + 12\sigma(n/6) & \text{if } n \text{ is even.} \end{cases}$$

Proof. Set $x = \exp(\pi i\tau)$ and $q = \exp(2\pi i\tau)$. Consider the following elliptic function:

$$\varphi(z) = \frac{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix} (z) \theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix} (z) \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}.$$

In the fundamental parallelogram, the pole of $\varphi(z)$ is $z = 0$, which implies that $\text{Res}(\varphi(z), 0) = 0$. Therefore, it follows that

$$\frac{1}{2} \frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} + \frac{1}{2} \frac{\theta'' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} + \frac{1}{2} \frac{\theta'' \begin{bmatrix} 1 \\ 0 \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}} - \frac{1}{2} \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} = 0.$$

The heat equation (2.5) and Jacobi's triple product identity (2.4) yield

$$\frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} = -2\pi i \frac{d}{d\tau} \log \frac{\eta(3\tau)\eta^3(2\tau)}{\eta^3(\tau)\eta(6\tau)},$$

which proves the theorem. □

Theorem 5.12. For each $n \in \mathbb{N}$, set

$$\delta^*(n) = d_{1,3}^*(n) - d_{2,3}^*(n), \text{ and } \epsilon^*(n) = d_{1,6}^*(n) + d_{2,6}^*(n) - d_{4,6}^*(n) - d_{5,6}^*(n).$$

Then, for $n \geq 3$,

$$\sum_{\substack{(k,l) \in \mathbb{N}^2 \\ 2k+l=n}} \delta^*(k)\delta^*(l) = \sigma(n/3) - \sigma(n/4) - \sigma(n/6) + \sigma(n/12),$$

$$\sum_{\substack{(k,l) \in \mathbb{N}^2 \\ 2k+l=n}} \delta^*(k)\epsilon^*(l) = \sigma(n/3) + \sigma(n/4) - 5\sigma(n/6) + 3\sigma(n/12).$$

Proof. For every $\tau \in \mathbb{H}^2$, set $x = \exp(\pi i \tau)$. Note that

$$\frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} = 4\sqrt{3}\pi \sum_{n=1}^{\infty} (d_{1,3}^*(n) - d_{2,3}^*(n))x^{2n}.$$

The theorem can be proved by considering

$$\left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} \right\} \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}, \text{ and } \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}} \right\} \frac{\theta' \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{2}{3} \end{bmatrix}}.$$

□

Remark

For each $n \in \mathbb{N}$, set $\delta(n) = d_{1,3}(n) - d_{2,3}(n)$. Using the theory of theta functions, Farkas [3] showed that for each $n \in \mathbb{N}_0$,

$$\sigma(3n+2) = 3 \sum_{k=0}^n \delta(3k+1)\delta(3(n-k)+1).$$

6 Case for $\Gamma_o(8)$

6.1 Theorem for $\epsilon = 1, \epsilon' = 1/4, 3/4$

Theorem 6.1. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta^3(8\tau)}{\eta^2(\tau)\eta(4\tau)} + \frac{1}{2\pi i \cdot 2} \left[\left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}(0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}(0, \tau)} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}(0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}(0, \tau)} \right\}^2 \right] = 0.$$

Proof. By Lemma 2.1, we first note that

$$\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix}(0, \tau) = \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix}(0, 2\tau) + \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix}(0, 2\tau),$$

and

$$\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau) = 2\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau),$$

which implies that

$$\theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau) = 4\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau).$$

The derivative formulas (3.4) and (3.5) yield

$$\left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix} (0, \tau)} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix} (0, \tau)} \right\}^2 = 6\pi^2\theta^4 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) + \pi^2\theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau).$$

By the proof of Theorems 4.2 and 4.4, we have

$$\begin{aligned} 6\pi^2\theta^4 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) + \pi^2\theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau) &= 6 \cdot 4\pi i \cdot \left(-\frac{1}{2}\right) \cdot \frac{d}{d\tau} \log \frac{\eta(8\tau)}{\eta(2\tau)} - 4\pi i \frac{d}{d\tau} \log \frac{\eta^3(2\tau)}{\eta^2(\tau)\eta(4\tau)} \\ &= 4\pi i \frac{d}{d\tau} \log \frac{\eta^2(\tau)\eta(4\tau)}{\eta^3(8\tau)}, \end{aligned}$$

which proves the theorem. \square

6.2 Theorem for $\epsilon = 0, \epsilon' = 1/4, 3/4$

Theorem 6.2. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{d}{d\tau} \log \frac{\eta^8(4\tau)}{\eta^2(\tau)\eta^3(2\tau)\eta^3(8\tau)} + \frac{1}{2\pi i \cdot 2} \left[\left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (0, \tau)} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix} (0, \tau)} \right\}^2 \right] = 0.$$

Proof. By Lemma 2.1, we first note that

$$\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, \tau) = \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) + \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau),$$

and

$$\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau) = 2\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau),$$

which implies that

$$\theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, \tau) = 4\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau).$$

The derivative formulas (3.6) and (3.7) yield

$$\left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (0, \tau)} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix} (0, \tau)}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix} (0, \tau)} \right\}^2 = \pi^2 \theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau) + 6\pi^2 \theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau).$$

By the proof of Theorem 4.4, we have

$$\begin{aligned} \pi^2 \theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau) + 6\pi^2 \theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) &= -4\pi i \frac{d}{d\tau} \log \frac{\eta^3(2\tau)}{\eta^2(\tau)\eta(4\tau)} - 6 \cdot 4\pi i \cdot \frac{1}{2} \cdot \frac{d}{d\tau} \log \frac{\eta^3(4\tau)}{\eta^2(2\tau)\eta(8\tau)} \\ &= 4\pi i \frac{d}{d\tau} \log \frac{\eta^2(\tau)\eta^3(2\tau)\eta^3(8\tau)}{\eta^8(4\tau)}, \end{aligned}$$

which proves the theorem. \square

6.3 Applications

In this subsection, for each $m \in \mathbb{N}$, we set

$$\left(\frac{8}{m} \right) = \begin{cases} +1, & \text{if } m \equiv \pm 1 \pmod{8}, \\ -1, & \text{if } m \equiv \pm 3 \pmod{8}, \\ 0, & \text{if } m \equiv 0 \pmod{2}. \end{cases}$$

6.3.1 Preliminary results

Proposition 6.3. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 4 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 = 0, \quad (6.1)$$

and

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} - 4 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 = 0. \quad (6.2)$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^2 \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix} (z) \theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}, \quad \psi(z) = \frac{\theta^2 \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix} (z) \theta \begin{bmatrix} 1 \\ -\frac{1}{2} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}.$$

In the fundamental parallelogram, the pole of $\varphi(z)$ or $\psi(z)$ is $z = 0$, which implies that $\text{Res}(\varphi(z), 0) = \text{Res}(\psi(z), 0) = 0$. The proposition follows from direct calculation. \square

Proposition 6.4. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 4 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 = 0, \quad (6.3)$$

and

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + 2 \frac{\theta'' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} - 4 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \cdot \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} + 2 \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 = 0. \quad (6.4)$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^2 \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (z) \theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}, \quad \psi(z) = \frac{\theta^2 \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix} (z) \theta \begin{bmatrix} 1 \\ -\frac{1}{2} \end{bmatrix} (z)}{\theta^3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}.$$

In the fundamental parallelogram, the pole of $\varphi(z)$ or $\psi(z)$ is $z = 0$, which implies that $\text{Res}(\varphi(z), 0) = \text{Res}(\psi(z), 0) = 0$. The proposition follows from direct calculation. \square

6.3.2 On $x^2 + y^2 + 2z^2 + 2w^2$

Theorem 6.5. *For each $n \in \mathbb{N}$, set*

$$S_{1,1,2,2}(n) = \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + y^2 + 2z^2 + 2w^2 = n\}.$$

Then, we have

$$S_{1,1,2,2}(n) = 4\sigma(n) - 4\sigma(n/2) + 8\sigma(n/4) - 32\sigma(n/8).$$

Proof. Set $q = \exp(2\pi i\tau)$. Summing both sides of equations (6.1) and (6.2) yields

$$\begin{aligned} & \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \\ & + 2 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \cdot \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} \right\} + \left[\left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 \right] = 0. \end{aligned}$$

The derivative formulas (3.1), (3.4) and (3.5) imply that

$$\begin{aligned} & \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + \left[\left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 \right] \\ & = 4\pi^2 \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau). \end{aligned}$$

The heat equation (2.5) and Jacobi's triple product identity (2.4) shows that

$$4\pi i \frac{d}{d\tau} \log \frac{\eta(8\tau)}{\eta(2\tau)} + \left[\left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 \right] = 4\pi^2 \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau).$$

Theorem 6.1 yields

$$4\pi i \frac{d}{d\tau} \log \frac{\eta^2(\tau)\eta(4\tau)}{\eta(2\tau)\eta^2(8\tau)} = 4\pi^2 \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau). \quad (6.5)$$

The theorem can be obtained by considering that

$$\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) = \left(\sum_{n \in \mathbb{Z}} q^{n^2} \right)^2 \left(\sum_{n \in \mathbb{Z}} q^{2n^2} \right)^2 = \sum_{x, y, z, w \in \mathbb{Z}} q^{x^2 + y^2 + 2z^2 + 2w^2}.$$

□

6.3.3 On $x^2 + 2y^2 + 4t_z + 4t_w$

Theorem 6.6. For each $n \in \mathbb{N}_0$, set

$$M_{1,2-4,4}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + 2y^2 + 4t_z + 4t_w = n\}.$$

Then, we have

$$M_{1,2-4,4}(n) = 4 \sum_{d|n+1} \frac{n+1}{d} \left(\frac{8}{d}\right).$$

Proof. Set $q = \exp(2\pi i\tau)$. By Lemma 2.1, we first note that

$$\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, \tau) = \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) + \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau).$$

Subtracting both sides of equations (6.1) and (6.2) yields

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta'' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} = -4\sqrt{2}\pi^2 \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau).$$

Jacobi's triple product identity (2.4) yields

$$\frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta'' \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} = -16\sqrt{2}\pi^2 \sum_{n=1}^{\infty} \left(\sum_{d|n} \frac{n}{d} \left(\frac{8}{d}\right) \right) q^n.$$

From the definition, it follows that

$$\begin{aligned} \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) &= \left(\sum_{n \in \mathbb{Z}} q^{n^2} \right) \left(\sum_{n \in \mathbb{Z}} q^{2n^2} \right) \left(q^{\frac{1}{2}} \sum_{n \in \mathbb{Z}} q^{4 \cdot \frac{n(n+1)}{2}} \right)^2 \\ &= \sum_{n=0}^{\infty} M_{1,2-4,4}(n) q^{n+1}, \end{aligned}$$

which proves the theorem. \square

6.3.4 On $x^2 + y^2 + 4t_z + 4t_w$

Theorem 6.7. For each $n \in \mathbb{N}_0$, set

$$M_{1,1-4,4}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + y^2 + 4t_z + 4t_w = n\}.$$

Then, we have

$$M_{1,1-4,4}(n) = 4 \left(\sigma(n+1) + \sigma\left(\frac{n+1}{2}\right) - 10\sigma\left(\frac{n+1}{4}\right) + 8\sigma\left(\frac{n+1}{8}\right) \right).$$

Proof. Set $q = \exp(2\pi i\tau)$. Summing both sides of equations (6.3) and (6.4) yields

$$\begin{aligned} & \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} \\ & + 2 \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} \cdot \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} \right\} + \left[\left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 \right] = 0. \end{aligned}$$

The derivative formulas (3.1), (3.6) and (3.7) imply that

$$\begin{aligned} & \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} + \frac{\theta'' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + \left[\left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 \right] \\ & = 4\pi^2 \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau). \end{aligned}$$

The heat equation (2.5) and Jacobi's triple product identity (2.4) shows that

$$4\pi i \frac{d}{d\tau} \log \frac{\eta^3(4\tau)}{\eta^2(2\tau)\eta(8\tau)} + \left[\left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} \right\}^2 + \left\{ \frac{\theta' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} \right\}^2 \right] = 4\pi^2 \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau).$$

Theorem 6.2 yields

$$4\pi i \frac{d}{d\tau} \log \frac{\eta^2(\tau)\eta(2\tau)\eta^2(8\tau)}{\eta^5(4\tau)} = 4\pi^2 \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau). \quad (6.6)$$

The theorem can be obtained by considering that

$$\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) = \left(\sum_{n \in \mathbb{Z}} q^{n^2} \right)^2 \left(q^{\frac{1}{2}} \sum_{n \in \mathbb{Z}} q^{4 \cdot \frac{n(n+1)}{2}} \right)^2 = \sum_{x, y, z, w \in \mathbb{Z}} q^{x^2 + y^2 + 4t_z + 4t_w + 1}.$$

□

6.3.5 On $x^2 + 2y^2 + 2z^2 + 4t_w$

Theorem 6.8. For each $n \in \mathbb{N}_0$, set

$$M_{1,2,2-4}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + 2y^2 + 2z^2 + 4t_w = n\}.$$

Then, we have

$$M_{1,2,2-4}(n) = 2 \sum_{d|2n+1} \frac{2n+1}{d} \left(\frac{8}{d} \right).$$

Proof. Set $x = \exp(\pi i \tau)$ and $q = \exp(2\pi i \tau)$. By Lemma 2.1, we first note that

$$\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, \tau) = \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) + \theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 2\tau).$$

Subtracting both sides of equations (6.3) and (6.4) yields

$$\frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta'' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} = -4\sqrt{2}\pi^2 \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau).$$

Jacobi's triple product identity (2.4) yields

$$\frac{\theta'' \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}} - \frac{\theta'' \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} = -8\sqrt{2}\pi^2 \sum_{n=0}^{\infty} \left(\sum_{d|2n+1} \frac{2n+1}{d} \left(\frac{8}{d} \right) \right) x^{2n+1}.$$

From the definition, it follows that

$$\begin{aligned} \theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau) \theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) &= \left(\sum_{n \in \mathbb{Z}} q^{n^2} \right) \left(\sum_{n \in \mathbb{Z}} q^{2n^2} \right)^2 \left(q^{\frac{1}{2}} \sum_{n \in \mathbb{Z}} q^{4 \cdot \frac{n(n+1)}{2}} \right) \\ &= \sum_{n=0}^{\infty} M_{1,2,2-4}(n) x^{2n+1}, \end{aligned}$$

which proves the theorem. □

6.4 More applications

In this subsection, for each $m \in \mathbb{N}$, we set

$$\left(\frac{8}{m}\right) = \begin{cases} +1, & \text{if } m \equiv \pm 1 \pmod{8}, \\ -1, & \text{if } m \equiv \pm 3 \pmod{8}, \\ 0, & \text{if } m \equiv 0 \pmod{2}. \end{cases}$$

Moreover we have the following lemma:

Lemma 6.9. *For every $(z, \tau) \in \mathbb{C} \times \mathbb{H}^2$, we have*

$$\left\{ \frac{\theta' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)}{\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)} \right\}^2 = \frac{\theta'' \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)}{\theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z)} - \frac{d^2}{dz^2} \log \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (z).$$

Proof. The lemma can be proved by direct calculation. □

6.4.1 On $x^2 + y^2 + z^2 + 2w^2$, and $x^2 + 2y^2 + 2z^2 + 2w^2$

Theorem 6.10. *For each $n \in \mathbb{N}$, set*

$$S_{1,1,1,2}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + y^2 + z^2 + 2w^2 = n\}.$$

Then, we have

$$S_{1,1,1,2}(n) = 8 \sum_{d|n} \frac{n}{d} \left(\frac{8}{d}\right) - 2 \sum_{d|n} d \left(\frac{8}{d}\right).$$

Proof. We first note that for each $n \in \mathbb{N}$,

$$\sum_{d|n, d:\text{odd}} d = \sigma(n) - 2\sigma\left(\frac{n}{2}\right).$$

The derivative formulas (3.6), (3.7), and equation (6.1) yield

$$\begin{aligned} & 4\sqrt{2}\pi^2\theta^3 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \\ &= -4\pi i \frac{d}{d\tau} \log \frac{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} \theta^4 \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix} \theta^4 \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \theta^4 \begin{bmatrix} 1 \\ \frac{3}{4} \end{bmatrix}} + 4\pi^2\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \\ & \quad + 2 \frac{d^2}{dz^2} \log \theta \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix} (z) \Big|_{z=0}. \end{aligned}$$

The theorem follows from equation (6.5) and Jacobi's triple product identity (2.4). □

Theorem 6.11. For each $n \in \mathbb{N}$, set

$$S_{1,2,2,2}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + 2y^2 + 2z^2 + 2w^2 = n\}.$$

Then, we have

$$S_{1,2,2,2}(n) = 4 \sum_{d|n} \frac{n}{d} \left(\frac{8}{d}\right) - 2 \sum_{d|n} d \left(\frac{8}{d}\right).$$

Proof. The derivative formulas (3.6), (3.7), and equation (6.1) yield

$$\begin{aligned} & -4\sqrt{2}\pi^2\theta^3 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) + 4\sqrt{2}\pi^2\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta^3 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau) \\ & = 4\pi i \frac{d}{d\tau} \log \frac{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ \frac{1}{4} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2\pi^2\theta^4 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 4\tau). \end{aligned}$$

The theorem follows from Theorems 4.2 and 6.10. □

6.4.2 On $x^2 + y^2 + z^2 + 4t_w$, and $x^2 + 4t_y + 4t_z + 4t_w$

Theorem 6.12. For each $n \in \mathbb{N}_0$, set

$$M_{1,1,1,4}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + y^2 + z^2 + 4t_w = n\}.$$

Then, we have

$$M_{1,1,1,4}(n) = 4 \sum_{d|2n+1} \frac{2n+1}{d} \left(\frac{8}{d}\right) - 2 \sum_{d|2n+1} d \left(\frac{8}{d}\right).$$

Proof. We first note that for each $n \in \mathbb{N}$,

$$3\sigma(n) - \sigma(2n) - 2\sigma\left(\frac{n}{2}\right) = 0.$$

The derivative formulas (3.6), (3.7), and equation (6.3) yield

$$\begin{aligned} & 4\sqrt{2}\pi^2\theta^3 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) \\ & = -4\pi i \frac{d}{d\tau} \log \frac{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} \theta^4 \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} \theta^4 \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix} \theta^4 \begin{bmatrix} 0 \\ \frac{3}{4} \end{bmatrix}} + 4\pi^2\theta^2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta^2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) \\ & \quad + 2 \frac{d^2}{dz^2} \log \theta \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix} (z) \Big|_{z=0}. \end{aligned}$$

The theorem follows from equation (6.6) and Jacobi's triple product identity (2.4). □

Theorem 6.13. For each $n \in \mathbb{N}_0$, set

$$M_{1-4,4,4}(n) := \#\{(x, y, z, w) \in \mathbb{Z}^4 \mid x^2 + 4t_y + 4t_z + 4t_w = n\}.$$

Then, we have

$$M_{1-4,4,4}(n) = 2 \sum_{d|2n+3} \frac{2n+3}{d} \left(\frac{8}{d}\right) - 2 \sum_{d|2n+3} d \left(\frac{8}{d}\right).$$

Proof. The derivative formulas (3.6), (3.7), and equation (6.3) yield

$$\begin{aligned} & -4\sqrt{2}\pi^2\theta^3 \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) + 4\sqrt{2}\pi^2\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} (0, 2\tau)\theta^3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau) \\ & = 4\pi i \frac{d}{d\tau} \log \frac{\theta \begin{bmatrix} 1 \\ \frac{1}{2} \end{bmatrix} \theta^2 \begin{bmatrix} 0 \\ \frac{1}{4} \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 2\pi^2\theta^4 \begin{bmatrix} 1 \\ 0 \end{bmatrix} (0, 4\tau). \end{aligned}$$

The theorem follows from Theorems 4.4 and 6.6. □

References

- [1] B. C. Berndt, *Number theory in the spirit of Ramanujan*, Stud. Math. Libr., **34** American Mathematical Society, Providence, RI, 2006.
- [2] L. E. Dickson, *Modern Elementary Theory of Numbers*, University of Chicago Press, Chicago, 1939.
- [3] H. M. Farkas, *Theta functions in complex analysis and number theory*. Surveys in number theory, Dev. Math., **17**, (2008), 57-87.
- [4] H. M. Farkas and Irwin Kra, *Theta constants, Riemann surfaces and the modular group*, AMS Grad. Studies in Math. **37** (2001).
- [5] K. Matsuda, *Analogues of Jacobi's derivative formula*, to appear in Ramanujan J, arXiv:1411.3361
- [6] K. Matsuda, *Analogues of Jacobi's derivative formula II*, preprint.
- [7] K. Matsuda, *Analogues of Jacobi's derivative formula III*, preprint.