

A NOTE ON AVERAGE OF ROOTS OF UNITY

CHATCHAWAN PANRAKSA AND PORNRAT RUENGROT

ABSTRACT. We consider the problem of characterizing all functions f defined on the set of integers modulo n with the property that an average of some n th roots of unity determined by f is always an algebraic integer. Examples of such functions with this property are linear functions. We show that, when n is a prime number, the converse also holds. That is, any function with this property is representable by a linear polynomial. Finally, we give an application of the main result to the problem of determining self perfect isometries for the cyclic group of prime order p .

1. INTRODUCTION

Let n be a positive integer. Denote by $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$ the ring of integers modulo n . Let $\omega = e^{2\pi i/n}$ be a primitive n th root of unity. In this work, we consider the following problem.

Problem 1. Suppose $f : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ is a function such that the average

$$(1) \quad \mu_f^{a,b} = \frac{1}{n} \sum_{x=0}^{n-1} \omega^{af(x)+bx} \quad \text{is an algebraic integer for every } a, b \in \mathbb{Z}_n.$$

What can be said about the function f ?

When n is a power of prime, such a problem is related to the problem of finding self perfect isometries (as defined in [1]) for cyclic p -groups, since the condition (1) is the integrality condition for perfect characters. More explanations on this relationship are given in the last section.

2. PRELIMINARIES

We shall use the symbol $(=)$ to denote ordinary equality. The symbol (\equiv) will be used to denote congruence $(\pmod n)$ (i.e., equality in \mathbb{Z}_n).

Definition 2. We say that a function $f : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ is a *polynomial function* if there exists a polynomial $F \in \mathbb{Z}_n[X]$ such that

$$f(x) \equiv F(x) \quad \text{for all } x = 0, 1, \dots, n-1.$$

First we show that any function representable by a linear polynomial function satisfies the condition (1).

theorem 3. Suppose that $f : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ is a linear polynomial function, then $\mu_f^{a,b}$ is an algebraic integer for every $a, b \in \mathbb{Z}_n$.

Date: January 26, 2020.

Proof. If f is given by a linear polynomial, then so is $af(x) + bx$ for any a, b . Suppose $af(x) + bx \equiv \alpha x + \beta$ for some $\alpha, \beta \in \mathbb{Z}_n$. If $\alpha \equiv 0$, then

$$\mu_f^{a,b} = \frac{1}{n} \sum_{x=0}^{n-1} \omega^\beta = \omega^\beta,$$

which is an algebraic integer. If $\alpha \not\equiv 0$, then $\omega^\alpha \neq 1$. Consequently,

$$\mu_f^{a,b} = \frac{1}{n} \sum_{x=0}^{n-1} \omega^{\alpha x + \beta} = \frac{\omega^\beta}{n} \left(\frac{1 - \omega^{\alpha n}}{1 - \omega^\alpha} \right) = 0,$$

which is also an algebraic integer. \square

The next result shows that it suffices to check (1) for $a \equiv 0$ (which is trivial) or $b \equiv 0$ or a, b relatively prime.

Proposition 4. *Let $f : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ be a function. Suppose that $\mu_f^{a,b}$ is an algebraic integer. Then $\mu_f^{ka,kb}$ is also an algebraic integer for any k relatively prime to n .*

Proof. Since k and n are coprime, we can define an automorphism $\sigma_k \in \text{Aut}(\mathbb{Q}(\omega)/\mathbb{Q})$ by $\sigma_k(\omega) = \omega^k$. Then

$$\mu_f^{ka,kb} = \frac{1}{n} \sum_{x=0}^{n-1} \omega^{k(af(x)+b)} = \frac{1}{n} \sum_{x=0}^{n-1} \sigma_k(\omega)^{af(x)+b} = \sigma_k(\mu_f^{a,b}).$$

Since $\mu_f^{a,b}$ is an algebraic integer and $\sigma_k \in \text{Aut}(\mathbb{Q}(\omega)/\mathbb{Q})$, it follows that $\mu_f^{ka,kb} = \sigma_k(\mu_f^{a,b})$ is also an algebraic integer. \square

Finally, we give a necessary and sufficient condition for the average of roots of unity to be an algebraic integer. This is a standard result in algebraic number theory.

theorem 5. *Let $\omega_1, \dots, \omega_n$ be roots of unity. Their average is an algebraic integer if and only if either $\omega_1 + \dots + \omega_n = 0$ or $\omega_1 = \dots = \omega_n$.*

Proof. The sufficiency is clear. Let μ denote the average of $\omega_1, \dots, \omega_n$ and assume that it is an algebraic integer. By the triangle inequality, $|\mu| \leq 1$ with equality if and only if $\omega_1 = \dots = \omega_n$. Moreover, $|\mu'| \leq 1$ for all algebraic conjugates μ' of μ . If not all ω_i 's are equal, then $|\mu| < 1$. As a result, we also have $|\alpha| < 1$, where α is the product of all algebraic conjugates of μ . But α is an integer, which implies that α must be 0. It follows that $\mu = 0$. \square

3. THE CASE WHERE n IS A PRIME

Let $n = p$ be a prime number. Our main result is to show that any function $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ satisfying (1) must be representable by a linear polynomial. To study a function $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$, it suffices to study polynomials of degree at most $p - 1$. The following lemma was proved (for a general finite field) by Dickson [2].

theorem 6. *For any function $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$, there exists a unique polynomial $F \in \mathbb{Z}_p[X]$ of degree at most $p - 1$ such that*

$$f(x) \equiv F(x) \pmod{p} \quad \text{for all } x = 0, 1, \dots, p - 1.$$

Henceforth, we shall identify a function $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ with its corresponding polynomial of degree at most $p - 1$. For the proof of our main result, we will need to consider the following set.

Definition 7. For a polynomial f in $\mathbb{Z}_p[X]$, define

$$W_f = \{\lambda \in \mathbb{Z}_p : x \mapsto (f(x) + \lambda x) \text{ is a permutation on } \mathbb{Z}_p\}.$$

theorem 8 (Stothers [3, Theorem 2]). *Let f be a polynomial in $\mathbb{Z}_p[X]$ of degree at most $p - 1$. If $|W_f| > (p - 3)/2$, then $\deg(f) \leq 1$.*

We now state our main result.

Theorem 9. *Let $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be a function. Suppose that the average*

$$\mu_f^{1,b} = \frac{1}{p} \sum_{x=0}^{p-1} \omega^{f(x)+bx}$$

is an algebraic integer for every $b \in \mathbb{Z}_p$. Then f is a linear polynomial function.

Proof. By Lemma 5, we have that, for each b , either $f(x) + bx$ is constant modulo p or $\mu_f^{1,b} = 0$. Suppose there is $b_0 \in \mathbb{Z}_p$ such that $f(x) + b_0x$ is constant modulo p . Then it is clear that f is of the form $f(x) \equiv \alpha x + \beta$ for all $x \in \mathbb{Z}_p$.

If there is no b with $f(x) + bx$ constant modulo p , then $\sum_{x=0}^{p-1} \omega^{f(x)+bx} = 0$ for all b . Since the minimal polynomial of ω is $1 + X + \dots + X^{p-1}$, we must have that $x \mapsto f(x) + bx$ is a permutation modulo p , for all b . This means that the set W_f has cardinality p . Hence, by Lemma 8, f is a linear polynomial. \square

Corollary 10. *Let $f : \mathbb{Z}_p \rightarrow \mathbb{Z}_p$ be a function. Suppose that the average*

$$\mu_f^{a,b} = \frac{1}{p} \sum_{x=0}^{p-1} \omega^{f(x)+bx}$$

is an algebraic integer for every $a, b \in \mathbb{Z}_p$. Then f is a linear polynomial function.

Proof. Take $a = 1$ and apply Theorem 9. \square

We have seen (in Lemma 3) that if a function $f : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ is representable by a linear polynomial, then it satisfies (1). Corollary 10 implies that the converse also holds for n prime. It is natural to make the following conjecture.

Conjecture 11. *Suppose $f : \mathbb{Z}_n \rightarrow \mathbb{Z}_n$ is a function such that the average $\mu_f^{a,b}$ is an algebraic integer for every $a, b \in \mathbb{Z}_n$. Then there exist $\alpha, \beta \in \mathbb{Z}_n$ such that $f(x) \equiv \alpha x + \beta$ for all x .*

Remark 12. It is possible that a function f may have a non-linear form and still satisfy (1). The conjecture says that such function should be representable by a linear polynomial in $\mathbb{Z}_n[X]$.

For example, when $n = 6$, it can be checked that $f(x) \equiv x^3 + x$ satisfies (1). However f is representable by a linear polynomial, since $x^3 + x \equiv 2x$ for all $x \in \mathbb{Z}_6$.

4. CONNECTION TO PERFECT ISOMETRIES

Corollary 10 has an application in representation theory of finite groups, especially in the problem of finding self perfect isometries for the cyclic group C_p of order prime p . For the purpose of illustrations, we will define perfect isometries specifically for this special case. Interested readers are referred to [1] for the definition of perfect isometries for general blocks of finite groups.

Throughout this section, let $G = C_p$. Denote by $\mathcal{R}(G)$ the free abelian group generated by $\text{Irr}(G)$, the set of all irreducible complex characters of G . We will regard $\mathcal{R}(G)$ as lying in $CF(G)^1$, the space of complex-valued class functions of G .

Let $I : \mathcal{R}(G) \rightarrow \mathcal{R}(G)$ be a linear map. Define a generalized character μ_I of $G \times G$ by

$$\mu_I(g, h) = \sum_{\chi \in \text{Irr}(G)} I(\chi)(g)\chi(h), \quad \text{for all } g, h \in G.$$

Definition 13. (Cf. Definition 1.1 in [1]) An isometry $I : \mathcal{R}(G) \rightarrow \mathcal{R}(G)$ is said to be a *perfect isometry* if μ_I satisfies the following two conditions.

- (i) (Integrality) For all $g, h \in G$, the number $\mu_I(g, h)/p$ is an algebraic integer.
- (ii) (Separation) If $\mu_I(g, h) \neq 0$, then both g and h are the identity element or both are not.

Let $\omega = e^{2\pi i/p}$. Suppose that G is generated by an element $u \in G$. For $x = 0, 1, \dots, p-1$, let χ_x be the irreducible complex character of G such that

$$\chi_x(u^a) = \omega^{ax}, \quad a = 0, 1, \dots, p-1.$$

In particular, χ_0 is the trivial character.

Any bijection f on the set $\{0, 1, \dots, p-1\}$ gives rise to an isometry $I_f : \mathcal{R}(G) \rightarrow \mathcal{R}(G)$ defined (on the basis) by

$$I_f(\chi_x) = \chi_{f(x)}, \quad x = 0, 1, \dots, p-1.$$

Proposition 14. *An isometry I_f is perfect if and only if f is a linear bijection.*

Proof. Since every element in G is of the form u^a for some a , we have

$$\mu_I(g, h) = \mu_I(u^a, u^b) = \sum_{x=0}^{p-1} I(\chi_x)(u^a)\chi_x(u^b) = \sum_{x=0}^{p-1} \omega^{af(x)+bx}.$$

Thus, we see that the condition in (1) is precisely the requirement that μ_I satisfies the integrality condition. This is the only condition to consider, as the separation condition is satisfied for any bijection f .

If f is a linear bijection, then by Lemma 3, the integrality condition is satisfied. Thus, I_f is a perfect isometry.

Conversely, if I_f is a perfect isometry, then $\mu_I(u^a, u^b)/p$ is an algebraic integer for all a, b . It follows from Corollary 10 that f must be linear. \square

Remark 15. The following actions on $\text{Irr}(G)$ are well known to give bijections on the set.

- (Multiplication by a linear character) For a fixed $\chi_\beta \in \text{Irr}(G)$, multiplication by χ_β gives a bijection

$$I_\beta : \text{Irr}(G) \rightarrow \text{Irr}(G), \quad I_\beta(\chi_x)(g) = (\chi_\beta\chi_x)(g) = \chi_{\beta+x}(g).$$

¹This is an inner product space with the standard inner product of group characters.

- (Automorphism action) For a fixed $\alpha \in \{1, 2, \dots, p-1\}$, the automorphism $g \mapsto g^\alpha$ induces a bijection

$$I_\alpha : \text{Irr}(G) \longrightarrow \text{Irr}(G), \quad I_\alpha(\chi_x)(g) = \chi_x(g^\alpha) = \chi_{\alpha x}(g).$$

Proposition 14 implies that, for an isometry induced by a bijection on $\text{Irr}(G)$ to be a perfect isometry, it must be a composition of the above two types of isometries.

REFERENCES

- [1] M. Broué, Isométries parfaites, types de blocs, catégories dérivées, *Astérisque* **181** no. 182 (1990) 61–192.
- [2] L. R. Dickson, The analytic representation of substitutions on a power of a prime number of letters with a discussion of the linear group, *Ann. of Math.* **11** no. 1/6 (1896) 65–120.
- [3] W. W. Stothers, On permutation polynomials whose difference is linear, *Glasgow Math. J.* **32** no. 2 (1990) 165–171.

(Chatchawan Panraksa) SCIENCE DIVISION, MAHIDOL UNIVERSITY INTERNATIONAL COLLEGE,
999 PHUTTHAMONTHON 4 ROAD, SALAYA, NAKHONPATHOM, THAILAND 73170

E-mail address: `chatchawan.pan@mahidol.ac.th`

(Pornrat Ruengrot) SCIENCE DIVISION, MAHIDOL UNIVERSITY INTERNATIONAL COLLEGE, 999
PHUTTHAMONTHON 4 ROAD, SALAYA, NAKHONPATHOM, THAILAND 73170

E-mail address: `pornrat.rue@mahidol.ac.th`