Wreaths, mixed wreaths and twisted coactions

Ross Street*

Centre of Australian Category Theory, Macquarie University, NSW 2109, Australia <ross.street@mq.edu.au>

July 26, 2021

2010 Mathematics Subject Classification: 18D10; 05A15; 18A32; 18D05; 20H30; 16T30

Key words and phrases: monad; comonad; wreath; Heisenberg product; convolution; mixed distributive law; twisted action; bialgebra.

Abstract

Distributive laws between two monads in a 2-category \mathcal{K} , as defined by Jon Beck in the case $\mathcal{K} = \operatorname{Cat}$, were pointed out by the author to be monads in a 2-category Mnd \mathcal{K} of monads. Steve Lack and the author defined wreaths to be monads in a 2-category EM \mathcal{K} of monads with different 2-cells from Mnd \mathcal{K} .

Mixed distributive laws were also considered by Jon Beck, Mike Barr and, later, various others; they are comonads in $\mathrm{Mnd}\mathcal{K}$. Actually, as pointed out by John Power and Hiroshi Watanabe, there are a number of dual possibilities for mixed distributive laws.

It is natural then to consider mixed wreaths as we do in this article; they are comonads in $EM\mathscr{K}$. There are also mixed opwreaths: comonoids in the Kleisli construction completion $Kl\mathscr{K}$ of \mathscr{K} . The main example studied here arises from a twisted coaction of a bimonoid on a monoid. Corresponding to the wreath product on the mixed side is wreath convolution, which is composition in a Kleisli-like construction. Walter Moreira's Heisenberg product of linear endomorphisms on a Hopf algebra, is an example of such convolution, actually involving merely a mixed distributive law. Monoidality of the Kleisli-like construction is also discussed.

 $^{^*}$ The author gratefully acknowledges the support of Australian Research Council Discovery Grant DP130101969.

Contents

1	Introduction	2
2	Review of wreaths	3
3	Mixed wreaths	9
4	Twisted coactions	13
5	Monoidality	18
6	Monoidal twisted coactions	21

1 Introduction

While trying to expose the categorical mechanism behind the Heisenberg product of endomorphisms, as defined and studied in [1, 26], we noticed that it has to do with distributive laws in the sense of Beck [2]; also see [30, 32] for the general setting and monad terminology. A distributive law $\xi \colon TS \Rightarrow ST$ of a monad T over a monad S on a category $\mathscr A$ gives rise to a monad structure on the composite ST. For a mixed distributive law $\zeta \colon SG \Rightarrow GS$ of a comonad G over a monad S, we do not expect GS to be a monad or comonad, so what takes its place? It is the (internalized) ζ -parametrized convolution of 2-cells $G\Rightarrow S$. To understand this to some extent (externally), consider the case where $T\dashv G$, and ξ and ζ are mates [21] under that adjunction. The adjunction gives an isomorphism

$$[\mathscr{A}, \mathscr{A}](1_\mathscr{A}, ST) \cong [\mathscr{A}, \mathscr{A}](G, S)$$

The monoid structure on the left-hand side, arising pointwise from the monad structure on ST determined by ξ , transports to a convolution-like monoid structure on the right-hand side, expressible in terms of ζ .

Rather than remain at the level of distributive law, since there are articles [14, 29, 13] which study that, we decided to generalize to the wreaths of [23].

This article begins with a review of wreaths as defined in [23]. We spend some time extending Example 3.2 of [23] to a wreath between monoids rather than groups: the use of fibrations is to bring out the cohomological aspects which permeate the paper.

As for mixed distributive laws [29], there are several possibilities for mixed wreaths. We look at those which are comonads in either the (limit) completion of the ambient 2-category under Eilenberg-Moore construction or the (colimit) completion under the Kleisli construction. The first are called mixed wreaths, the second mixed opwreaths. Mixed Eilenberg-Moore and mixed Kleisli constructions are described and their universal properties presented. Composition in a mixed Kleisli category is convolution parametrized by the mixed wreath.

Section 4 provides the construction of a mixed opwreath is a dual of the wreath construction appearing as Example 3.3 in [23] based on Sweedler's crossed product of Hopf algebras. We also generalize to bialgebras (bimonoids). The ingredient is a twisted coaction of a bimonoid on a monoid. Natural connections to cohomological structures are pursued.

Section 5 sets out when a mixed opwreath is opmonoidal. This is about a monoidal structure on the mixed Kleisli construction. The final section gives structure on a twisted coaction so that the associated mixed opwreath becomes opmonoidal.

We will use the string diagrams for monoidal categories as explained in [17]. However, we read the diagrams from top to bottom rather than the reverse. For example, if A is a monoid in any monoidal category \mathscr{V} , the multiplication $\mu = \mu_A$ and unit $\eta = \eta_A$ are respectively depicted as follows.




If we are dealing with a braided monoidal category, the braiding $c_{X,Y} \colon X \otimes Y \to Y \otimes X$ will be depicted as a crossing as follows.



2 Review of wreaths

The free completion EM(\mathfrak{K}) of a 2-category \mathfrak{K} (such as Cat) under the Eilenberg-Moore construction was identified in [23]. The objects of EM(\mathfrak{K}) are monads (\mathscr{A}, T) in \mathfrak{K} . That is, T is a monoid in the endomorphism category $\mathfrak{K}(\mathscr{A}, \mathscr{A})$, monoidal under composition as tensor product, so we can draw planar diagrams. A morphism $(F, \phi) \colon (\mathscr{A}, T) \to (\mathscr{B}, S)$ consists of a morphism $F \colon \mathscr{A} \to \mathscr{B}$ and a 2-cell $\phi \colon SF \Rightarrow FT$ in \mathfrak{K} compatible with the monad structures on T and S. A 2-cell $\rho \colon (F, \phi) \Rightarrow (G, \psi)$ is a 2-cell $\rho \colon F \Rightarrow GS$ such that the following equation holds.



Also in [23], wreaths were introduced and defined concisely as monads in the 2-category $EM(\mathfrak{K})$. The wreath product is the monad obtained as the Eilenberg-Moore construction in $EM(\mathfrak{K})$ on the wreath. Indeed, as always with a completion under limits, EM is a (dual) Kock-Zöberlein monad [22, 31] on the 2-category of 2-categories while the wreath product

$$\operatorname{wr} = \operatorname{wr}_{\mathfrak{K}} : \operatorname{EM}(\operatorname{EM}(\mathfrak{K})) \longrightarrow \operatorname{EM}(\mathfrak{K})$$

gives the multiplication for that monad; the unit

$$id = id_{\mathfrak{K}} \colon \mathfrak{K} \longrightarrow EM(\mathfrak{K})$$

simply takes each object \mathscr{A} to the identity monad on \mathscr{A} . When it exists, the Eilenberg-Moore construction for a 2-category \mathfrak{K} is a right adjoint


$$em = em_{\mathfrak{K}} : EM(\mathfrak{K}) \longrightarrow \mathfrak{K}$$

to id; we put $\mathscr{A}^T = \operatorname{em}_{\mathscr{A}} T = \operatorname{em}(\mathscr{A}, T)$.


We will now describe wreaths explicitly using string diagrams.

Let $T = (T, \mu, \eta)$ be a monad on an object \mathscr{A} of \mathscr{K} .

A wreath around T consists of an endomorphism S on \mathscr{A} , and 2-cells $\nu \colon SS \Longrightarrow ST$, $\sigma \colon 1_{\mathscr{A}} \Longrightarrow ST$ and $\lambda \colon TS \Longrightarrow ST$ satisfying seven axioms.



The product of the wreath S around T, or the wreath product, is the monad consisting of the endomorphism ST on $\mathscr A$ with the multiplication and unit as displayed in the diagram:



where the unlabelled nodes are the ternary and binary multiplications of T.

A distributive law [2] of a monad T over a monad S is a special case of a wreath around T consisting of the endofunctor S while the natural transformations ν and σ of the special form

$$\nu = \left(SS \xrightarrow{\mu} S \xrightarrow{S\eta} ST\right) \text{ and } \sigma = \left(1_{\mathscr{A}} \xrightarrow{\eta\eta} ST\right) , \qquad (2.1)$$

and λ remains arbitrary.

Example 1. We now generalise Example 3.2 of [23] from groups to monoids. We call a monoid morphism $p: E \to M$ (in the category Set of sets) a normal cloven lax fibration when it is equipped with a function $j: M \to E$ such that $p \circ j = 1_M$, j(1) = 1 and, for

$$A = \{a \in E \colon p(a) = 1\} = p^{-1}(1) ,$$

the function $h: M \times A \to E$, defined by h(x, a) = j(x)a, is invertible. This gives, for each $x \in M$, a pullback square.

$$\begin{array}{ccc}
A & \xrightarrow{h(x,-)} & E \\
\downarrow \downarrow & & \downarrow p \\
1 & \xrightarrow{x} & M
\end{array}$$
(2.2)

Generally, the kernel of a monoid morphism is a rather strange thing to consider, yet, because we have a fibration, all the fibres $p^{-1}(x)$ of p are isomorphic as sets. Unlike arbitrary fibres, the kernel has the advantage of being a submonoid of E.

We use the pullback (2.2) to obtain a function $\alpha \colon A \times M \to A$ of M on A; indeed, $\alpha(a,x) = a \cdot x \in A$ is characterized by the property

$$j(x)(a \cdot x) = aj(x) . (2.3)$$

In other words, this α measures the failure of the kernel to commute with the image of j. Using the pullback uniqueness clause, we see that each $-\cdot x \colon A \to A$ is a monoid morphism.

We also use the pullback (2.2) with x replaced by xy to obtain a function $\rho: M \times M \to A$ characterized by the property

$$j(xy)\rho(x,y) = j(x)j(y) . (2.4)$$

In other words, ρ measures the failure of j to be a monoid morphism. Indeed, for each $x, y \in M$, we have a 2-cell

$$\begin{array}{ccc}
A & \xrightarrow{-\cdot x} & A \\
& & & \\
-\cdot (xy) & & & \\
A & & & \\
\end{array}$$
(2.5)

in the 2-category Mon = MonSet of monoids; the "naturality" amounts to the equation

$$(a \cdot (xy))\rho(x,y) = \rho(x,y)((a \cdot x) \cdot y) \tag{2.6}$$


which shows that ρ also measures the failure of α to be an action of the monoid M on A. To prove (2.6), it suffices to prove we have equality after applying h(xy, -), which we do thus:

$$\begin{split} j(xy)(a\cdot(xy))\rho(x,y) &= aj(xy)\rho(x,y) \\ &= aj(x)j(y) \\ &= j(x)(a\cdot x)j(y) \\ &= j(x)j(y)((a\cdot x)\cdot y) \\ &= j(xy)\rho(x,y)((a\cdot x)\cdot y) \;. \end{split}$$

Let ΣM denote the category with one object 0 and hom $\Sigma M(0,0) = M$; composition is multiplication in M. What we are producing is a normal lax functor

$$P: \Sigma M^{\mathrm{op}} \longrightarrow \mathrm{Mon}$$
 (2.7)

with P0 = A and $Px = -\cdot x : A \to A$. The composition constraints are given by (2.5). Clearly $\rho(1,x) = 1 = \rho(x,1)$ so all that remains to prove is the coherence condition (2.8).




This amounts to the Schreier factor set or 2-cocycle condition (2.9).

$$\rho(xy,z)(\rho(x,y)\cdot z) = \rho(x,yz)\rho(y,z) \tag{2.9}$$

To prove this, it suffices to check after left multiplication by j(xyz), which we do thus:

$$\begin{split} j(xyz)\rho(xy,z)(\rho(x,y)\cdot z) &= j(xy)j(z)(\rho(x,y)\cdot z) \\ &= j(xy)\rho(x,y)j(z) \\ &= j(x)j(y)j(z) \\ &= j(x)j(yz)\rho(y,z) \\ &= j(xyz)\rho(x,yz)\rho(y,z) \;. \end{split}$$

We are now in a position to transport the multiplication of E to $M \times A$ across the isomorphism h.



The resultant multiplication on $M \times A$ is


$$(x,a)(y,b) = (xy, \rho(x,y)(a \cdot y)b)$$
 (2.10)

This gives an equivalence of categories between normal cloven lax fibrations over any monoid M and normal lax functors $P \colon \Sigma M^{\mathrm{op}} \longrightarrow \mathrm{Mon}$. This is essentially classical and is an interpretation theorem for the second cohomology of the monoid M: 2-cocycles equate to certain extensions $E \to M$.


Now we give the wreath. The category is Set. The remaining data all arise from data in Set by applying the strong monoidal functor Set \to [Set, Set] which takes K to $K \times -$. The monad T arises from the monoid A. The endofunctor S arises from the set M. The natural transformation ν arises from the function $M \times M \to M \times A$, $(x,y) \mapsto (xy,\rho(x,y))$. The natural transformation λ arises from the function $A \times M \to M \times A$, $(a,x) \mapsto (x,\alpha(a,x))$. The natural transformation σ arises from the function $\mathbf{1} \to M \times A$ which picks out (1,1).

The wreath product of course arises from the monoid $M \times A$ with product (2.10) and so recaptures E up to isomorphism.

Remark 2. 1. Here is the string diagram for (2.6).




2. Here is the string diagram for (2.9).



3. The structure on $p: E \to M$ of normal cloven (strict) fibration consists of a function $j: M \to E$ such that $p \circ j = 1_M$, j(1) = 1 and the square

$$\begin{array}{c} M \times E \xrightarrow{\mu \circ (j \times 1_E)} E \\ \downarrow^{1_M \times p} \downarrow & \downarrow^p \\ M \times M \xrightarrow{\mu} M \end{array}$$

is a pullback. Notice that we have the condition $M \times A \cong E$ for a lax fibration since we can paste two pullback squares as follows:



4. Similarly to Example 1, Example 3.3 of [23] can be generalised from Hopf algebras H to bimonoids M in a braided monoidal category. Moreover, there is no need for the convolution invertibility of ρ : however, the one axiom required for A to be a twisted M-module, which is stated in [23] in terms of that inverse of ρ , should be replaced by the naturality condition (2.11). We will discuss a dual of this in Section 4.

3 Mixed wreaths

There are several possibilities for mixed wreaths just as for mixed distributive laws; compare [29].

We will use the notation

$$\mathrm{EM}^{du}(\mathfrak{K}) = \mathrm{EM}(\mathfrak{K}^{du})^{du}$$

for any of the dualities $du \in \{op, co, coop\}$ (in the notation of [21]). We also put

$$KL(\mathfrak{K}) = EM^{op}(\mathfrak{K})$$

since it is the cocompletion of $\mathfrak K$ with respect to the Kleisli construction. This then leads to

$$KL^{co}(\mathfrak{K}) = EM^{coop}(\mathfrak{K})$$
.

Definition 3. Let T be a monad on \mathscr{A} in the 2-category \mathfrak{K} . A mixed wreath around the monad T is a comonad on (\mathscr{A},T) in the 2-category $\mathrm{EM}(\mathfrak{K})$.

More explicitly, a mixed wreath structure around T on an endomorphism G of \mathscr{A} consists of 2-cells $\delta \colon G \Rightarrow GGT$, $\varepsilon \colon G \Rightarrow T$ and $\xi \colon TG \Rightarrow GT$ satisfying four axioms which say that $(G,\xi) \colon (\mathscr{A},T) \to (\mathscr{A},T)$ is a morphism, and $\delta \colon (G,\xi) \Rightarrow (G,\xi)(G,\xi)$ and $\varepsilon \colon 1 \Rightarrow (G,\xi)$ are 2-cells, in $\mathrm{EM}(\mathfrak{K})$, and three axioms which say δ is coassociative with counit ε .

Suppose \mathfrak{K} admits the Eilenberg-Moore construction for both monads and comonads. Simply because em: $\mathrm{EM}(\mathfrak{K}) \longrightarrow \mathfrak{K}$ is a 2-functor, each mixed wreath $(G,\xi)\colon (\mathscr{A},T) \to (\mathscr{A},T)$ yields a comonad $G^{\xi}=\mathrm{em}(G,\xi)$ on \mathscr{A}^T in \mathfrak{K} . Define


$$mem(G, \xi, T) = (\mathscr{A}^T)^{G^{\xi}},$$

the Eilenberg-Moore construction for the comonad G_{ξ} . This gives the object function for a mixed Eilenberg-Moore construction

$$\operatorname{mem} \colon \operatorname{EM}^{\operatorname{co}}(\operatorname{EM}(\mathfrak{K})) \stackrel{\operatorname{EM}^{\operatorname{co}}(\operatorname{em})}{\longrightarrow} \operatorname{EM}^{\operatorname{co}}(\mathfrak{K}) \stackrel{\operatorname{em}^{\operatorname{co}}}{\longrightarrow} \mathfrak{K}$$
 (3.13)

with an obvious left adjoint.


We have the following description of mem (G, ξ, T) when $\mathfrak{K} = \operatorname{Cat}$. The objects (A, a, c) of the category consist of an Eilenberg-Moore T-algebra $a \colon TA \to A$ and a morphism $c \colon A \to GA$ satisfying the following three conditions.



To reinforce the limit nature of the mem (G, ξ, T) construction we next record its characterization as a representing object. This can be taken as the definition when \mathfrak{K} lacks the Eilenberg-Moore construction for monads or comonads in general.

Proposition 4. $\mathfrak{K}(\mathscr{X}, \text{mem}(G, \xi, T)) \cong \text{mem}(\mathfrak{K}(\mathscr{X}, G), \mathfrak{K}(\mathscr{X}, \xi), \mathfrak{K}(\mathscr{X}, T))$

Definition 5. Let T be a monoid on \mathscr{A} in the 2-category \mathfrak{K} . A mixed opwreath around the monad T is a comonad on (\mathscr{A},T) in the 2-category $\mathrm{KL}(\mathfrak{K})$. This consists of an endomorphism G of \mathscr{A} made into a morphism of $\mathrm{KL}(\mathfrak{K})$ by a 2-cell $\zeta \colon GT \Rightarrow TG$ and into a comonad by comultiplication $\delta \colon G \Rightarrow TGG$ and counit $\varepsilon \colon G \Rightarrow T$. The seven axioms are shown below in string form.



At the 2-category level Definition 5 really is just an example: a mixed opwreath in \mathfrak{K} is a mixed wreath in \mathfrak{K}^{op} . Indeed, in the presence of right adjoints, we will now point out how a mixed opwreath amounts to a wreath.

Recall from [21] the terminology and concept of mates under adjunction. Here is an exercise on mates using the string calculus.

Proposition 6. Suppose T is a monad on $\mathscr{A} \in \mathfrak{K}$. Suppose $G \dashv S$ are adjoint endomorphisms of \mathscr{A} . Mixed opwreath structures on G around T correspond under adjoint mateship to wreath structures on S around T.

In the situation of Proposition 6, the Eilenberg-Moore construction for the wreath product ST does not easily reinterpret in terms of G and T, rather, as you would expect, the Kleisli construction does. We shall now define this in general.

By applying Proposition 4 to \mathfrak{K}^{op} , and by defining the *mixed Kleisli construction* as the composite

mkl:
$$KL^{co}(KL(\mathfrak{K})) \xrightarrow{KL^{co}(kl)} KL^{co}(\mathfrak{K}) \xrightarrow{kl^{co}} \mathfrak{K}$$
, (3.14)

we obtain:

Proposition 7. $\mathfrak{K}(\mathrm{mkl}(G,\zeta,T),\mathscr{X}) \cong \mathrm{mem}(\mathfrak{K}(G,\mathscr{X}),\mathfrak{K}(\zeta,\mathscr{X}),\mathfrak{K}(T,\mathscr{X}))$

For $\mathfrak{K}=\mathrm{Cat}$, the category $\mathrm{mkl}(G,\zeta,T)$ has the same objects as \mathscr{A} and has homsets defined by

$$mkl(G, \zeta, T)(A, B) = \mathscr{A}(GA, TB)$$
.


Composition is defined by wreath convolution: the composite of $f: GA \to TB$ and $g: GB \to TC$ is $g \circ f = f *_{\zeta} g$ as in the commutative diagram (3.15).

$$GA \xrightarrow{\delta_A} TGGA \xrightarrow{TGf} TGTB$$

$$f *_{\zeta} g \downarrow \qquad \qquad \downarrow T\zeta_B$$

$$TC \xleftarrow{\mu_{3C}} TTTC \xleftarrow{}_{TTq} TTGB$$

$$(3.15)$$




- **Remark 8.** 1. For any monoidal category \mathscr{V} , the functor $\mathscr{V} \to [\mathscr{V}, \mathscr{V}]$, taking X to $X \otimes -$, is strong monoidal. So each monoid A = (A, m, j) in \mathscr{V} is taken to a monad $T = (T, \mu, \eta)$ on \mathscr{V} . We will speak of a mixed opwreath around A to mean a quadruple (C, d, w, z) consisting of an object C and morphisms $d \colon C \to A \otimes C \otimes C$, $w \colon C \to A$ and $z \colon C \otimes A \to A \otimes C$ satisfying the string diagram conditions of Definition 5 with $T, \mu, \eta, G, \delta, \varepsilon, \zeta$ replaced by A, m, j, C, d, w, z, respectively. Moreover, since mixed opwreaths are defined purely in terms of the monoidal structure, each mixed opwreath (C, d, w, z) around A in \mathscr{V} defines gives rise to a mixed opwreath $(G, \delta, \varepsilon, \zeta)$ around the monad $T = A \otimes -$. Furthermore, we write mkl(C, z, A) for the category $\text{mkl}(G, \zeta, T)$.
 - 2. For any monoidal category \mathscr{V} , the functor $\mathscr{V}^{\mathrm{rev}} \to [\mathscr{V}, \mathscr{V}]$, taking X to $-\otimes X$, is also strong monoidal. Thus the mixed opwreath around A as in Item 1 is taken to a mixed wreath around the monad $-\otimes A$ on \mathscr{V} . We write $\mathrm{mem}(C,z,A)$ for the mixed Eilenberg-Moore construction (3.13) applied to this mixed wreath.

Definition 9. The set $\text{mkl}(C, z, A)(I, I) \cong \mathcal{V}(C, A)$ of endomorphisms of I in the category mkl(C, z, A) is, of course, a monoid under composition. The multiplication might be called z-parametrized convolution on $\mathcal{V}(C, A)$.

Example 10 (The Heisenberg category). Suppose A = (A, m, j, c, e) is a bimonoid in the braided monoidal category \mathscr{V} . Denote the braiding by σ . We obtain a mixed opwreath on the monoid A = (A, m, j) (indeed it is a mixed opdistributive law) in \mathscr{V} by taking the comonoid C to be A = (A, c, e), z to be

$$z_{h} = \left(A \otimes A \overset{1_{A} \otimes \sigma_{A,A}^{-1} c}{\longrightarrow} A \otimes A \otimes A \overset{\sigma_{A,A}^{-1} \otimes 1_{A}}{\longrightarrow} A \otimes A \otimes A \overset{1_{A} \otimes m}{\longrightarrow} A \otimes A \right)$$
$$= \left(A \otimes A \overset{1_{A} \otimes c}{\longrightarrow} A \otimes A \otimes A \overset{\sigma_{A,A}^{-1} \otimes 1_{A}}{\longrightarrow} A \otimes A \otimes A \overset{1_{A} \otimes m}{\longrightarrow} A \otimes A \right) ,$$

d to be $j \otimes c : A \to A \otimes A \otimes A$, and w to be $j \circ e : A \to A$.



We put $\mathrm{Hb}(A) = \mathrm{mkl}(A, z_{\mathrm{h}}, A)$ and call it the *Heisenberg category* of the bimonoid A in \mathscr{V} . Here is the reason.

Proposition 11. Suppose $\mathcal{V} = \text{Vect}$ is the symmetric monoidal category of vector spaces over a fixed field and H is a Hopf algebra, then the z_h -parametrized convolution of linear endomorphisms of H is the Heisenberg product as defined in [26, 1].


4 Twisted coactions

The construction of a mixed opwreath explained in this section is a dual of the wreath construction appearing as Example 3.3 in [23] based on Sweedler's crossed product of Hopf algebras.

We begin by pointing out that, given a monoid A in any monoidal category \mathcal{V} , the representable functor $\mathcal{V}(-,A)\colon \mathcal{V}^{\mathrm{op}}\to \mathrm{Set}$ becomes monoidal when equipped with the natural family of functions

$$\mathscr{V}(X,A) \times \mathscr{V}(Y,A) \longrightarrow \mathscr{V}(X \otimes Y,A) ,$$
 (4.16)

defined by $(u, v) \mapsto u \bullet v$ as depicted below, and $\eta_A \in \mathcal{V}(I, A)$. The reason is that the Yoneda embedding $\mathcal{V} \to [\mathcal{V}^{\text{op}}, \text{Set}]$ is monoidal, where monoids in the codomain are precisely monoidal functors.



Here are the properties of the dot product:

(unitality)
$$\eta_A \bullet u = u = u \bullet \eta_A$$

(associativity) $(u \bullet v) \bullet w = u \bullet (v \bullet w)$
(naturality) $(u \bullet v) \circ (f \otimes g) = (u \circ f) \bullet (v \circ g)$

We also recall that, if \mathscr{V} is (lax) braided then the tensor product $A \otimes B$ of monoids A and B is again a monoid: the (lax) braiding gives a distributive law of A over B used in defining $\mu_{A \otimes B}$ in terms of the multiplications μ_A and μ_B .

Definition 12. Let A be a monoid and B be a bimonoid in a (lax) braided monoidal category \mathscr{V} . A twisted (right) coaction of B on A consists of a monoid morphism $\gamma \colon A \to A \otimes B$ and a morphism $\tau \colon I \to A \otimes B^{\otimes 2}$ such that

(counitality)
$$(1_A \otimes \varepsilon_B) \circ \gamma = 1_A$$


 $(\tau\text{-coassociativity}) \ \tau \bullet ((\gamma \otimes 1_B) \circ \gamma) = ((1_A \otimes \delta_B) \circ \gamma) \bullet \tau$

(2-cocyclicity)
$$((1_A \otimes \delta_B \otimes 1_B) \circ \tau) \bullet (\tau \otimes \eta_B) = ((1_{A \otimes B} \otimes \delta_B) \circ \tau) \bullet ((\gamma \otimes 1_{B \otimes B}) \circ \tau))$$

(normality) $(1_{A \otimes B} \otimes \varepsilon_B) \circ \tau = \eta_A \otimes \eta_B = (1_A \otimes \varepsilon_B \otimes 1_B) \circ \tau$.

Note that we do not require τ to be ullet-invertible.

Here in string form are the conditions on a twisted coaction.




Proposition 13. Given a twisted coaction of a monoid A on a bimonoid B in a braided monoidal category \mathcal{V} , using the notation of Definition 12, a mixed opwreath around A is defined by the comonoid B equipped with the morphisms

$$\zeta = (\eta_A \otimes 1_B) \bullet \gamma \colon B \otimes A \longrightarrow A \otimes B$$
$$\delta = (\eta_A \otimes \delta_B) \bullet \tau \colon B \longrightarrow A \otimes B \otimes B$$
$$\varepsilon = \eta_A \circ \varepsilon_B \colon B \longrightarrow A$$

as required by Definition 5.

Here are the string diagrams for these ζ , δ and ε .



In proving Proposition 13, a lemma will be useful.

Lemma 14. The following equations hold:

- (i) $\delta_B \bullet \delta_B = \delta_B \circ \mu_B$
- (ii) $(1 \otimes \delta_B) \bullet \tau = (1_A \otimes \eta_B \otimes \eta_B) \bullet \delta$
- (iii) $(\eta_A \otimes \delta_B) \bullet ((1 \otimes \delta_B) \circ \gamma) = ((1 \otimes \delta_B) \circ \zeta)$

Proof. Item (i) is a restatement of the bimonoid axiom for B asserting that μ_B preserves comultiplication. Item (ii) is immediate on drawing the string diagrams. Item (iii) is immediate from the string diagrams and using the bimonoid axiom. \square

Here now are some clues on proving Proposition 13. There are seven conditions satisfied by the twisted coaction. There are seven axioms to verify for the mixed opwreath. For condition 1, we can express the fact that γ preserves multiplication in the form $\gamma \circ \mu_A = \gamma \bullet \gamma$, then dot both sides on the left with $\eta_A \otimes 1_B$. Condition 2 follows by dotting on both sides by $\eta_A \otimes 1_B$ the equation expessing the fact that γ preserves unit. Condition 3 follows from counitality of γ and the bimonoid condition that μ_B preserves counit. Condition 4 is obtained by dotting both sides of the τ -coassociativity equation on the left by $\eta_A \otimes \delta_B$ and employing Lemma 14. Condition 5 follows by dotting both sides of the cocyclicity condition by $\eta_A \otimes \mu_{B3}$, where μ_{B3} is the ternary multiplication $\mu_B \circ (\mu_B \otimes 1_B) = \mu_B \circ (1_B \otimes \mu_B)$, and employing Lemma 14. Unsurprisingly by now, conditions 6 and 7 follow from the two equations of normality and that μ_B preserves counit.

Remark 15. If the 2-cocycle τ has the form $\eta_A \otimes \tau'$ for some $\tau' \colon I \to B \otimes B$ then the mixed opwreath of Proposition 13 is a mixed opdistributive law.

There is a 2-categorical viewpoint on twisted coactions. Recall (see [19] or Chapter 15 of [34], for example) that the category Mon $\mathscr V$ of monoids in the braided monoidal category $\mathscr V$ is a monoidal 2-category. If $f,g\colon M\to N$ are monoid morphisms then a 2-cell $\xi\colon f\Rightarrow g\colon M\to N$ is a morphism $\xi\colon I\to N$ in $\mathscr V$ satisfying the naturality condition $\xi\bullet f=g\bullet \xi$. The vertical composite of ξ with a 2-cell $\xi\colon g\Rightarrow h\colon M\to N$ is $\xi\bullet \xi$. The horizontal composite of ξ with a 2-cell $\xi'\colon f'\Rightarrow g'\colon N\to L$ is $\xi'\bullet (f'\circ \xi)=(g'\circ \xi)\bullet \xi'$. The tensor product in Mon $\mathscr V$ is the tensor product of monoids that we have been dealing with already (it uses the braiding of $\mathscr V$ yet is not itself a braided tensor product unless $\mathscr V$ is symmetric).

Now we can think of our bimonoid B as a comonoid in the 2-category Mon \mathscr{V} .

Proposition 16. A twisted right coaction of the bimonoid B on the monoid A in \mathcal{V} is precisely a normal lax right coaction of the comonoid B on the object A in the 2-category Mon \mathcal{V} .

Proof. We need to see what is involved in a normal lax right coaction. Indeed we have a morphism $\gamma \colon A \to A \otimes B$ in Mon \mathcal{V} , as required. We have a 2-cell

$$\begin{array}{ccc}
A & \xrightarrow{\gamma} & A \otimes B \\
\gamma \downarrow & \xrightarrow{\tau} & \downarrow_{1 \otimes \delta_B} \\
A \otimes B & \xrightarrow{\gamma \otimes 1} & A \otimes B \otimes B
\end{array} (4.17)$$

in Mon \mathscr{V} ; the 2-cell condition is precisely τ -coassociativity. A lax coaction also involves a 2-cell $\tau_0: 1_A \Rightarrow (1_A \otimes \varepsilon) \circ \gamma$ however the normality condition is that this should be an identity; this precisely amounts to counitality. The axioms on τ for a lax coaction are precisely cocyclicity and normality.

We now remind the reader of the role that variants of the (algebraist's) simplicial category Δ play as host to generic monoids, comonoids, actions and coactions (see [24]). We write $\Delta_{\perp, \top}$ for the strict monoidal category whose objects are the strictly positive finite ordinals, whose morphisms are order and first-and-last-element preserving functions; the tensor product is $m \oplus n = m + n - 1$ thought of, for the purposes of the value at morphisms, as identifying the last element of m with the first element of n. Similarly, Δ_{\top} denotes the category whose objects are the strictly positive finite ordinals, whose morphisms are order and last-element preserving functions. There is a strict right action

$$\oplus : \Delta_{\top} \otimes \Delta_{\perp, \top} \longrightarrow \Delta_{\top} \tag{4.18}$$

of $\Delta_{\perp,\top}$ on Δ_{\top} defined by $m \oplus n = m + n - 1$ as before except that on morphisms, the left morphism in the operation need not preserve the first element, so the result may not either.

Here is a picture of some generating morphisms of Δ_{\top} .

$$\Delta_{\top} : \qquad 1 \xrightarrow{\begin{array}{c} \partial_{0} \\ \hline \langle \sigma_{0} \\ \hline \rangle \end{array}} 2 \xrightarrow{\begin{array}{c} \partial_{0} \\ \hline \langle \sigma_{0} \\ \hline \langle \sigma_{1} \\ \hline \langle \sigma_{1} \\ \hline \rangle \end{array}} 3 \xrightarrow{\begin{array}{c} \partial_{0} \\ \hline \langle \sigma_{0} \\ \hline \langle \sigma_{1} \\ \hline \langle \sigma_{2} \\ \hline \langle \sigma_{2} \\ \hline \langle \sigma_{2} \\ \hline \rangle \end{array}} \cdots$$
 (4.19)

The corresponding picture for $\Delta_{\perp,\top}$ is obtained by deleting all the morphisms labelled ∂_0 . There is a canonical inclusion $\Delta_{\perp,\top} \rightarrowtail \Delta_{\top}$ which respects the right actions by $\Delta_{\perp,\top}$. The corresponding picture for Δ is obtained by adjoining the object 0 and morphisms $\partial_n \colon n \to (n+1)$. There is a canonical inclusion $\Delta_{\top} \rightarrowtail \Delta$. Moreover, $\Delta^{\text{op}} \cong \Delta_{\perp,\top}$.

A comonoid $B=(B,\delta_B,\varepsilon_B)$ in a monoidal category \mathscr{W} defines a strong monoidal functor $\bar{B}\colon \Delta_{\perp,\top} \longrightarrow \mathscr{W}$ whose value at n is $B^{\otimes (n-1)}$, whose value at $\sigma_r\colon (n+1)\to n$ is

$$\sigma_r = 1_{B^{\otimes r}} \otimes \varepsilon_B \otimes 1_{B^{\otimes (n-r-1)}} \colon B^{\otimes n} \to B^{\otimes (n-1)} , \qquad (4.20)$$

and whose value at $\partial_r : n \to (n+1)$ is

$$\partial_r = 1_{B^{\otimes r}} \otimes \delta_B \otimes 1_{B^{\otimes (n-r-2)}} \colon B^{\otimes (n-1)} \to B^{\otimes n} \ . \tag{4.21}$$

In fact this gives an equivalence of categories implying that, up to isomorphism, the comonoid B can be recaptured from the strong monoidal functor.

Suppose \mathcal{W} acts on a category \mathcal{A} via a functor $\star \colon \mathcal{A} \times \mathcal{W} \to \mathcal{A}$. The comonoid B in \mathcal{W} defines a comonad $-\star B$ on \mathcal{A} . We define a *right action* of B on an object $A \in \mathcal{A}$ to be the structure $\gamma \colon A \to A \star B$ of an Eilenberg-Moore $(-\star B)$ -coalgebra on A. There is a functor $\bar{A} \colon \Delta_{\top} \longrightarrow \mathcal{A}$ whose value at the object n is $A \star B^{\otimes (n-1)}$, whose value at $\partial_0 \colon n \to (n+1)$ is

$$\gamma \otimes 1_{B^{\otimes (n-1)}} \colon A \star B^{\otimes (n-1)} \longrightarrow A \star B^{\otimes n}$$

and whose value at the other morphisms ∂_{r+1} in (4.19) is $\partial_{r+1} = 1_A \star \partial_r$ where ∂_r comes from \bar{B} . Then \bar{A} and \bar{B} comprise an action morphism.

$$\begin{array}{cccc}
\Delta_{\top} \times \Delta_{\perp, \top} & \xrightarrow{\oplus} & \Delta_{\top} \\
\bar{A} \times \bar{B} \downarrow & & \downarrow \bar{A} \\
\mathscr{A} \times \mathscr{W} & \xrightarrow{\star} & \mathscr{A}
\end{array} (4.22)$$

Again, this is part of an equivalence of categories between action morphisms and pairs (A, B).

This is all standard material, albeit maybe not explicitly in the above dual version.

Now suppose \mathscr{A} is a 2-category and the action $\star \colon \mathscr{A} \times \mathscr{W} \to \mathscr{A}$ corresponds to a functor $\mathscr{W} \to [\mathscr{A}, \mathscr{A}]$ into the 2-functor 2-category. Suppose $A \in \mathscr{A}$ has merely a morphism $\gamma \colon A \to A \star B$. We can define \bar{A} on objects and generating morphisms as before but it is not quite a functor.

Proposition 17. A normal lax $(-\star B)$ -coalgebra structure on $\gamma \colon A \to A \star B$ amounts to a normal lax functor structure on $\bar{A} \colon \Delta_{\top} \to \mathscr{A}$ which has its constraints $\bar{A}(\zeta) \circ \bar{A}(\xi) \to \bar{A}(\zeta \circ \xi)$ identities unless neither ξ nor ζ is in $\Delta_{\perp,\top}$. In particular, \bar{A} restricts along the inclusion $\Delta_{\perp,\top} \to \Delta_{\top}$ to a strict functor, that is, a simplicial object of \mathscr{A} .

Proposition 18. In the situation of Proposition 17, suppose A is pointed by a morphism $\eta_A \colon I \to A$ in \mathscr{A} , the comonoid B is pointed by a comonoid morphism $\eta_B \colon I \to B$ in \mathscr{W} , and $\gamma \colon A \to A \star B$ respects the pointings, then each lax functor $\bar{A} \colon \Delta_{\top} \to \mathscr{A}$ extends along the inclusion $\Delta_{\top} \to \Delta$ to a lax functor $\hat{A} \colon \Delta \to \mathscr{A}$ by defining $\hat{A}(0) = I$, $\partial_0 = \eta_A \colon I \to A$, and

$$\partial_n \colon = 1_{A \otimes B^{\otimes (n-1)}} \otimes \eta_B \colon A \otimes B^{\otimes (n-1)} \to A \otimes B^{\otimes n}$$
.

In particular, for braided monoidal $\mathscr V$ and $\mathscr W=\mathscr A=\operatorname{Mon}\mathscr V$ (with the action on itself by its own tensor product), each twisted coaction of a bimonoid B on a monoid A determines a slightly lax (augmented) cosimplicial monoid $\hat A$ in $\mathscr V$ with $\hat A(n)=A\otimes B^{\otimes (n-1)}$. Our terminology that τ is a normalized 2-cocycle is justified by the formulas

$$(\partial_1 \tau) \bullet (\partial_3 \tau) = (\partial_2 \tau) \bullet (\partial_0 \tau)$$
, $\sigma_1 \tau = 1 = \sigma_0 \tau$.

5 Monoidality

The basis of this section is the pioneering work of Day [7, 8].

Suppose $T = (T, \mu, \eta)$ is a monoidal monad on the monoidal category \mathscr{A} . Then the Kleisli category \mathscr{A}_T is canonically monoidal: on objects, which are the same as for \mathscr{A} , the tensor product is that of \mathscr{A} ; on homs it is equal to

$$\mathscr{A}_{T}(X,Y) \times \mathscr{A}_{T}(X',Y') = \mathscr{A}(X,TY) \times \mathscr{A}(X',TY') \xrightarrow{\otimes} \mathscr{A}(X \otimes X',TY \otimes TY')$$

$$\xrightarrow{\mathscr{A}(1_{X \otimes X'},\phi_{Y,Y'})} \mathscr{A}(X \otimes X',T(Y \otimes Y')) = \mathscr{A}_{T}(X \otimes X',Y \otimes Y').$$

The canonical functor $\mathscr{A} \to \mathscr{A}_T$ is strict monoidal.

Definition 19. A mixed opwreath $(G, \zeta, \delta, \varepsilon)$ around the monoidal monad T on \mathscr{A} (see Definition 5) is *opmonoidal* when the lifted comonad $\bar{G} = (\bar{G}, \bar{\delta}, \bar{\varepsilon})$ on \mathscr{A}_T is equipped with opmonoidal structure.

By the dual of the fact that Kleisli categories of monoidal monads are canonically monoidal, for an opmonoidal opwreath, the category $\operatorname{mkl}(G,\zeta,T)$ is canonically monoidal since it is the Kleisli category for the opmonoidal comonoid \bar{G} on \mathcal{A}_T .

Let us spell out the data and axioms involved in Definition 19, and the monoidal structure on $\text{mkl}(G, \zeta, T)$.

The data are morphisms

$$\psi_{X,X'} \colon G(X \otimes X') \longrightarrow T(GX \otimes GX')$$
 (5.23)

indexed by pairs of objects $X, X' \in \mathcal{A}$. There are six axioms.

$$G(X \otimes X') \xrightarrow{\psi_{X,X'}} T(GX \otimes GX') \xrightarrow{T(Gf \otimes Gf')} T(GTY \otimes GTY')$$

$$G(f \otimes f') \downarrow \qquad \qquad \downarrow T(\zeta_Y \otimes \zeta_{Y'})$$


$$G(TY \otimes TY') \qquad \qquad T(TGY \otimes TGY')$$


$$G\phi_{Y,Y'} \downarrow \qquad \qquad \downarrow T\phi_{GY,GY'} \qquad (5.24)$$


$$GT(Y \otimes Y') \qquad \qquad TT(GY \otimes GY')$$


$$\zeta_{Y \otimes Y'} \downarrow \qquad \qquad \downarrow \mu_{GY \otimes GY'}$$

$$TG(Y \otimes Y') \xrightarrow{T\psi_{Y,Y'}} TT(GY \otimes GY') \xrightarrow{\mu_{GY \otimes GY'}} T(GY \otimes GY')$$












Diagram (5.24) expresses the naturality of ψ . Diagrams (5.25), (5.26), (5.27) express the opmonoidality of \bar{G} when equipped with ψ . Diagrams (5.28), (5.29) express the opmonoidality of $\bar{\delta}$, $\bar{\varepsilon}$, respectively. One of the other conditions is that the nullary piece ψ_0 of opmonoidal structure on \bar{G} must be ε_I . This means that the nullary conditions for $\bar{\delta}$, $\bar{\varepsilon}$ to be opmonoidal are automatically satisfied; the less trivial of these is the former, which amounts to the Diagram (5.30), yet that follows using 3 and 6 for a mixed opwreath.

$$TGTI \xrightarrow{T\zeta_I} TTGI \xrightarrow{TT\varepsilon_I} TTTI$$

$$TGGI \qquad \qquad \downarrow^{\mu_{TI}}$$

$$TGGI \qquad \qquad TTI$$

$$\delta_I \uparrow \qquad \qquad \downarrow^{\mu_I}$$

$$GI \xrightarrow{\varepsilon_I} TI$$

$$(5.30)$$

Now we come to the monoidal structure on $\operatorname{mkl}(G,\zeta,T)$. The tensor product of two objects X,X' is the tensor product $X\otimes X'$ of the objects as objects of \mathscr{A} . The tensor product of morphisms $f\colon X\to Y$ and $f'\colon X'\to Y'$ in $\operatorname{mkl}(G,\zeta,T)$, which are morphisms $f\colon GX\to TY$ and $f'\colon GX'\to TY'$ in \mathscr{A} , is the composite

$$G(X \otimes X') \stackrel{\psi_{X,X'}}{\longrightarrow} T(GX \otimes GX') \stackrel{T(f \otimes f')}{\longrightarrow} T(TY \otimes TY')$$

$$\stackrel{T\phi_{Y,Y'}}{\longrightarrow} TT(Y \otimes Y') \stackrel{\mu_{Y \otimes Y'}}{\longrightarrow} T(Y \otimes Y').$$

The unit of this monoidal structure is the unit I of \mathscr{A} . The Eckmann-Hilton argument [10] yields:


Corollary 20. If $(G, \zeta, \delta, \varepsilon)$ is an opmonoidal mixed opwreath around the monoidal monad T on $\mathscr A$ then the monoid $\operatorname{mkl}(G, \zeta, T)(I, I)$ of endomorphisms of I is commutative.


6 Monoidal twisted coactions

In this section we will show what structure on a twisted coaction leads to opmonoidality of the generated mixed opwreath of Proposition 13.

We work in a braided monoidal category \mathcal{V} .

Definition 21. A twisted coaction (γ, τ) of a bimonoid B on a monoid A (see Definition 12) is *monoidal* when it is equipped with a morphism $\mathfrak{d}: B \to A \otimes B \otimes B$ which satisfies the five conditions (6.31) to (6.34).






Proposition 22. Let \mathcal{V} be a braided monoidal category. Given a monoidal twisted coaction (γ, τ) of a bimonoid B on a commutative monoid A (see Definition 12), the mixed opwreath described in Proposition 13, equipped with the morphisms

$$\psi_{X,X'} = (1_A \otimes 1_B \otimes c_{X,B}^{-1} \otimes 1_{X'}) \circ (\mathfrak{d} \otimes 1_X \otimes 1_{X'}) ,$$

is opmonoidal.

Proof. Since A is commutative, the arising monad $T = A \otimes -$ is monoidal with $\phi_0 = \eta \otimes 1_I$ and $\phi_{X,X'} = \mu \circ (1_A \otimes c_{X,A} \otimes 1_{X'})$. Here are the string diagrams for ψ and ϕ .



With these data, and that of Proposition 13, draw the string diagrams for Diagrams (5.24) to (5.29). The remarkable fact is that the variables f and f' can be moved out of the top of Diagrams (5.24), while X and X' can be disconnected from all the diagrams. Diagram (5.29) follows from Diagram (6.33). Then Diagrams (6.31) to (6.34) are what remains. So indeed we obtain an opmonoidal mixed opwreath.

References

- [1] Marcelo Aguiar, Walter Ferrer Santos and Walter Moreira, The Heisenberg product: From Hopf algebras and species to symmetric functions arXiv:1504.06315..
- [2] Jon Beck, Distributive laws, Lecture Notes in Mathematics 80 (Springer, Berlin,1969) 119-140;
 <www.tac.mta.ca/tac/reprints/articles/18/tr18abs.html>.
- [3] Dimitri Chikhladze, Stephen Lack and Ross Street, *Hopf monoidal comonads*, Theory and Applications of Categories **24(19)** (2010) 554–563.
- [4] Brian J. Day, Construction of Biclosed Categories (PhD Thesis, UNSW, 1970) http://www.math.mq.edu.au/~street/DayPhD.pdf>.
- [5] Brian J. Day, On closed categories of functors, Lecture Notes in Mathematics 137 (Springer-Verlag, 1970) 1–38.
- [6] Brian J. Day, An embedding theorem for closed categories, in: "Category Seminar (Proc. Sem., Sydney, 1972/1973)", Lecture Notes in Math. 420 (Springer, Berlin, 1974) 55–64.
- [7] Brian J. Day, Note on monoidal monads, J. Austral. Math. Soc. Ser. A 23(3) (1977) 292–311.
- [8] Brian J. Day, *Promonoidal functor categories*, J. Austral. Math. Soc. Ser. A **23(3)** (1977) 312–328.

- [9] Brian J. Day and Ross Street, Monoidal bicategories and Hopf algebroids, Advances in Math. 129 (1997) 99–157.
- [10] Beno Eckmann and Peter Hilton, Group-like structures in general categories. I. Multiplications and comultiplications, Mathematische Annalen 145(3) (1962) 227–255.
- [11] Samuel Eilenberg and G. Max Kelly, *Closed categories*, Proceedings of the Conference on Categorical Algebra (La Jolla, 1965), (Springer-Verlag,1966) 421–562.
- [12] Richard Garner and Ross Street, Coalgebras governing both weighted Hurwitz products and their pointwise transforms, arXiv:1510.05323.
- [13] R. Harmer, M. Hyland and P.-A. Mellies, Categorical combinatorics for innocent strategies, Logic in Computer Science, Proc. 22nd Annual IEEE Symposium on Logic in Computer Science (LICS, IEEE, 2007) 379–388.
- [14] Daniela Hobst and Bodo Pareigis, *Double quantum groups*, J. Algebra 242 (2001) 460–494.
- [15] André Joyal, Une theory combinatoire des series formelles, Advances in Mathematics 42 (1981) 1–82.
- [16] André Joyal, Foncteurs analytiques et especes de structures, Lecture Notes in Mathematics **1234** (Springer 1986) 126–159.
- [17] André Joyal and Ross Street, *The geometry of tensor calculus I*, Advances in Math. **88** (1991) 55–112.
- [18] André Joyal and Ross Street, Tortile Yang-Baxter operators in tensor categories, Journal of Pure and Applied Algebra 71 (1991) 43–51.
- [19] André Joyal and Ross Street, *Braided tensor categories*, Advances in Mathematics **102** (1993) 20–78.
- [20] G. Max Kelly, Basic concepts of enriched category theory, London Mathematical Society Lecture Note Series 64 (Cambridge University Press, Cambridge, 1982).
- [21] G. Max Kelly and Ross Street, *Review of the elements of 2-categories*, Lecture Notes in Mathematics **420** (Springer-Verlag, 1974) 75–103.
- [22] Anders Kock, Monads for which structures are adjoint to units, Journal of Pure and Applied Algebra 104 (1995) 41–59.
- [23] Stephen Lack and Ross Street, *The formal theory of monads II*, J. Pure Appl. Algebra **175(1-3)** (2002) 243–265.

- [24] F. William Lawvere, Ordinal sums and equational doctrines, in: Sem. on Triples and Categorical Homology Theory, ETH, Zürich, 1966/1967, (Springer, Berlin, 1969) 141–155.
- [25] Saunders Mac Lane, Categories for the Working Mathematician, Graduate Texts in Mathematics 5 (Springer-Verlag, 1971).
- [26] Walter Moreira, Products of representations of the symmetric group and non-commutative versions (PhD Thesis, Texas A&M University, 2008).
- [27] Paddy McCrudden, *Opmonoidal monads*, Theory and Applications of Categories **10** (2002) 469–485.
- [28] Ieke Moerdijk, *Monads on tensor categories*, Journal of Pure Appl. Algebra **168** (2002) 189–208.
- [29] A. John Power and Hiroshi Watanabe, Combining a monad and a comonad, Theoretical Computer Science 280 (2002) 137–162.
- [30] Ross Street, The formal theory of monads, J. Pure Appl. Algebra 2 (1972) 149–168.
- [31] Ross Street, *Fibrations in bicategories*, Cahiers de topologie et géométrie différentielle **21** (1980) 111–160.
- [32] Ross Street, Categorical structures, Handbook of Algebra Volume 1 (editor M. Hazewinkel; Elsevier Science, Amsterdam 1996; ISBN 0 444 82212 7) 529–577.
- [33] Ross Street, Fusion operators and cocycloids in monoidal categories, Applied Categorical Structures **6(2)** (1998) 177–191.
- [34] Ross Street, Quantum Groups: a path to current algebra, Australian Math. Society Lecture Series 19 (Cambridge University Press; 18 January 2007; ISBN-978-0-521-69524-4).
- [35] Ross Street, Weak distributive laws, Theory and Applications of Categories **22(12)** (2009) 313–320.
- [36] Ross Street, Monoidal categories in, and linking, geometry and algebra, Bulletin of the Belgian Mathematical Society Simon Stevin **19(5)** (2012) 769–821.
- [37] Ross Street, Weighted tensor products of Joyal species, graphs, and charades, SIGMA (to appear 2016); arXiv:1503.02783.