MIXING ACTIONS OF 0-ENTROPY FOR COUNTABLE AMENABLE GROUPS

ALEXANDRE I. DANILENKO

ABSTRACT. It is shown that each discrete countable infinite amenable group admits a 0-entropy mixing action on a standard probability space.

0. Introduction

Let G be an amenable discrete infinite countable group. We recall that a measure preserving action $T = (T_g)_{g \in G}$ of G on a standard probability space (X, \mathfrak{B}, μ) is called mixing if $\mu(T_gA \cap B) \to \mu(A)\mu(B)$ as $g \to \infty$ for all measurable subsets $A, B \subset X$. The Kolmogoroff-Sinai entropy for measure-preserving transformations (i.e. \mathbb{Z} -actions) was extended to the actions of locally compact amenable groups in [OrWe]. Since 0-entropy actions are considered as "deterministic" while mixing actions are considered as "chaotic", it is natural to ask are there actions which possess the two properties simultaneously? Such examples for \mathbb{Z} -actions were found first in the class of Gaussian transformations (see [New]) and later in the class of rank-one transformations (see [Or], [Ad], [CrSi] and references therein). The rank-one analogues of the later family were constructed for $G = \mathbb{R}^{d_1} \times \mathbb{Z}^{d_2}$ in [DaSi], for G being a countable direct sum of finite groups in [Da2] and, more generally, for G being a locally normal countable group in [Da3]. Dan Rudolph asked¹ whether each amenable countable group G has a mixing action of zero entropy? The purpose of this work is to answer this question affirmatively.

Theorem 0.1. There is a 0-entropy mixing free² probability preserving action of G.

Since the proof of the theorem is based on the Poisson suspensions of infinite measure preserving actions we introduce the necessary definitions in the next section (see [Ne], [Roy], [Ja–Ru] for more detail). We note that the mixing property follows essentially from the construction elaborated in [Da4]. Thus we have only to show here how to modify that construction to achieve the freeness³ and 0-entropy of the action in question.

¹At an AMS meeting in early 90's.

²We stick to the free actions to avoid degeneracy and triviality like the following: if $(T_g)_{g \in G}$ is a 0-entropy mixing action of G, F is a finite group and $\widetilde{T}_{g,f} := T_g$ for $(g,f) \in G \times F$ then $(\widetilde{T}_{g,f})_{(g,f) \in G \times F}$ is a 0-entropy mixing action of $G \times F$. We consider this non-free action as degenerated and non-interesting.

³The freeness of the $H_3(\mathbb{R})$ -actions considered in [Da4] follows from some properties of group actions that are specific to actions of connected nilpotent Lie groups.

1. Poisson suspensions

Let X be a locally compact noncompact Cantor space (i.e. 0-dimensional, without isolated points). Denote by $C_{00}(X)$ the vector space of real valued functions on X with compact support. This space is endowed with the usual locally convex topology, i.e. the topology of uniform convergence on the compact subsets of X. The dual $C_{00}(X)'$ is called the space of (real) Radon measures on X. We are interested in the cone $X^* \subset C_{00}(X)'$ of nonnegative Radon measures on X. Furnish X^* with the Borel σ -algebra \mathfrak{B}^* generated by the *-weak topology related to the duality $\langle C_{00}(X), C_{00}(X)' \rangle$. We note that \mathfrak{B}^* is also the Borel σ -algebra generated by the strong topology related to this duality. Since the strong topology is Polish and X^* is closed in $C_{00}(X)'$, it follows that (X^*, \mathfrak{B}^*) is a standard Borel space.

Denote by K the set of all compact open subsets of X. Of course, K is infinite but countable. We also note that \mathfrak{B}^* is the smallest σ -algebra on X^* such that for each $K \in K$, the mapping $N_K : X^* \ni x^* \mapsto x^*(K) \in \mathbb{R}_+$ is measurable. Let μ^* be the only measure on (X^*, \mathfrak{B}^*) such that

- $\mu^* \circ N_K^{-1}$ is the Poisson distribution with parameter $\mu(K)$ and
- the random variables N_{K_1}, \ldots, N_{K_n} on (X^*, μ^*) are independent for each countable collection of mutually disjoint subsets $K_1, \ldots, K_m \in \mathcal{K}$.

Then $(X^*, \mathfrak{B}^*, \mu^*)$ is a standard probability space. To define μ^* rigorously we denote by \mathfrak{F} the set of all finite collections of mutually disjoint nonempty compact open subsets of X. For $\mathcal{F}_1, \mathcal{F}_2 \in \mathfrak{F}$, we write $\mathcal{F}_2 \succ \mathcal{F}_1$ if each element of \mathcal{F}_1 is a union of some elements from \mathcal{F}_2 . Then (\mathfrak{F}, \succ) is a directed partially ordered set. Given $K \in \mathcal{K}$, we define a probability measure ν_K on \mathbb{R}_+ by setting

$$\nu_K(A) := \sum_{i \in A \cap \mathbb{Z}_+} \frac{e^{-\mu(K)}\mu(K)^i}{i!}$$

for each Borel subset $A \subset \mathbb{R}_+$. We see, in particular, that ν_K is supported on \mathbb{Z}_+ . Since \mathbb{R}_+ is an additive semigroup, the convolution of probability measures on \mathbb{R}^* is well defined. It is easy to verify that if $K_1, K_2 \in \mathcal{K}$ and $K_1 \cap K_2 = \emptyset$ then $\nu_{K_1} * \nu_{K_2} = \nu_{K_1 \sqcup K_2}$. Given $\mathcal{F} \in \mathfrak{F}$, we denote by $\mathbb{R}_+^{\mathcal{F}}$ the set of all mappings $x : \mathcal{F} \ni K \mapsto x(K) \in \mathbb{R}_+$. We define a measure $\nu^{\mathcal{F}}$ on $\mathbb{R}_+^{\mathcal{F}}$ as the direct product $\nu^{\mathcal{F}} := \bigotimes_{K \in \mathcal{F}} \nu_K$. If $\mathcal{F}_1, \mathcal{F}_2 \in \mathfrak{F}$ and $\mathcal{F}_2 \succ \mathcal{F}_1$ we define a mapping $\pi_{\mathcal{F}_1}^{\mathcal{F}_2} : \mathbb{R}_+^{\mathcal{F}_2} \to \mathbb{R}_+^{\mathcal{F}_1}$ by setting

$$\left(\pi_{\mathcal{F}_1}^{\mathcal{F}_2}(x)\right)(K) = \left(\sum_{\{R \in \mathcal{F}_2 \mid R \subset K\}} x(R)\right) \quad \text{for each } K \in \mathcal{F}_1 \text{ and } x \in \mathbb{R}_+^{\mathcal{F}_2}.$$

The aforementioned convolution property of ν_K implies that $\nu^{\mathcal{F}_2} \circ (\pi_{\mathcal{F}_1}^{\mathcal{F}_2})^{-1} = \nu^{\mathcal{F}_1}$. Thus $\{(\mathbb{R}^{\mathcal{F}_1}, \nu^{\mathcal{F}_2}, \pi_{\mathcal{F}_1}^{\mathcal{F}_2})_{\mathcal{F}_2 \succ \mathcal{F}_1} \mid \mathcal{F}_1, \mathcal{F}_2 \in \mathfrak{F}\}$ is a projective system of probability spaces. Hence the projective limit $(Y, \kappa) := \operatorname{proj lim}_{(\mathfrak{F}, \succ)}(\mathbb{R}_+^{\mathcal{F}}, \nu^{\mathcal{F}})$ is well defined as a standard probability space. We now define a Borel map $\Phi : X^* \to Y$ by setting

$$\Phi(x^*) := (\Phi(x^*)_{\mathcal{F}})_{\mathcal{F} \in \mathfrak{F}}, \text{ where } (\Phi(x^*)_{\mathcal{F}} := (N_K(x^*))_{K \in \mathcal{F}}.$$

Then Φ is one-to-one and onto. Indeed, if two nonnegative Radon measures take the same values on every element of \mathcal{K} then these measures are equal. This proves

that Φ is one-to-one. On the other hand, each element y of Y can be interpreted as a finitely additive non-negative measure on \mathcal{K} , i.e. as a map $y:\mathcal{K}\to\mathbb{R}_+$ such that $y(K_1 \sqcup \cdots \sqcup K_s) = y(K_1) + \cdots + y(K_s)$ for every collection $\{K_1, \ldots, K_s\} \in \mathfrak{F}$. Then y extends uniquely to a σ -finite (non-negative) measure on X. Of course, the extension is a Radon measure. Hence Ψ is onto. Thus Ψ is a Borel isomorphism of X^* onto Y. It remains to export κ to X^* via Φ^{-1} and denote this measure by μ^* . It follows, in particular, that for each $K \in \mathcal{K}$ and $j \in \mathbb{Z}_+$,

(1-1)
$$\mu^*(\lbrace x^* \in X^* \mid x^*(K) = j \rbrace) = \frac{\mu(K)^j e^{-\mu(K)}}{i!}.$$

If X is partitioned into a union of mutually disjoint open non-compact subsets X_1, \ldots, X_m then the mapping

$$x^* \mapsto (x^* \upharpoonright X_1, \dots, x^* \upharpoonright X_m)$$

is a measure preserving Borel isomorphism of (X^*, μ^*) onto the direct product space $(X_1^* \times \cdots \times X_m^*, (\mu \upharpoonright X_1)^* \times \cdots \times (\mu \upharpoonright X_m)^*).$

Given a Borel σ -algebra $\mathfrak{F} \subset \mathfrak{B}$ generated by a topology which is weaker than the original topology on X, we denote by \mathfrak{F}^* the smallest sub- σ -algebra of \mathfrak{B}^* such that the mapping N_K is measurable for each $K \in \mathfrak{F} \cap \mathcal{K}$. For an increasing sequence of topologies $\tau_1 \prec \tau_2 \prec \cdots$ on X which are weaker than the original topology on X, we denote by $\mathfrak{F}_1 \subset \mathfrak{F}_2 \subset \cdots$ the corresponding increasing sequence of Borel sub- σ -algebras generated by these topologies. Since every compact open subset from $\bigvee_{n=1}^{\infty} \mathfrak{F}_n$ is contained in \mathfrak{F}_m for some m>0, it follows that

(1-2)
$$\left(\bigvee_{n=1}^{\infty} \mathfrak{F}_n\right)^* = \bigvee_{n=1}^{\infty} \mathfrak{F}_n^*.$$

Given an action $T = (T_g)_{g \in G}$ of G on X by μ -preserving homeomorphisms T_g , we associate a Borel action $T^* := (T_g^*)_{g \in G}$ on X^* by setting $T_g^* x^* := x^* \circ T_g^{-1}, g \in G$. It follows from (1-1) that T^* preserves μ^* . The dynamical system $(X^*, \mathfrak{B}^*, \mu^*, T^*)$ is called the *Poisson suspension* of $(X, \mathfrak{B}, \mu, T)$.

2. Proof of Theorem 0.1

We will proceed in several steps.

Step 1. First we construct a free strictly ergodic infinite measure preserving Gaction on a locally compact non-compact Cantor space. For that we will utilize the (C, F)-construction (see [Da1], [Da5]). Let $(F_n, C_{n+1})_{n=0}^{\infty}$ be a sequence of finite subsets in G such that $(F_n)_{n=0}^{\infty}$ is a Følner sequence in $G, F_0 = \{1\}$ and for each $n \ge 1$, the following three basic conditions are satisfied:

- $-1 \in F_n \text{ and } \#C_n > 1,$
- $-F_n^{-1}F_nF_nC_{n+1} \subsetneq F_{n+1},$ $-F_nc \cap F_nc' = \emptyset \text{ for all } c \neq c' \in C_{n+1}.$

We let $X_n := F_n \times C_{n+1} \times C_{n+2} \times \cdots$. Then X_n endowed with the infinite product of the discrete topologies on F_n and C_j , j > n, is a compact Cantor set. Moreover, the mapping

$$X_n \ni (f_n, c_{n+1}, c_{n+2}, \dots) \mapsto (f_n c_{n+1}, c_{n+2}, \dots) \in X_{n+1}$$

is a topological embedding of X_n into X_{n+1} . We now consider the union $X = \bigcup_{n \geq 1} X_n$ and endow it with the topology of inductive limit. Then X is a locally compact non-compact Cantor set and X_n is a compact open subset of X for each n. Given $g \in G$ and n > 0, we define a homeomorphism $T_g^{(n)}$ from a clopen subset $(g^{-1}F_n \cap F_n) \times C_{n+1} \times C_{n+2} \times \cdots$ of X_n to a clopen subset $(F_n \cap gF_n) \times C_{n+1} \times C_{n+2} \times \cdots$ of X_n by setting

$$T_g^{(n)}(f_n,c_{n+1},\ldots)\mapsto (gf_n,c_{n+1},\ldots).$$

It is easy to verify that the sequence $(T_g^{(n)})_{n\geq 1}$ determines uniquely a homeomorphism T_g of X such that $T_g \upharpoonright X_n = T_g^{(n)}$ for each n. It is straightforward to check that $T := (T_g)_{g \in G}$ is a free topological action of G on X. This action is minimal and uniquely ergodic, i.e. there is a unique T-invariant σ -finite Radon measure μ on X such that $\mu(X_0) = 1$. To define μ explicitly we consider for each $n \geq 0$ and $f \in F_n$, a subset $[f]_n := \{(f, c_{n+1}, c_{n+2}, \dots) \in X_n \mid c_i \in C_i \text{ for all } i > n\}$. Then $[f]_n$ is compact and open. We call it a *cylinder*. The family of all cylinders is a base for the topology on X. Every compact open subset of X is a union of finitely many mutually disjoint cylinders. Hence every Radon measure on X is determined uniquely by its values on the cylinders. It remains to note that $\mu([f]_n) = \frac{1}{\#C_1 \cdots \#C_n}$ for each $n \geq 0$ and $f \in F_n$.

In addition to the aforementioned three basic conditions on $(F_n, C_{n+1})_{n\geq 0}$, we will assume that

$$\lim_{n \to \infty} \frac{\#F_n}{\#C_1 \cdots \#C_n} = \infty.$$

This condition is equivalent to the fact that $\mu(X) = \infty$.

Step 2. On this step we obtain a free probability preserving G-action. For that we need two more conditions on $(F_n, C_{n+1})_{n=0}^{\infty}$:

- (\triangle) for each element g of infinite order in G, there are infinitely many n such that $g^{l_n}F_nC_{n+1} \subset F_{n+1} \setminus (F_nC_{n+1})$ for some $l_n > 0$.
- (\square) for each element g of finite order in G, there are infinitely many n such that $gF_n = F_n$.

It is easy to see that if (\triangle) is satisfied then

$$T_{g^{l_n}}X_n = T_{g^{l_n}} \bigsqcup_{f \in F_n} [f]_n = T_{g^{l_n}} \bigsqcup_{f \in F_n} \bigsqcup_{c \in C_{n+1}} [fc]_{n+1} = \bigsqcup_{f \in F_n} \bigsqcup_{c \in C_{n+1}} [g^{l_n}fc]_{n+1}.$$

We used the fact that $X_n = \bigsqcup_{f \in F_n} [f]_n$ and $T_s[f]_n = [sf]_n$ whenever $f, sf \in F_n$ and $s \in G$. Hence $T_{g^{l_n}} X_n \subset X_{n+1}$ and $T_{g^{l_n}} X_n \cap X_n = \emptyset$.

Let (X^*, μ^*, T^*) be the Poisson suspension of (X, μ, T) . Then μ^* is T^* -invariant and $\mu^*(X^*) = 1$. We now verify that T^* is free. For that we will check that for each $g \in G \setminus \{1\}$, the subset of fixed points of the transformation T_g^* is μ^* -null. Consider two cases. If g is of infinite order and A is a T_g -invariant subset of positive finite measure in X then in view of (Δ) there is n > 0 such that $\mu(A \cap X_n) > 0.9\mu(A)$ and $T_{g^{l_n}}X_n \cap X_n = \emptyset$. Hence

$$\mu(A\cap X_n) = \mu(T_{g^{l_n}}A\cap T_{g^{l_n}}X_n) = \mu(A\cap T_{g^{l_n}}X_n) \le \mu(A\setminus X_n) \le 0.1\mu(A),$$

a contradiction. Thus the transformation T_g has no invariant subsets of finite positive measure. Therefore the Poisson suspension T_g^* of T_g is weakly mixing

[Roy]. This yields that $\mu^*(\{x^* \in X^* \mid T_g^*x^* = x^*\}) = 0$. Consider now the case where g is of finite order. Let $H \subset G$ denote the cyclic subgroup generated by g. It follows from (\Box) that there is an open subset $Y \subset X$ of infinite measure such that the sets T_hY , $h \in H$, form an open partition of X. Indeed let $gF_{n_i} = F_{n_i}$ for an increasing sequence $n_1 < n_2 < \cdots$. Choose a subset $S_1 \subset F_{n_1}$ which meets each H-coset in F_{n_1} exactly once. If i > 1 choose a subset $S_i \subset F_{n_i}$ which meets each H-coset in $F_{n_i} \setminus (F_{n_{i-1}}C_{n_{i-1}+1}\cdots C_{n_i})$ exactly once. It remains to set $Y := \bigsqcup_{i=1}^{\infty} \bigsqcup_{s \in S_i} [s]_{n_i}$. Of course, Y is open and non-compact and $X = \bigsqcup_{h \in H} T_h Y$. Since $\infty = \mu(X) = \mu(\bigsqcup_{h \in H} T_h Y)$ and T preserves μ , it follows that $\mu(Y) = \infty$. Then the dynamical system $(X^*, \mu^*, (T_h^*)_{h \in H})$ is isomorphic to the finite direct product $(Y^*, (\mu \upharpoonright Y)^*)^H$ endowed with the natural shiftwise action of H (see §1). Since the measure $(\mu \upharpoonright Y)^*$ is nonatomic, this action is free.

Step 3. We now verify that $h(T^*) = 0$. Let \mathfrak{F}_n denote the σ -algebra on X generated by a single set $[1]_n$ and let \mathfrak{B}_n denote the σ -algebra on X generated by the compact open sets $[f]_n$, $f \in F_n$. Then $\mathfrak{B}_1 \subset \mathfrak{B}_2 \subset \cdots$ and $\bigvee_{n=1}^{\infty} \mathfrak{B}_n$ is the entire Borel σ -algebra \mathfrak{B} on X. Moreover, $\mathfrak{B}_n = \bigvee_{g \in F_n} T_g \mathfrak{F}_n$ and

$$\mathfrak{B}_n^* = \bigvee_{g \in F_n} T_g^* \mathfrak{F}_n^* \subset \bigvee_{g \in G} T_g^* \mathfrak{F}_n^* =: (\mathfrak{F}_n^*)^G$$

for each n. Therefore $\bigvee_{n=1}^{\infty} (\mathfrak{F}_n^*)^G \supset \bigvee_{n=1}^{\infty} \mathfrak{B}_n^* = \mathfrak{B}^*$. The latter equality follows from (1-2). Moreover, it is easy to verify that $(\mathfrak{F}_1^*)^G \subset (\mathfrak{F}_2^*)^G \subset \cdots$. Hence

$$h(T^*) = \lim_{n \to \infty} h\left(T^* \middle| (\mathfrak{F}_n^*)^G\right) \le \limsup_{n \to \infty} H(\mathfrak{F}_n^*).$$

We recall that \mathcal{K} denote the collection of all compact open subsets of X. Since $\mathfrak{F}_n \cap \mathcal{K} = \{[1]_n\}$, the σ -algebra \mathfrak{F}_n^* is generated by a countable partition of X^* into the sets $N_{[1]_n}^{-1}(r) = \{x^* \in X^* \mid x^*([1]_n) = r\}, r = 0, 1, \ldots$ This yields that

$$\limsup_{n \to \infty} H(\mathfrak{F}_n^*) = \lim_{n \to \infty} f(\mu([1]_n)),$$

where f(t) is the entropy of the Poisson distribution $(e^{-t}, e^{-t}t, e^{-t}t^2/2, ...)$. Since $\mu([1]_n) = 1/(\#C_1 \cdots \#C_n) \to 0$, it follows that $h(T^*) = 0$.

Step 4. On this step we show that some extra conditions on $(F_n, C_{n+1})_{n\geq 0}$ imply mixing for the dynamical system (X^*, μ^*, T^*) . Thus from now on we will assume that the following hold for each n (in addition to the conditions on $(F_n, C_{n+1})_{n\geq 0}$ listed above):

- (i) $F_n F_n^{-1} F_n C_{n+1} \subset F_{n+1}$,
- (ii) the sets $F_n c_1 c_2^{-1} F_n^{-1}$, $c_1 \neq c_2 \in C_{n+1}$, and $F_n F_n^{-1}$ are all pairwise disjoint and
- (iii) $\#C_n \to \infty$ as $n \to \infty$.

Denote by $U_T = (U_T(g))_{g \in G}$ the associated Koopman unitary representation of G in $L^2(X,\mu)$, i.e. $U_T(g)f := f \circ T_g^{-1}$ for each $f \in L^2(X,\mu)$. As was shown in [Da4, Theorem 5.1]⁴, the conditions (i)–(iii) imply that T is mixing as an infinite

⁴Though we considered in [Da4] mainly the actions of the Heisenberg group $H_3(\mathbb{Z})$, the proof of Theorem 5.1 there does not use any specific property of $H_3(\mathbb{Z})$. It holds for each amenable group.

measure preserving action, i.e. $U_T(g) \to 0$ weakly as $g \to \infty$. Let U_{T^*} stand for the Koopman unitary representation of G in $L^2(X^*, \mu^*)$ associated with the Poisson suspension (X^*, μ^*, T^*) . It is well known that U_{T^*} is unitarily equivalent to the Fock unitary representation of G generated by U_T in the Fock space generated by $L^2(X, \mu)$ (see [Ne]). It follows that $U_{T^*}(g)$ converges weakly to the orthogonal projection to the constants in $L^2(X^*, \mu^*)$ as $g \to \infty$. This is equivalent to the fact that T^* is mixing.

Summarizing, we obtain that T^* is mixing free action of G on (X^*, μ^*) and $h(T^*) = 0$. Thus Theorem 0.1 is proved.

References

- [Ad] T. Adams, Smorodinsky's conjecture on rank one systems, Proc. Amer. Math. Soc. 126 (1998), 739–744.
- [CrS] D. Creutz and C. E. Silva, Mixing on a class of rank-one transformations, Ergod. Th.
 & Dynam. Sys. 24 (2004), 407–440.
- [Da1] _____, (C, F)-actions in ergodic theory, Geometry and Dynamics of Groups and Spaces, Progress in Mathematics, vol. 265, Birkhäuser Verlag, Basel/Switzerland, 2008, pp. 325–351.
- [Da2] _____, Mixing rank-one actions for infinite sums of finite groups, Isr. J. Math. **156** (2006), 341–358.
- [Da3] _____, Uncountable collection of mixing rank-one actions for locally normal groups, Semin. et Congr. de la SMF **20** (2011), 253–266.
- [Da4] _____, Mixing actions of the Heisenberg group, Erg. Th. & Dyn. Syst. **34** (2014), 1142–1167.
- [Da5] _____, Actions of finite rank: weak rational ergodicity and partial rigidity, Ergod. Th. Dyn. Syst. (to appear).
- [DaSi] A. I. Danilenko and C. E. Silva, Mixing rank-one actions of locally compact Abelian groups, Ann. Inst. H. Poincaré, Probab. Statist. 43 (2007), 375–398.
- [Ja-Ru] E. Janvresse, T. Meyerovitch, E. Roy and T. de la Rue, *Poisson suspensions and entropy for infinite transformations*, Trans. Amer. Math. Soc. **362** (2010), 3069–3094.
- [Ne] Yu. Neretin, Categories of symmetries and infinite-dimensional groups, London Mathematical Society Monographs. New Series, 16. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1996.
- [New] D. Newton, On Gaussian processes with simple spectrum, Z. Wahrsch. Verw. Gebiete 5 (1966), 207–209.
- [Or] D. S. Ornstein, On the root problem in ergodic theory, Proc. Sixth Berkley Symp. Math. Stat. Prob. (Univ. California, Berkeley, Calif., 1970/1971), vol II, Univ. of California Press, Berkeley, Calif., 1972, pp. 347–356.
- [Or] D. S. Ornstein, On the root problem in ergodic theory, Proc. Sixth Berkley Symp. Math. Stat. Prob. (Univ. California, Berkeley, Calif., 1970/1971), vol. II, Univ. of California Press, Berkeley, Calif., 1972, pp. 347–356.
- [OrWe] D. S. Ornstein and B. Weiss, Entropy and isomorphism theorems for actions of amenable groups, J. Analyse Math. 48 (1987), 1–141.
- [Roy] E. Roy, *Poisson suspensions and infinite ergodic theory*, Ergod. Th. & Dynam. Systems (2009), 667–683.

Institute for Low Temperature Physics & Engineering of National Academy of Sciences of Ukraine, 47 Lenin Ave., Kharkov, 61164, UKRAINE

 $E ext{-}mail\ address: alexandre.danilenko@gmail.com}$