

# Weighted Hurwitz numbers and topological recursion: an overview

A. Alexandrov<sup>1,2,3\*</sup>, G. Chapuy<sup>4†</sup>, B. Eynard<sup>2,5‡</sup> and J. Harnad<sup>2,6§</sup>

<sup>1</sup>*Center for Geometry and Physics, Institute for Basic Science (IBS), Pohang 37673, Korea*

<sup>2</sup>*Centre de recherches mathématiques, Université de Montréal  
C. P. 6128, succ. centre ville, Montréal, QC H3C 3J7, Canada*

<sup>3</sup>*ITEP, Bolshaya Cheremushkinskaya 25, 117218 Moscow, Russia*

<sup>4</sup>*CNRS UMR 7089, Université Paris Diderot  
Paris 7, Case 7014, 75205 Paris Cedex 13, France*

<sup>5</sup>*Institut de Physique Théorique, CEA, IPhT  
CNRS URA 2306, F-91191 Gif-sur-Yvette, France*

<sup>6</sup>*Department of Mathematics and Statistics, Concordia University  
1455 de Maisonneuve Blvd. W. Montreal, QC H3G 1M8, Canada*

## Abstract

Multiparametric families of hypergeometric  $\tau$ -functions of KP or Toda type serve as generating functions for weighted Hurwitz numbers, providing weighted enumerations of branched covers of the Riemann sphere. A graphical interpretation of the weighting is given in terms of constellations mapped onto the covering surface. The theory is placed within the framework of topological recursion, with the Baker function at  $\mathbf{t} = \mathbf{0}$  shown to satisfy the quantum spectral curve equation, whose classical limit is rational. A basis for the space of formal power series in the spectral variable is generated that is adapted to the Grassmannian element associated to the  $\tau$ -function. Multicurrent correlators are defined in terms of the  $\tau$ -function and shown to provide an alternative generating function for weighted Hurwitz numbers. Fermionic VEV representations are provided for the adapted bases, pair correlators and multicurrent correlators. Choosing the weight generating function as a polynomial, and restricting the number of nonzero “second” KP flow parameters in the Toda  $\tau$ -function to be finite implies a finite rank covariant derivative equation with rational coefficients satisfied by a finite “window” of adapted basis elements. The pair correlator is shown to provide a Christoffel-Darboux type finite rank integrable kernel, and the WKB series coefficients of the associated adjoint system are computed recursively, leading to topological recursion relations for the generators of the weighted Hurwitz numbers.

---

\*e-mail: alexandrovsash@gmail.com

†e-mail: guillaume.chapuy@liafa.univ-paris-diderot.fr

‡e-mail: bertrand.eynard@cea.fr

§e-mail: harnad@crm.umontreal.ca

# Contents

<b>1</b>	<b>Weighted Hurwitz numbers and generating functions</b>	<b>3</b>
1.1	Geometric and combinatorial definitions of Hurwitz numbers . . . . .	4
1.2	Weighted Hurwitz numbers . . . . .	5
1.2.1	Weight generating function $G(z)$ . . . . .	5
1.2.2	Geometrical definition: weighted coverings . . . . .	5
1.2.3	Combinatorial definition: weighted constellations . . . . .	6
<b>2</b>	<b>Hypergeometric <math>\tau</math>-function, Baker function, recursions and correlators</b>	<b>9</b>
2.1	Hypergeometric $\tau$ -functions as generating functions for weighted Hurwitz numbers . . . . .	9
2.2	Baker function, adapted bases and recursions . . . . .	11
2.3	The pair correlator and its expansion . . . . .	13
2.4	The multicurrent correlator as generating function . . . . .	14
2.5	The Christoffel-Darboux relation . . . . .	16
<b>3</b>	<b>The spectral curve</b>	<b>17</b>
3.1	Quantum spectral curve . . . . .	17
3.2	Classical spectral curve . . . . .	18
<b>4</b>	<b>Linear first order system for the adapted basis</b>	<b>19</b>
4.1	The constant coefficient infinite system . . . . .	19
4.2	Finite dimensional first order system: folding . . . . .	20
4.3	Adjoint differential system . . . . .	21
4.4	$\beta$ -expansion of $M$ . . . . .	21
4.5	The current correlators $\tilde{W}_n$ . . . . .	22
<b>5</b>	<b>Fermionic representations</b>	<b>22</b>
5.1	The $\tau$ -function, Baker function and adapted bases . . . . .	23
5.2	The $n$ -pair correlator $K_n$ . . . . .	24
5.3	The multicurrent correlator $W_n$ . . . . .	24
<b>6</b>	<b>Topological recursion</b>	<b>24</b>
6.1	Definitions of $\tilde{\omega}_{g,n}$ and $\mathcal{W}_{g,n}$ . . . . .	24
6.2	Topological recursion for the $\tilde{\omega}_{g,n}$ 's . . . . .	25
<b>7</b>	<b>Some consequences of topological recursion</b>	<b>25</b>

# 1 Weighted Hurwitz numbers and generating functions

The study of weighted enumerations of branched coverings of the Riemann sphere was initiated by Hurwitz [31]. It was subsequently related to the character theory of the symmetric group by Frobenius [24], and computations of these enumerative invariants were obtained for low genus and simple branching structures. Interest was more recently revived by the work of Okounkov and Pandharipande [41,45], who showed that certain special forms of the Sato-Segal-Wilson [46,47,49] KP  $\tau$ -function and 2D-Toda  $\tau$ -functions, central to the modern theory of integrable systems, serve as generating functions for “simple” Hurwitz numbers, which enumerate branched covers in which all but one or two of the specified branching profiles are 2-cycles.

Further special cases, relating to the enumeration of monotonic paths in the Cayley graph of the symmetric group [25,26], three point branched covers [7–9,34,53], coverings of  $\mathbf{RP}^2$  [40] and certain cases of matrix integrals [4,8,14,37] were subsequently considered, and their generating functions similarly shown to be special cases of  $\tau$ -functions of hypergeometric type [42–44]. The most general case of weighted Hurwitz numbers, with weight generating functions depending on an infinite number of weighting parameters, was developed in [26] - [30].

The enumeration of branched coverings is classically equivalent to the enumeration of *constellations*, which are special types of bipartite graphs mapped onto Riemann surfaces and “decorated” in several possible ways [35]. The enumerative study of maps with respect to their genus invariant has been considerably developed by combinatorists since the pioneering work of Tutte in the planar case [51] and Bender and Canfield [11] for higher genus. (The reader is referred to [48] for an accessible introduction, historical references, and a modern entry to the rapidly growing literature on this topic.)

In recent years, the topological recursion program [20–22] has been a significant development in the study of enumerative, combinatorial, geometric and topological invariants associated to maps, Riemann surfaces, moduli spaces and knots. It has led to an explicit finite recursive algebraic scheme for computing such invariants. This begins with the construction of an associated “quantum curve” which, in general, is an ordinary differential or difference equation in the spectral parameter satisfied by the associated Baker function, whose WKB formal series solution, expanded in a suitable “small parameter” is studied recursively. Partial results on the inclusion of Hurwitz numbers in this scheme have been obtained in [5,14,18,19,34,38,39], mainly for the special cases mentioned above. In the present work, it is shown how a broad class of multiparametrically weighted Hurwitz numbers can be systematically placed into the topological recursion program, and the associated spectral curve, both quantum and classical, determined explicitly in terms of the weight generating function.

The aim of this work is to provide an accessible introduction that summarizes the main

results in this direction. All relevant constructions are indicated, and the principal results stated. Some proofs are provided in detail; others are only sketched in outline, with full details provided in one of two companion publications [2, 3]. The results include: explicit parametrization of both the classical and quantum spectral curves; the system of recursion relations satisfied by dual pairs of adapted bases for the underlying space of formal power series in a spectral parameter; construction of suitably defined multipair correlators and multicurrent correlators, which serve as generating functions for weighted Hurwitz numbers; expressions for all relevant quantities as fermionic vacuum state expectation values; a finite rank expression for the Fredholm integral kernel associated to the pair correlator, analogous to the Christoffel-Darboux kernel appearing in the theory of orthogonal polynomials and, finally, the topological recursion relations satisfied by the multiform weighted Hurwitz number generators.

## 1.1 Geometric and combinatorial definitions of Hurwitz numbers

Given a set of  $k$  partitions of  $N \in \mathbf{N}^+$ ,  $\{\mu^{(i)}\}_{i=1,\dots,k}$ ,  $|\mu^{(i)}| = N$ , we define the *pure Hurwitz number*  $H(\mu^{(1)}, \dots, \mu^{(k)})$  geometrically as the number of inequivalent  $N$ -sheeted branched coverings of the Riemann sphere with  $k$  branch points having ramification profiles  $\{\mu^{(i)}\}_{i=1,\dots,k}$ , normalized by the inverse of the order of the automorphism group of the cover. The Riemann-Hurwitz theorem expresses the Euler characteristic of the covering surface as

$$\chi = 2 - 2g = 2N - \sum_{i=1}^k \ell^*(\mu^{(i)}), \quad (1.1)$$

where  $\ell^*(\mu)$  denotes the *colength* of the partition  $\mu$  (i.e. the complement of its length)

$$\ell^*(\mu) := |\mu| - \ell(\mu). \quad (1.2)$$

Combinatorially, it may equivalently be defined as the number of distinct ways the identity element  $\mathbf{I} \in S_N$  in the symmetric group  $S_N$  may be factorized

$$\mathbf{I} = h_1 \cdots h_k \quad (1.3)$$

as a product of  $k$  elements  $h_i \in \text{cyc}(\mu^{(i)})$  in the conjugacy classes of cycle type  $\mu^{(i)}$ ; i.e., whose cycle lengths are the parts of the partition  $\mu^{(i)}$ , multiplied by the normalization factor  $1/N!$ . The Frobenius-Schur formula [35][Appendix A] expresses this in terms of the irreducible group characters  $\chi_\lambda(\mu)$  of  $S_N$ .

$$H(\mu^{(1)}, \dots, \mu^{(k)}) = \sum_{|\lambda|=n} h_\lambda^{k-2} \prod_{i=1}^k \frac{\chi_\lambda(\mu^{(i)})}{z_{\mu^{(i)}}}. \quad (1.4)$$

Here  $h_\lambda$  is the product of the hook lengths of the Young diagram of the partition  $\lambda$  determining the irreducible representation and  $z_\mu$  is the order of the stabilizer of an element of the conjugacy class of cycle type  $\text{cyc}(\mu)$ .

The equivalence of the two definitions follows from the homomorphism from the fundamental group of the sphere punctured at the branch points to the symmetric group  $S_N$  generated by monodromy, which equates the identity element to the product of monodromies of paths consisting of simple loops around a suitably ordered sequence of all branch points.

## 1.2 Weighted Hurwitz numbers

### 1.2.1 Weight generating function $G(z)$

Following refs. [26] - [30], we define multiparametric weighted Hurwitz numbers by first introducing a generating function  $G(z)$  for the weights. This admits either a formal power series representation

$$G(z) = 1 + \sum_{i=1}^{\infty} g_i z^i \quad (1.5)$$

or an infinite product one

$$G(z) = \prod_{i=1}^{\infty} (1 + c_i z), \quad (1.6)$$

(or a limiting form of the latter), where  $\mathbf{c} = (c_1, c_2, \dots)$  denotes an infinite sequence of indeterminates that may be evaluated to real or complex constants. If the two expansions are compared, with (1.5) viewed as a formal series, this may be interpreted as the generating function for the elementary symmetric functions in the indeterminates  $(c_1, c_2, \dots)$ .

$$g_i = e_i(\mathbf{c}). \quad (1.7)$$

### 1.2.2 Geometrical definition: weighted coverings

**Definition 1.1.** Picking a pair of partitions  $(\mu, \nu)$  of  $N$ , and a positive integer  $d \in \mathbf{N}^+$ , the *geometrically weighted single and double Hurwitz numbers*  $H_G^d(\mu)$ ,  $H_G^d(\mu, \nu)$  associated with the generating function  $G$  are defined to be the sums

$$H_G^d(\mu) := \sum_{k=0}^d \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ |\mu^{(i)}| = N \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} \mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu), \quad (1.8)$$

$$H_G^d(\mu, \nu) := \sum_{k=0}^d \sum'_{\substack{\mu^{(1)}, \dots, \mu^{(k)} \\ |\mu^{(i)}| = N \\ \sum_{i=1}^k \ell^*(\mu^{(i)}) = d}} \mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)}) H(\mu^{(1)}, \dots, \mu^{(k)}, \mu, \nu), \quad (1.9)$$

where  $\sum'$  denotes a sum over all  $k$ -tuples of partitions  $\{\mu^{(1)}, \dots, \mu^{(k)}\}$  of  $N$  other than the cycle type of the identity element ( $1^N$ ) and the weights  $\mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)})$  are given by

$$\mathcal{W}_G(\mu^{(1)}, \dots, \mu^{(k)}) := \frac{1}{k!} \sum_{\sigma \in \mathfrak{S}_k} \sum_{1 \leq b_1 < \dots < b_k} c_{b_{\sigma(1)}}^{\ell^*(\mu^{(1)})} \dots c_{b_{\sigma(k)}}^{\ell^*(\mu^{(k)})} = \frac{|\text{aut}(\lambda)|}{k!} m_\lambda(\mathbf{c}). \quad (1.10)$$

Here

$$m_\lambda(\mathbf{c}) = \frac{1}{|\text{aut}(\lambda)|} \sum_{\sigma \in \mathfrak{S}_k} \sum_{1 \leq b_1 < \dots < b_k} c_{b_{\sigma(1)}}^{\lambda_1} \dots c_{b_{\sigma(k)}}^{\lambda_k}, \quad (1.11)$$

is the monomial sum symmetric function [36] of the indeterminates  $\mathbf{c}$  corresponding to the partition  $\lambda$  of length  $k$  and weight  $|\lambda| = d$  whose parts  $\{\lambda_i\}$  are the colengths  $\{\ell^*(\mu^{(i)})\}$ , written in weakly descending order,

$$\{\lambda_i\}_{i=1, \dots, k} \sim \{\ell^*(\mu^{(i)})\}_{i=1, \dots, k} \quad (1.12)$$

with normalization factor

$$|\text{aut}(\lambda)| := \prod_i m_i(\lambda)! \quad (1.13)$$

equal to the order of the automorphism group of the partition  $\lambda$ , where  $m_i(\lambda)$  is the number of parts of  $\lambda$  equal to  $i$ .

If the sums in (1.8) and (1.9) are taken over connected coverings only (or, equivalently, the factors  $(h_1, \dots, h_k)$  are required to generate a subgroup of  $S_N$  that acts transitively), we denote the corresponding sums  $\tilde{H}_G^d(\mu)$  and  $\tilde{H}_G^d(\mu, \nu)$ , respectively.

### 1.2.3 Combinatorial definition: weighted constellations

Another way to arrive at weighted Hurwitz numbers is through weighted *constellations* [16, 17, 35]. In [35] a constellation of length  $k$  and degree  $N$  is defined as a factorization (1.3) of the identity element in  $\mathbf{I} \in S_N$  into a product  $h_1 \dots h_k$  such that the group  $\langle h_1, \dots, h_k \rangle$  they generate acts transitively on the set  $(1, \dots, N)$ . Here, we consider an enhanced version, in which a weighting is assigned to all vertices and edges, and the total weight of the constellation is the product of these. We further consider  $k + 2$  constellations of degree (or weight)  $N$ , since these correspond to  $N$ -sheeted branched covers of the Riemann sphere in ramification points over  $k$  of the branch points are assigned a distinct weight in the enumeration, while two of them (say, over 0 and  $\infty$ ) are weighted differently in the interpretation of the  $\tau$ -function (2.1), (2.10) below as a generating function. The graph theoretic equivalent of a  $k + 2$ -constellation of degree (or weight)  $N$  is a bipartite graph on a Riemann surface whose two types of vertices are either *coloured* or *star* vertices, with all faces homeomorphic to a disc. (Thus, it is a *map* with certain additional special structure.) There are  $N$  star vertices, numbered consecutively from 1 to  $N$  and at most  $(N - 1)(k + 2)$  coloured vertices




Figure 1: An example of a constellation with  $N = 5$ ,  $k = 3$ , corresponding to the values  $N = 5$ ,  $k = 3$  and to the factorization  $h_0 h_1 h_2 h_3 h_4 = 1$  with  $h_1 = (135)$ ,  $h_2 = (15)(23)$ ,  $h_3 = (14)$ ,  $h_0 = (321)$ ,  $h_4 = h_\infty = (14)$ , with corresponding partitions  $\mu^{(1)} = (3, 1, 1)$ ,  $\mu^{(2)} = (2, 2, 1)$ ,  $\mu^{(3)} = (2, 1, 1, 1)$ ,  $\mu := \mu^{(0)} = (3, 1, 1)$ ,  $\nu := \mu^{(4)} = (2, 1, 1, 1)$ .

carrying one of  $k$  colours, labelled by distinct positive integers  $b_i$  with  $i$  ranging from 1 to  $k$ , plus a further two, labelled 0 and  $\infty \equiv k + 1$ , denoting, in our adaptation, the special case of “white” and “black” vertices respectively. Each star vertex has an edge connecting it to all  $k + 2$  coloured vertices, consecutively labelled counter-clockwise. The number of coloured vertices of given colour  $i$  is equal to the number of parts  $\ell(\mu^{(i)})$  of the partition  $\mu^{(i)}$  (i.e., its *length*), or  $\ell(\mu)$  for the “white” vertices and  $\ell(\nu)$  for the “black” ones. As an illustration, see the example of Figure 1 where  $N = 5$ ,  $k = 3$ .

Associated to each coloured vertex is a cycle in  $S_N$  obtained by listing the star vertices connected to it by an edge in counterclockwise order. Taking the product of the distinct cyclic permutations associated to each colour  $i = 0, \dots, k$ , we obtain an element  $h_i \in S_N$ , with cycle type  $\text{cyc}(\mu^{(i)})$ , where  $\mu^{(i)}$  is the partition of  $N$  whose parts are the cycle lengths of  $h_i$ . The element associated to the white vertices is denoted  $h_0 = h$  and its cycle type denoted  $\mu := \mu^{(0)}$ , the one associated to the black vertices whose cycle type is denoted  $\nu := \mu^{(k+1)}$  is  $h_{k+1}$  and the other coloured vertices are associated to the elements  $\{h_i\}_{i=1, \dots, k}$ , with cycle types  $\{\mu^{(i)}\}_{i=1, \dots, k}$  respectively, and satisfy

$$\mathbf{I} = h_0 h_1 \cdots h_{k+1} \quad (1.14)$$

To obtain the constellation associated with a particular branched cover of  $\mathbf{CP}^1$ , we place a coloured vertex of colour  $i$  at each of the ramification points  $\{Q_j^{(i)}\}_{j=1, \dots, \ell(\mu^{(i)})}$  over a given branch point  $\{Q^{(i)}\}_{i=0, \dots, k+1}$ , and choose a single generic (non-branched) point  $P \in \mathbf{CP}^1$  over which we have, in some specified order, the  $N$  points  $(P_1, \dots, P_N)$ , which are identified as the star vertices. For every simple loop starting and ending at  $P_i$ , going round one of the branch points, we have the monodromy lift on the covering surface which permutes the points  $\{P_i\}$  in such a way that each ramification point of colour  $i$  defines a disjoint cycle,




Figure 2: Lifted loops and edges

with the product of all cycles over a given branch point  $Q^{(i)}$  equal to the monodromy element  $h_i$ . The edges connect the ramification points to those  $P_i$ 's that belong to the given cycle. (See Figure 2.)

In order to define the weighting, we add a further pair of indeterminates  $(\beta, \gamma)$ , that serve as expansion parameters, and may, for weighting purposes, be evaluated to real or complex numbers. We then assign, to each (non-white, non-black) coloured vertex  $Q_j^{(i)}$  of colour  $b_i$ , a weight  $\beta^{-1}c_{b_i}^{-1}$ ; to each edge connecting non-white, non-black vertex  $Q_j^{(i)}$  to a star vertex, the weight  $\beta c_{b_i}$  to each white vertex, a weight

$$p_{\mu_j} = \mu_j t_{\mu_j}, \quad j = 1, \dots, \ell(\mu) \quad (1.15)$$

equal to the power sum symmetric function in an arbitrary number of auxiliary variables, where  $\mu_i$  is equal to the number of edges connecting that white vertex to star vertices; to each black vertex, a weight

$$p_{\nu_j} = \nu_j s_{\nu_j}, \quad j = 1, \dots, \ell(\nu) \quad (1.16)$$

equal to the power sum symmetric function in a second set of auxiliary variables, where  $\nu_i$  is equal to the number of edges connecting that black vertex to star vertices; to the edges connecting the white or black ramification points  $\{Q_j^{(0)}, Q_j^{(k+1)}\}$  to any star vertex, the weight 1, and to each of the star vertices, the weight

$$\gamma \quad (1.17)$$

The doubly infinite set of parameters  $\mathbf{t} = (t_1, t_2, \dots)$  and  $\mathbf{s} = (s_1, s_2, \dots)$  are interpreted as the two independent KP flow parameters that appear in the  $2D$ -Toda  $\tau$ -function [47, 49, 50, 52].

Taking the product of these over all vertices, edges and faces gives the overall weight

$$\frac{\gamma^N \beta^d}{N!} \prod_{i=1}^k c_{b_i}^{\ell^*(\mu^{(i)})} p_\mu(\mathbf{t}) p_\nu(\mathbf{s}) \quad (1.18)$$

for the constellation, where the standard normalization factor  $\frac{1}{N!}$  is included in the enumeration,

$$d := \sum_{i=1}^k \ell^*(\mu^{(i)}) \quad (1.19)$$

is related to the Euler characteristic of the surface by the Riemann-Hurwitz relation

$$2 - 2g = 2N - d \quad (1.20)$$

and

$$p_\mu(\mathbf{t}) := \prod_{i=1}^{\ell(\mu)} p_{\mu_i}, \quad p_\nu(\mathbf{s}) := \prod_{i=1}^{\ell(\nu)} p'_{\nu_i}. \quad (1.21)$$

Summing these over all such weighted constellations, with all possible choices of  $k$  distinct colours, chosen from amongst the indices labelling the nonzero constants  $c_i$  in (1.6), corresponding to a factorization (1.14) with the same cycle types, permuted in all ways, normalized by  $\frac{1}{|\text{aut}(\lambda)|}$ , where  $\lambda$  is the partition of weight

$$|\lambda| = d := \sum_{i=1}^k \ell^*(\mu^{(i)}) \quad (1.22)$$

whose parts are the colengths  $\{\ell^*(\mu^{(i)})\}$ , suitably ordered, and summing over the integer  $k$ , we obtain the total weighting factor  $m_\lambda(c_1, c_2, \dots)$  of (1.11), multiplied by the Hurwitz number  $H(\mu, \mu^{(1)}, \dots, \mu^{(k)}, \nu)$  which enumerates the number of products (1.14) with factors in the conjugacy classes  $(\mu, \mu^{(1)}, \dots, \mu^{(k)}, \nu)$ , and recover the  $\tau$ -function (2.9) as generating function, as explained below.

## 2 Hypergeometric $\tau$ -function, Baker function, recursions and correlators

### 2.1 Hypergeometric $\tau$ -functions as generating functions for weighted Hurwitz numbers

Following refs. [26–28, 30], we define a family of 2D Toda  $\tau$ -functions of hypergeometric type [33, 43, 44] associated to the generating function  $G$  and a pair  $(\beta, \gamma)$  of complex parameters by the double Schur function [36] expansion:

$$\tau^{(G, \beta, \gamma)}(\mathbf{t}, \mathbf{s}) = \sum_{\lambda} \gamma^{|\lambda|} r_{\lambda}^{(G, \beta)} s_{\lambda}(\mathbf{t}) s_{\lambda}(\mathbf{s}), \quad (2.1)$$

where the Schur functions  $s_\lambda(\mathbf{t}), s_\lambda(\mathbf{s})$  are viewed as functions of two infinite sequences of 2-Toda flow variables  $\mathbf{t} = (t_1, t_2, \dots)$ ,  $\mathbf{s} = (s_1, s_2, \dots)$  and the *content product* coefficients  $r_\lambda^{(G,\beta)}$  are defined in terms of the weight generating function  $G$  as

$$r_\lambda^{(G,\beta)} := \prod_{(i,j) \in \lambda} r_{j-i}^G(\beta), \quad (2.2)$$

where

$$r_j^G(\beta) := G(j\beta). \quad (2.3)$$

Denote by  $\mathbb{K} = \mathbb{Q}[g_1, \dots, g_M, \dots]$  the algebra of polynomials in the  $g_k$ 's, with rational coefficients or, equivalently, the algebra of symmetric functions of the  $c_i$ 's. In the following, it will also be useful to define the quantities  $\{\rho_j, T_j\}_{j \in \mathbf{Z}}$  by

$$r_j^G(\beta) = \frac{\rho_j}{\gamma \rho_{j-1}}, \quad (2.4)$$

$$\rho_j := \gamma^j \prod_{i=1}^j G(i\beta) = e^{T_j}, \quad \rho_0 = 1, \quad (2.5)$$

$$\rho_{-j} := \gamma^{-j} \prod_{i=0}^{j-1} (G(-i\beta))^{-1} = e^{T_{-j}}, \quad j = 1, 2, \dots \quad (2.6)$$

Moreover, it is easily seen that there exists a formal power series  $T(x)$  such that

$$e^{T(x) - T(x-1)} = \gamma G(\beta x), \quad (2.7)$$

with  $T(0) = 0$ ,  $T(i) = T_i$ .

As shown in refs. [26–28, 30], identifying the flow parameters as

$$t_i = \frac{p_i}{i}, \quad s_i = \frac{p'_i}{i},$$

$$p_\mu(\mathbf{t}) := \prod_{i=1}^{\ell(\mu)} p_{\mu_i}, \quad p_\nu(\mathbf{s}) := \prod_{i=1}^{\ell(\nu)} p'_{\nu_i} \quad (2.8)$$

and changing to the basis of power sum symmetric functions [36], the  $\tau$ -function (2.1) serves as a generating function for the weighted Hurwitz numbers defined above.

**Theorem 2.1.**

$$\tau^{(G,\beta,\gamma)}(\mathbf{t}, \mathbf{s}) = \sum_{d=0}^{\infty} \beta^d \sum_{\substack{\mu, \nu \\ |\mu|=|\nu|}} \gamma^{|\mu|} H_G^d(\mu, \nu) p_\mu(\mathbf{t}) p_\nu(\mathbf{s}). \quad (2.9)$$

By standard combinatorial arguments, taking its logarithm gives the generating function for connected weighted Hurwitz numbers:

$$\ln(\tau^{(G,\beta,\gamma)}(\mathbf{t}, \mathbf{s})) = \sum_{d=0}^{\infty} \beta^d \sum_{\substack{\mu, \nu \\ |\mu|=|\nu|}} \gamma^{|\mu|} \tilde{H}_G^d(\mu, \nu) p_\mu(\mathbf{t}) p_\nu(\mathbf{s}). \quad (2.10)$$

All instances of weighted Hurwitz numbers to have appeared in the literature [1, 6–9, 14, 25–30, 34, 35, 41, 45, 53] have  $\tau$ -functions of hypergeometric type as generating functions and are either special cases of the above or are similarly derived from a second class of “dual” weight generating functions introduced in [27, 28], or a more general quantum weighting introduced in [29].

## 2.2 Baker function, adapted bases and recursions

Viewed purely as a KP  $\tau$ -function with respect to the first set of flow variables  $\mathbf{t} = (t_1, t_2, \dots)$ , with the remaining variables  $\mathbf{s} = (s_1, s_2, \dots)$  interpreted as auxiliary parameters, the corresponding parametric family of Baker functions  $\Psi_{(G,\beta,\gamma)}^-(\zeta, \mathbf{t}, \mathbf{s})$  and their duals  $\Psi_{(G,\beta,\gamma)}^+(\zeta, \mathbf{t}, \mathbf{s})$  are given by the Sato formula

$$\Psi_{(G,\beta,\gamma)}^\mp(\zeta, \mathbf{t}, \mathbf{s}) = e^{\pm \sum_{i=1}^{\infty} t_i \zeta^i} \frac{\tau^{(G,\beta,\gamma)}(\mathbf{t} \mp [x], \mathbf{s})}{\tau^{(G,\beta,\gamma)}(\mathbf{t}, \mathbf{s})} = e^{\pm \sum_{i=1}^{\infty} t_i \zeta^i} (1 + \mathcal{O}(1/\zeta)), \quad (2.11)$$

where

$$\zeta := x^{-1} \quad (2.12)$$

is the spectral parameter and  $[x]$  is the infinite vector whose components are equal to the terms of the Taylor expansion of  $-\ln(1-x)$  at  $x=0$

$$[x] := \left(x, \frac{x^2}{2}, \dots, \frac{x^n}{n}, \dots\right). \quad (2.13)$$

In the following, instead of considering the values of the Baker function and its dual at arbitrary KP flow parameter values, we shall be evaluating at  $\mathbf{t} = \mathbf{0}$ , and  $\beta^{-1}\mathbf{s}$

$$\Psi_{(G,\beta,\gamma)}^+(1/x, \mathbf{0}, \beta^{-1}\mathbf{s}) = \gamma \sum_{j=0}^{\infty} \rho_{j-1} h_j(\beta^{-1}\mathbf{s}) x^j, \quad (2.14)$$

$$\Psi_{(G,\beta,\gamma)}^-(1/x, \mathbf{0}, \beta^{-1}\mathbf{s}) = \sum_{j=0}^{\infty} \rho_{-j}^{-1} h_j(-\beta^{-1}\mathbf{s}) x^j, \quad (2.15)$$

where  $\{h_j\}_{j \in \mathbb{N}}$  are the complete symmetric functions, defined by

$$e^{\sum_{i=0}^{\infty} t_i \zeta^i} = \sum_{j=0}^{\infty} h_j(\mathbf{t}) \zeta^j. \quad (2.16)$$

More generally, we define the adapted basis  $\{\Psi_k^+(x)\}_{k \in \mathbb{Z}}$  and its dual  $\{\Psi_k^-(x)\}_{k \in \mathbb{Z}}$  by

$$\Psi_k^+(x) := \gamma \sum_{j=0}^{\infty} \rho_{j+k-1} h_j(\beta^{-1}\mathbf{s}) x^{j+k}, \quad (2.17)$$

$$\Psi_k^-(x) := \sum_{j=0}^{\infty} \rho_{-j-k}^{-1} h_j(-\beta^{-1}\mathbf{s}) x^{j+k} \quad (2.18)$$

for all  $k \in \mathbf{Z}$  so that, in particular

$$\Psi_0^-(x) = \Psi_{(G,\beta,\gamma)}^-(1/x, \mathbf{0}, \beta^{-1}\mathbf{s}), \quad \Psi_0^+(x) = \Psi_{(G,\beta,\gamma)}^+(1/x, \mathbf{0}, \beta^{-1}\mathbf{s}). \quad (2.19)$$

Note that we may view these as formal series in the parameters:

$$\Psi_k^+(x), \Psi_k^-(x) \in \gamma^k \mathbb{K}[x, x^{-1}, \mathbf{s}, \beta, \beta^{-1}][[\gamma]]. \quad (2.20)$$

**Remark 2.1.** The Baker function  $\Psi_{(G,\beta,\gamma)}^-(\zeta = 1/x, \mathbf{t}, \beta^{-1}\mathbf{s})$  and its dual  $\Psi_{(G,\beta,\gamma)}^+(\zeta = 1/x, \mathbf{t}, \beta^{-1}\mathbf{s})$ , as defined in (2.18), (2.17), are elements of the subspaces

$$W_{(G,\beta,\gamma,\mathbf{s})}^\pm := \text{span}\{\Psi_{-k}^\pm(\zeta = 1/x)\}_{k \in \mathbf{N}} \subset \mathbb{K}[x, x^{-1}, \mathbf{s}, \beta, \beta^{-1}][(\gamma)] \quad (2.21)$$

for all values of the KP flow parameters  $\mathbf{t}$ . These may be understood as elements of formal power series analogs of the Sato-Segal-Wilson Grassmannian, consisting of subspaces of the Hilbert space spanned by powers  $\{\zeta^k\}_{k \in \mathbf{Z}}$  of the spectral parameter  $\zeta = 1/x$ , commensurable with the subspace spanned by the monomials  $\{\zeta^k\}_{k=0,1,\dots}$ .

**Definition 2.1.** Define the recursion operators

$$R_\pm(x) := \gamma x G(\pm \beta D), \quad (2.22)$$

where

$$D := x \frac{d}{dx} \quad (2.23)$$

is the Euler operator.

**Lemma 2.2.** *These satisfy the commutation relations*

$$[D, R_\pm] = R_\pm. \quad (2.24)$$

Note also that the operators  $R_\pm$  can be expressed as

$$R_+ = e^{T(D-1)} \circ x \circ e^{-T(D-1)}, \quad (2.25)$$

$$R_- = e^{-T(-D)} \circ x \circ e^{T(-D)}. \quad (2.26)$$

By application of these operators to the series expansion (2.17) and (2.18), using (2.5) and (2.6), it follows that the basis elements  $\{\Psi_k^\pm(x)\}$  satisfy the recursion relations

**Proposition 2.3.**

$$\Psi_{k\pm 1}^+(x) := R_\pm^{\pm 1} \Psi_k^+(x), \quad (2.27)$$

$$\Psi_{k\pm 1}^-(x) := R_\pm^{\pm 1} \Psi_k^-(x). \quad (2.28)$$

From the identity

$$\sum_{j=0}^k h_j(\mathbf{t})h_{k-j}(-\mathbf{t}) = \delta_{k0}, \quad (2.29)$$

for complete symmetric functions, it also follows that the bases consisting of functions (2.17) and (2.18) are dual to each other

$$(\Psi_k^+, \Psi_m^-) = \gamma \delta_{k+m,1} \quad (2.30)$$

with respect to the pairing

$$(f, g) := \frac{1}{2\pi i} \oint_0 f(x)g(x) \frac{dx}{x^2}. \quad (2.31)$$

In particular, all  $\Psi_k^+$  for  $k \leq 0$  are orthogonal to all  $\Psi_l^-$  for  $l \leq 0$ , and this is equivalent to the Hirota bilinear relations for the KP hierarchy with respect to the times  $\mathbf{t}$ .

### 2.3 The pair correlator and its expansion

We now define the *pair correlation function*  $K(x, x')$  in terms of the  $\tau$ -function.

**Definition 2.2.** The single *pair correlator* is defined as

$$K(x; x') := \frac{1}{x - x'} \tau^{(G, \beta, \gamma)}([x] - [x'], \beta^{-1}\mathbf{s}), \quad (2.32)$$

while more generally, for  $n \geq 1$ , we define the  $n$ -pair correlator

$$K_n(x_1, \dots, x_n; x'_1, \dots, x'_n) := \det \left( \frac{1}{x_i - x'_j} \right)_{1 \leq i, j \leq n} \times \tau^{(G, \beta, \gamma)} \left( \sum_i [x_i] - [x'_i], \beta^{-1}\mathbf{s} \right). \quad (2.33)$$

As a formal series, for any  $n \geq 1$ ,  $K_n(x_1, \dots, x_n; x'_1, \dots, x'_n)$  may be viewed as belonging to  $\mathbb{K}(x_1, \dots, x_n, x'_1, \dots, x'_n)[\mathbf{s}, \beta, \beta^{-1}][[\gamma]]$ . It is a standard result, following from the Cauchy-Binet identity or, equivalently, the fermionic Wick theorem (cf. ref. [2]), that

**Proposition 2.4.**

$$K_n(x_1, \dots, x_n; x'_1, \dots, x'_n) = \det (K(x_i; x'_j)). \quad (2.34)$$

This should be interpreted as holding in  $\mathbb{K}(x_1, \dots, x_n, x'_1, \dots, x'_n)[\mathbf{s}, \beta, \beta^{-1}][[\gamma]]$ . We may also express  $K(x; x')$  as a formal double Taylor series with coefficients given by the complete symmetric functions and the weighting parameters  $\rho_i$ .

**Proposition 2.5.** *The function  $K(x, x')$ , can be expressed as:*

$$\begin{aligned} K(x, x') &= \frac{1}{x - x'} + \sum_{a, b \geq 0} \rho_a \rho_{-b-1}^{-1} x^a (-x')^b s_{(a|b)}(\beta^{-1}\mathbf{s}) \\ &= \frac{1}{x - x'} + \sum_{a=0}^{\infty} \sum_{b=0}^{\infty} \sum_{j=1}^{b+1} \rho_a h_{a+j}(\beta^{-1}\mathbf{s}) x^a \rho_{-b-1}^{-1} h_{b-j+1}(-\beta^{-1}\mathbf{s}) (x')^b. \end{aligned} \quad (2.35)$$

*Proof.* This follows from substituting the Schur function expansion (2.1) of the  $\tau$ -function  $\tau^{(G,\beta,\gamma)}(\mathbf{t}, \mathbf{s})$ , into the definition (2.32) of  $K(x, x')$  and noting that, making the special evaluation at  $[x] - [x']$ , this is an expansion in which only hook partitions appear, for which the Schur functions can be expressed as finite bilinear sums over the complete symmetric functions  $\{h_i\}$  [36].  $\square$

## 2.4 The multicurrent correlator as generating function

In terms of the indeterminates  $(x_1, \dots, x_n)$ , we define the following correlators

$$W_n(\mathbf{s}; x_1, \dots, x_n) := \left( \left( \prod_{i=1}^n \nabla(x_i) \right) \tau^{(G,\beta,\gamma)}(\mathbf{t}, \beta^{-1}\mathbf{s}) \right) \Big|_{\mathbf{t}=\mathbf{0}}, \quad (2.36)$$

$$\tilde{W}_n(\mathbf{s}; x_1, \dots, x_n) := \left( \left( \prod_{i=1}^n \nabla(x_i) \right) \ln \tau^{(G,\beta,\gamma)}(\mathbf{t}, \beta^{-1}\mathbf{s}) \right) \Big|_{\mathbf{t}=\mathbf{0}}, \quad (2.37)$$

where

$$\nabla(x) := \sum_{i=1}^{\infty} x^{i-1} \frac{\partial}{\partial t_i}. \quad (2.38)$$

**Remark 2.2.** As will be seen in the fermionic representation in Sec. 5 below, these may be viewed as *multicurrent correlators* and *connected multicurrent correlators*.

For  $g \geq 0$  we denote the genus- $g$  component of the function  $\tilde{W}_n(\mathbf{s}; x_1, \dots, x_n)$  as follows

$$\tilde{W}_{g,n}(\mathbf{s}; x_1, \dots, x_n) := [\beta^{2g-2+n}] \tilde{W}_n(\mathbf{s}; x_1, \dots, x_n). \quad (2.39)$$

We now introduce another type of generating function for weighted Hurwitz numbers, both for the connected and nonconnected case.

**Definition 2.3.**

$$F_n(\mathbf{s}; x_1, \dots, x_n) := \sum_{\mu, \nu, \ell(\mu)=n} \sum_d \gamma^{|\mu|} \beta^{d-\ell(\nu)} H_G^d(\mu, \nu) |\text{aut}(\mu)| m_\mu(x_1, \dots, x_n) p_\nu(\mathbf{s}) \quad (2.40)$$

$$\tilde{F}_n(\mathbf{s}; x_1, \dots, x_n) := \sum_{\mu, \nu, \ell(\mu)=n} \sum_d \gamma^{|\mu|} \beta^{d-\ell(\nu)} \tilde{H}_G^d(\mu, \nu) |\text{aut}(\mu)| m_\mu(x_1, \dots, x_n) p_\nu(\mathbf{s}) \quad (2.41)$$

$$\tilde{F}_{g,n}(\mathbf{s}; x_1, \dots, x_n) := \sum_{\mu, \nu, \ell(\mu)=n} \gamma^{|\mu|} \tilde{H}_G^d(\mu, \nu) |\text{aut}(\mu)| m_\mu(x_1, \dots, x_n) p_\nu(\mathbf{s}), \quad (2.42)$$

where  $m_\mu(x_1, \dots, x_n)$  denotes the monomial sum symmetric function corresponding to the partition  $\mu$ , and  $(g, d)$  are related by the Riemann-Hurwitz formula

$$2 - 2g = \ell(\mu) + \ell(\nu) - d. \quad (2.43)$$

Note that  $F_n, \tilde{F}_n$  belong to  $\mathbb{K}[x_1, \dots, x_n; \mathbf{s}; \beta, \beta^{-1}][[\gamma]]$  and  $\tilde{F}_{g,n}$  to  $\mathbb{K}[x_1, \dots, x_n; \mathbf{s}][[\gamma]]$ . These are related to the multicurrent correlators as follows

**Theorem 2.6.**

$$W_n(\mathbf{s}; x_1, \dots, x_n) = \frac{\partial^n}{\partial x_1 \cdots \partial x_n} F_n(\mathbf{s}; x_1, \dots, x_n), \quad (2.44)$$

$$\tilde{W}_n(\mathbf{s}; x_1, \dots, x_n) = \frac{\partial^n}{\partial x_1 \cdots \partial x_n} \tilde{F}_n(\mathbf{s}; x_1, \dots, x_n), \quad (2.45)$$

$$\tilde{W}_{g,n}(\mathbf{s}; x_1, \dots, x_n) = \frac{\partial^n}{\partial x_1 \cdots \partial x_n} \tilde{F}_{g,n}(\mathbf{s}; x_1, \dots, x_n). \quad (2.46)$$

*Proof.* Applying the operator  $\prod_{i=1}^n \nabla(x_i)$  to the symmetric function  $p_\mu(\mathbf{t})$  gives

$$\begin{aligned} \left( \prod_{i=1}^n \nabla(x_i) \right) p_\mu(\mathbf{t}) \Big|_{\mathbf{t}=\mathbf{0}} &= \prod_{i=1}^n \left( \sum_{j_i=1}^{\infty} x_i^{j_i-1} \frac{\partial}{\partial t_{j_i}} \right) \left( \prod_{k=1}^{\ell(\mu)} \mu_k t_{\mu_k} \right) \Big|_{\mathbf{t}=\mathbf{0}} \\ &= \delta_{\ell(\mu), n} \sum_{\sigma \in S_n} \prod_{k=1}^n \mu_k x_{\sigma(k)}^{\mu_k-1} = \delta_{\ell(\mu), n} |\text{aut}(\mu)| \frac{\partial^n m_\mu(x_1, \dots, x_n)}{\partial x_1 \cdots \partial x_n} \end{aligned} \quad (2.47)$$

by (1.11). Hence, applying it to  $\tau^{(G, \beta, \gamma)}(\mathbf{t}, \beta^{-1}\mathbf{s})$ , using the expansion (2.9), we obtain from the definition (2.36) of  $W_n(\mathbf{s}, x_1, \dots, x_n)$

$$\begin{aligned} W_n(\mathbf{s}, x_1, \dots, x_n) &= \sum_{\mu, \nu, \ell(\mu)=n} \sum_d \gamma^{|\mu|} \beta^{d-\ell(\nu)} H_G^d(\mu, \nu) |\text{aut}(\mu)| \frac{\partial^n m_\mu(x_1, \dots, x_n)}{\partial x_1 \cdots \partial x_n} p_\nu(\mathbf{s}) \\ &= \frac{\partial^n}{\partial x_1 \cdots \partial x_n} F_n(\mathbf{s}; x_1, \dots, x_n) \end{aligned} \quad (2.48)$$

by (2.40), proving (2.44). The same calculation proves the connected case (2.45), and (2.46) follows from equating like powers of  $\beta$  in the series expansion.  $\square$

Henceforth, we suppress the explicit  $\mathbf{s}$  dependence in the arguments and denote these quantities simply as  $W_n(x_1, \dots, x_n)$ , etc. It follows from (2.36) that  $W_n(x_1, \dots, x_n)$  may equivalently be expressed directly in terms of derivatives of special evaluations of the  $\tau$ -function  $\tau^{(G, \beta, \gamma)}(\mathbf{t}, \mathbf{s})$ .

**Lemma 2.7.**

$$W_n(x_1, \dots, x_n) = [\epsilon_1, \dots, \epsilon_n] \left( K_n(x_1 + \epsilon_1, \dots, x_n + \epsilon_n; x_1, \dots, x_n) / \det \left( \frac{1}{x_i - x_j + \epsilon_i} \right) \right), \quad (2.49)$$

where both sides are viewed as elements of  $\mathbb{K}[x_1, \dots, x_n, \mathbf{s}, \beta, \beta^{-1}][[\gamma]]$ .

For connected functions, we get the specially elegant relations:

**Proposition 2.8.**

$$\tilde{W}_1(x) = \lim_{x' \rightarrow x} \left( K(x, x') - \frac{1}{x - x'} \right), \quad (2.50)$$

$$\tilde{W}_2(x_1, x_2) = \left( -K(x_1, x_2)K(x_2, x_1) - \frac{1}{(x_1 - x_2)^2} \right), \quad (2.51)$$

and for  $n \geq 3$

$$\tilde{W}_n(x_1, \dots, x_n) = \sum_{\sigma \in \mathfrak{S}_n^{1\text{-cycle}}} (-1)^\sigma \prod_i K(x_i, x_{\sigma(i)}), \quad (2.52)$$

where the last sum is over all permutations in  $\mathfrak{S}_n$  having only one cycle. These equalities hold in  $\mathbb{K}(x, x_1, \dots, x_n)[\mathbf{s}, \beta, \beta^{-1}][[\gamma]]$ .

A detailed proof of this result is provided in the appendix to [3].

## 2.5 The Christoffel-Darboux relation

Define the functions  $\xi(x, \mathbf{s})$  and  $S(x, \mathbf{s})$  by

$$\xi(x, \mathbf{s}) := \sum_{k=1}^{\infty} s_k x^k, \quad (2.53)$$

$$S(x, \mathbf{s}) := x \frac{d}{dx} \xi(x, \mathbf{s}) = \sum_{k=1}^{\infty} k s_k x^k. \quad (2.54)$$

In the following, we assume that only a finite number of variables  $s_k$  are nonzero, so  $\xi(x, \mathbf{s})$  and  $S(x, \mathbf{s})$  are polynomials

$$\xi(x, \mathbf{s}) = \sum_{k=1}^L s_k x^k, \quad S(x, \mathbf{s}) = \sum_{k=1}^L k s_k x^k \quad (2.55)$$

of degree  $L$ . We also assume that only the first  $M$  parameters  $\{c_i\}$  are nonzero, so the weight generating function  $G(z)$  is also a polynomial, of degree  $M$

$$G(z) = \prod_{i=1}^M (1 + c_i z). \quad (2.56)$$

For brevity, the explicit dependence on the parameters  $\mathbf{s}$  will henceforth also be suppressed and  $S(x, \mathbf{s})$  and  $\xi(x, \mathbf{s})$  just denoted  $S(x)$ ,  $\xi(x)$ .

**Definition 2.4.** Define the operators

$$\Delta_{\pm}(x) := \pm \beta e^{\mp \beta^{-1} \xi(x)} \circ D \circ e^{\pm \beta^{-1} \xi(x)} = S(x) \pm \beta D \quad (2.57)$$

and

$$V_{\pm}(x) := \gamma^{-1} x^{-1} e^{\mp \beta^{-1} \xi(x)} \circ R_{\pm} \circ e^{\pm \beta^{-1} \xi(x)} = G(\Delta_{\pm}(x)). \quad (2.58)$$

In the following, we use boldface uppercase letters, such as  $\mathbf{A}$ ,  $\mathbf{F}$  or  $\mathbf{V}(x)$  to denote finite dimensional matrices.

**Definition 2.5.** Define the polynomial  $A(r, t)$  of degree  $LM - 1$  in each variable  $(r, t)$  which determines the  $LM \times LM$  nonsingular matrix  $\mathbf{A} = (\mathbf{A}_{ij})_{0 \leq i, j < LM}$  as follows

$$A(r, t) := (r V_-(t) - t V_+(r)) \left( \frac{1}{r - t} \right) = \sum_{i=0}^{LM-1} \sum_{j=0}^{LM-1} \mathbf{A}_{ij} r^i t^j. \quad (2.59)$$

The highest total degree term is

$$g_M (Ls_L)^M \frac{r t^{LM} - t r^{LM}}{r - t} = -g_M (Ls_L)^M \sum_{j=1}^{LM-1} r^j t^{LM-j} \quad (2.60)$$

so

$$\det \mathbf{A} = (-1)^{\frac{LM(LM-1)}{2}} g_M^{LM-1} (Ls_L)^{M(LM-1)}. \quad (2.61)$$

The following *Christoffel-Darboux* type relation, expressing  $K(x, x')$  as a finite sum of basis elements is proved in [2, 3].

**Theorem 2.9.**

$$K(x, x') = \frac{1}{x - x'} A(R_+, R_-) \Psi_0^+(x) \Psi_0^-(x'). \quad (2.62)$$

**Remark 2.3.** In the above, we made the assumption that  $G$  is a polynomial, of degree  $M$ . We keep this assumption in the following; however, because any given coefficient in the  $\tau$ -function is a polynomial in the coefficients of  $G$ , it is possible to drop it in many instances by taking projective limits. Indeed if  $G(z)$  is an infinite formal power series and  $G_M(z)$  is its degree- $M$  truncation, then all the correlators corresponding to  $G(z)$  are the limit (in the sense of convergence of individual coefficients) of those corresponding to  $G_M(z)$  as  $M \rightarrow \infty$ . This enables us, for example, to include the exponential weight generating function  $G(z) = e^z$  which gives the case of simple (single and double) Hurwitz numbers studied in [41, 45].

## 3 The spectral curve

### 3.1 Quantum spectral curve

Applying the operators  $\beta D$  and  $S(R_\pm)$  to the series expansions (2.14), (2.15) we obtain:

**Theorem 3.1.** *The function  $\Psi_0^+(x)$  satisfies*

$$(\beta D - S(R_+)) \Psi_0^+(x) = 0, \quad (3.1)$$

*and  $\Psi_0^-(x)$  satisfies the dual equation*

$$(\beta D + S(R_-)) \Psi_0^-(x) = 0. \quad (3.2)$$

The linear ordinary differential operators  $(\beta D - S(R_+))$  and  $(\beta D + S(R_-))$  with polynomial coefficients that annihilate  $\Psi_0^+(x)$ ,  $\Psi_0^-(x)$  in (3.1), (3.2) will be referred to as the *quantum spectral curve* and its dual.

### 3.2 Classical spectral curve

The classical spectral curve is obtained by replacing the Euler operator appearing in (3.1), (3.2) and (2.22) by the product  $\beta^{-1}xy$ , or by  $-\beta^{-1}xy$  in the dual case. Here  $y$  is viewed as the classical variable canonically conjugate to  $x$  (and  $\beta$  is identified with Planck's constant  $\hbar$ ). This gives

$$xy = S(\gamma x G(xy)) \quad (3.3)$$

as the classical spectral curve. This is a rational curve realized as an  $LM$ -sheeted branched cover of  $\mathbf{CP}^1$ . The significance of this is that  $y dx$ , viewed as a meromorphic form on this Riemann surface, is equal to the rational differential  $\tilde{W}_{0,1}(x)dx$ , as shown in ref. [3].

**Proposition 3.2.**

$$\tilde{F}_{0,1}(x) = \int y dx. \quad (3.4)$$

The spectral curve admits a rational parametrization by the functions  $X(z)$  and  $Y(z)$  defined as:

$$\begin{aligned} X(z) &:= \frac{z}{\gamma G(S(z))}, \\ Y(z) &:= \frac{S(z)}{z} \gamma G(S(z)). \end{aligned} \quad (3.5)$$

Note that we have

$$X(z)Y(z) = S(z). \quad (3.6)$$

The ramification points under the projection to the Riemann sphere  $(x, y) \rightarrow x$  are given by

$$\frac{dX}{dz} = 0, \quad \text{with} \quad \frac{dY}{dz} \neq 0. \quad (3.7)$$

Equivalently, the ramification points  $(X(z), Y(z))$  are given by  $z$ -values in the set  $\mathcal{A}$  of roots of the polynomial defined as follows

$$\sigma(z) := G(S(z)) - zS'(z)G'(S(z)), \quad (3.8)$$

and the point  $z = \infty$ , which we do not include in  $\mathcal{A}$ .

Assuming  $x$  is *not* a branch point and picking a point  $(X(z), Y(z))$  over  $x$  with uniformizing parameter  $z$ , over a neighbourhood of  $x = X(z)$  that contains no branch points, the equation  $X(z) = x$ , has  $LM$  solutions, which we label as follows:

$$\{z^{(0)}(z) := z, z^{(1)}(z), \dots, z^{(LM-1)}(z)\} \quad (3.9)$$

so

$$X(z^{(i)}(z)) = x, \quad i = 0, \dots, LM - 1, \quad (3.10)$$

The ordering for the roots  $\{z^{(1)}(z), \dots, z^{(LM-1)}(z)\}$  is fixed arbitrarily, once and for all, within this neighbourhood; and the functions  $z^{(i)}(z)$  are analytic in  $z$ , as are  $X(z)$ ,  $Y(z)$  and  $1/X'(z)$ .

**Definition 3.1.** For  $z$  as above let  $\mathbf{V}(z)$  denote the Vandermonde matrix with elements

$$\mathbf{V}(z)_{ij} = (z^{(j)}(z))^{i-1}. \quad (3.11)$$

## 4 Linear first order system for the adapted basis

### 4.1 The constant coefficient infinite system

Define two doubly infinite column vectors whose components are the functions  $\Psi_k^+(x)$  and  $\Psi_k^-(x)$ .

$$\vec{\Psi}_\infty^+ := \begin{pmatrix} \vdots \\ \Psi_{-1}^+ \\ \Psi_0^+ \\ \Psi_1^+ \\ \vdots \end{pmatrix}, \quad \vec{\Psi}_\infty^- := \begin{pmatrix} \vdots \\ \Psi_{-1}^- \\ \Psi_0^- \\ \Psi_1^- \\ \vdots \end{pmatrix}. \quad (4.1)$$

Here, and in what follows, the word *constant* refers to a quantity that does not depend on  $x$  or  $x'$ . Now define four doubly infinite constant matrices  $Q^\pm$  and  $P^\pm$  by their generating functions

$$\mathcal{P}^\pm(t, r) = \sum_{i, j=-\infty}^{\infty} P_{ij}^\pm t^i r^j = \Delta_\pm(r) \sum_{i=-\infty}^{\infty} (tr)^i, \quad (4.2)$$

$$\mathcal{Q}^\pm(t, r) = \sum_{i, j=-\infty}^{\infty} Q_{ij}^\pm t^i r^j = V_\pm(r) r^{-1} \sum_{i=-\infty}^{\infty} (tr)^i. \quad (4.3)$$

Note that  $P^\pm$  are upper triangular, with diagonal entries  $\mathcal{P}_{kk}^\pm = \pm k\beta$  and upper triangular entries  $\mathcal{P}_{kj}^\pm = (j - k)s_{j-k}$ , while  $Q^\pm$  are finite band matrices, with just one band below the principal diagonal, and  $LM - 2$  above it.

As shown in refs. [2, 3], the recursion relations (2.27), (2.28), together with the commutator relation (2.24) can be expressed equivalently as the following linear, constant coefficient system for  $\Psi_k^+(x)$  and  $\Psi_k^-(x)$  :

**Theorem 4.1.**

$$\beta D \vec{\Psi}_\infty^+ = P^+ \vec{\Psi}_\infty^+, \quad (4.4)$$

$$\frac{1}{\gamma x} \vec{\Psi}_\infty^+ = Q^+ \vec{\Psi}_\infty^+, \quad (4.5)$$

and

$$-\beta D\vec{\Psi}_\infty^- = P^-\vec{\Psi}_\infty^-, \quad (4.6)$$

$$\frac{1}{\gamma x}\vec{\Psi}_\infty^- = Q^-\vec{\Psi}_\infty^-. \quad (4.7)$$

## 4.2 Finite dimensional first order system: folding

Using the multiplicative recursion relations (4.5), (4.7) and the finite band structure of the matrices  $\mathcal{Q}^\pm$ , we can “fold” the constant coefficient terms on the RHS of eqs. (4.4) and (4.6) onto a finite “window” consisting of the first  $LM$  components of the vectors  $\vec{\Psi}_\infty^+$  and  $\vec{\Psi}_\infty^-$ , at the cost of introducing a polynomial dependence upon  $1/x$  in the coefficients. In fact, due to the special structure of eqs. (4.5), (4.7), this dependence turns out to just be linear in  $1/x$ .

Define the two column vectors  $\vec{\Psi}^+(x)$  and  $\vec{\Psi}^-(x)$  of dimension  $LM$  by

$$\vec{\Psi}^+ := \begin{pmatrix} \Psi_0^+ \\ \Psi_1^+ \\ \Psi_2^+ \\ \dots \\ \Psi_{LM-1}^+ \end{pmatrix}, \quad \vec{\Psi}^- := \begin{pmatrix} \Psi_0^- \\ \Psi_1^- \\ \Psi_2^- \\ \dots \\ \Psi_{LM-1}^- \end{pmatrix}. \quad (4.8)$$

Let the elements  $\tilde{\mathbf{E}}_{ij}(x)$  of the  $ML \times ML$  matrix  $\tilde{\mathbf{E}}(x)$  be defined by the generating function expression

$$\sum_{i,j=0}^{LM-1} \tilde{\mathbf{E}}(x)_{ij} r^i t^j := (\Delta_+(r)rV_-(t) - \Delta_-(t)tV_+(r)) \left( \frac{1}{r-t} \right) - \frac{1}{\gamma x} r t \frac{S(r) - S(t)}{r-t}, \quad (4.9)$$

and define a further pair of  $LM \times LM$  matrices  $\mathbf{E}(x)$  and  $\mathbf{E}'(x)$  by

$$\mathbf{E}(x) := (\mathbf{A}^T)^{-1} \tilde{\mathbf{E}}^T \quad (4.10)$$

$$\mathbf{E}'(x) := \mathbf{A}^{-1} \tilde{\mathbf{E}} \quad (4.11)$$

where  $\mathbf{A}$  is the  $LM \times LM$  matrix entering in the Christoffel-Darboux relation in Theorem 2.9. The matrices  $\mathbf{E}(x)$  and  $\mathbf{E}'(x)$  thus satisfy the following duality relation:

$$\mathbf{A}\mathbf{E}(x) - \mathbf{E}'(x)^T \mathbf{A} = 0. \quad (4.12)$$

The *folded* version of the linear relations (4.4) and (4.6) is then [3]:

**Theorem 4.2.** *The following finite dimensional differential systems follow from (4.4) and (4.6)*

$$\beta D\vec{\Psi}^+ = \mathbf{E}(x)\vec{\Psi}^+, \quad (4.13)$$

$$-\beta D\vec{\Psi}^- = \mathbf{E}'(x)\vec{\Psi}^-. \quad (4.14)$$

### 4.3 Adjoint differential system

**Definition 4.1.** Let  $\mathbf{M}(x)$  be the rank-1,  $LM \times LM$  matrix defined by

$$\mathbf{M}(x) := \vec{\Psi}^-(x) \vec{\Psi}^+(x)^T \mathbf{A}. \quad (4.15)$$

with entries

$$\mathbf{M}(x)_{ij} = \Psi_i^-(x) \sum_{k=0}^{LM-1} \Psi_k^+(x) \mathbf{A}_{k,j} \quad (4.16)$$

viewed as are elements of  $\mathbb{K}[x, s, \beta, \beta^{-1}][[\gamma]]$ .

The matrix  $\mathbf{M}(x)$  has the following properties:

**Proposition 4.3.** *The entries of  $\mathbf{M}(x)$  are elements of  $\mathbb{K}[x, s, \beta][[\gamma]]$ , i.e. they contain no negative power of  $\beta$ . Moreover,  $\mathbf{M}(x)$  is a rank 1 projector:*

$$\mathbf{M}(x)^2 = \mathbf{M}(x) \quad , \quad \text{Tr } \mathbf{M}(x) = 1. \quad (4.17)$$

and satisfies the adjoint differential system

$$\beta x \frac{d}{dx} \mathbf{M}(x) = [\mathbf{E}(x), \mathbf{M}(x)]. \quad (4.18)$$

### 4.4 $\beta$ -expansion of $M$

Let  $\mathbf{F}$  be the elementary  $LM \times LM$  matrix

$$\mathbf{F} := \text{diag}(1, 0, \dots, 0). \quad (4.19)$$

We then have [3]:

**Theorem 4.4.**

(I) *There exists a unique  $LM \times LM$  matrix  $\mathbf{M}^{\text{WKB}}(z)$  formal series expansion with the following properties:  $\mathbf{M}^{\text{WKB}}(z)$  is a rank 1 projector, its entries belong to  $\mathbb{K}[\mathbf{s}][[z, \beta]]$ , and  $\mathbf{M}^{\text{WKB}}(z)$  is a solution of the ODE*

$$\beta \frac{zG(S(z))}{\sigma(z)} \frac{d}{dz} \mathbf{M}^{\text{WKB}}(z) = [\mathbf{E}(X(z)), \mathbf{M}^{\text{WKB}}(z)] \quad (4.20)$$

with leading order:

$$\mathbf{M}^{\text{WKB}}(z) = \mathbf{V}(z) \mathbf{F} \mathbf{V}(z)^{-1} + O(\beta). \quad (4.21)$$

Moreover, for each  $k \geq 0$ , all entries of the matrix  $\mathbf{M}^{(k)}(z) = [\beta^k] \mathbf{M}^{\text{WKB}}(z)$  are rational functions of  $z$ , with poles only at the ramification points (elements of  $\mathcal{A}$ ) or at  $z = \infty$ .

(II) *For each  $k \geq 0$ , the following equality holds as formal power series of  $z$ :*

$$\mathbf{M}^{(k)}(z) = [\beta^k] \mathbf{M}(X(z)). \quad (4.22)$$

## 4.5 The current correlators $\tilde{W}_n$

It follows from the determinantal formula (2.8) and the Christoffel-Darboux Theorem 2.9 that [3]:

**Proposition 4.5.**

$$\tilde{W}_1(x) = \frac{1}{\beta x} \text{Tr} \mathbf{M}(x) \mathbf{E}(x), \quad (4.23)$$

$$\tilde{W}_2(x_1, x_2) = \frac{\text{Tr} \mathbf{M}(x_1) \mathbf{M}(x_2)}{(x_1 - x_2)^2} - \frac{1}{(x_1 - x_2)^2}, \quad (4.24)$$

and for  $n \geq 3$

$$\tilde{W}_n(x_1, \dots, x_n) = \sum_{\sigma \in \mathfrak{S}_n^{1\text{-cycle}}} (-1)^\sigma \frac{\text{Tr} \left( \prod_i \mathbf{M}(x_{\sigma(i)}) \right)}{\prod_i (x_{\sigma(i)} - x_{\sigma(i+1)})}, \quad (4.25)$$

It also follows that

**Proposition 4.6.** *For  $n \geq 0$ , let  $z_1, z_2, \dots, z_n$  be  $n$  generic distinct points on the Riemann sphere, the generating functions  $\tilde{W}_{g,n}$  evaluated at  $x_i = X(z_i)$ , are rational fractions of  $z_1, \dots, z_n$ :*

$$\sum_g \beta^{2g-2+n} \tilde{W}_{g,n}(X(z_1), \dots, X(z_n)) = \sum_{\sigma \in \mathfrak{S}_n^{1\text{-cycle}}} (-1)^\sigma \frac{\text{Tr} \left( \prod_i \mathbf{M}^{\text{WKB}}(z_{\sigma(i)}) \right)}{\prod_i (X(z_{\sigma(i)}) - X(z_{\sigma(i+1)}))}, \quad (4.26)$$

## 5 Fermionic representations

In the following, we use the standard free fermion representation, in which the creation and annihilation operators  $(\psi_i, \psi_i^*)_{i \in \mathbf{Z}}$  satisfy the anticommutation relations

$$[\psi_i, \psi_j]_+ = 0, \quad [\psi_i^*, \psi_j^*]_+ = 0, \quad [\psi_i, \psi_j^*]_+ = \delta_{ij}, \quad (5.1)$$

and the vacuum state  $|0\rangle$  is annihilated by the positive index annihilation operators and the negative creation operators

$$\psi_{-i}|0\rangle = 0, \quad i = 1, \dots, \infty, \quad \psi_i^*|0\rangle = 0, \quad i = 0, \dots, \infty. \quad (5.2)$$

The Fermi field and its adjoint are denoted

$$\psi(z) = \sum_{i=-\infty}^{\infty} \psi_i z^i, \quad \psi^*(z) = \sum_{i=-\infty}^{\infty} \psi_i^* z^{-i-1}, \quad (5.3)$$

and the current operator components are denoted

$$J_i := \sum_{j=-\infty}^{\infty} : \psi_j \psi_{i+j}^* : \quad i \in \mathbf{Z}. \quad (5.4)$$

The abelian groups  $\Gamma_{\pm}$  of *shift flows*, are defined as

$$\Gamma_+ := \{\hat{\gamma}_+(\mathbf{t})\}, \quad \Gamma_- := \{\hat{\gamma}_-(\mathbf{s})\} \quad (5.5)$$

where

$$\hat{\gamma}_+(\mathbf{t}) := e^{\sum_{i=1}^{\infty} t_i J_i}, \quad \hat{\gamma}_-(\mathbf{s}) := e^{\sum_{i=1}^{\infty} s_i J_{-i}}. \quad (5.6)$$

Another infinite abelian group that enters in the fermionic representation of our  $\tau$ -functions is the group  $\mathbf{C}$  of (generalized) convolution flows [30], consisting of diagonal elements

$$\mathbf{C} := \{\hat{C}_{\rho} := e^{\sum_{i=-\infty}^{\infty} T_i \psi_i \psi_i^*}\}, \quad (5.7)$$

where

$$\rho_i = e^{T_i}, \quad (5.8)$$

may be viewed as the Fourier coefficients of a function (or distribution) on the unit circle  $\{z : |z| = 1\}$ ,

$$\rho(z) = \sum_{i=-\infty}^{\infty} \rho_i z^{-i-1}. \quad (5.9)$$

The next three subsections 5.1, 5.2 and 5.3 give fermionic representations of all the quantities appearing above: the  $\tau$ -function, the Baker function, the adapted bases, the pair correlators, the multicurrent correlator. For proofs and further details of the relations between them, the reader should consult the companion paper [2].

## 5.1 The $\tau$ -function, Baker function and adapted bases

The  $\tau$ -function (2.1) has the following representation as a fermionic vacuum state expectation value (VEV)

$$\tau^{(G,\beta,\gamma)}(\mathbf{t}, \mathbf{s}) = \langle 0 | \hat{\gamma}_+(\mathbf{t}) \hat{C}_{\rho} \hat{\gamma}_-(\mathbf{s}) | 0 \rangle, \quad (5.10)$$

where the coefficients  $\rho_i$  are given by (2.5), (2.6).

Viewing this as a parametric family of KP  $\tau$ -functions (where  $\mathbf{s} = (s_1, s_2, \dots)$  are just viewed as additional parameters), the corresponding Baker function and its dual have the fermionic representation [2]

$$\Psi_{(G,\beta,\gamma)}^-(\zeta, \mathbf{t}, \mathbf{s}) = \frac{\langle 0 | \psi_0^* \hat{\gamma}_+(\mathbf{t}) \psi(\zeta) \hat{C}_{\rho} \hat{\gamma}_-(\mathbf{s}) | 0 \rangle}{\tau^{(G,\beta,\gamma)}(\mathbf{t}, \mathbf{s})}, \quad (5.11)$$

$$\Psi_{(G,\beta,\gamma)}^+(\zeta, \mathbf{t}, \mathbf{s}) = \frac{\langle 0 | \psi_{-1} \hat{\gamma}_+(\mathbf{t}) \psi^*(\zeta) \hat{C}_{\rho} \hat{\gamma}_-(\mathbf{s}) | 0 \rangle}{\tau^{(G,\beta,\gamma)}(\mathbf{t}, \mathbf{s})}. \quad (5.12)$$

The adapted basis defined by (2.17) has the fermionic representation [2]

$$\Psi_k^+(x) := \begin{cases} \gamma \langle 0 | \psi^*(1/x) \hat{C}_{\rho} \hat{\gamma}_-(\beta^{-1}\mathbf{s}) \psi_{k-1} | 0 \rangle & \text{if } k \geq 1, \\ \gamma \langle 0 | \psi_{k-1} \hat{\gamma}_-^{-1}(\beta^{-1}\mathbf{s}) \hat{C}_{\rho}^{-1} \psi^*(1/x) \hat{C}_{\rho} \hat{\gamma}_-(\beta^{-1}\mathbf{s}) | 0 \rangle & \text{if } k \leq 0, \end{cases} \quad (5.13)$$

and the dual basis defined by (2.18) is given [2] by

$$\Psi_k^-(x) := \begin{cases} \langle 0 | \psi(1/x) \hat{C}_\rho \hat{\gamma}_-(\beta^{-1}\mathbf{s}) \psi_{-k}^* | 0 \rangle & \text{if } k \geq 1 \\ \langle 0 | \psi_{-k}^* \hat{\gamma}_-^{-1}(\beta^{-1}\mathbf{s}) \hat{C}_\rho^{-1} \psi(1/x) \hat{C}_\rho \hat{\gamma}_-(\beta^{-1}\mathbf{s}) | 0 \rangle & \text{if } k \leq 0. \end{cases} \quad (5.14)$$

## 5.2 The $n$ -pair correlator $K_n$

The  $n$ -pair correlator (2.33) has the following fermionic representation [2]

$$K_n(x_1, \dots, x_n; x'_1, \dots, x'_n) = \frac{1}{\prod_{i=1}^n x_i x'_i} \langle 0 | \prod_{i=1}^n \psi(1/x'_i) \psi^*(1/x_i) \hat{C}_\rho \hat{\gamma}_-(\beta^{-1}\mathbf{s}) | 0 \rangle. \quad (5.15)$$

## 5.3 The multicurrent correlator $W_n$

It follows from the fermionic representation (5.10) of the  $\tau$ -function  $\tau^{(G, \beta, \gamma)}(\mathbf{t}, \mathbf{s})$  and the definition (2.36) of the multicurrent correlator  $W_n(x_1, \dots, x_n)$  that this may be expressed as the following fermionic vacuum expectation value [2]

$$W_n(x_1, \dots, x_n) = \frac{1}{\prod_{i=1}^n x_i} \langle 0 | \prod_{i=1}^n J_+(x_i) \hat{C}_\rho \hat{\gamma}_-(\beta^{-1}\mathbf{s}) | 0 \rangle, \quad (5.16)$$

where

$$J_+(x) := \sum_{i=1}^{\infty} x^i J_i. \quad (5.17)$$

# 6 Topological recursion

## 6.1 Definitions of $\tilde{\omega}_{g,n}$ and $\mathcal{W}_{g,n}$

The key invariants computed by topological recursion in this case are the forms  $\tilde{\omega}_{g,n}$  defined in terms of the correlators  $\tilde{W}_{g,n}$  as follows:

**Definition 6.1.** Let

$$\begin{aligned} \tilde{\omega}_{g,n}(z_1, \dots, z_n) := & \tilde{W}_{g,n}(X(z_1), \dots, X(z_n)) X'(z_1) \dots X'(z_n) dz_1 \dots dz_n \\ & + \delta_{g,0} \delta_{n,2} \frac{X'(z_1) X'(z_2) dz_1 dz_2}{(X(z_1) - X(z_2))^2}. \end{aligned} \quad (6.1)$$

It follows [3] that  $\tilde{\omega}_{g,n}$  for the stable cases are rational functions of all the  $z_i$ 's, with poles only at elements of  $\mathcal{A}$  (ramification points of the spectral curve) or at infinity. In fact we have the stronger property (see [3] for a detailed proof):

**Proposition 6.1.** *When  $2g-2+n > 0$ ,  $\tilde{\omega}_{g,n}(z_1, \dots, z_n)$  has poles only at ramification points, and no poles at  $z_i = \infty$ .*

In order to state our topological recursion in a more compact manner, it is convenient to introduce the following auxiliary quantity.

**Definition 6.2.**

$$\begin{aligned} \mathcal{W}_{g,n}(z, z', z_2, \dots, z_n) &:= \tilde{\omega}_{g-1, n+1}(z, z', z_2, \dots, z_n) \\ &+ \sum'_{g_1+g_2=g, I_1 \uplus I_2 = \{z_2, \dots, z_n\}} \tilde{\omega}_{g_1, 1+|I_1|}(z, I_1) \tilde{\omega}_{g_2, 1+|I_2|}(z', I_2), \end{aligned} \quad (6.2)$$

where  $\sum'$  means that we exclude the two terms  $(g_1, I_1) = (0, \emptyset)$  and  $(g_2, I_2) = (0, \emptyset)$ .

## 6.2 Topological recursion for the $\tilde{\omega}_{g,n}$ 's

As shown in [3], the  $\tilde{W}_n$ 's satisfy a set of equations called “loop equations” [10, 12]. The unique formal  $\beta$ -power series solution of the loop equations, having poles only at the ramification points, is the one given by the topological recursion relations (Theorem 4.5. of [20]; see [13] for another proof).

**Theorem 6.2.** *The  $\tilde{\omega}_{g,n}$ 's satisfy the topological recursion relations: If all ramification points  $a \in \mathcal{A}$  are simple, with local Galois involution  $z \mapsto \sigma_a(z)$ , we have for  $(g, n) \neq (0, 1), (0, 2)$ :*

$$\tilde{\omega}_{g,n}(z_1, \dots, z_n) = - \sum_{a \in \mathcal{A}} \text{Res}_{z \rightarrow a} \left[ \frac{dz_1}{z - z_1} - \frac{dz_1}{\sigma_a(z) - z_1} \right] \frac{\mathcal{W}_{g,n}(z, \sigma_a(z), z_2, \dots, z_n)}{2(Y(z) - Y(\sigma_a(z))) dX(z)}. \quad (6.3)$$

*If some ramification points have higher order, then the  $\tilde{\omega}_{g,n}$ s satisfy the higher order version of the topological recursion relation defined in [15].*

*For  $(g, n) = (0, 2)$  we have*

$$\tilde{\omega}_{0,2}(z_1, z_2) = \frac{dz_1 dz_2}{(z_1 - z_2)^2}. \quad (6.4)$$

A detailed proof of this main result is provided in ref. [3].

## 7 Some consequences of topological recursion

The first consequence of Theorem 6.2 is that  $\tilde{\omega}_{g,n}(z_1, \dots, z_n)$  can be computed explicitly for each  $g$  and  $n$ , which in turn gives an explicit rational parametrization of the function  $\tilde{W}_{g,n}(x_1, \dots, x_n)$ .

Theorem 6.2 has many other deep implications, of which a short glimpse is the following.

- **Structure properties:** When  $2g - 2 + n > 0$ , the  $\tilde{F}_{g,n}^{(G,\gamma)}(x_1, \dots, x_n)$  are not only formal series in  $\gamma$ , but are in fact algebraic functions of  $\gamma$  and the  $x_i$ 's, and of all parameters  $\mathbf{c} = (c_1, c_2, \dots)$  defining  $G$ , as well as the parameters  $\mathbf{s} = \{s_1, s_2, \dots\}$ . The radius of convergence in  $\gamma$  is given by the ramification point nearest the origin:

$$|\gamma x_i| < R \quad , \quad R = \min_j \{|X(a_j)| \mid X'(a_j) = 0\}. \quad (7.1)$$

The  $\tilde{F}_{g,n}(X(z_1), \dots, X(z_n))$ , with  $2g - 2 + n > 0$ , are rational functions of the arguments  $z_i$ , with poles only at ramification points, of order  $\leq 3(2g - 2 + n)$ . For  $(g, n) = (0, 1)$  explicit expressions are known involving logarithms, with the same radius of convergence, (See [20].)

- The first few  $\tilde{F}_{g,n}$  are easy to compute using topological recursion. For instance

$$\begin{aligned} \tilde{F}_{0,3} &= \frac{X(z_1)}{X'(z_1)S'(z_1)(z_1 - z_2)(z_1 - z_3)} + \text{sym}(z_1 \leftrightarrow z_2, z_3) \\ &\quad - \sum_{b, S'(b)=0} \frac{X(b)}{X'(b)S''(b)} \frac{1}{(z_1 - b)(z_2 - b)(z_3 - b)}. \end{aligned} \quad (7.2)$$

(For the case  $L = 1$  and  $M$  arbitrary, (7.2), was conjectured by John Irving (private communication to GC) in 2013. He also obtained the expression of  $\tilde{W}_{0,2}$  via elementary combinatorial manipulations, and conjectured that all the  $\tilde{W}_{0,n}$  are rational functions of the parameters  $z_i$ . We refer the reader to a paper in preparation [32] for these conjectures (which are here proved, and in [3], in greater generality) and for the combinatorial approach to the computation of  $\tilde{W}_{0,2}$ . Other combinatorial approaches to the computation of generating functions for constellations may be found in [17] and the thesis [23], which contains in particular an analog of the cut-and-join equation [23, Theorem 4.2].)

- Topological recursion implies a form-cycle duality formula for all deformations, for example

$$\frac{\partial}{\partial s_k} \tilde{F}_{g,n}(x_1, \dots, x_n) = \text{Res}_{z' \rightarrow \infty} \tilde{F}_{g,n+1}(x_1, \dots, x_n, X(z')) k z'^{k-1} dz', \quad (7.3)$$

$$\begin{aligned} \frac{\partial}{\partial g_k} \tilde{F}_{g,n}(x_1, \dots, x_n) &= \text{Res}_{z' \rightarrow \infty} \tilde{F}_{g,n+1}(x_1, \dots, x_n, X(z')) d \left( \frac{S'(z')}{G(z')} \right), \\ \frac{\partial}{\partial c_i} \tilde{F}_{g,n}(x_1, \dots, x_n) &= \text{Res}_{z' \rightarrow \infty} \tilde{F}_{g,n+1}(x_1, \dots, x_n, X(z')) d \left( \frac{S'(z')z'}{1 + c_i z'} \right). \end{aligned} \quad (7.4)$$

- It has many other deep consequences, which are well beyond this short summary. For example, topological recursion implies some ELSV-like formulas as in [22]. These will be developed in the longer version of this work [3].

*Acknowledgements.* The work of A. Alexandrov was supported by IBS-R003-D1, by RFBR grants 14-01-00547 and 15-52-50041YaF, and by the European Research Council (QUASIFT grant agreement 677368). G.Chapuy acknowledges support from the European Research Council, grant ERC-2016-STG 716083 “CombiTop”, from the Agence Nationale de la Recherche, grant ANR 12-JS02-001-01 “Cartaplus” and from the City of Paris, grant “Émergences 2013, Combinatoire à Paris”. B. Eynard was supported by the ERC Starting Grant no. 335739 “Quantum fields and knot homologies” funded by the European Research Council under the European Union’s Seventh Framework Programme, and also partly supported by the ANR grant Quantact : ANR-16-CE40-0017. The work of J. Harnad was partially supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) and the Fonds de recherche du Québec, Nature et technologies (FRQNT).

Much of this work was completed while G.C. was affiliated to the Centre de Recherches Mathématiques de Montréal as part of the CNRS UMI program; he wishes to thank the CRM for the very good working conditions provided. B.E. also wishes to thank the Centre de recherches mathématiques, Montréal, for the Aisenstadt Chair grant, and the FQRNT grant from the Québec government that partially supported this joint project. A.A. and J.H. wish to thank the Institut des Hautes Études Scientifiques for their kind hospitality for extended periods in 2017 - 2018, when much of this work was completed. The authors would also like to thank the organizers of the “Moduli spaces, integrable systems, and topological recursions” thematic semester program (June 2015 - Jan. 2016) at the CRM in Montréal, where part of this work was realized and the organizers of the January - March, 2017 thematic semester “Combinatorics and interactions” at the Institut Henri Poincaré, where they were participants during the completion of this work.

G.C. also thanks John Irving for sharing his results and Wenjie Fang, whose PhD thesis contains several insights that have helped to inspire this work. B.E. and A.A. thank Piotr Sułkowski and S. Shadrin for useful discussions.

Work supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), the Fonds de recherche du Québec - Nature et technologies (FRQNT), the European Research Council (ERC), and the Agence Nationale de la Recherche (ANR).

## References

- [1] A. Alexandrov, “Enumerative Geometry, Tau-Functions and Heisenberg-Virasoro Algebra,” *Commun. Math. Phys.* **338**, 195-249 (2015).
- [2] A. Alexandrov, G. Chapuy, B. Eynard and J. Harnad, “Fermionic approach to weighted Hurwitz numbers and topological recursion,” *Commun. Math. Phys.* **360** no.2, 777 (2018).
- [3] A. Alexandrov, G. Chapuy, B. Eynard and J. Harnad, “Weighted Hurwitz numbers and topological recursion”, [arXiv:1806.09738](https://arxiv.org/abs/1806.09738).
- [4] A. Alexandrov, “Matrix Models for Random Partitions,” *Nucl. Phys. B* **851** 620 (2011).
- [5] A. Alexandrov, D. Lewanski and S. Shadrin, “Ramifications of Hurwitz theory, KP integrability and quantum curves,” *JHEP* **1605**, 124 (2016).
- [6] A. Alexandrov, A. Mironov, A. Morozov and S. Natanzon, “Integrability of Hurwitz Partition Functions. I. Summary,” *J. Phys. A* **45**, 045209 (2012).

- [7] A. Alexandrov, A. Mironov, A. Morozov and S. Natanzon, “On KP-integrable Hurwitz functions,” *JHEP* **1411**, 080 (2014).
- [8] J. Ambjørn and L. Chekhov, “The matrix model for dessins d’enfants”, *Ann. Inst. Henri Poincaré, Comb. Phys. Interact.* **1**, 337-361 (2014).
- [9] J. Ambjørn and L. Chekhov, “A matrix model for hypergeometric Hurwitz numbers”, *Theor. Math. Phys.* **181**, 1486-1498 (2014).
- [10] R. Belliard, B. Eynard, O. Marchal, “Loop equations from differential systems”, “Integrable differential systems of topological type and reconstruction by the topological recursion”, *Ann. Henri Poincaré* **18** no. 10, 3193-3248 (2017).
- [11] E.A. Bender and E. Canfield, “The number of rooted maps on an orientable surface”, *J. Combin. Theory Ser. B* **53**, 293-299 (1991).
- [12] M. Bergere, G. Borot and B. Eynard, “Rational differential systems, loop equations, and application to the q-th reduction of KP”, *Ann. Henri Poincaré* **16**, 2713-2782 (2015).
- [13] G. Borot, B. Eynard, N. Orantin, “Abstract loop equations, topological recursion, and applications”, *Commun. Num. Theor. Phys.* **09** 51-187 (2015).
- [14] G. Borot, B. Eynard, M. Mulase and B. Safnuk, “A matrix model for Hurwitz numbers and topological recursion”, *J. Geom. Phys.* **61**, 522–540 (2011).
- [15] V. Bouchard and B. Eynard, “Think globally, Compute Locally”, *JHEP* **143**, 1302 (2013).
- [16] M. Bousquet-Mélou and G. Schaeffer, “Enumeration of planar constellations”. *Adv. in Appl. Math.* **24:4**, 337-368 (2000).
- [17] G. Chapuy, “Asymptotic enumeration of constellations and related families of maps on orientable surfaces”, *Combin. Probab. Comput.* **18**, 477-566 (2009).
- [18] N. Do, A. Dyer, and D.V. Mathews, “Topological recursion and a quantum curve for monotone Hurwitz numbers”, *J. Geom. Phys.* **120**, 19-36 (2017)
- [19] N. Do, O. Leigh, and P. Norbury, “Orbifold Hurwitz numbers and Eynard-Orantin invariants”, *Math. Res. Lett.* **23**, 1281-1327 (2016).
- [20] B. Eynard and N. Orantin, “Invariants of algebraic curves and topological expansion”, *Commun. Number Theor. Phys.* **1**, 347-452 (2007).
- [21] B. Eynard and N. Orantin, “Topological recursion in enumerative geometry and random matrices”, *J. Phys. A* **42**, 293001 (2009).
- [22] B. Eynard “Invariants of spectral curves and intersection theory of moduli spaces of complex curves”, *Commun. Num. Theor. Phys.* **8**, 541-588 (2014).
- [23] W. Fang, “Enumerative and bijective aspects of combinatorial maps: generalization, unification and application”, PhD thesis, Université Paris Diderot (2016).

- [24] G. Frobenius, “Über die Charaktere der symmetrischen Gruppe”, *Sitzber. Akad. Wiss., Berlin*, 516-534 (1900). Gesammelte Abhandlung III, 148-166; *ibid.* “Über die Charakterische Einheiten der symmetrischen Gruppe”, *Sitzber. Akad. Wiss., Berlin*, 328-358 (1903). Gesammelte Abhandlung III, 244-274.
- [25] I. P. Goulden, M. Guay-Paquet and J. Novak, “Monotone Hurwitz numbers and the HCIZ Integral”, *Ann. Math. Blaise Pascal* **21**, 71-99 (2014).
- [26] M. Guay-Paquet and J. Harnad, “2D Toda  $\tau$ -functions as combinatorial generating functions”, *Lett. Math. Phys.* **105**, 827-852 (2015).
- [27] M. Guay-Paquet and J. Harnad, “Generating functions for weighted Hurwitz numbers”, *J. Math. Phys.* **58**, 083503 (2017).
- [28] J. Harnad, “Weighted Hurwitz numbers and hypergeometric  $\tau$ -functions: an overview”, *AMS Proceedings of Symposia in Pure Mathematics* **93**, 289-333 (2016).
- [29] J. Harnad, “Quantum Hurwitz numbers and Macdonald polynomials”, *J. Math. Phys.* **57**, 113505 (2016.)
- [30] J. Harnad and A. Yu. Orlov, “Hypergeometric  $\tau$ -functions, Hurwitz numbers and enumeration of paths”, *Commun. Math. Phys.* **338**, 267-284 (2015).
- [31] A. Hurwitz, “Über Riemann’sche Fläche mit gegebene Verzweigungspunkten”, *Math. Ann.* **39**, 1-61 (1891); *Mathematische Werke I*, 321-384; *ibid.* “Über die Anzahl der Riemann’sche Fläche mit gegebene Verzweigungspunkten”, *Math. Ann.* **55**, 53-66 (1902); *Mathematische Werke I*, 42-505.
- [32] J. Irving, “On the enumeration of  $m$ -constellations”, (in preparation; private communication).
- [33] S. Kharchev, A. Marshakov, A. Mironov and A. Morozov, “Generalized Kazakov-Migdal-Kontsevich model: Group theory aspects,” *Int. J. Mod. Phys. A* **10**, 2015 (1995).
- [34] M. Kazarian and P. Zograf, “Virasoro constraints and topological recursion for Grothendieck’s dessin counting”, *Lett. Math. Phys.* **105**, 1057-1084 (2015).
- [35] S. K. Lando and A.K. Zvonkin *Graphs on Surfaces and their Applications*, Encyclopaedia of Mathematical Sciences, Volume **141**, with appendix by D. Zagier, Springer, N.Y. (2004).
- [36] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Clarendon Press, Oxford, (1995).
- [37] A. Morozov and S. Shakirov, “Generation of Matrix Models by W-operators,” *JHEP* **0904** 064 (2009).
- [38] M. Mulase, S. Shadrin and L. Spitz, “The spectral curve and the Schrödinger equation of double Hurwitz numbers and higher spin structures,” *Commun. Num. Theor Phys.* **07**, 125 (2013).
- [39] M. Mulase and P. Sulkowski, “Spectral curves and the Schrödinger equations for the Eynard-Orantin recursion,” *Adv. Theor. Math. Phys.* **19**, 955 (2015).

- [40] Sergey M. Natanzon and Aleksandr Yu. Orlov, “BKP and projective Hurwitz numbers” *Lett. Math. Phys.* **107**,1065-1109 (2017).
- [41] A. Okounkov, “Toda equations for Hurwitz numbers”, *Math. Res. Lett.* **7**, 447–453 (2000).
- [42] A. Yu. Orlov, “Hypergeometric functions as infinite-soliton  $\tau$ -functions”, *Theor. Math. Phys.* **146**, 183-206 (2006).
- [43] A. Y. Orlov and D. M. Scherbin, “Fermionic representation for basic hypergeometric functions related to Schur polynomials,” [arXiv:nlin/0001001](https://arxiv.org/abs/nlin/0001001).
- [44] A. Yu. Orlov and D. M. Scherbin, “Hypergeometric solutions of soliton equations”, *Theor. Math. Phys.* **128**, 906-926 (2001).
- [45] R. Pandharipande, “The Toda Equations and the Gromov-Witten Theory of the Riemann Sphere”, *Lett. Math. Phys.* **53**, 59-74 (2000).
- [46] M. Sato, “Soliton equations as dynamical systems on infinite dimensional Grassmann manifolds”, RIMS, Kyoto Univ. *Kokyuroku* **439**, 30-46 (1981). In “Nonlinear Partial Differential Equations in Applied Science”; Proceedings of The U.S.-Japan Seminar, Tokyo, 1982, (ed. Hiroshi Fujita, Peter D. Lax, Gilbert Strang).
- [47] M. Sato and Y. Sato. “Soliton equations as dynamical systems on infinite dimensional Grassmann manifold”, in: *Nonlinear PDE in Applied Science, Proc. U. S.-Japan Seminar*, Tokyo 1982, Kinokuniya, Tokyo, pp. 259-271 (1983).
- [48] G. Schaeffer, “Handbook of Enumerative Combinatorics”, Chapter 5 , CRC Press *Discrete Mathematics and its Applications*(ed. Miklos Bonà) (2015).
- [49] G. Segal and G. Wilson, “Loop groups and equations of KdV type ”, *Pub. Math. IHÉS* **61**, 5-65 (1985).
- [50] T. Takebe, “Representation theoretical meaning of the initial value problem for the Toda lattice hierarchy I”, *Lett. Math. Phys.* **21**, 77–84,(1991).
- [51] W. T. Tutte, “A census of planar maps”, *Canad. J. Math.* **15** 249–271 (1963).
- [52] K. Ueno and K. Takasaki, “Toda Lattice Hierarchy”, in: *Group Representation and Systems of Differential Equations, Adv. Stud. in Pure Math.* **4**, 1-95 (1984). Mathematical Society of Japan, Tokyo, Japan (ed. K. Okamoto).
- [53] P. Zograf, “Enumeration of Gröthendieck’s dessins and KP hierarchy”, *Int. Math. Res. Notices* **24**, 13533-13544 (2015).