

A GENERALIZATION OF THE NAKAYAMA FUNCTOR

SONDRE KVAMME

ABSTRACT. In this paper we introduce a generalization of the Nakayama functor for finite-dimensional algebras. This is obtained by abstracting its interaction with the forgetful functor to vector spaces. In particular, we characterize the Nakayama functor in terms of an ambidextrous adjunction of monads and comonads. In the second part we develop a theory of Gorenstein homological algebra for such Nakayama functor. We obtain analogues of several classical results for Iwanaga-Gorenstein algebras. One of our main examples is the module category $\Lambda\text{-Mod}$ of a k -algebra Λ , where k is a commutative ring and Λ is finitely generated projective as a k -module.

CONTENTS

1. Introduction	2
1.1. Introduction	2
1.2. Conventions	6
2. Background	6
2.1. Transformation of adjoints	6
2.2. Monads and comonads	8
2.3. Properties of subcategories	11
3. Generalization of the Nakayama functor	12
3.1. Derived functors without enough projectives or injectives	12
3.2. Definition of Nakayama functors	14
3.3. Properties of Nakayama functors	19
4. Gorenstein homological algebra for Nakayama functors relative to endofunctors	24
4.1. Gorenstein P -projective and I -injective objects	25
4.2. Iwanaga-Gorenstein functors	27
5. Functor categories	32
5.1. Reminder on the Hom and tensor functor	33

Date: July 22, 2019.

2010 Mathematics Subject Classification. 18E10, 16E65, 16D90;

Key words and phrases. Abelian category; Gorenstein homological algebra; Gorenstein ring; homological algebra; monad .

This is part of the authors PhD thesis. The author thanks Gustavo Jasso, Julian Külshammer, Rosanna Laking, and Jan Schröer for helpful discussions and comments and on a previous version of this paper. He would also like to thank the referee for helpful comments and suggestions. The work was made possible by the funding provided by the *Bonn International Graduate School in Mathematics*.

5.2. Properties of Hom-finite locally bounded categories	34
5.3. Nakayama functor for $i_! \dashv i^*$	35
References	37

1. INTRODUCTION

1.1. Introduction. Let Λ be a finite-dimensional k -algebra, and let $D(\Lambda) = \text{Hom}_k(\Lambda, k)$ be the k -dual. Consider the Nakayama functor

$$\nu = D(\Lambda) \otimes_{\Lambda} - : \Lambda\text{-mod} \rightarrow \Lambda\text{-mod}$$

It is used to define the Auslander-Reiten translation, which is a fundamental tool for doing Auslander-Reiten theory. It therefore plays an important role in determining the structure of $\Lambda\text{-mod}$. On the other hand, we also have the well developed theory of Gorenstein homological algebra, see for example [1, 2, 7, 12, 13, 14, 15, 17, 19]. This theory has been studied in the context of finite-dimensional algebras, see for example [3, 5, 8, 23], and in this case many of the statements are linked to the Nakayama functor. For example, a finite-dimensional algebra Λ is Iwanaga-Gorenstein if and only if

$$\text{proj. dim } D(\Lambda)_{\Lambda} < \infty \quad \text{and} \quad \text{proj. dim } {}_{\Lambda} D(\Lambda) < \infty.$$

and in this case it follows from a result by Zaks [26] that

$$\text{proj. dim } D(\Lambda)_{\Lambda} = \text{proj. dim } {}_{\Lambda} D(\Lambda).$$

where this number is the Gorenstein dimension of Λ . These are statements about the vanishing of the left and right derived functor of the Nakayama functor and its right adjoint, respectively.

The aim of this paper is twofold. We first give a generalization of the Nakayama functor. This is obtained by observing that it interacts nicely with the adjoint pair $f_! \dashv f^*$, where

$$f^* : \text{res}_k^{\Lambda} : \Lambda\text{-mod} \rightarrow k\text{-mod}$$

is the restriction functor, and $f_! := \Lambda \otimes_k - : k\text{-mod} \rightarrow \Lambda\text{-mod}$ is its left adjoint. More precisely, we make the following definition.

Definition 1.1.1 (Definition 3.2.1). Let \mathcal{D} be a preadditive category and \mathcal{A} an abelian category, and let $f^* : \mathcal{A} \rightarrow \mathcal{D}$ be a faithful functor with left adjoint $f_! : \mathcal{D} \rightarrow \mathcal{A}$. A **Nakayama functor** relative to $f_! \dashv f^*$ is a functor $\nu : \mathcal{A} \rightarrow \mathcal{A}$ with a right adjoint ν^- satisfying:

- (i) $\nu \circ f_!$ is right adjoint to f^* ;
- (ii) The unit of $\nu \dashv \nu^-$ induces an isomorphism $f_! \xrightarrow{\cong} \nu^- \circ \nu \circ f_!$ when precomposed with $f_!$.

The Nakayama functor is unique up to unique equivalence, see Theorem 3.3.4. Furthermore, a choice of a Nakayama functor is the same as a choice of an isomorphism of certain categories, or a choice of an isomorphism of certain monads, see Proposition 3.3.1 and Theorem 3.3.3.

We have the following example.

Example 1.1.2 (Example 3.2.5). Let k be a commutative ring, and let Λ be a k -algebra which is finitely generated projective as a k -module. Orders over complete regular local rings [18] are examples of such algebras. Consider the restriction functor

$$f^* := \text{res}_k^\Lambda : \Lambda\text{-Mod} \rightarrow k\text{-Mod}$$

with left adjoint $f_! := \Lambda \otimes_k - : k\text{-Mod} \rightarrow \Lambda\text{-Mod}$. Then the functor

$$\nu_{\Lambda\text{-Mod}} := D(\Lambda) \otimes_\Lambda - : \Lambda\text{-Mod} \rightarrow \Lambda\text{-Mod}$$

is a Nakayama functor relative to $f_! \dashv f^*$, where $D := \text{Hom}_k(-, k)$ is the algebraic dual.

This gives the classical Nakayama functor for a finite-dimensional algebra. For more examples see the end of Subsection 3.2. In fact, for any commutative ring k , any k -linear abelian category \mathcal{B} , and any small, k -linear, locally bounded and Hom-finite category \mathcal{C} , the functor category $\mathcal{A} = \mathcal{B}^{\mathcal{C}}$ has a Nakayama functor relative to an adjoint pair, see Theorem 5.3.3. Such categories \mathcal{C} are also studied in [10].

The notion of a Nakayama functor relative to an adjoint pair can also be phrased in terms of monads and comonads. In fact, if $f_! \dashv f^*$ has a Nakayama functor, then f^* is both faithful and exact. Hence, if \mathcal{D} is abelian, then by Beck's monadicity theorem (see Theorem 4.4.4 in [6] or Theorem 2.2.8 below) we have an equivalence $\mathcal{D}^\mathbb{T} \cong \mathcal{A}$ commuting up to isomorphism with f^* and the forgetful functor $U^\mathbb{T} : \mathcal{D}^\mathbb{T} \rightarrow \mathcal{D}$. Here $\mathcal{D}^\mathbb{T}$ is the Eilenberg-Moore category of the induced monad \mathbb{T} on \mathcal{D} coming from the adjunction, see Example 2.2.3 and Definition 2.2.5. We determine necessary and sufficient conditions on \mathbb{T} for the existence of a Nakayama functor.

Theorem 1.1.3 (Theorem 3.3.5). *Let \mathbb{T} be a monad on an abelian category \mathcal{B} , let $\mathcal{B}^\mathbb{T}$ be the Eilenberg-Moore category of \mathbb{T} , and let $U^\mathbb{T} : \mathcal{B}^\mathbb{T} \rightarrow \mathcal{B}$ be the forgetful functor with left adjoint $F^\mathbb{T} : \mathcal{B} \rightarrow \mathcal{B}^\mathbb{T}$. Then the adjoint pair $F^\mathbb{T} \dashv U^\mathbb{T}$ has a Nakayama functor if and only if there exists a comonad \mathbb{S} and an ambidextrous adjunction $\mathbb{S} \dashv \mathbb{T} \dashv \mathbb{S}$.*

See Definition 2.2.4 for the notion of adjunction between monads and comonads. Note that it is not sufficient to just assume there exists an ambidextrous adjunction $\mathbb{S} \dashv \mathbb{T} \dashv \mathbb{S}$ where \mathbb{T} is the underlying functor of \mathbb{T} , since the induced comonad structures on \mathbb{S} coming from the adjunctions $\mathbb{S} \dashv \mathbb{T}$ and $\mathbb{T} \dashv \mathbb{S}$ might be different. It follows from Beck's monadicity theorem and Theorem 1.1.3 that if \mathcal{D} is abelian and $f^* : \mathcal{A} \rightarrow \mathcal{D}$ is faithful and exact, then $f_! \dashv f^*$ has a Nakayama functor if and only if there exists a comonad \mathbb{S} and an ambidextrous adjunction $\mathbb{S} \dashv \mathbb{T} \dashv \mathbb{S}$, where \mathbb{T} is the induced monad on \mathcal{D} coming from $f_! \dashv f^*$.

The second aim of this paper is to develop a theory of Gorenstein homological algebra for such Nakayama functor. In fact, our constructions and

results in this part only depends on the endofunctor $P := f_! \circ f^*: \mathcal{A} \rightarrow \mathcal{A}$, and not on the adjoint pair itself.

Definition 1.1.4 (Definition 3.2.3). Let \mathcal{A} be an abelian category, and let $P: \mathcal{A} \rightarrow \mathcal{A}$ be a generating functor. A **Nakayama functor** relative to P is a functor $\nu: \mathcal{A} \rightarrow \mathcal{A}$ with a right adjoint ν^- , and satisfying the following:

- (i) $\nu \circ P$ is right adjoint to P ;
- (ii) The unit $\lambda: 1_{\mathcal{A}} \rightarrow \nu^- \circ \nu$ induces an isomorphism on objects of the form $P(A)$ for $A \in \mathcal{A}$.

Here P is generating if the the image of P is a generating subcategory of \mathcal{A} , see Definition 2.3.1.

We define the analogue of Gorenstein projective and injective modules.

Definition 1.1.5 (Definition 4.1.1). Let $P: \mathcal{A} \rightarrow \mathcal{A}$ be a generating functor with a Nakayama functor ν relative to P , and let $I := \nu \circ P$ be the right adjoint of P .

- (i) An object $G \in \mathcal{A}$ is **Gorenstein P -projective** if there exists an exact sequence

$$Q_{\bullet} = \cdots \xrightarrow{f_2} Q_1 \xrightarrow{f_1} Q_0 \xrightarrow{f_0} Q_{-1} \xrightarrow{f_{-1}} \cdots$$

in \mathcal{A} , where $Q_i = P(A_i)$ for $A_i \in \mathcal{A}$, such that the complex $\nu(Q_{\bullet})$ is exact, and with $Z_0(Q_{\bullet}) = \text{Ker } f_0 = G$;

- (ii) An object $G \in \mathcal{A}$ is **Gorenstein I -injective** if there exists an exact sequence

$$J_{\bullet} = \cdots \xrightarrow{g_2} J_1 \xrightarrow{g_1} J_0 \xrightarrow{g_0} J_{-1} \xrightarrow{g_{-1}} \cdots$$

in \mathcal{A} , where $J_i = I(A_i)$ for $A_i \in \mathcal{A}$, such that the complex $\nu^-(J_{\bullet})$ is exact, and with $Z_0(J_{\bullet}) = \text{Ker } g_0 = G$.

For a finite-dimensional algebra Λ with $P = f^* f_!$, a Λ -module is Gorenstein P -projective or I -injective if and only if it is Gorenstein projective or injective, respectively. Let $\mathcal{GP}_P(\mathcal{A})$ and $\mathcal{GI}_I(\mathcal{A})$ denote the subcategories of Gorenstein P -projective and I -injective objects. In Proposition 4.1.5 we show that these are resolving and coresolving subcategories, respectively. Hence, we can define the Gorenstein P -projective dimension and I -injective dimension $\dim_{\mathcal{GP}_P(\mathcal{A})}(-)$ and $\dim_{\mathcal{GI}_I(\mathcal{A})}(-)$ by taking resolutions of Gorenstein P -projective objects and coresolutions of Gorenstein I -injective objects, as described in Subsection 2.3.

We define an analog of Iwanaga-Gorenstein algebras.

Definition 1.1.6 (Definition 4.2.1). Let $P: \mathcal{A} \rightarrow \mathcal{A}$ be a generating functor with a Nakayama functor ν relative to P . We say that P is **Iwanaga-Gorenstein** if there exists an $m \geq 0$ such that $L_i \nu(A) = 0$ and $R^i \nu^-(A) = 0$ for all $A \in \mathcal{A}$ and $i > m$.

Note that the left derived functor of ν and the right derived functor of ν^- exists without assuming enough projectives or injectives, see Lemma 3.2.4.

For a finite-dimensional algebra Λ , the functor P is Iwanaga-Gorenstein if and only if Λ is Iwanaga-Gorenstein. More generally, the functor $P = f_! \circ f^*$ in Example 1.1.2 is Iwanaga-Gorenstein if and only if

$$\text{proj. dim } {}_{\Lambda}D(\Lambda) < \infty \quad \text{and} \quad \text{proj. dim } D(\Lambda)_{\Lambda} < \infty.$$

We show that if P is Iwanaga-Gorenstein, then there exists a simple description of $\mathcal{GP}_P(\mathcal{A})$ and $\mathcal{GI}_I(\mathcal{A})$, see Theorem 4.2.2. Our main result in this part is the following, which is an analogue of Iwanaga's result on the injective dimension of Iwanaga-Gorenstein algebras, and a generalization of [8, Theorem 2.3.3].

Theorem 1.1.7 (Theorem 4.2.6). *The following are equivalent:*

- (a) P is Iwanaga-Gorenstein;
- (b) $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) < \infty$;
- (c) $\dim_{\mathcal{GI}_I(\mathcal{A})}(\mathcal{A}) < \infty$.

Moreover, if this holds, then the following numbers coincide:

- (i) $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A})$;
- (ii) $\dim_{\mathcal{GI}_I(\mathcal{A})}(\mathcal{A})$;
- (iii) The smallest integer s such that $L_i\nu(A) = 0$ for all $i > s$ and $A \in \mathcal{A}$;
- (iv) The smallest integer t such that $R^i\nu^-(A) = 0$ for all $i > t$ and $A \in \mathcal{A}$.

We say that P is n -**Gorenstein** if this common number is n .

For a finite-dimensional algebra Λ , the endofunctor P is n -Gorenstein if and only if Λ is n -Gorenstein. For the endofunctor $P = f_! \circ f^*$ in Example 1.1.2 we get the following corollary.

Corollary 1.1.8 (See Theorem 5.3.4). *Let k be a commutative ring, and let Λ be a k -algebra which is finitely generated and projective as a k -module. Assume that*

$$\text{proj. dim } D(\Lambda)_{\Lambda} < \infty \quad \text{and} \quad \text{proj. dim } {}_{\Lambda}D(\Lambda) < \infty.$$

Then

$$\text{proj. dim } D(\Lambda)_{\Lambda} = \text{proj. dim } {}_{\Lambda}D(\Lambda).$$

Note that this dimension is n if and only if P is n -Gorenstein. We also have a version of this result for small, locally bounded, and Hom-finite categories, see Theorem 5.3.4. As far as we know, these results have not been written down before.

The paper is organized as follows. In Section 2 we recall the necessary preliminaries on adjunctions and monads that we need. In Section 3 we define Nakayama functors relative to adjoint pairs and endofunctors, and we provide examples. Furthermore, we give some different characterizations for when such Nakayama functors exist, see Proposition 3.3.1, Corollary 3.3.2 and Theorem 3.3.3, as well as proving a uniqueness result for such Nakayama functors, see Theorem 3.3.4. We end the section by proving Theorem 1.1.3 above. In Section 4 we develop a theory of Gorenstein homological algebra

for Nakayama functors relative to endofunctors. In particular, we define Gorenstein P -projective and I -injective objects and show that they form a resolving and coresolving subcategory, see Proposition 4.1.5. We also define the notion of P being Iwanaga-Gorenstein, and we prove Theorem 1.1.7 above. In Section 5 we show that the functor category $\mathcal{B}^{\mathcal{C}}$ has a Nakayama functors relative to an adjoint pair, where \mathcal{B} is an abelian category and \mathcal{C} is locally bounded Hom-finite category, see Theorem 5.3.3.

1.2. Conventions. All categories are assumed to be preadditive, and all functors are assumed to be additive. Also, all subcategories are assumed to be full. We always let \mathcal{A} and \mathcal{B} denote abelian categories. For a natural transformation $\eta: F \rightarrow F'$ we let $\eta_c: F(c) \rightarrow F'(c)$ denote the component of η at c . The natural transformation obtained by precomposing with a functor G is denoted by $\eta_G: F \circ G \rightarrow F' \circ G$. Unless otherwise specified, k denotes a commutative ring. We let $\text{proj } k$ denote the category of finitely generated projective k -modules. For a ring Λ we let $\Lambda\text{-Mod}$ and $\Lambda\text{-mod}$ denote the category of left Λ -modules and finitely presented left Λ -modules, respectively.

2. BACKGROUND

2.1. Transformation of adjoints. Let \mathcal{D} and \mathcal{E} be preadditive categories. An adjunction $(L, R, \phi, \alpha, \beta): \mathcal{D} \rightarrow \mathcal{E}$ is given by a pair of functors $L: \mathcal{D} \rightarrow \mathcal{E}$ and $R: \mathcal{E} \rightarrow \mathcal{D}$ together with an isomorphism

$$\phi: \mathcal{E}(L(D), E) \rightarrow \mathcal{D}(D, R(E))$$

for all pairs $D \in \mathcal{D}$ and $E \in \mathcal{E}$, which is natural in D and E . We also write $L \dashv R$ if there exists such an adjunction. Here $\alpha_D := \phi(1_{L(D)}): D \rightarrow RL(D)$ and $\beta_E := \phi^{-1}(1_{R(E)}): LR(E) \rightarrow E$ are the unit and the counit of the adjunction. They satisfy the *triangular identities*, i.e $R(\beta_E) \circ \alpha_{R(E)} = 1_{R(E)}$ and $\beta_{L(D)} \circ L(\alpha_D) = 1_{L(D)}$. By naturality we have that $\phi(f) = R(f) \circ \alpha_D$ and $\phi^{-1}(g) = \beta_E \circ L(g)$ for any morphisms $f: L(D) \rightarrow E$ and $g: D \rightarrow R(E)$. If we are working with several adjunctions, we write $\phi^{L \dashv R}$, $\alpha^{L \dashv R}$ and $\beta^{L \dashv R}$ for the isomorphism, the unit and the counit to indicate which adjunction we mean.

Assume $(L_1, R_1, \phi_1, \alpha_1, \beta_1): \mathcal{D} \rightarrow \mathcal{E}$ and $(L_2, R_2, \phi_2, \alpha_2, \beta_2): \mathcal{D} \rightarrow \mathcal{E}$ are adjunctions. Following [22], we say that two natural transformations $\sigma: L_1 \rightarrow L_2$ and $\tau: R_2 \rightarrow R_1$ are *conjugate* (for the given adjunctions) if the square

$$\begin{array}{ccc} \mathcal{E}(L_2(D), E) & \xrightarrow{\phi_2} & \mathcal{D}(D, R_2(E)) \\ \downarrow - \circ \sigma_D & & \downarrow \tau_E \circ - \\ \mathcal{E}(L_1(D), E) & \xrightarrow{\phi_1} & \mathcal{D}(D, R_1(E)) \end{array} \quad (2.1.1)$$

commutes for all pairs $D \in \mathcal{D}$ and $E \in \mathcal{E}$.

Proposition 2.1.2. *Let $(L_i, R_i, \phi_i, \alpha_i, \beta_i): \mathcal{D} \rightarrow \mathcal{E}$ be adjunctions for $1 \leq i \leq 2$. The following hold:*

- (i) *If $\sigma: L_1 \rightarrow L_2$ is a natural transformation, then there exists a unique natural transformation $\tau: R_2 \rightarrow R_1$ which is conjugate to σ ;*
- (ii) *If $\tau: R_2 \rightarrow R_1$ is a natural transformation, then there exists a unique natural transformation $\sigma: L_1 \rightarrow L_2$ which is conjugate to τ ;*

Proof. This follows from [22, Theorem IV.7.2] □

Proposition 2.1.3. *Assume we have adjunctions*

$$(L, G, \phi_1, \alpha_1, \beta_1): \mathcal{D} \rightarrow \mathcal{E} \quad \text{and} \quad (G, R, \phi_2, \alpha_2, \beta_2): \mathcal{E} \rightarrow \mathcal{D}.$$

Then α_1 and β_2 are conjugate, and β_1 and α_2 are conjugate.

Proof. This follows from [21, Theorem 2.15] and its dual. □

We need the following result later.

Proposition 2.1.4. *Let \mathcal{A} and \mathcal{B} be abelian categories. Assume*

$$(L_1, R_1, \phi_1, \alpha_1, \beta_1): \mathcal{B} \rightarrow \mathcal{A} \quad \text{and} \quad (L_2, R_2, \phi_2, \alpha_2, \beta_2): \mathcal{B} \rightarrow \mathcal{A}$$

are adjunctions, and $\sigma: L_1 \rightarrow L_2$ and $\tau: R_2 \rightarrow R_1$ are conjugate natural transformations. Then the functor $\text{Ker } \tau: \mathcal{A} \rightarrow \mathcal{B}$ is right adjoint to the functor $\text{Coker } \sigma: \mathcal{B} \rightarrow \mathcal{A}$.

Proof. Let $A \in \mathcal{A}$ and $B \in \mathcal{B}$ be arbitrary. We have exact sequences

$$\begin{aligned} L_1(B) &\xrightarrow{\sigma_B} L_2(B) \rightarrow \text{Coker } \sigma_B \rightarrow 0 \\ 0 &\rightarrow \text{Ker } \tau_A \rightarrow R_2(A) \xrightarrow{\tau_A} R_1(A). \end{aligned}$$

Applying $\mathcal{A}(-, A)$ to the first sequence and $\mathcal{B}(B, -)$ to the second sequence gives a diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & \mathcal{A}(\text{Coker } \sigma_B, A) & \longrightarrow & \mathcal{A}(L_2(B), A) & \xrightarrow{- \circ \sigma_B} & \mathcal{A}(L_1(B), A) \\ & & \downarrow \cong & & \downarrow \phi_2 & & \downarrow \phi_1 \\ 0 & \longrightarrow & \mathcal{B}(B, \text{Ker } \tau_A) & \longrightarrow & \mathcal{B}(B, R_2(A)) & \xrightarrow{\tau_A \circ -} & \mathcal{B}(B, R_1(A)) \end{array}$$

with exact rows. Since τ and σ are conjugate, the right square commutes. Therefore, we get an induced natural isomorphism

$$\mathcal{A}(\text{Coker } \sigma_B, A) \cong \mathcal{B}(B, \text{Ker } \tau_A)$$

and the result follows. □

2.2. Monads and comonads. We refer to chapter 4 in [6] and chapter VI in [22] for the general theory of monads and comonads. Let \mathcal{D} and \mathcal{E} be preadditive categories.

Definition 2.2.1.

- (i) A **monad** on \mathcal{D} is a tuple $\mathbb{T} = (T, \eta, \mu)$, where $T: \mathcal{D} \rightarrow \mathcal{D}$ is a functor and $\eta: 1_{\mathcal{D}} \rightarrow T$ and $\mu: T \circ T \rightarrow T$ are natural transformations such that the diagrams

$$\begin{array}{ccc} TTT & \xrightarrow{T(\mu)} & TT \\ \downarrow \mu_T & & \downarrow \mu \\ TT & \xrightarrow{\mu} & T \end{array} \quad \begin{array}{ccc} & T & \\ 1_T \nearrow & \mu \uparrow & \nwarrow 1_T \\ T & \xrightarrow{T(\eta)} & TT \xleftarrow{\eta_T} T \end{array}$$

commute.

- (ii) A **comonad** on \mathcal{D} is a tuple $\mathbb{S} = (S, \epsilon, \Delta)$, where $S: \mathcal{D} \rightarrow \mathcal{D}$ is a functor and $\epsilon: S \rightarrow 1_{\mathcal{D}}$ and $\Delta: S \rightarrow S \circ S$ are natural transformations such that the diagrams

$$\begin{array}{ccc} S & \xrightarrow{\Delta} & SS \\ \downarrow \Delta & & \downarrow \Delta_S \\ SS & \xrightarrow{S(\Delta)} & SSS \end{array} \quad \begin{array}{ccc} & S & \\ 1_S \swarrow & \Delta \downarrow & \searrow 1_S \\ S & \xleftarrow{S(\epsilon)} & SS \xrightarrow{\epsilon_S} S \end{array}$$

commute.

Note that a comonad on \mathcal{D} is the same as a monad on \mathcal{D}^{op} .

A **morphism of monads** $\delta: (T_1, \eta_1, \mu_1) \rightarrow (T_2, \delta_2, \mu_2)$ is given by a natural transformation $\delta: T_1 \rightarrow T_2$ satisfying $\mu_2 \circ \delta^2 = \delta \circ \mu_1$ and $\eta_2 = \delta \circ \eta_1$, where $\delta^2 = T_2(\delta) \circ \delta_{T_1} = \delta_{T_2} \circ T_1(\delta)$. Dually, a **morphism of comonads** $\zeta: (S_1, \epsilon_1, \Delta_1) \rightarrow (S_2, \epsilon_2, \Delta_2)$ is given by a natural transformation $\zeta: S_1 \rightarrow S_2$ satisfying $\Delta_2 \circ \zeta = \zeta^2 \circ \Delta_1$ and $\epsilon_1 = \epsilon_2 \circ \zeta$, where $\zeta^2 = S_2(\zeta) \circ \zeta_{S_1} = \zeta_{S_2} \circ S_1(\zeta)$. For a monad $\mathbb{T} = (T, \eta, \mu)$ on \mathcal{D} , an object D is called **\mathbb{T} -injective** if $\eta_D: D \rightarrow T(D)$ is a split monomorphism. These are precisely the summands of objects of the form $T(D)$. Dually, for a comonad $\mathbb{S} = (S, \epsilon, \Delta)$ on \mathcal{D} , an object D is called **\mathbb{S} -projective** if $\epsilon_D: S(D) \rightarrow D$ is a split epimorphism. These are precisely the summands of objects of the form $S(D)$.

Remark 2.2.2. The category of endofunctors on \mathcal{D} is a monoidal category. The product is given by composition, and the unit object is the identity functor. A monad and a comonad is just a monoid and a comonoid in this monoidal category.

Example 2.2.3. If $(L, R, \phi, \alpha, \beta): \mathcal{D} \rightarrow \mathcal{E}$ is an adjunction, then the tuple $\mathbb{T} = (R \circ L, \alpha, R(\beta_L))$ is a monad on \mathcal{D} , and the tuple $\mathbb{S} = (L \circ R, \beta, L(\alpha_R))$ is a comonad on \mathcal{E} .

We also have a notion of adjunctions between monads and comonads.

Definition 2.2.4. Let $\mathbb{T} = (T, \eta, \mu)$ and $\mathbb{S} = (S, \epsilon, \Delta)$ be a monad and comonad on \mathcal{D} .

- (i) We say that \mathbb{S} is **left adjoint** to \mathbb{T} (written $\mathbb{S} \dashv \mathbb{T}$) if S is left adjoint to T such that $\Delta: S \rightarrow SS$ and $\mu: TT \rightarrow T$ are conjugate, and $\epsilon: S \rightarrow 1_{\mathcal{D}}$ and $\eta: 1_{\mathcal{D}} \rightarrow T$ are conjugate.
- (ii) We say that \mathbb{S} is **right adjoint** to \mathbb{T} (written $\mathbb{T} \dashv \mathbb{S}$) if S is right adjoint to T such that $\Delta: S \rightarrow SS$ and $\mu: TT \rightarrow T$ are conjugate, and $\epsilon: S \rightarrow 1_{\mathcal{D}}$ and $\eta: 1_{\mathcal{D}} \rightarrow T$ are conjugate.

If $\mathbb{T} = (T, \eta, \mu)$ is a monad and S is a left or right adjoint of T , then by Proposition 2.1.2 there exist a unique comonad $\mathbb{S} = (S, \epsilon, \Delta)$ which is left or right adjoint to \mathbb{T} , respectively.

We recall the Eilenberg-Moore category of a monad.

Definition 2.2.5. Let $\mathbb{T} = (T, \eta, \mu)$ and $\mathbb{S} = (S, \epsilon, \Delta)$ be a monad and comonad on \mathcal{D} .

- (i) A **\mathbb{T} -module** is a morphism $s: T(D) \rightarrow D$ satisfying $s \circ \eta_D = 1_D$ and $s \circ \mu_D = s \circ T(s)$;
- (ii) A **\mathbb{S} -comodule** is a morphism $t: D \rightarrow S(D)$ satisfying $\epsilon_D \circ t = 1_D$ and $\Delta_D \circ t = S(t) \circ t$;
- (iii) The **Eilenberg-Moore category** $\mathcal{D}^{\mathbb{T}}$ of \mathbb{T} is the category where the objects are all \mathbb{T} -modules, and where a morphism between \mathbb{T} -modules $T(D_1) \xrightarrow{s_1} D_1$ to $T(D_2) \xrightarrow{s_2} D_2$ is a morphism $f: D_1 \rightarrow D_2$ in \mathcal{D} satisfying $f \circ s_1 = s_2 \circ T(f)$.
- (iv) The **Eilenberg-Moore category** $\mathcal{D}^{\mathbb{S}}$ of \mathbb{S} is the category where the objects are all \mathbb{S} -comodules, and where a morphism between \mathbb{S} -comodules $D_1 \xrightarrow{t_1} S(D_1)$ to $D_2 \xrightarrow{t_2} S(D_2)$ is a morphism $f: D_1 \rightarrow D_2$ in \mathcal{D} satisfying $t_2 \circ f = S(f) \circ t_1$.
- (v) The **Kleisli category** $\mathcal{D}_{\mathbb{T}}$ of \mathbb{T} is the category with objects being the objects in \mathcal{D} , and with morphisms being

$$\mathcal{D}_{\mathbb{T}}(D_1, D_2) := \mathcal{D}(D_1, T(D_2))$$

Composition of $D_1 \xrightarrow{g} T(D_2)$ and $D_2 \xrightarrow{h} T(D_3)$ in $\mathcal{D}_{\mathbb{T}}$ is given by the composite $D_1 \xrightarrow{g} T(D_2) \xrightarrow{T(h)} TT(D_3) \xrightarrow{\mu_{D_3}} T(D_3)$ in \mathcal{D} . The unit at $D \in \mathcal{D}_{\mathbb{T}}$ is $\eta_D: D \rightarrow T(D)$.

- (vi) The **co-Kleisli category** $\mathcal{D}_{\mathbb{S}}$ of \mathbb{S} is the category with objects being the objects in \mathcal{D} , and with morphisms being

$$\mathcal{D}_{\mathbb{S}}(D_1, D_2) := \mathcal{D}(S(D_1), D_2)$$

Composition of $S(D_1) \xrightarrow{g} D_2$ and $S(D_2) \xrightarrow{h} D_3$ in $\mathcal{D}_{\mathbb{S}}$ is given by the composite $S(D_1) \xrightarrow{\Delta_{D_1}} SS(D_1) \xrightarrow{S(g)} S(D_2) \xrightarrow{h} D_3$ in \mathcal{D} . The unit at $D \in \mathcal{D}_{\mathbb{S}}$ is $\epsilon_D: S(D) \rightarrow D$.

The forgetful functor $U^\top: \mathcal{D}^\top \rightarrow \mathcal{D}$ admits a left adjoint $F^\top: \mathcal{D} \rightarrow \mathcal{D}^\top$ sending D to $TT(D) \xrightarrow{\mu_D} T(D)$ and $g: D_1 \rightarrow D_2$ to $T(g): T(D_1) \rightarrow T(D_2)$, see [6, Proposition 4.1.4]. Similarly, we have functors

$$\begin{aligned} U_\top: \mathcal{D}_\top \rightarrow \mathcal{D} & \quad (D \mapsto T(D)) \quad ((g: D_1 \rightarrow T(D_2)) \mapsto \mu_{D_2} \circ T(g)) \\ F_\top: \mathcal{D} \rightarrow \mathcal{D}_\top & \quad (D \mapsto D) \quad ((f: D_1 \rightarrow D_2) \mapsto \eta_{D_2} \circ f) \end{aligned}$$

where F_\top is left adjoint to U_\top , see [6, Proposition 4.1.7]. For both of the adjunctions $F^\top \dashv U^\top$ and $F_\top \dashv U_\top$ the induced monad on \mathcal{D} is \top , see [6, Proposition 4.2.2]. The dual statements hold for the comonad S .

We need the following technical result to prove Theorem 3.3.3, which is used to prove Theorem 1.1.3 stated in the introduction. Here, for a functor $F: \mathcal{D} \rightarrow \mathcal{E}$, we let $\text{im } F$ denote the image of F . This is a category with same objects as \mathcal{D} , and where a morphism between D_1 and D_2 is a morphism $F(D_1) \rightarrow F(D_2)$ in \mathcal{E} .

Theorem 2.2.6. *Let*

$$(L_1, R_1, \phi_1, \alpha_1, \beta_1): \mathcal{D} \rightarrow \mathcal{E} \quad \text{and} \quad (L_2, R_2, \phi_2, \alpha_2, \beta_2): \mathcal{D} \rightarrow \mathcal{E}$$

be adjunctions, and let

$$\top_1 := (R_1 \circ L_1, \alpha_1, R_1((\beta_1)_{L_1})) \quad \text{and} \quad \top_2 := (R_2 \circ L_2, \alpha_2, R_2((\beta_2)_{L_2}))$$

be the induced monads on \mathcal{D} .

(i) *Given a morphism $\gamma: \top_1 \rightarrow \top_2$ of monads, we get a functor*

$$E_\gamma: \text{im } L_1 \rightarrow \text{im } L_2$$

satisfying $E_\gamma \circ L_1 = L_2$. It acts as identity on objects, and sends a morphism $f: L_1(A_1) \rightarrow L_1(A_2)$ to

$$(\beta_2)_{L_2(A_2)} \circ L_2(\gamma_{A_2}) \circ L_2 R_1(f) \circ L_2((\alpha_1)_{A_1}): L_2(A_1) \rightarrow L_2(A_2)$$

(ii) *Given a functor $E: \text{im } L_1 \rightarrow \text{im } L_2$ satisfying $E \circ L_1 = L_2$, there exists a unique morphism $\gamma: \top_1 \rightarrow \top_2$ of monads such that $E_\gamma = E$.*

Proof. Let \mathcal{D}_{\top_1} and \mathcal{D}_{\top_2} denote the Kleisli categories of \top_1 and \top_2 , and let $F_{\top_1}: \mathcal{D} \rightarrow \mathcal{D}_{\top_1}$ and $F_{\top_2}: \mathcal{D} \rightarrow \mathcal{D}_{\top_2}$ be the left adjoints in the Kleisli adjunctions. By [6, Proposition 4.2.1] there exists isomorphisms

$$\Phi_1: \text{im } L_1 \rightarrow \mathcal{D}_{\top_1} \quad \text{and} \quad \Phi_2: \text{im } L_2 \rightarrow \mathcal{D}_{\top_2}$$

satisfying $\Phi_1 \circ L_1 = F_{\top_1}$ and $\Phi_2 \circ L_2 = F_{\top_2}$. Explicitly, they act as identity on objects, and they send morphisms $f_1: L_1(A_1) \rightarrow L_1(A_2)$ and $f_2: L_2(A_1) \rightarrow L_2(A_2)$ to $\phi_1(f_1): A_1 \rightarrow R_1 L_1(A_2)$ and $\phi_2(f_2): A_1 \rightarrow R_2 L_2(A_2)$, respectively. Now any morphism $\gamma: \top_1 \rightarrow \top_2$ of monads induces a functor

$E'_\gamma: \mathcal{D}_{\top_1} \rightarrow \mathcal{D}_{\top_2}$ satisfying $E'_\gamma \circ F_{\top_1} = F_{\top_2}$. It acts as identity on objects and sends a morphism $A_1 \xrightarrow{f} R_1 L_1(A_1)$ to $A_1 \xrightarrow{f} R_1 L_1(A_1) \xrightarrow{\gamma_{A_1}} R_2 L_2(A_1)$. The composite $E_\gamma = (\Phi_2)^{-1} \circ E'_\gamma \circ \Phi_1$ gives the functor in part (i).

For Part (ii), we note that by [9, Proposition 2.4] (with J being the identity functor in their notation) the association $\gamma \rightarrow E'_\gamma$ gives a one-to-one correspondence between morphisms of monads $\gamma: \top_1 \rightarrow \top_2$ and functors

$E': \mathcal{D}_{\mathbb{T}_1} \rightarrow \mathcal{D}_{\mathbb{T}_2}$ satisfying $E' \circ F_{\mathbb{T}_1} = F_{\mathbb{T}_2}$. In particular, given a functor $E: \text{im } L_1 \rightarrow \text{im } L_2$ satisfying $E \circ L_1 = L_2$, the functor $E' := \Phi_2 \circ E \circ \Phi_1^{-1}$ satisfies $E' \circ F_{\mathbb{T}_1} = F_{\mathbb{T}_2}$ and therefore must be equal to E'_γ for some $\gamma: \mathbb{T}_1 \rightarrow \mathbb{T}_2$. Composing E' with the isomorphisms Φ_1 and Φ_2 then gives the result. \square

For a functor $E: \text{im } L_1 \rightarrow \text{im } L_2$ satisfying $E \circ L_1 = L_2$, one can easily check that the unique morphism of monads $\gamma: \mathbb{T}_1 \rightarrow \mathbb{T}_2$ with $E_\gamma = E$ is given by

$$\gamma := R_1 L_1 \xrightarrow{(\alpha_2)_{R_1 L_1}} R_2 L_2 R_1 L_1 = R_2 E L_1 R_1 L_1 \xrightarrow{R_2 E (\beta_1)_{L_1}} R_2 E L_1 = R_2 L_2. \quad (2.2.7)$$

A functor $R: \mathcal{D} \rightarrow \mathcal{E}$ is called *monadic* if there exists a monad \mathbb{T} on \mathcal{E} , an equivalence $J: \mathcal{D} \rightarrow \mathcal{E}^{\mathbb{T}}$, and an isomorphism of functors $R \cong U^{\mathbb{T}} \circ J$ where $U^{\mathbb{T}}: \mathcal{E}^{\mathbb{T}} \rightarrow \mathcal{E}$ is the forgetful functor. In the following we state a weak version of Beck's monadicity theorem, a result which characterizes when a functor is monad. See [6, Theorem 4.4.4] for the general version.

Theorem 2.2.8. *Let $R: \mathcal{A} \rightarrow \mathcal{B}$ be a functor between abelian categories. Then R is monadic if the following conditions hold:*

- (i) R has a left adjoint;
- (ii) R is faithful;
- (iii) R is right exact.

Proof. This follows from [6, Theorem 4.4.4]. \square

2.3. Properties of subcategories. Fix abelian categories \mathcal{A} and \mathcal{B} .

Definition 2.3.1. Let $\mathcal{X} \subset \mathcal{A}$ be a subcategory.

- (i) \mathcal{X} is *generating* if for all objects $A \in \mathcal{A}$ there exists $X \in \mathcal{X}$ and an epimorphism $X \rightarrow A$;
- (ii) \mathcal{X} is *cogenerating* if for all objects $A \in \mathcal{A}$ there exists $X \in \mathcal{X}$ and a monomorphism $A \rightarrow X$;
- (iii) \mathcal{X} is *resolving* if it is a generating subcategory which is closed under direct summands, extensions, and kernel of epimorphism;
- (iv) \mathcal{X} is *coresolving* if it is a cogenerating subcategory which is closed under direct summands, extensions, and cokernels of monomorphism.

Here we follow the conventions in [24] for the definition of resolving and coresolving. We can define the *resolution dimension* $\dim_{\mathcal{X}}(A)$ of any object $A \in \mathcal{A}$ with respect to a resolving subcategory \mathcal{X} . Explicitly, it is the smallest integer $n \geq 0$ such that there exists a long exact sequence

$$0 \rightarrow X_n \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 \rightarrow A \rightarrow 0$$

where $X_i \in \mathcal{X}$ for $0 \leq i \leq n$. We write $\dim_{\mathcal{X}}(A) = \infty$ if there doesn't exist such a resolution. If $\dim_{\mathcal{X}}(A) = n$, the dimension can be computed using

any resolution of A by objects in \mathcal{X} , i.e for any exact sequence

$$0 \rightarrow X'_n \rightarrow \cdots X'_1 \rightarrow X'_0 \rightarrow A \rightarrow 0$$

with $X'_i \in \mathcal{X}$ for all $0 \leq i \leq n-1$ we get that $X'_n \in \mathcal{X}$, see [24, Proposition 2.3]. The **global resolution dimension** $\dim_{\mathcal{X}}(\mathcal{A})$ of \mathcal{A} is defined as the supremum of $\dim_{\mathcal{X}}(A)$ over all $A \in \mathcal{A}$. We can define the **coresolution dimension** $\dim_{\mathcal{Y}}$ with respect to a coresolving category \mathcal{Y} dually.

Definition 2.3.2 (III.6.3 in [16]). Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor, and let \mathcal{X} be a subcategory of \mathcal{A} .

- (i) If F is left exact, we say that \mathcal{X} is **adapted** to F if \mathcal{X} is cogenerating and for any exact sequence $0 \rightarrow X_0 \rightarrow X_{-1} \rightarrow X_{-2} \rightarrow \cdots$ with $X_i \in \mathcal{X}$ for all $i \leq 0$ we have that $0 \rightarrow F(X_0) \rightarrow F(X_{-1}) \rightarrow F(X_{-2}) \rightarrow \cdots$ is exact;
- (ii) If F is right exact, we say that \mathcal{X} is **adapted** to F if \mathcal{X} is generating and for any exact sequence $\cdots \rightarrow X_2 \rightarrow X_1 \rightarrow X_0 \rightarrow 0$ with $X_i \in \mathcal{X}$ for all $i \geq 0$ we have that $\cdots \rightarrow F(X_2) \rightarrow F(X_1) \rightarrow F(X_0) \rightarrow 0$ is exact.

Theorem 2.3.3 (Theorem III.6.8 in [16]). *Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be an additive functor, and let \mathcal{X} be a subcategory of \mathcal{A} . The following holds:*

- (i) *If F is left exact and \mathcal{X} is adapted to F , then the right derived functor $RF: D^+(\mathcal{A}) \rightarrow D^+(\mathcal{B})$ exists, and $RF(X) \cong X$ for all $X \in \mathcal{X}$;*
- (ii) *If F is right exact and \mathcal{X} is adapted to F , then the left derived functor $LF: D^-(\mathcal{A}) \rightarrow D^-(\mathcal{B})$ exists, and $LF(X) \cong X$ for all $X \in \mathcal{X}$.*

The i th right and left derived functor is define to be $R^i F := H^i(RF)$ and $L_i F := H^{-i}(LF)$, respectively. Note that $R^i F(X) = 0$ and $L_i F(X) = 0$ for $X \in \mathcal{X}$ and $i > 0$ in these cases.

3. GENERALIZATION OF THE NAKAYAMA FUNCTOR

3.1. Derived functors without enough projectives or injectives. Our goal is to find easy criteria for when a subcategory is adapted to a functor when the subcategory is the image of another functor. We fix a preadditive category \mathcal{D} and abelian categories \mathcal{A} and \mathcal{B} . By abuse of notation, in the following definition we identify the image $\text{im } F$ of a functor $F: \mathcal{D} \rightarrow \mathcal{A}$ with the full subcategory of \mathcal{A} consisting of all objects of the form $F(D)$ for $D \in \mathcal{D}$.

Definition 3.1.1. Let $F: \mathcal{D} \rightarrow \mathcal{A}$ be a functor.

- (i) We say that F is **cogenerating** if $\text{im } F$ is cogenerating in \mathcal{A} ;
- (ii) We say that F is **generating** if $\text{im } F$ is generating in \mathcal{A} ;
- (iii) Let $G: \mathcal{A} \rightarrow \mathcal{B}$ be a left or right exact functor. We say that F is **adapted to G** if $\text{im } F$ is adapted to G ;

If $S = (S, \epsilon, \Delta)$ is a comonad on \mathcal{A} , then S is generating if and only if ϵ is an epimorphism. Dually, if $T = (T, \eta, \mu)$ is a monad on \mathcal{A} , then T is cogenerating if and only if η is a monomorphism.

Proposition 3.1.2. *Let $G: \mathcal{A} \rightarrow \mathcal{B}$ be a functor.*

- (i) *Assume G is left exact, $T = (T, \eta, \mu)$ is a monad on \mathcal{A} , and the functor $T: \mathcal{A} \rightarrow \mathcal{A}$ is cogenerating. If T and $G \circ T$ are exact functors, then T is adapted to G ;*
- (ii) *Assume G is right exact, $S = (S, \epsilon, \Delta)$ is comonad on \mathcal{A} , and the functor $S: \mathcal{A} \rightarrow \mathcal{A}$ is generating. If S and $G \circ S$ are exact functors, then S is adapted to G .*

Proof. We only show part (ii), part (i) follows dually. For $A \in \mathcal{A}$ consider the complex $C'(A) := \dots \xrightarrow{(\delta_2)_A} S^2(A) \xrightarrow{(\delta_1)_A} S(A) \xrightarrow{(\delta_0)_A} A \rightarrow 0$ where $\delta_n = \sum_{i=0}^n (-1)^i S^{n-i}(\epsilon_{S^i})$. By 4.2 in [4] the complex $\mathcal{A}(S(A'), C'(A))$ is acyclic for all objects A' . Since S is generating, the subcategory of objects of the form $S(A')$ is generating. Hence, any sequence $A_1 \rightarrow A_2 \rightarrow A_3$ is exact if the sequence $\mathcal{A}(S(A'), A_1) \rightarrow \mathcal{A}(S(A'), A_2) \rightarrow \mathcal{A}(S(A'), A_3)$ is exact for all $A' \in \mathcal{A}$. In particular, the complex $C'(A)$ is itself acyclic.

Let $C(A) := G(C'(A))$. We prove the following claim: Let $0 \rightarrow A_3 \xrightarrow{f_2} A_2 \xrightarrow{f_1} A_1 \rightarrow 0$ be an exact sequence, and assume the complexes $C(A_1)$ and $C(A_2)$ are acyclic. Then the following holds:

- (i) The sequence $0 \rightarrow G(A_3) \xrightarrow{G(f_2)} G(A_2) \xrightarrow{G(f_1)} G(A_1) \rightarrow 0$ is exact;
- (ii) The complex $C(A_3)$ is acyclic.

To show this, we consider the commutative diagram

$$\begin{array}{ccccc} GS^2(A_3) & \xrightarrow{GS^2(f_2)} & GS^2(A_2) & \xrightarrow{GS^2(f_1)} & GS^2(A_1) \\ \downarrow G((\delta_1)_{A_3}) & & \downarrow G((\delta_1)_{A_2}) & & \downarrow G((\delta_1)_{A_1}) \\ GS(A_3) & \xrightarrow{GS(f_2)} & GS(A_2) & \xrightarrow{GS(f_1)} & GS(A_1) \end{array}$$

with exact rows. Since $GS^3(f_1): GS^3(A_2) \rightarrow GS^3(A_1)$ is an epimorphism, and $C(A_1)$ and $C(A_2)$ are acyclic, it follows that the induced map

$$\text{Ker } G((\delta_1)_{A_2}) \rightarrow \text{Ker } G((\delta_1)_{A_1})$$

is an epimorphism. Hence, applying the snake lemma and using that the sequence $GS^2(A_3) \rightarrow GS(A_3) \rightarrow G(A_3) \rightarrow 0$ is exact since G is right exact, we get that the sequence $0 \rightarrow G(A_3) \xrightarrow{G(f_2)} G(A_2) \xrightarrow{G(f_1)} G(A_1) \rightarrow 0$ is exact. This proves part (i) of the claim. Also it follows that we have a short exact sequences

$$0 \rightarrow C(A_3) \xrightarrow{C(f_2)} C(A_2) \xrightarrow{C(f_1)} C(A_1) \rightarrow 0$$

of complexes. Since two of the complexes are acyclic, the third one is also acyclic. This proves part (ii) of the claim.

To prove the result in the proposition, it is sufficient to show that for an exact sequence

$$0 \rightarrow K \rightarrow A_n \xrightarrow{f_n} \cdots \xrightarrow{f_2} A_1 \xrightarrow{f_1} A_0 \rightarrow 0$$

with A_i being \mathbf{S} -projective for all i , we have that

$$0 \rightarrow G(K) \rightarrow G(A_n) \xrightarrow{G(f_n)} \cdots \xrightarrow{G(f_2)} G(A_1) \xrightarrow{G(f_1)} G(A_0) \rightarrow 0 \quad (3.1.3)$$

is exact. To this end, note that since A_i is \mathbf{S} -projective, the complex $C'(A_i)$ is contractible by the proof of Proposition 8.6.8 in [25] (the proposition only says that $C'(A_i)$ is acyclic, but in the proof they actually show that it is contractible). Hence, the complex $C(A_i)$ is also contractible and therefore acyclic. By part (ii) of the claim it follows that the complex $C(\text{Ker } f_i)$ is acyclic for all i . But then by part (i) of the claim it follows that the sequence (3.1.3) is exact. This proves the result. \square

Proposition 3.1.4. *Let $G: \mathcal{A} \rightarrow \mathcal{B}$ be a functor.*

- (i) *Assume G is left exact, $(L, R, \phi, \alpha, \beta): \mathcal{A} \rightarrow \mathcal{D}$ is an adjunction, and R is cogenerating. If $R \circ L$ and $G \circ R \circ L$ are exact functors, then R is adapted to G ;*
- (ii) *Assume G is right exact, $(L', R', \phi', \alpha', \beta'): \mathcal{D} \rightarrow \mathcal{A}$ is an adjunction, and L' is generating. If $L' \circ R'$ and $G \circ L' \circ R'$ are exact functors, then L' is adapted to G .*

Proof. From the triangle identities we have that $R(A)$ is a summand of $RLR(A)$. Hence, if R is cogenerating, then $R \circ L$ is cogenerating. Therefore, by Proposition 3.1.2 applied to the induced monad $\mathbb{T} = (R \circ L, \alpha, R(\beta_L))$, we get that $R \circ L$ is adapted to G . Finally, since $R(A)$ is a summand of $RLR(A)$ for any $A \in \mathcal{A}$, it follows that R itself is adapted to G . Part (ii) is proved dually. \square

3.2. Definition of Nakayama functors. Let Λ be a finite-dimensional algebra over a field k , and consider the restriction functor

$$f^* := \text{res}_k^\Lambda: \Lambda\text{-mod} \rightarrow k\text{-mod}.$$

It has a left adjoint $f_! := (\Lambda \otimes_k -): k\text{-mod} \rightarrow \Lambda\text{-mod}$, and it turns out that the adjoint pair $f_! \dashv f^*$ interact nicely with the Nakayama functor $\nu = \text{Hom}_k(\Lambda, k) \otimes_\Lambda -: \Lambda\text{-mod} \rightarrow \Lambda\text{-mod}$. More precisely, the composite $\nu \circ f_!$ is right adjoint to f^* , and there exists an isomorphism $\nu^- \circ \nu \circ f_! \cong f_!$ where ν^- is the right adjoint of ν . We generalize this in the following definition.

Definition 3.2.1. Let \mathcal{D} be an additive category and \mathcal{A} an abelian category, and let $f^*: \mathcal{A} \rightarrow \mathcal{D}$ be a faithful functor with left adjoint $f_!: \mathcal{D} \rightarrow \mathcal{A}$. A **Nakayama functor** relative to $f_! \dashv f^*$ is a functor $\nu: \mathcal{A} \rightarrow \mathcal{A}$ with a right adjoint ν^- satisfying:

- (i) $\nu \circ f_!$ is right adjoint to f^* ;
- (ii) The unit of $\nu \dashv \nu^-$ induces an isomorphism $f_! \xrightarrow{\cong} \nu^- \circ \nu \circ f_!$ when precomposed with $f_!$.

We also say that $f_! \dashv f^*$ has a Nakayama functor ν . In this case we let $f_* := \nu \circ f_!$ denote the right adjoint. Since the left adjoint of a functor is unique up to isomorphism, the definition of the Nakayama functor only depends on the functor f^* in the adjunction $f_! \dashv f^*$. It also turns out that the Nakayama functor is unique up to precomposing with an equivalence $\Phi: \mathcal{A} \rightarrow \mathcal{A}$ satisfying $\Phi \circ f_! = f_!$, see Theorem 3.3.4. Note that for a finite-dimensional algebra, the classical Nakayama functor is a Nakayama functor relative to the adjoint pair $(\Lambda \otimes_k -) \dashv \text{res}_k^\Lambda$ as described above.

We fix $(\nu, \nu^-, \theta, \lambda, \sigma): \mathcal{A} \rightarrow \mathcal{A}$ to be the adjunction of the Nakayama functor. The following holds for adjoint pairs with Nakayama functor.

Lemma 3.2.2. *Assume $f_! \dashv f^*$ admits a Nakayama functor ν . The following holds:*

- (i) $f_!$ is generating;
- (ii) f_* is cogenerating;
- (iii) The restriction $\nu: \text{im } f_! \rightarrow \text{im } f_*$ is an isomorphism of categories;
- (iv) $\sigma_{f_*}: \nu \circ \nu^- \circ f_* \rightarrow f_*$ is an isomorphism;
- (v) There are adjunctions $f^* \circ \nu \dashv f_! \dashv f^* \dashv f_* \dashv f^* \circ \nu^-$
- (vi) $f_!$ is adapted to ν ;
- (vii) f_* is adapted to ν^- .

Proof. Since f^* is faithful, it follows by [22, Theorem IV.3.1] and its dual that the counit of $f_! \dashv f^*$ is an epimorphism and the unit of $f^* \dashv f_*$ is a monomorphism. This shows that $f_!$ is generating and f_* is cogenerating. Part (iii) is obvious. For part (iv), note that $\sigma_{\nu \circ f_!} \circ \nu(\lambda_{f_!}) = 1_{\nu \circ f_!}$ by the triangle identities for $\nu \dashv \nu^-$. Since $\lambda_{f_!}$ is an isomorphism, it follows that $\sigma_{f_*} = \sigma_{\nu \circ f_!}$ is an isomorphism. For part (v), since $f^* \dashv \nu \circ f_!$, we have

$$f^* \circ \nu \dashv \nu^- \circ (\nu \circ f_!) \cong f_!.$$

Also, since $f_! \dashv f^*$, we have that $\nu \circ f_! \dashv f^* \circ \nu^-$. Part (vi) and (vii) follows immediately from Proposition 3.1.4. \square

Note that if ν is a Nakayama functor relative to $f_! \dashv f^*$, then ν^- is a Nakayama functor relative to $f_* \dashv f^*$ in the opposite categories.

For the constructions and results in Section 4 we only need to work with the endofunctor $P := f_! \circ f^*$, and not with the adjoint pair. We therefore introduce the following definition.

Definition 3.2.3. Let \mathcal{A} be an abelian category with a generating functor $P: \mathcal{A} \rightarrow \mathcal{A}$. A **Nakayama functor** relative to P is a functor $\nu: \mathcal{A} \rightarrow \mathcal{A}$ with a right adjoint $\nu^-: \mathcal{A} \rightarrow \mathcal{A}$ satisfying the following:

- (i) $\nu \circ P$ is right adjoint to P ;

- (ii) The unit of $\nu \dashv \nu^-$ induces an isomorphism $P \xrightarrow{\cong} \nu^- \circ \nu \circ P$ when precomposed with P .

We also say that P has a Nakayama functor ν . Note that if ν is a Nakayama functor for $f_! \dashv f^*$, then it is a Nakayama functor for the composite $P := f_! \circ f^*$.

In the following we let $I = \nu \circ P$. We also fix $(\nu, \nu^-, \theta, \lambda, \sigma): \mathcal{A} \rightarrow \mathcal{A}$ to be the adjunction of the Nakayama functor.

Lemma 3.2.4. *Let \mathcal{A} be an abelian category with a generating functor $P: \mathcal{A} \rightarrow \mathcal{A}$ and a Nakayama functor ν relative to P . The following holds:*

- (i) *The restriction $\nu: \text{im } P \rightarrow \text{im } I$ is an isomorphism of categories;*
- (ii) *$\sigma_I: \nu \circ \nu^- \circ I \rightarrow I$ is an isomorphism;*
- (iii) *There are adjunctions $P \circ \nu \dashv P \dashv I \dashv I \circ \nu^-$;*
- (iv) *P is adapted to ν ;*
- (v) *I is adapted to ν^- ;*
- (vi) *$I: \mathcal{A} \rightarrow \mathcal{A}$ is a cogenerating functor;*
- (vii) *P is faithful;*
- (viii) *I is faithful.*

Proof. Part (i)-(v) is proved similarly to part (iii)-(vii) in Lemma 3.2.2. For part (vi) note that the functor $\text{Ker } \alpha^{I \dashv I \circ \nu^-}$ is right adjoint to the functor $\text{Coker } \beta^{P \dashv I}$ by Proposition 2.1.3 and Proposition 2.1.4. Since P is generating, we have $\text{Coker } \beta^{P \dashv I} = 0$, and therefore $\text{Ker } \alpha^{I \dashv I \circ \nu^-} = 0$. Hence, I is cogenerating. For part (vii), note that we have an equality $\alpha_B^{P \dashv I} \circ f = IP(f) \circ \alpha_A^{P \dashv I}$ for any morphism $f: A \rightarrow B$. Since the functor I is cogenerating, the unit $\alpha_B^{P \dashv I}$ of the adjunction $P \dashv I$ is a monomorphism. Hence, the left hand side of the equality is nonzero. Therefore $IP(f) \circ \alpha_A^{P \dashv I} \neq 0$, hence $IP(f) \neq 0$, and so $P(f) \neq 0$. This shows that P is faithful. Part (viii) is proved dually. \square

Note that if ν is a Nakayama functor relative to P , then ν^- is a Nakayama functor relative to I in the opposite category.

We now give several examples of Nakayama functors for adjoints. They are all special cases of Theorem 5.3.3, obtained by identifying with functor categories via $\Lambda\text{-Mod} = (k\text{-Mod})^\Lambda$, $(\Lambda_1 \otimes_k \Lambda_2)\text{-Mod} = (\Lambda_2\text{-Mod})^{\Lambda_1}$ and $\mathcal{C}\text{-Mod} = (k\text{-Mod})^{\mathcal{C}}$ in Example 3.2.5, 3.2.6 and 3.2.7, respectively.

Example 3.2.5. Let k be a commutative ring, and let Λ be a k -algebra which is finitely generated and projective as a k -module. Furthermore, let

$$f^* := \text{res}_k^\Lambda: \Lambda\text{-Mod} \rightarrow k\text{-Mod} \quad \text{and} \quad f_! := \Lambda \otimes_k -: k\text{-Mod} \rightarrow \Lambda\text{-Mod}$$

Then the functor

$$\nu = \text{Hom}_k(\Lambda, k) \otimes_\Lambda -: \Lambda\text{-Mod} \rightarrow \Lambda\text{-Mod}$$

is a Nakayama functor relative to the adjoint pair $f_! \dashv f^*$. This follows from Theorem 5.3.3, but for the convenience of the reader we explain the

argument in more detail here: First note that the composite $\nu \circ f_!$ is given by

$$\nu \circ f_! \cong \text{Hom}_k(\Lambda, k) \otimes_k -: k\text{-Mod} \rightarrow \Lambda\text{-Mod}$$

Since Λ is finitely generated projective over k , the functor

$$\text{Hom}_k(\Lambda, -): k\text{-Mod} \rightarrow \Lambda\text{-Mod}$$

is exact and preserves coproducts, and hence there exists an isomorphism

$$\text{Hom}_k(\Lambda, -) \cong \text{Hom}_k(\Lambda, k) \otimes_k -: k\text{-Mod} \rightarrow \Lambda\text{-Mod}$$

Therefore, $\nu \circ f_!$ is right adjoint to f^* . Also, the right adjoint of ν is given by

$$\nu^- = \text{Hom}_\Lambda(\text{Hom}_k(\Lambda, k), -): \Lambda\text{-Mod} \rightarrow \Lambda\text{-Mod}$$

and we have

$$\nu^- \circ \nu \circ f_! \cong \text{Hom}_\Lambda(\text{Hom}_k(\Lambda, k), \text{Hom}_k(\Lambda, -)) \cong \text{Hom}_k(\text{Hom}_k(\Lambda, k), -)$$

where the second isomorphism follows from the Hom-tensor adjunction. Furthermore, since $\text{Hom}_k(\Lambda, k)$ is finitely generated projective over k , we have an isomorphism

$$\text{Hom}_k(\text{Hom}_k(\Lambda, k), -) \cong \text{Hom}_k(\text{Hom}_k(\Lambda, k), k) \otimes_k -$$

Finally, since Λ is finitely generated projective over k , the natural map $\Lambda \rightarrow \text{Hom}_k(\text{Hom}_k(\Lambda, k), k)$ is an isomorphism, and hence we have

$$\text{Hom}_k(\text{Hom}_k(\Lambda, k), k) \otimes_k - \cong \Lambda \otimes_k - = f_!$$

Composing these maps, we get an isomorphism $f_! \cong \nu^- \circ \nu \circ f_!$ which is given by the unit of the adjunction $\nu \dashv \nu^-$.

If we assume k is coherent, then $f_! \dashv f^*$ restrict to an adjoint pair with Nakayama functor ν between the category of finitely presented modules $\Lambda\text{-mod}$ and $k\text{-mod}$. Note that if k is a field, then we just obtain the classical Nakayama functor.

Example 3.2.6. Let k be a commutative ring, let Λ_1 be a k -algebra which is finitely generated and projective as a k -module, and let Λ_2 be a k -algebra. Furthermore, let

$$\begin{aligned} f^* &:= \text{res}_{\Lambda_2}^{\Lambda_1 \otimes_k \Lambda_2}: \Lambda_1 \otimes_k \Lambda_2\text{-Mod} \rightarrow \Lambda_2\text{-Mod} \\ f_! &:= \Lambda_1 \otimes_k -: \Lambda_2\text{-Mod} \rightarrow \Lambda_1 \otimes_k \Lambda_2\text{-Mod} \end{aligned}$$

Then the functor

$$\nu = \text{Hom}_k(\Lambda_1, k) \otimes_{\Lambda_1} -: \Lambda_1 \otimes_k \Lambda_2\text{-Mod} \rightarrow \Lambda_1 \otimes_k \Lambda_2\text{-Mod}$$

is a Nakayama functor for the adjoint pair $f_! \dashv f^*$. If we furthermore assume Λ_2 is left coherent, then $f_! \dashv f^*$ restrict to an adjoint pair with Nakayama functor ν between the category of finitely presented modules $\Lambda_1 \otimes_k \Lambda_2\text{-mod}$ and $\Lambda_2\text{-mod}$.

Example 3.2.7. Let k be a commutative ring, and let \mathcal{C} be a small, k -linear, Hom-finite and locally bounded category, see Definition 5.2.1. Then the evaluation functor

$$i^*: \mathcal{C}\text{-Mod} \rightarrow \prod_{c \in \mathcal{C}} k\text{-Mod} \quad i^*(F) = (F(c))_{c \in \mathcal{C}}$$

has a left adjoint $i_!$ given by

$$i_!: \prod_{c \in \mathcal{C}} k\text{-Mod} \rightarrow \mathcal{C}\text{-Mod} \quad i_!(M_c)_{c \in \mathcal{C}} = \bigoplus_{c \in \mathcal{C}} \mathcal{C}(c, -) \otimes_k M_c$$

Furthermore, we have a functor

$$\nu = \text{Hom}_k(\mathcal{C}, k) \otimes_{\mathcal{C}} -: \mathcal{C}\text{-Mod} \rightarrow \mathcal{C}\text{-Mod}$$

given by $\nu(M)(c) = \text{Hom}_k(\mathcal{C}(c, -), k) \otimes_{\mathcal{C}} M$, and it turns out that this is a Nakayama functor relative to $i_! \dashv i^*$. See Subsection 5.3 for more details.

Example 3.2.8. Let k be a commutative ring, and let \mathcal{C} be the k -linear category generated by the quiver

$$\cdots \xleftarrow{d_{i-1}} c_{i-1} \xleftarrow{d_i} c_i \xleftarrow{d_{i+1}} \cdots$$

with vertex set $\{c_i \mid i \in \mathbb{Z}/n\mathbb{Z}\}$ and relations $d_i \circ d_{i+1} = 0$. The category $\mathcal{B}^{\mathcal{C}}$ of k -linear functors from \mathcal{C} to \mathcal{B} can be identified with n -periodic complexes $B_{\bullet} = \cdots \xleftarrow{d_{i-1}} B_{i-1} \xleftarrow{d_i} B_i \xleftarrow{d_{i+1}} \cdots$ over \mathcal{B} (for $n = 0$ this is just unbounded complexes over \mathcal{B}). Let $[1]: \mathcal{B}^{\mathcal{C}} \rightarrow \mathcal{B}^{\mathcal{C}}$ be the shift functor defined by $(B_{\bullet}[1])_i = B_{i+1}$. The restriction functor

$$i^*: \mathcal{B}^{\mathcal{C}} \rightarrow \prod_{i \in \mathbb{Z}/n\mathbb{Z}} \mathcal{B} \quad i^*(B_{\bullet}) = (B_i)_{i \in \mathbb{Z}/n\mathbb{Z}}$$

has a left adjoint

$$i_!: \prod_{i \in \mathbb{Z}/n\mathbb{Z}} \mathcal{B} \rightarrow \mathcal{B}^{\mathcal{C}} \quad i_!(B_i)_{i \in \mathbb{Z}/n\mathbb{Z}} = \bigoplus_{i \in \mathbb{Z}/n\mathbb{Z}} C(B_i)[-i]$$

and a right adjoint

$$i_*: \prod_{i \in \mathbb{Z}/n\mathbb{Z}} \mathcal{B} \rightarrow \mathcal{B}^{\mathcal{C}} \quad i_*(B_i)_{i \in \mathbb{Z}/n\mathbb{Z}} = \bigoplus_{i \in \mathbb{Z}/n\mathbb{Z}} C(B_i)[-i-1]$$

where $C(B_i)$ is the complex with $C(B_i)_0 = C(B_i)_{-1} = B_i$, the differential $C(B_i)_0 \rightarrow C(B_i)_{-1}$ is the identity, and $C(B_i)_k = 0$ for $k \neq 0, -1$. It follows that $[-1]$ is a Nakayama functor relative to $i_! \dashv i^*$.

Example 3.2.9. Let $\mathbb{A}_2 = (1 \rightarrow 2)$ be the category with two objects and one morphism between them, and let $\mathcal{C} = k\mathbb{A}_2$ be the k -linearization of \mathbb{A}_2 . An object in $\mathcal{B}^{k\mathbb{A}_2}$ is then just a morphism $B_1 \xrightarrow{f} B_2$ in \mathcal{B} . The restriction functor

$$i^*: \mathcal{B}^{k\mathbb{A}_2} \rightarrow \mathcal{B} \prod \mathcal{B} \quad i^*(B_1 \xrightarrow{f} B_2) = (B_1, B_2)$$

has a left adjoint given by

$$i_! : \mathcal{B} \prod \mathcal{B} \rightarrow \mathcal{B}^{k\mathbb{A}_2} \quad i_!(B_1, B_2) = (B_1 \xrightarrow{1} B_1) \oplus (0 \rightarrow B_2).$$

The cokernel functor $\nu(B_1 \xrightarrow{f} B_2) = B_2 \rightarrow \text{Coker } f$ is then a Nakayama functor relative to $i_! \dashv i^*$, and its right adjoint ν^- is given by $\nu^-(B_1 \xrightarrow{f} B_2) = \text{Ker } f \rightarrow B_1$.

More generally, for $\mathbb{A}_n = (1 \rightarrow 2 \rightarrow \cdots \rightarrow n)$ the restriction functor

$$i^* : \mathcal{B}^{k\mathbb{A}_n} \rightarrow \prod_{1 \leq i \leq n} \mathcal{B} \quad i^*(B_1 \xrightarrow{f_1} B_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} B_n) = (B_1, B_2, \dots, B_n)$$

has a left adjoint $i_! : \prod_{1 \leq i \leq n} \mathcal{B} \rightarrow \mathcal{B}^{k\mathbb{A}_n}$ given by

$$i_!(B_1, B_2, \dots, B_n) = \bigoplus_{1 \leq i \leq n} (0 \rightarrow \cdots \rightarrow 0 \rightarrow B_i \xrightarrow{1} B_i \xrightarrow{1} \cdots \xrightarrow{1} B_i)$$

where the first $n+1-i$ terms of $(0 \rightarrow \cdots \rightarrow 0 \rightarrow B_i \xrightarrow{1} B_i \xrightarrow{1} \cdots \xrightarrow{1} B_i)$ are 0. The functor

$$\begin{aligned} \nu(B_1 \xrightarrow{f_1} B_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} B_n) = \\ (B_n \rightarrow \text{Coker } f_{n-1} \rightarrow \text{Coker}(f_{n-1}f_{n-2}) \rightarrow \cdots \rightarrow \text{Coker}(f_{n-1}f_{n-2} \cdots f_1)) \end{aligned}$$

is then a Nakayama functor relative to $i_! \dashv i^*$, and its right adjoint is given by

$$\begin{aligned} \nu^-(B_1 \xrightarrow{f_1} B_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} B_n) = \\ (\text{Ker}(f_{n-1}f_{n-2} \cdots f_1) \rightarrow \cdots \rightarrow \text{Ker}(f_2f_1) \rightarrow \text{Ker } f_1 \rightarrow B_1) \end{aligned}$$

3.3. Properties of Nakayama functors. Fix an abelian category \mathcal{A} and an additive category \mathcal{D} . Our main goal in this subsection is to prove Theorem 1.1.3 which was stated in the introduction.

If ν is a Nakayama functor relative to $f_! \dashv f^*$, then the restriction

$$E := \nu|_{\text{im } f_!} : \text{im } f_! \rightarrow \text{im } f_*$$

is an isomorphism of categories satisfying $E \circ f_! = f_*$, where $f_* = \nu \circ f_!$. We want to show that such an isomorphism is enough to have a Nakayama functor. To this end, we assume we have adjunctions $f_! \dashv f^* \dashv f_*$. We define functors

$$\begin{aligned} \text{mor}_{f_!} : \mathcal{A} &\rightarrow \text{Mor}(\text{im } f_!) & A &\rightarrow \text{mor}_{f_!}(A) \\ \text{mor}_{f_*} : \mathcal{A} &\rightarrow \text{Mor}(\text{im } f_*) & A &\rightarrow \text{mor}_{f_*}(A) \end{aligned}$$

where $\text{Mor}(\text{im } f_!)$ and $\text{Mor}(\text{im } f_*)$ are the categories of morphisms in $\text{im } f_!$ and $\text{im } f_*$, respectively. Explicitly, $\text{mor}_{f_!}$ and mor_{f_*} are defined by

$$\begin{aligned} \text{mor}_{f_!}(A) &:= i \circ \beta_{\text{Ker } \beta_A^{f_! \dashv f^*}}^{f_! \dashv f^*} : f_! f^*(\text{Ker } \beta_A^{f_! \dashv f^*}) \rightarrow f_! f^*(A) \\ \text{mor}_{f_*}(A) &:= \alpha_{\text{Coker } \alpha_A^{f^* \dashv f_*}}^{f^* \dashv f_*} \circ p : f_* f^*(A) \rightarrow f_* f^*(\text{Coker } \alpha_A^{f^* \dashv f_*}) \end{aligned}$$

for an object $A \in \mathcal{A}$, where $i: \text{Ker } \beta_A^{f_! \dashv f^*} \rightarrow f_! f^*(A)$ is the inclusion and $p: f_* f^*(A) \rightarrow \text{Coker } \alpha_A^{f^* \dashv f_*}$ is the projection.

Proposition 3.3.1. *Assume we have adjunctions $f_! \dashv f^* \dashv f_*$ where $f^*: \mathcal{A} \rightarrow \mathcal{D}$ is a faithful functor. Furthermore, let $E: \text{im } f_! \rightarrow \text{im } f_*$ be an isomorphism satisfying $E \circ f_! = f_*$. The following holds:*

- (i) *There exists a Nakayama functor ν relative to $f_! \dashv f^*$ which satisfies $\nu|_{\text{im } f_!} = E$;*
- (ii) *If ν' is any other right exact functor satisfying $\nu'|_{\text{im } f_!} = E$, then there exists a unique isomorphism $\zeta: \nu \rightarrow \nu'$ satisfying $\zeta_{f_!} = 1_{f_*}$.*

Proof. Let $E: \text{im } f_! \rightarrow \text{im } f_*$ be an isomorphism satisfying $E \circ f_! = f_*$, and let E^- be its inverse. These lifts to isomorphisms $E: \text{Mor}(\text{im } f_!) \rightarrow \text{Mor}(\text{im } f_*)$ and $E^-: \text{Mor}(\text{im } f_*) \rightarrow \text{Mor}(\text{im } f_!)$ defined pointwise. Note that the sequences

$$\begin{aligned} f_* f^*(\text{Ker } \beta_{f_!(D)}^{f_! \dashv f^*}) &\xrightarrow{E(\text{mor}_{f_!}(f_!(D)))} f_* f^* f_!(D) \xrightarrow{E(\beta_{f_!(D)}^{f_! \dashv f^*})} f_*(D) \rightarrow 0 \\ 0 \rightarrow f_!(D) &\xrightarrow{E^-(\alpha_{f_*(D)}^{f^* \dashv f_*})} f_! f^* f_*(D) \xrightarrow{E^-(\text{mor}_{f_*}(f_*(D)))} f_! f^*(\text{Coker } \alpha_{f_*(D)}^{f^* \dashv f_*}) \end{aligned}$$

are exact in \mathcal{A} . Hence, we can define functors

$$\begin{aligned} \nu &:= \text{Coker} \circ E \circ \text{mor}_{f_!} : \mathcal{A} \rightarrow \mathcal{A} \\ \nu^- &:= \text{Ker} \circ E^- \circ \text{mor}_{f_*} : \mathcal{A} \rightarrow \mathcal{A} \end{aligned}$$

such that $\nu|_{\text{im } f_!} = E$ and $\nu^-|_{\text{im } f_*} = E^-$, where $\text{Ker}: \text{Mor}(\text{im } f_!) \rightarrow \mathcal{A}$ and $\text{Coker}: \text{Mor}(\text{im } f_*) \rightarrow \mathcal{A}$ are the kernel and cokernel functors. We claim that $\nu \dashv \nu^-$. Let $h: \nu(A_1) \rightarrow A_2$ be a morphism in \mathcal{A} . By definition, we have an exact sequence

$$f_* f^*(\text{Ker } \beta_{A_1}^{f_! \dashv f^*}) \xrightarrow{E(\text{mor}_{f_!}(A_1))} f_* f^*(A_1) \rightarrow \nu(A_1) \rightarrow 0$$

Consider the composite

$$\bar{h} := f_* f^*(A_1) \rightarrow \nu(A_1) \xrightarrow{h} A_2 \xrightarrow{\alpha_{A_2}^{f^* \dashv f_*}} f_* f^*(A_2)$$

It satisfies $\bar{h} \circ E(\text{mor}_{f_!}(A_1)) = 0$ and $\text{mor}_{f_*}(A_2) \circ \bar{h} = 0$. Applying E^- to \bar{h} therefore gives a morphism $E^-(\bar{h}): f_! f^*(A_1) \rightarrow f_! f^*(A_2)$ satisfying

$E^-(\bar{h}) \circ \text{mor}_{f_!}(A_1) = 0$ and $E^-(\text{mor}_{f_*}(A_2)) \circ E^-(\bar{h}) = 0$. Since we have exact sequences

$$f_!f^*(\text{Ker } \beta_{A_1}^{f_! \dashv f^*}) \xrightarrow{\text{mor}_{f_!}(A_1)} f_!f^*(A_1) \xrightarrow{\beta_{A_1}^{f_! \dashv f^*}} A_1 \rightarrow 0$$

and

$$0 \rightarrow \nu^-(A_2) \rightarrow f_!f^*(A_2) \xrightarrow{E^-(\text{mor}_{f_*}(A_2))} f_!f^*(\text{Coker } \alpha_{A_2}^{f^* \dashv f_*})$$

it follows that the morphism $E^-(\bar{h})$ induces a morphism $\theta(h): A_1 \rightarrow \nu^-(A_2)$. Obviously, the map $h \rightarrow \theta(h)$ is bijective and natural, and therefore we have an adjunction $\nu \dashv \nu^-$. Furthermore, under this bijection the unit $1_{\mathcal{A}} \rightarrow \nu^- \circ \nu$ is the identity on $\text{im } f_!$. Hence, ν is a Nakayama functor which satisfies $\nu|_{\text{im } f_!} = E$. The uniqueness of ν is obvious. \square

We get the following corollary from the proof of Proposition 3.3.1, which is useful for computing the Nakayama functor and its adjoint in examples.

Corollary 3.3.2. *Assume we have adjunctions $f_! \dashv f^* \dashv f_*$ where $f^*: \mathcal{A} \rightarrow \mathcal{D}$ is a faithful functor. The following holds:*

- (i) *If $\nu: \mathcal{A} \rightarrow \mathcal{A}$ is a right exact functor such that $\nu \circ f_! = f_*$ and $\nu|_{\text{im } f_!}: \text{im } f_! \rightarrow \text{im } f_*$ is an isomorphism, then ν is a Nakayama functor relative to $f_! \dashv f^*$;*
- (ii) *If $\nu^-: \mathcal{A} \rightarrow \mathcal{A}$ is a left exact functor such that $\nu^- \circ f_* = f_!$ and $\nu^-|_{\text{im } f_*}: \text{im } f_* \rightarrow \text{im } f_!$ is an inverse to $\nu|_{\text{im } f_!}$, then $\nu \dashv \nu^-$;*
- (iii) *The unit and counit of $\nu \dashv \nu^-$ are the unique natural transformations $\lambda: 1_{\mathcal{A}} \rightarrow \nu^- \circ \nu$ and $\sigma: \nu \circ \nu^- \rightarrow 1_{\mathcal{A}}$ which satisfies $\lambda_{f_!} = 1_{f_!}$ and $\sigma_{f_*} = 1_{f_*}$.*

Combining Proposition 3.3.1 with Theorem 2.2.6 gives the following.

Theorem 3.3.3. *Let $f^*: \mathcal{A} \rightarrow \mathcal{D}$ be a faithful functor, and assume there exists adjunctions $f_! \dashv f^* \dashv f_* \dashv f^\times$.*

- (i) *If $\nu: \mathcal{A} \rightarrow \mathcal{A}$ is a Nakayama functor relative to $f_! \dashv f^*$ satisfying $\nu \circ f_! = f_*$, then $\gamma_\nu := \chi_{f_*} \circ f^*(\lambda_{f_!})$ induces an isomorphism of monads*

$$\gamma_\nu: (f^*f_!, \alpha^{f_! \dashv f^*}, f^*(\beta_{f_!}^{f_! \dashv f^*})) \xrightarrow{\cong} (f^\times f_*, \alpha^{f^* \dashv f^\times}, f^\times(\beta_{f_*}^{f^* \dashv f^\times}))$$

where $\chi: f^* \circ \nu^- \xrightarrow{\cong} f^\times$ is the natural transformation which is conjugate to the identity $f_* = \nu \circ f_!$;

- (ii) *If $\gamma: (f^*f_!, \alpha^{f_! \dashv f^*}, f^*(\beta_{f_!}^{f_! \dashv f^*})) \xrightarrow{\cong} (f^\times f_*, \alpha^{f^* \dashv f^\times}, f^\times(\beta_{f_*}^{f^* \dashv f^\times}))$ is an isomorphism of monads, then there exists a Nakayama functor $\nu: \mathcal{A} \rightarrow \mathcal{A}$ relative to $f_! \dashv f^*$ satisfying $\nu \circ f_! = f_*$ and $\gamma_\nu = \gamma$. Furthermore, if ν' is another Nakayama functor relative to $f_! \dashv f^*$ satisfying $\nu' \circ f_! = f_*$ and $\gamma_{\nu'} = \gamma$, then there exists a unique isomorphism $\zeta: \nu \rightarrow \nu'$ with $\zeta_{f_!} = 1_{f_*}$.*

Proof. We only need to show that the natural transformation in (2.2.7) coincide with the one given for γ_ν above. The rest follows from Proposition 3.3.1 and Theorem 2.2.6. By (2.2.7) we have that

$$\gamma_\nu := f^* f_! \xrightarrow{(\alpha^{f^* \dashv f^\times})_{f^* f_!}} f^\times f_* f^* f_! = f^\times \nu f_! f^* f_! \xrightarrow{f^\times \nu (\beta^{f_! \dashv f^*})_{f_!}} f^\times \nu f_! = f^\times f_*$$

Since

$$\phi^{f^* \dashv f^\times} : \mathcal{A}(\nu \circ f_!, -) \xrightarrow{\theta} \mathcal{A}(f_!, \nu^-) \xrightarrow{\phi^{f_! \dashv f^*}} \mathcal{D}(-, f^* \circ \nu^-) \xrightarrow{\chi^{\circ-}} \mathcal{D}(-, f^\times).$$

it follows that

$$\begin{aligned} \alpha^{f^* \dashv f^\times} &= \phi^{f^* \dashv f^\times}(1_{f_*}) = \chi_{f_*} \circ \phi^{f_! \dashv f^*}(\theta(1_{\nu \circ f_!})) = \chi_{f_*} \circ \phi^{f_! \dashv f^*}(\lambda_{f_!}) \\ &= \chi_{f_*} \circ f^*(\lambda_{f_!}) \circ \alpha^{f_! \dashv f^*} \end{aligned}$$

Hence, it follows that

$$\begin{aligned} \gamma_\nu &= f^\times \nu (\beta^{f_! \dashv f^*})_{f_!} \circ (\alpha^{f^* \dashv f^\times})_{f^* f_!} \\ &= f^\times \nu (\beta^{f_! \dashv f^*}) \circ \chi_{f_* f^* f_!} \circ f^*(\lambda_{f_! f^* f_!}) \circ \alpha_{f^* f_!}^{f_! \dashv f^*} \\ &= \chi_{f_*} \circ f^* \nu^- \nu (\beta_{f_!}^{f_! \dashv f^*}) \circ f^*(\lambda_{f_! f^* f_!}) \circ \alpha_{f^* f_!}^{f_! \dashv f^*} \\ &= \chi_{f_*} \circ f^*(\lambda_{f_!}) \circ f^*(\beta_{f_!}^{f_! \dashv f^*}) \circ \alpha_{f^* f_!}^{f_! \dashv f^*} \\ &= \chi_{f_*} \circ f^*(\lambda_{f_!}) \end{aligned}$$

by naturality and the triangle identities. \square

We leave it to the reader to state the dual of Theorem 3.3.3.

We have the following uniqueness result for Nakayama functors.

Theorem 3.3.4. *Assume we have adjunctions $f_! \dashv f^* \dashv f_*$ where $f^* : \mathcal{A} \rightarrow \mathcal{D}$ is a faithful functor. Let ν_1 and ν_2 be Nakayama functors relative to $f_! \dashv f^*$ satisfying $\nu_1 \circ f_! = f_* = \nu_2 \circ f_!$. The following holds:*

- (i) *There exists an equivalence $\Phi : \mathcal{A} \rightarrow \mathcal{A}$ satisfying $\Phi \circ f_! = f_!$ and a unique natural isomorphism $\xi : \nu_1 \circ \Phi \rightarrow \nu_2$ satisfying $\xi_{f_!} = 1_{f_*}$. Furthermore, if Ψ is another such equivalence, then there exists a unique natural isomorphism $\zeta : \Phi \rightarrow \Psi$ satisfying $\zeta_{f_!} = 1_{f_!}$.*
- (ii) *There exists an equivalence $\Phi' : \mathcal{A} \rightarrow \mathcal{A}$ satisfying $\Phi' \circ f_* = f_*$ and a unique natural isomorphism $\xi' : \Phi' \circ \nu_1 \rightarrow \nu_2$ satisfying $\xi'_{f_!} = 1_{f_*}$. Furthermore, if Ψ' is another such equivalence, then there exists a unique natural isomorphism $\zeta : \Phi' \rightarrow \Psi'$ satisfying $\zeta_{f_!} = 1_{f_*}$.*

Proof. We only prove part (i), part (ii) is proved similarly. Let $E_1 := \nu_1|_{\text{im } f_!} : \text{im } f_! \rightarrow \text{im } f_*$ and $E_2 := \nu_2|_{\text{im } f_!} : \text{im } f_! \rightarrow \text{im } f_*$, and let $H := E_1^{-1} \circ E_2 : \text{im } f_! \rightarrow \text{im } f_!$. Then obviously $H \circ f_! = f_!$. We define $\Phi := \text{Coker} \circ H \circ \text{mor}_{f_!}$, where we choose the cokernels such that $\Phi|_{\text{im } f_!} = H$. Since $\text{im } f_!$ is generating, Φ is an equivalence on \mathcal{A} . Furthermore, $\Phi \circ f_! = f_!$. Applying ν_1 to the exact sequence

$$H f_! f^*(\text{Ker } \beta_A^{f_! \dashv f^*}) \rightarrow H f_! f^*(A) \rightarrow \Phi(A) \rightarrow 0$$

and using that $\nu_1|_{\text{im } f_!} = E_1$ and $E_1 H = E_2$, we get a commutative diagram with exact rows

$$\begin{array}{ccccccc} E_1 H f_! f^*(\text{Ker } \beta_A^{f_! \dashv f^*}) & \longrightarrow & E_1 H f_! f^*(A) & \longrightarrow & \nu_1 \Phi(A) & \longrightarrow & 0 \\ \downarrow = & & \downarrow = & & \downarrow \xi_A & & \\ E_2 f_! f^*(\text{Ker } \beta_A^{f_! \dashv f^*}) & \longrightarrow & E_2 f_! f^*(A) & \longrightarrow & \nu_2(A) & \longrightarrow & 0 \end{array}$$

This defines ξ_A for each $A \in \mathcal{A}$. Obviously, ξ satisfies $\xi_{f_!} = 1_{f_*}$, and is the unique such natural isomorphism.

Now let $\Psi: \mathcal{A} \rightarrow \mathcal{A}$ be another such equivalence. Then the composite $\zeta': \nu_1 \circ \Phi \xrightarrow{\cong} \nu_2 \xrightarrow{\cong} \nu_1 \circ \Psi$ gives a natural isomorphism satisfying $\zeta'_{f_!} = 1_{f_*}$. Restricting to $\text{im } f_!$ and using that $\nu_1|_{\text{im } f_!}: \text{im } f_! \rightarrow \text{im } f_*$ is an isomorphism, we get that $\Phi|_{\text{im } f_!} = \Psi|_{\text{im } f_!}$. Hence, we have a commutative diagram

$$\begin{array}{ccccccc} f_! f^*(\text{Ker } \beta_A^{f_! \dashv f^*}) & \xrightarrow{\Phi(\text{mor}_{f_!}(A))} & f_! f^*(A) & \longrightarrow & \Phi(A) & \longrightarrow & 0 \\ \downarrow = & & \downarrow = & & \downarrow \zeta_A & & \\ f_! f^*(\text{Ker } \beta_A^{f_! \dashv f^*}) & \xrightarrow{\Psi(\text{mor}_{f_!}(A))} & f_! f^*(A) & \longrightarrow & \Psi(A) & \longrightarrow & 0 \end{array}$$

which defines ζ_A for each $A \in \mathcal{A}$. Obviously ζ is the unique isomorphism satisfying $\zeta_{f_!} = 1_{f_!}$. \square

Let $f^*: \mathcal{A} \rightarrow \mathcal{B}$ be a faithful functor between abelian categories with a left adjoint $f_!$, and assume $f_! \dashv f^*$ has a Nakayama functor as in Definition 3.2.1. Then f^* is faithful and exact. Let $\mathbb{T} = (f^* \circ f_!, \alpha^{f_! \dashv f^*}, f^*(\beta_{f_!}^{f_! \dashv f^*}))$ be the induced monad on \mathcal{B} , and let $\mathcal{B}^{\mathbb{T}}$ be the Eilenberg-Moore category of \mathbb{T} , see Definition 2.2.5. By Beck's monadicity theorem we have an equivalence of categories $J: \mathcal{A} \xrightarrow{\cong} \mathcal{B}^{\mathbb{T}}$ such that $U^{\mathbb{T}} \circ J$ is isomorphic to f^* , see Theorem 2.2.8. It is therefore natural to ask for necessary and sufficient conditions on a monad \mathbb{T} on an abelian category \mathcal{B} so that the induced adjoint pair $F^{\mathbb{T}} \dashv U^{\mathbb{T}}$ on $\mathcal{B}^{\mathbb{T}}$ has a Nakayama functor.

Theorem 3.3.5. *Let \mathbb{T} be a monad on an abelian category \mathcal{B} , let $\mathcal{B}^{\mathbb{T}}$ be the Eilenberg-Moore category of \mathbb{T} , and let $U^{\mathbb{T}}: \mathcal{B}^{\mathbb{T}} \rightarrow \mathcal{B}$ be the forgetful functor with left adjoint $F^{\mathbb{T}}: \mathcal{B} \rightarrow \mathcal{B}^{\mathbb{T}}$. Then the adjoint pair $F^{\mathbb{T}} \dashv U^{\mathbb{T}}$ has a Nakayama functor if and only if there exists a comonad \mathbb{S} and an ambidextrous adjunction $\mathbb{S} \dashv \mathbb{T} \dashv \mathbb{S}$.*

Proof. We set $f^* := U^{\mathbb{T}}$ and $f_! := F^{\mathbb{T}}$. If $f_! \dashv f^*$ has a Nakayama functor, then by Lemma 3.2.2 part (v) we have adjunctions $f^? \dashv f_! \dashv f^* \dashv f_*$. Let $\mathbb{S}_1 = (f^? \circ f_!, \beta^{f^? \dashv f_!}, f^?(\alpha_{f_!}^{f^? \dashv f_!}))$ and $\mathbb{S}_2 = (f^* \circ f_*, \beta^{f^* \dashv f_*}, f^*(\alpha_{f_*}^{f^* \dashv f_*}))$ be the induced comonads on \mathcal{B} . It follows from [21, Theorem 2.15] that $\mathbb{S}_1 \dashv \mathbb{T} \dashv \mathbb{S}_2$. Finally, there exists an isomorphism $\mathbb{S}_1 \cong \mathbb{S}_2$ of comonads by

the dual of Theorem 3.3.3 part (i). This shows that we have an ambidextrous adjunction.

Now assume there exists an ambidextrous adjunction $S \dashv T \dashv S$ for some comonad S on \mathcal{B} . It follows from [11, Proposition 5.3] that \mathcal{B}^T is abelian. Furthermore, since $T \dashv S$, we have an isomorphism $J: \mathcal{B}^S \xrightarrow{\cong} \mathcal{B}^T$ of the Eilenberg-Moore categories which commute with the forgetful functors, see [21, Theorem 2.14]. Hence, if we put $f_* := J \circ F^S$, we get an adjoint triple $f_! \dashv f^* \dashv f_*$ such that $S = (f^* \circ f_*, \beta^{f^* \dashv f_*}, f^*(\alpha_{f_*}^{f^* \dashv f_*}))$. Now by the dual of [6, Lemma 4.3.3] we have that for every $A \in \mathcal{B}^T$ there exists an exact sequence $0 \rightarrow A \xrightarrow{\alpha_{f_*}^{f^* \dashv f_*}} f_* f^*(A) \xrightarrow{f_* f^*(\alpha_A^{f^* \dashv f_*}) - \alpha_{f_* f^*(A)}^{f^* \dashv f_*}} f_* f^* f_* f^*(A)$. Hence, in particular it follows that

$$f_! = \text{Ker}(f_* f^* f_! \xrightarrow{f_* f^*(\alpha_{f_!}^{f^* \dashv f_*}) - \alpha_{f_* f^* f_!}^{f^* \dashv f_*}} f_* f^* f_* f^* f_!)$$

Since $S \dashv T$, we have that $f^* f_* \dashv f^* f_!$. It follows by Proposition 2.1.2 that there exists a natural transformation $s: f^* f_* f^* f_! f^* \rightarrow f^* f_* f^*$ such that s and $f_* f^*(\alpha_{f_!}^{f^* \dashv f_*}) - \alpha_{f_* f^* f_!}^{f^* \dashv f_*}$ are conjugate. If we let $f^?: \mathcal{B}^T \rightarrow \mathcal{B}$ denote the cokernel of s , then it follows that $f^? \dashv f_!$ by Proposition 2.1.4. Now consider the comonad $S' := (f^? \circ f_!, \beta^{f^? \dashv f_!}, f^?(\alpha_{f_!}^{f^? \dashv f_!}))$. By [21, Theorem 2.15] we have that $S' \dashv T$. But since the left adjoint of a functor is unique up to an isomorphism and the conjugate of a natural transformation is unique, it follows that the left adjoint of a monad is unique up to an isomorphism of comonads. Therefore, since $S \dashv T$, we get that $S \cong S'$. Hence, it follows that

$$(f^? \circ f_!, \beta^{f^? \dashv f_!}, f^?(\alpha_{f_!}^{f^? \dashv f_!})) \cong (f^* \circ f_*, \beta^{f^* \dashv f_*}, f^*(\alpha_{f_*}^{f^* \dashv f_*}))$$

Finally, by the dual of Theorem 3.3.3 part (ii) it follows that $f_! \dashv f^*$ has a Nakayama functor. \square

4. GORENSTEIN HOMOLOGICAL ALGEBRA FOR NAKAYAMA FUNCTORS RELATIVE TO ENDOFUNCTORS

In this section we construct analogues of Gorenstein projective modules, Gorenstein injectives modules, and Iwanaga-Gorenstein algebras in the setting of Nakayama functors relative to endofunctors. Furthermore, we prove an analogue of Zak's result on the equality of the injective dimension for Iwanaga-Gorenstein algebras. We make the following assumption throughout the section.

Setting 4.0.1. Let $P: \mathcal{A} \rightarrow \mathcal{A}$ be a generating functor and let $\nu: \mathcal{A} \rightarrow \mathcal{A}$ be a Nakayama functor relative to P . We fix the notation $I := \nu \circ P$, $T = P \circ \nu$, and $S := I \circ \nu^-$.

Note that $T \dashv P \dashv I \dashv S$.

4.1. Gorenstein P -projective and I -injective objects. We call an object $A \in \mathcal{A}$ for P -*projective* or I -*injective* if it is a summand of an object $P(A')$ or $I(A')$, respectively.

Definition 4.1.1. Let $A \in \mathcal{A}$ be an object.

- (i) An object $A \in \mathcal{A}$ is **Gorenstein P -projective** if there exists an exact sequence

$$A_{\bullet} = \cdots \xrightarrow{s_2} A_1 \xrightarrow{s_1} A_0 \xrightarrow{s_0} A_{-1} \xrightarrow{s_{-1}} \cdots$$

with $A_i \in \mathcal{A}$ being P -projective for all $i \in \mathbb{Z}$, such that

$$\cdots \xrightarrow{\nu(s_2)} \nu(A_1) \xrightarrow{\nu(s_1)} \nu(A_0) \xrightarrow{\nu(s_0)} \nu(A_{-1}) \xrightarrow{\nu(s_{-1})} \cdots$$

is exact, and with $Z_0(A_{\bullet}) = \text{Ker } s_0 = A$. The subcategory of \mathcal{A} consisting of all Gorenstein P -projective objects is denoted by $\mathcal{GP}_P(\mathcal{A})$

- (ii) An object $A \in \mathcal{A}$ is **Gorenstein I -injective** if there exists an exact sequence

$$A_{\bullet} = \cdots \xrightarrow{s_2} A_1 \xrightarrow{s_1} A_0 \xrightarrow{s_0} A_{-1} \xrightarrow{s_{-1}} \cdots$$

with $A_i \in \mathcal{A}$ being I -injective for all $i \in \mathbb{Z}$, such that

$$\cdots \xrightarrow{\nu^-(s_2)} \nu^-(A_1) \xrightarrow{\nu^-(s_1)} \nu^-(A_0) \xrightarrow{\nu^-(s_0)} \nu^-(A_{-1}) \xrightarrow{\nu^-(s_{-1})} \cdots$$

is exact, and with $Z_0(A_{\bullet}) = \text{Ker } s_0 = A$. The subcategory of \mathcal{A} consisting of all Gorenstein I -injective objects is denoted by $\mathcal{GI}_I(\mathcal{A})$

In Example 3.2.5 with k a field and $P = f_! \circ f^*$, a module is Gorenstein P projective or I -injective if and only if it is Gorenstein projective or injective, respectively. In Example 3.2.8 with $P = f_! \circ f^*$ all objects are Gorenstein P -projective and I -injective. In Example 3.2.9 with $P = f_! \circ f^*$ a straightforward computation shows that a morphism is Gorenstein P -projective or I -injective if and only if it is a monomorphism or an epimorphism, respectively.

The following result shows that $\mathcal{GP}_P(\mathcal{A})$ and $\mathcal{GI}_I(\mathcal{A})$ are equivalent categories.

Proposition 4.1.2. *Let $A \in \mathcal{A}$ be arbitrary. The following holds:*

- (i) *If $A \in \mathcal{GP}_P(\mathcal{A})$, then $\nu(A) \in \mathcal{GI}_I(\mathcal{A})$;*
- (ii) *If $A \in \mathcal{GI}_I(\mathcal{A})$, then $\nu^-(A) \in \mathcal{GP}_P(\mathcal{A})$;*
- (iii) *If $A \in \mathcal{GP}_P(\mathcal{A})$, then $\lambda_A: A \rightarrow \nu^-\nu(A)$ is an isomorphism;*
- (iv) *If $A \in \mathcal{GI}_I(\mathcal{A})$, then $\sigma_A: \nu\nu^-(A) \rightarrow A$ is an isomorphism.*

In particular, the restriction $\nu|_{\mathcal{GP}_P(\mathcal{A})}: \mathcal{GP}_P(\mathcal{A}) \rightarrow \mathcal{GI}_I(\mathcal{A})$ is an equivalence with quasi-inverse $\nu^-|_{\mathcal{GI}_I(\mathcal{A})}: \mathcal{GI}_I(\mathcal{A}) \rightarrow \mathcal{GP}_P(\mathcal{A})$.

Proof. Let $Q_{\bullet} = \cdots \xrightarrow{s_2} Q_1 \xrightarrow{s_1} Q_0 \xrightarrow{s_0} Q_{-1} \xrightarrow{s_{-1}} \cdots$ be an exact sequence as in the definition of Gorenstein P -projective. Applying ν gives an exact sequence

$$\nu(Q_{\bullet}) = \cdots \xrightarrow{\nu(s_2)} \nu(Q_1) \xrightarrow{\nu(s_1)} \nu(Q_0) \xrightarrow{\nu(s_0)} \nu(Q_{-1}) \xrightarrow{\nu(s_{-1})} \cdots$$

Since the components of Q_\bullet are P -projective, we have an isomorphism $\nu^- \nu(Q_\bullet) \cong Q_\bullet$, and therefore $\nu^- \nu(Q_\bullet)$ is exact. Also, the complex $\nu(Q_\bullet)$ has I -injective components since ν sends P -projective objects to I -injective objects. Hence, if $Z_0(Q_\bullet) = A$, then $Z_0(\nu(Q_\bullet)) = \nu(A) \in \mathcal{GI}_I(\mathcal{A})$. This shows (i). Now consider the exact sequence $0 \rightarrow \nu(A) \rightarrow \nu(Q_0) \xrightarrow{\nu(s_0)} \nu(Q_{-1})$. Applying ν^- gives a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \longrightarrow & Q_0 & \xrightarrow{s_0} & Q_{-1} \\ & & \downarrow \lambda_A & & \downarrow \lambda_{Q_0} & & \downarrow \lambda_{Q_{-1}} \\ 0 & \longrightarrow & \nu^- \nu(A) & \longrightarrow & \nu^- \nu(Q_0) & \xrightarrow{\nu^- \nu(s_0)} & \nu^- \nu(Q_{-1}) \end{array}$$

where the lower row is exact since ν^- is left exact. Since λ_{Q_0} and $\lambda_{Q_{-1}}$ are isomorphisms, it follows that λ_A is an isomorphism. This proves part (iii). Part (ii) and (iv) are proved dually. \square

In the following, note that if $A \in \mathcal{GP}_P(\mathcal{A})$, then $L_i \nu(A) = 0$ for all $i > 0$. Dually, if $A \in \mathcal{GI}_I(\mathcal{A})$, then $R^i \nu^-(A) = 0$ for all $i > 0$.

Proposition 4.1.3. *Let $A \in \mathcal{A}$. Then $A \in \mathcal{GP}_P(\mathcal{A})$ if and only if the following holds:*

- (i) $L_i \nu(A) = 0$ for all $i > 0$;
- (ii) $R^i \nu^-(\nu(A)) = 0$ for all $i > 0$;
- (iii) The unit $\lambda_A: A \rightarrow \nu^- \nu(A)$ is an isomorphism.

Proof. Assume $A \in \mathcal{GP}_P(\mathcal{A})$, and let A_\bullet be the complex in the definition of Gorenstein P -projective. Consider the exact sequence

$$0 \rightarrow \text{Ker } s_1 \rightarrow A_1 \rightarrow A \rightarrow 0$$

Applying ν to this gives a long exact sequence

$$\cdots \rightarrow L_1 \nu(A_1) \rightarrow L_1 \nu(A) \rightarrow \nu(\text{Ker } s_1) \rightarrow \nu(A_1) \rightarrow \nu(A) \rightarrow 0$$

But by assumption the sequence $0 \rightarrow \nu(\text{Ker } s_1) \rightarrow \nu(A_1) \rightarrow \nu(A) \rightarrow 0$ is exact and $L_i \nu(A_1) = 0$ for all $i > 0$. Hence, it follows that $L_1 \nu(A) = 0$. Since $\text{Ker } s_1$ is also Gorenstein P -projective and $L_i \nu(\text{Ker } s_1) \cong L_{i+1} \nu(A)$ for all $i > 0$, it follows by induction that $L_i \nu(A) = 0$ for all $i > 0$. Since $\nu(A) \in \mathcal{GI}_I(\mathcal{A})$, a dual argument shows that $R^i \nu^-(\nu(A)) = 0$ for all $i > 0$. Finally, $\lambda_A: A \rightarrow \nu^- \nu(A)$ is an isomorphism by Proposition 4.1.2.

Assume A satisfies (i), (ii), and (iii). Choose an exact sequence $0 \rightarrow \nu(A) \rightarrow J_{-1} \rightarrow J_{-2} \rightarrow \cdots$ with J_i being I -injective for all i . Applying ν^- and using that $A \cong \nu^- \circ \nu(A)$ and $R^i \nu^-(\nu(A)) = 0$ for all i , we get an exact sequence

$$0 \rightarrow A \rightarrow \nu^-(J_{-1}) \rightarrow \nu^-(J_{-2}) \rightarrow \cdots$$

which is still exact when one applies ν . Now choose an exact sequence $\cdots \rightarrow Q_1 \rightarrow Q_0 \rightarrow A \rightarrow 0$ with Q_i being P -projective for all i , and consider

the exact sequence

$$Q_\bullet = \cdots \rightarrow Q_1 \rightarrow Q_0 \rightarrow \nu^-(J_{-1}) \rightarrow \nu^-(J_{-2}) \rightarrow \cdots$$

obtained by gluing the two sequence together at A . Since $L_i\nu(A) = 0$ for all $i > 0$, this sequence is still exact when one applies ν . Furthermore, it has P -projective components and satisfy $Z_0(Q_\bullet) = A$. This shows that $A \in \mathcal{GP}_P(\mathcal{A})$. \square

We have the following dual of Proposition 4.1.3.

Proposition 4.1.4. *Let $A \in \mathcal{A}$. Then $A \in \mathcal{GI}(\mathcal{A})$ if and only if the following holds:*

- (i) $R^i\nu^-(A) = 0$ for all $i > 0$;
- (ii) $L_i\nu(\nu^-(A)) = 0$ for all $i > 0$;
- (iii) The counit $\sigma_A: \nu\nu^-(A) \rightarrow A$ is an isomorphism.

Proposition 4.1.5. *The following holds:*

- (i) $\mathcal{GP}_P(\mathcal{A})$ is a resolving subcategory of \mathcal{A} ;
- (ii) $\mathcal{GI}(\mathcal{A})$ is a coresolving subcategory of \mathcal{A} .

Proof. This follows from the description of $\mathcal{GP}_P(\mathcal{A})$ and $\mathcal{GI}(\mathcal{A})$ in Proposition 4.1.3 and Proposition 4.1.4. \square

Hence, we can define the resolving dimension $\dim_{\mathcal{GP}_P(\mathcal{A})}(-)$ and the coresolving dimension $\dim_{\mathcal{GI}(\mathcal{A})}(-)$ as described in Subsection 2.3.

We end this subsection with the following result

Lemma 4.1.6. *The following holds:*

- (i) If $A_0 \xrightarrow{i} A_1 \rightarrow A_2 \rightarrow 0$ is an exact sequence where $A_0, A_1 \in \mathcal{GP}_P(\mathcal{A})$ and such that $\nu(i)$ is a monomorphism, then i is a monomorphism and $A_2 \in \mathcal{GP}_P(\mathcal{A})$.
- (ii) If $0 \rightarrow A_0 \rightarrow A_1 \xrightarrow{p} A_2$ is an exact sequence with $A_1, A_2 \in \mathcal{GI}(\mathcal{A})$ and such that $\nu^-(p)$ is an epimorphism, then p is an epimorphism and $A_0 \in \mathcal{GI}(\mathcal{A})$.

Proof. We prove part (i), part (ii) is proved dually. Since $\nu(i)$ is a monomorphism, it follows that $\nu^-\nu(i)$ is a monomorphism. Since $\nu^-\nu(i)$ corresponds to i under the isomorphisms $A_0 \cong \nu^-\nu(A_0)$ and $A_1 \cong \nu^-\nu(A_1)$, it follows that i is a monomorphism. The fact that $A_2 \in \mathcal{GP}_P(\mathcal{A})$ follows from Proposition 4.1.3. \square

4.2. Iwanaga-Gorenstein functors.

Definition 4.2.1. Assume P is a generating functor with Nakayama functor ν relative to P . Then P is called **Iwanaga-Gorenstein** if there exists an $m \geq 0$ such that $L_i\nu(A) = 0$ and $R^i\nu^-(A) = 0$ for all $A \in \mathcal{A}$ and $i > m$.

We have a simpler description of $\mathcal{GP}_P(\mathcal{A})$ and $\mathcal{GI}(\mathcal{A})$ when P is Iwanaga-Gorenstein.

Theorem 4.2.2. *Assume P is Iwanaga-Gorenstein. The following holds:*

- (i) $A \in \mathcal{GP}_P(\mathcal{A})$ if and only if $L_i\nu(A) = 0$ for all $i > 0$;
- (ii) $A \in \mathcal{GI}_I(\mathcal{A})$ if and only if $R^i\nu^-(A) = 0$ for all $i > 0$.

Proof. If $A \in \mathcal{GP}_P(\mathcal{A})$, then $L_i\nu(A) = 0$ for all $i > 0$ by Proposition 4.1.3. For the converse, fix a number m such that $L_i\nu = 0$ and $R^i\nu^- = 0$ for all $i > m$, and note that by Proposition 4.1.3 we only need to show that $R^i\nu^-(\nu(A)) = 0$ and $\lambda_A: A \rightarrow \nu^-\nu(A)$ is an isomorphism. To this end, choose an exact sequence

$$\cdots \xrightarrow{s_3} Q_2 \xrightarrow{s_2} Q_1 \xrightarrow{s_1} A \rightarrow 0$$

with Q_i being P -projective. Applying ν gives an exact sequence

$$\cdots \xrightarrow{\nu(s_3)} \nu(Q_2) \xrightarrow{\nu(s_2)} \nu(Q_1) \xrightarrow{\nu(s_1)} \nu(A) \rightarrow 0$$

since $L_i\nu(A) = 0$ for all $i > 0$. Also, since $\nu(Q_i)$ is I -injective, it follows by dimension shifting that $R^i\nu^-(\nu(A)) \cong R^{i+m}\nu^-(\text{Ker } \nu(s_m)) = 0$ and $R^i\nu^-(\text{Ker } \nu(s_j)) \cong R^{i+m}\nu^-(\text{Ker } \nu(s_{j+m})) = 0$ for all $i > 0$ and $j > 0$. Therefore, we have a commutative diagram

$$\begin{array}{ccccccc} \cdots & \xrightarrow{s_3} & Q_2 & \xrightarrow{s_2} & Q_1 & \xrightarrow{s_1} & A \longrightarrow 0 \\ & & \downarrow \lambda_{Q_2} & & \downarrow \lambda_{Q_1} & & \downarrow \lambda_A \\ \cdots & \xrightarrow{\nu^-\nu(s_3)} & \nu^-\nu(Q_2) & \xrightarrow{\nu^-\nu(s_2)} & \nu^-\nu(Q_1) & \xrightarrow{\nu^-\nu(s_1)} & \nu^-\nu(A) \longrightarrow 0 \end{array}$$

where the rows are exact. Hence, the morphism $\lambda_A: A \rightarrow \nu^-\nu(A)$ is an isomorphism. This proves part (i). Part (ii) is proved dually. \square

It follows from Theorem 4.2.2 that if P is Iwanaga-Gorenstein, then

$$\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) < \infty \quad \text{and} \quad \dim_{\mathcal{GI}_I(\mathcal{A})}(\mathcal{A}) < \infty.$$

Our goal is to prove that when P is Iwanaga-Gorenstein the following numbers are equal:

- 1) $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A})$;
- 2) $\dim_{\mathcal{GI}_I(\mathcal{A})}(\mathcal{A})$;
- 3) The smallest integer $s \geq 0$ such that $L_i\nu(A) = 0$ for all $i > s$ and $A \in \mathcal{A}$;
- 4) The smallest integer $t \geq 0$ such that $R^i\nu^-(A) = 0$ for all $i > t$ and $A \in \mathcal{A}$.

We also show that if the numbers in 1) or 2) are finite, then P is Iwanaga-Gorenstein. This generalizes the finite-dimensional case, see [3, Proposition 3.1 part b)] and [3, Proposition 4.2]. It is also analogous to other results in Gorenstein homological algebra, see for example [12, Theorem 2.28].

In order to prove this we need some preparation.

Lemma 4.2.3. *Let $A \in \mathcal{A}$. The following holds:*

- (i) $A \cong \nu(A')$ for some $A' \in \mathcal{A}$ if and only if there exists an exact sequence $J_0 \rightarrow J_1 \rightarrow A \rightarrow 0$ with J_0 and J_1 being I -injective;
(ii) $A \cong \nu^-(A')$ for some $A' \in \mathcal{A}$ if and only if there exists an exact sequence $0 \rightarrow A \rightarrow Q_0 \rightarrow Q_1$ with Q_0 and Q_1 being P -projective.

Proof. For any object $A' \in \mathcal{A}$ choose an exact sequence $Q_0 \rightarrow Q_1 \rightarrow A' \rightarrow 0$ with Q_0 and Q_1 being P -projective. By applying ν and using that it is right exact and sends P -projective objects to I -injective objects, we get one direction of part (i). For the converse, assume we have an exact sequence $J_0 \xrightarrow{s} J_1 \rightarrow A \rightarrow 0$ with J_0 and J_1 being I -injective. Since $\sigma_{J_i}: \nu\nu^-(J_i) \rightarrow J_i$ is an isomorphism, it follows that

$$A = \text{Coker } s \cong \text{Coker } \nu\nu^-(s) \cong \nu(\text{Coker } \nu^-(s)).$$

This proves part (i). Part (ii) is proved dually. \square

Let $\Omega := \text{Ker } \beta^{P+I}: \mathcal{A} \rightarrow \mathcal{A}$ and $\Sigma := \text{Coker } \alpha^{T+P}: \mathcal{A} \rightarrow \mathcal{A}$. We then have exact sequences

$$\begin{aligned} 0 \rightarrow \Omega(A) \rightarrow PI(A) \xrightarrow{\beta_A^{P+I}} A \rightarrow 0 \\ A \xrightarrow{\alpha_A^{T+P}} PT(A) \rightarrow \Sigma(A) \rightarrow 0 \end{aligned}$$

By Proposition 2.1.3 and Proposition 2.1.4 we have that $\Sigma \dashv \Omega$. Hence, for $r \geq 0$ the functor $\Omega^r \circ \nu^-$ is right adjoint to $\nu \circ \Sigma^r$.

Lemma 4.2.4. *Let $A \in \mathcal{A}$ and $r \geq 0$ be an integer. The following holds:*

- (i) *If $A \cong \nu\Sigma^r(A')$ for an object $A' \in \mathcal{A}$, then there exists an exact sequence*

$$J_0 \rightarrow J_1 \rightarrow \cdots \rightarrow J_{r+1} \rightarrow A \rightarrow 0$$

with J_i being I -injective;

- (ii) *If $A \in \mathcal{GP}_P(\mathcal{A})$, then $\nu\Sigma^r(A) \in \mathcal{GI}_I(\mathcal{A})$;*

- (iii) *If $A \in \mathcal{GI}_I(\mathcal{A})$, then $\Omega^r \nu^-(A) \in \mathcal{GP}_P(\mathcal{A})$.*

Proof. We prove (i). Consider the sequence

$$\begin{aligned} A' \xrightarrow{\alpha_{A'}^{T+P}} PT(A') \xrightarrow{s_0} PT\Sigma(A') \xrightarrow{s_1} \cdots \xrightarrow{s_{r-1}} PT\Sigma^{r-1}(A') \\ \xrightarrow{p_{r-1}} \Sigma^r(A') \rightarrow 0 \end{aligned} \quad (4.2.5)$$

where p_i is the canonical projection and s_i is the composite

$$PT\Sigma^i(A') \xrightarrow{p_i} \Sigma^{i+1}(A') \xrightarrow{\alpha_{\Sigma^{i+1}(A')}^{T+P}} PT\Sigma^{i+1}(A').$$

Note that $T(\alpha^{T+P}): T \rightarrow T \circ P \circ T$ is a split monomorphism by the triangle identities. Since $T = P \circ \nu$ and P is faithful by Lemma 3.2.4, it follows that

$\nu(\alpha^{T+P}): \nu \rightarrow \nu \circ P \circ T$ is a monomorphism. Hence, applying ν to (4.2.5) gives an exact sequence

$$0 \rightarrow \nu(A') \xrightarrow{\nu(\alpha_{A'}^{T+P})} IT(A') \xrightarrow{\nu(s_0)} IT\Sigma(A') \xrightarrow{\nu(s_1)} \dots \xrightarrow{\nu(s_{r-1})} IT\Sigma^{r-1}(A') \rightarrow A \rightarrow 0.$$

This proves part (i). Finally, note that if $A \in \mathcal{GP}_P(\mathcal{A})$, then $\Omega(A) \in \mathcal{GP}_P(\mathcal{A})$ since $\mathcal{GP}_P(\mathcal{A})$ is resolving, and $\Sigma(A) \in \mathcal{GP}_P(\mathcal{A})$ by Lemma 4.1.6. Since $\nu(A) \in \mathcal{GI}(\mathcal{A})$ and $\nu^-(A') \in \mathcal{GP}_P(\mathcal{A})$ for $A \in \mathcal{GP}_P(\mathcal{A})$ and $A' \in \mathcal{GI}(\mathcal{A})$ by Proposition 4.1.2, we get part (ii) and (iii). \square

We now prove the main result of this subsection.

Theorem 4.2.6. *The following are equivalent:*

- (i) P is Iwanaga-Gorenstein;
- (ii) $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) < \infty$;
- (iii) $\dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A}) < \infty$.

Moreover, if this holds, then the following numbers coincide:

- (a) $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A})$;
- (b) $\dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A})$;
- (c) The smallest integer s such that $L_i \nu(A) = 0$ for all $i > s$ and $A \in \mathcal{A}$;
- (d) The smallest integer t such that $R^i \nu^-(A) = 0$ for all $i > t$ and $A \in \mathcal{A}$.

If this common number is n , we say that P is n -**Gorenstein**.

Proof. We prove that $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) = \dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A})$ by showing that

$$\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) \leq n \quad \text{if and only if} \quad \dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A}) \leq n$$

for any number $n \geq 0$. This together with Theorem 4.2.2 shows the equivalence of statement (i), (ii), and (iii), as well as the equality of the numbers in (a), (b), (c), and (d).

First assume $n \geq 2$ and $\dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A}) \leq n$. Let $A \in \mathcal{A}$ and consider the exact sequence

$$0 \rightarrow \Omega^n(A) \xrightarrow{i_n} PI\Omega^{n-1}(A) \xrightarrow{s_{n-1}} \dots \xrightarrow{s_3} PI\Omega^2(A) \xrightarrow{s_2} PI\Omega(A) \xrightarrow{s_1} PI(A) \xrightarrow{\beta_A^{P-I}} A \rightarrow 0$$

where s_j is the composition

$$PI\Omega^j(A) \xrightarrow{\beta_{\Omega^j(A)}^{P-I}} \Omega^j(A) \xrightarrow{i_j} PI\Omega^{j-1}(A)$$

and i_j is the inclusion. By Lemma 4.2.3 part (ii) there exists an object $A' \in \mathcal{A}$ such that $\Omega^2(A) \cong \nu^-(A')$. This implies that

$$\Omega^n(A) \cong \Omega^{n-2} \nu^-(A').$$

By Lemma 4.2.4 part (i) and our assumption we know that $\nu\Sigma^{n-2}(A'') \in \mathcal{G}\mathcal{I}_I(\mathcal{A})$ for all $A'' \in \mathcal{A}$. Hence, by Lemma 4.2.4 part (iii) it follows that

$$\Omega^{n-2}\nu^-\nu\Sigma^{n-2}\Omega^{n-2}\nu^-(A') \in \mathcal{G}\mathcal{P}_P(\mathcal{A}).$$

By the triangle identities for $\nu \circ \Sigma^{n-2} \dashv \Omega^{n-2} \circ \nu^-$, we get that $\Omega^n(A) \cong \Omega^{n-2}\nu^-(A')$ is a direct summand of $\Omega^{n-2}\nu^-\nu\Sigma^{n-2}\Omega^{n-2}\nu^-(A')$. It follows that

$$\Omega^n(A) \in \mathcal{G}\mathcal{P}_P(\mathcal{A})$$

since $\mathcal{G}\mathcal{P}_P(\mathcal{A})$ is closed under direct summands. This shows that

$$\dim_{\mathcal{G}\mathcal{P}_P(\mathcal{A})}(\mathcal{A}) \leq n.$$

Now assume $\dim_{\mathcal{G}\mathcal{I}_I(\mathcal{A})}(\mathcal{A}) \leq 1$. By the argument above we know that $\dim_{\mathcal{G}\mathcal{P}_P(\mathcal{A})}(\mathcal{A}) \leq 2$. Let $A \in \mathcal{A}$ by arbitrary, and choose an exact sequence

$$0 \rightarrow \text{Ker } s \xrightarrow{i} Q_0 \xrightarrow{s} Q_1 \xrightarrow{p} A \rightarrow 0$$

with Q_0, Q_1 being P -projective. Since $\dim_{\mathcal{G}\mathcal{P}_P(\mathcal{A})}(\mathcal{A}) \leq 2$, we get that $\text{Ker } s \in \mathcal{G}\mathcal{P}_P(\mathcal{A})$. Consider the exact sequence $0 \rightarrow \text{Ker } s \xrightarrow{i} Q_0 \xrightarrow{q} \text{im } s \rightarrow 0$. Applying ν to this gives an exact sequence

$$\nu(\text{Ker } s) \xrightarrow{\nu(i)} \nu(Q_0) \xrightarrow{\nu(q)} \nu(\text{im } s) \rightarrow 0.$$

Hence, we have an epimorphism $\nu(\text{Ker } s) \xrightarrow{p'} \text{Ker } \nu(q) \rightarrow 0$. Since $\nu(\text{Ker } s) \in \mathcal{G}\mathcal{I}_I(\mathcal{A})$ and $\dim_{\mathcal{G}\mathcal{I}_I(\mathcal{A})}(\mathcal{A}) \leq 1$, it follows that $\text{Ker } \nu(q) \in \mathcal{G}\mathcal{I}_I(\mathcal{A})$. Furthermore, we have a commutative diagram

$$\begin{array}{ccccc} \text{Ker } s & \xrightarrow{i} & Q_0 & \xrightarrow{s} & Q_1 \\ \downarrow \lambda_{\text{Ker } s} & & \downarrow \lambda_{Q_0} & & \downarrow \lambda_{Q_1} \\ \nu^-\nu(\text{Ker } s) & \xrightarrow{\nu^-\nu(i)} & \nu^-\nu(Q_0) & \xrightarrow{\nu^-\nu(s)} & \nu^-\nu(Q_1) \end{array}$$

The vertical morphisms are isomorphisms by Proposition 4.1.2 part (iii).

Hence, the morphism $\nu^-\nu(\text{Ker } s) \xrightarrow{\nu^-\nu(i)} \nu^-\nu(Q_0)$ is the kernel of $\nu^-\nu(s)$. In particular, it is a monomorphism. On the other hand, $\nu^-\nu(i)$ is also equal to the composition

$$\nu^-\nu(\text{Ker } s) \xrightarrow{\nu^-(p')} \nu^-(\text{Ker } \nu(q)) \xrightarrow{\nu^-(j)} \nu^-\nu(Q_0)$$

where $j: \text{Ker } \nu(q) \rightarrow \nu(Q_0)$ is the inclusion. Since $\nu^-\nu(s) \circ \nu^-(j) = 0$ and $\nu^-(j)$ is a monomorphism, it follows that $\nu^-(p')$ is an isomorphism. Now consider the commutative diagram

$$\begin{array}{ccc}
\nu\nu^{-}\nu(\text{Ker } s) & \xrightarrow{\nu\nu^{-}(p')} & \nu\nu^{-}(\text{Ker } \nu(q)) \\
\downarrow \sigma_{\nu(\text{Ker } s)} & & \downarrow \sigma_{\text{Ker } \nu(q)} \\
\nu(\text{Ker } s) & \xrightarrow{p'} & \text{Ker } \nu(q)
\end{array}$$

Since the vertical maps and the upper horizontal map are isomorphisms, it follows that p' is an isomorphism. Hence, the exact sequence $0 \rightarrow \text{Ker } s \xrightarrow{i} Q_0 \xrightarrow{q} \text{im } s \rightarrow 0$ is still exact after one applies ν . By Lemma 4.1.6 it follows that $\text{im } s \in \mathcal{GP}_P(\mathcal{A})$. This implies that $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) \leq 1$, and since A was arbitrary we get that $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) \leq 1$.

Finally, we consider the case when $\dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A}) = 0$. This implies that ν^{-} is exact. Also, $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) \leq 1$ by the argument above. Let $A \in \mathcal{A}$ be arbitrary, and choose a right exact sequence

$$Q_0 \xrightarrow{s} Q_1 \xrightarrow{p} A \rightarrow 0$$

with Q_0, Q_1 being P -projective. Since ν^{-} is exact and ν is right exact, the sequence $\nu^{-}\nu(Q_0) \xrightarrow{\nu^{-}\nu(s)} \nu^{-}\nu(Q_1) \xrightarrow{\nu^{-}\nu(p)} \nu^{-}\nu(A) \rightarrow 0$ is exact. Hence we have a commutative diagram

$$\begin{array}{ccccccc}
Q_0 & \xrightarrow{s} & Q_1 & \xrightarrow{p} & A & \longrightarrow & 0 \\
\downarrow \lambda_{Q_0} & & \downarrow \lambda_{Q_1} & & \downarrow \lambda_A & & \\
\nu^{-}\nu(Q_0) & \xrightarrow{\nu^{-}\nu(s)} & \nu^{-}\nu(Q_1) & \xrightarrow{\nu^{-}\nu(p)} & \nu^{-}\nu(A) & \longrightarrow & 0
\end{array}$$

with right exact rows. Since λ_{Q_0} and λ_{Q_1} are isomorphisms, it follows that λ_A is an isomorphism. Since $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) \leq 1$, it follows by Lemma 4.2.3 part (ii) that $\nu(A) \in \mathcal{GI}(\mathcal{A})$, and hence $A \cong \nu^{-}\nu(A) \in \mathcal{GP}_P(\mathcal{A})$. Since A was arbitrary, we get that $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) = 0$.

The dual of the above argument shows that if $\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) \leq n$, then $\dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A}) \leq n$. Hence, it follows that

$$\dim_{\mathcal{GP}_P(\mathcal{A})}(\mathcal{A}) = \dim_{\mathcal{GI}(\mathcal{A})}(\mathcal{A}).$$

and we are done. \square

Note that in Example 3.2.8 the functor $P = f_! \circ f^*$ is 0-Gorenstein, and in Example 3.2.9 it is 1-Gorenstein.

5. FUNCTOR CATEGORIES

Let k be a commutative ring, let \mathcal{B} be a k -linear abelian category, and let \mathcal{C} be a small, k -linear, locally bounded, and Hom-finite category. Our

goal in this section is to show that there exists an adjoint pair $i_! \dashv i^*$ with Nakayama functor ν where

$$i^*: \mathcal{B}^{\mathcal{C}} \rightarrow \prod_{c \in \mathcal{C}} \mathcal{B}$$

is the evaluation functor.

5.1. Reminder on the Hom and tensor functor. Let k be a commutative ring. The *tensor product* of two k -linear categories \mathcal{D} and \mathcal{E} is a k -linear category $\mathcal{D} \otimes \mathcal{E}$ with objects pairs (D, E) with $D \in \mathcal{D}$ and $E \in \mathcal{E}$. The set morphisms between (D, E) and (D', E') is $\mathcal{D}(D, D') \otimes_k \mathcal{E}(E, E')$, and composition is given by $(h_1 \otimes g_1) \circ (h_2 \otimes g_2) = (h_1 \circ h_2) \otimes (g_1 \circ g_2)$. The identity at (D, E) is $1_{D \otimes E} = 1_D \otimes 1_E$.

Let \mathcal{C} be a small k -linear category. Recall that the Yoneda lemma gives a fully faithful functor $h_{\mathcal{C}}: \mathcal{C} \rightarrow \text{Mod-}\mathcal{C}$, where $\text{Mod-}\mathcal{C}$ is the category of right \mathcal{C} -modules. The image $h_{\mathcal{C}}(c) = \mathcal{C}(-, c)$ is a projective \mathcal{C} -module. A right \mathcal{C} -module M is called *finitely presented* if there exists an exact sequence

$$\bigoplus_{i=1}^m \mathcal{C}(-, c_i) \rightarrow \bigoplus_{j=1}^n \mathcal{C}(-, d_j) \rightarrow M \rightarrow 0$$

in $\text{Mod-}\mathcal{C}$ for objects $c_i, d_j \in \mathcal{C}$. The category of finitely presented right \mathcal{C} -modules is denoted by $\text{mod-}\mathcal{C}$. This is an additive category with cokernels.

Let \mathcal{B} be a k -linear abelian category, and let $\mathcal{B}^{\mathcal{C}}$ denote the category of k -linear functors from \mathcal{C} to \mathcal{B} . Since \mathcal{B} is finitely complete and cocomplete, it follows from chapter 3 in [20] that there exist functors

$$\begin{aligned} - \otimes_{\mathcal{C}} -: (\text{mod-}\mathcal{C}) \otimes \mathcal{B}^{\mathcal{C}} &\rightarrow \mathcal{B} \\ \text{Hom}_{\mathcal{C}}(-, -): (\text{mod-}\mathcal{C}^{\text{op}})^{\text{op}} \otimes \mathcal{B}^{\mathcal{C}} &\rightarrow \mathcal{B} \end{aligned}$$

satisfying the following:

- (i) The functor $- \otimes_{\mathcal{C}} F: \text{mod-}\mathcal{C} \rightarrow \mathcal{B}$ is right exact for all $F \in \mathcal{B}^{\mathcal{C}}$.
- (ii) There exists an isomorphism $\mathcal{C}(-, c) \otimes_{\mathcal{C}} F \cong F(c)$ natural in c and F .
- (iii) The functor $\text{Hom}_{\mathcal{C}}(-, F): (\text{mod-}\mathcal{C}^{\text{op}})^{\text{op}} \rightarrow \mathcal{B}$ is left exact for all $F \in \mathcal{B}^{\mathcal{C}}$.
- (iv) There exists an isomorphism $\text{Hom}_{\mathcal{C}}(\mathcal{C}(c, -), F) \cong F(c)$ natural in c and F .

In [20] $M \otimes_{\mathcal{C}} F$ and $\text{Hom}_{\mathcal{C}}(N, F)$ is called the *colimit of F indexed by M* and the *limit of F indexed by N* , respectively. Note that (i) and (ii) determines $- \otimes_{\mathcal{C}} -$ uniquely, and (iii) and (iv) determines $\text{Hom}_{\mathcal{C}}(-, -)$ uniquely. If $\mathcal{C} = k$ we get functors $- \otimes_k -: (\text{mod-}k) \otimes \mathcal{B} \rightarrow \mathcal{B}$ and $\text{Hom}_k(-, -): (\text{mod-}k)^{\text{op}} \otimes \mathcal{B} \rightarrow \mathcal{B}$, respectively.

Now let \mathcal{C}_1 and \mathcal{C}_2 be small k -linear categories. Assume $M \in \text{Mod-}(\mathcal{C}_1 \otimes \mathcal{C}_2^{\text{op}})$ satisfies $M(c_1, -) \in \text{mod-}\mathcal{C}_2^{\text{op}}$ and $M(-, c_2) \in \text{mod-}\mathcal{C}_1$ for all $c_1 \in \mathcal{C}_1$ and $c_2 \in \mathcal{C}_2$. We then have functors

$$M \otimes_{\mathcal{C}_1} -: \mathcal{B}^{\mathcal{C}_1} \rightarrow \mathcal{B}^{\mathcal{C}_2} \quad \text{and} \quad \text{Hom}_{\mathcal{C}_2}(M, -): \mathcal{B}^{\mathcal{C}_2} \rightarrow \mathcal{B}^{\mathcal{C}_1}$$

and an isomorphism

$$\mathcal{B}^{\mathcal{C}_2}(M \otimes_{\mathcal{C}_1} F, G) \cong \mathcal{B}^{\mathcal{C}_1}(F, \text{Hom}_{\mathcal{C}_2}(M, G)). \quad (5.1.1)$$

natural in $F \in \mathcal{B}^{\mathcal{C}_1}$, $G \in \mathcal{B}^{\mathcal{C}_2}$ and M . It follows that $M \otimes_{\mathcal{C}_1} -$ is left adjoint to $\text{Hom}_{\mathcal{C}_2}(M, -)$. Finally, by (3.23) in [20] we have an isomorphism

$$N \otimes_{\mathcal{C}_2}(M \otimes_{\mathcal{C}_1} F) \cong (N \otimes_{\mathcal{C}_2} M) \otimes_{\mathcal{C}_1} F \quad (5.1.2)$$

natural in $F \in \mathcal{B}^{\mathcal{C}_1}$, $N \in \text{mod } \mathcal{C}_2$ and M .

5.2. Properties of Hom-finite locally bounded categories. Here we use the same terminology as in [10].

Definition 5.2.1. Let \mathcal{C} be a small k -linear category.

- (i) \mathcal{C} is **locally bounded** if for any object $c \in \mathcal{C}$ there are only finitely many objects in \mathcal{C} mapping nontrivially in and out of c . This means that for each $c \in \mathcal{C}$ we have

$$\mathcal{C}(c, c') \neq 0 \quad \text{for only finitely many } c' \in \mathcal{C}$$

and

$$\mathcal{C}(c'', c) \neq 0 \quad \text{for only finitely many } c'' \in \mathcal{C};$$

- (ii) \mathcal{C} is **Hom-finite** if $\mathcal{C}(c, c') \in \text{proj } k$ for all $c, c' \in \mathcal{C}$.

Lemma 5.2.2. Let \mathcal{C} be a small, k -linear, locally bounded, and Hom-finite category. Assume that $M \in \text{Mod } \mathcal{C}$ satisfy

$$\begin{aligned} M(c) &\in \text{proj } k \quad \forall c \in \mathcal{C} \\ M(c) &\neq 0 \quad \text{for only finitely many } c \in \mathcal{C}. \end{aligned}$$

Then there exists an exact sequence

$$\cdots \rightarrow Q_2 \rightarrow Q_1 \rightarrow Q_0 \rightarrow M \rightarrow 0$$

where Q_i is a finitely generated projective right \mathcal{C} -module for all i .

Proof. Choose an epimorphism $p^c: k^{n_c} \rightarrow M(c) \rightarrow 0$ for each $c \in \mathcal{C}$ with $M(c) \neq 0$, where $n_c \in \mathbb{N}$. Via the adjunction in (5.1.1) with $\mathcal{C}_1 = k$, $\mathcal{C} = \mathcal{C}_2^{\text{op}}$ and $\mathcal{B} = \text{mod } k$ this corresponds to a morphism $\mathcal{C}(-, c) \otimes_k k^{n_c} \xrightarrow{p^c} M$ in $\text{Mod } \mathcal{C}$. The induced map

$$\bigoplus_{c \in \mathcal{C}, M(c) \neq 0} \mathcal{C}(-, c) \otimes_k k^{n_c} \xrightarrow{\bigoplus p^c} M$$

is then an epimorphism. Let K be the kernel of this map. Then $K(c') \neq 0$ for only finitely many $c' \in \mathcal{C}$ since the same holds for $\bigoplus_{c \in \mathcal{C}, M(c) \neq 0} \mathcal{C}(-, c) \otimes_k k^{n_c}$. Also, $K(c')$ is the kernel of the epimorphism

$$\bigoplus_{c \in \mathcal{C}, M(c) \neq 0} \mathcal{C}(c', c) \otimes_k k^{n_c} \xrightarrow{\bigoplus q^c} M(c')$$

and since $M(c') \in \text{proj } k$ and $\bigoplus_{c \in \mathcal{C}, M(c) \neq 0} \mathcal{C}(c', c) \otimes_k k^{n_c} \in \text{proj } k$, we get that $K(c') \in \text{proj } k$. Hence, K satisfies the same properties as M . We can therefore repeat this construction, which proves the claim. \square

In the following we set $D := \text{Hom}_k(-, k)$. For $B \in \mathcal{B}$ the functors

$$D(-) \otimes_k B: (\text{proj } k)^{\text{op}} \rightarrow \mathcal{B} \quad \text{and} \quad \text{Hom}_k(-, B): (\text{proj } k)^{\text{op}} \rightarrow \mathcal{B}$$

both send k to B . Hence, we get an isomorphism

$$D(V) \otimes_k B \cong \text{Hom}_k(V, B) \tag{5.2.3}$$

in \mathcal{B} when V is a finitely generated projective k -module.

Lemma 5.2.4. *Assume \mathcal{C} is a small, k -linear, locally bounded and Hom-finite category. Choose an object $c \in \mathcal{C}$. The following holds:*

- (i) $D(\mathcal{C}(c, -))$ is a finitely presented right \mathcal{C} -module;
- (ii) We have an isomorphism

$$D(\mathcal{C}(-, c)) \otimes_k B \cong \text{Hom}_k(\mathcal{C}(-, c), B)$$

in $\mathcal{B}^{\mathcal{C}}$ for all $B \in \mathcal{B}$;

- (iii) We have an isomorphism

$$\mathcal{C}(c, -) \otimes_k B \cong \text{Hom}_k(D(\mathcal{C}(c, -)), B)$$

in $\mathcal{B}^{\mathcal{C}}$ for all $B \in \mathcal{B}$.

Proof. Statement (i) follows from Lemma 5.2.2. Statement (ii) and (iii) follow from the isomorphism in (5.2.3). \square

5.3. Nakayama functor for $i_! \dashv i^*$. In this subsection we fix a small, k -linear, locally bounded, and Hom-finite category \mathcal{C} . Let $k(\text{ob-}\mathcal{C})$ be the category with the same objects as \mathcal{C} , and with morphisms

$$k(\text{ob-}\mathcal{C})(c_1, c_2) = \begin{cases} 0 & \text{if } c_1 \neq c_2, \\ k & \text{if } c_1 = c_2. \end{cases}$$

The functor category $\mathcal{B}^{k(\text{ob-}\mathcal{C})}$ is just a product of copies of \mathcal{B} , indexed over the objects of \mathcal{C} . Let $i: k(\text{ob-}\mathcal{C}) \rightarrow \mathcal{C}$ be the inclusion. We have functors

$$\begin{aligned} i_!: \mathcal{B}^{k(\text{ob-}\mathcal{C})} &\rightarrow \mathcal{B}^{\mathcal{C}} & i_!((B^c)_{c \in \mathcal{C}}) &= \bigoplus_{c \in \mathcal{C}} \mathcal{C}(c, -) \otimes_k B^c \\ i^*: \mathcal{B}^{\mathcal{C}} &\rightarrow \mathcal{B}^{k(\text{ob-}\mathcal{C})} & i^*(F) &= (F(c))_{c \in \mathcal{C}} \\ i_*: \mathcal{B}^{k(\text{ob-}\mathcal{C})} &\rightarrow \mathcal{B}^{\mathcal{C}} & i_*((B^c)_{c \in \mathcal{C}}) &= \prod_{c \in \mathcal{C}} \text{Hom}_k(\mathcal{C}(-, c), B^c). \end{aligned}$$

Note that the functors $i_!$ and i_* are well defined since \mathcal{C} is locally bounded. In fact, evaluating $i_!((B^c)_{c \in \mathcal{C}})$ and $i_*((B^c)_{c \in \mathcal{C}})$ on an object in \mathcal{C} gives a finite sum, and since limits are taken pointwise in $\mathcal{B}^{\mathcal{C}}$, it follows that

$$i_!((B^c)_{c \in \mathcal{C}}) = \bigoplus_{c \in \mathcal{C}} \mathcal{C}(c, -) \otimes_k B^c = \prod_{c \in \mathcal{C}} \mathcal{C}(c, -) \otimes_k B^c \tag{5.3.1}$$

and

$$i_*((B^c)_{c \in \mathcal{C}}) = \prod_{c \in \mathcal{C}} \text{Hom}_k(\mathcal{C}(c, -), B^c) = \bigoplus_{c \in \mathcal{C}} \text{Hom}_k(\mathcal{C}(-, c), B^c). \quad (5.3.2)$$

Also, $\text{Hom}_{\mathcal{C}}(\mathcal{C}(c, -), F) = F(c) = \mathcal{C}(-, c) \otimes_{\mathcal{C}} F$, and hence by (5.1.1) we get that i^* is right adjoint to $i_!$ and left adjoint to i_* . Finally, we have functors

$$\begin{aligned} \nu: \mathcal{B}^{\mathcal{C}} &\rightarrow \mathcal{B}^{\mathcal{C}} & \nu(F) &= D(\mathcal{C}) \otimes_{\mathcal{C}} F \\ \nu^-: \mathcal{B}^{\mathcal{C}} &\rightarrow \mathcal{B}^{\mathcal{C}} & \nu^-(F) &= \text{Hom}_{\mathcal{C}}(D(\mathcal{C}), F) \end{aligned}$$

where

$$\begin{aligned} (D(\mathcal{C}) \otimes_{\mathcal{C}} F)(c) &= D(\mathcal{C}(c, -)) \otimes_{\mathcal{C}} F \\ \text{Hom}_{\mathcal{C}}(D(\mathcal{C}), F)(c) &= \text{Hom}_{\mathcal{C}}(D(\mathcal{C}(-, c)), F). \end{aligned}$$

It follows from (5.1.1) that ν is left adjoint to ν^- .

Theorem 5.3.3. *The functor $\nu: \mathcal{B}^{\mathcal{C}} \rightarrow \mathcal{B}^{\mathcal{C}}$ is a Nakayama functor relative to $i_! \dashv i^*$*

Proof. For $(B^c)_{c \in \mathcal{C}} \in \mathcal{B}^{k(\text{ob-}\mathcal{C})}$ we have natural isomorphisms

$$\begin{aligned} \nu \circ i_!((B^c)_{c \in \mathcal{C}}) &= \nu\left(\bigoplus_{c \in \mathcal{C}} \mathcal{C}(c, -) \otimes_k B^c\right) \cong \bigoplus_{c \in \mathcal{C}} D(\mathcal{C}(-, c)) \otimes_k B^c \\ &\cong \bigoplus_{c \in \mathcal{C}} \text{Hom}_k(\mathcal{C}(-, c), B^c) \cong i_*((B^c)_{c \in \mathcal{C}}) \end{aligned}$$

where the second last isomorphism follows from Lemma 5.2.4 part (ii). Hence, $\nu \circ i_! \cong i_*$, and $\nu \circ i_!$ is therefore right adjoint to i^* . Also, we have natural isomorphisms

$$\begin{aligned} \nu^- \circ \nu \circ i_!((B^c)_{c \in \mathcal{C}}) &= \text{Hom}_{\mathcal{C}}(D(\mathcal{C}), \bigoplus_{c \in \mathcal{C}} \text{Hom}_k(\mathcal{C}(-, c), B^c)) \\ &\cong \bigoplus_{c \in \mathcal{C}} \text{Hom}_{\mathcal{C}}(D(\mathcal{C}), \text{Hom}_k(\mathcal{C}(-, c), B^c)) \\ &\cong \bigoplus_{c \in \mathcal{C}} \text{Hom}_k(D(\mathcal{C}(c, -)), B^c) \cong \bigoplus_{c \in \mathcal{C}} \mathcal{C}(c, -) \otimes_k B^c = i_!((B^c)_{c \in \mathcal{C}}). \end{aligned}$$

This gives an inverse to the unit map $\alpha_{i_!}^{\nu^- \circ \nu}: i_! \rightarrow \nu^- \circ \nu \circ i_!$. Hence, the claim follows. \square

See Example 3.2.8 and Example 3.2.9 for concrete choices of \mathcal{C} . In fact there are interesting examples even when \mathcal{C} has only one object. In particular, if $\mathcal{C} = \Lambda$ is a finite-dimensional algebra, then the theorem above applies to the module categories

$$\Lambda\text{-mod} = (k\text{-mod})^{\Lambda} \quad \text{and} \quad \Lambda\text{-Mod} = (k\text{-Mod})^{\Lambda}$$

see Example 3.2.5. More generally, if k is a commutative ring and Λ is a k -algebra which is finitely generated projective as a k -module, then Theorem

5.3.3 holds for $\Lambda\text{-Mod} = (k\text{-Mod})^\Lambda$, see Example 3.2.5. If in addition Λ' is a k -algebra, then the theorem also applies to

$$\Lambda' \otimes_k \Lambda\text{-Mod} = (\Lambda'\text{-Mod})^\Lambda$$

see Example 3.2.6. If we furthermore assume that k is coherent or Λ' is left coherent then it also holds for the categories of finitely presented modules

$$\Lambda\text{-mod} = (k\text{-mod})^\Lambda \quad \text{or} \quad \Lambda' \otimes_k \Lambda\text{-mod} = (\Lambda'\text{-mod})^\Lambda$$

see Example 3.2.5 and Example 3.2.6, respectively.

We now apply Theorem 4.2.6 to Example 3.2.7.

Theorem 5.3.4. *Let k be a commutative ring, and let \mathcal{C} be a small, k -linear, locally bounded, and Hom-finite category. Assume*

$$\sup_{c \in \mathcal{C}} (\text{proj. dim } D(\mathcal{C}(-, c))) < \infty \quad \text{and} \quad \sup_{c \in \mathcal{C}} (\text{proj. dim } D(\mathcal{C}(c, -))) < \infty$$

as left and right \mathcal{C} -modules respectively. Then

$$\sup_{c \in \mathcal{C}} (\text{proj. dim } D(\mathcal{C}(-, c))) = \sup_{c \in \mathcal{C}} (\text{proj. dim } D(\mathcal{C}(c, -))).$$

and this number is equal to the Gorenstein dimension of the functor $P = i_! \circ i^$ on $\mathcal{C}\text{-Mod}$.*

Proof. By assumption the functor $P = i_! \circ i^*$ on $\mathcal{C}\text{-Mod} = (k\text{-Mod})^\mathcal{C}$ is Iwanaga-Gorenstein. Hence, by Theorem 4.2.6 we have that P is n -Gorenstein for some $n \in \mathbb{N}$. It follows that

$$\begin{aligned} \text{Ext}_{\mathcal{C}\text{-Mod}}^n(D\mathcal{C}, -) \neq 0 \quad \text{and} \quad \text{Ext}_{\mathcal{C}\text{-Mod}}^i(D\mathcal{C}, -) = 0 \text{ for } i > n \\ \text{Tor}_n^{\mathcal{C}}(D\mathcal{C}, -) \neq 0 \quad \text{and} \quad \text{Tor}_i^{\mathcal{C}}(D\mathcal{C}, -) = 0 \text{ for } i > n. \end{aligned}$$

We therefore get that

$$\sup_{c \in \mathcal{C}} (\text{proj. dim } D(\mathcal{C}(-, c))) = n = \sup_{c \in \mathcal{C}} (\text{flat. dim } D(\mathcal{C}(c, -))).$$

On the other hand, by Lemma 5.2.2 there exists an n th syzygy of $D(\mathcal{C}(c, -))$ which is finitely presented. Since finitely presented flat modules are projective, it follows that $\text{flat. dim } D(\mathcal{C}(c, -)) = \text{proj. dim } D(\mathcal{C}(c, -))$. This proves the claim. \square

Remark 5.3.5. Locally bounded Hom-finite categories are one of the main object of study [10]. In [10, Theorem 4.6] they assume that \mathcal{C} has a Serre functor relative to k . This implies that the functor P on $\mathcal{C}\text{-Mod}$ is 0-Gorenstein.

REFERENCES

- [1] Maurice Auslander and Mark Bridger. *Stable module theory*. Memoirs of the American Mathematical Society, No. 94. American Mathematical Society, Providence, R.I., 1969.
- [2] Maurice Auslander and Ragnar-Olaf Buchweitz. The homological theory of maximal Cohen-Macaulay approximations. *Mém. Soc. Math. France (N.S.)*, (38):5–37, 1989. Colloque en l'honneur de Pierre Samuel (Orsay, 1987).

- [3] Maurice Auslander and Idun Reiten. Cohen-Macaulay and Gorenstein Artin algebras. In *Representation theory of finite groups and finite-dimensional algebras (Bielefeld, 1991)*, volume 95 of *Progr. Math.*, pages 221–245. Birkhäuser, Basel, 1991.
- [4] Michael Barr and Jon Beck. Homology and standard constructions. In *Sem. on Triples and Categorical Homology Theory (ETH, Zürich, 1966/67)*, pages 245–335. Springer, Berlin, 1969.
- [5] Apostolos Beligiannis and Henning Krause. Thick subcategories and virtually Gorenstein algebras. *Illinois J. Math.*, 52(2):551–562, 2008.
- [6] Francis Borceux. *Handbook of categorical algebra. 2*, volume 51 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1994. Categories and structures.
- [7] Ragnar-Olaf Buchweitz. Maximal Cohen-Macaulay modules and Tate-cohomology over Gorenstein rings. Unpublished manuscript, 1986.
- [8] Xiao-Wu Chen. Gorenstein homological algebra of artin algebras. Postdoctoral Report, USTC, 2010.
- [9] J. Climent Vidal and J. Soliveres Tur. Kleisli and Eilenberg-Moore constructions as parts of biadjoint situations. *Extracta Math.*, 25(1):1–61, 2010.
- [10] Ivo Dell’Ambrogio, Greg Stevenson, and Jan Stovicek. Gorenstein homological algebra and universal coefficient theorems. *Math. Z.*, 287(3-4):1109–1155, 2017.
- [11] Samuel Eilenberg and John C. Moore. Adjoint functors and triples. *Illinois J. Math.*, 9:381–398, 1965.
- [12] E. Enochs, S. Estrada, and J. R. Garcia-Rozas. Gorenstein categories and Tate cohomology on projective schemes. *Math. Nachr.*, 281(4):525–540, 2008.
- [13] Edgar E. Enochs and Overtoun M. G. Jenda. Gorenstein injective and projective modules. *Math. Z.*, 220(4):611–633, 1995.
- [14] Edgar E. Enochs and Overtoun M. G. Jenda. *Relative homological algebra. Volume 1*, volume 30 of *De Gruyter Expositions in Mathematics*. Walter de Gruyter GmbH & Co. KG, Berlin, extended edition, 2011.
- [15] Edgar E. Enochs and Overtoun M. G. Jenda. *Relative homological algebra. Volume 2*, volume 54 of *De Gruyter Expositions in Mathematics*. Walter de Gruyter GmbH & Co. KG, Berlin, 2011.
- [16] Sergei I. Gelfand and Yuri I. Manin. *Methods of homological algebra*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, second edition, 2003.
- [17] Henrik Holm. Gorenstein homological dimensions. *J. Pure Appl. Algebra*, 189(1-3):167–193, 2004.
- [18] Osamu Iyama. Representation theory of orders. In *Algebra—representation theory (Constanta, 2000)*, volume 28 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 63–96. Kluwer Acad. Publ., Dordrecht, 2001.
- [19] Peter Jørgensen. Existence of Gorenstein projective resolutions and Tate cohomology. *J. Eur. Math. Soc. (JEMS)*, 9(1):59–76, 2007.
- [20] G. M. Kelly. Basic concepts of enriched category theory. *Repr. Theory Appl. Categ.*, (10):vi+137, 2005. Reprint of the 1982 original [Cambridge Univ. Press, Cambridge; MR0651714].
- [21] Aaron D. Lauda. Frobenius algebras and ambidextrous adjunctions. *Theory Appl. Categ.*, 16:No. 4, 84–122, 2006.
- [22] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [23] Dawei Shen. A description of Gorenstein projective modules over the tensor products of algebras, 2016. arXiv:1602.00116.
- [24] Jan Stovicek. Derived equivalences induced by big cotilting modules. *Adv. Math.*, 263:45–87, 2014.
- [25] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.

- [26] Abraham Zaks. Injective dimension of semi-primary rings. *J. Algebra*, 13:73–86, 1969.

LABORATOIRE DE MATHÉMATIQUES D'ORSAY, UNIVERSITÉ PARIS-SUD, 91405 ORSAY,
FRANCE

E-mail address: `sondre.kvamme@u-psud.fr`