

Koszul duality between Higgs and Coulomb categories \mathcal{O}

BEN WEBSTER¹

Department of Pure Mathematics
University of Waterloo &
Perimeter Institute for Mathematical Physics
Waterloo, ON, Canada
Email: ben.webster@uwaterloo.ca

Abstract. We prove a Koszul duality theorem between the category of weight modules over the quantized Coulomb branch (as defined by Braverman, Finkelberg and Nakajima) attached to a group G and representation V and a category of G -equivariant D-modules on the vector space V . This is proven by relating both categories to an explicit, combinatorially presented category.

These categories are related to generalized categories \mathcal{O} for symplectic singularities. Letting $\mathcal{O}_{\text{Coulomb}}$ and $\mathcal{O}_{\text{Higgs}}$ be these categories for the Coulomb and Higgs branches associated to V and G , we obtain a functor $\mathcal{O}_{\text{Coulomb}}^! \rightarrow \mathcal{O}_{\text{Higgs}}$ from the Koszul dual of one to the other. This functor is an equivalence in the special cases where the hyperkähler quotient of T^*V by G is a Nakajima quiver variety or a smooth hypertoric variety. This includes as special cases the parabolic-singular Koszul duality of category \mathcal{O} in type A, the categorified rank-level duality proposed by Chuang and Miyachi and proven by Shan, Vasserot and Varagnolo, and the hypertoric Koszul duality proven by Braden, Licata, Proudfoot and the author.

We also show that this equivalence intertwines the so-called twisting and shuffling functors. Together with the Koszul duality discussed above, this confirms the most important components of the symplectic duality conjecture of Braden, Licata, Proudfoot, and the author in this case.

1. INTRODUCTION

Let V be a complex vector space, and let G be a connected reductive algebraic group with a fixed faithful linear action on V . Attached to these data, we have two interesting spaces, which physicists call the **Higgs** and **Coulomb** branches (of the associated 3-dimensional $\mathcal{N} = 4$ supersymmetric gauge theory):

- The Higgs branch is well-known to mathematicians: it is given by an algebraic symplectic reduction of the cotangent bundle T^*V . That is, we have

$$\mathfrak{M}_H := \mu^{-1}(0) // G = \text{Spec}(\mathbb{C}[\mu^{-1}(0)]^G)$$

where $\mu: T^*V \rightarrow \mathfrak{g}$ is the moment map.

¹Supported by the NSF under Grant DMS-1151473, the Alfred P. Sloan Foundation and an NSERC Discovery Grant. This research was supported in part by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported in part by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Colleges and Universities.

- The Coulomb branch \mathfrak{M}_C has only been precisely defined in a recent series of papers by Nakajima, Braverman and Finkelberg [Nak16; BFN18]. It is defined as the spectrum of a ring, which we denote by $\mathcal{A}_0^{\text{sph}}$ in Section 3, constructed as a convolution algebra in the homology of the affine Grassmannian. The choice of representation V is incorporated as certain “quantum corrections” to convolution in homology, which are kept track of by an auxiliary vector bundle. To readers unhappy with the terms that appear in the sentences above: in order to prove our results, we will use what we believe to be the first purely algebraic description of the Coulomb branch for general G and V ; the geometric description given above will only be used to show that this algebraic presentation is correct, so readers can safely set the affine Grassmannian to one side if they desire.

Both of these spaces have Poisson structures. Bellamy has recently confirmed that Coulomb branches have symplectic singularities [Bel23], and they always possess a \mathbb{C}^\times -action with respect to which the Poisson structure has degree -2. The pair (G, V) is called **good** if this action is conical, that is, only the constant functions have degree ≤ 0 . As discussed in [BFN18, §2(iii)], whether a theory is good is easily tested combinatorially. Many pairs are good, but pairs that are not good are common as well; in general, larger representations are more likely to be good.

There are also many interesting cases where the Higgs branch also defines a conical symplectic singularity. In contrast to the Coulomb case, the Higgs branch is always conical, but its scheme structure may be badly behaved: for example, the ring $\mathbb{C}[\mu^{-1}(0)]^G$ may not be radical.

Since they are often conical symplectic singularities, a conjecture of Braden, Licata, Proudfoot, and the author suggests a surprising relationship between these spaces: they should be *symplectic dual* [BLPW16, §10.1]. The construction of Higgs and Coulomb branches discussed above cover most of the examples discussed in [BLPW16, §10.2], with the important exception of the nilcones (and more generally, S3 varieties) of simple Lie algebras which are not type A.

This conjecture requires a number of different geometric and representation-theoretic properties, the most important of which is a Koszul duality between generalizations of the category \mathcal{O} over quantizations of these varieties. The existence of such a duality has been proven in several special cases (see [BLPW16, §10.2]), but in this paper, we will give a uniform construction. Since we are working in somewhat different generality from [BLPW16], we will not perfectly match the conjecture given there—as indicated, there are symplectic dual pairs which are not expected to arise from a representation (G, N) as above.

First, let us be a bit more precise about what we mean by Koszul duality. For each algebra A over a field \mathbb{k} graded by the nonnegative integers with A_0 finite-dimensional and semi-simple, we can define a Koszul dual $A^!$ which is a quadratic algebra with the

same properties (see Section 4.2). By [MOS09, Thm. 30], the Koszul complex yields an equivalence between the derived categories of these algebras if and only if A is Koszul in the usual sense. For a graded category \mathcal{C} equivalent to A -gmod for A as above, the category $\mathcal{C}^! \cong A^!$ -gmod only depends on \mathcal{C} up to canonical equivalence.

In order to construct category \mathcal{O} 's, we need to choose auxiliary data, which determine finiteness conditions: we must choose a flavor ϕ (a \mathbb{C}^\times -action on \mathfrak{M}_H with weight 1 on the symplectic form and commuting with the \mathbb{C}^\times -action induced by scaling on T^*V), and a stability parameter $\xi \in (\mathfrak{g}^*)^G$. Note that the choice of ξ allows us to define the GIT quotient $\mathfrak{M}_{H,\xi} = \mu^{-1}(0) //_{\xi} G$ with ξ as the stability condition. Taking the unique closed orbit in the closure of a semi-stable orbit defines a map $\mathfrak{M}_{H,\xi} \rightarrow \mathfrak{M}_H$. There are interesting cases where this is a resolution of singularities, although $\mathfrak{M}_{H,\xi}$ may not be smooth, and even if smooth, its map to \mathfrak{M}_H may not be dominant.

We'll define a geometric category \mathfrak{D}_g , which should be considered as version of modules over a natural quantization, constructed via Hamiltonian reduction of microlocal differential operators on T^*V as in Kashiwara and Rouquier [KR08, §2.5]. This category follows the approach of McGerty and Nevins [MN14a], by defining \mathfrak{D}_g as a quotient of the category of strongly G -equivariant D -modules on V , which we can then identify with a category of DQ-modules over a quantization in favorable cases. See Section for more details.

Associated to the data of (G, V, ϕ, ξ) , we have two versions of category \mathcal{O} :

- (1) We let $\mathcal{O}_{\text{Higgs}}$ be the geometric category \mathcal{O} inside the category \mathfrak{D}_g , associated to the flavor ϕ . See Definition 2.44.
- (2) We let $\mathcal{O}_{\text{Coulomb}}$ be the algebraic category \mathcal{O} for the quantization of \mathfrak{M}_C defined by the flavor ϕ with integral weights. The element ξ induces an inner grading on this algebra which we use to define the category \mathcal{O} . See Definition 4.12.

There is a small asymmetry here, since one of these categories is a category of sheaves and the other a category of modules, but the difference is smaller than it may appear. We can compare algebraic and geometric category \mathcal{O} 's by taking sections of sheaves, and in favorable circumstances, the section functor will be an equivalence, generalizing the famous theorem of Beilinson and Bernstein [BB81]. We'll discuss this in more detail at the end of Section 2.9.

The category $\mathcal{O}_{\text{Higgs}}$ has an intrinsically defined graded lift $\tilde{\mathcal{O}}_{\text{Higgs}}$, which uses the category of mixed Hodge modules on V (Def. 2.58); the category $\mathcal{O}_{\text{Coulomb}}$ has a graded lift, for which we will give an explicit algebraic definition (Def. 4.7).

Two special cases will be of particular interest to us:

- (1) The group G and representation V correspond to a finite ADE or affine type A quiver gauge theory, so $\mathfrak{M}_{H,\xi}$ is a Nakajima quiver variety.
- (2) The group G is a torus and the representation V corresponds to a unimodular hyperplane arrangement, so $\mathfrak{M}_{H,\xi}$ is a smooth hypertoric variety.

Theorem A *There is a fully faithful functor $\tilde{\mathcal{O}}_{\text{Coulomb}}^! \rightarrow \tilde{\mathcal{O}}_{\text{Higgs}}$. In cases (1) and (2) above, if the variety $\mathfrak{M}_{H,\xi}$ is smooth, then this functor is an equivalence for every choice of ϕ .*

There is a general geometric property (\dagger) which assures the equivalences above. We expect this to hold in all cases where \mathfrak{M}_H is smooth and is proven in cases (1) and (2) in [Web17b] (see the discussion in Section 2.11), but at the moment, we lack the tools to prove it in full generality. For hypertoric varieties, Theorem A is proven in [Bra+12]. For the quiver cases, the connection to Coulomb branches was only recently made precise, so this version of the theorem was not proved before, but the results of [SVV14; Web17b] were very suggestive for the affine type A case; in particular, our work gives a new proof of the Chuang-Miyachi conjecture on Koszul duality between blocks of category \mathcal{O} on Cherednik algebras. Since the case of finite-type quiver varieties is the most novel and interesting case of this result, it is covered in more detail in [Kam+24]. It is also closely related to the study of shifted Yangians as in [KWWY14; Kam+19a]. In particular, it leads to a proof of the main conjectures of the latter paper [Kam+19b].

In Theorem A above, we have striven to make the simplest possible statement of this result, but we can weaken its hypotheses in various ways (see Theorem 4.15).

- In certain other cases, such as non-smooth hypertoric varieties, this functor is an equivalence onto a block of $\tilde{\mathcal{O}}_{\text{Higgs}}$ but there are other blocks.
- There is also a version of this theorem where we weaken the integrality hypothesis on ϕ ; the quantizations of \mathfrak{M}_C are parameterized by an affine space in which the cocharacters ϕ form a lattice. For more generic quantizations, we have an analogous functor from $\mathcal{O}_{\text{Coulomb}}^!$ to the category \mathcal{O} attached to a Higgs branch, but one associated to a subspace of V as a representation over a Levi of G . This phenomenon is a generalization of the theorem proved in [Web17c; Web19a] relating blocks of the Cherednik category \mathcal{O} to KLRW algebras. The details of this theory for quiver gauge theories are worked out in more detail by the author and collaborators in [Kam+24, Th. 1.4].

Theorem A depends on explicit calculation. For arbitrary (G, V, ϕ, ξ) , we give two presentations of the endomorphisms of the projective generators in $\mathcal{O}_{\text{Coulomb}}$. This in turn depends on a presentation of the algebra quantizing \mathfrak{M}_C itself. This presentation is simpler and easier to see if we consider the Coulomb branch in the context of a larger “extended category.” This category is natural from the geometric perspective of the affine Grassmannian and also seems to have a physical interpretation as a specific category of line defects in the associated gauge theory, with the original BFN algebra corresponding to a trivial defect.

Using a standard representation-theoretic construction, this presentation of the extended category gives us a presentation of the endomorphisms of projective generators

in $\mathcal{O}_{\text{Coulomb}}$. However, we wish to focus more on a second presentation that has the advantage of being graded, allowing us to define the graded lift $\tilde{\mathcal{O}}_{\text{Coulomb}}$. After this paper circulated as a preprint, H. Nakajima pointed out to us that the connection between these presentations has a geometric explanation, using the concentration map to the fixed points of a complexified cocharacter. This generalizes the work of Varagnolo and Vasserot [VV10, §2], which concerns the case of the adjoint representation in connection with double affine Hecke algebras. This will be explained in more detail in forthcoming work [Nak].

This second presentation also appears naturally in the Ext algebra of certain semi-simple G -equivariant D-modules on V , which makes the functor $\tilde{\mathcal{O}}_{\text{Coulomb}}^! \rightarrow \tilde{\mathcal{O}}_{\text{Higgs}}$ manifest. Instead of category \mathcal{O} , we may consider the category $\mathcal{W}_{\text{Coulomb}}$ of all integral weight modules, which has a graded lift $\tilde{\mathcal{W}}_{\text{Coulomb}}$, defined using the same presentation. We obtain a fully faithful functor $\tilde{\mathcal{W}}_{\text{Coulomb}}^! \rightarrow \mathcal{D}\text{-mod}$ to the category of strongly G -equivariant D-modules on V , independent of any properties of V or G . The functor $\tilde{\mathcal{O}}_{\text{Coulomb}}^! \rightarrow \tilde{\mathcal{O}}_{\text{Higgs}}$ is induced by this functor, and the hypertoric or quiver hypothesis is only needed to ensure that the quotient functor from $\mathcal{D}(V/G)\text{-mod}$ to modules over the quantization of $\mathfrak{M}_{H,\xi}$ has the correct properties.

This geometric interpretation shows a remarkable positivity in the structure of the weight representations of the Coulomb branch, based on the Hodge theory of the D-modules discussed above. In particular, using the standard relation between mixed categories and canonical bases axiomatized in [Web15], we obtain the result:

Theorem B (Cor. 4.8) *The classes of simple modules in $\tilde{\mathcal{W}}_{\text{Coulomb}}$ correspond to a dual canonical basis in its Grothendieck group.*

This result generalizes the Kazhdan-Lusztig conjecture for type A and Rouquier’s conjecture for category \mathcal{O} for the Cherednik algebra [Rou08b, §6.5]. More precisely, it extends these results to the categories of Gelfand-Tsetlin modules and Dunkl-Opdam modules, respectively, which correspond to weight modules under the isomorphisms of $U(\mathfrak{gl}_n)$ and $\mathbf{H}^{\text{sph}}(G(r, 1, \ell))$ to Coulomb branches shown in [WWY20] and [KN18]. Since making these connections carefully is a calculation of non-trivial length, we leave discussion of these connections to other work [Kam+19b; Web20; Web19a].

Thus, Theorem A can be strengthened to give not just an equivalence between these categories, but in fact a combinatorial description of both of them. The algebras that appear are an interesting generalization of KLRW algebras. Considering the richness of the theory developed around KLR algebras, there is reason to think that these new algebras will also prove quite interesting from the perspective of combinatorial representation theory.

In this paper, we will mainly limit our attention to studying the representation theory of these algebras in characteristic 0, but a similar approach can be applied over a field

of characteristic p . In this case, there is a natural relationship between quantizations, tilting bundles, and coherent sheaves, which we consider in more detail in [Web; Web22].

Because of the nature of our proof of Theorem A, it is easily extended to show that these equivalences are compatible with certain natural autoequivalences of derived categories, called shuffling and twisting functors. See [BLPW16, §8] for more on these functors.

Theorem C *Under the hypothesis (\dagger) , the functor of Theorem A induces an equivalence of graded derived categories $D^b(\tilde{\mathcal{O}}_{\text{Coulomb}}) \rightarrow D^b(\tilde{\mathcal{O}}_{\text{Higgs}})$ which intertwines twisting functors with shuffling functors and vice versa.*

This verifies two of the most important predictions of the conjecture that the Higgs and Coulomb branches of a single theory are symplectic dual to each other in the sense of [BLPW16, Def. 10.1]; it remains to confirm the more geometric aspects of this duality, such as a bijection between special strata.

Acknowledgements. We would like to thank Hiraku Nakajima for pointing out the connection to Varagnolo and Vasserot’s past work, as well as his forthcoming work. Many thanks to Tom Braden, Alexander Braverman, Kevin Costello, Tudor Dimofte, Joel Kamnitzer, Anthony Licata, Nick Proudfoot, Alex Weekes, and Oded Yacobi for many useful discussions on these topics.

2. THE HIGGS SIDE

Let us begin by setting up the basic data we use to define and analyze the Higgs and Coulomb sides of our construction. Let V be a complex vector space and let G be a connected reductive algebraic group with a fixed linear action on V ; we will call G the **gauge group** and V the **matter representation** following the standard practice in physics. Throughout, we will let d denote the dimension of V . Let Δ_0 be the set of roots of the corresponding Lie algebra.

Let $H = N_{GL(V)}^\circ(G)$ be the connected component of the normalizer of G in $GL(V)$ and $C = C_H(G) = \text{Aut}_G(V)$. There is a natural map of $H \rightarrow \text{Aut}(G)$ given by conjugation. The group C is the kernel of this map and the product GC surjects to $\text{Inn}(G)$, the group of inner automorphisms. This shows that H/GC injects into the outer automorphism group of G . Since H/GC is connected and the latter group is finite, this shows that $H = GC$ is generated by the connected subgroups G and C . Note that C has a natural action on each multiplicity space $M_\lambda = \text{Hom}(V_\lambda, V)$ where V_λ ranges over representatives of \hat{G} , the isomorphism classes of finite-dimensional G -irreps. This induces an isomorphism $C = \prod_{\lambda \in \hat{G}} GL(M_\lambda)$.

We choose a compact real form $G_{\mathbb{R}}$ of G . The representation V carries a $G_{\mathbb{R}}$ -invariant inner product. If we fix a $G_{\mathbb{R}}$ -invariant inner product on V_λ for each λ , this is equivalent to choosing such an inner product on each multiplicity space; these are related

by requiring $V \cong \oplus_\lambda M_\lambda \otimes V_\lambda$ to be an isometry of Hilbert spaces. We let $H_{\mathbb{R}}$ be the subgroup of H preserving this inner product. This is generated by $G_{\mathbb{R}}$ and the orthogonal groups in $GL(M_\lambda)$ with respect to our chosen inner product. Note that this shows that any two choices of $H_{\mathbb{R}}$ for different inner products are conjugate in H , and this conjugacy is unique up to inner automorphism (since the same is true for maximal compact subgroups of $GL(M_\lambda)$). We let $F = H/G$, and call this the **flavor group**.

We let $\tau: \mathbb{C}^\times \rightarrow GL(T^*V)$ be the cocharacter that acts trivially on V and with weight -1 on each cotangent fiber T_v^*V . This induces an action of the product $\tilde{H} = H \times \mathbb{C}^\times$ on T^*V , with the first factors acting by the unique symplectic action extending their action on V . Let $\tilde{F} = \tilde{H}/G$. Note that the usual symplectic form Ω transforms under the character $\nu: H \times \mathbb{C}^\times \rightarrow \mathbb{C}^\times$ given by $(g, t) \cdot \Omega = \nu(g, t)\Omega = t\Omega$.

We let T be a fixed maximal torus of G , and let $T_{\tilde{H}}$ be a maximal torus of \tilde{H} which contains T ; this induces tori $T_{\tilde{F}}, T_H, T_C$, etc. of the other groups we have considered. Letting \dagger stand for one of these groups G, H, F , etc., we let \mathfrak{t}_\dagger be the corresponding Lie algebra, with $\mathfrak{t}_{\dagger, \mathbb{Z}}$ denoting the derivatives of cocharacters of T_\dagger and $\mathfrak{t}_{\dagger, \mathbb{Q}}, \mathfrak{t}_{\dagger, \mathbb{R}}, \mathfrak{t}_{\dagger, \mathbb{C}}$ the span of these over the corresponding fields.

We'll also want to consider the subgroup $Q \subset H$ generated by G and the maximal torus T_H fixed earlier. This is the same as the preimage in H of the maximal torus $T_F \subset F$. More explicitly, this means that we only consider elements of H that preserve a particular decomposition of V into simples, given by the weight spaces of T_C . This group is denoted \tilde{G}' in [BFN18, S 3(ix)].

We fix a homomorphism $\phi: \mathbb{C}^\times \cong \mathbb{T} \rightarrow T_{\tilde{F}}$ such that $\phi(t) \cdot \Omega = t\Omega$; of course, any such cocharacter $\mathbb{T} \rightarrow \tilde{F}$ can be conjugated to have image in $T_{\tilde{F}}$. Let

$$\tilde{G} = \{(h, t) \in \tilde{H} \times \mathbb{C}^\times \mid hG = \phi(t) \in \tilde{F}\}.$$

Note that $\phi(t) = (\phi_0(t), t)$ for some $\phi_0: \mathbb{C}^\times \rightarrow T_F$. However, we prefer not to think of these components separately so as to emphasize that this splitting induced by the isomorphism $\tilde{H} \cong H \times \mathbb{C}^\times$ is not unique. For example, Braverman-Finkelberg-Nakajima [BFN18] prefer to use a different splitting, which is only defined on the level of Lie algebras; they use the splitting $\phi_0(t) \mapsto (t^{1/2}\phi_0(t), t)$, so it becomes symplectic. This only becomes a well-defined map of groups if we pass to a double cover, which is part of why we prefer to avoid it, along with the desire to avoid a blizzard of $1/2$'s appearing in formulas. However, our less symmetric choice will generate combinatorial complications of its own.

It would be more faithful to our original inspiration in physics to consider T^*V as a symplectic representation, without necessarily fixing an invariant Lagrangian subspace. Indeed, if we choose representations $V \not\cong V'$ such that $T^*V \cong T^*V'$, then our constructions will generally be equivalent via Fourier transform equivalences.

2.1. Lifts and chambers. In this section, we make some combinatorial definitions needed in order to understand this category \mathcal{O} .

The cocharacter ϕ has target \tilde{F} ; thus it naturally acts on any quotient of a \tilde{H} -space by G , but not on the vector space V itself. In order to act on V , we must choose a cocharacter into \tilde{H} whose composition with the projection is ϕ . For simplicity, we will only consider such lifts to the torus $T_{\tilde{H}}$, since any other lift will be conjugate to one of these. The set

$$\mathfrak{t}_{1,\mathbb{Z}} = \{\gamma: \mathbb{C}^\times \rightarrow T_{\tilde{H}} \mid \pi_{\tilde{H} \rightarrow \tilde{F}}(\gamma) = \phi\}$$

of such lifts is a torsor for the cocharacter lattice $X_*(T)$.

By differentiating, we can view these lifts as elements of the Lie algebra $\mathfrak{t}_{\tilde{H}}$ satisfying an integrality condition, as the notation above suggests; they are the intersection of the cocharacter lattice $\mathfrak{t}_{\tilde{H},\mathbb{Z}}$ with the preimage of the derivative of ϕ . If we consider affine linear combinations of these with coefficients in $\mathbb{C}, \mathbb{R}, \mathbb{Q}$, etc. we call these complex, real, rational, etc. lifts of ϕ , and use $\mathfrak{t}_{1,\dagger}$, etc. to denote the space of these lifts with $\dagger = \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$, etc.

Consider the action of T_H on V and choose a decomposition $V \cong \bigoplus_{i=1}^d V_i$ such that $d = \dim V$, $\dim V_i = 1$ and each V_i is invariant under T_H . Let φ_i be the weight of V_i over T_H . Note that each weight space for the subtorus T_C on V is an irreducible representation of G ; we define an equivalence relation on indices by $i \sim j$ if V_i and V_j have the same weight under T_C . Note that the Weyl group W of G acts naturally on the set of weights; we can upgrade this to a group action of W on the set $[1, d]$ such that $\varphi_{w \cdot i} = w \cdot \varphi_i$; this action is not canonical (for example, it cannot come from choosing the subspaces V_i so that $w \cdot V_i = V_{w \cdot i}$, as the adjoint representation of \mathfrak{sl}_n shows).

Example 2.i. We'll use the example of $G = GL(2)$ with $V \cong \mathbb{C}^2 \oplus \mathbb{C}^2$ as our standard example throughout. Let γ_1, γ_2 be the weights of the defining representation of $GL(2)$ on \mathbb{C}^2 . In this case, $d = 4$, with $\varphi_1 = \varphi_3 = \gamma_1$ and $\varphi_2 = \varphi_4 = \gamma_2$. The centralizer of $GL(2)$ is another copy of $GL(2)$, which acts by the vector representation on the multiplicity space $\text{Hom}_{GL(2)}(\mathbb{C}^2, V)$. The normalizer H is thus the image of $GL(2) \times GL(2)$ acting on the tensor product $\mathbb{C}^2 \otimes \mathbb{C}^2$. This is the same as the conformal orthogonal group acting on $\mathbb{C}^4 \cong \mathbb{C}^2 \otimes \mathbb{C}^2$, where \mathbb{C}^2 is endowed with a symmetric bilinear form making the coordinate vectors isotropic.

Choosing the usual torus in $F \cong PGL(2)$, we obtain the relation $1 \sim 2$ and $3 \sim 4$. We let ϕ be the cocharacter with weight -1 on the spaces V_{φ_1} and V_{φ_2} and weight 1 on the spaces V_{φ_3} and V_{φ_4} . The condition that $\phi(t) \cdot \Omega = t\Omega$ forces it to have weight 0 and -2 on the duals of these spaces. Thus, in this basis, $\widetilde{GL(2)} \cong GL(2) \times \mathbb{T}$ acts on V by the matrices

$$\left[\begin{array}{c|c} t^{-1}A & 0 \\ \hline 0 & tA \end{array} \right] \quad A \in GL(2), t \in \mathbb{T} \cong \mathbb{C}^\times$$

Our basic building blocks for category $\mathcal{O}_{\text{Higgs}}$ will be D-modules on V , to which we can apply Hamiltonian reduction. The construction of these begins with considering the D-modules given by pushforward of functions from subspaces. Of course, we must

choose these subspaces carefully to ultimately arrive at objects in category \mathcal{O} . We'll always want to consider subspaces of the following form:

Definition 2.1 For a sign sequence $\sigma \in \{+, 0, -\}^d$, we let V_σ be the sum of the subspaces V_i with $\sigma_i = +$. We let $(T^*V)_\sigma$ be the sum of V_i with $\sigma_i = +$ and V_i^* with $\sigma_i = -$.

If we have any other set \mathcal{I} equipped with a map $\iota: \mathcal{I} \rightarrow \{+, 0, -\}^d$, then we can denote $V_x = V_{\iota(x)}$ and $(T^*V)_x = (T^*V)_{\iota(x)}$ for each $x \in \mathcal{I}$. This notation leaves ι implicit, but in all the examples we will consider, this map will be unambiguous.

Definition 2.2 We call σ compatible with a Borel $\tilde{B} \subset \tilde{G}$ containing $T_{\tilde{G}}$ if $(T^*V)_\sigma$ is \tilde{B} -invariant. We let \mathcal{K} be the set of pairs of sign vectors in $\{+, 0, -\}^d$ and compatible Borels.

If we fix a preferred Weyl chamber \mathcal{C} of \tilde{G} (and thus a standard Borel \tilde{B}), we have a bijection of all the Weyl chambers with the Weyl group W of G (which is also the Weyl group of \tilde{G}), and thus can think of \mathcal{K} as a subset of $\{+, 0, -\}^d \times W$.

Example 2.ii. In our running example, if \tilde{B} is the standard Borel, then the non-compatible sign vectors are of the form

$$(-, +, *, *) \quad (-, 0, *, *) \quad (0, +, *, *) \quad (*, *, -, +) \quad (*, *, -, 0) \quad (*, *, 0, +).$$

If we consider the opposite Borel (the only other), then +'s and -'s exchange places.

Now, we let

$$(2.1) \quad \varphi_i^+ = \varphi_i \quad \varphi_i^- = -\varphi_i - \nu \quad \varphi_i^{\text{mid}} = \frac{1}{2}(\varphi_i^+ - \varphi_i^-) = \varphi_i + \frac{1}{2}\nu.$$

Together, φ_i^\pm give the weights of $\mathfrak{t}_{\tilde{H}}$ acting on T^*V . As we mentioned before, the papers of Braverman-Finkelberg-Nakajima [Nak16; BFN18] use the action on V with weights φ_i^{mid} .

Example 2.iii. In our example, $\mathfrak{t}_{\tilde{H}}$ is 4-dimensional and is identified with the diagonal matrices of the form $\text{diag}(a+s, b+s, a+t, b+t)$; passing to \mathfrak{t}_1 means considering these with $s = -1, t = 1$. The weights φ_i^+ are the entries of this diagonal matrix; the weights φ_i^- are the weights on V^* , which are the negatives of these weights minus ν .

One question we will often want to address is

Definition 2.3 For a sign sequence $\sigma \in \{+, -\}^d$, we let

$$c_\sigma = \{\gamma \in \mathfrak{t}_{1, \mathbb{Z}} \mid \varphi_i^{\sigma_i}(\gamma) \geq 0\}$$


$$C_\sigma = \{\gamma \in \mathfrak{t}_{1, \mathbb{R}} \mid \sigma_i \varphi_i^{\text{mid}}(\gamma) \geq 0\}$$

That is, these are the subsets of $\mathfrak{t}_{1, \mathbb{Z}}$ and $\mathfrak{t}_{1, \mathbb{R}}$ respectively where the sign of $\varphi_i^{\text{mid}}(\gamma)$ corresponds to σ_i . We let $C_{\sigma, w} = C_\sigma \cap w \cdot \mathcal{C}$ be the intersection of C_σ with the open

Weyl chamber attached to w , and similarly $c_{\sigma,w} = c_{\sigma} \cap w \cdot \mathcal{C}$. Note that if $C_{\sigma,w} \neq \emptyset$, then σ is compatible with wBw^{-1} .

We can extend this notation to sequences in $\{+, 0, -\}^d$ by requiring $\varphi_i^{\pm}(\gamma) \in (-\nu(\phi), 0)$ if $\sigma_i = 0$. Note that if we have $\phi \in \mathfrak{t}_{F,\mathbb{R}}$, then the chambers C_{σ} for all the sign vectors $\sigma \in \{+, -\}^d$ cover the whole space $\mathfrak{t}_{1,\mathbb{R}}$, with the only overlap between two chambers being a polytope in the intersection of the hyperplanes on which they have different sign. On the other hand, if $\nu(\phi) > 0$, then there will be a gap between chambers, as though they are separated by a fat hyperplane. If $\nu(\phi) < 0$, then the chambers will have larger overlaps, but we will not be interested in this case.

Example 2.iv. Thus, if we use a and b as our coordinates on $\mathfrak{t}_{1,\mathbb{R}}$, we obtain the hyperplane arrangement:



The side of a hyperplane carrying a fringe indicates the non-negative side (which thus includes the hyperplane itself).

Given a cocharacter $\gamma \in \mathfrak{t}_{\tilde{H},\mathbb{Z}}$, we let


$$V_{\gamma} = \{x \in V \mid \lim_{t \rightarrow 0} \gamma(t) \cdot x \text{ exists}\} \quad (T^*V)_{\gamma} = \{x \in T^*V \mid \lim_{t \rightarrow 0} \gamma(t) \cdot x \text{ exists}\}$$

be the sum of the non-negative weight spaces for γ . Using the notation above, we have $V_{\gamma} = V_{\sigma}$ for all $\gamma \in c_{\sigma}$.

The space $(T^*V)_{\gamma}$ is Lagrangian and thus the conormal to V_{γ} for any (integral) lift with $\nu(\gamma) = 1$; for a real or rational lift with $\nu(\gamma) > 0$, this space may be isotropic (and not Lagrangian) if $\gamma \notin c_{\sigma}$ for all $\sigma \in \{+, -\}^d$. Furthermore, $(T^*V)_{\gamma} = (T^*V)_{\gamma'}$ for lifts

γ and γ' if and only if both lie in c_σ for some $\sigma \in \{+, 0, -\}^d$, in which case, both are equal to $(T^*V)_\sigma$.

In the diagram below, we've marked the chambers corresponding to sign vectors in $\{+, -\}^d$ with V_σ (represented in terms of which coordinates are non-zero).



An interesting special case to consider is when φ_i is trivial on T_G ; it will never be trivial on T_H , since this group contains the scalar multiplication on V . In this case, the functions φ_i^+ , φ_i^- , φ_i^{mid} have constant values on $\mathfrak{t}_{1,\mathbb{R}}$ given by, so the sets c_σ, C_σ can only be non-empty for $\sigma_i = +$ if the weight of ϕ on V_i is ≥ 0 , and $\sigma_i = -$ if this weight is < 0 . Thus, in either case, the weight 0 will not contribute to our arrangement.

Example 2.v. For example, if we consider the adjoint representation $V = \mathfrak{sl}_2$ of $G = SL_2$ with basis $\{E, H, F\}$, then we have $H = Q = SL_2 \times \mathbb{C}^\times$, with the second factor acting by scalar multiplication. If we take ϕ_0 to be the identity map $\mathbb{C}^\times \rightarrow \mathbb{C}^\times$, then we can identify $\mathfrak{t}_{1,\mathbb{R}} \cong \mathbb{R}$ sending $a \in \mathbb{R}$ to the matrix $\text{diag}(1 + 2a, 1, 1 - 2a)$. Thus, we have

$$\varphi_1^{\text{mid}} = \frac{3}{2} + 2a \quad \varphi_2^{\text{mid}} = \frac{3}{2} \quad \varphi_3^{\text{mid}} = \frac{3}{2} - 2a$$

and the non-empty chambers C_σ correspond to the sign vectors:

$$\begin{aligned} C_{(+,+, -)} &= \{a \mid a \geq \frac{3}{4}\} & V_\gamma &= \text{span}(E, H) \\ C_{(+,+, +)} &= \{a \mid \frac{3}{4} \geq a \geq -\frac{3}{4}\} & V_\gamma &= \mathfrak{sl}_2 \\ C_{(-,+, +)} &= \text{span}\{a \mid a \leq -\frac{3}{4}\} & V_\gamma &= \text{span}(H, F) \end{aligned}$$

On the other hand if we take ϕ_0 to be the inverse map, then

$$\varphi_1^{\text{mid}} = -\frac{1}{2} + 2a \quad \varphi_2^{\text{mid}} = -\frac{1}{2} \quad \varphi_3^{\text{mid}} = -\frac{1}{2} - 2a$$

and the non-empty chambers C_σ correspond to the sign vectors:

$$\begin{aligned} C_{(+,-,-)} &= \left\{ a \mid a \geq \frac{1}{4} \right\} & V_\gamma &= \text{span}(E) \\ C_{(+,-,+)} &= \left\{ a \mid \frac{1}{4} \geq a \geq -\frac{1}{4} \right\} & V_\gamma &= \{0\} \\ C_{(-,-,+)} &= \left\{ a \mid a \leq -\frac{1}{4} \right\} & V_\gamma &= \text{span}(F) \end{aligned}$$

2.2. The Steinberg algebra and category. Throughout the rest of the paper, we will frequently use induction of group actions: given groups $\mathbb{G} \subset \mathbb{H}$, and a left \mathbb{G} -space X , we let $\mathbb{H} \times^{\mathbb{G}} X$ denote the quotient $(\mathbb{H} \times X)/\mathbb{G}$ by the action $g \cdot (h, x) = (hg^{-1}, gx)$. This is a fiber bundle with fiber X over \mathbb{H}/\mathbb{G} . Note that if we can \mathbb{G} -equivariantly embed X inside a \mathbb{H} -space Y , then we have a bundle embedding

$$\mathbb{H} \times^{\mathbb{G}} X \rightarrow \mathbb{H}/\mathbb{G} \times Y \quad (h, x) \mapsto (hG, hx)$$

which is an isomorphism to the subbundle $\{(hG, x' \mid h^{-1}x' \in X\} \subset \mathbb{H}/\mathbb{G} \times Y$.

For each pair $(\sigma, w) \in \mathcal{K}$, we have an attached G -space

$$(2.2) \quad X_{\sigma,w} = G \times^{wBw^{-1}} V_\sigma = \{(gwBw^{-1}, v) \in G/wBw^{-1} \times V \mid g^{-1}v \in V_\sigma\}$$

with the shorthand $X_\sigma = X_{\sigma,1}$, with the induced map $p_{\sigma,w}: X_{\sigma,w} \rightarrow V$ induced by projection to the second factor in the description of (2.2).

For each collection of these pairs $I \subset \mathcal{K}$, we can define a Steinberg variety by taking the fiber product of each pair of them over V :

$$(2.3) \quad \mathbb{X}_I := \bigsqcup_{\substack{(\sigma,w) \in I \\ (\sigma',w') \in I}} X_{\sigma,w} \times_V X_{\sigma',w'}$$

with a natural G action. As in the introduction, fix a field \mathbb{k} ; we assume for simplicity that its characteristic is large enough that $H_G^*(pt; \mathbb{k})$ is torsion free². Throughout, we'll let H_*^{BM} denote the Borel-Moore homology of a space with coefficients in \mathbb{k} .

Definition 2.4 *The G -equivariant Borel-Moore homology $H_*^{BM,G}(\mathbb{X}_I)$ equipped with its convolution multiplication is called the **Steinberg algebra** in [Sau13].*

Equivalently, we can think of the **Steinberg category** \mathcal{X}_I whose objects are elements of I and where morphisms $(\sigma', w') \rightarrow (\sigma, w)$ are given by $H_*^{BM,G}(X_{\sigma,w} \times_V X_{\sigma',w'})$, with composition given by convolution. The Steinberg algebra is simply the sum of all the morphisms in this category; modules over the Steinberg algebra are naturally equivalent

²This will hold for any field if $G = \prod_i GL_{n_i}$ and for fields of characteristic $\notin \{2, 3, 5\}$ for all G .

to the category of modules over the category \mathcal{X}_I (that is, functors from this category to the category of \mathbb{k} -vector spaces).

The homology $H_*^{BM,G}(\mathbb{X}_I)$ has its natural geometric grading, but the convolution multiplication is not homogeneous in the grading. Instead, we consider the grading where $H_i^{BM,G}(X_{\sigma,w} \times_V X_{\sigma',w'})$ has degree $-i + \frac{\dim X_{\sigma,w} + \dim X_{\sigma',w'}}{2}$. An easy way to remember this convention is that if $\sigma = \sigma', w = w'$, then this will place the fundamental class of the diagonal in degree 0.

Let $S = \text{Sym}(\mathfrak{t}_{\mathbb{k}}^*) = H_G^*(G/B; \mathbb{k}) \cong H_T^*(pt; \mathbb{k})$. Recall that $H_G^*(pt; \mathbb{k}) = S^W$. We'll also want to consider the analogous rings $S_Q = \text{Sym}(\mathfrak{t}_Q^*; \mathbb{k})$, $S_F = \text{Sym}(\mathfrak{t}_F^*; \mathbb{k})$.

The space $X_{\sigma,w}$ also has an action of Q , since if $(\sigma, w) \in \mathcal{K}$, then V_{σ} is invariant under wB_Qw^{-1} where $B_Q = BT_Q$ and

$$X_{\sigma,w} = Q \times {}^{wB_Qw^{-1}} V_{\sigma}.$$

We can also construct a **deformed Steinberg category** \mathcal{X}_I^Q on the same object set and morphisms $H_*^{BM,Q}(X_{\sigma,w} \times_V X_{\sigma',w'})$. The ring $H_*^{BM,Q}(\mathbb{X}_I)$ is a flat deformation of $H_*^{BM,G}(\mathbb{X}_I)$ over the base ring S_F .

Example 2.vi. If $V = 0$, then there is only the empty sign vector, and $X_{\emptyset} = 0$. Thus, $A_{\{\emptyset\}} = H_*^G(G/B \times G/B)$. This is isomorphic to $\text{Hom}_{S^W}(S, S) \cong M_{\#W \times \#W}(S^W)$ via the convolution action on S , which is a free module of rank $\#W$ over S^W .

Let us state some of the basic facts that we will need about this category. For a fixed pair $(\sigma, w), (\sigma', w')$, let $\mathbb{X}(v)$ be the subset of $X_{\sigma,w} \times_V X_{\sigma',w'}$ where the two flags have relative position v .

Lemma 2.5 *All the results below hold in \mathcal{X}_I , and in its deformation over S_F with S replaced by S_H .*

- (1) *Every object carries an action of the cohomology ring S . The action of S^W is central.*
- (2) *The subspace $\mathbb{X}(v)$ is isomorphic to a vector bundle over G/B . Thus, as a S -module under left or right multiplication, $H_*^{BM,G}(\mathbb{X}(v))$ is free of rank 1 generated by the fundamental class.*
- (3) *Thus, the homology $H_*^{BM,G}(X_{\sigma,w} \times_V X_{\sigma',w'})$ is a free S -module of rank $\#W$ and $X_{\sigma,w} \times_V X_{\sigma',w'}$ is equivariantly formal.*
- (4) *Let \mathcal{X}_I° be the category obtained after tensor product with the fraction field K of S^W . In \mathcal{X}_I° , the objects $(\sigma, w), (\sigma', w')$ are isomorphic if $V_{\sigma} \cap V^T = V_{\sigma'} \cap V^T$, and the endomorphisms of every object are isomorphic to*

$$K \otimes_{S^W} H_*^G(G/B \times G/B) \cong M_{\#W \times \#W}(K)$$

In particular, in $\mathcal{X}_I^{Q,\circ}$, all objects are isomorphic after base extension to the fraction field K_H of S_H .

- (5) There is a representation of \mathcal{X}_I given by $Y_X(\boldsymbol{\sigma}, w) = H_*^{BM,G}(X_{\boldsymbol{\sigma},w}) \cong S$, with the action by convolution product. If I has the property that $V_{\boldsymbol{\sigma}}^T = V_{\boldsymbol{\sigma}'}$ for all $\boldsymbol{\sigma}, \boldsymbol{\sigma}' \in I$, then this representation is faithful. Again, note that this property is automatic in \mathcal{X}_I^Q .

Proof. For simplicity in the proofs, we assume that $w = w' = 1$.

- (1) This action is given by the homology of the diagonal

$$H_*^{BM,G}(X_{\boldsymbol{\sigma},w}) \subset H_*^{BM,G}(X_{\boldsymbol{\sigma},w} \times_V X_{\boldsymbol{\sigma},w}).$$

- (2) This follows from [Sau13, Lem. 11(a)].

- (3) This follows from [Sau13, Lem. 12].

(4) This follows because in these cases, the quotient vector bundle $G \times^B (V_{\boldsymbol{\sigma}} / (V_{\boldsymbol{\sigma}} \cap V_{\boldsymbol{\sigma}'}))$ has non-zero Euler class, which is thus invertible in K . The pushforward $G \times^B (V_{\boldsymbol{\sigma}} \cap V_{\boldsymbol{\sigma}'}) \rightarrow X_{\boldsymbol{\sigma}}$ is thus an isomorphism on homology after tensor product with K , and similarly with $\boldsymbol{\sigma}$ and $\boldsymbol{\sigma}'$ reversed. In \mathcal{X}_I^Q , we just use the fact that $V^{TH} = 0$.

Finally, we need to calculate the endomorphism algebra. We have a unique B -equivariant projection map $V_{\boldsymbol{\sigma}} \rightarrow V_{\boldsymbol{\sigma}}^T$, and this induces a map $X_{\boldsymbol{\sigma}} \rightarrow G/B \times (V_{\boldsymbol{\sigma}}^T)$; pushforward by this map is an isomorphism after tensoring with K , and

$$H_*^{BM,G}(G/B \times G/B \times (V_{\boldsymbol{\sigma}}^T)) \cong H_*^{BM,G}(G/B \times G/B)$$

as algebras by the Thom isomorphism. Thus, we can reduce the case where $V = 0$.

- (5) This representation follows from the general construction of convolution product [CG97, p. 2.7.5], with

$$M_1 = X_{\boldsymbol{\sigma},w} \quad M_2 = X_{\boldsymbol{\sigma}',w'} \quad M_3 = pt \quad Z_{12} = X_{\boldsymbol{\sigma},w} \times_V X_{\boldsymbol{\sigma}',w'} \quad Z_{23} = X_{\boldsymbol{\sigma}',w'}.$$

It's enough to prove faithfulness after tensor product with K ; since all objects in \mathcal{X}_I° are isomorphic, this reduces to the fact that $H_*^G(G/B \times G/B)$ acts faithfully on S , which is clear from the isomorphism $H_*^G(G/B \times G/B) \cong \text{Hom}_{S^w}(S, S)$.

□

This category has a sheaf-theoretic interpretation as well. By [CG97, Thm. 8.6.7], we have that

$$H_*^{BM,G}(X_{\boldsymbol{\sigma},w} \times_V X_{\boldsymbol{\sigma}',w'}) \cong \text{Ext}^\bullet((p_{\boldsymbol{\sigma},w})_* \mathbb{k}_{X_{\boldsymbol{\sigma},w}}, (p_{\boldsymbol{\sigma}',w'})_* \mathbb{k}_{X_{\boldsymbol{\sigma}',w'}})$$

with convolution product matching Yoneda product. The argument in [CG97] shows that this can be enhanced to a dg-functor $\mathcal{X}_I \rightarrow D_{\text{dg}}^b(V)$, where \mathcal{X}_I is made into a dg-category by replacing $H_*^{BM,G}(X_{\boldsymbol{\sigma},w} \times_V X_{\boldsymbol{\sigma}',w'})$ with the Borel-Moore chain complex on $X_{\boldsymbol{\sigma},w} \times_V X_{\boldsymbol{\sigma}',w'}$.

Example 2.vii. If we return to our usual example of GL_2 acting on $\mathbb{C}^2 \times \mathbb{C}^2$, then the sheaves $(p_{\boldsymbol{\sigma},w})_* \mathbb{k}_{X_{\boldsymbol{\sigma},w}}$ from different chambers depend on the structure of the corresponding subspace:

- (1) The subspaces $(*, *, *, *)$, $(0, 0, *, *)$, $(0, 0, 0, 0)$ are $GL(2)$ subrepresentations, and so the bundles $X_{\sigma,w}$ are trivial in these cases, and the pushforward is the sum of two copies of the functions on the subspace.
- (2) The subspaces $(0, *, *, *)$, $(*, 0, *, *)$, $(0, 0, *, 0)$, $(0, 0, 0, *)$ are not $GL(2)$ -invariant, and the map from $X_{\sigma,w}$ to its image is modeled on the map from the total space of $\mathfrak{S}(-1)$ on \mathbb{P}^1 , mapping to \mathbb{C}^2 . The pushforward sheaf is $(p_{\sigma,w})_* \mathbb{k}_{X_{\sigma,w}}$ is the sum of the constant sheaf on the image (which is the subspace $(*, *, *, *)$ or $(0, 0, *, *)$) and the constant sheaf on the image of the exceptional locus (which is $(0, 0, *, *)$ or $(0, 0, 0, 0)$).
- (3) Finally, the cases $(0, *, 0, *)$, $(*, 0, *, 0)$ will both have image $\{(x, y, z, w) \mid xw - yz = 0\}$ and exceptional locus $(0, 0, 0, 0)$. Unlike the cases above, the pushforward will be simple, since the map $p_{\sigma,w}$ is small.

Note that these correspond to the 3 cases in [Web20, Cor. 3.8 & 4.13].

Of course, we can define the same space, algebra, or category when I is a set with a map to \mathcal{K} . The Steinberg category \mathcal{X}_I attached to a set with such a map is equivalent to the category attached to its image (so the corresponding algebras are Morita equivalent). Furthermore, the spaces $X_{\sigma,1}$ and $X_{w \cdot \sigma, w}$ are isomorphic through the action of any lift of w to \tilde{G} , so the graph of this isomorphism provides an isomorphism between the objects $(\sigma, 1)$ and $(w \cdot \sigma, w)$ in the Steinberg category.

Definition 2.6 We'll let $f \mapsto f^*$ denote the equivalence of the Steinberg category to its opposite induced by flipping the order of the tensor factors.

2.3. A presentation of the Steinberg category. We will give an explicit presentation of Steinberg algebras for certain sets which generalize both the KLR algebras of [KL09; Rou08a] and the hypertoric algebras of [Bra+10; Bra+12].

Consider the space $\mathfrak{t}_{1,\mathbb{R}}$ of real lifts of ϕ . We can think of the weights φ_i and roots α as affine functions on this affine space, and so their vanishing loci are hyperplanes.

Definition 2.7 The **matter hyperplane** H_i for φ_i a weight as before, and **Coxeter hyperplane** h_α for $\alpha \in \Delta_0$ are defined by

$$H_i = \{\gamma \mid \varphi_i^{\text{mid}}(\gamma) = 0\} \quad h_\alpha = \{\gamma \mid \alpha(\gamma) = 0\}.$$

We'll draw matter hyperplanes with solid lines and Coxeter hyperplanes with dotted lines in diagrams.

Definition 2.8 We let I (resp. I') be the set of sign vectors $\sigma \in \{+, -\}^d$ such that there exists a choice of flavor ϕ such that $c_{\sigma,1} \neq \emptyset$ (resp. $C_{\sigma,1} \neq \emptyset$); note that $I' \supseteq I$. For a fixed flavor ϕ , we denote the corresponding sets I_ϕ and I'_ϕ .

Example 2.viii. In our running example, we have

$$I'_\phi = I_\phi = \{(+, +, +, +), (+, -, +, +), (+, -, +, -), (-, -, +, +), (-, -, +, -), (-, -, -, -)\}.$$

These sets are not always equal; choose your favorite example of a polytope defined by equations with integral coefficients but containing no lattice points. For example, if \mathbb{C}^\times acts on \mathbb{C}^2 with weights 2 and 3, and ϕ is the usual scalar multiplication, then $I_\phi = \{(+, +), (-, -)\}$, but $I'_\phi = \{(+, +), (-, +), (-, -)\}$.

If $C_\sigma \neq \emptyset$, then there is a unique sign vector $w\sigma$ such that $C_{w\sigma} = w \cdot C_\sigma$. This is the unique permutation of σ such that each φ_i is switched with $\varphi_j = w\varphi_i$ such that $i \sim j$. This is well-defined since if $\varphi_i = \varphi_k$ and $i \sim k$, then these have the same sign (since $C_\sigma \neq \emptyset$). In particular, if $\sigma \in I'$, the translate $w\sigma$ is well-defined.

Definition 2.9 Given a pair (σ, σ') , we let $\varphi(\sigma, \sigma')$ be the product of the weights φ_i such that $\sigma_i = +$ and $\sigma'_i = -$. Given a triple $(\sigma, \sigma', \sigma'')$, we let $\varphi(\sigma, \sigma', \sigma'')$ be the product of the weights φ_i such that $\sigma_i = \sigma''_i = -\sigma'_i$.

Remark 2.10. It is a possibility that $\varphi_i = 0$ upon restriction to $\mathfrak{t}_{1, \mathbb{R}}$. However, note that in this case, if $\sigma, \sigma' \in I'_\phi$, then $\sigma_i = \sigma'_i$ is the sign of the weight of ϕ_0 on V_i . Thus, $\varphi_i = 0$ never appears as a factor in $\varphi(\sigma, \sigma')$ in this case.

However, there will be situations where we want to consider $\varphi_i \in I'_\phi$ and $\varphi'_i \in I''_\phi$, where ϕ_0 and ϕ'_0 have opposite signs on V_i . In this case, we can have $\varphi(\sigma, \sigma') = 0$. This will manifest in some of the proofs below as needing to consider the deformed category $\mathcal{X}_{I'}^Q$, since the action of Q on V will never have non-trivial weight vectors of weight 0. Note that exactly the same phenomenon appears in [Tel21, §3.4].




FIGURE 1. In our running example, the resulting arrangement of matter and Coxeter hyperplanes is given above.

Note that for $\sigma, \sigma', \sigma'' \in \{+, -\}^d$, these expressions are related by the equation

$$\varphi(\sigma, \sigma', \sigma'')\varphi(\sigma, \sigma'') = \varphi(\sigma, \sigma')\varphi(\sigma', \sigma'').$$

Let $\partial_\alpha(f) = \frac{s_\alpha \cdot f - f}{\alpha}$ be the usual BGG-Demazure operator on S .

Definition 2.11 We let A_I' denote the category with objects given by the sign vectors $\sigma \in I'$, and morphisms freely generated by

- The identity morphism $e(\sigma)$ on each object σ .
- A morphism $f: \sigma \rightarrow \sigma$ for each $f \in S$ and each object σ , which satisfy the usual relations of product in S .
- Wall-crossing elements $w(\sigma; \sigma'): \sigma' \rightarrow \sigma$.
- Elements $\psi_\alpha(\sigma): \sigma \rightarrow \sigma$ for roots α such that $s_\alpha \cdot \sigma = \sigma$.

subject to the “codimension 1” relations:

$$(2.4a) \quad w(\sigma'', \sigma')w(\sigma', \sigma) = \varphi(\sigma, \sigma', \sigma'')w(\sigma, \sigma'') \quad : \sigma \rightarrow \sigma''$$

$$(2.4b) \quad \mu w(\sigma', \sigma) = w(\sigma', \sigma)\mu \quad : \sigma \rightarrow \sigma'$$


$$(2.4c) \quad \psi_\alpha(\sigma)^2 = 0 \quad : \sigma \rightarrow \sigma$$

$$(2.4d) \quad \psi_\alpha(\sigma) \cdot \mu - (s_\alpha \mu) \cdot \psi_\alpha(\sigma) = \partial_\alpha(\mu) \quad : \sigma \rightarrow \sigma$$

with $\sigma, \sigma', \sigma'' \in I'$, $\mu \in \mathfrak{t}_k^*$ and $\alpha \in \Delta_0$ a root with $s_\alpha \cdot C_\sigma = C_\sigma$ and the “codimension 2” relations (2.4e–2.4g) below. We get one of these for every codimension 2 intersection of hyperplanes which forms a face of $C_{\sigma,1}$. There are 3 possible types of these intersections, depending on whether the system of roots vanishing on the face is of rank 2, 1 or 0.

- (1) The codimension 2 subspace is the intersection of 2 Coxeter hyperplanes h_α and h_β which form the simple roots of a rank 2 subsystem: No matter hyperplane contains this intersection, since if H_i did, then φ_i would have to lie in the span of α and β , so φ_i^\pm would also vanish on this subspace. It follows that the value of φ_i^{mid} lies in $\mathbb{Z} + 1/2$ and thus is certainly not 0.


In this case, for every chamber $C_{\sigma,1}$ adjacent to these hyperplanes, we have the usual Coxeter relations for $m = \alpha^\vee(\beta) \cdot \beta^\vee(\alpha)$:



$$(2.4e) \quad \underbrace{\psi_\alpha(\sigma)\psi_\beta(\sigma)\psi_\alpha(\sigma)\cdots}_{m \text{ times}} = \underbrace{\psi_\beta(\sigma)\psi_\alpha(\sigma)\psi_\beta(\sigma)\cdots}_{m \text{ times}}$$

- (2) The codimension 2 subspace lies in a single Coxeter hyperplane h_α : In this case, the codimension 2 subspace lies in some number of matter hyperplanes $H_{i_1}, \dots, H_{i_{m-1}}$,

ordered cyclically; note that some of these weights will be equal or multiples of each other, since the representation V might have weights of multiplicity > 1 . These are closed under action by s_α , and this action reverses the cyclic ordering so we have $s_\alpha \cdot H_{i_j} = H_{i_{m-j}}$. We label the chamber between $H_{i_{j-1}}$ and H_{i_j} on the positive side of h_α by σ_j (with the convention that $H_{i_0} = H_{i_m} = h_\alpha$). Note that some of these may be empty.



Going from σ_j to $s_\alpha \cdot \sigma_{m-j+1}$, there are two minimal length paths which go around the codimension 2 locus in the two opposite ways. These don't necessarily agree, but they differ up to "lower order terms."


$$(2.4f) \quad \mathbb{w}(\sigma_{m-j+1}, \sigma_1) \psi_\alpha(\sigma_1) \mathbb{w}(\sigma_1, \sigma_j) - \mathbb{w}(\sigma_{m-j+1}, \sigma_m) \psi_\alpha(\sigma_m) \mathbb{w}(\sigma_m, \sigma_j) \\ = \partial_\alpha(\varphi_1 \cdot s_\alpha(\varphi_2)) \mathbb{w}(\sigma_{m-j+1}, \sigma_j)$$

where

$$\varphi_1 = \varphi(\sigma_1, \sigma_j) = \varphi(s_\alpha \cdot \sigma_{m-j+1}, s_\alpha \cdot \sigma_m) \\ \varphi_2 = \varphi(\sigma_m, \sigma_{m-j+1}) = \varphi(s_\alpha \cdot \sigma_m, s_\alpha \cdot \sigma_1).$$

- (3) The codimension 2 subspace does not lie in a root hyperplane, and thus is the intersection of some number of matter hyperplanes. The resulting relation here is a consequence of (2.4a), but we include it for completeness. The relation (2.4a) says that the two paths that join the chambers opposite the codimension 2 subspace that go around it in the two opposite ways are equal.

If there are exactly two hyperplanes, we label the adjacent chambers π, ρ, σ, τ as shown.



We then have the relation

$$(2.4g) \quad \mathbb{w}(\pi, \rho) \mathbb{w}(\rho, \sigma) = \mathbb{w}(\pi, \tau) \mathbb{w}(\tau, \sigma)$$

We let A_P for P a set equipped with a map $\iota: P \rightarrow I'$ be the category with the morphisms

$$\mathrm{Hom}_{A_P}(p, p') := \mathrm{Hom}_{A_{I'}}(\iota(p), \iota(p')).$$

One particularly interesting case is when $P = I'_\phi$.

Definition 2.12 *Let A_I^Q be the deformed version of this category; this is the category where replace S with the larger polynomial algebra S_Q , and keep all other morphisms and relations the same. This is a S_F -algebra, and base change to \mathbb{C} , we recover A_I .*

Remark 2.13. Just as with the Steinberg algebra and category, we can equally well think of A as the algebra given by the sum of its Hom spaces, and we will sometimes refer to it as an algebra. For example, we use $A_I e(\boldsymbol{\sigma})$ to denote the principal left module $\boldsymbol{\sigma}' \mapsto \mathrm{Hom}(\boldsymbol{\sigma}, \boldsymbol{\sigma}')$.

The categories A_I and A_I^Q have a grading with respect to which the relations (2.4a–2.4g) are homogeneous:

$$\begin{aligned} \deg \mu &= 2 & (\mu \in \mathfrak{t}_T \text{ or } \mathfrak{t}_H) & & \deg e(\boldsymbol{\sigma}) &= 0 \\ \deg w(\boldsymbol{\sigma}; \boldsymbol{\sigma}') &= \deg \varphi(\boldsymbol{\sigma}, \boldsymbol{\sigma}') + \deg \varphi(\boldsymbol{\sigma}', \boldsymbol{\sigma}) & & & \deg \psi_\alpha(\boldsymbol{\sigma}) &= -2 \end{aligned}$$

Thus, S or S_H is given twice its usual grading, which of course, matches the homological grading on $H_T^*(pt)$. Note that $\deg w(\boldsymbol{\sigma}; \boldsymbol{\sigma}')$ is simply the number of weights with opposite signs in $\boldsymbol{\sigma}$ and $\boldsymbol{\sigma}'$.

Example 2.ix. In our running example, the resulting category is well known: we can represent the positive Weyl chamber as a pair of points on the real line giving the coordinates (a, b) . Since we are in the positive Weyl chamber $a > b$, there is no ambiguity. We cross a hyperplane when these points meet or when they cross $x = 1$ or $x = -1$. Thus, if we add red points at $x \in \{1, -1\}$, we'll obtain a bijection between chambers in I'_ϕ and configurations of points up to isotopy leaving the red points in place.

We'll represent morphisms $\boldsymbol{\sigma} \rightarrow \boldsymbol{\sigma}'$ by Stendhal diagrams (as defined in [Web17a, §4]) that match $\boldsymbol{\sigma}$ at the bottom and $\boldsymbol{\sigma}'$ at the top (with composition given by stacking, using isotopies to match the top and bottom if possible). We send the

- identity on $\boldsymbol{\sigma}$ to a diagram with all strands vertical,
- the action of $\mathbb{C}[\gamma_1, \gamma_2]$ to a polynomial ring placing dots on the two strands,
- $w(\boldsymbol{\sigma}; \boldsymbol{\sigma}')$ to a diagram with straight lines interpolating between the top and bottom
- $\psi_\alpha(\boldsymbol{\sigma})$ is only well-defined if there is no red line separating the two black lines; we send this to a crossing of the two black strands.





FIGURE 2. The correspondence between chambers and Stendhal diagrams



The relations (2.4a–2.4g) exactly match those of \tilde{T}_{-2}^2 as defined in [Web16, Def. 2.3] (a special case of the algebras defined in [Web17a, §4]). Note the notation mismatch with our usual notation of T for a maximal torus of G . Thus, A_{I_ϕ} is equivalent to the category of projective modules over this category. This is a special case of a much more general result, which we will discuss in Section 2.5.

Given a simple root α_i , let P_i be the unique minimal parabolic containing the Borel B and the root SL_2 for α_i . Let $G_i \subset G/B \times G/B$ be the preimage of the diagonal in $G/P_i \times G/P_i$. Given a P_i -representation Q , we let $L_{P_i}(Q)$ be the pullback of the associated bundle on G/P_i to G_i , and if Q is a representation of the Borel, then let $L(Q)$ be the associated vector bundle on the diagonal inside $G/B \times G/B$. Note that a sign vector $\sigma \in I'$ satisfies $s_{\alpha_i} \cdot \sigma = \sigma$ if and only if V_σ is a subrepresentation for P_i .

Theorem 2.14 *We have a natural equivalence $A_{I'} \cong \mathcal{X}_{I'}$ and $A_{I'}^Q \cong \mathcal{X}_{I'}^Q$ which matches objects in the obvious way, and sends*

- (1) $\mu: \sigma \rightarrow \sigma$ to the Euler class $e(L(\mu))$ of the associated bundle on the diagonal copy of X_σ in $\mathbb{X}_{I'}$.

(2) $\mathbb{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}')$ to the fundamental class of the variety $[L(V_{\boldsymbol{\sigma}} \cap V_{\boldsymbol{\sigma}'})]$ embedded naturally as

$$L(V_{\boldsymbol{\sigma}} \cap V_{\boldsymbol{\sigma}'}) \cong \{(gB, gB, v) \in G/B \times G/B \times V \mid g^{-1}v \in V_{\boldsymbol{\sigma}} \cap V_{\boldsymbol{\sigma}'}\} \subset X_{\boldsymbol{\sigma}} \times_V X_{\boldsymbol{\sigma}'}$$

(3) $\psi_{\alpha_i}(\boldsymbol{\sigma})$ to the fundamental class of the associated variety $[L_{P_i}(V_{\boldsymbol{\sigma}})]$ embedded naturally in $X_{\boldsymbol{\sigma}} \times_V X_{\boldsymbol{\sigma}} \subset G/B \times G/B \times V$.

This isomorphism intertwines the anti-automorphism \star to the anti-isomorphism:

$$\mathbb{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}')^{\star} = \mathbb{w}(\boldsymbol{\sigma}', \boldsymbol{\sigma}) \quad \psi(\boldsymbol{\sigma})^{\star} = \psi(\boldsymbol{\sigma}) \quad \mu^{\star} = \mu.$$

We will prove this theorem below, once we have developed some of the theory of these categories.

Lemma 2.15 *The category A_I has a natural representation Y which sends each object $\boldsymbol{\sigma}$ to the polynomial ring S . The action is defined by the formulae:*

$$(2.5a) \quad \mathbb{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}') \cdot f = \varphi(\boldsymbol{\sigma}, \boldsymbol{\sigma}')f$$

$$(2.5b) \quad \psi_{\alpha}(\boldsymbol{\sigma}) \cdot f = \partial_{\alpha}(f)$$

$$(2.5c) \quad \mu \cdot f = \mu f$$

Proof. The codimension 1 relations are simple to check. For the codimension 2 relations, (2.4e) is a standard relation between BGG-Demazure operators. Applying the twisted Leibnitz rule

$$\partial_{\alpha}(gf) = s_{\alpha}(g)\partial_{\alpha}(f) + \partial_{\alpha}(g)f,$$

the relation (2.4f) follows from:

$$\begin{aligned} & \varphi(\boldsymbol{\sigma}_{m-j+1}, \boldsymbol{\sigma}_1)\partial_{\alpha}(\varphi(\boldsymbol{\sigma}_1, \boldsymbol{\sigma}_j)f) - \varphi(\boldsymbol{\sigma}_{m-j+1}, \boldsymbol{\sigma}_m)\partial_{\alpha}(\varphi(\boldsymbol{\sigma}_m, \boldsymbol{\sigma}_j)f) \\ &= (\varphi_1\varphi_3\partial_{\alpha}(\boldsymbol{\sigma}_1)\varphi_2 - \varphi_2\partial_{\alpha}\varphi_1\varphi_3)f = \partial_{\alpha}(\varphi_1 \cdot s_{\alpha}(\varphi_2))\varphi(\boldsymbol{\sigma}_{m-j+1}, \boldsymbol{\sigma}_j)f \end{aligned}$$

where, by definition:

$$\begin{aligned} \varphi_1 &= \varphi(\boldsymbol{\sigma}_1, \boldsymbol{\sigma}_j) = \varphi(s_{\alpha} \cdot \boldsymbol{\sigma}_{m-j+1}, s_{\alpha} \cdot \boldsymbol{\sigma}_m) \\ \varphi_2 &= \varphi(\boldsymbol{\sigma}_m, \boldsymbol{\sigma}_{m-j+1}) = \varphi(s_{\alpha} \cdot \boldsymbol{\sigma}_m, s_{\alpha} \cdot \boldsymbol{\sigma}_1). \\ \varphi_3 &= \varphi(\boldsymbol{\sigma}_{m-j+1}, \boldsymbol{\sigma}_j). \end{aligned} \quad \square$$

For each pair $(\boldsymbol{\sigma}, \boldsymbol{\sigma}') \in I' \times I'$, and $w \in W$, we fix a minimal length path (that is, crossing a minimal number of hyperplanes) from $C_{\boldsymbol{\sigma}', 1}$ to $C_{w\boldsymbol{\sigma}, w}$. Now, fold this path so that it lies in the positive Weyl chamber: the first time it crosses a root hyperplane, apply the corresponding simple reflection to what remains of the path. Then follow this new path until it strikes another wall, and apply that simple reflection to the remaining path, etc. The result is a sequence $\beta_1, \beta_2, \dots, \beta_p$ of simple root hyperplanes and sign vectors $\boldsymbol{\sigma}_1, \dots, \boldsymbol{\sigma}_p$ corresponding to the chambers where we reflect. Now, consider the morphism $\tilde{\mathbb{w}}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', w): \boldsymbol{\sigma}' \rightarrow \boldsymbol{\sigma}$ defined by the product:

$$(2.6) \quad \tilde{\mathbb{w}}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', w) = \mathbb{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_p)\psi_{\beta_p}(\boldsymbol{\sigma}_p)\mathbb{w}(\boldsymbol{\sigma}_p, \boldsymbol{\sigma}_{p-1})\psi_{\beta_{p-1}}(\boldsymbol{\sigma}_{p-1}) \cdots \psi_{\beta_1}(\boldsymbol{\sigma}_1)\mathbb{w}(\boldsymbol{\sigma}_1, \boldsymbol{\sigma}').$$





FIGURE 3. The paths from Example 2.x

Example 2.x. In our running example, this is given by the diagrams without dots which join the black strands with no crossing if $w = 1$ and with a crossing if $w = s_\alpha$, and a minimal number of red/black crossings possible.

In Figure 3, we show one possible path $\sigma' = (+, -, +, -) \rightarrow s_\alpha \sigma = (-, +, +, +)$, and its reflection. The resulting element $\tilde{w}((+, -, +, +), (+, -, +, -), s_\alpha)$ is given by:

$$\mathbb{w}((+, -, +, +), (-, -, +, +))\psi_\alpha((-, -, +, +))\mathbb{w}((-, -, +, +), (+, -, +, -))$$

and represented by the diagram



Theorem 2.16 *The elements $\tilde{w}(\sigma, \sigma', w)$ are a basis of the morphisms in A_I as a right module over S .*

Proof. These elements span: To show that they span, it suffices to show that their span contains the identity of each object, which is $\tilde{w}(\sigma, \sigma, 1)$ and is closed under right multiplication by the generators $\mathbb{w}(-, -)$ and $\psi(-)$. Note that

$$\tilde{w}(\sigma, \sigma', w)\psi_\alpha(\sigma') = \begin{cases} \tilde{w}(\sigma, \sigma', ws_\alpha) & ws_\alpha > w \\ 0 & ws_\alpha < w \end{cases}$$

$$\tilde{w}(\sigma, \sigma', w)\mathbb{w}(\sigma', \sigma'') = \tilde{w}(\sigma, \sigma'', w)\varphi(\sigma_p, \sigma', \sigma'')$$

so this shows that these vectors span.

Linear independence: Now, consider the action of these morphisms in the representation Y localized over the fraction field K of S . By construction, the image of every morphism $\sigma \rightarrow \sigma'$ lies in the subalgebra of \mathbb{C} -linear endomorphisms of K generated by

the action of elements of W and multiplication by elements of K . This is, of course, a copy of the twisted group algebra W over the field K . The action of $\tilde{w}(\sigma, \sigma', w)$ is given by the element w , times a non-zero rational function, plus elements which are shorter in Bruhat order. Thus, the morphisms $\sigma \rightarrow \sigma'$ have the same K -span as the element W . Since this span is a vector space of dimension $\#W$ over K , this is only possible if the elements $\tilde{w}(\sigma, \sigma', w)$ are linearly independent over S . \square

We can think about this proof a little differently if we think a bit more explicitly about paths. Let $\pi: [0, 1] \rightarrow \mathfrak{t}_{1, \mathbb{R}}$ be a generic path between generic points (i.e. any point where this path meets a hyperplane is distinct from all other hyperplanes, and it is transverse at these points), and let π' be the path obtained when we reflect in the Coxeter hyperplanes that π reaches in order so that we stay in the Weyl chamber \mathcal{C} .

Definition 2.17 Let w_π be the product of the morphisms $w(\sigma, \sigma')$ when π' crosses a matter hyperplane passing from the chamber C'_σ to C_σ and $\psi_\alpha(\sigma)$ when π' strikes and reflects off the corresponding Coxeter hyperplane, in the order we reach them along π' .

Note that we can extend this definition to paths bearing coupons with elements of S on them at generic points, where this simply means that we insert multiplication by that element of S into the product above. In these terms, we can rewrite the relations of Definition 2.11 in geometric terms: (2.4a) and (2.4b) around the hyperplane $\varphi_i^{\text{mid}} = 0$ become

$$(2.7a) \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \quad \begin{array}{c} \boxed{f} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \boxed{f} \end{array}$$

while (2.4c) and (2.4d), with the dotted line given by the Coxeter hyperplane $\alpha = 0$ becomes

$$(2.7b) \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = 0 \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} - \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array}$$

We have similar forms of the codimension 2 relations; the only one which is not a straightforward isotopy without corrections is (2.4f) which in the generic case where $m = 3$ becomes

$$(2.7c) \quad \begin{array}{c} s_\alpha \cdot \varphi_i^{\text{mid}} = 0 \\ \varphi_i^{\text{mid}} = 0 \\ \text{---} \\ \text{---} \\ \text{---} \end{array} - \begin{array}{c} s_\alpha \cdot \varphi_i^{\text{mid}} = 0 \\ \varphi_i^{\text{mid}} = 0 \\ \text{---} \\ \text{---} \\ \text{---} \end{array} = \begin{array}{c} s_\alpha \cdot \varphi_i^{\text{mid}} = 0 \\ \varphi_i^{\text{mid}} = 0 \\ \text{---} \\ \text{---} \\ \text{---} \\ \boxed{\partial_\alpha(\varphi_i)} \end{array}$$

As usual, any two such paths with the same endpoints can be joined by a combination of isotopies that preserve genericity and move only through “subgeneric” configurations. By “subgeneric,” we mean configurations where we have a single tangency of the path and a hyperplane, which allows us to create or cancel a pair of intersection points, or the path goes through a generic point in the intersection of two hyperplanes, which allows us to relate the paths around this codimension 2 subspace in the two different ways.

The relations (2.4a) and (2.4c) show that if a path has intersection points which we can cancel, then we can write \tilde{w}_π in terms of shorter paths, and if two paths differ by passing through a codimension 2 locus, the “codimension 2” relations (2.4e–2.4g) show that the corresponding \tilde{w}_π ’s differ by an element in the span of shorter paths. You can visualize these moves using the graphical rewriting of these relations.

The argument above can be rephrased as noting that ranging over all generic paths π , the elements \tilde{w}_π for all different paths span, and the relations above are enough to write every path in terms of a single homotopy representative of the minimal paths between each pair of endpoints, given by $\tilde{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', w)$.

Proof of Theorem 2.14. **There is a faithful, essentially surjective functor** $A_{I'}^Q \rightarrow \mathcal{X}_{I'}^Q$: We will first prove this for the deformed category; the statement for the undeformed category holds by base change. The functor described in the statement matches the action of $A_{I'}^Q$ on Y with that Y_X^Q of \mathcal{X}_I^Q sending $(\boldsymbol{\sigma}, w) \mapsto H_*^{BM,G}(X_{\boldsymbol{\sigma},w})$ described in Lemma 2.5(5), as simple computations with pushforward and pullback confirm (for example, as in [VV11, Prop. 2.23]):

- The equation (2.5a) is simply the fact that the pushforward in question gives multiplication by the normal bundle to $V_\sigma \cap V_{\sigma'}$ inside V_σ , which is the sum of the weight spaces with $\sigma_i = +$ and $\sigma'_i = -$. Thus, the Euler class of this normal bundle is the product of these weights.
- The equation (2.5b) is simply the formula for integration over a \mathbb{P}^1 bundle by Atiyah-Bott.
- The equation (2.5c) is clear from the definition.

The action on Y_X^Q is faithful by Lemma 2.5(5), so this shows that we have a faithful functor $A_{I'}^Q \rightarrow \mathcal{X}_{I'}^Q$, which is essentially surjective by construction.

This functor is full: Let $\mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', w)$ be the subset of the space $X_{\sigma'} \times_V X_\sigma$ where the relative position of the two flags is $w \in W$, and $\mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', < w), \mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', \leq w)$ where it is $< w$ or $\leq w$ in Bruhat order. By Lemma 2.5(2), the relative Borel-Moore homology $H_*^{BM,G}(\mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', \leq w), \mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', < w))$ is a free left or right $\text{Sym } \mathfrak{t}^*$ -module generated by the fundamental class $[\mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', w)]$. The homology class $\tilde{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', w)$ is the pushforward under the projection to $\mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}', \leq w)$ of the fundamental class of the fiber product

$$(2.8) \quad \mathbb{X}(\boldsymbol{\sigma}, \boldsymbol{\sigma}_p, \leq s_{\beta_p}) \times_{X_{\sigma_p}} \cdots \times_{X_{\sigma_1}} \mathbb{X}(\boldsymbol{\sigma}_1, \boldsymbol{\sigma}', \leq s_{\beta_1}).$$

where β_i, σ_i are as in (2.6). This projection map is an isomorphism over $\mathbb{X}(\sigma, \sigma', w)$, so in $H_*^{BM,G}(\mathbb{X}(\sigma, \sigma', \leq w), \mathbb{X}(\sigma, \sigma', < w))$, the class $\tilde{w}(\sigma, \sigma', w)$ is precisely the fundamental class. By Lemma 2.5(3), ranging over w , these give a basis as desired. This shows that the functor is faithful.

Compatibility with duality: The compatibility with duality is manifest from the definitions of the homology classes in Theorem 2.14(1-3). \square

One slightly annoying aspect of the structure of the category $A_{I'}$ is that it is not immediately apparent how to index its simple modules, or equivalently, its indecomposable projectives. We can consider the identity morphism $e(\sigma)$ as an idempotent element of $A_{I'}$, and study the projective $A_{I'}e(\sigma)$. If G is non-abelian, this is typically not be indecomposable.

In general, finding the set of simple modules might be quite challenging, but we can at least reduce it to studying a subset of the chambers I' . Choose a W -invariant inner product on \mathfrak{t} ; we'll use the distance function $d(-, -)$ induced by this inner product below.

Definition 2.18 *The defect of a chamber in I' is*

$$\text{def}(\sigma) = \min\{d(x, p) \mid x \in C_\sigma, p \in \mathfrak{t}_1^W\}.$$

In particular, the defect of a chamber is 0 if and only if it contains a fixed point.

If we let W_σ be the stabilizer of $\sigma \in I'$, then the projective module $A_{I'}e(\sigma)$ has an action of W_σ has an action of the nilHecke algebra of W_σ by right multiplication. Since the nilHecke algebra of W_σ is a matrix algebra of rank $\#W_\sigma$, we have that

$$A_{I'}e(\sigma) \cong P_\sigma^{\oplus \#W_\sigma}$$

where $P_\sigma = (A_{I'}e(\sigma))^{W_\sigma}$ is the invariants of this action.

We call an indecomposable summand of P_σ **novel** if it is not isomorphic to a summand of $P_{\sigma'}$ with lower defect or with $\#W_{\sigma'} > \#W_\sigma$, and **boring** if it is.

Proposition 2.19 *For each sign vector $\sigma \in I'$, the projective P_σ has at most one novel indecomposable summand.*

Building on this observation, we call σ itself **novel** if P_σ has a novel summand and **boring** if it does not.

Proof. Reduction to locality: Let n be the defect of σ and $m = \#W_\sigma$. Since $A_{I'}$ -gmod is a Krull-Schmidt category, the summands of P_σ are controlled by the quotient of $\text{End}(P_\sigma)$ modulo its Jacobson radical, which will be a sum of matrix algebras, one for each decomposable summand. Furthermore, the summand is isomorphic to a summand of $P_{\sigma'}$ with lower defect if the corresponding matrix algebra lies in the two-sided ideal $J_{<n,m}$ of morphisms factoring through $P_{\sigma'}$ with defect $< n$ or with

$\#W_{\sigma'} > m$. Thus, the result will follow if we show that $\text{End}(P_{\sigma})/J_{<n,m}$ is a local ring when it is non-trivial.

Preliminaries to locality: Now, fix σ with $C_{\sigma} \neq \emptyset$. Since both $C_{\sigma} \neq \emptyset$ and \mathfrak{t}_1^W are convex and close, there are unique points $x_0 \in C_{\sigma}, p_0 \in \mathfrak{t}_1^W$ which minimize the distance $d(x_0, p_0)$. Let x be generic in the interior of C_{σ} in a ball of radius ϵ around x_0 .

By uniqueness, the stabilizer W_{x_0} contains the stabilizer W_{σ} , but there is one situation where these are not equal: it could be that W_{σ} is trivial, but x_0 lies on a single root hyperplane. Otherwise, x_0 would have to lie in the intersection of two root hyperplanes, and thus could not lie on the intersection of two matter hyperplanes; in this case, the reflections for the root hyperplanes will preserve C_{σ} as well.

$\tilde{w}(\sigma, \sigma, w) \in J_{<n,m}$ if $w\sigma \neq \sigma$: Note that $\text{End}_{A_I}(A_I e(\sigma)) = e(\sigma)A_I e(\sigma)$ is spanned over S by the elements $\tilde{w}(\sigma, \sigma, w)$. We wish to show that whenever $w\sigma \neq \sigma$, we have $\tilde{w}(\sigma, \sigma, w) \in J_{<n,m}$. Constructing this element depends on a choice of a path joining x to the point $wx \in C_{w\sigma}$.

First, consider the case where $x_0 \neq wx_0$. Of course, the distance function $d(-, p_0)$ on the straight line that joins x_0 to wx_0 decreases from x_0 to the midpoint and then increases. Choosing our path sufficiently close to this straight line, we can guarantee that it passes through a chamber with lower defect than σ . Thus $\tilde{w}(\sigma, \sigma, w) \in J_{<n,m}$ in this case.

Next, we need to consider the case where $W_{\sigma} = \{1\}, W_{x_0} = \{1, s_{\alpha}\}$, and $w = s_{\alpha}$. Note, in this case, $m = 1$. In this case, since we have chosen x generically, the midpoint of the straight line from x to $s_{\alpha}x$ will lie in another chamber σ' with $s_{\alpha} \in W_{\sigma'}$. This means that $\tilde{w}(\sigma, \sigma, s_{\alpha})$ factors through this chamber with $\#W_{\sigma'} > 1$, and so $\tilde{w}(\sigma, \sigma, s_{\alpha}) \in J_{<n,m}$.

Completing the proof: Thus, we see that $\text{End}(A_I e(\sigma))/J_{<n,m}$ is spanned over S by $\tilde{w}(\sigma, \sigma, w)$ for $w \in W_{\sigma}$. Since we pass to W_{σ} -invariants, this means that it is a quotient of $S^{W_{\sigma}}$. Any graded finite-dimensional quotient of this polynomial ring is local, completing the proof. \square

2.4. Variations. As mentioned above, we can generalize these categories by taking any set P with a map $\iota: P \rightarrow I'$, and considering the category \mathcal{X}_P with objects given by P where

$$\text{Hom}_{\mathcal{X}_P}(p, p') := \text{Hom}_{\mathcal{X}_{I'}}(\iota(p), \iota(p')).$$

2.4.1. The set J . We let J (resp. J') be the subset of \mathcal{K} such that $c_{\sigma,w} \neq \emptyset$ (resp. $C_{\sigma,w} \neq \emptyset$); note that $J' \supseteq J$. In this case, the map $J' \rightarrow I'$ is given by $(\sigma, w) \mapsto (w^{-1}\sigma, 1)$. The category $\mathcal{X}_{J'}$ is Morita equivalent to $\mathcal{X}_{I'}$. However, it is a convenient framework for understanding this category, because we can define certain special elements of it. We let $w: (\sigma, w') \rightarrow (w\sigma, ww')$ to be the image of the identity on $(w^{-1}\sigma, 1)$ under the isomorphism

$$\text{Hom}_{\mathcal{X}_{J'}}((w\sigma, ww'), (\sigma, w')) := \text{Hom}_{\mathcal{X}_{I'}}(((w')^{-1}\sigma, 1), ((w')^{-1}\sigma, 1)).$$

These obviously satisfy the relations of W . It's more natural to think of the S action on $(\boldsymbol{\sigma}, w)$ to be the conjugate by w of that on $(w^{-1}\boldsymbol{\sigma}, 1)$. For each pair of pairs $(\boldsymbol{\sigma}, w)$ and $(\boldsymbol{\sigma}', w')$, we have a well-defined morphism $\mathbb{w}(\boldsymbol{\sigma}', w'; \boldsymbol{\sigma}, w)$ defined using the folding of a minimal path from $C'_{w^{-1}\boldsymbol{\sigma}, 1}$ to $C'_{w^{-1}\boldsymbol{\sigma}', w^{-1}w'}$ (using the same notation as (2.6)) by

$$\mathbb{w}(\boldsymbol{\sigma}', w'; \boldsymbol{\sigma}, w) = w' \mathbb{w}(\boldsymbol{\sigma}', \boldsymbol{\sigma}_p) s_{\beta_p} \mathbb{w}(\boldsymbol{\sigma}_p, \boldsymbol{\sigma}_{p-1}) \cdots s_{\beta_1} \mathbb{w}(\boldsymbol{\sigma}_1, \boldsymbol{\sigma}) w^{-1}.$$

When we extend the polynomial representation Y to this category, we thus still send every object to a copy of S with the action given by

$$(2.9) \quad \begin{aligned} \mathbb{w}(\boldsymbol{\sigma}, w; \boldsymbol{\sigma}', w') \cdot f &= \varphi(\boldsymbol{\sigma}, \boldsymbol{\sigma}') f \\ \psi_\alpha(\boldsymbol{\sigma}', w') \cdot f &= \partial_\alpha(f) \\ w \cdot f &= f^w \\ \mu \cdot f &= \mu f \end{aligned}$$

2.4.2. *The set K .* Note that if a sign vector $\boldsymbol{\sigma}$ is compatible with wBw^{-1} and $w'B(w')^{-1}$, then $\mathbb{w}(\boldsymbol{\sigma}, w; \boldsymbol{\sigma}, w')$ gives an isomorphism between these objects. Thus, we can reduce the size of our category by only choosing one object per sign vector $\boldsymbol{\sigma}$, and identifying it with any others via the elements $\mathbb{w}(\boldsymbol{\sigma}, w; \boldsymbol{\sigma}, w')$. This is the category \mathcal{X}_K attached to the set K of sign vectors with $c_\sigma \neq \emptyset$ for some ϕ (similarly, we can define K'), with the map to I' associating a sign vector to the unique Weyl translate compatible with $1 \in W$. Note that in this category, if $s_\alpha \boldsymbol{\sigma} = \boldsymbol{\sigma}$, then s_α is an endomorphism of this object. A calculation in the polynomial representation confirms the relation $s_\alpha = \alpha \psi_\alpha + 1$.

In \mathcal{X}_K , we have morphisms $\mathbb{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}')$, $\psi_\alpha(\boldsymbol{\sigma})$, $w \in W$, $\mu \in \mathfrak{t}_K^*$ as above, labeled by sign vectors $\boldsymbol{\sigma}, \boldsymbol{\sigma}' \in I'$, and these act as in (2.9). This category contains as a subcategory $\mathcal{X}_K^{\text{ab}}$, the category attached to the representation V and the torus $T_{\tilde{G}} \subset \tilde{G}$. This is generated over S by the elements $\mathbb{w}(\boldsymbol{\sigma}, \boldsymbol{\sigma}')$.

2.4.3. *The set \mathcal{C} .* Finally, we consider the extended arrangement on $\mathfrak{t}_{1, \mathbb{R}}$ defined by the hyperplanes $\varphi_i^{\text{mid}}(\xi) = n$ for $n \in \mathbb{Z}$. The chambers of this arrangement are defined not by sign vectors, but rather by integer vectors: associated to $\mathbf{a} = (a_1, \dots, a_d)$ we have the chamber

$$(2.10) \quad C_{\mathbf{a}} = \{\xi \in \mathfrak{t}_{1, \mathbb{R}} \mid a_i < \varphi_i^{\text{mid}}(\xi) < a_i + 1 \text{ for all } i\}.$$

As usual, we call \mathbf{a} feasible if this set is non-empty.

Definition 2.20 *Let \mathcal{C} be the set of feasible elements of \mathbb{Z}^d , identified with chambers as in (2.10).*

Considering the inclusion of chambers induces a map $\eta: \mathcal{C} \rightarrow K'$, which gives us a category $\mathcal{X}_{\mathcal{C}}$. Since the map $\mathcal{C} \rightarrow K'$ is surjective, $\mathcal{X}_{\mathcal{C}}$ is equivalent to $\mathcal{X}_{I'}$, but it will be useful to have this category for comparison to the Coulomb case. As before, we can

generate the morphisms of this category with morphisms $w(\mathbf{a}, \mathbf{a}')$ and copies of $\mathbb{k}[W]$ and S . These act in the polynomial representation by

$$(2.11a) \quad w(\mathbf{a}; \mathbf{a}') \cdot f = \varphi(\eta(\mathbf{a}), \eta(\mathbf{a}'))f$$

$$(2.11b) \quad \psi_\alpha(\mathbf{a}) \cdot f = \partial_\alpha(f)$$

$$(2.11c) \quad w \cdot f = f^w$$

$$(2.11d) \quad \mu \cdot f = \mu f.$$

2.5. The quiver and hypertoric cases. If G is abelian, then no relations involving ψ are relevant, since there are no Coxeter hyperplanes. We are left with the relations (2.4a, 2.4b, 2.4g), which appeared in [Bra+10; Bra+12]. The result is the algebra $A_{\text{pol}}^1(\phi, -)$ from [Bra+12, §8.6].

Now we fix a quiver Γ (possibly with edge loops) with vertex set \mathcal{V} and consider $V = \bigoplus_{i \rightarrow j} \text{Hom}(\mathbb{C}^{d_i}, \mathbb{C}^{d_j})$ as a module over $G = \prod GL_{d_i}$ as usual. In this case, we obtain the relations of a weighted KLR algebra as defined in [Web19b]. Let us explain this connection in a bit more detail. In this case, the group $H = N_{GL(V)}(G)$ is generated by G and by $GL(\mathbb{C}^{\chi_{i,j}})$ where $\chi_{i,j}$ is the number of edges $i \rightarrow j$, acting by taking linear combinations of the maps along these edges, that is, via the isomorphism

$$\bigoplus_{i \rightarrow j} \text{Hom}(\mathbb{C}^{v_i}, \mathbb{C}^{v_j}) \cong \bigoplus_{(i,j) \in \mathcal{V} \times \mathcal{V}} \text{Hom}(\mathbb{C}^{v_i}, \mathbb{C}^{v_j}) \otimes \mathbb{C}^{\chi_{i,j}}.$$

The subgroup H is generated by G and just the diagonal matrices in $GL(\mathbb{C}^{\chi_{i,j}})$, that is, those that act by rescaling the maps along the edges. In particular, a flavor ϕ_0 can be thought of as just assigning an integer ϕ_e to each edge. Two such cocharacters define the same homomorphism to F if and only if they are cohomologous.

The weight spaces of the representation V are given by the matrix coefficients of the map along each edge e . That is, they are indexed by an edge $e: i \rightarrow j$, and indices $k \leq v_i$ and $m \leq v_j$.

Letting $z_{i,1}, \dots, z_{i,v_i}$ be the usual coordinates on the diagonal matrices in $\mathfrak{gl}(\mathbb{C}^{v_i})$, we can transfer these to coordinates on \mathfrak{t}_1 so that the weight of the matrix coefficient for $(e; k, m)$ is $\varphi_{e;k,m} = z_{j,m} + \phi_e - z_{i,k}$.

In this case, the chambers C_σ are thus defined by inequalities of the form

$$\begin{aligned} z_{i,k} &\leq z_{j,m} + \phi_e - \frac{1}{2} & \sigma_{e;k,m} &= + \\ z_{i,k} &\geq z_{j,m} + \phi_e - \frac{1}{2} & \sigma_{e;k,m} &= -. \end{aligned}$$

We can interpret the coordinates $z_{i,k}$ as giving us a loading ([Web19b]), that is, a finite subset of \mathbb{R} , labeled with elements of \mathcal{V} . Considered this way, the chambers $C_{\sigma,1}$ are precisely the equivalence classes of loadings for the KLR algebra with the weighting $\vartheta_e = \phi_e - \frac{1}{2}$ by [Web19b, p. 2.12]. We can choose one representative generic point in the interior of each chamber $C_{\sigma,1}$.

Thus, we can consider the weighted KLR algebra $W_{I'_\phi}^\vartheta$ for the quiver Γ , weighting ϑ and loading set I'_ϕ , that is the collection of weighted KLR diagrams ([Web19b, Def. 2.3]) whose top and bottom match one of the representative points in each chamber. We'll use the convention that the polynomial $Q_e(u, v)$ which appears in the relations of the weighted KLR algebra assigned to every edge is $Q_e(u, v) = u - v$.

Furthermore, we can visualize a path $\pi: [0, 1] \rightarrow \mathfrak{t}_1$ by superimposing the plots of the path $t \mapsto (z_{i,k}(t), t)$ for each pair (i, k) in $\mathbb{R} \times [0, 1]$. We can consider this collection of curves as a weighted KLR diagram by labeling the curve for each (i, k) with the node i . In addition, for each edge e and each $k \leq h(e)$, we add a ghost strand on the curve $(z_{h(e),k}(t) + \vartheta_e, t)$. We call this diagram \mathbb{D}_π .

Finally, for each of these loadings, we can identify the ring $S = \mathbb{k}[z_{i,k}]_{i \in \mathcal{V}, k \in [1, v_i]}$ with the polynomial ring of dots on the strands of this loading by sending $z_{i,k}$ to the dot y_p where the p th strand from the left (counting those with all labels) is the k th strand from the left with label i . This defines an isomorphism $p_\sigma: S \rightarrow \mathbb{k}[\mathbf{y}]$.

Theorem 2.21 *There is an isomorphism $A_{I'_\phi} \rightarrow W_{I'_\phi}^\vartheta$ sending $w_\pi \mapsto \mathbb{D}_\pi$ and S to the polynomial ring $\mathbb{k}[y_1, \dots, y_n]$ via p_σ .*

If, instead, we worked with the deformed category $A_{I'_\phi}^Q$, this will correspond to the deformations introduced in [Web19b, §2.7].

Proof. Let us study the polynomial representation given in Lemma 2.15 for $A_{I'_\phi}$ and [Web19b, Prop. 2.7] for $W_{I'_\phi}^\vartheta$. It's enough to check that associating the action of \mathbb{D}_π to the action of w_π on this representation defines an action of $W_{I'_\phi}^\vartheta$. It will be more convenient if for each σ , we apply the corresponding isomorphism p_σ to transfer this action to a sum of some number of copies of $\mathbb{k}[\mathbf{y}]$.

(1) The idempotent $e(\sigma)$ corresponds to a constant path at some loading, which is sent under the corresponding idempotent in $W_{I'_\phi}^\vartheta$ with straight vertical strands. Thus, this idempotent acts by projection to the corresponding copy of $\mathbb{k}[\mathbf{y}]$.

(2) The endomorphisms S act on the appropriate copy of S by left multiplication, and thus the same is true of the dots $\mathbb{k}[\mathbf{y}]$ on the corresponding copy of $\mathbb{k}[\mathbf{y}]$.

(3) A simple root α of G corresponds to $i \in \mathcal{V}$ and consecutive indices $k, k + 1$, and the morphism ψ_α acts by $f \mapsto \frac{f^{s_\alpha} - f}{\alpha}$. This corresponds to the path that goes straight from our fixed point to the wall $\alpha = 0$, “bounces off” and then returns. Now, let us consider the corresponding diagram \mathbb{D}_π . Assume that the k and $k + 1$ th strands with label i are the r th and s th from the left. The diagram \mathbb{D}_π crosses these strands, while leaving all other strands straight vertical. Precomposing with p_σ^{-1} and postcomposing with p_σ , we find that the induced action is:

$$(2.12a) \quad \psi_\alpha \cdot f(\mathbf{y}) = \frac{f(\dots, y_r \dots, y_r, \dots) - f(\mathbf{y})}{y_s - y_r}$$

(4) The wall-crossing element $w(\boldsymbol{\sigma}; \boldsymbol{\sigma}')$ for two adjacent chambers changes the sign of one weight $\varphi_{e;k,m} = z_{j,m} + \phi_e - z_{i,k}$. The action of this element is given by

$$w(\boldsymbol{\sigma}; \boldsymbol{\sigma}') \cdot f = \begin{cases} f & \boldsymbol{\sigma}'_{e;k,m} = +, \boldsymbol{\sigma}_{e;k,m} = - \\ (z_{j,m} - z_{i,k})f & \boldsymbol{\sigma}'_{e;k,m} = -, \boldsymbol{\sigma}_{e;k,m} = + \end{cases}$$

The corresponding diagram \mathbb{D}_π has a single crossing of a strand with some label i over the ghost for an edge $e: i \rightarrow j$ and some number of crossings of solid strands with different labels, but none of strands with the same label. In fact, since $\varphi_{e;k,m}$ is the weight which changes sign, the strand with label i is the k th from the left with this label, and the strand with label j is the m th left with this label. Considering strands with all labels, assume that these strands are the r th and s th from the left, respectively. Let w denote the permutation of solid strands induced by this diagram.

Note that $\varphi_{e;k,m}$ is positive if the solid strand is left of the ghost and negative if it is to the right. Precomposing with $p_{\boldsymbol{\sigma}'}$ and postcomposing with $p_{\boldsymbol{\sigma}}$, we find that the induced action is:

$$(2.12b) \quad f(Y_1, \dots, Y_n) \mapsto f(Y_{w(1)}, \dots, Y_{w(n)}) \cdot (Y_s - Y_r)$$

if the solid strand goes from right to left of the ghost and by

$$(2.12c) \quad f(Y_1, \dots, Y_n) \mapsto f(Y_{w(1)}, \dots, Y_{w(n)})$$

if the solid strand goes from left to right of the ghost.

The formulas (2.12a–2.12c) almost match the action of [Web19b, Prop. 2.7], but annoyingly, the conventions of these papers don't quite match. The representation we desire will result if we switch since “left” and “right” in the first and second bullet points of that definition. Exactly the same proof shows that this is a faithful representation. Equivalently, we would obtain a representation of $A_{I'_\phi}$ matching that of [Web19b, Prop. 2.7] if we swapped $+$ and $-$ in the definition of $\varphi(\boldsymbol{\sigma}, \boldsymbol{\sigma}')$. The proof of Lemma 2.15 is easily modified to show that this is a faithful representation.

By faithfulness, this shows that we have an injective homomorphism $A_{I'_\phi} \rightarrow W_{I'_\phi}^\vartheta$. This map is surjective, since, by definition, the diagrams \mathbb{D}_π and $\mathbb{k}[\mathbf{y}]$ generate $W_{I'_\phi}^\vartheta$. This completes the proof. \square

2.6. Kirwan-Ness characters. Throughout this section, we fix a flavor ϕ and stability parameter ξ . We'll need for some results later in the paper that the category $A_{I'}$ has an additional structure: a graded triangular basis in the sense of Brundan [Bru23]. We can also consider this as a purely algebraic counterpart of the categorical Kirwan-Ness stratification of McGerty and Nevins' [MN14b], and indeed its combinatorics matches that of the Kirwan-Ness stratification on $\mu^{-1}(0) \subset T^*V$ which measures “unstable” a given point is.

We begin with the combinatorics of chambers. We call a chamber C_σ **weakly ξ -bounded** if ξ attains a maximum on C_σ and **strongly ξ -bounded** if the subset on which the maximum is attained is itself bounded; of course, these are equivalent if ξ is a generic linear function, but we cannot always assume this. In linear programming, “bounded” is usually used to mean in the weak sense, but both will be relevant for us at different times. Let us state a result we will use later:

Lemma 2.22 *There is a choice of σ such that C_σ is weakly ξ -bounded but not strongly ξ -bounded if and only if ξ is in a proper subspace spanned by a subset of the weights φ_i .*

In particular, there is a choice of $\xi \in (\mathfrak{t}^)^W$ such that all weakly ξ -bounded chambers are strongly ξ -bounded if and only if $(\mathfrak{t}^*)^W$ does not lie in any proper subspace spanned by φ_i 's.*

Proof. Assume that there is a C_σ is weakly ξ -bounded but not strongly ξ -bounded, and let A be the subset of points on which ξ attains a maximum. The asymptotic cone A_\circ of A is the set of points on which the corresponding homogeneous linear program attains a maximum. This cone is exactly the vectors $x \in \mathfrak{t}$ such that $\sigma_i \varphi_i(x) \geq 0$ and $\xi(x) = 0$. Since C_σ is weakly bounded and non-empty, Farkas' lemma implies that $-\xi = \sum a_i \varphi_i$ with $a_i \in \mathbb{R}_{>0}$. Of course, if $\varphi_i(x) \neq 0$, then this implies that $a_i = 0$, and thus $-\xi$ lies in the span of the weights φ_i such that $\varphi_i(x) = 0$ for any $x \in A_\circ$. Thus, if any non-zero x exists, then ξ lies in a proper subspace spanned by φ_i 's which is perpendicular to x . On the other hand, if ξ lies in such a proper space, we can choose any x in its perpendicular. For some σ , the polyhedron C_σ contains a ray parallel to x which shows that it is not strongly ξ -bounded. \square

Note that if (G, V) is a quiver theory, then the span U of any set of weights induces an equivalence relation on the set of coordinates $\Omega = \{(i, k) \mid i \in \mathcal{V}, k \in [1, v_i]\}$ where $(i, k) \sim (j, \ell)$ if $z_{i,k} - z_{j,\ell}$ is in the span. We can divide Ω into two subsets $\Omega = \Omega_0 \cup \Omega_1$, where $\Omega_0 = \{(i, k) \mid z_{i,k} \in U\}$ and Ω_1 is the union of all the other equivalence classes. One can easily check that U will be exactly the vectors such that the sum of the coordinates on each Ω_1 is 0. Thus, U cannot contain a point all of whose coordinates are positive unless $U = \mathfrak{t}^*$.

Thus, in the cases of greatest interest to us, there will always be a choice of ξ where the notion of strongly and weakly ξ -bounded coincide:

Corollary 2.23 (1) *If G is abelian, then $W = 1$, so \mathfrak{t}^* cannot lie in a proper subspace, and any generic ξ will have the notion of strongly and weakly ξ -bounded coincide.*

(2) *If (G, V) is a quiver theory, then any ξ whose coordinates are all positive will have the notion of strongly and weakly ξ -bounded coincide.*

If C_σ is not weakly ξ -bounded, we call it ξ -unbounded. This means that all the level sets of ξ on C_σ will be bounded if C_σ is strongly ξ -bounded.

Choose a W -invariant inner product on \mathfrak{t} . This allows us to define cocharacters $\xi^\vee, \varphi_i^\vee \in \mathfrak{t}$ such that $\langle \varphi_i^\vee, \nu \rangle = \varphi_i(\nu)$ for all cocharacters $\nu \in \mathfrak{t}$. Consider the convex subset:

$$(2.13) \quad \text{KN}_\sigma = \left\{ \xi^\vee + \sum_{i=1}^d a_i \sigma_i \varphi_i^\vee \mid a_i \in \mathbb{R}_{\geq 0} \right\} \subset \mathfrak{t}$$

Since the set KN_σ is convex, there is a unique point in this set with minimal norm which we denote β_σ . This is a measure of “how far from being stable” a generic point in $(T^*V)_\sigma$ is, or equivalently, how “far” from weakly ξ -bounded C_σ is.

Lemma 2.24 *The chamber C_σ is weakly ξ -bounded iff $\beta_\sigma = 0$.*

Proof. This is Farkas’ lemma again: The equation $\sum_{i=1}^d a_i \sigma_i \varphi_i^\vee = -\xi^\vee$ has a solution with $a_i \geq 0$ if and only if there is no element of $X \in \mathfrak{t}$ such that $\sigma_i \varphi_i(X) \geq 0$ for all i and $\xi(X) > 0$. Of course, any ray contained in the chamber C_σ is parallel to such an X , and the condition $\xi(X) \leq 0$ ensures that ξ is bounded above on this ray. Since C_σ is a polyhedron, it is weakly ξ -bounded if and only if ξ is bounded above on each ray in the chamber. \square

Let $\mathcal{KN} = \{\beta_\sigma \mid \sigma \in I'_\phi\}$. We endow the set I'_ϕ with a pre-order $\sigma \leq \sigma'$ if $\beta_{\sigma'} \in \text{KN}_\sigma$. We can define a corresponding partial order on \mathcal{KN} coarsening this one by declaring $\beta \leq \beta'$ if $\langle \beta, \beta \rangle \leq \langle \beta', \beta \rangle$.

If $a_i \neq 0$, the point $\beta_\sigma = \xi^\vee + \sum_{i=1}^d a_i \sigma_i \varphi_i^\vee$ must be a critical point of $|\xi^\vee + \sum_{i=1}^d a_i \sigma_i \varphi_i^\vee|$ with respect to varying a_i . Thus, if $a_i \neq 0$, then we must have that $\langle \varphi_i^\vee, \beta_\sigma \rangle = 0$. In particular, the orthogonal projection of ξ^\vee to the span of β_σ is β_σ itself. One can easily check that:

Lemma 2.25 *For any $x \in C_\sigma$, then $x + a\beta_\sigma \in C_\sigma$ for $a \in \mathbb{R}_{\geq 0}$.*

Note that for any sign vector σ and any β , there is a new sign vector $\sigma_{+,\beta}$ defined by

$$(\sigma_{+,\beta})_i = \begin{cases} + & \varphi_i(\beta) > 0 \\ - & \varphi_i(\beta) < 0 \\ \sigma_i & \varphi_i(\beta) = 0 \end{cases}$$

This sign vector is uniquely characterized by the property that $x + a\beta \in C_{\sigma_{+,\beta}}$ for $a \gg 0$ and $x \in C_\sigma$. Note that for $x \in C_{\sigma,1}$ dominant, we might have that $x + a\beta$ is no longer dominant, if β is not. However, there is some w of minimal length so that $\sigma_{++,\beta} = w^{-1}\sigma_{+,\beta}$ is dominant. We define the **standard morphism**

$$(2.14) \quad \mathfrak{s}(\sigma, \beta) = \tilde{w}(\sigma_{++,\beta}, \sigma, w)$$

for σ, β . In the formalism that associates paths to morphisms, this corresponds to starting at a generic point of C_σ and traveling a large distance in the direction of β , and then folding the path to stay in the dominant Weyl chamber. If $x \in C_{\sigma,1} \neq \emptyset$, then β_σ must be dominant, so the same is true of $x + a\beta_\sigma$, and so $\mathfrak{s}(\sigma, \beta_\sigma) = e(\sigma)$ in this case.

Let W_β be the stabilizer of β , G_β the corresponding Levi subgroup and

$$\mathbb{V}_\beta = \text{span}\{V_i \mid \varphi_i(\beta) = 0\};$$

note that \mathbb{V}_β is a G_β -module. We'll consistently use underline to denote objects corresponding to this choice of gauge group and matter representation. Thus, we let $\underline{\sigma}$ denote the restriction of the sign vector σ , and $e(\underline{\sigma})$ denote the corresponding idempotent. Furthermore, we let \underline{A} denote the category A constructed with respect to the group and representation $(G_\beta, \mathbb{V}_\beta)$.

Let $\xi_\beta = \xi^\vee - \beta$ and $\xi_\sigma = \xi_{\beta_\sigma}$; by definition, we have that $\xi_\sigma = -\sum_{\varphi_i(\beta_\sigma)=0} a_i \sigma_i \varphi_i^\vee$ with $a_i \geq 0$. Thus, the chamber $\underline{C}_{\underline{\sigma}}$ of the matter arrangement for \mathbb{V}_β is weakly ξ_β -bounded.

More generally, keeping σ and $\beta = \beta_\sigma$ fixed as above, we can consider an arbitrary σ' , and the corresponding restricted sign vector $\underline{\sigma}'$. We can now analyze the Kirwan-Ness character for $\underline{\sigma}'$ for $\xi_\beta = \xi_\sigma$ above.

Lemma 2.26 *The chamber $\underline{C}_{\underline{\sigma}'}$ is weakly ξ_β -bounded if and only if $\beta_{\sigma'+\beta} \leq \beta$. If $\beta_{\sigma'+\beta} = \beta$ then $\beta_{\sigma'} \leq \beta$.*

Proof. Let $\sigma'' = \sigma'_{+\beta}$. By assumption, we have that $\sigma''_i \varphi_i(\beta) \geq 0$, so for $x = \xi^\vee + \sum_{i=1}^d a_i \sigma_i \varphi_i^\vee \in \text{KN}_{\sigma''}$, we have

$$\langle x, \beta \rangle = \langle \beta, \beta \rangle + \sum_{i=1}^d a_i \sigma'_i \varphi_i(\beta) \geq \langle \beta, \beta \rangle.$$

Thus, we have $\beta_{\sigma''} \geq \beta$.

We've already discussed the fact that if $\beta = \beta_{\sigma''}$ then $\underline{C}_{\underline{\sigma}'}$ is weakly ξ_β -bounded. On the other hand, if $\underline{C}_{\underline{\sigma}'} = \underline{C}_{\underline{\sigma}''}$ is weakly ξ_β -bounded, then we must have $\xi_\beta = \sum_{\varphi_i(\beta)=0} -a_i \sigma'_i \varphi_i^\vee$ for some $a_i \geq 0$, which implies that $\beta \in \text{KN}_{\sigma''}$ so $\beta_{\sigma''} \leq \beta$. This implies $\beta_{\sigma''} = \beta$.

If $\beta'' = \beta$, then we have an expression $\beta = \xi + \sum_{\varphi_i(\beta)=0}^d a_i \sigma''_i \varphi_i^\vee$. For any i such that $\varphi_i(\beta) = 0$, we have $\sigma'_i = \sigma''_i$, so $\beta \in \text{KN}_{\sigma'}$ and $\beta_{\sigma'} \leq \beta$. \square

For a dominant $\beta \in \mathcal{KN}$, let \mathcal{I}_β be the ideal of A_I generated by $e(\sigma)$ for $\beta_\sigma = \beta$. Let $\mathcal{I}_{\geq \beta} = \sum_{\beta' \geq \beta} \mathcal{I}_{\beta'}$ and $\mathcal{I}_{> \beta} = \sum_{\beta' > \beta} \mathcal{I}_{\beta'}$.

Lemma 2.27 *If σ and σ' satisfy $\beta = \beta_\sigma = \beta_{\sigma'}$, then we have an injective homomorphism*

$$\iota_\beta: e(\underline{\sigma})\underline{A}e(\underline{\sigma}') \rightarrow e(\sigma)Ae(\sigma')$$

sending w_π to $w_{\pi+a\beta}$ for $a \gg 0$.

Proof. Since $\beta = \beta_\sigma = \beta_{\sigma'}$, the sign vectors σ and σ' only differ for indices i such that $\varphi_i(\beta) = 0$. Furthermore, the path $w_{\pi+a\beta}$ will only cross matter hyperplanes of this type for a sufficiently large. In particular, this shows that when we take the relations of (2.7a–2.7c) in \underline{A} and add $a\beta$ to them, we obtain the relations in A , and thus we have a homomorphism. \square

Let $\hat{A}^{(\beta)}$ be the completion of $A_{I'_\phi}$ at the ideal in S^W defined by the orbit of β . Each idempotent $e(\sigma)$ in this completion carries an action of the corresponding completion of S . This is a semi-local commutative ring, with maximal ideals corresponding to the elements of $W \cdot \beta$.

Let $e(\sigma, w\beta)$ denote the projection to the stable kernel of the maximal ideal $\mathfrak{m}_{w\beta}$ for $w\beta$, that is, the unique element of the completion congruent to 1 modulo any power of $\mathfrak{m}_{w\beta}$ and congruent to 0 modulo any power of another maximal ideal. Note that $e(\sigma) = \sum_{\beta' \in W \cdot \beta} e(\sigma, \beta')$.

Let $\hat{A}^{(0)}$ be the completion of the corresponding category $\underline{A}_{I'_\phi}$ for (G_β, V_β) at the origin.

Lemma 2.28 *The categories $\hat{A}^{(0)}$ and $\hat{A}^{(\beta)}$ are Morita equivalent.*

Proof. Let $\underline{e} = \sum_{\tau \in I'_\phi} e(\tau, \beta)$.

There is an isomorphism $\underline{e}\hat{A}^{(\beta)}\underline{e} \cong \hat{A}^{(0)}$: Note that for any element of the image of ι_β , we have $\underline{e}\iota_\beta(a) = \iota_\beta(a)\underline{e}$. This shows that $a \mapsto \underline{e}\iota_\beta(a)\underline{e}$ is an algebra homomorphism. Precomposing with the algebra homomorphism that maps $\mu \mapsto \mu - \langle \mu, \beta \rangle$ results in a map that is continuous in the appropriate ideal-adic topologies, and thus induces a homomorphism $\hat{\iota}: \hat{A}^{(0)} \rightarrow \underline{e}\hat{A}^{(\beta)}\underline{e}$.

The space $\hat{A}^{(\beta)}e(\tau, \beta)$ is the completion of $Ae(\tau)$ with respect to the maximal ideal for β in the right action of S . The action on the left side decomposes into subspaces $e(\sigma, w\beta)\hat{A}^{(\beta)}e(\tau, \beta)$ for β' ranging over $W\beta$ and σ ranging over I'_ϕ . Furthermore, from the relations of Definition 2.11, we can see that $\tilde{w}(\sigma, \tau, w)$ lies in the span of $e(\sigma, w'\beta)\hat{A}^{(\beta)}e(\tau, \beta)$ for $w' \leq w$ in Bruhat order, with non-trivial image in the projection to $e(\sigma, w\beta)\hat{A}^{(\beta)}e(\tau, \beta)$. This is only possible if the elements $\tilde{w}(\sigma, \tau, w'')$ for $w'' \in wW_\beta$ project to a basis of $e(\sigma, w\beta)\hat{A}^{(\beta)}e(\tau, \beta)$, and in particular $e(\sigma, \beta)\hat{A}^{(\beta)}e(\tau, \beta)$ has a basis given by $\tilde{w}(\sigma, \tau, w'')$ for $w'' \in W_\beta$, which are precisely the image of the same basis for $\hat{A}^{(0)}$ in the image of $\hat{\iota}$. This shows that $\hat{\iota}$ induces an isomorphism $\underline{e}\hat{A}^{(\beta)}\underline{e} \cong \hat{A}^{(0)}$ as desired.

The proof of Morita equivalence: Morita equivalence will hold if \underline{e} generates the algebra $\hat{A}^{(\beta)}$ as a 2-sided ideal, that is, if $e(\sigma, w\beta) \in \hat{A}^{(\beta)}\underline{e}\hat{A}^{(\beta)}$ for all $\sigma \in I'_\phi, w \in W$.

Consider the morphism $\mathbb{w}(\sigma_{+,w\beta}, \sigma)$. By (2.4a), we see that we have an equality

$$\mathbb{w}(\sigma, \sigma_{+,w\beta})\mathbb{w}(\sigma_{+,w\beta}, \sigma) = \varphi(\sigma, \sigma_{+,w\beta}, \sigma).$$

The right-hand side is the product of the weights on which σ and $\sigma_{+,w\beta}$ have different signs. Of course, this is only possible if this weight is non-zero on $w\beta$, so this product is invertible in the completion of S . This shows that

$$e(\sigma, w\beta) = \frac{1}{\varphi(\sigma, \sigma_{+,w\beta}, \sigma)}\mathbb{w}(\sigma, \sigma_{+,w\beta})e(\sigma_{+,w\beta}, w\beta)\mathbb{w}(\sigma_{+,w\beta}, \sigma),$$

so $e(\sigma, w\beta)$ is in the 2-sided ideal generated by $e(\sigma_{+,w\beta}, w\beta)$.

On the other hand $e(\sigma_{+,w\beta}, w\beta) = we(\sigma'_{+, \beta}, \beta)w^{-1}$ where $\sigma' = w^{-1}\sigma$. If W_β is non-trivial, then σ' is only unique up to the action of W_β , and we can choose the unique element of this orbit such that $\underline{\sigma'}$ is dominant, so $e(\sigma'_{+, \beta}, \beta)$ is a summand of \underline{e} . This shows that $e(\sigma, w\beta)$ is in the ideal generated by \underline{e} , and thus the desired Morita equivalence. \square

Lemma 2.29 *The subspace of morphisms $\sigma' \rightarrow \sigma$ which lie in the ideal \mathcal{I}_β is generated as a left S -module by elements of the form $\mathfrak{s}(\sigma, w\beta)^*\mathbb{w}_\pi\mathfrak{s}(\sigma', w\beta)$ for π a path from $x' \in C_{\sigma'_{++}, \beta, 1}$ to $x \in C_{\sigma_{++}, \beta, 1}$ in the image of ι_β .*

Proof. By Theorem 2.16, any element of \mathcal{I}_β is in the S -span of morphisms of the form $\mathbb{w}(\sigma, \sigma'', w)\mathbb{w}(\sigma'', \sigma', w')$ for different choices of w, w' .

Furthermore, we have some freedom in the construction of these morphisms, since they depend on a choice of a minimal length path from z' to $w'z''$ to $w'wz$ for any $z \in C_{\sigma, 1}, z' \in C_{\sigma', 1}, z'' \in C_{\sigma'', 1}$. We can freely replace z'' with $z'' + a\beta$ for $a \geq 0$. If we choose a sufficiently large, then the straight-line path from z' to $w'z''$ must pass through $C_{\sigma'_{+, w'\beta}, w_1}$ for some w_1 ; let $w_2 = w_1^{-1}w'$. We can thus write

$$\mathbb{w}(\sigma'', \sigma', w') = \mathbb{w}(\sigma'', \sigma'_{+, w'\beta}, w_2)\mathfrak{s}(\sigma', w'\beta).$$

Similarly, we can write

$$\mathbb{w}(\sigma, \sigma'', w) = \mathfrak{s}(\sigma, w'\beta)^*\mathbb{w}(\sigma_{++, w\beta}, \sigma'', w_3)$$

for some w_3 , and we can take

$$\mathbb{w}_\pi = \mathbb{w}(\sigma'', \sigma_{++, w\beta}, w_3)\mathbb{w}(\sigma'_{+, w'\beta}, \sigma'', w_2).$$

The element \mathbb{w}_π comes from folding a portion of the path z' to $w'z''$ to $w'wz$ which by construction only passes through chambers for ρ with $\rho_{+, w'\beta} = \rho$. This folding is unchanged by applying an element of the Weyl group, so we can instead consider a portion of the path $(w')^{-1}z'$ to z'' to wz . The corresponding morphism is thus unchanged by translating by a large positive multiple of β . In particular, this shows that it only crosses the walls for roots such that $\alpha(\beta) = 0$. This shows that it is in the image of ι_β . \square

Note that any $\beta \in \mathcal{KN}$ will be dominant, since $C_{\sigma,1}$ must be non-empty for some σ with $\beta = \beta_\sigma$. Thus, we will often want to consider the saturation of this set under W . For any $\beta' = w\beta$ and $\sigma \in I'_\phi$, we can consider the chamber $\sigma_{++,\beta'}$ and the morphism $\mathfrak{s}(\sigma, \beta'): \sigma \rightarrow \sigma_{++,\beta}$. By construction $\beta_{\sigma_{++,\beta'}} = \beta$, so we can consider the image $\iota_\beta(\underline{A})$ and multiply this on the right with $\mathfrak{s}(\sigma, \beta')$ and on the left with a mirror image $\mathfrak{s}(\sigma', \beta')^*$ for some other chamber β' .

Let \underline{A}_0 be a vector space complement to $\mathcal{I}_{>0}$ inside \underline{A} .

As mentioned previously, we want to prove that we have a graded triangular basis in the sense of [Bru23, Def. 1.1]. The data of this basis are:

- The set \mathbf{S} of special idempotents and the set of distinguished idempotents is just the set $e(\sigma)$ for $\sigma \in I'_\phi$.
- The poset (Λ, \leq) is the poset \mathcal{KN} , with the function $I'_\phi \rightarrow \mathcal{KN}$ given by $\sigma \mapsto \beta_\sigma$.
- The subset $H(\sigma, \sigma')$ for $\beta = \beta_\sigma = \beta_{\sigma'}$ is a fixed basis of $\iota_\beta(\underline{A}_0)$.
- The subset $Y(\sigma', \sigma)$ is the set $\mathfrak{s}(\sigma, \beta')$ for $\beta' \in W \cdot \beta_{\sigma'}$ such that $\sigma' = \sigma_{++,\beta'}$, and $X(\sigma, \sigma') = Y(\sigma', \sigma)^*$.

For each $\beta \in \mathcal{KN}$, let E_β be the span of $\mathfrak{s}(\sigma', \beta')^* \iota_\beta(\underline{A}_0) \mathfrak{s}(\sigma, \beta')$ for all pairs $\sigma, \sigma' \in I'_\phi$ and all $\beta' \in W\beta$. Note that this is the same as the space of the products $X(\sigma, \tau)H(\tau, \tau')Y(\tau', \sigma')$ for σ, σ' arbitrary and $\beta = \beta_\tau = \beta_{\tau'}$.

Lemma 2.30 *The data above define a graded triangular basis. In particular, as a vector space, the algebra A is the direct sum of the subspaces E_β for $\beta \in \mathcal{KN}$.*

Proof. **The subspaces E_β span:** We'll prove this by induction. Our inductive hypothesis will be that:

(a_γ) The ideal $\mathcal{I}_{\geq\gamma}$ is the span of the subspaces E_β with $\beta \geq \gamma$.

We'll prove that (a_γ) holds if (a_β) holds for all $\beta > \gamma$. The base case of our induction is $\gamma = \xi$, in which case $\mathcal{I}_{\geq\xi} = E_\xi$ and the result holds.

By Lemma 2.29, the ideal $\mathcal{I}_{\geq\gamma}$ is spanned by $\mathfrak{s}(\sigma', \beta')^* \iota_\beta(\underline{A}) \mathfrak{s}(\sigma, \beta')$ for $\beta \geq \gamma$ and $\sigma, \sigma' \in I'_\phi$ and $\beta' \in W\beta$ as before. On the other hand, by Lemma 2.27, we have that

$$\mathfrak{s}(\sigma', \beta')^* \iota_\beta(\mathcal{I}_{>0}) \mathfrak{s}(\sigma, \beta') \subset \mathcal{I}_{>\beta}.$$

Thus, E_γ and $\mathcal{I}_{>\gamma}$ span $\mathcal{I}_{\geq\gamma}$. Since $\mathcal{I}_{>\gamma}$ is spanned by $\mathcal{I}_{\geq\beta}$ for $\beta > \gamma$, applying the inductive hypothesis (a_β) for all such β , we obtain the conclusion (a_γ).

The algebra A follows by the case where $\beta = 0$.

The subspaces are independent: Given a linear dependence between these subspaces, we can assume that β is chosen to have minimal norm amongst the terms appearing. That is, we can assume that x is an element of E_β which also lies in the sum of E_γ for $\gamma \neq \beta$ and $|\gamma| \geq |\beta|$. By Lemma 2.26, for any σ' with such $\gamma = \beta_{\sigma'}$, the corresponding chamber $\underline{\sigma}'$ in the arrangement $(G_\beta, \mathbb{V}_\beta)$ is ξ_β -unbounded.

Consider the image of these elements in the completion $\hat{A}^{(\beta)}$; this map is injective, so x remains non-zero. The subspaces E_γ have image in the ideal corresponding to $\underline{\mathcal{I}}_{>0}$ in

$\hat{\underline{A}}^{(0)}$ by the unboundedness discussed above, thus, the same is true of x . On the other hand, we can check that the element $e(\sigma_{++}, w\beta, \beta)\mathfrak{s}(\sigma, \beta')e(\sigma, w\beta)$ is an isomorphism between these idempotents, since $\mathfrak{s}(\sigma, \beta')$ spans the elements that act on the polynomial ring by a linear combination of elements $\leq w$ in Bruhat order, modulo those which are a linear combination of elements $< w$. In fact, one can explicitly check that the rational function in front of w in the expression for the action of $\mathfrak{s}(\sigma, \beta')$ in the polynomial representation is invertible in this completion. This means that we can reduce to the case where x is in the image of $\iota_\beta(\underline{A}_0)$, but this is impossible, since \underline{A}_0 is complementary to $\mathcal{I}_{>0}$ and they cannot have any non-trivial vectors in common.

This shows that the products $X(\sigma, \tau)H(\tau, \tau')Y(\tau', \sigma')$ for $\sigma, \sigma', \tau, \tau'$ arbitrary give a basis of A_I .

The other conditions: We need that $X(\sigma, \sigma) = \{e(\sigma)\} = Y(\sigma, \sigma)$. This follows immediately from Lemma 2.26, since if $\sigma = \sigma_{++}, \beta'$, we must have $\beta' = \beta_\sigma$, so $\mathfrak{s}(\sigma, \beta') = e(\sigma)$. Similarly, Lemma 2.26 shows that $X(\sigma, \tau) = Y(\tau, \sigma)$ can only be non-empty if $\beta_\tau \geq \beta_\sigma$, and $H(\sigma, \sigma')$ by definition is only non-empty if $\beta_\sigma = \beta_{\sigma'}$. The finiteness condition is automatic since all the sets involved are finite. \square

We call the modules $\Delta(\sigma) = Ae(\sigma)/A\mathcal{I}_{>\beta_\sigma}e(\sigma)$ the **standard modules** over A . The results above show that the modules $\Delta(\sigma)$ are a stratifying system. That is:

Lemma 2.31

- (1) *If there is a non-zero homomorphism $Ae(\sigma) \rightarrow \Delta(\sigma')$, then we must have $\beta_\sigma = \beta_{\sigma'}$ or $|\beta_\sigma| < |\beta_{\sigma'}|$.*
- (2) *Every simple module is a quotient of $\Delta(\sigma)$ for some σ .*
- (3) *The kernel of the map $Ae(\sigma) \rightarrow \Delta(\sigma)$ has a filtration with each subquotient equal to $\Delta(\sigma')$ with $|\beta_{\sigma'}| > |\beta_\sigma|$.*

Proof. (1) The space of such homomorphisms is $e(\sigma)\Delta(\sigma')$, which is spanned by the image of $E_{\beta_{\sigma'}}$. This can only be non-zero if the restriction of the sign vector σ with respect to $\beta_{\sigma'}$ is weakly ξ_β -bounded. Lemma 2.26 shows that this can only happen if $\beta_\sigma = \beta_{\sigma'}$ or $|\beta_\sigma| < |\beta_{\sigma'}|$.

(2) For any simple, there must be a sign vector σ such that $e(\sigma)L \neq 0$, and $|\beta_\sigma|$ is maximal with respect to this property. This shows that $\mathcal{I}_{>\beta}$ acts trivially on this module and so we have an induced map $\Delta(\sigma) \rightarrow L$, which is surjective by simplicity.

(3) Filter $Ae(\sigma)$ by the submodules $\mathcal{I}_{\geq\beta}e(\sigma)$. In the subquotient $\mathcal{I}_{\geq\beta}e(\sigma)/\mathcal{I}_{>\beta}e(\sigma)$, the vectors $\mathfrak{s}(\sigma, \beta')$ for $\beta' \in W\beta$ each generate a copy of $\Delta(\sigma_{++}, \beta')$, and $\mathcal{I}_{\geq\beta}e(\sigma)/\mathcal{I}_{>\beta}e(\sigma)$ is a direct sum of these submodules. \square

If M, N are $A/\mathcal{I}_{>\beta}$ -modules, then we can consider Ext between these modules over $A/\mathcal{I}_{>\beta}$ or by inflating them to modules over A . This inflation functor is denoted i in [Bru23, §7]. In general, these can be quite different, but in this case, a standard argument (see [Bru23, Lem. 7.4]) shows that:

Lemma 2.32 $\mathrm{Ext}_{A/\mathcal{F}_{>\beta}}^i(M, N) \cong \mathrm{Ext}_A^i(M, N)$.

This stratification property will be important for us below.

2.7. (Dual) canonical bases. In this section, we'll assume that $k = \mathbb{C}$. In this case, the Steinberg algebra has an interpretation in terms of G -equivariant D-modules on V . This corresponds to the sheaf-theoretic interpretation discussed before by the Riemann-Hilbert correspondence.

Let D_V be the ring of differential operators on V , and $\mu_q: U(\mathfrak{g}) \rightarrow D_V$ the ring homomorphism sending X to the corresponding vector field on V . A **strongly equivariant D-module** M on V is one where the action of \mathfrak{g} on M via the homomorphism μ_q integrates to a G -equivariant structure on M . A strongly equivariant D-module is the same thing as a D-module on the Artin stack V/G , and more generally, if for some character $\xi: \mathfrak{g} \rightarrow \mathbb{C}$, we have that the Lie algebra action by $\mu_q - \xi$ integrates to our equivariant structure, then this is a twisted D-modules on V/G , with twist specified by the image of ξ under the Kirwan map. It will be important for use that we work in the derived category of D-modules on the quotient. This has the effect that anywhere cohomology appears in the calculation of Hom spaces between D-modules, equivariant cohomology appears instead.

Consider the union $X_{I'} = \sqcup_{\sigma \in I'} X_\sigma$ and let $p: X \rightarrow V$ be the projection to the second factor. Let $L = p_* \mathfrak{S}_X$ be the pushforward of the structure sheaf on X considered as a D-module by this proper map and $L_\sigma = p_* \mathfrak{S}_{X_\sigma}$, considered as a complex of D-modules on V/G , that is an object in the derived category \mathcal{D} of strongly G -equivariant D-modules on V , which we consider with its usual dg-enhancement. We will sometimes want to consider a set P equipped with a map $P \rightarrow I'$, and the corresponding object $L_P = \bigoplus_{\sigma \in I'} L_\sigma$.

As discussed earlier, the isomorphism [CG97, Thm. 8.6.7] together with the Riemann-Hilbert correspondence shows that:

Proposition 2.33 *We have a isomorphism of dg-algebras $A_P \cong \mathrm{Ext}^\bullet(L_P, L_P)^{\mathrm{op}}$ where the left hand side is thought of as a dg-algebra with trivial differential.*

Remark 2.34. It's also true that $A_P \cong \mathrm{Ext}^\bullet(L_P, L_P)$ by reversing the two factors of the fiber product, or from a sheaf-theoretic perspective, applying Verdier duality. We prefer to think of it as an opposite algebra, so L is a right module object over A_P .

Since L is a sum of shifts of simple D-modules by the Decomposition Theorem [BBD82], this shows that:

Corollary 2.35 *The graded algebra A_P is graded Morita equivalent to an algebra ${}^+A$ which is semi-simple commutative in degree 0, and non-negatively graded.*

In fact, [Web15, Lem. 1.18] implies that graded projective A_P -modules are a **mixed humorous category** in the sense of [Web15, Def. 1.2 & 1.11]. Furthermore, Proposition 2.19 shows that the set of novel sign vectors σ supply a collection of objects satisfying [Web15, Lemma 1.6(1)].

Thus by [Web15, Lemma 1.13], we have that:

Corollary 2.36 ([Web15, Lem. 1.13 & Cor. 2.4])

- (1) *The classes of indecomposable projectives over A_P form a canonical basis in the Grothendieck group of graded projective finitely generated A_P -modules $K^0(A_P\text{-gpm})$; in particular, they are uniquely characterized by being bar-invariant, almost orthogonal, and having positive virtual dimension.*
- (2) *The classes of simple modules over A_P form the dual canonical basis of the Grothendieck group of finite-dimensional graded A_P -modules $K^0(A_P\text{-fgm})$; that is, they are uniquely characterized by the same properties, but for the dual precanonical structure.*

In particular, this means that the graded Cartan matrix of the algebra A_P can be found using a generalization of the Krull-Schmidt algorithm: just as you can orthonormalize a collection of vectors “one at a time,” you can similarly make a collection of vectors in a $\mathbb{Z}[q, q^{-1}]$ -module with a bar involution almost orthogonal and bar-invariant.

2.8. Consequences of Hodge theory. We can also interpret A_P -modules in terms of D-modules. We have a dg-functor

$$L_P \overset{L}{\otimes}_{A_P} -: A_P\text{-dg-mod} \rightarrow \mathcal{D}.$$

This functor is fully faithful, since it induces an isomorphism

$$\text{Ext}_{A_P}^\bullet(A_P, A_P) \cong A_P \cong \text{Ext}_{\mathcal{D}}^\bullet(L_P, L_P)^{\text{op}}.$$

Its image is, by definition, the subcategory $\langle L \rangle \subset \mathcal{D}$ generated by L .

We can strengthen this result by incorporating Hodge theory. The object L has a unique pure Hodge structure of weight 0 induced by that for the D-module of functions on X , defined by the usual variation of Hodge structure on the trivial line bundle. This induces a Hodge structure on $A_P = \text{Ext}^\bullet(L, L)^{\text{op}} \cong H_*^{BM,G}(\mathbb{X}_P)$ which agrees with the geometric Hodge structure on $H_*^{BM,G}(\mathbb{X}_P)$, using the identification of $H_k^{BM}(Y) \cong H_{n-k}^*(Y', Y' - Y)$ where $Y \hookrightarrow Y'$ is an inclusion into a smooth n -dimensional variety Y' .

Lemma 2.37 *The induced Hodge structure on $A_P = \text{Ext}^\bullet(L, L)^{\text{op}}$ is Tate and pure of weight 0.*

Proof. Consider the singular variety $X_\sigma \times_V X_{\sigma'}$. This has a natural map $X_\sigma \times_V X_{\sigma'} \rightarrow G/B \times G/B$, which over each G -orbit is a G -equivariant vector bundle of the form

$\tilde{H}_w = \{(gB, gwB, x) \mid x \in gV_\sigma \cap gwV_{\sigma'}\}$. Letting w range over W gives all the distinct G -orbits $H_w = \{(gB, gwB) \mid g \in G\}$ on $G/B \times G/B$. We can divide $G/B \times G/B$ into subvarieties Z_m given by the union of the G -orbits of dimension m , and consider the corresponding partition of $X_\sigma \times_V X_{\sigma'}$ by the preimages \tilde{Z}_m given of these. Considering the filtration by $\bigcup_{k \leq m} \tilde{Z}_k$, we have a spectral sequence computing the equivariant Borel-Moore homology of $X_\sigma \times_V X_{\sigma'}$ whose E_2 -page is given by

$$E_2^{pq} = H_{p+q}^{BM,G} \left(\bigcup_{m \leq p} \tilde{Z}_m, \bigcup_{m \leq p-1} \tilde{Z}_m \right) \cong H_{p+q}^{BM,G}(\tilde{Z}_p).$$

This sequence collapses at the E_2 page for parity reasons: we have

$$H_{p+q}^{BM,G}(\tilde{H}_w) \cong H_{p+q}^{BM,B \cap wBw^{-1}}(V_\sigma \cap wV_{\sigma'})$$

and this is only non-zero for $p+q$ even since $B \cap wBw^{-1}$ is homotopy equivalent to its Levi complement T , which is reductive. In fact, the equivariant Borel-Moore homology $H_{p+q}^{BM,B \cap wBw^{-1}}(V_\sigma \cap wV_{\sigma'})$ is free of rank 1 as a module over $H^*(BT)$, generated by the fundamental class. Note that this shows that $H_{p+q}^{BM,G}(\tilde{Z}_p)$ is free as a $H^*(BG)$ module. Since the equivariant cohomology of a point $H_B^*(pt)$ is pure of weight 0 and Tate, the same is true of $H_{p+q}^{BM,G}(\tilde{Z}_p)$. The same follows for $H_*^{BM,G}(\mathbb{X}_P)$ using the spectral sequence above and the fact that any extension of pure Tate mixed Hodge modules of a given weight is again pure Tate of the same weight. \square

In more informal terms, this means that we lose nothing by thinking of this Hodge structure as a second grading on A_P , which coincides with the homological grading, which we call the **Hodge grading**. It might seem odd to have another name for this grading, but we will be interested in bigraded modules for this pair of gradings, on which the two gradings will typically not coincide.

More precisely, recall that a **differential-graded-graded algebra** (dgg-algebra) is an algebra with a \mathbb{Z}^2 grading, equipped with a differential whose degree is $(1, 0)$; that is, it is a dg-algebra with an additional grading in which the differential is homogeneous. A dgg-module over a dgg-algebra is defined similarly.

The Hodge grading makes A_P into a differential-graded-graded algebra. A dgg-module over A_P can also be thought of as a complex of graded A_P -modules by sending an element of grading (r, s) to one of internal grading s and homological grading $r - s$. This is chosen so that the action of A_P preserves the new homological grading, while keeping the differential degree 1 in this grading.

Certain constructions become easier if we replace A_P by the unique Morita equivalent basic graded algebra ${}^+A_P$ of Corollary 2.35. Note that A_P only depends up to Morita equivalence on the image of P in I' , and thus ${}^+A_P$ only depends on this image. In particular, the grading on ${}^+A_P$ is non-negative, and $({}^+A_P)_0$ is semi-simple, and isomorphic to the sum of one copy of each simple module.

This Hodge structure has an important consequence for the structure of the category \mathcal{D} . Thus far, we have only discussed $\text{Ext}^\bullet(L_P, L_P)^{\text{op}}$ as a usual graded algebra, but in

fact it is the cohomology of a canonical (up to quasi-equivalence) dgg-algebra which we will denote $\mathrm{Hom}_{\mathcal{D}}^{\bullet, \bullet}(L, L)$.

First note that in the category of mixed Hodge modules, we can obtain a **mixed Hodge dg-algebra** that is, a dg-algebra internal to the abelian tensor category of mixed Hodge structures (on vector spaces). Since mixed Hodge modules don't have enough projectives as an abelian category, this is most canonically done by considering the derived category as the dg-quotient of the dg-category of all complexes by the subcategory of exact complexes as described by Drinfeld [Dri04, §3.1]. Next, we can apply the Deligne splitting to each term in this dg-algebra. The Deligne splitting of any mixed Hodge structure gives us a bigraded vector space, but since all Hodge structures appearing in our story are Tate, we can simplify by collapsing to a single weight grading.

The Deligne splitting applied to this mixed Hodge dg-algebra gives a dgg-algebra $\mathrm{Hom}_{\mathcal{D}}^{\bullet, \bullet}(L, L)$ by [CS22, Prop. 5.4], with the first grading the usual homological grading and the second given by the weight grading. The result *loc. cit.* also shows that the A_{∞} -structure on the cohomology of this dgg-algebra is homogeneous of degree 0 in the weight grading. In particular:

Theorem 2.38 *If the Hodge structure on A_P is pure of weight 0, then the dgg-algebra $\mathrm{Hom}_{\mathcal{D}}^{\bullet, \bullet}(L, L)$ is formal, that is, quasi-isomorphic to its cohomology as a dgg-algebra.*

In fact, the construction of such a quasi-isomorphism is given in [Sch11, Prop. 4]: the subalgebra of spanned by elements of degree (k, ℓ) with $\ell > k$ and closed elements of degree (k, k) is a subalgebra quasi-isomorphic to the full algebra, with the cohomology algebra as a quotient.

Remark 2.39. This argument does *not* show that the corresponding Hodge dg-algebra is formal. Carlson, Clemens, and Morgan [CCM81] show that there are Kähler manifolds whose cohomology is not formal as a Hodge dg-algebra, while of course, the corresponding dgg-algebra is pure by the Hodge theorem, as shown in [Del+75]. It seems likely to the author that in our case, the Hodge structure is in fact formal, due to the presence of so many nice properties (parity vanishing, in particular) in $H_*^{BM, G}(\mathbb{X}_P)$, but proving this will require the intervention of someone with greater expertise in Hodge theory.

2.9. Quantum Hamiltonian reduction. Now, we wish to pass from the category \mathcal{D} to one which is closer to the Higgs branch itself. Morally, we would wish to consider the category of modules over a quantization of the GIT quotient $\mathfrak{M}_{H, \xi}$, but as discussed in the introduction, for many pairs (G, V) , this GIT quotient is badly behaved.

We will need a bit of preparation to describe the subcategory of interest to us. Consider the left ideal $J_{\xi'} = \sum D_V(\mu_q(X) - \xi'(X))$. For a given $\xi: \mathfrak{g} \rightarrow \mathbb{C}$, we can define the functor

$$K_{\xi'}(M) = \{m \in M \mid \mu_q(X)m = \xi'(X)m \forall X \in \mathfrak{g}\} = \mathrm{Hom}(D_V/J_{\xi'}, M).$$

Since the G -equivariant structure on M is strong, $K_0(M) = M^G$, and $K_{k\xi}(M) = M^{G,k\xi}$, the space of semi-invariants for this character. Note that $K_\xi(M)$ is naturally a module over the Hamiltonian reduction $A_{\xi'} = (D_V/J_{\xi'})^G$.

Definition 2.40 Let $K(M) = \bigoplus_{k \geq 0} K_{k\xi}(M)$ considered as a module over the \mathbb{Z} -algebra

$$Z = \bigoplus_{m \geq k \geq 0} (D_V/J_{k\xi})^{G,(m-k)\xi}.$$

Let $\mathcal{D}^{\text{ured}}$ be the full subcategory of objects such that for some integer m , the reduction $K_{k\xi}(M) = 0$ for all $k > m$; that is, those M where $K(M)$ is bounded in the sense of [BPW16, §5].

Definition 2.41 Let $\mathfrak{D}_g = \mathfrak{D}_g(G, V)$ be the quotient $\mathcal{D}/\mathcal{D}^{\text{ured}}$.

Since $\mathcal{D}^{\text{ured}}$ is a Serre subcategory, we can also write the (dg-enhanced) derived category $D^b(\mathfrak{D}_g(G, V))$ as a dg-quotient $D^b(\mathcal{D})/D^b(\mathcal{D}^{\text{ured}})$.

Lemma 2.42 The functor $M \mapsto \bigoplus_{k \geq 0} K_{k\xi}(M)$ is an equivalence

$$\mathcal{D}/\mathcal{D}^{\text{ured}} \cong Z\text{-mod}/Z\text{-mod}_{\text{bd}}$$

to the quotient of the category of Z -modules by the subcategory of bounded modules.

Proof. Consider $\mathbf{D} = \bigoplus_{k \geq 0} D_V/J_{k\xi}$ as a right Z -module. We can write the functor K as the Hom-space $\text{Hom}_{D_V}(\mathbf{D}, -)$. Note that by definition, $\text{Hom}_{D_V}(\mathbf{D}, -)$ sends $\mathcal{D}^{\text{ured}}$ to $Z\text{-mod}_{\text{bd}}$, and so induces a functor $\mathcal{D}/\mathcal{D}^{\text{ured}} \rightarrow Z\text{-mod}/Z\text{-mod}_{\text{bd}}$.

We wish to show that this is invertible. To do this, we consider the tensor product $K_1(N) = D \otimes_Z N$, which is the left adjoint to $\text{Hom}_{D_V}(\mathbf{D}, -) = \bigoplus_{k \geq 0} K_{k\xi}(-)$.

For a Z -module N and D_V -module M , we have natural maps:

$$\iota: N \rightarrow \text{Hom}_{D_V}(\mathbf{D}, \mathbf{D} \otimes_Z N) \quad \epsilon: \mathbf{D} \otimes \text{Hom}_{D_V}(\mathbf{D}, M) \rightarrow M$$

given by the unit and counit of the usual adjunction.

The kernel of ι is bounded: There are finitely many elements $n_i \in N_{m_i}$ which generate N and thus $\mathbf{D} \otimes_Z N$ as well. Let j be the maximal degree of these generators. By the Noetherian property, N is finitely presented in terms of the relations $\sum_i r_{pi} n_i$ for $p = 1, \dots, s$. The module $\mathbf{D} \otimes_Z N$ is generated by the same generators, subject to the same relations. Let j' be the maximum of the degrees of these relations.

If $n \in N_k$ is in the kernel of this map, then we must have that $1 \otimes n = 0 \in \mathbf{D} \otimes_Z N$. Thus, one must be able to write $1 \otimes n = \sum_{p,i} a_p r_{pi} \otimes n_i$ for some $a_p \in D_V$. We can replace a_p by their projection a'_p to the semi-invariants, since n and n_i are themselves semi-invariant. Thus, if $k \geq j$, the element a'_p is semi-invariant for a positive multiple of ξ , so we can move it over the tensor and find $n = \sum_{p,i} a'_p r_{pi} n_i = 0$. Thus, the kernel of ι is bounded above by j' .

The cokernel of the map ι is bounded: On the other hand, the elements in degree k of the cokernel of ι are sums $a = \sum a_i \otimes n_i$. As above, we can assume that a_i

are semi-invariants; if $k \geq j$, then they are all semi-invariant for non-negative multiples of ξ , and so $a = \iota(\sum a_i n_i)$. Thus, the cokernel is bounded.

Completing the proof: The functor $\mathbf{D} \otimes_Z -$ sends $Z\text{-mod}_{\text{bd}}$ to $\mathcal{D}^{\text{ured}}$ and thus induces a functor $Z\text{-mod}/Z\text{-mod}_{\text{bd}} \rightarrow \mathcal{D}/\mathcal{D}^{\text{ured}}$. This functor is fully faithful by adjunction. Thus, this functor is an equivalence if and only if any object in \mathcal{D} whose image is in $Z\text{-mod}_{\text{bd}}$ lies in $\mathcal{D}^{\text{ured}}$, but this is just the definition. This completes the proof. \square

Following the proof of [BPW16, Th. 5.8], we can interpret the quotient $\mathcal{D}/\mathcal{D}^{\text{ured}}$ as the modules over the sheaf of algebras on $\mathfrak{M}_{H,\xi}$ defined by quantum Hamiltonian reduction as in [BPW16, §3.4]. This is the sense in which this gives a “geometric category” but we will not use this interpretation extensively. Note that if the quotient is badly behaved, this sheaf of algebras may not be a quantization of the structure sheaf in the sense of [BK04]. This endows an object in $\mathcal{D}/\mathcal{D}^{\text{ured}}$ with an important structure: its **support** and more generally **characteristic cycle**.

- This is thought of most naturally as the support of the corresponding sheaf on $\mathfrak{M}_{H,\xi}$.
- Readers more comfortable with projective geometry can follow the approach of Gordon-Stafford, and take the associated graded of a good filtration on the Z -module $K(M)$, arrive at a module over the projective coordinate ring of $\mathfrak{M}_{H,\xi}$ (i.e. the semi-invariants) and localize this to a coherent sheaf.
- Probably most familiar to readers familiar with D-modules, we can take the usual singular support for a D-module in \mathcal{D} ; the singular support of any strongly equivariant D-module will lie in $\mu^{-1}(0)$, and will always contain the preimage of the support as defined above. If G acts freely on $\mu^{-1}(0)$, then the intersection of the singular support with the semi-stable locus will exactly be the preimage of this support, but in general, we cannot rule out that it is larger.

A related category $\mathcal{D}^{\text{uns}} \subset \mathcal{D}$ is the subcategory of D -modules whose singular support is contained in the unstable locus $\mu^{-1}(0)^{\text{uns}}$. In [MN14a], McGerty and Nevins use $\mathcal{D}/\mathcal{D}^{\text{uns}}$ as their version of the reduced category, and in [MN14a, Prop. 4.9], they show that if the action of G on the semi-stable locus $\mu^{-1}(0)^{\text{ss}}$ is free, the moment map is flat, and $\mathfrak{M}_{H,\xi} \rightarrow \mathfrak{M}_H$ is a symplectic resolution, then this category can be interpreted as a category of sheaves over a quantization of $\mathfrak{M}_{H,\xi}$. This latter set of conditions is needed to ensure that the quantum Hamiltonian reduction will be a quantization of the structure sheaf, and in this case, we can see that we recover McGerty and Nevins’s result:

Lemma 2.43 *We have a containment $\mathcal{D}^{\text{uns}} \subset \mathcal{D}^{\text{ured}}$. If the action of G on the semi-stable locus $\mu^{-1}(0)^{\text{ss}}$ is free, then $\mathcal{D}^{\text{uns}} = \mathcal{D}^{\text{ured}}$.*

Proof. Containment $\mathcal{D}^{\text{uns}} \subset \mathcal{D}^{\text{ured}}$: Choose any good filtration on M . If $M \in \mathcal{D}^{\text{uns}}$, we have that any element of the associated graded is killed by some power of the ideal generated by $\mathbb{C}[\mu^{-1}(0)]^{G,k\xi}$ for $k > 0$. If we consider the semi-invariants $\bigoplus_{k \in \mathbb{Z}} \text{gr } M^{G,k\xi}$ as a module over $\bigoplus_{k \in \mathbb{Z}} \mathbb{C}[\mu^{-1}(0)]^{G,k\xi}$, it will be finitely generated, since $\text{gr } M$ is finitely generated and G is reductive. Thus, if m' is maximal such that $\text{gr } M^{G,m'\xi}$ contains a generator, and m'' is maximal such that an element of $\mathbb{C}[\mu^{-1}(0)]^{G,m''\xi}$ has non-zero action on $\text{gr } M$, then $K_{k\xi}(M) = 0$ for all $k > m = m' + m''$.

Opposite containment: On the other hand, assume that the action on the semi-stable set is free. If $M \notin \mathcal{D}^{\text{uns}}$, there is an element of M^G whose image in the associated graded of M , considered as a coherent sheaf on $\mu^{-1}(0)$, is non-zero on a stable orbit. Furthermore, since this point is stable, there are functions in $\mathbb{C}[\mu^{-1}(0)]^{G,k\xi}$ for infinitely many $k \geq 0$ which do not vanish on this point. This shows that $K_{k\xi}(M)$ has non-zero associated graded for infinitely many k . \square

One more question worth addressing is that of localization. In terms of the Z -algebras above, this is the question:

- is there an integer k such that the functor $K_{k\xi}: \mathcal{D} \rightarrow A_{k\xi}\text{-mod}$ induces an equivalence $\mathfrak{D}_g \cong A_{k\xi}\text{-mod}$?

If the action of G on the semi-stable locus $\mu^{-1}(0)^{\text{ss}}$ is free, the moment map is flat, and $\mathfrak{M}_{H,\xi} \rightarrow \mathfrak{M}_H$ is a symplectic resolution, then the existence of such a parameter follows from [BPW16, Cor. B.1]. It seems likely that this will hold for all (V, G) , but we have not investigated this point in detail.

Finding the precise values of k where this equivalence holds is a much more subtle point. See [Los21, §1] for an excellent summary of the state of the art on these questions.

2.10. Category \mathcal{O} . We will be interested in a subcategory $\mathcal{O}_g \subset \mathfrak{D}_g$ which plays the role of a category \mathcal{O} . As in [BLPW16], this definition depends on a choice of \mathbb{C}^\times -action on the quotient $\mathfrak{M}_{H,\xi}$.

In this paper, we will specifically consider \mathbb{C}^\times -actions which factor through the group \tilde{F} which acts naturally on the Higgs branch $\mathfrak{M}_{H,\xi}$ for every character ξ . Of course, such an action is exactly the choice of a flavor $\phi: \mathbb{T} \rightarrow \tilde{F}$. Note that unlike in [BLPW16], such an action will have weight 1 on the symplectic form, not weight 0 like the Hamiltonian actions considered in that paper.

The group \mathbb{T} acts on the G -invariant functions of $\mu^{-1}(0)$.

Definition 2.44 *Let $p\mathcal{O}_g \subset \mathcal{D}$ be the full subcategory of objects M which have a good filtration such that each G -invariant function on $\mu^{-1}(0)$ with positive \mathbb{T} -weight acts trivially on $\text{gr } M$. Let $\mathcal{O}_g \subset \mathfrak{D}_g$ be the image of this category in the quotient.*

The category \mathcal{O}_g is what we call $\mathcal{O}_{\text{Higgs}}$ in the introduction.

Lemma 2.45 *For a cocharacter $\phi_0: \mathbb{T} \rightarrow F = H/G$, the category \mathcal{O}_g for ϕ_0 in [BLPW16, Def. 3.15] is the same as that of Definition 2.44 for the pointwise product of ϕ_0^ℓ and the action of \mathbb{S} for $\ell \gg 0$.*

Proof. This follows immediately from [BLPW16, Prop. 3.18]: the G invariant functions with positive weight under this pointwise product for $\ell \gg 0$ are those where ϕ_0 has positive weight or where ϕ_0 has weight 0 and \mathbb{S} has positive weight. Thus, these are precisely the functions in the ideal J defined in [BLPW16, §3.1]. \square

By the same argument as [Web17b, Thm. 2.18], we have that:

Lemma 2.46 *The D -module $L_{I'_\phi}$ lies in $p\mathcal{O}_g$.*

Definition 2.47 *For a given character ξ , we call a sign vector σ **weakly unsteady** if there is a non-zero cocharacter $\nu \in \mathfrak{t}_\mathbb{R}$ with $\langle \xi, \nu \rangle \geq 0$, and $(T^*V)_\nu \supset (T^*V)_\sigma$, and **unsteady** if there exists such a ν with $\langle \xi, \nu \rangle > 0$.*

For our purposes, we wish to have a more user-friendly characterization of instability.

Lemma 2.48 *If $C_\sigma \neq \emptyset$, then the sign vector σ is unsteady if and only if C_σ is ξ -unbounded, and weakly unsteady if and only if C_σ is not strongly ξ -bounded.*

Proof. We have that $\langle \xi, \nu \rangle > 0$ if and only if ξ does not attain a maximum on any ray parallel to ν . Furthermore, we have that $(T^*V)_\nu \supset (T^*V)_\sigma$ if and only if C_σ contains a ray parallel to ν . By a standard result of linear programming, C_σ is ξ -unbounded if and only if ξ does not attain a maximum on some ray in the chamber.

On the other hand, if $\langle \xi, \nu \rangle \geq 0$, then ξ might attain a maximum on a ray parallel to ν , but only by being constant on the ray. Thus we have that C_σ is not strongly ξ -bounded. \square

2.11. Character sheaves. We now define a category of G -equivariant D -modules on V which are analogous to character sheaves/ D -modules on Lie algebras. Let the **nilcone** \mathcal{N} of $\mu^{-1}(0)$ be the set of elements which are GIT unstable for the action of \tilde{Q} , with respect to the cocharacter $\nu: \tilde{Q} \rightarrow \mathbb{C}^\times$ corresponding to the action of \tilde{Q} on the line spanned by the symplectic form. That is, the elements $v \in T^*V$ where there is a cocharacter $\beta: \mathbb{C}^\times \rightarrow \tilde{Q}$ such that the limit $\lim_{t \rightarrow 0} \beta(t)v$ exists and $\nu(\beta(t)) = t^n$ for $n > 0$.

This space is naturally stratified by the Kirwan-Ness stratification. That is, every point $v \in \mathcal{N}$ has a unique choice of β as above with minimal norm. Following the notation of [MN14b, §2.1], we consider the fixed points and points with limits

$$(2.15) \quad Z_\beta = V^{\beta(\mathbb{C}^\times)} \qquad Y_\beta = V_\beta$$

$$(2.16) \quad T^*Z_\beta = (T^*V)^{\beta(\mathbb{C}^\times)} \qquad N^*Y_\beta = (T^*V)_\beta$$

Of course, Z_β, T^*Z_β are modules over the Levi subgroup L_β centralizing β , and Y_β, N^*Y_β over P_β , the parabolic in G whose Lie algebra is the sum of non-negative weight spaces for β .

We thus have maps

$$V \xleftarrow{p} G \times^{P_\beta} Y_\beta \xrightarrow{q} G \times^{L_\beta} Z_\beta$$

Definition 2.49 *The slanted induction functor $\text{Ind}_\beta: \mathcal{D}(Z_\beta/L_\beta) \rightarrow \mathcal{D}$ is*

$$\text{Ind}_\beta(M) = p_*q^*M[\dim Y_\beta - \dim Z_\beta].$$

The slanted restriction functor $\text{Res}_\beta: \mathcal{D} \rightarrow \mathcal{D}(Z_\beta/L_\beta)$ is

$$\text{Res}_\beta(M) = q_*p^*M[\dim G/P_\beta + \dim Y_\beta - \dim V].$$

This is a generalization of the notion of “spiral induction” introduced by Lusztig and Yun [LY17, §4].

We call a regular simple D-module M with singular support in \mathcal{N} **cuspidal** if $\text{Res}_\beta(M) = 0$ for any β with $(G_\beta, Z_\beta) \neq (G, V)$.

Conjecture 2.50 *Every regular simple D-module M with support in \mathcal{N} is a summand of $\text{Ind}_\beta(L')$ for a sheaf L' which is cuspidal for some β . If M is such a summand for two β, β' then for some $g \in G$, we have $G_\beta = gG_{\beta'}g^{-1}$ and $Z_\beta = gZ_{\beta'}$.*

This is proven by the adjoint representation of a semi-simple Lie algebra in [LY17, Th. 0.6]. We have thus far had no luck in proving this conjecture in full generality. It will be useful to us to prove a weaker version of this result, however:

Lemma 2.51 *If the singular support of L is contained in the unstable locus of character ν , then L is a summand of $\text{Ind}_\beta(M)$ for $\langle \nu, \beta \rangle > 0$.*

Proof. Let $Y \subset V$ be the usual support of L . The conormal bundle of the smooth locus Y° of Y lies in the support of L and thus in the unstable locus. By [Kem18, Th. 6] a generic point of this conormal has an attached optimal cocharacter β satisfying $\langle \nu, \beta \rangle > 0$, and for this β , we must have $Y = \overline{G \cdot Y_\beta}$. In this case, the map $G \times^{P_\beta} Y_\beta \rightarrow Y$ is generically an isomorphism.

This shows that L is a summand of the induction $\text{Ind}_\beta(M)$ where M is the intermediate extension of the local system on $Y^{\text{circ}} \cap Z_\beta$ induced by the restriction of L . \square

Thus, we will need to give a name to versions of it which we can prove as special cases, which we will use as hypotheses for some of our later results. The strongest of these is:

- (†) Each simple module in $p\mathcal{O}_g$ is a summand of a shift of L , every simple with unstable support is a summand of L_σ for σ unsteady, and every weakly unsteady σ is unsteady.

Note that whether (\dagger) holds depends on the choice of P . If it holds for some choice of P , then it holds for $P = I'_\phi$. A slightly weaker assumption is:

(\dagger') Each simple module M in $p\mathcal{O}_g$ such that $\text{Ext}^\bullet(L, M) \neq 0$ is a summand of a shift of L , and every such simple with unstable support is a summand of L_γ for $\gamma \in B$ unsteady.

Proposition 2.52 *If Conjecture 2.50 holds for (V, G) , then (\dagger') holds for $P = I'_\phi$. If, in addition, there are no cuspidal D -modules except in the case where $Z_\beta = 0$ and every weakly unsteady sign vector σ is unsteady, then (\dagger) holds for $P = I'_\phi$.*

Proof. By assumption, M is a summand of $\text{Ind}_\beta(M')$ for some cuspidal M' . Thus, if the Ext-group $\text{Ext}^\bullet(L, M)$ is non-zero, then $\text{Ext}(L_{\beta'}, \text{Ind}_\beta(M)) \neq 0$ where β' satisfies $Z_{\beta'} = 0$. This is, up to homological shift, the same as $\text{Res}_{\beta'}(\text{Ind}_\beta(M))$ which is zero by the cuspidality of M unless $Z_\beta = 0$ as well. In this case, we find that M is a summand of L .

Now, assume that M has unstable support and is a summand of L . By Lemma 2.51, we have that M is a summand of $\text{Ind}_{\beta'}(M')$ where $\langle \nu, \beta' \rangle > 0$. By uniqueness, we must have that M' is a summand of $\text{Ind}_{\beta''}(M'')$ where M'' is the skyscraper sheaf of $Z_{\beta''}$ for some β'' . Thus, M is a summand of $\text{Ind}_{\beta'}(\text{Ind}_{\beta''}(M'')) \cong \text{Ind}_\beta(M'')$ where $\beta = k\beta' + \beta''$ for some large $k \in \mathbb{Z}_{>0}$. For $k \gg 0$, we have $\langle \nu, \beta \rangle > 0$, so this shows that M is a summand of L_γ for $\gamma \in B$ unsteady. \square

It's shown that (\dagger) holds for the ADE quiver case in [Web17b, Th. 5.1] and in the affine type A case [Web17b, Th. 5.11]. As discussed in [Web17b, §4.3], it would follow in the general quiver case if we knew a theorem of Ginzburg and Baranovsky which was communicated privately to the author, but has not yet appeared in published form.

Theorem 2.53 *If G is abelian, then Conjecture 2.50 holds.*

Proof. In the abelian case, the group Q is just the full subgroup of diagonal matrices in some basis, and the nilcone \mathcal{N} is the union of the conormal subvarieties for the coordinate subspaces in V (that is, the subspaces which can be written as a sum of V_i 's). A perverse sheaf on V will have singular support in this variety if it is the perverse extension of a local system on a coordinate space $V_S = \sum_{i \in S} V_i$ minus the hyperplanes where any coordinate vanishes. For each $i \in S$, we have the monodromy $\alpha_i \in \mathbb{C}^\times$ around this hyperplane; since the fundamental group of this complement is $\mathbb{Z}^{|S|}$, generated by loops around these hyperplanes, the irreducible local systems are classified by α_i , and there is a local system for each choice of α_i . Let L be the perverse extension of this local system. We can divide $S = S_1 \cup S_2$ where $i \in S_1$ if $\alpha_i = 1$, and $i \in S_2$ if $\alpha_i \neq 1$.

The sheaf L is cuspidal if its pushforward by the projection $V_S \rightarrow V_{S \setminus \{i\}}$ is trivial for all i . This projection is trivial if and only if $\alpha_i \neq 1$. Thus, L is cuspidal if and only if $S = S_2$.

On the other hand, if $S_1 \neq \emptyset$, then we can choose β to have weight 1 on V_i for $i \in S_1$ and weight 0 on V_j for $j \in S_2$, and write $L = \text{Ind}_\beta(L')$, where L' is the local system on V_{S_2} with monodromy around the hyperplanes in S_2 unchanged, and obviously there is no other cuspidal simple that L is the induction of. \square

This shows that (\dagger') holds in the general hypertoric case. On the other hand, there are fewer cuspidals than the above discussion might lead you to expect if one considers G -equivariant sheaves.

First, consider the case where V' has a dense torus orbit, which necessarily consists of the elements with non-zero coordinates in the coordinates induced by any weight basis. Any simple D-module with singular support in \mathcal{N} whose support in V' is dense must be the intermediate extension of a strongly equivariant local system on this torus. Equivariant local systems are classified by representations of the stabilizer of a point in the dense orbit, and the strong equivariance condition requires that the Lie algebra of the stabilizer acts trivially, i.e. the representation factors through the component group of the stabilizer. This stabilizer is the Pontryagin dual of the quotient of weight lattice of T by the weights in V' , and so the representations of the component group are isomorphic to the torsion in this quotient.

If the GIT quotient $\mathfrak{M}_{H,\xi}$ is a smooth resolution of \mathfrak{M}_H for some ξ , then this implies that the action of T on V is unimodular, which simply means that if $V' \subset V$ is an invariant subspace with a dense torus orbit, the stabilizer of that orbit is connected (see [BD00, pp. 3.2 & 3.3]).

Theorem 2.54 *Any simple object in $p\mathcal{O}_g$ for G abelian is of the form $L = \text{Ind}_\gamma(M)$ where $\gamma \in \mathfrak{t}_{\mathbb{Q},1,\mathbb{R}}$ and the action of T on Z_γ has a dense orbit.*

By the discussion above, this implies that there are no cuspidals and (\dagger) holds if $\mathfrak{M}_{H,\xi}$ is a smooth hypertoric variety.

Proof. Let $I \subset [1, d]$ is the set of i such that $\alpha_i \neq 1$. We know from Theorem 2.53 that L is induced from a cuspidal sheaf on the subspace V_I where only the coordinates in I are non-zero. We can always Fourier transform so that the zero section is a component of the singular support, i.e. L has dense support. We must be able to choose γ so that $\varphi_i^\perp(\gamma) > 0$ for all i . Furthermore, if the monodromy around a hyperplane is non-trivial, then Fourier transforming to the conormal, we still have a sheaf with dense support, and so we must have a choice of γ' with opposite sign on φ_i^\perp . More generally, different choices of γ must realize all possible signs φ_i^\perp in \mathfrak{t}_1 .

This is only possible if the weights φ_i are all linearly independent in \mathfrak{t}^* , since a relation between them would forbid at least one pattern of signs (by Farkas' lemma, again). This shows that the action of T on V_I has a dense orbit. \square

If any non-empty c_σ has non-empty interior, then we can assume that $\gamma \in \mathfrak{t}_{1,\mathbb{R}}$; in fact, we can take it to be a vertex of some c_σ .

2.12. An algebraic description of category \mathcal{O} . Now we turn to understanding the category \mathcal{O}_g when the hypotheses (\dagger) and (\dagger') hold.

Definition 2.55 *Let $\mathcal{I}_\xi \subset A_P$ be the ideal generated by the idempotents $e(\sigma)$ with C_σ not strongly ξ -bounded.*

Remark 2.56. This ideal is very close to the ideal $\mathcal{I}_{>0}$ defined in Section 2.6, but that ideal is generated by the idempotents $e(\sigma)$ for σ which are ξ -unbounded; that is, it does not include $e(\sigma)$ for σ which are weakly ξ -bounded. Of course, if (\dagger') holds, then these ideals coincide.

This allows us to prove one of the key steps on the way to Theorem A:

Theorem 2.57 *If (\dagger') holds, then $\mathrm{RHom}^*(\mathfrak{r}(L), \mathfrak{r}(L))^{\mathrm{op}} \cong A_\xi = A_P / \mathcal{I}_\xi$ as a formal dg-algebra. That is, we have a commutative diagram of functors:*

$$\begin{array}{ccc}
 A\text{-dg-mod} & \xrightarrow{A_\xi \overset{L}{\otimes}_A -} & A_\xi\text{-dg-mod} \\
 \downarrow \overset{L}{\otimes}_A - & & \downarrow \mathfrak{r} \overset{L}{\otimes}_{A_\xi} - \\
 D_P \mathcal{O}_g & \xrightarrow{\mathfrak{r}} & D_{\mathcal{O}_g}
 \end{array}$$

with the vertical functors fully faithful.

Proof. Proof that $\mathrm{RHom}^(\mathfrak{r}(L), \mathfrak{r}(L))^{\mathrm{op}} \cong A_\xi$:* We have that the A -module $M = \mathrm{RHom}^*(L, \mathfrak{r}_*(\mathfrak{r}(L)))$ is equal to A_ξ . Since (\dagger') holds, the object $\mathfrak{r}_*(\mathfrak{r}(L))$ is uniquely characterized by:

- (1) $\mathrm{RHom}(L_\sigma, \mathfrak{r}_*(\mathfrak{r}(L))) = 0$ for σ unsteady.
- (2) There is a map $L \rightarrow \mathfrak{r}_*(\mathfrak{r}(L))$ whose cone is killed by \mathfrak{r} .

This means that M is uniquely characterized by:

- (1) $e(\sigma)M = 0$ for σ unsteady.
- (2) There is a map $A \rightarrow M$ whose cone K satisfies $\mathrm{RHom}_A(K, N) = 0$ for any N the inflation of an A_ξ -module.

The module A_ξ obviously satisfies (1). The obvious map $A \rightarrow A_\xi$ has cone $\mathcal{I}_\xi[-1]$. As discussed in Remark 2.56, this is the same at $\mathcal{I}_{>0}$. If N is the inflation of an A_ξ -module, then $\mathrm{RHom}_A(\mathcal{I}_\xi, N) = 0$ since $\mathrm{RHom}_A(A, N) \cong \mathrm{RHom}_{A_\xi}(A_\xi, N)$ is isomorphic to $\mathrm{RHom}_A(A_\xi, N)$ by Lemma 2.32. This shows that A_ξ satisfies (2).

The vertical maps are fully faithful: This follows if we have an induced isomorphism on RHom 's for a generating object. This follows for the left-hand map with L by Proposition 2.33, and for the right-hand map with $\mathfrak{r}(L)$ by the isomorphism $\mathrm{RHom}^*(\mathfrak{r}(L), \mathfrak{r}(L))^{\mathrm{op}} \cong A_\xi$.

The diagram commutes: Again, it's enough to check this commutation on A as a generating object. Going right and then down gives $A \mapsto A_\xi \mapsto \mathfrak{r}(L)$, while going down and then right gives $A \mapsto L \mapsto \mathfrak{r}(L)$, so the diagram commutes. \square

In this case, the dg-category $D_{\mathcal{O}_g}$ is a quotient of the dg-category $D_{p\mathcal{O}_g}$ by the subcategory generated by unsteady objects.

We can define a graded lift of the image of this functor in category \mathcal{O} . We consider A_ξ as a formal pure dgg-algebra. By the full faithfulness of Theorem 2.57, the category of dgg-modules over this algebra provides a graded lift of category \mathcal{O} .

Definition 2.58 *A grading on an object $M \in \mathcal{O}_g$ is a choice of a dgg-module N over A_ξ and an isomorphism $M \cong \mathfrak{r}L \overset{L}{\otimes}_{A_\xi} N$ and similarly for $p\mathcal{O}_g$ and the dgg-algebra A . Let $\tilde{\mathcal{O}}_g$ (resp. $\widetilde{p\mathcal{O}}_g$) be the category of graded objects in \mathcal{O}_g (resp. $p\mathcal{O}_g$) with morphisms given by morphisms of dgg-algebras.*

Remark 2.59. If the Hodge dg-algebra lifting A_ξ is formal, we can give a more geometric definition of a grading as a Hodge, or perhaps more naturally twistor, structure on the D-module $\mathfrak{r}_!M$, the image of M under the right adjoint. There can be many different choices of Hodge or twistor structure that give the same dgg-module, since non-Tate structures exist. Formality would allow us to define a “most Tate” structure corresponding to any dgg-module, but since this formality is uncertain, we leave the details of this approach to another time.

Theorem 2.57 is a “Koszul dual” description of the category \mathcal{O}_g , in that we have described the Ext algebra of simples. We can also give a dual description in terms of projectives. In this case, we let ${}^+A$ be the positively graded algebra from Corollary 2.35 and ${}^+A^!$ its Koszul dual. The category of graded modules over ${}^+A^!$ is equivalent to the category of linear projective complexes of ${}^+A$. Of course, this equivalence is realized by a projective object \mathbf{P} whose endomorphisms are ${}^+A^!$. Let $P = L \overset{L}{\otimes}_{{}^+A} \mathbf{P}$ be the corresponding projective in $p\mathcal{O}_g$. Let \mathbf{P}_ξ, P_ξ be the corresponding projectives in $\mathrm{LPC}({}^+A_\xi)$ and \mathcal{O}_g .

Consider the idempotent $e_\xi \in {}^+A_0$ whose image in A_0 is the simples L_i which are summands of L_σ for σ unsteady. Let $e'_\xi = 1 - e_\xi$ be the complementary idempotent. Note that we can use the same symbols to denote idempotents in the quadratic dual ${}^+A^!$. Note that we have an obvious map of the quadratic dual $A_\xi^! \rightarrow e'_\xi {}^+A e'_\xi$.

Theorem 2.60 *If (\dagger') holds, then the map above induces an isomorphism ${}^+A_\xi^! = e_\xi'^+ A e_\xi'$. Furthermore, we have a commutative diagram of functors:*

$$\begin{array}{ccc} {}^+A^!-\text{mod} & \xrightarrow{e_{\xi^-}} & {}^+A_\xi^!-\text{mod} \\ P \otimes_{{}^+A^!} - \downarrow & & \downarrow P_\xi \otimes_{{}^+A_\xi^!} - \\ p\mathcal{O}_g & \xrightarrow{\tau} & \mathcal{O}_g \end{array}$$

with the vertical functors fully faithful. This induces equivalences ${}^+A_\xi^!-\text{gmod} \cong \tilde{\mathcal{O}}_g$ and ${}^+A^!-\text{gmod} \cong \widetilde{p\mathcal{O}_g}$.

Proof. The vertical functors are fully faithful since the functor $L \otimes_A -$ is fully faithful. Thus, we have $\text{End}(P) = {}^+A^!$ and $\text{End}(P_\xi) = {}^+A_\xi^!$.

Since τ is an exact functor, its left adjoint sends projectives to projectives, and so $\tau_!(P_\xi)$ is a projective object in $p\mathcal{O}_g$, and its head is the sum of the simple modules which are not killed by τ . Thus, $\tau_!(P_\xi)$ is a summand of P , which is the projective cover of the sum of all simples, and in fact, it is isomorphic to the image of e_ξ' . Thus,

$${}^+A_\xi^! = \text{Hom}(P_\xi, P_\xi) = \text{Hom}(\tau_!(P_\xi), \tau_!(P_\xi)) = e_\xi'^+ A^! e_\xi'. \quad \square$$

Let us discuss how these results apply in the hypertoric and quiver cases. In the hypertoric case, we recover the results of [Bra+12], though from the Koszul dual perspective, i.e. the ring $A_\xi^!$ is denoted $A(\phi, \xi)$ in [Bra+12], since we are computing the endomorphisms of projectives in that paper.

In the quiver case, we have already computed the algebra A_P as a weighted KLR algebra. We can identify the unsteady sign vectors σ with unsteady idempotents in the sense of [Web19b, Def. 2.21], as we explain below. The function ξ must be a linear combination of the determinant characters on the factors \mathfrak{gl}_{v_i} with some weights ξ_i . In the case where $w_i = 0$ for all i , the copy of \mathbb{C} given by scalar matrices for a single fixed scalar λ on each vertex acts trivially, and so we must choose ξ to vanish on this subgroup, so $\sum_i \xi_i v_i = 0$.

This translates into a function $\mathcal{V} \rightarrow \mathbb{C}_+$ (called a **charge** in [Web19b]) sending $j \mapsto \xi_j + \iota \in \mathbb{C}$, where we use $\iota = \sqrt{-1}$ to avoid confusion with the use of i as an index.

Theorem 2.61 *The ideal \mathcal{I}_ξ coincides with the kernel of the map from the weighted KLR algebra $A_P \cong W_P^\vartheta$ under the isomorphism of Theorem 2.21.*

Proof. **\mathcal{I}_ξ maps trivially to the steadied quotient:** Assume that σ is unsteady, and choose a ν such that $\langle \xi, \nu \rangle \geq 0$, and $(T^*V)_\nu \supset (T^*V)_\sigma$. This means that the equivalence class of the corresponding loading is unchanged when we add the value $\nu_{i,k}$ of ν on the k th basis vector of \mathbb{C}^{v_i} to the x -coordinate of the corresponding point in the loading.

After adding $q\nu_{i,k}$ for some $q \geq 0$, this means that the points of the loading will be separated into groups, according to their weight for $\nu_{i,k}$. For each value $\nu_{i,k} = r$, we let $\Xi(r)$ be the sum of the factors ξ_i , plus $-\delta_{r,0} \sum_i \xi_i \nu_i$ to account for the red strands (i.e. the Crawley-Boevey vertex). Thus, we have that $\sum_{r \in \mathbb{Z}} r \Xi(r) = \langle \xi, \nu \rangle$. We can write:

$$\sum_{r \in \mathbb{Z}} r \Xi(r) = \sum_{r \in \mathbb{Z}} \sum_{r \leq s} \Xi(s).$$

Since the left-hand side of this equality is positive, one of the terms on the right-hand side must be positive. So, for some r , the sum $\sum_{r \leq s} \Xi(s) > 0$ and $\sum_{r > s} \Xi(s) < 0$

We can separate the points of the loading (including the red strands) into two sets:

- The set S_1 corresponds to the points where $\nu_{i,k} < r$.
- The set S_2 corresponds to those where $\nu_{i,k} \geq r$.

These sets are both non-empty since the sums of ξ_i over them are both non-zero. This decomposition shows that $c(\text{wt}(S_1)) = (\sum_{r > s} \Xi(s)) + \iota \cdot \#S_1$ has negative real part, and $c(\text{wt}(S_2)) = (\sum_{r \leq s} \Xi(s)) + \iota \cdot \#S_2$ has positive real part, so the former must have greater argument, and $\text{wt}(S_1) >_c \text{wt}(S_2)$. This shows that the idempotent $e(\sigma)$ is in the kernel of the map to the steadied quotient, as desired.

Opposite inclusion of ideals: We need to show that each unsteady idempotent in the sense of [Web19b, Def. 2.21] lies in \mathcal{I}_ξ . In this case, we have a decomposition of the points in the loading into S_1 and S_2 with all strands in S_1 and their ghosts left of all strands in S_2 and their ghosts such that $\text{wt}(S_1) >_c \text{wt}(S_2)$.

Exactly one of the subsets, either S_1 or S_2 , must contain all the red strands. If S_1 contains the red strands, we choose ν so that all elements of S_1 have $\nu_{*,*} = 0$, and those in S_2 have $\nu_{*,*} = 1$. On the other hand, if the red strands are in S_2 , we choose $\nu_{*,*} = -1$ and $\nu_{*,*} = 0$ respectively. In either case, we have $c(\text{wt}(S_1)) + c(\text{wt}(S_2)) = \iota(\#S_1 + \#S_2)$, so the real parts of $c(\text{wt}(S_1))$ and $c(\text{wt}(S_2))$ must have opposite sign, and $\text{wt}(S_1) >_c \text{wt}(S_2)$ if and only if

$$\langle \xi, \nu \rangle = \Re(c(\text{wt}(S_2)) - c(\text{wt}(S_1))) > 0,$$

because of our choice of ν . Thus, this idempotent lies in \mathcal{I}_ξ . □

3. THE COULOMB SIDE

3.1. Coulomb branch preliminaries. The Coulomb side of our correspondence is given by a remarkable recent construction of Braverman, Finkelberg, and Nakajima [Nak16; BFN18]. As we mentioned in the introduction, a more algebraic-minded reader could ignore this geometric construction and take Theorem 3.13 as a definition. We'll wish to modify this construction somewhat, so let us describe it in some detail. As before, let G be a connected reductive algebraic group over \mathbb{C} , with $G((t)), G[[t]]$ its points over $\mathbb{C}((t)), \mathbb{C}[[t]]$. For a fixed Borel $B \subset G$, we let \mathcal{J} be the associated Iwahori subgroup

$$\mathcal{J} = \{g(t) \in G[[t]] \mid g(0) \in B\} \subset G[[t]].$$

The **affine flag variety** $\mathcal{F}\ell = G((t))/\mathcal{J}$ is just the quotient by this Iwahori.

Let V be the G -representation fixed in the previous section, and $U \subset V((t))$ a subspace invariant under \mathcal{J} . We equip $V((t))$ with a loop \mathbb{C}^\times -action such that vt^a has weight a . As in Section 2, we'll consider the group $\tilde{G} = G \times \mathbb{C}^\times$ which acts on $V((t))$ by $(g, s) \cdot vt^a = \phi_0(s)gv(st)^a$. This is compatible with the standard loop action on $G((t))$.

We'll be interested in the infinite-dimensional vector bundle on $\mathcal{F}\ell$ given by

$$\mathcal{X}_U := G((t)) \times^{\mathcal{J}} U.$$

Note that we have a natural $G((t))$ -equivariant action map $\mathcal{X}_U \rightarrow V((t))$.

Definition 3.1 *The flag BFN space is the fiber product $\mathcal{X}_{V[[t]]} \times_{V((t))} \mathcal{X}_{V[[t]]}$.*

We'll consider this as a set of triples $\mathcal{X}_{V[[t]]} \times_{V((t))} \mathcal{X}_{V[[t]]} \subset V((t)) \times \mathcal{F}\ell \times \mathcal{F}\ell$. Recall that $H/C_{\text{GL}(V)}(G) \cong G/Z(G)$, so we can define an action of $H((t))$ on $V((t)) \times \mathcal{F}\ell \times \mathcal{F}\ell$, with the action on the second two terms factoring through this quotient. Furthermore, we have a natural action of the loop rotation \mathbb{C}^\times which commutes past this in the usual way, so we have an action of $H((t)) \rtimes \mathbb{C}^\times$. The group $\tilde{H} = H \times \mathbb{C}^\times$ has a natural inclusion in $H((t)) \rtimes \mathbb{C}^\times$ induced by the inclusion $H \hookrightarrow H((t))$ on the first factor and the identity on the factors of \mathbb{C}^\times . This induced a further inclusion $\tilde{G} \subset \tilde{H} \hookrightarrow H((t)) \rtimes \mathbb{C}^\times$.

Let $\widetilde{G((t))}$ be the subgroup of $H((t)) \times \mathbb{C}^*$ generated by $G((t))$ and the image of \tilde{G} .

We'll want to consider the equivariant homology $H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_{V[[t]]} \times_{V((t))} \mathcal{X}_{V[[t]]})$. Defining this properly is a finicky technical issue, since the space $\mathcal{X}_{V[[t]]} \times_{V((t))} \mathcal{X}_{V[[t]]}$ can be thought of as a union of affine spaces which are both infinite-dimensional and infinite-codimensional, making it hard to define their degree in homology. First, we note that it is technically more convenient to consider the space

$${}_{V[[t]]}\mathcal{X}_{V[[t]]} = \{(g\mathcal{J}, v(t)) \in \mathcal{F}\ell \times V[[t]] \mid g^{-1} \cdot v(t) \in V[[t]]\}.$$

This is the intersection of the trivial bundle $\mathcal{F}\ell \times V[[t]]$ with the image of the associated bundle $G((t)) \times^{\mathcal{J}} V[[t]]$ in $\mathcal{F}\ell \times V((t))$. Basic properties of equivariant homology lead us to expect that

$$H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_{V[[t]]} \times_{V((t))} \mathcal{X}_{V[[t]]}) \cong H_*^{BM, T_{\tilde{G}}}({}_{V[[t]]}\mathcal{X}_{V[[t]]});$$

we will use this as a definition of the left-hand side. The preimage in ${}_{V[[t]]}\mathcal{X}_{V[[t]]}$ of a Schubert cell in $\mathcal{F}\ell$ is a cofinite dimensional affine subbundle of $\mathcal{X}_{V[[t]]}$; thus, using both the dimension of the Schubert cell and the codimension of the affine bundle, we can make sense of the difference between the dimensions of these cells. With a bit more work, this allows us to make precise the notion of this homology, as in [BFN18, §2(ii)]. For our purposes, we can use their construction as a black-box, only knowing that basic properties of pushforward and pullback operate as expected.

Definition 3.2 *The Iwahori Coulomb branch algebra \mathcal{A} is the equivariant Borel-Moore homology $H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_{V[[t]]} \times_{V((t))} \mathcal{X}_{V[[t]]})$.*

An important special case of this algebra has also been considered in [APM20, §4], when the representation is of quiver type (as discussed in Section 2.5). As usual, we let h be the equivariant parameter corresponding to the character ν . Note that this algebra contains a copy of $S_h = S[h] \cong \mathbb{k}[\tilde{\mathfrak{t}}]$, the coordinate ring of $\tilde{\mathfrak{t}}$, embedded as $H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_{V[[t]]}) \cong H_*^{\tilde{\mathfrak{t}}}(*)$. The algebra \mathcal{A} also possesses a natural action on this cohomology ring.

The original BFN quantum Coulomb branch algebra \mathcal{A}^{sph} is defined in essentially the same way, using $\mathcal{Y}_{V[[t]]} := G((t)) \times^{G[[t]]} V[[t]]$. Pullback by the natural map $\mathcal{X}_{V[[t]]} \rightarrow \mathcal{Y}_{V[[t]]}$ defines a homomorphism $\mathcal{A}^{\text{sph}} \rightarrow \mathcal{A}$.

Theorem 3.3 *The algebras \mathcal{A}^{sph} and \mathcal{A} are Morita equivalent. In fact, the latter is a matrix algebra over the former of rank $\#W$.*

We can geometrically realize the \mathcal{A}^{sph} - \mathcal{A} - and \mathcal{A} - \mathcal{A}^{sph} -bimodules of this Morita equivalence as the homologies

$$(3.1) \quad \begin{aligned} H_*^{BM, \widetilde{G((t))}}(\mathcal{Y}_{V[[t]]} \times_{V((t))} \mathcal{X}_{V[[t]]}) &\cong H_*^{BM, \tilde{G}}(V[[t]] \mathcal{X}_{V[[t]]}) \\ H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_{V[[t]]} \times_{V((t))} \mathcal{Y}_{V[[t]]}) &\cong H_*^{BM, T_{\tilde{G}}}(V[[t]] \mathcal{Y}_{V[[t]]}). \end{aligned}$$

Proof. Consider a fiber bundle $p: X \rightarrow Y$ such that the pushforward of \mathbb{k}_X to Y is a sum of shifted constant sheaves: $p_*\mathbb{k}_X \cong \bigoplus_{i \in \mathbb{Z}} \mathbb{k}_Y[-i]^{\oplus k_i}$. In this case, the rank k_i is given by the i th Betti number of the fiber. In this case, for any map $q: Y \rightarrow Z$, we have that $q_*p_*\mathbb{k}_X \cong q_*\mathbb{k}_Y^{\oplus k}$. In this case the convolution algebra

$$H^*(X \times_Z X) \cong \text{Ext}^*(q_*p_*\mathbb{k}_X, q_*p_*\mathbb{k}_X) \cong \text{Ext}^*(q_* \bigoplus_{i \in \mathbb{Z}} \mathbb{k}_Y[-i]^{\oplus k_i}, q_* \bigoplus_{i \in \mathbb{Z}} \mathbb{k}_Y[-i]^{\oplus k_i})$$

is a matrix algebra over $H^*(Y \times_Z Y) \cong \text{Ext}^*(q_*\mathbb{k}_Y, q_*\mathbb{k}_Y)$ of rank $\sum_{i \in \mathbb{Z}} k_i$, with the grading shifted to account for the homological shifts. In particular, $H^*(X \times_Z X)$ and $H^*(Y \times_Z Y)$ are Morita equivalent via the bimodules $H^*(X \times_Z Y)$ and $H^*(Y \times_Z X)$.

We have a natural homomorphism $H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_{V[[t]]} \times_{\mathcal{Y}_{V[[t]]}} \mathcal{X}_{V[[t]]}) \rightarrow \mathcal{A}$. The map $\mathcal{X}_{V[[t]]} \rightarrow \mathcal{Y}_{V[[t]]}$ is a fiber bundle with fiber G/B , and the pullback of the fiber bundle $\mathcal{F}\ell \rightarrow G((t))/G[[t]]$ over the affine Grassmannian. Since this pushforward is equivariant for $G((t))$ and the group $G[[t]]$ is connected, it must be a sum of constant sheaves. Those readers who are made nervous by the indirectness in the definition of $H_*^{BM, \widetilde{G((t))}}$ from [BFN18] for these infinite-dimensional vector bundles can reassure themselves by checking that this fiber bundle property holds modulo any power of t as well.

Thus, the convolution algebra $H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_{V[[t]]} \times_{\mathcal{Y}_{V[[t]]}} \mathcal{X}_{V[[t]]})$ is a matrix algebra over $H_*^{BM, \widetilde{G((t))}}(\mathcal{Y}_{V[[t]]}) \cong S_h$ of rank $\#W$. As a consequence, we can write the identity in this algebra as a sum of $\#W$ orthogonal and isomorphic idempotents $e_1, \dots, e_{\#W}$;

we can assume that the image of e_1 in $H_*^{BM, \widetilde{G}((t))}(\mathcal{X}_{V[[t]]})$ is the image under pullback of $H_*^{BM, \widetilde{G}((t))}(\mathcal{Y}_{V[[t]]})$. The image e of e_1 in $H_*^{BM, \widetilde{G}((t))}(\mathcal{X}_{V[[t]]} \times_{\mathcal{Y}_{V[[t]]}} \mathcal{X}_{V[[t]]})$ gives an idempotent $e \in \mathcal{A}$ such that $\mathcal{A}e\mathcal{A} = \mathcal{A}$ and $\mathcal{A}^{\text{sph}} \cong e\mathcal{A}e$, thus these algebras are Morita equivalent. \square

Let \mathcal{A}_{ab} be the BFN Steinberg algebra for the pair (T, V) . This is the equivariant homology of ${}_{V[[t]]}\mathcal{X}_{V[[t]]}^{\text{ab}} = \{(g, v(t)) \in T((t))/T[[t]] \times V[[t]] \mid g^{-1}v(t) \in V[[t]]\}$, which we can view as a subspace of ${}_{V[[t]]}\mathcal{X}_{V[[t]]}$. By [BFN18, Lem. 5.5], pushforward induces an action of left action of this algebra on the homology group Eq. (3.1). Since \mathcal{A} is the full endomorphism algebra of that right module structure, this implies that:

Lemma 3.4 *Pushforward induces an injective algebra map making the diagram*

$$\begin{array}{ccc} \mathcal{A}_{\text{ab}} & \longrightarrow & \mathcal{A} \\ \downarrow & & \downarrow \\ H_*^{BM, T_{\tilde{G}}}({}_{V[[t]]}\mathcal{X}_{V[[t]]}^{\text{ab}}) & \longrightarrow & H_*^{BM, T_{\tilde{G}}}({}_{V[[t]]}\mathcal{X}_{V[[t]]}) \end{array}$$

commute.

3.2. The extended category. While the Coulomb branch is our focus, it is easier to study it in a larger context: there is an extended category in which it appears as the endomorphisms of one object. This extended category has a simpler set of relations and thus is easier to get a handle on than the Coulomb branch on its own.

Before giving this definition, let us say a word or two about motivation. The Coulomb branch naturally arises in a 3-dimensional quantum field theory as an algebra of local operators of a topological twist. This is the Hilbert space of S^2 in the twist, as shown by usual cut-and-glue techniques.

In physical terms, our extended category considers these local operators as the endomorphisms of a trivial line defect and studies such operators in the context of homomorphisms to other line defects. Line defects in these categories are discussed in more detail in work of Dimofte, Garner, Geracie, and Hilburn [Dim+20] and forthcoming work of Hilburn and Yoo [HY].

Definition 3.5 *Let $\mathfrak{t}_{1, \tilde{H}} = \{\psi \mid d\nu(\psi) = 1\} \subset \mathfrak{t}_{\tilde{H}, \mathbb{R}}$ be the set of real cocharacters which act with weight 1 on the loop parameter t . Note that $\mathfrak{t}_{1, \mathbb{R}} \subset \mathfrak{t}_{1, \tilde{H}}$; we'll also be interested in the coset $\mathfrak{t}_{\tau} = \tau + \mathfrak{t}_{\mathbb{R}} \subset \mathfrak{t}_{1, \tilde{H}}$.*

Given any $\eta \in \mathfrak{t}_{1, \tilde{H}}$, we can consider the induced action on the vector space $V((t))$.


Definition 3.6 *Let \mathcal{J}_{η} be the subgroup whose Lie algebra is the sum of non-negative weight spaces for the adjoint action of η on $\mathfrak{g}((t))$. This only depends on the alcove in which η lies, i.e. which chamber of the arrangement given by the hyperplanes $\{\alpha(\eta) =$*

$n \mid \alpha \in \Delta_0, n \in \mathbb{Z}$ contains η ; the subgroup \mathcal{J}_η is an Iwahori if η does not lie on any of these hyperplanes.

Let U_η be the subspace of elements of weight $\geq -1/2$ under η . This subspace is closed under the action of \mathcal{J}_η . This depends only on the vector \mathbf{a} such that $\eta \in C_{\mathbf{a}}$, as defined in (2.10).

We call η **unexceptional** if it does not lie on the hyperplanes $\{\varphi_i^{\text{mid}}(\eta) = n \mid n \in \mathbb{Z}\}$ and **generic** if it is unexceptional and does not lie on any of the hyperplanes $\{\alpha(\eta) = n \mid \alpha \in \Delta_0, n \in \mathbb{Z}\}$. We'll call the hyperplanes defined above which the generic points avoid the **unrolled hyperplane arrangement**. Given any unexceptional point η , it has a neighborhood in the classical topology on which $U_{\eta'} = U_\eta$; this neighborhood necessarily contains a generic point.

Example 3.i. We illustrate this arrangement in the case of our running example:



The spaces U_χ which arise will be of the form:

$$U_\chi = \{(f_1 t^a, f_2 t^b, f_3 t^a, f_4 t^b) \in \mathbb{C}^4((t)) \mid f_i \in \mathbb{C}[[t]]\}$$

for $a, b \in \mathbb{Z}$. The region where $a = b = 0$ is shaded in the diagram above.

For each generic $\eta \in \mathfrak{t}_{1, \tilde{H}}$, we can consider $\mathcal{X}_\eta := \mathcal{X}_{U_\eta} := G((t)) \times_{\mathcal{J}_\eta} U_\eta$, the associated vector bundle. The space $\mathfrak{t}_{1, \tilde{H}}$ has a natural adjoint action of $\widehat{W} = N_{G((t))}(T_G)/T_G$. In this case, we have $U_{w \cdot \eta} = w \cdot U_\eta$. Note that since $\tau(s)t^\lambda \tau(s^{-1}) = (st)^\lambda$, we have that

$$t^\lambda \eta t^{-\lambda} = \eta - \lambda.$$

The ring S_h carries an action of the extended affine Weyl group induced by the identification with $H_*^{I \times \mathbb{C}^\times}(\ast) \cong \mathfrak{t}_{\mathbb{C}} \oplus \mathbb{C} \cdot h$. On this space, the subgroup $\mathfrak{t}_{\mathbb{Z}} \subset \widehat{W}$ acts by translation times h and $W \subset \widehat{W}$ acts as usual. We let

$$\eta \mathcal{X}_{\eta'} = \{(g, v(t)) \in G((t)) \times U_{\eta'} \mid g^{-1} \cdot v(t) \in U_\eta\} / \mathcal{J}_\eta.$$

As with $v_{[[t]]}\mathcal{X}_{V[[t]]}$, we can interpret this as the intersection of the trivial bundle $G((t))/\mathcal{J}_\eta \times U_{\eta'}$ with the associated bundle $G((t)) \times^{\mathcal{J}_\eta} U_\eta$.

Definition 3.7 *Let the extended BFN category \mathcal{B} be the category whose objects are unexceptional cocharacters $\eta \in \mathfrak{t}_{1, \tilde{H}}$, and with morphisms given by*

$$\mathrm{Hom}_{\mathcal{B}}(\eta, \eta') = H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_\eta \times_{V((t))} \mathcal{X}_{\eta'}) \cong H_*^{BM, \tilde{T}}({}_\eta \mathcal{X}_{\eta'}).$$

Let \mathcal{B}_τ be the subcategory where we consider only objects in \mathfrak{t}_τ .

Remark 3.8. Without an enormous digression into quantum field theory, let us try to put this construction into a framework more familiar in that context. We can consider the D-module of functions on \mathcal{X}_η , and push this forward to form a $G((t))$ -equivariant D-module on $V((t))$. Readers who prefer stacks can interpret this as the pushforward by the map $U_\eta/\mathcal{J}_\eta \rightarrow V((t))/G((t))$. Note that we can interpret the space $V[[t]]/G[[t]]$ as the moduli space of G -principal bundles on a formal disk with a holomorphic section of the associated bundle for V .

If η is in the fundamental alcove, the space U_η/\mathcal{J}_η has a similar interpretation as the space of G -principal bundles which have a Borel reduction at the origin and a meromorphic section whose zero or pole at the origin is controlled by η . That is, the Borel reduction fixes a filtration on the fiber over the origin, such that the coefficients of poles of order $< k$ are concentrated in the elements of weight λ such that $\langle \lambda, \eta \rangle \geq k - \frac{1}{2}$. Note that while individual weight spaces in this fiber are only well-defined up to the action of the Borel, the whole direct sum is B -invariant.

As in [CG97], we can interpret $H_*^{BM, \widetilde{G((t))}}(\mathcal{X}_\eta \times_{V((t))} \mathcal{X}_{\eta'})$ as the Ext group of the D-modules constructed by the pushforward of functions via the map $U_\eta/\mathcal{J}_\eta \rightarrow V((t))/G((t))$, and convolution as the Yoneda product in this Ext group. This quotient $V((t))/G((t))$ can be interpreted as the loop space of the stack V/G ; it is argued in [Dim+20, (1.4)] that the D-modules on this space should be interpreted as line defects in the associated TQFT.

As before, this homology is defined using the techniques in [BFN18, §2(ii)].

Remark 3.9. In several theorems below, we'll only consider generic points in this category. These are easier to work with for presenting endomorphisms, and there is a very little loss of generality in doing so. As discussed above, for each unexceptional point $\eta \in \mathfrak{t}_{1, \mathbb{R}}$, there is a nearby generic point $\eta' \in \mathfrak{t}_{1, \mathbb{R}}$ with $U_{\eta'} = U_\eta$. Theorem 3.3 can be extended with the same proof to show that $\mathrm{Hom}(\eta, \eta')$ gives a Morita equivalence between $\mathrm{End}(\eta)$ and $\mathrm{End}(\eta')$. This shows that after passing to the Karoubi envelope, the categories where we have all unexceptional objects and the subcategory where we allow only generic η are equivalent.

Note that $U_\tau = V[[t]]$. Thus, there must exist a nearby generic element $o \in \mathfrak{t}_\tau$ of the fundamental alcove such that $U_o = V[[t]]$. In this case, we have that \mathcal{J}_o is the standard

Iwahori so

$$(3.2) \quad \mathcal{A}^{\text{sph}} = \text{Hom}_{\mathcal{B}}(\tau, \tau) \quad \mathcal{A} = \text{Hom}_{\mathcal{B}}(o, o).$$

Thus, this extended category encodes the structure of both of these algebras and more.

Furthermore, the category of representations of \mathcal{A} (or equivalently, \mathcal{A}^{sph}) is closely related to that of \mathcal{B} . Let M be a representation of \mathcal{B} , that is, a functor from \mathcal{B} to the category of \mathbb{k} -vector spaces. The vector space $N := M(o)$ has an induced \mathcal{A} -module structure. Since $\text{Hom}(\eta, o)$ and $\text{Hom}(o, \eta)$ are finitely generated as \mathcal{A} -modules, this functor preserves finite generation, and is in fact a quotient functor, with left adjoint given by

$$N \mapsto \mathcal{B} \otimes_{\mathcal{A}} N(\eta) := \text{Hom}(\eta, o) \otimes_{\mathcal{A}} N.$$

Note that there is a natural subcategory \mathcal{B}_{ab} (with the same objects), where the morphisms are given by

$$\text{Hom}_{\text{ab}}(\eta, \eta') \cong H_*^{BM, T \times \mathbb{C}^\times}(\{(g, v(t)) \in T((t)) \times U_{\eta'} \mid g \cdot v(t) \in U_{\eta'}\} / T[[t]]).$$

The inclusion is induced by pushforward in homology.

3.3. A presentation of the extended category. In this section, we'll define several morphisms in the category \mathcal{B} by explicit homology classes. All of these classes are defined by subspaces in ${}_{\eta}\mathcal{X}_{\eta'}$, each of which is the preimage of the closure of an \mathcal{J}_{η} -orbit in $G((t))/\mathcal{J}_{\eta'}$ or of a T -fixed point. Thus, each gives a well-defined Borel-Moore homology class.

Note that any lift of $w \in \widehat{W}$ to $G((t))$ induces an isomorphism $\mathcal{X}_{\eta} \cong \mathcal{X}_{w \cdot \eta}$ which intertwines the action maps to $V((t))$, given by $(g, v(t)) \mapsto (gw^{-1}, w \cdot v(t))$. We denote the homology class of the graph of this isomorphism by y_w (this class is independent of the choice of lift). As a cycle in ${}_{\eta}\mathcal{X}_{w \cdot \eta}$, this is given by

$$(3.3) \quad y_w = [\{(w\mathcal{J}_{\eta}, v(t)) \mid v(t) \in U_{\eta'}\} / \mathcal{J}_{\eta}]$$

Consider two generic cocharacters $\eta, \eta' \in \mathfrak{t}_{1, \mathbb{R}}$. Let

$$(3.4) \quad r(\eta, \eta') = [\{(e, v(t)) \in T((t)) \times U_{\eta'} \mid v(t) \in U_{\eta'}\} / T[[t]]] \in \text{Hom}_{\text{ab}}(\eta', \eta).$$

If $\mathcal{J}_{\eta} = \mathcal{J}_{\eta'}$ (that is, the chambers are in the same alcove), this is sent to the class in $\text{Hom}(\eta', \eta)$ of the space

$$Y(\eta, \eta') = \{(e\mathcal{J}_{\eta}, v(t)) \in G((t)) \times U_{\eta'} \mid v(t) \in U_{\eta'}\} / \mathcal{J}_{\eta}$$

but this is not the case for η, η' in different alcoves. We also have the morphism $y_{\zeta} \in \text{Hom}_{\text{ab}}(\eta, \eta - \zeta)$ for $\zeta \in \mathfrak{t}_{\mathbb{Z}}$ (thought of as a translation in the extended affine Weyl group); this is just the graph of multiplication by t^{ζ} . The element denoted r^{λ} in [BFN18] is represented in our notation by $y_{-\lambda} r(o, \lambda + o) = r(-\lambda + o, o) y_{-\lambda}$.

Definition 3.10 Let $\Phi(\eta, \eta')$ be the product of the terms $\varphi_i^+ - nh$ over pairs $(i, n) \in [1, d] \times \mathbb{Z}$ such that the inequalities

$$\varphi_i(\eta) > n - \frac{1}{2} \quad \varphi_i(\eta') < n - \frac{1}{2}$$

hold. Note that we could write this using (2.1) as

$$\varphi_i^{\text{mid}}(\eta) > n \quad \varphi_i^{\text{mid}}(\eta') < n,$$

and similarly with the inequalities (3.5a-3.5b) below.

Let $\Phi(\eta, \eta', \eta'')$ be the product of the terms $\varphi_i^+ - nh$ over pairs $(i, n) \in [1, d] \times \mathbb{Z}$ such that we have the inequalities

$$(3.5a) \quad \varphi_i(\eta'') > n - \frac{1}{2} \quad \varphi_i(\eta') < n - \frac{1}{2} \quad \varphi_i(\eta) > n - \frac{1}{2}$$

or the inequalities

$$(3.5b) \quad \varphi_i(\eta'') < n - \frac{1}{2} \quad \varphi_i(\eta') > n - \frac{1}{2} \quad \varphi_i(\eta) < n - \frac{1}{2}.$$

These terms correspond to the hyperplanes that a path $\eta \rightarrow \eta' \rightarrow \eta''$ must cross twice.

Note that if h is specialized to 0, then we just get each weight φ_i raised to a power given by the number of corresponding unrolled hyperplanes crossed.

Remark 3.11. Since it has been absorbed into the notation, we should emphasize that we still have an underlying fixed flavor ϕ , used to define \tilde{G} . This flavor is implicit in the definition of φ_i^+ as elements of S_h , and thus in the products $\Phi(-, -, -)$; this is the **only** place it will appear in our presentation of the BFN category.

Proposition 3.12 The morphisms $\text{Hom}_{\text{ab}}(\eta', \eta)$ between two generic cocharacters η, η' have a basis over S_h of form $y_\zeta \cdot r(\eta + \zeta, \eta')$ for $\zeta \in \mathfrak{t}_{\mathbb{Z}}$, with the relations in the category \mathcal{B}_{ab} generated by:

$$(3.6a) \quad y_\zeta \cdot y_{\zeta'} = y_{\zeta + \zeta'}$$

$$(3.6b) \quad y_\zeta \cdot r(\eta, \eta') \cdot y_{-\zeta} = r(\eta - \zeta, \eta' - \zeta)$$

$$(3.6c) \quad \mu \cdot r(\eta, \eta') = r(\eta, \eta') \cdot \mu$$

$$(3.6d) \quad y_\zeta \cdot \mu \cdot y_{-\zeta} = \mu + h\langle \zeta, \mu \rangle$$

$$(3.6e) \quad r(\eta, \eta')r(\eta'', \eta''') = \delta_{\eta', \eta''} \Phi(\eta, \eta', \eta''')r(\eta, \eta''')$$

for arbitrary $\zeta, \zeta' \in \mathfrak{t}_{\mathbb{Z}}, \mu \in \mathfrak{t}_{\tilde{G}}^*$ and generic cocharacters $\eta, \eta', \eta'', \eta'''$.

Proof. In this case, the closed points of $T((t))/T[[t]]$ are the set t^ζ for $\zeta \in \mathfrak{t}_{\mathbb{Z}}$, and so $\text{Hom}_{\text{ab}}(\eta', \eta)$ is the direct sum of the Borel-Moore homology of the preimages of these points. Each point contributes a rank 1 free module over $H_{T_{\tilde{G}}}^*(pt) = S_h$, which is generated by the fundamental class $y_\zeta \cdot r(\eta + \zeta, \eta')$. Thus, these classes give a basis as

claimed. Equation (3.6a) is just the fact that $t^\zeta t^{\zeta'} = t^{\zeta+\zeta'}$. Equation (3.6b) is the fact that

$$t^\zeta Y(\eta, \eta') t^{-\zeta} = \{(e, v(t)) \in G((t)) \times t^{-\zeta} U_\eta \mid t^\zeta v(t) \in U_{\eta'}\} / \mathcal{J}_\eta = Y(\eta - \zeta, \eta' - \zeta).$$

Equations (3.6c) and (3.6d) both follow from fixing a representation of $T_{\tilde{G}}$, and comparing the Chern classes of the associated line bundles on \mathcal{F} pulled back to ${}_\eta \mathcal{X}_{\eta'}$ with the trivial line bundle with the induced equivariant structure. On $Y(\eta, \eta')$, this is obvious, since we are just considering the fiber over the identity coset in \mathcal{F} , which shows (3.6c). For (3.6d), we must make a more careful comparison. The argument is the same as [BFN18, Lemma 3.20]: the associated line bundle pulled back from \mathcal{F} is T -equivariantly trivial on each component, but the loop \mathbb{C}^\times acts by with weight $\langle \zeta, \mu \rangle$ on the component of t^ζ .

Finally, the proof of (3.6e) is the same as that of [BFN18, Sec. 4(i–iii)], although we need to correct a mistake in their statement: [BFN18, (4.7)] should read

$$r^\lambda r^\mu = \prod_{i=1}^n A_i(\lambda, \mu) \cdot r^{\lambda+\mu}.$$

The product $\Phi(\eta, \eta', \eta'')$ is the product of the Euler classes of the associated bundles for $U_\eta \cap U_{\eta''} / (U_\eta \cap U_{\eta'} \cap U_{\eta''})$ (which corresponds to the factors satisfying (3.5a)) and $(U_\eta + U_{\eta'} + U_{\eta''}) / (U_\eta + U_{\eta''})$ (which corresponds to the factors satisfying (3.5b)). The former is spanned by the vectors of the form $t^{-n}v$ where v is a weight vector of weight φ_i that satisfies

$$\varphi_i^+(\eta') - n \geq -1/2 \quad \varphi_i^+(\eta') - n \leq -1/2 \quad \varphi_i^+(\eta'') - n \geq -1/2$$

which are equivalent to (3.5a), and similarly the latter to

$$\varphi_i^+(\eta') - n \leq -1/2 \quad \varphi_i^+(\eta') - n \geq -1/2 \quad \varphi_i^+(\eta'') - n \leq -1/2$$

which are equivalent to (3.5b). Since the Euler class of the corresponding line bundle is $\varphi_i - nh$, the product of these Euler classes is $\Phi(\eta, \eta', \eta'')$. \square

If we draw $r(\eta', \eta)$ as a straight line path in $\mathfrak{t}_{1, \tilde{H}}$, and thus compositions of these elements as piecewise linear paths, with the unrolled arrangement drawn in, we can visualize the relation (3.6e) as saying that when we remove two crossings of the hyperplane $\varphi_i^{\text{mid}}(\eta) = n$ from the path, we do so at the cost of multiplying by $\varphi_i^+ - nh$. We can thus represent elements of $\text{Hom}_{\text{ab}}(\eta, \eta)$ as paths that start at η and go to any other chamber of the form $\eta - \zeta$ (we implicitly follow these with translation y_ζ). Composition of two paths p and q is thus accomplished by translating p so that its start matches the end of q , and then straightening using the relation (3.6e).

Example 3.ii. In our running example, let us fix $\eta = \tau$. We let ξ_1, ξ_2 be the usual coordinate cocharacters of the diagonal 2×2 matrices. The algebra $\mathcal{A}_{\text{ab}} = \text{Hom}_{\text{ab}}(\tau, \tau)$ is generated over S_h by

$$w_1 = y_{\xi_1} \cdot r(\tau + \xi_1, \tau) \quad w_2 = y_{\xi_2} \cdot r(\tau + \xi_2, \tau) \quad z_1 = y_{-\xi_1} \cdot r(\tau - \xi_1, \tau) \quad z_2 = y_{-\xi_2} \cdot r(\tau - \xi_2, \tau)$$

with the relations

$$[z_1, w_1] = [z_1, w_2] = [z_2, w_1] = [z_2, w_2] = 0$$

$$z_1 w_1 = \gamma_1(\gamma_1 - 2h) \quad w_1 z_1 = (\gamma_1 + h)(\gamma_1 - h)$$

$$z_2 w_2 = \gamma_2(\gamma_2 - 2h) \quad w_2 z_2 = (\gamma_2 + h)(\gamma_2 - h)$$

since

$$\varphi_1^+ = \gamma_1 + h \quad \varphi_2^+ = \gamma_2 + h \quad \varphi_1^- = \gamma_1 - h \quad \varphi_2^- = \gamma_2 - h.$$

Now, we turn to generalizing this presentation to the nonabelian case. Let $\widehat{\Delta}$ denote the set of real affine roots of G ; that is $\widehat{\Delta} = \Delta_0 + \mathbb{Z}\delta$. We can easily check that the relations (3.6a–3.6d) hold in \mathcal{B} for all elements of the extended affine Weyl group:

$$(3.7a) \quad y_w \cdot y_{w'} = y_{ww'}$$

$$(3.7b) \quad y_w r(\eta', \eta) y_w^{-1} = r(w \cdot \eta', w \cdot \eta)$$

$$(3.7c) \quad y_w \mu y_w^{-1} = w \cdot \mu$$

Finally, if $\alpha(\eta) = n$ for some finite root α but no other weights or roots vanish, then we can make this generic in two different ways: $\eta_{\pm} := \eta \pm \epsilon \alpha^{\vee}$ for some small $\epsilon > 0$. Let \mathcal{J}_{\pm} be the corresponding Iwahoris. These two Iwahoris together generate a parahoric \mathcal{J}_{η} ; this is also generated by one of \mathcal{J}_{\pm} and the root SL_2 for $\alpha - n\delta$. The identity coset $e\mathcal{J}_{\pm}$ in $G((t))/\mathcal{J}_{\pm}$ thus lies in an orbit X of \mathcal{J}_{η} which is isomorphic to $\mathbb{P}^1 \cong \mathcal{J}_{\eta}/\mathcal{J}_{\pm}$. Note that for $\epsilon \ll 1$, we have $U_{\eta} = U_{\eta_{\pm}}$. Since U_{η} is invariant under η , the preimage of X in ${}_{\eta}\mathcal{X}_{\eta} = {}_{\eta_{\pm}}\mathcal{X}_{\eta_{\mp}}$ is precisely $X \times U_{\eta} \cong \mathcal{J}_{\eta} \times^{\mathcal{J}_{\pm}} U_{\eta}$. Let $u_{\alpha}(\eta) = [\overline{X \times U_{\alpha}}] \in \text{Hom}(\eta_{\pm}, \eta_{\mp})$.

We wish to also consider the BGG-Demazure operators for an affine root of the form $\alpha - n\delta$. Note that on $\mathfrak{t}_{1, \mathbb{R}}$, the equation $\alpha(\eta) = n$ is the vanishing set of $\alpha - n\delta$. Since we consider a loop group which is not centrally extended, but which does include the loop action, we cannot see the difference between such affine roots, but we do have a difference between the coroots. Thus, we can define the affine reflections

$$s_{\alpha - n\delta} \cdot \mu = \mu - 2(\alpha^{\vee}(\mu) - nh)\alpha \quad \partial_{\alpha - n\delta}(f) = \frac{s_{\alpha - n\delta} f - f}{\alpha}.$$

Theorem 3.13 *The morphisms in the extended BFN category are generated by*

- (1) y_w for $w \in \widehat{W}$,
- (2) $r(\eta, \eta')$ for $\eta, \eta' \in \mathfrak{t}_{1, \widehat{H}}$ generic,
- (3) the polynomials in S_h ,
- (4) $u_{\alpha'}(\eta) = u_{\alpha - n\delta}(\eta)$ for η_{\pm} affine chambers adjacent across $\alpha'(\eta) = 0$ for $\alpha' \in \widehat{\Delta}$ an affine root (i.e. $\alpha' = \alpha - n\delta$ for some finite root α').

This category has a faithful polynomial representation where each object η is assigned to $H_*^{BM, \widehat{G}^{(t)}}(\mathcal{X}_\eta) \cong S_h \cdot [\mathcal{X}_\eta]$, and the generators above act by:

$$(3.8a) \quad r(\eta, \eta') \cdot f[\mathcal{X}_{\eta'}] = \Phi(\eta, \eta') f \cdot [\mathcal{X}_\eta]$$

$$(3.8b) \quad u_\alpha \cdot f[\mathcal{X}_{\eta_\pm}] = \partial_\alpha(f) \cdot [\mathcal{X}_{\eta_\mp}]$$

$$(3.8c) \quad y_w \cdot f[\mathcal{X}_\eta] = (w \cdot f)[\mathcal{X}_{w \cdot \eta}]$$

$$(3.8d) \quad \mu \cdot f[\mathcal{X}_\eta] = \mu f \cdot [\mathcal{X}_\eta]$$

The relations between these operators are given by

(1) the relations (3.6a–3.7c), which are the image of relations in the abelianized category

(2) the relations:

$$(3.9a) \quad u_\alpha^2 = 0$$


$$(3.9b) \quad \underbrace{u_\alpha u_{s_\alpha \beta} u_{s_\alpha s_\beta \alpha} \cdots}_{m_{\alpha\beta}} = \underbrace{u_\beta u_{s_\beta \alpha} u_{s_\beta s_\alpha \beta} \cdots}_{m_{\alpha\beta}}$$

$$(3.9c) \quad y_w u_\alpha y_{w^{-1}} = u_{w \cdot \alpha}$$

$$(3.9d) \quad u_\alpha \mu - (s_\alpha \cdot \mu) u_\alpha = r(\eta_\mp, \eta_\pm) \partial_\alpha(\mu)$$

whenever these morphisms are well-defined

(3) If we have a codimension 2 intersection of root and matter hyperplanes, then chambers around it will be as shown below:



That is, we will have two pairs η'_\pm and η''_\pm which are opposite across $\alpha(\eta) = 0$ and adjacent to the intersection. We consider a pair of chambers η, η'' which differ by a 180° rotation around the corresponding codimension 2 subspace. We have the following relation:

$$(3.9e) \quad r(\eta''', \eta'_-) u_\alpha r(\eta'_+, \eta) - r(\eta''', \eta''_-) u_\alpha r(\eta''_+, \eta) \\ = \partial_\alpha (\Phi(\eta'_+, \eta) \cdot s_\alpha \Phi(\eta, \eta'_-)) r(\eta''', s_\alpha \eta) s_\alpha.$$

As in the abelian case, we can represent morphisms in our category by paths $\pi: [0, 1] \rightarrow \mathfrak{t}_{1, \mathbb{R}}$ which are suitably generic. Let β_1, \dots, β_k be the list of affine roots ($\beta_i \in \widehat{\Delta}$) whose

hyperplanes that the path π crosses in order, and let $\eta_{\pm}^{(i)}$ be generic points in π on the positive and negative side of the hyperplane for β_i . Let $\eta = \pi(0)$ and $\eta' = \pi(1)$.

Definition 3.14 *Let $\eta = \pi(0)$ and $\eta' = \pi(1)$. Consider the morphism*

$$(3.10) \quad \tilde{r}_{\pi} = r(\eta', \eta_{\pm}^{(k)}) u_{\beta_k} r(\eta_{\mp}^{(k)}, \eta_{\pm}^{(k-1)}) u_{\beta_{k-1}} \cdots u_{\beta_1} r(\eta_{\mp}^{(1)}, \eta).$$

For a given η', η , choose a fixed path π crossing a minimal number of hyperplanes between them, and let $\tilde{r}(\eta', \eta) = \tilde{r}_{\pi}$ for this path. We can represent the relations of Theorem 3.13 locally in terms of this path. For example, (3.6c) and (3.6e) can be represented as:

$$(3.10a) \quad \begin{array}{c} \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \end{array} \quad \begin{array}{c} \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \end{array}$$

while (3.9a) and (3.9d), with the dotted line given by the Coxeter hyperplane $\alpha = 0$ for an affine root $\alpha \in \hat{\Delta}$ becomes

$$(3.10b) \quad \begin{array}{c} \text{---} \\ \text{---} \end{array} = 0 \quad \begin{array}{c} \text{---} \\ \text{---} \end{array} - \begin{array}{c} \text{---} \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \text{---} \end{array}$$

where the green arrow at the right side of the last diagram indicates applying s_{α} . We won't carefully write out the graphical version of (3.9e), but it is similar in structure to (2.7c).

Proof of Theorem 3.13. The formulas (3.8a–3.8d) define an action: The verification of the action is straightforward:

- (1) The action of the affine Weyl group is that induced on S_h by the conjugation action of the torus, which is the action we defined before.
- (2) The action of $r(\eta, \eta')$ is multiplication by the Euler class of $U_{\eta}/(U_{\eta'} \cap U_{\eta})$ (as in the proof of [BFN18, Thm 4.1]). That Euler class is the product of the weights of this space, which is exactly $\Phi(\eta, \eta')$.
- (3) This holds essentially by definition.
- (4) The effect of u_{α} is the same as the pushforward $\mathcal{X}_{\eta_{\pm}} \rightarrow G((t)) \times^{J_{\eta}} U_{\eta}$, which is a bundle with fiber $\mathbb{P}^1 = \mathcal{J}_{\eta}/\mathcal{J}_{\eta_{\pm}}$ followed by pullback to $\mathcal{X}_{\eta_{\mp}}$. Thus, the calculation reduces to a standard calculation for GL_2 -equivariant integration over \mathbb{P}^1 . See, for example, [VV11, Prop. 2.23] or [SW14, Prop. 3.4].

The action is faithful: We have a map

$$\zeta: H_*^{BM, \tilde{T}}(\eta \mathcal{X}_{\eta'}) \rightarrow H_*^{BM, \tilde{T}}(\mathcal{X}_{\eta'}) \cong H_*^{BM, \tilde{T}}(\mathcal{F}\ell)$$

analogous to the map \mathbf{z}^* defined in [BFN18, §5(iv)]. This is an algebra homomorphism by the argument of [BFN18, Lem. 5.11]: the pushforward map

$$\oplus_{\eta'} H_*^{BM, \tilde{T}}(\eta \mathcal{X}_{\eta'}) \rightarrow \oplus_{\eta'} H_*^{BM, \tilde{T}}(\mathcal{X}_{\eta'})$$

is a map of right modules, and sends $1 \mapsto [e\mathcal{J} \times U_\eta]$, which in turn maps to $1 = [e\mathcal{J}] \in H_*^{BM, \tilde{T}}(\mathcal{F}\ell)$. Thus,

$$\zeta(ab) = \zeta(1 \cdot a \cdot b) = \zeta(1)\zeta(a)\zeta(b) = \zeta(a)\zeta(b).$$

The same argument shows that this map intertwines the action of (3.8a–3.8d) with that of $H_*^{BM, \tilde{T}}(\mathcal{F}\ell)$ on $H_*^{BM, \tilde{T}}(G((t))) \cong S_h$.

This latter action is the polynomial representation of the degenerate affine Hecke algebra, which is faithful; we can also see this by localization to T -fixed points. Thus, it suffices to check that ζ is injective. As in Lemma 2.5(5), this follows because ${}_\eta \mathcal{X}_{\eta'}$ is equivariantly formal for \tilde{T} and $({}_\eta \mathcal{X}_{\eta'})^{\tilde{T}} = G \times^J V^T = \mathcal{X}_{\eta'}^{\tilde{T}}$; the analogous result for Coulomb branches is [BFN18, Lem. 5.13].

The relations (3.6a–3.9e) hold: Given the representation and its faithfulness, the reader can readily verify that the relations (3.6a–3.9e) are satisfied. The most interesting of these relations is (3.9e), so let us verify this relation in more detail. The action of the LHS in the polynomial rep on a polynomial f is:

$$\begin{aligned} (3.11) \quad & \Phi(s_\alpha \eta, \eta'_\pm) \partial_\alpha(\Phi(\eta'_\mp, \eta) f) - \Phi(s_\alpha \eta, \eta''_\pm) \partial_\alpha(\Phi(\eta''_\mp, \eta) f) \\ &= \Phi(s_\alpha \eta, \eta'_\pm) \Phi(\eta'_\mp, \eta) \partial_\alpha(f) + \Phi(s_\alpha \eta, \eta''_\pm) \partial_\alpha(\Phi(\eta''_\mp, \eta)) f^{s_\alpha} \\ & \quad - \Phi(s_\alpha \eta, \eta''_\pm) \Phi(\eta''_\mp, \eta) \partial_\alpha(f) - \Phi(s_\alpha \eta, \eta''_\pm) \partial_\alpha(\Phi(\eta''_\mp, \eta)) f^{s_\alpha}. \end{aligned}$$

Since η' and η'' are both on the minimal-length paths, neither is separated from both $s_\alpha \eta$ and η by any given unrolled hyperplane. Thus, we have that

$$\Phi(s_\alpha \eta, \eta'_\pm) \Phi(\eta'_\mp, \eta) = \Phi(s_\alpha \eta, \eta''_\pm) \Phi(\eta''_\mp, \eta) = \Phi(s_\alpha \eta, \eta).$$

It follows that the first positive and negative terms in (3.11) cancel, and we obtain the RHS of (3.9e). This confirms the relation.

The relations (3.6a–3.9e) are complete: Now, we need to show that no other relations are needed to present the category. Using the action of the elements of \widehat{W} , we can reduce to the case where η and η' are in the same alcove. The space $\text{Hom}(\eta, \eta')$ has a filtration by the length of the relative position of the two affine flags. Let $\text{Hom}^{\leq w}(\eta, \eta')$ be the homology classes supported on the pairs of relative distance $\leq w$. By basic algebraic topology, $\text{Hom}^{\leq w}(\eta, \eta') / \text{Hom}^{< w}(\eta, \eta')$ is a free module of rank 1 over S_h , since this space is isomorphic to the J -equivariant Borel-Moore homology of an (infinite-dimensional) affine space.

We'll prove that

$$(*) \text{ the } S_h\text{-module } \text{Hom}^{\leq w}(\eta, \eta') / \text{Hom}^{< w}(\eta, \eta') \text{ is generated by the element } \tilde{r}(\eta', w\eta)w.$$

The element $\tilde{r}(\eta', w\eta)w$ is the pushforward of the fundamental class by the map

$$Y(\eta', \eta_{\pm}^{(k)}) \times_{\mathcal{J}_{\eta_{\pm}^{(k)}}} \times_{\mathcal{J}_{\eta_{\mp}^{(k)}}} Y(\eta_{\mp}^{(k)}, \eta_{\pm}^{(k-1)}) \times_{\mathcal{J}_{\eta_{\pm}^{(k-1)}}} \times_{\mathcal{J}_{\eta_{\mp}^{(k-1)}}} \dots \times_{\mathcal{J}_{\eta_{\pm}^{(1)}}} \times_{\mathcal{J}_{\eta_{\mp}^{(1)}}} Y(\eta_{\mp}^{(1)}, \eta) \rightarrow \mathcal{X}_{\xi'} \times_{V((t))} \mathcal{X}_{\xi}.$$

This map is an isomorphism on the set of affine flags of relative position w , since the map

$$\mathcal{J}_{\eta_{\pm}^{(k)}} \mathcal{J}_{\eta_{\pm}^{(k-1)}} \times_{\mathcal{J}_{\eta_{\pm}^{(k-1)}}} \mathcal{J}_{\eta_{\pm}^{(k-1)}} \mathcal{J}_{\eta_{\pm}^{(k-2)}} \times_{\mathcal{J}_{\eta_{\pm}^{(k-2)}}} \dots \times_{\mathcal{J}_{\eta_{\pm}^{(2)}}} \mathcal{J}_{\eta_{\pm}^{(2)}} \mathcal{J}_{\eta_{\pm}^{(1)}} \rightarrow \mathcal{J}_{\eta_{\pm}^{(k)}} \mathcal{J}_{\eta_{\pm}^{(1)}}$$

is an isomorphism by standard results in Bruhat theory. Thus, the elements $\tilde{r}(\eta', w\eta)w$ give a free basis of the associated graded for this filtration. This implies that they are a basis of the original module; in particular, this implies that the elements (1-4) above are generators.

On the other hand, we can also easily show that the relations displayed are enough to bring any element into the form of a sum of elements $\tilde{r}(\eta', w\eta)w$. We can pull all elements of the Weyl group to the right using (3.7b, 3.7c, 3.9c), all elements of S_h to the right using the relation (3.9d), and rewrite any crossing of a Coxeter hyperplane by $r(-, -)$ using the relation $r(\eta_{\pm}, \eta_{\mp}) = s_{\alpha} - \alpha u_{\alpha}$, which can be verified using the polynomial representation. This shows that these relations suffice, since there can be no further relations within our basis. \square

Note that in the course of this proof, we have shown that:

Corollary 3.15 *The elements $\tilde{r}(\eta', w\eta)w$ for $w \in \widehat{W}$ is a basis of $\text{Hom}_{\mathcal{O}}(\eta, \eta')$ as a right module (or left module) over the ring S_h .*

We can extend this to the case where η and η' are not generic but merely unexceptional. First, consider the case where η and η' are not separated by any matter hyperplanes and lie in closed faces of the Coxeter complex with non-trivial intersection. In this case, $U_{\eta} = U_{\eta'}$ and the parahorics \mathcal{J}_{η} and $\mathcal{J}_{\eta'}$ generate a larger parahoric \mathcal{J} (corresponding to their intersection as a face in the Coxeter complex). In this case, we let

$$\tilde{r}(\eta', \eta) = \left[\overline{\left\{ ((g\mathcal{J}_{\eta'}, u), (g\mathcal{J}_{\eta}, u)) \in \mathcal{X}_{\eta'} \times \mathcal{X}_{\eta} \mid g \in G((t)), u \in U_{\eta} = U_{\eta'} \right\}} \right].$$

Note that it is a standard result of Schubert calculus that if η and η' are generic, this agrees with Definition 3.14. With this formula in hand, we can rewrite Definition 3.14 as

$$(3.12) \quad \tilde{r}(\eta', \eta) = \tilde{r}(\eta', \eta_p) \tilde{r}(\eta_p, \eta_{p-1}) \cdots \tilde{r}(\eta_1, \eta)$$

where $\eta_i = t_i \eta + (1 - t_i) \eta'$ for $0 < t_1 < \cdots < t_p < 1$ real numbers such that any consecutive pair η_i and η_{i+1} are either in the same alcove or are not separated by any matter hyperplanes and lie in closed faces of the Coxeter complex with non-trivial

intersection. Equation (3.12) defines $\tilde{r}(\eta', \eta)$ for every unexceptional pair and is easily extended to any piecewise linear path with unexceptional vertices.

Unlike the generic case, the morphisms $\tilde{r}(\eta', w\eta)w$ are not distinct, but only depend on the double coset of w with respect to the stabilizers \widehat{W}_η and $\widehat{W}_{\eta'}$ of η and η' in \widehat{W} . Furthermore, in order to obtain a basis, it's not enough to multiply on the left or right with polynomials (since we can only multiply by the invariants $S_h^{\widehat{W}_\eta}$ and $S_h^{\widehat{W}_{\eta'}}$); instead we have to “dress” these morphisms with polynomials forming a basis of $S_h^{\widehat{W}_{\eta'} \cap w^{-1}\widehat{W}_\eta w}$ by inserting these at a generic point in the path. The resulting morphism depends on the point of insertion.

In the case where $\eta = \eta' = 0$, we are thus taking double cosets for the finite Weyl group inside the affine, so each double coset contains a unique translation by a dominant coweight. Thus, the basis obtained is the basis of dressed monopole operators discussed in [Cre+14]:

$$m_\lambda(f) = y_\lambda \tilde{r}(-\lambda, -\lambda\epsilon) f \tilde{r}(-\lambda\epsilon, 0).$$

As mentioned before, this is not a truly canonical definition, since we choose other insertion points for f . We thus obtain that:

Proposition 3.16 *As λ runs over integral dominant coweights, and f over any fixed basis of $S_h^{W_\lambda}$, the dressed monopoles $m_\lambda(f)$ give a basis of \mathcal{A}^{sph} .*

While the existence of such a basis is clear from the degeneration of [BFN18, §6], the author could not find an explicit construction of one for a general Coulomb branch.

3.4. The deformed category and Hamiltonian reduction. Having fixed G and V , we have defined a larger group H acting on V , and we can also consider the Coulomb branch for this larger group. It is more convenient to consider the subgroup $Q \subset H$; note that the tori $T_H = T_Q$ coincide by definition.

Consider the affine Grassmannian of Q ; this has a natural map to the affine Grassmannian of T_F , which we identify with the coweight lattice $X(F) \cong \pi_1(F)$. Alternately, we can think of this as induced by the tropicalization homomorphism $\mathfrak{q}: Q((t)) \rightarrow T_F((t)) \rightarrow X(F)$. If $T_F = \mathbb{G}_m$, then the second map is the homomorphism sending a Laurent power series to its order of 0 at $t = 0$.

Let \mathcal{A}_Q denote the quantum Coulomb branch of the action of Q on V . Since the affine Grassmannian is the disjoint union of the preimages of the points in $X(F)$, we obtain a grading of \mathcal{A}_Q as a vector space; we can equivalently think of this as an action of the Pontryagin dual $T_F^\vee \cong \text{Hom}(X(F), \mathbb{C}^\times)$. The algebra \mathcal{A}_Q contains a copy of $S_h^Q \cong H_{T_Q}^*(pt)$, the equivariant cohomology of a point for $T_Q \cong T_Q \times \mathbb{C}^\times$.

Lemma 3.17 ([BFN18, 3(vii)(d)]) *The action of T_F^\vee on \mathcal{A}_Q is by algebra automorphisms, and after specializing $h = 1$ has a non-commutative moment map induced by the natural map $U(\mathfrak{t}_F^\vee) \cong \mathbb{C}[\mathfrak{t}_F] \hookrightarrow S_1^Q \cong \mathbb{C}[\mathfrak{t}_Q] \rightarrow \mathcal{A}_Q$.*

Note that the isomorphism $S_1^Q \cong \mathbb{C}[\mathfrak{t}_Q]$ depends on a choice of splitting $T_{\tilde{Q}} \cong T_Q \times \mathbb{C}^\times$. This paper and [BFN18] make different choices; we let \mathbb{C}^\times act on V trivially, and in [BFN18], it acts with weight $-1/2$. However, these different choices all lead to non-commutative moment maps for the same action.

Applying quantum Hamiltonian reduction gives us a standard package of a deformation and collection of bimodules attached to it. If we consider the invariants $\mathcal{A}_Q^{T_F^\vee}$, then the image of $U(\mathfrak{t}_F^\vee)$ is central in these invariants, and we obtain a family over the base \mathfrak{t}_F which is essentially \mathcal{A} with its flavor parameters considered as formal elements of the base ring $U(\mathfrak{t}_F^\vee)$.

The additional ingredient we obtain from this approach is that for each character $\chi: T_F^\vee \rightarrow \mathbb{G}_m$, the semi-invariants of \mathcal{A}_Q^X transforming according to this character form a bimodule over $\mathcal{A}_Q^{T_F^\vee}$. While $U(\mathfrak{t}_F^\vee)$ is central in $\mathcal{A}_Q^{T_F^\vee}$, the non-commutative moment map condition guarantees that the left and right actions on \mathcal{A}_Q^X differ by a shift by $h\chi$. Thus, if we specialize our flavor parameters to scalars, we will obtain a bimodule relating two different choices of flavor. We'll show below that in the setting where it makes sense to compare them, these agree with bimodules that have played an important role in the representation theory of symplectic singularities, and induce **twisting** or **wall-crossing** functors, as discussed in [BPW16, §6].

We can easily extend all of these structures to the category \mathcal{B} .

Definition 3.18 *The deformed extended BFN category \mathcal{B}^{def} is the category with the same objects as \mathcal{B} and*

$$\text{Hom}(\eta, \eta') = H_*^{BM, Q((t)) \times \mathbb{C}^\times}(\mathcal{X}_\eta \times_{V((t))} \mathcal{X}_{\eta'}) \cong H_*^{BM, T_{\tilde{Q}}}(\eta \mathcal{X}_{\eta'}).$$

The results above, such as Theorem 3.13, carry over in a straightforward way to this category. The only difference is that we must interpret the products of weights $\Phi(-, -)$ as weights of $T_{\tilde{Q}}$, and rather than an action of S_h , we have one of S_h^Q .

The category \mathcal{B}^Q has a natural action of the torus $(T_F)^\vee$ on the morphisms between any two objects with $r(-, -), \mathbb{k}[\tilde{\mathfrak{t}}_Q]$ and \widehat{W} having weight 0, and the copy of $\mathbb{k}[T_H]$ having the obvious action. The classes of weight χ under this action (which is a coweight of F) correspond to homology classes concentrated on the components of the affine flag variety whose corresponding loop has a homotopy class hitting χ under the map $\pi_1(Q) \rightarrow X(F)$. Furthermore, this action is induced by the same non-commutative moment map induced by the inclusion of S_h^F into the endomorphism ring of each object.

We can construct a faithful functor $\mathcal{B}^{\text{def}} \rightarrow \mathcal{B}^Q$ to the extended BFN category for the group Q acting on V . It is the subcategory where we only consider cocharacters in $\mathfrak{t}_{1, \mathbb{R}}$ as objects and only allow ourselves to use \mathfrak{t}_Z in the extended affine Weyl group, rather than all of $(\mathfrak{t}_Q)_\mathbb{Z}$. This is precisely the morphisms which are invariant under F^\vee . Thus, we have that: This shows that:

Lemma 3.19 *The deformed extended BFN category \mathcal{B}^{def} is equivalent to the subcategory of $\mathcal{B}^{\mathcal{Q}}$ equipped with only F^{\vee} -invariant morphisms.*

Of course, if we instead fix $\chi \in X(F)$ and look only at morphisms with this weight, we obtain a $\mathcal{B}^{\text{def}} - \mathcal{B}^{\text{def}}$ bimodule, which we denote $\mathcal{T}(\chi)$.

As we discussed in the case of a single algebra, this bimodule has different left and right actions of S_h^F , and thus if we impose a single flavor, we will arrive at a $\mathcal{B}_{\phi+\chi} - \mathcal{B}_{\phi}$ -bimodule for extended categories associated to flavors whose difference is χ .

Definition 3.20 *Let ${}_{\phi+\chi}\mathcal{T}_{\phi}$ be the $\mathcal{B}_{\phi+\chi} - \mathcal{B}_{\phi}$ bimodule which is the quotient of $\mathcal{T}(\chi)$ by $I(\mathfrak{t}_{1,\mathbb{R}})$ acting on the right or $I(\mathfrak{t}_{1,\mathbb{R}} + \chi)$ acting on the left.*

Let ${}_{\phi+\chi}T_{\phi} = {}_{\phi+\chi}\mathcal{T}_{\phi}(o, o)$ be the corresponding bimodule over $\mathcal{A}_{\phi+\chi}$ and \mathcal{A}_{ϕ} .

As discussed above, a notion of twisting bimodules is defined in [BPW16, Def. 6.21]. The framework on that paper (conical symplectic resolutions) does not apply to all Coulomb branches. The usual homological grading makes $\mathbb{C}[\mathfrak{M}]$ into a graded algebra; as in [BPW16], we use \mathbb{S} to denote the copy of \mathbb{G}_m which acts on \mathfrak{M} according to this grading. Since the equivariant parameter h has degree 2, this means that the induced Poisson structure has degree -2 in this grading. However, many Coulomb branches are not conical with respect to this grading (there are non-constant functions of non-positive degree). In physics, theories where all non-constant homogeneous functions on \mathfrak{M} have positive degree are called *good* and those where this property fails *bad* or *ugly*. Similarly, many Coulomb branches do not possess a symplectic resolution, though as discussed earlier, they are now known to have symplectic singularities by [Bel23].

In [BFN18, Prop. 3.25], Braverman-Finkelberg-Nakajima define a partial resolution of \mathfrak{M} as a GIT quotient, using the cocharacter $\nu: \mathbb{C}^{\times} \rightarrow T_F$ as stability parameter. If \mathfrak{M}' is a symplectic resolution of singularities, then we call it a BFN resolution of \mathfrak{M} .

Lemma 3.21 *Assume that the homological grading on $\mathbb{C}[\mathfrak{M}]$ is conical (i.e. it corresponds to a good theory) and \mathfrak{M} has a BFN resolution. The bimodule ${}_{\phi+\chi}T_{\phi}$ is isomorphic to the bimodule denoted by the same symbol in [BPW16, Def. 6.21].*

Proof. Both of these bimodules ${}_{\phi+\chi}T_{\phi}$, as defined above or in [BPW16] are specializations of the quantization of a deformation of the Coulomb branch \mathfrak{M} . The only difference is that in [BPW16], we assumed that we were using the universal deformation of this singularity (and that this deformation was generically smooth), and in this paper we use the universal flavor deformation $\mathfrak{M}^{\mathcal{Q}}/T_F^{\vee}$. This deformation has the property of being smooth and universal in many interesting cases, but we cannot assume it is in general.

By [Nam11, Thm. 5.1], the functor of deforming \mathfrak{M} as a Poisson variety is unobstructed. As discussed in [Nam11, §5.4], the conical structure shows that \mathfrak{M} has a universal Poisson deformation \mathcal{M} over a polynomial ring $B = \mathbb{C}[y_1, \dots, y_d]$ (following

the notation of [Nam11, §5], and the flavor deformation over the base \mathfrak{t}_F induced by a ring homomorphism $\alpha: B \rightarrow U(\mathfrak{t}_F^\vee)$. From deformation theory, we only obtain this result after completion, but we can obtain varieties of finite type over \mathbb{k} by passing to spaces of \mathbb{S} -finite vectors.

As above, we consider the family $\tilde{\mathcal{M}}$ defined over the base \mathfrak{t}_F as the GIT quotient with respect to ν of the Coulomb branch \mathfrak{M}^Q by the action of T_F^\vee , with the map to \mathfrak{t}_F being the residual moment map. By assumption, there is a cocharacter ν such that this is a resolution of \mathcal{M} . The fiber over 0 in this family is the symplectic resolution $\tilde{\mathfrak{M}}$ of \mathfrak{M} . The cocharacter χ , though of as a character of T_F^\vee , thus defines an associated line bundle \mathcal{L}_χ on $\tilde{\mathcal{M}}$ and its restriction \mathcal{L}_χ to $\tilde{\mathfrak{M}}$. We can consider the line $\mathbb{A}^1 \subset \mathfrak{t}_F$ spanned by ν , and as in [BLPW16], it is useful to consider the 1-dimensional family $\tilde{\mathcal{M}}_\nu$ inside $\tilde{\mathcal{M}}$ defined over this line, with the pullback line bundle $\mathcal{L}_\chi^{(\nu)}$. We can also construct this family by reducing the Coulomb branch $\mathfrak{M}^{\tilde{G}}$ for (\tilde{G}, V) , where we use $\phi_0 = \nu$. By [Kal08, Lemma 2.5], all fibers of this family other than $\tilde{\mathfrak{M}}$ are smooth and affine.

In the proof of [BPW16, Prop. 6.26], we show that bimodule defined in [BPW16, Def. 6.21] can also be constructed by considering the \mathbb{S} -invariant sections of the unique quantization of the line bundle $\mathcal{L}_\chi^{(\nu)}$ (constructed in [BPW16, Prop. 5.2]), and specializing the deformation parameters as we do in Definition 3.20. On the other hand, we can construct this quantized line bundle by considering the quantization of the structure sheaf on the Coulomb branch $\mathfrak{M}^{\tilde{G}}$, pushing forward its restriction to stable locus to $\tilde{\mathcal{M}}_\nu$ and taking semi-invariants for ν . This defines a map of $\phi_{+\chi}T_\phi$ to the sections of this quantized line bundle; this is an isomorphism since after passing to associated graded, we obtain the natural map from semi-invariants in $\mathbb{k}[\mathfrak{M}^{\tilde{G}}]$ to $\Gamma(\tilde{\mathcal{M}}_\nu, \mathcal{L}_\chi^{(\nu)})$.

Thus, to complete the proof, we need only establish that this map is an isomorphism. The issue is that $\Gamma(\tilde{\mathcal{M}}_\nu, \mathcal{L}_\chi^{(\nu)})$ is equal to the semi-invariants of the function ring of the semi-stable locus, which in general might be larger than $\mathbb{k}[\mathfrak{M}^{\tilde{G}}]$. Note that the unstable locus in $\mathfrak{M}^{\tilde{G}}$ must lie only in the fiber over $0 \in \mathbb{A}^1$ (since the other fibers are smooth and affine) and is a proper subvariety in this fiber. Thus, it has codimension ≥ 2 in $\mathfrak{M}^{\tilde{G}}$. Since $\mathfrak{M}^{\tilde{G}}$ is normal by [BFN18, Prop. 6.12], this means that the functions on the semistable locus are just $\mathbb{k}[\mathfrak{M}^{\tilde{G}}]$. This establishes the desired isomorphism. \square

Remark 3.22. The reader will note that we used the universal deformation of the Coulomb branch and will likely have wondered how this relates to the deformation arising from \mathcal{B}^{def} . This will happen when a version of Kirwan surjectivity holds: the natural map $(\mathfrak{g}^*)^G \rightarrow H^2(\tilde{\mathfrak{M}}; \mathbb{C})$ is surjective. In the cases of greatest interest to us, such as hypertoric varieties and affine type A quiver varieties, this map is indeed surjective. Thus, these deformations coincide, but it is unclear how generally this will hold.

3.5. Representation theory. Throughout this section, we specialize $h = 1$; we let $S_1 = S_h/(h - 1) \cong \mathbb{C}[\mathfrak{t}_{1,\mathbb{k}}]$. Furthermore, we assume that \mathbb{k} has characteristic 0.

Definition 3.23 We call a finitely generated \mathcal{B} -module M (resp. \mathcal{A} -module N) a **weight module** if for every η , we have that $M(\eta)$ (resp. N) is locally finite as a module over S_1 with finite dimensional generalized weight spaces.

Obviously, if M is a weight module, then $N = M(o)$ is as well. The left adjoint functor $\mathcal{B} \otimes_{\mathcal{A}} -$ also sends weight modules to weight modules, since the adjoint action of S_1 on $\text{Hom}(\eta, o)$ is semi-simple with eigenspaces finitely generated over S_1 .

For each $v \in \mathfrak{t}_{1,\mathbb{k}}, \eta \in \mathfrak{t}_{1,\mathbb{R}}$, we can consider the functor $\mathcal{W}_{v,\eta}: \mathcal{B}\text{-mod}_W \rightarrow \mathbb{k}\text{-mod}$ defined by

$$(3.13) \quad \mathcal{W}_{v,\eta}(M) = \{m \in M(\eta) \mid \mathfrak{m}_v^N m = 0 \text{ for } N \gg 0\},$$

with \mathfrak{m}_v for $v \in \mathfrak{t}_{1,\mathbb{k}}$ be the corresponding maximal ideal in S_1 . These functors are exact and prorepresentable. If we let $\mathcal{A}_\eta := \text{Hom}(\eta, \eta)$, then they are represented by the projective limit

$$P_{v,\eta} := \varprojlim \mathcal{B} \otimes_{\mathcal{A}_\eta} (\mathcal{A}_\eta / \mathcal{A}_\eta \mathfrak{m}_v^N)$$

as $N \rightarrow \infty$. Note that this is now a module which carries a topology compatible with the action of \mathcal{A}_η .

Thus, as in [MV98, Prop. 3.5.6], we can present the category of weight modules as the set of modules, continuous in the discrete topology, over the topological algebra $\text{End}(\oplus P_{v,\eta})$ of continuous module endomorphisms of this sum. Since this is a highly infinite sum, it is perhaps more conducive to package these Hom spaces into the morphisms of a category:

Definition 3.24 Let $\widehat{\mathcal{B}}$ be the category whose objects are the set \mathcal{J} of pairs of generic $\eta \in \mathfrak{t}_{1,\mathbb{R}}$ and any $v \in \mathfrak{t}_{1,\mathbb{k}}$, with morphisms defined by

$$\begin{aligned} \text{Hom}_{\widehat{\mathcal{B}}}((\eta', v'), (\eta, v)) &= \text{Hom}(P_{v',\eta'}, P_{v,\eta}) \\ &= \varprojlim \text{Hom}_{\mathcal{B}}(\eta', \eta) / (\mathfrak{m}_v^N \text{Hom}_{\mathcal{B}}(\eta', \eta) + \text{Hom}_{\mathcal{B}}(\eta', \eta) \mathfrak{m}_{v'}^N). \end{aligned}$$

We let $\widehat{\mathcal{B}}_{v'}$ be the subcategory where we only allow objects (η, v) with $\eta \in \mathfrak{t}_\tau, v \in v' + \mathfrak{t}_\mathbb{k}$.

We let $S_1^v = \varprojlim S_1 / \mathfrak{m}_v^N$; this ring naturally acts by endomorphisms on (η, v) in $\widehat{\mathcal{B}}$ for each η .

Lemma 3.25 The morphism space $\text{Hom}_{\widehat{\mathcal{B}}}((\eta', v'), (\eta, v))$ is freely spanned as a left module over S_1^v by $w \cdot \tilde{r}(w^{-1}\eta, \eta')$ for $w \in \widehat{W}$ satisfying $w \cdot v' = v$.

In particular, if $v \notin \widehat{W} \cdot v'$, then $\text{Hom}_{\widehat{\mathcal{B}}}((\eta', v'), (\eta, v)) = 0$.

Proof. Consider the quotient $\text{Hom}_{\mathcal{B}}(\eta', \eta) / \mathfrak{m}_v^N \text{Hom}_{\mathcal{B}}(\eta', \eta)$; since under the left action of S_1 , the identity of η has weight v , an element $f \in \text{Hom}(\eta', \eta)$ such that $\mathfrak{m}_v f \supset f \mathfrak{m}_{v'}^N$ for some N maps to an element killed by the right action of $\mathfrak{m}_{v'}^N$.

Now, note that the space of morphisms in \mathcal{B} have a filtration by the number of root hyperplanes crossed by the corresponding path (and elements of \widehat{W} have filtration degree 0). This is closely related to the filtration of [BFN18, §6(i)], though it is slightly different, since it corresponds to \mathcal{J} -orbits instead of $G[[t]]$ orbits. Note that we have $w\tilde{r}(w^{-1}\eta, \eta')f = w(f) \cdot w\tilde{r}(w^{-1}\eta, \eta')$ modulo lower order terms in this filtration for $f \in S_1$. Thus, we have that if $w \cdot v' \neq v$, then the projection of $w\tilde{r}(w^{-1}\eta, \eta')$ to $\text{Hom}_{\mathcal{B}}(\eta', \eta)/\mathfrak{m}_v^N \text{Hom}_{\mathcal{B}}(\eta', \eta)$ is in the span of lower order terms in the filtration. This shows that the elements with $w \cdot v' = v$ must span. \square

It might seem more natural to consider the larger category where we allow $v \in \widehat{W} \cdot v'$, but the resulting categories are equivalent, since $(\eta, v) \cong (w\eta, wv)$ for $w \in W$ in the finite Weyl group, and \mathfrak{t}_τ is invariant under \widehat{W} .

The results above establish:

Lemma 3.26 *The category of weight modules over \mathcal{B} is equivalent to the category of representations of $\widehat{\mathcal{B}}$ in the category of finite dimensional vector spaces. The category of weight modules over \mathcal{B}_τ with weights in $\widehat{W} \cdot v'$ is equivalent to the category of representations of $\widehat{\mathcal{B}}_{v'}$ in the category of finite dimensional vector spaces.*

The category $\widehat{\mathcal{B}}_{v'}$ contains a subcategory $\widehat{\mathcal{A}}_{v'}$ given by the objects of the form (o, v) for $v \in v' + \mathfrak{t}_\mathbb{Z}$. Let $\mathcal{A}\text{-mod}_{v'}$ denote the representations of this category, which are equivalent to the category of weight modules over \mathcal{A} with weights in $\widehat{W} \cdot v'$.

A useful notion for understanding this representation theory is that of **clans**:

Definition 3.27 *Consider the equivalence relation $(\eta, v) \sim (\eta', v')$ if the corresponding objects in $\widehat{\mathcal{B}}$ are isomorphic. We call the equivalence classes of this relation **clans**.*

The clans are thus the sets of weight spaces which are naturally isomorphic in every representation.

4. HIGGS AND COULOMB

4.1. The isomorphism. Assume that \mathbb{k} has characteristic 0; we are still specializing $h = 1$. Consider $\rho \in \mathfrak{t}_{1, \mathbb{k}}$, and let $\mathcal{W} = \rho + \mathfrak{t}_\mathbb{Z}$. For simplicity, let us assume for now that no element of $\widehat{W} \cdot \rho$ is fixed by any non-trivial length 0 element of \widehat{W} . This holds if $\rho \in \mathfrak{t}_{1, \mathbb{Z}}$, for all ρ if G is simply connected (since in this case, only $1 \in \widehat{W}$ has length 0) or G is isomorphic to a product of GL_m 's (since in this case, every non-trivial element of length 0 of \widehat{W} has no fixed points), but not for any non-trivial quotient of SL_n , for example. We'll discuss how to deal with other cases in Remark 4.4.

We'll call a weight φ_i of V or root α_i of \mathfrak{g} **relevant** if it has integral value on ρ , and **irrelevant** if it does not. The relevant roots form the root system of a pseudo-Levi subalgebra $\mathfrak{g}_\rho \subset \mathfrak{g}$, and the sum V_ρ of relevant weight spaces carries an \mathfrak{g}_ρ action. By

replacing ρ by another element of its coset, we can assume without loss of generality that $\varphi_i^+(\rho) = 0$ for all φ_i relevant and that ρ is stabilized by W_ρ .

Our assumption on length 0 elements means that W_ρ is precisely the stabilizer of ρ —in particular, $\#W_\rho$ is the rank of the Hom space between any two objects of $\widehat{\mathcal{B}}_\rho$.

We now turn to considering the constructions of Section 2 for the group L acting on the vector space V_ρ . Let

$$(4.1) \quad rC_{\mathbf{a}} = \{\xi \in \mathfrak{t}_{1,\mathbb{R}} \mid a_i < \varphi_i^{\text{mid}}(\xi) < a_i + 1 \text{ for all } i \text{ relevant}\}.$$

be the affine chambers associated to this smaller group. Let $\eta_{\mathbf{a}}$ be an element of \mathfrak{t}_τ satisfying the equalities as in equation (4.1). Such an element exists because there is an isomorphism $\mathfrak{t}_{1,\mathbb{R}} \cong \mathfrak{t}_\tau$ sending $\rho \rightarrow \tau$ while preserving φ_i^+ for i relevant.

Note that we cannot make this choice W_ρ -equivariantly if we insist on choosing generic elements. However, if a chamber $rC_{\mathbf{a}}$ has non-trivial stabilizer, a standard averaging argument and convexity show that the interior of $rC_{\mathbf{a}}$ contains a fixed point of this stabilizer, and we can choose $\eta_{\mathbf{a}}$ arbitrarily close to this fixed point. In particular, for $w \in \text{Stab}(\mathbf{a})$, we have that $w \cdot \eta_{\mathbf{a}}$ and $\eta_{\mathbf{a}}$ are not separated by any matter hyperplanes. Thus, the elements $r(w \cdot \eta_{\mathbf{a}}, \eta_{\mathbf{a}})$ and $r(\eta_{\mathbf{a}}, w \cdot \eta_{\mathbf{a}}) = w^{-1}r(w \cdot \eta_{\mathbf{a}}, \eta_{\mathbf{a}})$ give inverse isomorphisms in \mathcal{B} .

We can always choose $\eta_{w \cdot \mathbf{a}}$ to be the image of $\eta_{\mathbf{a}}$ under some element of the Weyl group W_ρ . This implies that $r(\eta_{w \cdot \mathbf{a}}, w \cdot \eta_{\mathbf{a}})$ is always an isomorphism with obvious inverse.

Lemma 4.1

$$r(\eta_{w'w'' \cdot \mathbf{a}}, w' \cdot \eta_{w'' \cdot \mathbf{a}})w' \cdot r(\eta_{w'' \cdot \mathbf{a}}, w'' \cdot \eta_{\mathbf{a}})w'' = r(\eta_{w'w'' \cdot \mathbf{a}}, w'w'' \cdot \eta_{\mathbf{a}})w'w''$$

We can consider the categories $\mathcal{X}_\ell, \mathcal{X}_X$ as discussed in Section 2.4. Let $\widehat{\mathcal{X}}_\ell, \widehat{\mathcal{X}}_X$ be the completions of these categories with respect to their gradings.

Let $\Phi_0(\eta, \eta', \rho)$ be the product of the terms $\varphi_i^+ - n$ over pairs $(i, n) \in [1, d] \times \mathbb{Z}$ such that we have the inequalities

$$(4.2) \quad \varphi_i^{\text{mid}}(\eta) > n \quad \varphi_i^{\text{mid}}(\eta') < n \quad \langle \varphi_i^+, \rho \rangle \neq n.$$

Note that if $\mu \in \mathfrak{t}_{\mathbb{k}}^*$, then μ does not have a canonical extension to $\mathfrak{t}_{1,\mathbb{k}}$, but the expression $\mu - \langle \mu, \rho \rangle$ is well-defined on $\mathfrak{t}_{1,\mathbb{k}}$, giving the same answer for every extension of μ to $\tilde{\mathfrak{t}}_{\mathbb{k}}$.

Lemma 4.2 *There is a unique functor*

$$\gamma: \widehat{\mathcal{X}}_\ell \rightarrow \widehat{\mathcal{B}}_\rho$$

which sends each $\mathbf{a} \in \mathcal{C}$ to $(\xi_{\mathbf{a}}, \rho)$, for $\xi_{\mathbf{a}}$ a generic element of $rC_{\mathbf{a}}$, and which acts on morphisms by

$$(4.3a) \quad \gamma(\mathfrak{w}(\mathbf{a}, \mathbf{b})) = \frac{1}{\Phi_0(\eta_{\mathbf{a}}, \eta_{\mathbf{b}}, \rho)} r(\eta_{\mathbf{a}}, \eta_{\mathbf{b}})$$

$$(4.3b) \quad \gamma(\psi_i(\mathbf{a})) = s_{\alpha_i} u_{\alpha_i}$$

$$(4.3c) \quad \gamma(w) = r(\eta_{w \cdot \mathbf{a}}, w \cdot \eta_{\mathbf{a}}) y_w$$

$$(4.3d) \quad \gamma(\mu) = \mu - \langle \mu, \rho \rangle$$

As mentioned in the introduction, these formulas can be explained geometrically. In particular, $\Phi_0(\eta_{\mathbf{a}}, \eta_{\mathbf{b}}, \rho)$ can be interpreted as the Euler class of a normal bundle, just as in [VV10, Proposition 2.4.7].

Proof. In order to check that this functor is well-defined, we use the natural faithful representation of the category $\widehat{\mathcal{B}}_{\rho}$ sending (η, v) to S_1^v using the representation of (3.8a–3.8d). As in many previous proofs, we’ll prove the equivalence by comparing this with the polynomial representation of $\widehat{\mathcal{X}}_{\mathcal{C}}$ given in (2.11a–2.11d). We consider the completion of this polynomial representation with respect to the grading.

Consider the induced isomorphism $\mathfrak{s}_{\rho}: \mathbb{C}[[\mathfrak{t}_{\mathbb{k}}]] \mapsto S_{\rho}$ by shifting, that is, the image of a linear function $\mu \in \mathfrak{t}_{\mathbb{k}}^*$ is $\mu - \langle \rho, \mu \rangle$. Simple calculations show that (2.11b) and (3.8b) match via (4.3b), (2.11c) and (3.8c) match via (4.3c) and Lemma 4.1, and (2.11d) and (3.8d) match via (4.3d), but perhaps we should say a bit more about (4.3a). The image of $\varphi(\mathbf{a}, \mathbf{b})$ is the product of the linear factors $\varphi_i^+ - \langle \rho, \varphi_i \rangle$ over φ_i^+ which are relevant and which have different signs on \mathbf{a} and \mathbf{b} . This always divides $\Phi(\xi_{\mathbf{a}}, \xi_{\mathbf{b}})$, and the remaining factors are precisely $\Phi_0(\xi_{\mathbf{a}}, \xi_{\mathbf{b}}, \rho)$, which shows the compatibility of (2.11a) and (4.3a) with (3.8a). This confirms that the functor is well-defined. \square

As the formulas above suggest, this isomorphism is essentially induced by that for the abelianizations. It can be understood more clearly if we visualize $\mathfrak{w}(\mathbf{a}, \mathbf{b})$ and $r(\eta_{\mathbf{a}}, \eta_{\mathbf{b}})$ as paths as discussed in Sections 2.3 and 3.3; we work in the case where G is abelian, so $\tilde{\mathfrak{w}} = \mathfrak{w}$ and $\tilde{r} = r$. The action of $\mathfrak{w}(\mathbf{a}, \mathbf{b})$ depends on whether it crosses certain hyperplanes defined by a single level set of φ_i for each i ; on the other hand, the action of $r(\eta_{\mathbf{a}}, \eta_{\mathbf{b}})$ plays a similar role with all integral translates of this level set. However, these translates have different linear factors attached to them, and on any given weight space, at most one of these translates acts non-invertibly. The product $\Phi_0(\eta, \eta', \rho)$ precisely captures the non-invertible factors.

Theorem 4.3 *The functor γ is an equivalence $\widehat{\mathcal{X}}_{\mathcal{C}} \cong \widehat{\mathcal{B}}_{\rho}$ which induces an equivalence $\widehat{\mathcal{X}}_X \cong \widehat{\mathcal{A}}_{\rho}$ sending $\chi \mapsto (o, \rho - \chi)$.*

Before discussing the proof, let us discuss a very simple, but illustrative example:

Example 4.i. Consider the case of $V = \mathbb{C}$ with $G = \mathbb{C}^\times$ acting naturally, and flavor ϕ giving weight $-a$ on \mathbb{C} and $a - 1$ on its dual space.

We can parameterize the space $\mathfrak{t}_{1, \mathbb{R}}$ by the scalar x with which it acts on \mathbb{C} . In these terms, $\varphi_1^{\text{mid}} = x + 1/2$. Thus, we have chambers C_k for each integer k , given by the inequalities $k - 1/2 < x < k + 1/2$. Representatives of these chambers are given by η_k , the point where $x = k$.

We now fix a complex number ρ , and consider $\widehat{\mathcal{B}}_\rho$. The objects in this category are identified up to obvious isomorphism with \mathbb{Z} by considering (η_k, ρ) . From this perspective, our completion just means that we allow power series in $t - \rho$ on each object, where $t \in S_h$ corresponds to the obvious character.

Morphisms $\text{Hom}_{\widehat{\mathcal{B}}_\rho}((\eta_k, \rho), (\eta_m, \rho))$ are a rank-one module over $\mathbb{k}[[t - \rho]]$, generated by $r(\eta_m, \eta_k)$. Let us consider how the isomorphism of Theorem 4.3 behaves in this case. If $k > m$, then $\Phi_0(\eta_m, \eta_k, \rho) = 1$ in all cases (no real number satisfies the first two inequalities of (4.2)), so $\gamma(\mathfrak{w}(m, k)) = r(m, k)$. On the other hand, $k < m$, then

$$\Phi_0(\eta_m, \eta_k, \rho) = \prod_{\substack{k \leq n < m \\ \rho \neq a+n}} t - a - n.$$

Thus, if $\rho - a$ is not between k and m and in their coset of the integers, we have that $r(m, k)r(k, m) = \Phi_0(\eta_m, \eta_k, \rho)$, that is, $\gamma(\mathfrak{w}(m, k))\gamma(\mathfrak{w}(k, m)) = 1$, as the relation (2.4a) requires.

On the other hand, if $k - \rho + a$ and $m - \rho + a$ are integers of opposite sign, then

$$\gamma(\mathfrak{w}(m, k))\gamma(\mathfrak{w}(k, m)) = \frac{\Phi(\eta_m, \eta_k)}{\Phi_0(\eta_m, \eta_k, \rho)} = t - \rho,$$

again, following (2.4a), since $\gamma(t) = t - \rho$. This confirms the isomorphism in this case.

Proof of Theorem 4.3. To show that the functor γ is full, we need to use the fact that the morphism space $\text{Hom}_{\widehat{\mathcal{B}}_\rho}((\eta_{\mathbf{a}}, \rho), (\eta_{\mathbf{b}}, \rho))$ is spanned by the elements $\tilde{r}(\eta_{\mathbf{b}}, w\eta_{\mathbf{a}})w$ for each $w \in W_\rho$, by Corollary 3.15. Using the elements w , we can reduce to showing that $\tilde{r}(\eta_{\mathbf{b}}, \eta_{\mathbf{a}})$ is in this image for all pairs \mathbf{a}, \mathbf{b} are in this image. Let us prove this using induction on the number of root hyperplanes separating \mathbf{a} and \mathbf{b} .

First, note that if there are no such hyperplanes, then

$$r(\eta_{\mathbf{a}}, \eta_{\mathbf{b}}) = \gamma(\gamma^{-1}(\Phi_0(\eta_{\mathbf{a}}, \eta_{\mathbf{b}}, \rho))\mathfrak{w}(\mathbf{a}, \mathbf{b}))$$

lies in the image by (4.3a). Otherwise, we have

$$\tilde{r}(\eta_{\mathbf{b}}, \eta_{\mathbf{a}}) = r(\eta_{\mathbf{b}}, \eta_{\mathbf{b}'})u_\alpha \tilde{r}(\eta_{\mathbf{b}'}, \eta_{\mathbf{a}})$$

where $C'_{\mathbf{b}}$ is the chamber where the path from $C_{\mathbf{a}}$ to $C_{\mathbf{b}}$ crosses the last root hyperplane. If α is a relevant root of the finite system, then u_α is in the image by (4.3b) and we are done. If not, then $\alpha(\rho) \neq 0$, so the action of u_α can be written as $\frac{1}{\alpha}(s_\alpha - 1)$. The image in the Hom space $\text{Hom}_{\widehat{\mathcal{B}}_\rho}((\eta_{\mathbf{a}}, \rho), (s_\alpha \eta_{\mathbf{a}}, \rho))$ is thus $\frac{1}{\alpha}(s_\alpha m_1 + m_2)$ where

$$m_1 \in \text{Hom}_{\widehat{\mathcal{B}}_\rho}((\eta_{\mathbf{a}}, \rho), (s_\alpha \eta_{\mathbf{b}}, s_\alpha \rho)) \quad m_2 \in \text{Hom}_{\widehat{\mathcal{B}}_\rho}((\eta_{\mathbf{a}}, \rho), (\eta_{\mathbf{b}}, \rho))$$

and m_2 is the projection to this Hom space of $r(\eta_{\mathbf{b}}, \eta_{\mathbf{b}'})\tilde{r}(\eta_{\mathbf{b}'}, \eta_{\mathbf{a}})$. Thus, m_2 is in the image by construction. On the other hand, $s_\alpha \mathbf{b}$ is separated from $\eta_{\mathbf{a}}$ by strictly fewer root hyperplanes than $\eta_{\mathbf{b}}$, so $s_\alpha m_1$ is in the image by induction.

Thus, the map $\text{Hom}_{\widehat{\mathcal{X}}_\epsilon}(\mathbf{a}, \mathbf{b}) \rightarrow \text{Hom}_{\widehat{\mathcal{B}}_\rho}(\eta_{\mathbf{a}}, \eta_{\mathbf{b}})$ is surjective. These are both free modules over $\mathbb{C}[[t_k]]$ with rank equal to $\#W_\rho$, so a surjective map between them must be an isomorphism.

Finally, we note that if $rC_{\mathbf{a}}$ contains a point of $X_1^*(T)$, then we can take this to be $\eta_{\mathbf{a}}$ and in $\widehat{\mathcal{B}}_\rho$, we have an isomorphism $(\eta_{\mathbf{a}}, \rho) \cong (0, \rho - \eta_{\mathbf{a}})$ induced by thinking of $\eta_{\mathbf{a}}$ as an element of the affine Weyl group and applying $y_{\eta_{\mathbf{a}}}$. The latter is an object in $\widehat{\mathcal{A}}_\rho$, and every one is isomorphic to one of this form. \square

Remark 4.4. Lemma 4.2 holds even if our assumption on length 0 elements fails, but the functor γ will no longer be full. In this case, we have a finite subgroup Z of length 0 elements fixing ρ , which act on W_ρ by diagram automorphisms. We have an action of $Z \times G_\rho$ on V_ρ . In this case, we need to replace G throughout the construction of the Steinberg category, and instead consider $X_\sigma = (Z \times G_\rho) \times^{B_\rho} V_{\rho, \sigma}$, and the $Z \times G_\rho$ equivariant homology of the fiber products of these spaces.

Of course, the homology of the spaces should have a combinatorial description similar to Theorem 4.3, but this is a sufficiently rare case that we will leave the details of it to another time, or perhaps an ambitious reader.

This theorem also gives us a strong understanding of clans in this context:

Corollary 4.5 *If $rC_{\mathbf{a}}$ and $rC_{\mathbf{a}'}$ correspond to the same element of I' , that is, if their W -images in the dominant chamber lie in the same chamber of the matter arrangement, then $(\eta_{\mathbf{a}}, \rho)$ and $(\eta_{\mathbf{a}'}, \rho)$ lie in the same clan. In particular, $(o, \rho - \chi)$ and $(o, \rho - \chi')$ are in the same clan if and only if χ and χ' are in the same chamber.*

As before, let I be the set of sign vectors σ such that $c_{\sigma, 1} \neq \emptyset$, that is, whose chamber $C_{\sigma, 1}$ contains an element of X . The sum of morphisms in the category A_I gives a finite dimensional algebra; we'll abuse notation and let the same symbol denote this finite-dimensional algebra. The identity 1_σ on σ can be thought of as an idempotent in this algebra.

Corollary 4.6 *The category $\mathcal{A}\text{-mod}_\rho$ of weight modules with weights in $\widehat{W} \cdot \rho$ is equivalent to the category of representations of A_I in finite-dimensional vector spaces where S acts nilpotently; this functor matches the weight space for each weight in $-C_{\sigma, 1}$ with the image of the idempotent 1_σ .*

As discussed in the introduction, this result is very powerful in understanding the representation theory of the Coulomb branch in its own terms; unspooling its consequences will require a series of papers; the case of the Cherednik algebra of $G(\ell, 1, n)$ is covered

in [Web19a], of ADE type quivers in [Kam+19b; Kam+24], and its consequences for Gelfand-Tsetlin modules in [Web20].

This isomorphism also allows us to define a graded lift:

Definition 4.7 *Let $\widetilde{\mathcal{A}\text{-mod}}_\rho$ be the category of pairs (M, g) of a weight module M and a grading g on each of its weight spaces such that $\mathcal{X}_\mathcal{G}$ acts homogeneously.*

Corollary 2.35 shows that this category has a strong positive grading property; that is, it is mixed humorous. It is a bit awkward to talk about projective objects in this graded lift, since these correspond to pro-weight modules, but we can apply [Web15, Lemma 2.4] to instead relate simple modules to the dual canonical basis:

Corollary 4.8 *The classes of the self-dual simple modules in $K^0(\widetilde{\mathcal{A}\text{-mod}}_\rho)$ are the dual canonical basis of the induced pre-canonical structure as defined in [Web15, §2].*

We can also extend this isomorphism to the bimodule ${}_{\phi+\chi}\mathcal{T}_\phi$. This has a natural completion ${}_{\phi+\chi}\widehat{\mathcal{T}}_\phi$ to a bimodule over the categories $\widehat{\mathcal{B}}$ associated to the flavors $\phi + \chi$ and ϕ . Applying Theorem 4.3 to the action of Q on V , we find an isomorphism:

Corollary 4.9 ${}_{I_{\phi+\chi}}\widehat{A}_{I_\phi} \cong {}_{\phi+\chi}\widehat{\mathcal{T}}_\phi$.

4.2. Koszul duality. Assume that A is an algebra over a field \mathbb{k} graded by the non-negative integers with A_0 finite-dimensional and semi-simple. The **Koszul dual** of A is, by definition, the algebra $A^! \cong T_{A_0}A_1^*/R^\perp$ where $R \subset A_1 \otimes_{A_0} A_1$ is the space of quadratic relations, the kernel of the map to A_2 . The representation category of $A^!$ is equivalent to the abelian category $\text{LPC}(A)$ of linear complexes of projectives over A . If an abelian category is equivalent to the modules over an algebra A as above, then the Koszul dual of the category is the category of representations of $A^!$.

Fix a flavor ϕ . Recall from Section 2.7 that we defined a pushforward:

$$L_{I_\phi} = \bigoplus_{C_{\sigma,1} \neq \emptyset} (p_\sigma)_* \mathfrak{S}_{X_\sigma}.$$

Since we will only want to consider this sheaf for the rest of the paper, let us shorten this to $L = L_{I_\phi}$. Combining Theorems 2.60 & 4.3 shows that:

Proposition 4.10 *The Koszul dual of the category $\widetilde{\mathcal{A}\text{-mod}}_\rho$ is the category $p\widetilde{\mathcal{O}}_\mathfrak{g}$.*

Remark 4.11. In general, $D^b(p\mathcal{O}_\mathfrak{g})$ may not be a full subcategory of the derived category of the abelian category \mathcal{D} , so $\text{Ext}_{\mathcal{D}}^\bullet(L, L)^{\text{op}} \cong A_{I_\phi}$ is not the same as the Ext-algebra in the subcategory $\langle L \rangle$. For example, in the “pure gauge field” case of $V = 0$, we find that A_I is the cohomology of BG_ρ , which we can think of as symmetric functions on $\mathfrak{t}_\mathbb{k}$ for the action of the integral Weyl group W_ρ . The algebra \mathcal{A} in this case is just the smash

product $S_h \rtimes W$, and the subcategory $\mathcal{A}\text{-mod}_\rho$ corresponds to the modules which are the sum of their weight spaces over S_1 for the W -translates of ρ .

The ρ -weight space has a natural action of the stabilizer W_ρ , and considering the W_ρ invariants defines a functor from $\mathcal{A}\text{-mod}_\rho$ to $H^*(BG_\rho)$ -modules where we let $H^*(BG_\rho) \cong \mathbb{C}[\mathfrak{t}_k]^{W_\rho}$ act by the ρ -shifted action. This gives the equivalence induced by Theorem 4.3.


In this case, since $H^*(BG_\rho)$ has no elements of degree 1, the only linear complexes over this ring are those with trivial differentials. Thus, the category $\langle L \rangle$ is equivalent to the category of vector spaces.

Example 4.ii. As before, consider the case of $V = \mathbb{C}$ with $G = \mathbb{C}^\times$ acting naturally and flavor ϕ giving weight $-a$ on \mathbb{C}^\times and $a - 1$ on its dual space. In this case $S_1 \cong \mathbb{C}[t]$ with t the natural cocharacter. The algebra $\mathcal{A} = \mathcal{A}^{\text{sph}}$ has generators r^+ and r^- with

$$r^-r^+ = t - a + 1 \qquad r^+r^- = t - a.$$

Note that the elements r^\pm give an isomorphism between the k and $k - 1$ weight spaces unless $k = a$, the action of $t - a$ on the k weight space $\mathcal{W}_k(M)$ has semi-simple part given by the scalar $k - a$ and nilpotent part $t - k$. Thus, if the weights of t are not in $a + \mathbb{Z}$, then all the weight spaces are isomorphic, and we are equivalent to the pure gauge situation, since the only weight of the representation is irrelevant.

If we take weight spaces of the form $a + \mathbb{Z}$, then we have the picture below:



We argued above that the composition around any loop except that connecting a and $a - 1$ is an isomorphism, so there are two isomorphism classes, represented by a and $a - 1$. Thinking of taking this weight space as a functor, the elements r^\pm give morphisms in both directions between them, with the composition in either direction acting by the nilpotent part of t . Thus, we obtain the completed path algebra of an oriented 2-cycle as $\text{End}(P_{0,a} \oplus P_{0,a-1})$. The Koszulity of this path algebra algebra is easily verified directly (since every simple has a length 2 linear projective resolution).

Since this path algebra has no quadratic relations, its quadratic dual is given by imposing all (two) possible quadratic relations: it is the path algebra of an oriented 2-cycle with all length-2 paths set to 0. This is the endomorphism ring of the projective generator in the category of strongly \mathbb{C}^\times -equivariant D -modules on \mathbb{A}^1 generated by the functions and the δ -functions at the origin. The two indecomposable projective D -modules in this category are $D_{\mathbb{A}^1}/D_{\mathbb{A}^1}(z\frac{\partial}{\partial z})$ and $D_{\mathbb{A}^1}/D_{\mathbb{A}^1}(\frac{\partial}{\partial z}z)$; their sum has the desired endomorphism algebra. The untruncated path algebra appears as the Ext-algebra of the sum of simple D -modules $D_{\mathbb{A}^1}/D_{\mathbb{A}^1}z \oplus D_{\mathbb{A}^1}/D_{\mathbb{A}^1}\frac{\partial}{\partial z}$.

Definition 4.12 *Let \mathcal{O}_ρ be the intersection of $\mathcal{A}\text{-mod}_\rho$ with category \mathcal{O} for $\xi \in (\mathfrak{g}^*)^G \cong (\mathfrak{t}^*)^W \subset S_1$, that is, the modules such that the eigenspaces for ξ are finite dimensional and the spectrum is bounded above in real part.*

This is the category we called $\mathcal{O}_{\text{Coulomb}}$ in the introduction.

The category \mathcal{O}_a as defined in [BLPW16, Def. 3.10] will be the direct sum of the categories \mathcal{O}_ρ for all \widehat{W} -orbits. Note that this category can only be nonempty if the weights of V_ρ span \mathfrak{t}^* since otherwise, there will be chambers which are not strongly ξ -bounded. In particular, for generic chambers, V_ρ will be trivial.

Applying the equivalence of Corollary 4.6, we can describe \mathcal{O}_ρ as a subcategory of modules over A_I . Recall that we defined an ideal \mathcal{I}_ξ generated by $e(\sigma)$ such that C_σ is not strongly ξ -bounded.

Theorem 4.13 *The category \mathcal{O}_ρ is equivalent to the category of modules over the quotient $A_{-\xi}$.*

Proof. Because of the minus sign in the isomorphism of Theorem 4.3, a chamber C_σ on which $-\xi$ corresponds to the weight spaces of weights in the chamber $-C_\sigma$.

Since all weight spaces for weights in $-C_\sigma$ are isomorphic, if ξ does not obtain a maximum on $-C_\sigma$, and any of these weight spaces are non-zero, then the corresponding object is not in \mathcal{O} . Thus, indeed, in this case $e(\sigma)$ acts trivially on the A_I -module corresponding to an object in \mathcal{O} . That is, any module coming from category \mathcal{O} factors through the quotient $A_{-\xi}$.

On the other hand, if a module does factor through this quotient, its weight diagram is a union of strongly ξ -bounded chambers. This shows that any level set of ξ on each of these chambers is bounded and therefore has finitely many points in $\widehat{W} \cdot \rho$. Since there are only finitely many such chambers, we find that ξ has a maximum on the weight diagram, and there are only finitely many integral points in this weight diagram for each fixed value of ξ . Thus, the corresponding module lies in category \mathcal{O} . \square

Remark 4.14. An important special case is when $\xi = 0$. In this case, $\mathcal{O}_{\text{Coulomb}}$ is the category of finite-dimensional modules in $\mathcal{A}\text{-mod}_\rho$. Of course, the function 0 achieves its maximum everywhere, a chamber is strongly ξ -bounded if and only if the chamber is itself bounded. The ideal \mathcal{I}_0 is thus generated by $e(\sigma)$ for all the chambers which are not strongly ξ -bounded, since the corresponding clans are infinite and must have trivial weight space for any finite-dimensional representation.

Since $\mathcal{I}_{-\xi}$ is a homogeneous ideal, this induces a graded lift $\tilde{\mathcal{O}}_\rho$ of this category, defined as modules in \mathcal{O}_ρ endowed with a grading on which the induced action of \mathcal{X}_ξ is homogeneous.

Thus, combining Theorems 2.60 and Theorem 4.13, we have that:

$D(\mathcal{O}_\xi^\phi)$ to the category \mathcal{O} on \mathfrak{M}_ξ with the left or right adjoint of this functor for ξ' .

In order to compare with the definition [BLPW16, §8.1], we need to assume that for some k , the functor $K_{k\xi}$ is an equivalence from \mathcal{O}_ξ^ϕ to category \mathcal{O}_a for $A_{k\xi}$, and similarly with ξ' replacing ξ . In this case, $K_{k\xi'} \circ K_{k\xi, \star}^*: D^-(A_{k\xi'}\text{-mod}) \rightarrow D^-(A_{k\xi}\text{-mod})$ is easily seen to be isomorphic to the functor $K_{k\xi'}(D_v/J_{k\xi}) \overset{L}{\otimes} -$. In the case where the moment map is flat and $\mathfrak{M}_{H,\xi} \rightarrow \mathfrak{M}_\xi$ is a symplectic resolution, $K_{k\xi}, K_{k\xi'}$ are equivalences for $k \gg 0$ by [BPW16, Cor. B.1], and $K_{k\xi'}(D_v/J_{k\xi}) \overset{L}{\otimes} -$ is the twisting functor for $k \gg 0$ by [BPW16, Lem. 6.28]. See [BL21, (4.10)] and [Web17b, §4.4] for further discussion of this comparison.

Note that if $\mathfrak{M}_{H,\xi}, \mathfrak{M}_{H,\xi'}$ are both symplectic resolutions and the moment map is flat, then this functor will be an equivalence by [BPW16, Cor. 6.32].

- The pure shuffling functors are generated by composing the inclusion functor i^ϕ of \mathcal{O}_ξ^ϕ into $D^b(\mathfrak{D}_g)$ with its left or right adjoint $i_\star^{\phi'}$, as defined in [BLPW16, §8.2].

On \mathfrak{M}_C :

- The pure twisting functors are generated by tensor product with ${}_\phi T_\phi$ for ϕ and ϕ' both generic flavors and the adjoints of these functors; this is the definition given in [BLPW16, §8.1] when (G, V) is good and a BFN resolution exists, since we have already compared the bimodules in Lemma 3.21. Again, [BPW16, Cor. 6.32] implies that this is an equivalence of categories for generic ϕ, ϕ' .
- The pure shuffling functors are generated by composing the inclusion functor i^ξ of $\mathcal{O}_{\text{Coulomb}}$ into $\mathcal{A}\text{-mod}$ with its left or right adjoint i_\star^ξ in the derived category (i.e. the derived functor of taking the largest quotient or submodule in category \mathcal{O}), as above.

Theorem 4.16 *The Koszul duality of Theorem 4.15 switches pure twisting and shuffling functors matching $\tau^{-\xi'} \circ \tau_\star^{-\xi}$ with $i_\star^{\xi'} \circ i^\xi$ and ${}_{\phi'} T_\phi \overset{L}{\otimes}_{A_\phi} -$ with $i_\star^{\phi'} \circ i^\phi$.*

Proof. The proof of this fact is roughly the same as in [Bra+12, p. 8.24]. The shuffling functors come from inclusion of a projection to a subcategory, and the twisting functors come from projection to an adjoint inclusion of a quotient category; these naturally interchange under Koszul duality.

Now, let us be more precise. Given two subsets $P, P' \subset I'$, we define the $A_{P'} - A_P$ -bimodule ${}_P A_{P'}$ (or similarly a $\mathcal{X}_{P'} - \mathcal{X}_P$ -bimodule ${}_{P'} \mathcal{X}_P$) by simply associating to the pair $(p', p) \in P' \times P$ the vector space $\text{Hom}_{A_{I'}}(p, p')$. This extends in an obvious way to P, P' simply mapping to I' (or to K' , etc.).

Let

$$A_P^\xi := A_P / \mathcal{I}_\xi \quad {}_{P'} A_P^\xi := {}_{P'} A_P / (\mathcal{I}_\xi {}_{P'} A_P^\xi + {}_{P'} A_P^\xi \mathcal{I}_\xi).$$

- Under the equivalence of $D(p\mathcal{O}^\phi)$ to A_{I_ϕ} -dg-mod and $D(\mathcal{O}_\phi^\xi)$ with $A_{I_\phi}^\xi$ -dg-mod, the functor \mathfrak{r}_* is intertwined with inflation of an $A_{I_\phi}^\xi$ -module to an A_{I_ϕ} -module, and thus \mathfrak{r} with its left adjoint $A_{I_\phi}^{-\xi'} \otimes_{A_{I_\phi}}^L -$. Since $\mathfrak{r}^{-\xi'} \circ \mathfrak{r}_*^{-\xi}$ is an equivalence, its left and right adjoints agree and $\mathfrak{r}^{-\xi'} \circ \mathfrak{r}_*^{-\xi}$ is intertwined with $\mathrm{RHom}_{A_{I_\phi}}(A_{I_\phi}^{-\xi'}, -)$.
- The categories \mathcal{O}_ρ for different choices of ξ are equivalent to the modules over $A^{-\xi}$, and the inclusion i^ξ corresponds to the pullback of $A_{I_\phi}^{-\xi}$ -modules to A_{I_ϕ} -modules by the quotient map. Thus, the shuffling functors $i_!^{\xi'} \circ i^\xi$ is intertwined with $A_{I_\phi}^{-\xi'} \otimes_{A_{I_\phi}}^L -$ and $i_*^{\xi'} \circ i^\xi$ with its adjoint.

This shows the first desired match of functors.

- We have an equivalence of the subcategory $\langle \mathfrak{r}(L_{I'}) \rangle \subset \mathfrak{D}_g$ with $A^{-\xi}$ -dg-mod by the same argument as Theorem 2.57. The shuffling functors are determined by taking Ext of $\mathfrak{r}(M)$ and $\mathfrak{r}(M')$ for $M \in \mathcal{O}_\xi^\phi, M' \in \mathcal{O}_\xi^{\phi'}$. From the equivalence above, we find that

$$\mathrm{Ext}^\bullet(\mathfrak{r}(M'), \mathfrak{r}(M)) \cong \mathrm{Ext}^\bullet(\mathfrak{r}(M'), i_*^{\phi'} \circ i^\phi(\mathfrak{r}(M))) \cong {}_{I_{\phi'}} A_{I_\phi}^{-\xi}.$$

Thus, we have that $i_*^{\phi'} \circ i^\phi$ corresponds to ${}_{I_{\phi'}} A_{I_\phi}^{-\xi} \otimes_{A_{I_\phi}^{-\xi}}^L -$ and $i_!^{\phi'} \circ i^\phi$ to $\mathrm{RHom}_{A_{I_\phi}^{-\xi}}({}_{I_\phi} A_{I_{\phi'}}^{-\xi}, -)$.

- Under the isomorphism of Theorem 4.3, the tensor product with ${}_{\phi'} T_\phi$ corresponds to ${}_{I_{\phi'}} A_{I_\phi}^{-\xi} \otimes_{A_{I_\phi}^\xi}^L -$.

This shows the second desired match. □

INDEX OF NOTATION

This index of notation gives a brief description of the main notation used in the paper, together with the section and page where the notation is defined.

§	Symbol	Description	Page
1.0	\mathfrak{M}_H	The Higgs branch $\mu^{-1}(0)//G = \mathrm{Spec}(\mathbb{C}[\mu^{-1}(0)]^G)$.	1
	\mathfrak{M}_C	The Coulomb branch $\mathfrak{M}_C = \mathrm{Spec}(\mathcal{A}_0^{\mathrm{sph}})$.	2
	$\mathfrak{M}_{H,\xi}$	The GIT quotient $\mu^{-1}(0)//_\xi G$ with $\xi \in (\mathfrak{g}^*)^G$ as stability parameter.	3
2.0	G	The gauge group	6
	V	The matter representation	6
	F	The flavor group H/G .	7
	T_\dagger	The maximal torus of the group $\dagger \in \{G, H, Q, F, \dots\}$ induced by the choice of $T_{\tilde{H}}$.	7
	Q	The subgroup of H generated by G and the maximal torus T_H .	7

§	Symbol	Description	Page
	ϕ, ϕ_0	The flavor cocharacter $\phi = (\phi_0(t), t): \mathbb{C}^\times \cong \mathbb{T} \rightarrow T_{\tilde{F}}$ satisfying $\phi(t) \cdot \Omega = t\Omega$.	7
	\tilde{G}	The pullback group $\tilde{G} = \{(h, t) \in \tilde{H} \times \mathbb{C}^\times \mid hG = \phi(t) \in \tilde{F}\}$.	7
2.1	$\mathfrak{t}_{1, \dagger}$	The space of elements of $\mathfrak{t}_{\tilde{H}, \dagger}$ with coefficients in $\dagger = \mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ which agree with ϕ under projection of $\mathfrak{t}_{\tilde{F}}$.	8
	V_i	A 1-dimensional space of V , compatible with the weight decomposition.	8
	φ_i	The weight of the subspace V_i .	8
	V_σ	The sum of the subspaces V_i with $\sigma_i = +$ for a sign sequence σ .	8
	\mathcal{K}	The set of pairs of sign vectors in $\{+, 0, -\}^d$ and compatible Borels.	9
	φ_i^{mid}	$\varphi_i^{\text{mid}} = \frac{1}{2}(\varphi_i^+ - \varphi_i^-) = \varphi_i + \frac{1}{2}\nu$	9
	$C_\sigma, C_{\sigma, w}$	The subset of $\mathfrak{t}_{1, \mathbb{R}}$ where the value of $\varphi_i^{\text{mid}}(\gamma)$ has the sign of σ_i , and its intersection with a Weyl chamber: $C_{\sigma, w} = C_\sigma \cap w \cdot \mathcal{C}$.	10
	V_σ, V_γ	The subspaces $V_\sigma \cong \bigoplus_{\sigma_i=+} V_i$ and $V_\gamma = \{x \in V \mid \lim_{t \rightarrow 0} \gamma(t) \cdot x \text{ exists}\}$.	10
2.2	$\mathbb{H} \times^{\mathbb{G}} X$	The bundle with fiber X associated to the principal \mathbb{H} -bundle \mathbb{G} .	12
	$X_{\sigma, w}, X_\sigma$	The vector bundle $G \times^{wBw^{-1}} V_\sigma$.	12
	\mathbb{X}_I	The Steinberg variety (2.3) for the index set $I \subset \mathcal{K}$.	12
	\mathcal{X}_I	The Steinberg category associated to the set I	12
	S, S_H, S_F	The polynomial rings $S = \text{Sym}(\mathfrak{t}_{\mathbb{k}}^*) \cong H_G^*(G/B; \mathbb{k})$, $S_Q = \text{Sym}(\mathfrak{t}_Q^*; \mathbb{k})$ and $S_F = \text{Sym}(\mathfrak{t}_F^*; \mathbb{k})$.	13
2.3	I, I'	The sign vectors such that $c_{\sigma, 1} \neq \emptyset$ (resp. $C_{\sigma, 1} \neq \emptyset$) for some flavor.	15
	I_ϕ, I'_ϕ	The sign vectors such that $c_{\sigma, 1} \neq \emptyset$ (resp. $C_{\sigma, 1} \neq \emptyset$) for a fixed flavor ϕ .	15
	$\varphi(\sigma, \sigma')$	Product of the weights such that $\sigma_i = +$ and $\sigma'_i = -$.	16
	$\varphi(\sigma, \sigma', \sigma'')$	Product of the weights such that $\sigma_i = \sigma''_i = -\sigma'_i$.	16
	$e(\sigma)$	The identity morphism $e(\sigma)$ on each object $\sigma \in I'$; we can also consider this as an idempotent in $A_{I'}$.	17
	$A_{I'}, A_P$	The combinatorial Steinberg category (Definition 2.11) and its variation with object set P .	19
	$\tilde{w}(\sigma, \sigma', w)$	The morphism defined in (2.6).	21
2.4	J, J'	Subset of \mathcal{K} such that $c_{\sigma, w} \neq \emptyset$ (resp. $C_{\sigma, w} \neq \emptyset$).	26
	\mathcal{C}	The set of chambers of the arrangement $\varphi_i^{\text{mid}}(\xi) = n$ for $n \in \mathbb{Z}$ (Definition 2.20).	27

§	Symbol	Description	Page
2.6	β_σ	The cocharacter which measures the ξ -unboundedness of the chamber C_σ .	32
	\mathcal{KN}	The set of cocharacters of the form β_σ for $\sigma \in I'_\phi$.	32
	$\mathfrak{s}(\sigma, \beta)$	The standard morphism $\mathfrak{s}(\sigma, \beta) = \tilde{\mathfrak{w}}(\sigma_{++}, \beta, \sigma, w)$.	33
	$W_\beta, G_\beta, \mathbb{V}_\beta$	The stabilizer of β in W , the corresponding Levi, and fixed space of β .	33
2.7	L_σ, L	The pushforward D-module $L_\sigma = p_* \mathfrak{S}_{X_\sigma}$, and the sum $L = \bigoplus_{\sigma \in I'} L_\sigma$.	38
2.8	${}^+A_P, {}^+A_P^!$	The unique basic graded algebra Morita equivalent to A_P and its quadratic dual.	40
2.9	$\mathcal{D}^{\text{ured}}$	The full subcategory of objects such that $K_{k\xi}(M) = \{m \in M \mid \mu_q(X)m = \xi(X)m \forall X \in \mathfrak{g}\} = 0$ for all $k \gg 0$.	42
	\mathfrak{D}_g	The quotient $\mathcal{D}/\mathcal{D}^{\text{ured}}$.	42
	\mathcal{D}^{uns}	The subcategory of D -modules in \mathcal{D} whose singular support is contained in the unstable locus $\mu^{-1}(0)^{\text{uns}}$.	43
2.10	$p\mathcal{O}_g$	The subcategory in \mathcal{D} of objects M which have a good filtration such that each G -invariant function on $\mu^{-1}(0)$ with positive \mathbb{T} -weight acts trivially on $\text{gr } M$.	44
	\mathcal{O}_g	The image of $p\mathcal{O}_g$ in the quotient \mathfrak{D}_g .	44
2.12	\mathcal{I}_ξ	The ideal in A_P generated by the idempotents $e(\sigma)$ with C_σ not strongly ξ -bounded.	49
	$\tilde{\mathcal{O}}_g, \widetilde{p\mathcal{O}}_g$	The graded lifts of \mathcal{O}_g and $p\mathcal{O}_g$ induced by Theorem 2.57.	50
3.1	$\mathcal{J}, \mathcal{F}\ell$	The standard Iwahori subgroup and its space of cosets, the affine flag variety.	53
	$\mathcal{X}_U, \mathcal{X}_\eta$	The infinite-dimensional vector bundle $\mathcal{X}_U = G((t)) \times^J U$ or $\mathcal{X}_\eta = G((t)) \times^J U_\eta$.	53
	${}_U\mathcal{X}_{U'}$	${}_U\mathcal{X}_{U'} = \{(g\mathcal{J}, v(t)) \in \mathcal{F}\ell \times U' \mid g^{-1} \cdot v(t) \in U\}$ and ${}_\eta\mathcal{X}_{\eta'} = {}_{U_\eta}\mathcal{X}_{U_{\eta'}}$	53
	$\mathcal{A}, \mathcal{A}^{\text{sph}}$	The Iwahori and original BFN quantized Coulomb branch algebras (Definition 3.2).	54
3.2	\mathcal{J}_η	Parahoric subgroup whose Lie algebra is non-negative weight spaces for the adjoint action of η on $\mathfrak{g}((t))$ (Definition 3.6).	55
	U_η	The subspace in $V((t))$ of elements of weight $\geq -1/2$ under η (Definition 3.6).	55
	\mathcal{B}	The extended BFN category (Definition 3.7).	57
3.3	y_w	The morphism in \mathcal{B} defined by the graph of an affine Weyl group element (3.3).	58

§	Symbol	Description	Page
	$r(\eta, \eta')$	The morphism (3.4).	58
	$\Phi(\eta, \eta')$	The product of the terms $\varphi_i^+ - nh$ over pairs $(i, n) \in [1, d] \times \mathbb{Z}$ such that the inequalities $\varphi_i(\eta) > n - \frac{1}{2}$, $\varphi_i(\eta') < n - \frac{1}{2}$ hold (Definition 3.10).	58
	$\Phi(\eta, \eta', \eta'')$	The product of the terms $\varphi_i^+ - nh$ over pairs $(i, n) \in [1, d] \times \mathbb{Z}$ such that (3.5a) or (3.5b) (Definition 3.10).	59
	$\tilde{r}_\pi, \tilde{r}(\eta', \eta)$	The morphism associated to a path by (3.10).	63
3.4	\mathcal{A}_Q	The quantum Coulomb branch of the action of Q on V .	66
	\mathcal{B}^{def}	The deformed extended BFN category (Definition 3.18).	67
	$\phi+\chi T_\phi$	The twisting bimodule (Definition 3.20).	68
3.5	$\mathcal{W}_{v, \eta}$	The weight functor (3.13).	70
	$\widehat{\mathcal{B}}$	The completed category that describes the weight modules over \mathcal{B} .	70
4.1	$rC_{\mathbf{a}}$	The chamber of the unrolled relevant arrangement (4.1).	72
	$\mathcal{A}\text{-mod}_\rho$	The category of weight modules with weights in $\widehat{W} \cdot \rho$.	75
	$\widetilde{\mathcal{A}\text{-mod}_\rho}$	The graded lift of $\mathcal{A}\text{-mod}_\rho$ (Definition 4.7).	75

REFERENCES

- [APM20] Alexander Braverman, Pavel Etingof, and Michael Finkelberg. “Cyclo-tomic double affine Hecke algebras”. *Annales scientifiques de l’École normale supérieure* 53.5 (2020), pp. 1249–1312.
- [BB81] Alexander Beilinson and Joseph Bernstein. “Localisation de g -modules”. *C. R. Acad. Sci. Paris Sér. I Math.* 292.1 (1981), pp. 15–18.
- [BBD82] A. A. Beilinson, J. Bernstein, and P. Deligne. “Faisceaux pervers”. *Analysis and topology on singular spaces, I (Luminy, 1981)*. Vol. 100. Astérisque. Paris: Soc. Math. France, 1982, pp. 5–171.
- [BD00] Roger Bielawski and Andrew S. Dancer. “The geometry and topology of toric hyperkähler manifolds”. *Communications in Analysis and Geometry* 8.4 (2000), pp. 727–760.
- [Bel23] Gwyn Bellamy. *Coulomb Branches have symplectic singularities*. Apr. 18, 2023. arXiv: 2304.09213 [math]. URL: <http://arxiv.org/abs/2304.09213>. Pre-published.
- [BFN18] Alexander Braverman, Michael Finkelberg, and Hiraku Nakajima. “Towards a mathematical definition of Coulomb branches of 3-dimensional $\mathcal{N} = 4$ gauge theories, II”. *Advances in Theoretical and Mathematical Physics* 22.5 (2018), pp. 1071–1147.
- [BK04] Roman Bezrukavnikov and Dmitry Kaledin. “Fedosov quantization in algebraic context”. *Moscow Mathematical Journal* 4.3 (2004), pp. 559–592, 782.

- [BL21] Roman Bezrukavnikov and Ivan Losev. “Etingof’s conjecture for quantized quiver varieties”. *Inventiones mathematicae* 223.3 (Mar. 2021), pp. 1097–1226.
- [BLPW16] Tom Braden, Anthony Licata, Nicholas Proudfoot, and Ben Webster. “Quantizations of conical symplectic resolutions II: category \mathcal{O} and symplectic duality”. *Astérisque* 384 (2016), pp. 75–179.
- [BPW16] Tom Braden, Nicholas Proudfoot, and Ben Webster. “Quantizations of conical symplectic resolutions I: local and global structure”. *Astérisque* 384 (2016), pp. 1–73.
- [Bra+10] Tom Braden, Anthony Licata, Nicholas Proudfoot, and Ben Webster. “Gale duality and Koszul duality”. *Advances in Mathematics* 225.4 (Nov. 2010), pp. 2002–2049.
- [Bra+12] Tom Braden, Anthony Licata, Nicholas Proudfoot, and Ben Webster. “Hypertoric category \mathcal{O} ”. *Advances in Mathematics* 231.3-4 (Oct. 2012), pp. 1487–1545.
- [Bru23] Jonathan Brundan. *Graded triangular bases*. May 8, 2023. arXiv: 2305.05122 [math]. URL: <http://arxiv.org/abs/2305.05122>. Pre-published.
- [CCM81] J Carlson, H Clemens, and J Morgan. “On the mixed Hodge structure associated to 3 of a simply connected complex projective manifold” (1981).
- [CG97] Neil Chriss and Victor Ginzburg. *Representation theory and complex geometry*. Boston, MA: Birkhäuser Boston Inc., 1997, pp. x+495.
- [Cre+14] Stefano Cremonesi, Giulia Ferlito, Amihay Hanany, and Noppadol Mekareeya. “Coulomb branch and the moduli space of instantons”. *Journal of High Energy Physics* 2014.12 (2014), p. 103.
- [CS22] Joana Cirici and Anna Sopena-Gilboy. “Filtered A-infinity structures in complex geometry”. *Proceedings of the American Mathematical Society* 150.9 (June 10, 2022), pp. 4067–4082.
- [Del+75] Pierre Deligne, Phillip Griffiths, John Morgan, and Dennis Sullivan. “Real homotopy theory of Kähler manifolds”. *Inventiones Mathematicae* 29.3 (1975), pp. 245–274.
- [Dim+20] Tudor Dimofte, Niklas Garner, Michael Geracie, and Justin Hilburn. “Mirror symmetry and line operators”. *Journal of High Energy Physics* 2020.2 (Feb. 2020), p. 75.
- [Dri04] Vladimir Drinfeld. “DG quotients of DG categories”. *Journal of Algebra* 272.2 (2004), pp. 643–691.
- [HY] Justin Hilburn and Philsang Yoo. *Symplectic duality and geometric Langlands II: The local case*.
- [Kal08] Dmitry Kaledin. “Derived equivalences by quantization”. *Geometric and Functional Analysis* 17.6 (2008), pp. 1968–2004.
- [Kam+19a] Joel Kamnitzer, Peter Tingley, Ben Webster, Alex Weekes, and Oded Yacobi. “Highest weights for truncated shifted Yangians and product monomial crystals”. *Journal of Combinatorial Algebra* 3.3 (2019), pp. 237–303.
- [Kam+19b] Joel Kamnitzer, Peter Tingley, Ben Webster, Alex Weekes, and Oded Yacobi. “On category \mathcal{O} for affine Grassmannian slices and categorified

- tensor products”. *Proceedings of the London Mathematical Society* 119.5 (Nov. 2019), pp. 1179–1233.
- [Kam+24] Joel Kamnitzer, Ben Webster, Alex Weekes, and Oded Yacobi. “Lie algebra actions on module categories for truncated shifted yangians”. *Forum of Mathematics, Sigma* 12 (2024), e18.
- [Kem18] George Kempf. *Instability in invariant theory*. July 8, 2018. arXiv: 1807.02890 [math]. URL: <http://arxiv.org/abs/1807.02890>. Pre-published.
- [KL09] Mikhail Khovanov and Aaron D. Lauda. “A diagrammatic approach to categorification of quantum groups. I”. *Representation Theory* 13 (2009), pp. 309–347.
- [KN18] Ryosuke Kodera and Hiraku Nakajima. “Quantized Coulomb branches of Jordan quiver gauge theories and cyclotomic rational Cherednik algebras”. *String-Math 2016*. Vol. 98. Proc. Sympos. Pure math. Amer. Math. Soc., Providence, RI, 2018, pp. 49–78.
- [KR08] Masaki Kashiwara and Raphaël Rouquier. “Microlocalization of rational Cherednik algebras”. *Duke Mathematical Journal* 144.3 (2008), pp. 525–573.
- [KWWY14] Joel Kamnitzer, Ben Webster, Alex Weekes, and Oded Yacobi. “Yangians and quantizations of slices in the affine Grassmannian”. *Algebra & Number Theory* 8.4 (2014), pp. 857–893.
- [Los21] Ivan Losev. *Localization theorems for quantized symplectic resolutions*. Mar. 20, 2021. arXiv: 2103.11193 [math]. URL: <http://arxiv.org/abs/2103.11193>. Pre-published.
- [LY17] George Lusztig and Zhiwei Yun. “ \mathbb{Z}/m -graded Lie algebras and perverse sheaves, I”. *Representation Theory of the American Mathematical Society* 21.12 (Sept. 14, 2017), pp. 277–321.
- [MN14a] Kevin McGerty and Thomas Nevins. “Derived equivalence for quantum symplectic resolutions”. *Selecta Mathematica. New Series* 20.2 (2014), pp. 675–717.
- [MN14b] Kevin McGerty and Thomas Nevins. *Morse decomposition for D-module categories on stacks*. Feb. 28, 2014. arXiv: 1402.7365 [math]. URL: <http://arxiv.org/abs/1402.7365>. Pre-published.
- [MOS09] Volodymyr Mazorchuk, Serge Ovsienko, and Catharina Stroppel. “Quadratic duals, Koszul dual functors, and applications”. *Transactions of the American Mathematical Society* 361.3 (2009), pp. 1129–1172.
- [MV98] Ian M. Musson and Michel Van den Bergh. “Invariants under tori of rings of differential operators and related topics”. *Memoirs of the American Mathematical Society* 136.650 (1998), pp. viii+85.
- [Nak] Hiraku Nakajima. *Modules of quantized Coulomb branches*.
- [Nak16] Hiraku Nakajima. “Towards a mathematical definition of Coulomb branches of 3-dimensional $N=4$ gauge theories, I”. *Advances in Theoretical and Mathematical Physics* 20.3 (2016), pp. 595–669.
- [Nam11] Yoshinori Namikawa. “Poisson deformations of affine symplectic varieties”. *Duke Mathematical Journal* 156.1 (2011), pp. 51–85.

-
- [Rou08a] Raphael Rouquier. *2-Kac-Moody algebras*. Dec. 30, 2008. arXiv: 0812.5023 [math]. URL: <http://arxiv.org/abs/0812.5023>. Pre-published.
- [Rou08b] Raphael Rouquier. “q-Schur algebras and complex reflection groups”. *Moscow Mathematical Journal* 8.1 (2008), pp. 119–158, 184.
- [Sau13] Julia Sauter. *Generalized quiver Hecke algebras*. July 3, 2013. arXiv: 1306.3892 [math]. URL: <http://arxiv.org/abs/1306.3892>. Pre-published.
- [Sch11] Olaf M. Schnürer. “Equivariant sheaves on flag varieties”. *Mathematische Zeitschrift* 267.1-2 (2011), pp. 27–80.
- [SVV14] P. Shan, M. Varagnolo, and E. Vasserot. “Koszul duality of affine Kac-Moody algebras and cyclotomic rational double affine Hecke algebras”. *Advances in Mathematics* 262 (2014), pp. 370–435.
- [SW14] Catharina Stroppel and Ben Webster. *Quiver Schur algebras and q-Fock space*. July 15, 2014. arXiv: 1110.1115 [math]. URL: <http://arxiv.org/abs/1110.1115>. Pre-published.
- [Tel21] Constantin Teleman. “The rôle of Coulomb branches in 2D gauge theory”. *Journal of the European Mathematical Society* 23.11 (June 9, 2021), pp. 3497–3520.
- [VV10] Michela Varagnolo and Eric Vasserot. “Double affine Hecke algebras and affine flag manifolds, I”. *Affine flag manifolds and principal bundles*. Trends math. Birkhäuser/Springer Basel AG, Basel, 2010, pp. 233–289.
- [VV11] Michela Varagnolo and Eric Vasserot. “Canonical bases and KLR-algebras”. *Journal Fur Die Reine Und Angewandte Mathematik* 659 (2011), pp. 67–100.
- [Web] Ben Webster. *Coherent sheaves and quantum Coulomb branches I: tilting bundles from integrable systems*. URL: <https://arxiv.org/abs/1905.04623>. Pre-published.
- [Web15] Ben Webster. “Canonical bases and higher representation theory”. *Compositio Mathematica* 151.1 (2015), pp. 121–166.
- [Web16] Ben Webster. “Tensor product algebras, Grassmannians and Khovanov homology”. *Physics and mathematics of link homology*. Contemp. Math. 680 (2016), pp. 23–58.
- [Web17a] Ben Webster. “Knot invariants and higher representation theory”. *Memoirs of the American Mathematical Society* 250.1191 (2017), p. 141.
- [Web17b] Ben Webster. “On generalized category \mathcal{O} for a quiver variety”. *Mathematische Annalen* 368.1 (June 1, 2017), pp. 483–536.
- [Web17c] Ben Webster. “Rouquier’s conjecture and diagrammatic algebra”. *Forum of Mathematics. Sigma* 5 (2017), e27, 71.
- [Web19a] Ben Webster. “Representation theory of the cyclotomic Cherednik algebra via the Dunkl-Opdam subalgebra”. *New York Journal of Mathematics* 25 (2019), pp. 1017–1047.
- [Web19b] Ben Webster. “Weighted Khovanov-Lauda-Rouquier algebras”. *Documenta Mathematica* 24 (2019), pp. 209–250.

- [Web20] Ben Webster. *Gelfand-Tsetlin modules in the Coulomb context*. Dec. 16, 2020. arXiv: 1904.05415 [math]. URL: <http://arxiv.org/abs/1904.05415>. Pre-published.
- [Web22] Ben Webster. *Coherent sheaves and quantum Coulomb branches II: quiver gauge theories and knot homology*. Nov. 3, 2022. arXiv: 2211.02099 [math-ph]. URL: <http://arxiv.org/abs/2211.02099>. Pre-published.
- [WWY20] Ben Webster, Alex Weekes, and Oded Yacobi. “A quantum Mirković-Vybornov isomorphism”. *Representation Theory of the American Mathematical Society* 24.2 (Jan. 16, 2020), pp. 38–84.