

# A well-scaling natural orbital theory

Ralph Gebauer<sup>\*</sup>, Morrel H. Cohen<sup>†‡</sup>, and Roberto Car<sup>‡§</sup>

<sup>\*</sup>ICTP – The Abdus Salam International Centre for Theoretical Physics, Strada Costiera 11, 34151 Trieste, Italy, <sup>†</sup>Department of Physics and Astronomy, Rutgers University, Piscataway, NJ 08854, <sup>‡</sup>Department of Chemistry, Princeton University, Princeton, NJ 08544, and <sup>§</sup>Department of Physics, Princeton University, Princeton, NJ 08544

Published in Proceedings of the National Academy of Sciences of the United States of America, vol. 113 no. 46 p. 12913 (2016)

**We introduce a new energy functional for ground-state electronic structure calculations. Its variables are the natural spin-orbitals of singlet many-body wave functions and their joint occupation probabilities deriving from controlled approximations to the two-particle density matrix that yield algebraic scaling in general, and Hartree-Fock scaling in its seniority-zero version. Results from the latter version for small molecular systems are compared with those of highly accurate quantum-chemical computations. The energies lie above full configuration interaction calculations, close to doubly occupied configuration interaction calculations. Their accuracy is considerably greater than that obtained from current density-functional theory approximations and from current functionals of the one-particle density-matrix.**

Electronic structure | correlation | density matrix

## Introduction

Computing the ground-state energy of  $N$  interacting electrons is central to quantum chemistry, condensed-matter physics, and related sciences. Reducing its complexity significantly below that of the many-body wave function is a major goal. Density-functional theory (DFT)<sup>1</sup> [1, 2] achieved maximal reduction using electron density as the basic variable. DFT transformed many sciences and technologies, but finding accurate, parameter-free approximations to its exchange-correlation energy functional that avoid self-interaction and capture strong electron correlation is difficult. One-particle density-matrix (1-DM) functional theories [3] have one more degree of complexity. In them, the 1-DM is often represented by its eigenvalues, the occupation numbers, and the corresponding eigenvectors, the natural spin-orbitals (NSOs), e.g. [4, 5]. While avoiding the mean-field form of the 1-DM of DFT [2], the approximations to the exchange-correlation functional of the 1-DM have difficulties like those of the DFT approximations. Two-particle density-matrix (2-DM) functional theories, e.g. [6–8], are less reduced. The ground-state energy is a known, explicit functional of the 2-DM in Coulombic systems. However, while useful complete conditions are known for the  $N$ -representability of the 1-DM, the form of the known complete conditions [7] for the 2-DM renders them unsuitable for practical application. Nevertheless, major progress has been made towards necessary conditions for  $N$ -representability that can be systematically refined [9, 10]. While not variational, the resulting calculations are almost as accurate as full configuration interaction (FCI) calculations [9, 11, 12]. Their computational cost scales as at least the 6th power of the basis-set size, significantly worse than the asymptotic 3rd power scaling of Hartree-Fock theory, DFT, and 1-DM theories. In our new natural-orbital-functional theory, OP-NOFT, the basic variables are the NSOs, their occupation numbers, and their joint occupation probabilities (OP). That allows us to represent the 2-DM accurately, transcending the limitations of the 1-DM theories. Its general form contains single-NSO through 4-NSO joint-occupation probabilities (n-OPs) and scales as the 5th power of the basis-set size. Its simplest formulation, for seniority 0, OP-NOFT-0, approximates doubly-occupied configuration interaction (DOCI) [13]. It contains only 1- and 2-natural-orbital (NO) OPs and retains the 3rd-power scaling of Hartree-Fock energy-functional minimization with a higher prefactor. It describes the dissociation of simple diatomic molecules and multi-atom chains with accuracy that can be comparable to that of

DOCI, which uses a compact basis of Slater determinants (SD) but retains combinatorial scaling. In 4-electron systems, illustrated with the Paldus  $H_4$  test [14], it yields results identical to those of DOCI. OP-NOFT-0 is powerful at high correlation, i.e. for static correlation at intermediate and large interatomic separations where Hartree-Fock fails due to the multi-reference character of the ground-state wavefunction. There, OP-NOFT-0 outperforms Hartree-Fock, DFT with standard approximations, and quantum-chemistry methods such as (single-reference) coupled cluster with single, double and perturbative triple electron-hole excitations (CCSD(T)), a standard of accuracy near equilibrium separations. This introduction of higher-order OPs as variational parameters, with closure of the theory at their level, is the essential novelty of our work and is responsible for its favorable scaling.

## OP-NOFT, general formulation

**The NSO basis.** We consider time-reversal invariant saturated systems with non-degenerate, singlet ground states. The inverse approach [6–8] starts from the  $N$ -representability conditions on the 2-DM. Instead, we take a forward approach: we introduce a specific form for the trial wavefunction and derive the 2-DM explicitly. Our starting point is that of conventional FCI, except that our one-particle basis is the complete set of NSOs of the trial function  $\Psi$ ,  $\psi_k(x) = \phi_k(\mathbf{r})\chi_k(\sigma)$ , with  $\mathbf{r}$  space and  $\sigma$  spin coordinates. The NOs  $\phi_k(\mathbf{r})$  can be real and are independent of the spin function. The complete set of  $N$ -electron orthonormal SDs  $\Phi_{\mathbf{n}}(x_1, x_2, \dots, x_N)$ ,  $\mathbf{n} = k_1, k_2, \dots, k_N$ , formed from its NSOs supports representation

### Significance

Computations of locations of nuclei and movement of electrons within molecules and materials are widely used in science and technology. Direct computation of a system's wave function for that purpose becomes impractical as system size grows. Current alternative methods can have difficulty with strongly-correlated electron motion or spurious electron self-interaction. By using "natural spin orbitals" to describe the motion of individual electrons, solving for them together with their joint and individual probabilities of occurrence within the system, we are able to account better for electron correlation when strong while avoiding self-interaction and maintaining the growth of computation cost with system size at the level of Hartree-Fock theory. Our numerical results for some small test molecules are very good.

## Reserved for Publication Footnotes

<sup>1</sup>Section S8 of the online Supporting Information is a list of acronyms.

of any trial wavefunction  $\Psi(x_1, x_2, \dots, x_N)$  as the expansion

$$\Psi(x_1, \dots, x_N) = \sum_{\mathbf{n}} C_{\mathbf{n}} \Phi_{\mathbf{n}}(x_1, \dots, x_N). \quad [1]$$

As the ground-state wave function can be chosen to be real, so can be the trial functions and the normalized  $C_{\mathbf{n}}$  ( $\sum_{\mathbf{n}} C_{\mathbf{n}}^2 = 1$ ). The NSOs vary with the trial function or the coefficients in the search for the ground state. The combinatorial complexity of determining the ground-state energy by variation of the  $C_{\mathbf{n}}$  is composed of the separate combinatorial complexities of the signs and magnitudes of the coefficients  $C_{\mathbf{n}}$ . We use distinct reductive approximations for their signs and magnitudes. The signs depend on the sign convention chosen for the SDs. We use the Leibniz form for the SDs,

$$\Phi_{\mathbf{n}}(x_1, \dots, x_N) = \frac{1}{\sqrt{N!}} \sum_p \text{sgn}\{P_p\} P_p \psi_{k_1}(x_1) \cdots \psi_{k_N}(x_N). \quad [2]$$

The sum is over the elements of the symmetric group of order  $N$ , the permutations  $P_p$ . The sign of  $\Phi_{\mathbf{n}}$  is fixed by the ordering  $k_1 < k_2 < \dots < k_N$  in the product of the NSOs  $\psi_{k_i}$  in **2**. The SDs and their coefficients can then be specified by listing the NSOs occupied in the SDs, i.e. by the index  $\mathbf{n}$ .

**The 1-DM, the orthogonality constraint, the PDC.** The 1-DM of  $\Psi$ ,

$$\rho(x', x) = N \int dx_2 \cdots dx_N \Psi(x', x_2, \dots, x_N) \Psi(x, x_2, \dots, x_N), \quad [3]$$

becomes

$$\rho(x', x) = \sum_{i \neq j} \left[ \sum_{\mathbf{m} \not\ni i, j} C_{i, \mathbf{m}} C_{j, \mathbf{m}} \right] \psi_i(x') \psi_j(x) \quad [4]$$

after **1** and **2** are inserted into **3**. In **4** the subindex  $\mathbf{m}$  specifies the  $N - 1$  NSOs present in  $\Phi_{i, \mathbf{m}}$  and  $\Phi_{j, \mathbf{m}}$ , excluding  $\psi_i$  and  $\psi_j$ . As the  $\psi$  are the NSOs of  $\Psi$ , the eigenfunctions of  $\rho(x', x)$ , the bracketed quantity in **4** must vanish for  $i \neq j$ . Regard the coefficients  $C_{i, \mathbf{m}}$  and  $C_{j, \mathbf{m}}$  as the components of vectors  $\mathbf{C}_i$  and  $\mathbf{C}_j$  and the bracket as their scalar product  $\mathbf{C}_i \cdot \mathbf{C}_j$ , which must vanish. There are two realizations of this orthogonality constraint. In the first, and most general form (OC), the presence of  $\Phi_{i, \mathbf{m}}$  in  $\Psi$  does not exclude the presence of  $\Phi_{j, \mathbf{m}}$ . The individual terms in the scalar product need not vanish. The second, the pair-difference constraint (PDC), is a special case of the OC, in which the presence of  $\Phi_{i, \mathbf{m}}$  excludes  $\Phi_{j, \mathbf{m}}$  so that either  $C_{i, \mathbf{m}}$  or  $C_{j, \mathbf{m}}$  is zero for each  $\mathbf{m}$ , and the sum vanishes term by term. Under the PDC, those  $\Phi$  present in the expansion of  $\Psi$  must differ from one another by at least two NSOs. The OC is necessary and sufficient for  $N$ -representability, whereas the PDC is only sufficient. We impose the PDC on the  $\Phi$  as a simplifying variational approximation. The PDC proved well satisfied in the FCI result for  $\text{H}_8$  using the minimal basis set STO-6G.

Under the OC or PDC,  $\rho(x', x)$  takes the diagonal form

$$\rho(x', x) = \sum_k p_1(k) \psi_k(x') \psi_k(x). \quad [5]$$

Here, the  $p_1(k) = \sum_{\mathbf{n}} C_{\mathbf{n}}^2 \nu_{k, \mathbf{n}}$ , where  $\nu_{k, \mathbf{n}} = 1$  if  $k \in \mathbf{n}$  and 0 otherwise, are the eigenvalues of  $\rho(x', x)$ , the occupation numbers or occupation probabilities (1-OP) of its eigenfunctions  $\psi_k$ . They satisfy the necessary and sufficient conditions  $0 \leq p_1(k) \leq 1$  and  $\sum_k p_1(k) = N$ . In general, only  $M > N$  occupation numbers  $p_1(k)$  are non-negligible, and only the corresponding *active* NSOs need be included in the representation of any trial function, providing a natural cutoff. The 1-DM is thus of algebraic complexity in the 1-OPs and the NSOs.

**The 2-DM, the sign conjecture, the  $\xi$ -approximation.** The 2-DM of  $\Psi$ ,

$$\pi(x'_1 x'_2; x_1 x_2) = N(N-1) \int dx_3 \cdots dx_N \Psi(x'_1, x'_2, x_3, \dots, x_N) \Psi^*(x_1, x_2, x_3, \dots, x_N),$$

becomes

$$\pi(x'_1 x'_2; x_1 x_2) = \sum_{\substack{i < i', j < j', \mathbf{m} \\ i, i', j, j' \notin \mathbf{m}}} C_{ii' \mathbf{m}} C_{jj' \mathbf{m}} (\psi_i(x'_1) \psi_{i'}(x'_2) - \psi_{i'}(x'_1) \psi_i(x'_2)) (\psi_j(x_1) \psi_{j'}(x_2) - \psi_{j'}(x_1) \psi_j(x_2)). \quad [6]$$

$\pi$  separates into a part  $\pi^d$  diagonal in the indices, i.e. with  $ii' = jj'$ , and an off-diagonal part,  $\pi^{od}$ , with  $ii' \neq jj'$ :

$$\pi^d(x'_1 x'_2; x_1 x_2) = \frac{1}{2} \sum_{i \neq j} p_{11}(ij) (\psi_i(x'_1) \psi_j(x'_2) - \psi_j(x'_1) \psi_i(x'_2)) (\psi_i(x_1) \psi_j(x_2) - \psi_j(x_1) \psi_i(x_2)) \quad [7]$$

$$\pi^{od}(x'_1 x'_2; x_1 x_2) = \sum_{\substack{i < i' \neq j < j', \mathbf{m} \\ i, i', j, j' \notin \mathbf{m}}} C_{ii' \mathbf{m}} C_{jj' \mathbf{m}} (\psi_i(x'_1) \psi_{i'}(x'_2) - \psi_{i'}(x'_1) \psi_i(x'_2)) (\psi_j(x_1) \psi_{j'}(x_2) - \psi_{j'}(x_1) \psi_j(x_2)) \quad [8]$$

Electron correlation is expressed through  $\pi^{od}$ . The analogous off-diagonal part of  $\rho(x', x)$  is suppressed by the OC, an advantage of the NSO basis. Note that the PDC has eliminated 3-index terms from  $\pi^{od}$  in **8**.

The  $p_{11}(ij) = \sum_{\mathbf{n}} C_{\mathbf{n}}^2 \nu_{i, \mathbf{n}} \nu_{j, \mathbf{n}}$  in  $\pi^d$  are joint 2-state occupation probabilities (2-OPs). Mazziotti has reported [10, 15] necessary and sufficient conditions on the 2-OPs that arise from the positivity conditions on the q-OPs, i.e. the  $p_{11 \dots 1}(i_1, i_2, \dots, i_q) = \sum_{\mathbf{n}} C_{\mathbf{n}}^2 \nu_{i_1, \mathbf{n}} \nu_{i_2, \mathbf{n}} \cdots \nu_{i_q, \mathbf{n}}$ , at any order  $2 \leq q \leq N$ . These conditions derive from the positivity conditions [15] on the diagonal elements of the 2-DM [16]. Limiting ourselves to the (2, 2) and (2, 3) conditions, the following conditions for the 2-OPs hold:

$$\sup(p_1(i) + p_1(j) - 1, 0) \leq p_{11}(ij) \leq p_1(<) \quad [9]$$

$$\sup(p_1(i) + p_1(j) + p_1(k) - 1, 0) \leq p_{11}(ij) + p_{11}(ik) + p_{11}(jk) \quad [10]$$

$p_1(<)$  is the lesser of  $p_1(i)$  and  $p_1(j)$ . In addition the sum rule

$$\sum_{j(\neq i)} p_{11}(ij) = (N-1)p_1(i) \quad [11]$$

must be satisfied. Conditions **9–11** were first established by Weinhold and Bright Wilson [17]. They are necessary but not sufficient conditions for  $N$ -representability [7, 10, 16, 18]. Establishing a complete set of conditions is QMA-hard in  $N$  because the number of (2,  $q$ ) positivity conditions increases combinatorially with increasing  $q \leq N$ . Fortunately numerical calculations on atoms and molecules indicate that sufficiently accurate lower-bound ground-state energies often result by imposing (2,  $q$ )-positivity conditions with  $q \leq 3$  [9, 19]. This suggests that even in the most difficult situations, fermionic problems in atoms and molecules should require only a finite and small set of

positivity conditions. Here we shall limit ourselves to conditions **9–11**, as we found in our numerical calculations that they are sufficient to produce accurate lower-bounds. If higher-order conditions were found to be necessary, it would not be hard for us to add a few more.

The  $\pi^d$  of **7** contains only 2-OPs and products of 2 distinct NSOs; it has at most algebraic complexity  $\sim M^3$  deriving from condition **10**. Thus when only conditions **9–11** are imposed, the combinatorial complexity of the ground-state problem resides entirely in the  $\pi^{od}$  of **8**. We extract the sign  $s(ii'\mathbf{m})$  of the coefficient  $C_{ii'\mathbf{m}}$  in **8** and, relating its magnitude to the joint  $N$ -OP  $p_{11\dots 1}(ii'\mathbf{m}) \equiv C_{ii'\mathbf{m}}^2$ , we rewrite **8** as

$$\begin{aligned} \pi^{od}(x'_1 x'_2; x_1 x_2) &= \sum_{\substack{i < i' \neq j < j' \\ i, i', j, j' \notin \mathbf{m}}} s(ii'\mathbf{m})s(jj'\mathbf{m}) \\ & p_{11\dots 1}^{1/2}(ii'\mathbf{m})p_{11\dots 1}^{1/2}(jj'\mathbf{m}) \\ & (\psi_i(x'_1)\psi_{i'}(x'_2) - \psi_{i'}(x'_1)\psi_i(x'_2)) \\ & (\psi_j(x_1)\psi_{j'}(x_2) - \psi_{j'}(x_1)\psi_j(x_2)). \quad [12] \end{aligned}$$

We suppose that a variational approximation exists in which

$$s(ii'\mathbf{m})s(jj'\mathbf{m}) = s(ii')s(jj'), \forall \mathbf{m}. \quad [13]$$

This *sign conjecture* reduces the sign complexity to algebraic, scaling as  $M^2$ .  $\pi^{od}$  simplifies to

$$\begin{aligned} \pi^{od}(x'_1 x'_2; x_1 x_2) &= \sum_{i < i' \neq j < j'} s(ii')s(jj') \\ & \left[ \sum_{\substack{\mathbf{m} \\ i, i', j, j' \notin \mathbf{m}}} p_{11\dots 1}^{1/2}(ii'\mathbf{m})p_{11\dots 1}^{1/2}(jj'\mathbf{m}) \right] \\ & (\psi_i(x'_1)\psi_{i'}(x'_2) - \psi_{i'}(x'_1)\psi_i(x'_2)) \\ & (\psi_j(x_1)\psi_{j'}(x_2) - \psi_{j'}(x_1)\psi_j(x_2)). \quad [14] \end{aligned}$$

The quantities  $p_{11\dots 1}^{1/2}(ii'\mathbf{m})$  and  $p_{11\dots 1}^{1/2}(jj'\mathbf{m})$  are  $\mathbf{m}$ -th components of vectors  $\mathbf{p}_{11\dots 1}^{1/2}(ii')$  and  $\mathbf{p}_{11\dots 1}^{1/2}(jj')$ . The bracketed quantity in **14** is their scalar product. Express it as

$$\begin{aligned} \sum_{\substack{\mathbf{m} \\ i, i', j, j' \notin \mathbf{m}}} p_{11\dots 1}^{1/2}(ii'\mathbf{m})p_{11\dots 1}^{1/2}(jj'\mathbf{m}) &= \\ p_{1100}(ii'jj')p_{0011}(ii'jj')\xi(ii'jj'), & \quad [15] \end{aligned}$$

where  $p_{1100}(ii'jj')$  is the square magnitude of the vector  $\mathbf{p}_{11\dots 1}^{1/2}(ii')$  and  $p_{0011}(ii'jj')$  that of  $\mathbf{p}_{11\dots 1}^{1/2}(jj')$ .  $p_{1100}(ii'jj')$  is the probability that  $\psi_i$  and  $\psi_{i'}$  are occupied while  $\psi_j$  and  $\psi_{j'}$  are not:

$$\begin{aligned} p_{1100}(ii'jj') &= \sum_{\substack{\mathbf{m} \\ i, i', j, j' \notin \mathbf{m}}} p_{11\dots 1}(ii'\mathbf{m}) \\ &= \sum_{\mathbf{n}} C_{\mathbf{n}}^2 \nu_{i, \mathbf{n}} \nu_{i', \mathbf{n}} (1 - \nu_{j, \mathbf{n}}) (1 - \nu_{j', \mathbf{n}}), \end{aligned}$$

and the reverse is true for  $p_{0011}(ii'jj')$ .

The Schwarz inequality  $0 \leq \xi(ii'jj') \leq 1$  imposes bounds on  $\xi(ii'jj')$ , the cosine of the hyper-angle between the vectors. The upper bound  $\xi = 1$  is exact for  $N = 2$ . Substituting **15** into **14** yields

$$\begin{aligned} \pi^{od}(x'_1 x'_2; x_1 x_2) &= \sum_{i < i' \neq j < j'} s(ii')s(jj') \\ & [p_{1100}(ii'jj') p_{0011}(ii'jj')]^{1/2} \xi(ii'jj') \\ & (\psi_i(x'_1)\psi_{i'}(x'_2) - \psi_{i'}(x'_1)\psi_i(x'_2)) \\ & (\psi_j(x_1)\psi_{j'}(x_2) - \psi_{j'}(x_1)\psi_j(x_2)), \quad [16] \end{aligned}$$

in which only  $\xi(ii'jj')$  retains combinatorial complexity:

$$\xi(ii'jj') = \frac{\sum' p_{11\dots 1}^{1/2}(ii'\mathbf{m})p_{11\dots 1}^{1/2}(jj'\mathbf{m})}{(\sum' p_{11\dots 1}(ii'\mathbf{m}) \sum' p_{11\dots 1}(jj'\mathbf{m}))^{1/2}},$$

where the primed sums are over all  $\mathbf{m}$  with  $i, i', j, j' \notin \mathbf{m}$ .

Inserting 4-OPs like

$$p_{1111}(ii'kl) = \sum_{\mathbf{n}} C_{\mathbf{n}}^2 \nu_{ii', \mathbf{n}} \nu_{kl, \mathbf{n}}; \quad k < l \neq i, i', j, j'$$

in place of the  $N$ -OPs in  $\xi$  reduces the complexity of  $\pi^{od}$  to algebraic. The resulting approximation,

$$\xi(ii'jj') \approx \frac{\sum''_{k < l} p_{1111}^{1/2}(ii'kl)p_{1111}^{1/2}(jj'kl)}{(\sum''_{k < l} p_{1111}(ii'kl) \sum''_{k < l} p_{1111}(jj'kl))^{1/2}}, \quad [17]$$

is not variational, but obeys the 0,1 bounds of the Schwarz inequality. It is exact for  $N = 4$ , and scales as  $M^4$ . In **17** the doubly-primed sums are over the indices  $k < l$ , which must differ from  $i, i', j, j'$ . Bounds on the  $p_{1111}$  that are the generalizations of **9–11** for 3-OPs and 4-OPs can be formulated.

**The OP-NOFT energy functional.** The trial energy  $E[\Psi] = \langle \Psi | \hat{H} | \Psi \rangle$ , the expectation value of the Hamiltonian  $\hat{H}$ , is an explicit functional of the 1- and 2-DM:

$$E[\Psi] = E[\rho, \pi] = \text{tr} \{ \rho \hat{h} \} + \text{tr} \{ \pi \hat{w} \}.$$

Here  $\hat{h}$  is the single-particle kinetic-energy operator plus the external potential, and  $\hat{w}$  is the 2-electron Coulomb interaction.  $E[\rho, \pi]$  splits into two parts,  $E^d$  diagonal and  $E^{od}$  off-diagonal in the SD:

$$\begin{aligned} E &= E^d + E^{od} \\ E^d &= \text{tr} \{ \rho \hat{h} \} + \text{tr} \{ \pi^d \hat{w} \} \\ E^{od} &= \text{tr} \{ \pi^{od} \hat{w} \}. \quad [18] \end{aligned}$$

The Hartree-Fock wave function minimizes  $E^d$ ;  $\pi^{od}$  introduces electron correlation into  $E^{od}$ . The explicit forms of  $E^d$  and  $E^{od}$  follow from **5, 7**, and **16**:

$$E^d = \sum_i p_1(i) h_{ii} + \sum_{i < j} p_{11}(ij) [\mathcal{J}_{ij} - \mathcal{K}_{ij}], \quad [19]$$

where  $h_{ii} = \langle \psi_i | \hat{h} | \psi_i \rangle$ , and  $\mathcal{J}_{ij} = \langle \psi_i \psi_j | \hat{w} | \psi_j \psi_i \rangle$  and  $\mathcal{K}_{ij} = \langle \psi_i \psi_j | \hat{w} | \psi_i \psi_j \rangle$  are the Coulomb and exchange integrals, respectively. The second term on the rhs of **19** originates from  $\pi^d$ , the form of which is represented exactly in our theory. It contains only positive contributions and is essential; the integral relation connecting  $\pi$  and  $\rho$  depends only on  $\pi^d$  and guarantees that the  $E$  is self-interaction free.

$$\begin{aligned} E^{od} &= \sum_{i < i' \neq j < j'} s(ii')s(jj') p_{1100}^{1/2}(ii'jj') p_{0011}^{1/2}(ii'jj') \\ & \xi(ii'jj') [\mathcal{K}_{ii'jj'} - \mathcal{K}_{ii'j'j}], \quad [20] \end{aligned}$$

where  $\mathcal{K}_{ii'jj'} = \langle \psi_i \psi_{i'} | \hat{w} | \psi_j \psi_{j'} \rangle$ . **18 – 20** define the OP-NOFT energy functional within the PDC. Including the complexity of efficient evaluation of the matrix elements, it scales as  $M^5$  if the  $N$ -representability conditions for the 3- and 4-OPs can be limited to those deriving from the (3,  $q$ ) and (4,  $q$ ) positivity conditions with  $q \leq 4$ .

**Proof of the sign conjecture.** A variational sign approximation must be a statement about the sign  $s(ii'\mathbf{m})$  or  $s(jj'\mathbf{m})$  of each coefficient appearing in **8**. To prove **13**, we must find at least one statement in which the  $\mathbf{m}$ -dependences of  $s(ii'\mathbf{m})$  and  $s(jj'\mathbf{m})$  cancel. We have found two and present one here and one in section S1 of the SI. The former is valid for the general case of matrix elements  $[\mathcal{K}_{ii',jj'} - \mathcal{K}_{ii',j'j}]$  of arbitrary sign, the latter only for positive ones.

Assigning each index  $l$  in  $C_n$  a sign  $s(l)$  and taking  $s(\mathbf{n})$  as their product to form  $s(\mathbf{n}) = \prod_{l \in \mathbf{n}} s(l)$  is a variational approximation. Consequently  $s(ii'\mathbf{m}) = s(i)s(i')s(\mathbf{m})$ , and

$$s(ii'\mathbf{m})s(jj'\mathbf{m}) = s(i)s(i')s(j)s(j') \quad [21]$$

so that **13** is proved, with  $s(ii') = s(i)s(i')$ .

This approximation treats the form and phase, 0 or  $\pi$ , of each NSO as independent variables. The choice of signs for each index is not specified in **21**. Most energy minimization schemes start with random initial NSOs; similarly the choice of signs in **21** should be random, half positive and half negative. The number of the initial NSOs should be greater than the anticipated value of  $M$  to allow for unequal numbers of positive and negative signs of the active NSOs.

This complete factorization of  $s(\mathbf{n})$  and thence of  $s(ii')$  is a restrictive approximation. A variational approximation yielding unfactorized  $s(ii')$  could be more accurate. In Section S1 of the SI we have introduced a different variational approximation and rule for the signs which leads to **13** without factorization for positive matrix elements. Random assignment of signs in **13** and the sign rule of Section S1 yield identical results where tested, the significance of which is discussed there.

#### OP-NOFT-0

The SD's in **1** can be classified by their seniority, the number  $A$  of singly-occupied one-particle states they contain. For  $N$  even and for a global spin singlet ( $S = 0$ ) state, the  $N$ -particle Hilbert space divides into sectors of increasing even seniority starting with  $A = 0$ , where all SD's contain only doubly occupied states. For molecular systems CI expansions converge rapidly with seniority, and DOCI  $A = 0$  calculations describe static correlation rather well, as demonstrated in [20].

The PDC is equivalent to a restriction on seniorities in that seniorities differing only by 4 are allowed. Recent CI calculations for systems with even numbers of electrons showed that the seniority 2 sector largely decouples from the seniority 0 sector, supporting the accuracy of the PDC [20]. These considerations also apply to systems with an odd number of electrons, in which seniorities would be odd but still differ only by 4.

We now formulate OP-NOFT explicitly in the  $A = 0$  sector to illustrate further how an OP-NOFT functional is constructed and to prepare for numerical implementation; it becomes OP-NOFT-0, in which the PDC is automatically satisfied. Tracing out the spins, **5** becomes:

$$\rho(\mathbf{r}', \mathbf{r}) = 2 \sum_k p_1(k) \phi_k(\mathbf{r}') \phi_k(\mathbf{r}). \quad [22]$$

$k$  now labels  $M (> N/2)$  active doubly-occupied NO states, and the following conditions hold:

$$0 \leq p_1(k) \leq 1 \text{ and } 2 \sum_k p_1(k) = N. \quad [23]$$

In **22** and **23**  $p_1(k)$  is the occupation number of either of the paired NSOs having the NO  $\phi_k$ .

In the 2-DM, double occupancy results in a major simplification of the structure of  $\pi^{od}$ . We make the orbital and spin components of the NSO indices explicit. They take the form  $is$ , with  $i$  now the orbital index and  $s = \pm$  the spin index. The only index pairs that can enter

$\pi^{od}$  in **16** are  $i+, i-$  and  $j+, j-$ . The only sets of two index pairs that can enter the rhs of **17** are  $i+, i-, k+, k-$  and  $j+, j-, k+, k-$ . The occupation numbers  $\nu_{i+}$  and  $\nu_{i-}$  are equal, with values 0 or 1, so that all 4-NSO OPs in **17** and **16** are identical to the corresponding spin independent 2-NO OPs, e.g.  $p_{1111}(i+, i-, k+, k-) = p_{11}(ik)$ . The signs in **16** depend on a single orbital index,  $s(i+, i-) = s(i)$ , and the  $\xi$  depend on two-orbital indices,  $\xi(i+, i-, j+, j-) = \xi(ij)$ . With these simplifications, the 2-DM of **7** and **16** becomes

$$\pi(\mathbf{r}'_1 \mathbf{r}'_2; \mathbf{r}_1 \mathbf{r}_2) = \pi^d(\mathbf{r}'_1 \mathbf{r}'_2; \mathbf{r}_1 \mathbf{r}_2) + \pi^{od}(\mathbf{r}'_1 \mathbf{r}'_2; \mathbf{r}_1 \mathbf{r}_2), \quad [24]$$

after tracing out the spins, with

$$\begin{aligned} \pi^d(\mathbf{r}'_1 \mathbf{r}'_2; \mathbf{r}_1 \mathbf{r}_2) &= 2 \sum_{ij} p_{11}(ij) \\ &\quad (2\phi_i(\mathbf{r}'_1)\phi_j(\mathbf{r}'_2)\phi_i(\mathbf{r}_1)\phi_j(\mathbf{r}_2) - \\ &\quad \phi_i(\mathbf{r}'_1)\phi_j(\mathbf{r}'_2)\phi_j(\mathbf{r}_1)\phi_i(\mathbf{r}_2)) \end{aligned} \quad [25]$$

$$\begin{aligned} \pi^{od}(\mathbf{r}'_1 \mathbf{r}'_2; \mathbf{r}_1 \mathbf{r}_2) &= 2 \sum_{i \neq j} s(i)s(j) \\ &\quad [p_{10}(ij)p_{01}(ij)]^{1/2} \xi(ij) \\ &\quad \phi_i(\mathbf{r}'_1)\phi_i(\mathbf{r}'_2)\phi_j(\mathbf{r}_1)\phi_j(\mathbf{r}_2). \end{aligned} \quad [26]$$

The sum in **25** includes the term  $i = j$ , for which  $p_{11}(ii) = p_1(i)$ , and  $\xi(ij)$  in **26** is now

$$\xi(ij) \approx \frac{\sum_{k(\neq i,j)} p_{11}^{1/2}(ik)p_{11}^{1/2}(jk)}{\left[ \sum_{k(\neq i,j)} p_{11}(ik) \sum_{k(\neq i,j)} p_{11}(jk) \right]^{1/2}}. \quad [27]$$

The one- and two-orbital OPs of OP-NOFT-0 lie within the same bounds as in the general case, **9–10**, and their sum rules become, respectively, **23** and

$$2 \sum_{j(\neq i)} p_{11}(ij) = (N - 2)p_1(i). \quad [28]$$

The  $\pi$  of **24** satisfies two important sum rules

$$\begin{aligned} \int d\mathbf{r}_2 \pi(\mathbf{r}\mathbf{r}_2; \mathbf{r}'\mathbf{r}_2) &= (N - 1)\rho(\mathbf{r}, \mathbf{r}') \\ \int d\mathbf{r}_1 d\mathbf{r}_2 \pi(\mathbf{r}_1\mathbf{r}_2; \mathbf{r}_1\mathbf{r}_2)w(r_{12}) &\geq 0. \end{aligned}$$

The OP-NOFT-0 form for  $\pi$ , **24–27**, is exact when  $N = 2$  with  $\xi = 1$ . It is equivalent to DOCI for  $N = 4$  if the signs are correct. We show numerically that this is the case for the sign choice of Section S1 for the Paldus  $H_4$  test, as reported in Section S5 of the SI. For all the  $H_4$  configurations studied, the OP-NOFT-0 correlation energy coincides with that of DOCI to numerical precision. That the signs are correct for  $H_2$  and  $H_4$  confirms the validity of the sign rule in those cases and suggests a broader utility. When  $N > 4$ , the  $\xi$ -approximation of **27** and the limitation to the (2,2) and (2,3) positivity conditions break the equivalence to DOCI.

The expectation value  $E = \langle \Psi | \hat{H} | \Psi \rangle$  becomes:

$$\begin{aligned} E &= 2 \sum_i p_1(i) \langle \phi_i | \hat{h} | \phi_i \rangle + \\ &\quad \sum_{ij} p_{11}(ij) (2J_{ij} - K_{ij}) + \\ &\quad \sum_{i \neq j} s(i)s(j) p_{10}^{1/2}(ij) p_{01}^{1/2}(ij) \xi(ij) K_{ij}, \end{aligned} \quad [29]$$

where  $J_{ij}$  and  $K_{ij}$  are positive Hartree and exchange integrals defined in terms of the NOs,  $J_{ij} = \langle \phi_i \phi_i | \hat{w} | \phi_j \phi_j \rangle$  and  $K_{ij} =$

$\langle \phi_i \phi_j | \hat{w} | \phi_j \phi_i \rangle$ . With 27 for  $\xi(ij)$ ,  $E$  in 29 is a functional of the NOs and the 1- and 2-state OPs. Kollmar introduced a similar J-K functional but simplified the 2-DM [21]. The signs are chosen by a sign rule and are not variables.  $p_{10}$  is related to  $p_{11}$  and  $p_1$  by  $p_1(i) = p_{11}(ij) + p_{10}(ij)$  and is eliminated from the functional. Each sum in the denominator of  $\xi(ij)$  in 27 is simplified by the sum rule of 28 to, e.g.,

$$\sum_{k(\neq i, j)} p_{11}(ik) = \frac{1}{2}(N-2)p_1(i) - p_{11}(ij).$$

As stated above, we assume that the (2, 2) and (2, 3) positivity conditions are sufficient in practice. Under this circumstance, the infimum of  $E$  with respect to the NOs and the OPs, subject to the constraints 23 and 9, 10, 28, yields a variational approximation to the ground-state energy, apart from the  $\xi$ -approximation 27, for  $N > 4$ .

29 is a generalization of the NOFT formulations of 1-DM functional theories, which require only 1-state OPs. The extra complexity from 2-state OPs and implicit 4-state OPs is more than compensated by the substantial gain in accuracy it makes possible. The computational cost of calculating  $E$  from 29 scales like Hartree-Fock energy-functional minimization with a greater prefactor ( $M^3$  vs  $(N/2)^3$ ) due to fractional occupation of NOs.

### Numerical results for simple molecular systems

To test OP-NOFT-0, we studied several diatomic molecules, the Paldus  $H_4$  test, and linear chains of H atoms with open boundary conditions. We included all electrons (core and valence) and expanded the NOs in the Gaussian 6-31G\*\*, STO-6G, and cc-pVTZ bases. The constrained minimization was performed by damped Car-Parrinello dynamics [22], as detailed in Section S2 of the SI.

We started the minimization from NOs and OPs obeying the constraints but otherwise random. The signs were taken from the sign rule of the Table S2 in Sec. S1 of the SI. They were kept fixed during optimization.

At convergence, the *active* subset of NOs had  $p_1 \geq 10^{-3}$ . The remaining NOs contributed negligibly to the energy. The same active NOs and signs were found for several test cases starting instead from a sufficiently large set of random NOs, half with positive and half with negative signs<sup>2</sup>. It is significant for the rule of 21 for arbitrary matrix-element signs that for the systems tested, the Brillouin-Wigner perturbation-theory based rule of Table S1, and the alternative of random initial assignment of signs to pairs yield the same results for positive matrix elements. The procedure of 21 also yields half positive and half negative signs for the pairs when signs are assigned randomly to the individual NOs with no reference to the matrix-element signs.

We report the dissociation energy curves of the dimers  $H_2$ , LiH and HF in Figs. S1, S2, and S3 in Section S3 of the SI. We performed restricted Hartree-Fock, DFT (PBE [24] and/or PBE0 [25]), CASSCF, and CCSD(T) calculations with the same basis. For  $H_2$ , our functional depends only on 1-state OPs and the signs  $s(i)$ ; it reduces to the exact expression of Löwdin and Shull [26]. When the  $s(i)$  are chosen from Table S2 of the SI, the OP-NOFT-0 dissociation energy curve coincides with that of CASSCF at all interatomic separations, implying that this sign rule is exact for  $H_2$ . Even in a system as simple as  $H_2$ , spin-restricted Hartree-Fock and DFT fail badly at dissociation because these single-reference theories cannot recover the Heitler-London form of the wavefunction.

OP-NOFT-0 becomes identical to DOCI for  $N = 4$  when the sign choice is correct. The conditions 9, 10, and 28 simplify in this case as discussed in Section S4 of the SI. Expression 27 for  $\xi$  is exact, but the  $A = 0$  restriction is not. We performed the  $H_4$  Paldus test using a minimal 1s basis set for OP-NOFT-0, DOCI, and FCI. The DOCI/OP-NOFT-0 equivalence and the accuracy of the OP-NOFT-0 signs are confirmed by the results reported in Section S5. While DOCI only captures about 25-90% of the configuration-dependent




Fig. 1. Symmetric dissociation curve of a linear  $H_8$  chain. The squares indicate one half of the energy of a  $H_{16}$  chain (black square: Hartree-Fock energy; green squares: OP-NOFT-0 energy). All Hartree-Fock results are from spin-restricted calculations.

correlation energy in the Paldus test, the OP-NOFT-0 dissociation curve of LiH almost coincides with CASSCF in Fig. S2, which indicates that its 1s-electrons are nearly inert so that higher seniorities contribute negligibly to its ground-state energy.

HF, a 10-electron system, provides the first complete test of the relation of OP-NOFT-0 to DOCI. The energies obtained with the basis set 6-31G\*\* are shown in Fig. S3 of the SI. OP-NOFT-0 and DOCI are above both CASSCF and CCSD(T) though OP-NOFT-0 lies below DOCI because, while size consistent, the  $\xi$ -approximation is non-variational here. The OP-NOFT-0 signs are correct. The results are discussed further in Section S3 of the SI.

Linear H chains are relatively simple systems whose energy surfaces present a serious challenge for single reference methods. Fig. 1 shows symmetric dissociation energy curves of  $H_8$ . OP-NOFT-0 provides a consistent description of the energy close to and everywhere




Fig. 2. Dissociation curve of the  $N_2$  molecule.

<sup>2</sup>For  $H_2$  the sign rule ( $s(i \leq N) = +1$  and  $s(i > N) = -1$ ) holds near equilibrium, but a more complex pattern emerges at large separation where additional positive signs are needed for the van der Waals tail of the interaction potential [23]. In principle, these positive signs could be obtained with our minimization procedure, but their effect is beyond the accuracy of the present calculations

<sup>3</sup>We do not give the CASSCF energies in this case, as the dimension of the active subspace would make these calculations very expensive.

above the CASSCF reference. The breakdown of CCSD(T) at large separations is caused by its single-reference character. The deviation of OP-NOFT-0 from CASSCF should be attributed mainly to the restriction to the  $A = 0$  sector, a conclusion supported by the seniority-restricted CI calculations of Ref. [20]. Close comparison with those calculations is not entirely straightforward, as Ref. [20] used the slightly smaller 6-31G basis and a fixed, symmetric or broken symmetry, molecular orbital (MO) basis, whereas we used self-consistent NOs.

The OP-NOFT-0 1-DM displays the entanglement due to correlation through variation of the occupation numbers and the Von Neumann entanglement entropy with interatomic separation shown in Fig. S5 in Section S6 of the SI. The increase of entanglement entropy with separation signals a dramatic increase of correlation corresponding to multi-reference character. The OP-NOFT-0 2-DM gives access to electron pair correlations.

To test the dependence of the accuracy of OP-NOFT-0 on electron number, we studied the symmetric dissociation of  $H_{16}$ .<sup>3</sup> Results for the energy of  $H_{16}$  divided by 2 are shown as squares in Fig. 1. OP-NOFT-0 works equally well for this longer chain. The total energy at dissociation is twice that of  $H_8$ , and the slightly increased binding energy per atom at equilibrium arises from an increase in the correlation energy, as expected from more effective screening in the larger system.

The  $N_2$  molecule is a severe test because of its triple bond. OP-NOFT-0, DOCI, and FCI results obtained with the minimal basis set STO-6G are compared in Fig. 2. They are similar to those for HF, with OP-NOFT-0 and DOCI both above FCI but OP-NOFT-0 below DOCI. The accuracy of our DOCI results was improved by use of the optimized OP-NOFT-0 NOs for the DOCI basis.

It is interesting to note that in all the systems studied, the positivity condition (2, 2) was found to be sufficient at near equilibrium up to intermediate separations dominated by dynamic correlation because the (2, 3) condition was automatically satisfied there. Moreover, only in the case of  $H_8$ ,  $H_{16}$  and  $N_2$  at large separations did inclusion of the

(2, 3) positivity condition turn out to be essential to prevent runaway from the ground-state solution.

## Discussion

We have introduced a new method for correlated electronic-structure calculations, OP-NOFT, that scales algebraically. Its DOCI-like simplification, OP-NOFT-0, scales favorably with system size, with Hartree-Fock energy-minimization scaling. The close correspondence of the energies calculated via OP-NOFT-0 with DOCI calculations support the accuracy of limiting the positivity conditions to **9–10** and also of the  $\xi$ -approximation **27**. OP-NOFT-0 is restricted to the  $A = 0$  sector of the Hilbert space. It provides an accurate description of single-bond breaking and is a considerable improvement over single-reference methods in all cases studied.

Adding the computation of interatomic forces to the OP-NOFT-0 energy-minimization methodology would make possible the use of the theory for structural optimization and ab-initio molecular dynamics [22].

From the practical point of view, minimization of the functional **29** is significantly more laborious than minimization of the Hartree-Fock or the DFT functional. We attribute this difficulty to the need to include in **29** small occupation numbers. In damped dynamics minimization the forces acting on the corresponding NOs are thus very weak compared to the forces acting on the NOs with occupation numbers close to 1, slowing down considerably the entire procedure. This difficulty is common to all NO-based methods including those based on the 1-DM. Solving it is essential to making OP-NOFT methods applicable in practice. Care must be taken to avoid spurious minima, as in other nonlinear optimization problems.

**ACKNOWLEDGMENTS.** The authors acknowledge illuminating discussions with Paul Ayers and Kieron Burke. Refs. [13] and [17] were brought to the authors' attention by Paul Ayers. The authors further wish to thank J. E. Moussa for important comments. M.H.C. and R.C. acknowledge support from the DOE under grant DE-FG02-05ER46201.

- Hohenberg P, Kohn W (1964) Inhomogeneous Electron Gas. *Physical Review* 136:B864–B871.
- Kohn W, Sham LJ (1965) Self-Consistent Equations Including Exchange and Correlation Effects. *Physical Review* 140:A1133–A1138.
- Gilbert TL (1975) Hohenberg-Kohn theorem for nonlocal external potentials. *Physical Review B* 12:2111–2120.
- Gritsenko O, Pernal K, Baerends EJ (2005) An improved density matrix functional by physically motivated repulsive corrections. *The Journal of chemical physics* 122:204102.
- Lathiotakis NN, Marques MAL (2008) Benchmark calculations for reduced density-matrix functional theory. *The Journal of chemical physics* 128:184103.
- Coleman AJ (1963) Structure of Fermion Density Matrices. *Reviews of Modern Physics* 35:668–686.
- Garrod C, Percus JK (1964) Reduction of the N-Particle Variational Problem. *Journal of Mathematical Physics* 5:1756.
- Percus J (2013) On the trail of the 2-body reduced density matrix. *Computational and Theoretical Chemistry* 1003:2–7.
- Zhao Z, Braams BJ, Fukuda M, Overton ML, Percus JK (2004) The reduced density matrix method for electronic structure calculations and the role of three-index representability conditions. *The Journal of chemical physics* 120:2095 – 2104.
- Mazziotti DA (2012) Structure of Fermionic Density Matrices: Complete N-Representability Conditions. *Physical Review Letters* 108:263002.
- Nakata M, et al. (2001) Variational calculations of fermion second-order reduced density matrices by semidefinite programming algorithm. *The Journal of Chemical Physics* 114:8282.
- Gidofalvi G, Mazziotti DA (2008) Active-space two-electron reduced-density-matrix method: complete active-space calculations without diagonalization of the N-electron Hamiltonian. *The Journal of chemical physics* 129:134108.
- Weinhold F, Bright Wilson E (1967) Reduced Density Matrices of Atoms and Molecules. I. The 2 Matrix of Double-Occupancy, Configuration-Interaction Wavefunctions for Singlet States. *The Journal of Chemical Physics* 46:2752.
- Jankowski K, Paldus J (1980) Applicability of coupled-pair theories to quasidegenerate electronic states: A model study. *International Journal of Quantum Chemistry* 18:1243–1269.
- Mazziotti DA (2012) Significant conditions for the two-electron reduced density matrix from the constructive solution of N representability. *Physical Review A* 85:062507.
- Ayers PW, Davidson ER (2007) Linear Inequalities for Diagonal Elements of Density Matrices. *Advances in Chemical Physics* 134:443 – 483.
- Weinhold F, Bright Wilson E (1967) Reduced Density Matrices of Atoms and Molecules. II. On the N-Representability Problem. *The Journal of Chemical Physics* 47:2298.
- Davidson ER (1969) Linear Inequalities for Density Matrices. *Journal of Mathematical Physics* 10:725.
- Mazziotti DA (2012) Two-electron reduced density matrix as the basic variable in many-electron quantum chemistry and physics. *Chemical reviews* 112:244–62.
- Bytautas L, Henderson TM, Jiménez-Hoyos CA, Ellis JK, Scuseria GE (2011) Seniority and orbital symmetry as tools for establishing a full configuration interaction hierarchy. *The Journal of chemical physics* 135:044119.
- Kollmar C (2004) The "JK-only" approximation in density matrix functional and wave function theory. *The Journal of chemical physics* 121:11581–6.
- Car R, Parrinello M (1985) Unified Approach for Molecular Dynamics and Density-Functional Theory. *Physical Review Letters* 55:2471–2474.
- Sheng XW, Mentel AM, Gritsenko OV, Baerends EJ (2013) A natural orbital analysis of the long range behavior of chemical bonding and van der Waals interaction in singlet H2: The issue of zero natural orbital occupation numbers. *The Journal of Chemical Physics* 138:164105.
- Perdew JP, Burke K, Ernzerhof M (1996) Generalized Gradient Approximation Made Simple. *Physical Review Letters* 77:3865–3868.
- Adamo C, Barone V (1999) Toward reliable density functional methods without adjustable parameters: The PBE0 model. *The Journal of Chemical Physics* 110:6158.
- Löwdin PO, Shull H (1956) Natural Orbitals in the Quantum Theory of Two-Electron Systems. *Physical Review* 101:1730–1739.