

## Heterogeneous Reciprocal Graphical Models

Yang Ni<sup>1,\*</sup>, Peter Müller<sup>2</sup>, Yitan Zhu<sup>3</sup>, and Yuan Ji<sup>3,4</sup>

<sup>1</sup>Department of Statistics and Data Sciences, The University of Texas at Austin

<sup>2</sup>Department of Mathematics, The University of Texas at Austin

<sup>3</sup>Program for Computational Genomics & Medicine, NorthShore University HealthSystem

<sup>4</sup>Department of Public Health Sciences, The University of Chicago

\**email*: yangni87@gmail.com

**SUMMARY:** We develop novel hierarchical reciprocal graphical models to infer gene networks from heterogeneous data. In the case of data that can be naturally divided into known groups, we propose to connect graphs by introducing a hierarchical prior across group-specific graphs, including a correlation on edge strengths across graphs. Thresholding priors are applied to induce sparsity of the estimated networks. In the case of unknown groups, we cluster subjects into subpopulations and jointly estimate cluster-specific gene networks, again using similar hierarchical priors across clusters. We illustrate the proposed approach by simulation studies and three applications with multiplatform genomic data for multiple cancers.

**KEY WORDS:** Dirichlet-multinomial allocation; hierarchical model; model-based clustering; multiplatform genomic data; Pitman-Yor process; thresholding prior.

## 1. Introduction

We develop a heterogeneous reciprocal graphical model (hRGM) to infer gene networks in heterogeneous populations. Traditional graphical model approaches (Dempster, 1972; Lauritzen, 1996; Meinshausen and Bühlmann, 2006; Whittaker, 2009; Wang and West, 2009; Dobra et al., 2011; Green and Thomas, 2013; Wang et al., 2013; Peterson et al., 2015) assume i.i.d. sampling. However, many applications to inference for biomedical data, including the applications in this paper, include highly heterogeneous populations, and understanding and characterizing such heterogeneity is an important inference goal. We therefore propose an approach that admits statistical inference for potentially heterogeneous gene regulatory relationships across *known as well as unknown* groups/subpopulations. In particular, for the case of known groups, we model the related graphs under a Bayesian hierarchical model framework and allow the information to be shared across different groups. Borrowing of strength is implemented for the graph structure as well as for the edge strengths. For the case of unknown groups, we propose to cluster the subjects into subpopulations with meaningfully different graphs, that is, with group-specific graphs that differ in ways that allow biologically meaningful interpretation. Importantly, we implement clustering based on differences in the networks, in contrast to earlier proposed clustering methods (Dahl, 2006; Quintana, 2006; Lau and Green, 2007; Rodriguez et al., 2008; Müller et al., 2011; Lijoi et al., 2014), which are mostly based on cluster-specific mean/location.

Our work is motivated by three cancer genomic applications. In the first two applications, we construct gene networks for different cancer types. Recent pan-cancer studies (Hoadley et al., 2014) find both differences and commonalities across cancers. Traditional methods that assume one homogeneous population are not suitable in this case. Novel statistical methods accounting for data heterogeneity as well as adaptively borrowing strength across subtypes are needed. Our third application finds motivation in breast cancer which is molec-

ularly heterogeneous. Current classification systems based on three biomarkers are argued to be suboptimal as a means of directing therapeutic decisions for breast cancer patients (Di Leo et al., 2015). Improving the classification system is particularly important and yet challenging. This calls for better ways of clustering breast cancer patients.

Graphical models are commonly used to probabilistically model gene regulations. We build on a less commonly used class of graphical models, namely reciprocal graphical models (RGMs, Koster 1996). RGMs are a flexible class of models that allow for undirected edges, directed edges and directed cycles. These features are important for modeling biological feedback loops. RGMs strictly contain Markov random fields (MRFs) and directed acyclic graphs (DAGs) as special cases. However, RGMs are surprisingly underused in the biostatistics and bioinformatics literature. We prefer to use RGMs over other graphical models, because the inclusion of directed edges and possible cycles is critical for the three motivating applications.

Although graphical models for homogeneous data have been studied extensively in the literature, until recently only few approaches were proposed for heterogeneous data. When the groups are known, it is natural to “borrow strength” across different sample groups via a Bayesian hierarchical model for group-specific graphs. We provide a brief review of recent Bayesian methods and discuss the need and possibility for improvement. Mitra et al. (2016) consider MRFs for  $K = 2$  groups, identifying one group as the reference group and the other as the differential group. They assign a uniform prior to the reference graph and construct a mixture prior for the differential graph. Similarly, Oates et al. (2016) develop a multiple DAG approach for  $K > 2$ . They penalize the difference between graphs based on structural Hamming distance and utilize integer linear programming to find the posterior mode. Peterson et al. (2015) couple undirected MRF graphs by assuming an MRF prior on the edges. Computation can be challenging when  $K$  is moderate because of an analytically intractable normalization constant in the MRF model. One common limitation of these

methods is that they only borrow strength in the graph space, leaving the strength of selected edges (e.g. partial correlation) to be modeled/estimated independently. One exception is Yajima et al. (2015). They propose a multiple DAG which correlates the edge strength across groups. However, similarly to Mitra et al. (2016), they only consider  $K = 2$  groups and inference depends on the choice of reference/differential groups. Generalization to  $K > 2$  is not straightforward. Several non-Bayesian methods (Guo et al., 2011; Danaher et al., 2014; Mohan et al., 2014; Lee and Liu, 2015; Ma and Michailidis, 2016) have also been developed to learn MRFs across multiple groups.

There are few approaches for unknown groups. Rodriguez et al. (2011) propose a Dirichlet process mixture of MRFs to simultaneously cluster samples into homogeneous groups and infer undirected relationships between variables for each group. Similarly, Ickstadt et al. (2011) propose a multinomial-Dirichlet-Poisson mixture of DAGs to study directed relationships within each cluster. Mukherjee and Rodriguez (2016) recently develop a GPU-based stochastic search algorithm to improve the computation in Rodriguez et al. (2011).

In this article, we propose a hierarchical extension of RGMs to heterogeneous RGMs (hRGMs) as a model for both cases, known and unknown groups. When groups are known, the hRGM borrows strength across groups for inference on the group-specific graphs as well as the strength of the included edges. For unknown groups, the hRGM clusters a heterogeneous population into homogeneous subpopulations and allows each cluster to have its own network.

## 2. Reciprocal graphs

A graph  $\mathcal{G} = (V, E)$  consists of a set of vertices  $V = \{1, \dots, p\}$  usually representing a set of random variables and a set of edges  $E$  connecting these vertices. A reciprocal graph admits both directed edges  $i \rightarrow j$  and undirected edges  $i - j$ . Moreover, it explicitly allows for directed cycles, which is useful for modeling feedback mechanisms, which are a common motif in gene networks. Markov properties (i.e. conditional independence relationships) can

be read off from the RGM through moralization. Statistically, RGMs are a strictly larger class of probability models than MRFs and DAGs in terms of conditional independence (Koster, 1996). We provide a brief review of reciprocal graphs in Web Appendix A.

RGMs are only identifiable up to a Markov equivalence class in which all RGMs encode the same conditional independence relationships; see a series of work by Richardson (1996a,b, 1997) for characterization and identification of Markov equivalence class of cyclic graphs. Beyond this interventional or time-course data or other external information are needed to learn the structure of RGMs and determine the directionality of the edges. Recently, several approaches (Cai et al., 2013; Zhang and Kim, 2014; Ni et al., 2016) have proposed to infer directionality with observational data by integrating multi-platform data and fixing some edge directions based on biological knowledge. In this paper, we follow the approach of Ni et al. (2016) to include external information about known directional edges, by exploiting the central dogma of molecular biology, implying that DNA methylation and DNA copy number can affect mRNA gene expression, but not vice versa. Based on this consideration, we integrate DNA copy number and DNA methylation with mRNA gene expression and fix the direction of potential edges between DNA measurements and mRNA gene expressions. For illustration, let vertices  $\alpha$  and  $\beta$  in Figure 1 denote mRNA gene expressions. With observational data, we can not differentiate between graphs 1(a), 1(b), 1(c) and 1(d) because they all define the same conditional independence relationships between vertices  $\alpha$  and  $\beta$  (i.e. none). The identifiability issue can be resolved by incorporating additional information. Let vertices  $\gamma$  and  $\delta$  represent copy number or methylation of the corresponding genes. We introduce fixed directed edges  $\gamma \rightarrow \alpha$  and  $\delta \rightarrow \beta$ . Then the relationship between vertices  $\alpha$  and  $\beta$  are identifiable because each of graphs 1(e), 1(f), 1(g) and 1(h) encode different conditional independence relationships. A theoretical justification for this framework is discussed in Logsdon and Mezey (2010) and Oates et al. (2016).

[Figure 1 about here.]

We briefly review the mapping between RGMs and simultaneous equation models (SEMs). Let  $\mathbf{y}_i \in \mathbb{R}^p$  denote gene expressions for  $p$  genes and  $\mathbf{x}_i \in \mathbb{R}^q$  denote  $q$  DNA level measurements (copy number and methylation in our case study) for subject  $i = 1, \dots, n$ . Consider  $(\mathbf{y}_i, \mathbf{x}_i)$  to be jointly normal and satisfy the following SEM

$$\mathbf{y}_i = \mathbf{A}\mathbf{y}_i + \mathbf{B}\mathbf{x}_i + \mathbf{e}_i \quad (1)$$

where  $\mathbf{A} = (a_{jj'}) \in \mathbb{R}^{p \times p}$  with zeros on the diagonal,  $\mathbf{B} = (b_{jj'}) \in \mathbb{R}^{p \times q}$ ,  $\mathbf{e}_i \sim N_p(0, \boldsymbol{\Sigma})$  with diagonal covariance matrix  $\boldsymbol{\Sigma} = \text{diag}(\sigma_1, \dots, \sigma_p)$  and  $\mathbf{e}_i$ . The linearity and normality assumptions of the SEM in (1) can be relaxed by using, for example, the approaches in Mooij et al. (2011) and Lacerda et al. (2012). Assuming that  $\mathbf{I} - \mathbf{A}$  is invertible, model (1) can be written as

$$\mathbf{y}_i | \mathbf{x}_i \sim N_p \left\{ (\mathbf{I} - \mathbf{A})^{-1} \mathbf{B} \mathbf{x}_i, (\mathbf{I} - \mathbf{A})^{-1} \boldsymbol{\Sigma} (\mathbf{I} - \mathbf{A})^{-T} \right\}. \quad (2)$$

We define an RGM,  $\mathcal{G} = (V, E)$ , associated with the SEM (1) by letting vertices  $V = \{1, \dots, p + q\}$  denote the variables in  $(\mathbf{y}_i, \mathbf{x}_i)$ . That is, the variables corresponding to  $\mathbf{y}_i$  are indexed  $1, \dots, p$ , and the variables corresponding to  $\mathbf{x}_i$  are indexed  $p + 1, \dots, p + q$ . We allow directed edges terminating in  $y$ -vertices,  $j = 1, \dots, p$ , and undirected edges between  $x$ -vertices. Specifically, we draw a directed edge  $j' \rightarrow j$  from a  $y$ -vertex  $j' = 1, \dots, p$ , to another  $y$ -vertex  $j = 1, \dots, p$  if  $a_{jj'} \neq 0$ . And we draw directed edges  $h \rightarrow j$  from  $x$ -vertices  $h = p + 1, \dots, p + q$  to  $y$ -vertices if  $b_{j,h-p} \neq 0$ . Finally, we draw undirected edges  $h - h'$  between all  $x$ -vertices,  $h = p + 1, \dots, p + q$  and  $h' = p + 1, \dots, p + q$ ,  $h \neq h'$ . If needed also the inclusion of undirected edges between  $x$ -vertices could be explicitly modeled. However, we do not pursue this direction since the focus of this paper is on learning gene regulations (i.e. directed edges between  $y$ -vertices).

The distribution of  $(\mathbf{y}_i, \mathbf{x}_i)$  is global Markov with respect to the resulting RGM  $\mathcal{G}$  (Koster,

1996), i.e. the presence and absence of edges encode the conditional independence relationships of the vertices. The construction defines a mapping between SEMs of the type (1) and RGMs. In this paper,  $\mathbf{B}$  is block diagonal because copy number and methylation of gene  $j$ , in principle, only affect the expression of gene  $j$ .

We remark that SEM/RGM also allows for potential causal interpretation (Koster, 1996; Moerkerke et al., 2015): the presence of directed edges indicates possible direct causal effects whereas the absence of directed edges suggests exclusion of such possibility. However, the directed edges and cycles should not be over-interpreted beyond the scope of SEM without further causal assumptions. We refer to Robins (1999); Pearl (2003); Goetgeluk et al. (2008) for detailed discussion on the connection among causal inference, graphical models and SEMs.

### 3. hRGMs with known groups

#### 3.1 Sampling model

Suppose the data arises from a heterogeneous population that is divided into  $K$  known groups. Let  $\mathbf{y}_k$  and  $\mathbf{x}_k$  be the  $n_k \times p$  and  $n_k \times q$  matrices of observations for group  $k$ ,  $k = 1, \dots, K$ , with  $n = \sum_{k=1}^K n_k$ . We assume that the samples are exchangeable within each group and the probability model for subject  $i$  in group  $k$  is given by

$$\mathbf{y}_{ki} \mid \mathbf{x}_{ki}, \mathbf{A}^{(k)}, \mathbf{B}^{(k)}, \boldsymbol{\Sigma}^{(k)} \sim \text{N}_p \left\{ (\mathbf{I} - \mathbf{A}^{(k)})^{-1} \mathbf{B}^{(k)} \mathbf{x}_{ki}, (\mathbf{I} - \mathbf{A}^{(k)})^{-1} \boldsymbol{\Sigma}^{(k)} (\mathbf{I} - \mathbf{A}^{(k)})^{-T} \right\} \quad (3)$$

for  $k = 1, \dots, K$  and  $i = 1, \dots, n_k$  with  $\mathbf{A}^{(k)} = \left( a_{jj'}^{(k)} \right)$ ,  $\mathbf{B}^{(k)} = \left( b_{jj'}^{(k)} \right)$  and  $\boldsymbol{\Sigma}^{(k)} = \text{diag} \left( \sigma_1^{(k)}, \dots, \sigma_p^{(k)} \right)$ . Let  $\boldsymbol{\theta}^{(k)} = (\mathbf{A}^{(k)}, \mathbf{B}^{(k)}, \boldsymbol{\Sigma}^{(k)})$  denote the group-specific parameters. By the earlier described mapping between an SEM (1) and an RGM model, the  $\boldsymbol{\theta}^{(k)}$  define group-specific RGM models  $\mathcal{G}_k$ . The structural zeros of  $\mathbf{A}^{(k)}$  and  $\mathbf{B}^{(k)}$  correspond to missing edges in the RGM  $\mathcal{G}_k$  for group  $k$ .

Next we introduce a model feature to induce sparsity in  $\mathbf{A}^{(k)}$  and  $\mathbf{B}^{(k)}$ . We use a non-local

prior, defined as follows. First, we expand each entry as

$$a_{jj'}^{(k)} = \tilde{a}_{jj'}^{(k)} I(|\tilde{a}_{jj'}^{(k)}| > t_{jj'}) \quad \text{and} \quad b_{jj'}^{(k)} = \tilde{b}_{jj'}^{(k)} I(|\tilde{b}_{jj'}^{(k)}| > t_{jj'}) \quad (4)$$

where  $\tilde{a}_{jj'}^{(k)}$  and  $\tilde{b}_{jj'}^{(k)}$  are latent variables and the threshold parameters  $t_{jj'}$  define a minimum effect sizes of  $a_{jj'}^{(k)}$  and  $b_{jj'}^{(k)}$ . That is, we represent  $a_{jj'}^{(k)}$  and  $b_{jj'}^{(k)}$  by thresholding latent  $\tilde{a}_{jj'}^{(k)}$  and  $\tilde{b}_{jj'}^{(k)}$ . A theoretical motivation of the construction in (4) is discussed in Web Appendix B. The advantages of this thresholding mechanism over standard Bayesian variable selection framework, for example, a spike-and-slab prior will become clear when we discuss the priors that link multiple groups. Notice that  $t_{jj'}$  is edge-specific but shared across different groups, which is the key for inducing the desired dependence of graphs across similar groups and will be discussed in more detail in Section 3.2.

### 3.2 Priors linking multiple groups

We first introduce the prior for  $\tilde{a}_{jj'}^{(k)}$ . We assume multivariate normal priors on  $\tilde{\mathbf{a}}_{jj'} = \left( \tilde{a}_{jj'}^{(1)}, \dots, \tilde{a}_{jj'}^{(K)} \right)^T$ ,

$$\tilde{\mathbf{a}}_{jj'} \sim p(\tilde{\mathbf{a}}_{jj'}) = \text{N}(0, \tau_{jj'} \mathbf{\Omega}) \quad (5)$$

where  $\tau_{jj'}$  is an edge-specific variance component. The  $K \times K$  relatedness matrix  $\mathbf{\Omega} = [\Omega_{kk'}]$  links edge strength (effect sizes) and by the thresholding in (4) also edge inclusion across  $K$  different groups, which in turn also links the graphs across groups. The strength of the dependence between groups  $k, k'$  is characterized by correlation  $\Omega_{kk'} / \sqrt{\Omega_{kk} \Omega_{k'k'}}$ . When  $\Omega_{kk'}$  is significantly away from zero,  $\tilde{a}_{jj'}^{(k)}$  and  $\tilde{a}_{jj'}^{(k')}$  are likely to be of similar magnitude *a priori* and since the thresholding parameters  $t_{jj'}$  are shared across groups, there is a high probability that  $\tilde{a}_{jj'}^{(k)}$  and  $\tilde{a}_{jj'}^{(k')}$  are either both non-zero or both shrunk to zero. Therefore, graphs  $\mathcal{G}_k$  and  $\mathcal{G}_{k'}$  are more likely to share common edges. On the other hand, when  $\Omega_{kk'}$  is negligible (i.e. close to 0), groups  $k$  and  $k'$  are unrelated. We do not constrain  $\Omega_{kk'}$  to be non-negative, which is imposed by Peterson et al. (2015). This additional flexibility allows edge strength



to have different signs for different groups, which is desirable in estimating gene networks because potentially gene regulations can switch from activation to inactivation across groups (e.g. case vs control). Note that  $\tau_{jj'}$  and  $\mathbf{\Omega}$  are not identifiable because  $\tau_{jj'}\mathbf{\Omega} = c\tau_{jj'}\mathbf{\Omega}/c$  for any  $c > 0$ , which can be resolved by fixing  $\Omega_{11} = 1$ . The priors for  $\tilde{b}_{jj'}^{(k)}$  are defined similarly,

$$\tilde{\mathbf{b}}_{jj'} = \left( \tilde{b}_{jj'}^{(1)}, \dots, \tilde{b}_{jj'}^{(K)} \right)^T \sim N(0, \lambda_{jj'}\mathbf{\Omega}). \quad (6)$$

The priors in (5) and (6) share the same relatedness matrix  $\mathbf{\Omega}$  that describes pairwise similarity of the graphs for each group. The uncertainty of each edge strength is captured by the edge-specific variance components  $\tau_{jj'}$  and  $\lambda_{jj'}$ . We assume a simple uniform prior for  $t_{jj'} \sim \text{Unif}(0, b_t)$ . We remark that the induced prior for  $\mathbf{A}^{(k)}$  should be restricted to the cone of invertible  $\mathbf{I} - \mathbf{A}^{(k)}$ , which practically is not a constraint given that any random matrix is almost surely invertible (Rudelson, 2008).

### 3.3 Priors for the relatedness matrix and hyperparameters

We assign a conjugate inverse-Wishart prior for the relatedness matrix  $\mathbf{\Omega}$  subject to  $\Omega_{11} = 1$  which controls the relatedness of different groups as discussed in Section 3.2,  $\mathbf{\Omega}^* \sim \text{IW}(\nu, \mathbf{\Phi})$  and  $\mathbf{\Omega} = \mathbf{\Omega}^*/\Omega_{11}^*$ , with degrees of freedom  $\nu$  and a scale matrix  $\mathbf{\Phi}$ . Alternatively, one could introduce a hyper MRF graph  $\mathcal{H}$  and put a G-Wishart prior on  $\mathbf{\Omega}^{-1} \sim W_{\mathcal{H}}(\cdot, \cdot)$  if one wants to learn the conditional independence relationships between groups. Since in our case study, we only have  $K = 3$  groups and the correlations between groups are quite significant (as shown in Section 5.1), we are not pursuing this direction in this paper. We complete the model by assuming conjugate priors  $\sigma_j^{(k)} \sim \text{IG}(a_\sigma, b_\sigma)$ ,  $\tau_{jj'} \sim \text{IG}(a_\tau, b_\tau)$  and  $\lambda_{jj'} \sim \text{IG}(a_\lambda, b_\lambda)$ . For our simulations and case studies, we use noninformative hyperpriors. But if desired, informative priors can be used, for example, to introduce prior knowledge about a reference network.

#### 4. hRGMs with unknown groups

We introduce a clustering approach to split heterogeneous samples into (unknown) homogeneous groups. Under a Bayesian approach, the number  $K$  of clusters need not be fixed *a priori* and can be inferred from the data. We first introduce a latent cluster membership indicator  $s_i$  with  $s_i = k$  when subject  $i$  belongs to group  $k$ . Conditional on  $s_i = k$  the likelihood is the same as (3). For reference we restate it here, now with conditioning on  $s_i$ ,

$$\mathbf{y}_i \mid \mathbf{x}_i, \boldsymbol{\theta}^{(k)}, s_i = k \sim N_p \left\{ (\mathbf{I} - \mathbf{A}^{(k)})^{-1} \mathbf{B}^{(k)} \mathbf{x}_i, (\mathbf{I} - \mathbf{A}^{(k)})^{-1} \boldsymbol{\Sigma}^{(k)} (\mathbf{I} - \mathbf{A}^{(k)})^{-T} \right\}. \quad (7)$$

We use the same priors for cluster-specific parameters,  $\tilde{\mathbf{a}}_{jj'} \sim p(\tilde{\mathbf{a}}_{jj'}) = N(0, \tau_{jj'} \boldsymbol{\Omega})$ ,  $\tilde{\mathbf{b}}_{jj'} \sim N(0, \lambda_{jj'} \boldsymbol{\Omega})$  and  $\sigma_j^{(k)} \sim \text{IG}(a_\sigma, b_\sigma)$  as in Section 3. We complete the model with prior distributions for  $s_i$  and  $K$ . We use a Dirichlet-multinomial (DM) allocation model,

$$s_i \mid \boldsymbol{\pi}, K \sim \text{Multinomial}(1, \pi_1, \dots, \pi_K) \text{ and } \boldsymbol{\pi} \mid K \sim \text{Dir}(\eta, \dots, \eta),$$

and a geometric prior for  $K \sim \text{Geo}(\rho)$ . We refer to model (7) with a DM prior as the hRGM-DM. Alternatively, we consider nonparametric Bayesian priors that give rise to exchangeable random partitions such as the Poisson-Dirichlet process, also known as Pitman-Yor (PY) process priors (Pitman and Yor, 1997). A PY process induces a prior distribution on  $s_i$ 's and  $K$  which is characterized by a (modified) Chinese restaurant process  $\text{CRP}(\alpha, d)$  with total mass parameter  $\alpha$  and discount parameter  $d$ ,

$$p(s_i = k \mid s_1, \dots, s_{i-1}) \propto \begin{cases} n_k^{(-i)} - d & \text{for } k = 1, \dots, K^{(-i)} \\ \alpha + dK^{(-i)} & \text{for } k = K^{(-i)} + 1 \end{cases} \quad (8)$$

where  $n_k^{(-i)}$  and  $K^{(-i)}$  are the size of cluster  $k$  and the total number of clusters after removing the  $i$ th sample. The admissible values for  $(\alpha, d)$  are  $d \in [0, 1)$  with  $\alpha > -d$  or  $d < 0$  with  $\alpha = |md|$  for some integer  $m$ . The popular Dirichlet process (Ferguson, 1973; Blackwell and MacQueen, 1973) is a special case of a PY process when  $d = 0$ . The extra parameter  $d$  in the

PY process makes it more flexible than the Dirichlet process prior (De Blasi et al., 2015). We refer to model (7) with the PY process prior as the hRGM-PY. De Blasi et al. (2015) and Barrios et al. (2013) argue for the PY prior as more flexible prior for random partitions than DP prior and a parametric DM prior.

The choice of  $\mathbf{\Omega}$  is motivated by the following consideration. The goal is to divide subjects into subpopulations with distinct networks, rather than induce dependence between clusters. We therefore set the relatedness matrix  $\mathbf{\Omega} = \text{diag}(\Omega_1, \dots, \Omega_K)$  to be diagonal, i.e. networks are independent across clusters.

Implementing posterior inference under the hRGM-DM or the hRGM-PY model is straightforward by MCMC simulation. Details are provided in Web Appendix C, including how we report point estimates for networks.

## 5. Case studies

We validate inference in proposed hRGM with extensive simulation studies in Web Appendix D. In this section, we discuss application of the proposed inference to three TCGA genomic datasets.

### 5.1 *p53 pathway across three cancer subtypes*

A recent pan-cancer genomic study (Hoadley et al., 2014) has identified 11 major cancer subtypes based on molecular characterizations instead of tissue-of-origin. This provides independent information for clinical prognosis. Although the subtypes are correlated with tissue-of-origin, several distinct cancer types are classified into common subtypes. For example, head and neck squamous cell carcinoma (HNSC) and lung squamous cell carcinoma (LUSC) are classified into subtype “C2-Squamous-like” by molecular alterations including p53 whereas invasive breast cancer (BRCA) is identified as subtype “C3-BRCA/Luminal” by itself.

Using inference under the proposed hRGM, we explore related dependencies of gene net-

works in the same three cancer types, HNSC, LUSC and BRCA. That is, we construct a gene network for each cancer and borrow strength across cancer types adaptively, depending on how similar the cancers are. We expect the networks of HNSC and LUSC to have stronger association than those of HNSC and BRCA or of LUSC and BRCA since HNSC and LUSC belong to the same molecular subtype.

We use the software package TCGA-Assembler (Zhu et al., 2014) to retrieve mRNA gene expression, DNA copy number and DNA methylation for HNSC, LUSC and BRCA from The Cancer Genome Atlas (TCGA). We focus on genes that are mapped to the core members of the p53 pathway ( $p = 10$ ) which respond to stresses that can cause errors in DNA replication and cell division (Harris and Levine, 2005) and plays a critical role in all three cancers (Gasco et al., 2002; Leemans et al., 2011; Perez-Moreno et al., 2012). A more detailed description of these 10 genes appears in Web Appendix E. We use the same samples of HNSC, LUSC and luminal BRCA that were analyzed in Hoadley et al. (2014) and match genomic data from different platforms, including methylation and copy number variation corresponding to each gene. That is, we record  $q = 2p = 20$  DNA level variables. The resulting dataset includes  $p + q = 30$  variables for  $n_1 = 298$  HNSC samples,  $n_2 = 100$  LUSC samples and  $n_3 = 432$  luminal BRCA samples. We run MCMC simulation for 100,000 iterations, discard the first 50% as burn-in and thin the chain to every 5th sample. MCMC diagnostics and model checking show no evidence for lack of practical convergence and lack of fit (see Web Appendix E for details).

Figure 2 shows point estimates for the gene networks, based on controlling posterior expected FDR (Newton et al., 2004; Müller et al., 2006) at 1%. The edges between genes and corresponding copy number and methylation are omitted for clarity. These associations are shown separately in Web Appendix E. Most gene expressions are associated with the corresponding copy number whereas only a few genes are correlated to their methylation.

In Figure 2, edges that are shared across subtypes are represented as solid lines. Differential edges are represented as dashed lines. Arrowheads denote stimulatory regulations, whereas small perpendicular bars indicate inhibitory regulations. The number of edges in each of three gene networks and the number of shared edges between each pair of networks are given in the upper triangular part of Table 1. Due to the molecular similarity of HNSC and LUSC, they share many edges: 10 out of 11 edges in the LUSC network are also found in the HNSC network. And as expected, BRCA shares fewer, but still a reasonable number of edges with HNSC and LUSC. In summary, inference under the proposed hRGM model recognizes difference in association between cancer types and borrows strength adaptively in a way that confirms the subtypes found by Hoadley et al. (2014). The estimated levels of adaptive borrowing of strength are reflected in the estimated relatedness matrix  $\hat{\Omega}$  (given in the lower triangular part of Table 1). The moderate to strong correlations suggest that the joint analysis under the hRGM is more appropriate than separate inference under separate RGMs. In fact, separate RGMs detect much fewer edges and obtain networks with almost no shared edges. The results are provided in Web Appendix E.

[Table 1 about here.]

A noticeable feature in all three estimated networks in Figure 2 is the central role of E2F1. E2F1 is an important transcription factor gene across cancer types which interacts with multiple genes in all networks. Such highly connected genes are known as hub genes and are often more involved in multiple regulatory activities than non-hub genes. In fact, E2F1 has a pivotal role in controlling cell cycle progression and induces apoptosis. It has been found that E2F1 is mutated in most, if not all, human tumors (Polager and Ginsberg, 2009). Our finding that E2F1 plays important roles in all networks is consistent with the fact that hub genes are more likely to be conserved across species, diseases and tumor (sub-)types (Casci, 2006). In addition, some edges that we find in our analysis are well studied in the biological

literature. For example, ATM phosphorylates and stabilizes E2F1 in response to various stresses including DNA damage. Our study confirms this positive regulatory relationship across all cancers.

[Figure 2 about here.]

## 5.2 KEGG colorectal cancer pathway cross two cancers

In this application, we investigate the genomic patterns of colon adenocarcinoma (COAD) and rectum adenocarcinoma (READ). COAD and READ are closely related cancers in terms of the copy number alteration, DNA methylation and gene expression patterns (TCGA, 2012). However, they are also known to exhibit substantial difference in genetic characterization (Frattini et al., 2004), the process of carcinogenesis (Kapiteijn et al., 2001) and chemosensitivity (Kornmann et al., 2014). We download DNA copy number, methylation and mRNA gene expression using TCGA-Assembler (Zhu et al., 2014) for genes that are mapped to the KEGG colorectal cancer pathway (<http://www.genome.jp/>). The dataset contains  $n_1 = 276$  COAD samples,  $n_2 = 91$  READ samples and totally 174 genomic measurements (58 in each of copy number, methylation and gene expression).

The full networks with 174 vertices for COAD and READ are shown in Figures 3(a) and 3(b), respectively. They share 77 edges which are represented by solid lines. Each cancer has moderate number of differential edges as well, indicated by dashed lines: COAD has 60 distinctive edges whereas READ has 27 unique edges.

For better visualization, we focus on the gene networks (i.e. without copy number and methylation) in Figures 3(c) and 3(d). We find in COAD, PIK3R1 inactivates GSK3B. The biological literature confirms this finding that PIK3R1 phosphorylates and activates AKT which in turn phosphorylates and inactivates GSK3B (Danielsen et al., 2015; Lin et al., 2015). Surprisingly, however, this link is missing in the READ gene network. This unexpected result deserves further experimental validation.

PIK3CG, a catalytic subunit of phosphatidylinositide 3-OH kinase, seems to play a critical role in both COAD and READ: it has 8 and 7 links to other genes, respectively. In fact, down-regulation of PIK3CG expression inhibits PI3K-AKT signaling pathway responsible for carcinogenesis and progression of colorectal cancers, which occurs frequently in primary colorectal cancers (Semba et al., 2002).

[Figure 3 about here.]

### 5.3 Clustering breast cancer subtypes

Breast cancer is known to be a highly heterogeneous disease. Based on three biomarkers (ER/PR/HER2), breast cancer is traditionally classified into three groups: luminal, HER2 and basal. However, this classification system is also known to be suboptimal and in need of improvement for better diagnostics and prognostics (Di Leo et al., 2015).

Instead of three biomarkers, we consider  $p = 10$  core genes of a critical pathway, RAS-MAPK, which transmits and amplifies signals involved in cell proliferation and cell death in breast cancer (Santen et al., 2002). These 10 genes form the key component of the RAS-MAPK pathway known as the MAPK cascade which is explained in detail in Web Appendix E. We use inference under the proposed hRGM-DM to find subtypes based on breast cancer data ( $n = 720$  samples) retrieved from TCGA as described in Section 5.1. Our goal is to simultaneously cluster patients and estimate gene networks for each cluster. We implement inference using MCMC posterior simulation, using the same setup as in Section 5.1. MCMC diagnostics and model checking show no evidence for lack of practical convergence and lack of fit (see Web Appendix E for details).

We identify two major clusters with 600 and 107 samples. To explore the clinical relevance of the reported clusters we evaluate Kaplan-Meier (KM) estimates of patients' survivals for the two clusters. These are shown in Figure 4(a). We find a substantial difference in median survival between the two clusters: a difference of 928 days. For comparison, we also apply

RLD (Rodriguez et al., 2011) and the K-means algorithm with  $K = 2$  to the same data including gene expression, copy number and methylation. Their KM estimators are shown in Figures 4(b) and 4(c). RLD finds one major cluster with 687 patients. The second largest cluster has only 22 patients. The difference of median survival is 120.5 days between the two clusters. The two clusters identified by K-means have 635 and 85 samples with median difference in survival of 399 days. In addition, we also compute KM estimates, shown in Figure 4(d), for the luminal/HER2 group versus basal group; the latter usually has much worse prognosis. The difference of median survival is 497 days.

To formally compare the clinical relevance in terms of significantly different overall survival for the reported clusters, we carried out log-rank tests (p-values shown in Figure 4) for the difference of the corresponding survival distributions. In fact, only hRGM-DM identifies clusters that have significantly different survival distributions. The hRGM-DM model detects the most distinctive clusters in terms of prognosis. Such clusters or subtypes may be useful in refining subtypes of breast cancer and developing new treatment options. In addition, we quantify the difference between clusters estimated from hRGM-DM, RLD and K-means by the adjusted Rand index and the adjusted variation of information (Meilă, 2003) in the Web Appendix E. Similarly to what the KM estimates suggest, the clusters obtained from each method are quite different.

[Figure 4 about here.]

In terms of the estimated network structure, we find that all genes are associated with their respective copy number for both clusters except for *SOS1* and *KRAS* in cluster 1 and *SOS1* and *MAPK3* in cluster 2. In addition, *NRAS* and *KRAS* in both clusters and *SOS2* and *BRAF* in cluster 2 are found to be associated with their methylation. Figure 5 shows the estimated gene networks for the two clusters when controlling posterior expected FDR at 1%. As



before, shared edges across both clusters are represented by solid lines and differential edges are shown as dashed lines. Arrowheads denote stimulatory regulations, whereas horizontal bars indicate inhibitory regulations. We find 27 edges for cluster 1 and 41 edges for cluster 2. The two networks share 18 edges but are otherwise quite different from each other. For example, the well-known cascade  $SOS1 \rightarrow KRAS \rightarrow MAPK1 \rightarrow MAP2K2$  (Santen et al., 2002) is found in cluster 2 but not in cluster 1. Furthermore, we find MAP2K2 inhibits SOS1 in cluster 2, which completes the negative feedback loop, a commonly observed motif in gene network (Krishna et al., 2006), as shown in Figure 5(c). Interestingly, it is recently discovered that MAP2K2 phosphorylates and inhibits SOS, therefore reducing MAP2K2 activation (Mendoza et al., 2011).

[Figure 5 about here.]

## 6. Discussion

We have developed the hRGM(-DM,-PY) model for inference on gene networks for heterogeneous samples. The hRGM connects graphs across known groups by introducing correlation on edge strength and edge inclusion. In the case of unknown groups, the hRGM together with a Dirichlet-multinomial or Pitman-Yor process prior for the cluster arrangement allows to learn the unknown groups and estimate group-specific networks at the same time. Bayesian model selection is implemented with a thresholding prior and used to obtain sparse networks. Simulation studies demonstrate the advantages of hRGM over a comparable approach with separate inference for group-specific networks. In the first application, we find both common and differential network structures for different cancer types. In the second application, we are able to identify clusters that differ significantly in network structures. The clinical significance of the discovered clusters is validated by significantly different cluster-specific survival functions. Inference provides important information for refining subtypes of breast

cancer and for developing new therapeutic options.

In this paper, we only consider continuous random variables and assume normal distributions. This is because the mapping between an RGM and an SEM is only valid for Gaussian distributions (Koster, 1996). Although we have empirically demonstrated the robustness of our method to the distribution misspecification, it would be desirable to extend the model to non-Gaussian data. For example, one may want to integrate data on single nucleotide polymorphisms (SNPs) and mutation status (MUs) into the gene network reconstruction. However, SNP and MU are recorded as discrete variables. One possible implementation is the use of latent Gaussian variables. In principle, RGMs can be embedded in a more general exponential family graphical model framework (Yang et al., 2015). However, without the mapping between RGM and SEM, computation is a major bottleneck for such extensions.

## 7. Supplementary Materials

Web Appendices, Tables, and Figures referenced in Sections 2, 4 and 5 and Matlab program implementing our methods, are available with this paper at the Biometrics website on Wiley Online Library.

## References

- Barrios, E., Lijoi, A., Nieto-Barajas, L. E., and Prünster, I. (2013). Modeling with normalized random measure mixture models. *Statistical Science* **28**, 313–334.
- Blackwell, D. and MacQueen, J. B. (1973). Ferguson distributions via Pólya urn schemes. *The Annals of Statistics* **1**, 353–355.
- Cai, X., Bazerque, J. A., and Giannakis, G. B. (2013). Inference of gene regulatory networks with sparse structural equation models exploiting genetic perturbations. *PLoS Comput Biol* **9**, e1003068.
- Casci, T. (2006). Network fundamentals, via hub genes. *Nature Reviews Genetics* **7**, 664–665.

- Dahl, D. B. (2006). Model-based clustering for expression data via a Dirichlet process mixture model. *Bayesian Inference for Gene Expression and Proteomics* pages 201–218.
- Danaher, P., Wang, P., and Witten, D. M. (2014). The joint graphical lasso for inverse covariance estimation across multiple classes. *J R Stat Soc B* **76**, 373–397.
- Danielsen, S. A., Eide, P. W., Nesbakken, A., Guren, T., Leithe, E., and Lothe, R. A. (2015). Portrait of the pi3k/akt pathway in colorectal cancer. *Biochimica et Biophysica Acta (BBA)-Reviews on Cancer* **1855**, 104–121.
- De Blasi, P., Favaro, S., Lijoi, A., Mena, R. H., Prünster, I., and Ruggiero, M. (2015). Are Gibbs-type priors the most natural generalization of the Dirichlet process? *IEEE Trans Pattern Anal Mach Intell* **37**, 212–229.
- Dempster, A. P. (1972). Covariance selection. *Biometrics* **28**, 157–175.
- Di Leo, A., Curigliano, G., Diéras, V., Malorni, L., Sotiriou, C., Swanton, C., Thompson, A., Tutt, A., and Piccart, M. (2015). New approaches for improving outcomes in breast cancer in Europe. *The Breast* **24**, 321–330.
- Dobra, A., Lenkoski, A., and Rodriguez, A. (2011). Bayesian inference for general gaussian graphical models with application to multivariate lattice data. *Journal of the American Statistical Association* **106**, 1418–1433.
- Ferguson, T. S. (1973). A Bayesian analysis of some nonparametric problems. *The Annals of Statistics* **1**, 209–230.
- Frattini, M., Balestra, D., Suardi, S., Oggionni, M., Alberici, P., Radice, P., Costa, A., Daidone, M. G., Leo, E., Pilotti, S., et al. (2004). Different genetic features associated with colon and rectal carcinogenesis. *Clinical Cancer Research* **10**, 4015–4021.
- Gasco, M., Shami, S., and Crook, T. (2002). The p53 pathway in breast cancer. *Breast Cancer Research* **4**, 70.
- Goetgeluk, S., Vansteelandt, S., and Goetghebeur, E. (2008). Estimation of controlled direct

- effects. *J R Stat Soc B* **70**, 1049–1066.
- Green, P. J. and Thomas, A. (2013). Sampling decomposable graphs using a Markov chain on junction trees. *Biometrika* **100**, 91–110.
- Guo, J., Levina, E., Michailidis, G., and Zhu, J. (2011). Joint estimation of multiple graphical models. *Biometrika* **98**, 1–15.
- Harris, S. L. and Levine, A. J. (2005). The p53 pathway: positive and negative feedback loops. *Oncogene* **24**, 2899–2908.
- Hoadley, K. A. et al. (2014). Multiplatform analysis of 12 cancer types reveals molecular classification within and across tissues of origin. *Cell* **158**, 929–944.
- Ickstadt, K., Bornkamp, B., Grzegorzczak, M., Wieczorek, J., Sheriff, M. R., Grecco, H. E., and Zamir, E. (2011). Nonparametric Bayesian networks. *Bayesian Statistics 9* **9**, 283.
- Kapiteijn, E., Liefers, G., Los, L., Klein Kranenbarg, E., Hermans, J., Tollenaar, R., Moriya, Y., Van De Velde, C., and Van Krieken, J. (2001). Mechanisms of oncogenesis in colon versus rectal cancer. *The Journal of Pathology* **195**, 171–178.
- Kornmann, M., Link, K.-H., and Formentini, A. (2014). Differences in colon and rectal cancer chemosensitivity. *Colorectal Cancer* **3**, 93–105.
- Koster, J. T. (1996). Markov properties of nonrecursive causal models. *The Annals of Statistics* **24**, 2148–2177.
- Krishna, S., Andersson, A. M., Semsey, S., and Sneppen, K. (2006). Structure and function of negative feedback loops at the interface of genetic and metabolic networks. *Nucleic Acids Research* **34**, 2455–2462.
- Lacerda, G., Spirtes, P. L., Ramsey, J., and Hoyer, P. O. (2012). Discovering cyclic causal models by independent components analysis. *arXiv preprint arXiv:1206.3273* .
- Lau, J. W. and Green, P. J. (2007). Bayesian model-based clustering procedures. *Journal of Computational and Graphical Statistics* **16**, 526–558.


- Lauritzen, S. L. (1996). *Graphical models*, volume 17. Oxford University Press, USA.
- Lee, W. and Liu, Y. (2015). Joint estimation of multiple precision matrices with common structures. *J Mach Learn Res* **16**, 1035–1062.
- Leemans, C. R., Braakhuis, B. J., and Brakenhoff, R. H. (2011). The molecular biology of head and neck cancer. *Nature Reviews Cancer* **11**, 9–22.
- Lijoi, A., Nipoti, B., and Prünster, I. (2014). Dependent mixture models: clustering and borrowing information. *Computational Statistics & Data Analysis* **71**, 417–433.
- Lin, Y., Yang, Z., Xu, A., Dong, P., Huang, Y., Liu, H., et al. (2015). Pik3r1 negatively regulates the epithelial-mesenchymal transition and stem-like phenotype of renal cancer cells through the akt/gsk3 $\beta$ /ctnnb1 signaling pathway. *Scientific Reports* **5**, 8997.
- Logsdon, B. A. and Mezey, J. (2010). Gene expression network reconstruction by convex feature selection when incorporating genetic perturbations. *PLOS Computational Biology* **6**, 1–13.
- Ma, J. and Michailidis, G. (2016). Joint structural estimation of multiple graphical models. *J Mach Learn Res* **17**, 1–48.
- Meilă, M. (2003). *Comparing Clusterings by the Variation of Information*, pages 173–187. Springer Berlin Heidelberg, Berlin, Heidelberg.
- Meinshausen, N. and Bühlmann, P. (2006). High-dimensional graphs and variable selection with the lasso. *The Annals of Statistics* **34**, 1436–1462.
- Mendoza, M. C., Er, E. E., and Blenis, J. (2011). The Ras-ERK and PI3K-mTOR pathways: cross-talk and compensation. *Trends in Biochemical Sciences* **36**, 320–328.
- Mitra, R., Müller, P., and Ji, Y. (2016). Bayesian graphical models for differential pathways. *Bayesian Analysis* **11**, 99–124.
- Moerkerke, B., Loeys, T., and Vansteelandt, S. (2015). Structural equation modeling versus marginal structural modeling for assessing mediation in the presence of posttreatment

- confounding. *Psychological methods* **20**, 204.
- Mohan, K., London, P., Fazel, M., Witten, D., and Lee, S.-I. (2014). Node-based learning of multiple gaussian graphical models. *J Mach Learn Res* **15**, 445–488.
- Mooij, J. M., Janzing, D., Heskes, T., and Schölkopf, B. (2011). On causal discovery with cyclic additive noise models. In *Advances in neural information processing systems*, pages 639–647.
- Mukherjee, C. and Rodriguez, A. (2016). Gpu-powered shotgun stochastic search for dirichlet process mixtures of gaussian graphical models. *Journal of Computational and Graphical Statistics* **25**, 762–788.
- Müller, P., Parmigiani, G., and Rice, K. (2006). FDR and Bayesian Multiple Comparisons Rules.
- Müller, P., Quintana, F., and Rosner, G. L. (2011). A product partition model with regression on covariates. *Journal of Computational and Graphical Statistics* **20**, 260–278.
- Newton, M. A., Noueir, A., Sarkar, D., and Ahlquist, P. (2004). Detecting differential gene expression with a semiparametric hierarchical mixture method. *Biostatistics* **5**, 155–176.
- Ni, Y., Ji, Y., and Mueller, P. (2016). Reciprocal graphical models for integrative gene regulatory network analysis. *arXiv preprint arXiv:1607.06849*.
- Oates, C. J., Smith, J. Q., and Mukherjee, S. (2016). Estimating causal structure using conditional DAG models. *Journal of Machine Learning Research* **17**, 1–23.
- Oates, C. J., Smith, J. Q., Mukherjee, S., and Cussens, J. (2016). Exact estimation of multiple directed acyclic graphs. *Statistics and Computing* **26**, 797–811.
- Pearl, J. (2003). Causality: models, reasoning and inference. *Econometric Theory* **19**, 46.
- Perez-Moreno, P., Brambilla, E., Thomas, R., and Soria, J.-C. (2012). Squamous cell carcinoma of the lung: molecular subtypes and therapeutic opportunities. *Clinical Cancer Research* **18**, 2443–2451.


- Peterson, C., Stingo, F. C., and Vannucci, M. (2015). Bayesian inference of multiple Gaussian graphical models. *Journal of the American Statistical Association* **110**, 159–174.
- Pitman, J. and Yor, M. (1997). The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator. *The Annals of Probability* **25**, 855–900.
- Polager, S. and Ginsberg, D. (2009). p53 and E2f: partners in life and death. *Nature Reviews Cancer* **9**, 738–748.
- Quintana, F. A. (2006). A predictive view of Bayesian clustering. *Journal of Statistical Planning and Inference* **136**, 2407–2429.
- Richardson, T. (1996a). A discovery algorithm for directed cyclic graphs. In *Proceedings of the Twelfth international conference on Uncertainty in artificial intelligence*, pages 454–461. Morgan Kaufmann Publishers Inc.
- Richardson, T. (1996b). A polynomial-time algorithm for deciding markov equivalence of directed cyclic graphical models. In *Proceedings of the Twelfth international conference on Uncertainty in artificial intelligence*, pages 462–469. Morgan Kaufmann Publishers Inc.
- Richardson, T. (1997). A characterization of markov equivalence for directed cyclic graphs. *International Journal of Approximate Reasoning* **17**, 107–162.
- Robins, J. M. (1999). Testing and estimation of direct effects by reparameterizing directed acyclic graphs with structural nested models. *Computation, Causation, and Discovery* pages 349–405.
- Rodriguez, A., Dunson, D. B., and Gelfand, A. E. (2008). The nested dirichlet process. *Journal of the American Statistical Association* **103**, 1131–1154.
- Rodriguez, A., Lenkoski, A., and Dobra, A. (2011). Sparse covariance estimation in heterogeneous samples. *Electronic Journal of Statistics* **5**, 981.
- Rudelson, M. (2008). Invertibility of random matrices: norm of the inverse. *Annals of*

- Mathematics* **168**, 575–600.
- Santen, R. J., Song, R. X., McPherson, R., Kumar, R., Adam, L., Jeng, M.-H., and Yue, W. (2002). The role of mitogen-activated protein (MAP) kinase in breast cancer. *The Journal of Steroid Biochemistry and Molecular Biology* **80**, 239–256.
- Semba, S., Itoh, N., Ito, M., Youssef, E. M., Harada, M., Moriya, T., Kimura, W., and Yamakawa, M. (2002). Down-regulation of pik3cg, a catalytic subunit of phosphatidylinositol 3-oh kinase, by cpg hypermethylation in human colorectal carcinoma. *Clinical Cancer Research* **8**, 3824–3831.
- TCGA (2012). Comprehensive molecular characterization of human colon and rectal cancer. *Nature* **487**, 330–337.
- Wang, H. and West, M. (2009). Bayesian analysis of matrix normal graphical models. *Biometrika* **96**, 821–834.
- Wang, W., Baladandayuthapani, V., Holmes, C. C., and Do, K.-A. (2013). Integrative network-based Bayesian analysis of diverse genomics data. *BMC Bioinformatics* **14**, S8.
- Whittaker, J. (2009). *Graphical Models in Applied Multivariate Statistics*. Wiley Publishing, New York.
- Yajima, M., Telesca, D., Ji, Y., and Müller, P. (2015). Detecting differential patterns of interaction in molecular pathways. *Biostatistics* **16**, 240–251.
- Yang, E., Ravikumar, P., Allen, G. I., and Liu, Z. (2015). Graphical models via univariate exponential family distributions. *Journal of Machine Learning Research* **16**, 3813–3847.
- Zhang, L. and Kim, S. (2014). Learning gene networks under SNP perturbations using eQTL datasets. *PLoS Comput Biol* **10**, e1003420.
- Zhu, Y., Qiu, P., and Ji, Y. (2014). TCGA-assembler: open-source software for retrieving and processing TCGA data. *Nature Methods* **11**, 599–600.







**Figure 1.** Graphs for illustration. Graphs in (a)-(d) are Markov equivalent reciprocal graphs. Each graph in (e)-(h) encodes different conditional independence relationships.




**Figure 2.** Inferred gene networks for HNSC, LUSC and luminal BRCA. Shared edges across all subtypes are represented by solid lines; differential edges are shown as dashed lines. Arrowheads represent stimulatory interactions, whereas horizontal bars denote inhibitory regulations.



**Figure 3.** Inferred networks for COAD and READ. Shared edges across cancer types are represented by solid lines; differential edges are shown as dashed lines. Edges with arrowheads represent stimulatory interactions, whereas edges with horizontal bars denote inhibitory regulations. (a) Full network in COAD with copy number, methylation and gene expression. (b) Full network in READ. (c) Gene network in COAD. (d) Gene network in READ. In (a) and (b), the suffixes represent: c=copy number, m=methylation and g=gene. In (c) and (d), disconnected genes are omitted.



**Figure 4.** Kaplan-Meier estimators for cluster 1 (solid) and cluster 2 (dashed). P-values of log-rank test are shown on the top of each figure.



**Figure 5.** Inferred gene networks for breast cancer. Top two figures: networks for clusters 1 and 2 with shared edges represented by solid lines and differential edges represented by dashed lines. Bottom figure: negative feedback loop of gene network for cluster 2. Arrowheads represent stimulatory interactions, whereas horizontal bars denote inhibitory regulations.

**Table 1**

*Network summaries for HNSC (H), LUSC (L) and BRCA (B). Upper triangular: the number of edges in each of the three gene networks and the number of shared edges between each pair of networks. Lower triangular: the estimated relatedness matrix  $\hat{\Omega}$ .*

	H	L	B
H	16	10	9
L	0.77	11	8
B	0.63	0.57	13