Nonreversible Langevin Samplers: Splitting Schemes, Analysis and Implementation

A.B. Duncan
School of Mathematical and Physical Sciences
University of Sussex
Falmer
and
G.A. Pavliotis
Department of Mathematics
Imperial College London
London SW7 2AZ, UK
and
K.C. Zygalakis
School of Mathematics
University of Edinburgh
Edinburgh, EH9 3FD, UK

January 17, 2017

Abstract

For a given target density $\pi:\mathbb{R}^d\mapsto\mathbb{R}$, there exist an infinite number of diffusion processes which have unique invariant density π . As observed in a number of papers [7, 37, 38] samplers based on nonreversible diffusion processes can significantly outperform their reversible counterparts both in terms of asymptotic variance and rate of convergence to equilibrium. In this paper, we take advantage of this in order to construct efficient sampling algorithms based on the Lie-Trotter decomposition of a nonreversible diffusion process into reversible and nonreversible components. We show that samplers based on this scheme can significantly outperform standard MCMC methods, at the cost of introducing some controlled bias. In particular, we prove that numerical integrators constructed according to this decomposition are geometrically ergodic and characterize fully their asymptotic bias and variance, showing that the sampler inherits the good mixing properties of the underlying nonreversible diffusion. This is illustrated further with a number of numerical examples ranging from highly correlated low dimensional distributions, to logistic regression problems in high dimensions as well as inference for spatial models with many latent variables.

1 Introduction

Consider the problem of computing expectations with respect to a probability distribution with smooth density $\pi(x)$, known only up to a normalization constant, i.e. we wish to evaluate

$$\pi(f) = \int_{\mathbb{R}^d} f(x)\pi(x) dx. \tag{1.1}$$

For high dimensional distributions, deterministic techniques are no longer tractable. On the other hand, probabilistic methods do not suffer the same curse of dimensionality and thus are often the method of choice. One such approach is *Markov Chain Monte Carlo* (MCMC) which is based on the construction of a Markov process on \mathbb{R}^d whose unique invariant distribution is $\pi(x)$. Due to their simplicity and wide applicability, Markov chains based on Metropolis-Hastings (MH) transition kernels [13, 28] and their numerous variants remain the most widely used scheme for sampling from a general target probability distribution, despite having been introduced over 60 years ago. As there are infinitely many Markov processes which are ergodic with respect to a given target distribution

 π , a natural question is whether a Markov process can be chosen which is more efficient, in terms of convergence to equilibrium and mixing. Metropolized schemes are reversible Markov chains by construction, i.e. they satisfy *detailed balance*. It is a well documented fact that nonreversible chains might convergence to equilibrium faster than reversible ones [34, 5, 32]. Various MCMC schemes have been proposed which are based on the general idea of breaking reversibility by introducing an augmented target measure on an extended state space, along with dynamics which are invariant with respect to the augmented target measure. For discrete state spaces, the lifting method [5, 15, 47] is one such approach, where the Markov chain is "lifted" from the state space E to $E \times \{1, -1\}$. The transition probabilities in each copy of E are modified to introduce transitions between the copies to preserve the invariant distribution but now promote the sampler to generate long trajectories. For continuous state spaces, analogous approaches involve augmenting the state space with a velocity/momentum variable and constructing Makovian dynamics which are able to mix more rapidly in the augmented state space. Such methods include Hybrid Monte Carlo (HMC) methods, inspired by Hamiltonian dynamics. While the standard construction of HMC [6, 35] is reversible, it is straightforward to construct dynamics based on the Generalized HMC scheme [14] which will not be reversible, see also [36] and more recently [23].

Deferring issues of simulation until later, another candidate Markov process for sampling from π is the diffusion $(X_t)_{t>0}$ defined by the following Itô stochastic differential equation (SDE):

$$dX_t = b(X_t) dt + \sqrt{2} dW_t, \tag{1.2}$$

where W_t is a standard \mathbb{R}^d -valued Brownian motion and $b: \mathbb{R}^d \to \mathbb{R}^d$ is a smooth vector field which satisfies

$$b(x) = \nabla \log \pi(x) + \gamma(x), \quad \nabla \cdot (\pi(x)\gamma(x)) = 0, \tag{1.3}$$

for some smooth vector field γ on \mathbb{R}^d satisfying some mild assumptions (c.f. Proposition 2.2). The process X_t is nonreversible if and only if $\gamma \neq 0$. By the Birkhoff ergodic theorem,

$$\lim_{T \to \infty} \frac{1}{T} \int_0^T f(X_s) \, ds = \mathbb{E}_{\pi}[f], \quad f \in L^1(\pi),$$

and thus one can use

$$\pi_T(f) := \frac{1}{T} \int_0^T f(X_s) \, ds$$

as an estimator for $\mathbb{E}_{\pi}[f]$, for T sufficiently large. A natural way to measure the efficiency of such estimator is the mean square error (MSE) given by

$$MSE(T) := \mathbb{E}|\pi_T(f) - \pi(f)|^2.$$
 (1.4)

Under appropriate conditions on X_t and f, the estimator $\pi_T(f)$ will satisfy a *central limit theorem*, i.e.

$$\lim_{T \to +\infty} \sqrt{T} \left(\pi_T(f) - \mathbb{E}_{\pi}[f] \right) = \mathcal{N}(0, 2\sigma^2(f)), \tag{1.5}$$

where $\sigma^2(f)$ is the asymptotic variance of the estimator $\pi_T(f)$ which can be expressed by

$$\sigma^2(f) := \langle \phi, (-\mathcal{L})\phi \rangle_{\pi}, \tag{1.6}$$

where \mathcal{L} is the infinitesimal generator of (1.2) and ϕ is the mean zero solution of the following Poisson equation on \mathbb{R}^d ,

$$-\mathcal{L}\phi = f - \pi(f). \tag{1.7}$$

This relationship can be used to simplify the expression for the MSE (1.4) and decompose it in terms of bias $\mu_T(f)$ and variance $\sigma_T^2(f)$ as follows

$$\mathbb{E}|\pi_T(f) - \pi(f)|^2 = (\mathbb{E}\pi_T(f) - \pi(f))^2 + \mathbb{E}(\pi_T(f) - \mathbb{E}\pi_T(f))^2 = (\mu_T(f))^2 + \sigma_T^2(f).$$

For large T, the variance satisfies $\sigma_T^2(f) \simeq T^{-1}\sigma^2(f)$, while $\mu_T(f)^2 = o(T^{-1})$. Since $\gamma(x)$ is not uniquely defined in (1.3), a natural question is how it should be chosen to ensure that for a given time T, the MSE in (1.4) is as small as possible. This can be achieved in two manners, the first by maximising the $L^2(\pi)$ -spectral gap associated with (1.2) as studied in [20, 48] and hence increasing the speed with which μ_T converges to zero. In

general, maximising the $L^2(\pi)$ -spectral gap is challenging. An alternative is to choose $\gamma(x)$ in such a way so as to reduce the asymptotic variance $\sigma^2(f)$. It should be emphasised that the optimal choice will be different for each case. In particular in [7, 37, 38], it was shown that the choice $\gamma(x)=0$, which corresponds to using reversible dynamics, gives the maximum value of asymptotic variance for a given choice of diffusion tensor. In particular, introducing a nonreversible perturbation will never decrease the performance of an estimator based on Langevin dynamics, both in terms of convergence to equilibrium and asymptotic variance.

In general (1.2) cannot be simulated exactly, and one typically resorts to a discretisation of the SDE, denoted by $\widehat{X}_{n}^{\Delta t}$, in order to approximate $\pi(f)$. In particular, the following ergodic average is used

$$\widehat{\pi}_T^{\Delta t}(f) := \frac{1}{N} \sum_{k=0}^N f(\widehat{X}_k^{\Delta t}), \quad N\Delta t = T.$$
(1.8)

Extra caution has to be taken in order to ensure that the above quantity converges in the limit of $T \to \infty$ since even if (1.2) is ergodic (or even exponentially ergodic), this will not necessarily be the case for its numerical discretisation [39, 43, 44]. In addition, even when the numerical discretization is ergodic and thus

$$\lim_{T \to \infty} \widehat{\pi}_T(f) = \widehat{\pi}^{\Delta t}(f) = \int_{\mathbb{R}^d} f(x)\widehat{\pi}^{\Delta t}(x)dx, \tag{1.9}$$

it is not true in general that $\widehat{\pi}^{\Delta t} = \pi$, since the underlying numerical discretization introduces bias in the estimation of $\pi(f)$ (see [45, 1, 2]). One way to eliminate such bias is through Metropolization [42, 46], *i.e.* the introduction of an accept-reject step that ensures that the corresponding Markov chain is ergodic with respect to the target distribution π . However, such bias elimination might not be advantageous in practice since the Metropolised chain will be reversible by construction, thus eliminating any benefit introduced by the nonreversible perturbation γ .

When computing expectations of distributions with expensive likelihoods, it might be too costly to sample a long Markov chain trajectory. If an appropriate nonreversible Langevin dynamics (1.2) can be introduced which does give rise to a dramatic reduction in asymptotic variance, then it might be advantageous to permit a controlled amount of bias in exchange for needing to sample fare less. This bias-variance tradeoff, in the context of numerical discretisations of (1.2) is the subject of study of this paper. In particular, we will consider discretizations based on a Lie-Trotter splitting between the reversible and the nonreversible part of the dynamics. More specifically, we consider integrators of the form

$$\widehat{X}_{n+1}^{\Delta t} = \Theta_{\Delta t} \circ \Phi_{\Delta t}(\widehat{X}_n^{\Delta t}), \tag{1.10}$$

where $\Phi_{\Delta t}(x)$ is a integrator that approximates the flow map corresponding to the deterministic dynamics

$$\frac{dx_t}{dt} = \gamma(x_t),\tag{1.11}$$

and $\Theta_{\Delta t}(x)$ which approximates the reversible dynamics

$$dx_t = \nabla \log \pi(x_t)dt + \sqrt{2}dW_t. \tag{1.12}$$

The choice of $\Phi_{\Delta t}$, $\Theta_{\Delta t}$ has a fundamental influence on the bias, asymptotic variance and stability of the resulting sampler. In particular, if one chooses $\Phi_{\Delta t}$ to be a Metropolised integrator [3] then, similarly to the result in [2], the order of convergence of the deterministic integrator $\Phi_{\Delta t}$ provides a lower bound for the difference between expectations with respect to $\widehat{\pi}^{\Delta t}$ and π . However, this is not the case for the numerical asymptotic variance $\widehat{\sigma}_{\Delta t}^2(f)$, since even though we can show that it is a perturbation of $\sigma^2(f)$ the difference will depend crucially on the choice of $\Theta_{\Delta t}$. These results are important as they allow to choose the correct combination of dynamics and numerical scheme that drastically reduces the computational cost required to achieve a given tolerance of error.

In summary, the main of the contributions of this paper are

- 1. proving geometric ergodicity for the Markov chain given by (1.10) for a variety of different numerical integrators applied to the reversible part;
- 2. a complete characterisation of the asymptotic bias of (1.10);

- 3. showing that, by completely characterising the asymptotic variance, numerical integrators of the type (1.10) inherit the asymptotic variance benefits of the non reversible SDE (1.2);
- 4. exhibiting the potential of using nonreversible integrators for sampling as illustrated from a number of different numerical experiments on inference for spatial models as well as real data sets.

The rest of the paper is organised as follows. In Section 2 we describe some known theoretical results for the SDE (1.2) which are necessary for the development of this paper. In Section 3 we identifity sufficient conditions to guarantee geometric ergodicity of the Lie-Trotter splitting scheme (1.10) on \mathbb{R}^d . In Section 4 we study the asymptotic properties of a class of numerical integrators for (1.2) for which the Lie-Trotter scheme is a special case. In particular we derive perturbative expansions for the asymptotic bias and variance. In Section 5 we apply these results to characterise the asymptotic bias and variance of the Lie-Trotter scheme on the bounded domain \mathbb{T}^d . In Section 6, we focus on the case where the target distribution is Gaussian and study analytically the trade-off between the asymptotic bias and asymptotic variance in this case. To demonstrate the efficacy of these schemes, in Section 7 we present a number of numerical experiments on inference for spatial models as well as on Bayesian logistic regression. Proofs of the main results of this paper are deferred to Section 8 as well as the Appendices. Finally, a discussion of the results presented in this paper and potential future research directions can be found in Section 9.

2 Properties of Overdamped Langevin Diffusions

In this section we discuss different known theoretical results that are useful for understanding the main results of the paper. We start by listing the assumptions we shall make on π and the SDE (1.2) to ensure ergodicity.

Assumptions 2.1.

- 1. The measure π possesses a positive smooth density $\pi(x) > 0$, known up to a normalizing constant, such that $\pi \in L^1(\mathbb{R}^d)$.
- 2. The drift vector $b: \mathbb{R}^d \to \mathbb{R}^d$ of (1.2) is smooth and satisfies

$$b(x) = \nabla \log \pi(x) + \gamma(x), \tag{2.1}$$

where $\gamma: \mathbb{R}^d \to \mathbb{R}^d$ is a smooth vector field with components in $L^1(\pi)$ such that

$$\nabla \cdot (\pi(x)\gamma(x)) = 0. \tag{2.2}$$

The following result provides necessary and sufficient conditions on the coefficients of (1.2) to ensure that X_t possesses a unique stationary distribution π .

Proposition 2.2. Suppose that Assumptions 2.1 hold. Then the diffusion process X_t defined by (1.2) possesses a strongly continuous semigroup $(P_t)_{t>0}$ on $L^2(\pi)$ defined by

$$P_t f(x) = \mathbb{E}[f(X_t) | X_0 = x]. \tag{2.3}$$

The associated infinitesimal generator is an an extension of

$$\mathcal{L} = \frac{1}{\pi} \nabla \cdot (\pi \nabla \cdot) + \gamma \cdot \nabla \tag{2.4}$$

with core $C_c^{\infty}(\mathbb{R}^d)$. Moreover, P_t has unique invariant distribution π . Conversely, given a diffusion process of the form (1.2) which is invariant with respect to π , then the drift b necessarily satisfies (2.1) and (2.2).

Proof. The first part of this result is a direct application of [21, Thm 8.1.26]. The converse implication can be checked using integration by parts. \Box

While many choices for γ are possible (see [22] for a more complete recipe) a natural family of vector fields is given by $\gamma(x) = J \nabla \Phi(\pi(x))$, where Φ is a smooth function satisfying $\nabla \Phi(\pi(\cdot)) \in L^1(\pi)$ and J is $d \times d$ skew-symmetric matrix. We shall focus specifically on the following three choices:

1. If π satisfies $\int_{\mathbb{R}^d} |\nabla \log \pi(x)| \pi(dx) < \infty$, then the vector field

$$\gamma(x) = J\nabla \log \pi(x), \quad J = -J^{\top}, \tag{2.5}$$

satisfies condition (2.2). This was the choice which was studied specifically in [7].

2. If $\int_{\mathbb{R}^d} |\nabla \log \pi(x)| \pi^{1+\alpha}(dx) < \infty$ for some $\alpha > 0$ then another natural choice for the vector field is given by

$$\gamma(x) = J \nabla \pi^{\alpha}(x), \quad J = -J^{\top}. \tag{2.6}$$

Although (2.6) introduces an additional tuning parameter α , one might prefer this choice as it coincides with the intuition that when far away from the modes the sampler should move towards the modes as quickly as possible, and should only undergo these deterministic meanders in regions of high probability.

3. Let $\Psi: \mathbb{R} \to \mathbb{R}$ be a smooth, compactly supported function. Then

$$\gamma(x) = J\nabla \log \pi(x)\Psi(\pi(x)), \quad J = -J^{\top}, \text{ and } \beta \in \mathbb{R},$$
 (2.7)

will always satisfy (2.2). Moreover, if π has compact level sets, then γ will also be compactly supported on \mathbb{R}^d .

Applying the results detailed in [11, 31], we shall assume that the process X_t possesses a Lyapunov function, which is sufficient to ensure the exponential ergodicity of X_t , as detailed in the subsequent proposition.

Assumptions 2.3 (Foster–Lyapunov Criterion). There exists a function $V: \mathbb{R}^d \to \mathbb{R}$ and constants c > 0 and $b \in \mathbb{R}$ such that

$$\mathcal{L}V(x) \le -cV(x) + b\mathbf{1}_C$$
, and $V(x) \ge 1$, $x \in \mathbb{R}^d$, (2.8)

where $\mathbf{1}_C$ is the indicator function over a petite set.

For the definition of a petite set we refer the reader to [30]. For the generator \mathcal{L} corresponding to the process (1.2) compact sets are always petite. The exponential ergodicity of X_t follows from the following proposition (see also [24, 30]).

Proposition 2.4. Suppose that Assumption 2.3 holds, then there exist constants C > 0 and $\lambda > 0$ such that:

$$|P_t f(x) - \pi(f)| \le CU(x)e^{-\lambda t}, \quad x \in \mathbb{R}^d, \tag{2.9}$$

for all f satisfying $|f| \leq U$.

Moreover, the Foster-Lyapunov criterion also provides a sufficient condition for the Poisson equation (1.7) to be well-posed, and thus for the central limit theorem (1.5) to hold.

Proposition 2.5. Suppose that Assumption 2.3 holds and that $\pi(U^2) < \infty$, then for any function f such that $|f| \leq U$, the central limit theorem (1.5) holds, i.e. $\sqrt{T}(\pi_T(f) - \pi(f))$ converges weakly to a $\mathcal{N}(0, \sigma^2(f))$ -distributed random variable, with

$$\sigma^{2}(f) = \int_{\mathbb{R}^{d}} \phi(x)(-\mathcal{L})\phi(x)\pi(x) dx,$$

where ϕ is the unique mean zero solution to the Poisson equation (1.7). Moreover the solution ϕ can be expressed as

$$\phi = \int_0^\infty \left[P_t f - \pi(f) \right] dt.$$

The following lemma provides a sufficient condition on π for (1.2) to possess a Lyapunov function. It is a slight generalisation of a similar result from [40], extended to apply also in the case of nonreversible diffusion processes.

Lemma 2.6. [40, Theorem 2 .3] Consider the process X_t defined by (1.2) with drift coefficient b satisfying (2.1). Suppose that π is bounded, there exists $0 < \delta < 1$ such that,

$$\lim_{|x| \to \infty} \inf \left((1 - \delta) |\nabla \log \pi(x)|^2 + \Delta \log \pi(x) \right) > 0, \tag{2.10}$$

and the vector field γ satisfies

$$\nabla \cdot \gamma(x) = 0, \quad x \in \mathbb{R}^d, \tag{2.11}$$

then the Foster-Lyapunov criterion holds for (1.2) with $U(x) = \pi^{-\delta}(x)$ and moreover $\pi(U) < \infty$.

Remark 2.7. Note that when $\gamma(x) = J\nabla\Phi(\pi(x))$ equation (2.11) is automatically satisfied. Hence the choices of choices of γ specified by (2.5), (2.6) and (2.7) all satisfy (2.11).

3 Geometric ergodicity of the splitting scheme on \mathbb{R}^d

In this section we identify sufficient conditions under which the Lie-Trotter scheme on \mathbb{R}^d is geometrically ergodic with respect to an invariant distribution $\widehat{\pi}^{\Delta t}$ which will be a perturbation of π . In general, a discretization of the ergodic diffusion process (1.2) need not to be ergodic, geometric or otherwise, see [40]. For the splitting scheme we shall show that provided the approximate nonreversible flow $\Phi_{\Delta t}$ is sufficiently weak away from the origin, the process (1.10) will inherit the geometric ergodicity from the reversible dynamics.

We follow the Meyn and Tweedie [30] recipe to demonstrate geometric ergodicity of $\left(\widehat{X}_n^{\Delta t}\right)_{n\in\mathbb{N}}$. Consider the reversible process defined by

$$Z_{n+1}^{\Delta t} = \Theta_{\Delta t} Z_n^{\Delta t},\tag{3.1}$$

and $\widetilde{P}_{\Delta t}$ be the corresponding transition semigroup. We shall assume that the reversible dynamics are a Metropolis-Hastings chain, with proposal kernel $q_{\Delta t}(\cdot|x)$, more specifically, given $x \in \mathbb{R}^d$, $\Theta_{\Delta t}(x)$ is constructed as follows

- 1. Sample $y \sim q_{\Delta t}(\cdot \mid x)$.
- 2. With probability

$$\alpha(x,y) = \min\left(1, \frac{\pi(y)q_{\Delta t}(x|y)}{\pi(x)q_{\Delta t}(y|x)}\right),\,$$

set $\Theta_{\Delta t}x := y$ otherwise $\Theta_{\Delta t}x := x$.

It is well known that the target distribution π is invariant under the map $\Theta_{\Delta t}$ [28, 13].

Denote by $\widehat{P}_{\Delta t}(x,\cdot)$ and $\widetilde{P}_{\Delta t}(x,\cdot)$ the transition distribution functions of the splitting scheme (1.10) and (3.1) respectively. Then clearly

$$\widehat{P}_{\Delta t}f(x,A) = (\widetilde{P}_{\Delta t}f)(\Phi_{\Delta t}(x),A), \quad A \in \mathcal{B}(\mathbb{R}^d).$$

Following the approach of [26] we first show that (1.10) is a π -irreducible, aperiodic Markov chain. Moreover, we will show that all compact sets are small, i.e. for every compact set C, there exists a $\delta > 0$ and n > 0 such that

$$\widehat{P}_{\Delta t}^{n}(x,\cdot) \ge \delta \nu(\cdot), \quad x \in C.$$

Finally, we will show that if a Foster-Lyapunov condition holds for the reversible dynamics $\widetilde{P}_{\Delta t}$, then it also holds for $\widehat{P}_{\Delta t}$. To this end, we shall make the following assumptions.

Assumptions 3.1. For Δt sufficiently small, we assume that

I The reversible chain (3.1) satisfies a Foster-Lyapunov condition, i.e. there exists a continuous function $V \ge 1$, a compact set $C \subset \mathbb{R}^d$ and constants $\lambda \in (0,1)$ and $b \ge 0$ such that

$$\widetilde{P}_{\Delta t}V(x) \le \lambda V(x) + b\mathbf{1}_C(x), \quad x \in \mathbb{R}^d.$$
 (3.2)

2 The nonreversible flow map $\Phi_{\Delta t}$ satisfies the following condition,

$$\lim \sup_{|x| \to \infty} \frac{V(\Phi_{\Delta t}(x)) - V(x)}{V(x)} < \frac{1}{\lambda} - 1.$$
(3.3)

3 The preimage $\Phi_{\Delta t}^{-1}(C)$ is bounded.

The main theorem of this section establishes the geometric ergodicity of (1.10).

Theorem 3.2. Suppose that Assumptions 3.1 hold, and that π and $q_{\Delta t}(y|x)$ are positive and continuous for all $x,y\in\mathbb{R}^d$. Then for Δt sufficiently small, the process $\widehat{X}_n^{\Delta t}$ is geometrically ergodic, i.e. there exists $\rho\in(0,1)$ and K>0 such that

$$\sup_{|g| \le V} \left| \int_{\mathbb{R}^d} g(y) \left(P_{\Delta t}(x, y) - \pi(y) \right) \, dy \right| \le KV(x) \rho^n, \quad n \in \mathbb{N}.$$

We now focus on the case when the reversible dynamics are simulated using MALA (Metropolis-Adjusted Langevin Algorithm), i.e. using a proposal of the form

$$q_{\Delta t}(\cdot|x) \sim \mathcal{N}(x + \nabla \log \pi(x)\Delta t, 2\Delta t),$$
 (3.4)

for a stepsize $\Delta t > 0$. The following result is an application of Theorem 3.2 for the proposal (3.4).

Corollary 3.3 (Geometric Ergodicity of Lie-Trotter scheme with MALA dynamics). Consider the Lie-Trotter splitting scheme $\widehat{X}_n^{\Delta t}$ where the reversible dynamics (1.12) are simulated using a MALA scheme with proposal defined by (3.4). Suppose that the conditions on π and $q_{\Delta t}$ specified in [40, Theorem 4.1] hold and moreover that

$$\lim_{|x| \to \infty} (|\Phi_{\Delta t}(x)| - |x|) = 0, \tag{3.5}$$

for Δt sufficiently small. Then $\widehat{X}_n^{\Delta t}$ is geometrically ergodic.

In particular, suppose that $\lim_{|x|\to\infty} \pi(x) \to 0$, and that, given $\alpha > 0$, there exist positive constants α' , K_1 and K_2 such that

$$|\nabla \pi^{\alpha}(x)| \le K_1 \pi^{\alpha'}(x), \quad |\nabla \nabla \pi^{\alpha}(x)|_{max} \le K_2, \quad x \in \mathbb{R}^d,$$
 (3.6)

where $|\cdot|_{max}$ denotes the max norm. If $\gamma = J\nabla\pi^{\alpha}$ for J antisymmetric, then condition (3.5) will hold if $\Phi_{\Delta t}(x)$ is simulated using an explicit Euler or Runge-Kutta scheme. A similar result holds for γ given by (2.7).

4 Asymptotic Bias and Variance Estimates for general integrators

In this section we consider the asymptotic behaviour of the estimator (1.8) for $\pi(f)$, obtained for a general numerical scheme $(\widehat{X}_k^{\Delta t})_{k\geq 0}$. In particular, we shall derive estimates for the asymptotic bias and asymptotic variance of the estimator $\widehat{\pi}_{\Delta t}(f)$. For simplicity we shall focus on the case where the domain is \mathbb{T}^d , i.e. the unit hypercube with periodic boundary conditions. As in [25] this set-up greatly simplifies the derivation of expressions for bias and variance, particularly since remainder terms arising from Taylor expansions can be easily controlled. We expect that extending these results to unbounded domains should be possible by following analogous approaches in [17]. Throughout this section, we shall assume that the numerical integrator $\widehat{X}_k^{\Delta t}$ is ergodic, with unique invariant distribution $\widehat{\pi}^{\Delta t}$.

4.1 Notation

We first introduce the notation which will be used in this section and the remainder of the paper. Given a probability measure μ on $(\mathbb{T}^d,\mathcal{B}(\mathbb{T}^d))$ define $L^2(\mu)$ to be the Hilbert space of square integrable functions on \mathbb{T}^d , equipped with inner product $\langle\cdot,\cdot\rangle_{\mu}$ and norm $\|\cdot\|_{L^2(\pi)}$. The subspace $L^2_0(\mu)$ of $L^2(\mu)$ is defined to be

$$L_0^2(\mu) = \{ f \in L^2(\mu) : \mu(f) = 0 \}, \tag{4.1}$$

We define $L^{\infty}(\mu)$ (also denoted $L^{\infty}(\mathbb{T}^d)$) to be the Banach space of essentially bounded functions on \mathbb{T}^d equipped with norm $\|\cdot\|_{L^{\infty}(\mathbb{T}^d)}$. The subspace $L^{\infty}_0(\mu)$ of $L^{\infty}(\mu)$ is defined analogously to (4.1). Finally, given a (signed) measure ν on $(\mathbb{T}^d,\mathcal{B}(\mathbb{T}^d))$ we denote the total variation norm of ν by $\|\nu\|_{TV}$.

4.2 Backward error analysis for ODEs

Backward error analysis is a powerful tool for the analysis of numerical integrators for differential equations [41, 19, 12]. In particular, it is the main ingredient for the proof of the good energy conservation (without drift) of symplectic Runge-Kutta methods when applied to deterministic Hamiltonian systems over exponentially long time intervals [12]. In our context it is useful to characterize the infinitesimal generator of the numerical flow $\Phi_{\Delta t}$ approximating the solution of the ODE (1.11). Indeed, given a consistent integrator $z_{n+1} = \Phi_{\Delta t}(z_n)$ for the ODE

$$\frac{dz(t)}{dt} = f(z(t)),\tag{4.2}$$

the idea of backward error analysis is to search for a modified differential equation written as a formal series in powers of the stepsize Δt ,

$$\frac{d\widetilde{z}}{dt} = f(\widetilde{z}) + \Delta t f_1(\widetilde{z}) + \Delta t^2 f_2(\widetilde{z}) + \dots, \quad \widetilde{z}(0) = z_0$$
(4.3)

such that (formally) $z_n = \widetilde{z}(t_n)$, where $t_n = n\Delta t$ (in the above differential equation, we omit the time variable for brevity). The numerical solution can this be interpreted as a higher order approximation of the exact solution of a modified ODE. For all reasonable integrators, the vector fields f_j can be constructed inductively [19, 12], starting from $f_0 = f$. In general, the series in (4.3) will diverge for nonlinear systems, and thus needs to be truncated. We thus consider the truncated modified ODE at order s

$$\frac{d\widetilde{z}}{dt} = f(\widetilde{z}) + \Delta t f_1(\widetilde{z}) + \Delta t^2 f_2(\widetilde{z}) + \dots + \Delta t^s f_s(\widetilde{z}), \quad \widetilde{z}(0) = z_0$$
(4.4)

we have $z_n = \widetilde{z}(t_n) + \mathcal{O}(\Delta t^{s+1})$ for $\Delta t \to 0$ for bounded times $t_n = n\Delta t \le T$. We note that the flow $\widetilde{\Phi}_{\Delta t}(z)$ of the modified differential equation (4.4) satisfies

$$\phi \circ \widetilde{\Phi}_{\Delta t} = \left(\sum_{k=0}^{M} \frac{\Delta t^k \widetilde{\mathcal{L}}_D^k}{k!}\right) \phi + \mathcal{O}(\Delta t^{M+1}), \qquad \widetilde{\mathcal{L}}_D = F_0 + \Delta t F_1 + \Delta t^2 F_2 + \ldots + \Delta t^s F_s, \tag{4.5}$$

for all $M \geq 0$, and smooth test functions ϕ , and where $F_j \phi = f_j \cdot \nabla \phi$, $j = 0, \dots, s$ and $f_0 = f$. Note that the $\mathcal{O}(\Delta t^{M+1})$ terms in (4.5) are independent of $\Delta t \to 0$ but depend on M, s and ϕ^1 .

4.3 Asymptotic bias of numerical integrators

The aim of this subsection is to describe the conditions on a numerical integrator for (1.2) which are sufficient for the numerical invariant distribution $\widehat{\pi}^{\Delta t}$ to approximate π to order r in the weak sense. These conditions relate directly to the expansion of one-step numerical expectations in powers of Δt . In particular, denote by $\widehat{P}_{\Delta t}$ the transition semigroup associated with $\widehat{X}^{\Delta t}$, i.e.

$$\widehat{P}_{\Delta t}f := \mathbb{E}\left[f(\widehat{X}_1^{\Delta t})|X_0 = x\right].$$

and assume that the following expansion holds

$$\widehat{P}_{\Delta t}f = f + \Delta t A_0 f + \dots + \Delta t^k A_{k-1} f + \Delta t^{k+1} A_k f + \Delta t^q Q_{f,\Delta t}, \quad q > k+1$$

$$\tag{4.6}$$

where $A_i, i=0,1,\cdots k$ are linear differential operators with coefficients depending smoothly on $\pi(x)$ and its derivatives, as well as on the choice of the numerical integrator. In addition $Q_{f,\Delta t}$ is a smooth remainder term depending both on f and Δt while being uniformly bounded with respect to Δt . The following theorem provides sufficient conditions for expectations with respect to $\widehat{\pi}^{\Delta t}$ to approximate expectations with respect to π to order r.

Theorem 4.1. Consider equation (1.2) solved by an numerical scheme which is ergodic with respect to some probability measure $\hat{\pi}_{\Delta t}$ and such that

$$A_i^* \pi = 0, \quad \text{for} \quad j = 1, \dots, r - 1,$$
 (4.7)

where q > r, then one obtains

$$\int_{\mathbb{T}^d} f(x)\widehat{\pi}^{\Delta t}(dx) = \int_{\mathbb{T}^d} f(x)\pi(dx) + \Delta t^r \int_{\mathbb{T}^d} A_r(-\mathcal{L})^{-1}(f - \pi(f))\pi(dx) + \Delta t^q R_{f,\Delta t}, \tag{4.8}$$

where the remainder term $R_{f,\Delta t}$ is uniformly bounded with respect to Δt , for Δt sufficiently small.

Remark 4.2. Integrators $\widehat{X}_n^{\Delta t}$ which have weak error order r will automatically satisfy condition (4.7) for $j=0,\ldots,r-1$. However, the converse is not necessarily true, see [1] for further discussion.

An immediate corollary of Theorem (4.1) is that, if (4.7) holds, then for Δt sufficiently small, the estimator $\hat{\pi}_T$ given by (1.8) satisfies

$$\lim_{N \to \infty} \widehat{\pi}_{N\Delta t}(f) = \pi(f) + \Delta t^r \int_{\mathbb{T}^d} A_r(-\mathcal{L})^{-1} (f - \pi(f)) \pi(dx).$$

¹For all Δt small enough, the sum in (4.5) can be shown to converge for $M \to \infty$ in the case of analytic vector fields f_j (and analytic test functions ϕ), which permits to remove the $\mathcal O$ remainder.

4.4 Asymptotic variance of numerical integrators

The aim of this subsection is to derive a perturbation expansion in the small timestep regime for the asymptotic variance of an arbitrary ergodic numerical integrator for the dynamics (1.2). To this end, we consider a diffusion X_t for which the central limit theorem (1.5) holds. Moreover, we shall make the following assumption, which implies that the corresponding numerical scheme $\widehat{X}_k^{\Delta t}$ converges to equilibrium exponentially fast in $L^{\infty}(\mathbb{T}^d)$, with rate which is uniform with respect to Δt .

Assumptions 4.3. There exist constants C > 0 and $\lambda > 0$ independent of Δt such that, for Δt sufficiently small,

$$\left\|\widehat{P}_{\Delta t}^{k}f - \widehat{\pi}^{\Delta t}(f)\right\|_{L^{\infty}(\mathbb{T}^{d})} \leq Ce^{-\lambda k\Delta t} \left\|f - \widehat{\pi}^{\Delta t}(f)\right\|_{L^{\infty}(\mathbb{T}^{d})}, \quad f \in L^{\infty}(\mathbb{T}^{d}).$$

Remark 4.4. This condition is nontrivial to verify in general. For the specific case of the Lie-Trotter integrator (1.10), when the reversible component of the dynamics is integrated using MALA, in Theorem C.3 we prove that Assumption 4.3 holds.

Given an observable $f \in C^{\infty}(\mathbb{T}^d)$ we consider $\widehat{\pi}_T^{\Delta t}$ as in (1.8). We define the rescaled asymptotic variance of the estimator $\widehat{\pi}_T^{\Delta t}$ as follows

$$\widehat{\sigma}_{\Delta t}^{2}(f) = \Delta t \lim_{N \to \infty} N \operatorname{Var}_{\widehat{\pi}^{\Delta t}} \left[\frac{1}{N} \sum_{k=0}^{N-1} f(\widehat{X}_{k}^{\Delta t}) \right]. \tag{4.9}$$

Note here that we rescale the asymptotic variance with Δt , to guarantee a well-defined limit when $\Delta t \to 0$. Assumption 4.3 implies that there exists a constant K > 0, independent of Δt such that

$$\left\| \left[\frac{I - \widehat{P}_{\Delta t}}{\Delta t} \right]^{-1} \right\|_{L_0^{\infty}(\widehat{\pi}_{\Delta t})} < K, \tag{4.10}$$

for Δt sufficiently small. In particular, we can express (4.9) as

$$\widehat{\sigma}_{\Delta t}^{2}(f) = 2\Delta t \left\langle \left(f - \widehat{\pi}^{\Delta t}(f) \right), \left(I - \widehat{P}_{\Delta t} \right)^{-1} \left(f - \widehat{\pi}^{\Delta t}(f) \right) \right\rangle_{\widehat{\pi}^{\Delta t}} - \Delta t \operatorname{Var}_{\widehat{\pi}^{\Delta t}}[f]. \tag{4.11}$$

It should be clear from (4.11) that there will be two contributions to the error between $\widehat{\sigma}_{\Delta t}^2(f)$ and $\sigma^2(f)$: one arising from the order of weak convergence of the numerical method, and one from the time discreteness of the process $\widehat{X}_k^{\Delta t}$. Indeed, even when one considers the exact discrete time dynamics defined by

$$X_n^{\Delta t} = X(n\Delta t), \quad n \in \mathbb{N},$$

the error between the corresponding asymptotic variance $\sigma^2_{\Delta t}(f)$ and $\sigma^2(f)$ will be non-zero, despite the fact that both discrete and continuous time Markov processes have the same invariant distribution. To isolate the different sources of error, we present first Proposition 4.5 which quantifies the effect of the time-discreteness on the asymptotic variance. In Theorem 4.6 we then quantify the error between the asymptotic variances $\sigma^2_{\Delta t}(f)$ and $\widehat{\sigma}^2_{\Delta t}(f)$ of $X_n^{\Delta t}$ and $\widehat{X}_n^{\Delta t}$, respectively.

Proposition 4.5. For all $\phi \in C^{\infty}(\mathbb{T}^d)$, such that $\pi(\phi) = 0$ there exists a smooth function R_{ϕ} such that for Δt sufficiently small,

$$\left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} \phi(x) = (-\mathcal{L})^{-1} \phi(x) + \frac{\Delta t}{2} \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} (-\mathcal{L}) \phi(x) - \frac{\Delta t^2}{6} \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} (-\mathcal{L})^2 \phi + \Delta t^3 R_{\phi}, \tag{4.12}$$

where R_{ϕ} is bounded, independent of Δt . In particular, for $f \in C^{\infty}(\mathbb{T}^d)$,

$$\sigma_{\Delta t}^{2}(f) = \sigma^{2}(f) + \frac{\Delta t^{2}}{6} \left\langle \left(-\mathcal{L}\right) \left(f - \pi(f)\right), f - \pi(f)\right\rangle_{\pi} + o(\Delta t^{2}).$$

Proof. The proof can be found in Section 8.2.

Define the operator $M_{\Delta t}$ to be the projector onto functions with mean zero with respect to $\widehat{\pi}_{\Delta t}$, i.e.

$$M_{\Delta t}\phi(x) = \phi(x) - \int_{\mathbb{T}^d} \phi(y)\widehat{\pi}_{\Delta t}(y) \, dy.$$

The following theorem characterises the difference between the asymptotic variance arising from the exact discrete time dynamics $X_n^{\Delta t}$ and the numerical integrator $\widehat{X}_n^{\Delta t}$.

Theorem 4.6. Suppose that, for some $k \in \mathbb{N}$, $k \geq 1$, there exist operators A_0, \ldots, A_k on $C^{\infty}(\mathbb{T}^d)$, bounded uniformly with respect to Δt , where $A_i = \frac{\mathcal{L}^{i+1}}{(i+1)!}$, $i = 0, \cdots, k-1$ and such that for all $\psi \in C^{\infty}(\mathbb{T}^d)$ the semigroup $\widehat{P}_{\Delta t}$ satisfies (4.6). Suppose that the corresponding invariant distribution $\widehat{\pi}^{\Delta t}$ satisfies

$$\int_{\mathbb{T}^d} \psi(x) \widehat{\pi}^{\Delta t}(x) \, dx = \int_{\mathbb{T}^d} \psi(x) \pi(x) \, dx + \Delta t^r R_{\psi},$$

where r > k and R_{ψ} is a smooth remainder term, uniformly bounded with respect to Δt . Moreover, suppose that $\widehat{P}_{\Delta t}$ satisfies (4.10). Then for all $f, g \in C^{\infty}(\mathbb{T}^d)$ such that $\pi(f) = \pi(g) = 0$, we have the expansion

$$\left\langle g, \left(\frac{I - P_{\Delta t}}{\Delta t} \right)^{-1} f \right\rangle_{\pi} = \left\langle M_{\Delta t} g, \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t} \right)^{-1} M_{\Delta t} f \right\rangle_{\widehat{\pi}_{\Delta t}} + \Delta t^{k} R_{1}(f, g) + o(\Delta t^{k}), \tag{4.13}$$

where

$$R_1(f,g) = \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t} \right)^{-1} M_{\Delta t} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_k \right) \left(\frac{I - P_{\Delta t}}{\Delta t} \right)^{-1} f, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}.$$
 (4.14)

In particular

$$\widehat{\sigma}_{\Delta t}^2(f) = \sigma_{\Delta t}^2(f) + 2\Delta t^k R_1(f, f) + o(\Delta t^k). \tag{4.15}$$

If moreover

$$\pi\left(A_k\psi\right) = 0,\tag{4.16}$$

holds for for all $\psi \in C^{\infty}(\mathbb{T}^d)$, then we can write

$$R_1(f,g) = \left\langle (-\mathcal{L})^{-1} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_k \right) (-\mathcal{L})^{-1} f, g \right\rangle_{\pi} + o(\Delta t^k). \tag{4.17}$$

Proof. The proof can be found at Section 8.2.

Remark 4.7. It is interesting to note that contrary to the case of the asymptotic bias in Theorem 4.1, the order of error for the discrete asymptotic variance in Theorem 4.6 depends crucially on the order of the weak convergence of the underlying numerical integrator. Furthermore, we see that if the weak order of the integrator is than two then the leading order error term between $\hat{\sigma}_{\Delta t}^2(f)$ and the asymptotic variance of the continuous process $\sigma^2(f)$ equals to the leading order term of difference between $\sigma_{\Delta t}^2(f)$ and $\sigma^2(f)$.

To complete this analysis we shall consider the asymptotic variance arising from a perturbed diffusion process \widetilde{X}_t having infinitesimal generator $\widetilde{\mathcal{L}}_{\Delta t}$ such that, for Δt sufficiently small

$$\widetilde{\mathcal{L}}_{\Delta t} f = \mathcal{L} f + \Delta t^k \mathcal{L}_k f + \Delta t^{q-1} R_f, \quad f \in C^{\infty}(\mathbb{T}^d), \tag{4.18}$$

where q > k+1. We shall also assume that $(\widetilde{L}_{\Delta t})^{-1}$ is bounded in $L_0^{\infty}(\widehat{\pi}_{\Delta t})$ uniformly with respect to Δt . More specifically there exists K > 0, independent of Δt such that

$$\left\| \left(-\widetilde{\mathcal{L}}_{\Delta t} \right)^{-1} \right\|_{L_{\infty}^{\infty}(\widehat{\pi}_{\Delta t})} < K, \tag{4.19}$$

for Δt sufficiently small. The following result characterises the influence of this perturbation on the asymptotic variance for small Δt . For numerical approximations of X_t for which a modified SDE [49] is known, the following result combined with Proposition 4.5 provide a convenient means of obtaining an expression for the asymptotic variance $\tilde{\sigma}_{\Delta t}^2$ of the numerical scheme in terms of $\sigma^2(f)$.

Proposition 4.8. Consider a diffusion process \widetilde{X}_t on \mathbb{T}^d with smooth coefficients and generator $\widetilde{\mathcal{L}}_{\Delta t}$ which satisfies (4.18) and (4.19). Suppose that \widetilde{X}_t has unique invariant distribution $\widehat{\pi}_{\Delta t}$ which satisfies

$$\int \psi(x)\widehat{\pi}_{\Delta t}(x) dx = \int \psi(x)\pi(x) dx + \Delta t^r R_{\psi}, \tag{4.20}$$

where r > k, and R_{ψ} is a smooth remainder term, uniformly bounded with respect to Δt . Then for all $f \in C^{\infty}(\mathbb{T}^d)$ with $\pi(f) = 0$:

$$\widetilde{\sigma}_{\Delta t}^2(f) = \sigma_{\Delta t}^2(f) + 2\Delta t^k R_f + o(\Delta t^k). \tag{4.21}$$

where

$$R_f = \left\langle \left(-\widetilde{\mathcal{L}}_{\Delta t} \right)^{-1} M_{\Delta t} (-\mathcal{L}_k) (-\mathcal{L})^{-1} f, M_{\Delta t} f \right\rangle_{\widehat{\pi}_{\Delta t}}.$$
 (4.22)

If moreover

$$\pi\left(\mathcal{L}_k\psi\right) = 0,\tag{4.23}$$

holds for for all $\psi \in C^{\infty}(\mathbb{T}^d)$, then we can write

$$R_f = \left\langle \left(-\mathcal{L} \right)^{-1} \left(-\mathcal{L}_k \right) \left(-\mathcal{L} \right)^{-1} f, f \right\rangle_{\pi} + o(\Delta t^k). \tag{4.24}$$

The result follows from an argument similar to that of Theorem 4.6.

5 Asymptotic Bias and Variance Estimates for the splitting scheme

In this section we derive asymptotic bias and variance estimates for the Lie-Trotter splitting scheme (1.10) on \mathbb{T}^d by applying the general results derived in Section 4. In Section 5.1 we apply Theorem 4.1 to obtain an asymptotic bias estimate for the splitting scheme, while in Section 5.2 we obtain estimates for the asymptotic variance, in the particular case where a MALA scheme is used to integrate the reversible part of the dynamics.

5.1 Asymptotic bias of the splitting scheme

We now consider the Lie-Trotter scheme (1.10) on \mathbb{T}^d . In this section we obtain estimates for the asymptotic bias of the scheme by applying Theorem 4.1.

Theorem 5.1. Suppose that the integrator $\Theta_{\Delta t}$ used for the reversible dynamics is invariant with respect to π and that that the deterministic flow $\Phi_{\Delta t}$ satisfies a modified backward equation of the form (4.3) where the vector fields f_j satisfy

$$\nabla \cdot (f_j(x)\pi(x)) = 0, \quad j = 1, \dots, r - 1. \tag{5.1}$$

Then, assuming ergodicity, the Lie-Trotter splitting (1.10) has order r of accuracy for the invariant measure. More precisely, for all $\phi \in C^2(\mathbb{T}^d)$ and Δt sufficiently small

$$\int_{\mathbb{T}^d} \phi(x) \widehat{\pi}^{\Delta t}(dx) = \int_{\mathbb{T}^d} \phi(x) \pi(dx) + \Delta t^r C_{r,\phi} + \Delta t^{r+1} R_{\phi,\Delta t}, \tag{5.2}$$

where $C_{r,\phi}$ and $R_{\phi,\Delta t}$ are uniformly bounded and

$$C_{r,\phi} = \langle f_r, (-\mathcal{L})^{-1}(\phi - \pi(\phi)) \rangle_{\pi}.$$

Remark 5.2. From standard elliptic energy estimates, the remainder term $C_{r,\phi}$ in (5.2) satisfies the a priori bound

$$|C_{r,\phi}| \le 2||f_r||_{L^2(\pi)} ||\phi||_{L^2(\pi)}.$$

Theorem 5.1 follows from a direct application of Theorem 4.1 and is proved in Section 8.3. Suppose that the nonreversible dynamics is determined by (1.11) where $\gamma(x) = \beta \widetilde{\gamma}(x)$, for $\beta \in \mathbb{R}$ and for some smooth vector field $\widetilde{\gamma}$. If $\Psi_{\Delta t}$ is an integrator for the flow with error order r, then it is straightforward to show that $\Psi_{\Delta t}$ will satisfy a modified backward equation of the form (4.3) where the vector fields f_j satisfy the scaling $f_j = |\beta|^{j+1} \widetilde{f}_j$, with $\|\widetilde{f}_j\|_{L^2(\pi)} \sim O(1)$ for $j=0,\ldots,r-1$. It follows that if the conditions of Theorem 5.1 hold, then the leading order term of the bias is of the form $C\Delta t^r |\beta|^{r+1}$, where C is independent of Δt and β . This estimate provides a

rule of thumb for choosing the magnitude of the nonreversible perturbation β . Clearly, this should be as large as is possible while maintaining a given tolerance ϵ for the bias. To this end, for $\Delta t \ll 1$, β must satisfy

$$|\beta| \simeq \epsilon^{\frac{1}{r+1}} \Delta t^{-\frac{r}{r+1}}.$$

In particular, assuming that $|\beta| \simeq \Delta t^{-\kappa}$ where $\kappa \in \mathbb{R}$, we obtain an upper bound

$$\kappa \le -\frac{1}{r+1} \frac{\log \epsilon}{\log \Delta t} + \frac{r}{r+1}. \tag{5.3}$$

For $\epsilon \asymp \Delta t$, this rule suggests that β should have been chosen to be O(1) with respect to Δt if a first order integrator is used to simulate the nonreversible dynamics. Employing a higher order integrator however, permits larger values of $|\beta|$, in particular $|\beta| \asymp \Delta t^{-0.6}$ for a fourth order scheme as considered in the examples of Section 7. We emphasise that unless we have explicit control on the growth of the remainder term in (4.5) as a function of β , then (5.3) is only heuristic. Moreover, we are assuming that the integrator $\Psi_{\Delta t}$ is stable for this parameter regime. In practice, the stiffness of the ODE (1.11) would impose additional constraints on β .

5.2 Asymptotic variance of the splitting scheme

Contrary to Theorem 5.1 we shall focus on the case of MALA for the integrator $\Theta_{\Delta t}$ for which we are able to verify that Assumption 4.3 holds. As before we shall assume that the integrator $\Phi_{\Delta t}$ for the nonreversible flow satisfies the following expansion

$$\Phi_{\Delta t}\phi = \phi + \Delta t \mathcal{A}_1 \phi + \Delta t^2 \mathcal{A}_2 \phi + \Delta t^3 R_{\phi}, \quad \phi \in C^{\infty}(\mathbb{T}^d),$$

where $A_1 = \gamma(x) \cdot \nabla$ is the antisymmetric part of \mathcal{L} in $L^2(\pi)$ and $R_{\phi} \in C^{\infty}(\mathbb{T}^d)$ is bounded independently of Δt . We make the following assumption.

Assumptions 5.3. The numerical flow Φ_h is a consistent scheme for (1.10) and that there exists $\Delta t_0 > 0$ and L > 0 independent of Δt such that

$$|\Phi_{\Delta t}(z_1) - \Phi_{\Delta t}(z_2)| \le L|z_1 - z_2|, \qquad z_1, z_2 \in \mathbb{T}^d,$$
 (5.4)

for all $\Delta t < \Delta t_0$.

Provided that (5.4) holds, Theorem A.2 in the Appendix implies that the reversible integrator $\Theta_{\Delta t}$ satisfies the following perturbation expansion

$$\Theta_{\Delta t}\phi = \phi + \Delta t \mathcal{G}_1 \phi + \Delta t^2 \mathcal{G}_2 \phi + \Delta t^{5/2} R_{\phi}, \quad \phi \in C^{\infty}(\mathbb{T}^d), \tag{5.5}$$

where $\mathcal{G}_1 = \mathcal{S}$ is the symmetric part of \mathcal{L} in $L^2(\pi)$ and \mathcal{G}_2 is given by (A.4), and R_{ϕ} is a smooth remainder term bounded independently with respect to Δt . The following theorem then characterises the asymptotic variance of the Lie-Trotter splitting scheme (1.10) when the reversible dynamics are integrated with MALA. It is a direct application of Theorem 4.6 and is proved in Section 8.4.

Theorem 5.4. Consider the Lie-Trotter splitting scheme defined by (1.10) where $\Theta_{\Delta t}$ is integrated using MALA and suppose that the nonreversible dynamics preserves the invariant distribution up to order 2 and satisfies Assumption 5.3. Then for all $f \in C^{\infty}(\mathbb{T}^d)$ we have

$$\widehat{\sigma}_{\Delta t}^{2}(f) = \sigma^{2}(f) + \Delta t \left\langle (-\mathcal{L})^{-1} (\mathcal{L}^{2} - 2(\mathcal{A}_{2} + \mathcal{G}_{1}\mathcal{A}_{1} + \mathcal{G}_{2})(-\mathcal{L})^{-1} (f - \pi(f), f - \pi(f)) \right\rangle_{\pi} + o(\Delta t).$$

If moreover, the nonreversible dynamics is integrated using a second order scheme then the $O(\Delta t)$ term can be written as

$$\left\langle (-\mathcal{L})^{-1} \left((\mathcal{S}^2 - 2\mathcal{G}_2) + [\mathcal{S}, \mathcal{A}] \right) (-\mathcal{L})^{-1} (f - \pi(f), f - \pi(f) \right\rangle_{\pi},$$

where S and A are the symmetric and antisymmetric parts of L in $L^2(\pi)$, respectively.

From the point of view of tuning the nonreversible Langevin sampler defined by (1.10) the main conclusion of Theorem 5.4 is that, for Δt sufficiently small, the asymptotic variance of (1.10) is, to leading order, equal to the asymptotic variance of the exact dynamics (1.2). In particular, given an observable f, this result implies that a choice of flow γ which reduces the variance of a sampler based on (1.2) will have a similarly beneficial effect on (1.10). One can thus leverage the theory detailed in [7] and [20] to design efficient samplers for a given target distribution π and observable f.

Gaussian target distributions 6

In Sections 5.1 and 5.2, the asymptotic bias and variance for estimators based on Lie-Trotter splitting scheme (1.10) were characterised in terms of stepsize Δt and magnitude of the nonreversible perturbation β . This detailed analysis was however restricted to the case of \mathbb{T}^d -valued diffusions, as a similar analysis for \mathbb{R}^d would be significantly more involved (see for example [17]). To demonstrate that analogous expressions for the asymptotic variance and bias can be derived in the \mathbb{R}^d case, in this section we consider the class of linear SDEs given by

$$dX_t = -AX_t dt + dW_t (6.1)$$

where $X_t \in \mathbb{R}^d$, W_t is a standard d-dimensional Brownian motion.

In the case where -A is stable the dynamics generated by (6.1) are ergodic with respect to $\mathcal{N}(0, \Sigma_{\infty})$ where Σ_{∞} satisfies the Lyapunov equation [9]:

$$A\Sigma_{\infty} + \Sigma_{\infty}A^T = I. \tag{6.2}$$

We shall consider a vector field γ satisfying (2.2) which is given by

$$\gamma(z) = \beta J A z,$$

where J is a skew symmetric matrix, and β is a free parameter. Hence (1.2) becomes

$$dX_t = -(I - \beta J)AX_t dt + dW_t. \tag{6.3}$$

The fact that equation (6.3) is linear implies that is amenable to very detailed analysis, as for certain classes of numerical schemes, one can find another linear SDE that the numerical method solves exactly in the weak sense. We explain this idea further in Section 6.1, while in Section 6.2 we extend the formula for the asymptotic variance from [7] to linear diffusions with a general positive definite diffusion tensor. This allows the use of the modified equation analysis presented in Section 6.1 not just to study the infinite time bias of numerical schemes applied to (6.3), but also the asymptotic variance. This is discussed further in Section 6.3 in the context of a simple two dimensional example.

6.1 **Exact modified equation**

Consider a one step method applied to (6.3)

$$\widehat{X}_{n+1}^{\Delta t} = B(\Delta t)\widehat{X}_n^{\Delta t} + f(\Delta t, \omega), \quad \widehat{X}_0^{\Delta t} = x_0, \tag{6.4}$$

where $f(\Delta t, \omega)$ is the flow map for the noise process and $B(\Delta t) \in \mathbb{R}^{d \times d}$ satisfies B(0) = I. For an Euler-Maruyama discretisation of (6.3),

$$B(\Delta t) = I - \Delta t (I - \beta J) A,$$

$$f(\Delta t, \omega) = \sqrt{\Delta t} \xi,$$

where $\xi \in \mathbb{R}^d$ satisfies $\xi \sim \mathcal{N}(0,I)$. The fact that (6.4) remains linear imply that the solution $\widehat{X}_n^{\Delta t}$ remains Gaussian at all times, assuming a deterministic initial condition x_0 . This implies [49], that the numerical solution (6.4) satisfies exactly in the weak sense at all times the following stochastic differential equation

$$d\widetilde{X}_t = \widetilde{B}\widetilde{X}_t + \widetilde{\Sigma}^{1/2}dW_t \tag{6.5}$$

where $\widetilde{B} \in \mathbb{R}^{d \times d}$ and $\widetilde{\Sigma} \in \mathbb{R}^{d \times d}_{sym}$ are defined by

$$\widetilde{B} = \frac{\log(B(\Delta t))}{\Delta t},$$
 (6.6a)

$$\widetilde{B} = \frac{\log(B(\Delta t))}{\Delta t}, \tag{6.6a}$$

$$B(\Delta t)\widetilde{\Sigma}B(\Delta t)^T - \widetilde{\Sigma} = \widetilde{B}L + L\widetilde{B}^T, \tag{6.6b}$$

where $L = \mathbb{E}(ff^T)$. For sufficiently small Δt one can show that (6.5) is ergodic with respect to $\mathcal{N}(0, \widetilde{K})$ where K satisfies a Lyapunov equation similar to (6.2). Thus, by solving this equation we can obtain an expression for the invariant measure that the numerical scheme is ergodic with respect to, and hence have an explicit expression for the asymptotic bias of the numerical method. We study this further in Section 6.3, in the context of a two dimensional example.

6.2 Asymptotic variance

By extending the results from [7] one can calculate the asymptotic variance for (6.5). In particular if we consider the SDE (6.1) our objective is to derive an explicit expression for the asymptotic variance $\sigma^2(f)$ of

$$I_t = \frac{1}{t} \int_0^t f(\widetilde{X}_s) \, ds,$$

where f is a function of the form

$$f(x) = x \cdot Mx + L \cdot x + K,$$

for some $M \in \mathbb{R}^{d \times d}_{sym}$, $L \in \mathbb{R}^d$ and $K \in \mathbb{R}$. In particular we have the following proposition

Proposition 6.1. Consider the linear diffusion defined by the SDE,

$$dX_t = -AX_t dt + \sigma dW_t,$$

where W_t is a m-dimensional Brownian motion, $\sigma \in \mathbb{R}^{d \times m}$ such that $\Sigma = \sigma \sigma^{\top}$ is positive definite and -A is stable. Then, for

$$f(x) = x \cdot Mx + L \cdot x + K,\tag{6.7}$$

the asymptotic variance $\sigma^2(f)$ is given by

$$\sigma^2(f) = 2Tr \left[\left(\int_0^\infty e^{-A^\top t} M e^{-At} \, dt \right) M_\Sigma \right] + 2L_\Sigma \cdot A^{-1} \Sigma_\infty A^{-\top} L_\Sigma,$$

where $M_{\Sigma} = \Sigma^{1/2} M \Sigma^{1/2}$ and $L_{\Sigma} = \sqrt{2} \Sigma^{-1/2} L$.

Proof. The proof of this proposition can be found in Appendix B.

6.3 Example

We now consider the linear diffusion (6.3) where

$$A = \left(\begin{array}{cc} \alpha & 0 \\ 0 & \alpha \end{array}\right),$$

for which we know that the stationary covariance satisfies

$$\Sigma_{\infty} = \begin{pmatrix} \frac{1}{2\alpha} & 0\\ 0 & \frac{1}{2\alpha} \end{pmatrix}$$

We now study the properties of integrators where the $\Phi_{\Delta t}$ and $\Theta_{\Delta t}$ in (1.10) are given by

$$\Phi_{\Delta t}(z) = \left(I + \Delta t \beta J A + \frac{\Delta t^2}{2} (\beta J A)^2 + \dots + \frac{\Delta t^p}{p!} (\beta J A)^p\right) z \tag{6.8a}$$

$$\Theta_{\Delta t}(z) = e^{-A\Delta t}z + \sigma_{\Delta t}\xi \tag{6.8b}$$

where

$$\sigma_{\Delta t} \sigma_{\Delta t}^T = \int_0^{\Delta t} e^{-A(\Delta t - s)} \sum e^{-A^T(\Delta t - s)} ds = \frac{1}{2\alpha} \left[1 - e^{-2\alpha \Delta t} \right] I.$$

More precisely we solve the reversible part of the dynamics exactly, while we apply a Taylor-based method of order p to the nonreversible part of the dynamics. We note here that the exact solution of the reversible part of the dynamics is only possible because the dynamics are linear. A further consequence of the linearity of the dynamics is that it is possible to conserve the invariant measure for the reversible part with using the θ method with $\theta = \frac{1}{2}$, see [1]. Hence we will also consider the integrator $\widetilde{\Theta}_{\Delta t}(z)$ given by

$$\widetilde{\Theta}_{\Delta t}(z) = \left(I + \frac{\Delta t}{2}A\right)^{-1} \left[\left(I - \frac{\Delta t}{2}A\right)z + \sqrt{\Delta t}\xi \right]$$
(6.9)

The other interesting feature of (6.9) is that even though not exact like (6.8), when metropolised, proposals generated from by (6.9) will be accepted almost surely. For nonlinear problems, the reversible dynamics cannot be integrated exactly, and

it is impossible to construct an exact solution and (6.9) does not conserve the invariant measure. Hence one would replace these integrator with one that conserves the invariant measure by introducing a Metropolisation step, and Theorem 5.1 would still hold.

	Reversible first	Non reversible first
p = 1	$\frac{\left(1 - e^{-2\alpha\Delta t}\right)\left(1 + \alpha^2\beta^2\Delta t^2\right)}{2\alpha\left[1 - e^{-2\alpha\Delta t}\left(1 + \alpha^2\beta^2\Delta t^2\right)\right]}$	$\frac{1 - e^{-2\alpha\Delta t}}{2\alpha[1 - e^{-2\alpha\Delta t}(1 + \alpha^2\beta^2\Delta t^2)]}$
p = 2	$\frac{\left(1 - e^{-2\alpha\Delta t}\right)\left(\alpha^4\beta^4\Delta t^4 + 4\right)}{2\alpha(e^{-2\alpha\Delta t}\left(\alpha^4\beta^4\Delta t^4 + 4\right) - 4)}$	$\frac{2(1-e^{-2\alpha\Delta t})}{\alpha[4-e^{-2\alpha\Delta t}(4+\alpha^4\beta^4\Delta t^4)]}$

Table 1: Numerical invariant measure when the reversible part is solved exactly

	Reversible first	Non reversible first
p=1	$\frac{4+4\alpha^2\beta^2\Delta t^2}{8\alpha-4\alpha^2\beta^2\Delta t+4\alpha^3\beta^2\Delta t^2-\alpha^4\beta^2\Delta t^3}$	$\frac{4}{8\alpha - 4\alpha^2\beta^2\Delta t + 4\alpha^3\beta^2\Delta t^2 - \alpha^4\beta^2\Delta t^3}$
p=2	$\frac{4(4+\alpha^4\beta^4\Delta t^4)}{\alpha(32-\alpha^3\beta^4\Delta t^3(2-\alpha\Delta t)^2)}$	$\frac{16}{\alpha(32-\alpha^3\beta^4\Delta t^3(2-\alpha\Delta t)^2)}$

Table 2: Numerical invariant measure when the reversible part is solved by θ -method for $\theta = \frac{1}{2}$.

Study of the invariant measure bias We now study the properties of the numerical invariant measure using (6.6). We use Mathematica to symbolically calculate the solutions to (6.6) and then obtain an expression for the numerical invariant measure, when a first and a second order numerical method is used to solve the nonreversible part of the diffusion. In particular, in Tables 1,2 we present exact expressions for the numerical invariant measure based on the Lie-Trotter splitting (1.10), for different ordering of the splitting and different choices of integrators for the reversible and nonreversible part. Furthermore, in Figure 1 we plot the 2-norm of the difference between the covariance matrix of the numerical method and the true covariance matrix Σ_{∞} when the nonreversible part is solved first and then the θ -method with $\theta = 1/2$ is used for the reversible part². As we can see the order of convergence is always odd. This was also observed in [2] and it relates with the fact that for the deterministic methods used here, the coefficient f_p in Theorem 5.1 is always zero when p is even hence giving the extra order of convergence observed in Figure 1. Additionally in Figure 2 we plot the asymptotic bias of Δt when a numerical integrator of order 1 is used to solve the nonreversible part for different values of β . As we can see, the larger the value of β the larger the asymptotic bias.




Figure 1: Accuracy of the numerical invariant measure (covariance matrix error) of the Lie-Trotter splitting The lines corresponds to explicit deterministic integrators for the nonreversible part of orders $p=1,2,3,\ldots,9$ (from top to bottom), respectively. The orders of accuracy for the invariant measure are always odd.

²We have not included any of the other possible combinations of ordering of splitting and numerical integrators for the reversible part as the results are qualitatively the same




Figure 2: Accuracy of the numerical invariant measure (covariance matrix error) of the Lie-Trotter splitting for different values of β when a first order numerical method is used for solving the nonreversible part.

Study of the asymptotic variance We now study the properties of the asymptotic variance using (6.6). In particular the idea is that since our numerical solution satisfies exactly in the weak sense the corresponding modified equation then it is enough to look at Proposition 6.1 where A and σ are now replaced with the modified coefficients (6.6). Similarly to the case of the invariant measure bias we use Mathematica to symbolically calculate the solutions to (6.6) and then obtain an expression for the asymptotic variance, when a first and a second order numerical method is used to solve the nonreversible part of the diffusion. In particular, we take K=0, L=0 and M the two-by-two identity in (6.7) we find that when the reversible part is solved exactly that for p=1, we have

$$\widetilde{\sigma}_{\Delta t}^{2}(f) = \frac{2+\beta^{2}}{2\alpha\left(1+\beta^{2}\right)} + \frac{\left(2\beta^{2}+\beta^{4}+\beta^{6}\right)\Delta t}{4\left(1+\beta^{2}\right)^{2}} + \mathcal{O}(\Delta t^{2}),$$

independently of the ordering of the splitting, while for p = 2 we have

$$\widetilde{\sigma}_{\Delta t}^2(f) = \frac{2+\beta^2}{2\alpha\left(1+\beta^2\right)} - \frac{\alpha\beta^4\Delta t^2}{6\left(1+\beta^2\right)^2} + \mathcal{O}(\Delta t^3),$$

again independently of the ordering of the splitting. The expressions above change to

$$\widetilde{\sigma}_{\Delta t}^{2}(f) = \frac{2+\beta^{2}}{2\alpha\left(1+\beta^{2}\right)} + \frac{\left(2\beta^{2}+\beta^{4}+\beta^{6}\right)\Delta t}{4\left(1+\beta^{2}\right)^{2}} + \mathcal{O}(\Delta t^{2}),$$

for p = 1, and

$$\widetilde{\sigma}_{\Delta t}^{2}(f) = \frac{2 + \beta^{2}}{2\alpha (1 + \beta^{2})} - \frac{\alpha(-\beta^{2} + 2\beta^{4})\Delta t^{2}}{12 (1 + \beta^{2})^{2}} + \mathcal{O}(\Delta t^{3}),$$

when the reversible part of the dynamics is solved by the θ -method for $\theta=1/2$, again independently of the ordering of the splitting. We note here that these results agree with Proposition 4.8, since for p=1 the leading order perturbation in terms of the continuous time variance is $\mathcal{O}(\Delta t)$ while for p=2 is $\mathcal{O}(\Delta t^2)$.


Mean Square Error Having obtained analytical expressions for the asymptotic bias of the invariant measure as well as for the asymptotic variance of the corresponding numerical schemes, we combine them in order to study the mean square error. More precisely, decomposing the MSE into bias and variance,

$$\mathbb{E}|\widehat{\pi}_T(f) - \pi(f)|^2 = (\mathbb{E}\widehat{\pi}_T(f) - \pi(f))^2 + \mathbb{E}(\widehat{\pi}_T(f) - \mathbb{E}\widehat{\pi}_T(f)) = \widehat{\mu}_T^2 + \widehat{\sigma}_T^2,$$

we approximate $\hat{\mu}_T$ by the invariant measure bias, while on the other hand

$$\widehat{\sigma}_T^2 \simeq \frac{\widehat{\sigma}^2(f)}{T}.$$

We now plot in Figure 3 the MSE when a first and a second order numerical method is used to solve the nonreversible part and the reversible part is solved exactly. In particular, we choose our timestep $\Delta t = 10^{-4}$, $\alpha = 1$, $T = 10^3$ and we study the influence of β on the MSE. As can be seen in both cases there is a range of values of the parameter β for which the MSE is reduced almost to $\frac{1}{2}$ which is the theoretical minimal variance attainable using this choice of dynamics [7]. Increasing the magnitude of β beyond this point, eventually the bias term will dominate the mean-square error which will rapidly increase. Using a second order integrator for the nonreversible dynamics mitigates this increase in bias, and a significant reduction in MSE is possible for a much wider range of β . Indeed, in Figure 3 we see that, in this case, the asymptotic bias is $\mathcal{O}(\Delta t^3)$ and will not dominate the MSE for a wider range of β values.






Figure 3: MSE for two different methods applied to the nonreversible part.

7 Numerical experiments

In this section, we perform a number of different numerical investigations that illustrate the superiority of the nonreversible Langevin samplers over standard Metropolis-Hastings algorithms for a fixed computational budget. In particular, we define computational cost here in terms of number of density evaluations which is the dominating cost in high dimensions. To this end we ensure that every comparison is made for the same computational cost, *i.e.*, same number of density evaluations.

7.1 Warped Gaussian distribution

As a first numerical we consider the expectation of an observable with respect to the following two dimensional distribution

$$\pi(x) \propto \exp\left(-\frac{x_1^2}{100} - (x_2 + bx_1^2 - 100b)^2\right)$$
 (7.1)

where $x=(x_1,x_2)$. The parameter b>0 controls the degree of warpedness, and is chosen to be b=0.05. The log density is plotted in Figure 4a. Our objective is to estimate $\pi(f)$ where $f(x)=|x|^2$. The nonreversible flow γ is chosen as follows:

$$\gamma(x) = J\nabla \log \pi(x), \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

In Figure 4, we plot characteristic trajectories of MALA as well its nonreversible counterpart (for $\beta=25$) starting from the initial point x=(15,2). The figure suggests superior mixing of the nonreversible samplers, which improves further with increasing β values. In Figure 5 the mean-square error is plotted as a function of




Figure 4: Typical trajectories for MALA and Lie-Trotter splitting scheme applied to the warped Gaussian distribution (7.1), with computational budget of 3200 density evaluations. Both schemes started from x=(15,2) depicted by a blue dot.

stepsize for different values of flow strength β . The reversible part of the Lie-Trotter scheme is simulated using MALA, RWMH and Barker rule in Figures 5a,5b and 5c, respectively. The "exact" value of $\pi(f)$ used to compute the MSE is obtained via adaptive Gaussian quadrature, accurate up to 10^{-10} . In accordance with the results of Theorems 5.1 and 5.4, the MSE is a tradeoff between bias and variance. For a fixed computational budget as Δt decreases, the bias arising from the discretisation of the nonreversible flow decreases. However, the variance simultaneously increases as the total simulated time $T=N\Delta t$ is reduced. This competion between bias and variance suggest an optimal choice of timestep Δt which minimises the MSE. This tradeoff is further exacerbated when β is increased. Nevertheless, for an appropriate choice of β the MSE can be up to an order of magnitude lower than that of MALA, at the same computational cost.




Figure 5: Comparison of the MSE between MALA and different nonreversible samplers applied to the warped Gaussian distribution (7.1). The computational budget is set to $N=3.5\cdot 10^3$ density evaluations, and 4^{th} order Runge-Kutta method is used for the nonreversible component.

7.2 Logistic Regression

Let X be a $m \times d$ design matrix comprising m samples with d covariates and a binary response variable $Y \in \{-1,1\}^m$. A Bayesian logistic regression model of the binary response is obtained by the introduction of the regression coefficient $\theta \in \mathbb{R}^d$. For the sake of exposition, we shall assume a Gaussian prior of θ , i. e., $\beta \sim \mathcal{N}(0,\Sigma)$. The posterior distribution $\pi(\theta|X,Y)$ is given by

$$\pi(\theta|(X,Y)) \propto \exp\left(\sum_{i=1}^{m} Y_i \theta^T X_i - \log\left(1 + e^{\theta^T X_i}\right) - \frac{1}{2} \theta^T \Sigma^{-1} \theta\right)$$
 (7.2)

In Figure 6 we investigate the use of the Lie Trotter sampler applied to this problem for the Pima indians 3 dataset obtained from the UCI machine learning repository. The skew symmetric matrix J is chosen by generating a random permutation $\sigma(1), \ldots, \sigma(d)$ and setting

$$J_{\sigma(i),\sigma(i+1)}=1$$
 and $J_{\sigma(i+1),\sigma(i)}=-1,$

for $i=1,\ldots,d-1$, and zero elsewhere. In Figure 6a we plot the first estimator $\widehat{\pi}_T^{\Delta t}(\theta_1)$ with 95% confidence intervals for different values of β and stepsize. Each point in the plot cost $3.5\cdot 10^3$ density evaluations. To provide a comparison against the truth, an optimally tuned MALA scheme was integrated over 10^7 timesteps. In Figure 6b we plot the effective sample size (ESS) of the Lie-Trotter scheme for different values of β and Δt . The markers denote the median value of the ESS with the markers denoting the 5% and 95% percentiles. We note however that there typically be a very small number of observables for which the nonreversible scheme offers no advantage. This agrees with the theory detailed in [7] which characterises the minimum attainable variance reduction in terms of the projection of the observable f on the nullspace of the operator $J\nabla V(x)\cdot\nabla$. As J is chosen randomly, there will always been a number of observables which are close to this subspace, and thus the nonreversible dynamics offer no advantage. One possible remedy around this is to periodically resample the nonreversible matrix J, but we do not investigate this here.




Figure 6: Confidence Interval for the first covariate and ESS for estimators for π (θ_i), i = 1, ..., 9 for logistic regression of the Pima Indians data set. Each data point in these plots is set to $3.5 \cdot 10^3$ density evaluations. The results are compared to an optimally tuned MALA simulation run for 10^7 density evaluations.

7.3 Spatial model

We now consider a high dimensional target distribution related to inference for a log-Gaussian Cox point process previously considered in [33]. In particular, given the location of 126 Scots pine saplings in a natural forest in Finland, we wish to infer the average intensity of a corresponding Poisson point process. Following [4], we consider a discretised version of the model where the spatial region is discretised to a 64×64 regular grid. For each

 $^{^{3}}$ Here m = 768, d = 9.

i,j $X_{i,j}$ is the random variable counting the number of observations in the (i,j)-cell and hence the dimension of the problem is $d=64^2=4096$. The observations are assumed to be generated by a Poisson point process with unobserved intensity $\Lambda_{i,j}, i,j=1,\cdots,64$. Given the $\Lambda_{i,j}$ the random variables $X_{i,j}$ are assumed to be conditional independent with Poisson distributed mean $m\Lambda_{i,j}$, where m=1/4096 is the area of a single cell. We impose a log-Gaussian prior on $\Lambda_{i,j}$, more specifically

$$\Lambda_{i,j} = \exp(Y_{i,j})$$

where $Y = (Y_{i,j}, i, j = 1, \cdots 64) \sim \mathcal{N}(\mu \mathbf{1}, \Sigma)$ where

$$\Sigma_{i,j,i',j'} = \sigma^2 \left[-\frac{-\{(i-i')^2 + (j-j')^2\}^{1/2}}{64\beta} \right], \quad i,j,i',j' = 1 \cdots, 64.$$

The posterior distribution is thus given by

$$f(y|x) \propto \prod_{i,j=1}^{64} \exp\{(x_{i,j}y_{i,j}) - m\exp(y_{i,j})\} \exp\{-0.5(y-\mu\mathbf{1})^T\Sigma^{-1}(y-\mu\mathbf{1})\}$$

Due to the poor scaling of the posterior distribution in [4] a reparametrization of y is introduced to improve the mixing of the Metropolis-Hastings scheme. This procedure is expensive with a computational cost of $\mathcal{O}(d^3)$. However, in the case of the nonreversible samplers, the nonreversible perturbation compensates for the poor scaling, thus rendering this reparametrisation unnecessary.

In Figures 7 we plot an estimator of $\mathbb{E}(\Lambda \mid x)$ using MALA and its nonreversible counterpart respectively. For this computation the skew-symmetric matrix J was generated randomly as in the logistic regression example. Due to the large number of covariates, for any given random choice of J, there would be a small number of covariates for which the nonreversible scheme does not offer significant advantage over MALA, as described in [7]. To better understand the effect the nonreversible flow for an average covariate, we thus generate 10 independent random skew-symmetric matrices, and compute the average ESS over J. The results are presented in Figure 8. In Figure 8c a histogram of the ESS over all covariates is plotted for both MALA and the splitting scheme for specific choices of Δt and β . We observe that the ESS for the nonreversible scheme is orders of magnitude better than MALA. To illustrate the dependence of ESS on timestep, similarly to the case of logistic regression, in Figure 8b we plot the median ESS for different choices of timestep. It is clear that increasing β and Δt as much as possible increases the ESS. However, this comes at the cost of increasing bias as can be observed in Figure 8a. Nonetheless, it is evident that the nonreversible sampler significantly outperforms the MALA scheme.




Figure 7: Average inferred Poisson intensity using the different schemes. The computational budget is set to $N=3.5\cdot 10^3$ gradient evaluations.




Figure 8: Results for the inference of the log-Gaussian cox process. The computational budget is set to $N=3.5\cdot 10^3$ density evaluations. A reference MALA simulation run for 10^7 density evaluations is provided for comparison.

8 Proofs of the main results

In this section we prove the main results of the paper. In particular, in Section 8.1 we prove the geometric ergodicity of the splitting scheme (1.10), while in Sections 8.3 and 8.4 we prove the results related to the asymptotic bias and variance of the splitting method.

8.1 Ergodicity of the splitting scheme

Here we prove the theorems and corollaries stated in Section 3.

Proof of Theorem 3.2. We verify the criteria for geometric ergodicity formulated in Chapters 15 and 16 of [29].

1. We show that $P_{\Delta t}(x,\cdot)$ is π -irreducible. Let $A \subset \mathbb{R}^d$ such that $\pi(A) > 0$, then

$$P_{\Delta t}(x,A) = \int_A q_{\Delta t}(y | \Phi_{\Delta t}(y)) \alpha(\Phi_{\Delta t}(x), y) dy$$
$$+ \mathbf{1}_A(x) \int_{\mathbb{R}^d} q_{\Delta t}(z | \Phi_{\Delta t}(x)) (1 - \alpha(z, \Phi_{\Delta t}(x))) dz > 0,$$

which implies that $P_{\Delta t}$ is π -irreducible.

2. We now show that every compact set C of positive measure is small. To this end, let C be such a set and B a measurable subset of C. Then $D = C \cup \Phi_{\Delta t}(C)$ is also a compact set of positive measure. Since the target density π and proposal $q_{\Delta t}(y|x)$ are positive and continuous for all x, y, applying [26, Lemma 1.2] implies that there exists $\eta > 0$ such that

$$\widetilde{P}_{\Delta t}(x,B) \ge \eta \pi(B), \quad B \subset D, x \in D.$$

In particular,

$$\widehat{P}_{\Delta t}(x,B) = \widetilde{P}_{\Delta t}(\Phi_{\Delta t}(x),B) \ge \eta \pi(B), \quad B \subset C, x \in C, \tag{8.1}$$

so that C is small. Aperiodicity of the chain follows immediately from (8.1).

3. To complete the proof we show that $P_{\Delta t}$ satisfies a Foster-Lyapunov condition for the Lyapunov function V. To this end using (3.2), given $x \in \mathbb{R}^d$:

$$\begin{split} \widehat{P}_{\Delta t} V(x) &\leq \lambda V(\Phi_{\Delta t}(x)) + b \mathbf{1}_C(\Phi_{\Delta t}(x)) \\ &\leq \lambda V(\Phi_{\Delta t}(x)) + b \mathbf{1}_{\Phi_{\Delta t}^{-1}(C)}(x) \\ &\leq \lambda V(x) + \lambda \left(V(\Phi_{\Delta t}(x)) - V(x) \right) + b \mathbf{1}_{\Phi_{\Delta t}^{-1}(C)}(x). \end{split}$$

By Assumptions 3.1(2) and 3.1(3), there exists a compact set $D \subset \mathbb{R}^d$ and 0 < c < 1 such that $\Phi_{\Delta t}^{-1}(C) \subset D$ and moreover

$$\lambda V(x) + \lambda \left(V(\Phi_{\Delta t}(x)) - V(x) \right) \le cV(x), \quad x \in \mathbb{R}^d \setminus D,$$

which implies that

$$\widehat{P}_{\Delta t}V(x) \le cV(x) + b\mathbf{1}_D(x),$$

as required.

Proof of Corollary 3.3. Provided that the conditions of [40, Theorem 4.1] hold, then the MALA chain satisfies a Foster-Lyapunov condition for $V(x) = e^{s|x|}$ for s > 0 sufficiently small. If we consider

$$\frac{V(\Phi_{\Delta t}(x)) - V(x)}{V(x)} = e^{s(|\Phi(x)| - |x|)} \le e^{s(|\Phi(x)| - |x|)},$$

then Assumption 3.1(2) follows immediately. Finally, we note that (3.5) implies that there exists K > 0 such that

$$|\Phi_{\Delta t}(x)| - K \le |x| \le |\Phi_{\Delta t}(x)| + K, \quad x \in \mathbb{R}^d,$$

from which 3.1(3) follows immediately. Hence, the conditions of Theorem 3.2 all hold, and thus the process $\widehat{X}_n^{\Delta t}$ is geometrically ergodic.

Suppose now that $\gamma = J \nabla \pi^{\alpha}$, where $J = -J^{\top}$ and $\alpha > 0$. Suppose $\Phi_{\Delta t}(x)$ is an explicit Runge-Kutta discretisation of the nonreversible dynamics having s stages. Then we can write

$$\Phi_{\Delta t}(x) = x + h \sum_{i=1}^{s} b_i k_i(x),$$
(8.2)

where

$$k_1(x) = \gamma(x)$$

$$k_2(x) = \gamma (x + hw_{2,1}k_1(x))$$

$$k_3(x) = \gamma (x + h(w_{3,1}k_1(x) + w_{3,2}k_2(x)))$$

$$\vdots$$

$$k_s(x) = \gamma \left(x + h \sum_{i=1}^{s-1} w_{s,i} k_i(x) \right),$$

where $(w_{i,j})$ is the Runge-Kutta matrix associated with the discretisation. By (3.6) there exist positive constants α' , K' and K_1 such that

$$k_1(x) \le |\gamma(x)| \le K' |\nabla \pi^{\alpha}(x)| \le K_1 \pi^{\alpha'}(x).$$

Suppose now that there exists constants K_2, \ldots, K_{i-1} such that

$$|k_j(x)| \le K_j \pi^{\alpha'}(x), \quad x \in \mathbb{R}^d, \quad j = 1, \dots, i - 1.$$

By (3.6) the matrix $\nabla \gamma = (\partial_{x_i} \gamma_j(x))_{i,j}$ has bounded components in \mathbb{R}^d and so applying the mean value theorem to every component of γ , it follows that

$$|k_i(x)| \le |\gamma(x)| + h\left(\sup_{x \in \mathbb{R}^d} |\nabla \gamma(x)|_{max}\right) \sum_{j=1}^{i-1} |w_{i,j}| |k_j(x)| \le K_i \pi^{\alpha'}(x),$$

for some constant K_i . It follows by induction that $|\Phi_{\Delta t}(x) - x| \le K\pi^{\alpha'}(x)$, for all $x \in \mathbb{R}^d$, which implies (3.5). The corresponding result for γ given by (2.7) follows similarly.

8.2 Asymptotic variance of numerical integrators

Here we prove Proposition 4.5 and Theorem 4.6 which characterises the error in the asymptotic variance for an arbitrary numerical integrator

Proof of Proposition 4.5. It follows from standard elliptic regularity that the operator $(-\mathcal{L})^{-1}$ is bounded on $L_0^{\infty}(\pi)$. Similarly the operator $\Delta t(I-P_{\Delta t})^{-1}$ is bounded on $L_0^{\infty}(\pi)$, uniformly with respect to Δt .

Let $\psi \in C^{\infty}(\mathbb{T}^d)$ with $\pi(\psi) = 0$. There exists R_{ψ} , smooth and bounded uniformly with respect to Δt such that

$$\left(\frac{I - P_{\Delta t}}{\Delta t}\right)\psi = -\mathcal{L}\psi - \frac{\Delta t}{2}\mathcal{L}^2\psi - \frac{\Delta t^2}{6}\mathcal{L}^3\psi + \Delta t^3R_{\psi},$$
(8.3)

provided that Δt is sufficiently small. Hence using (8.3) we obtain

$$(-\mathcal{L})^{-1}\psi = \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} \left(\frac{I - P_{\Delta t}}{\Delta t}\right) (-\mathcal{L})^{-1}\psi$$

$$= \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} \psi + \frac{\Delta t}{2} \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} \mathcal{L}\psi + \frac{\Delta t^{2}}{6} \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} \mathcal{L}^{2}\psi$$

$$+ \Delta t^{3} \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} R_{(-\mathcal{L})^{-1}\psi}.$$
(8.4)

Since both sides of (8.3) has mean zero and $\pi(\mathcal{L}^i\psi) = 0$ for $i \geq 0$, it follows that $\pi(R_{(-\mathcal{L})^{-1}\psi}) = 0$. Thus the remainder term in (8.4) is well-defined and uniformly bounded with respect to Δt .

Now let $f \in C^{\infty}(\mathbb{T}^d)$, then similar to (4.9), the asymptotic variance of the estimator $N^{-1}\sum_{n=0}^{N-1} f(X_n^{\Delta t})$ for the discretized exact process is given by

$$\sigma_{\Delta t}^{2}(f) = 2 \left\langle \left(\frac{I - P_{\Delta t}}{\Delta t} \right)^{-1} (f - \pi(f)), f - \pi(f) \right\rangle_{\pi} - \Delta t \operatorname{Var}_{\pi}[f].$$

By (8.4) it follows that

$$\begin{split} \sigma_{\Delta t}^2(f) &= 2 \left< (-\mathcal{L})^{-1}(f - \pi(f)), f - \pi(f) \right>_{\pi} \\ &+ \Delta t \left< \left(\frac{I - P_{\Delta t}}{\Delta t} \right)^{-1} (-\mathcal{L})(f - \pi(f)), f - \pi(f) \right>_{\pi} \\ &- \frac{\Delta t^2}{3} \left< \left(\frac{I - P_{\Delta t}}{\Delta t} \right)^{-1} (-\mathcal{L})^2 (f - \pi(f)), f - \pi(f) \right>_{\pi} + \Delta t^3 R_f - \Delta t \mathrm{Var}_{\pi}[f], \end{split}$$

where R_f is a remainder term depending on f. Since f is smooth, we can iteratively apply (8.4) to the second term and third terms on the RHS obtaining

$$\begin{split} \sigma_{\Delta t}^2(f) &= 2 \left\langle (-\mathcal{L})^{-1}(f - \pi(f)), f - \pi(f) \right\rangle_{\pi} \\ &+ \Delta t \left\langle (-\mathcal{L})^{-1}(-\mathcal{L})(f - \pi(f)), f - \pi(f) \right\rangle_{\pi} - \Delta t \mathrm{Var}_{\pi}[f] \\ &+ \frac{\Delta t^2}{6} \left\langle (-\mathcal{L})(f - \pi(f)), f - \pi(f) \right\rangle_{\pi} \\ &+ \Delta t^3 R_f \\ &= \sigma^2(f) + \frac{\Delta t^2}{6} \left\langle (-\mathcal{L})(f - \pi(f)), f - \pi(f) \right\rangle_{\pi} + \Delta t^3 \widetilde{R}_f, \end{split}$$

as required.

Proof of Theorem 4.6. The proof of this result follows closely that of [18, Theorem 2.9]. To this end, given $f,g \in C^{\infty}(\mathbb{T}^d)$ such that $\pi(f) = \pi(g) = 0$, consider

$$\left\langle \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, g \right\rangle_{\pi}$$
.

Since $\left(\frac{I-P_{\Delta t}}{\Delta t}\right)^{-1}f$ has mean zero with respect to π , then

$$\left\langle \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, g \right\rangle_{\pi} = \left\langle \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, M_{\Delta t} g \right\rangle_{\pi}$$
$$= \left\langle \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}} + \Delta t^{r} R_{f,g},$$

for a smooth remainder term $R_{f,g}$ bounded uniformly with respect to Δt . Using the expansion (4.6) for the semigroup $\widehat{P}_{\Delta t}$:

$$\left\langle \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}} = \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right) \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}$$

$$= \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} \left(\frac{I - P_{\Delta t}}{\Delta t} + \Delta t^{k} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_{k}\right)\right) \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}$$

$$+ \Delta t^{q-1} \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} R_{f}, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}$$

$$+ \Delta t^{k} \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_{k}\right) \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}$$

$$+ \Delta t^{q-1} \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} R_{f}, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}$$

$$+ \Delta t^{q-1} \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} R_{f}, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}$$

$$+ \Delta t^{q-1} \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} R_{f}, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}}$$

where R_f is a smooth function depending on f, bounded uniformly with respect to Δt . By Assumption 4.3, the coefficients of the Δt^k and Δt^{q-1} terms are bounded uniformly with respect to Δt . Equation (4.13) then follows immediately, and thus (4.15). Assume now that (4.16) holds, then by applying (8.5) with

$$f = \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_k\right) \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f$$
, and $g = g$,

we obtain

$$\begin{split} R_1(f,g) &= \left\langle \left(\frac{I - \widehat{P}_{\Delta t}}{\Delta t}\right)^{-1} M_{\Delta t} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_k\right) \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, M_{\Delta t} g \right\rangle_{\widehat{\pi}_{\Delta t}} \\ &= \left\langle \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_k\right) \left(\frac{I - P_{\Delta t}}{\Delta t}\right)^{-1} f, g \right\rangle_{\pi} + \Delta t^{q-1} R_2(f,g), \end{split}$$

for some smooth, uniformly bounded remainder term R_2 . We now apply (4.12) to the discrete generator $\Delta t^{-1}(I-P_{\Delta t})$ to obtain

$$\left\langle \left(\frac{I - P_{\Delta t}}{\Delta t} \right)^{-1} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_k \right) \left(\frac{I - P_{\Delta t}}{\Delta t} \right)^{-1} f, g \right\rangle_{\pi} = \left\langle (-\mathcal{L})^{-1} \left(\frac{\mathcal{L}^{k+1}}{(k+1)!} - A_k \right) (-\mathcal{L})^{-1} f, g \right\rangle_{\pi} + \Delta t R_3(f, g),$$

for a smooth bounded remainder term R_3 , from which (4.14) follows.

8.3 Asymptotic bias of the splitting scheme

Here we prove the results from Section 5.1

Proof of Theorem 5.1. Assume that the transition semigroup associated with $\widehat{X}^{\Delta t}$ satisfies the expansion (4.6). In order to prove the first part of Theorem 5.1 it is enough to show that

$$A_i^* \pi = 0 \quad \text{for} \quad j = 1, \dots r - 1, \qquad A_r^* \pi = \text{div}(f_r \pi).$$
 (8.6)

The result then follows immediately from Theorem 4.1 using the identity

$$\int_{\mathbb{T}^d} A_r \psi(z) \pi(z) dz = -\int_{\mathbb{T}^d} \psi(z) \operatorname{div}(f_r(z) \pi(z)) dz. \tag{8.7}$$

We now start with the calculation of A_j . In particular, given $\phi \in C^{\infty}(\mathbb{T}^d)$ and $x \in \mathbb{R}^N$, using the semigroup property of the Markov process we have

$$\mathbb{E}\left[\phi\left(\widehat{X}_{1}^{\Delta t}\right)|\widehat{X}_{0}^{\Delta t}=x\right]=\mathbb{E}\left[\phi\left(\Phi_{\Delta t}\circ\Theta_{\Delta t}(x)\right)\right]=e^{\Delta t\mathcal{L}_{S,num}}(\phi\circ\Phi_{\Delta t})(x),\tag{8.8}$$

where $e^{\Delta t \mathcal{L}_{S,num}} \phi$ denotes the numerical flow generated by the numerical method applied to the reversible part of the dynamics (1.12). We next recall the generator (4.5) of the truncated modified equation (4.4) of the integrator $\Phi_{\Delta t}$,

$$\widetilde{\mathcal{L}}_D \phi = F_0 + \Delta t F_1 \phi + \ldots + \Delta t^r F_r \phi + \Delta t^{r+1} R_{\phi},$$

where R_{ϕ} is a smooth remainder term bounded uniformly with respect to Δt and where we define the differential operators $F_j \phi = f_j \cdot \nabla \phi$ (with $f_0 = f$). We then have

$$\mathbb{E}\left[\phi\left(\widehat{X}_{1}^{\Delta t}\right)|\widehat{X}_{0}^{\Delta t}=x\right] = \left(\sum_{k=0}^{r} \frac{\Delta t^{k} \mathcal{L}_{S,num}^{k}}{k!}\right) \left(\sum_{k=0}^{r} \frac{\Delta t^{k} \widetilde{\mathcal{L}}_{D}^{k}}{k!}\right) \phi(x) + \Delta t^{r+2} R_{\phi}'$$

$$= \phi(x) + \Delta t \mathcal{L}\phi(x) + \sum_{k=1}^{r} \Delta t^{k+1} A_{k} \phi(x) + \Delta t^{r+2} R_{\phi}',$$

for a smooth remainder term R_ϕ^\prime and where

$$A_k = \sum_{j=0}^{k+1} \mathcal{L}_{S,num}^{k+1-j} \Big(\sum_{\substack{1 \le i \le j \\ n_1 + n_2 + \dots + n_i = j-i}} \frac{1}{i!(k+1-j)!} F_{n_1} \cdots F_{n_i} \Big),$$

where the second sum above is over integers $n_1, \ldots, n_i \ge 0$ and is equal to the identity I when j = 0. We obtain for all $k \ge 1$,

$$A_k^*\pi = \sum_{j=0}^{k+1} \Big(\sum_{\substack{1 \le i \le j \\ n_1 + n_2 + \dots + n_k = j-i}} \frac{1}{i!(k+1-j)!} F_{n_i}^* \cdots F_{n_1}^* \Big) (\mathcal{L}_{S,num}^*)^{k+1-j} \pi.$$

Now since the integrator applied to the reversible part preserves the invariant measure we have $\mathcal{L}_{S,num}^*\pi=0$ which together with $F_i^*\pi=0,\ i=1,\cdots r-1$ implies that for $k\leq r$, the only possibly non-zero term in the above sum is obtained for j=r+1, k=r, i=1, i.e., $F_r^*\pi=\mathrm{div}(f_r\pi)$. Hence, we deduce (8.6) which permits to conclude the proof.

8.4 Asymptotic variance of the splitting scheme

Here we prove the results from Section 5.2.

Proof of Theorem 5.4. Clearly, Assumption 4.3 holds immediately from Theorem C.3 in the Appendix. Consider the one step semigroup $\widehat{P}_{\Delta t} = \Theta_{\Delta t} \widehat{\Phi}_{\Delta t}$ be the one-step semigroup corresponding to the Lie-Trotter splitting scheme (1.10), where $\Theta_{\Delta t}$ is the one-step semigroup integrated by MALA. By A.2 one obtains

$$\widehat{P}_{\Delta t}\phi = \phi + \Delta t A_0 \phi + \Delta t^2 A_1 \phi + \Delta t^{5/2} R_{\phi},$$

where

$$A_0 = \mathcal{A}_1 + \mathcal{G}_1 = \mathcal{L},$$

$$A_1 = \mathcal{A}_2 + \mathcal{G}_1 \mathcal{A}_1 + \mathcal{G}_2,$$

and where R_{ϕ} is a smooth remainder term, bounded uniformly with respect to Δt . Since the integrator $\widehat{\Phi}_{\Delta t}$ is assumed to preserve the invariant distribution up to order 2, and $\widehat{\Theta}_{\Delta t}$ preserves π it follows that

$$\pi\left(\left(\mathcal{A}_2 + \mathcal{G}_1 \mathcal{A}_1 + \mathcal{G}_2\right) \phi\right) = 0, \quad \phi \in C^{\infty}(\mathbb{T}^d).$$

Applying Theorem 4.6, it follows that for $f \in C^{\infty}(\mathbb{T}^d)$,

$$\widehat{\sigma}_{\Delta t}^{2}(f) = \sigma_{\Delta}^{2}(f) + \Delta t R_{f} + o(\Delta t),$$

where

$$R_f = 2 \left\langle (-\mathcal{L})^{-1} (\mathcal{L}^2 / 2 - (\mathcal{A}_2 + \mathcal{G}_1 \mathcal{A}_1 + \mathcal{G}_2) (-\mathcal{L})^{-1} (f - \pi(f), f - \pi(f)) \right\rangle_{\pi}.$$

Finally, invoking Theorem 4.1 we obtain

$$\widehat{\sigma}_{\Delta t}^{2}(f) = \sigma^{2}(f) + \Delta t R_{f} + o(\Delta t),$$

as required.

9 Discussion

In this paper sampling methods based on nonreversible diffusions have been proposed and evaluated on a range of different inference problems. The development of these methods is an attempt to improve on existing MCMC methodology in the case of target densities that might be of high dimension and exhibit strong correlations. The key idea behind these samplers is the exploitation of the irreversibility of an underlying diffusion process, which leads to reduced asymptotic variance. This becomes possible through a careful discretisation of the underlying SDE that introduces a controllable bias, but more importantly mimics the reduced asymptotic variance of the non-reversible diffusion.

From a practical point of view, the careful balancing of the bias and variance achieved by the nonreversible samplers leads to much more efficient sampling than MALA. In particular, across all our experiments we observe improvements of two orders of magnitude in terms of effective sample size. Moreover, all our comparisons are being made on the basis of the same number of density evaluations used in the nonreversible samplers and MALA. Furthermore, in the case of the log- Gaussian Cox model the nonreversible samplers are able to achieve this dramatic improvement in terms of the ESS without the need of an expensive $\mathcal{O}(d^3)$ reparametrisation, which is also the computational bottleneck in high dimensions for more sophisticated sampling algorithms such as MMALA [10].

There exist a number of different directions that one could extend this work. In particular, when dealing with the nonreversible part of the dynamics further computational benefits may be achieved with the use of adaptive integration. Furthermore, one could replace the Metropolis-Hasting scheme used for simulating the reversible part of the dynamics by appropriate numerical schemes [1] that preserve the invariant measure to high order. In this situation one would expected the results of our analysis to still hold which is important as the corresponding nonreversible samplers would allow for greater flexibility in the presence of big data, where traditional MCMC methods might become prohibitively expensive.

Acknowledgements

G.A. Pavliotis is supported by the Engineering and Physical Sciences Research Council of the UK through Grants Nos. EP/L020564, EP/L024926 and EP/L025159. A. B Duncan ackowledges the EPSRC for support under EP/J009636/1, EP/L020564/1, EP/K009788/2 (Network on Computational Statistics and Machine Learning). K.

C. Zygalakis was supported by a grant from the Simons Foundation and by the Alan Turing Institute under the EPSRC grant EP/ N510129/1. Part of this work was done during the author's stay at the Newton Institute for the program Stochastic Dynamical Systems in Biology: Numerical Methods and Applications. The authors would like to thank Mark Girolami and Andrew Stuart for various discussions about the paper.

A Expansions for the Generator of the Reversible Dynamics

In this section we present the expansion of the generator for a variety of different Metropolised integrators

A.1 Expansion of the Generator for MALA

Consider the MALA scheme with proposal distribution $^4\mathcal{N}(x-\nabla U(x)\Delta t,2\Delta t)$, having density

$$q_{\Delta t}(x' \mid x) \propto \exp \left[-\frac{\left\langle x' - \left(x - \nabla U(x) \Delta t \right), \left(x' - \left(x - \nabla U(x) \Delta t \right) \right) \right\rangle}{4 \Delta t} \right],$$

where Δt is the stepsize. The acceptance probability is given by

$$\alpha(x', x) = \min\left(1, r(x', x)\right),\,$$

where

$$r(x',x) = \frac{\pi(x')q_{\Delta t}(x \mid x')}{\pi(x)q_{\Delta t}(x' \mid x)} = e^{-U(x')+U(x)-\frac{1}{4\Delta t}\langle x-x'+\nabla U(x')\Delta t, x-x'+\nabla U(x')\Delta t\rangle + \frac{1}{4\Delta t}\langle x'-x+\nabla U(x)\Delta t, x'-x+\nabla U(x)\Delta t\rangle}$$

$$= e^{-U(x')+U(x)+\frac{\Delta t}{4}|\nabla U(x)|^2 - \frac{\Delta t}{4}|\nabla U(x')|^2 + \frac{1}{2}\langle x'-x,\nabla U(x)+\nabla U(x')\rangle}$$

$$= e^{-\lambda(x',x)}.$$

where

$$\lambda(x',x) = U(x') - U(x) - \frac{\Delta t}{4} |\nabla U(x)|^2 + \frac{\Delta t}{4} |\nabla U(x')|^2 - \frac{1}{2} \langle x' - x, \nabla U(x) + \nabla U(x') \rangle$$

We now Taylor expand U(x') around x up to fourth order, using integral remainders, to obtain

$$U(x') - U(x) = \langle \nabla U(x), x' - x \rangle + \frac{1}{2} \langle x' - x, \nabla \nabla U(x)(x' - x) \rangle$$
$$+ \frac{1}{6} \nabla \nabla \nabla U(x) : (x' - x)^{\otimes 3}$$
$$+ \frac{1}{6} \int_0^1 (1 - t)^3 \nabla^4 U((1 - t)x + tx') : (x' - x)^{\otimes 4} dt,$$

and similarly

$$\nabla U(x') = \nabla U(x) + \langle \nabla \nabla U(x), x' - x \rangle + \frac{1}{2} \langle \nabla \nabla \nabla U(x), x' - x \rangle$$
$$+ \frac{1}{2} \int_0^1 (1 - t)^2 \nabla^4 U((1 - t)x + tx') : (x' - x)^{\otimes 3} dt.$$

⁴Here $U(x) = -\log \pi(x)$.

Substituting the above expansions in $\lambda(x',x)$, that a number of terms cancel out, leaving

$$\begin{split} \lambda(x',x) = & \frac{1}{6} \nabla \nabla \nabla U(x) : (x'-x)^{\otimes 3} + \frac{1}{6} \int_{0}^{1} (1-t)^{3} \nabla^{4} U((1-t)x + tx') : (x'-x)^{\otimes 4} \, dt \\ & - \frac{1}{2} \left\langle x' - x, \frac{1}{2} \nabla^{3} U(x) : (x'-x)^{\otimes 2} + \frac{1}{2} \int_{0}^{1} (1-t)^{2} \nabla^{4} U(x) ((1-t)x + tx') : (x'-x)^{\otimes 3} \, dt \right\rangle \\ & - \frac{\Delta t}{4} \left| \nabla U(x) \right|^{2} \\ & + \frac{\Delta t}{4} \left| \nabla U(x) + \nabla \nabla U(x) : (x'-x) + \frac{1}{2} \nabla^{3} U(x) : (x'-x)^{\otimes 2} \right. \\ & \left. + \frac{1}{2} \int_{0}^{1} (1-t)^{2} \nabla^{4} U((1-t)x + tx') : (x'-x)^{\otimes 3} \, dt \right|^{2}. \end{split}$$

Our objective is to obtain explicit expressions for the leading terms in the expansion of $\lambda(x',x)$, in the specific case where

$$x' = x - \nabla U(x)\Delta t + \sqrt{2\Delta t}G,$$

where $G \sim \mathcal{N}(0, I)$ and Δt is small. Indeed, we have that

$$\lambda \left(x - \nabla U(x) \Delta t + \sqrt{2\Delta t} G, x \right) = \Delta t^{3/2} T(x, G) + \Delta t^2 \xi(x, G),$$

where

$$T(x,G) = -\frac{\sqrt{2}}{6}\nabla^3 U(x) : G^{\otimes 3} + \frac{1}{\sqrt{2}} \left\langle \nabla U(x), \nabla \nabla U(x)G \right\rangle,$$

and $\xi(x,G)$ collects all terms of order Δt^2 are higher. Note that, since $\nabla^k U(q)$ is bounded for all $k \geq 0$, we have:

$$|\xi(x,G)| \le C(1+|G|^6),$$

for some constant C independent of q and G and uniformly on $0 \le \Delta t \le 1$. We use this lemma from [8].

Lemma A.1. For $a \in \mathbb{R}$, let $a_+ = 0 \vee a$. Then we have the following relationship:

$$x_{+} - \frac{x_{+}^{2}}{2} \le 1 - 1 \wedge e^{-x} \le x_{+}.$$
 (A.1)

As a consequence of this lemma, we have that

$$\alpha(x - \nabla U(x)\Delta t + \sqrt{2\Delta t}G, x) = \min\left(1, e^{-\alpha(x', x)}\right)$$

$$= 1 - \Delta t^{3/2}T_{+}(x, G) + \Delta t^{2}\widetilde{\xi}(x, G),$$
(A.2)

where $|\widetilde{\xi}(x,G)| \leq \widetilde{C}(1+|G|^{12})$. Given a smooth observable ψ , we now consider the effect of the semigroup on ψ over a short time δ . First note that the transition density of the MALA chain is given by

$$p_{MALA}(x'|x) = \alpha(x',x) + \delta_x(x') \int (1 - \alpha(z,x)) dz,$$

and so the semigroup for a single step of size Δt is given by

$$\widetilde{P}_{\Delta t}\psi(x) = \psi(x) + \mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\alpha(x - \nabla U(x)\Delta t + \sqrt{2\Delta t}G, x) \left(\psi(x - \nabla U(x)\Delta t + \sqrt{2\Delta t}G) - \psi(x) \right) \right].$$

We split the dynamics into two parts, a part which arises from the proposal, and a part which arises from the acceptance/rejection term. What we shall observe is that the second term does not contribute to the leading order

term. Indeed, the accept/reject part only has an $O(\Delta t^2)$ contribution.

$$\mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\alpha(x - \Delta t \nabla U(x) + \sqrt{2\Delta t}G, x) \left(\psi(x - \Delta t \nabla U(x) + \sqrt{2\Delta t}G) - \psi(x) \right) \right]$$

$$= \underbrace{\mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\psi(x - \Delta t \nabla U(x) + \sqrt{2\Delta t}G) - \psi(x) \right]}_{A} + \underbrace{\mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\left(\alpha(x - \Delta t \nabla U(q) + \sqrt{2\Delta t}G, q) - 1 \right) \left(\psi(q - \Delta t \nabla U(x) + \sqrt{2\Delta t}G) - \psi(x) \right) \right]}_{B}.$$

For the first term, we obtain after Taylor expansion of ψ :

$$A = \Delta t \left[-\nabla U(x) \cdot \nabla \psi(x) + \Delta \psi(x) \right]$$

+ $\Delta t^2 \left[\frac{1}{2} \nabla U(x) \cdot \nabla \nabla \psi(x) \nabla U(x) - \frac{1}{3} \nabla U(x) \nabla \Delta \psi + 2\Delta^2 \psi(x) \right] + r_1(x),$

where we use the fact that,

$$\mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\nabla U(x) \cdot \nabla \nabla \nabla \psi(x) : GG \right] = \mathbb{E}_{g \sim \mathcal{N}(0,I)} \left[\partial_{x_j} U(x) \partial_{jkl} \psi(x) G_k G_l \right]$$
$$= \partial_{x_j} U(x) \partial_{jkk} \psi(x)$$
$$= \nabla U(x) \cdot \nabla \Delta \psi(x),$$

and

$$\mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\nabla^4 \psi(x) : G^{\otimes 4} \right] = \mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\partial_{jklm} \psi(x) G_j G_k G_l G_m \right]$$
$$= 12 \sum_{j,k} \partial_{jjkk} \psi(x)$$
$$= 12 \Delta^2 \psi(x),$$

and where $|r_1(x)| \leq C\Delta t^{5/2}$. For the second term, using $x' = x - \nabla U(x)\Delta t + \sqrt{2\Delta t}G$,

$$\begin{split} B &= \mathbb{E}_{G \sim \mathcal{N}(0,I)} \left[\left(-\Delta t^{3/2} T_+(x,G) + \Delta t^2 \widetilde{\xi}(x,G) \right) \right. \\ & \left. \left(\left\langle -\Delta t \nabla U(x) + \sqrt{2\Delta t} G, \nabla \psi(x) \right\rangle + \int_0^1 (1-t) \nabla \nabla \psi((1-t)x + tx') : (x'-x)^{\otimes 2} \, dt \right) \right] \\ &= -\sqrt{2} \Delta t^2 \int \left(1 \wedge T(x,g) \right) \left\langle \nabla \psi(x), g \right\rangle \frac{e^{-|g|^2/2}}{\sqrt{(2\pi)^d}} \, dg + \Delta t^{5/2} r(x) \end{split}$$

Therefore we have that

$$P_{\Delta t}\psi(q) - \psi(q) = \Delta t \mathcal{G}_1\psi(q) + \Delta t^2 \mathcal{G}_2\psi(q) + \Delta t^{5/2}r(q),$$

where G_1 is the reversible part of the infinitesimal generator (2.4), i.e.

$$\mathcal{G}_1 = -\nabla U \cdot \nabla + \Delta,\tag{A.3}$$

and

$$\mathcal{G}_{2}\psi(q) = \frac{1}{2}\nabla U(q) \cdot \nabla \nabla \psi(q) \nabla U(q) - \frac{1}{3}\nabla U(q) \cdot \nabla \Delta \psi(q)$$

$$+ 2\Delta^{2}\psi - \sqrt{2}\int (1 \wedge T(q,g)) \langle \nabla \psi(q), g \rangle \frac{e^{-g \cdot g/2}}{\sqrt{(2\pi)^{d}}} dg,$$
(A.4)

and $|r(q)| \leq C$, uniformly in $0 < \Delta t \leq 1$.

The conclusion of the above is summarised in the following proposition.

Proposition A.2. Let $P_{\Delta t}$ denote the evolution operator corresponding to the MALA scheme, then for all smooth $\psi : \mathbb{T}^d \to \mathbb{R}$:

$$(P_{\Delta t} - I)\psi(q) = \Delta t \mathcal{G}_1 \psi(q) + \Delta t^2 \mathcal{G}_2 \psi + \Delta t^{5/2} r(q),$$

where G_1 , G_2 are given by (A.3) and (A.4), respectively and where r(q) are as given above. In particular, the infinitesimal generator $G_{\Delta t}$ corresponding to the MALA scheme satisfies

$$\mathcal{G}_{\Delta t}\psi(q) = \mathcal{G}_1\psi(q) + \Delta t \mathcal{G}_2\psi + \Delta t^{3/2}r(q), \quad \forall \psi \in C^{\infty}(\mathbb{T}^d).$$

B Analysis for Gaussian Distributions

In this section we will study the specific example where the dynamics (1.2) are linear and of the form

$$dX_t = -AX_t dt + \sqrt{2}\sigma dW_t, \tag{B.1}$$

where W_t is a standard m-dimensional Brownian motion, $A \in \mathbb{R}^{d \times d}$ and $\sigma \in \mathbb{R}^{d \times m}$ such that $\Sigma = \sigma \sigma^{\top}$ is positive definite. Provided that -A is stable, and Σ is positive definite, X_t is ergodic with unique invariant distribution $\pi(x) \propto \exp(-x \cdot \Sigma_{\infty} x/2)$, where the stationary covariance Σ_{∞} is the solution of the Lyapunov equation

$$A\Sigma_{\infty} + \Sigma_{\infty}A^{\top} = 2\Sigma,$$

which can be written explicitly as

$$\Sigma_{\infty} = 2 \int_{0}^{\infty} e^{-As} \Sigma e^{-A^{\top}s} \, ds$$

Our objective is to derive an explicit expression for the asymptotic variance $\sigma^2(f)$ of

$$I_t = \frac{1}{t} \int_0^t f(X_s) \, ds,$$

where f is a function of the form

$$f(x) = x \cdot Mx + L \cdot x + K,$$

for some $M \in \mathbb{R}^{d \times d}_{sym}$, $L \in \mathbb{R}^d$ and $K \in \mathbb{R}$. Taking a different approach to [7], we shall obtain this expression via the Green-Kubo formula, i.e.

$$\sigma^{2}(f) = 2 \int_{0}^{\infty} \langle P_{t}f - \pi(f), f - \pi(f) \rangle_{\pi} dt = 2 \int_{0}^{\infty} \left(\int_{\mathbb{R}^{d}} P_{t}(f - \pi(f))(x) (f(x) - \pi(f)) \pi(dx) \right) dt, \quad (B.2)$$

where P_t is the semigroup corresponding to (B.1). We note that for l>0, the process X_t satisfies the Foster-Lyapunov condition (2.8) with Lyapunov function $V_l(x)=1+|x|^{2l}$. In particular, by Proposition 2.5, a CLT for the estimator $\pi_T(f)=T^{-1}\int_0^T f(X_t)\,dt$ will hold for all observables f having algebraic growth, and moreover (B.2) is well defined and finite. We shall first prove the result for $\Sigma=I$, and then obtain the general case via a simple linear transformation. In this case $\Sigma_\infty=C^{-1}$, where $C=\mathrm{Sym}[A]=\frac{1}{2}\left[A+A^\top\right]$. To obtain this result we shall make use of the following form of Mehler's formula.

Lemma B.1. Let P_t be the semigroup corresponding to the dynamics

$$dX_t = -AX_t dt + \sqrt{2}dW_t.$$

where W_t is a standard d-dimensional Brownian motion. Then, for all $f \in L^2(\pi)$ we have

$$P_t f(x) = \mathbb{E}\left[f(e^{-At}x + \sqrt{I - e^{-2Ct}}Z)\right] = \int_{\mathbb{R}^d} f(e^{-At}x + C^{-1}\sqrt{I - e^{-2Ct}}z) e^{-|z|^2/2} dz$$

where $Z \sim \mathcal{N}(0, C^{-1})$.

First consider the observable $f_1(x) = x \cdot Mx$ for $M \in \mathbb{R}^{d \times d}_{sym}$, then

$$\begin{split} P_t f_1(x) &= x \cdot e^{-A^\top t} M e^{-At} x + \operatorname{Tr} \left[C^{-1} \sqrt{I - e^{-2Ct}} M \sqrt{I - e^{-2Ct}} \right] \\ &= x \cdot e^{-A^\top t} M e^{-At} x + \operatorname{Tr} \left[C^{-1} M \left(I - e^{-2Ct} \right) \right]. \end{split}$$

Now, using the fact that $\pi(f_1) = \text{Tr}[C^{-1}M]$, we can write

$$\int_0^\infty P_t[f_1(x) - \pi(f_1)] dt = x \cdot \left[\int_0^\infty e^{-A^\top t} M e^{-At} dt \right] x - \frac{1}{2} \text{Tr}[C^{-1} M C^{-1}].$$

Similarly, if $f_2(x) = L \cdot x$, then $\pi(f_2) = 0$ and

$$P_t f_2(x) = x \cdot e^{-A^{\top} t} L,$$

so that

$$\int_{0}^{\infty} P_{t} f_{2}(x) - \pi(f_{2}) = x \cdot (A^{\top})^{-1} L.$$

Thus, it follows that the unique, mean-zero solution of the Poisson equation

$$-Ax \cdot \nabla \phi(x) + \Delta \phi(x) = f(x) - \pi(f),$$

is given by

$$\phi(x) = x \cdot \left[\int_0^\infty e^{-A^\top t} M e^{-At} \, dt \right] x - \operatorname{Tr} \left[C^{-1} \int_0^\infty e^{-A^\top t} M e^{-At} \, dt \right] + x \cdot \left(A^\top \right)^{-1} L.$$

We now use the following Green-Kubo type formula to compute the asymptotic variance for $f=f_1+f_2$:

$$\begin{split} \frac{1}{2}\sigma^2(f) &= \int_{\mathbb{R}^d} \int_0^\infty \left[P_t f(x) - \pi(f) \right] \left[f(x) - \pi(f) \right] \, dt \, \pi(dx) \\ &= \int \left[\left(x \cdot \Pi x \right) \left(x \cdot M x \right) - \mathrm{Tr} [MC^{-1}] x \cdot \Pi x + \left(x \cdot A^{-\top} L \right)^2 \right] \pi(dx) \\ &= 2 \mathrm{Tr} \left[C^{-1} \Pi C^{-1} M \right] + L \cdot A^{-1} C^{-1} A^{-\top} L, \end{split}$$

where

$$\Pi = \int_0^\infty e^{-A^{\mathsf{T}}t} M e^{-At} \, dt.$$

In the case when $A = I + \alpha J$, for $J^{\top} = -J$, so that C = I, we obtain

$$\frac{1}{2}\sigma^2(f) = 2\operatorname{Tr}\left[\left(\int_0^\infty e^{-A^\top t} M e^{-At} \, dt\right) M\right] + L\cdot (I + \alpha^2 J J^\top)^{-1} L,$$

which is precisely the formula derived in [7] using a different approach.

Now suppose that $\Sigma \neq I$, so that the Poisson equation we must solve becomes

$$-Ax \cdot \nabla \phi(x) + \Sigma : \nabla \nabla \phi(x) = f(x) - \pi(f).$$

Writing $\phi(x) = \psi(\Sigma^{-1/2}x)$, then we have that

$$-Ax\cdot \Sigma^{-1/2}\nabla \psi(\Sigma^{-1/2}x) + \Delta \psi(\Sigma^{-1/2}x) = f(x) - \pi(f),$$

so that

$$- \Sigma^{-1/2} A \Sigma^{1/2} x \cdot \nabla \psi(x) + \Delta \psi(x) = f(\Sigma^{1/2} x) - \pi(f).$$
 (B.3)

The linear operator defined on the left hand side of (B.3) corresponds to a linear diffusion with stationary distribution $\widetilde{\pi} = \mathcal{N}(0, \widetilde{\Sigma}_{\infty})$ where

$$\widetilde{\Sigma}_{\infty} = \int_0^{\infty} e^{A_{\Sigma}t} e^{A_{\Sigma}^{\top}t} dt = \Sigma^{-1/2} \Sigma_{\infty} \Sigma^{-1/2}.$$

where $A_{\Sigma} = \Sigma^{-1/2} A \Sigma^{1/2}$. Computing the asymptotic variance $\sigma^2(f)$:

$$2\int_{\mathbb{R}^d} \phi(x)(f(x) - \pi(f))\pi(x) = 2\int_{\mathbb{R}^d} \psi(\Sigma^{-1/2}x)(f(x) - \pi(f))\pi(x) dx$$
$$= 2\int_{\mathbb{R}^d} \psi(x)[f(\Sigma^{1/2}x) - \pi(f)]\widetilde{\pi}(x) dx.$$

Applying the previous result, it follows that

$$\frac{1}{2}\sigma^2(f) = 2\mathrm{Tr}\left[\left(\int_0^\infty e^{-A_\Sigma^\top t} M_\Sigma e^{-A_\Sigma t}\,dt\right)M\right] + L\cdot A_\Sigma^{-1}\widetilde{\Sigma}_\infty^{-1}A_\Sigma^{-\top}L,$$

where $M_{\Sigma} = \Sigma^{1/2} M \Sigma^{1/2}$. Noting that $e^{-A_{\Sigma}t} = \Sigma^{-1/2} e^{-At} \Sigma^{1/2}$, it follows that

$$2\mathrm{Tr}\left[\left(\int_0^\infty \Sigma^{1/2} e^{-A^\top t} M e^{-At} \Sigma^{1/2} \, dt\right) M\right] = 2\mathrm{Tr}\left[\left(\int_0^\infty e^{-A^\top t} M e^{-At} \, dt\right) M_\Sigma\right]$$

and moreover,

$$L \cdot \Sigma^{-1/2} A^{-1} \Sigma^{1/2} \widetilde{\Sigma}_{\infty} \Sigma^{1/2} A^{-\top} \Sigma^{-1/2} L = L \cdot \Sigma^{-1/2} A^{-1} \Sigma_{\infty} A^{-\top} \Sigma^{-1/2} L = L_{\Sigma} \cdot A^{-1} \Sigma_{\infty} A^{-\top} L_{\Sigma},$$

where $L_{\Sigma} = \Sigma^{-1/2}L$. In summary we have the following result.

Proposition B.2. Consider the linear diffusion defined by the SDE,

$$dX_t = -AX_t dt + \sqrt{2}\sigma dW_t$$

where W_t is a m-dimensional Brownian motion, $\sigma \in \mathbb{R}^{d \times m}$ such that $\Sigma = \sigma \sigma^{\top}$ is positive definite and -A is stable. Then, for

$$f(x) = x \cdot Mx + L \cdot x + K,$$

the asymptotic variance $\sigma^2(f)$ is given by

$$\frac{1}{2}\sigma^2(f) = 2Tr \left[\left(\int_0^\infty e^{-A^\top t} M e^{-At} dt \right) M_\Sigma \right] + L_\Sigma \cdot A^{-1} \Sigma_\infty A^{-\top} L_\Sigma,$$

where $M_{\Sigma} = \Sigma^{1/2} M \Sigma^{1/2}$ and $L_{\Sigma} = \Sigma^{-1/2} L$.

Remark B.3. Note that there is no impediment to deriving the asymptotic variance for observables involving higher powers, e.g. a third order tensor of the form $\sum_{i,j,k} K_{i,j,k} x_i x_j x_k$, but we only provide the result up to second order for the sake of clarity. A more general approach would potentially be possible by considering the decomposition of an observable f with respect to the eigenbasis of the Ornstein Uhlenbeck operator \mathcal{L} , which can be shown to be Hermite polynomials [27].

C Spectral Gap estimate for the Splitting Scheme

In this section we shall focus specifically on the splitting scheme where the reversible component is simulated using MALA where we show that Assumption 4.3 holds in this case. The approach we follow is strongly based on arguments found in [8, 3]. The method depends strongly on the fact that the proposal of the MALA scheme is a first order approximation of the corresponding SDE. In this section, we shall assume that Assumption 5.3 holds.

Define $Q_{\Delta t}(x,y)$ to be the transition kernel for the exact dynamics

$$dY_t = (\nabla \log \pi(Y_t) + \gamma(Y_t)) dt + \sqrt{2} dW_t, \tag{C.1}$$

and $\widetilde{P}_{\Delta t}(x,y)$ the transition kernel of the unadjusted Lie-Trotter scheme defined by

$$\widetilde{Z}_{n+1} = \Phi_{\Delta t}(\widetilde{Z}_n) + \nabla \log \pi(\Phi_{\Delta t}(\widetilde{Z}_n))\Delta t + \sqrt{2\Delta t}\,\xi_n,\tag{C.2}$$

where $\xi_n \sim \mathcal{N}(0, I)$, and $\widetilde{Q}_{\Delta t}(x, y)$ to be the transition kernel of the Euler-Maruyama discretisation of (C.1), i.e.

$$\widetilde{Y}_{n+1} = \widetilde{Y}_n + \nabla \log \pi(\widetilde{Y}_n) \Delta t + \gamma(\widetilde{Y}_n) \Delta t + \sqrt{2\Delta t} \, \xi_n. \tag{C.3}$$

Lemma C.1. Consider a coupling of \widetilde{Z}_n and \widetilde{Y}_n such that they are driven by the same noise W_t , and $\widetilde{Z}_0 = \widetilde{Y}_0 = x$. Then, for $t = n\Delta t$ there exists a constant C(t) > 0 independent of Δt such that

$$\mathbb{E}_x \left| \widetilde{Y}_n - \widetilde{Z}_n \right| \le C(t) \Delta t \tag{C.4}$$

for Δt sufficiently small.

Proof. Using the fact that $\Phi_{\Delta t}(x) = x + \gamma(x)\Delta t + K_1(x)\Delta t^2$ for some function K_1 bounded uniformly on \mathbb{T}^d for Δt sufficiently small, we have that

$$\begin{split} \mathbb{E}_{x}|\widetilde{Y}_{n+1} - \widetilde{Z}_{n+1}| &\leq \mathbb{E}_{x}|\Phi_{\Delta t}(\widetilde{Z}_{n}) + \nabla \log \pi(\Phi_{\Delta t}(\widetilde{Z}_{n}))\Delta t - \widetilde{Y}_{n} - \nabla \log \pi(\widetilde{Y}_{n})\Delta t - \gamma(\widetilde{Y}_{n})\Delta t| \\ &\leq \mathbb{E}_{x}|\widetilde{Z}_{n} - \widetilde{Y}_{n}| + \Delta t \mathbb{E}_{x}|\gamma(\widetilde{Y}_{n}) - \gamma(\widetilde{Z}_{n})| + \Delta t \mathbb{E}_{x}|\nabla \log \pi(\widetilde{Y}_{n}) - \nabla \log \pi(\widetilde{Z}_{n})| + C\Delta t^{2} \\ &\leq (1 + K\Delta t)\mathbb{E}_{x}|\widetilde{Z}_{n} - \widetilde{Y}_{n}| + C\Delta t^{2}, \end{split}$$

where C > 0 is a constant, from which the result follows.

Remark C.2. It follows automatically from (C.1) and standard estimates for Euler-Maruyama discretisation of SDEs with additive noise that, for Δt sufficiently small, there exists C(t) > 0 such that

$$\mathbb{E}_x |Z_n - Y_{n\Delta t}| \le C(t)\Delta t,$$

where $t = n\Delta t$.

Theorem C.3. Suppose that Assumption 5.3 holds, then the Lie-Trotter scheme (1.10) possesses a unique invariant distribution $\pi_{\Delta t}$ and moreover, there exists $\Delta t_0 > 0$, $C, \lambda > 0$ such that, for any $0 < \Delta t \le \Delta t_0$ and all $n \in \mathbb{N}$

$$||P_{\Delta t}^n f - \pi_{\Delta t}(f)||_{\infty} \le Ce^{-\lambda n\Delta t} ||f - \pi_{\Delta t}(f)||_{\infty}, \quad f \in L^{\infty}(\mathbb{T}^d).$$

As a consequence, there exists a constant K>0, independent of Δt such that

$$\left\| \frac{I - P_{\Delta t}}{\Delta t} \right\|_{L_0^{\infty}(\pi_{\Delta t})} < K, \tag{C.5}$$

for Δt sufficiently small.

Proof. Denote by $P_{\Delta t}^n(x,y)$ the transition kernel density corresponding to (1.10). Since the domain is compact, we need only verify that we have the following uniform minorisation condition

$$\left\| P_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) - P_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(y,\cdot) \right\|_{TV} \le 2(1-\gamma), \qquad x,y \in \mathbb{T}^d,$$

for some $\gamma \in (0,1)$, uniformly on Δt . Then

$$\begin{split} \sup_{x,y \in \mathbb{T}^d} \left\| P_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) - P_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(y,\cdot) \right\|_{TV} &\leq 2 \sup_{x \in \mathbb{T}^d} \left\| P_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) \right\|_{TV} \\ &+ \sup_{x,y \in \mathbb{T}^d} \left\| Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(y,\cdot) \right\|_{TV}. \end{split}$$

Now, since the domain is compact, and the diffusion process Z_t is uniformly elliptic, we know that the transition kernel $Q_{\Delta t}(\cdot,\cdot)$ satisfies a minorisation condition, and thus there exists $\gamma_1 > 0$ such that

$$\sup_{x,y \in \mathbb{T}^d} \left\| Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(y,\cdot) \right\|_{TV} \le 2(1-\gamma_1). \tag{C.6}$$

Following [3, 8] we introduce the transition kernel of the un-adjusted splitting scheme (C.2), denoted by $\widetilde{P}_{\Delta t}(x,y)$. Then we have

$$\sup_{x \in \mathbb{T}^d} \left\| P_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x, \cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x, \cdot) \right\|_{TV} \leq \sup_{x \in \mathbb{T}^d} \left\| \widetilde{P}_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x, \cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x, \cdot) \right\|_{TV} + \sup_{x \in \mathbb{T}^d} \left\| \widetilde{P}_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x, \cdot) - P_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x, \cdot) \right\|_{TV}$$
(C.7)

To control the second term on the RHS of (C.7) we apply a coupling argument, identical to that of [3, Lemma 3.2] and [8, Lemma 2]. To this end, consider the processes Z_n and \widetilde{Z}_n defined by (1.10) and (C.2) respectively, and assume that they are driven by the same noise process, starting from $Z_0 = \widetilde{Z}_0 = x$. Using the coupling characterization of total variation

$$\sup_{x \in \mathbb{T}^d} \left\| \widetilde{P}_{\Delta t}^n(x, \cdot) - P_{\Delta t}^n(x, \cdot) \right\|_{TV} \le 2\mathbb{P} \left[Z_n \neq \widetilde{Z}_n | Z_0 = \widetilde{Z}_0 = x \right] \le 2\sum_{i=1}^n \mathbb{E} \left[\left(1 - \alpha(\Phi_h(Z_i), Z_{i+1}) \right) | Z_0 = x \right],$$

where $\alpha(x,y)$ is the probability of the standard MALA scheme of accepting a transition from x to y. From (A.2), (see also [8, Lemma 1]), there exists $C_1 > 0$ such that

$$\mathbb{E}\left[\left(1 - \alpha(\Phi_{\Delta t}(Z_i), Z_{i+1})\right) \mid Z_0 = x\right] \le C_1 \Delta t^{3/2},$$

for Δt sufficiently small. We can rewrite the first term on the RHS of (C.7) as

$$\begin{split} \sup_{x \in \mathbb{T}^d} \left\| \widetilde{P}_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) \right\|_{TV} & \leq \sup_{x \in \mathbb{T}^d} \left\| \widetilde{P}_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) - \widetilde{P}_{\Delta t} \circ Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor - 1}(x,\cdot) \right\|_{TV} \\ & + \sup_{x \in \mathbb{T}^d} \left\| \widetilde{P}_{\Delta t} \circ Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor - 1}(x,\cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x,\cdot) \right\|_{TV}. \end{split}$$

Noting that

$$\left\|\widetilde{P}_{\Delta t}^{\lfloor 1/\Delta t\rfloor}(x,\cdot) - \widetilde{P}_{\Delta t} \circ Q_{\Delta t}^{\lfloor 1/\Delta t\rfloor - 1}(x,\cdot)\right\|_{TV} = \mathbb{E}_x \left\|\widetilde{P}_{\Delta t}(\widetilde{Z}_{\lfloor 1/\Delta t\rfloor - 1},\cdot) - \widetilde{P}_{\Delta t}(Y_{\lfloor 1/\Delta t\rfloor - 1},\cdot)\right\|_{TV}.$$

From Pinkser's inequality, we get

$$\begin{split} \left\| \widetilde{P}_{\Delta t}(x,\cdot) - \widetilde{P}_{\Delta t}(y,\cdot) \right\|_{TV} &\leq \frac{\left| \Phi_{\Delta t}(x) + \nabla \log \pi(\Phi_{\Delta t}(x)) \Delta t - \Phi_{\Delta t}(y) - \nabla \log \pi(\Phi_{\Delta t}(y)) \Delta t \right|}{\sqrt{2\Delta t}} \\ &\leq \frac{1}{\sqrt{2\Delta t}} \left| \Phi_{\Delta t}(x) - \Phi_{\Delta t}(y) \right| + \sqrt{\frac{\Delta t}{2}} \left| \nabla \log \pi(\Phi_{\Delta t}(x)) - \nabla \log \pi(\Phi_{\Delta t}(y)) \right| \\ &\leq K \left(\Delta t^{1/2} + \Delta t^{-1/2} \right) \left| x - y \right|, \end{split}$$

using (5.4) and the fact that $\nabla\nabla \log \pi$ is bounded uniformly on \mathbb{T}^d . Therefore, from the remark following Lemma C.1,

$$\mathbb{E}_{x} \left\| \widetilde{P}_{\Delta t}(\widetilde{Z}_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) - \widetilde{P}_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) \right\|_{TV} \leq K \Delta t^{-1/2} \mathbb{E}_{x} \left\| \widetilde{Z}_{\lfloor 1/\Delta t \rfloor - 1} - Y_{\lfloor 1/\Delta t \rfloor - 1} \right\| \leq C \Delta t^{1/2},$$

where C is a constant independent of Δt . We introduce an intermediate continuous time process \widetilde{Y}_t defined on $[0, \Delta t]$ by

$$d\widetilde{Y}_t = (\gamma(x) + \nabla \log \pi(x)) dt + \sqrt{2} dW_t, \quad \widetilde{Y}_0 = x,$$

with corresponding transition kernel $\widetilde{Q}_t(x,\cdot)$. Then we have

$$\begin{split} \left\| \widetilde{P}_{\Delta t} \circ Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor - 1}(x, \cdot) - Q_{\Delta t}^{\lfloor 1/\Delta t \rfloor}(x, \cdot) \right\|_{TV} = & \mathbb{E}_{x} \left\| \widetilde{P}_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) - Q_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) \right\|_{TV} \\ \leq & \mathbb{E}_{x} \left\| \widetilde{Q}_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) - Q_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) \right\|_{TV} \\ & + \mathbb{E}_{x} \left\| \widetilde{P}_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) - \widetilde{Q}_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) \right\|_{TV}. \end{split}$$
(C.8)

From Pinsker's inequality we have

$$\left\|\widetilde{P}_{\Delta t}(x,\cdot) - \widetilde{Q}_{\Delta t}(x,\cdot)\right\|_{TV} \leq \frac{\left|\Phi_{\Delta t}(x) + \nabla \log \pi (\Phi_{\Delta t}(x))\Delta t - x - \nabla \log \pi (x)\Delta t - \gamma(x)\Delta t\right|}{\sqrt{2\Delta t}}.$$

Using the fact that

$$\Phi_{\Delta t}(x) = x + \gamma(x)\Delta t + K_1(x)\Delta t^2,$$

for some function K_1 bounded uniformly on \mathbb{T}^d for Δt sufficiently small, and applying Taylor's theorem for $\nabla \log \pi$, we obtain the bound

$$\|\widetilde{P}_{\Delta t}(x,\cdot) - \widetilde{Q}_{\Delta t}(x,\cdot)\|_{TV} \le K\Delta t^{3/2},$$

for some constant K independent of Δt . Denote by $\widetilde{\mathbb{Q}}_{\Delta t}$ and $\mathbb{Q}_{\Delta t}$ the path measures on $C[0, \Delta t]$ induced by the processes \widetilde{Y}_t and Y_t , respectively. Then by Girsanov's theorem [16, Ch. 3, Cor. 5.16] we obtain

$$\frac{d\mathbb{Q}_{\Delta t}}{d\widetilde{\mathbb{Q}}_{\Delta t}}(\widetilde{Y}_t) = \exp\left(\frac{1}{2} \int_0^{\Delta t} \left\langle \gamma(\widetilde{Y}_s) + \nabla \log \pi(\widetilde{Y}_s) - \gamma(x) - \nabla \log \pi(x), d\widetilde{Y}_s \right\rangle - \frac{1}{4} \int_0^{\Delta t} |\gamma(\widetilde{Y}_s) + \nabla \log \pi(\widetilde{Y}_s)|^2 - |\gamma(x) + \nabla \log \pi(x)|^2 ds \right).$$

By Pinsker's inequality, it follows that

$$\begin{split} \|\widetilde{Q}_{\Delta t}(x,\cdot) - Q_{\Delta t}(x,\cdot)\|_{TV}^2 &\leq \frac{1}{2} \mathbb{E}_x \int_0^{\Delta t} \left| \gamma(\widetilde{Y}_s) + \nabla \log \pi(\widetilde{Y}_s) - \gamma(x) - \nabla \log \pi(x) \right|^2 \, ds \\ &\leq \frac{1}{2} \mathbb{E}_x \int_0^{\Delta t} \left| \gamma(\widetilde{Y}_s) + \nabla \log \pi(\widetilde{Y}_s) - \gamma(x) - \nabla \log \pi(x) \right|^2 \, ds \\ &\leq C \int_0^{\Delta t} \left| \mathbb{E}_x \left| \widetilde{Y}_s - x \right|^2 \, ds \right. \\ &\leq C \int_0^{\Delta t} \left| \gamma(x) + \nabla \log \pi(x) \right|^2 s^2 + s \, ds \\ &\leq C \left(|\gamma(x) + \nabla \log \pi(x)|^2 \frac{\Delta t^3}{3} + \frac{\Delta t^2}{2} \right), \end{split}$$

and so, there exists a constant K>0 independent of Δt such that

$$\mathbb{E}_{x} \left\| \widetilde{Q}_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) - Q_{\Delta t}(Y_{\lfloor 1/\Delta t \rfloor - 1}, \cdot) \right\|_{TV} \le K\Delta t. \tag{C.9}$$

Collecting the terms together, it follows that for Δt sufficiently small, condition (C.6) holds. The bound (C.5) then follows immediately.

References

- [1] A. Abdulle, G. Vilmart, and K. C. Zygalakis. High order numerical approximation of the invariant measure of ergodic SDEs. *SIAM J. Numer. Anal.*, 52(4):1600–1622, 2014.
- [2] A. Abdulle, G. Vilmart, and K. C. Zygalakis. Long time accuracy of Lie–Trotter splitting methods for langevin dynamics. *SIAM Journal on Numerical Analysis*, 53(1):1–16, 2015.
- [3] N. Bou-Rabee and M. Hairer. Nonasymptotic mixing of the MALA algorithm. *IMA Journal of Numerical Analysis*, 2012.
- [4] O. F. Christensen, G. O. Roberts, and J. S. Rosenthal. Scaling limits for the transient phase of local Metropolis–Hastings algorithms. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 67(2):253–268, 2005.
- [5] P. Diaconis, S. Holmes, and R. M. Neal. Analysis of a nonreversible Markov chain sampler. *The Annals of Applied Probability*, 10(3):726–752, 2000.
- [6] S. Duane, A. D. Kennedy, B. J. Pendleton, and D. Roweth. Hybrid monte carlo. *Physics letters B*, 195(2):216–222, 1987.

- [7] A. B. Duncan, T Lelièvre, and G. A. Pavliotis. Variance reduction using nonreversible Langevin samplers. *Journal of Statistical Physics*, 163(3):457–491, 2016.
- [8] M. Fathi, A.-A. Homman, and G. Stoltz. Error analysis of the transport properties of metropolized schemes. *ESAIM: Proceedings and Surveys*, 48:341–363, 2015.
- [9] C. W. Gardiner. Handbook of stochastic methods. Springer-Verlag, Berlin, second edition, 1985.
- [10] M. Girolami and B. Calderhead. Riemann Manifold Langevin and Hamiltonian Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(2):123–214, 2011.
- [11] P. W. Glynn and S. P. Meyn. A Liapounov bound for solutions of the Poisson equation. *The Annals of Probability*, 24(2):916–931, 1996.
- [12] E. Hairer, C. Lubich, and G. Wanner. *Geometric Numerical Integration. Structure-Preserving Algorithms for Ordinary Differential Equations*. Springer Series in Computational Mathematics 31. Springer-Verlag, Berlin, second edition, 2006.
- [13] W. K. Hastings. Monte carlo sampling methods using Markov chains and their applications. *Biometrika*, 57(1):97–109, 1970.
- [14] A. M. Horowitz. A generalized guided Monte Carlo algorithm. *Physics Letters B*, 268(2):247–252, 1991.
- [15] K. Hukushima and Y. Sakai. An irreversible Markov-chain Monte Carlo method with skew detailed balance conditions. In *Journal of Physics: Conference Series*, volume 473, page 012012. IOP Publishing, 2013.
- [16] I. Karatzas and S. Shreve. *Brownian motion and stochastic calculus*, volume 113. Springer Science & Business Media, 2012.
- [17] M. Kopec. Weak backward error analysis for overdamped Langevin processes. *IMA Journal of Numerical Analysis*, page dru016, 2014.
- [18] B. Leimkuhler, C. Matthews, and G. Stoltz. The computation of averages from equilibrium and nonequilibrium Langevin molecular dynamics. *IMA Journal of Numerical Analysis*, 2015.
- [19] B. Leimkuhler and S. Reich. *Simulating Hamiltonian Dynamics*. Cambridge Monographs on Applied and Computational Mathematics 14. Cambridge University Press, Cambridge, 2004.
- [20] T. Lelièvre, F. Nier, and G. A. Pavliotis. Optimal non-reversible linear drift for the convergence to equilibrium of a diffusion. *Journal of Statistical Physics*, 152(2):237–274, 2013.
- [21] L. Lorenzi and M. Bertoldi. Analytical methods for Markov semigroups. CRC Press, 2006.
- [22] Y.-A. Ma, T. Chen, and E. Fox. A complete recipe for stochastic gradient MCMC. In *Advances in Neural Information Processing Systems*, pages 2899–2907, 2015.
- [23] Y.-A. Ma, T. Chen, L. Wu, and E. B. Fox. A unifying framework for devising efficient and irreversible MCMC samplers. *arXiv preprint arXiv:1608.05973*, 2016.
- [24] J. C. Mattingly, A. M. Stuart, and D. J. Higham. Ergodicity for SDEs and approximations: locally Lipschitz vector fields and degenerate noise. *Stochastic processes and their applications*, 101(2):185–232, 2002.
- [25] J. C. Mattingly, A. M. Stuart, and M. V. Tretyakov. Convergence of numerical time-averaging and stationary measures via Poisson equations. *SIAM Journal on Numerical Analysis*, 48(2):552–577, 2010.
- [26] K. L. Mengersen and R. L. Tweedie. Rates of convergence of the Hastings and Metropolis algorithms. *The Annals of Statistics*, 24(1):101–121, 1996.
- [27] G. Metafune, D. Pallara, and E. Priola. Spectrum of Ornstein-Uhlenbeck operators in L_p spaces with respect to invariant measures. *Journal of Functional Analysis*, 196(1):40–60, 2002.
- [28] N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller. Equation of state calculations by fast computing machines. *The journal of chemical physics*, 21(6):1087–1092, 1953.

- [29] S. P. Meyn and R. L. Tweedie. Markov chains and stochastic stability. communication and control engineering series. *Springer-Verlag London Ltd.*, *London*, 1:993, 1993.
- [30] S. P. Meyn and R. L. Tweedie. Stability of Markovian processes III: Foster-Lyapunov criteria for continuous-time processes. *Advances in Applied Probability*, 25(3):518–548, 1993.
- [31] S. P. Meyn and R. L. Tweedie. A survey of Foster-Lyapunov techniques for general state space Markov processes. In *Proceedings of the Workshop on Stochastic Stability and Stochastic Stabilization, Metz, France*. Citeseer, 1993.
- [32] A. Mira and C. J. Geyer. On non-reversible Markov chains. *Monte Carlo Methods, Fields Institute/AMS*, pages 95–110, 2000.
- [33] J. Møller, A. R. Syversveen, and R. P. Waagepetersen. Log Gaussian Cox processes. *Scandinavian journal of statistics*, 25(3):451–482, 1998.
- [34] R. M. Neal. Improving asymptotic variance of MCMC estimators: Nonreversible chains are better. *arXiv* preprint math/0407281, 2004.
- [35] R. M Neal. MCMC using Hamiltonian dynamics. In S. Brooks, A. Gelman, G. Jones, and X.-L. Meng, editors, *Handbook of Markov Chain Monte Carlo*, pages 113–162. CRC press, Boca Raton, 2011.
- [36] M. Ottobre, N. S. Pillai, F. J. Pinski, and A. M. Stuart. A function space hmc algorithm with second order Langevin diffusion limit. *Bernoulli*, 22(1):60–106, 2016.
- [37] L. Rey-Bellet and K. Spiliopoulos. Irreversible Langevin samplers and variance reduction: a large deviations approach. *Nonlinearity*, 28(7):2081, 2015.
- [38] L. Rey-Bellet and K. Spiliopoulos. Variance reduction for irreversible langevin samplers and diffusion on graphs. *Electronic Communications in Probability*, 20:16 pp., 2015.
- [39] G. O. Roberts and O. Stramer. Langevin diffusions and Metropolis-Hastings algorithms. *Methodology and computing in applied probability*, 4(4):337–357, 2002.
- [40] G. O. Roberts and R. L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341–363, 1996.
- [41] J. M. Sanz-Serna and M. P. Calvo. *Numerical Hamiltonian problems*, volume 7 of *Applied Mathematics and Mathematical Computation*. Chapman & Hall, London, 1994.
- [42] A. F. M. Smith and G. O. Roberts. Bayesian computation via the Gibbs sampler and related Markov chain Monte Carlo methods. *Journal of the Royal Statistical Society. Series B (Methodological)*, pages 3–23, 1993.
- [43] O Stramer and R. L. Tweedie. Langevin-type models I: Diffusions with given stationary distributions and their discretizations. *Methodology and Computing in Applied Probability*, 1(3):283–306, 1999.
- [44] O. Stramer and R. L. Tweedie. Langevin-type models II: self-targeting candidates for MCMC algorithms. *Methodology and Computing in Applied Probability*, 1(3):307–328, 1999.
- [45] D. Talay and L. Tubaro. Expansion of the global error for numerical schemes solving stochastic differential equations. *Stochastic Anal. Appl.*, 8(4):483–509 (1991), 1990.
- [46] L. Tierney. Markov chains for exploring posterior distributions. *the Annals of Statistics*, pages 1701–1728, 1994.
- [47] K. S. Turitsyn, M. Chertkov, and M. Vucelja. Irreversible Monte Carlo algorithms for efficient sampling. *Physica D: Nonlinear Phenomena*, 240(4):410–414, 2011.
- [48] S.-J. Wu, C.-R. Hwang, and M. T. Chu. Attaining the optimal gaussian diffusion acceleration. *Journal of Statistical Physics*, 155(3):571–590, 2014.
- [49] K. C. Zygalakis. On the existence and the applications of modified equations for stochastic differential equations. *SIAM J. Sci. Comput.*, 33(1):102–130, 2011.