

GEOMETRIC REALIZATIONS OF THE ACCORDION COMPLEX OF A DISSECTION

THIBAUT MANNEVILLE AND VINCENT PILAUD

ABSTRACT. Consider $2n$ points on the unit circle and a reference dissection D_\circ of the convex hull of the odd points. The accordion complex of D_\circ is the simplicial complex of non-crossing subsets of the diagonals with even endpoints that cross an accordion of the dissection D_\circ . In particular, this complex is an associahedron when D_\circ is a triangulation and a Stokes complex when D_\circ is a quadrangulation. In this paper, we provide geometric realizations (by polytopes and fans) of the accordion complex of any reference dissection D_\circ , generalizing known constructions arising from cluster algebras.

KEYWORDS. Permutahedra · Zonotopes · Associahedra · \mathbf{g} -, \mathbf{c} - and \mathbf{d} -vectors.

The $(n - 3)$ -dimensional *associahedron* is a polytope whose boundary complex is isomorphic to the reverse inclusion poset of non-crossing subsets of diagonals of a convex n -gon. Introduced in early works of D. Tamari [Tam51] and J. Stasheff [Sta63], it was first realized as a convex polytope by M. Haiman [Hai84] and C. Lee [Lee89], and later constructed by more systematic methods developed by several authors, in particular [GKZ08, Lod04, HL07, CSZ15]. Various relevant generalizations of the associahedron were introduced and studied, in particular secondary polytopes and fiber polytopes [GKZ08, BFS90], generalized associahedra [FZ03b, CFZ02, HLT11, Ste13, Hoh] in connection to cluster algebras [FZ02, FZ03a], graph associahedra [CD06, Pos09, FS05, Zel06, Pil13, MP16], or brick polytopes [PS12, PS15].

In a different context, Y. Baryshnikov [Bar01] introduced the simplicial complex of crossing-free subsets of the set of diagonals of a polygon that are in some sense compatible with a reference quadrangulation Q_\circ . Although the precise definition of compatibility is a bit technical in [Bar01], it turns out that a diagonal is compatible with Q_\circ if and only if it crosses a connected subset of diagonals of Q_\circ that we call *accordion* of Q_\circ . We thus call Y. Baryshnikov's simplicial complex the *accordion complex* $\mathcal{AC}(Q_\circ)$. A polytopal realization of $\mathcal{AC}(Q_\circ)$ was announced in [Bar01], but the explicit construction and its proof were never published as far as we know. Revisiting some combinatorial and algebraic properties of $\mathcal{AC}(Q_\circ)$, F. Chapoton [Cha16] raised three explicit challenges: first prove that the oriented dual graph of $\mathcal{AC}(Q_\circ)$ has a lattice structure extending the Tamari and Cambrian lattices [MHPS12, Rea06]; second construct geometric realizations of $\mathcal{AC}(Q_\circ)$ as fans and polytopes generalizing the known constructions of the associahedron; third show that the facets of $\mathcal{AC}(Q_\circ)$ are in bijection with other combinatorial objects called serpent nests [Cha16].

In [GM16], A. Garver and T. McConville defined and studied the accordion complex $\mathcal{AC}(D_\circ)$ of any reference dissection D_\circ (their presentation slightly differs as they use a compatibility condition on the dual tree of the dissection D_\circ , but the simplicial complex is the same). In this context, they settled F. Chapoton's lattice question, using lattice quotients of a lattice of biclosed sets. In this paper, we present geometric realizations of $\mathcal{AC}(D_\circ)$ for any reference dissection D_\circ , providing in particular an answer to F. Chapoton's geometric question. In fact, we present three methods to realize $\mathcal{AC}(D_\circ)$ based on constructions of the classical associahedron.

Our first method is based on the \mathbf{g} -vector fan. It belongs to a series of constructions of the (generalized) associahedra initiated by S. Shnider and S. Sternberg [SS93], popularised by J.-L. Loday [Lod04], developed by C. Hohlweg, C. Lange and H. Thomas [HL07, HLT11] using works of N. Reading and D. Speyer [Rea06, Rea07, RS09], and revisited by S. Stella [Ste13] and by V. Pilaud, F. Santos, and C. Stump [PS12, PS15]. It was recently extended by C. Hohlweg, V. Pilaud, and S. Stella [HPS17] to construct an associahedron parametrized by any initial triangulation. Here, we first extend to the D_\circ -accordion complex $\mathcal{AC}(D_\circ)$ the \mathbf{g} -vectors and \mathbf{c} -vectors defined in the context of cluster algebras by S. Fomin and A. Zelevinski [FZ07]. When D_\circ is a triangulation, our definitions coincide with those given in terms of triangulations and laminations for cluster algebras from surfaces by S. Fomin and D. Thurston [FT12]. We then show that the \mathbf{g} -vectors with respect to the dissection D_\circ support a complete simplicial fan $\mathcal{F}^{\mathbf{g}}(D_\circ)$ realizing the D_\circ -accordion

complex $\mathcal{AC}(D_\circ)$. Finally, we construct a D_\circ -accordiohedron $\text{Acco}(D_\circ)$ realizing the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(D_\circ)$ by deleting inequalities from the facet description of the D_\circ -zonotope $\text{Zono}(D_\circ)$ obtained as the Minkowski sum of all \mathbf{c} -vectors. See Figure 6 for an illustration of D_\circ -accordiohedra.

Our second method is based on the \mathbf{d} -vector fan. This construction is inspired from the original cluster fan of S. Fomin and A. Zelevinsky [FZ03a] later realized as a polytope by F. Chapoton, S. Fomin and A. Zelevinsky [CFZ02], and from the generalization of F. Santos [CSZ15] to construct a compatibility fan and an associahedron from any initial triangulation. For any reference dissection D_\circ , we associate to each diagonal a \mathbf{d} -vector which records the crossings of this diagonal with those of D_\circ . We show that the \mathbf{d} -vectors support a complete simplicial fan realizing the D_\circ -accordion complex $\mathcal{AC}(D_\circ)$ if and only if D_\circ contains no even interior cell. The polytopality of the resulting fan remains open in general, but was shown for arbitrary triangulations in [CSZ15].

Finally, our third method is based on projections of associahedra. Namely, for any dissection D_\circ and triangulation T_\circ such that $D_\circ \subseteq T_\circ$, the accordion complex $\mathcal{AC}(D_\circ)$ is a subcomplex of the simplicial associahedron $\mathcal{AC}(T_\circ)$. It turns out that the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(D_\circ)$ is then a section of the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(T_\circ)$ by a coordinate subspace. Therefore, the accordion complex $\mathcal{AC}(D_\circ)$ is realized by a projection of the associahedron $\text{Asso}(T_\circ)$ of [HPS17]. This point of view provides a complementary perspective on accordion complexes that leads on the one hand to more concise but less instructive proofs of combinatorial and geometric properties of the accordion complex (pseudomanifold, \mathbf{g} -vector fan, accordiohedron), and on the other hand to natural extensions to coordinate sections of the \mathbf{g} -vector fan in arbitrary cluster algebras.

The paper is organized as follows. Section 1 introduces the accordion complex and accordion lattice of a dissection D_\circ . We essentially follow the definitions and arguments of A. Garver and T. McConville [GM16], except that we prefer to work on the dissection D_\circ rather than on its dual graph. Section 2 is devoted to the generalization of the \mathbf{g} -vector fan and the associahedra of [HL07, HPS17]. Section 3 discusses the generalization of the construction of the \mathbf{d} -vector fan and associahedra of [FZ03a, CSZ15]. Finally, Section 4 shows that the accordion complex is realized by a projection of a well-chosen associahedron and presents related conjectures on cluster algebras, subcomplexes of the cluster complex, and sections of the \mathbf{g} -vector fan.

1. THE ACCORDION COMPLEX AND THE ACCORDION LATTICE

In this section, we define the accordion complex $\mathcal{AC}(D_\circ)$ of a dissection D_\circ , show that it is a pseudo-manifold, and define an orientation of its dual graph. Our definitions and proofs are essentially translations of the arguments of A. Garver and T. McConville [GM16] given in terms of the dual tree of the dissection D_\circ . However our presentation in terms of dissections is more convenient for our latter purposes.

1.1. The accordion complex. Let P be a convex polygon. We call *diagonals* of P the segments connecting two vertices of P . This includes both the internal diagonals and the external diagonals (or boundary edges) of P . A *dissection* of P is a set D of non-crossing internal diagonals of P . The *cells* of D are the closures of the connected components of P minus the diagonals of D . We denote by \bar{D} the dissection D together with all boundary edges of P . An *accordion* of D is a subset of \bar{D} which contains either no or two consecutive diagonals in each cell of D . A *subaccordion* of D is a subset of D formed by the diagonals between two given internal diagonals in an accordion of D . A *zigzag* of D is a subset $\{\delta_0, \dots, \delta_{p+1}\}$ of D where δ_i shares distinct endpoints with and




FIGURE 1. A dissection D (left) and three accordions whose zigzags are bolded (middle and right).




FIGURE 2. A hollow dissection D_o^{ex} , a solid D_o^{ex} -accordion diagonal whose corresponding hollow accordion is bolded, and two maximal solid D_o^{ex} -accordion dissections.

separates δ_{i-1} and δ_{i+1} for any $i \in [p]$. The *zigzag* of an accordion A is the subset of the diagonals of A that disconnect A . Note that we include boundary edges of P in the accordions of D , but not in the subaccordions nor in the zigzags of D . See Figure 1.

We consider $2n$ points on the unit circle labeled clockwise by $1_o, 2_\bullet, 3_o, 4_\bullet, \dots, (2n-1)_o, (2n)_\bullet$. We say that $1_o, \dots, (2n-1)_o$ are the *hollow vertices* while $2_\bullet, \dots, (2n)_\bullet$ are the *solid vertices*. The *hollow polygon* is the convex hull P_o of $1_o, \dots, (2n-1)_o$ while the *solid polygon* is the convex hull P_\bullet of $2_\bullet, \dots, (2n)_\bullet$. We simultaneously consider *hollow diagonals* δ_o (with two hollow vertices) and *solid diagonals* δ_\bullet (with two solid vertices), but we never consider diagonals with one hollow vertex and one solid vertex. Similarly, we consider *hollow dissections* D_o (of the hollow polygon, with only hollow diagonals) and *solid dissections* D_\bullet (of the solid polygon, with only solid diagonals), but never mix hollow and solid diagonals in a dissection. To help distinguishing them, hollow (resp. solid) vertices and diagonals appear red (resp. blue) in all pictures.

We fix an arbitrary reference hollow dissection D_o . A solid diagonal δ_\bullet is a *D_o -accordion diagonal* if the hollow diagonals of \bar{D}_o crossed by δ_\bullet form an accordion of D_o . In other words, δ_\bullet cannot enter and exit a cell of D_o using two non-incident diagonals. For example, note that for any hollow diagonal $i_o j_o \in \bar{D}_o$, the solid diagonals $(i-1)_\bullet(j-1)_\bullet$ and $(i+1)_\bullet(j+1)_\bullet$ are D_o -accordion diagonals (here and throughout, labels are considered modulo $2n$). In particular, all boundary edges of the solid polygon are D_o -accordion diagonals. A *D_o -accordion dissection* is a set of non-crossing internal D_o -accordion diagonals. We call *D_o -accordion complex* the simplicial complex $\mathcal{AC}(D_o)$ of D_o -accordion dissections.

Example 1. As a running example, we consider the reference dissection D_o^{ex} of Figure 2 (left). Examples of maximal D_o^{ex} -accordion dissections are given in Figure 2 (right). The D_o^{ex} -accordion complex is illustrated in Figure 3 (left).

Remark 2. Special reference hollow dissections D_o give rise to special accordion complexes $\mathcal{AC}(D_o)$:

- ◊ If D_o is the empty dissection with the whole hollow polygon as unique cell, then the D_o -accordion complex $\mathcal{AC}(D_o)$ is reduced to the empty D_o -accordion dissection.
- ◊ If D_o has a unique internal diagonal, then the D_o -accordion complex $\mathcal{AC}(D_o)$ is a segment.
- ◊ For a hollow triangulation T_o , all solid diagonals are T_o -accordions, so that the T_o -accordion complex $\mathcal{AC}(T_o)$ is the simplicial associahedron.
- ◊ For a hollow quadrangulation Q_o , a solid diagonal is a Q_o -accordion if and only if it does not cross two opposite edges of a quadrangle of Q_o . The Q_o -accordion complex $\mathcal{AC}(Q_o)$ is thus the Stokes complex defined by Y. Baryshnikov [Bar01] and studied by F. Chapoton [Cha16].

Remark 3. Following the original definition of the non-crossing complex of A. Garver and T. McConville [GM16], the accordion complex could equivalently be defined in terms of the dual tree D_o^* of D_o (with one node in each cell of D and one edge connecting two adjacent cells). For example, a diagonal $u_\bullet v_\bullet$ is a D_o -accordion diagonal if and only if any two consecutive edges of the (unique) path between the leaves u_\bullet^* and v_\bullet^* in D_o^* belong to the boundary of a face of the complement of D_o^* in the unit disk. The *g*-, *c*- and *d*-vectors defined in Section 2.1 could as well be defined in terms of D_o^* , but we find more convenient to work directly with dissections, in particular in Section 3.




FIGURE 3. The D_0^{ex} -accordion complex (left) and the D_0^{ex} -accordion lattice (right), oriented from bottom to top, for the reference hollow dissection D_0^{ex} of Figure 2 (left).

Remark 4. Assume that D_0 has a cell C_0 containing p boundary edges of the hollow polygon P_0 . Let C_0^1, \dots, C_0^p denote the p (possibly empty) connected components of the hollow polygon minus C_0 . For $i \in [p]$, let D_0^i denote the dissection formed by the cell C_0 together with the cells of D_0 in C_0^i . Since no D_0 -accordion can contain internal diagonals from distinct dissections D_0^i and D_0^j (with $i \neq j$), the D_0 -accordion complex is the join of the D_0^i -accordion complexes: $\mathcal{AC}(D_0) = \mathcal{AC}(D_0^1) * \dots * \mathcal{AC}(D_0^p)$. In particular, we can do the following reductions:

- (i) If a non-triangular cell of D_0 has two consecutive boundary edges γ_0, δ_0 of the hollow polygon, then contracting γ_0 and δ_0 to a single boundary edge preserves the D_0 -accordion complex.
- (ii) If a cell of D_0 has two non-consecutive boundary edges of the hollow polygon, then the D_0 -accordion complex is a join of smaller accordion complexes.

In all the examples of the paper, we therefore only consider dissections where any non-triangular cell of D_0 has at most one boundary edge. All our constructions work in general, but are just obtained as products or joins of the non-degenerate situation.

Remark 5. The links in an accordion complex are joins of accordion complexes. Namely, consider a D_0 -accordion dissection D_\bullet with cells $C_\bullet^1, \dots, C_\bullet^p$. Let D_\bullet^i denote the hollow dissection obtained from D_0 by contracting all hollow boundary edges which do not cross C_\bullet^i . Then the link of D_\bullet in $\mathcal{AC}(D_0)$ is isomorphic to the join $\mathcal{AC}(D_\bullet^1) * \dots * \mathcal{AC}(D_\bullet^p)$.

1.2. Pseudo-manifold. We now prove that the accordion complex $\mathcal{AC}(D_0)$ is a *pseudo-manifold*, i.e. that it is:

- (i) *pure*: all maximal D_0 -accordion dissections have as many diagonals as D_0 , and
- (ii) *thin*: any codimension 1 simplex of $\mathcal{AC}(D_0)$ is contained in exactly two maximal D_0 -accordion dissections.

We follow the arguments of A. Garver and T. McConville [GM16] (except that they work on the dual tree of the dissection D_0). A much more concise but less instructive proof of the pseudomanifold property will be derived from geometric considerations in Remark 56.

Recall that we denote by \bar{D}_0 the set formed by D_0 together with all boundary edges of the hollow polygon. An *angle* $u_0 v_0 w_0$ of \bar{D}_0 is a pair $\{u_0 v_0, v_0 w_0\}$ of two consecutive diagonals of \bar{D}_0 around a common vertex v_0 , called *apex*. Note that \bar{D}_0 has $2|D_0| + n = 2|\bar{D}_0| - n$ angles. We say that a solid vertex p_\bullet belongs to a hollow angle $u_0 v_0 w_0$ if it lies in the cone generated by the edges $v_0 u_0$ and $v_0 w_0$ of the angle. The main observation is given in the following statement.




FIGURE 4. Two maximal D_0 -accordion dissection $D_•$ (left) and $D'_•$ (right) related by the flip of $\delta_•$ to $\delta'_•$. The angles of D_0 closed by $\delta_•$ and $\delta'_•$ are shaded. The flip is oriented from $D_•$ to $D'_•$.

Lemma 6. *Let $D_•$ be a maximal D_0 -accordion dissection, and let $p_•, q_•, r_•, s_•$ denote four consecutive vertices of a cell $C_•$ of $D_•$ (with possibly $p_• = s_•$ if $C_•$ is a triangle). Then $p_•$ and $s_•$ belong to the same angle of the accordion of \bar{D}_0 crossed by $q_•r_•$.*

Proof. Let A_0 be the accordion of \bar{D}_0 crossed by $q_•r_•$. Assume that $p_•$ and $s_•$ belong to distinct angles of A_0 . Then they are separated by a diagonal ε_0 of A_0 . Therefore, there are two boundary edges $q_•r_•$ and $u_•v_•$ of $C_•$ with distinct vertices such that the hollow diagonal ε_0 separates the vertices $q_•, u_•$ from the vertices $r_•, v_•$. Let $\gamma_0^1, \dots, \gamma_0^i = \varepsilon_0, \dots, \gamma_0^a$ (resp. $\delta_0^1, \dots, \delta_0^j = \varepsilon_0, \dots, \delta_0^b$) denote the diagonals of D_0 crossed by $q_•r_•$ from $q_•$ to $r_•$ (resp. crossed by $u_•v_•$ from $u_•$ to $v_•$). Then the hollow diagonals $\gamma_0^1, \dots, \gamma_0^i = \varepsilon_0 = \delta_0^j, \dots, \delta_0^b$ which are crossed by $q_•v_•$ also form an accordion. It follows that $D_•$ is not maximal as we can still include $q_•v_•$. \square

Consider now an angle $u_0v_0w_0$ of \bar{D}_0 . In any maximal D_0 -accordion dissection $D_•$, the set $X_•$ of diagonals of $\bar{D}_•$ that cross both u_0v_0 and v_0w_0 is non-empty (since it contains the boundary edge $(v-1)_•(v+1)_•$) and totally ordered (since the diagonals of $D_•$ do not cross). We say that the angle $u_0v_0w_0$ is *closed* by the farthest diagonal of $X_•$ from v_0 in the dissection $\bar{D}_•$. Note that each angle of \bar{D}_0 is closed by precisely one diagonal of $\bar{D}_•$. The following lemma is stated and proved in [GM16] in terms of the dual tree D_0^* of the dissection D_0 .

Lemma 7 ([GM16]). *For any maximal D_0 -accordion dissection $D_•$, each internal diagonal $\delta_•$ of $D_•$ closes two angles of \bar{D}_0 (one apex on each side of $\delta_•$) while each boundary edge of the solid polygon closes one angle of \bar{D}_0 . Therefore the accordion complex $\mathcal{AC}(D_0)$ is pure of dimension $|D_0|$.*

Proof. The first sentence is a consequence of Lemma 6: for any four consecutive vertices $p_•, q_•, r_•, s_•$ of a cell of $\bar{D}_•$, the diagonal $q_•r_•$ closes the unique angle of the accordion of \bar{D}_0 crossed by $q_•r_•$ that contains the vertices $p_•$ and $s_•$. Therefore, $q_•r_•$ closes precisely two angles (resp. one angle) of D_0 if it is an internal diagonal (resp. a boundary edge of the solid polygon). We finally obtain by double-counting that $2|D_0| + n = |\{\text{angles of } \bar{D}_0\}| = 2|D_•| + n$ and thus $|D_•| = |D_0|$ for any maximal D_0 -accordion dissection $D_•$. \square

We are now ready to prove that the D_0 -accordion complex is thin, *i.e.* that each internal diagonal of a maximal D_0 -accordion dissection can be flipped into a unique other internal diagonal to form a new maximal D_0 -accordion dissection. The following statement is illustrated in Figure 4.

Lemma 8 ([GM16]). *Let $D_•$ be a maximal D_0 -accordion dissection and $\delta_•$ be a diagonal of $D_•$. Let u_0 and v_0 be the apices of the angles of D_0 closed by $\delta_•$, let $\mu_•$ and $\nu_•$ denote the edges of the cells of $D_•$ containing $\delta_•$, which separate $\delta_•$ from u_0 and v_0 respectively, and let $Q_•$ denote the quadrilateral defined by the four vertices of $\mu_•$ and $\nu_•$. Note that $\delta_•$ is a diagonal of $Q_•$, and let $\delta'_•$ denote the other diagonal. Then $D'_• := D_• \triangle \{\delta_•, \delta'_•\}$ is a maximal D_0 -accordion dissection, and $D_•$ and $D'_•$ are the only maximal D_0 -accordion dissections containing $D_• \setminus \{\delta_•\}$. In other words, the accordion complex $\mathcal{AC}(D_0)$ is thin.*

Proof. We first observe that δ'_\bullet is a D_\circ -accordion diagonal, since the edges of \bar{D}_\circ crossed by δ'_\bullet are obtained by merging three subaccordions of D_\circ : the subaccordion formed by the diagonals of \bar{D}_\circ crossed by μ_\bullet but not δ_\bullet nor ν_\bullet , the subaccordion formed by the diagonals of \bar{D}_\circ crossed by δ_\bullet , μ_\bullet and ν_\bullet , and the subaccordion formed by the diagonals of \bar{D}_\circ crossed by ν_\bullet but not δ_\bullet nor μ_\bullet . Moreover, δ_\bullet and δ'_\bullet are the only D_\circ -accordion diagonals compatible with $D_\bullet \setminus \{\delta_\bullet\}$. Indeed, any other such diagonal would cross δ_\bullet and δ'_\bullet (by maximality of D_\bullet and D'_\bullet), and thus also the subaccordion A_\circ of D_\circ crossed by δ_\bullet and δ'_\bullet (because it cannot cross μ and ν). But it would then improperly intersect the two cells of D_\circ containing precisely one diagonal of A_\circ . \square

The *D_\circ -accordion flip graph* is the dual graph $\mathcal{AFG}(D_\circ)$ of the D_\circ -accordion complex: its vertices are the maximal D_\circ -accordion dissections, and its edges are the *flips* between them, *i.e.* the pairs $\{D_\bullet, D'_\bullet\}$ of maximal D_\circ -accordion dissections with $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$. See Figure 3 (right).

1.3. The accordion lattice. We now define a natural orientation on the D_\circ -accordion flip graph. We use the notations of Lemma 8, where $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ and $\delta_\bullet, \delta'_\bullet$ are the two diagonals of the quadrilateral defined by μ_\bullet, ν_\bullet . Observe that one of the path $\mu_\bullet \delta_\bullet \nu_\bullet$ and $\mu_\bullet \delta'_\bullet \nu_\bullet$ forms a Σ while the other forms a Z , see Figure 4. We then orient the flip from the dissection containing the Σ to that containing the Z . See Figure 3 (right) for an illustration of D_\circ -accordion oriented flip graph (where the graph is oriented from bottom to top).

A. Garver and T. McConville introduced a natural closure on sets of D_\circ -subaccordions, and showed that the inclusion poset of biclosed sets of D_\circ -subaccordions is a well-behaved lattice (namely, semidistributive, congruence-uniform and polygonal). Then, they introduced a lattice congruence map from biclosed sets of D_\circ -subaccordions to maximal D_\circ -accordion dissections, which imply the following statement.

Theorem 9 ([GM16]). *The D_\circ -accordion oriented flip graph is the Hasse diagram of a lattice, that we call the D_\circ -accordion lattice and denote by $\mathcal{AL}(D_\circ)$.*

In particular, the D_\circ -accordion oriented flip graph is connected and acyclic, and has a unique source $D_\bullet^- := \{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$ (obtained by slightly rotating D_\circ counterclockwise) and a unique sink $D_\bullet^+ := \{(i+1)_\bullet(j+1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$ (obtained by slightly rotating D_\circ clockwise).

Remark 10. Following Remark 2, note that special reference hollow dissections D_\circ give rise to special accordion lattices $\mathcal{AL}(D_\circ)$, as it was already observed in [GM16]:

- ◊ For a fan triangulation F_\circ (*i.e.* where all internal diagonals are incident to a common vertex), the F_\circ -accordion lattice $\mathcal{AL}(F_\circ)$ is the famous Tamari lattice [Tam51, MHPS12] defined equivalently by slope increasing flips on triangulations of a convex polygon, by right rotations on binary trees, or by flips on Dyck paths.
- ◊ In general, accordion lattices of accordion triangulations (*i.e.* with no interior triangle) precisely correspond to type A Cambrian lattices defined by N. Reading [Rea06].
- ◊ For an arbitrary triangulation T_\circ (with or without interior triangle), the T_\circ -accordion oriented flip graph $\mathcal{AFG}(A_\circ)$ was defined by T. Brüstle, G. Dupont and M. Pérotin [BDP14].
- ◊ For a quadrangulation Q_\circ , the Q_\circ -accordion lattice $\mathcal{AL}(Q_\circ)$ is the Stokes poset on Q_\circ -compatible quadrangulations studied by F. Chapoton [Cha16].

Remark 11. Following Remark 4, assume that D_\circ has a cell containing p boundary edges of the hollow polygon, and consider the dissections $D_\circ^1, \dots, D_\circ^p$ as in Remark 4. Then the D_\circ -accordion lattice is the Cartesian product of the D_\circ^i -accordion lattices: $\mathcal{AL}(D_\circ) = \mathcal{AL}(D_\circ^1) \times \dots \times \mathcal{AL}(D_\circ^p)$. In particular, if two consecutive boundary edges $\gamma_\circ, \delta_\circ$ of the hollow polygon belong to the same non-triangular cell of D_\circ , then contracting γ_\circ and δ_\circ to a single boundary edge preserves the D_\circ -accordion lattice. This shows in particular that the D_\circ -accordion lattice of a ribbon dissection D_\circ is a Cambrian lattice, as conjectured for quadrangulations in [Cha16] and proved in [BMP16].

Remark 12. Call *cell-sequence* of a dissection the sequence whose i th entry is its number of $(i+2)$ -cells. For example, the dissection of Figure 2 (left) has cell-sequence $3, 1, 0^\infty$ and all $(p+2)$ -angulations of a $(pm+2)$ -gon have cell-sequence $0^{p-1}, m, 0^\infty$. Observe that the flip preserves the cell-sequence. Therefore, all D_\circ -accordion dissections have the same cell-sequence as D_\circ .

We conclude this section with a reciprocity result on accordion dissections.

Proposition 13. *Let D_\circ be a hollow dissection and D_\bullet be a solid dissection. Then D_\bullet is a maximal D_\circ -accordion dissection if and only if D_\circ is a maximal D_\bullet -accordion dissection.*

Proof. Since $D_\bullet^- := \{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$ and $D_\bullet^+ := \{(i+1)_\bullet(j+1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$ are both D_\circ -accordion dissections, we already know that D_\circ is a D_\bullet^- -accordion dissection. Observe now in Figure 4 that if D_\bullet and D'_\bullet are maximal D_\circ -accordion dissections connected by a flip, then D_\circ is a D_\bullet -accordion dissection if and only if it is a D'_\bullet -accordion dissection. Indeed, if δ_\bullet belongs to the zigzag of the D_\bullet -accordion A_\bullet of a hollow diagonal δ_\circ , then δ_\circ crosses both μ_\bullet and ν_\bullet , but then it also crosses δ'_\bullet and thus the D'_\bullet -accordion $A_\bullet \triangle \{\delta_\bullet, \delta'_\bullet\}$. Since the D_\circ -accordion flip graph is connected, we obtain that D_\circ is a D_\bullet -accordion dissection for any maximal D_\circ -accordion dissection D_\bullet . Finally, maximality follows since all maximal D_\circ -accordion dissections have $|D_\circ|$ diagonals. The equivalence follows by symmetry. \square

2. THE \mathbf{g} -VECTOR FAN

In this Section, we construct accordiohedra using \mathbf{g} - and \mathbf{c} -vectors. Our construction is in the same spirit as the Cambrian fans of N. Reading and D. Speyer [Rea06, Rea07, RS09] and their polytopal realizations by C. Hohlweg, C. Lange and H. Thomas [HL07, HLT11], recently extended in [HPS17] to any initial triangulation, acyclic or not. A different approach to the \mathbf{g} -vector fan together with an alternative polytopal realization will be presented in Section 4.

2.1. \mathbf{g} - and \mathbf{c} -vectors. Consider a hollow dissection D_\circ and a solid dissection D_\bullet that are maximal accordion dissection of each other (see Proposition 13), and let $\delta_\circ \in D_\circ$ and $\delta_\bullet \in D_\bullet$. When δ_\circ crosses δ_\bullet , we let μ_\circ and ν_\circ be the other diagonals of \bar{D}_\circ crossed by δ_\bullet in the two cells of D_\circ containing δ_\circ . We say that δ_\bullet *slaloms* on δ_\circ if $\mu_\circ \delta_\circ \nu_\circ$ forms a path, and we define $\varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet)$ to be 1, -1, or 0 depending on whether $\mu_\circ \delta_\circ \nu_\circ$ forms a \mathbf{Z} , a $\mathbf{\Sigma}$, or a $\mathbf{\Psi}$. Similarly we let μ_\bullet and ν_\bullet be the other diagonals of \bar{D}_\bullet crossed by δ_\circ in the two cells of D_\bullet containing δ_\bullet , we say that δ_\circ *slaloms* on δ_\bullet if $\mu_\bullet \delta_\bullet \nu_\bullet$ forms a path, and we define $\varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet)$ to be 1, -1, or 0 depending on whether $\mu_\bullet \delta_\bullet \nu_\bullet$ forms a $\mathbf{\Sigma}$, a \mathbf{Z} , or a $\mathbf{\Psi}$. Note that the sign convention for $\varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet)$ and $\varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet)$ is opposite: the reciprocity already observed in Proposition 13 naturally reverses the orientation. More informally, we exchange the role of hollow and solid dissections by looking at the picture from the opposite side of the blackboard, which of course reverses the orientation. Finally, if δ_\circ and δ_\bullet do not cross, then we let $\varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet) = \varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet) = 0$. Let $(\mathbf{e}_{\delta_\circ})_{\delta_\circ \in D_\circ}$ denote the canonical basis of \mathbb{R}^{D_\circ} . As in [HPS17], we define the following vectors:

- (i) the **\mathbf{g} -vector** of δ_\bullet with respect to D_\circ is $\mathbf{g}(D_\circ \mid \delta_\bullet) := \sum_{\delta_\circ \in D_\circ} \varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet) \mathbf{e}_{\delta_\circ}$. We also define $\mathbf{g}(D_\circ \mid D_\bullet) := \{\mathbf{g}(D_\circ \mid \delta_\bullet) \mid \delta_\bullet \in D_\bullet\}$.
- (ii) the **\mathbf{c} -vector** of $\delta_\bullet \in D_\bullet$ with respect to D_\circ is $\mathbf{c}(D_\circ \mid \delta_\bullet \in D_\bullet) := \sum_{\delta_\circ \in D_\circ} \varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet) \mathbf{e}_{\delta_\circ}$. We denote by $\mathbf{c}(D_\circ \mid D_\bullet) := \{\mathbf{c}(D_\circ \mid \delta_\bullet \in D_\bullet) \mid \delta_\bullet \in D_\bullet\}$ the set of \mathbf{c} -vectors of the diagonals of D_\bullet and by $\mathbf{C}(D_\circ) := \bigcup_{D_\bullet} \mathbf{c}(D_\circ \mid D_\bullet)$ the set of all \mathbf{c} -vectors with respect to D_\circ .

Example 14. Consider the hollow dissection $D_\circ^{\text{ex}} = \{3_\circ 7_\circ, 3_\circ 13_\circ, 9_\circ 13_\circ\}$ and the rightmost solid dissection $D_\bullet^{\text{ex}} = \{2_\bullet 6_\bullet, 2_\bullet 10_\bullet, 10_\bullet 14_\bullet\}$ of Figure 2. Then we have for example

- $\diamond \varepsilon_\circ(3_\circ 13_\circ \in D_\circ^{\text{ex}} \mid 2_\bullet 10_\bullet) = 1$ since the path $1_\circ - 3_\circ - 13_\circ - 9_\circ$ forms a \mathbf{Z} ,
- $\diamond \varepsilon_\circ(9_\circ 13_\circ \in D_\circ^{\text{ex}} \mid 2_\bullet 10_\bullet) = -1$ since the path $3_\circ - 13_\circ - 9_\circ - 11_\circ$ forms a $\mathbf{\Sigma}$, and
- $\diamond \varepsilon_\circ(3_\circ 13_\circ \in D_\circ^{\text{ex}} \mid 2_\bullet 6_\bullet) = 0$ since 3_\circ connects $1_\circ, 13_\circ, 7_\circ$ as a $\mathbf{\Psi}$.

Moreover, we have

$$\begin{aligned} \mathbf{g}(D_\circ^{\text{ex}} \mid 2_\bullet 6_\bullet) &= \mathbf{e}_{3_\circ 7_\circ}, & \mathbf{c}(D_\circ^{\text{ex}} \mid 2_\bullet 6_\bullet \in D_\bullet^{\text{ex}}) &= \mathbf{e}_{3_\circ 7_\circ}, \\ \mathbf{g}(D_\circ^{\text{ex}} \mid 2_\bullet 10_\bullet) &= \mathbf{e}_{3_\circ 13_\circ} - \mathbf{e}_{9_\circ 13_\circ}, & \mathbf{c}(D_\circ^{\text{ex}} \mid 2_\bullet 10_\bullet \in D_\bullet^{\text{ex}}) &= \mathbf{e}_{3_\circ 13_\circ}, \\ \mathbf{g}(D_\circ^{\text{ex}} \mid 10_\bullet 14_\bullet) &= -\mathbf{e}_{9_\circ 13_\circ}, & \mathbf{c}(D_\circ^{\text{ex}} \mid 10_\bullet 14_\bullet \in D_\bullet^{\text{ex}}) &= -\mathbf{e}_{3_\circ 13_\circ} - \mathbf{e}_{9_\circ 13_\circ}. \end{aligned}$$

Example 15. For any hollow diagonal $i_\circ j_\circ \in D_\circ$, we have

$$\begin{aligned} \mathbf{g}(D_\circ \mid (i-1)_\bullet(j-1)_\bullet) &= -\mathbf{e}_{i_\circ j_\circ}, & \mathbf{c}(D_\circ \mid (i-1)_\bullet(j-1)_\bullet \in D_\bullet^-) &= -\mathbf{e}_{i_\circ j_\circ}, \\ \mathbf{g}(D_\circ \mid (i+1)_\bullet(j+1)_\bullet) &= \mathbf{e}_{i_\circ j_\circ}, & \mathbf{c}(D_\circ \mid (i+1)_\bullet(j+1)_\bullet \in D_\bullet^+) &= \mathbf{e}_{i_\circ j_\circ}. \end{aligned}$$

Remark 16. For a hollow triangulation T_\circ , our definitions of \mathbf{g} - and \mathbf{c} -vectors coincide with the shear coordinates of S. Fomin and D. Thurston [FT12], defined in the much more general context of cluster algebras on surfaces [FST08].

Remark 17. Consider the quiver $Q(D_\circ)$ of the reference dissection D_\circ , with one node on each internal diagonal of D_\circ and one arrow between two diagonals counter-clockwise consecutive around a cell of D_\circ . Let $W(D_\circ)$ be the reflection group with Dynkin diagram $Q(D_\circ)$. Then all \mathbf{g} -vectors of the D_\circ -accordion diagonals are weights of $W(D_\circ)$ and all \mathbf{c} -vectors of $\mathbf{C}(D_\circ)$ are roots of $W(D_\circ)$.

Remark 18. Informally, the \mathbf{g} - and \mathbf{c} -vectors can be interpreted as follows:

- (i) The \mathbf{g} -vector $\mathbf{g}(D_\circ | \delta_\bullet)$ has coordinate 1 and -1 alternating along the zigzag of the accordion crossed by δ_\bullet in D_\circ , and coordinate 0 on all other diagonals of D_\circ .
- (ii) The \mathbf{c} -vector $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$ is, up to a sign, the characteristic vector of the diagonals of the subaccordion of D_\circ crossed by both μ_\bullet and ν_\bullet of Lemma 8 (see Figure 4). Thus, any \mathbf{c} -vector is either *positive* (only non-negative coordinates) or *negative* (only non-positive coordinates).

In fact, the \mathbf{g} -vectors are clearly in bijection with the accordions and with the zigzags in D_\circ . In contrast, many $\delta_\bullet \in D_\bullet$ produce the same \mathbf{c} -vector $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$. For example, if two dissections D_\bullet, D'_\bullet contain δ_\bullet and have the same cells incident to δ_\bullet , then $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) = \mathbf{c}(D_\circ | \delta_\bullet \in D'_\bullet)$. The set of \mathbf{c} -vectors $\mathbf{C}(D_\circ)$ without repetitions can be understood as follows.

Lemma 19. *There are bijections between:*

- ◊ the negative (resp. positive) \mathbf{c} -vectors of $\mathbf{C}(D_\circ)$,
- ◊ the subaccordions of D_\circ ,
- ◊ the D_\circ -accordion diagonals not in the source dissection $D_\bullet^- := \{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$ (resp. not in the sink dissection $D_\bullet^+ := \{(i+1)_\bullet(j+1)_\bullet \mid i_\circ j_\circ \in D_\circ\}$).

Proof. By Remark 18 (ii), the support of any \mathbf{c} -vector is a subaccordion of D_\circ . Reciprocally, let A_\circ be a subaccordion of D_\circ , let C_\circ and C'_\circ denote the two cells of D_\circ containing exactly one diagonal of A_\circ , and let $p_\circ, q_\circ, r_\circ, s_\circ$ (resp. $p'_\circ, q'_\circ, r'_\circ, s'_\circ$) denote the four consecutive vertices in clockwise order around C_\circ (resp. around C'_\circ) such that $q_\circ r_\circ$ (resp. $q'_\circ r'_\circ$) is the diagonal of A_\circ in C_\circ (resp. in C'_\circ). Let $\delta_\bullet := (s-1)_\bullet(s'-1)_\bullet$, $\mu_\bullet := (p+1)_\bullet(s'-1)_\bullet$ and $\nu_\bullet := (p'+1)_\bullet(s-1)_\bullet$ and consider any D_\circ -accordion dissection D_\bullet containing $\{\mu_\bullet, \delta_\bullet, \nu_\bullet\}$. Then A_\circ is precisely the support of the negative \mathbf{c} -vector $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$. Finally, we have associated to the subaccordion A_\circ of D_\circ a D_\circ -diagonal $\delta_\bullet = (s-1)_\bullet(s'-1)_\bullet$ which cannot be in D_\bullet^- as otherwise $s_\circ s'_\circ$ would cross $q_\circ r_\circ$. Reciprocally, A_\circ is precisely the set of diagonals of D_\circ crossed by δ_\bullet and not incident to s_\circ or s'_\circ . \square

The \mathbf{g} -vectors and \mathbf{c} -vectors are connected in the following two statements, inspired and motivated by classical analogues in cluster algebra theory.

Proposition 20. *For any maximal D_\circ -accordion dissection D_\bullet , the set of \mathbf{g} -vectors $\mathbf{g}(D_\circ | D_\bullet)$ and the set of \mathbf{c} -vectors $\mathbf{c}(D_\circ | D_\bullet)$ form dual bases.*

Proof. Given two solid diagonals $\gamma_\bullet, \delta_\bullet$ of D_\bullet , we want to compute $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) \rangle$. By Remark 18 (i), the \mathbf{g} -vector $\mathbf{g}(D_\circ | \gamma_\bullet)$ has coordinate ± 1 alternating along the zigzag Z_\circ of the accordion crossed by γ_\bullet in D_\circ , and coordinate 0 on all other diagonals of D_\circ . Moreover, by Remark 18 (ii), the \mathbf{c} -vector $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$ has coordinate ± 1 on the diagonals of D_\circ which slalom on δ_\bullet in D_\bullet , and coordinate 0 on all other diagonals of D_\circ . We thus need to understand how the diagonals of Z_\circ slalom on δ_\bullet in D_\bullet . Observe that there is an even (resp. odd) number of hollow diagonals of Z_\circ that slalom on δ_\bullet when $\delta_\bullet \neq \gamma_\bullet$ (resp. when $\delta_\bullet = \gamma_\bullet$). Moreover, since they are non-crossing, all hollow diagonals of Z_\circ slaloming on δ_\bullet do it the same way (either all as a Σ or all as a Z). Finally, when $\gamma_\bullet = \delta_\bullet$, consider the first hollow diagonal δ_\circ of the zigzag Z_\circ which slaloms on δ_\bullet . Then δ_\circ slaloms on δ_\bullet in the opposite way as δ_\bullet slaloms on δ_\circ . This shows that

$$\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet) \rangle = \sum_{\delta_\circ \in D_\circ} \varepsilon_\circ(\delta_\circ \in D_\circ \mid \gamma_\bullet) \cdot \varepsilon_\bullet(\delta_\circ \mid \delta_\bullet \in D_\bullet) = \mathbb{1}_{\gamma_\bullet = \delta_\bullet},$$

since we sum an even number of alternating ± 1 when $\gamma_\bullet \neq \delta_\bullet$, and an odd number of alternating ± 1 starting by a 1 when $\gamma_\bullet = \delta_\bullet$. In other words, $\mathbf{g}(D_\circ | D_\bullet)$ and $\mathbf{c}(D_\circ | D_\bullet)$ form dual bases. \square

Proposition 21. *Let D_\circ be a hollow dissection and D_\bullet be a solid dissection such that D_\circ and D_\bullet are maximal accordion dissection of each other (see Proposition 13). Then*

$$\mathbf{g}(D_\circ | D_\bullet) = -\mathbf{c}(D_\bullet | D_\circ)^t \quad \text{and} \quad \mathbf{c}(D_\circ | D_\bullet) = -\mathbf{g}(D_\bullet | D_\circ)^t,$$

where we consider the sets of \mathbf{g} -vectors $\mathbf{g}(D_\circ | D_\bullet)$ and \mathbf{c} -vectors $\mathbf{c}(D_\circ | D_\bullet)$ as matrices in $\mathbb{R}^{D_\circ \times D_\bullet}$, and M^t denotes the transpose of a matrix M .

Proof. We immediately derive from the definitions that for any $\delta_\circ \in D_\circ$ and $\delta_\bullet \in D_\bullet$,

$$\mathbf{g}(D_\circ | D_\bullet)_{(\delta_\circ, \delta_\bullet)} = \varepsilon_\circ(\delta_\circ \in D_\circ | \delta_\bullet) = -\varepsilon_\bullet(\delta_\bullet | \delta_\circ \in D_\circ) = -\mathbf{c}(D_\bullet | D_\circ)_{(\delta_\bullet, \delta_\circ)},$$

which shows $\mathbf{g}(D_\circ | D_\bullet) = -\mathbf{c}(D_\bullet | D_\circ)^t$. The other equality follows by exchanging D_\circ and D_\bullet . \square

Corollary 22. *For any maximal D_\circ -accordion dissection D_\bullet , we have the following [sign coherence](#):*

- (i) *for any $\delta_\bullet \in D_\bullet$, all coordinates of the \mathbf{c} -vector $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$ have the same sign,*
- (ii) *for any $\delta_\circ \in D_\circ$, the δ_\circ -coordinates of all \mathbf{g} -vectors $\mathbf{g}(D_\circ | \delta_\bullet)$ for $\delta_\bullet \in D_\bullet$ have the same sign.*

Proof. Point (i) was already seen in Remark 18(ii), and Point (ii) follows by Proposition 21. \square

2.2. \mathbf{c} -vector fan and D_\circ -zonotope. Call \mathbf{c} -vector fan of D_\circ the complete polyhedral fan $\mathcal{F}^c(D_\circ)$ defined by the arrangement of the linear hyperplanes orthogonal to the \mathbf{c} -vectors of $\mathbf{C}(D_\circ)$. Be careful: contrarily to the \mathbf{g} - and \mathbf{d} -vector fans defined later, the \mathbf{c} -vectors are not the rays of $\mathcal{F}^c(D_\circ)$ but the normal vectors of the hyperplanes supporting the facets of $\mathcal{F}^c(D_\circ)$.

We call D_\circ -zonotope the Minkowski sum $\text{Zono}(D_\circ)$ of all \mathbf{c} -vectors:

$$\text{Zono}(D_\circ) := \sum_{\mathbf{c} \in \mathbf{C}(D_\circ)} \mathbf{c}.$$

The normal fan of the D_\circ -zonotope $\text{Zono}(D_\circ)$ is the \mathbf{c} -vector fan $\mathcal{F}^c(D_\circ)$. Note that the \mathbf{c} -vector fan is not always simplicial, and thus the D_\circ -zonotope $\text{Zono}(D_\circ)$ is not always simple. See Figure 6.

Example 23. Consider an accordion dissection $A_\circ = \{\delta_\circ^1, \dots, \delta_\circ^{|A_\circ|}\}$, with diagonals labeled such that δ_\circ^k and δ_\circ^{k+1} belong to the same cell of A_\circ for all k . Identifying $\mathbf{e}_{\delta_\circ^k}$ to the simple root $\mathbf{f}_k - \mathbf{f}_{k+1}$ of type $A_{|A_\circ|}$, the \mathbf{c} -vectors of $\mathbf{C}(A_\circ)$ are all roots $\pm(\mathbf{f}_i - \mathbf{f}_j) = \pm \sum_{i \leq k \leq j} \mathbf{e}_{\delta_\circ^k}$ of type $A_{|A_\circ|}$. Therefore, the \mathbf{c} -vector fan is the type $A_{|A_\circ|}$ Coxeter fan and the A_\circ -zonotope is the classical permutahedron $\text{Perm}(|A_\circ|) := \text{conv} \{ \sum_{i \in [|A_\circ|+1]} \sigma(i) \mathbf{f}_i \mid \sigma \in \mathfrak{S}_{|A_\circ|+1} \}$.

The vertices of $\text{Zono}(D_\circ)$ correspond to separable subsets of $\mathbf{C}(D_\circ)$. Although we could work out all facets of $\text{Zono}(D_\circ)$, we will only need the following specific inequalities.

Proposition 24. *For any D_\circ -accordion diagonal γ_\bullet , the D_\circ -zonotope $\text{Zono}(D_\circ)$ has a facet defined by the inequality*

$$\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{x} \rangle \leq \omega(D_\circ | \gamma_\bullet),$$

where $\omega(D_\circ | \gamma_\bullet)$ is the [D \$\circ\$ -height](#) of γ_\bullet , i.e. the number of D_\circ -accordion diagonals that cross γ_\bullet .

Proof. Let $\omega(D_\circ | \gamma_\bullet)$ denote the maximum of $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{x} \rangle$ over $\text{Zono}(D_\circ)$. As $\text{Zono}(D_\circ)$ is the Minkowski sum of all \mathbf{c} -vectors, we have

$$\omega(D_\circ | \gamma_\bullet) = \sum_{\substack{\mathbf{c} \in \mathbf{C}(D_\circ) \\ \langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle > 0}} \langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle.$$

By Remark 18, we have $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle \in \{-1, 0, 1\}$ for any $\mathbf{c} \in \mathbf{C}(D_\circ)$. We thus just need to count the distinct \mathbf{c} -vectors \mathbf{c} such that $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle > 0$. It turns out that it is more convenient and equivalent (since $\mathbf{C}(D_\circ) = -\mathbf{C}(D_\circ)$) to count the distinct \mathbf{c} -vectors \mathbf{c} such that $\langle \mathbf{g}(D_\circ | \gamma_\bullet) \mid \mathbf{c} \rangle < 0$. For that, let Z_\circ denote the zigzag of the accordion crossed by γ_\bullet in D_\circ , and decompose $Z_\circ = Z_\circ^- \sqcup Z_\circ^+$ such that $\mathbf{g}(D_\circ | \gamma_\bullet) = \mathbb{1}_{Z_\circ^+} - \mathbb{1}_{Z_\circ^-}$ (where $\mathbb{1}_{X_\circ} := \sum_{\delta_\circ \in X_\circ} \mathbf{e}_{\delta_\circ}$ for $X_\circ \subseteq D_\circ$). Let δ_\bullet be a D_\circ -accordion diagonal. Let A_\circ^- (resp. A_\circ^+) denote the accordion crossed by $\delta_\bullet = u_\bullet v_\bullet$ in D_\circ and not incident to $(u+1)_\circ$ or $(v+1)_\circ$ (resp. to $(u-1)_\circ$ or $(v-1)_\circ$). Let $\mathbf{c}^-(\delta_\bullet) := -\mathbb{1}_{A_\circ^-}$ and $\mathbf{c}^+(\delta_\bullet) := \mathbb{1}_{A_\circ^+}$. Recall from Lemma 19 that the negative (resp. positive) \mathbf{c} -vectors of $\mathbf{C}(D_\circ)$ are given by $\mathbf{c}^-(\delta_\bullet)$ (resp. $\mathbf{c}^+(\delta_\bullet)$) for all D_\circ -accordion diagonal δ_\bullet not in D_\bullet^- (resp. D_\bullet^+). We let the reader check that:

- ◊ If γ_\bullet and δ_\bullet do not cross and have no common endpoint, both $|Z_\circ \cap A_\circ^-|$ and $|Z_\circ \cap A_\circ^+|$ are even. Thus $\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}^-(\delta_\bullet) \rangle = \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}^+(\delta_\bullet) \rangle = 0$.
- ◊ If γ_\bullet and δ_\bullet have a common endpoint, and $\gamma_\bullet \delta_\bullet$ form a counterclockwise angle, then $|Z_\circ \cap A_\circ^-|$ is even while $Z_\circ \cap A_\circ^+$ is empty or starts and ends in Z_\circ^+ . Thus $\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}^-(\delta_\bullet) \rangle = 0$ while $\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}^+(\delta_\bullet) \rangle \geq 0$. The situation is similar if $\gamma_\bullet \delta_\bullet$ form a clockwise angle.
- ◊ If γ_\bullet and δ_\bullet cross, $Z_\circ \cap A_\circ^-$ and $Z_\circ \cap A_\circ^+$ are empty or start and end both in Z_\circ^- or both in Z_\circ^+ . Thus, either $\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}^-(\delta_\bullet) \rangle < 0$ and $\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c}^+(\delta_\bullet) \rangle \geq 0$ or conversely.

We conclude from this case analysis that

$$\omega(D_\circ | \gamma_\bullet) = |\{\mathbf{c} \in \mathbf{C}(D_\circ) \mid \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{c} \rangle < 0\}| = |\{D_\circ\text{-accordion diagonals crossing } \gamma_\bullet\}|.$$

Finally, the inequality $\langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{x} \rangle \leq \omega(D_\circ | \gamma_\bullet)$ defines a priori a face $\mathbf{F}(\gamma_\bullet)$ of the zonotope $\text{Zono}(D_\circ)$. This face $\mathbf{F}(\gamma_\bullet)$ is the Minkowski sum of the \mathbf{c} -vectors of $\mathbf{C}(D_\circ)$ orthogonal to $\mathbf{g}(D_\circ | \gamma_\bullet)$. Proposition 20 ensures that any D_\circ -accordion dissection D_\bullet containing γ_\bullet already provides $|D_\bullet| - 1$ linearly independent such \mathbf{c} -vectors $\mathbf{c}(D_\circ | \delta_\bullet \in D_\bullet)$ for $\delta_\bullet \in D_\bullet \setminus \{\gamma_\bullet\}$. We obtain that $\mathbf{F}(\gamma_\bullet)$ has dimension $|D_\bullet| - 1 = |D_\circ| - 1$ and is therefore a facet of the zonotope $\text{Zono}(D_\circ)$. \square

Define the half-space and the hyperplane corresponding to a solid D_\circ -accordion diagonal γ_\bullet by

$$\begin{aligned} \mathbf{H}^\leq(D_\circ | \gamma_\bullet) &:= \{\mathbf{x} \in \mathbb{R}^{D_\circ} \mid \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{x} \rangle \leq \omega(D_\circ | \gamma_\bullet)\}, \\ \text{and } \mathbf{H}^=(D_\circ | \gamma_\bullet) &:= \{\mathbf{x} \in \mathbb{R}^{D_\circ} \mid \langle \mathbf{g}(D_\circ | \gamma_\bullet) | \mathbf{x} \rangle = \omega(D_\circ | \gamma_\bullet)\}. \end{aligned}$$

2.3. g-vector fan and D_\circ -accordiohedron. In this section, we give a geometric realization of the D_\circ -accordion complex. We start by realizing this simplicial complex as a complete simplicial fan in \mathbb{R}^{D_\circ} . We denote by $\mathbb{R}_{\geq 0}\mathbf{R}$ the positive span of a set \mathbf{R} of vectors in \mathbb{R}^{D_\circ} .

Theorem 25. *The collection of cones*

$$\mathcal{F}^{\mathbf{g}}(D_\circ) := \{\mathbb{R}_{\geq 0}\mathbf{g}(D_\circ | D_\bullet) \mid D_\bullet \text{ any } D_\circ\text{-accordion dissection}\}$$

*forms a complete simplicial fan, that we call the **g-vector fan** of D_\circ .*

The proof uses the following characterization of complete simplicial fans [DRS10, Corollary 4.5.20]. We will provide as well an alternative proof in Remark 56 based on sections of Cambrian fans.

Proposition 26. *Consider a pseudomanifold Δ with vertex set X and a set of vectors $\mathbf{R} := (\mathbf{r}_x)_{x \in X}$ of \mathbb{R}^d . For $\triangle \in \Delta$, define $\mathbf{R}_\triangle := \{\mathbf{r}_x \mid x \in \triangle\}$. Then the collection of cones $\{\mathbb{R}_{\geq 0}\mathbf{R}_\triangle \mid \triangle \in \Delta\}$ forms a complete simplicial fan if and only if*

- (1) *there exists a facet \triangle of Δ such that \mathbf{R}_\triangle is a basis of \mathbb{R}^d and such that the open cones $\mathbb{R}_{> 0}\mathbf{R}_\triangle$ and $\mathbb{R}_{> 0}\mathbf{R}_{\triangle'}$ are disjoint for any facet \triangle' of Δ distinct from \triangle ;*
- (2) *for two adjacent facets \triangle, \triangle' of Δ with $\triangle \setminus \{x\} = \triangle' \setminus \{x'\}$, there is a linear dependence*

$$\alpha \mathbf{r}_x + \alpha' \mathbf{r}_{x'} + \sum_{y \in \triangle \cap \triangle'} \beta_y \mathbf{r}_y = 0$$

on $\mathbf{R}_{\triangle \cup \triangle'}$ where the coefficients α and α' have the same sign. (When these conditions hold, these coefficients do not vanish and the linear dependence is unique up to rescaling.)

Proof of Theorem 25. By Corollary 22, the cone $\mathbb{R}_{\geq 0}\mathbf{g}(D_\circ | D_\bullet^-)$ is the only cone of $\mathcal{F}^{\mathbf{g}}(D_\circ)$ intersecting the interior of the positive orthant $(\mathbb{R}_{\geq 0})^{D_\circ}$. Consider now two adjacent maximal D_\circ -accordion dissections D_\bullet, D'_\bullet . Let $\delta_\bullet \in D_\bullet$ and $\delta'_\bullet \in D'_\bullet$ be such that $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$, and let μ_\bullet and ν_\bullet be the other diagonals of Figure 4 as defined in Lemma 8. Note that a diagonal of D_\circ crosses none of (resp. one of, resp. both) the diagonals $\delta_\bullet, \delta'_\bullet$ if and only if it crosses none of (resp. one of, resp. both) the diagonals μ_\bullet, ν_\bullet . The same holds for a Z or a Σ of D_\circ . Therefore, we have the linear dependence $\mathbf{g}(D_\circ | \delta_\bullet) + \mathbf{g}(D_\circ | \delta'_\bullet) = \mathbf{g}(D_\circ | \mu_\bullet) + \mathbf{g}(D_\circ | \nu_\bullet)$. This shows that $\mathcal{F}^{\mathbf{g}}(D_\circ)$ satisfies the two conditions of Proposition 26, and thus concludes the proof. \square

Remark 27. The linear dependence $\mathbf{g}(D_\circ | \delta_\bullet) + \mathbf{g}(D_\circ | \delta'_\bullet) = \mathbf{g}(D_\circ | \mu_\bullet) + \mathbf{g}(D_\circ | \nu_\bullet)$ relating the \mathbf{g} -vectors of two adjacent maximal D_\circ -accordion dissections D_\bullet, D'_\bullet with $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$ shows that $\det(\mathbf{g}(D_\circ | D_\bullet)) = -\det(\mathbf{g}(D_\circ | D'_\bullet))$. Since the initial cone $\mathbb{R}_{\geq 0}\mathbf{g}(D_\circ | D_\bullet^-)$ is generated by the coordinate vectors (see Example 15), we obtain that $\det(\mathbf{g}(D_\circ | D_\bullet)) = \pm 1$ for all D_\circ -accordion dissection D_\bullet , so that the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(D_\circ)$ is always *smooth*.

Remark 28. By Proposition 20, any non-maximal cone of $\mathcal{F}^g(D_\circ)$ is supported by an hyperplane orthogonal to a \mathbf{c} -vector of $\mathbf{C}(D_\circ)$. The \mathbf{g} -vector fan $\mathcal{F}^g(D_\circ)$ thus coarsens the \mathbf{c} -vector fan $\mathcal{F}^c(D_\circ)$.

Remark 29. Following Remark 2, we observe that special reference dissections give rise to the following relevant fans:

- ◊ For an accordion triangulation A_\circ (*i.e.* with no interior triangle), the \mathbf{g} -vector fan $\mathcal{F}^g(A_\circ)$ coincides with a type A Cambrian fan of N. Reading and D. Speyer [RS09].
- ◊ For an arbitrary triangulation T_\circ (with or without interior triangle), the \mathbf{g} -vector fan $\mathcal{F}^g(T_\circ)$ was recently constructed in [HPS17].

Example 30. Figure 5 illustrates the \mathbf{g} -vector fans $\mathcal{F}^g(D_\circ)$ for various reference dissections D_\circ : the fan, the snake, and the cyclic triangulation of the hexagon, and a dissection of the heptagon. More precisely, we have represented the stereographic projection of the fans from the point $[1, 1, 1]$. Therefore, the external face of the projection corresponds to the D_\circ -accordion dissection D_\circ^- . We have labeled all vertices of the projection (*i.e.* the rays of the fan) by the corresponding D_\circ -accordion diagonals.




FIGURE 5. Stereographic projections of the \mathbf{g} -vector fans $\mathcal{F}^g(D_\circ)$ for various reference hollow dissections D_\circ . See Figure 8 for alternative simplicial fan realizations of these accordion complexes.

We now provide a first polytopal realization of the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathbf{D}_\circ)$ (see also Section 4). This fan has a maximal cone for each maximal \mathbf{D}_\circ -accordion dissection and a ray for each \mathbf{D}_\circ -accordion diagonal. For a maximal \mathbf{D}_\circ -accordion dissection \mathbf{D}_\bullet , we define a point $\mathbf{p}(\mathbf{D}_\circ | \mathbf{D}_\bullet) \in \mathbb{R}^{\mathbf{D}_\circ}$ by

$$\mathbf{p}(\mathbf{D}_\circ | \mathbf{D}_\bullet) := \sum_{\delta_\bullet \in \mathbf{D}_\bullet} \omega(\mathbf{D}_\circ | \delta_\bullet) \cdot \mathbf{c}(\mathbf{D}_\circ | \delta_\bullet \in \mathbf{D}_\bullet),$$

where $\omega(\mathbf{D}_\circ | \delta_\bullet)$ still denotes the \mathbf{D}_\circ -height of δ_\bullet defined as the number of \mathbf{D}_\circ -accordion diagonals that cross δ_\bullet . We will need the following two technical lemmas in the proof of Theorem 33.

Lemma 31. *For any maximal \mathbf{D}_\circ -accordion dissection \mathbf{D}_\bullet , the point $\mathbf{p}(\mathbf{D}_\circ | \mathbf{D}_\bullet)$ is the intersection of the hyperplanes $\mathbf{H}^=(\mathbf{D}_\circ | \delta_\bullet)$ for $\delta_\bullet \in \mathbf{D}_\bullet$.*

Proof. Since $\mathbf{g}(\mathbf{D}_\circ | \mathbf{D}_\bullet)$ and $\mathbf{c}(\mathbf{D}_\circ | \mathbf{D}_\bullet)$ form dual bases by Proposition 20, we have for any $\gamma_\bullet \in \mathbf{D}_\bullet$:

$$\begin{aligned} \langle \mathbf{g}(\mathbf{D}_\circ | \gamma_\bullet) | \mathbf{p}(\mathbf{D}_\circ | \mathbf{D}_\bullet) \rangle &= \sum_{\delta_\bullet \in \mathbf{D}_\bullet} \omega(\mathbf{D}_\circ | \delta_\bullet) \cdot \langle \mathbf{g}(\mathbf{D}_\circ | \gamma_\bullet) | \mathbf{c}(\mathbf{D}_\circ | \delta_\bullet \in \mathbf{D}_\bullet) \rangle \\ &= \sum_{\delta_\bullet \in \mathbf{D}_\bullet} \omega(\mathbf{D}_\circ | \delta_\bullet) \cdot \mathbb{1}_{\gamma_\bullet = \delta_\bullet} = \omega(\mathbf{D}_\circ | \gamma_\bullet). \end{aligned} \quad \square$$

Lemma 32. *If $\mathbf{D}_\bullet, \mathbf{D}'_\bullet$ are two adjacent maximal \mathbf{D}_\circ -accordion dissections, and $\delta_\bullet \in \mathbf{D}_\bullet$ and $\delta'_\bullet \in \mathbf{D}'_\bullet$ are such that $\mathbf{D}_\bullet \setminus \{\delta_\bullet\} = \mathbf{D}'_\bullet \setminus \{\delta'_\bullet\}$, then*

$$\mathbf{c}(\mathbf{D}_\circ | \delta_\bullet \in \mathbf{D}_\bullet) = -\mathbf{c}(\mathbf{D}_\circ | \delta'_\bullet \in \mathbf{D}'_\bullet) \quad \text{and} \quad \mathbf{p}(\mathbf{D}_\circ | \mathbf{D}'_\bullet) - \mathbf{p}(\mathbf{D}_\circ | \mathbf{D}_\bullet) \in \mathbb{Z}_{<0} \cdot \mathbf{c}(\mathbf{D}_\circ | \delta_\bullet \in \mathbf{D}_\bullet).$$

Proof. Let $\mathbf{D}_\bullet, \mathbf{D}'_\bullet$ be two adjacent maximal \mathbf{D}_\circ -accordion dissections, let $\delta_\bullet \in \mathbf{D}_\bullet$ and $\delta'_\bullet \in \mathbf{D}'_\bullet$ be such that $\mathbf{D}_\bullet \setminus \{\delta_\bullet\} = \mathbf{D}'_\bullet \setminus \{\delta'_\bullet\}$, and let μ_\bullet and ν_\bullet be the other diagonals of Figure 4 as defined in Lemma 8. A quick case analysis then shows that

$$\mathbf{c}(\mathbf{D}_\circ | \gamma_\bullet \in \mathbf{D}'_\bullet) = \begin{cases} \mathbf{c}(\mathbf{D}_\circ | \gamma_\bullet \in \mathbf{D}_\bullet) & \text{for all diagonal } \gamma_\bullet \in \mathbf{D}_\bullet \setminus \{\delta_\bullet, \mu_\bullet, \nu_\bullet\}, \\ -\mathbf{c}(\mathbf{D}_\circ | \delta_\bullet \in \mathbf{D}_\bullet) & \text{if } \gamma_\bullet = \delta'_\bullet, \\ \mathbf{c}(\mathbf{D}_\circ | \gamma_\bullet \in \mathbf{D}_\bullet) + \mathbf{c}(\mathbf{D}_\circ | \delta_\bullet \in \mathbf{D}_\bullet) & \text{if } \gamma_\bullet \in \{\mu_\bullet, \nu_\bullet\}. \end{cases}$$

Summing the contribution of all \mathbf{c} -vectors with their coefficients $\omega(\mathbf{D}_\circ | \gamma_\bullet)$, we obtain

$$\mathbf{p}(\mathbf{D}_\circ | \mathbf{D}'_\bullet) - \mathbf{p}(\mathbf{D}_\circ | \mathbf{D}_\bullet) = (\omega(\mathbf{D}_\circ | \mu_\bullet) + \omega(\mathbf{D}_\circ | \nu_\bullet) - \omega(\mathbf{D}_\circ | \delta_\bullet) - \omega(\mathbf{D}_\circ | \delta'_\bullet)) \cdot \mathbf{c}(\mathbf{D}_\circ | \delta_\bullet \in \mathbf{D}_\bullet).$$

Finally, note that any diagonal of \mathbf{P}_\bullet that crosses one of (resp. both) the diagonals μ_\bullet, ν_\bullet also crosses one of (resp. both) the diagonals $\delta_\bullet, \delta'_\bullet$. Moreover, δ_\bullet and δ'_\bullet cross each other but do not cross μ_\bullet and ν_\bullet . It follows that $\omega(\mathbf{D}_\circ | \mu_\bullet) + \omega(\mathbf{D}_\circ | \nu_\bullet) - \omega(\mathbf{D}_\circ | \delta_\bullet) - \omega(\mathbf{D}_\circ | \delta'_\bullet) \leq -2 < 0$. \square

Theorem 33. *The two sets given by*

- ◊ the convex hull of the points $\mathbf{p}(\mathbf{D}_\circ | \mathbf{D}_\bullet)$ for all maximal \mathbf{D}_\circ -accordion dissection \mathbf{D}_\bullet ,
- ◊ the intersection of the half-spaces $\mathbf{H}^{\leq}(\mathbf{D}_\circ | \gamma_\bullet)$ for all \mathbf{D}_\circ -accordion diagonals γ_\bullet ,

define the same polytope, that we called \mathbf{D}_\circ -accordiohedron and denote by $\text{Acco}(\mathbf{D}_\circ)$. Its normal fan is the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathbf{D}_\circ)$. Thus, it is a polytopal realization of the \mathbf{D}_\circ -accordion complex $\mathcal{AC}(\mathbf{D}_\circ)$.

The proof of Theorem 33 is based on the following characterization of polytopal realizations of a complete simplicial fan, whose proof can be found e.g. in [HLT11, Theorem 4.1].

Theorem 34. *Given a complete simplicial fan \mathcal{F} in \mathbb{R}^d , consider for each ray \mathbf{r} of \mathcal{F} a half-space $\mathbf{H}_\mathbf{r}^{\leq}$ of \mathbb{R}^d containing the origin and defined by a hyperplane $\mathbf{H}_\mathbf{r}^-$ orthogonal to \mathbf{r} . For each maximal cone C of \mathcal{F} , let $\mathbf{a}(C) \in \mathbb{R}^d$ be the intersection of the hyperplanes $\mathbf{H}_\mathbf{r}^-$ for $\mathbf{r} \in C$. Then the following assertions are equivalent:*

- (i) *The vector $\mathbf{a}(C') - \mathbf{a}(C)$ points from C to C' for any two adjacent maximal cones C, C' of \mathcal{F} .*
- (ii) *The polytopes*

$$\text{conv}\{\mathbf{a}(C) \mid C \text{ maximal cone of } \mathcal{F}\} \quad \text{and} \quad \bigcap_{\mathbf{r} \text{ ray of } \mathcal{F}} \mathbf{H}_\mathbf{r}^{\leq}$$

coincide and their normal fan is \mathcal{F} .

Proof of Theorem 33. The \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_o)$ has a ray $\mathbf{g}(\mathcal{D}_o | \delta_\bullet)$ for each \mathcal{D}_o -accordion diagonal δ_\bullet and a maximal cone $\mathcal{C}(\mathcal{D}_o) = \mathbb{R}_{\geq 0} \mathbf{g}(\mathcal{D}_o | \mathcal{D}_\bullet)$ for each maximal \mathcal{D}_o -accordion dissection \mathcal{D}_\bullet . Consider the half-spaces $\mathbf{H}^{\leq}(\mathcal{D}_o | \gamma_\bullet)$ for all \mathcal{D}_o -accordion diagonals γ_\bullet . Lemma 31 ensures that the point $\mathbf{a}(\mathcal{C}(\mathcal{D}_o))$ coincides with $\mathbf{p}(\mathcal{D}_o | \mathcal{D}_\bullet)$ for each maximal \mathcal{D}_o -accordion dissection \mathcal{D}_\bullet . Finally, Lemma 32 shows that the conditions of application of Theorem 34 are fulfilled. \square

Remark 35. Following Remark 2, observe that special reference hollow dissections give rise to the following relevant polytopes, illustrated in Figure 6:

- ◊ For a fan triangulation \mathcal{T}_o , the \mathcal{T}_o -accordiohedron $\text{Acco}(\mathcal{T}_o)$ is the classical associahedron constructed by S. Shnider and S. Sternberg [SS93] and J.-L. Loday [Lod04].
- ◊ The \mathcal{A}_o -accordiohedra $\text{Acco}(\mathcal{A}_o)$ for all accordion triangulations \mathcal{A}_o are precisely the associahedra constructed by C. Hohlweg and C. Lange in [HL07].
- ◊ For a triangulation \mathcal{T}_o with an interior triangle, the \mathcal{T}_o -accordiohedron $\text{Acco}(\mathcal{T}_o)$ was recently constructed in [HPS17]. For example, for the triangulation of the hexagon with an interior triangle, this associahedron appeared as a mysterious realization in [CSZ15].
- ◊ For a quadrangulation \mathcal{Q}_o , the \mathcal{Q}_o -accordiohedron $\text{Acco}(\mathcal{Q}_o)$ is a realization of the Stokes polytope announced by Y. Baryshnikov [Bar01] and discussed by F. Chapoton in [Cha16].

2.4. Some properties of $\text{Acco}(\mathcal{D}_o)$. We conclude this section by pointing out some relevant combinatorial and geometric properties and observations on the \mathcal{D}_o -accordiohedron.

Proposition 36. *The graph of the \mathcal{D}_o -accordiohedron $\text{Acco}(\mathcal{D}_o)$ linearly oriented in the direction $-\mathbb{1} := -\sum_{\delta_o \in \mathcal{D}_o} \mathbf{e}_{\delta_o}$ is the Hasse diagram of the accordion lattice $\mathcal{AL}(\mathcal{D}_o)$.*

Proof. Consider two adjacent maximal \mathcal{D}_o -accordion dissections $\mathcal{D}_\bullet, \mathcal{D}'_\bullet$ such that the flip from \mathcal{D}_\bullet to \mathcal{D}'_\bullet is increasing. Let $\delta_\bullet \in \mathcal{D}_\bullet$ and $\delta'_\bullet \in \mathcal{D}'_\bullet$ be such that $\mathcal{D}_\bullet \setminus \{\delta_\bullet\} = \mathcal{D}'_\bullet \setminus \{\delta'_\bullet\}$. As observed in Remark 18 (ii), the \mathbf{c} -vector $\mathbf{c}(\mathcal{D}_o | \delta_\bullet \in \mathcal{D}_\bullet)$ is the characteristic vector $\mathbb{1}_{\mathcal{A}_o}$ of the set \mathcal{A}_o of diagonals of \mathcal{D}_o crossed by both δ_\bullet and δ'_\bullet . Applying Lemma 32, we therefore obtain that

$$\langle -\mathbb{1} | \mathbf{p}(\mathcal{D}_o | \mathcal{D}'_\bullet) - \mathbf{p}(\mathcal{D}_o | \mathcal{D}_\bullet) \rangle = \langle -\mathbb{1} | \lambda \cdot \mathbf{c}(\mathcal{D}_o | \delta_\bullet \in \mathcal{D}_\bullet) \rangle = \lambda \cdot \langle -\mathbb{1} | \mathbb{1}_{\mathcal{A}_o} \rangle = -\lambda \cdot |\mathcal{A}_o|,$$

for some $\lambda \in \mathbb{Z}_{<0}$. Thus, the linear functional $-\mathbb{1}$ indeed orients the edge $[\mathbf{p}(\mathcal{D}_o | \mathcal{D}_\bullet), \mathbf{p}(\mathcal{D}_o | \mathcal{D}'_\bullet)]$ from $\mathbf{p}(\mathcal{D}_o | \mathcal{D}_\bullet)$ to $\mathbf{p}(\mathcal{D}_o | \mathcal{D}'_\bullet)$. \square

Remark 37. Since the \mathbf{c} -vector fan $\mathcal{F}^{\mathbf{c}}(\mathcal{D}_o)$ refines the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_o)$, there is a natural projection π from the vertices of the \mathcal{D}_o -zonotope $\text{Zono}(\mathcal{D}_o)$ to that of the \mathcal{D}_o -accordiohedron $\text{Acco}(\mathcal{D}_o)$. In analogy to the acyclic case, one could hope to obtain the accordion lattice as a lattice quotient through this projection. However, the transitive closure of the graph of the \mathcal{D}_o -zonotope $\text{Zono}(\mathcal{D}_o)$ oriented in the direction $-\mathbb{1}$ is not a lattice in general (the first counter-example is the dissection with a central square surrounded by 4 triangles). As shown in [GM16], the right objects are not the separable subsets of \mathbf{c} -vectors (*i.e.* the vertices of $\text{Zono}(\mathcal{D}_o)$) but the biclosed subsets of \mathbf{c} -vectors.

Proposition 38. *The accordiohedron $\text{Acco}(\mathcal{D}_o)$ has precisely $|\mathcal{D}_o|$ pairs of parallel facets.*

Proof. Two facets of $\text{Acco}(\mathcal{D}_o)$ are parallel if and only if the corresponding \mathbf{g} -vectors are opposite. We therefore want to prove that the pairs of opposite coordinate vectors are the only pairs of opposite \mathbf{g} -vectors. Assume by contradiction that there exist two hollow diagonals $\delta_o, \delta'_o \in \mathcal{D}_o$ and two solid \mathcal{D}_o -diagonals $\delta_\bullet, \delta'_\bullet$ such that $\mathbf{g}(\mathcal{D}_o | \delta_\bullet)$ and $\mathbf{g}(\mathcal{D}_o | \delta'_\bullet)$ have non-zero opposite coordinate vectors both on δ_o and δ'_o . Then both δ_\bullet and δ'_\bullet cross both δ_o and δ'_o . But this implies that they both slalom on δ_o (and on δ'_o) in the same way. Contradiction. \square

Recall from Example 15 that the \mathbf{g} -vectors of the diagonals of \mathcal{D}_\bullet^- (resp. \mathcal{D}_\bullet^+) are the coordinate vectors (resp. negative of the coordinate vectors). Consider the *\mathcal{D}_o -parallelepiped*

$$\text{Para}(\mathcal{D}_o) := \{\mathbf{x} \in \mathbb{R}^{\mathcal{D}_o} \mid \langle \mathbf{g}(\mathcal{D}_o | \delta_\bullet) | \mathbf{x} \rangle \leq \omega(\mathcal{D}_o | \delta_\bullet) \text{ for all } \delta_\bullet \in \mathcal{D}_\bullet^- \cup \mathcal{D}_\bullet^+\}$$

defined by the inequalities of the \mathcal{D}_o -zonotope $\text{Zono}(\mathcal{D}_o)$ corresponding to the positive and negative basis vectors. Our next statement follows from Proposition 38 and is illustrated in Figure 6.

Corollary 39. *For any \mathcal{D}_o , we have matriochka polytopes: $\text{Zono}(\mathcal{D}_o) \subseteq \text{Acco}(\mathcal{D}_o) \subseteq \text{Para}(\mathcal{D}_o)$.*

In fact, each polytope in this chain is obtained by deleting facets from the previous one.




FIGURE 6. The zonotope $\text{Zono}(D_o)$, D_o -accordiohedron $\text{Acco}(D_o)$ and parallelepiped $\text{Para}(D_o)$ for different reference dissections D_o . The first column is J.-L. Loday's associahedron [Lod04], the second column is one of C. Hohlweg and C. Lange's associahedra [HL07], the third column appeared in a discussion in C. Ceballos, F. Santos and G. Ziegler's survey on associahedra [CSZ15, Figure 3] and was explained in C. Hohlweg, V. Pilaud and S. Stella's recent paper [HPS17], and the last column is a Stokes complex discussed by F. Chapoton in [Cha16] and illustrated in Figure 3.

Consider now an isometry σ of the plane that preserves the hollow polygon P_o and the solid polygon P_\bullet . For any diagonals and dissections $\delta_\bullet \in D_\bullet$ and $\delta_o \in D_o$, we have

- ◇ δ_\bullet is a D_o -accordion diagonal $\iff \sigma(\delta_\bullet)$ is a $\sigma(D_o)$ -accordion diagonal,
- ◇ D_\bullet is a D_o -accordion dissection $\iff \sigma(D_\bullet)$ is a $\sigma(D_o)$ -accordion dissection,
- ◇ if $\Sigma : \mathbb{R}^{D_o} \rightarrow \mathbb{R}^{\sigma(D_o)}$ denotes the isometry defined by $(\Sigma(\mathbf{x}))_{\sigma(\delta_o)} := \varepsilon(\sigma) \cdot \mathbf{x}_{\delta_o}$, (where $\varepsilon(\sigma) = 1$ if σ is direct and -1 if σ is indirect), then we have

$$\begin{aligned} \mathbf{g}(\sigma(D_o) \mid \sigma(\delta_\bullet)) &= \Sigma(\mathbf{g}(D_o \mid \delta_\bullet)), & \mathbf{c}(\sigma(D_o) \mid \sigma(\delta_\bullet) \in \sigma(D_\bullet)) &= \Sigma(\mathbf{c}(D_o \mid \delta_\bullet \in D_\bullet)), \\ \omega(\sigma(D_o) \mid \sigma(\delta_\bullet)) &= \omega(D_o \mid \delta_\bullet), & \text{and} & \quad \mathbf{p}(\sigma(D_o) \mid \sigma(D_\bullet)) = \Sigma(\mathbf{p}(D_o \mid D_\bullet)). \end{aligned}$$

This immediately implies the following statement.

Proposition 40. *Any P_o -preserving isometry $\sigma : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ induces an isometry $\Sigma : \mathbb{R}^{D_o} \rightarrow \mathbb{R}^{\sigma(D_o)}$ with $\Sigma(\text{Zono}(D_o)) = \text{Zono}(\sigma(D_o))$, $\Sigma(\text{Acco}(D_o)) = \text{Acco}(\sigma(D_o))$ and $\Sigma(\text{Para}(D_o)) = \text{Para}(\sigma(D_o))$.*

We say that a dissection D is *σ -invariant* when $\sigma(D) = D$. Assume now that σ is a rotation and D_\circ is σ -invariant. We call *σ -invariant D_\circ -accordion complex* the simplicial complex $\mathcal{AC}^\sigma(D_\circ)$ whose vertices are the crossing-free σ -orbits of D_\circ -accordion diagonals, and whose faces are sets of such orbits whose union is crossing-free. In other words, the faces of $\mathcal{AC}^\sigma(D_\circ)$ are σ -invariant D_\circ -accordion dissections, seen as sets of σ -orbits of diagonals.

Lemma 41. *The σ -invariant D_\circ -accordion complex $\mathcal{AC}^\sigma(D_\circ)$ is a pseudomanifold.*

Proof. Assume first that σ is the central symmetry. In this case, there are two possible types of orbits: the long D_\circ -accordion diagonals and the centrally symmetric pairs of D_\circ -accordion diagonals. One can check that any facet of $\mathcal{AC}^\sigma(D_\circ)$ has a long diagonal if and only if D_\circ has, and has as many centrally symmetric pairs of diagonals as D_\circ . Finally, any orbit in any facet of $\mathcal{AC}^\sigma(D_\circ)$ can be flipped: long diagonals can already be flipped in $\mathcal{AC}(D_\circ)$, and a centrally symmetric pair of diagonals can be flipped by flipping one after the other its two diagonals in $\mathcal{AC}(D_\circ)$.

Finally, the general statement follows from this special case. Indeed, if σ is not a central symmetry, let C_\circ denote the cell of D_\circ containing the center of P_\circ , let u_\circ be a vertex of C_\circ , let \underline{D}_\circ be the set of diagonals of D_\circ whose endpoints are between u_\circ and $\sigma(u_\circ)$, and let ρ be the central symmetry around the middle of $u_\circ\sigma(u_\circ)$. Then $\mathcal{AC}^\sigma(D_\circ)$ is isomorphic to $\mathcal{AC}^\rho(\underline{D}_\circ \cup \rho(\underline{D}_\circ))$. \square

Let $\Sigma : \mathbb{R}^{D_\circ} \rightarrow \mathbb{R}^{D_\circ}$ denote the isometry defined by $(\Sigma(\mathbf{x}))_{\sigma(\delta_\circ)} := \mathbf{x}_{\delta_\circ}$ and $\text{Fix}(\Sigma)$ denote the linear subspace of fixed points of Σ . According to the previous discussion, a maximal D_\circ -accordion dissection D_\bullet is σ -invariant if and only if $\mathbf{p}(D_\circ | D_\bullet) \in \text{Fix}(\Sigma)$. We obtain the following statement.

Proposition 42. *For a σ -invariant dissection D_\circ , the polytope $\text{Acco}^\sigma(D_\circ)$ defined equivalently as*

- \diamond *the convex hull of $\mathbf{p}(D_\circ | D_\bullet)$ for all σ -invariant maximal D_\circ -accordion dissections D_\bullet ,*
- \diamond *the intersection of the D_\circ -accordiohedron $\text{Acco}(D_\circ)$ with the fixed space $\text{Fix}(\Sigma)$,*

is a polytopal realization of the σ -invariant accordion complex $\mathcal{AC}^\sigma(D_\circ)$.

Proof. Denote by $P = \text{conv} \{ \mathbf{p}(D_\circ | D_\bullet) \mid \sigma\text{-invariant maximal } D_\circ\text{-accordion dissections } D_\bullet \}$ and by $Q = \text{Acco}(D_\circ) \cap \text{Fix}(\Sigma)$. The inclusion $P \subseteq Q$ is clear since D_\bullet is σ -invariant if and only if $\mathbf{p}(D_\circ | D_\bullet) \in \text{Fix}(\Sigma)$. We now prove the reverse inclusion. For that, consider an arbitrary σ -invariant maximal D_\circ -accordion dissection D_\bullet . Its corresponding point $\mathbf{p}(D_\circ | D_\bullet)$ is a common vertex of P and Q . Moreover, any edge e of Q incident to $\mathbf{p}(D_\circ | D_\bullet)$ is the intersection of $\text{Fix}(\Sigma)$ with a face F of $\text{Acco}(D_\circ)$ that corresponds to a σ -invariant D_\circ -dissection. Since $\mathcal{AC}^\sigma(D_\circ)$ is a pseudomanifold, this dissection can be refined into another maximal σ -invariant D_\circ -accordion dissection D'_\bullet . The point $\mathbf{p}(D_\circ | D'_\bullet)$ belongs to F and to $\text{Fix}(\Sigma)$ and thus to e . We conclude that if v is a common vertex of P and Q , then so are all neighbors of v in the graph of Q . Propagating this property, we obtain that all vertices of Q are also vertices of P , so that $P = Q$. Finally, there is a clear injection from the σ -invariant accordion complex $\mathcal{AC}^\sigma(D_\circ)$ to the boundary complex of $P = Q$, thus a bijection (since these complexes are two spheres with the same vertex set). \square

3. THE \mathbf{d} -VECTOR FAN

In this section, we discuss the generalization to the D_\circ -accordion complex of another classical geometric realization of the associahedron coming from the theory of cluster algebras [FZ02, FZ03a, CFZ02, CSZ15]. Namely, we define compatibility vectors in analogy with the denominator vectors of cluster variables, and we characterize the reference dissections D_\circ for which these vectors support a complete simplicial fan realizing the D_\circ -accordion complex.

3.1. \mathbf{d} -vectors. Fix a dissection D_\circ of the hollow n -gon. For a hollow diagonal $\delta_\circ = i_\circ j_\circ$ and a solid diagonal δ_\bullet , we denote by

$$(\delta_\circ | \delta_\bullet) := \begin{cases} -1 & \text{if } \delta_\bullet = (i-1)_\bullet(j-1)_\bullet, \\ 0 & \text{if } \delta_\bullet \text{ and } (i-1)_\bullet(j-1)_\bullet \text{ do not cross,} \\ 1 & \text{if } \delta_\bullet \text{ and } (i-1)_\bullet(j-1)_\bullet \text{ cross.} \end{cases}$$

For any D_\circ -accordion diagonal δ_\bullet , the *\mathbf{d} -vector* of δ_\bullet with respect to D_\circ is the vector

$$\mathbf{d}(D_\circ | \delta_\bullet) = \sum_{\delta_\circ \in D_\circ} (\delta_\circ | \delta_\bullet) \mathbf{e}_{\delta_\circ}.$$

In other words, our \mathbf{d} -vector $\mathbf{d}(D_o | \delta_\bullet)$ records the compatibility of the diagonal δ_\bullet with the dissection D_\bullet^- . For a D_o -accordion dissection D_\bullet , we define $\mathbf{d}(D_o | D_\bullet) := \{\mathbf{d}(D_o | \delta_\bullet) \mid \delta_\bullet \in D_\bullet\}$.

Example 43. Consider the hollow dissection $D_o^{\text{ex}} = \{3_o 7_o, 3_o 13_o, 9_o 13_o\}$ and the rightmost solid dissection $D_\bullet^{\text{ex}} = \{2_\bullet 6_\bullet, 2_\bullet 10_\bullet, 10_\bullet 14_\bullet\}$ of Figure 2. Its \mathbf{d} -vectors are given by

$$\mathbf{d}(D_o^{\text{ex}} | 2_\bullet 6_\bullet) = -\mathbf{e}_{3_o 7_o}, \quad \mathbf{d}(D_o^{\text{ex}} | 2_\bullet 10_\bullet) = \mathbf{e}_{9_o 13_o}, \quad \text{and} \quad \mathbf{d}(D_o^{\text{ex}} | 10_\bullet 14_\bullet) = \mathbf{e}_{3_o 13_o} + \mathbf{e}_{9_o 13_o}.$$

3.2. \mathbf{d} -vector fan. We now consider the set of cones

$$\{\mathbb{R}_{\geq 0} \mathbf{d}(D_o | D_\bullet) \mid D_\bullet \text{ any } D_o\text{-accordion dissection}\}$$

generated by the \mathbf{d} -vectors of the D_o -accordion dissections. We want to characterize the reference hollow dissections D_o for which these cones form a complete simplicial fan realizing the D_o -accordion complex. We start with a negative result.

Remark 44. Assume that the reference hollow dissection D_o contains an *even interior cell* C_o , with an even number of edges which are all internal diagonals of D_o . Denote its vertices by i_o^1, \dots, i_o^{2p} (in clockwise order) and its edges $\delta_o^k := i_o^k i_o^{k+1}$ for $k \in [2p]$ (where $i_o^{2p+1} = i_o^1$ by convention). Denote by D_o^k the set of diagonals of D_o separated from C_o by δ_o^k (including δ_o^k itself), and let $D_\bullet^k := \{(i-1)_\bullet(j-1)_\bullet \mid i_o j_o \in D_o^k\}$. Consider the solid diagonals $\delta_\bullet^k := (i^k+1)_\bullet(i^{k+1}+1)_\bullet$ for $k \in [2p]$. Observe that δ_\bullet^k only crosses diagonals of D_\bullet^{k-1} and D_\bullet^k , and that δ_\bullet^k and δ_\bullet^{k+1} cross precisely the same diagonals of D_\bullet^k . Since the cell is even, it ensures that the \mathbf{d} -vectors of the diagonals δ_\bullet^k for $k \in [2p]$ satisfy the linear dependence

$$\sum_{\substack{k \in [2p] \\ k \text{ even}}} \mathbf{d}(D_o | \delta_\bullet^k) = \sum_{\substack{k \in [2p] \\ k \text{ odd}}} \mathbf{d}(D_o | \delta_\bullet^k).$$

However, as already mentioned in Section 1.3, the diagonals δ_\bullet^k for $k \in [2p]$ all belong to the D_o -accordion dissection $D_\bullet^+ := \{(i+1)_\bullet(j+1)_\bullet \mid i_o j_o \in D_o\}$. Therefore, the cone $\mathbb{R}_{\geq 0} \mathbf{d}(D_o | D_\bullet^+)$ is degenerate, so that the \mathbf{d} -vectors cannot realize the D_o -accordion complex.

Example 45. Consider a hollow octagon and the reference dissection $D_o := \{1_o 5_o, 5_o 9_o, 9_o 13_o, 13_o 1_o\}$ with an interior square cell $1_o 5_o 9_o 13_o$. Then we have

$$\begin{aligned} \mathbf{d}(D_o | 2_\bullet 6_\bullet) &= \mathbf{e}_{1_o 5_o} + \mathbf{e}_{5_o 9_o} & \mathbf{d}(D_o | 6_\bullet 10_\bullet) &= \mathbf{e}_{5_o 9_o} + \mathbf{e}_{9_o 13_o} \\ \mathbf{d}(D_o | 10_\bullet 14_\bullet) &= \mathbf{e}_{9_o 13_o} + \mathbf{e}_{13_o 1_o} & \mathbf{d}(D_o | 14_\bullet 2_\bullet) &= \mathbf{e}_{13_o 1_o} + \mathbf{e}_{1_o 5_o} \end{aligned}$$

so that there is already a linear dependence

$$\mathbf{d}(D_o | 2_\bullet 6_\bullet) + \mathbf{d}(D_o | 10_\bullet 14_\bullet) = \mathbf{d}(D_o | 6_\bullet 10_\bullet) + \mathbf{d}(D_o | 14_\bullet 2_\bullet)$$

among the \mathbf{d} -vectors of the D_o -accordion dissection $D_\bullet^+ = \{2_\bullet 6_\bullet, 6_\bullet 10_\bullet, 10_\bullet 14_\bullet, 14_\bullet 2_\bullet\}$.

On the negative side, we have seen that even interior cells are redhibitory for the \mathbf{d} -vector fan. The positive side is that even interior cells are the only obstructions to this construction.

Theorem 46. *The collection of cones*

$$\mathcal{F}^{\mathbf{d}}(D_o) := \{\mathbb{R}_{\geq 0} \mathbf{d}(D_o | D_\bullet) \mid D_\bullet \text{ any } D_o\text{-accordion dissection}\}$$

forms a complete simplicial fan, that we call the \mathbf{d} -vector fan of D_o , if and only if D_o contains no even interior cell.

Proof. We use the characterization of complete simplicial fans presented in Proposition 26.

Observe first that $\mathbf{d}(D_o | D_\bullet^-) = (\mathbb{R}_{\leq 0})^{D_o}$ is the only cone of $\mathcal{F}^{\mathbf{d}}(D_o)$ intersecting the interior of the negative orthant $(\mathbb{R}_{\leq 0})^{D_o}$. Therefore, $\mathcal{F}^{\mathbf{d}}(D_o)$ fulfills Condition (1) in Proposition 26.

To check Condition (2), consider two adjacent maximal D_o -accordion dissections D_\bullet and D'_\bullet and let $\delta_\bullet \in D_\bullet$ and $\delta'_\bullet \in D'_\bullet$ be such that $D_\bullet \setminus \{\delta_\bullet\} = D'_\bullet \setminus \{\delta'_\bullet\}$. Let μ_\bullet and ν_\bullet be the diagonals of $\bar{D}_\bullet \cap \bar{D}'_\bullet$ as defined in Lemma 8. In other words, μ_\bullet and ν_\bullet are incident to both δ_\bullet and δ'_\bullet , and they are crossed by the hollow diagonal which intersect δ_\bullet and δ'_\bullet . Let $\gamma_o = i_o j_o$ be such a hollow diagonals crossing $\delta_\bullet, \delta'_\bullet, \mu_\bullet$ and ν_\bullet , and let $\gamma_\bullet = (i-1)_\bullet(j-1)_\bullet$. We now distinguish three cases:




FIGURE 7. Illustration of the notations and of the different cases in the proof of Theorem 46.

- ◇ Assume that γ_\bullet still crosses μ_\bullet and ν_\bullet . In this case, any diagonal of D_\bullet^- crossing both (resp. either) δ_\bullet and (resp. or) δ'_\bullet also crosses both (resp. either) μ_\bullet and (resp. or) ν_\bullet . See Figure 7 (left). Therefore, the \mathbf{d} -vectors of $D_\bullet \cup D'_\bullet$ satisfy the linear dependence

$$\mathbf{d}(D_\bullet | \delta_\bullet) + \mathbf{d}(D_\bullet | \delta'_\bullet) = \mathbf{d}(D_\bullet | \mu_\bullet) + \mathbf{d}(D_\bullet | \nu_\bullet).$$

- ◇ Assume that γ_\bullet crosses neither μ_\bullet nor ν_\bullet . Then γ_\bullet is incident to both μ_\bullet and ν_\bullet , and therefore is either δ_\bullet or δ'_\bullet , say $\gamma_\bullet = \delta_\bullet$. Then $\mathbf{d}(\gamma_\bullet | \delta_\bullet) = -1$ while $\mathbf{d}(\gamma_\bullet | \delta'_\bullet) = 1$ (since δ'_\bullet crosses $\delta_\bullet = \gamma_\bullet$), so that $\mathbf{d}(\gamma_\bullet | \delta_\bullet) + \mathbf{d}(\gamma_\bullet | \delta'_\bullet) = 0$. Moreover, we have $\mathbf{d}(\gamma_\bullet | \delta'_\bullet) = 0$ for any diagonal $\varepsilon_\bullet \in D_\bullet \cap D'_\bullet$ since $\delta_\bullet = \gamma_\bullet$ cannot cross ε_\bullet as they both belong to D_\bullet . Therefore, the set $\{\mathbf{d}(D_\bullet | \delta_\bullet) + \mathbf{d}(D_\bullet | \delta'_\bullet)\} \cup \mathbf{d}(D_\bullet | D_\bullet \cap D'_\bullet)$ contains $|D_\bullet|$ vectors of \mathbb{R}^{D_\bullet} whose γ_\bullet -coordinate all vanish, so that it admits a linear dependence.
- ◇ Otherwise, we can assume that γ_\bullet crosses μ_\bullet but not ν_\bullet . Then γ_\bullet has a common endpoint with ν_\bullet and δ_\bullet (or δ'_\bullet , but we then permute notations). Changing our initial choice of γ_\bullet , we can assume that no diagonal of D_\bullet^- separates γ_\bullet from δ_\bullet . We now denote clockwise
 - by $\nu_\bullet =: \lambda_\bullet^0, \lambda_\bullet^1, \dots, \lambda_\bullet^\ell := \delta_\bullet$ the edges of the cell C_\bullet of D_\bullet containing ν_\bullet and δ_\bullet ,
 - by $\gamma_\bullet =: \gamma_\bullet^0, \gamma_\bullet^1, \dots, \gamma_\bullet^k$ the edges of the cell C_\bullet^- of D_\bullet^- containing γ_\bullet and crossed by δ_\bullet .
 These notations are illustrated on Figure 7. We still distinguish two subcases as in Figure 7:
 - If γ_\bullet^i crosses λ_\bullet^i for all i as in Figure 7 (middle), then $\ell = k$ and we have the linear dependence

$$2\mathbf{d}(D_\bullet | \delta_\bullet) + \mathbf{d}(D_\bullet | \delta'_\bullet) = \mathbf{d}(D_\bullet | \mu_\bullet) + \sum_{i \in [\ell-1]} (-1)^{(i-1)} \mathbf{d}(D_\bullet | \lambda_\bullet^i).$$

It is essential here that $\ell = k$ is even. This is guaranteed by the assumption that D_\bullet (and thus D_\bullet^-) has no even interior cell, since C_\bullet^- is an interior cell of D_\bullet^- of size k .

- Otherwise, we are in a situation similar to Figure 7 (right). Considering the maximal index m such that γ_\bullet^i crosses λ_\bullet^i for all $i \leq m$, and we have the linear dependence

$$\mathbf{d}(D_\bullet | \delta_\bullet) + \mathbf{d}(D_\bullet | \delta'_\bullet) = \mathbf{d}(D_\bullet | \mu_\bullet) + \sum_{i \in [m]} (-1)^{(i-1)} \mathbf{d}(D_\bullet | \lambda_\bullet^i). \quad \square$$

Remark 47. Following Remark 2, we observe that special reference dissections give rise to the following relevant fans:

- ◇ For a snake triangulation Σ_\bullet , the \mathbf{d} -vector fan $\mathcal{F}^{\mathbf{d}}(\Sigma_\bullet)$ coincides with the type A cluster fan of S. Fomin and A. Zelevinsky [FZ03a].
- ◇ For any triangulation T_\bullet , the \mathbf{d} -vector fan $\mathcal{F}^{\mathbf{d}}(T_\bullet)$ was already constructed in [CSZ15].
- ◇ For a quadrangulation Q_\bullet with no interior quadrangle (equivalently, with no cross), we obtain an alternative realization of the Stokes complexes studied in [Bar01, Cha16]. This was observed by A.-H. Bateni, T. Manneville and V. Pilaud in [BMP16].

Figure 8 illustrates the \mathbf{d} -vector fans $\mathcal{F}^{\mathbf{d}}(D_\bullet)$ for the same reference dissections D_\bullet as in Figure 5. More precisely, we have represented the stereographic projection of the fans from the




FIGURE 8. Stereographic projections of the \mathbf{d} -vector fans $\mathcal{F}^{\mathbf{d}}(D_{\circ})$ for various reference hollow dissections D_{\circ} . See Figure 5 for alternative simplicial fan realizations of these accordion complexes.

point $[-1, -1, -1]$. Therefore, the external face of the projection corresponds to the D_{\circ} -accordion dissection D_{\bullet}^{-} . We have labeled all vertices of the projection (*i.e.* the rays of the fan) by the corresponding D_{\circ} -accordion diagonals. Compare with Figure 5.

Remark 48. To prove that the \mathbf{d} -vector fan $\mathcal{F}^{\mathbf{d}}(D_{\circ})$ is polytopal, we would need to find suitable hyperplanes orthogonal to their rays in order to apply Theorem 34. For the \mathbf{g} -vector fan, these hyperplanes were defined using the height function $\omega(D_{\circ} \mid \delta_{\bullet})$. It would be natural to use the same height function for the \mathbf{d} -vector fan as well. Unfortunately, for this choice of height function, we can only prove Condition (i) of Theorem 34 when D_{\circ} is a triangulation (see also [CSZ15]). We were not able to find suitable right hand sides for any dissection D_{\circ} .

Remark 49. Our \mathbf{d} -vectors record the compatibility with the dissection D_{\bullet}^{-} . A priori, we could compute compatibility vectors with respect to any other maximal D_{\circ} -accordion dissection D_{\bullet}^{ini} . Experiments suggest that the \mathbf{d} -vector construction provides a complete simplicial fan as soon as either D_{\circ} or D_{\bullet}^{ini} contain no even interior cell. We checked it for reference quadrangulations with at most 5 diagonals. The linear dependences involved seem however much more complicated than those of the proof of Theorem 46 (in particular, they may involve \mathbf{d} -vectors of diagonals not included in the cells containing δ_{\bullet} and δ'_{\bullet}).

4. SECTIONS AND PROJECTIONS

Recall that for a fan \mathcal{F} of \mathbb{R}^d and a linear subspace V of \mathbb{R}^d , the [section](#) of \mathcal{F} by V is the fan $\mathcal{F}|_V := \{C \cap V \mid C \in \mathcal{F}\}$. For a polytope $P \subseteq \mathbb{R}^d$ and a projection $\pi : \mathbb{R}^d \rightarrow V$, the normal fan of the projected polytope $\pi(P)$ is the section of the normal fan of P by V [Zie95, Lemma 7.11]. We now consider sections of the \mathbf{g} - and \mathbf{d} -vector fans by coordinate subspaces. For two dissections $D_\circ \subset D'_\circ$, we naturally identify \mathbb{R}^{D_\circ} with the subspace spanned by $\{\mathbf{e}_{\delta_\circ} \mid \delta_\circ \in D_\circ\}$ in $\mathbb{R}^{D'_\circ}$.

4.1. Coordinate sections of the \mathbf{d} -vector fan. We start by sections of the \mathbf{d} -vector fan which are not very surprising. The following lemma is immediate from the definition of \mathbf{d} -vectors.

Lemma 50. *Consider two dissections $D_\circ \subset D'_\circ$, and a D'_\circ -accordion diagonal δ_\bullet . Then we have $\mathbf{d}(D_\circ \mid \delta_\bullet) \in \mathbb{R}^{D_\circ}$ if and only if δ_\bullet does not cross any diagonal of $\{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D'_\circ \setminus D_\circ\}$.*

Corollary 51. *For two dissections $D_\circ \subset D'_\circ$, the section of the \mathbf{d} -vector fan $\mathcal{F}^{\mathbf{d}}(D'_\circ)$ by \mathbb{R}^{D_\circ} has the combinatorics of the link of the dissection $\{(i-1)_\bullet(j-1)_\bullet \mid i_\circ j_\circ \in D'_\circ \setminus D_\circ\}$ in the D'_\circ -accordion complex $\mathcal{AC}(D'_\circ)$, thus of a join of smaller accordion complexes (see Remark 5).*

4.2. Coordinate sections of the \mathbf{g} -vector fan. More relevant are the sections of the \mathbf{g} -vector fan. They provide an alternative approach to polytopal realizations of the accordion complex based on projected associahedra. This approach relies on the following crucial observation.

Lemma 52. *Consider two dissections $D_\circ \subset D'_\circ$, and a D'_\circ -accordion diagonal δ_\bullet . Then we have $\mathbf{g}(D'_\circ \mid \delta_\bullet) \in \mathbb{R}^{D_\circ}$ if and only if δ_\bullet is a D_\circ -accordion diagonal. Moreover, in this case, the \mathbf{g} -vectors $\mathbf{g}(D_\circ \mid \delta_\bullet)$ and $\mathbf{g}(D'_\circ \mid \delta_\bullet)$ coincide.*

Proof. Let $\delta_\circ \in D'_\circ \setminus D_\circ$. By definition, a D'_\circ -accordion diagonal δ_\bullet slaloms on δ_\circ if and only if $\mathbf{g}(D'_\circ \mid \delta_\bullet)_{\delta_\circ} = \varepsilon_\circ(\delta_\circ \in D_\circ \mid \delta_\bullet) \neq 0$. Thus, δ_\bullet is a D_\circ -accordion diagonal if and only if it slaloms on none of the diagonals of $D'_\circ \setminus D_\circ$, i.e. if and only if $\mathbf{g}(D'_\circ \mid \delta_\bullet)_{\delta_\circ} = 0$ for all $\delta_\circ \in D'_\circ \setminus D_\circ$. \square

Based on this lemma, we obtain in the following statement an alternative realization on the \mathbf{g} -vector fan, which is illustrated on Figure 9.

Theorem 53. *Consider two dissections $D_\circ \subset D'_\circ$. Then the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(D_\circ)$ is given by $\mathcal{F}^{\mathbf{g}}(D_\circ) = \{C \in \mathcal{F}^{\mathbf{g}}(D'_\circ) \mid C \subset \mathbb{R}^{D_\circ}\}$ and coincides with the section of the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(D'_\circ)$ by \mathbb{R}^{D_\circ} . Thus $\mathcal{F}^{\mathbf{g}}(D_\circ)$ is realized by the orthogonal projection of the D'_\circ -accordiohedron $\text{Acco}(D'_\circ)$ on \mathbb{R}^{D_\circ} , which is equivalently described by:*

- ◇ the convex hull of the points $\sum_{\delta_\bullet \in D_\bullet} \omega(D'_\circ \mid \delta_\bullet) \cdot \mathbf{c}(D_\circ \mid \delta_\bullet \in D_\bullet)$ for all D_\circ -accordion dissections D_\bullet ,
- ◇ the intersection of the half-spaces $\{\mathbf{x} \in \mathbb{R}^{D_\circ} \mid \langle \mathbf{g}(D_\circ \mid \gamma_\bullet) \mid \mathbf{x} \rangle \leq \omega(D'_\circ \mid \delta_\circ)\}$ for all D_\circ -accordion diagonals γ_\bullet .




FIGURE 9. Projecting accordiohedra on coordinate planes yields smaller accordiohedra.

Proof. Lemma 52 immediately implies that $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_\circ) = \{C \in \mathcal{F}^{\mathbf{g}}(\mathcal{D}'_\circ) \mid C \subset \mathbb{R}^{\mathcal{D}_\circ}\}$. A priori, it is a subfan of the section $\mathcal{F}^{\mathbf{g}}(\mathcal{D}'_\circ)|_{\mathbb{R}^{\mathcal{D}_\circ}} = \{C \cap \mathbb{R}^{\mathcal{D}_\circ} \mid C \in \mathcal{F}^{\mathbf{g}}(\mathcal{D}'_\circ)\}$. However, since $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_\circ)$ is already a complete simplicial fan of $\mathbb{R}^{\mathcal{D}_\circ}$, it coincides with $\mathcal{F}^{\mathbf{g}}(\mathcal{D}'_\circ)|_{\mathbb{R}^{\mathcal{D}_\circ}}$. Since $\mathcal{F}^{\mathbf{g}}(\mathcal{D}'_\circ)$ is the normal fan of $\text{Acco}(\mathcal{D}'_\circ)$, this shows that $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_\circ) = \mathcal{F}^{\mathbf{g}}(\mathcal{D}'_\circ)|_{\mathbb{R}^{\mathcal{D}_\circ}}$ is the normal fan of the orthogonal projection of $\text{Acco}(\mathcal{D}'_\circ)$ on $\mathbb{R}^{\mathcal{D}_\circ}$ [Zie95, Lemma 7.11].

To conclude, we prove the given vertex and facet descriptions of this projection. First, since $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_\circ) = \mathcal{F}^{\mathbf{g}}(\mathcal{D}'_\circ)|_{\mathbb{R}^{\mathcal{D}_\circ}}$, the inequalities of the projection of $\text{Acco}(\mathcal{D}'_\circ)$ on $\mathbb{R}^{\mathcal{D}_\circ}$ are just the inequalities of $\text{Acco}(\mathcal{D}'_\circ)$ whose normal vectors are in $\mathbb{R}^{\mathcal{D}_\circ}$. Finally, the vertex description follow from the inequality description using the same argument as in Lemma 31. \square

Remark 54. The projection of the accordiohedron $\text{Acco}(\mathcal{D}'_\circ)$ on $\mathbb{R}^{\mathcal{D}_\circ}$ differs from the accordiohedron $\text{Acco}(\mathcal{D}_\circ)$: they have both $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_\circ)$ as normal fan, but their precise geometry is different.

Corollary 55. *For any hollow dissection \mathcal{D}_\circ , the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(\mathcal{D}_\circ)$ is realized by a projection of an associahedron of [HPS17].*

Proof. Apply Theorem 53 to any triangulation \mathcal{T}_\circ that refines \mathcal{D}_\circ . \square

Remark 56. Approaching accordion complexes as coordinate sections of \mathbf{g} -vector fans actually provides more concise (but also less instructive) proofs for Sections 1.2 and 2.3. Namely, consider any dissection \mathcal{D}_\circ and let \mathcal{T}_\circ be a triangulation that refines \mathcal{D}_\circ . The sign coherence property for triangulations (see Corollary 22) shows that the section $\mathcal{F}^{\mathbf{g}}(\mathcal{T}_\circ)|_{\mathbb{R}^{\mathcal{D}_\circ}} = \{C \cap \mathbb{R}^{\mathcal{D}_\circ} \mid C \in \mathcal{F}^{\mathbf{g}}(\mathcal{T}_\circ)\}$ actually coincides with $\{C \in \mathcal{F}^{\mathbf{g}}(\mathcal{T}_\circ) \mid C \subset \mathbb{R}^{\mathcal{D}_\circ}\}$. Therefore, this gives an alternative concise proof that the collection of cones $\{C \in \mathcal{F}^{\mathbf{g}}(\mathcal{T}_\circ) \mid C \subset \mathbb{R}^{\mathcal{D}_\circ}\}$ forms a complete simplicial fan. Moreover, this fan has the same combinatorics as the \mathcal{D}_\circ -accordion complex $\mathcal{AC}(\mathcal{D}_\circ)$ by Lemma 52. We conclude directly that $\mathcal{AC}(\mathcal{D}_\circ)$ is a pseudomanifold realized by the fan $\{C \in \mathcal{F}^{\mathbf{g}}(\mathcal{T}_\circ) \mid C \subset \mathbb{R}^{\mathcal{D}_\circ}\}$ and by the orthogonal projection of the associahedron $\text{Asso}(\mathcal{T}_\circ)$ on $\mathbb{R}^{\mathcal{D}_\circ}$.

4.3. Cluster algebra analogues. The perspective on accordion complexes developed in this section also opens the door to generalizations on arbitrary cluster algebras (finite type or not). Namely, consider an arbitrary cluster $X_\circ = (x_\circ^1, \dots, x_\circ^m)$ in an arbitrary cluster algebra \mathcal{A} . For any cluster variable $y \in \mathcal{A}$, we denote by $\mathbf{g}(X_\circ | y) \in \mathbb{R}^m$ and $\mathbf{d}(X_\circ | y) \in \mathbb{R}^m$ the \mathbf{g} - and \mathbf{d} -vectors of y computed with respect to X_\circ , see [FZ02, FZ07]. Fix a non-empty proper subset I of $[m]$. We consider two natural subcomplexes of the cluster complex of \mathcal{A} :

- ◊ the subcomplex $\Delta^{\mathbf{d}}(X_\circ, I)$ induced by the variables y such that $\mathbf{d}(X_\circ | y)_i = 0$ for all $i \in I$,
- ◊ the subcomplex $\Delta^{\mathbf{g}}(X_\circ, I)$ induced by the variables y such that $\mathbf{g}(X_\circ | y)_i = 0$ for all $i \in I$.

It is well-known that the subcomplex $\Delta^{\mathbf{d}}(X_\circ, I)$ is the cluster complex obtained by freezing all variables x_i for $i \in I$. For example in type A , it is a join of simplicial associahedra and it can therefore be realized by a product of smaller associahedra. In contrast, we are not aware that the subcomplex $\Delta^{\mathbf{g}}(X_\circ, I)$ be investigated. The present paper dealt with the type A situation.

Example 57. Let \mathcal{T}_\circ be a triangulation, with internal diagonals labeled by $1, \dots, m$. Consider the corresponding type A_m cluster X_\circ . Then for any non-empty proper subset I of $[m]$, the subcomplex $\Delta^{\mathbf{g}}(X_\circ, I)$ is isomorphic to the \mathcal{D}_\circ -accordion complex, where \mathcal{D}_\circ is the dissection obtained by deleting in \mathcal{T}_\circ the diagonals labeled by I .

Example 58. Example 57 extends to cluster algebras on surfaces [FST08, FT12], using accordions of dissections of surfaces.

The following statement extends Theorem 53 to arbitrary cluster algebras.

Theorem 59. *The subset $\{C \in \mathcal{F}^{\mathbf{g}}(X_\circ) \mid C \subseteq \mathbb{R}^{[m] \setminus I}\}$ of the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(X_\circ)$ of X_\circ coincides with the section $\mathcal{F}^{\mathbf{g}}(X_\circ)|_{\mathbb{R}^{[m] \setminus I}} = \{C \cap \mathbb{R}^{[m] \setminus I} \mid C \in \mathcal{F}^{\mathbf{g}}(X_\circ)\}$.*

Proof. The inclusion $\{C \in \mathcal{F}^{\mathbf{g}}(X_\circ) \mid C \subseteq \mathbb{R}^{[m] \setminus I}\} \subseteq \mathcal{F}^{\mathbf{g}}(X_\circ)|_{\mathbb{R}^{[m] \setminus I}}$ is clear. For the reverse inclusion, we use the sign coherence property of \mathbf{g} -vectors in cluster algebras, which was conjectured

in [FZ07, Conjecture 6.13] and proved in [GHKK14, Theorem 5.1] in general. This property implies that the coordinate plane $\mathbb{R}^{[m] \setminus I}$ intersects any cone C of $\mathcal{F}^{\mathbf{g}}(X_o)$ in a face C' . This shows that $C \cap \mathbb{R}^{[m] \setminus I} = C'$ belongs to $\{C \in \mathcal{F}^{\mathbf{g}}(X_o) \mid C \subseteq \mathbb{R}^{[m] \setminus I}\}$. \square

Corollary 60. *The subcomplex $\Delta^{\mathbf{g}}(X_o, I)$ induced by the variables y such that $\mathbf{g}(X_o|y)_i = 0$ for all $i \in I$ is a pseudomanifold.*

Moreover, extending the result of C. Hohlweg, C. Lange and H. Thomas [HLT11] in the acyclic case, C. Hohlweg, V. Pilaud and S. Stella recently constructed a polytope $\text{Asso}(X_o)$ realizing the \mathbf{g} -vector fan $\mathcal{F}^{\mathbf{g}}(X_o)$ in [HPS17]. We can use this associahedron to realize the subcomplex $\Delta^{\mathbf{g}}(X_o, I)$ as a convex polytope.

Corollary 61. *The orthogonal projection of $\text{Asso}(X_o)$ on $\mathbb{R}^{[m] \setminus I}$ is a realization of $\Delta^{\mathbf{g}}(X_o, I)$.*

Finally, when oriented in the suitable direction v (the sum of the positive roots, or equivalently the sum of the fundamental weights), the graph of the generalized associahedron $\text{Asso}(X_o)$ is the Hasse diagram of a Cambrian lattice [Rea06]. One can similarly orient the graph of the projection of $\text{Asso}(X_o)$ on $\mathbb{R}^{[m] \setminus I}$ in the direction of the projection of v on $\mathbb{R}^{[m] \setminus I}$. Is the resulting graph the Hasse diagram of a lattice? Combining the results of [GM16] with that of the present paper shows that this property holds in type A . We also computationally verified the statement in types B_4 , B_5 , D_4 and D_5 . Following [GM16] it seems promising to construct first a lattice structure on biclosed sets of \mathbf{c} -vectors, and to obtain then the graph of the projection of $\text{Asso}(X_o)$ on $\mathbb{R}^{[m] \setminus I}$ as the Hasse diagram of a lattice quotient.

ACKNOWLEDGEMENTS

We thank C. Hohlweg and S. Stella for many helpful discussions on realizations of the associahedron [HPS17] which were the starting point of this paper. We are grateful to F. Chapoton for various conversations on quadrangulations and Stokes posets, and to A. Garver and T. McConville for introducing us with the accordion complexes during FPSAC'16. Their works [Cha16, GM16] were highly inspiring and motivating. We also thank N. Thiery for a question which led to the approach of Section 4.2, and to P.-G. Plamondon for discussions on the generalization to cluster algebras presented in Section 4.3.

REFERENCES

- [Bar01] Yuliy Baryshnikov. On Stokes sets. In *New developments in singularity theory (Cambridge, 2000)*, volume 21 of *NATO Sci. Ser. II Math. Phys. Chem.*, pages 65–86. Kluwer Acad. Publ., Dordrecht, 2001.
- [BDP14] Thomas Brüstle, Grégoire Dupont, and Matthieu Pérotin. On maximal green sequences. *Int. Math. Res. Not. IMRN*, (16):4547–4586, 2014.
- [BFS90] Louis J. Billera, Paul Filliman, and Bernd Sturmfels. Constructions and complexity of secondary polytopes. *Adv. Math.*, 83(2):155–179, 1990.
- [BMP16] Amir-Hossein Bateni, Thibault Manneville, and Vincent Pilaud. A note on quadrangulations and Stokes complexes. In preparation, 2016.
- [CD06] Michael P. Carr and Satyan L. Devadoss. Coxeter complexes and graph-associahedra. *Topology Appl.*, 153(12):2155–2168, 2006.
- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. *Canad. Math. Bull.*, 45(4):537–566, 2002.
- [Cha16] Frédéric Chapoton. Stokes posets and serpent nests. *Discrete Math. Theor. Comput. Sci.*, 18(3), 2016.
- [CSZ15] Cesar Ceballos, Francisco Santos, and Günter M. Ziegler. Many non-equivalent realizations of the associahedron. *Combinatorica*, 35(5):513–551, 2015.
- [DRS10] Jesus A. De Loera, Jörg Rambau, and Francisco Santos. *Triangulations: Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics*. Springer Verlag, 2010.
- [FS05] Eva Maria Feichtner and Bernd Sturmfels. Matroid polytopes, nested sets and Bergman fans. *Port. Math. (N.S.)*, 62(4):437–468, 2005.
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces I. Cluster complexes. *Acta Math.*, 201(1):83–146, 2008.
- [FT12] Sergey Fomin and Dylan Thurston. Cluster algebras and triangulated surfaces. part II: Lambda lengths. Preprint, [arXiv:1210.5569](https://arxiv.org/abs/1210.5569), 2012.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. I. Foundations. *J. Amer. Math. Soc.*, 15(2):497–529, 2002.

- [FZ03a] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. II. Finite type classification. *Invent. Math.*, 154(1):63–121, 2003.
- [FZ03b] Sergey Fomin and Andrei Zelevinsky. Y -systems and generalized associahedra. *Ann. of Math. (2)*, 158(3):977–1018, 2003.
- [FZ07] Sergey Fomin and Andrei Zelevinsky. Cluster algebras. IV. Coefficients. *Compos. Math.*, 143(1):112–164, 2007.
- [GHKK14] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich. Canonical bases for cluster algebras. Preprint, [arXiv:1411.1394](https://arxiv.org/abs/1411.1394), 2014.
- [GKZ08] Israel Gelfand, Mikhail Kapranov, and Andrei Zelevinsky. *Discriminants, resultants and multidimensional determinants*. Modern Birkhäuser Classics. Birkhäuser Boston Inc., Boston, MA, 2008. Reprint of the 1994 edition.
- [GM16] Alexander Garver and Thomas McConville. Oriented flip graphs and noncrossing tree partitions. Preprint, [arXiv:1604.06009](https://arxiv.org/abs/1604.06009), 2016.
- [Hai84] Mark Haiman. Constructing the associahedron. Unpublished manuscript, 11 pages, available at <http://www.math.berkeley.edu/~mhaiman/ftp/assoc/manuscript.pdf>, 1984.
- [HL07] Christophe Hohlweg and Carsten Lange. Realizations of the associahedron and cyclohedron. *Discrete Comput. Geom.*, 37(4):517–543, 2007.
- [HLT11] Christophe Hohlweg, Carsten Lange, and Hugh Thomas. Permutohedra and generalized associahedra. *Adv. Math.*, 226(1):608–640, 2011.
- [Hoh] Christophe Hohlweg. Permutohedra and associahedra. Pages 129–159 in [MHPS12].
- [HPS17] Christophe Hohlweg, Vincent Pilaud, and Salvatore Stella. Associahedra from cyclic seeds. Preprint, [arXiv:1703.09551](https://arxiv.org/abs/1703.09551), 2017.
- [Lee89] Carl W. Lee. The associahedron and triangulations of the n -gon. *European J. Combin.*, 10(6):551–560, 1989.
- [Lod04] Jean-Louis Loday. Realization of the Stasheff polytope. *Arch. Math. (Basel)*, 83(3):267–278, 2004.
- [MHPS12] Folkert Müller-Hoissen, Jean Marcel Pallo, and Jim Stasheff, editors. *Associahedra, Tamari Lattices and Related Structures. Tamari Memorial Festschrift*, volume 299 of *Progress in Mathematics*. Springer, New York, 2012.
- [MP16] Thibault Manneville and Vincent Pilaud. Compatibility fans for graphical nested complexes. Preprint [arXiv:1501.07152](https://arxiv.org/abs/1501.07152), to appear in *J. Combin. Theory Ser. A*, 2016.
- [Pil13] Vincent Pilaud. Signed tree associahedra. Preprint, [arXiv:1309.5222](https://arxiv.org/abs/1309.5222), 2013.
- [Pos09] Alexander Postnikov. Permutohedra, associahedra, and beyond. *Int. Math. Res. Not. IMRN*, (6):1026–1106, 2009.
- [PS12] Vincent Pilaud and Francisco Santos. The brick polytope of a sorting network. *European J. Combin.*, 33(4):632–662, 2012.
- [PS15] Vincent Pilaud and Christian Stump. Brick polytopes of spherical subword complexes and generalized associahedra. *Adv. Math.*, 276:1–61, 2015.
- [Rea06] Nathan Reading. Cambrian lattices. *Adv. Math.*, 205(2):313–353, 2006.
- [Rea07] Nathan Reading. Sortable elements and Cambrian lattices. *Algebra Universalis*, 56(3-4):411–437, 2007.
- [RS09] Nathan Reading and David E. Speyer. Cambrian fans. *J. Eur. Math. Soc.*, 11(2):407–447, 2009.
- [SS93] Steve Shnider and Shlomo Sternberg. *Quantum groups: From coalgebras to Drinfeld algebras*. Series in Mathematical Physics. International Press, Cambridge, MA, 1993.
- [Sta63] Jim Stasheff. Homotopy associativity of H -spaces I, II. *Trans. Amer. Math. Soc.*, 108(2):293–312, 1963.
- [Ste13] Salvatore Stella. Polyhedral models for generalized associahedra via Coxeter elements. *J. Algebraic Combin.*, 38(1):121–158, 2013.
- [Tam51] Dov Tamari. *Monoides préordonnés et chaînes de Malcev*. PhD thesis, Université Paris Sorbonne, 1951.
- [Zel06] Andrei Zelevinsky. Nested complexes and their polyhedral realizations. *Pure Appl. Math. Q.*, 2(3):655–671, 2006.
- [Zie95] Günter M. Ziegler. *Lectures on Polytopes*, volume 152 of *Graduate texts in Mathematics*. Springer-Verlag, New York, 1995.

(Thibault Manneville) LIX, ÉCOLE POLYTECHNIQUE
 E-mail address: thibault.manneville@lix.polytechnique.fr
 URL: <http://www.lix.polytechnique.fr/~manneville/>

(Vincent Pilaud) CNRS & LIX, ÉCOLE POLYTECHNIQUE, PALAISEAU
 E-mail address: vincent.pilaud@lix.polytechnique.fr
 URL: <http://www.lix.polytechnique.fr/~pilaud/>