

Periodic solutions of semilinear Duffing equations with impulsive effects

Yanmin Niu^a, Xiong Li^{1a}

^a*School of Mathematical Sciences, Beijing Normal University, Beijing 100875, P.R. China.*

Abstract

In this paper we are concerned with the existence of periodic solutions for semilinear Duffing equations with impulsive effects. Firstly for the autonomous one, basing on Poincaré-Birkhoff twist theorem, we prove the existence of infinitely many periodic solutions. Secondly, as for the nonautonomous case, the impulse brings us great challenges for the study, and there are only finitely many periodic solutions, which is quite different from the corresponding equation without impulses. Here, taking the autonomous one as an auxiliary equation, we find the relation between these two equations and then obtain the result also by Poincaré-Birkhoff twist theorem.

Keywords: Impulsive differential equations, Periodic solutions, Poincaré-Birkhoff twist theorem

1. Introduction

We are concerned in this paper with the existence of periodic solutions for the second order impulsive differential equation

$$\begin{cases} x'' + g(x) = p(t), & t \neq t_j; \\ \Delta x|_{t=t_j} = 0, \\ \Delta x'|_{t=t_j} = -2x'(t_j-), & j = \pm 1, \pm 2, \dots, \end{cases} \quad (1.1)$$

where $0 \leq t_1 < 2\pi$, $\Delta x|_{t=t_j} = x(t_j+) - x(t_j-)$, $\Delta x'|_{t=t_j} = x'(t_j+) - x'(t_j-)$, $g(x)$, $p(t) \in C(\mathbb{R}, \mathbb{R})$ and $p(t)$ is 2π -periodic. In addition, we

¹Partially supported by the NSFC (11571041) and the Fundamental Research Funds for the Central Universities. Corresponding author.

URL: nym1216@163.com (Yanmin Niu), xli@bnu.edu.cn (Xiong Li)

assume that the impulsive time is 2π -periodic, that is, $t_{j+1} = t_j + 2\pi$ for $j = \pm 1, \pm 2, \dots$.

This problem comes from Duffing equation

$$x'' + g(x) = p(t), \quad (1.2)$$

and there is a wide literature dealing with the existence of periodic solutions for the above Duffing equation, not only because of its physical significance, but also the application of various mathematical techniques on it, such as Poincaré-Birkhoff twist theorem in [11], [14], [15], the variational method in [1], [18] and topological degree or index theories in [4], [5]. Under different assumptions on the function g , for example being superlinear, sublinear, semilinear and so on, there are many interesting results on the existence and multiplicity of periodic solutions of (1.2), see [10], [24], [25], [28], [29] and the references therein. Among these, the existence problem of periodic solutions for semilinear Duffing equations challenges more attentions for its special resonance phenomenon. At resonance, equation (1.2) may have no bounded solutions, therefore the crucial point of solving this problem is to exclude the resonance, and there are many studies on it, see [6], [8], [9] and [30].

Recently, as impulsive equations widely arise in applied mathematics, they attract a lot of attentions and many authors study the general properties of them in [2], [17], along with the existence of periodic solutions of impulsive differential equations via fixed point theory in [20], [21], topological degree theory in [13], [27], and the variational method in [22], [31]. However, different from the extensive study for second order differential equations without impulsive terms, there are only a few results on the existence and multiplicity of periodic solutions for second order impulsive differential equations.

In [26], Qian etc considered the superlinear impulsive differential equation

$$\begin{cases} x'' + g(x) = p(t, x, x'), & t \neq t_j; \\ \Delta x|_{t=t_j} = I_j(x(t_j-), x'(t_j-)), \\ \Delta x'|_{t=t_j} = J_j(x(t_j-), x'(t_j-)), & j = \pm 1, \pm 2, \dots, \end{cases} \quad (1.3)$$

where $0 \leq t_1 < \dots < t_k < 2\pi$, $I_j, J_j : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous maps, $t_{j+k} = t_j + 2\pi$ for $j = \pm 1, \pm 2, \dots$, and g is a continuous function with the superlinear growth condition

$$(g'_0) : \lim_{|x| \rightarrow +\infty} \frac{g(x)}{x} = +\infty.$$

The authors proved via Poincaré-Birkhoff twist theorem the existence of infinitely many periodic solutions of (1.3) with $p = p(t)$, and also the existence of periodic solutions for non-conservative case with degenerate impulsive terms by developing a new twist fixed point theorem.

In [23], we discussed the existence of periodic solutions for the sublinear impulsive differential equation

$$\begin{cases} x'' + g(x) = p(t, x, x'), & t \neq t_j; \\ \Delta x|_{t=t_j} = ax(t_j-), \\ \Delta x'|_{t=t_j} = ax'(t_j-), & j = \pm 1, \pm 2, \dots, \end{cases} \quad (1.4)$$

where $0 \leq t_1 < \dots < t_k < 2\pi$, $a > 0$ is a constant, $t_{j+k} = t_j + 2\pi$ for $j = \pm 1, \pm 2, \dots$, and g is a continuous function with the sublinear growth condition

$$(g_0) : \lim_{|x| \rightarrow +\infty} \frac{g(x)}{x} = 0.$$

Basing on the Poincaré-Bohl fixed point theorem and a new twist fixed point theorem recently established by Qian etc in [26], we obtained the existence of harmonic solutions and subharmonic solutions, respectively. The impulsive function is especially chosen to keeping the arguments of trajectories unchanged under the polar coordinate, such that the obstacle causing by the impulse can be solved.

In this article, we discuss semilinear Duffing equation (1.1) with the impulse, which is different from the superlinear or sublinear case and there is few papers studying on it up to now. As we all know, the existence of impulses, even the simplest impulsive function, may cause complicated dynamic phenomena and bring great difficulties to study. The behavior of solutions with impulsive effects may have great changes compared with solutions without impulses. Here we take the simplest linear equation as an example and consider

$$x'' + x = 0, \quad (1.5)$$

with the impulsive condition

$$\Delta x|_{t=t_j} = 0, \quad \Delta x'|_{t=t_j} = -2x'(t_j-), \quad j = \pm 1, \pm 2, \dots, \quad (1.6)$$

where $0 < t_1 < 2\pi$, $t_{j+1} = t_j + 2\pi$ for $j = \pm 1, \pm 2, \dots$. It is easy to see that without the impulse, all solutions of (1.5) are 2π -periodic and has the form

$$x(t) = C_1 \cos t + C_2 \sin t,$$

where C_1 and C_2 are arbitrary constants. However, with $t_1 = \frac{\pi}{2}$ in (1.6), the solution starting from $(x(0), x'(0)) = (x_0, y_0)$ has the form

$$x(t) = \begin{cases} x_0 \cos t + y_0 \sin t, & t \in [0, \frac{\pi}{2}), \\ y_0 \cos(t - \frac{\pi}{2}) + x_0 \sin(t - \frac{\pi}{2}), & t \in [\frac{\pi}{2}, \frac{5\pi}{2}), \\ y_0 \cos(t - \frac{5\pi}{2}) - x_0 \sin(t - \frac{5\pi}{2}), & t \in [\frac{5\pi}{2}, 4\pi], \end{cases} \quad (1.7)$$

from which one has $(x(2\pi), x'(2\pi)) = (-x_0, y_0)$ and $(x(4\pi), x'(4\pi)) = (x_0, y_0)$. Therefore for $x_0 \neq 0$, the solutions of (1.5) have the least period 4π , which implies that the impulse breaks the rule of periods of solutions. On the other hand, if $x_0 = 0$,

$$x(t) = y_0 \sin t, \quad t \in \mathbb{R},$$

the period of which is 2π . In this case, the impulse exists in name only and has no any effects on solutions. More specific discussions about this linear impulsive equation are in [2] Chapter II.

The construction of the impulsive function in (1.1) is also inspired by the oscillator with obstacle, which has the form

$$\begin{cases} x'' + g(x) = f(t), & x(t) > 0; \\ x(t) \geq 0; \\ x(t_0) = 0 \Rightarrow x'(t_0+) = -x'(t_0-), \end{cases} \quad (1.8)$$

where g and f are continuous functions. This equation has actual physical backgrounds in classical billiard, in electric and magnetic fields, three body problem and so on, see [3], [7], [16]. Equation (1.8) can be thought of the model of the motion of a particle which is attached to a spring that pushes the particle against a rigid wall situated at $x = 0$, and at this barrier the particle bounces elastically. At the bouncing time t_0 satisfying $x(t_0) = 0$, the velocity change $x'(t_0+) = -x'(t_0-)$ is same as the impulsive function in (1.1). In both cases, the motions are actually continuous and just the velocity changes the sign at the impulsive/bouncing times. However, in (1.1) the impulsive times are given in advance while in (1.8) the bouncing times depend on the specific solutions, even any positive solutions never meet the wall.

It is well known that the solutions of general autonomous Duffing equations are a family of closed curves in the phase plane. Due to the existence of impulses, the closed curves may not be preserved and even be broken.

For this point, the impulsive function in (1.1) guarantees the motion of trajectories still on the closed curves, which reduces the difficulty causing by impulses.

The rest is organized as follows. The main results about the existence of periodic solutions of (1.1) and (2.5), together with some basic lemmas and symbols, are given in Section 2. In Section 3, we consider the autonomous equation with the same impulsive condition as (1.1), then obtain infinitely many 2π -periodic solutions by Poincaré-Birkhoff twist theorem. In the last section, from the discussion about the autonomous equation, the relation between arguments of the two equations under the polar coordinate is established. Again it follows from Poincaré-Birkhoff twist theorem that there are finitely many 2π -periodic solutions of (1.1).

2. The main results

For convenience, we introduce some basic lemmas and symbols which are used in the sequel. Firstly consider Duffing equation without impulses and perturbation

$$x'' + g(x) = 0, \quad (2.1)$$

and its equivalent system

$$x' = y, \quad y' = -g(x). \quad (2.2)$$

This is a planar autonomous system whose orbits are curves determined by the following equation

$$V(x, y) = \frac{1}{2}y^2 + G(x) = c, \quad (2.3)$$

where $G(x) = \int_0^x g(u)du$ and $c > 0$ is a parameter. Then $V^{-1}(c)$ is star-shaped, the proof of which we omit and one can refer to [8].

Lemma 2.1. (*[8]*) *If the condition (H_2) below holds, then $V^{-1}(c)$ is a closed curve for $c \geq c_0$ which is star-shaped about the origin, where c_0 is a positive constant.*

Denote the curve $V^{-1}(c)$ by Γ_c . It follows from Lemma 2.1 that each curve Γ_c ($c \geq c_0$) intersects the x -axis at two points: $(h(c), 0)$ and $(-h_1(c), 0)$, where $h(c) > 0$ and $h_1(c) > 0$ are uniquely determined by the formula

$$G(h(c)) = G(-h_1(c)) = c. \quad (2.4)$$

Let $(x(t), y(t))$ be any solutions of (2.2) whose orbit is Γ_c ($c \geq c_0$). Obviously, this solution is periodic and denote by $\tau(c)$ the least positive period of it. It can be induced from equation (2.3) that

$$\tau(c) = \sqrt{2} \int_{-h_1(c)}^{h(c)} \frac{du}{\sqrt{c - G(u)}}.$$

Generally speaking, Poincaré-Birkhoff twist theorem is a powerful tool to obtain the existence of periodic solutions. Here we briefly restate a generalized form of it in [12].

Let \mathcal{D} denote an annular region in the (x, y) -plane. The boundary of \mathcal{D} consists of two simple closed curves: the inner boundary curve C_1 and the outer boundary curve C_2 . Let \mathcal{D}_1 denote the simple connected open set bounded by C_1 . Consider an area-preserving mapping $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$. Suppose that $T(\mathcal{D}) \subset \mathbb{R}^2 - 0$, where 0 is the origin. Suppose (γ, θ) be the polar coordinate of (x, y) , that is, $x = \gamma \cos \theta, y = \gamma \sin \theta$. Assume the restriction $T|_{\mathcal{D}}$ is given by

$$\gamma^* = f(\gamma, \theta), \quad \theta^* = \theta + g(\gamma, \theta),$$

where f and g are continuous in (γ, θ) and 2π -periodic in θ .

Lemma 2.2. ([12]) *Besides the above mentioned assumptions, we assume that*

1. C_1 is star-shaped about the origin;
2. $0 \in T(\mathcal{D}_1)$;
3. $g(\gamma, \theta) > 0 (< 0)$, $(\gamma \cos \theta, \gamma \sin \theta) \in C_1$;
 $g(\gamma, \theta) < 0 (> 0)$, $(\gamma \cos \theta, \gamma \sin \theta) \in C_2$.

Then T has at least two fixed points in \mathcal{D} .

Before stating the main results, we give the following hypotheses:

(H_1) Let $g(x) \in C^1(\mathbb{R}, \mathbb{R})$, and K be a positive constant, such that

$$|g'(x)| \leq K, \quad x \in \mathbb{R};$$

(H_2) There exist two constants $A_0 > 0$ and $M_0 > 0$, such that

$$x^{-1}g(x) \geq A_0, \quad |x| \geq M_0.$$

(H_3) There exist a constant $\alpha > 0$, an integer $m > 0$, and two sequences a_k and b_k , such that $a_k \rightarrow \infty$ and $b_k \rightarrow \infty$ as $k \rightarrow \infty$; and moreover

$$\tau(a_k) < \frac{2\pi}{m} - \alpha, \quad \tau(b_k) > \frac{2\pi}{m} + \alpha.$$

These assumptions are completely same as that in [8], and Ding in this paper constructed a function g satisfying $(H_1) - (H_3)$ very skillfully.

Our results on equation (1.1) are different for $p(t) = 0$ and $p(t) \neq 0$. Then we will discuss the two cases respectively in the following parts. The autonomous impulsive equation has the form

$$\begin{cases} z'' + g(z) = 0, & t \neq t_j; \\ \Delta z|_{t=t_j} = 0, \\ \Delta z'|_{t=t_j} = -2z'(t_j-), & j = \pm 1, \pm 2, \dots, \end{cases} \quad (2.5)$$

where $0 \leq t_1 < 2\pi$, $t_{j+1} = t_j + 2\pi$ for $j = \pm 1, \pm 2, \dots$.

Theorem 2.3. *Assume that $(H_1) - (H_3)$ hold. Then equation (2.5) has infinitely many 2π -periodic solutions.*

As for (1.1), due to the existence of the impulse, we only guarantee that the number of 2π -periodic solutions is finite, which is not totally same as the result on semilinear Duffing equation without the impulse in [8].

Theorem 2.4. *Assume that $(H_1) - (H_3)$ hold and $p(t)$ is a 2π -periodic continuous function. Then equation (1.1) has finitely many 2π -periodic solutions.*

3. Autonomous Duffing impulsive equation

We start with equation (2.5), which is a special case of (1.1) with $p(t) = 0$, and its equivalent system is

$$\begin{cases} z' = w, & w' = -g(z), & t \neq t_j; \\ \Delta z|_{t=t_j} = 0, \\ \Delta w|_{t=t_j} = -2w(t_j-), & j = \pm 1, \pm 2, \dots. \end{cases} \quad (3.1)$$

Let $(\bar{z}(t, z, w), \bar{w}(t, z, w))$ be the solution of (3.1) with the initial point $(\bar{z}(0), \bar{w}(0)) = (z, w)$. It is not hard to show that every such solution exists on the whole t -axis under the condition (H_1) (see [2]). Then the Poincaré map $P_1 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is well defined by

$$(z, w) \rightarrow (\bar{z}(2\pi, z, w), \bar{w}(2\pi, z, w)).$$

Next, we take the transform $\bar{z}(t) = \rho(t) \cos \varphi(t)$, $\bar{w}(t) = \rho(t) \sin \varphi(t)$ for system (3.1). Then the resulting equations for $\rho(t)$ and $\varphi(t)$ are

$$\begin{cases} \rho' = \rho \cos \varphi \cdot \sin \varphi - g(\rho \cos \varphi) \sin \varphi, \\ \varphi' = -\sin^2 \varphi - \frac{1}{\rho} g(\rho \cos \varphi) \cos \varphi, & t \neq t_j; \\ \rho(t_j+) = \rho(t_j-), \\ \varphi(t_j+) = \arctan(-\tan \varphi(t_j-)), & j = \pm 1, \pm 2, \dots \end{cases} \quad (3.2)$$

Let $(\bar{\rho}(t, \rho, \varphi), \bar{\varphi}(t, \rho, \varphi))$ be the solution of (3.2) with the initial point $(\bar{\rho}(0), \bar{\varphi}(0)) = (\rho, \varphi)$. Now we estimate the variation of $\bar{\varphi}(t, \rho, \varphi)$ during the time interval $[0, 2\pi]$.

Lemma 3.1. *Let $\Phi(\rho, \varphi) = \bar{\varphi}(2\pi, \rho, \varphi) - \varphi$. Then for k sufficiently large, there exist two positive constants β_1, β_2 , such that*

$$\begin{cases} \Phi(\rho, \varphi) \leq -2m\pi - \beta_1, & (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{a_k}; \\ \Phi(\rho, \varphi) \geq -2m\pi + \beta_2, & (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{b_k}, \end{cases} \quad (3.3)$$

where m, a_k, b_k are given in (H_3) .

Proof. Suppose $(z, w) = (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{a_k}$. Consider the solution $(\bar{\rho}(t, \rho, \varphi), \bar{\varphi}(t, \rho, \varphi))$ of (3.2). Without loss of generality, let the initial point (ρ, φ) be on the positive x -axis, that is, $(\rho, \varphi) = (\rho, 0)$. From (H_2) , there exist two constants $c_0 > 0$ and $A_1 > 0$ such that if $c \geq c_0$, then

$$w^2 + zg(z) \geq A_1(z^2 + w^2), \quad (z, w) \in \Gamma_c. \quad (3.4)$$

It follows from (3.4) and the second equation of (3.2) that for k large enough satisfying $a_k \geq c_0$,

$$\bar{\varphi}'(t, \rho, \varphi) \leq -A_1. \quad (3.5)$$

Since the solution of (2.2) lying in Γ_{a_k} has the least period $\tau(a_k)$, we know that the time in which $\bar{\varphi}(t)$ has a decrement 2π without the impulse is just $\tau(a_k)$. Denote

$$\Phi(\rho, \varphi) = \bar{\varphi}(2\pi, \rho, \varphi) - \varphi = \bar{\varphi}(2\pi) - \bar{\varphi}(0) = -2l\pi - \sigma,$$

where $l \geq 0$ is an integer, and $0 \leq \sigma < 2\pi$. Let t_σ denote the time in which $\bar{\varphi}(t)$ decreases from $\varphi - 2l\pi$ to $\varphi - 2l\pi - \sigma$. Then there are three possible cases according to different positions of $\bar{\varphi}(t)$ when the impulse occurs.

Case i The impulsive time t_1 occurs during the time interval in which $\bar{\varphi}(t)$ travels the front $l - 1$ loops, that is, in the time interval $[t_1, 2\pi]$, $\bar{\varphi}(t)$ can travel at least one loop. Then there exists $0 < \tau^* \leq \tau(a_k)$ such that

$$(l - 1)\tau(a_k) + \tau^* + t_\sigma = 2\pi,$$

and the impulsive time t_1 is out of the time interval in which $\bar{\varphi}(t)$ decreases from $\varphi - 2l\pi$ to $\varphi - 2l\pi - \sigma$. Assume the plane coordinate of the solution at $t_1 \pm$ be $(z(t_1 \pm), w(t_1 \pm))$, then τ^* stands for the time of $\bar{\varphi}(t)$ traveling the argument of 2π under the impulsive effect and has the following form

$$\tau^* = \tau(a_k) - \sqrt{2} \int_{z(t_1-)}^{h(a_k)} \frac{du}{\sqrt{a_k - G(u)}},$$

when $(z(t_1-), w(t_1-))$ is in the first or the second quadrant, and has the form

$$\tau^* = \tau(a_k) - \sqrt{2} \int_{-h_1(a_k)}^{z(t_1+)} \frac{du}{\sqrt{a_k - G(u)}},$$

when $(z(t_1-), w(t_1-))$ is in the third or the fourth quadrant, where h, h_1 are defined in (2.4).

In this case we infer $l \geq 1$. Since $0 \leq t_\sigma < \tau(a_k)$, by the condition (H_3) it holds that

$$\begin{aligned} 2\pi &= (l - 1)\tau(a_k) + \tau^* + t_\sigma \\ &< (l - 1)\tau(a_k) + \tau(a_k) + \tau(a_k) \\ &= (l + 1)\tau(a_k) \\ &\leq (l + 1) \left(\frac{2\pi}{m} - \alpha \right), \end{aligned}$$

which implies $l \geq m$. If $l \geq m + 1$, we have

$$\Phi(\rho, \varphi) = -2l\pi - \sigma \leq -2l\pi \leq -2(m + 1)\pi. \quad (3.6)$$

Now, assume $l = m$. Then we have

$$\begin{aligned} t_\sigma &= 2\pi - (m - 1)\tau(a_k) - \tau^* \\ &> 2\pi - (m - 1)\tau(a_k) - \tau(a_k) \\ &= 2\pi - m\tau(a_k) \\ &\geq 2\pi - m \left(\frac{2\pi}{m} - \alpha \right) = m\alpha. \end{aligned} \quad (3.7)$$

By (3.5) and (3.7), we obtain

$$-\sigma = \int_{(l-1)\tau(a_k)+\tau^*}^{(l-1)\tau(a_k)+\tau^*+t_\sigma} \bar{\varphi}'(t, \rho, \varphi) dt \leq -A_1 t_\sigma \leq -m\alpha A_1.$$

Thus

$$\Phi(\rho, \varphi) = -2l\pi - \sigma \leq -2m\pi - m\alpha A_1. \quad (3.8)$$

Combining (3.6) and (3.8) yields the first inequality of (3.3), where $\beta_1 = \min\{2\pi, m\alpha A_1\}$.

Case ii The impulsive time t_1 occurs at the l th loop and in the time interval $[t_1, 2\pi]$, $\bar{\varphi}(t)$ travels no more than one loop. Then there exists $0 < \tau^* \leq \tau(a_k)$ such that

$$(l-1)\tau(a_k) + \tau^* + t_\sigma = 2\pi,$$

and the impulsive time t_1 has impact on the time interval in which $\bar{\varphi}(t)$ decreases from $\varphi - 2l\pi$ to $\varphi - 2l\pi - \sigma$. More specifically, this case happens when $(z(t_1-), w(t_1-))$ is in the first quadrant or the second quadrant, such that τ^* has the form

$$\tau^* = \tau(a_k) - \frac{\sqrt{2}}{2} \int_{z(t_1-)}^{h(a_k)} \frac{du}{\sqrt{a_k - G(u)}}.$$

Also we infer $l \geq 1$ and similarly to the proof of case (i), we obtain $l \geq m$, together with (3.6) and (3.7). By (3.5) and (3.7), we have

$$\begin{aligned} -\sigma &= \frac{1}{2} (\bar{\varphi}'(t_1+, \rho, \varphi) - \bar{\varphi}'(t_1+, \rho, \varphi)) + \int_{(l-1)\tau(a_k)+\tau^*}^{(l-1)\tau(a_k)+\tau^*+t_\sigma} \bar{\varphi}'(t, \rho, \varphi) dt \\ &\leq \int_{(l-1)\tau(a_k)+\tau^*}^{(l-1)\tau(a_k)+\tau^*+t_\sigma} \bar{\varphi}'(t, \rho, \varphi) dt \\ &\leq -A_1 t_\sigma \\ &\leq -m\alpha A_1. \end{aligned}$$

Thus (3.8) holds. Again combining (3.6) and (3.8) yields the first inequality of (3.3), where $\beta_1 = \min\{2\pi, m\alpha A_1\}$.

Case iii The movement of $\bar{\varphi}(t)$ during the front l loops is out of the effect of the impulsive time t_1 . Then

$$l\tau(a_k) + t_\sigma = 2\pi$$

and the impulsive time t_1 is in the time interval in which $\bar{\varphi}(t)$ decreases from $\varphi - 2l\pi$ to $\varphi - 2l\pi - \sigma$. In this case,

$$t_1 \in (l \cdot \tau(a_k), 2\pi).$$

Since $0 \leq t_\sigma < \tau(a_k)$, by the condition (H_3) it holds that

$$2\pi = l \cdot \tau(a_k) + t_\sigma < (l+1)\tau(a_k) \leq (l+1) \left(\frac{2\pi}{m} - \alpha \right),$$

which implies $l \geq m$. If $l \geq m+1$, (3.6) holds. Assume $l = m$. Then we have

$$t_\sigma = 2\pi - m\tau(a_k) \geq 2\pi - m \left(\frac{2\pi}{m} - \alpha \right) = m\alpha. \quad (3.9)$$

By (3.5) and (3.9), we obtain

$$\begin{aligned} -\sigma &= \int_{l \cdot \tau(a_k)}^{t_1^-} \bar{\varphi}'(t, \rho, \varphi) dt + (\bar{\varphi}'(t_1+, \rho, \varphi) - \bar{\varphi}'(t_1-, \rho, \varphi)) + \int_{t_1+}^{2\pi} \bar{\varphi}'(t, \rho, \varphi) dt \\ &\leq -A_1(t_1 - l \cdot \tau(a_k)) - A_1(2\pi - t_1) \\ &= -A_1 t_\sigma \\ &\leq -m\alpha A_1. \end{aligned}$$

Thus

$$\Phi(\rho, \varphi) = -2l\pi - \sigma \leq -2m\pi - m\alpha A_1.$$

Similarly, we obtain the first inequality of (3.3), where $\beta_1 = \min\{2\pi, m\alpha A_1\}$.

We are now in a position to prove the second inequality of (3.3). Suppose $(z, w) = (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{b_k}$, where k is sufficiently large such that $b_k \geq c_0$. We see that the time in which $\bar{\varphi}(t)$ has a decrement 2π without the impulse is just $\tau(b_k)$. Denote

$$\Phi(\rho, \varphi) = \bar{\varphi}(2\pi, \rho, \varphi) - \varphi = \bar{\varphi}(2\pi) - \bar{\varphi}(0) = -2q\pi + \xi,$$

where $q \geq 1$ is an integer, and $0 < \xi \leq 2\pi$. Let t_ξ denote the time in which $\bar{\varphi}(t)$ decreases from $\varphi - 2q\pi + \xi$ to $\varphi - 2q\pi$. In the following proof, we always assume that during the time interval from 2π to $2\pi + t_\xi$, there is under no influence of another impulsive time. In fact, if there is another impulsive time t_2 in $(2\pi, 2\pi + t_\xi)$, the proof below is similar and we just need to change some signs in (3.14). Under these assumptions, the three cases discussed above on Γ_{a_k} can be simplified into one case: among the q loops, only one loop is effected by the impulse. Then there exists $\tau^\wedge \in (0, \tau(b_k)]$ such that

$$(q-1)\tau(b_k) + \tau^\wedge - t_\xi = 2\pi,$$

where

$$\tau^\wedge = \tau(b_k) - \sqrt{2} \int_{z(t_1^-)}^{h(a_k)} \frac{du}{\sqrt{b_k - G(u)}},$$

when $(z(t_1-), w(t_1-))$ is in the first or the second quadrant, and

$$\tau^\wedge = \tau(b_k) - \sqrt{2} \int_{-h_1(b_k)}^{z(t_1+)} \frac{du}{\sqrt{b_k - G(u)}},$$

when $(z(t_1-), w(t_1-))$ is in the third or the fourth quadrant.

Since $0 < \tau^\wedge \leq \tau(b_k)$, let $\tau^\wedge = \mu\tau(b_k)$, then $0 < \mu \leq 1$. By the condition (H_3) and $0 < t_\xi \leq \tau(b_k)$, it holds that

$$2\pi = (q-1)\tau(b_k) + \tau^\wedge - t_\xi \geq (q-2+\mu)\tau(b_k) > (q-2+\mu) \left(\frac{2\pi}{m} + \alpha \right), \quad (3.10)$$

provided $q > 1$. Notice that if $q = 1$,

$$\Phi(\rho, \varphi) = -2\pi + \xi \geq -2m\pi + \xi,$$

then letting $\beta_2 = \xi$, we obtain the result. We just consider the case $q > 1$. It follows from (3.10) that $q \leq m$. Notice that if m given in (H_3) equals to 1, then $q = 1$. Similarly to the discussion above, we obtain the result. Next assume $m > 1$. If $q \leq m - 1$, we have

$$\Phi(\rho, \varphi) = -2q\pi + \xi > -2q\pi = -2(m-1)\pi. \quad (3.11)$$

Now if $q = m$, writing $\tau^\wedge = \tau(b_k) - \eta$, we have

$$\begin{aligned} t_\xi &= -2\pi + (m-1)\tau(b_k) + \tau^\wedge \\ &= -2\pi + m\tau(b_k) - \eta \\ &\geq -2\pi + m \left(\frac{2\pi}{m} + \alpha \right) - \eta \\ &= m\alpha - \eta. \end{aligned} \quad (3.12)$$

Since $\xi > 0$, $t_\xi > \frac{1}{2}t_\xi > 0$, then there exists $\varepsilon_\xi > 0$, such that

$$t_\xi \geq \max\{\varepsilon_\xi, m\alpha - \eta\} \triangleq M_\xi > 0. \quad (3.13)$$

By (3.5) and (3.13), we have

$$\begin{aligned} \xi &= \int_{(q-1)\tau(b_k) + \tau^\wedge}^{(q-1)\tau(b_k) + \tau^\wedge - t_\xi} \overline{\varphi}'(t, \rho, \varphi) dt \\ &= - \int_{(q-1)\tau(b_k) + \tau^\wedge - t_\xi}^{(q-1)\tau(b_k) + \tau^\wedge} \overline{\varphi}'(t, \rho, \varphi) dt \\ &\geq -(-A_1 t_\xi) \\ &\geq M_\xi A_1. \end{aligned} \quad (3.14)$$

Thus

$$\Phi(\rho, \varphi) = -2q\pi + \xi \geq -2m\pi + M_\xi A_1. \quad (3.15)$$

Combining (3.11) and (3.15) yields the second inequality of (3.3), where $\beta_2 = \min\{2\pi, M_\xi A_1\}$.

The proof of Lemma 3.1 is then completed. \square

Remark 3.2. β_2 is related to k . Indeed, for each k sufficiently large, $\beta_2 = \min\{2\pi, M_\xi A_1\} > 0$, where $M_\xi \triangleq \max\{\varepsilon_\xi, m\alpha - \eta\} > 0$ may depend on k . However, $\beta_1 = \min\{2\pi, m\alpha A_1\}$ is independent of k .

Under the polar coordinate, the Poincaré map P_1 can be written in the form

$$\rho^* = \bar{\rho}(2\pi, \rho, \varphi), \quad \varphi^* = \bar{\varphi}(2\pi, \rho, \varphi) + 2k\pi, \quad (3.16)$$

where k is an arbitrary integer. It can be easily seen that if (ρ, φ) satisfies

$$\bar{\rho}(t, \rho, \varphi) > 0, \quad t \in [0, 2\pi], \quad (3.17)$$

then $\bar{\varphi}(2\pi, \rho, \varphi)$ is well defined and continuous in (ρ, φ) , moreover,

$$\bar{\varphi}(2\pi, \rho, \varphi + 2\pi) = \bar{\varphi}(2\pi, \rho, \varphi) + 2\pi.$$

Now, let Γ_{a_k} and Γ_{b_k} be the curves given by Lemma 2.1, where the specified parameters $a_k, b_k \geq c_0$ are given by (H_3) , for $k \geq n_0$. We can rearrange a_k and b_k such that $a_k < b_k < a_{k+1}$ for $k \geq n_0$. Then Γ_{a_k} and Γ_{b_k} bound an annular region \mathcal{A}_k , and Γ_{b_k} and $\Gamma_{a_{k+1}}$ bound another annular region \mathcal{B}_k , for $k \geq n_0$.

It is well known that each fixed point of P_1 corresponds to a 2π -periodic solution of (2.5). In the sequel, we will apply Lemma 2.2 to show that P_1 has at least two fixed points in each \mathcal{A}_k and \mathcal{B}_k for sufficiently large k . As a consequence, (2.5) has an infinite class of 2π -periodic solutions.

We turn to prove Theorem 2.3 by the twist theorem.

Proof of Theorem 2.3 Assume that $a_k, b_k \geq c_0$ for $k \geq n_0$, where n_0 is large enough and c_0 is given in Lemma 2.1. Denote by \mathcal{A}_k the region bounded by Γ_{a_k} and Γ_{b_k} . Thus the restriction $P_1|_{\mathcal{A}_k}$ can be written in (3.16), where we put the integer $k = m$. Then (3.16) can be rewritten in the form of

$$\rho^* = \bar{\rho}(2\pi, \rho, \varphi), \quad \varphi^* = \varphi + \Phi_1(\rho, \varphi),$$

with $\Phi_1(\rho, \varphi) = \bar{\varphi}(2\pi, \rho, \varphi) - \varphi + 2m\pi = \Phi(\rho, \varphi) + 2m\pi$. By Lemma 3.1, we obtain

$$\begin{cases} \Phi_1(\rho, \varphi) \leq -\beta_1 < 0, & (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{a_k}; \\ \Phi_1(\rho, \varphi) \geq \beta_2 > 0, & (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{b_k}, \end{cases}$$

for $k \geq n_0$.

This proves the validity of condition 3 of Lemma 2.2 for the restriction $P_1|_{\mathcal{A}_k}$ ($k \geq n_0$). Since k can be chosen large enough such that a_k, b_k can be sufficiently large, then condition 2 of Lemma 2.2 can be easily verified. By Lemma 2.1, condition 1 of Lemma 2.2 also holds. In addition, we need to clarify P_1 is an area-preserving mapping. Indeed, the Poincaré map $P_1 : (z, w) \rightarrow (\bar{z}(2\pi, z, w), \bar{w}(2\pi, z, w))$ can be expressed by

$$P_1 = P^1 \circ I^1 \circ P^0,$$

where

$$\begin{aligned} P^0 : (z, w) &\mapsto (\bar{z}(t_1-, z, w), \bar{w}(t_1-, z, w)), \\ I^1 : (\bar{z}(t_1-), \bar{w}(t_1-)) &\mapsto (\bar{z}(t_1-) + \Delta z(t_1), \bar{w}(t_1-) + \Delta w(t_1)), \\ P^1 : (\bar{z}(t_1+), \bar{w}(t_1+)) &\mapsto (\bar{z}(2\pi), \bar{z}(2\pi)), \end{aligned}$$

and $\Delta z(t_1), \Delta w(t_1)$ are given in (3.1). Since $P^j, j = 0, 1$ are symplectic by equation $z'' + g(z) = 0$ being conservative and I^1 is area-preserving with $\det(I^1) = 1$, the Poincaré map P_1 is area-preserving.

Therefore, we can apply Lemma 2.2 to ensure the existence of at least two fixed points of P_1 in \mathcal{A}_k ($k \geq n_0$). This means that (2.5) has at least two 2π -periodic solutions with initial points in \mathcal{A}_k . Similarly, we can prove that P_1 has at least two fixed points in \mathcal{B}_k which correspond to two 2π -periodic solutions of (2.5). Since each periodic solutions of (2.5) is bounded by Γ_{a_k} and Γ_{b_k} , then the above specified 2π -periodic solutions of (2.5) constitute an infinite class.

The proof of Theorem 2.3 is thus completed.

4. Nonautonomous Duffing impulsive equation

We write (1.1) as an equivalent system of the form

$$\begin{cases} x' = y, & y' = -g(x) + p(t), & t \neq t_j; \\ \Delta x|_{t=t_j} = 0, \\ \Delta y|_{t=t_j} = -2y(t_j-), & j = \pm 1, \pm 2, \dots \end{cases} \quad (4.1)$$

Let $(\bar{x}(t, x, y), \bar{y}(t, x, y))$ be the solution of (4.1) with the initial point $(\bar{x}(0), \bar{y}(0)) = (x, y)$. Also it follows from the condition (H_1) that every such solution exists on the whole t -axis (see [2]). Then the Poincaré map $P_2 : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is well defined by

$$(x, y) \rightarrow (\bar{x}(2\pi, x, y), \bar{y}(2\pi, x, y)).$$

By applying the transform $\bar{x}(t) = \gamma(t) \cos \theta(t), \bar{y}(t) = \gamma(t) \sin \theta(t)$ to (4.1), we get the equations for $\gamma(t)$ and $\theta(t)$,

$$\begin{cases} \gamma' = \gamma \cos \theta \cdot \sin \theta - g(\gamma \cos \theta) \sin \theta + p(t) \sin \theta, \\ \theta' = -\sin^2 \theta - \frac{1}{\gamma} [g(\gamma \cos \theta) \cos \theta - p(t) \cos \theta], & t \neq t_j; \\ \gamma(t_j+) = \gamma(t_j-), \\ \theta(t_j+) = \arctan(-\tan \theta(t_j-)), & j = \pm 1, \pm 2, \dots \end{cases} \quad (4.2)$$

Let $(\bar{\gamma}(t, \gamma, \theta), \bar{\theta}(t, \gamma, \theta))$ be the solution of (4.2) through the initial point $(\bar{\gamma}(0), \bar{\theta}(0)) = (\gamma, \theta)$. Then the map P_2 also can be written in the polar coordinate form

$$\gamma^* = \bar{\gamma}(2\pi, \gamma, \theta), \quad \theta^* = \bar{\theta}(2\pi, \gamma, \theta) + 2l\pi, \quad (4.3)$$

where l is an arbitrary integer. It can be easily seen that if (γ, θ) satisfies

$$\bar{\gamma}(t, \gamma, \theta) > 0, \quad t \in [0, 2\pi], \quad (4.4)$$

then $\bar{\theta}(2\pi, \gamma, \theta)$ is well defined and continuous in (γ, θ) , moreover,

$$\bar{\theta}(2\pi, \gamma, \theta + 2\pi) = \bar{\theta}(2\pi, \gamma, \theta) + 2\pi.$$

Based on Lemma 3.1 in Section 3, we first will study the intimate relation between equations (1.1) and (2.5), especially the motions of their arguments under the polar coordinates. Let

$$\Theta(\gamma, \theta) = \bar{\theta}(2\pi, \gamma, \theta) - \theta.$$

The following Lemma will estimate the difference between $\Theta(\gamma, \theta)$ and $\Phi(\gamma, \theta)$ with the same initial point.

Lemma 4.1. *For any $\varepsilon > 0$, there exists $\gamma^* > 0$ such that, for $\gamma \geq \gamma^*$,*

$$|\Theta(\gamma, \theta) - \Phi(\gamma, \theta)| = |\bar{\theta}(2\pi, \gamma, \theta) - \bar{\varphi}(2\pi, \gamma, \theta)| < \varepsilon.$$

Proof. For $0 < \varepsilon < \pi$, let $(\bar{z}(t, x, y), \bar{w}(t, x, y))$ be the solution of (3.1) with the initial point $(\bar{z}(0), \bar{w}(0)) = (x, y)$. Denote

$$\begin{cases} u(t) = u(t, x, y) = \bar{x}(t, x, y) - \bar{z}(t, x, y), \\ v(t) = v(t, x, y) = \bar{y}(t, x, y) - \bar{w}(t, x, y). \end{cases}$$

Then we have

$$\frac{du}{dt} = v, \quad \frac{dv}{dt} = p(t) - g'(\vartheta(t))u(t), \quad t \neq t_j, \quad j = \pm 1, \pm 2, \dots,$$

where $\vartheta(t) = \bar{x}(t) + \lambda(t)(\bar{z}(t) - \bar{x}(t))$, $0 \leq \lambda(t) \leq 1$. Since the impulsive functions in (3.1) and (4.1) are same and linear, then

$$u(t_j+) = u(t_j), \quad v(t_j+) = -v(t_j), \quad j = \pm 1, \pm 2, \dots. \quad (4.5)$$

Let $\eta(t) = (u^2(t) + v^2(t))^{\frac{1}{2}}$. Then for $t \neq t_j$, we have

$$\eta \frac{d\eta}{dt} = uv + p(t)v - g'(\vartheta(t))uv.$$

It follows from (H_1) that

$$\left| \frac{d\eta}{dt} \right| \leq \frac{1}{2}(1 + K)\eta + B, \quad (4.6)$$

where $B = \sup_{t \in [0, 2\pi]} |p(t)|$.

First we consider the difference between the arguments on $[0, t_1]$. The differential inequality (4.6) together with $\eta(0) = 0$ yields

$$\eta(t) \leq \frac{2B}{1 + K} \left[e^{(1+K)\pi} - 1 \right] \equiv H_0$$

for $t \in [0, t_1]$. Denote

$$\psi(t) = \psi(t, \gamma, \theta) = \bar{\theta}(t, \gamma, \theta) - \bar{\varphi}(t, \gamma, \theta),$$

where (γ, θ) is the polar coordinate of (x, y) , that is, $(\gamma \cos \theta, \gamma \sin \theta) = (x, y)$. It is clear that if $|\psi(t)| < \pi$, then $\psi(t)$ is just the angle between the vectors $(\bar{x}(t), \bar{y}(t))$ and $(\bar{z}(t), \bar{w}(t))$. By the law of cosines, we have

$$\cos \psi(t) = \frac{\bar{\gamma}^2(t) + \bar{\rho}^2(t) - \bar{\eta}^2(t)}{2\bar{\gamma}(t)\bar{\rho}(t)} \geq 1 - \frac{H_0^2}{2\bar{\gamma}(t)\bar{\rho}(t)}, \quad t \in [0, t_1].$$

On the other hand, $\bar{\gamma}(t) \geq \bar{\rho}(t) - \eta(t) \geq \bar{\rho}(t) - H_0$. Therefore, under the assumption that $|\psi(t)| < \pi$ and $\bar{\rho}(t) - H_0 > 0$, we have

$$\cos \psi(t) \geq 1 - \frac{H_0^2}{2\bar{\rho}(t)(\bar{\rho}(t) - H_0)}, \quad t \in [0, t_1]. \quad (4.7)$$

Note that

$$F(\gamma, \theta) = \inf_{t \in [0, 2\pi]} \bar{\rho}(t) = \inf_{t \in [0, 2\pi]} \bar{\rho}(t, \gamma, \theta)$$

becomes arbitrarily large if γ is sufficiently large (see [26]). Then there is a constant $\gamma_0 > 0$ such that, for $\gamma \geq \gamma_0$ and $t \in [0, t_1]$,

$$\bar{\rho}(t) - H_0 > 0, \quad \frac{H_0^2}{2\bar{\rho}(t)(\bar{\rho}(t) - H_0)} < 1 - \cos \delta, \quad (4.8)$$

where $\delta = \min\{\frac{\pi}{2}, \frac{\varepsilon}{2}\}$. From (4.7) and (4.8), we conclude that if $|\psi(t)| < \pi$ on $[0, t_1]$, then the inequality

$$\cos \psi(t) > \cos \delta$$

holds, which implies

$$|\psi(t)| < \delta, \quad t \in [0, t_1].$$

Now we just need to verify

$$|\psi(t)| < \pi, \quad t \in [0, t_1]. \quad (4.9)$$

Since $\psi(0) = 0$ and $\psi(t)$ is continuous on $[0, t_1]$, there exists $a > 0$ such that

$$|\psi(t)| < \pi, \quad t \in [0, a]. \quad (4.10)$$

To prove (4.9), we need to verify $a > t_1$. If it is not true, then the a given above belongs to $(0, t_1]$ such that (4.10) holds and $|\psi(a)| = \pi$. Note that inequality (4.7), together with (4.8) and (4.10) implies

$$|\psi(t)| < \delta, \quad t \in [0, a].$$

Thus we obtain $|\psi(a)| \leq \delta \leq \frac{\pi}{2} < \pi$, which contradicts with $|\psi(a)| = \pi$, hence (4.9) holds and then for $t \in [0, t_1]$,

$$|\psi(t)| < \delta \leq \frac{\varepsilon}{2} < \varepsilon. \quad (4.11)$$

Next we turn to consider $\psi(t)$ on $(t_1, 2\pi]$. By (4.5), it follows that

$$\eta(t_1+) = \eta(t_1) \triangleq \eta_1.$$

Integrating the inequality (4.6) on $(t_1, 2\pi]$, we have

$$\eta(t) \leq \left| \frac{2B_1}{1+K} \left[e^{(1+K)\pi} - \frac{B}{B_1} \right] \right| \equiv H_1, \quad t \in (t_1, 2\pi],$$

where $B_1 = \frac{1+K}{2}\eta_1 + B$. In the same way with the discussion on $[0, t_1]$, we obtain that there is a constant $\gamma_1 > 0$, such that for $\gamma \geq \gamma_1$ and $t \in (t_1, 2\pi]$,

$$\cos \psi(t) \geq 1 - \frac{H_1^2}{2\bar{\rho}(t)(\bar{\rho}(t) - H_1)}, \quad (4.12)$$

and

$$\bar{\rho}(t) - H_1 > 0, \quad \frac{H_1^2}{2\bar{\rho}(t)(\bar{\rho}(t) - H_1)} < 1 - \cos \delta, \quad (4.13)$$

where δ is the same as before, provided $|\psi(t)| < \pi$ on $(t_1, 2\pi]$. It follows that if $|\psi(t)| < \pi$ on $(t_1, 2\pi]$, then the inequality

$$|\psi(t)| < \delta$$

holds.

Similarly, we just need to prove

$$|\psi(t)| < \pi, \quad t \in (t_1, 2\pi]. \quad (4.14)$$

By (4.11), $\psi(t_1) = \psi(t_1-) < \varepsilon$. Since the impulsive functions in (4.2) and (3.2) are same and $\psi(t)$ is continuous in t_1 , $\psi(t_1+)$ can be small enough. Then there exists $a > 0$ such that

$$|\psi(t)| < \pi, \quad t \in (t_1, a). \quad (4.15)$$

We just need to verify $a > 2\pi$. Otherwise, there is an $a \in (t_1, 2\pi]$ such that (4.15) holds and $|\psi(a)| = \pi$. Note that inequality (4.15), together with (4.12) and (4.13) implies

$$|\psi(t)| < \delta, \quad t \in (t_1, a).$$

Thus we obtain $|\psi(a)| \leq \delta \leq \frac{\pi}{2} < \pi$, which contradicts with $|\psi(a)| = \pi$. Then (4.14) holds and for $t \in (t_1, 2\pi]$,

$$|\psi(t)| < \delta \leq \frac{\varepsilon}{2} < \varepsilon.$$

In particular, $|\psi(2\pi)| < \varepsilon$.

Consequently, choosing $\gamma^* = \max\{\gamma_0, \gamma_1\}$, for $\gamma \geq \gamma^*$, we have

$$|\Theta(\gamma, \theta) - \Phi(\gamma, \theta)| = |\bar{\theta}(2\pi, \gamma, \theta) - \bar{\varphi}(2\pi, \gamma, \theta)| = |\psi(2\pi)| < \varepsilon.$$

The proof of Lemma 4.1 is then completed. \square

By Lemma 3.1, together with the small difference between $\bar{\theta}(2\pi)$ and $\bar{\varphi}(2\pi)$, we can obtain the existence of 2π -periodic solutions. However, according to Remark 3.2, we cannot guarantee the application of the twist theorem for all sufficiently large k . Then the number of 2π -periodic solutions of (1.1) is limited instead of infinity.

Proof of Theorem 2.4 Let $c_1 \geq c_0$ be sufficiently large, such that for $c \geq c_1$, $(\gamma, \theta) \in \Gamma_c$ implies $\gamma \geq \gamma^*$, where γ^* is specified in Lemma 4.1. There is no loss of generality to assume that $a_k, b_k \geq c_1$ for $k \geq n_0$, where n_0 is large enough. It follows that

$$\begin{cases} \bar{\gamma}(t, \gamma, \theta) \geq \bar{\rho}(t) - H_0 > 0, & t \in [0, t_1], \\ \bar{\gamma}(t, \gamma, \theta) \geq \bar{\rho}(t) - H_1 > 0, & t \in (t_1, 2\pi], \end{cases} \quad (4.16)$$

provided that $(\gamma, \theta) \in \mathcal{A}_k$ for $k \geq n_0$. Thus the restriction $P_2|_{\mathcal{A}_k}$ can be written in (4.3), where we put the integer $l = m$. Rewritten (4.3) in the following form

$$\gamma^* = \bar{\gamma}(2\pi, \gamma, \theta), \quad \theta^* = \theta + \Theta_1(\gamma, \theta),$$

with $\Theta_1(\gamma, \theta) = \Theta(\gamma, \theta) + 2m\pi$.

From Remark 3.2, we first choose arbitrary finitely many integers k_1, k_2, \dots, k_n with $k_i \geq n_0$, $i = 1, 2, \dots, n$. Correspondingly, for each k_i , denote the curve by $\Gamma_{a_{k_i}}, \Gamma_{b_{k_i}}$ and the annulus by $\mathcal{A}_{k_i}, \mathcal{B}_{k_i}$. To be precise, in Lemma 3.1, we denote $\beta_2 = \beta_2^i$ for each k_i . Let $\varepsilon_0 = \min\{\beta_1, \beta_2^1, \beta_2^2, \dots, \beta_2^n\}$. Since $\beta_1 > 0$ and $\beta_2^i > 0$, $i = 1, 2, \dots, n$, then $\varepsilon_0 > 0$. By Lemma 4.1, choosing $\varepsilon = \varepsilon_0$, we obtain

$$|\Theta_1(\gamma, \theta) - \Phi_1(\gamma, \theta) - 2m\pi| < \varepsilon_0,$$

which together with (3.3) yields

$$\begin{cases} \Theta_1(\gamma, \theta) < 0, & (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{a_{k_i}}; \\ \Theta_1(\gamma, \theta) > 0, & (\rho \cos \varphi, \rho \sin \varphi) \in \Gamma_{b_{k_i}}. \end{cases}$$

This proves the validity of condition 3 of Lemma 2.2 for the restriction $P_2|_{\mathcal{A}_{k_i}}$ ($k_i \geq n_0$, $i = 1, 2, \dots, n$). Since (4.16) holds, then condition 2 of

Lemma 2.2 can be easily verified. By Lemma 2.1, condition 1 of Lemma 2.2 also holds. Similar to the proof of Theorem 2.3, area-preserving property of P_2 is obtained.

Therefore, we can apply Lemma 2.2 to ensure the existence of at least two fixed points of P_2 in \mathcal{A}_{k_i} ($k_i \geq n_0$, $i = 1, 2, \dots, n$). This means that (1.1) has at least two 2π -periodic solutions with initial points in \mathcal{A}_{k_i} ($k_i \geq n_0$, $i = 1, 2, \dots, n$). Similarly, we can prove that P_2 has at least two fixed points in \mathcal{B}_{k_i} ($k_i \geq n_0$, $i = 1, 2, \dots, n$) which correspond to two 2π -periodic solutions of (1.1). Since each period solutions of (1.1) is bounded by $\Gamma_{a_{k_i}}, \Gamma_{b_{k_i}}$, then the above specified 2π -periodic solutions of (1.1) constitute a finite class of solutions.

The proof of Theorem 2.4 is thus completed.

References

References

- [1] A. Bahri, H. Berestycki, Existence of forced oscillations for some nonlinear differential equations, *Comm. Pure Appl. Math.* 37 (1984) 403-442.
- [2] D. Bainov, P. Simenov, *Impulsive Differential Equations: Periodic Solutions and Applications*, Longman Scientific and Technical, Essex, England, 1993.
- [3] P. Boyland, Dual billiards, twist maps and impact oscillators, *Nonlinearity*, 9 (1996), 1411-1438.
- [4] A. Capietto, J. Mawhin, F. Zanolin, Continuation theorems for periodic perturbations of autonomous systems, *Trans. Amer. Math. Soc.* 329 (1992) 41-72.
- [5] A. Capietto, J. Mawhin, F. Zanolin, A continuation approach for superlinear periodic value problems, *J. Differential Equations* 88 (1990) 347-395.
- [6] A. C. Lazer, D. E. Leach, Bounded perturbations of forced harmonic oscillations at resonance, *Ann. Mat. Pure Appl.* 82 (1969) 49-68.
- [7] M. Corbera, J. Llibre, Periodic orbits of a collinear restricted three body problem, *Celestial. Mech. Dynam. Astronom.* 86 (2003) 163-183.

- [8] T. Ding, An infinite class of periodic solutions of periodically perturbed Duffing equations at resonance, Proc. Amer. Math. Soc. 86 (1982) 47-54.
- [9] T. Ding, Nonlinear oscillations at a point of resonance, Sci. Sinica (Chinese), Series A, 1 (1982) 1-13.
- [10] T. Ding, F. Zanolin, On periodic solutions of sublinear Duffing equations, J. Math. Anal. Appl. 158 (1991) 316-332.
- [11] T. Ding, F. Zanolin, Periodic solutions of Duffing's equations with superquadratic potential, J. Differential Equations 97 (1992) 328-378.
- [12] W. Y. Ding, A generalization of the Poincaré-Birkhoff theorem, Proc. Amer. Math. Soc. 88 (1983), 341-346.
- [13] Y. Dong, Sublinear impulse effects and solvability of boundary value problems for differential equations with impulses, J. Math. Anal. Appl. 264 (2001) 32-48.
- [14] P. Hartman, On boundary value problems for second order differential equations, J. Differential Equations 26 (1977) 37-53.
- [15] H. Jacobowitz, Periodic solutions of $x'' + f(x, t) = 0$ via the Poincaré-Birkhoff theorem, J. Differential Equations 20 (1976) 37-52.
- [16] H. Lamba, Chaotic, regular and unbounded behaviour in the elastic impact oscillator, Physica D, 82 (1995) 117-135.
- [17] V. Lakshmikantham, D. Bainov, P. Simeonov, Theory of Impulsive Differential Equations, World Scientific, Singapore, 1989.
- [18] Y. Long, Multiple solutions of perturbed superquadratic second order Hamiltonian systems, Trans. Amer. Math. Soc. 311 (1989) 749-780.
- [19] J. Mawhin, An extension of a theorem of A. C Lazer on forced nonlinear oscillations, J. Math. Anal. Appl. 40 (1972) 20-29.
- [20] J. J. Nieto, Basic theory for nonresonance impulsive periodic problems of first order, J. Math. Anal. Appl. 205 (1997) 423-433.
- [21] J. J. Nieto, Impulsive resonance periodic problems of first order, Appl. Math. Lett. 15 (2002) 480-493.

- [22] J. J. Nieto, D. O'Regan, Variational approach to impulsive differential equations, *Nonlinear Anal. Real World Appl.* 10 (2009) 680-690.
- [23] Y. M. Niu, X. Li, Periodic solutions of sublinear impulsive differential equations, Preprint.
- [24] M. del Pino, R. Manasevich, Infinitely many 2π -periodic solutions for a problem arising in nonlinear elasticity, *J. Differential Equations* 103 (1993) 260-277.
- [25] D. Qian, Infinity of subharmonics for asymmetric Duffing equations with the Lazer-Leach-Dancer condition, *J. Differential Equations* 171 (2001) 233-250.
- [26] D. Qian, L. Chen, X. Sun, Periodic solutions of superlinear impulsive differential equations: A geometric approach, *J. Differential Equations* 258 (2015) 3088-3106.
- [27] D. Qian, X. Li, Periodic solutions for ordinary differential equations with sublinear impulsive effects, *J. Math. Anal. Appl.* 303 (2005) 288-303.
- [28] D. Qian, P. J. Torres, Periodic motions of linear impact oscillators via the successor map, *SIAM J. Math. Anal.* 36 (2005) 1707-1725.
- [29] C. Rebelo, F. Zanolin, Multiplicity results for periodic solutions of second order ODEs with asymmetric nonlinearities, *Trans. Amer. Math. Soc.* 348 (1996) 2349-2389.
- [30] R. Reissing, Contractive mappings and periodically perturbed, non-conservative systems, *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* 58 (1975) 696-702.
- [31] J. Sun, H. Chen, J. J. Nieto, Infinitely many solutions for second-order Hamiltonian system with impulsive effects, *Math. Comput. Modelling* 54 (2011) 544-555.