# Deformations of infinite-dimensional Lie algebras, exotic cohomology, and integrable nonlinear partial differential equations

Oleg I. Morozov

Faculty of Applied Mathematics, AGH University of Science and Technology, Al. Mickiewicza 30, Cracow 30-059, Poland

## Abstract

The important unsolved problem in theory of integrable systems is to find conditions guaranteeing existence of a Lax representation for a given PDE. The exotic cohomology of the symmetry algebras opens a way to formulate such conditions in internal terms of the PDEs under the study. In this paper we consider certain examples of infinite-dimensional Lie algebras with nontrivial second exotic cohomology groups and show that the Maurer–Cartan forms of the associated extensions of these Lie algebras generate Lax representations for integrable systems, both known and new ones.

Keywords: exotic cohomology, Lie pseudo-groups, Maurer–Cartan forms, symmetries of differential equations, Lax representations
2000 MSC: 58H05, 58J70, 35A30, 37K05, 37K10
Subject Classification: integrable PDEs, symmetries of PDEs, cohomology of Lie algebras

# 1. Introduction

The existence of a Lax representation is the key property of integrable equations, [34, 7], and a starting setting for a number of techniques to study nonlinear partial differential equations (PDEs) such as Bäcklund transformations, nonlocal symmetries and conservation laws, recursion operators,

Preprint submitted to Journal of Geometry and Physics

Email address: morozov@agh.edu.pl (Oleg I. Morozov)

Darboux transformations, etc. Although these structures are of great significance in the theory of integrable PDEs, up to now the problem of finding conditions for a PDE to admit a Lax representation is open. In [26] we propose an approach for solving this problem in internal terms of the PDE under the study. We show there that for some PDEs their Lax representations can be derived from the second exotic<sup>1</sup> cohomology of the symmetry pseudogroups of the PDEs. The main advantage of this approach is that it allows one to get rid of apriori assumptions about the defining equations of the Lax representation. In this paper we generalize the constructions of [26]. We consider a deformation of the tensor product of the Lie algebra of vector fields on a line and the algebra of truncated polynomials as well as certain extensions of this deformation and show that at some values of the deformation parameter the Maurer–Cartan forms of the obtained Lie algebras produce Lax representations for some known as well as some new integrable systems.

#### 2. Preliminaries

All considerations in this paper are local. All functions are assumed to be real-analytic.

#### 2.1. Coverings of PDEs

The coherent geometric formulation of Lax representations, Wahlquist– Estabrook prolongation structures, Bäcklund transformations, recursion operators, nonlocal symmetries, and nonlocal conservation laws is based on the concept of differential covering of a PDE [14, 15]. In this subsection we closely follow [16, 17] to present the basic notions of the theory of differential coverings.

Let  $\pi: \mathbb{R}^n \times \mathbb{R}^m \to \mathbb{R}^n$ ,  $\pi: (x^1, \ldots, x^n, u^1, \ldots, u^m) \mapsto (x^1, \ldots, x^n)$  be a trivial bundle, and  $J^{\infty}(\pi)$  be the bundle of its jets of the infinite order. The local coordinates on  $J^{\infty}(\pi)$  are  $(x^i, u^{\alpha}, u_I^{\alpha})$ , where  $I = (i_1, \ldots, i_n)$  is a multiindex, and for every local section  $f: \mathbb{R}^n \to \mathbb{R}^n \times \mathbb{R}^m$  of  $\pi$  the corresponding infinite jet  $j_{\infty}(f)$  is a section  $j_{\infty}(f): \mathbb{R}^n \to J^{\infty}(\pi)$  such that  $u_I^{\alpha}(j_{\infty}(f)) = \frac{\partial^{\#I} f^{\alpha}}{\partial x^I} = \frac{\partial^{i_1 + \ldots + i_n} f^{\alpha}}{(\partial x^1)^{i_1} \ldots (\partial x^n)^{i_n}}$ . We put  $u^{\alpha} = u_{(0,\ldots,0)}^{\alpha}$ . Also, in the case of

<sup>&</sup>lt;sup>1</sup>Unlike in [26], in this paper we follow [27] and use the term "exotic cohomology" instead of "deformed cohomology", since here we discuss deformations of Lie algebras which are not related to "deformed cohomology" in the sense of [26].

m = 1 and, e.g., n = 4 we denote  $x^1 = t$ ,  $x^2 = x$ ,  $x^3 = y$ ,  $x^4 = z$ , and  $u_{(i,j,k,l)}^1 = u_{t\dots tx\dots xy\dots yz\dots z}$  with *i* times *t*, *j* times *x*, *k* times *y*, and *l* times *z*. The vector fields

$$D_{x^k} = \frac{\partial}{\partial x^k} + \sum_{\#I \ge 0} \sum_{\alpha=1}^m u_{I+1_k}^{\alpha} \frac{\partial}{\partial u_I^{\alpha}}, \qquad k \in \{1, \dots, n\},$$

with  $I + 1_k = (i_1, ..., i_k, ..., i_n) + 1_k = (i_1, ..., i_k + 1, ..., i_n)$  are referred to as total derivatives. They commute everywhere on  $J^{\infty}(\pi)$ :  $[D_{x^i}, D_{x^j}] = 0$ .

A system of PDEs  $F_r(x^i, u_I^{\alpha}) = 0, \#I \leq s, r \in \{1, \ldots, \sigma\}$ , of the order  $s \geq s$ 1 with  $\sigma \geq 1$  defines the submanifold  $\mathcal{E} = \{(x^i, u_I^{\alpha}) \in J^{\infty}(\pi) \mid D_K(F_r(x^i, u_I^{\alpha})) = 0\}$  $0, \ \#K \ge 0\}$  in  $J^{\infty}(\pi)$ .

Denote  $\mathcal{W} = \mathbb{R}^{\infty}$  with coordinates  $w^a, a \in \mathbb{N} \cup \{0\}$ . Locally, an (infinitedimensional) differential covering over  $\mathcal{E}$  is a trivial bundle  $\tau: J^{\infty}(\pi) \times \mathcal{W} \to$  $J^{\infty}(\pi)$  equipped with the extended total derivatives

$$\tilde{D}_{x^k} = D_{x^k} + \sum_{a=0}^{\infty} T_k^a(x^i, u_I^\alpha, w^b) \frac{\partial}{\partial w^a}$$
(1)

such that  $[\tilde{D}_{x^i}, \tilde{D}_{x^j}] = 0$  for all  $i \neq j$  whenever  $(x^i, u_I^{\alpha}) \in \mathcal{E}$ . For the partial derivatives of  $w^a$  which are defined as  $w^a_{x^k} = \tilde{D}_{x^k}(w^a)$  we have the system of covering equations

$$w_{x^k}^a = T_k^a(x^i, u_I^\alpha, w^b).$$

This over-determined system of PDEs is compatible whenever  $(x^i, u_I^{\alpha}) \in \mathcal{E}$ .

Dually the covering with extended total derivatives (1) is defined by the differential ideal generated by the Wahlquist-Estabrook forms, [7, p. 81],

$$\varpi^a = dw^a - \sum_{k=1}^n T_k^a(x^i, u_I^\alpha, w^b) \, dx^k.$$

This ideal is integrable on  $\mathcal{E}$ , that is,

$$d\varpi^a \equiv \sum_b \eta^a_b \wedge \varpi^b \mod \langle \vartheta_I \rangle,$$

where  $\eta_b^a$  are some 1-forms on  $\mathcal{E} \times \mathcal{W}$  and  $\vartheta_I = (du_I^\alpha - \sum_k u_{I+1_k}^\alpha dx^k)|_{\mathcal{E}}$ .

# 2.2. Exotic cohomology

Let  $\mathfrak{g}$  be a Lie algebra over  $\mathbb{R}$  and  $\rho: \mathfrak{g} \to \operatorname{End}(V)$  be its representation. Let  $C^k(\mathfrak{g}, V) = \operatorname{Hom}(\Lambda^k(\mathfrak{g}), V), \ k \ge 1$ , be the space of all k-linear skewsymmetric mappings from  $\mathfrak{g}$  to V. Then the Chevalley-Eilenberg differential complex

$$V = C^{0}(\mathfrak{g}, V) \stackrel{d}{\longrightarrow} C^{1}(\mathfrak{g}, V) \stackrel{d}{\longrightarrow} \dots \stackrel{d}{\longrightarrow} C^{k}(\mathfrak{g}, V) \stackrel{d}{\longrightarrow} C^{k+1}(\mathfrak{g}, V) \stackrel{d}{\longrightarrow} \dots$$

is is generated by the differential defined by the formula

$$d\theta(X_1, ..., X_{k+1}) = \sum_{q=1}^{k+1} (-1)^{q+1} \rho(X_q) \left(\theta(X_1, ..., \hat{X}_q, ..., X_{k+1})\right) + \sum_{1 \le p < q \le k+1} (-1)^{p+q} \theta([X_p, X_q], X_1, ..., \hat{X}_p, ..., \hat{X}_q, ..., X_{k+1}).$$
(2)

The cohomology groups of the complex  $(C^*(\mathfrak{g}, V), d)$  are referred to as the cohomology groups of the Lie algebra  $\mathfrak{g}$  with coefficients in the representation  $\rho$ . For the trivial representation  $\rho_0: \mathfrak{g} \to \mathbb{R}$ ,  $\rho_0: X \mapsto 0$ , the complex and its cohomology are denoted by  $C^*(\mathfrak{g})$  and  $H^*(\mathfrak{g})$ , respectively.

Consider a Lie algebra  $\mathfrak{g}$  over  $\mathbb{R}$  with non-trivial first cohomology group  $H^1(\mathfrak{g})$  and take a closed 1-form  $\alpha$  on  $\mathfrak{g}$ . Then for any  $\lambda \in \mathbb{R}$  define new differential  $d_{\lambda\alpha}: C^k(\mathfrak{g}, \mathbb{R}) \to C^{k+1}(\mathfrak{g}, \mathbb{R})$  by the formula

$$d_{\lambda\alpha}\theta = d\theta + \lambda\,\alpha \wedge \theta.$$

From  $d\alpha = 0$  it follows that

$$d_{\lambda\alpha}^2 = 0. \tag{3}$$

The cohomology groups of the complex

$$C^{1}(\mathfrak{g},\mathbb{R}) \xrightarrow{d_{\lambda\alpha}} \dots \xrightarrow{d_{\lambda\alpha}} C^{k}(\mathfrak{g},\mathbb{R}) \xrightarrow{d_{\lambda\alpha}} C^{k+1}(\mathfrak{g},\mathbb{R}) \xrightarrow{d_{\lambda\alpha}} \dots$$

are referred to as the *exotic cohomology groups* of  $\mathfrak{g}$  and denoted by  $H^*_{\lambda\alpha}(\mathfrak{g})$ .

REMARK 1. Cohomology  $H^*_{\lambda\alpha}(\mathfrak{g})$  coincides with cohomology of  $\mathfrak{g}$  with coefficients in the one-dimensional representation  $\rho_{\lambda\alpha}: \mathfrak{g} \to \mathbb{R}, \rho_{\lambda\alpha}: X \mapsto \lambda \alpha(X)$ . In particular, when  $\lambda = 0$ , cohomology  $H^*_{\lambda\alpha}(\mathfrak{g})$  coincides with  $H^*(\mathfrak{g})$ .  $\diamond$ REMARK 2. In all the cases considered in this paper  $H^1(\mathfrak{g}) = Z^1(\mathfrak{g})$  due to  $C^0(\mathfrak{g}) = \{0\}$  and  $B^1(\mathfrak{g}) = \{0\}$ , so a closed 1-form  $\alpha$  can be identified with its cohomology class.  $\diamond$ 

# 3. The Lax representation for the potential Khokhlov-Zabolotskaya equation through the second exotic cohomology of the symmetry pseudogroup

A relation between the exotic cohomology of symmetry pseudogroups and Lax representations for integrable systems was established in [26]. In a slightly different notation one of the results of that paper can be presented as follows.

Consider the potential Khokhlov–Zabolotskaya equation  $\mathcal{E}_1$  (or Lin–Reissner–Tsien equation), [24, 32],

$$u_{yy} = u_{tx} + u_x \, u_{xx}.\tag{4}$$

The infinite normal prolongation, [4, 5, 31, 30], of the structure equations for the pseudogroup  $Sym(\mathcal{E}_1)$  of contact symmetries of this equation has the form

$$d\alpha = 0, \tag{5}$$

$$d\Theta_0 = \nabla \Theta_0 \wedge \Theta_0, \tag{6}$$

$$d\Theta_1 = \alpha \wedge \Theta_1 + \nabla \Theta_1 \wedge \Theta_0 + \frac{2}{3} \nabla \Theta_0 \wedge \Theta_1, \tag{7}$$

$$d\Theta_2 = 2\alpha \wedge \Theta_2 + \nabla\Theta_2 \wedge \Theta_0 + \frac{2}{3}\nabla\Theta_1 \wedge \Theta_1 + \frac{1}{3}\nabla\Theta_0 \wedge \Theta_2, \tag{8}$$

$$d\Theta_3 = 3\alpha \wedge \Theta_3 + \nabla\Theta_3 \wedge \Theta_0 + \frac{2}{3}\nabla\Theta_2 \wedge \Theta_1 + \frac{1}{3}\nabla\Theta_1 \wedge \Theta_2, \tag{9}$$

$$d\Theta_4 = 4 \alpha \wedge \Theta_4 + \nabla \Theta_4 \wedge \Theta_0 + \frac{2}{3} \nabla \Theta_3 \wedge \Theta_1 + \frac{1}{3} \nabla \Theta_2 \wedge \Theta_2 -\frac{1}{3} \nabla \Theta_0 \wedge \Theta_4,$$
(10)

where

$$\Theta_m = \sum_{j=0}^{\infty} \frac{h^j}{j!} \theta_{m,j}, \qquad \nabla \Theta_m = \frac{\partial}{\partial h} \Theta_m = \sum_{j=0}^{\infty} \frac{h^j}{j!} \theta_{m,j+1}$$
(11)

for  $0 \le m \le 4$ , while dh = 0 and  $\theta_{3,0} = 0$ . We have

$$\begin{aligned} \alpha &= p^{-1} dp, \qquad \theta_{0,0} = q dt, \\ \theta_{1,0} &= p q^{2/3} (dy + a_1 dt), \\ \theta_{2,0} &= p^2 q^{1/3} (dx + \frac{2}{3} a_1 dy + a_2 dt), \\ \theta_{4,0} &= p^4 q^{-1/3} (du - u_t dt - u_x dx - u_y dy), \end{aligned}$$

where  $p, q, a_1, a_2 \in \mathbb{R}, p \neq 0, q \neq 0$ , are parameters.

REMARK 3. All the other forms  $\theta_{i,j}$  can be found inductively from the series of equations (5) – (10). For example, the first equation from the series of equations (6),  $d\theta_{0,0} = \theta_{0,1} \wedge \theta_{0,0}$ , has the form  $dq \wedge dt = \theta_{0,1} \wedge q \, dt$ , therefore we get  $\theta_{0,1} = q^{-1} \, dq + b_1 \, dt$  with  $b_1 \in \mathbb{R}$ . Now forms  $\theta_{0,0}$  and  $\theta_{0,1}$  in the second equation  $d\theta_{1,0} = \theta_{0,2} \wedge \theta_{0,0}$  from the series (6) are known, and we obtain  $\theta_{0,2} = db_1 + b_2 \, dt$ ,  $b_2 \in \mathbb{R}$ . Likewise, we can compute all the forms  $\theta_{0,j}$ . When  $\theta_{0,j}$  and  $\theta_{1,0}$  are known, from (7) it is possible to find  $\theta_{1,j}$ , etc.

Equations (5) – (10) imply  $H^1(Sym(\mathcal{E}_1)) = \mathbb{R} \alpha$ . The following theorem describes the structure of the second exotic cohomology group for  $Sym(\mathcal{E}_1)$ , for the proof of a more general result see Theorem 2 below.

THEOREM 1.

$$H^2_{\lambda\alpha}(Sym(\mathcal{E}_1)) = \begin{cases} \mathbb{R} \ [\Omega], & \lambda = -3, \\ \{0\}, & \lambda \neq -3, \end{cases}$$

where  $\Omega = \theta_{3,1} \wedge \theta_{0,0} + \frac{2}{3} \theta_{2,1} \wedge \theta_{1,0} + \frac{1}{3} \theta_{1,1} \wedge \theta_{2,0}$ .

COROLLARY. Equation

$$d\omega = 3\,\alpha \wedge \omega + \Omega \tag{12}$$

is compatible with the structure equations (5) – (10) of  $Sym(\mathcal{E}_1)$ .

REMARK 4. If we rename  $\omega = \theta_{3,0}$ , then (12) gets the form

$$d\theta_{3,0} = 3\alpha \wedge \theta_{3,0} + \theta_{3,1} \wedge \theta_{0,0} + \frac{2}{3}\theta_{2,1} \wedge \theta_{1,0} + \frac{1}{3}\theta_{1,1} \wedge \theta_{2,0}.$$
 (13)

Therefore, if we add  $\theta_{3,0}$  as the coefficient at  $h^0$  in the series for  $\Theta_3$  in (11), then (9) remains valid.

Since equation (13) is compatible with system (5)–(10), Lie's third inverse fundamental theorem in Cartan's form, [4, 5, 31, 30], ensures existence of a solution  $\theta_{3,0}$  to (13). We integrate this equation and put  $a_1 = v_x$ . This yields

$$\theta_{3,0} = p^3 \left( dv - v_x \, dx - \left( \frac{1}{3} \, v_x^3 - u_x \, v_x - u_y \right) \, dt - \left( \frac{1}{2} \, v_x^2 - u_x \right) \, dy \right).$$

This is the Wahlquist–Estabrook form of the Lax representation, [23, 9, 18, 35],

$$\begin{cases} v_t = \frac{1}{3}v_x^3 - u_x v_x - u_y, \\ v_y = \frac{1}{2}v_x^2 - u_x \end{cases}$$
(14)

for equation (4).

# 4. A deformation of the tensor product of the Lie algebra of vector fields on a line and the algebra of truncated polynomials

The structure equations (5)-(10), (13) can be written in the form

$$d\alpha = 0,$$
  

$$d\Theta_k = k \alpha \wedge \Theta_k + \sum_{m=0}^k \left(1 - \frac{1}{3}m\right) \nabla \Theta_{k-m} \wedge \Theta_m, \qquad 0 \le k \le 4.$$

Then the results of Section 3 admit the following generalization.

DEFINITION. For  $n \in \mathbb{N}$  and  $\varepsilon \in \mathbb{R}$  denote by  $\mathfrak{G}(n, \varepsilon)$  the Lie algebra with the structure equations

$$d\alpha = 0,$$
  

$$d\Theta_k = k \alpha \wedge \Theta_k + \sum_{m=0}^k (1 + \varepsilon m) \nabla \Theta_{k-m} \wedge \Theta_m, \quad 0 \le k \le n, \quad (15)$$

where equations (11) hold for each  $m \in \{0, 1, \ldots, n\}$ .

REMARK 5. The Lie algebra  $\mathfrak{G}(n,\varepsilon)$  has the following description. Consider the tensor product  $\mathbb{R}_{n+1}[s] \otimes C^{\omega}(\mathbb{R})$  of the algebra  $\mathbb{R}_{n+1}[s] = \mathbb{R}[s]/\langle s^{n+1} \rangle$ of truncated polynomials of degree less that n + 1 and the algebra of realanalytic functions of  $t \in \mathbb{R}$ . This is a vector space over  $\mathbb{R}$  generated by functions  $X_{k,m} = \frac{1}{m!} s^k t^m, k \in \{0, \ldots, n\}, m \in \mathbb{N} \cup \{0\}$ . This vector space is equipped with the Lie bracket

$$[f,g]_{\varepsilon} = fg_t - gf_t + \varepsilon s (f_s g_t - g_s f_t).$$
(16)

The vector field  $Y = s \partial_s$  is an outer derivative of the above Lie algebra. Then  $\mathfrak{G}(n, \varepsilon)$  is the semi-direct sum of  $\mathbb{R}_{n+1}[s] \otimes C^{\omega}(\mathbb{R})$  and one-dimensional Lie algebra generated by Y:

$$\mathfrak{G}(n,\varepsilon) = (\mathbb{R}_{n+1}[s] \otimes C^{\omega}(\mathbb{R})) \rtimes \mathbb{R} Y.$$

This Lie algebra is a deformation of  $\mathfrak{G}(n,0)$ . For  $\varepsilon = 0$  the bracket (16) is the Lie bracket in the algebra of analytic vector fields on a line.

Consider 1-forms  $\theta_{i,j}$ ,  $\alpha$  dual to  $X_{i,j}$ , Y, that is, 1-forms on  $\mathfrak{G}(n, \varepsilon)$  such that equations

$$\theta_{i,j}(X_{k,m}) = \delta_{ik}\,\delta_{jm}, \quad \theta_{i,j}(Y) = 0, \quad \alpha(X_{k,m}) = 0, \quad \alpha(Y) = 1 \tag{17}$$

hold for all  $i, j, k, m \in \mathbb{N} \cup \{0\}$ . Then (17), (2), and (11) imply the structure equations (15).

System (15) is the infinite normal prolongation of the system

$$d\alpha = 0, \tag{18}$$

$$d\theta_{k,0} = k \alpha \wedge \theta_{k,0} + \sum_{m=0}^{k} (1 + \varepsilon m) \ \theta_{k-m,1} \wedge \theta_{m,0}, \quad 0 \le k \le n.$$
(19)

This system is involutive, [4, §6], [28, Def. 11.7], therefore the third inverse fundamental Lie's theorem in Cartan's form implies existence of 1-forms  $\alpha$ ,  $\theta_{k,j}$ ,  $k \in \{0, \ldots, n\}$ ,  $j \in \{0, 1\}$ , that satisfy (19). Forms  $\alpha$ ,  $\theta_{0,0}$ ,  $\ldots$ ,  $\theta_{n,0}$ define a Lie pseudo-group on  $\mathbb{R}^{n+1}$ . While for the purposes of the present paper we need explicit expressions for the forms  $\theta_{k,0}$  and  $\theta_{k,1}$  only, all the other forms  $\theta_{k,j}$  can be found inductively by integration of equations (15). We need the whole system (15) to prove Theorem 2 below.

We note that system (19) has the following specific structure. For each fixed  $k \in \{0, \ldots, n\}$  the first k + 1 equations from (19) satisfy the conditions of the Frobenius theorem, see e.g. [12, Th. 1.3.4], for forms  $\theta_{0,0}, \ldots, \theta_{k,0}$ . We consider  $\alpha$  and  $\theta_{j,0}, \theta_{j,1}$  for  $j \in \{0, \ldots, k\}$  as 1-forms on  $\mathbb{R}^N$  for  $N \geq 3(n+1)+1$ . Eq. (18) implies that there is a function  $p \neq 0$  on  $\mathbb{R}$  such that<sup>2</sup>  $\alpha = dp/p$ . Then the Frobenius theorem implies existence of independent functions (coordinates)  $x_0, \ldots, x_k$  on  $\mathbb{R}^N$  such that the differential ideal of forms  $\theta_{0,0}, \ldots, \theta_{k,0}$  is generated algebraically by forms  $dx_0, \ldots, dx_k$ , that

is, 
$$\theta_{0,0} = A_0 \, dx_0$$
 and  $\theta_{i,0} = A_i \left( dx_i + \sum_{j=0}^{i-1} B_{ij} dx_j \right), \ i \in \{1, \dots, k\}$ , for some

functions  $A_j \neq 0$ ,  $B_{ij}$  on  $\mathbb{R}^N$ . Substituting these forms into (19) yields a triangular system for 1-forms  $\theta_{i,1}$ , so it is easy to find them.

EXAMPLE 1. For k = 2 from first three equations of (19) we have

$$\begin{aligned} \theta_{0,0} &= a_0 \, dx_0, \\ \theta_{0,1} &= p \, a_0^{1+\varepsilon} \, (dx_1 + a_1 \, dx_0), \\ \theta_{0,2} &= p^2 \, a_0^{1+2\varepsilon} \, (dx_2 + (1+\varepsilon) \, a_1 \, dx_1 + a_2 \, dx_0), \\ \theta_{0,1} &= da_0/a_0 + b_0 \, dx_0, \end{aligned}$$

 $<sup>^2 {\</sup>rm Here}$  and below we put  $\alpha = dp/p$  instead of the natural choice  $\alpha = dp$  to simplify the further computations

$$\begin{aligned} \theta_{1,1} &= p \, a_0^{\varepsilon} \, (da_1 + (1 + \varepsilon) \, b_0 \, dx_1 + b_1 \, dx_0), \\ \theta_{2,1} &= p^2 \, a_0^{2\varepsilon} \, (da_2 - (1 + \varepsilon) \, a_1 da_1 + (1 + 2 \, \varepsilon) \, b_0 dx_2 \\ &+ (1 + \varepsilon) \, (b_1 + \varepsilon \, a_1 b_0) \, dx_1 + b_2 \, dx_2) \end{aligned}$$

with  $a_0 \neq 0$ . Then substituting for  $\theta_{3,0} = A_3 (dx_3 + B_{30}dx_0 + B_{31}dx_1 + B_{32}dx_2)$ into the equation for  $d\theta_{3,0}$  from (19) allows one to define coefficients  $A_3$ ,  $B_{3j}$ . After this, the equation for  $d\theta_{3,0}$  defines  $\theta_{3,1}$  up to adding the form  $b_3 dx_0$ with the new coordinate  $b_3$  on  $\mathbb{R}^N$ .

The Lie algebra  $\mathfrak{G}(n,\varepsilon)$  has the following important feature: when  $\varepsilon = -\frac{1}{r}$  for  $r \in \mathbb{N}$  and  $n \geq r$ , 1-form  $\theta_{r,0}$  has only two entries in the whole system (15). Indeed, the first equation from the series for  $d\Theta_r$  acquires the form

$$d\theta_{r,0} = r \,\alpha \wedge \theta_{r,0} + \Psi_r,\tag{20}$$

while neither 2-form  $\Psi_r$  nor the other equations from (15) include  $\theta_{r,0}$ . Therefore in order to generalize results of Section 3 we consider the Lie algebra  $\mathfrak{G}(r+1, -1/r)$  and proceed in the following steps. First, we find forms  $\theta_{i,0}$  and  $\theta_{i,1}$  for  $0 \leq i \leq r-1$  from equations (15) in the similar way as it was described above. Second, we assume that  $\theta_{r+1,0} = A_{r+1} (du - u_{x_0} dx_0 - \ldots - u_{x_{r-1}} dx_{r-1})$ ,  $A_{r+1} \neq 0$ , that is, we consider  $\theta_{r+1,0}$  as the zeroth order contact form on the jet bundle  $J^{\infty}(\pi)$  for the bundle  $\pi: (x_0, x_1, \ldots, x_{r-1}, u) \mapsto (x_0, x_1, \ldots, x_{r-1})$ . This implies some expressions for the parameters  $B_{i,j}$  in terms of  $u_{x_0}, \ldots$ ,  $u_{x_{r-1}}$ . Third, we integrate equation (20) to find  $\theta_{r,0} = b_r (dv - T_0 dx_0 - \ldots - T_{r-1} dx_{r-1})$  and then consider the system of PDEs  $v_{x_i} = T_i, 0 \leq i \leq r-1$ , generated by equation  $\theta_{r,0} = 0$ .

# 5. Lie algebras $\mathfrak{G}(r+1, -1/r)$ , their extensions, and associated integrable systems

5.1. Integrable systems generated by  $\mathfrak{G}(r+1, -1/r)$ 

In this subsection we present some results of computations discussed in the end of Section 4.

EXAMPLE 2. For  $\mathfrak{G}(5, -1/4)$  we have<sup>3</sup>

 $\theta_{0,0} = \frac{1}{4} q^4 dt,$ 

<sup>&</sup>lt;sup>3</sup>Here and below we rescale parameters in forms  $\theta_{j,0}$  in order to simplify the resulting PDE and its Lax representation.

$$\begin{aligned} \theta_{1,0} &= \frac{1}{3} p \, q^3 \, (dy + a \, dt), \\ \theta_{2,0} &= \frac{1}{2} p^2 \, q^2 \, (dx + 2 \, a \, dy + (3 \, a^2 - 2 \, u_z) \, dt), \\ \theta_{3,0} &= p^3 \, q \, (dz + a \, dx + (a^2 - u_z) \, dy + (a^3 - 2 \, a \, u_z - u_x) \, dt), \\ \theta_{5,0} &= p^5 \, q^{-1} \, (du - u_t \, dt - u_x \, dx - u_y \, dy - u_z \, dz), \end{aligned}$$

Then we integrate (20) with r = 4 and put  $a = v_z$ . This gives

$$\theta_{4,0} = -p^4 \left( dv - \left( \frac{1}{4} v_z^4 - u_z v_z^2 - u_x v_z - u_y + \frac{1}{2} u_z^2 \right) dt - \left( \frac{1}{2} v_z^2 - u_z \right) dx - \left( \frac{1}{3} v_z^3 - u_z v_z - u_x \right) dy - v_z dz \right)$$

Then equation  $\theta_{4,0} = 0$  yields the system

$$\begin{cases} v_x = \frac{1}{2}v_z^2 - u_z, \\ v_y = \frac{1}{3}v_z^3 - u_z v_z - u_x, \\ v_t = \frac{1}{4}v_z^4 - u_z v_z^2 - u_x v_z - u_y + \frac{1}{2}u_z^2. \end{cases}$$

This system is compatible whenever equations

$$\begin{cases}
 u_{xx} = u_{yz} + u_z u_{zz}, \\
 u_{xy} = u_{tz} + u_z u_{xz} + u_x u_{zz}, \\
 u_{yy} = u_{tx} + u_x u_{xz} + u_z^2 u_{zz}
\end{cases}$$
(21)

hold. System (21) is the second system from the dKP hierarchy, [33, 13, 6, 18, 22, 35, 3, 10].  $\diamond$ 

EXAMPLE 3. The Lie algebra  $\mathfrak{G}(6, -1/5)$  provides the third system from the dKP hierarchy. In this case integration of (15) gives

$$\begin{aligned} \theta_{0,0} &= \frac{1}{5} q^5 dt, \\ \theta_{1,0} &= p q^4 \left( \frac{1}{4} dy + a dt \right), \\ \theta_{2,0} &= p^2 q^3 \left( \frac{1}{3} dx + a dy + (2 a^2 - u_s) dt \right), \\ \theta_{3,0} &= p^3 q^2 \left( \frac{1}{2} dz + a dx + (3 a^2 - 2 u_s) dy + 2 (2 a^3 - 3 a u_s - u_z) dt \right), \\ \theta_{4,0} &= p^4 q \left( ds + a dz + (a^2 - u_s) dx + (a^3 - 2 u_s a - u_z) dy \right) \end{aligned}$$

$$+(a^4 - 3 u_s a^2 - 2 u_z a - u_x + u_s^2) dt),$$
  
$$\theta_{6,0} = p^6 q^{-1} (du - u_t dt - u_x dx - u_y dy - u_z dz - u_s ds).$$

Then from (20) we get

$$\theta_{5,0} = -p^5 \left( dv - \left(\frac{1}{5} v_s^5 - u_s v_s^3 - u_z v_s^2 + (u_s^2 - u_x) v_s + u_z u_s - u_y \right) dt - \left(\frac{1}{3} v_s^3 - u_s v_s - u_z\right) dx - \left(\frac{1}{4} v_s^4 - u_s v_s^2 - u_z v_s - u_x + \frac{1}{2} u_s^2\right) dy - \left(\frac{1}{2} v_s^2 - u_s\right) dz - v_s ds \right)$$

with  $a = v_s$ . This form generates the system

$$\begin{cases} v_z &= \frac{1}{2} v_s^2 - u_s, \\ v_x &= \frac{1}{3} v_s^3 - u_s v_s - u_z, \\ v_y &= \frac{1}{4} v_s^4 - u_s v_s^2 - u_z v_s - u_x + \frac{1}{2} u_s^2, \\ v_t &= \frac{1}{5} v_s^5 - u_s v_s^3 - u_z v_s^2 + (u_s^2 - u_x) v_s + u_z u_s - u_y. \end{cases}$$

This is the Lax representation for the third system from the dKP hierarchy

$$\begin{cases} u_{zz} = u_{xs} + u_s u_{ss}, \\ u_{xz} = u_{ys} + u_s u_{zs} + u_z u_{ss}, \\ u_{yz} = u_{ts} + u_s u_{xs} + u_z u_{zs} + u_x u_{ss}, \\ u_{xx} = u_{yz} + u_z u_{zs} + u_s^2 u_{ss}, \\ u_{xy} = u_{tz} + u_z u_{xs} + (u_x + u_s^2) u_{zs} + 2 u_z u_s u_{ss}, \\ u_{yy} = u_{tx} + u_x u_{xs} + 2 u_z u_s u_{zs} + (u_z^2 + u_s^3) u_{ss}. \end{cases}$$

 $\diamond$ 

5.2. Right extensions of  $\mathfrak{G}(n, -1/r)$ 

While it is natural to expect that Maurer-Cartan forms of  $\mathfrak{G}(r+1, -1/r)$  with r > 5 produce higher elements of the dKP hierarchy, an interesting question is to consider the case r = 2. But the integration scheme from the end of Section 4 does not give any Lax representation of a PDE in the case

of  $\mathfrak{G}(3, -1/2)$ . Therefore we consider an extension of  $\mathfrak{G}(n, \varepsilon)$ . For  $n \ge r \ge 1$  we denote by  $\mathfrak{H}(n, r)$  the Lie algebra with the structure equations

$$d\alpha = 0,$$
  

$$d\beta = r \alpha \wedge \beta,$$
  

$$d\Theta_{k'} = k' \alpha \wedge \Theta_{k'} + \sum_{m=0}^{k'} \left(1 - \frac{m}{r}\right) \nabla \Theta_{k'-m} \wedge \Theta_m,$$
  

$$d\Theta_{k''} = k'' \alpha \wedge \Theta_{k''} + \sum_{m=0}^{k''} \left(1 - \frac{m}{r}\right) \nabla \Theta_{k''-m} \wedge \Theta_m + \beta \wedge \nabla \Theta_{k''-r}, \quad (22)$$

where  $k' \in \{0, \ldots, r-1\}$  and  $k'' \in \{r, \ldots, n\}$ . Then  $\mathfrak{H}(n, r)$  is a onedimensional right extension, [11, § 1.4.4], of  $\mathfrak{G}(n, -1/r)$ . Indeed, from the structure equations (22) it follows that the dual element associated to the new 1-form  $\beta$  is a derivative  $Z: \mathfrak{G}(n, -1/r) \to \mathfrak{G}(n, -1/r)$  such that  $Z: X_{k,m} \mapsto X_{k+r,m-1}$  for  $m \geq 1$  and  $k+r \leq n$ ,  $Z: X_{k,m} \mapsto 0$  for m = 0 or k+r > n, and Z(Y(f)) - Y(Z(f)) = -r Y(f) for every  $f \in \mathfrak{G}(n, -1/r)$ , while there is no function  $g \in \mathbb{R}_{n+1}[s] \otimes C^{\omega}(\mathbb{R})$  such that  $[g, X_{k,m}]_{-1/r} = X_{k+r,m-1}$  for all  $m \geq 1$  and  $k+r \leq n$ .

EXAMPLE 4. Consider the Lie algebra  $\mathfrak{H}(3,2)$ . We have

$$H^2_{\lambda\alpha}(\mathfrak{H}(3,2)) = \begin{cases} \mathbb{R} [\alpha \land \beta], & \lambda = -2, \\ \{0\}, & \lambda \neq -2. \end{cases}$$

This assertion can be proven similarly to Theorem 2 below. Integration yields

$$\begin{split} \beta &= p^2 \, dx, \\ \theta_{0,0} &= -\frac{1}{2} \, q^2 \, dt, \\ \theta_{1,0} &= p \, q \, \left( dy - u_x \, dt \right), \\ \theta_{3,0} &= p^3 \, q^{-1} \left( du - u_t \, dt - u_x \, dx - u_y \, dy \right). \end{split}$$

Then instead of equation  $d\theta_{2,0} = 2 \alpha \wedge \theta_{2,0} + \Phi$  with  $\Phi = \theta_{2,1} \wedge \theta_{0,0} + \frac{1}{2} \theta_{1,1} \wedge \theta_{1,0} + \beta \wedge \theta_{0,1}$  from (22) we consider equation

$$d\hat{\theta}_{2,0} = 2\,\alpha \wedge \hat{\theta}_{2,0} + \Phi + 2\,\alpha \wedge \beta.$$

We rename  $q = p \exp(v_x)$  and get

$$\tilde{\theta}_{2,0} = p^2 \left( dv + \left( u_y - \frac{1}{2} u_x^2 \right) dt - v_x \, dx + u_x \, dy \right)$$

This form generates the system

$$\begin{cases} v_t = \frac{1}{2}u_x^2 - u_y, \\ v_y = -u_x. \end{cases}$$
(23)

This system is compatible whenever the equation [8, 1, 29]

$$u_{yy} = u_{tx} + u_x \, u_{xy}. \tag{24}$$

holds. Thus system (23) defines an Abelian covering, [2], or a conservation law, for equation (24). To obtain non-Abelian covering for (24) we consider the form

$$\omega = \tilde{\theta}_{2,0} + \theta_{0,0} - \theta_{1,0} = p^2 \left( dv + \left( u_y - \frac{1}{2} u_x^2 - \frac{1}{2} e^{2v_x} + u_x e^{v_x} \right) dt - v_x dx + \left( u_x - e^{v_x} \right) dy \right).$$
(25)

This form is a solution to the equation  $d\omega = 2\alpha \wedge \omega + \Psi$ , where  $\Psi = \Phi + 2\alpha \wedge \beta + d_{-2\alpha}(\theta_{0,0} - \theta_{1,0})$  is  $d_{-2\alpha}$ -cohomologous to the cocycle  $\Phi + 2\alpha \wedge \beta$ . Form (25) generates the Lax representation

$$\begin{cases} v_t = \frac{1}{2} (e^{v_x} - u_x)^2 - u_y, \\ v_y = e^{v_x} - u_x \end{cases}$$

for (24). This system was derived in [29].

 $\diamond$ 

EXAMPLE 5. In the same way, for the Lie algebra  $\mathfrak{H}(4,3)$  we have

$$H^2_{\lambda\alpha}(\mathfrak{H}(4,3)) = \begin{cases} \mathbb{R}\left[\alpha \land \beta\right], & \lambda = -3, \\ \{0\}, & \lambda \neq -3, \end{cases}$$

and

$$\beta = p^{3} dx,$$
  

$$\theta_{0,0} = \frac{1}{3} q^{3} dt,$$
  

$$\theta_{1,0} = -\frac{1}{2} p q^{2} (dy - 2 u_{x} dt),$$
  

$$\theta_{2,0} = p^{2} q (dz - u_{x} dy - (u_{z} - u_{x}^{2}) dt),$$

$$\theta_{4,0} = p^4 q^{-1} (du - u_t dt - u_x dx - u_y dy - u_z dz).$$

Instead of equation for  $d\theta_{3,0}$  from (22) we consider equation

$$d\tilde{\theta}_{3,0} = 3\alpha \wedge \tilde{\theta}_{3,0} + \theta_{3,1} \wedge \theta_{0,0} + \frac{2}{3}\theta_{2,1} \wedge \theta_{1,0} + \frac{1}{3}\theta_{1,1} \wedge \theta_{2,0} + \beta \wedge \theta_{0,1} + 3\alpha \wedge \beta.$$

Then for  $q = p \exp(v_x)$  we get the Wahlquist-Estabrook form

$$\tilde{\theta}_{3,0} - \theta_{0,0} + \theta_{1,0} - \theta_{2,0} = p^3 \left( dv - \left(\frac{1}{3} \left( e^{v_x} - u_x \right)^3 - u_z \left( e^{v_x} - u_x \right) - u_y \right) dt - v_x \, dx - \left(\frac{1}{2} \left( e^{v_x} - u_x \right)^2 - u_z \right) \, dy - \left( e^{v_x} - u_x \right) \, dz \right)$$

for the Lax representation

$$\begin{cases} v_t = \frac{1}{3} (e^{v_x} - u_x)^3 - u_z (e^{v_x} - u_x) - u_y, \\ v_y = \frac{1}{2} (e^{v_x} - u_x)^2 - u_z, \\ v_z = e^{v_x} - u_x \end{cases}$$

of the system

$$\begin{cases} u_{yy} = u_{tz} + u_z u_{zz}, \\ u_{zz} = u_{xy} + u_x u_{xz}, \\ u_{yz} = u_{tx} + u_x u_{xy} + u_z u_{xz} \end{cases}$$

The first and the second equations of this system differ from the potential Khokhlov–Zabolotskaya equation (4) and equation (24), respectively, by notation.  $\diamond$ 

We suppose that for algebras  $\mathfrak{H}(r+1, r)$  the same computations will give higher elements of the integrable hierarchy generated by equation (24).

# 6. Second exotic cohomology group of $\mathfrak{G}(n,\varepsilon)$

To obtain further generalizations of the constructions from Section 5 we study the second exotic cohomology group of the Lie algebra  $\mathfrak{G}(n, \varepsilon)$ .

THEOREM 2. Suppose  $n \ge 2$ , then for each fixed  $r \in \{2, \ldots, n\}$ 

$$H^2_{-r\,\alpha}(\mathfrak{G}(n,-2/r)) = \mathbb{R}\,[\Phi_r],$$

where

$$\Phi_r = \sum_{m=0}^{[r/2]} (r - 2m) \ \theta_{r-m,0} \wedge \theta_{m,0}.$$

For other values of  $\lambda$  and  $\varepsilon$  we have

$$H^2_{\lambda\alpha}(\mathfrak{G}(n,\varepsilon)) = \{0\}.$$

Proof. For forms  $\eta, \omega_1, \ldots, \omega_k$  such that deg  $\eta \ge \deg \omega_i$  denote by  $\eta|_{\omega_1=0,\ldots,\omega_k=0}$  the result of replacement of each entry of  $\omega_1, \ldots, \omega_k$  in  $\eta$  by 0. The condition  $\eta = \eta|_{\omega_1=0,\ldots,\omega_k=0}$  will be denoted as  $\eta = o(\omega_1,\ldots,\omega_k)$ .

From (15) it follows that for each  $k \ge 0, m \ge 0$ 

$$d\theta_{k,m} = (k \alpha + (k \varepsilon + 1 - m) \theta_{0,1}) \wedge \theta_{k,m} + o(\alpha, \theta_{0,1}),$$

hence for  $U = -\varepsilon Y + X_{0,1} \in \mathfrak{G}(n,\varepsilon)$  we have

$$[U, X_{k,m}] = (m-1) X_{k,m}.$$

Therefore U defines an inner grading in  $\mathfrak{G}(n,\varepsilon)$ , [11, § 1.5.2], that is,

$$\mathfrak{G}(n,\varepsilon) = \bigoplus_{j=-1}^{\infty} \mathfrak{g}_j, \qquad [\mathfrak{g}_i,\mathfrak{g}_j] \subseteq \mathfrak{g}_{i+j},$$

where  $\mathfrak{g}_j = \{X \in \mathfrak{G}(n,\varepsilon) \mid [U,X] = jX\}$  for  $j \geq -1$  and  $\mathfrak{g}_j = \{0\}$  for j < -1. We have  $\mathfrak{g}_0 = \langle Y, X_{0,1} \rangle$  and  $\mathfrak{g}_j = \langle X_{k,j+1} \mid 0 \leq k \leq n \rangle$  for  $j \in \mathbb{N} \cup \{-1\}$ , so dim  $\mathfrak{g}_j \leq n$ . From Remark 1 it follows that  $H^*_{\lambda\alpha}(\mathfrak{G}(n,\varepsilon))$  is the cohomology of  $\mathfrak{G}(n,\varepsilon)$  with coefficients in the module  $A = \bigcup_{\mu \in \mathbb{R}} A_{\mu}$ , where

$$A_{\mu} = \{a \in \mathbb{R} \mid \lambda \alpha(U) \cdot a = \mu \cdot a\} = \{a \in \mathbb{R} \mid -\varepsilon \lambda a = \mu a\} = \begin{cases} \mathbb{R}, & \mu = \varepsilon \lambda, \\ \{0\}, & \mu \neq \varepsilon \lambda. \end{cases}$$

For  $m \geq 0$  and  $\nu \in \mathbb{R}$  denote

 $C^m_{(\nu)}(\mathfrak{G}(n,\varepsilon),A) = \{ \omega \in C^m(\mathfrak{G}(n,\varepsilon),A) \mid \omega(W_1,\ldots,W_m) \in A_{j_1+\ldots+j_m-\nu} \text{ for } W_k \in \mathfrak{g}_{j_k} \}.$ In particular,

$$C^{2}_{(0)}(\mathfrak{G}(n,\varepsilon),A) = \bigoplus_{\substack{i+j = -\varepsilon\lambda, \\ i \ge -1, j \ge -1}} \operatorname{Hom}(\mathfrak{g}_{i} \wedge \mathfrak{g}_{j},\mathbb{R}).$$
(26)

In accordance with [11, Th. 1.5.2a] the inclusion  $C^*_{(0)}(\mathfrak{G}(n,\varepsilon),A) \to C^*(\mathfrak{G}(n,\varepsilon),A)$ induces a cohomological isomorphism. Hence, from (26) it follows that  $H^2_{\lambda\alpha}(\mathfrak{G}(n,\varepsilon))$  is finite-dimensional for each fixed pair  $\lambda, \varepsilon$ , and for an arbitrary solution  $\Omega$  of equation  $d\Omega + \lambda \alpha \wedge \Omega = 0$  we can assume without loss of generality that  $M = \min \{j \in \mathbb{N} \cup \{0\} \mid \Omega = o(\theta_{0,j+1}, \theta_{1,j+1}, \dots, \theta_{n,j+1})\} < \infty$ . Then  $\Omega = \sum_{j=0}^{n} \gamma_j \wedge \theta_{j,M}$  with  $\gamma_j = o(\theta_{j,M}, \theta_{j+1,M}, \dots, \theta_{n,M})$ . Suppose  $M \ge 1$ .

From the structure equations (15) it follows that

$$d\theta_{k,M} = \sum_{m=0}^{k} (1+\varepsilon m) \,\theta_{k-m,M+1} \wedge \theta_{m,0} + o(\theta_{0,M+1}, \theta_{1,M+1}, \dots, \theta_{n,M+1}).$$

Hence  $d\Omega + \lambda \alpha \wedge \Omega = \gamma_n \wedge \theta_{n,M+1} \wedge \theta_{0,0} + o(\theta_{n,M+1}) = 0$  yields  $\gamma_n = c \theta_{0,0}$ with  $c \in \mathbb{R}$ . Thus  $\Omega - d_{\lambda \alpha}(c \theta_{n,M-1}) = o(\theta_{n,M})$ . In other words, we can put  $\gamma_n = 0$  without loss of generality. In the same way we can consequently put  $\gamma_{n-1} = 0$ , then  $\gamma_{n-2} = 0$ , etc.,  $\gamma_0 = 0$ . So we have M = 0 and

$$\Omega = \sum_{0 \le i < j \le n} a_{ij} \,\theta_{i,0} \wedge \theta_{j,0} + \sum_{j=0}^n b_j \,\alpha \wedge \theta_{j,0}$$

with  $a_{ij}, b_j \in \mathbb{R}$ .

Then we have  $d\Omega + \lambda \alpha \wedge \Omega = -b_j \alpha \wedge \theta_{j,1} \wedge \theta_{0,0} + o(\alpha \wedge \theta_{j,1} \wedge \theta_{0,0})$  for each  $j \in \{0, 1, \ldots, n\}$ , so  $b_0 = b_1 = \ldots = b_n = 0$ . We assume that  $a_{ij} \neq 0$  for some *i*, *j*, then equation  $d\Omega + \lambda \alpha \wedge \Omega = a_{ij} (i + j + \lambda) \alpha \wedge \theta_{i,0} \wedge \theta_{j,0} + o(\theta_{i,0} \wedge \theta_{j,0})$ yields i + j = r, where  $r \in \mathbb{N}$  is a constant, and  $\lambda = -r$ . Therefore,  $\Omega = \sum_{j=0}^{\lfloor r/2 \rfloor} c_j \theta_{j,0} \wedge \theta_{r-j,0} \text{ for some } c_j \in \mathbb{R}.$ 

We claim that  $r \leq n$ . Indeed, assume r > n and put  $r_0 = \min \{j \in$  $\mathbb{N} \mid \Omega = o(\theta_{i+1,0})$ . Since  $r_0 \leq n$ , we have  $r - r_0 \geq 1$ . Then from  $\Omega = 0$  $c_{r-r_0} \theta_{r-r_0,0} \wedge \theta_{r_0,0} + o(\theta_{r_0,0})$  we have  $c_{r-r_0} \neq 0$ . But  $d\Omega - r \alpha \wedge \Omega = -c_{r-r_0} \theta_{r-r_0,0} \wedge \theta_{r-r_0,0}$  $\theta_{r_0,1} \wedge \theta_{0,0} + o(\theta_{r_0,1})$  implies  $c_{r-r_0} = 0$ . The contradiction proves the claim.

Consider the case of even r and put r = 2m for  $m \in \mathbb{N}$ . Then  $\Omega =$  $\sum_{i=0}^{m-1} c_j \theta_{j,0} \wedge \theta_{2m-j,0} \text{ and }$ 

$$d\Omega - r \alpha \wedge \Omega = \sum_{j=1}^{m-1} (c_j - c_0 (1 + \varepsilon j)) \theta_{0,0} \wedge \theta_{r-j,1} \wedge \theta_{j,0} + o(\theta_{0,0} \wedge \theta_{m+1,1}, \theta_{0,0} \wedge \theta_{m+2,1}, \dots, \theta_{0,0} \wedge \theta_{2m,1})$$

So we get  $c_j = c_0 (1 + \varepsilon j)$  and thus

$$\Omega = c_0 \sum_{j=0}^{m-1} (1 + \varepsilon j) \,\theta_{j,0} \wedge \theta_{r-j,0}.$$
(27)

Finally we have

$$d\Omega - r \, \alpha \wedge \Omega = (1 + \varepsilon \, m) \, \left( d\theta_m \wedge \theta_m + 2 \, \sum_{j=0}^{m-1} \, d\theta_{j,0} \wedge \theta_{2m-j,0} \right),$$

and hence  $\varepsilon = -\frac{1}{m} = -\frac{2}{r}$ . Substituting this into (27) with  $c_0 = -r$  yields  $\Omega = \Phi_r$ .

In the case r = 2m + 1 the proof is similar.

COROLLARY. For each  $r \in \{2, \ldots, n\}$  the equation

$$d\omega = r \,\alpha \wedge \omega + \Phi_r. \tag{28}$$

is compatible with the structure equations of  $\mathfrak{G}(n, -2/r)$ .

EXAMPLE 6. In the case of n = r = 3 integration of the structure equations of  $\mathfrak{G}(3, -\frac{2}{3})$  gives

$$\begin{aligned} \theta_{0,0} &= -\frac{1}{3} q^3 dt, \\ \theta_{1,0} &= -p q \left( dy + u_x dt \right), \\ \theta_{2,0} &= p^2 q^{-1} \left( dx - u_x dy - (u_y + u_x^2) dt \right), \\ \theta_{3,0} &= p^3 q^{-2} \left( du - u_t dt - u_x dx - u_y dy \right), \end{aligned}$$

while equation (28) acquires the form

$$d\omega = 3\alpha \wedge \omega + 3\theta_{3,0} \wedge \theta_{0,0} + \theta_{2,0} \wedge \theta_{1,0}.$$

We integrate this to obtain the form  $\omega$ , then rename  $q = p^{-1} v_x^{-1}$  and consider

$$\omega + \theta_{2,0} = p^3 \left( dv - \left( u - \left( u_x^2 + u_y \right) v_x \right) dt - v_x \, dx - \left( x - u_x \, v_x \right) dy \right).$$

This form defines the Lax representation

$$\begin{cases} v_t = u - (u_x^2 + u_y) v_x, \\ v_y = x - u_x v_x \end{cases}$$

for the equation

$$u_{yy} = u_{tx} + (u_y - u_x^2) \, u_{xx} - 3 \, u_x \, u_{xy}.$$

This Lax representation was found in [25].

EXAMPLE 7. In the case n = r = 4 instead of  $\mathfrak{G}(4, -1/2)$  we consider the Lie algebra  $\mathfrak{H}(4, 2)$ . We have

$$H^2_{\lambda\alpha}(\mathfrak{H}(4,2)) = \begin{cases} \mathbb{R} [\alpha \wedge \beta], & \lambda = -2, \\ \mathbb{R} [\tilde{\Phi}_4], & \lambda = -4, \\ \{0\}, & \lambda \notin \{-2, -4\}, \end{cases}$$

where

$$\tilde{\Phi}_4 = \frac{1}{2} \Phi_4 + \beta \wedge \theta_{2,0} = 2 \theta_{4,0} \wedge \theta_{0,0} + \theta_{3,0} \wedge \theta_{1,0} + \beta \wedge \theta_{2,0}.$$

Consider the case  $\lambda = -4$ . From the structure equations of  $\mathfrak{H}(4,2)$  we obtain

$$\begin{split} \beta &= \frac{1}{2} p^2 \, dx, \\ \theta_{0,0} &= -\frac{1}{2} \, q^2 \, dt, \\ \theta_{1,0} &= p \, q \, \left( dy - u_z \, dt \right), \\ \theta_{2,0} &= p^2 \, \left( dw - \left( u_x - u_z^2 \right) dt - \ln |q| \, dx + u_z \, dy \right), \\ \theta_{3,0} &= p^3 \, q^{-1} \, \left( dz + \left( u_y - u_x u_z - \frac{1}{2} \, u_z^3 \right) \, dt - u_z \, dx + \left( u_x + \frac{1}{2} \, u_z \right) \, dy \right), \\ \theta_{4,0} &= p^4 \, q^{-2} \left( du - u_t \, dt - u_x \, dx - u_y \, dy - u_z \, dz \right). \end{split}$$

Then we integrate the equation

$$d\omega = 4\,\alpha \wedge \omega + \tilde{\Phi}_4$$

and get

$$\omega = p^4 \left( dv - u \, dt + z \, dy - w \, dx \right).$$

After the change of notation  $q = p^{-1} v_z^{-1}$ ,  $w = v_x + u_z v_z$  we have

$$\omega - \theta_{3,0} = p^4 \left( dv - \left( u + \left( u_y - u_x \, u_z - \frac{1}{2} \, u_z^3 \right) \, v_z \right) \, dt - v_x \, dx$$

$$-\left(\left(u_x+\frac{1}{2}u_z\right) v_z-z\right) dy-v_z dz\right).$$

This is the Wahlquist–Estabrook form for the Lax representation

$$\begin{cases} v_t = u + (u_y - u_x u_z - \frac{1}{2} u_z^3) v_z \\ v_y = z - (u_x + \frac{1}{2} u_z) v_z. \end{cases}$$

of the equation

$$u_{yy} = u_{tx} + u_z \, u_{xy} - u_y \, u_{xz} + u_z \, u_{tz} + 2 \, (u_x + u_z^2) \, u_{yz}$$
$$- \left(u_x^2 + u_x \, u_z^2 + u_y \, u_z + \frac{1}{4} \, u_z^4\right) \, u_{zz}.$$

# 7. Concluding remarks

In the present paper we have shown that both known (Examples 2, 3, 4, and 6) and new (Examples 5 and 7) Lax representations for integrable PDEs can be derived from the second exotic cohomology group of symmetry pseudogroups of the PDEs under the study. This approach gives the solution to the problem of existence of a Lax representation in internal terms of the PDE and allows one to eliminate apriori assumptions about the possible form of the Lax representation. The approach is universal and can be used to analyze a lot of equations or Lie algebras with nontrivial second exotic cohomology group. Quite natural this leads to the question of generalization of the Lie algebras considered above. In particular, it seems to be important to study the adjoint cohomology group  $H^1(\mathfrak{G}(n,\varepsilon),\mathfrak{G}(n,\varepsilon))$  in order to describe right extensions of  $\mathfrak{G}(n,\varepsilon)$  and to generalize the Lie algebras  $\mathfrak{H}(n,r)$ . Another challenge is to replace the Lie algebra of vector fields on  $\mathbb{R}$  by Lie algebras of vector fields on  $\mathbb{R}^n$  in the constructions of  $\mathfrak{G}(n,\varepsilon)$  and  $\mathfrak{H}(n,r)$ , to consider exotic cohomology of the resulting Lie algebras and to study whether there exist related integrable systems. For example, the symmetry pseudo-groups of the heavenly equations and their deformations are right extensions of Lie algebras of the form  $\mathcal{A} \otimes \mathfrak{h}$ , where  $\mathfrak{h}$  is the Lie algebra of Hamiltonian vector fields on  $\mathbb{R}^2$  and  $\mathcal{A}$  are algebras of truncated polynomials, [19, 20]. Other interesting examples include equations discussed in [21]. It is natural to elucidate the relationship among the Lax representations and the structure of symmetry pseudogroups for the above-mentioned integrable equations. We intend to address this issue in our future work.

#### Acknowledgments

This research was supported in part by the Polish Ministry of Science and Higher Education. I am grateful to the organizers of the conference "Geometry and Algebra of PDEs" held at the Arctic University of Norway, Tromsø, 6–10 June 2017, for financial support.

I am pleased to thank V.V. Lychagin, I.S. Krasil'shchik, P.J. Olver, B.S. Kruglikov, D.V. Alekseevsky, B.A. Khesin and P. Zusmanovich for useful and stimulating discussions, and to V.S. Gerdjikov for pointing out the reference [8] and providing the copy thereof.

# References

# References

- M. Błaszak. Classical R-matrices on Poisson algebras and related dispersionless systems. Phys. Lett. A 297 (2002), 191–195
- [2] A.V. Bocharov et al., Symmetries of Differential Equations in Mathematical Physics and Natural Sciences, edited by A.M. Vinogradov and I.S. Krasil'shchik. Factorial Publ. House, 1997 (in Russian). English translation: Amer. Math. Soc., 1999
- [3] R. Carroll. Y. Kodama. Solution of the dispersionless Hirota equations. J. Phys. A, 28 (1995), 6373–6387
- [4] E. Cartan, Sur la structure des groupes infinis de transformations, *Œuvres Complètes*, Part II, V. 2, 571–714, Paris, Gauthier - Villars, 1953.
- [5] E. Cartan, La structure des groupes infinis, *Œuvres Complètes*, Part II, V. 2, 1335–1384, Paris, Gauthier - Villars, 1953.
- [6] B.A. Dubrovin, S.P. Novikov. Hydrodynamics of weakly deformed soliton lattices. Differential geometry and Hamiltonian theory. Russian Mathematical Surveys, 44:6 (1989), 35–124
- [7] Geometrical Approaches to Differential Equations, R. Martini (Ed.) Lecture Notes in Mathematics, Springer, 1980

- [8] V. S. Gerdjikov.  $Z_N$ -reductions and new integrable versions of derivative nonlinear Schrödinger equations. In: Nonlinear Evolution Equations: Integrability and Spectral Methods, Ed. A. P. Fordy, A. Degasperis, M. Lakshmanan, Manchester University Press, 1990, 367–372.
- [9] J. Gibbons. The Zabolotskaya–Khokhlov equation and the inverse scattering problem of classical mechanics. In: *Dynamical Problems in Soliton Systems*, Ed. S. Takeno, Springer 1985, 36–41
- [10] J. Gibbons, S.P. Tsarev. Reductions of the Benney equations. Phys. Lett. A, 211 (1996), 19–24
- [11] D.B. Fuks. Cohomology of Infinite-Dimensional Lie Algebras. Consultants Bureau, N.Y., 1986
- [12] T.A. Ivey, J.M. Landsberg. Cartan for Beginners: Differential Geometry via Moving Frames and Exterior Differential Systems. AMS, Providence, Rhode Island, 2003
- [13] Y. Kodama. A method for solving the dispersionless KP equation and its exact solutions. Phys. Lett. A 129 (1988), 223–226
- [14] I.S. Krasil'shchik, A.M. Vinogradov, Nonlocal symmetries and the theory of coverings. Acta Appl. Math. 2 (1984) 79–86
- [15] I.S. Krasil'shchik, A.M. Vinogradov. Nonlocal trends in the geometry of differential equations: symmetries, conservation laws, and Bäcklund transformations. Acta Appl. Math. 15 (1989), 161–209
- [16] I.S. Krasil'shchik, A.M. Verbovetsky, R. Vitolo, A unified approach to computation of integrable structures, Acta Appl. Math. 120 (2012) 199– 218
- [17] J. Krasil'shchik, A. Verbovetsky, Geometry of jet spaces and integrable systems, J. Geom. Phys. 61 (2011) 1633–1674
- [18] I.M. Krichever. Method of averaging for two-dimensional "integrable" equations. Functional Analysis and Its Applications 22:3 (1988) 200– 213
- [19] B.S. Kruglikov, O.I. Morozov. SDiff(2) and uniqueness of the Plebański equation. J. Math. Phys. 53 (2012), 083506, 11 pp.

- [20] B.S. Kruglikov, O.I. Morozov. Integrable dispersionless PDEs in 4D, their symmetry pseudogroups and deformations. Lett. Math. Phys. 105 (2015) 1703–1723
- [21] B.S. Kruglikov, O.I. Morozov. A Bäcklund transformation between the four-dimensional Martínez Alonso–Shabat and Ferapontov–Khusnutdinova equations. Theor. Math. Phys. 188(3) (2016) 1358–1360
- [22] B.A. Kupershmidt. The quasiclassical limit of the modified KP hierarchy. J. Phys. A, 23 (1990), 871–886
- [23] G.M. Kuz'mina, On a possibility to reduce a system of two first-order partial differential equations to a single equation of the second order, Proc. Moscow State Pedagogical Institute 271 (1967), 67–76 (in Russian).
- [24] C.C. Lin, E. Reissner, H.S. Tsien, On two-dimensional non-steady motion of a slender body in a compressible fluid, J. Math. Phys. 27 (1948) 220–231
- [25] O.I. Morozov. Contact integrable extensions of symmetry pseudo-groups and coverings of (2 + 1) dispersionless integrable equations. J. Geom. Phys. **59** (2009), 1461–1475
- [26] O.I. Morozov. Deformed cohomologies of symmetry pseudo-groups and coverings of differential equations. J. Geom. Phys. 113 (2017), 215–225
- [27] S.P. Novikov. On the exotic De-Rham cohomology. Perturbation theory as a spectral sequence. arXiv:math-ph/0201019
- [28] P.J. Olver. Equivalence, Invariants, and Symmetry. Cambridge University Press, Cambridge, 1995.
- [29] M.V. Pavlov. Modified dispersionless Veselov-Novikov equations and corresponding hydrodynamic chains. arXiv:nlin/0611022
- [30] O. Stormark, Lie's Structural Approach to PDE Systems, Encyclopedia Math. Appl., 80, Cambridge University Press, Cambridge, 2000
- [31] M.V. Vasil'eva, Structure of Infinite Lie Groups of Transformations, Moscow State Pedagogical Institute, Moscow, 1972 (in Russian)

- [32] E.A. Zabolotskaya, R.V. Khokhlov, Quasi-plane waves in the nonlinear acoustics of confined beams, Sov. Phys. Acoust. 15 (1969) 35–40
- [33] V.E. Zakharov. Benney equations and quasiclassical approximation in the method of the inverse problem. Functional Analysis and Its Applications, 1980, 14:2, 8998
- [34] What is integrability, V.E. Zakharov (Ed.) Springer Series in Nonlinear Dynamics, Springer, 1991
- [35] V.E. Zakharov, Dispersionless limit of integrable systems in 2+1 dimensions, in: Singular Limits of Dispersive Waves, Ed. N.M. Ercolani et al., Plenum Press, NY (1994) 165–174