

General N -Dark Soliton Solutions of the Multi-Component Mel'nikov System

Zhong Han^a, Yong Chen^{a,b*}

^aShanghai Key Laboratory of Trustworthy Computing, East China Normal University, Shanghai, 200062, People's Republic of China

^bMOE International Joint Lab of Trustworthy Software, East China Normal University, Shanghai, 200062, People's Republic of China

Abstract

A general form of N -dark soliton solutions of the multi-component Mel'nikov system is presented. Taking the coupled Mel'nikov system comprised of two-component short waves and one-component long wave as an example, its general N -dark-dark soliton solutions in Gram determinant form are constructed through the KP hierarchy reduction method. The dynamics of single dark-dark soliton and two dark-dark solitons are discussed in detail. It can be shown that the collisions of dark-dark solitons are elastic and energies of the solitons in different components completely transmit through. In addition, the dark-dark soliton bound states including both stationary and moving cases are also investigated. An interesting feature for the coupled Mel'nikov system is that the stationary dark-dark soliton bound states can exist for all possible combinations of nonlinearity coefficients including all-positive, all-negative and mixed types, while the moving case are possible when they take opposite signs or they are both negative. The dynamics and several interesting structures of the solutions are illustrated through some figures.

Keywords: Multi-component Mel'nikov system; Dark soliton; KP hierarchy reduction method; τ function

1. Introduction

During the past decades, many studies have been done on coupled systems describing the interaction of short wave packets with long waves in nonlinear dispersive media as they are frequently used in the fields of plasma physics, fluid dynamics and solid state physics [1–3]. What's more, the nonlinear interaction of multiple waves results in several interesting new physical processes [4–6]. On the other hand, the studies of multi-component nonlinear systems have received much attention in recent years [7–9]. Of particular interest is the multi-component generalization of the nonlinear Schrödinger (NLS) equation, the so-called vector NLS equation [8–13]. Another interesting example is the multi-component long-wave-short-wave resonance interaction (LSRI) system [14–16], or the so called multi-component Yajima-Oikawa (YO) system [17–20].

In the current paper, we concentrate on the Mel'nikov system [21–24]

$$i\Phi_y = \Phi_{xx} + u\Phi, \quad (1)$$

$$u_{xt} + u_{xxxx} + 3(u^2)_{xx} - 3u_{yy} + \sigma(\Phi\Phi^*)_{xx} = 0, \quad (2)$$

where $\sigma = \pm 1$, $\Phi \equiv \Phi(x, y, t)$ is the complex short wave amplitude and $u \equiv u(x, y, t)$ is the real long wave amplitude, the subscripts denote partial differentiation and the asterisk means complex conjugate hereafter. This system is used to describe the interaction of long waves with short wave packets propagating on the x - y plane at an angle to each other. It may be considered either as a generalization of the Kadomtsev-Petviashvili (KP) equation with the addition of a complex scalar field or as a generalization of the NLS equation with a real scalar field [25]. Its two-component generalization consisting of two short wave components and one long wave component is given

*Corresponding author.

Email address: ychen@sei.ecnu.edu.cn (Yong Chen^{a,b})

by

$$i\Phi_y^{(1)} = \Phi_{xx}^{(1)} + u\Phi^{(1)}, \quad (3)$$

$$i\Phi_y^{(2)} = \Phi_{xx}^{(2)} + u\Phi^{(2)}, \quad (4)$$

$$u_{xt} + u_{xxxx} + 3(u^2)_{xx} - 3u_{yy} + (\sigma_1\Phi^{(1)}\Phi^{(1)*} + \sigma_2\Phi^{(2)}\Phi^{(2)*})_{xx} = 0, \quad (5)$$

where $\sigma_1, \sigma_2 = \pm 1$. Actually, the above coupled system can be generalized to a multi-component system consisting of M short wave components and one long wave component, which is written in the following form

$$i\Phi_y^{(k)} = \Phi_{xx}^{(k)} + u\Phi^{(k)}, \quad k = 1, 2, \dots, M, \quad (6)$$

$$u_{xt} + u_{xxxx} + 3(u^2)_{xx} - 3u_{yy} + \left(\sum_{k=1}^M \sigma_k \Phi^{(k)} \Phi^{(k)*} \right)_{xx} = 0, \quad (7)$$

where $\sigma_k = \pm 1$ for $k = 1, 2, \dots, M$.

The Mel'nikov system (1)-(2) admits boomeron type solutions which can be realized from an asymptotic analysis of the two soliton solutions [23], and its multi-soliton solutions are derived in Ref. [24] via the theory of matrices. Soliton solutions of bright- and dark-types have been obtained from the Wronskian solutions of the KP hierarchy equations [26]. Its Painlevé property and some exponentially localized dromion type solutions are studied in Ref. [25]. More recently, its rogue wave solutions are also derived by virtue of the Hirota's bilinear method [27]. However, as far as we know, the general multi-dark soliton solutions of the multi-component Mel'nikov system have not been reported yet. Actually, general multi-dark soliton solutions of two-dimensional multi-component integrable systems are rather rare excepted that the multi-dark soliton solutions of the two-dimensional (2D) multi-component Yajima-Oikawa (YO) systems are reported in Ref. [18].

The KP hierarchy reduction method for deriving soliton solutions of integrable systems is firstly developed by the Kyoto school [29, 30], and later used to get solutions of the NLS equation, the modified KdV equation and the Davey-Stewartson (DS) equation. Indeed, the pseudo-reduction of the two-dimensional Toda lattice hierarchy to constrained KP systems with dark soliton solutions is established in Ref. [31], while the reduction to constrained KP systems with bright soliton solutions from the multi-component KP hierarchy is developed in Ref. [32]. Using this reduction method, the general N -dark-dark solitons of a two-coupled focusing-defocusing NLS equations are obtained in Ref. [28] by Ohta et al. Also by virtue of this method, the general bright-dark N -soliton solutions of the vector NLS equations with all possible combinations of nonlinearities containing all focusing, all defocusing and mixed types are constructed in Ref. [33] by Feng. More recently, this method has been applied to derive the N -dark soliton [18] and mixed N -soliton [19] solutions of multi-component YO system. In addition, the KP hierarchy reduction method has also been developed to construct rational solutions (lump and rogue wave solutions) of soliton equations [34–36], see also Refs. [20, 37]

In the present paper, the general N -dark-dark soliton solutions of Eqs.(3)-(5) are obtained through the KP hierarchy reduction method and their dynamics are also studied in detail. Following the KP theory, we derive the general N -dark-dark soliton solutions in terms of Gram determinants from the τ -function solutions of the KP hierarchy. For single dark-dark solitons, we show that the degrees of "darkness" in the two components are different in general. For the collisions of two dark-dark solitons, it can be shown that there is no energy transfer in two components of each soliton, hence they are elastic collisions. As a matter of fact, the soliton bound state is a fascinating subject in the soliton theory. The dark-dark soliton bound states in the 2D coupled YO system and the coupled NLS equations have been reported in Refs. [18] and [28] respectively. For those two models, it has been shown that the bound states exist only when the coefficients of nonlinear terms take opposite signs. Different from Refs. [18] and [28], it can be shown that for the coupled Mel'nikov system (3)-(5), the stationary dark-dark soliton bound states can exist for all possible combinations of nonlinearity coefficients including all-positive, all-negative and mixed types, while the moving case are possible when they take opposite signs or they are both negative.

The rest of this paper is organized as follows. In section 2, the general N -dark-dark soliton solutions in the Gram determinant form of the two-component Mel'nikov system are obtained via the KP hierarchy reduction

method. Section 3 devotes to analysis the dynamics of single and two dark-dark solitons. In section 4, the dark-dark soliton bound states including both the stationary and the moving cases are investigated in detail. In section 5, the general N -dark soliton solutions of the multi-component Mel'nikov system are presented without a detail derivation. We summarize the paper in section 6.

2. Dark-Dark Soliton Solutions of the Coupled Mel'nikov system

By virtue of the dependent variable transformation

$$\Phi^{(1)} = \rho_1 e^{i\theta_1} \frac{g}{f}, \quad \Phi^{(2)} = \rho_2 e^{i\theta_2} \frac{h}{f}, \quad u = 2(\log f)_{xx}, \quad (8)$$

where $f \equiv f(x, y, t)$ is a real function, $g \equiv g(x, y, t)$ and $h \equiv h(x, y, t)$ are two complex functions, ρ_1 and ρ_2 are two positive constants. Meanwhile, $\theta_i = \alpha_i x + \alpha_i^2 y + \beta_i(t)$ for $i = 1, 2$, where α_i are real constants and $\beta_i(t)$ are arbitrary real functions. Then, Eqs.(3)-(5) can be converted into the following bilinear forms

$$(D_x^2 + 2i\alpha_1 D_x - iD_y)g \cdot f = 0, \quad (9)$$

$$(D_x^2 + 2i\alpha_2 D_x - iD_y)h \cdot f = 0, \quad (10)$$

$$(D_x^4 + D_x D_t - 3D_y^2)f \cdot f = \sigma_1 \rho_1^2 (f^2 - gg^*) + \sigma_2 \rho_2^2 (f^2 - hh^*), \quad (11)$$

where D is Hirota's bilinear differential operator defined as

$$D_x^l D_y^m D_t^n f(x, y, t) \cdot g(x, y, t) = \left(\frac{\partial}{\partial x} - \frac{\partial}{\partial x'}\right)^l \left(\frac{\partial}{\partial y} - \frac{\partial}{\partial y'}\right)^m \left(\frac{\partial}{\partial t} - \frac{\partial}{\partial t'}\right)^n f(x, y, t) \cdot g(x', y', t')|_{x=x', y=y', t=t'}. \quad (12)$$

If a new independent variable s is introduced, Eq.(11) can be decoupled into

$$(D_x^4 + D_x D_s - 3D_y^2)f \cdot f = 0, \quad (13)$$

$$(D_x D_t - D_x D_s)f \cdot f = \sigma_1 \rho_1^2 (f^2 - gg^*) + \sigma_2 \rho_2^2 (f^2 - hh^*). \quad (14)$$

Following the KP theory, the equations under study are considered as a reduction of the KP hierarchy, then their soliton solutions in the Gram type or Wronski type determinants can be derived directly from the τ function of the KP hierarchy under this reduction. To this end, the Gram determinant solutions of the KP hierarchy are presented here.

Lemma 1 The following bilinear equations in the KP hierarchy [18, 26, 28]

$$(D_{x_1}^2 + 2aD_{x_1} - D_{x_2})\tau(k+1, l) \cdot \tau(k, l) = 0, \quad (15)$$

$$\left(\frac{1}{2}D_{x_1} D_{x_{-1}} - 1\right)\tau(k, l) \cdot \tau(k, l) = -\tau(k+1, l)\tau(k-1, l), \quad (16)$$

$$(D_{x_1}^2 + 2bD_{x_1} - D_{x_2})\tau(k, l+1) \cdot \tau(k, l) = 0, \quad (17)$$

$$\left(\frac{1}{2}D_{x_1} D_{y_{-1}} - 1\right)\tau(k, l) \cdot \tau(k, l) = -\tau(k, l+1)\tau(k, l-1), \quad (18)$$

$$(D_{x_1}^4 - 4D_{x_1} D_{x_3} + 3D_{x_2}^2)\tau(k, l) \cdot \tau(k, l) = 0, \quad (19)$$

where a and b are complex constants, k and l are integers, have the Gram determinant solutions

$$\tau(k, l) = |m_{ij}(k, l)|_{1 \leq i, j \leq N}, \quad (20)$$

where the entries of the determinant are given by

$$\begin{aligned} m_{ij}(k, l) &= c_{ij} + \int \varphi_i(k, l) \psi_j(k, l) dx_1, \\ \varphi_i(k, l) &= (p_i - a)^k (p_i - b)^l \exp(\theta_i), \\ \psi_j(k, l) &= \left(-\frac{1}{q_j + a}\right)^k \left(-\frac{1}{q_j + b}\right)^l \exp(\tilde{\theta}_j), \end{aligned}$$

with

$$\begin{aligned} \theta_i &= \frac{1}{p_i - a} x_{-1} + \frac{1}{p_i - b} y_{-1} + p_i x_1 + p_i^2 x_2 + p_i^3 x_3 + \theta_{i0}, \\ \tilde{\theta}_j &= \frac{1}{q_j + a} x_{-1} + \frac{1}{q_j + b} y_{-1} + q_j x_1 - q_j^2 x_2 + q_j^3 x_3 + \tilde{\theta}_{j0}, \end{aligned}$$

where $c_{ij}, p_i, q_j, \theta_{i0}$ and $\tilde{\theta}_{j0}$, ($i, j = 1, 2, \dots, N$) are arbitrary complex constants.

Firstly, it is easy to verify that the φ_i and ψ_j given in Lemma 1 satisfy the following differential and difference relations

$$\partial_{x_2} \varphi_i(k, l) = \partial_{x_1}^2 \varphi_i(k, l), \quad (21)$$

$$\partial_{x_3} \varphi_i(k, l) = \partial_{x_1}^3 \varphi_i(k, l), \quad (22)$$

$$\varphi_i(k + 1, l) = (\partial_{x_1} - a) \varphi_i(k, l), \quad (23)$$

$$\partial_{x_2} \psi_i(k, l) = -\partial_{x_1}^2 \psi_i(k, l), \quad (24)$$

$$\partial_{x_3} \psi_i(k, l) = \partial_{x_1}^3 \psi_i(k, l), \quad (25)$$

$$\psi_i(k - 1, l) = -(\partial_{x_1} + a) \psi_i(k, l). \quad (26)$$

Next, take the determinant

$$\tau(k, l) = \det_{1 \leq i, j \leq N} (m_{ij}(k, l)) = |m_{ij}(k, l)|_{1 \leq i, j \leq N},$$

with the matrix elements $m_{ij}(k, l)$ defined as

$$m_{ij}(k, l) = c_{ij} + \int \varphi_i(k, l) \psi_j(k, l) dx_1.$$

Then it is easy to verify that $m_{ij}(k, l)$ satisfy

$$\partial_{x_{-1}} m_{ij}(k, l) = -\varphi_i(k - 1, l) \psi_j(k + 1, l), \quad (27)$$

$$\partial_{x_1} m_{ij}(k, l) = \varphi_i(k, l) \psi_j(k, l), \quad (28)$$

$$\partial_{x_2} m_{ij}(k, l) = (\partial_{x_1} \varphi_i(k, l)) \psi_j(k, l) - \varphi_i(k, l) \partial_{x_1} \psi_j(k, l), \quad (29)$$

$$\partial_{x_3} m_{ij}(k, l) = (\partial_{x_1}^2 \varphi_i(k, l)) \psi_j(k, l) - (\partial_{x_1} \varphi_i(k, l)) \partial_{x_1} \psi_j(k, l) + \varphi_i(k, l) \partial_{x_1}^2 \psi_j(k, l), \quad (30)$$

$$m_{ij}(k + 1, l) = m_{ij}(k, l) + \varphi_i(k, l) \psi_j(k + 1, l). \quad (31)$$

Note that Eqs.(15) and (16) are independent of the variable x_3 , the differential and difference relations with respect to the variables x_{-1}, x_1 and x_2 are the same with the Lemma 2.1 in Ref. [18] and the Lemma 1 in Ref. [28], hence they hold true automatically. Here, we only need to prove that the determinant (20) satisfies the bilinear KP equation (19).

Proof Lemma 1: By virtue of the differential of determinant and the expansion formula of bordered determi-

nant [38], the derivatives of the τ function can be expressed by the following bordered determinants

$$\partial_{x_1} \tau(k, l) = - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix}, \quad (32)$$

$$\partial_{x_2} \tau(k, l) = - \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix} + \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix}, \quad (33)$$

$$\partial_{x_3} \tau(k, l) = - \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1}^2 \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix} + \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1}^2 \psi_j(k, l) & 0 \end{vmatrix}, \quad (34)$$

$$\partial_{x_1}^2 \tau(k, l) = - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix}, \quad (35)$$

$$\partial_{x_1}^3 \tau(k, l) = - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1}^2 \psi_j(k, l) & 0 \end{vmatrix} - 2 \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1}^2 \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix}, \quad (36)$$

$$\partial_{x_1} \partial_{x_3} \tau(k, l) = \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1}^3 \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1}^3 \psi_j(k, l) & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \psi_j(k, l) & 0 & 0 \\ \partial_{x_1} \psi_j(k, l) & 0 & 0 \end{vmatrix}, \quad (37)$$

$$\begin{aligned} \partial_{x_2}^2 \tau(k, l) &= \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \partial_{x_1}^2 \psi_j(k, l) & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1}^3 \psi_j(k, l) & 0 \end{vmatrix} + \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1}^2 \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix} \\ &\quad - \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1}^3 \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix} + 2 \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \psi_j(k, l) & 0 & 0 \\ \partial_{x_1} \psi_j(k, l) & 0 & 0 \end{vmatrix}, \end{aligned} \quad (38)$$

$$\begin{aligned} \partial_{x_1}^4 \tau(k, l) &= - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1}^3 \psi_j(k, l) & 0 \end{vmatrix} - \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1}^3 \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix} - 3 \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \partial_{x_1}^2 \psi_j(k, l) & 0 \end{vmatrix} \\ &\quad - 3 \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1}^2 \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix} + 2 \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \psi_j(k, l) & 0 & 0 \\ \partial_{x_1} \psi_j(k, l) & 0 & 0 \end{vmatrix}, \end{aligned} \quad (39)$$

where the bordered determinants are defined as

$$\begin{vmatrix} m_{ij} & \varphi_i \\ \psi_j & 0 \end{vmatrix} = \begin{vmatrix} m_{11} & m_{12} & \cdots & m_{1N} & \varphi_1 \\ m_{21} & m_{22} & \cdots & m_{2N} & \varphi_2 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ m_{N1} & m_{N2} & \cdots & m_{NN} & \varphi_N \\ \psi_1 & \psi_2 & \cdots & \psi_N & 0 \end{vmatrix}. \quad (40)$$

By virtue of the above expresses and the Jacobi formula of determinants, it is easy to verify that the τ function satisfies the bilinear KP equation (19)

$$\begin{aligned} &(D_{x_1}^4 - 4D_{x_1} D_{x_3} + 3D_{x_2}^2) \tau(k, l) \cdot \tau(k, l) \\ &= 2 \left[\left(\partial_{x_1}^4 \tau(k, l) - 4 \partial_{x_1} \partial_{x_3} \tau(k, l) + 3 \partial_{x_2}^2 \tau(k, l) \right) \tau(k, l) - 4 \left(\partial_{x_1}^3 \tau(k, l) - \partial_{x_3} \tau(k, l) \right) \partial_{x_1} \tau(k, l) \right. \\ &\quad \left. + 3 \left(\partial_{x_1}^2 \tau(k, l) - \partial_{x_2} \tau(k, l) \right) \left(\partial_{x_1}^2 \tau(k, l) + \partial_{x_2} \tau(k, l) \right) \right] \\ &= 24 \left(\begin{vmatrix} m_{ij}(k, l) \\ \psi_j(k, l) \end{vmatrix} \cdot \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \psi_j(k, l) & 0 & 0 \\ \partial_{x_1} \psi_j(k, l) & 0 & 0 \end{vmatrix} + \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix} \cdot \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix} \right. \\ &\quad \left. - \begin{vmatrix} m_{ij}(k, l) & \varphi_i(k, l) \\ \psi_j(k, l) & 0 \end{vmatrix} \cdot \begin{vmatrix} m_{ij}(k, l) & \partial_{x_1} \varphi_i(k, l) \\ \partial_{x_1} \psi_j(k, l) & 0 \end{vmatrix} \right) \\ &= 0. \end{aligned} \quad (41)$$

This completes the proof of Lemma 1. \square

Now we perform a reduction of the above bilinear equations to derive the general formulae for N -dark-dark soliton solutions of Eqs.(3)-(5). Assuming x_{-1}, y_{-1}, x_1, x_3 are real, $x_2, a(= i\alpha_1), b(= i\alpha_2)$ are pure imaginary and $q_i = p_i^*, \tilde{\theta}_{j0} = \theta_{j0}^*, c_{ji} = c_{ij}^* = \delta_{ij}$, then we can get

$$\tilde{\theta}_j = \theta_j^*, \quad m_{ji}(k, l) = m_{ij}^*(-k, -l), \quad \tau(k, l) = \tau^*(-k, -l). \quad (42)$$

Moreover, by defining

$$f = \tau(0, 0), \quad g = \tau(1, 0), \quad h = \tau(0, 1), \quad g^* = \tau(-1, 0), \quad h^* = \tau(0, -1), \quad (43)$$

the bilinear equations (15)-(19) can be recast into

$$(D_{x_1}^2 + 2i\alpha_1 D_{x_1} - D_{x_2})g \cdot f = 0, \quad (44)$$

$$\left(\frac{1}{2}D_{x_1}D_{x_{-1}} - 1\right)f \cdot f = -gg^*, \quad (45)$$

$$(D_{x_1}^2 + 2i\alpha_2 D_{x_1} - D_{x_2})h \cdot f = 0, \quad (46)$$

$$\left(\frac{1}{2}D_{x_1}D_{y_{-1}} - 1\right)f \cdot f = -hh^*, \quad (47)$$

$$(D_{x_1}^4 - 4D_{x_1}D_{x_3} + 3D_{x_2}^2)f \cdot f = 0. \quad (48)$$

By virtue of the following independent variable transformation

$$x_1 = x, \quad x_2 = -iy, \quad x_3 = -4(t + s), \quad x_{-1} = \frac{1}{2}\sigma_1\rho_1^2t, \quad y_{-1} = \frac{1}{2}\sigma_2\rho_2^2t, \quad (49)$$

the bilinear equations (44)-(48) are recast into the bilinear form (9)-(10) and (13)-(14). Thus, applying the independent variable transformation (49) to the f, g, h in (43) and neglecting the s dependence, the following theorem for the general N -dark-dark soliton solutions of Eqs.(3)-(5) is immediately obtained.

Theorem 1 The N -dark-dark soliton solutions of the coupled Mel'nikov system (3)-(5) are

$$\Phi^{(1)} = \rho_1 e^{i[\alpha_1 x + \alpha_1^2 y + \beta_1(t)]} \frac{g}{f}, \quad (50)$$

$$\Phi^{(2)} = \rho_2 e^{i[\alpha_2 x + \alpha_2^2 y + \beta_2(t)]} \frac{h}{f}, \quad (51)$$

$$u = 2(\log f)_{xx}, \quad (52)$$

where f, g and h are Gram determinants given by

$$f = \left| \delta_{ij} + \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$

$$g = \left| \delta_{ij} + \left(-\frac{p_i - i\alpha_1}{p_j^* + i\alpha_1} \right) \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$

$$h = \left| \delta_{ij} + \left(-\frac{p_i - i\alpha_2}{p_j^* + i\alpha_2} \right) \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$

with

$$\xi_j = p_j x - ip_j^2 y + \frac{1}{2} \left(\frac{\sigma_1 \rho_1^2}{p_j - i\alpha_1} + \frac{\sigma_2 \rho_2^2}{p_j - i\alpha_2} - 8p_j^3 \right) t + \xi_{j0},$$

where p_j and ξ_{j0} are complex constants, δ_{ij} is the Kronecker symbol (δ_{ij} is 1 when $i = j$ and 0 otherwise).

3. Dynamics of Dark-Dark Solitons

3.1. Single dark-dark solitons

To get single dark-dark soliton solution of Eqs.(1)-(3), we take $N = 1$ in the formula (50)-(52). The Gram determinants read

$$f = 1 + \frac{1}{p_1 + p_1^*} e^{\xi_1 + \xi_1^*}, \quad (53)$$

$$g = 1 - \frac{1}{p_1 + p_1^*} \frac{p_1 - i\alpha_1}{p_1^* + i\alpha_1} e^{\xi_1 + \xi_1^*}, \quad (54)$$

$$h = 1 - \frac{1}{p_1 + p_1^*} \frac{p_1 - i\alpha_2}{p_1^* + i\alpha_2} e^{\xi_1 + \xi_1^*}, \quad (55)$$

and the single dark-dark soliton solution can be written as

$$\Phi^{(1)} = \frac{\rho_1}{2} e^{i[\alpha_1 x + \alpha_1^2 y + \beta_1(t)]} \times \left[1 + K_1^{(1)} + (K_1^{(1)} - 1) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right) \right], \quad (56)$$

$$\Phi^{(2)} = \frac{\rho_2}{2} e^{i[\alpha_2 x + \alpha_2^2 y + \beta_2(t)]} \times \left[1 + K_1^{(2)} + (K_1^{(2)} - 1) \tanh\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right) \right], \quad (57)$$

$$u = \frac{1}{2} (p_1 + p_1^*)^2 \operatorname{sech}^2\left(\frac{\xi_1 + \xi_1^* + \Theta_1}{2}\right), \quad (58)$$

with

$$e^{\Theta_1} = \frac{1}{p_1 + p_1^*} = \frac{1}{2a_1},$$

$$K_1^{(1)} = -\frac{p_1 - i\alpha_1}{p_1^* + i\alpha_1} = -\frac{a_1 + i(b_1 - \alpha_1)}{a_1 - i(b_1 - \alpha_1)},$$

$$K_1^{(2)} = -\frac{p_1 - i\alpha_2}{p_1^* + i\alpha_2} = -\frac{a_1 + i(b_1 - \alpha_2)}{a_1 - i(b_1 - \alpha_2)},$$

$$\xi_1 + \xi_1^* = 2a_1 x + 4a_1 b_1 y + \left[\frac{\sigma_1 a_1 \rho_1^2}{a_1^2 + (b_1 - \alpha_1)^2} + \frac{\sigma_2 a_1 \rho_2^2}{a_1^2 + (b_1 - \alpha_2)^2} - 8a_1(a_1^2 - 3b_1^2) \right] t + 2\xi_{10R},$$

where $p_1 = a_1 + ib_1$, and a_1, b_1, ξ_{10R} are real constants, ξ_{10R} is the real part of ξ_{10} .

From (56)-(58), it is easy to know that the intensity functions $|\Phi^{(1)}|, |\Phi^{(2)}|$ of the short wave components and the long wave component u moving at velocity $-\frac{1}{2} \left(\frac{\sigma_1 \rho_1^2}{a_1^2 + (b_1 - \alpha_1)^2} + \frac{\sigma_2 \rho_2^2}{a_1^2 + (b_1 - \alpha_2)^2} \right) + 4(a_1^2 - 3b_1^2)$ along the x -direction. In addition, $|\Phi^{(1)}| \rightarrow \rho_1, |\Phi^{(2)}| \rightarrow \rho_2, u \rightarrow 0$ when $x, y \rightarrow \pm\infty$.

Taking $K_1^{(1)} = \exp(2i\phi_1^{(1)})$ and $K_1^{(2)} = \exp(2i\phi_1^{(2)})$, i.e., $2\phi_1^{(1)}$ and $2\phi_1^{(2)}$ are the phases of the constants $K_1^{(1)}$ and $K_1^{(2)}$ respectively. As x and y vary from $-\infty$ to $+\infty$, the phases of the short wave components $\Phi^{(1)}$ and $\Phi^{(2)}$ acquire shifts in the amount of $2\phi_1^{(1)}$ and $2\phi_1^{(2)}$ while the long wave component u phase shifts is zero. Without loss of generality, we can restrict $-\pi < 2\phi_1^{(1)}, 2\phi_1^{(2)} \leq \pi$, i.e., $-\frac{\pi}{2} < \phi_1^{(1)}, \phi_1^{(2)} \leq \frac{\pi}{2}$. Then at the center of the solitons where $\xi_1 + \xi_1^* + \Theta_1 = 0$, intensities of the components are $|\Phi^{(1)}|_{\text{center}} = \rho_1 \cos \phi_1^{(1)}, |\Phi^{(2)}|_{\text{center}} = \rho_2 \cos \phi_1^{(2)}, u = 2a_1^2$. For the short wave components $\Phi^{(1)}$ and $\Phi^{(2)}$, the fact that the center intensities are lower than the background intensities ρ_1 and ρ_2 , thus these solitons are dark-dark solitons. Further more, the intensity dips at the centers of the $\Phi^{(1)}$ and $\Phi^{(2)}$ components are controlled by the phase shifts $2\phi_1^{(1)}$ and $2\phi_1^{(2)}$ respectively, hence these phase shifts dictate how "dark" the center is.

According to the values of α_1 and α_2 , there exist the following two different cases:

(1) $\alpha_1 = \alpha_2$, then $K_1^{(1)} = K_1^{(2)}$, therefor $\phi_1^{(1)} = \phi_1^{(2)}$. For this case, the short wave components $\Phi^{(1)}$ and $\Phi^{(2)}$ are proportional to each other, and they have the same degrees of darkness at the center. In this situation, the dark-dark solitons of the coupled Mel'nikov system is equivalent to a scalar dark soliton of the single-component Mel'nikov system, thus it is viewed as a degenerate case similar to the coupled YO system [18] and the coupled NLS equations [28]. These degenerate solitons are illustrated in Fig.1, from which it can be seen that both the $\Phi^{(1)}$ and $\Phi^{(2)}$ components are zero intensity at the soliton center.

(2) $\alpha_1 \neq \alpha_2$, then $K_1^{(1)} \neq K_1^{(2)}$, thus $\phi_1^{(1)} \neq \phi_1^{(2)}$. This suggests that the $\Phi^{(1)}$ and $\Phi^{(2)}$ components in these solitons are not proportional to each other, thus can not be reducible to scalar single dark soliton. In this non-degenerate case, the $\Phi^{(1)}$ and $\Phi^{(2)}$ components have different degrees of darkness at the center. As is shown in Fig.2, the intensity of the component $\Phi^{(1)}$ is zero intensity, while the intensity of the component $\Phi^{(2)}$ is nonzero intensity at its center.

3.2. Two dark-dark solitons

Two dark-dark solitons of Eqs.(1)-(3) correspond to $N = 2$ in the formula (50)-(52). In this case, we have

$$\Phi^{(1)} = \rho_1 e^{i[\alpha_1 x + \alpha_1^2 y + \beta_1(t)]} \frac{g_2}{f_2}, \quad (59)$$

$$\Phi^{(2)} = \rho_2 e^{i[\alpha_2 x + \alpha_2^2 y + \beta_2(t)]} \frac{h_2}{f_2}, \quad (60)$$

$$u = 2(\log f_2)_{xx}, \quad (61)$$

with

$$f_2 = 1 + e^{\xi_1 + \xi_1^* + \Theta_1} + e^{\xi_2 + \xi_2^* + \Theta_2} + \Omega_{12} e^{\xi_1 + \xi_1^* + \xi_2 + \xi_2^* + \Theta_1 + \Theta_2}, \quad (62)$$

$$g_2 = 1 + K_1^{(1)} e^{\xi_1 + \xi_1^* + \Theta_1} + K_2^{(1)} e^{\xi_2 + \xi_2^* + \Theta_2} + \Omega_{12} K_1^{(1)} K_2^{(1)} e^{\xi_1 + \xi_1^* + \xi_2 + \xi_2^* + \Theta_1 + \Theta_2}, \quad (63)$$

$$h_2 = 1 + K_1^{(2)} e^{\xi_1 + \xi_1^* + \Theta_1} + K_2^{(2)} e^{\xi_2 + \xi_2^* + \Theta_2} + \Omega_{12} K_1^{(2)} K_2^{(2)} e^{\xi_1 + \xi_1^* + \xi_2 + \xi_2^* + \Theta_1 + \Theta_2}, \quad (64)$$

and

$$e^{\Theta_j} = \frac{1}{p_j + p_j^*} = \frac{1}{2a_j},$$

$$K_j^{(1)} = -\frac{p_j - i\alpha_1}{p_j^* + i\alpha_1} = -\frac{a_j + i(b_j - \alpha_1)}{a_j - i(b_j - \alpha_1)},$$

$$K_j^{(2)} = -\frac{p_j - i\alpha_2}{p_j^* + i\alpha_2} = -\frac{a_j + i(b_j - \alpha_2)}{a_j - i(b_j - \alpha_2)},$$

$$\Omega_{12} = \left| \frac{p_1 - p_2}{p_1 + p_2^*} \right|^2 = \frac{(a_1 - a_2)^2 + (b_1 - b_2)^2}{(a_1 + a_2)^2 + (b_1 - b_2)^2},$$

$$\begin{aligned} \xi_j + \xi_j^* &= 2a_j x + 4a_j b_j y + \left[\frac{\sigma_1 \rho_1^2 a_j}{a_j^2 + (b_j - \alpha_1)^2} + \frac{\sigma_2 \rho_2^2 a_j}{a_j^2 + (b_j - \alpha_2)^2} - 8a_j(a_j^2 - 3b_j^2) \right] t + 2\xi_{j0R} \\ &= k_{x,j} x + k_{y,j} y + \omega_j t + 2\xi_{j0R} \end{aligned}$$

where $p_j = a_j + ib_j$, a_j, b_j, ξ_{j0R} , ($j = 1, 2$) are real constants.

In the case of $a_2 = -a_1$ and $b_2 = b_1$, i.e., $p_2 = -p_1^*$, the denominator of Ω_{12} is zero. The soliton interaction possess Y-shape on this particular wave number. This Y-shape type soliton solution is found in the KP equation and is also known as the resonant soliton solution. Analogous to the two-soliton solution of the KP equation, the above two soliton interaction solutions are classified into two different types [39–42]:

(1) If $a_1 a_2 < 0$, then $\Omega_{12} > 1$. This case is called the O-type soliton interaction. In this situation, the two asymptotic soliton amplitudes $2a_1^2$ and $2a_2^2$ (in the long wave component u) are equivalent when $a_1 = -a_2$. The interaction peak (the maximum of u) is always greater than the sum of the asymptotic soliton amplitudes.

(2) If $a_1 a_2 > 0$, then $0 < \Omega_{12} < 1$. This case is known as the P-type soliton interaction. In this situation, the two asymptotic soliton amplitudes $2a_1^2$ and $2a_2^2$ (also in u) cannot be equivalent. The interaction peak (the maximum of u) is always less than the sum of the asymptotic soliton amplitudes.

It is obvious that the types of soliton interaction do not depend on the parameters b_1 and b_2 (the imaginary parts of p_1 and p_2). The resonant Y-shape soliton solution can be obtained via taking the limit $b_2 \rightarrow b_1$ in the equal-amplitude O-type two-soliton solution ($a_1 = -a_2$). In addition, the interaction coefficient Ω_{12} in the two-soliton solution of the coupled Mel'nikov system is always non-negative while it can be negative in the two-soliton

solution of the KP equation.

The collision of two dark-dark solitons is illustrated in Fig.3. It can be seen that after collision, the two solitons pass through each other without any change of shape, darkness or velocity in both components. Therefore, there is no energy exchange between the two solitons or between the $\Phi^{(1)}$ and $\Phi^{(2)}$ components after collision. This complete transmission of energy of dark-dark solitons in both components after collision is a common phenomenon of the coupled Mel'nikov system which occurs for all possible combinations of σ_1 and σ_2 values (all-positive, all-negative and mixed types).

The reason of the complete energy transmission in dark-dark soliton collisions is that the intensity profile of each dark-dark soliton is completely determined by the background parameters ρ_i, α_i and the soliton parameters a_i, b_i . For the colliding solitons, these parameters are the same, hence do not change through collisions. Therefore, the intensity profile of each dark-dark soliton does not change before and after collision.

4. Dark-Dark Soliton Bound States

In the study of dark solitons, multi-dark-soliton bound states is a fascinating subject. To get two dark-dark soliton bound states of the coupled Mel'nikov system, the two solitons should have the same velocity in both short and long wave components, so that the two constituent dark solitons can stay together for all times. This means that the parameters need to satisfy $\frac{\omega_1}{k_{x,1}} = \frac{\omega_2}{k_{x,2}}$ and $\frac{\omega_1}{k_{y,1}} = \frac{\omega_2}{k_{y,2}}$.

4.1. The stationary dark-dark soliton bound states

The stationary dark-dark soliton bound states suggest that the common velocity is zero. This requires the coefficients of t in the solution (59)-(61) to be zero, i.e.,

$$\frac{\sigma_1 \rho_1^2}{a_i^2 + (b_i - \alpha_1)^2} + \frac{\sigma_2 \rho_2^2}{a_i^2 + (b_i - \alpha_2)^2} - 8(a_i^2 - 3b_i^2) = 0 \quad \text{for } i = 1, 2. \quad (65)$$

These constraints can be satisfied for all possible combinations of σ_1 and σ_2 values. However, for the coupled YO system [18] and the coupled NLS system [28], the corresponding constraints are possible only when nonlinearity coefficients take opposite signs. Hence, for the coupled Mel'nikov system, the stationary dark-dark soliton bound states are expected to exist for all possible combinations of nonlinearity coefficients including all-positive, all-negative and mixed types.

Two different types of bound states are illustrated in Figs.4 and 5 respectively. A nontrivial case of $\frac{k_{y,1}}{k_{x,1}} \neq \frac{k_{y,2}}{k_{x,2}}$ is displayed in Fig.4, which corresponds to an oblique bound state. Meanwhile, a trivial case of $\frac{k_{y,1}}{k_{x,1}} = \frac{k_{y,2}}{k_{x,2}}$ is displayed in Fig.5, which corresponds to a quasi-one-dimensional one.

4.2. The moving dark-dark soliton bound states

If the common velocity being nonzero, i.e., $\omega_1 \neq 0$ and $\omega_2 \neq 0$, we can get the moving dark-dark soliton bound states. Thus the parameters need to satisfy the following conditions:

$$b_1 = b_2, \quad (66)$$

$$\frac{\sigma_1 \rho_1^2}{a_1^2 + (b_1 - \alpha_1)^2} + \frac{\sigma_2 \rho_2^2}{a_1^2 + (b_1 - \alpha_2)^2} - 8a_1^2 = \frac{\sigma_1 \rho_1^2}{a_2^2 + (b_2 - \alpha_1)^2} + \frac{\sigma_2 \rho_2^2}{a_2^2 + (b_2 - \alpha_2)^2} - 8a_2^2. \quad (67)$$

These constraints are not possible when σ_1 and σ_2 are both positive. The reason is that when $\sigma_1 = \sigma_2 = 1$, the function on the left-hand (also right-hand) side of Eq.(67) is a decreasing function of a_j^2 , thus it can not be satisfied for two different positive values a_1 and a_2 for the same values $b_1 = b_2$. However, when σ_1 and σ_2 take opposite signs or they are both negative, the function on the left-hand (also right-hand) side of Eq.(67) may become non-monotone in a_j^2 , hence it becomes possible for Eq.(67) to admit two different positive values a_1 and a_2 for the same values $b_1 = b_2$. For the coupled NLS equations and the coupled YO system, the corresponding constraints can be

satisfied only when the coefficients of nonlinear terms take opposite signs. To demonstrate these moving dark-dark soliton bound states, the following parameters are chosen

$$\sigma_1 = \sigma_2 = -1, \rho_1 = 1, \rho_2 = 2, \alpha_1 = 1, \alpha_2 = \frac{1}{2}, p_1 = 1 + i, p_2 = 0.6012 + i, \xi_{10R} = \xi_{20R} = 0 \quad (68)$$

and the corresponding profiles are illustrated in Figs.6-8 at different times.

An important feature of the bound states is that, for both stationary and moving bound states, as x and y move from $-\infty$ to $+\infty$, the short wave components acquire non-zero phase shifts while the long wave component admits no phase shift. Actually, let $2\phi_j^{(1)}$ and $2\phi_j^{(2)}$ are the phases of constants $K_j^{(1)}$ and $K_j^{(2)}$ respectively, then the phase shifts of the components are $\Phi_{\text{phase shift}}^{(1)} = 2\phi_1^{(1)} + 2\phi_2^{(1)}$, $\Phi_{\text{phase shift}}^{(2)} = 2\phi_1^{(2)} + 2\phi_2^{(2)}$ and $u_{\text{phase shift}} = 0$. It can be seen that the total phase shifts of each short wave component are equal to the sum of the individual phase shifts of the two constituent dark solitons, and are generally non-zero, while the phase shifts of the long wave component are always zero. For instance, the total phase shift of the $\Phi^{(1)}$ component is 2π , and the total phase shift of the $\Phi^{(2)}$ component is -3.9684 , as can be calculated from the above formula.

5. General N Dark Soliton Solutions of the Multi-component Mel'nikov system

In this section, the previous analysis is extended to the Multi-component Mel'nikov system (6)-(7) to get its general N dark soliton solutions. As is known that the multi-bright soliton solutions can be obtained through the reduction of multi-component KP hierarchy, and the multi-dark soliton solutions can be derived from the reduction of single KP hierarchy but with multiple copies of shifted singular points. Consequently, the general N dark soliton solutions of the multi-component Mel'nikov system can be obtained in the same manner as the two-component Mel'nikov system. Here, only the results are presented without a detail calculation.

The multi-component Mel'nikov system (6)-(7) consisting of M short wave components and one long wave component can be converted to the following bilinear form

$$(D_x^2 + 2i\alpha_k D_x - iD_y)g_k \cdot f = 0, \quad k = 1, 2, \dots, M, \quad (69)$$

$$(D_x^4 + D_x D_t - 3D_y^2)f \cdot f = \sum_{k=1}^M \sigma_k \rho_k^2 (f_k^2 - hh^*), \quad (70)$$

by virtue of the dependent variable transformations

$$\Phi^{(k)} = \rho_k e^{i[\alpha_k x + \alpha_k^2 y + \beta_k(t)] \frac{g_k}{f}}, \quad k = 1, 2, \dots, M, \quad (71)$$

$$u = 2(\log f)_{xx}, \quad (72)$$

where $f \equiv f(x, y, t)$ is a real function, $g_k \equiv g_k(x, y, t)$ are complex functions, ρ_k are positive constants, α_k are real constants and $\beta_k(t)$ are arbitrary real functions for $k = 1, 2, \dots, M$.

Also introduce a new independent variable s , Eq.(70) can be decoupled into

$$(D_x^4 + D_x D_s - 3D_y^2)f \cdot f = 0, \quad (73)$$

$$(D_x D_t - D_x D_s)f \cdot f = \sum_{k=1}^M \sigma_k \rho_k^2 (f_k^2 - hh^*). \quad (74)$$

Consider the Gram determinant solutions of the KP hierarchy, in the similar way, we can get the following N -dark soliton solutions of Eqs.(6)-(7)

$$f = \left| \delta_{ij} + \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$

$$g_k = \left| \delta_{ij} + \left(-\frac{p_i - i\alpha_k}{p_j^* + i\alpha_k} \right) \frac{1}{p_i + p_j^*} e^{\xi_i + \xi_j^*} \right|_{N \times N},$$

with

$$\xi_j = p_j x - i p_j^2 y + \frac{1}{2} \left(\sum_{k=1}^M \frac{\sigma_k \rho_k^2}{p_j - i \alpha_k} - 8 p_j^3 \right) t + \xi_{j0},$$

where p_j and ξ_{j0} are complex constants, δ_{ij} is the Kronecker symbol.

6. Conclusions

In this paper, the N -dark-dark soliton solutions of the coupled Mel'nikov system containing two short wave components and one long wave component are constructed in Gram determinant form by virtue of the KP hierarchy reduction method. The derivation of N -dark-dark soliton solutions relies on the connections of the Mel'nikov system with other integrable equations in the KP hierarchy. In addition, the similar expression of general N -dark soliton solutions of multi-component Mel'nikov system comprised of M short wave components and one long wave component are also obtained by simply inserting more copies of the shifts of singular points. As far as we know, general N -dark soliton solutions of two-dimensional (2D) multi-component soliton equations are rather rare except that the N -dark soliton solutions of the 2D multi-component Yajima-Oikawa (YO) systems are studied in Ref. [18].

The dynamics of single and two dark-dark solitons of the coupled Mel'nikov system with two short wave components are studied in detail, and several interesting structures of the solutions have been illustrated through some figures. For single dark-dark solitons, we have shown that the degrees of "darkness" in the two components are different in general. For two dark-dark solitons, in contrast with bright-bright solitons, it has been shown that the collisions of dark-dark solitons are elastic and there is no energy transfer in two components of each soliton. What's more, the dark-dark soliton bound states including both stationary and moving cases are also discussed. For the stationary case, the constraints can be satisfied for all possible combinations of nonlinearity coefficients including all-positive, all-negative and mixed types. For the moving case, the corresponding constraints can be satisfied when they take opposite signs or they are both negative. Whereas, for the coupled YO system [18] and the coupled NLS system [28], the dark-dark soliton bound states are possible only under the condition that the coefficients of nonlinear terms take opposite signs.

Recently, the general mixed N -soliton solutions of the one-dimensional multi-component YO system and the vector NLS equations are obtained in Refs. [19] and [33] respectively. Motivated by the remarkable work in Refs. [19] and [33], the general N -bright and N -bright-dark solitons of the multi-component Mel'nikov system may also be obtained by the KP hierarchy reduction method. However, the reductions for bright and bright-dark solitons are different from the ones for dark solitons in the current paper. On the other hand, rogue waves, as a special phenomenon of solitary waves originally occurring in the deep ocean, have drawn more and more attentions in many other fields [43–45]. As mentioned in Section 1, the KP hierarchy reduction method can also be applied to derive lump and rogue wave solutions of soliton equations. Although the rogue waves of the Mel'nikov system (1)-(2) have been investigated in Ref. [27], the general rational solutions (lump and rogue wave solutions) of the multi-component Mel'nikov system (6)-(7) have not been reported so far (to our knowledge). These questions are interesting and meaningful, we will study them in our future papers.

Acknowledgment

We would like to express our sincere thanks to S.Y. Lou, J.C. Chen and other members of our discussion group for their valuable comments and suggestions. The project is supported by the Global Change Research Program of China (No.2015CB953904), National Natural Science Foundation of China (No.11675054 and 11435005), and Shanghai Collaborative Innovation Center of Trustworthy Software for Internet of Things (No. ZF1213).

References

References

- [1] Zakharov VE. Collapse of Langmuir waves. *Sov Phys JETP* 1972;35:908.
- [2] Nishikawa K, Hojo H, Mima K, Ikezi H. Coupled nonlinear electron-plasma and ion-acoustic waves. *Phys Rev Lett* 1974;33:148.
- [3] Yajima N, Satsuma J. Soliton solutions in a diatomic lattice system. *Prog Theor Phys* 1979;62:370-378.
- [4] Kivshar YS, Agrawal GP. *Optical Solitons: From Fibers to Photonic Crystals*. San Diego: Academic Press; 2003.
- [5] Akhmediev N, Ankiewicz A. *Solitons: Nonlinear Pulses and Beams*. London: Chapman Hall; 1997.
- [6] Scott AC. *Nonlinear Science: Emergence and Dynamics of Coherent Structures*. Oxford: Oxford University Press; 1999.
- [7] Hasegawa A, Kodama Y. *Solitons in optical communications*. Oxford: Clarendon Press; 1995.
- [8] Ablowitz MJ, Prinari B, Trubatch AD. *Discrete and continuous nonlinear Schrödinger systems*. Cambridge: Cambridge Univ. Press; 2004.
- [9] Manakov SV. On the theory of two-dimensional stationary self-focusing of electromagnetic waves. *Sov Phys JETP* 1974;38:248-253.
- [10] Hisakado M, Wadati M. Integrable multi-component hybrid nonlinear Schrödinger equations. *J Phys Soc Jpn* 1995;64:408-413.
- [11] Yan Z. Integrable PT-symmetric local and nonlocal vector nonlinear Schrödinger equations: A unified two-parameter model. *Appl Math Lett* 2015;47:61-68.
- [12] Ling L, Zhao LC, Guo B. Darboux transformation and multi-dark soliton for N -component nonlinear Schrödinger equations. *Nonlinearity* 2015;28:3243.
- [13] Ling L, Zhao LC, Guo B. Darboux transformation and classification of solution for mixed coupled nonlinear Schrödinger equations. *Commun Nonlinear Sci Numer Simulat* 2016;32:285-304.
- [14] Kanna T, Vijayajayanthi M, Sakkaravarthi K, Lakshmanan M. Higher dimensional bright solitons and their collisions in a multicomponent long wave-short wave system. *J Phys A Math Theor* 2009;42:115103.
- [15] Sakkaravarthi K, Kanna T. Dynamics of bright soliton bound states in (2+1)-dimensional multicomponent long wave-short wave system. *Eur Phys J Spec Top* 2013;222:641-653.
- [16] Kanna T, Vijayajayanthi M, Lakshmanan M. Mixed solitons in a (2+1)-dimensional multicomponent long-wave-short-wave system. *Phys Revi E* 2014;90:042901.
- [17] Liu QP. Bi-Hamiltonian structures of the coupled AKNS hierarchy and the coupled Yajima-Oikawa hierarchy. *J Math Phys* 1996;37:2307-2314.
- [18] Chen J, Chen Y, Feng B, Maruno K. Multi-dark soliton solutions of the two-dimensional multi-component Yajima-Oikawa systems. *J Phys Soc Jpn* 2015;84:034002.
- [19] Chen J, Chen Y, Feng B, Maruno K. General mixed multi-soliton solutions to one-dimensional multi-component Yajima-Oikawa system. *J Phys Soc Jpn* 2015;84:074001.
- [20] Chen J, Chen Y, Feng B, Maruno K. Rational solutions to two-and one-dimensional multicomponent Yajima-Oikawa systems. *Phys Lett A*, 2015;379:1510-1519.
- [21] Mel'nikov VK. On equations for wave interactions. *Lett Math Phys* 1983;7: 129-136.
- [22] Mel'nikov VK. Wave emission and absorption in a nonlinear integrable system. *Phys Lett A* 1986;118:22-24.
- [23] Mel'nikov VK. Reflection of waves in nonlinear integrable systems. *J Math Phys* 1987;28:2603-2609.
- [24] Mel'nikov VK. A direct method for deriving a multi-soliton solution for the problem of interaction of waves on the x, y plane. *Comm Math Phys* 1987;112:639-652.
- [25] Kumar CS, Radha R, Lakshmanan M. Exponentially localized solutions of Mel'nikov equation. *Chaos Solitons Fractals* 2004;22:705-712.
- [26] Hase Y, Hirota R, Ohta Y. Soliton Solutions to the Mel'nikov equations. *J Phys Soc Jpn* 1989;58:2713-2720.
- [27] Mu G, Qin Z. Two spatial dimensional N -rogue waves and their dynamics in Mel'nikov equation. *Nonlinear Analysis: Real World Appl* 2014;18:1-13.
- [28] Ohta Y, Wang D, Yang J. General N -dark-dark Solitons in the coupled nonlinear Schrödinger equations. *Stud Appl Math* 2011;127:345-371.
- [29] Jimbo M, Miwa T. Solitons and infinite dimensional Lie algebras. *Publ RIMS Kyoto Univ* 1983;19:943-1001.
- [30] Ohta Y. Wronskian solutions of soliton equations. *RIMS kokyuroku* 1989;684.
- [31] Willox R, Loris I. An algebraic description of generalized k -constraints. *J Phys A Math Gener*, 1999;32:2027.
- [32] Willox R, Loris I. KP constraints from reduced multi-component hierarchies. *J Math Phys* 1999;40:6501.
- [33] Feng BF. General N -soliton solution to a vector nonlinear Schrödinger equation. *J Phys A Math Theor* 2014;47:355203.
- [34] Ohta Y, Yang J. General high-order rogue waves and their dynamics in the nonlinear Schrödinger equation. *Proc R Soc A The Royal Society*, 2012;468:1716-1740.

- [35] Ohta Y, Yang J. Rogue waves in the Davey-Stewartson I equation. *Phys Revi E* 2012;86:036604.
- [36] Ohta Y, Yang J. Dynamics of rogue waves in the Davey-Stewartson II equation. *J Phys A Math Theor* 2013;46:105202.
- [37] Shi Y, Zhang Y. Rogue waves of a (3+1)-dimensional nonlinear evolution equation. *Commun Nonlinear Sci Numer Simulat* 2017;44:120-129.
- [38] Hirota R. *The direct method in soliton theory*. Cambridge: Cambridge Univ. Press; 2004.
- [39] Chakravarty S, Kodama Y. Classification of the line-soliton solutions of KP II. *J Phys A Math Theor* 2008;41:275209.
- [40] Chakravarty S, Kodama Y. Soliton solutions of the KP equation and application to shallow water waves. *Stud Appl Math* 2009;123:83-151.
- [41] Kodama Y. KP solitons in shallow water. *J Phys A Math Theor* 2010;43:434004.
- [42] Chakravarty S, Lewkow T, Maruno KI. On the construction of the KP line-solitons and their interactions. *Applicable Analysis* 2010;89:529-545.
- [43] Zhang Y, Guo L, Xu S, Wu Z, He J. The hierarchy of higher order solutions of the derivative nonlinear Schrödinger equation. *Commun Nonlinear Sci Numer Simulat* 2014;19:1706-1722.
- [44] Yuan F, Qiu D, Liu W, Porsezian, He J. On the evolution of a rogue wave along the orthogonal direction of the (t, x) -plane. *Commun Nonlinear Sci Numer Simulat* 2017;44:245-257.
- [45] Wen XY, Yan Z. Higher-order rational solitons and rogue-like wave solutions of the (2+1)-dimensional nonlinear fluid mechanics equations. *Commun Nonlinear Sci Numer Simulat* 2017;43:311-329.