

ALMA REVEALS METALS YET NO DUST WITHIN MULTIPLE COMPONENTS IN CR7

J. MATTHEE,¹ D. SOBRAL,^{2,1} F. BOONE,³ H. RÖTTGERING,¹ D. SCHAEERER,^{4,3} M. GIRARD,⁴ A. PALLOTTINI,^{5,6,7,8}
L. VALLINI,⁹ A. FERRARA,^{8,10} B. DARVISH,¹¹ AND B. MOBASHER¹²

¹*Leiden Observatory, Leiden University, PO Box 9513, NL-2300 RA, Leiden, The Netherlands*

²*Department of Physics, Lancaster University, Lancaster, LA1 4YB, UK*

³*Université de Toulouse; UPS-OMP; IRAP; Toulouse, France*

⁴*Observatoire de Genève, Université Genève, 51 Ch. des Maillettes, 1290 Versoix, Switzerland*

⁵*Centro Fermi, Museo Storico della Fisica e Centro Studi e Ricerche “Enrico Fermi”, Piazza del Viminale 1, Roma, 00184, Italy*

⁶*Cavendish Laboratory, University of Cambridge, 19 J. J. Thomson Ave., Cambridge CB3 0HE, UK*

⁷*Kavli Institute for Cosmology, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK*

⁸*Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy*

⁹*Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden*

¹⁰*Kavli IPMU, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa 277-8583, Japan*

¹¹*Cahill Center for Astrophysics, California Institute of Technology, 1216 East California Boulevard, Pasadena, CA 91125, USA*

¹²*Department of Physics and Astronomy, University of California, Riverside, CA 92521, USA*

ABSTRACT

We present spectroscopic follow-up observations of CR7 with ALMA, targeted at constraining the infrared (IR) continuum and [CII]_{158 μ m} line-emission at high spatial resolution matched to the *HST*/WFC3 imaging. CR7 is a luminous Ly α emitting galaxy at $z = 6.6$ that consists of three separated UV-continuum components. Our observations reveal several well-separated components of [CII] emission. The two most luminous components in [CII] coincide with the brightest UV components (A and B), blue-shifted by ≈ 150 km s⁻¹ with respect to the peak of Ly α emission. Other [CII] components are observed close to UV clumps B and C and are blue-shifted by ≈ 300 and ≈ 80 km s⁻¹ with respect to the systemic redshift. We do not detect FIR continuum emission due to dust with a 3σ limiting luminosity $L_{\text{IR}}(T_d = 35 \text{ K}) < 3.1 \times 10^{10} L_{\odot}$. This allows us to mitigate uncertainties in the dust-corrected SFR and derive SFRs for the three UV clumps A, B and C of 28, 5 and 7 M $_{\odot}$ yr⁻¹. All clumps have [CII] luminosities consistent within the scatter observed in the local relation between SFR and L_[CII], implying that strong Ly α emission does not necessarily anti-correlate with [CII] luminosity. Combining our measurements with the literature, we show that galaxies with blue UV slopes have weaker [CII] emission at fixed SFR, potentially due to their lower metallicities and/or higher photoionisation. Comparison with hydrodynamical simulations suggests that CR7’s clumps have metallicities of $0.1 < Z/Z_{\odot} < 0.2$. The observed ISM structure of CR7 indicates that we are likely witnessing the build up of a central galaxy in the early Universe through complex accretion of satellites.

Keywords: galaxies: formation — galaxies: high-redshift — galaxies: ISM — galaxies: kinematics and dynamics — dark ages, reionization, first stars

1. INTRODUCTION

Characterising the properties of the interstellar medium (ISM) of the first generations of galaxies is one of the prime goals of observational astrophysics. With the advent of the Atacama Large Millimetre Array (ALMA), direct measurements of the ISM are now becoming possible for typical star-forming galaxies in the early Universe ($z > 6$, e.g. [Maiolino et al. 2015](#); [Watson et al. 2015](#); [Knudsen et al. 2017](#)), in addition to studies of bright quasar hosts and galaxies with extreme bursts of star-formation (e.g. [Swinbank et al. 2012](#); [Riechers et al. 2013](#); [Wang et al. 2013](#); [Decarli et al. 2017](#); [Riechers et al. 2017](#)). These measurements are very valuable in constraining models of early galaxy formation (e.g. [Ceverino et al. 2010](#); [Hopkins et al. 2014](#); [Pallottini et al. 2017b](#)).

Rest-frame far-infrared continuum measurements (redshifted to sub-millimetre wavelengths detectable by ALMA) can provide a direct determination of the dust mass, temperature and attenuation (e.g. [da Cunha et al. 2015](#); [Bouwens et al. 2016](#); [Scoville et al. 2016, 2017](#)), and constrain the rate at which dust has been produced (e.g. [Hirashita et al. 2014](#); [Michałowski 2015](#); [Mancini et al. 2015, 2016](#)). Combining measurements of emission lines in the far-infrared such as the fine-structure lines $[\text{CII}]_{158\mu\text{m}}$, $[\text{NII}]_{205\mu\text{m}}$ and $[\text{OIII}]_{88\mu\text{m}}$ allows to constrain the star formation rate (SFR), metallicity, density and ionisation state of the gas, both at low redshift (e.g. [Ferkinnhoff et al. 2010](#); [De Looze et al. 2014](#); [Herrera-Camus et al. 2015](#)) and at high-redshift (e.g. [Inoue et al. 2016](#); [Carniani et al. 2017](#)). Furthermore, spatially resolved emission-line measurements can probe the dynamical structure of the gas (e.g. [Jones et al. 2017](#); [Smit et al. 2017](#)).

Besides quasar-hosts, dust continuum at $z > 6$ has been detected in a few sources ([Watson et al. 2015](#); [Laporte et al. 2017b](#)), but most sources show very little dust (e.g. [Schaerer et al. 2015](#); [Bouwens et al. 2016](#)). Observations of the $[\text{CII}]$ fine-structure cooling line indicate a large scatter in $[\text{CII}]$ luminosities at fixed SFR, particularly when compared to estimates based on the UV (e.g. [De Looze et al. 2014](#)). While some sources have similar $[\text{CII}]$ luminosities as those on the local relation between SFR(UV) and $L_{[\text{CII}]}$, (e.g. [Capak et al. 2015](#); [Smit et al. 2017](#); [Jones et al. 2017](#)), others have significantly fainter $[\text{CII}]$ luminosities at fixed UV SFR (e.g. [Ouchi et al. 2013](#); [Ota et al. 2014](#)). A potential explanation is that this is due to a selection bias. A large number of early sources observed by the ALMA were selected based on their $\text{Ly}\alpha$ emission needed to measure redshifts. Strong $\text{Ly}\alpha$ emission is typically associated with a high ionisation state and/or lower metallicity

(e.g. [Nakajima et al. 2016](#); [Trainor et al. 2016](#); [Matthee et al. 2017a](#); [Stark et al. 2017](#)), which should result in a deficit in $[\text{CII}]$ luminosity due to photo-dissociation (e.g. [Vallini et al. 2015](#)). This can be tested with deeper $[\text{CII}]$ observations of sources that span a wider parameter space (e.g. [Knudsen et al. 2017](#); [Bradač et al. 2017](#)).

Here we present deep spectroscopic observations of the COSMOS Redshift 7 galaxy (CR7) with ALMA, targeting the far-infrared dust continuum emission and the $[\text{CII}]$ line. CR7 has been identified as the most luminous $\text{Ly}\alpha$ emitter (LAE) at $z = 6.6$ based on narrow-band imaging with Subaru/Suprime-Cam ([Matthee et al. 2015](#)), with $L_{\text{Ly}\alpha} = 8.5 \times 10^{43} \text{ erg s}^{-1}$ and $\text{EW}_{0,\text{Ly}\alpha} = 210 \text{ \AA}$ ([Sobral et al. 2015](#)). Independently, [Bowler et al. \(2012, 2014\)](#) identified CR7 as a candidate luminous Lyman-break galaxy at $z \sim 6 - 7$ with ground-based near-infrared imaging. Hence, CR7 is also among the most UV-luminous galaxies known at $z \sim 7$, with $M_{1500} = -22.2 \pm 0.1$ (see [Matthee et al. 2017b](#) for even more luminous ones).

Besides strong $\text{Ly}\alpha$ emission, [Sobral et al. \(2015\)](#) identified a narrow HeII emission-line in near-infrared spectroscopic follow-up observations, and obtained limits on UV metal lines such as CIV and CIII]. *HST*/WFC3 near-infrared imaging revealed three separate rest-frame UV components, of which the brightest component (A) is closest to the peak of $\text{Ly}\alpha$ surface brightness on which spectroscopic observations were centred. The other two components (B and C) have photometric redshifts consistent with $z \gtrsim 6.5$.

These properties led to spectacular interpretations; clump A could contain low metallicity, hot ($T_{\text{eff}} \gtrsim 10^5 \text{K}$) PopIII-like stars (e.g. [Sobral et al. 2015](#); [Visbal et al. 2016](#)) or a direct collapse black hole (e.g. [Pallottini et al. 2015](#); [Dijkstra et al. 2016](#); [Hartwig et al. 2016](#); [Agarwal et al. 2016](#); [Smidt et al. 2016](#); [Smith et al. 2016](#); [Agarwal et al. 2017](#)). However, in independent analyses, [Bowler et al. \(2017\)](#) presented evidence of the possible presence of $[\text{OIII}]$ line-emission inferred from deblended *Spitzer*/IRAC photometry. While the [Bowler et al. \(2017\)](#) measurements are still consistent with a direct collapse or low metallicity AGN (e.g. [Agarwal et al. 2017](#); [Pacucci et al. 2017](#)), the lower claimed significance of the HeII line by [Shibuya et al. \(2017\)](#) removes most of the evidence for an AGN. As presented in detail in [Sobral et al. \(2017a\)](#), a re-analysis of old spectra and new near-infrared spectroscopy with the *HST*/WFC3 grism shows that the HeII line is indeed at lower significance. Moreover, if present, it is at lower luminosity (by a factor ≈ 3) and does not spatially coincide with clump A, but rather is emitted in the direction of clump C. No UV metal lines are detected in clump A. This points

towards a moderately low metallicity galaxy that is actively forming stars, without clear evidence for AGN activity (but see further details in Sobral et al. 2017a).

In this paper, we answer to the question of the presence of metals in CR7, and test whether the metallicity and dust content varies between different UV components. We summarise the UV properties of CR7 in §2. ALMA observations, data reduction, astrometry and sensitivity are discussed in §3. We investigate the resolved [CII] emission in §4. Dust continuum measurements and their implications for the SFRs are detailed in §5. In §6, we discuss where the different clumps of CR7 are located in the SFR- $L_{\text{[CII]}}$ relation, and how this compares with other sources. We discuss the implication of our results in light of recent simulations in §7, where we also use these observations to update the interpretation of the nature of CR7. The conclusions of this work are summarised in §8. Throughout the paper, we assume a Λ CDM cosmology with $\Omega_M = 0.70$, $\Omega_\Lambda = 0.30$ and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, and assume a Salpeter (1955) initial mass function (IMF) with mass limits $0.1 M_\odot$ and $100 M_\odot$.

2. UV PROPERTIES OF CR7

In the rest-frame UV, CR7 consists of three clumps (A, B and C, Sobral et al. 2015), of which the brightest (A) coincides with the peak of $\text{Ly}\alpha$ emission, and is also spectroscopically confirmed at $z_{\text{Ly}\alpha} = 6.604$. UV components are separated by $\sim 1''$, corresponding to projected distances of $\sim 5 \text{ kpc}$. The latest photometry on the *HST*/WFC3 imaging in the F110W and F160W filters has been performed by Bowler et al. (2017). We use this photometry to compute UV slopes of the three different clumps individually. The contribution of $\text{Ly}\alpha$ to the flux observed in the F110W filter is based on our $\text{Ly}\alpha$ narrow-band imaging as follows: we first correct the Subaru/S-Cam NB921 image for the contribution from the UV continuum by subtracting the z' image (which is calibrated such that a colour of $z' - \text{NB921} = 0$ corresponds to a line flux of zero, see Matthee et al. 2015) and then use this $\text{Ly}\alpha$ image to measure the $\text{Ly}\alpha$ flux at the positions of clumps A, B and C. The $\text{Ly}\alpha$ flux is multiplied by a factor two to take into account that the NB921 filter transmission is 50 % at the wavelength of CR7's $\text{Ly}\alpha$. We then convolve the apertures that have been used for *HST* photometry ($1''$, $0.4''$, $0.4''$, for clumps A, B and C respectively) with the PSF of the NB921 imaging ($0.6''$) and measure the $\text{Ly}\alpha$ flux in those PSF-convolved apertures. We measure $\text{Ly}\alpha$ fluxes of $8.3, 2.7, 1.3 \times 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2}$ in these apertures. Correcting the flux-density observed in the F110W for this line-flux contribution results in corrections of $+0.15,$




Figure 1. ALMA [CII] narrow-band image of CR7, collapsed for frequencies with -614 to $+96 \text{ km s}^{-1}$ with respect to the $\text{Ly}\alpha$ reference frame ($z = 6.604$). The image ($54 \times 54 \text{ kpc}$) is centred on the peak of $\text{Ly}\alpha$ position and the ALMA astrometry is shifted by $0.25''$ to the west to align it with *HST* imaging as described in §3.2. Contours are shown at the $3, 4, 5\sigma$ significance levels (with $1\sigma = 0.016 \text{ Jy beam}^{-1} \text{ km s}^{-1}$). Negative contours are shown as dashed lines. The beam size is shown in the bottom left corner and has FWHM $0.82'' \times 0.77''$. The red crosses mark the positions of the UV clumps of CR7. [CII] is clearly detected in CR7, indicating the presence of carbon.

$+0.16$ and $+0.05$ to the F110W magnitude (and M_{1500}) for clumps A, B and C, respectively. We measure UV slopes of $\beta = -2.3 \pm 0.4, -1.0 \pm 1.0$ and -2.3 ± 0.8 , for the three clumps respectively (see Table 1).¹ Due to the shallow depth of the observations, the UV slopes of clumps B and C are only poorly constrained, and deeper observations particularly in the F160W filter are required to improve them.

Based on the rest-frame UV luminosities and UV slopes, we derive SFR_{UV} , where dust attenuation is estimated using the Meurer et al. (1999) attenuation law. We take the measurement uncertainties in the rest-UV magnitudes into account by perturbing these values

¹ Without correcting the F110W photometry for the contribution from $\text{Ly}\alpha$, we would obtain UV slopes of $\beta = -2.8 \pm 0.4, -1.5 \pm 1.0$ and -2.4 ± 0.8 for clump A, B and C. While the errors are still dominated by measurement uncertainty, systematic uncertainties due to ignoring potentially important contributions from $\text{Ly}\alpha$ could bias UV slope measurements to artificially bluer values.

10,000 times (assuming the uncertainties are gaussian). In each realisation, we re-compute the UV luminosity and UV slope, and use these to derive dust-corrected SFR. We then obtain the median and the 1σ percentiles and use these to derive the asymmetric uncertainties. With this method, the SFRs are 29_{-2}^{+23} , 38_{-32}^{+182} and 7_{-1}^{+19} $M_{\odot} \text{ yr}^{-1}$ for clumps A, B and C, respectively. The uncertainties in the dust corrected SFRs are large due to the propagation of errors in β . However, as we show in §5, these uncertainties are mitigated by constraints on the IR luminosity from our deep ALMA observations, which place firm limits on the dust-obscured SFR. Because of these constraints, our final results would also only change marginally if an SMC-like attenuation law is used. The results are listed in Table 1. The F110W imaging has a 3σ sensitivity of 27.3 AB magnitude in a $0.4''$ diameter, which corresponds to a limiting UV magnitude $M_{1500} > -19.5$ and a $\text{SFR}_{\text{UV}} < 4 M_{\odot} \text{ yr}^{-1}$.

3. ALMA DATA

3.1. Observations & data reduction

We observed CR7 (10:00:58.00 +01:48:15.3, J2000) in band 6 with ALMA during cycle 3 with configuration C43-4, aimed to achieve a $0.3''$ angular resolution (program ID #2015.1.00122.S). The target has been observed with between 35-43 antennas for a total on source integration time of 6.0h on 22, 23, 24 May and 4, 8 November 2016, with precipitable water vapor ranging from 0.3-1.4 mm. We observed in four spectral windows centred at 249.94592, 247.94636, 234.94917 and 232.9496 GHz with a bandwidth of 1875 MHz, of which the first is centred at [CII] line emission ($\nu_{z=6.604} = 249.9395$ GHz). The central frequencies of the spectral windows correspond to rest-frame frequencies between 1771.7 GHz and 1901.0 GHz, with a velocity resolution of 19.4 km s^{-1} . The phase calibrations have been performed on the source J0948+0022. The quasar J1058+0133 has been observed regularly as a bandpass and flux calibrator, resulting in typical flux calibration errors of $\sim 10\%$. Data have been reduced using CASA version 4.7.0 (McMullin et al. 2007) with natural weighting and channel averaging resulting in a 38.8 km s^{-1} velocity resolution. We use uv tapering of the visibilities with a Gaussian width FWHM= $0.7''$ to optimise the S/N ratio. The observational set-up and data reduction results in a beam size of $0.35 \times 0.33''$ (gaussian σ ; $0.82 \times 0.77''$ FWHM), with a position angle of -49.5° . We have also performed the reduction with uv tapering with different smoothing kernels ranging from $0.4 - 0.8''$, but find that all changes in luminosity, line-width and size measurements are consistent within the error-bars. A simple estimate of the noise level of the pixels within




Figure 2. Integrated spectrum over the area within the 3σ [CII] contours in Fig. 1, in two observed spectral windows. The grey shaded region indicates the 1σ error estimated using apertures with the same geometry as the 3σ contour area, as a function of frequency. The blue dashed line indicates where [CII] would be detected at $z = 6.604$, the Ly α redshift of CR7. Ly α is redshifted by $\approx -170 \text{ km s}^{-1}$ with respect to the systemic [CII] redshift.

a radius of $20''$ of the center results in $\text{rms}=0.06 \text{ mJy beam}^{-1}$ in 38.8 km s^{-1} channels.

3.2. Astrometry

As described in §4, we detect two separate clumps at high significance in the [CII] image collapsed over all velocities at which significant line-emission is detected, see Fig. 1. These [CII] clumps are offset by $0.25''$ to the east with respect to the UV positions of clumps A and B, while the offset in declination is $< 0.10''$, respectively. The alignment of both clumps, and their separation is also similar in the UV and in [CII]. As these two [CII] clumps resemble the geometry of clumps A and B, we assume that the offsets between the UV positions and [CII] positions are due to errors in the astrometry. This offset is similar in magnitude as reported in e.g. Dunlop et al. (2017), but in the R.A. direction instead of the Dec. direction (see also Carniani et al. 2017 and references therein for offsets in $z \sim 6 - 7$ galaxies). For the rest of the analysis, we align the ALMA data with *HST* and Subaru narrow-band data by applying an offset of $0.25''$ to the west. We note that the positions of two serendipitous IR continuum-detections of foreground galaxies are also in agreement with this offset (see §5.1 and Fig. 8).

3.3. Line-sensitivity measurements

As the noise is correlated spatially and spectrally, we measure the sensitivity of the observations and significance of detections as follows. In each collapsed image




Figure 3. Histogram of the integrated [CII] fluxes measured over circular, beam-sized scales at random positions in the collapsed cube (with velocities from -614 to $+96$ km s^{-1} with respect to $\text{Ly}\alpha$). The blue histogram shows positive fluxes, while the red histogram shows negative fluxes. We indicate the [CII] flux measured at the positions of *HST* clumps A, B and C with vertical lines. These correspond to $\approx 5, 4, 3 \sigma$ detections, respectively. The grey curve shows the expected noise properties if the noise is gaussian with a 1σ dispersion. Note that our final measurement of clump A uses a larger aperture (with radius $0.55''$), resulting in a similar significance but higher flux. Also note that the detections are at slightly higher significance in narrower-velocity slices optimised for each clumps specifically.

(which has a size of $51.2'' \times 51.2''$, centred on CR7), we measure the noise over the same spatial scales as those used for measurements (typically a beam-size, $r = 0.35''$). We place circular apertures on 100,000 random positions on the image and integrate the flux over the scales confined by these apertures. As the image size is much larger than the source size, we do not mask any central region. We then compute the 1σ detection significance using the r.m.s., which results in 16.9 $\text{mJy beam}^{-1} \text{ km s}^{-1}$ for beam-sized apertures on the collapsed image shown in Fig. 1.

As illustrated in Fig. 3, the distribution of fluxes measured on random positions approximately follows a gaussian. It can be seen that the negative image (where we have inverted all counts) has slightly lower number counts than the positive image at high flux levels. This is due to random positions around the centre of CR7, or real sources around CR7. These sources (such as a $\sim 4\sigma$ detection $4''$ from the south-east of CR7, see Fig. 1) are discussed and investigated in a future paper. When integrating over a larger spatial scale, the noise estimate is slightly higher due to spatially correlated noise. The noise level is lower when data is collapsed over a smaller velocity range. In individual veloc-

ity channels (with resolution 38.8 km s^{-1}), we measure a 1σ sensitivity between 30 - $40 \mu\text{Jy}$ when integrating flux over beam-sized scales. Throughout the paper, we convert measured fluxes to [CII] luminosities following e.g. Solomon et al. (1992) and Carilli & Walter (2013):

$$L_{[\text{CII}]} / L_{\odot} = 1.04 \times 10^{-3} \times S_{\nu} \Delta v \times \nu_{\text{obs}} \times D_{\text{L}}^2, \quad (1)$$

where $S_{\nu} \Delta v$ is the flux in Jy km s^{-1} , ν_{obs} the observed frequency in GHz (≈ 250.0 GHz for CR7) and D_{L} is the luminosity distance in Mpc (64457.8 Mpc at $z = 6.60$).

4. RESOLVED [CII] EMISSION

4.1. Total luminosity

In order to search for [CII] emission around CR7, we inspect the data-cube of the spectral tuning centered at the frequency where [CII] is expected to be observed. As shown in Fig. 1, resolved [CII] emission is clearly detected.² We show the collapsed data-cube with velocities between -614 and $+96 \text{ km s}^{-1}$ with respect to the $\text{Ly}\alpha$ redshift, which includes the full velocity range over which we detect [CII] emission. Within the 3σ significance contours, we measure $L_{[\text{CII}]} = 2.17 \pm 0.36 \times 10^8 L_{\odot}$ in the collapsed image.³ The spectrum extracted over this region is shown in Fig. 2 and is best-fitted with a line-width of $v_{\text{FWHM}} = 299 \pm 27 \text{ km s}^{-1}$ that is offset by $-167 \pm 27 \text{ km s}^{-1}$ with respect to the $\text{Ly}\alpha$ redshift, see Table 1. The 3σ significance contours span a total area of 2.2 arcsec^2 and have a maximum diameter of $1.8''$, which corresponds to 9.7 kpc at $z = 6.60$. This detection of [CII] emission in a large region in/around CR7 clearly indicates the presence of metals, ruling out the possibility that a large fraction of the gas is primordial.

4.2. Multiple separated [CII] components

As illustrated in Fig. 1, the [CII] emission is clearly not homogeneous over the 3σ contour and reveals already two local peaks, resembling the rest-frame UV morphology as observed by *HST* (see Fig. 5). In order to show structure in both the spatial and spectral dimensions, we create collapsed [CII] maps in three different velocity ranges of 100 km s^{-1} each and show these in Fig. 4. As the noise level varies slightly at different velocities and is smaller for images that are collapsed over a smaller velocity range, we compute the noise level in each image independently as described in §3.3.

² The individual [CII] channel maps with widths $\Delta v = 38.8 \text{ km s}^{-1}$ and frequencies from 249.83 - 250.42 GHz (corresponding to $z_{[\text{CII}]} = 6.607 - 6.589$) are shown in Fig. 12.

³ This error has been estimated by measuring the flux in randomly located apertures with the same geometry as the 3σ contours.




Figure 4. [CII] map in three velocity channels of 100 km s^{-1} , zoomed in on CR7 (with a scale of $32 \times 32 \text{ kpc}$) and with velocities with respect to the $\text{Ly}\alpha$ redshift of $z = 6.604$. Contours mark the 3, 4, 5 σ levels, with sensitivities $\sigma \approx 5 - 6 \text{ mJy beam}^{-1} \text{ km s}^{-1}$. Red crosses mark the positions of the UV clumps, while black crosses mark the positions of additional [CII] components B-2 and C-2. The most luminous [CII] components coincide with UV clumps A and B and have a similar peak velocity. Fainter [CII] flux is detected at $> 4\sigma$ in between clumps A, B and C offset by -120 km s^{-1} (clump C-2), and close to clump B at -290 km s^{-1} (clump B-2).




Figure 5. Rest-frame UV image (from *HST*/WFC3 F110W imaging) of CR7, overlaid with ALMA [CII] 3, 4, 5 σ contours at a scale of $54 \times 54 \text{ kpc}$. The names of the three UV clumps are annotated in HST imaging. Relatively compact components of [CII] emission are detected around clumps A and B, while there is diffuse [CII] emission covering clump C, although at lower significance.

The three velocity channels reveal that the [CII] emission in CR7 consists of multiple, well separated components. Around the systemic velocity (-130 km s^{-1} with respect to $\text{Ly}\alpha$), two sources are clearly detected at the UV positions of clumps A and B. At a blue-shift

of $\approx 250 \text{ km s}^{-1}$ a relatively faint, resolved component is detected in between the three UV clumps, but closest to clump C. No significant UV emission is detected at that ALMA position. We name this component C-2. Another compact sub-component is detected at an even larger blue-shift of $\approx 420 \text{ km s}^{-1}$ close to the position of clump B (well within the beam radius), and hence named B-2. In these narrower velocity slices, all detections are at the $\gtrsim 4\sigma$ significance level, see Fig. 4.

Hence, we detect several components of [CII] emission in the CR7 system that are significantly offset in both the spatial and spectral dimension. Before discussing the implications of these detections in §7.1, we first measure their luminosities and [CII] line-profiles.

4.2.1. Measurements of individual clumps

Except for clumps A and C-2, all clumps are unresolved (see §4.3). For unresolved sources, we measure the [CII] luminosity within a circular aperture-region with radius of $0.35''$ (corresponding to the beam semi-major axis). For clumps A and C-2, we use $0.55''$ and $0.50''$ apertures respectively to account for the fact that these sources are slightly extended. For these clumps, we find that these apertures retrieve the same fraction of the flux as is retrieved for modelled point sources with a $0.35''$ aperture. Measurements at *HST*/UV positions are done on the collapsed image over the full velocity range (e.g. Fig. 1), while measurements at ALMA positions (B-2 and C-2) are done in the collapsed velocity range in which these clumps are detected (e.g. Fig. 4).

In Fig. 6 we show the ALMA spectra of the different clumps (extracted over their respective apertures) in the rest-frame of clump A ($z = 6.601$) and compare these

to the Ly α profile measured with VLT/X-SHOOTER at the peak of Ly α emission (close to clump A). We measure the velocity offset between [CII] and Ly α and the [CII] line-width and amplitude by fitting gaussian profiles. All measurements are summarised in Table 1.

We find that clump A, which is the brightest in the UV and closest to the peak of Ly α emission, is also the brightest in [CII] line-emission, with $L_{[\text{CII}]} = (0.96 \pm 0.20) \times 10^8 L_{\odot}$. We caution that we measure a factor two lower luminosity if we measure the luminosity of clump A with a beam-sized aperture. Clump A is blue-shifted by $146 \pm 27 \text{ km s}^{-1}$ with respect to Ly α . This is slightly smaller than typical offsets between Ly α and nebular lines for sources with similar UV luminosities (e.g. Stark et al. 2017), although Carniani et al. (2017) shows a wide spread of velocity offsets in a compilation of ALMA detections (i.e. $-150 < \Delta v < +400 \text{ km s}^{-1}$ for galaxies with $10 < \text{SFR} < 100 M_{\odot} \text{ yr}^{-1}$). The [CII] line-width is $259 \pm 24 \text{ km s}^{-1}$, similar to the Ly α line-width (see e.g. Fig. 6).

At the UV position of clump B, we measure [CII], with $L_{[\text{CII}]} = (0.34 \pm 0.09) \times 10^8 L_{\odot}$, which contains the contribution from both the ALMA detections B and B-2 (with $\approx 33\%$ of the flux being due to the sub-component B-2, see Table 1). The brightest [CII] component of clump B is at $z = 6.600 \pm 0.001$, similar to the redshift of clump A, but has a narrower line-width. Clump B-2 is blue-shifted by $290 \pm 30 \text{ km s}^{-1}$ with respect to the systemic redshift and has a relatively narrow-line width of $92 \pm 22 \text{ km s}^{-1}$.

At the position of clump C, we measure relatively faint [CII] emission at the $\approx 3\sigma$ level, see Fig. 1 and Fig. 5. The aperture at the position of clump C is contaminated slightly by this clump B-2 (which may also be associated with C), explaining the bluest line, but the redder [CII] component C-2 is offset spatially (see Fig. 4). At the position of C-2, we measure $L_{[\text{CII}]} = (0.27 \pm 0.07) \times 10^8 L_{\odot}$ with a line-width of $181 \pm 30 \text{ km s}^{-1}$ and an offset of $-232 \pm 27 \text{ km s}^{-1}$ with respect to Ly α . This offset is bluer than clumps A and B, but redder than B-2. Due to its large blue-shift, component C-2 also dominates the velocity map shown in Fig. 7. Due to this strong contribution of component C-2 to the velocity map, we find that there is no clear evidence for ordered rotation over the full [CII] extent. This shows that the spatial resolution of our observations is crucial, as we may have not been able to distinguish clumps C-2 and B/B-2 with lower resolution, and could have mis-interpreted the velocity map as rotation. It is challenging to investigate whether ordered rotation is present inside individual clumps because they are either unresolved or only marginally resolved. Similarly as for clump A, we




Figure 6. Flux densities of Ly α (from X-SHOOTER, centered at clump A) and [CII] at the HST positions of clumps A, B and C as a function of velocity with respect to the Ly α peak velocity at $z = 6.604$. We also measure [CII] line-profiles at the positions B-2 and C-2 (shown in green), based on [CII] detections, see Fig. 4. Gaussian fits are shown in black, and we highlight the central velocities with vertical dotted lines. Errors have been estimated for each velocity-slice and aperture as described in §3.3.

Table 1. Properties of different components in CR7. Clumps B-2 and C-2 are based on [CII] detections, while A, B and C are based on *HST* detections (see §4.2.1). The ‘Full’ uses ground-based imaging and the 3σ contour levels from Fig. 1. $\Delta R.A._{HST}$ and $\Delta Dec._{HST}$ are the difference between the *HST* and ALMA positions before shifting the ALMA R.A. by $0.25''$ to the east. For B-2 and C-2 the differences are between *HST* clumps B and C, and ALMA detections B-2 and C-2, respectively. M_{1500} is estimated from F110W imaging after correcting for the contribution of Ly α emission. SFRs assume a Salpeter IMF following Kennicutt (1998). Dust-attenuation of the UV SFR is estimated based on the UV slope β following Meurer et al. (1999). We provide [CII] luminosities measured from the collapsed image in a beam-sized aperture and [CII] luminosities obtained from the fitted gaussian line-profiles. Dynamical masses are estimated following Wang et al. (2013).

| | Clump A | Clump B | Clump B-2 | Clump C | Clump C-2 | Full |
|---|-------------------|-------------------|-------------------|-----------------|---------------------|-------------------|
| $\Delta R.A._{HST}$ | $-0.25''$ | $-0.24''$ | $-0.07''$ | - | $+0.14''$ | - |
| $\Delta Dec._{HST}$ | $-0.12''$ | $-0.18''$ | $-0.04''$ | - | $+0.40''$ | - |
| M_{1500} | -21.6 ± 0.1 | -19.6 ± 0.2 | - | -20.1 ± 0.1 | > -19.5 | -22.2 ± 0.1 |
| β | -2.3 ± 0.4 | -1.0 ± 1.0 | - | -2.3 ± 0.8 | - | -2.2 ± 0.4 |
| $SFR_{UV, \text{nodust}}/M_{\odot} \text{ yr}^{-1}$ | 27 ± 1 | 4 ± 1 | - | 7 ± 1 | < 4 | 44 ± 2 |
| $SFR_{UV, \text{Meurer}}/M_{\odot} \text{ yr}^{-1}$ | 29^{+23}_{-2} | 38^{+182}_{-32} | - | 7^{+19}_{-1} | - | 50^{+55}_{-6} |
| $SFR_{UV+IR}/M_{\odot} \text{ yr}^{-1}$ | 28^{+1}_{-1} | 5^{+2}_{-1} | - | 7^{+1}_{-1} | < 4 | 45^{+2}_{-2} |
| $\Delta v_{Ly\alpha}/\text{km s}^{-1}$ | -146 ± 27 | -152 ± 21 | -422 ± 22 | - | -232 ± 27 | -167 ± 27 |
| $z_{[CII]}$ | 6.601 ± 0.001 | 6.600 ± 0.001 | 6.593 ± 0.001 | - | 6.598 ± 0.001 | 6.600 ± 0.001 |
| $v_{FWHM, [CII]}/\text{km s}^{-1}$ | 259 ± 24 | 130 ± 21 | 92 ± 22 | - | 181 ± 30 | 299 ± 26 |
| $S_{\nu, \text{fit}} \Delta v / \text{mJy km s}^{-1}$ | 81.1 ± 20.7 | 25.0 ± 5.3 | 10.0 ± 3.9 | - | 35.4 ± 10.9 | 185.2 ± 39.0 |
| $L_{[CII], \text{fit}}/10^8 L_{\odot}$ | 0.88 ± 0.23 | 0.26 ± 0.07 | 0.12 ± 0.03 | - | 0.27 ± 0.11 | 2.00 ± 0.43 |
| $L_{[CII], \text{aperture}}/10^8 L_{\odot}$ | 0.96 ± 0.20 | 0.34 ± 0.09 | 0.11 ± 0.03 | 0.26 ± 0.09 | 0.27 ± 0.07 | 2.17 ± 0.36 |
| $r_{1/2, [CII]}/\text{kpc}$ | 3.0 ± 0.8 | < 2.2 | < 2.2 | - | $3.8^{+1.1}_{-0.7}$ | - |
| $M_{\text{dyn}}/(\sin i)^2 10^{10} M_{\odot}$ | 3.9 ± 1.7 | < 0.7 | < 0.4 | - | 2.4 ± 1.9 | - |
| $L_{UV}/10^{10} L_{\odot}$ | 9.1 | 1.4 | - | 2.3 | < 1.3 | 15.8 |
| $\text{rms}_{\text{cont}, \lambda_0=160\mu\text{m}}/\mu\text{Jy beam}^{-1}$ | 7 | 7 | 7 | 7 | 7 | 7 |
| $L_{IR} (T_d = 35\text{K})/10^{10} L_{\odot}$ | < 3.14 | < 3.14 | < 3.14 | < 3.14 | < 3.14 | < 3.14 |
| $SFR_{IR} (T_d = 35\text{K})/M_{\odot} \text{ yr}^{-1}$ | < 5.4 | < 5.4 | < 5.4 | < 5.4 | < 5.4 | < 5.4 |
| $M_{\text{dust}}(T_d = 35\text{K})/10^6 M_{\odot}$ | < 8.1 | < 8.1 | < 8.1 | < 8.1 | < 8.1 | < 8.1 |

note that the [CII] luminosity of clump C-2 would be reduced by a factor ≈ 2 if we use a smaller aperture of $0.35''$, which is used for unresolved sources. Therefore, the luminosities of clumps A and C-2 should be interpreted with caution. The total [CII] luminosity that is associated to clumps is $(1.57 \pm 0.23) \times 10^8 L_{\odot}$, which is the sum of the aperture measurements at the *HST* positions. This corresponds to a fraction of 0.72 ± 0.18 of the total observed [CII] luminosity.

4.3. Sizes and dynamical masses

Starting from the measured velocity width and size of the emitting regions, it is possible to obtain a rough estimate of the dynamical mass of the various [CII] components (e.g. Wang et al. 2013). First, we measure the size by performing two dimensional gaussian fits in the collapsed 2D image. Clumps A and B are simultane-

ously fitted in the image that is collapsed over the full velocity range, while clumps B-2 and C-2 are fitted in the collapsed image in the velocity range as shown in Fig. 4. In our fitting procedure, we fixed the position angle and the ratio between the dispersion in both dimensions corresponding to the those of the beam semi-major and semi-minor axis. We also fix the positions to those indicated in Fig. 4. Hence, the free parameters are the amplitudes and widths of the gaussian profiles. In order to take the (correlated) noise into account, we add the noise from a randomly selected cut-out region in the collapsed image, avoiding the source itself, before performing the fit. This is repeated 5000 times in order to estimate the statistical reliability of the fit. We find that more than $> 50\%$ of the measurements for clumps B and B-2 result in a size that is equal to or smaller




Figure 7. [CII] velocity map (at a 22×22 kpc scale) in the rest-frame of clump A, based on the first moment map collapsed for frequencies with -614 to $+96$ km s^{-1} with respect to the Ly α redshift (see Fig. 1 for the corresponding [CII] flux map). Contours show the 3, 4 and 5 σ threshold and velocity maps are shown for $> 3\sigma$ detections. The map is driven by the the strongly blue-shifted component C-2.

than the beam, meaning that these clumps are unresolved and hence have a size $r_{1/2, [\text{CII}]} < 2.2$ kpc. We measure an observed size of $r_{1/2, [\text{CII}], \text{obs}} = 3.7^{+0.6}_{-0.6}$ kpc for clump A (4 % of the measurements result in an unresolved size, meaning that clump A is resolved at $\approx 2\sigma$ significance) and $r_{1/2, [\text{CII}], \text{obs}} = 4.4^{+0.9}_{-0.6}$ kpc for clump C-2 (resolved at $\approx 3.5\sigma$ significance). We deconvolve the sizes of A and C-2 to obtain the intrinsic size with $r_{\sigma, [\text{CII}]} = \sqrt{r_{\text{obs}}^2 - 2.2^2}$. Resulting sizes are listed in Table 1. Dynamical masses are computed following Wang et al. (2013):

$$M_{\text{dyn}}/M_{\odot} (\sin i)^2 = 1.94 \times 10^5 \times v_{\text{FWHM}, [\text{CII}]}^2 \times r_{1/2, [\text{CII}]}, \quad (2)$$

where M_{dyn} is the dynamical mass in M_{\odot} , i is the inclination angle, $v_{\text{FWHM}, [\text{CII}]}$ the line-width in km s^{-1} and $r_{1/2, [\text{CII}]}$ the size in kpc (half-light radius). This results in dynamical masses (uncorrected for inclination) ranging from $(3.9 \pm 1.7) \times 10^{10} M_{\odot}$ for component A, $(2.4 \pm 1.9) \times 10^{10} M_{\odot}$ for clump C-2 and $< 0.7 \times 10^{10} M_{\odot}$ for $< 0.4 \times 10^{10} M_{\odot}$ for clumps B and B-2, respectively. These dynamical mass estimates are lower than typical quasar host galaxies at $z \approx 6$ (Wang et al. 2013), and comparable to star-forming galaxies at $z \approx 7$ with




Figure 8. IR continuum map at 230-250 GHz centered on the position of CR7. The black contours show the 2, 3 σ level, where $1\sigma = 7\mu\text{Jy beam}^{-1}$. We also show *HST* rest-frame UV (F814W+F110W+F160W) contours at the 2, 3, 4 σ levels to highlight the positions of known (foreground) sources. Sources are annotated with their photometric redshifts estimated by Laigle et al. (2016). No dust continuum is detected at CR7, although a $\approx 2\sigma$ signal is detected around potential clump D. Dust continuum is also clearly detected in a foreground source at $z = 0.84$.

similar SFRs as CR7 (e.g. Pentericci et al. 2016; Smit et al. 2017).

5. IR CONTINUUM

5.1. Blind detections

We combine the flux in all four spectral windows from our ALMA coverage to search for dust continuum emission. In the entire image, we find two detections with $S/N > 3$, but they are not associated with CR7. One detection is $3.5''$ north-east of CR7 (associated with ID number 339509 in the catalog from Laigle et al. 2016, $photo-z = 0.84$ and visible in Fig. 8), while the other is $18.5''$ to the south-west (ID number 335753 in Laigle et al. 2016, $photo-z = 3.10$, not visible in the image). The positions of these foreground galaxies confirm the astrometric correction described in §3.2. We note that we detect a tentative (3σ) line at 250.484 GHz at the position of ID 339509 that is identified as CO(4-3) with $\nu_0 = 461.041$ GHz at $z = 0.841$, perfectly consistent with its photometric redshift.

As visible in Fig. 8, there is a $\approx 2.7\sigma$ continuum detection $\approx 2''$ to the south-east of CR7, nearby a faint

HST detection that we name potential clump D. This faint *HST* detection has $F110W=27.2 \pm 0.2$ in a $0.4''$ aperture, similar to clump B. It is also detected at $\approx 3\sigma$ in the ACS/F814W filter ($F814W=27.5 \pm 0.2$) and F160W filter ($F160W=27.0 \pm 0.3$). It is not detected in NB921 and z' imaging, which have 3σ depths of 25.8 AB magnitude, hence consistent with the *HST* photometry. Potential clump D is not detected in the Subaru S-cam imaging in the B, V, R filters with 2σ upper limits of $\lesssim 28.5-29.0$. Therefore, the photometry is consistent with the Lyman-break falling in the F814W filter, such that $4.8 < z_{\text{phot}} < 6.9$. There is no significant [CII] detection at its position in any of the spectral tunings observed with ALMA. Further IFU observations are required to measure the redshift of clump D and test whether it is associated to CR7.

5.2. Upper limits for CR7

There is no IR continuum detected at the position of CR7's clumps A, B and C, see Fig. 8. Around the position of CR7, we measure an rms of $7 \mu\text{Jy beam}^{-1}$ at $\lambda_0 \approx 160 \mu\text{m}$. Note that we do not remove contamination from the [CII] line as we expect that this is negligible, as a result our upper limit is therefore somewhat on the conservative side. We follow Schaerer et al. (2015) to convert this limit into a 3σ upper limit on the infrared luminosity (between 8-1000 μm) of $L_{\text{IR}} < 3.14 \times 10^{10} L_{\odot}$, under the assumption that the dust temperature is 35 K and a modified black body SED with a power-law slope of 2.9 in the Wien regime, $\beta_{\text{IR}} = 1.5$ and after removing the contribution of the CMB to the dust heating, see da Cunha et al. (2013). In the case that the dust temperature is 25 (45) K, the limit on the IR luminosity is $L_{\text{IR}} < 1.68(6.15) \times 10^{10} L_{\odot}$. Combined with the total [CII] luminosity, we find a lower limit of the the ratio $\log_{10}(L_{[\text{CII}]} / L_{\text{IR}}) > -2.1$, which is similar to the highest values measured in local star-forming galaxies (e.g. Malhotra et al. 2001) and significantly higher than sub-mm galaxies and quasar host galaxies at $z > 3$ (see Ota et al. 2014 for a compilation). This could potentially indicate a lower dust-to-metal ratio, which in turn could result in a higher dust temperature due to the lower dust opacity (e.g. Cen & Kimm 2014).

The ratio of the $\text{IR}_{160\mu\text{m}}$ to $\text{UV}_{0.15\mu\text{m}}$ flux density is constrained to $(\nu F_{\nu})_{\text{IR}} / (\nu F_{\nu})_{\text{UV}} < 0.04$ (3σ). This limit is almost an order of magnitude stricter than other $z \sim 7$ sources observed by Maiolino et al. (2015). In the local Universe, this ratio is related to metallicity (see the compilation and fitted relation in Maiolino et al. 2015), likely as a result of dust content increasing with metallicity (e.g. Rémy-Ruyer et al. 2014). If such a relation would be unchanged at $z = 7$ (e.g. Popping

et al. 2017), the measurements imply a metallicity of $12 + \log(\text{O}/\text{H}) \lesssim 7.5$ (corresponding to $\lesssim 0.1 Z_{\odot}$), which is only found in dwarf galaxies in the local Universe with masses $M_{\text{star}} \lesssim 10^8 M_{\odot}$ (e.g. Izotov et al. 2015; Guseva et al. 2017).

We use the upper limits on the IR continuum to place an upper limit on the dust mass. We follow the method outlined in Ota et al. (2014), which assumes a dust mass absorption coefficient $k_{\nu} = 1.875(\nu/239.84)^{\beta_{\text{IR}}} \text{m}^2 \text{kg}^{-1}$ where ν is in GHz and $\beta_{\text{IR}} = 1.5$ and removes the contribution from the CMB. This results in $M_{\text{dust}} < 8.1(27.5) \times 10^6 M_{\odot}$ if the dust temperature is 35 (25) K. These limits are consistent with expectations for CR7's UV luminosity, based on post-processed hydrodynamical simulations by Mancini et al. (2016), who find dust masses between $\approx 1 - 10 \times 10^5 M_{\odot}$.

Following the prescription from Kennicutt (1998), we measure a 3σ upper limit of the dust obscured $\text{SFR}_{\text{IR}} < 5.4 M_{\odot} \text{yr}^{-1}$ for a Salpeter (1955) IMF. This allows us to put a stronger constraint on the total SFR of different clumps. Because of these tight limits, the large errors in β (which for example allow relatively red UV slopes and hence relatively large dust obscurations) are mitigated. Combining these measurements (by constraining that the difference between dust-corrected SFR and dust-free SFR is $< 5.4 M_{\odot} \text{yr}^{-1}$ in each of the 10,000 realisations used to self-consistently compute the uncertainties as described in §2) results in $\text{SFR}_{\text{UV+IR}} = 28_{-1}^{+1}, 5_{-1}^{+2}, 7_{-1}^{+1} M_{\odot} \text{yr}^{-1}$ for clump A, B and C, respectively. For the total (ground-based) photometry, we find $\text{SFR}_{\text{UV+IR}} = 45_{-2}^{+2} M_{\odot} \text{yr}^{-1}$, which is roughly consistent with the sum of the *HST* detected clumps and indicates no other significant source of star-formation.

6. THE SFR- $L_{[\text{CII}]}$ RELATION

Due to its luminosity, the [CII] fine-structure cooling line has been proposed as a SFR tracer (see e.g. De Looze et al. 2014 and references therein), which is observable with ALMA at high-redshift, while other tracers such as H α are currently unavailable. However, while [CII] is insensitive to dust attenuation, unlike (for example) UV emission, its luminosity is sensitive to metallicity (e.g. Vallini et al. 2013; Hemmati et al. 2017; Olsen et al. 2017). The [CII] luminosity is also dependent on the ionisation state of the gas and saturates at high temperatures (Kaufman et al. 1999; Decataldo et al. 2017; Vallini et al. 2017).

Early ALMA observations of galaxy samples at $z \approx 5-6$ (e.g. Capak et al. 2015; Willott et al. 2015) found that these galaxies have similar $L_{[\text{CII}]}$ to SFR(UV) ratios as local star-forming galaxies (De Looze et al. 2014). Galaxies at $z \approx 6-7$ were typically selected based on




Figure 9. Dust corrected SFR(UV) versus [CII] luminosity for a compilation of galaxies at $z \sim 6$ (blue circles; Capak et al. 2015; Willott et al. 2015) and $z \sim 7$ (green pentagons; Kanekar et al. 2013; Ouchi et al. 2013; Ota et al. 2014; Maiolino et al. 2015; Schaerer et al. 2015; Knudsen et al. 2016; Pentericci et al. 2016; Bradač et al. 2017; Knudsen et al. 2017; Smit et al. 2017). We show the individual UV components of CR7 with red stars, and the sum of these component as an orange star. The integrated measurement over the full [CII] emitting region around CR7 is shown with an open star. Downward pointing triangles show galaxies with upper limits on the [CII] luminosity, while left-ward pointing triangles show [CII] detections with upper limits on SFR(UV). The black line shows the relation for local star-forming galaxies measured by De Looze et al. (2014), where the grey region shows the observed dispersion. The dotted lines are model-predictions from Vallini et al. (2015) on how the [CII] luminosity varies with metallicity. We find that the clumps in CR7 are within the scatter from the local relation. These measurements indicate metallicities $0.1 < Z/Z_{\odot} < 0.2$. Note that we have corrected published SFRs (including those from the local relation) to the Salpeter IMF.

their Ly α redshifts and were initially observed at fainter [CII] luminosities or luminosity limits (e.g. Kanekar et al. 2013; Ouchi et al. 2013; Ota et al. 2014; Maiolino et al. 2015; Schaerer et al. 2015). This was later confirmed in observations of lensed galaxies (Knudsen et al. 2016; Bradač et al. 2017). However, sources that have been selected independently of strong Ly α emission show a wider range in [CII] luminosities (e.g. Pentericci et al. 2016; Knudsen et al. 2017; Smit et al. 2017), although some may still be relatively luminous in Ly α (e.g. Laporte et al. 2017a).

In Fig. 9, we show how our [CII] measurements of the UV components of CR7, their sum, and the integrated CR7 measurements compare to samples at $z \sim 5-7$ and to the local relation. As we use the aperture measurements centred at the UV clumps, the [CII] luminosity of clump B includes that of sub-component B-2. We also

show the upper limit of the UV SFR of clump C-2, if we do not associate its [CII] luminosity with the UV luminosity of clump C. Most sources that currently have been targeted are UV luminous (with SFR $\approx 20-30 M_{\odot} \text{ yr}^{-1}$, similar to clump A), while the two lensed sources have much lower SFRs.⁴ CR7’s clumps B and C lie roughly in the middle of the probed parameter space. All individual clumps in CR7, and the total luminosity are consistent within the observed scatter and their error-bars with the local relation, with only a marginal offset. This is in contrast with previously targeted luminous LAEs such as Himiko (Ouchi et al. 2013) and IOK-1 Ota et al. (2014).

⁴ Note that we have also rescaled all published SFRs, including the local relation from De Looze et al. (2014), to a Salpeter IMF.

Combining our measurements of clump A and CR7s total luminosity with sources from the literature, we confirm a large spread of an order of magnitude in [CII] luminosities at fixed UV SFR, indicating that these galaxies exhibit a range in metallicities and/or ionisation states (the dotted lines illustrate how the [CII] strength changes with metallicity in the model from Vallini et al. 2015). Using these models, we find that the components in CR7 would have gas-metallicities $0.1 < Z/Z_{\odot} < 0.2$, with relatively little variation between different components, although components A and C are consistent with being slightly more metal poor than B.

6.1. Are UV continuum, [CII] and Ly α related?

Because of the initial non-detections of [CII] emission in galaxies with strong Ly α emission (e.g. Ouchi et al. 2013; Ota et al. 2014), it has been speculated that the [CII] luminosity at fixed SFR is related to the strength of Ly α emission (e.g. Knudsen et al. 2016; Smit et al. 2017), as [CII] emission may be reduced in ISM conditions that favour high Ly α escape. Indeed, conditions that may lower the strength of [CII] emission are typically found in Ly α emitters. Compared to the general galaxy population, strong Ly α emitters at $z \approx 2 - 3$ are observed to have low metallicities and high ionisation states (e.g. Erb et al. 2016; Nakajima et al. 2016; Trainor et al. 2016; Kojima et al. 2017). Our results however do not agree with such scenario, as CR7 has an extremely high Ly α luminosity and EW, but has [CII] luminosities similar to those expected for the local relation (both integrated, and for individual clumps). Therefore, the strength of Ly α emission may not be the most important property driving the scatter in the SFR- $L_{\text{[CII]}}$ relation. Hence, indications of a potential relation between Ly α strength and [CII] luminosity may have reflected a more fundamental correlation.

A property that is related to the strength of Ly α emission in star-forming galaxies across $z \approx 0 - 5$ is the UV slope, with bluer galaxies being brighter in Ly α (e.g. Atek et al. 2008; Matthee et al. 2016; Oyarzún et al. 2017). The UV slope β is closely related to the age, metallicity and dust attenuation law (e.g. Bouwens et al. 2012; Duncan & Conselice 2015; Mancini et al. 2016; Narayanan et al. 2017), and therefore indirectly with properties that determine the [CII] luminosity at fixed SFR. We illustrate this in Fig. 10, where we show the [CII] luminosity as a function of the UV slope for galaxies from the literature with SFR between 15 and 35 $M_{\odot} \text{ yr}^{-1}$. This is the range of SFRs in Fig. 9 that includes most galaxies at $z \approx 5 - 7$ that have currently been observed. We note that we explicitly investigate a relatively narrow range in SFR to prevent being diluted by




Figure 10. The relation between [CII] luminosity and UV slope, β , for galaxies with SFRs between 15 – 35 $M_{\odot} \text{ yr}^{-1}$. Downward pointing triangles show galaxies with upper limits on the [CII] luminosity. Galaxies with redder colours are typically more evolved (older, higher metallicity), and consequently have higher [CII] luminosities. For reference, we show the measurements of clumps B, C and the total of CR7 in fainter colours.

relations between SFR and galaxy properties themselves (for example, it has been established that fainter galaxies have bluer colours, Bouwens et al. 2012). Although the error-bars on UV slope measurements are large⁵, it can be seen that at fixed SFR, galaxies with redder colours (which are typically more evolved) have higher [CII] luminosities. Since β depends on several parameters, direct measurements of these parameters (such as metallicity) are required to fully understand the origin of this trend, and which parameter is most important.

It is possible to compare our results in more detail to the results obtained for Himiko in Ouchi et al. (2013). Himiko consists of three UV components, with similar total UV and Ly α luminosity as CR7. Himiko has similar limits on its dust mass, but has not been detected in [CII], with a limiting luminosity $L_{\text{[CII]}} \lesssim 5 \times 10^7 L_{\odot}$ (Ouchi et al. 2013), which is well below the total [CII] luminosity from CR7. However, unlike CR7, the individual components in Himiko have SFRs of $\approx 7, 5$ and $8 M_{\odot} \text{ yr}^{-1}$ (Ouchi et al. 2013), and are hence likely of lower mass than CR7s clump A. The SFRs of Himiko’s components are comparable to CR7 clumps B and C, that have lower [CII] luminosities than the existing [CII] limits for Himiko (and the clumps in Himiko are all very

⁵ Note that for the sample at $z \approx 5 - 6$, we used the most recent UV slopes as measured by Barisic et al. (2017), which differ significantly from the ground-based estimates from Capak et al. (2015).




Figure 11. Rest-frame Ly α image (obtained by subtracting the (weak) contribution from the continuum in the NB921 image using the Subaru/S-Cam z' band image.) of CR7, overlaid with ALMA [CII] contours, with a scale of 54×54 kpc. The Ly α surface brightness is closely related to the [CII] surface brightness. At clump A, Ly α is redshifted with respect to [CII] by 127 ± 27 km s $^{-1}$.

blue, with $\beta \lesssim -2.0$). Deeper [CII] observations at the resolution we obtained for CR7 could perhaps be able to detect those clumps. Another explanation could be that Himiko has a lower dynamical mass, such that supernova explosions disperse [CII] emitting gas over larger areas, decreasing its observability.

6.2. Spatial and spectral Ly α -[CII] connection in CR7

As shown in Fig. 11, the [CII] emission in CR7 coincides spatially with the Ly α emission. This may not be unexpected if both Ly α and [CII] originate from star-formation. Although [CII] is also emitted from gas that is (partly) neutral (due to its slightly lower ionisation energy than hydrogen; for example photo-dissociating regions), Ly α emission scatters in these neutral gas regions, such that it may also be observed from these regions.

Interestingly, the [CII] line-width and velocity offset of clump A are surprisingly similar to the best fitted-shell model to CR7's Ly α profile (Dijkstra et al. 2016), which has an intrinsic line-width of 259 km s $^{-1}$ and outflow velocity of 230 km s $^{-1}$ (compare to Table 1). Their best fitted shell-model assumes a negligible dust content (consistent with our constraints on the IR luminosity), and the resulting shell-model parameters are similar to

Ly α emitters and analogue galaxies at lower redshifts. Therefore, the resemblance of the shell-model parameters based on the Ly α profile with the observed velocity offsets indicates feedback processes are already present in CR7. Further detailed studies on how the Ly α and [CII] line-profiles are related spatially require high resolution Ly α IFU observations with e.g. MUSE, as our current Ly α spectral information is limited to clump A.

7. DISCUSSION

7.1. Multiple velocity components: merger/satellites?

As shown in Figures 4 and 6, CR7 consists of several different components of [CII] emission, at different velocities. Similar clumpy structures are also observed in [OIII] $_{88\mu\text{m}}$ and [CII] emission in the $z = 7.1$ galaxy BDF3299 (Carniani et al. 2017), and also in [CII] emission in a relatively massive galaxy at $z = 6.1$ (Jones et al. 2017). Such relatively complex [CII] structures and the observed velocity offsets are also observed in hydrodynamical simulations of galaxies at $z \approx 6 - 7$ with halo masses of $\sim 10^{11} M_{\odot}$ (e.g. Vallini et al. 2013; Pallottini et al. 2017b). The line-profiles do not show evidence for broad-line components due to outflows (e.g. Gallerani et al. 2016). Therefore, we are likely observing satellites (B, B-2, C, C-2) falling into a more massive central galaxy. These satellites have velocity widths similar to those of satellites in simulations by Pallottini et al. (2017a,b), although simulated satellites typically have lower masses than those estimated for CR7's clumps. Clump A is the component that has the highest dynamical mass estimate (see Table 1, although we stress that these are rough estimates) and likely the central galaxy. The dynamical mass estimate of clump C-2 is on the same order of magnitude, potentially indicating a major merger. Observations at higher resolution and with improved sensitivity are required to robustly resolve different components and perform more detailed kinematical/dynamical analyses.

7.2. On the nature of CR7

Here, we investigate what our new measurements mean for the interpretation of CR7. For a more in depth discussion on the updated detection significance of HeII and limits on other high-ionisation UV lines we refer to Sobral et al. (2017a). The main conclusion from Sobral et al. (2017a) is that the most luminous component of CR7 is likely undergoing a recent burst of star-formation, with no clear evidence for AGN activity and no convincing detection of UV metal lines in clump A.

Initially, Sobral et al. (2015) argued that different UV components in CR7 were in different evolutionary stages, with the majority of the stellar mass possibly

being found in clumps B and C (see also Agarwal et al. 2016). This interpretation was mostly based on the different UV slopes, with clump A being the bluest and hence the youngest. However, there are two uncertainties related to this interpretation. The first is that UV slopes are relatively uncertain (see Table 1), and thus consistent with being the same for all sources. The second is that it is challenging to estimate where the majority of stellar mass is from *Spitzer*/IRAC observations, which has a relatively large PSF. For example, contrary to Sobral et al. (2015), Bowler et al. (2017) found that the majority of stellar mass is likely associated to clump A using a deconvolution of the IRAC data based on UV detections as a prior. The new observations point towards a scenario where most of the mass is in clump A as the dynamical mass estimate of clump A is higher than that of clumps B and C. Simultaneously, the constraints from the UV slope, IR continuum and the Ly α luminosity indicates that clump A is also the youngest.

As the SFR-[CII] ratio of different clumps are similar, there is evidence that all clumps have similar metallicities. However, [CII] luminosity may be reduced due to photo-evaporation by far-UV (6 – 13.6 eV) and ionising (> 13.6eV) photons emitted from young stars in the vicinity, which typically happens a few tens of Myrs after the burst of star-formation (e.g. Decataldo et al. 2017; Vallini et al. 2017), which can affect different clumps differently. The metallicities that we would infer from [CII] ($\approx 0.1 - 0.2Z_{\odot}$) are inconsistent with the metallicity inferred by Bowler et al. (2017), which is $\approx 0.005Z_{\odot}$. However, their result depends on a HeII strength that is now ruled out (Sobral et al. 2017a). Based on upper limits on UV metal lines, the SFR and photoionization modelling, Sobral et al. (2017a) inferred a metallicity of $\approx 0.05 - 0.2 Z_{\odot}$, which is consistent with the metallicity indicated by the [CII] luminosity. Therefore, it is ruled out that the luminosity in all clumps in CR7 is dominated by PopIII-like stars, but note that small amounts of PopIII-like stars in unpolluted pockets of gas could still be present (e.g. Pallottini et al. 2015). The fact that the SFR-[CII] ratio is similar to other $z \approx 6 - 7$ galaxies and galaxies in the local Universe also indicates that photo-evaporation does not play a major role in lowering the [CII] luminosity.

Due to its extreme Ly α luminosity ($\approx 10L^*$, Matthee et al. 2015), CR7 could be powered by an AGN, particularly as the AGN fraction of Ly α emitters with similar luminosities at $z = 2 - 5$ approaches 100 % (Calhau et al. 2017). However, these galaxies typically have (much) broader Ly α lines and CIV luminosities exceeding the observational limits for CR7 (e.g. Sobral et al. 2017b), such that they are not fully comparable. The

data presented in this paper do not favour an AGN explanation, particularly for clump A, because AGN typically have much lower $\log_{10}(L_{[\text{CII}]}/L_{\text{IR}})$ ratios (e.g. Ota et al. 2014). Furthermore, Bowler et al. (2017) show that the UV emission in clump A is slightly extended, which could also be at odds with an AGN-dominated scenario. Therefore, it is likely that clump A in CR7 is powered by a burst of star-formation with moderately low metallicity. It is remarkable that there is no UV nor IR continuum emission observed at the position of the ALMA detection C-2, which has a similar dynamical mass estimate as clump A. If potential UV emission is obscured, the dust is likely at a higher temperature (such that our IR continuum luminosity limit would be weak).

In order to improve our understanding of the CR7 system and other similar systems at $z \sim 7$, it is most important to obtain more accurate constraints on the stellar masses, ages and metallicities of different clumps. While deeper *HST* imaging results in more strongly constrained UV slopes, the majority of progress will be made with integral field spectroscopic observations in the IR which can trace the rest-frame optical. Future adaptive optics assisted observations with MUSE can further map out the relation between the Ly α and [CII] kinematics, and simultaneously probe the large scale environment on $\sim 0.3 - 7$ Mpc scales.

8. CONCLUSIONS

In this paper, we have presented deep follow-up observations of the CR7 galaxy in band 6 with ALMA targeted at the [CII] fine-structure cooling line and the FIR dust continuum emission (on source integration time 6.0h). After the discovery of CR7 as the most luminous Ly α emitter at $z \sim 7$ (Matthee et al. 2015), the spectroscopic confirmation of Ly α (Sobral et al. 2015), the evidence for a narrow and strong HeII line and the existence of several clumps seen in the *HST*/WFC3 imaging (Sobral et al. 2015), numerous spectacular interpretations emerged. While a high HeII/Ly α ratio is now ruled out for clump A (Sobral et al. 2017a), our ALMA observations provide independent measurements of the properties of the structure of the ISM around the three UV clumps observed in CR7 and furthermore reveal the presence of metals in its ISM. Our main results are:

1. We detect extended [CII] $_{158\mu\text{m}}$ line-emission around CR7, with a total luminosity of $(2.17 \pm 0.36) \times 10^8 L_{\odot}$ (Fig. 1). The [CII] emission is observed over an 2.2 arcsec^2 area with a typical blue-shift of $-167 \pm 27 \text{ km s}^{-1}$ with respect to Ly α , consistent with simple shell-model parameters based on an outflowing shell of low column

- density hydrogen. A fraction of 0.72 ± 0.18 of the [CII] flux is associated with separate [CII] emitting clumps, see §4.
2. Within CR7, four separate components can be identified in the three dimensional [CII] data-cube, see §4.2.1. UV components A and B are associated to clumpy [CII] emission at a similar blue-shift of $\approx 130 \text{ km s}^{-1}$ with respect to the Ly α redshift, see Table 1. A separate [CII] component, C-2, is blue-shifted by $\approx 100 \text{ km s}^{-1}$ with respect to the systemic redshift. This component is not clearly associated to a UV clump, but closest to clump C. Around clump B we detect another [CII] component blue-shifted by $\approx 300 \text{ km s}^{-1}$ compared to the systemic.
 3. The observed rich dynamical structure resembles the structure found in recent hydrodynamical simulations of $z \sim 6 - 7$ galaxies with halo masses $\sim 10^{11} M_{\odot}$, and leads to the interpretation that these components are likely inflowing satellites. The measured line-widths and dynamical mass estimates (Table 1) indicate that clump A contains most of the mass and is likely the central galaxy of the halo, although the dynamical mass estimate of clump C-2 is on the same order of magnitude, potentially indicating a major merger.
 4. We do not detect FIR continuum emission, resulting in a 3σ upper limit of $L_{\text{IR}} < 2.8 \times 10^{10} L_{\odot}$ (under the assumption that the dust temperature is 35 K), corresponding to $M_{\text{dust}} < 2.7 \times 10^6 M_{\odot}$, see §5. The limiting FIR-to-[CII] ratio indicates a low dust-to-metal ratio, much higher than typical sub-mm and quasar host galaxies. We use the FIR constraints to mitigate large uncertainties in the SFRs (due to uncertainties in their UV slopes) and find SFRs of 28_{-1}^{+1} , 5_{-1}^{+2} and $7_{-1}^{+1} M_{\odot} \text{ yr}^{-1}$ for clumps A, B and C, respectively.
 5. Based on spatially resolved [CII] measurements, we compare the positions of clumps A, B and C and CR7 in total on the SFR- $L_{\text{[CII]}}$ diagram with the literature, see §6 and Fig. 9. We find that all clumps and the total CR7 luminosity are consistent with the local relation (and other recent detections at $z \approx 6 - 7$) within the scatter. Comparison with hydrodynamical simulations (and with the caveat in mind that poorly constrained ionisation conditions may play an important role), these measurements indicate gas metallicities of $0.1 < Z/Z_{\odot} < 0.2$.

6. We find that for galaxies with SFRs between 15 and $35 M_{\odot} \text{ yr}^{-1}$, the [CII] luminosity increases strongly with increasing UV slope, which is a tracer of dust, age and metallicity (Fig. 10). This explains why several Ly α emitters are offset towards lower [CII] luminosities, as the Ly α escape fraction increases towards bluer UV slopes.

The picture that emerges is that in CR7 we are likely witnessing the build up of a central galaxy in the early Universe through complex accretion of satellites. The strong Ly α emission is likely powered by a star-burst with young, metal poor stars. Major improvements will be made with resolved spectroscopy that can measure gas metallicities and ionisation states, which can be used to constrain the properties of stellar populations in the different components.

We thank the referee for their constructive comments which have helped improving the quality and clarity of this work. We thank Raffaella Schneider for comments on an earlier version of this paper. We thank Leindert Boogaard, Steven Bos, Rychard Bouwens and Renske Smit for discussions. JM acknowledges the support of a Huygens PhD fellowship from Leiden University. DS acknowledges financial support from the Netherlands Organisation for Scientific research (NWO) through a Veni fellowship and from Lancaster University through an Early Career Internal Grant A100679. AF acknowledges support from the ERC Advanced Grant INTERSTELLAR H2020/740120. BD acknowledges financial support from NASA through the Astrophysics Data Analysis Program (ADAP), grant number NNX12AE20G. Based on observations made with ESO Telescopes at the La Silla Paranal Observatory under programme ID 294.A-5018. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2015.1.00122.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan) and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ.

Facilities: ALMA (band 6), *HST* (WFC3), VLT (X-SHOOTER), Keck (DEIMOS), VISTA (Vircam), Subaru (S-Cam).




Figure 12. Channel maps of the [CII] emission around CR7 (54x54 kpc). Contours show the $\pm 2, 3, 4, 5\sigma$ levels, where $1\sigma \approx 0.06$ mJy beam $^{-1}$. Red crosses mark the positions of the UV clumps. Channels have widths $\Delta v = 38.8$ km s $^{-1}$ or 30 MHz. The displayed channels range from $z_{\text{[CII]}} = 6.589 - 6.607$.

REFERENCES

- Agarwal, B., Johnson, J. L., Khochfar, S., et al. 2017, *MNRAS*, 469, 231
- Agarwal, B., Johnson, J. L., Zackrisson, E., et al. 2016, *MNRAS*, 460, 4003
- Atek, H., Kunth, D., Hayes, M., Östlin, G., & Mas-Hesse, J. M. 2008, *AAP*, 488, 491
- Barisic, I., et al. 2017, *ApJ*, 845, 41
- Bouwens, R. J., et al. 2012, *ApJ*, 754, 83
- . 2016, *ApJ*, 833, 72
- Bowler, R. A. A., McLure, R. J., Dunlop, J. S., et al. 2017, *MNRAS*, 469, 448
- Bowler, R. A. A., Dunlop, J. S., McLure, R. J., et al. 2012, *MNRAS*, 426, 2772
- . 2014, *MNRAS*, 440, 2810
- Bradač, M., et al. 2017, *ApJL*, 836, L2
- Calhau, J., et al. 2017, in prep
- Capak, P. L., et al. 2015, *Nature*, 522, 455
- Carilli, C. L., & Walter, F. 2013, *ARAA*, 51, 105
- Carniani, S., et al. 2017, *AAP*, 605, A42
- Cen, R., & Kimm, T. 2014, *ApJ*, 782, 32
- Ceverino, D., Dekel, A., & Bournaud, F. 2010, *MNRAS*, 404, 2151
- da Cunha, E., et al. 2013, *ApJ*, 766, 13
- . 2015, *ApJ*, 806, 110
- De Looze, I., et al. 2014, *AAP*, 568, A62
- Decarli, R., et al. 2017, *Nature*, 545, 457
- Decataldo, D., Ferrara, A., Pallottini, A., Gallerani, S., & Vallini, L. 2017, *MNRAS*, 471, 4476
- Dijkstra, M., Gronke, M., & Sobral, D. 2016, *ApJ*, 823, 74
- Duncan, K., & Conselice, C. J. 2015, *MNRAS*, 451, 2030
- Dunlop, J. S., et al. 2017, *MNRAS*, 466, 861
- Erb, D. K., Pettini, M., Steidel, C. C., et al. 2016, *ApJ*, 830, 52
- Ferkinhoff, C., Hailey-Dunsheath, S., Nikola, T., et al. 2010, *ApJL*, 714, L147
- Gallerani, S., Pallottini, A., Feruglio, C., et al. 2016, *ArXiv e-prints*, arXiv:1604.05714
- Guseva, N. G., Izotov, Y. I., Fricke, K. J., & Henkel, C. 2017, *AAP*, 599, A65
- Hartwig, T., Latif, M. A., Magg, M., et al. 2016, *MNRAS*, 462, 2184
- Hemmati, S., Yan, L., Diaz-Santos, T., et al. 2017, *ApJ*, 834, 36
- Herrera-Camus, R., et al. 2015, *ApJ*, 800, 1
- Hirashita, H., Ferrara, A., Dayal, P., & Ouchi, M. 2014, *MNRAS*, 443, 1704
- Hopkins, P. F., Kereš, D., Oñorbe, J., et al. 2014, *MNRAS*, 445, 581
- Inoue, A. K., et al. 2016, *Science*, 352, 1559
- Izotov, Y. I., Guseva, N. G., Fricke, K. J., & Henkel, C. 2015, *MNRAS*, 451, 2251
- Jones, G. C., Willott, C. J., Carilli, C. L., et al. 2017, *ApJ*, 845, 175
- Kanekar, N., Wagg, J., Chary, R. R., & Carilli, C. L. 2013, *ApJL*, 771, L20
- Kaufman, M. J., Wolfire, M. G., Hollenbach, D. J., & Luhman, M. L. 1999, *ApJ*, 527, 795
- Kennicutt, Jr., R. C. 1998, *ARAA*, 36, 189
- Knudsen, K. K., Richard, J., Kneib, J.-P., et al. 2016, *MNRAS*, 462, L6
- Knudsen, K. K., Watson, D., Frayer, D., et al. 2017, *MNRAS*, 466, 138
- Kojima, T., Ouchi, M., Nakajima, K., et al. 2017, *PASJ*, 69, 44
- Laigle, C., et al. 2016, *ApJS*, 224, 24
- Laporte, N., Nakajima, K., Ellis, R. S., et al. 2017a, *ArXiv e-prints*, arXiv:1708.05173
- Laporte, N., et al. 2017b, *ApJL*, 837, L21
- Maiolino, R., et al. 2015, *MNRAS*, 452, 54
- Malhotra, S., et al. 2001, *ApJ*, 561, 766
- Mancini, M., Schneider, R., Graziani, L., et al. 2016, *MNRAS*, 462, 3130
- . 2015, *MNRAS*, 451, L70
- Matthee, J., Sobral, D., Best, P., et al. 2017a, *MNRAS*, 465, 3637
- Matthee, J., Sobral, D., Darvish, B., et al. 2017b, *MNRAS*, 472, 772
- Matthee, J., Sobral, D., Oteo, I., et al. 2016, *MNRAS*, 458, 449
- Matthee, J., Sobral, D., Santos, S., et al. 2015, *MNRAS*, 451, 400
- McMullin, J. P., Waters, B., Schiebel, D., Young, W., & Golap, K. 2007, in *Astronomical Society of the Pacific Conference Series*, Vol. 376, *Astronomical Data Analysis Software and Systems XVI*, ed. R. A. Shaw, F. Hill, & D. J. Bell, 127
- Meurer, G. R., Heckman, T. M., & Calzetti, D. 1999, *ApJ*, 521, 64
- Michałowski, M. J. 2015, *AAP*, 577, A80
- Nakajima, K., Ellis, R. S., Iwata, I., et al. 2016, *ApJL*, 831, L9
- Narayanan, D., Dave, R., Johnson, B., et al. 2017, *ArXiv e-prints*, arXiv:1705.05858
- Olsen, K. P., Greve, T. R., Narayanan, D., et al. 2017, *ArXiv e-prints*, arXiv:1708.04936
- Ota, K., et al. 2014, *ApJ*, 792, 34
- Ouchi, M., et al. 2013, *ApJ*, 778, 102

- Oyarzún, G. A., Blanc, G. A., González, V., Mateo, M., & Bailey, III, J. I. 2017, ArXiv e-prints, arXiv:1706.01886
- Pacucci, F., Pallottini, A., Ferrara, A., & Gallerani, S. 2017, MNRAS, 468, L77
- Pallottini, A., Ferrara, A., Bovino, S., et al. 2017a, MNRAS, 471, 4128
- Pallottini, A., Ferrara, A., Gallerani, S., et al. 2017b, MNRAS, 465, 2540
- Pallottini, A., Ferrara, A., Pacucci, F., et al. 2015, MNRAS, 453, 2465
- Pentericci, L., et al. 2016, ApJL, 829, L11
- Popping, G., Somerville, R. S., & Galametz, M. 2017, MNRAS, 471, 3152
- Rémy-Ruyer, A., et al. 2014, AAP, 563, A31
- Riechers, D. A., et al. 2013, Nature, 496, 329
- . 2017, ArXiv e-prints, arXiv:1705.09660
- Salpeter, E. E. 1955, ApJ, 121, 161
- Schaerer, D., Boone, F., Zamojski, M., et al. 2015, AAP, 574, A19
- Scoville, N., et al. 2016, ApJ, 820, 83
- . 2017, ApJ, 837, 150
- Shibuya, T., et al. 2017, ArXiv e-prints, arXiv:1705.00733
- Smidt, J., Wiggins, B. K., & Johnson, J. L. 2016, ApJL, 829, L6
- Smit, R., et al. 2017, ArXiv e-prints, arXiv:1706.04614
- Smith, A., Bromm, V., & Loeb, A. 2016, MNRAS, 460, 3143
- Sobral, D., Matthee, J., Darvish, B., et al. 2015, ApJ, 808, 139
- Sobral, D., et al. 2017a, ArXiv e-prints, arXiv:1710.08422
- . 2017b, in prep
- Solomon, P. M., Downes, D., & Radford, S. J. E. 1992, ApJL, 398, L29
- Stark, D. P., et al. 2017, MNRAS, 464, 469
- Swinbank, A. M., et al. 2012, MNRAS, 427, 1066
- Trainor, R. F., Strom, A. L., Steidel, C. C., & Rudie, G. C. 2016, ApJ, 832, 171
- Vallini, L., Ferrara, A., Pallottini, A., & Gallerani, S. 2017, MNRAS, 467, 1300
- Vallini, L., Gallerani, S., Ferrara, A., & Baek, S. 2013, MNRAS, 433, 1567
- Vallini, L., Gallerani, S., Ferrara, A., Pallottini, A., & Yue, B. 2015, ApJ, 813, 36
- Visbal, E., Haiman, Z., & Bryan, G. L. 2016, MNRAS, 460, L59
- Wang, R., et al. 2013, ApJ, 773, 44
- Watson, D., Christensen, L., Knudsen, K. K., et al. 2015, Nature, 519, 327
- Willott, C. J., Carilli, C. L., Wagg, J., & Wang, R. 2015, ApJ, 807, 180