

TOWARDS AN UNIFIED THEORY FOR TESTING STATISTICAL HYPOTHESES: MULTIVARIATE MEAN WITH NUISANCE COVARIANCE MATRIX

BY MING-TIEN TSAI

Academia Sinica

Under a multinormal distribution with arbitrary unknown covariance matrix, the main purpose of this paper is to propose a framework to achieve the goal of reconciliation of Bayesian, frequentist and Fisherian paradigms for the problems of testing mean against restricted alternatives (closed convex cones). Combine fiducial inference and Wald decision theory via d -admissibility into an unified approach, the goal can then be achieved. To proceed, the tests constructed via the union-intersection principle (Roy, 1953) are studied. The difficulty for likelihood ratio principle is mentioned.

AMS 1991 subject classification: 62C15, 62H15.

Key words and phrases: Bayesian paradigm, d -admissibility, fiducial inference, Fisherian paradigm, Neyman-Pearson approach, union-intersection test.

1. Introduction and preliminary notions. Let $\{\mathbf{X}_i; 1 \leq i \leq n\}$ be independent and identically distributed random vectors (i.i.d.r.v.) having a p -variate normal distribution with mean vector $\boldsymbol{\mu}$ and dispersion matrix $\boldsymbol{\Sigma}$. Consider the null hypothesis

$$H_0^* : \boldsymbol{\mu} \in \Gamma_1 = \{\boldsymbol{\mu} | \mathbf{B}_1 \boldsymbol{\mu} = \mathbf{0}\} \quad (1.1)$$

against a restricted (convex polyhedral cone) alternative

$$H_1^* : \boldsymbol{\mu} \in \Gamma_2 = \{\boldsymbol{\mu} | \mathbf{B}_2 \boldsymbol{\mu} \geq \mathbf{0}\}, \quad (1.2)$$

where $\mathbf{B}_i \in \mathcal{C}(m_i, p)$, the set of $m_i \times p$ matrices of rank m_i ($1 \leq m_i \leq p$), $i=1,2$, and Γ_1 is assumed to be the linear hull of $\tilde{\Gamma}_2 = \{\boldsymbol{\mu} | \mathbf{B}_1 \boldsymbol{\mu} = \mathbf{0}, \mathbf{B}_2 \boldsymbol{\mu} \geq \mathbf{0}\}$. An impasse in a general formulation of optimal tests for this problem is the lack of invariance of the model (as well as the likelihood ratio or allied tests) under suitable groups of transformations which map the sample space onto itself. Specifically, if we consider an affine transformation $\mathbf{X} \rightarrow \mathbf{Y} = \mathbf{a} + \mathbf{A}\mathbf{X}$, where \mathbf{A} is non-singular, this hypothesis testing problem is not invariant in general although it is invariant under two special groups of linear transformations (positive diagonal matrices and permutation matrices) when $\mathbf{B}_1 = \mathbf{B}_2 = \mathbf{I}$. As such, a canonical reduction of the noncentrality to a single coordinate may not work out, and if lacking this invariance, the usual techniques fail to provide an optimality property of the tests. For this reason, often, a hypothesis related transformation is used on the basic r.v.'s, and on this transformed r.v.'s suitable tests are formulated. Towards this, we may consider $\mathbf{Y}_i = \mathbf{B}_1 \mathbf{X}_i, i \geq 1, \boldsymbol{\theta} = \mathbf{B}_1 \boldsymbol{\mu}$ and $\mathbf{B} = \mathbf{B}_2 \mathbf{B}_1' (\mathbf{B}_1 \mathbf{B}_1')^{-1}$, so that based on the \mathbf{Y}_i , the hypotheses in (1.1) and (1.2) can be expressed as

$$\bar{H}_0 : \boldsymbol{\theta} = \mathbf{0} \quad \text{vs.} \quad \bar{H}_1 : \boldsymbol{\theta} \in \Gamma_3 = \{\boldsymbol{\theta} | \mathbf{B}\boldsymbol{\theta} \geq \mathbf{0}, \|\boldsymbol{\theta}\| > 0\}, \quad (1.3)$$

(Recall that $\boldsymbol{\Sigma} \rightarrow \boldsymbol{\Sigma}_1 = \mathbf{B}_1 \boldsymbol{\Sigma} \mathbf{B}_1'$ and need not be diagonal). Further reduction to a positive orthant model is feasible by the transformations $\mathbf{Z}_i = \mathbf{B}\mathbf{Y}_i, \boldsymbol{\beta} = \mathbf{B}\boldsymbol{\theta}$ and $\boldsymbol{\Sigma} \rightarrow \mathbf{B}\boldsymbol{\Sigma}_1 \mathbf{B}'$. Situations are similar for the cases when the conditions $m_i \leq p, i = 1, 2$ are relaxed. Also note that a specific halfspace can be transformed into another halfspace by a non-singular linear transformation. Hence without loss of generality it suffices to consider the following models:

Let $\mathbf{X}_1, \dots, \mathbf{X}_n$ be i.i.d.r.v.'s with the $N_p(\boldsymbol{\theta}, \boldsymbol{\Sigma})$ d.f., where $\boldsymbol{\Sigma}$ is unknown and assumed to be positive definite (p.d.). Consider the hypotheses

$$H_0 : \boldsymbol{\theta} = \mathbf{0} \quad \text{vs.} \quad H_1 : \boldsymbol{\theta} \in \mathcal{C} \setminus \{\mathbf{0}\}, \quad (1.4)$$

where \mathcal{C} denotes a closed convex cone containing a p -dimensional open set. Denote the positive orthant space by $\mathcal{O}_p^+ = \{\boldsymbol{\theta} \in R^p | \boldsymbol{\theta} \geq \mathbf{0}\}$. Notice that when \mathcal{C} is a proper set contained in a

halfspace, under a suitable linear transformation the problem in (1.4) can be reduced to the problem for testing against the positive orthant space with another unknown positive definite covariance matrix. When \mathcal{C} is a specific halfspace, then it can be transformed into another halfspace by a non-singular linear transformation. Hence without loss of generality it suffices enough to study the cases that \mathcal{C} is the positive orthant space \mathcal{O}_p^+ and \mathcal{C} is the halfspace $\mathcal{H}_p^* = \{\boldsymbol{\theta} \in R^p \mid \theta_p \geq 0\}$ throughout this paper.

For testing against global alternative, $H_0^g : \boldsymbol{\theta} = \mathbf{0}$ vs. $H_1^g : \boldsymbol{\theta} \neq \mathbf{0}$, the likelihood ratio test (LRT) and the union-intersection test (UIT) are isomorphic, and are well-known as the Hotelling's T^2 -test (Anderson, 2003). Kiefer and Schwartz (1965) showed that the Hotelling's T^2 -test is a proper Bayes test. The Hotelling's T^2 -test is also known as a version of integrated LRT with respect to objective priors. As such, for testing against global alternative, there are several ways to establish the equivalent relationships of the Hotelling's T^2 -test with Bayes tests.

On the other hand, for the problems of testing against restricted alternatives considered in (1.4), the LRT and the UIT are different (Perlman, 1969, Sen and Tsai, 1999) when the covariance matrix $\boldsymbol{\Sigma}$ is totally unknown. The problem (1.4) indeed provides us some partial informations of interesting parameters. However, due to the difficulty of integration over high-dimensional restricted parameter space the explicit forms of Bayes tests are hardly obtained, though the numerical values of them can be obtained via the method of Markov chain Monte Carlo. The same difficulty arises for the approach of Bayes factor. In Section 2, we show that the UIT are no longer the Bayes tests for the problem (1.4). The likelihood integrated (Berger, Liseo and Wolpert, 1999) with respect to a relevant Haar measure for the nuisance covariance matrix while remaining neutral with respect to the parameter of interest, which fiducial inference was intended to be, can be viewed in a objective Bayesian light. It is shown that for the problems of testing against restricted alternatives both the LRT and the UIT are the versions of integrated LRT and integrated UIT, respectively. These result in the reconciliation of frequentist and Bayesian paradigms.

For testing against global alternative, the Hotelling's T^2 -test enjoys many optimal properties of Neyman-Pearson testing hypothesis theory such as similarity, unbiasedness, power monotonicity, most stringent, uniformly most powerful invariant and α -admissibility (Anderson, 2003). For the problem of testing against the positive orthant space, both the LRT and the UIT are shown not unbiased (Perlman, 1969, Sen and Tsai, 1999). And it is easy to see that both the LRT and the UIT are power dominated by the corresponding LRT and UIT for the problem of testing against the halfspace \mathcal{H}_p^* , respectively. As such, for the problem of testing against the positive orthant space both the LRT and the UIT are α -inadmissible. The related domination problems are also studied in

Section 3, some of these domination results are against the common statistical sense. The situation in Neyman-Pearson theory is beautifully peaceful and compelling for the unrestricted alternative, while it is not in some restricted alternatives world. One of the major reasons for these phenomena is that for the problem of testing against the positive orthant space the distribution functions of both the LRT statistic and the UIT statistic are not free from the dependence on the unknown nuisance covariance matrix under null hypothesis. Because of this unpleasant feature, the Fisher's approach of reporting p -values may also not work well for the problem of testing against the positive orthant space. Several methods have been proposed to get rid of the unpleasant feature, we mention some in Section 3.

In the literature, there are many ways to eliminate the nuisance parameters (Basu, 1977). Most of them can be applied to the problem of testing against the positive orthant space. As we know from Section 3 that the domination problems of hypothesis testing heavily depend on the choice of the critical points which are used to report the p -values. Sometimes the danger stems from too narrow definitions of what is meant by optimality and the strongest intuition can sometimes go astray too. Hence, the question naturally raised is whether there exists any satisfactory, unified approach to overcome the difficulties. Neither Fisher's approach of reporting p -values alone nor Neyman-Pearson's optimal theory for power function alone is a well satisfactory criterion for evaluating the performance of tests. The spirit of compromise between Fisher's approach and Neyman-Pearson's optimal theory without detailed consideration of power may shed light on the testing hypothesis theory. Imposing on the balance between type I error and power, Wald's decision theory still paves a unified way to combine the best features of both Neyman-Pearson's and Fisher's ideas. In Section 4, we show that the Hotelling's T^2 -test is inadmissible for the problem (1.4). And further show that the UIT is α -admissible and d -admissible for the problem of testing against the halfspace, and it is α -inadmissible but d -admissible for the problem of testing against the positive orthant space, respectively. We fail to claim the same results for LRT in the problem (1.4) due to the facts that the acceptance region of LRT is not convex.

2. A reconciliation of frequentist and Bayesian paradigms. For every $n (\geq 2)$, let

$$\bar{\mathbf{X}}_n = n^{-1} \sum_{i=1}^n \mathbf{X}_i \quad \text{and} \quad \mathbf{S}_n = (n-1)^{-1} \sum_{i=1}^n (\mathbf{X}_i - \bar{\mathbf{X}}_n)(\mathbf{X}_i - \bar{\mathbf{X}}_n)', \quad (2.1)$$

the Hotelling's T^2 -test is expressed as

$$T_n^2 = n\bar{\mathbf{X}}_n' \mathbf{S}_n^{-1} \bar{\mathbf{X}}_n. \quad (2.2)$$

For testing against global alternative, Kiefer and Schwartz (1965) showed that the Hotelling's T^2 -test

is a proper Bayes test. The main goal of this section is to see whether the reconciliation of frequentist and Bayesian paradigms can be established for the problem (1.4). First, we establish the necessary conditions for the Bayes tests.

PROPOSITION 2.1. *Let \mathcal{A} denote the acceptance region of a size- α Bayes test for the problem of testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{C} \setminus \{\mathbf{0}\}$, where \mathcal{C} is either the positive orthant space \mathcal{O}_p^+ or the halfspace \mathcal{H}_p^* , and \mathcal{L} be any line of support of \mathcal{A} . Denote $\bar{\mathcal{A}} = \mathcal{A} \cup \partial\mathcal{A}$, $\partial\mathcal{A}$ being the boundary set of \mathcal{A} . Then either $\mathcal{L} \subseteq \partial\mathcal{A}$ or $\mathcal{L} \cap \bar{\mathcal{A}} = \{\mathbf{a}\}$, where $\mathbf{a} \in \partial\mathcal{A}$.*

PROOF. The density of $\mathbf{X}_1, \dots, \mathbf{X}_n$ is

$$\frac{e^{-\frac{1}{2}n\boldsymbol{\theta}'\boldsymbol{\Sigma}^{-1}\boldsymbol{\theta}}}{(2\pi)^{\frac{1}{2}pn}|\boldsymbol{\Sigma}|^{\frac{1}{2}n}} \exp \left[n\boldsymbol{\theta}'\boldsymbol{\Sigma}^{-1}\bar{\mathbf{X}}_n + \text{tr} \left(-\frac{1}{2}\boldsymbol{\Sigma}^{-1} \sum_{j=1}^n \mathbf{X}_j \mathbf{X}_j' \right) \right]. \quad (2.3)$$

The vector $\mathbf{y} = (\mathbf{y}^{(1)'}, \mathbf{y}^{(2)'})'$ is composed of $\mathbf{y}^{(1)} = \bar{\mathbf{X}}_n$ and $\mathbf{y}^{(2)} = (v_{11}, 2v_{12}, \dots, 2v_{1p}, v_{22}, \dots, v_{pp})$, where $(v_{ij}) = \mathbf{V}_n = \sum_{j=1}^n \mathbf{X}_j \mathbf{X}_j'$. The vector $\boldsymbol{\omega} = (\boldsymbol{\omega}^{(1)'}, \boldsymbol{\omega}^{(2)'})'$ is composed of $\boldsymbol{\omega}^{(1)} = n\boldsymbol{\Sigma}^{-1}\boldsymbol{\theta}$ and $\boldsymbol{\omega}^{(2)} = -\frac{1}{2}(\sigma^{11}, \sigma^{12}, \dots, \sigma^{1p}, \sigma^{22}, \dots, \sigma^{pp})'$, where $(\sigma^{ij}) = \boldsymbol{\Sigma}^{-1}$. Write $\boldsymbol{\Sigma}^{-1} = \boldsymbol{\Gamma}(\boldsymbol{\omega}^{(2)})$, thus the density of \mathbf{Y} becomes

$$\frac{|\boldsymbol{\Gamma}(\boldsymbol{\omega}^{(2)})|^{\frac{1}{2}n} e^{-\frac{1}{2}\boldsymbol{\omega}^{(1)'}\boldsymbol{\omega}^{(1)}}}{(2\pi)^{\frac{1}{2}pn}} \exp\{\boldsymbol{\omega}'\mathbf{y}\} \quad (2.4)$$

Since $\boldsymbol{\Sigma}$ is arbitrary positive definite, $\boldsymbol{\theta} \in \mathcal{O}_p^+$ implies that $\boldsymbol{\omega}^{(1)} \in \mathcal{H}_p^+$, where $\mathcal{H}_p^+ = \{\boldsymbol{\theta} \in R^p \mid \sum_{i=1}^p \theta_i \geq 0\}$. Similarly, let \mathcal{S} be the set so that $\boldsymbol{\omega}^{(1)} \in \mathcal{S}$ whenever $\boldsymbol{\theta} \in \mathcal{H}_p^*$. Thus for simplicity and without loss of generality, it suffices to work the case that $\boldsymbol{\omega}^{(1)} \in \mathcal{H}_p^+$. Let \mathcal{M} be the group of positive definite matrices and $G(\boldsymbol{\omega})$ denote an arbitrary prior distribution function $G(\boldsymbol{\omega})$ on the set $\mathcal{N} = \{\boldsymbol{\omega} \mid \boldsymbol{\omega}^{(1)} \in \mathcal{H}_p^+ \text{ and } \boldsymbol{\Gamma}(\boldsymbol{\omega}^{(2)}) \in \mathcal{M}\}$. Let $G_0(\boldsymbol{\omega})$ denote the prior distribution function on the set $\mathcal{N}_0 = \{\boldsymbol{\omega} \mid \boldsymbol{\omega}^{(1)} = \mathbf{0} \text{ and } \boldsymbol{\Gamma}(\boldsymbol{\omega}^{(2)}) \in \mathcal{M}\}$. Then the size- α Bayes test of H_0 vs. H_1 against this prior distribution rejects H_0 if

$$t(\mathbf{y}) = \int_{\mathcal{N}} \exp\{\boldsymbol{\omega}^{(1)'}\mathbf{y}^{(1)}\} dG^*(\boldsymbol{\omega}) \geq c, \quad (2.5)$$

where

$$dG^*(\boldsymbol{\omega}) = \frac{\exp\{\boldsymbol{\omega}^{(2)'}\mathbf{y}^{(2)}\} |\boldsymbol{\Gamma}(\boldsymbol{\omega}^{(2)})|^{\frac{1}{2}n} e^{-\frac{1}{2}\boldsymbol{\omega}^{(1)'}\boldsymbol{\omega}^{(1)}} dG(\boldsymbol{\omega})}{\int_{\mathcal{N}_0} \exp\{\boldsymbol{\omega}^{(2)'}\mathbf{y}^{(2)}\} |\boldsymbol{\Gamma}(\boldsymbol{\omega}^{(2)})|^{\frac{1}{2}n} dG_0(\boldsymbol{\omega})}, \quad (2.6)$$

and c is a constant. Let P^* be the probability measure induced by G^* on the parameter space \mathcal{N} , and $\mathbf{u}, \mathbf{v} \in \partial\mathcal{A}$. Also let $\mathcal{L} = \{\mathbf{z} \mid \mathbf{z} = \rho\mathbf{u} + (1-\rho)\mathbf{v}, -\infty < \rho < \infty\}$. If $P^*\{\boldsymbol{\omega}'(\mathbf{u} - \mathbf{v}) \neq 0\} \neq 0$, then by the inequality

$$\exp(\rho m_1 + (1-\rho)m_2) \leq \rho e^{m_1} + (1-\rho) e^{m_2}, \quad 0 < \rho < 1, \quad (2.7)$$

for all real m_1 and m_2 with equality if and only if $m_1 = m_2$. Thus $t(\mathbf{z}) < c$. And hence the line \mathcal{L} is not a support of \mathcal{A} unless the intersection of \mathcal{L} and the closure of \mathcal{A} is a single boundary point. If $P^*\{\boldsymbol{\omega}'(\mathbf{u} - \mathbf{v}) \neq 0\} = 0$, then

$$t(\mathbf{z}) = \int_{\mathcal{N}} \exp\{\boldsymbol{\omega}'\mathbf{v} + \rho\boldsymbol{\omega}'(\mathbf{u} - \mathbf{v})\} dG^*(\boldsymbol{\omega}) = c. \quad (2.8)$$

Thus $\mathcal{L} \subseteq \partial\mathcal{A}$. Q.E.D.

Obviously, Proposition 2.1 can be extended to the more general set-up, though it is designed for the problem (1.4) in this paper. Let $P = \{1, \dots, p\}$, and for every $a: \emptyset \subseteq a \subseteq P$, let a' be its complement and $|a|$ its cardinality. For each a , we partition $\bar{\mathbf{X}}_n$ and \mathbf{S}_n as

$$\bar{\mathbf{X}}_n = \begin{pmatrix} \bar{\mathbf{X}}_{na} \\ \bar{\mathbf{X}}_{na'} \end{pmatrix} \quad \text{and} \quad \mathbf{S}_n = \begin{pmatrix} \mathbf{S}_{naa} & \mathbf{S}_{naa'} \\ \mathbf{S}_{na'a} & \mathbf{S}_{na'a'} \end{pmatrix}, \quad (2.9)$$

and write

$$\bar{\mathbf{X}}_{na:a'} = \bar{\mathbf{X}}_{na} - \mathbf{S}_{naa'}\mathbf{S}_{na'a'}^{-1}\bar{\mathbf{X}}_{na'}, \quad (2.10)$$

$$\mathbf{S}_{naa:a'} = \mathbf{S}_{naa} - \mathbf{S}_{naa'}\mathbf{S}_{na'a'}^{-1}\mathbf{S}_{na'a}. \quad (2.11)$$

Further, let

$$I_{na} = 1\{\bar{\mathbf{X}}_{na:a'} > \mathbf{0}, \mathbf{S}_{na'a'}^{-1}\bar{\mathbf{X}}_{na'} \leq 0\}, \quad (2.12)$$

for $\emptyset \subseteq a \subseteq P$, where $1\{\cdot\}$ denotes the indicator function. Then for the problem of testing $H_0: \boldsymbol{\theta} = \mathbf{0}$ against the positive orthant space $H_{1O^+}: \boldsymbol{\theta} \in \mathcal{O}_p^+ \setminus \{\mathbf{0}\}$, from the results of Perlman (1969) and Sen and Tsai (1999) the LRT and the UIT statistics are of the forms

$$L_n = \sum_{\emptyset \subseteq a \subseteq P} \left\{ \frac{n\bar{\mathbf{X}}_{na:a'}'\mathbf{S}_{naa:a'}^{-1}\bar{\mathbf{X}}_{na:a'}}{1 + n\bar{\mathbf{X}}_{na'a'}'\mathbf{S}_{na'a'}^{-1}\bar{\mathbf{X}}_{na'a'}} \right\} I_{na} \quad (2.13)$$

and

$$U_n = \sum_{\emptyset \subseteq a \subseteq P} \{n\bar{\mathbf{X}}_{na'a'}'\mathbf{S}_{naa:a'}^{-1}\bar{\mathbf{X}}_{na'a'}\} I_{na} \quad (2.14)$$

respectively.

Side by side, for testing $H_0: \boldsymbol{\theta} = \mathbf{0}$ against the halfspace $H_{1H^*}: \boldsymbol{\theta} \in \mathcal{H}_p^* \setminus \{\mathbf{0}\}$ the LR and the UI test statistics are of the forms

$$L_n^* = \sum_{P_1 \subseteq a \subseteq P} \left\{ \frac{n\bar{\mathbf{X}}_{na:a'}'\mathbf{S}_{naa:a'}^{-1}\bar{\mathbf{X}}_{na:a'}}{1 + n\bar{\mathbf{X}}_{na'a'}'\mathbf{S}_{na'a'}^{-1}\bar{\mathbf{X}}_{na'a'}} \right\} I_{na}^* \quad (2.15)$$

and

$$U_n^* = \sum_{P_1 \subseteq a \subseteq P} \{n\bar{\mathbf{X}}'_{na:a'} \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'}\} I_{na}^* \quad (2.16)$$

respectively, where $P_1 = P - \{p\}$, $I_{nP}^* = 1\{\bar{X}_{np} > 0\}$ and $I_{nP_1}^* = 1\{\bar{X}_{np} \leq 0\}$.

It is clear from (2.13)-(2.16) that

$$L_n^* \geq L_n, \quad \text{and} \quad U_n^* \geq U_n, \quad (2.17)$$

with probability one.

THEOREM 2.1. *For the problem of testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{C} \setminus \{\mathbf{0}\}$, where \mathcal{C} is either the positive orthant space \mathcal{O}_p^+ or the halfspace \mathcal{H}_p^* , the UIT is not a Bayes test.*

PROOF. The UIT statistics in (2.14) and in (2.16) have a similar structure, hence without loss of generality it suffices to consider the problem of testing against the positive orthant space. Let u_α be the critical point of the level of significance α for the UIT, which can be obtained from the right hand side of (3.9). Let $\mathcal{A}_U = \cup_{\emptyset}^P \mathcal{A}_{Ua}$ be the acceptance region of the UIT for testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{O}_p^+ \setminus \{\mathbf{0}\}$, where $\mathcal{A}_{Ua} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n\bar{\mathbf{X}}'_{na:a'} \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'} \leq u_\alpha\} I_{na}$, $\emptyset \subseteq a \subseteq P$. For the situation $|a| = 1$, we may note that there exists a support \mathcal{L}_a of the set \mathcal{A}_U such that $(\mathcal{L}_a \cap I_{na}) \subseteq \partial \mathcal{A}_{Ua}$ and $\mathcal{L}_a \not\subseteq \partial \mathcal{A}_U$, since $\mathcal{L}_a \cap I_{na}$ is easily seen to be a half-line. Thus theorem follows by Proposition 2.1. Q.E.D.

Note that for the problem of testing against the positive orthant space or the problem of testing against the half-space, the LRT is isomorphic to the UIT when the covariance matrix is known ($\boldsymbol{\Sigma} = \boldsymbol{\Sigma}_0$, with \mathbf{S}_n being replaced by $\boldsymbol{\Sigma}_0$) or is partially unknown, ($\boldsymbol{\Sigma} = \sigma^2 \boldsymbol{\Sigma}_0$, where σ^2 is an unknown scalar and $\boldsymbol{\Sigma}_0$ is a known positive definite matrix, with \mathbf{S}_n being replaced by $s^2 \boldsymbol{\Sigma}_0$). Hence, similar arguments as in the proof of theorem 2.1, we may conclude that the LRT is not a Bayes when the covariance matrix is known or partially unknown for the problems of testing against restricted alternatives. As such, when the covraiance matrix is totally unknown the LRT is asymptotic equivalent to the UIT as the sample size is sufficiently large. Thus, the LRT is also not a Bayes test when sample size is sufficiently large for the problems of testing against restricted alternatives..

Next, consider for the finite union-intersection test (FUIT, Roy et al., 1972) based on the one-sided coordinatedwise Student t-tests. Define $\mathbf{S}_n = (S_{nij})$ as in (2.9) and let

$$t_j = \frac{\sqrt{n} \bar{X}_{nj}}{\sqrt{S_{njj}}}, \quad j = 1, \dots, p. \quad (2.18)$$

Corresponding to a given significance level α , define

$$\alpha^* : p\alpha^* = \alpha. \quad (2.19)$$

Let t_{n-1, α^*} be the upper $100\alpha^*\%$ point of the Studentn t-distribution with $n - 1$ degrees of freedom. Consider the critical region $\mathcal{W}_j = \{t_j | t_j \geq t_{n-1, \alpha^*}\}$ for $j = 1, \dots, p$. Then the critical region of the FUIT is

$$\mathcal{W}^* = \cup_{j=1}^p \mathcal{W}_j \quad (2.20)$$

and the acceptance region is

$$\mathcal{A}^* = \cap_{j=1}^p \bar{\mathcal{A}}_j, \quad (2.21)$$

where $\mathcal{A}_j = R \setminus \mathcal{W}_j$ and $\bar{\mathcal{A}}_j$ denote the closure of $\mathcal{A}_j, j = 1, \dots, p$. Therefore, by Proposition 2.1 we have the following.

THEOREM 2.2. *For the problem of testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{C} \setminus \{\mathbf{0}\}$, where \mathcal{C} is either the positive orthant space \mathcal{O}_p^+ or the halfspace \mathcal{H}_p^* , the FUIT is not a Bayes test.*

In passing, we note that the explicit forms of the Bayes tests are hardly obtained due to the difficulty of integration over restricted parameter spaces. Hence, for the problems of testing against restricted alternatives (1.4) we will study the Bayesian tests only in terms of the non-informative prior distribution for the nuisance parameter $\boldsymbol{\Sigma}$

$$g(\boldsymbol{\Sigma}) \propto |\boldsymbol{\Sigma}|^{-\frac{1}{2}(p+1)} d\nu(\boldsymbol{\Sigma}), \quad \boldsymbol{\Sigma} \in \mathcal{M} \quad (2.22)$$

for some measure $\nu(\cdot)$, where \mathcal{M} is the set of positive definite matrices. First, consider the unique Haar measure in $\boldsymbol{\Sigma}$ over the set \mathcal{M} , i.e., $\nu(\boldsymbol{\Sigma}) = \boldsymbol{\Sigma}$. Then the marginal density of $\bar{\mathbf{X}}_n$ and \mathbf{S}_n is given by

$$P_1(\bar{\mathbf{X}}_n, \mathbf{S}_n | \boldsymbol{\theta}) \propto |\mathbf{S}_n|^{\frac{1}{2}(n-p-2)} |\mathbf{S}_n + n(\bar{\mathbf{X}}_n - \boldsymbol{\theta})(\bar{\mathbf{X}}_n - \boldsymbol{\theta})'|^{-\frac{1}{2}(n-1)}, \quad (2.23)$$

namely,

$$P_1(\bar{\mathbf{X}}_n, \mathbf{S}_n | \boldsymbol{\theta}) \propto |\mathbf{S}_n|^{-\frac{1}{2}(p+1)} [1 + n(\bar{\mathbf{X}}_n - \boldsymbol{\theta})' \mathbf{S}_n^{-1} (\bar{\mathbf{X}}_n - \boldsymbol{\theta})]^{-\frac{1}{2}(n-1)}. \quad (2.24)$$

Based on the density in (2.20), then

$$\begin{aligned}
& \frac{\sup_{\boldsymbol{\theta} \in \mathcal{O}_p^+} P_1(\boldsymbol{\theta} | \bar{\mathbf{X}}_n, \mathbf{S}_n)}{P_1(\mathbf{0} | \bar{\mathbf{X}}_n, \mathbf{S}_n)} \\
&= \left[\frac{1 + n \bar{\mathbf{X}}_n' \mathbf{S}_n^{-1} \bar{\mathbf{X}}_n}{1 + \inf_{\boldsymbol{\theta} \in \mathcal{O}_p^+} n(\bar{\mathbf{X}}_n - \boldsymbol{\theta})' \mathbf{S}_n^{-1} (\bar{\mathbf{X}}_n - \boldsymbol{\theta})} \right]^{\frac{1}{2}(n-1)} \\
&= \left[1 + \|\pi_{\mathbf{S}_n}(n^{1/2} \bar{\mathbf{X}}_n; \mathcal{O}_p^+)\|_{\mathbf{S}_n}^2 \{1 + \|n^{1/2} \bar{\mathbf{X}}_n - \pi_{\mathbf{S}_n}(n^{1/2} \bar{\mathbf{X}}_n; \mathcal{O}_p^+)\|_{\mathbf{S}_n}^2\}^{-1} \right]^{\frac{1}{2}(n-1)} \\
&= (1 + L_n)^{\frac{1}{2}(n-1)}.
\end{aligned} \tag{2.25}$$

For any $\mathbf{b} \in R^p \setminus \{\mathbf{0}\}$, the following density can be obtained from (2.19)

$$P_2(\bar{\mathbf{X}}_n, \mathbf{S}_n, \mathbf{b} | \boldsymbol{\theta}) \propto |\mathbf{b}' \mathbf{S}_n \mathbf{b}|^{-\frac{1}{2}(p+1)} [1 + n \mathbf{b}' (\bar{\mathbf{X}}_n - \boldsymbol{\theta}) (\bar{\mathbf{X}}_n - \boldsymbol{\theta})' \mathbf{b} / \mathbf{b}' \mathbf{S}_n \mathbf{b}]^{-\frac{1}{2}(n-1)}. \tag{2.26}$$

Based on the density in (2.22), for each $\mathbf{b} \in R^p \setminus \{\mathbf{0}\}$, we have

$$\begin{aligned}
& \frac{\sup_{\boldsymbol{\theta} \in \mathcal{O}_p^+} P_2(\boldsymbol{\theta} | \bar{\mathbf{X}}_n, \mathbf{S}_n, \mathbf{b})}{P_2(\mathbf{0} | \bar{\mathbf{X}}_n, \mathbf{S}_n, \mathbf{b})} \\
&= \left[\frac{1 + n \mathbf{b}' \bar{\mathbf{X}}_n \bar{\mathbf{X}}_n' \mathbf{b} / \mathbf{b}' \mathbf{S}_n \mathbf{b}}{1 + \inf_{\boldsymbol{\theta} \in \mathcal{O}_p^+} n(\bar{\mathbf{X}}_n - \boldsymbol{\theta})' \mathbf{b} \mathbf{b}' (\bar{\mathbf{X}}_n - \boldsymbol{\theta}) / \mathbf{b}' \mathbf{S}_n \mathbf{b}} \right]^{\frac{1}{2}(n-1)}.
\end{aligned} \tag{2.27}$$

Partition $\mathbf{b}' = (\mathbf{b}'_a, \mathbf{b}'_{a'})$ the same as in (2.9) and for each a , $\emptyset \subseteq a \subseteq P$, let $\mathbf{G}_a = \begin{bmatrix} \mathbf{I}_a & -\mathbf{S}_{naa'} \mathbf{S}_{na'a'}^{-1} \\ \mathbf{0} & \mathbf{I}_{a'} \end{bmatrix}$.

Further multiply \mathbf{G}_a on the left of $\boldsymbol{\theta}$ and $\bar{\mathbf{X}}_n$ and its inverse on the right of \mathbf{b}' . Next by using Lemma 1.1 of Nüesch (1966) and choosing $\mathbf{b}_{a'} = -\mathbf{S}_{na'a'} \mathbf{S}_{naa'} \mathbf{b}_a$, then after some simplifications the *r.h.s.* of (2.27) reduces to

$$\left[1 + \sum_{\emptyset \subseteq a \subseteq P} \sup_{\mathbf{b}_a \neq \mathbf{0}} \left\{ \frac{n(\mathbf{b}'_a \bar{\mathbf{X}}_{na:a'})^2}{\mathbf{b}'_a \mathbf{S}_{naa:a'} \mathbf{b}_a} \right\} I_{na} \right]^{\frac{1}{2}(n-1)}. \tag{2.28}$$

Since (2.28) holds for all $\mathbf{b}_a \neq \mathbf{0}$, $\emptyset \subseteq a \subseteq P$, further note that

$$\begin{aligned}
& \sum_{\emptyset \subseteq a \subseteq P} \sup_{\mathbf{b}_a \neq \mathbf{0}} \left\{ \frac{n(\mathbf{b}'_a \bar{\mathbf{X}}_{na:a'})^2}{\mathbf{b}'_a \mathbf{S}_{naa:a'} \mathbf{b}_a} \right\} I_{na} \\
&= \sum_{\emptyset \subseteq a \subseteq P} \{n \bar{\mathbf{X}}_{na:a'}' \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'}\} I_{na} \\
&= U_n.
\end{aligned} \tag{2.29}$$

If a Bernardo reference prior (Chang and Eaves, 1990) of the form

$$g_1(\boldsymbol{\Sigma}) \propto |\boldsymbol{\Sigma}|^{-\frac{1}{2}(p+1)} |\mathbf{I} + \boldsymbol{\Sigma} * \boldsymbol{\Sigma}^{-1}|^{-\frac{1}{2}} d\boldsymbol{\Sigma}, \quad \boldsymbol{\Sigma} \in \mathcal{M} \tag{2.30}$$

is considered, where $\mathbf{C} * \mathbf{D}$ denotes the Hadamard product of the matrices \mathbf{C} and \mathbf{D} , then the marginal density is

$$h(\mathbf{W}_n) = c \int_{\Sigma \in \mathcal{M}} |\Sigma|^{-\frac{1}{2}(p+1)} |\mathbf{I} + \Sigma * \Sigma^{-1}|^{-\frac{1}{2}} e^{-\frac{1}{2}\text{tr}(\Sigma^{-1}\mathbf{W}_n)} d\Sigma, \quad (2.31)$$

where $\mathbf{W}_n = \mathbf{S}_n + n(\bar{\mathbf{X}}_n - \boldsymbol{\theta})(\bar{\mathbf{X}}_n - \boldsymbol{\theta})'$ and c is the norming constant. By the facts that $d|\mathbf{W}_n| = |\mathbf{W}_n|\text{tr}(\mathbf{W}_n^{-1})(d\mathbf{W}_n)$ and $d\text{tr}(\Sigma^{-1}\mathbf{W}_n) = \text{tr}(\Sigma^{-1})(d\mathbf{W}_n)$, where $(d\mathbf{W}_n)$ denotes the exterior differential form of \mathbf{W}_n . Then

$$\begin{aligned} \frac{dh(\mathbf{W}_n)}{d|\mathbf{W}_n|} &= \frac{-c}{2} \int_{\Sigma \in \mathcal{M}} |\mathbf{W}_n|^{-1} (\text{tr} \mathbf{W}_n^{-1})^{-1} \text{tr}(\Sigma^{-1}) |\Sigma|^{-\frac{1}{2}(p+1)} \\ &\quad \times |\mathbf{I} + \Sigma * \Sigma^{-1}|^{-\frac{1}{2}} e^{-\frac{1}{2}\text{tr}(\Sigma^{-1}\mathbf{W}_n)} d\Sigma \\ &< 0, \end{aligned} \quad (2.32)$$

hence $h(\mathbf{W}_n)$ is strictly decreasing in $|\mathbf{W}_n|$. Thus the corresponding integrated LRT and integrated UIT can be constructed based on $|\mathbf{W}_n|$ only.

Next, we consider the case if Σ is assigned a prior distribution $W_p^{-1}(\Gamma, m)$, where $W_p^{-1}(\Gamma, m)$ denoting the p -dimensional inverted Wishart distribution with m degrees of freedom and expectation $m\Gamma$ which is positive definite. Then by Theorem 7.7.2 of Anderson (2003), the marginal density is

$$\begin{aligned} h_1(\bar{\mathbf{X}}_n, \mathbf{S}_n | \boldsymbol{\theta}) &\propto |\Gamma|^{\frac{m}{2}} |\mathbf{S}_n|^{\frac{1}{2}(n-p-2)} |\mathbf{S}_n + n(\bar{\mathbf{X}}_n - \boldsymbol{\theta})(\bar{\mathbf{X}}_n - \boldsymbol{\theta})'| + \Gamma|^{-\frac{1}{2}(n+m-1)} \\ &= |\Gamma|^{\frac{m}{2}} |\mathbf{S}_n|^{\frac{1}{2}(n-p-2)} |\mathbf{W}_n + \Gamma|^{-\frac{1}{2}(n+m-1)}, \end{aligned} \quad (2.33)$$

which is a monotone function of $|\mathbf{W}_n|$.

Similarly, by (2.25) and (2.29) the L_n^* and U_n^* can be exactly obtained with respect to those non-informative prior differentials mentioned above over the space $\Sigma \in \mathcal{M}$ for the problem of testing against a halfspace \mathcal{H}_p^* .

Let \mathcal{P}_p be the space of positive definite matrices. Riemannian geometry yields an invariant volume element dv on \mathcal{P}_p which is of the form $dv = (\det \Sigma)^{-(p+1)/2} (d\Sigma)$, where $(d\Sigma) = \prod_{1 \leq j \leq i \leq p} d\sigma_{ij}$ with $d\sigma_{ij}$ being the Lebesgue measure on R . We may naturally adopt this relevant Haar measure dv as the underline measure of the a prior of Σ . The likelihood integrated with respect to this relevant Haar measure dv for the nuisance covariace matrix can be viewed as in a Bayesian light. It is not hard to note that for testing against global alternative, the Hotelling's T^2 -test is the version of integrated LRT. Moreover, for the problem (1.4) the corresponding integrated LRT and integrated UIT also have the same forms as those of the corresponding LRT and UIT, respectively. These will reconcile frequentist with Bayesian paradigms via the fiducial inference for the problem (1.4).

THEOREM 2.3. *For the problem of testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{C} \setminus \{\mathbf{0}\}$, where \mathcal{C} is either the positive orthant space \mathcal{O}_p^+ or the halfspace $\mathcal{H}_p^* = \{\boldsymbol{\theta} \mid \theta_p \geq 0\}$, then both the LRT and the UIT are the versions of integrated LRT and integrated UIT respectively.*

3. Power comparison or reporting P -value? Historically, when it comes to hypothesis testing problems, two different approaches are adopted to evaluate the tests for frequentist paradigm: one is Fisher approach to report p -value of the test, the other is Neyman-Pearson approach to carry out the power of test at a fixed significance level. Both Fisher approach and Neyman-Pearson approach are well adopted to measure the performance of the Hotelling's T^2 -test for the problem of testing against the unrestricted alternatives. We would like to know whether the Fisher approach and/or the Neyman-Pearson approach can be well adopted for the problem (1.4). First, consider the problem of testing against the half-space.

Let χ_m^2 be the central chisquare distribution with $m(\geq 0)$ degrees of freedom. Denoted by

$$G_{a,b}(u) = P\{\chi_a^2/\chi_b^2 \leq u\}, \quad u \in R^+, \quad (3.1)$$

and $\bar{G}_{a,b}(u) = 1 - G_{a,b}(u)$, where R^+ denotes the positive real number. Also let

$$\bar{G}_{n,a,p}^*(u) = \int_0^\infty \bar{G}_{a,n-p}\left(\frac{u}{1+t}\right) dG_{p-a,n-p+a}(t), \quad u \in R^+, \quad (3.2)$$

which is the convolution of the d.f.'s of χ_a^2/χ_{n-p}^2 and $1+\chi_{p-a}^2/\chi_{n-p+a}^2$. By (2.15) and (2.16), as such for the problem of testing against the halfspace $\mathcal{H}_p^* \setminus \{\mathbf{0}\}$, it is easy to show that when $\boldsymbol{\Sigma} \in \mathcal{M}$, $\forall c > 0$

$$P_{\mathbf{0},\boldsymbol{\Sigma}}\{L_n^* \geq c\} = \frac{1}{2} [\bar{G}_{p-1,n-p}(c) + \bar{G}_{p,n-p}(c)], \quad (3.3)$$

and

$$P_{\mathbf{0},\boldsymbol{\Sigma}}\{U_n^* \geq c\} = \frac{1}{2} [\bar{G}_{n,p-1,p}^*(c) + \bar{G}_{n,p,p}^*(c)]. \quad (3.4)$$

Notice that (3.3) and (3.4) are free from the nuisance parameter $\boldsymbol{\Sigma}$, namely, the p -values of LRT and UIT do not depend on the nuisance parameter. By the result of Tang (1994) and similar arguments in Theorems 2.1 and 3.1 of Sen and Tsai (1999) we then obtain that both LRT and UIT for testing against a halfspace are similar and unbiased. Hence, for the problem of testing against a halfspace space both approaches of Fisher and Neyman-Pearson can well explain the performance of the LRT and the UIT.

As for the problem of testing against the positive orthant space, it follows from the arguments of

Perlman (1969) and Sen and Tsai (1999) that for every $c > 0$,

$$\begin{aligned} P\{L_n \geq c | H_0, \Sigma\} &= P_{\mathbf{0}, \Sigma}\{L_n \geq c\} \\ &= \sum_{k=1}^p w(p, k; \Sigma) P\{\chi_k^2 / \chi_{n-p}^2 \geq c\} \end{aligned} \quad (3.5)$$

and

$$\begin{aligned} P\{U_n \geq c | H_0, \Sigma\} &= P_{\mathbf{0}, \Sigma}\{U_n \geq c\} \\ &= \sum_{k=1}^p w(p, k; \Sigma) \bar{G}_{n,k,p}^*(c), \end{aligned} \quad (3.6)$$

respectively, where for each $k (=0, 1, \dots, p)$

$$w(p, k; \Sigma) = \sum_{\{a \subseteq P: |a|=k\}} P\{\mathbf{Z}_{a:a'} > \mathbf{0}, \Sigma_{a'a'}^{-1} \mathbf{Z}_{a'} \leq \mathbf{0}\}, \quad (3.7)$$

and $\mathbf{Z} \sim N_p(\mathbf{0}, \Sigma)$ and the partitions are made as in (2.9)-(2.11) (with $(\bar{\mathbf{X}}_n, \mathbf{S}_n)$ being replaced by (\mathbf{Z}, Σ)).

By virtue of (3.5), (3.6) and (3.7), we encounter the problem for which both the null distributions of LRT and UIT statistics depend on the unknown covariance matrix, though the dependence of (3.5) and (3.6) on Σ is only through the $w(p, k; \Sigma)$. Hence, we have difficulty to adopt Fisher approach to report the p -value. On the other hand, according to the definition of level of significance, Perlman (1969) and Sen and Tsai (1999) allowed Σ to vary over the entire class \mathcal{M} and obtained that for every $c > 0$,

$$\sup_{\{\Sigma \in \mathcal{M}\}} P_{\mathbf{0}, \Sigma}\{L_n \geq c\} = \frac{1}{2} [\bar{G}_{p-1, n-p}(c) + \bar{G}_{p, n-p}(c)] \quad (3.8)$$

and

$$\sup_{\{\Sigma \in \mathcal{M}\}} P_{\mathbf{0}, \Sigma}\{U_n \geq c\} = \frac{1}{2} [\bar{G}_{n, p-1, p}^*(c) + \bar{G}_{n, p, p}^*(c)], \quad (3.9)$$

respectively. If the right hand side of (3.8) and (3.9) are equated to α , we then get the critical values of LRT and UIT respectively. However, based on them we will obtain the results which are against our common statistical sense. To proceed them, some notations are needed.

Let $\pi_{\mathbf{A}}(\mathbf{x}; \mathcal{C})$ be the orthogonal projection of \mathbf{x} onto \mathcal{C} with respect to the inner product $\langle, \rangle_{\mathbf{A}}$, then

$$U_n = \|\pi_{\mathbf{S}_n}(n^{1/2} \bar{\mathbf{X}}_n; \mathcal{O}_p^+)\|_{\mathbf{S}_n}^2 \quad (3.10)$$

and

$$U_n^* = \|\pi_{\mathbf{S}_n}(n^{1/2} \bar{\mathbf{X}}_n; \mathcal{H}_p^*)\|_{\mathbf{S}_n}^2. \quad (3.11)$$

PROPOSITION 3.1. *The UIT for testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{C} \setminus \{\mathbf{0}\}$, where \mathcal{C} is a closed convex cone such that $\mathcal{U} \subseteq \mathcal{C} \subseteq \mathcal{H}_p^*$ with \mathcal{U} being a p -dimensional open set and being contained in the halfspace \mathcal{H}_p^* , is dominated by the UIT for testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{H}_p^* \setminus \{\mathbf{0}\}$.*

PROOF. First note that $\mathcal{C} \subseteq \mathcal{H}_p^*$, hence

$$\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^*)\|_{\mathbf{S}_n}^2 \geq \|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{C})\|_{\mathbf{S}_n}^2.$$

Then for every $c > 0$,

$$P_{\boldsymbol{\theta}, \boldsymbol{\Sigma}}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^*)\|_{\mathbf{S}_n}^2 \geq c\} \geq P_{\boldsymbol{\theta}, \boldsymbol{\Sigma}}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{C})\|_{\mathbf{S}_n}^2 \geq c\}.$$

Let $\boldsymbol{\Sigma}^{-1} = \mathbf{B}'\mathbf{B}$, and $\bar{\mathbf{Y}}_n = \mathbf{B}\bar{\mathbf{X}}_n$, and $\mathbf{T}_n = \mathbf{B}\mathbf{S}_n\mathbf{B}'$. Since $\mathbf{B}\mathcal{H}_p^* = \mathcal{H}_p^+$ (another halfspace), thus $P_{\mathbf{0}, \boldsymbol{\Sigma}}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^+)\|_{\mathbf{S}_n}^2 \geq c\} = P_{\mathbf{0}, \mathbf{I}}\{\|\pi_{\mathbf{T}_n}(n^{1/2}\bar{\mathbf{Y}}_n; \mathcal{H}_p^+)\|_{\mathbf{T}_n}^2 \geq c\}$. Notice that $\mathcal{E}(\mathbf{Y}_n) = (n-1)\mathbf{I}$, thus $P_{\mathbf{0}, \boldsymbol{\Sigma}}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^+)\|_{\mathbf{S}_n}^2 \geq c\}$ is independent of the unknown $\boldsymbol{\Sigma}$. Consider the sequence $\{\boldsymbol{\Sigma}_k\}$ and let $\boldsymbol{\Sigma}_k^{-1} = \mathbf{B}'_k\mathbf{B}_k$, for all $k \geq 1$, such that $\mathbf{B}_k(\mathcal{C}_\lambda) \subseteq \mathbf{B}_{k+1}(\mathcal{C}_\lambda)$ and $\cup_{k=1}^\infty \mathbf{B}_k(\mathcal{C}_\lambda) = \text{int}(\mathcal{H}_p^+)$, where \mathcal{C}_λ with $0 < \lambda < 1$ is a right circular cone defined as in 2.3^o of Perlman (1969). The same proof as in Theorem 6.2 of Perlman (1969), we obtain

$$\begin{aligned} P_{\mathbf{0}, \boldsymbol{\Sigma}_k}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^+)\|_{\mathbf{S}_n}^2 \geq c\} &\geq P_{\mathbf{0}, \boldsymbol{\Sigma}_k}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{C})\|_{\mathbf{S}_n}^2 \geq c\} \\ &\geq P_{\mathbf{0}, \boldsymbol{\Sigma}_k}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{C}_\lambda)\|_{\mathbf{S}_n}^2 \geq c\} = P_{\mathbf{0}, \mathbf{I}}\{\|\pi_{\mathbf{S}_n^*}(n^{1/2}\bar{\mathbf{X}}_n^*; \mathbf{B}_k(\mathcal{C}_\lambda))\|_{\mathbf{S}_n^*}^2 \geq c\}, \end{aligned}$$

where $\bar{\mathbf{X}}_n^* = \mathbf{B}_k\bar{\mathbf{X}}_n$, and $\mathbf{S}_n^* = \mathbf{B}_k\mathbf{S}_n\mathbf{B}_k'$. By Lemma 2.8 of Brown (1986), we may conclude that the power functions of the UIT for testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ vs. $H_1 : \boldsymbol{\theta} \in \mathcal{C}$ are analytic on the space $\{\boldsymbol{\theta} \in \mathcal{C}, \boldsymbol{\Sigma} > \mathbf{0}\}$, and hence they are continuous. Thus $\lim_{k \rightarrow \infty} P_{\mathbf{0}, \mathbf{I}}\{\|\pi_{\mathbf{S}_n^*}(n^{1/2}\bar{\mathbf{X}}_n^*; \mathbf{B}_k(\mathcal{C}_\lambda))\|_{\mathbf{S}_n^*}^2 \geq c\} = P_{\mathbf{0}, \mathbf{I}}\{\|\pi_{\mathbf{S}_n^*}(n^{1/2}\bar{\mathbf{X}}_n^*; \mathcal{H}_p^+)\|_{\mathbf{S}_n^*}^2 \geq c\}$, and hence

$$\sup_{\{\boldsymbol{\Sigma} \in \mathcal{M}\}} P_{\mathbf{0}, \boldsymbol{\Sigma}}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{C})\|_{\mathbf{S}_n}^2 \geq c\} = P_{\mathbf{0}, \boldsymbol{\Sigma}}\{\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^+)\|_{\mathbf{S}_n}^2 \geq c\}. \quad (3.12)$$

Thus, the proposition follows.

Q.E.D.

As for the LRT, first, note that the LRT statistics in (2.13) and (2.15) can be represented as

$$L_n = \|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{O}_p^+)\|_{\mathbf{S}_n}^2 \{1 + \|n^{1/2}\bar{\mathbf{X}}_n - \pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{O}_p^+)\|_{\mathbf{S}_n}^2\}^{-1} \quad (3.13)$$

and

$$L_n^* = \|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^*)\|_{\mathbf{S}_n}^2 \{1 + \|n^{1/2}\bar{\mathbf{X}}_n - \pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^*)\|_{\mathbf{S}_n}^2\}^{-1} \quad (3.14)$$

respectively. Also note that for any $\mathcal{C} \subseteq \mathcal{H}_p^*$, $\|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^*)\|_{\mathbf{S}_n}^2 \geq \|\pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{C})\|_{\mathbf{S}_n}^2$, and hence $\|n^{1/2}\bar{\mathbf{X}}_n - \pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{H}_p^*)\|_{\mathbf{S}_n}^2 \leq \|n^{1/2}\bar{\mathbf{X}}_n - \pi_{\mathbf{S}_n}(n^{1/2}\bar{\mathbf{X}}_n; \mathcal{C})\|_{\mathbf{S}_n}^2$. Thus, by similar argument as in the proof of proposition 3.1, we also have the following.

PROPOSITION 3.2. *The LRT for testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{C} \setminus \{\mathbf{0}\}$, where \mathcal{C} is a closed convex cone such that $\mathcal{U} \subseteq \mathcal{C} \subseteq \mathcal{H}_p^*$ with \mathcal{U} being a p -dimensional open set and being contained in the halfspace \mathcal{H}_p^* , is dominated by the LRT for testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_1 : \boldsymbol{\theta} \in \mathcal{H}_p^* \setminus \{\mathbf{0}\}$.*

Proposition 3.2 implies the phenomenon first noted by Tang (1994) that the LRT for testing against the positive orthant space is dominated by that of testing against the halfspace. Note the propositions 3.1 and 3.2 are no longer to be true when the sample size is sufficiently large.

To overcome these unpleasant phenomena, instead of using the conservative maximum principle to define the level of significance α , i.e., taking the supremum of rejected probability under null hypothesis over the set \mathcal{M} , we may use the Bayesian notions, by taking the average of rejected probability under null hypothesis over the set \mathcal{M} with respect to the weight function $g(\boldsymbol{\Sigma})$ of $\boldsymbol{\Sigma}$, to define the level of significance α .

The joint density of $(\bar{\mathbf{X}}_n, \mathbf{S}_n)$ under the null hypothesis is

$$f(\bar{\mathbf{X}}_n, \mathbf{S}_n) = k_0 |\boldsymbol{\Sigma}|^{-\frac{1}{2}n} |\mathbf{S}_n|^{\frac{1}{2}(n-p-2)} e^{-\frac{1}{2}\text{tr}(\boldsymbol{\Sigma}^{-1}\mathbf{V}_n)}, \quad (3.15)$$

where k_0 is the norming constant and $\mathbf{V}_n = \mathbf{S}_n + n\bar{\mathbf{X}}_n\bar{\mathbf{X}}_n'$. First consider the inverted Wishart distribution $W^{-1}(\boldsymbol{\Gamma}, m)$ (Anderson, 2003), which is a proper prior of $\boldsymbol{\Sigma}$, as the weight function. Then we have

$$\begin{aligned} h_1(\bar{\mathbf{X}}_n, \mathbf{S}_n) &= \int_{\boldsymbol{\Sigma} \in \mathcal{M}} f(\bar{\mathbf{X}}_n, \mathbf{S}_n) g(\boldsymbol{\Sigma}) d\boldsymbol{\Sigma} \\ &= k_2 |\boldsymbol{\Gamma}|^{\frac{1}{2}m} |\mathbf{S}_n|^{\frac{1}{2}(n-p-2)} |\mathbf{V}_n + \boldsymbol{\Gamma}|^{-\frac{1}{2}(n+m-1)}, \end{aligned} \quad (3.16)$$

where the constant k_2 can be determined by the equation $\int_{\mathcal{D}} h_1(\bar{\mathbf{X}}_n, \mathbf{S}_n) dx_1 \cdots dx_p = 1$ with $\mathcal{D} = \{(x_1, \dots, x_p) | \mathbf{S}_n \in \mathcal{M} \text{ in probability}\}$. Let

$$b_1(k, n, p) = \sum_{|a|=k} \int_{I_{na} \cap \mathcal{D}} h_1(\bar{\mathbf{X}}_n, \mathbf{S}_n) dx_1 \cdots dx_p, \quad (3.17)$$

then it is easy to note that $b_1(k, n, p) > 0, \forall k = 1, 2, \dots, p$ and $\sum_{k=0}^p b_1(k, n, p) = 1$. Thus by Fubini theorem the critical values of the LRT and the UIT for testing against the positive orthant space can then be determined via the formula

$$\alpha = \sum_{k=0}^p b_1(k, n, p) \bar{G}_{k, n-p}(d_{n, \alpha}^1) \quad (3.18)$$

and

$$\alpha = \sum_{k=0}^p b_1(k, n, p) \bar{G}_{n, k, p}^*(c_{n, \alpha}^1) \quad (3.19)$$

respectively. If the critical values of LRT and UIT are determined by the equations (3.18) and (3.19), then obviously both the LRT and the UIT are similar and unbiased. These conclusions are contrary to the ones made by Perlman (1969) and Sen and Tsai (1999) that both the LRT and the UIT are not similar and biased.

The level of significance α may be defined by using the noninformative priors such as the Haar measure or the Bernardo reference prior, in which the corresponding posterior densities exist, as the weight functions. Suppose that the Haar measure of Σ is taken as the weight function, then we have

$$\begin{aligned} h_2(\bar{\mathbf{X}}_n, \mathbf{S}_n) &= k_0 \int_{\Sigma \in \mathcal{M}} f(\bar{\mathbf{X}}_n, \mathbf{S}_n) |\Sigma_n|^{-\frac{1}{2}(p+1)} d\Sigma \\ &= k_3 |\mathbf{S}_n|^{\frac{1}{2}(n-p-2)} |\mathbf{V}_n|^{-\frac{1}{2}n}, \end{aligned} \quad (3.20)$$

where the constant k_3 can be determined by the equation $\int_{\mathcal{D}} h_2(\bar{\mathbf{X}}_n, \mathbf{S}_n) dx_1 \cdots dx_p = 1$. Let $b_2(k, n, p)$ be defined the same as in (3.5) with $h_2(\bar{\mathbf{X}}_n, \mathbf{S}_n)$ instead of $h_1(\bar{\mathbf{X}}_n, \mathbf{S}_n)$. As such, we may obtain another critical points for the LRT and the UIT. Further note that condition on \mathbf{V}_n , then (3.20) reduces to that of Wang and McDermott (1998) for finding the critical value of semi-conditional LRT.

From the above arguments, we may conclude that the phenomena whether the LRT and the UIT are similar or not depend on the ways of defining the level of significance α . In passing, we note that different weight functions result in different critical values for the LRT and the UIT respectively, as such the powers of the LRT and the UIT heavily depend on the choice of weight functions for the nuisance parameter Σ . Therefore, for a specific test one may have several different power functions if different weight functions are adopted. On the other hand, we may base on a specific weight function to study the power properties and power dominances for some different tests. The phenomena of power dominance for the interesting tests might be different under different weight functions. The crux of using Bayesian notion to define the level of significance lies how to decide what is the optimal weight function for the nuisance parameter. Thus, one of the disadvantages arises: using Bayesian notions to define the level of significance may require extensive numerical computations to get the critical values when $p \geq 4$.

As a consequence, for a specific test when the rejected probability under null hypothesis depends on the nuisance parameter Σ , its power properties such as similarity, unbiasedness and uniformly more powerful phenomenon relative to other tests may be completely different if different ways of obtaining critical values are adopted. In this paper, we regard these different ways of obtaining critical values for a specific test as different kinds of definitions of significance level for that specific test. To give a new definition of the level of significance can be viewed as a way to dig up the

insight information of null hypothesis. Of course, there are other approaches to define the level of significance other than using Bayesian notions and maximum principle.

In passing, by virtue of (3.5) and (3.6) we note that the Fisher approach for p -values of the LRT and the UIT are hardly reported because their null distributions depend on the nuisance parameter. For the problem of testing against the positive orthant space, the main issue of reporting p -value still lies in how to deal with the nuisance parameter. Several studies for handling the nuisance parameter in reporting p -value, which can also be applied to the present problem, have been well investigated in the literature. We may refer to Berger (1999) for the details. One of the main difficulties is how to find an unified way to handle the problems. We are afraid that neither Fisher approach for reporting p -value alone nor Neyman-Pearson approach alone is sound enough to take care the problem of testing against the positive orthant space. Most people think that these two approaches disagree with each other, however Lehmann (1993) asserted that these two approaches are complementary rather than contradictory. In passing, we may note that the essential point of Fisher reporting p -value is to find the optimal ways to obtain the insight information under null hypothesis, and Neyman-Pearson theory essentially shows us how to find the tests that enjoy some optimal power properties. As such, we may agree with Lehmann that there are essentially only one theory, not two, for both Fisher and Neyman-Pearson approaches.

4. D -admissibility. For the problem of testing against the positive orthant space, the difficulty is that both the distribution functions of the LRT and the UIT under null hypothesis are dependent on the nuisance parameter. In the literature, many authors heavily rely on Neyman-Pearson optimal theory to construct new tests so that their new tests dominate the corresponding LRT in the sense of uniformly more powerful. For the details, we refer to Wang and McDermott (1998), Berger (1989), Menendez and Salvador (1991), Menendez et al. (1992) and the references therein. Sen and Tsai (1999) adopted Stein's concept (1945) to establish the two-stage LRT, which enjoys some optimal properties, however the cost is that the complicated distribution function of the test statistic arises. The method proposed by Sen and Tsai (1999) is also essentially based on Neyman-Pearson optimal theory. On the other hand, Perlman and Wu (1999) encouraged to abandon the criteria of α -admissibility, similarity and unbiasedness of Neyman-Pearson theory, and subjectively suggested to use the LRT. Berger (1999) criticized Perlman and Wu's suggestion for not providing any elegant support for the LRT. One of the main goals in this section is to seek a possible resolution for the arguments between Perlman and Wu (1999) and Berger (1999).

For testing against a global shift alternative the Hotelling's T^2 -test is uniformly most power-

ful invariant (UMPI), and hence, is also admissible (Simika, 1941). Stein (1956) established the admissibility of the Hotelling's T^2 -test by using the exponential structure of the parameter space. The UMPI character, or even the admissibility of the Hotelling's T^2 -test may not generally hold for restricted alternatives, such as $H_{1\mathcal{O}^+}$ in (1.4). The affine-invariance structure of the parameter space $\Theta = \{\boldsymbol{\theta} \in R^p\}$ does not hold for $H_{1\mathcal{O}^+}$, and hence, when $\boldsymbol{\Sigma}$ is arbitrary p.d. restriction to invariant tests makes little sense. As such, it is conjectured, though not formally established, that possibly some other non-(affine) invariant tests dominate Hotelling's T^2 -tests, and hence, the latter is inadmissible. Our contention is to establish that for testing H_0 vs $H_{1\mathcal{O}^+}$, the Hotelling's T^2 -test is inadmissible. For testing H_0 against $H_{1\mathcal{O}^+}$, the set of proper Bayes tests and their weak limits only constitute a proper subset of essentially complete class of tests (Marden, 1982). Marden's minimal complete classes consist of proper Bayes tests and the tests with convex and decreasing acceptance regions for the density functions which are more general than exponential family. For the problem (1.4), Theorem 2.1 tells us that the UIT is not a Bayes test. Furthermore, the acceptance regions of the UIT may not be decreasing, though it can be shown as the convex set. Hence, the conditions of Marden's minimal complete classes of tests are also not satisfied for the UIT.

We appraise Eaton's (1970) basic result on essentially complete class of test functions for testing against restricted alternatives, when the underlying density belongs to an exponential family as in our present context. Let Φ be Eaton's essentially complete class of tests, so for any test $\varphi^* \notin \Phi$ there exists a test $\varphi \in \Phi$ such that φ is at least as good as φ^* .

THEOREM 4.1. *For the problem of testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_{1\mathcal{O}^+} : \boldsymbol{\theta} \in \mathcal{O}_p^+ \setminus \{\mathbf{0}\}$, the Hotelling's T^2 -test is inadmissible.*

PROOF. Note that the present testing hypothesis problem is invariant under the group of transformations of positive diagonal matrices, hence, for simplicity, $\boldsymbol{\Sigma}$ may be treated as the correlation matrix. Following Eaton (1970), we define

$$\Omega_1 = \{\boldsymbol{\Sigma}^{-1}\boldsymbol{\theta} \mid \boldsymbol{\theta} \in \mathcal{O}_p^+ \setminus \{\mathbf{0}\}\}. \quad (4.1)$$

Let $\mathcal{V} \subseteq R^p$ be the smallest closed convex cone containing Ω_1 . Then the dual cone of \mathcal{V} is defined as

$$\mathcal{V}^- = \{\mathbf{w} \mid \langle \mathbf{w}, \mathbf{x} \rangle \leq 0, \forall \mathbf{x} \in \mathcal{V}\}. \quad (4.2)$$

Then, by (4.1) we have that \mathcal{V} becomes a halfspace and its dual cone is

$$\mathcal{V}^- = \{\boldsymbol{\theta} \in R^p \mid \sum_{i=1}^p \theta_i = 0\}, \quad (4.3)$$

which is an unbounded hyperplane.

The acceptance region of the Hotelling's T^2 -test is given by

$$\mathcal{A}_{T^2} = \{(\bar{\mathbf{X}}, \mathbf{S}) \mid T^2 \leq t_\alpha^2\}, \quad (4.4)$$

where t_α^2 is the upper $100\alpha\%$ point of the null hypothesis distribution of T^2 (which is linked to a F-distribution). Since \mathcal{A}_{T^2} is a hyperellipsoidal set with origin $\mathbf{0}$, it is bounded while \mathcal{V}^- , as shown above, is unbounded. Therefore, Eaton's (1970) condition is not tenable. Hence the Hotelling T^2 -test is not a member of essentially complete class. Q.E.D.

Next, for testing H_0 against $H_{1\mathcal{O}^+}$ we will show that the finite union-intersection test (FUIT) (Tsai and Sen, 2004) and the union-intersection test (UIT) (Sen and Tsai, 1999) are the members of Eaton's (1970) essentially complete class though they are not the proper Bayes. A test which is a member of Eaton's (1970) essentially complete class of tests can be established by showing the acceptance region contains the subspace \mathcal{V}^- . Corresponding to a preassigned α ($0 < \alpha < 1$), let c_α be the critical level, obtained by equating the right hand side of (3.9) to α , thus the UIT is a size- α test for H_0 vs $H_{1\mathcal{O}^+}$. Let \mathcal{A}_U be the acceptance region formed by letting $U_n \leq c_\alpha$ in (2.14). Partition the sample space R^p into $\cup_a I_{na}$, and for each a , $\emptyset \subseteq a \subseteq P$, let $\mathcal{A}_a = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n\bar{\mathbf{X}}'_{na:a'} \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'} \leq c_\alpha\} I_{0na}$. Treating $\mathcal{V}_n^- = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \mathbf{S}_n^{-1} \bar{\mathbf{X}}_n \leq \mathbf{0}\} = I_{0n\emptyset}$ as the skelton (pivotal set), then by (2.17) we have that $\mathcal{A}_U = \mathcal{V}_n^- \cup_{attach} \mathcal{A}_a$, where \cup_{attach} means that for each a , $\emptyset \subset a \subseteq P$, the hyperspace \mathcal{A}_a is attached to the boundary of \mathcal{V}_n^- on the subspace I_{0na} . By the property that \mathbf{S}_n is positive definite with probability one, we have $\mathcal{V}^- \subseteq \mathcal{V}_n^- \subseteq \mathcal{A}_U - \mathbf{a}^0$ for each $\mathbf{a}^0 \in \partial\mathcal{A}_U$. Similarly, for the FUIT we may note that its acceptance set \mathcal{A}^* [see (2.21)] is a closed convex set and $R_p^- = \{\mathbf{x} \in R^p \mid \mathbf{x} \leq \mathbf{0}\} \subseteq \mathcal{A}^*$ as long as $t_{n-1, \alpha^*} \geq 0$ for $\alpha^* \leq \frac{1}{2}$. Thus, the FUIT is a size- α test for H_0 vs $H_{1\mathcal{O}^+}$ and $\mathcal{V}^- \subseteq \mathcal{A}^*$.

THEOREM 4.2. *In the same setup of Theorem 4.1, both the UIT and the FUIT belong to Eaton's essentially complete class of tests.*

For the problem of testing H_0 against $H_{1\mathcal{O}^+}$, by virtues of Theorems 4.1 and 4.2 it is interesting to see whether the Hotelling's T^2 -test is power dominated by the UIT. For the case that both the Hotelling's T^2 -test and the UIT are unbiased, Tsai and Sen (2004) provided an affirmed answer for it when $\Sigma = \sigma^2 \Delta$, where σ^2 is an unknown scalar parameter and Δ is a known M -matrix. However, the situation is still open when Σ is totally unknown. Theorem 4.2 does not guarantee that the UIT (based on the test statistic U_n) is admissible. First, we examine the admissibility for the corresponding UIT test based on the test statistic U_n^* for the problem of testing against the halfspace.

THEOREM 4.3. *For the problem of testing $H_0 : \theta = \mathbf{0}$ against $H_{1\mathcal{H}^*} : \theta \in \mathcal{H}_p^* \setminus \{\mathbf{0}\}$, the UIT is d -admissible and α -admissible.*

PROOF. The acceptance region of UIT is

$$\mathcal{A}_{U^*} = \left\{ (\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \sum_{P_1 \subseteq a \subseteq P} \{n\bar{\mathbf{X}}'_{na:a'} \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'}\} I_{na}^* \leq k_1, \mathbf{S}_n \text{ positive definite} \right\}, \quad (4.5)$$

where $P_1 = P - \{p\}$ for a suitable k_1 . Note that

$$\mathcal{A}_{U^*} = \mathcal{A}_{U^*P} \cup \mathcal{A}_{U^*P_1}, \quad (4.6)$$

where

$$\mathcal{A}_{U^*P} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n\bar{\mathbf{X}}'_n \mathbf{S}_n^{-1} \bar{\mathbf{X}}_n \leq k_1, \bar{X}_{np} \geq 0\}, \quad (4.7)$$

and

$$\mathcal{A}_{U^*P_1} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n\bar{\mathbf{X}}'_{nP_1:p} \mathbf{S}_{nP_1P_1:p}^{-1} \bar{\mathbf{X}}_{nP_1:p} \leq k_1, \bar{X}_{np} \leq 0\} \quad (4.8)$$

respectively. For each $a, P_1 \subseteq a \subseteq P$, Theorem 2.1 of Tsai (2003) tells us that the function $\{n\bar{\mathbf{X}}'_{na:a'} \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'}\}$ is convex in $(\bar{\mathbf{X}}_n, \mathbf{S}_n)$. Thus the sets \mathcal{A}_{U^*P} and $\mathcal{A}_{U^*P_1}$ are convex. Notice that the set \mathcal{A}_{U^*P} is a half-ball in the space $\{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \bar{X}_{np} \geq 0\}$ and the set $\mathcal{A}_{U^*P_1}$ is an unbounded closed cylinder-type convex cone in the space $\{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \bar{X}_{np} \leq 0\}$. Also the extreme set (Rockafellar, 1972) of \mathcal{A}_{U^*P} on the hyperplane $\{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \bar{X}_{np} = 0\}$ is identical to the one of $\mathcal{A}_{U^*P_1}$ on the hyperplane $\{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \bar{X}_{np} = 0\}$. The sets \mathcal{A}_{U^*P} and $\mathcal{A}_{U^*P_1}$ are not disjoint since $\mathcal{A}_{U^*P_1} = \mathcal{A}_{U^*P}$ on the hyperplane $\{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \bar{X}_{np} = 0\}$, actually the intersection of these two sets is identical to the extreme set of \mathcal{A}_{U^*P} on the hyperplane $\{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \bar{X}_{np} = 0\}$. The set \mathcal{A}_{U^*} is the convex hull of $\mathcal{A}_{U^*P_1}$ and \mathcal{A}_{U^*P} . Thus the set \mathcal{A}_{U^*} is convex, the probability of $\partial\mathcal{A}_{U^*}$ is zero.

Since \mathcal{A}_{U^*} is convex on $(\bar{\mathbf{X}}_n, \mathbf{S}_n)$, we assume \mathcal{A}_{U^*} is disjoint with the halfspace

$$\mathcal{H}_c = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \boldsymbol{\omega}' \mathbf{y} = \boldsymbol{\nu}' \bar{\mathbf{X}}_n - \frac{1}{2} \text{tr} \boldsymbol{\Lambda} \mathbf{V}_n > c, \} \quad (4.9)$$

where $\boldsymbol{\Lambda}$ is a symmetric matrix and \mathbf{V}_n is p.d.. Let $\mathcal{H}_p^- = \{\boldsymbol{\theta} \in R^p \mid \sum_{i=1}^p \theta_i \leq 0\}$ and \mathcal{H}_p^\perp be the halfspace which contains the hyperplane $\mathcal{H}_p^0 = \{\boldsymbol{\theta} \in R^p \mid \theta_1 = \dots = \theta_{p-1} = 0, \theta_p < 0\}$ and is perpendicular to the hyperplane $\mathcal{H}_p^1 = \{\boldsymbol{\theta} \in R^p \mid \sum_{i=1}^p \theta_i = 0\}$. For $\boldsymbol{\Sigma}$ being any arbitrarily positive definite matrices, $\boldsymbol{\Sigma}^{-1} : R^{p(p+1)/2} \rightarrow R^{p(p+1)/2}$ is a continuous, linear and open mapping on the general linear group $GL(R^{p(p+1)})$. Let $\mathcal{S} = \mathcal{H}_p^- \cap \mathcal{H}_p^\perp$, thus $\boldsymbol{\theta} \in \mathcal{H}_p^*$ implies that $\boldsymbol{\nu} = n\boldsymbol{\Sigma}^{-1}\boldsymbol{\theta} \in \overline{R^p \setminus \mathcal{S}}$. Here we want to show that $\boldsymbol{\omega}_0 + \lambda\boldsymbol{\omega} \in \overline{(R^p \setminus \mathcal{S})} \setminus \{\mathbf{0}\}$ holds, namely to show $\mathbf{I} + \lambda\boldsymbol{\Lambda}$ is positive definite (it suffices to show that $\boldsymbol{\Lambda}$ is positive semidefinite) and $\boldsymbol{\nu}_0 + \lambda\boldsymbol{\nu} \in \overline{(R^p \setminus \mathcal{S})} \setminus \{\mathbf{0}\}$ for $\lambda > 0$.

(I). To show that $\boldsymbol{\Lambda}$ is positive semidefinite. Suppose that $\boldsymbol{\Lambda}$ is not, and then assume that it is

$$\boldsymbol{\Lambda} = \mathbf{D} \begin{bmatrix} \mathbf{I}_{b_1} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & -\mathbf{I}_{b_2} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0}_{b_3} \end{bmatrix} \mathbf{D}',$$

where \mathbf{D} is nonsingular, and $1 \leq b_2 < p, 1 \leq b_1 + b_2 < p$ and $\sum_{i=1}^3 b_i = p$. Take $\bar{\mathbf{X}}_n = \frac{1}{\lambda} \bar{\mathbf{Z}}_n$ and

$$\mathbf{V}_n = \lambda(\mathbf{D}')^{-1} \begin{bmatrix} \mathbf{I}_{b_1} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_{b_2} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_{b_3} \end{bmatrix} \mathbf{D}^{-1} \quad (4.10)$$

for positive λ . Then

$$\boldsymbol{\omega}' \mathbf{y} = \frac{1}{\lambda} \boldsymbol{\nu}' \bar{\mathbf{Z}}_n - \frac{\lambda}{2} \text{tr} \begin{bmatrix} \mathbf{I}_{b_1} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & -\mathbf{I}_{b_2} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0}_{b_3} \end{bmatrix},$$

which is greater than c , a positive constant. Let the space $\mathcal{D}^* = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \mathbf{S}_n \text{ is p.d. and } \mathbf{V}_n \text{ has the form in (4.10)}\}$. Thus, the halfspace in (4.9) becomes

$$\mathcal{H}_c \cap \mathcal{D}^* = \{\bar{\mathbf{Z}}_n \mid \boldsymbol{\nu}' \bar{\mathbf{Z}}_n > \lambda c - \frac{\lambda^2}{2}(b_1 - b_2)\}. \quad (4.11)$$

Let

$$\mathcal{A}_{T^2, k_1} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n \bar{\mathbf{X}}_n' \mathbf{S}_n^{-1} \bar{\mathbf{X}}_n \leq k_1\}. \quad (4.12)$$

Note that

$$\mathcal{A}_{U^*} \supseteq \mathcal{A}_{T^2, k_1} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n \bar{\mathbf{X}}_n' \mathbf{V}_n^{-1} \bar{\mathbf{X}}_n \leq k_1^*\} \quad (4.13)$$

for a suitable positive constant k_2 . Thus,

$$\mathcal{A}_{T^2, k_1} \cap \mathcal{D}^* = \{\bar{\mathbf{Z}}_n \mid n \bar{\mathbf{Z}}_n' \mathbf{D} \mathbf{D}' \bar{\mathbf{Z}}_n \leq \lambda^3 k_1^*\}. \quad (4.14)$$

By virtue of (4.11), (4.13) and (4.14), the intersection of $\mathcal{A}_{U^*} \cap \mathcal{D}^*$ and $\mathcal{H}_c \cap \mathcal{D}^*$ is nonempty for sufficiently large λ . This contradicts the fact that (4.5) and (4.9) are disjoint. Thus $\mathbf{\Lambda}$ is positive semidefinite.

(II). To show that $\boldsymbol{\nu}_0 + \lambda \boldsymbol{\nu} \in (\overline{R^p \setminus \mathcal{S}}) \setminus \{\mathbf{0}\}$ for $\lambda > 0$. Our aim is to show that $\boldsymbol{\nu} \in (\overline{R^p \setminus \mathcal{S}}) \setminus \{\mathbf{0}\}$.

To proceed, we shall work with another halfspace

$$\mathcal{H}_c^a = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n \boldsymbol{\nu}' \bar{\mathbf{X}}_n - \frac{1}{2} \text{tr}[(\mathbf{I} + \mathbf{\Lambda})(\mathbf{S}_n + n \bar{\mathbf{X}}_n \bar{\mathbf{X}}_n')]\} > c\}. \quad (4.15)$$

Note that \mathbf{S}_n is p.d. with probability one, thus $(\boldsymbol{\Gamma}, \mathbf{S}_n) \in \mathcal{H}_c^a$ implies that $(\bar{\mathbf{X}}_n, \mathbf{S}_n) \in \mathcal{H}_c$. Hence, we have

$$\mathcal{H}_c^a \subset \mathcal{H}_c. \quad (4.16)$$

Consequently, the intersection of \mathcal{A}_{U^*} and \mathcal{H}_c^a is empty too. Note that $\mathbf{I} + \mathbf{\Lambda}$ is p.d., thus we may denote its inverse by $\mathbf{\Gamma}$, i.e., $(\mathbf{I} + \mathbf{\Lambda})^{-1} = \mathbf{\Gamma}$. Let $\tilde{\boldsymbol{\nu}} = \mathbf{\Gamma}\boldsymbol{\nu}$ and $\bar{\mathbf{Z}}_n = \mathbf{\Gamma}^{-1}\bar{\mathbf{X}}_n$, then \mathcal{H}_c^a becomes

$$\mathcal{H}_c^a = \{(\bar{\mathbf{Z}}_n, \mathbf{S}_n) \mid n\tilde{\boldsymbol{\nu}}'\bar{\mathbf{Z}}_n - \frac{1}{2}n\bar{\mathbf{Z}}_n'\mathbf{\Gamma}\bar{\mathbf{Z}}_n - \frac{1}{2}\text{tr}(\mathbf{\Gamma}^{-1}\mathbf{S}_n) > c\}. \quad (4.17)$$

And then $\boldsymbol{\nu} \in (\overline{R^p \setminus \mathcal{S}}) \setminus \{\mathbf{0}\}$ implies that $\tilde{\boldsymbol{\nu}} \in R^p \setminus \{\mathbf{0}\}$, hence our aim becomes to show that $\tilde{\boldsymbol{\nu}} \in R^p \setminus \{\mathbf{0}\}$. Suppose that the statement is not true, then we have to show that $\tilde{\boldsymbol{\nu}} = \mathbf{0}$ leads to a contradiction.

Take $\mathbf{S}_n = \mathbf{\Gamma}$ and let $\tilde{\boldsymbol{\nu}} = \mathbf{0}$, then we have

$$\mathcal{H}_c^a = \{\bar{\mathbf{Z}}_n \mid n\bar{\mathbf{Z}}_n'\mathbf{\Gamma}\bar{\mathbf{Z}}_n \leq -(p + 2c)\}. \quad (4.18)$$

Also note that given $\mathbf{S}_n = \mathbf{\Gamma}$, \mathcal{A}_{T^2, k_1} becomes

$$\mathcal{A}_{T^2, k_1} = \{\bar{\mathbf{Z}}_n \mid n\bar{\mathbf{Z}}_n'\mathbf{\Gamma}\bar{\mathbf{Z}}_n \leq 2k_1^*\}. \quad (4.19)$$

It is obvious that $\mathcal{H}_c^a \cap \mathcal{A}_{T^2, k_1} \neq \emptyset$, and hence $\mathcal{H}_c^a \cap \mathcal{A}_{U^*} \neq \emptyset$. This contradicts the fact that (4.5) and (4.9) are disjoint. Hence the condition that $\boldsymbol{\omega}_1 + \lambda\boldsymbol{\omega} \in \mathcal{H}_p^*$ for an arbitrarily large λ is satisfied, namely the conditions of Theorem 8 of Lehmann (1986) hold. Thus by Theorem 8 of Lehmann (1986), we conclude that for testing against the halfspace the UIT is d -admissible.

With similar arguments as in Theorems 2.1 and 3.1 of Sen and Tsai (1999), we may show that the UIT is similar and unbiased. Therefore by Corollary 2 of Lehmann (1986), the UIT is α -admissible. Q.E.D.

For the problem of testing H_0 against $H_{1\mathcal{O}^+}$, it is interesting to see whether the UIT are admissible in decision-theoretic sense. Our problems are under the set up of exponential family, Theorem 8 of Lehmann (1986), which states the conditions for tests to be d -admissible for the problems of testing against restricted alternatives, can be viewed as a generalization of Birnbaum (1955) and Stein (1956). Based on the results of Theorem 8, Corollary 2 of Lehmann (1986) and Theorem 2.1 of Tsai (2003) we prove the d -admissibilities of UIT for the problem (1.4). With arguments parallel to those in the proof of Theorem 4.3, we also have the following.

THEOREM 4.4. *For the problem of testing $H_0 : \boldsymbol{\theta} = \mathbf{0}$ against $H_{1\mathcal{O}^+} : \boldsymbol{\theta} \in \mathcal{O}_p^+ \setminus \{\mathbf{0}\}$, the UIT is d -admissible.*

OUTLINE OF THE PROOF. The acceptance region of UIT is

$$\mathcal{A}_U = \left\{ (\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \sum_{\emptyset \subseteq a \subseteq P} \{n\bar{\mathbf{X}}_{na:a'}'\mathbf{S}_{naa:a'}^{-1}\bar{\mathbf{X}}_{na:a'}\} I_{na} \leq k_2, \mathbf{S}_n \text{ positive definite} \right\}, \quad (4.20)$$

for a suitable k_2 . Note that

$$\mathcal{A}_U = \cup_{\emptyset \subseteq a \subseteq P} \mathcal{A}_{Ua}, \quad (4.21)$$

where

$$\mathcal{A}_{Ua} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid n\bar{\mathbf{X}}'_{na:a'} \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'} \leq k_2, \bar{\mathbf{X}}_{na:a'} > \mathbf{0}, \mathbf{S}_{na'a'}^{-1} \bar{\mathbf{X}}_{na'} \leq \mathbf{0}\}. \quad (4.22)$$

For each a , $\emptyset \subseteq a \subseteq P$, the function $\{n\bar{\mathbf{X}}'_{na:a'} \mathbf{S}_{naa:a'}^{-1} \bar{\mathbf{X}}_{na:a'}\} I_{na}$ is convex in $(\bar{\mathbf{X}}_n, \mathbf{S}_n)$ for \mathbf{S}_n positive definite. Thus each \mathcal{A}_{Ua} is convex. Notice that any two sets \mathcal{A}_{Ua} and \mathcal{A}_{Ub} are not disjoint, where $\emptyset \subseteq a, b \subseteq P$ with $\|a\| - \|b\| = 1$ and the intersection of these two sets is identical to their common extreme set. Treating $\mathcal{A}_{U\emptyset} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) \mid \mathbf{S}_n^{-1} \bar{\mathbf{X}}_n \leq \mathbf{0}\}$ as the skeleton, we may note that $\mathcal{A}_U = \mathcal{A}_{U\emptyset} \cup_{attach} \mathcal{A}_{Ua}$, where \cup_{attach} means that for each a , $\emptyset \subset a \subseteq P$, the hyperspace \mathcal{A}_{Ua} is attached to the common extreme set with the set of $\mathcal{A}_{U\emptyset}$. The set \mathcal{A}_U is the convex hull of \mathcal{A}_{Ua} , $\emptyset \subset a \subseteq P$. Thus the set \mathcal{A}_U is convex; the probability of $\partial\mathcal{A}_U$ is zero.

Assume \mathcal{A}_U is disjoint with the halfspace in (4.9). Note that $\boldsymbol{\theta} \in \mathcal{O}_p^+$ implies that $\boldsymbol{\nu} = n\boldsymbol{\Sigma}^{-1}\boldsymbol{\theta} \in \overline{R^p \setminus \mathcal{H}_p^-}$, where $\mathcal{H}_p^- = \{\boldsymbol{\theta} \in R^p \mid \sum_{i=1}^p \theta_i \leq 0\}$. Here we want to show that $\boldsymbol{\omega}_1 + \lambda\boldsymbol{\omega} \in (\overline{R^p \setminus \mathcal{H}_p^-}) \setminus \{\mathbf{0}\}$ holds, i.e., $\boldsymbol{\nu}_1 + \lambda\boldsymbol{\nu} \in (\overline{R^p \setminus \mathcal{H}_p^-}) \setminus \{\mathbf{0}\}$ and $\boldsymbol{\Lambda}$ is positive semidefinite for $\lambda > 0$. We shall take $\boldsymbol{\nu}_1 = \mathbf{0}$. With similar arguments as in (1) of Theorem 4.3, we can claim that $\boldsymbol{\Lambda}$ is positive semidefinite. We also adopt the notations $\tilde{\boldsymbol{\nu}}$, $\bar{\mathbf{Z}}_n$ and \mathcal{H}_c^a from Theorem 4.3. Thus, $\boldsymbol{\nu} \in (\overline{R^p \setminus \mathcal{H}_p^-})$ implies that $\tilde{\boldsymbol{\nu}} \in \overline{R_p^-}$, where $R_p^- = \{\mathbf{x} \in R^p \mid \mathbf{x} \leq \mathbf{0}\}$.

Take $\mathbf{S}_n = \boldsymbol{\Gamma}$, then we have

$$\begin{aligned} \mathcal{H}_c^a &= \{\bar{\mathbf{Z}}_n \mid 2n\tilde{\boldsymbol{\nu}}'\bar{\mathbf{Z}}_n - n\bar{\mathbf{Z}}_n'\boldsymbol{\Gamma}\bar{\mathbf{Z}}_n \geq p + 2c\} \\ &= \{\bar{\mathbf{Z}}_n \mid n(\bar{\mathbf{Z}}_n - \boldsymbol{\Gamma}^{-1}\tilde{\boldsymbol{\nu}})'\boldsymbol{\Gamma}(\bar{\mathbf{Z}}_n - \boldsymbol{\Gamma}^{-1}\tilde{\boldsymbol{\nu}}) \leq n\tilde{\boldsymbol{\nu}}'\boldsymbol{\Gamma}^{-1}\tilde{\boldsymbol{\nu}} - (p + 2c)\} \end{aligned} \quad (4.23)$$

Our aim is to show that $\tilde{\boldsymbol{\nu}} \in (\overline{R^p \setminus \mathcal{H}_p^-}) \setminus \{\mathbf{0}\}$. Suppose it is not true, then $\tilde{\boldsymbol{\nu}} \in R_p^-$ and hence \mathcal{H}_c^a contains an open ball in the space $\{\bar{\mathbf{Z}}_n \mid \bar{\mathbf{Z}}_n \leq \mathbf{0}\}$. Given $\mathbf{S}_n = \boldsymbol{\Gamma}$, it is easy to see that $\mathcal{A}_{U\emptyset} = \{\bar{\mathbf{Z}}_n \mid \bar{\mathbf{Z}}_n \leq \mathbf{0}\}$. Thus, we have $\mathcal{H}_c^a \cap \mathcal{A}_{T^2, k_1} \neq \emptyset$, and hence $\mathcal{H}_c^a \cap \mathcal{A}_{U^*} \neq \emptyset$. This contradicts the fact that (4.5) and (4.9) are disjoint. Hence the condition that $\boldsymbol{\omega}_1 + \lambda\boldsymbol{\omega} \in (\overline{R^p \setminus \mathcal{H}_p^-}) \setminus \{\mathbf{0}\}$ for an arbitrarily large λ is satisfied, namely the conditions of Theorem 8 of Lehmann (1986) hold. Therefore by Theorem 8 of Lehmann (1986), we conclude that for testing against the positive orthant space the UIT is d -admissible. Q.E.D.

For the problem of testing H_0 against $H_{1\mathcal{O}^+}$, Proposition 3.1 shows that the UIT based on the test statistic U_n is α -inadmissible because its power is dominated by that of the corresponding UIT based on the test statistic U_n^* , this result is against our common statistical sense. This is due to

the fact that the power function of UIT based on U_n under null hypothesis depends on the nuisance parameter Σ as discussed in above section. To avoid this unpleasant phenomeno in this situation, the spirits lies in the compromise of Fisher approach emphasizing on the type I erros and Neyman-Pearson optimal theory without detailed consideration of power and power comparisions of tests would make sense only in the situation when the tests are similar. As such, Wald's decision theory via d -admissibility and Neyman-Pearson's optimal theory is essentially one theory, not two. This way can also provide a resolution for the arguments between Perlman and Wu (1999) and Berger (1999). Therefore for the problem of testing H_0 against $H_{1\mathcal{O}^+}$, we suggest use the unified conservative maximum principle to define the level of significance and then adopt the criterion in decision-theoretic sense (d -admissibility) for the power dominance problems.

5. The difficulty of LRT. When Σ is totally unknown, Proposition 2.1 fails to characterize whether or not the LRT is the generalized Bayes test. Also, let l_α be the critical point of the level of significance α for the LRT for the problem of testing against the positive orthant space, which can be obtained from the right hand sides of (3.8), and $\mathcal{A}_L = \cup_{\emptyset}^P \mathcal{A}_{La}$ be the acceptance region of the LRT, where $\mathcal{A}_{La} = \{(\bar{\mathbf{X}}_n, \mathbf{S}_n) | n\bar{\mathbf{X}}'_{na:a'}\mathbf{S}_{naa:a'}^{-1}\bar{\mathbf{X}}_{na:a'}/(1+n\bar{\mathbf{X}}'_{na'a'}\mathbf{S}_{na'a'}^{-1}\bar{\mathbf{X}}_{na'a'}) \leq l_\alpha\}I_{na}$, $\emptyset \subseteq a \subseteq P$. For simplicity, we take $p = 2$ and \mathbf{S}_n be the diagonal matrix, then it is easy to see that the acceptance region \mathcal{A}_L is not convex. Similar arguments, it is true that the acceptance region of the LRT for testing against the half-space is not convex. In the literature, most of the results of admissibility for tests are established based on the fundamental assumption that the acceptance region is convex. The nonconvexity of acceptance region of the LRT may cause the difficulty for the question whether the LRT is d -admissible for problem (1.4). We will put it as the project for future study.

REFERENCES

- ANDERSON, T. W. (2003). *An Introduction to Multivariate Statistical Analysis*, 3rd edition. Wiely, New York.
- BASU, D. (1977). On the elimination of nuisance parameters. *J. Amer. Statist. Assoc.* **72** 355-366.
- BERGER, J. O., LISEO, B. and WOLPERT, R. L. (1999). Integrated likelihood methods for eliminating nuisance parameters. *Statist. Sci.* **14** 1-28.
- BERGER, R. L. (1989). Uniformly more powerful tests for hypotheses concerning linear inequalities and normal means. *J. Amer. Statist. Assoc.* **84** 192-199.
- BERGER, R. L. (1999). Comment on "The emperor's new tests", *Statist. Sci.* **14** 370-373.

- BIRNBAUM, A. (1955). Characterization of complete classes of tests of some multiparametric hypothesis, with applications to likelihood ratio tests. *Ann. Math. Statist.* **26** 21-36.
- BROWN, L. D. (1986). *Fundamentals of Statistical Exponential families, with Application in Statistical Decision Theory*. Institute of Mathematical Statistics, Lecture Notes-Monograph Series, Volume **9**. Hayward, California.
- CHANG, T. and EAVES, D. (1990). Reference priors for the orbit in a group model. *Ann. Statist.* **18** 1595-1614.
- EATON, M. L. (1970). A complete class theorem for multidimensional one sided alternatives. *Ann. Math. Statist.* **41** 1884-1888.
- KIEFER, J. and SCHWARTZ, R. (1965). Admissible Bayes character of T^2- , R^2- , and other fully invariant tests for classical multivariate normal problems. *Ann. Math. Statist.* **36** 747-770.
- LEHMANN, E. L. (1986). *Testing Statistical Hypotheses*, 2nd edition. Wiley, New York.
- LEHMANN, E. L. (1993). The Fisher, Neyman-Pearson theories of testing hypotheses: One theory or two? *J. Amer. Statist. Assoc.* **88** 1242-1249.
- MARDEN, J. I. (1982). Minimal complete classes of tests of hypothesis with multivariate one-sided alternatives. *Ann. Statist.* **10** 962-970.
- MENENDEZ, J. A. and SALVADOR, B. (1991). Anomalies of the likelihood ratio test for testing restricted hypotheses. *Ann. Statist.* **19** 889-898.
- MENENDEZ, J. A., RUEDA, C. and SALVADOR, B. (1992). Dominance of likelihood ratio tests under cone constraints. *Ann. Statist.* **20** 2087-2099.
- NÜESCH, P. E. (1966). On the problem of testing location in multivariate problems for restricted alternatives. *Ann. Math. Statist.* **37** 113-119.
- PERLMAN, M. D. (1969). One-sided problems in multivariate analysis. *Ann. Math. Statist.* **40** 549-567.
- PERLMAN, M. D. and WU, L. (1999). The emperor's new tests. *Statist. Sci.* **14** 355-369.
- ROCKAFELLAR, R. T. (1972). *Convex Analysis*. Princeton University Press.
- ROY, S. N. (1953). On a heuristic method of test construction and its use in multivariate analysis. *Ann. Math. Statist.* **24** 220-238.

- ROY, S. N., GNANADESIKAN, R. and SRIVASTAVA, J.N. (1972). *Analysis and Deign of Certain Quantitative Multiresponse Experiments*. Pergamon Press, Oxford.
- SEN, P. K. and TSAI, M. T. (1999). Two-stage LR and UI tests for one-sided alternatives: multivariate mean with nuisance dispersion matrix. *J. Multivariate Anal.* **68** 264-282.
- SIMAIIKA, J. B. (1941). On an optimum property of two important statistical tests. *Biometrika* **32** 70-80.
- STEIN, C. (1945). A two-sample test for a linear hypothesis whose power function is independent of σ . *Ann. Math. Statist.* **16** 243-258.
- STEIN, C. (1956). The admissibility of Hotelling's T^2 -test. *Ann. Math. Statist.* **27** 616-623.
- TANG, D. I. (1994). Uniformly more powerful tests in a one-sided multivariate problem. *J. Amer. Statist. Assoc.* **89** 1006-1011.
- TSAI, M. T. (2003). On the invariant tests for means with covariates. *TR 2003-05*. Technical Report, Institute of Statistical Science, Academia Sinica.
- TSAI, M. T. and SEN, P. K. (2004). On inadmissibility of Hotelling's T^2 -test for restricted alternatives. *J. Multivariate Anal.* **89** 87-96.
- WALD, A. (1950). *Statistical Decision Functions*. Wiely, New York.
- WANG, Y. and MCDERMOTT, M. P. (1998). Conditional likelihood ratio test for a nonnegative normal mean vector. *J. Amer. Statist. Assoc.* **93** 380-386.

INSTITUTE OF STATISTICAL SCIENCE
ACADEMIA SINICA
TAIPEI 11529, TAIWAN
E-mail: mtttsai@stat.sinica.edu.tw