

On the connection between a skew product IFS and the ergodic optimization for a finite family of potentials

Elismar R. Oliveira* ¹

¹Universidade Federal do Rio Grande do Sul, Instituto de Matemática e Estatística
- UFRGS, Av. Bento Gonçalves, 9500, Porto Alegre - 91500 - RS -Brasil

August 24, 2018

Abstract

We study the skew product IFS on the cylinder $\mathbf{S}^1 \times \mathbb{R}$ defined by the maps $G_c(x, y) = (T(x), A_c(x) + \lambda y)$, $c \in \mathcal{C}$ for a finite family \mathcal{C} of $m \geq 2$ potential functions $A_c : \mathbf{S}^1 \rightarrow \mathbb{R}$ where $T : \mathbf{S}^1 \rightarrow \mathbf{S}^1$ is the doubling map. We show that there exists a compact invariant set $\Lambda \subseteq \mathbf{S}^1 \times \mathbb{R}$ which has an attractive behavior. Then, we introduce the random SRB measure $\mu \in \text{Prob}(\mathbf{S}^1 \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}})$ which is ergodic with respect to the skew dynamical system $G(x, y, \bar{c}) = (G_{c_0}(x, y), \sigma(\bar{c}))$, having the property that μ a.e.

$$\frac{1}{N} \sum_{j=0}^{N-1} g(G^j(x, y, \bar{b})) \xrightarrow{N \rightarrow \infty} \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} g(x, y, \bar{b}) d\mu(x, y, \bar{b}),$$

for all $g \in L^1(\mu)$. In the second part, we consider the IFS $R = (\mathbf{S}^1, \tau_a)_{a \in \mathcal{I}}$ where $\mathcal{I} = \{0, 1\}$ and $\tau_i(x) = \frac{1}{2}x + \frac{i}{2}$, $i \in \mathcal{I}$, are the inverse branches of $T(x)$. In this setting we study the IFS ergodic optimization problem for a finite family of potentials, consisting in finding the value $m(R) = \sup_{\mu \in \mathcal{H}} \int A_c(\tau_a x) d\mu(x, c, a)$, over the set of holonomic probabilities \mathcal{H} and also the optimal measures $\mu_\infty \in \mathcal{H}$, such that, $m(R) = \int A_c(\tau_a x) d\mu_\infty(x, c, a)$. As a tool to understand it, we study, for each $\lambda < 1$, the discounted IFS ergodic optimization problem for a finite family of potentials consisting in to find the value $m_\lambda(R) = \sup \int A_c(\tau_a x) d\mu(x, c, a)$, over the set of discounted holonomic probabilities $\mathcal{H}^\lambda(\nu)$ and the optimal measures. Moreover, we show that if $\mu = \mu_{x_0, \bar{c}, \bar{a}}^\lambda$ is an empirical discounted holonomic measure, then $\int A_c(\tau_a x) d\mu(x, c, a) = (1 - \lambda) S_{x_0}(\bar{c}, \bar{a})$. This shows the connection between this maximization problem and the superior boundary of the set Λ which is the graph $\{(x, v_\lambda^+(x)) \mid x \in X\}$ where $v_\lambda^+(x) = \max_{(\bar{c}, \bar{a}) \in \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}}} S_{x_0}(\bar{c}, \bar{a})$. Additionally, when the discounted holonomic measures has trace $\nu = \delta_{x_0}$ supported in a point of maximum of v_λ^+ we show that $m_\lambda(R) \xrightarrow{\lambda \rightarrow 1} m(R)$ and any cluster point of optimal discounted holonomic measures is a optimal holonomic measure.

*Email: elismar.oliveira@ufrgs.br

⁰2010 *Mathematics Subject Classification*: 37-XX, 28Dxx, 37B10, 37Hxx, 37B55, 49Lxx, 49L20, 90C39, 37L40.

⁰*Keywords*: Iterated function system, Ergodic Theory, Ergodic Optimization, Dynamic Programming, Bellman Equation, Discounted Ergodic Averages, SRB measures

1 Introduction

In [BKRLU06] the authors study, from a topological point of view, the dynamics of the skew product $G : X \times \mathbb{R} \rightarrow X \times \mathbb{R}$ given by $G(x, y) = (T(x), A(x) + \lambda y)$, where $0 < \lambda < 1$, $X = \mathbb{R}/\mathbb{Z} = \mathbf{S}^1$, T is the multiplication by some $l \geq 2$ and $A : X \rightarrow \mathbb{R}$ is a Lipschitz potential function. That maps are also called linear Baker maps. The goal of that paper was to prove that there exists a topological attractor and that is homeomorphic to an annulus when the discount parameter λ vanishes, that is, for $\lambda \rightarrow 1$. In [Tsu01] this map was studied from a measure theoretical point of view giving a description of the SRB measure. This kind of skew products and its connection with iterated function systems (IFS) theory received a lot of attention in the last few years (see [DaGR17] for skew products involving diffeomorphisms in manifolds, [Ram03] for non linear baker maps and [Tsu01], [BKRLU06], [LO17] for the linear ones).

Analyzing the results in [BKRLU06] we address three main questions:

- a) What happens when we increase the number of skew maps, that is, if we consider a skew product IFS on the cylinder $X \times \mathbb{R}$ formed by maps $G_j(x, y) = (T(x), A_j(x) + \lambda y)$ for a finite family of $m \geq 2$, potential functions $\{A_j\}_{0 \leq j \leq m-1}$? We still have an attractor? How to characterize it?
- b) The second question is when the discount parameter λ vanishes, that is, for λ close enough to 1. Can we recover some information via ergodic optimization for a finite family of potential functions $\{A_j\}_{0 \leq j \leq m-1}$?
- c) The third question is on the existence and properties of the SRB measure for such systems.

In [LO17], question (b) was studied and a partial answer to the conjecture on the structure of the boundary of the attractor was given. The authors proves that the upper boundary of the attractor is a piecewise differentiable graph $(x, v_\lambda(x))$ under some hypothesis. Other result was to describe the relation between the upper boundary of the attractor and the maximizing measures for the potential. This led us to believe that we can extend this ideas for the IFS problem. In the next sections we explain the motivation and our strategy.

We point out that our choice of a simple setting is just to make easy to understand de ideas. However, several generalizations are trivially derived from our arguments. In the skew IFS problem we can replace \mathbf{S}^1 by the a n -torus \mathbb{T}^n (or other suitable compact metric space) and consider the skew IFS in $\mathbb{T}^n \times \mathbb{R}$, or to keep \mathbf{S}^1 and consider the map T as the multiplication by $2 \leq l \in \mathbb{N}$ or even a expanding automorphism of degree $l \geq 2$. In this case, we just change the Lebesgue measure ℓ by some measure ergodic for T , obtaining the same results for random SRB measures.

For the second part, we can easily replace the IFS $\tau_i(x) = \frac{1}{2}x + \frac{i}{2}$, $i \in \mathcal{I} = \{0, 1\}$, by the inverse branches of an expanding automorphism $T(x)$ or even a continuous IFS $R = (X, \tau_a)_{a \in \mathcal{I}}$ for a compact set \mathcal{I} of maps on a compact metric space X . The results regarding to IFS ergodic optimization still true but we lose the connection with the invariant set of the skew IFS because the IFS maps are not inverse branches of the map T that defines the skew IFS.

The structure of the paper is the following:
In the second section we consider the skew IFS problem showing the existence of a fractal compact

invariant set Λ and characterize its boundary using dynamic programming. We also study the random SRB measure associated to this system. In the third section we analyze the IFS ergodic optimization problem for a finite family of potentials. Initially, for ergodic averages, linking this problem with the superior boundary of Λ , and after for discounted ergodic averages. Additionally, we observe that we can approximate solutions of the IFS ergodic optimization problem when the discount vanishes. In the fourth section we present, a duality result of discounted holonomic measures for IFS to give some additional insight on the problems of Section 3.

2 The skew product IFS

Consider a fixed number $0 < \lambda < 1$. We set the topology in $X \times \mathbb{R}$ by considering it as a complete metric space with the distance induced by the quotient $\mathbb{R}/\mathbb{Z} \times \mathbb{R}$ over \mathbb{R}^2 . Therefore $X \times \mathbb{R}$ is a complete metric space.

Any shift space with N symbols $\{q_0, \dots, q_{N-1}\}^{\mathbb{N}}$ used here will be endowed with the product topology induced by the distance $d(\bar{a}, \bar{b}) = \lambda^n$, where $a_k = b_k$ for $k = 0, \dots, n-1$ and $a_n \neq b_n$. The shift map σ and the concatenation map $*$ are given by

$$\sigma(\bar{b}) = (b_1, b_2, \dots) \text{ and } e * \bar{b} = (e, b_0, b_1, \dots),$$

where $\bar{b} = (b_0, b_1, \dots) \in \{q_0, \dots, q_{N-1}\}^{\mathbb{N}}$ (we will denote \bar{b} the sequence and b_n its elements).

Consider $A_c : X \rightarrow \mathbb{R}$ for $c \in \mathcal{C} = \{0, \dots, m-1\}$ a family of Lipschitz potentials

$$\text{Lip}(A_c) = \sup_{x \neq y} \frac{|A_c(x) - A_c(y)|}{d(x, y)} < \infty,$$

$T(x) = 2x$, the multiplication on $X = \mathbb{R}/\mathbb{Z}$ and the IFS $R = (X \times \mathbb{R}, G_c(x, y))_{c \in \mathcal{C}}$ where,

$$G_c(x, y) = (T(x), A_c(x) + \lambda y).$$

Our purpose here is to investigate dynamical and ergodic properties of the skew IFS R in $X \times \mathbb{R}$.

2.1 The invariant set Λ

We may ask if there exists some invariant set or some attractor for the IFS R and how this set can be characterized. Since each map $G_c(x, y)$ expand by a factor 2 in the x direction and contract by a factor λ in the y direction, it is not possible to employ the classical methods for contractive IFS. In the case $m = 1$, [BKRLU06] shows that there exists a solenoidal attractor and study its topological properties.

The orbit of a point (x', y') by the IFS R , controlled by the sequence $\bar{c} = (c_0, c_1, \dots) \in \mathcal{C}^{\mathbb{N}}$, is the projection on $X \times \mathbb{R}$ of the orbit of the skew dynamical system

$$G(x', y', \bar{c}) = (G_{c_0}(x', y'), \sigma(\bar{c})) = (T(x'), A_{c_0}(x') + \lambda y', (c_1, c_2, \dots)).$$

The n -th iterate $G^n(x', y', \bar{c})$ will be $(G_{c_{n-1}}(\dots(G_{c_0}(x', y'))), \sigma^n(\bar{c}))$ or, more precisely

$$\left(T^n(x'), \sum_{i=0}^{n-1} \lambda^i A_{c_{n-1-i}}(T^{n-1-i}(x')) + \lambda^n y', (c_n, c_{n+1}, \dots) \right).$$

Therefore, the projection (x, y) on $X \times \mathbb{R}$ is $x = T^n(x')$ and $y = \sum_{i=0}^{n-1} \lambda^i A_{c_{n-1-i}}(T^{n-1-i}(x')) + \lambda^n y'$.

Remark 2.1. We denote $\text{Per}_n(T) = \{x \in X \mid T^n(x) = x\}$ the set of periodic points of T with period n . Using the formulas for the n -th iterate $G^n(x', y', \bar{c})$ we can see that each Baker map G_c has periodic points projecting in $\text{Per}_n(T)$, that is,

$$\text{Per}_n(G_c) = \left\{ (x', y') \in X \times \mathbb{R} \mid x' \in \text{Per}_n(T) \text{ and } y' = \frac{1}{1 - \lambda^n} \sum_{i=0}^{n-1} \lambda^i A_c(T^{n-1-i}(x')) \right\}.$$

In particular, $|y'| \leq \frac{1}{1-\lambda} \|A_c\|_\infty$ for any $(x', y') \in \text{Per}_n(G_c)$.

We recall that the Hutchinson-Barnsley (HB) operator is given by $F(U) = \bigcup_{c \in \mathcal{C}} G_c(U)$ for any $U \in 2^{X \times \mathbb{R}}$. Considering $\mathcal{K}^*(X \times \mathbb{R})$ the family of all nonempty compact subsets of $X \times \mathbb{R}$, we endow it with the Hausdorff metric and restrict F to this family.

A compact set Λ is self-similar (invariant or fractal) if $F(\Lambda) = \Lambda$. We said that Λ is an attractor if there exists U_0 a neighborhood of Λ , called the basin of attraction, such that, for any $U \in \mathcal{K}^*(U_0)$ we have $F^n(U) \rightarrow \Lambda$. When $U_0 = X \times \mathbb{R}$ we say that the attractor is a global attractor. From [BV11], Lemma 2, if Λ is an attractor with basin of attraction U_0 , then $\Lambda = \lim_{N \rightarrow \infty} \bigcup_{n \geq N} F^n(U)$ for any $U \in \mathcal{K}^*(U_0)$.

Theorem 2.2. Consider any $T_0 > \frac{1}{1-\lambda} \max_{c \in \mathcal{C}} \|A_c\|_\infty$ and the annulus $U_0 = X \times (-T_0, T_0)$ then

- a) $F(\overline{U_0}) \subset U_0$.
- b) U_0 is globally attracting, that is, for any $M > 0$ we have $F^n(X \times [-M, M]) \subset U_0$ for some $n \in \mathbb{N}$ depending only on M .
- c) If $\Lambda \in \mathcal{K}^*(X \times \mathbb{R})$ is self similar, then $\Lambda \subset U_0$.

Proof. (a) We know that $G_{c_0}(x', y') = (T(x'), A_{c_0}(x') + \lambda y') = (x, y)$ so

$$\begin{aligned} |y| &\leq |A_{c_0}(x')| + \lambda |y'| \leq \max_{c \in \mathcal{C}} \|A_c\|_\infty + \lambda |y'| = \\ &= (1 - \lambda) \frac{1}{1 - \lambda} \max_{c \in \mathcal{C}} \|A_c\|_\infty + \lambda |y'| < (1 - \lambda) T_0 + \lambda |y'| \leq (1 - \lambda) T_0 + \lambda T_0 = T_0. \end{aligned}$$

If $(x', y') \in \overline{U_0}$ then $G_{c_0}(x', y') \in U_0$ for any $c_0 \in \mathcal{C}$, thus $F(\overline{U_0}) \subset U_0$, because it is a finite union.

(b) For a fixed $(x', y') \in X \times \mathbb{R}$ and for any $\bar{c} = (c_0, c_1, \dots) \in \mathcal{C}^{\mathbb{N}}$ the projection on $X \times \mathbb{R}$ of $G^n(x', y', \bar{c})$ will be in U_0 for n big enough. We need to prove that

$$\left| \sum_{i=0}^{n-1} \lambda^i A_{c_{n-1-i}}(T^{n-1-i}(x')) + \lambda^n y' \right| < T_0,$$

when n is big enough uniformly in \bar{c} . Indeed,

$$\left| \sum_{i=0}^{n-1} \lambda^i A_{c_{n-1-i}}(T^{n-1-i}(x')) + \lambda^n y' \right| \leq \frac{1 - \lambda^n}{1 - \lambda} \max_{c \in \mathcal{C}} \|A_c\|_\infty + \lambda^n |y'|$$

$$\leq \frac{1}{1-\lambda} \max_{c \in \mathcal{C}} \|A_c\|_\infty + \lambda^n \left(|y'| - \frac{1}{1-\lambda} \max_{c \in \mathcal{C}} \|A_c\|_\infty \right) < T_0,$$

because $\lambda^n \left(|y'| - \frac{1}{1-\lambda} \max_{c \in \mathcal{C}} \|A_c\|_\infty \right) \xrightarrow{n \rightarrow \infty} 0$, $T_0 > \frac{1}{1-\lambda} \max_{c \in \mathcal{C}} \|A_c\|_\infty$ and this limit is independent of \bar{c} , depending only on $|y'|$. From this, we can conclude that for any $M > 0$ we have $F^n(X \times [-M, M]) \subset U_0$, for some $n \in \mathbb{N}$, depending only on M .

(c) From (b), taking n large enough, we get that Λ should be a subset of the “attracting basin” U_0 because $\Lambda = F^n(\Lambda)$. \blacksquare

To characterize the points of Λ , we need to evaluate the images of arbitrarily large order of a point $(x', y') \in X \times \mathbb{R}$ because $F^n(\Lambda) = \Lambda$, for all $n \in \mathbb{N}$, when Λ is self similar. We can write $F^n(\Lambda)$ as

$$\{(x, y) = G_{c_{n-1}}(\cdots(G_{c_0}(x', y')) \mid c_0, \dots, c_{n-1} \in \mathcal{C}, (x', y') \in \Lambda) = \left\{ \left(T^n(x'), \sum_{i=0}^{n-1} \lambda^i A_{c_{n-1-i}}(T^{n-1-i}(x')) + \lambda^n y' \right) \mid c_0, \dots, c_{n-1} \in \mathcal{C}, (x', y') \in \Lambda \right\}.$$

Since X is compact it is natural to assume, by analogy with [BKRLU06], that the y coordinate is bounded in every infinite pre-orbit of Λ , so we define

$$\Lambda = \{(x, y) \in X \times \mathbb{R} \mid \forall \bar{c} \in \mathcal{C}^{\mathbb{N}}, \exists M > 0, |y_n| \leq M, \text{ when } G^n(x_n, y_n, \bar{c}) = (x, y), \forall n \in \mathbb{N}\}.$$

Consider $\mathcal{I} = \{0, 1\}$, $\bar{a} = (a_0, a_1, \dots) \in \mathcal{I}^{\mathbb{N}}$ and $\tau_i(x) = \frac{1}{2}x + \frac{i}{2}$, $i \in \mathcal{I}$, the inverse branches of $T(x)$. We also define $\tau_{i, \bar{a}}(x) = \tau_{a_{i-1}} \circ \tau_{a_{i-2}} \cdots \circ \tau_{a_0}(x)$.

Inspired by [Tsu01] equation (2), we introduce the function $S : X \times \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}} \rightarrow \mathbb{R}$ by $S_x(\bar{c}, \bar{a}) = \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x))$ to study the iteration of points in Λ .

Proposition 2.3. *The set Λ can be written as*

$$\Lambda = \{(x, S_x(\bar{c}, \bar{a})) \in X \times \mathbb{R} \mid \forall \bar{c} \in \mathcal{C}^{\mathbb{N}}, \forall \bar{a} \in \mathcal{I}^{\mathbb{N}}\}.$$

Proof. Let us to compute some pre-images of a point (x, y) :

If $G_{c_0}(x_1, y_1) = (x, y)$, then $T(x_1) = x$ and $A_{c_0}(x_1) + \lambda y_1 = y$. Thus, $\exists a_0 \in \mathcal{I}$, such that, $x_1 = \tau_{a_0}(x)$ and $y = A_{c_0}(\tau_{a_0}(x)) + \lambda y_1$. Therefore $y_1 = \frac{1}{\lambda}(y - A_{c_0}(\tau_{a_0}(x)))$.

Following this procedure we obtain that if $G_{c_1}(x_2, y_2) = (x_1, y_1)$, then $x_2 = \tau_{a_1} \tau_{a_0}(x)$ and $y_1 = A_{c_1}(x_2) + \lambda y_2$. So $y_2 = \frac{1}{\lambda^2}(y - (A_{c_0}(\tau_{a_0}(x)) - \lambda A_{c_1}(\tau_{a_1} \tau_{a_0}(x))))$. By proceeding a formal induction we obtain $x_n = \tau_{a_{n-1}} \circ \cdots \circ \tau_{a_0}(x)$ and $y_n = \frac{1}{\lambda^n} \left(y - \sum_{i=0}^{n-1} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x)) \right)$.

From this computations we conclude that

$$\Lambda = \left\{ (x, y) \in X \times \mathbb{R} \mid \forall \bar{c} \in \mathcal{C}^{\mathbb{N}}, \exists M > 0, \exists \bar{a} \in \mathcal{I}^{\mathbb{N}}, \left| y - \sum_{i=0}^{n-1} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x)) \right| \leq M \lambda^n, \forall n \in \mathbb{N} \right\}.$$

Taking $n \rightarrow \infty$, we obtain that $\Lambda = \{(x, \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x))) \in X \times \mathbb{R} \mid \forall \bar{c} \in \mathcal{C}^{\mathbb{N}}, \forall \bar{a} \in \mathcal{I}^{\mathbb{N}}\}$. \blacksquare

In the next proposition we show that Λ is self similar.

Proposition 2.4. *The set Λ is self similar, that is, $F(\Lambda) = \Lambda$.*

Proof. Indeed, to show that $F(\Lambda) \subseteq \Lambda$ we take $(x', y') \in \Lambda$ and $\bar{c} \in \mathcal{C}^{\mathbb{N}}$, $M > 0$, $\bar{a} \in \mathcal{I}^{\mathbb{N}}$, such that,

$$\left| y' - \sum_{i=0}^{n-1} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x')) \right| \leq M \lambda^n, \text{ for all } n \in \mathbb{N}. \text{ We need to show that } G_b(x', y') \in \Lambda, \text{ for any } b \in \mathcal{C}.$$

If $G_b(x', y') = (x, y)$, then $T(x') = x$ and $A_b(x') + \lambda y' = y$. Thus, $\exists e \in \mathcal{I}$, such that, $x' = \tau_e(x)$ and $y = A_b(\tau_e(x)) + \lambda y'$, therefore $y' = \frac{1}{\lambda} (y - A_b(\tau_e(x)))$.

Substituting x' and y' in $\left| y' - \sum_{i=0}^{n-1} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x')) \right| \leq M \lambda^n$, we obtain

$$\left| \frac{1}{\lambda} (y - A_b(\tau_e(x))) - \sum_{i=0}^{n-1} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(\tau_e(x))) \right| \leq M \lambda^n,$$

which is equivalent to $\left| y - \sum_{i=0}^n \lambda^i A_{b_i}(\tau_{i, \bar{e}}(x)) \right| \leq M \lambda^{n+1}$, where $\bar{b} = b * \bar{c} = (b, c_0, c_1, \dots)$ and $\bar{e} = e * \bar{a} = (e, a_0, a_1, \dots)$. Thus, $(x, y) \in \Lambda$.

To prove the opposite inequality, $\Lambda \subseteq F(\Lambda)$, we consider $\left(x, \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x)) \right) \in \Lambda$ and $x' = \tau_{a_0}(x)$. Then, $x = T(x')$, $x' = \tau_{a_0}(T(x'))$ and

$$\left(x, \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x)) \right) = G_{c_0} \left(x', \sum_{i=0}^{\infty} \lambda^i A_{c_{i+1}}(\tau_{i, \sigma \bar{a}}(x')) \right).$$

This means that $(x, y) \in F(\Lambda)$ because $\left(x', \sum_{i=0}^{\infty} \lambda^i A_{c_{i+1}}(\tau_{i, \sigma \bar{a}}(x')) \right) \in \Lambda$. ■

Lemma 2.5. *The function $S : X \times \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}} \rightarrow \mathbb{R}$ given by $S_x(\bar{c}, \bar{a}) = \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x))$ is Lipschitz continuous, more precisely*

$$|S_x(\bar{c}, \bar{a}) - S_{x'}(\bar{b}, \bar{e})| \leq \frac{2}{2-\lambda} \max_{c \in \mathcal{C}} \text{Lip}(A_c) d(x, x') + \frac{2}{1-\lambda} \max_{c \in \mathcal{C}} \|A_c\|_{\infty} (d(\bar{c}, \bar{b}) + d(\bar{a}, \bar{e})),$$

where each distance is taken in the respective space.

Proposition 2.6. *Λ is closed, and in particular it is a compact set.*

Proof. To see that $\bar{\Lambda} = \Lambda$ we take $(x^k, S_{x^k}(\bar{c}^k, \bar{a}^k)) \rightarrow (x, y)$, when $k \rightarrow \infty$. We notice that $\mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}}$ is a compact space thus we can assume that $(\bar{c}^k, \bar{a}^k) \rightarrow (\bar{b}, \bar{e})$, possibly by taking a subsequence. We claim that $y = S_x(\bar{b}, \bar{e})$. Indeed, from Lemma 2.5 we know that $S_x(\bar{b}, \bar{e}) = \lim_{k \rightarrow \infty} S_{x^k}(\bar{c}^k, \bar{a}^k) = y$. Thus, $(x, y) \in \Lambda$. The compactness follow from the fact that Λ is bounded. ■

2.2 Dynamic programming and the boundary of Λ

The notation and the main results in dynamic programming presented here are from [JMN14] and the results on discounted limits are from [CO17]. We consider a decision-making process $S = \{X, A, \psi, f, u, \delta\}$ given by:

- a) the state space $(X = \mathbf{S}^1, d)$ is a compact metric space;
- b) the action space $(\mathcal{C} \times \mathcal{I}, d_{\mathcal{C} \times \mathcal{I}})$ is a compact metric space where \mathcal{C} (describe the maps on the IFS) and \mathcal{I} (describe the injective domains of T) are both finite sets;
- c) the action function $\psi(x) = \mathcal{C} \times \mathcal{I}, \forall x \in X$;
- d) the dynamics is given by the contractive IFS $f(x, c, a) = \tau_a(x)$ for $a \in \mathcal{I}$ (the dynamics does not depends on c);
- e) the immediate return is $u(x, c, a) = A_c(\tau_a(x))$, $c \in \mathcal{C}$, $a \in \mathcal{I}$; where A_c is Lipschitz with respect to x , uniformly in c ;
- f) the discount function is linear, $\delta(t) = \lambda t$ for $0 < \lambda < 1$.

Assuming such hypothesis we can show that for each fixed $0 < \lambda < 1$ there exists an unique Lipschitz continuous function v_λ^+ which satisfies the Bellman equation

$$v_\lambda^+(x) = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) + \lambda v_\lambda^+(\tau_a(x)),$$

where $v_\lambda^+(x) = \max_{\bar{c} \in \mathcal{C}^{\mathbb{N}}, \bar{a} \in \mathcal{I}^{\mathbb{N}}} \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}} x)$ is attained for some optimal pair (\bar{c}, \bar{a}) . The same is true

for $v_\lambda^-(x) = \min_{\bar{c} \in \mathcal{C}^{\mathbb{N}}, \bar{a} \in \mathcal{I}^{\mathbb{N}}} \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}} x)$ but we are interested in maximization thus, we will use just

the notation v_λ^+ and denote v_λ if there is no risk of misunderstanding.

One can show that $(1 - \lambda) \max_x v_\lambda \rightarrow \bar{u}$ and $v_\lambda(x) - \max_x v_\lambda$ converges up to subsequence, to a continuous function b , such that, $b(x) = \max_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) - \bar{u} + b(\tau_a(x))$ that can be rewritten as the calibrated sub-action equation in ergodic optimization (see [GL08] and [GLT09]) or the analogous of Hamilton-Jacobi's equation for a discrete Lagrangian (see [Gom05])

$$\bar{u} = \max_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) + b(\tau_a(x)) - b(x) = \max_{c, a \in \mathcal{C} \times \mathcal{I}} d_x b(a) + A_c(\tau_a(x)),$$

where $d_x b(a) = b(f(x, c, a)) - b(x) = b(\tau_a x) - b(x)$ is the discrete differential. \bar{u} is uniquely determined by

$$\bar{u} = \sup_{\mu \in \mathcal{H}} \int A_c(\tau_a(x)) d\mu(x, c, a),$$

where the set \mathcal{H} of holonomic probabilities is

$$\mathcal{H} = \left\{ \mu \in \text{Prob}(X \times \mathcal{C} \times \mathcal{I}) \mid \int d_x g(a) d\mu(x, c, a) = 0, \forall g \in C^0(X, \mathbb{R}) \right\}.$$

Remark 2.7. We point out that, in the case where the potential A is not changing at each iteration, the set \mathcal{C} is a single point then, $A_c(x) = A(x)$ and $\bar{u} = \sup_{\mu \in \mathcal{H}} \int A_c(\tau_a(x)) d\mu(x, c, a) = \sup_{\mu \in \mathcal{H}} \int d_x A(a) + A(x) d\mu(x, c, a) = \sup_{\mu \in \mathcal{H}} \int A(x) d\mu(x, c, a)$, because $A_c(\tau_a(x)) = A(\tau_a(x)) = A(\tau_a(x)) - A(x) + A(x) = d_x A(a) + A(x)$ and $\int d_x A(a) d\mu(x, c, a) = 0$. Thus, we recover exactly the classical setting of ergodic optimization for an IFS as in [LO17].

The next proposition characterizes the boundary of Λ .

Proposition 2.8. For each $(x, y) \in \Lambda$ we have $v_\lambda^-(x) \leq y \leq v_\lambda^+(x)$. Moreover, the graphs $\{(x, v_\lambda^-(x)) \mid x \in X\}$ and $\{(x, v_\lambda^+(x)) \mid x \in X\}$ are subsets of Λ .

Proof. We determinate the boundaries of Λ using dynamic programming. Evaluating the above equations we can pick at each step an infinite backwards orbit proving that $(x, v_\lambda^+(x)) \in \Lambda$. Thus, the graphs of v_λ^+ and v_λ^- are subsets of Λ . As a consequence, $\Lambda \subseteq \{(x, y) \mid v_\lambda^-(x) \leq y \leq v_\lambda^+(x)\}$, in particular Λ will be a Jordan curve only if $v_\lambda^-(x) = v_\lambda^+(x)$ for all x . ■

2.3 Approximation of Λ

As a motivation, we consider a numerical example to provide some insight on what happens when we draw an orbit of G .

Example 2.9. Consider $m = 2$, $\lambda = 0.48$ and the potentials $A_0, A_1 : X \rightarrow \mathbb{R}$ given by $A_0(x) = (x - \frac{1}{2})^2$ and $A_1(x) = 2x$, $0 \leq x \leq \frac{1}{2}$, $A_1(x) = -2x + 2$, $\frac{1}{2} \leq x \leq 1$. Let $R = (X \times \mathbb{R}, G_j(x, y))$ be the IFS given by $G_j(x, y) = (T(x), A_j(x) + \lambda y)$. In this example we choose an initial point $(x_0, y_0) = (0.2472135954, 0.1)$ and draw its orbit using a chosen sequence $\bar{c} = (c_0, c_1, \dots)$.

We consider three different situations:

- When $\bar{c} = (0, 0, 0, \dots)$, then iterate only $G_0(x, y) = (T(x), A_0(x) + 0.48y)$ beginning in (x_0, y_0) .
- When we choose, in a random way, a sequence $\bar{c} = (c_0, c_1, \dots) \in \{0, 1\}^N$ and iterate $G_{c_i}(x, y) = (T(x), A_{c_i}(x) + 0.48y)$ beginning in (x_0, y_0) .
- When $\bar{c} = (1, 1, 1, \dots)$ and iterate only $G_1(x, y) = (T(x), A_1(x) + 0.48y)$ beginning in (x_0, y_0) .

The associated pictures are showed in the Figure 1.

We can see in (b) that the a finite family of potential system(red) presents some mixing of the two autonomous (a) and (c) (blue and black) studied by [BKRLU06]. This example makes us to conjecture that closure of a typically orbit of the IFS drawn the picture of the invariant set Λ .

Finally, in the Figure 2 we draw a picture, describing the iteration using a random orbit with 10.000 iterations after the iterate 1.000. We also show the approximations of $v_{0.48}^+(x)$ (green) and $v_{0.48}^-(x)$ (yellow) obtained by iteration of the contractive operator

$$\mathcal{L}^+(f)(x) = \max_{c \in \{0,1\}, a \in \{0,1\}} A_c(\tau_a x) + 0.48f(\tau_a x),$$

(resp. \mathcal{L}^-) because $\mathcal{L}^+(v_{0.48}^+) = v_{0.48}^+$ (resp. $\mathcal{L}^-(v_{0.48}^-) = v_{0.48}^-$).




Figure 1: Approximating the attractor: a, b and c, from the left to the right




Figure 2: Approx. of Λ and the upper and lower boundary $v_{0.48}^+(x)$ and $v_{0.48}^-(x)$.

In the same setting of the Example 2.9, we consider a periodic point $x_0 = \frac{1}{3}$ and $(x_0, y_0) = (0.33\dots, 1.4)$. In the Figure 3, we choose a sequence $\bar{c} = (c_0, c_1, \dots) \in \{0, 1\}^{\mathbb{N}}$, iterate $G_{c_i}(x, y) = (T(x), A_{c_i}(x) + 0.48y)$ by 2000 times beginning in (x_0, y_0) and we plot it for $i \geq 500$.




Figure 3: Iteration of the compact $\{(0.33\dots, 1.4)\}$ by F .

Proposition 2.10. Λ is not an attractor with any basin of attraction U_0 , in the IFS sense.

Proof. Consider an arbitrary point $\{(1/3, y')\}$ and calculate $F^n(\{(1/3, y')\})$. A simple computation shows that

$$G_{b_{-1}}(1/3, y') = (T(1/3), A_{b_{-1}}(1/3) + \lambda y') = (2/3, A_{b_{-1}}(1/3) + \lambda y')$$

and

$$\begin{aligned} G_{b_{-2}}G_{b_{-1}}(1/3, y') &= (T(2/3), A_{b_{-2}}(2/3) + \lambda A_{b_{-1}}(1/3) + \lambda^2 y') = \\ &= (1/3, A_{b_{-2}}(2/3) + \lambda A_{b_{-1}}(1/3) + \lambda^2 y'). \end{aligned}$$

The same behavior occurs for $G_{b_{-n}} \cdots G_{b_{-1}}(1/3, y')$ depending only if n is even or odd.

We can see that Λ is not an attractor with basin of attraction U_0 , in the IFS sense, because

$$\pi_x \left(\lim_{n \rightarrow \infty} F^n(\{(1/3, y')\}) \right) = \left\{ \frac{1}{3}, \frac{2}{3} \right\},$$

thus it is impossible to obtain $\lim_{n \rightarrow \infty} F^n(\{(1/3, y')\}) = \Lambda$. This contradicts the fact that $\{(1/3, y')\}$ is a compact set in the basin of attraction U_0 . \blacksquare

2.4 The dynamic in Λ and the random SRB measure

As Λ is a compact subset and $F(\Lambda) = \Lambda$ we can restrict the IFS $R = (X \times \mathbb{R}, G_c(x, y))_{c \in \mathcal{C}}$ to Λ , that is, $R' = (\Lambda, G_c(x, y))_{c \in \mathcal{C}}$.

We recall that $\Lambda = \{(x, S_x(\bar{c}, \bar{a})) \in X \times \mathbb{R} \mid \forall \bar{c} \in \mathcal{C}^{\mathbb{N}}, \forall \bar{a} \in \mathcal{I}^{\mathbb{N}}\}$, where the function $S : X \times \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}} \rightarrow \mathbb{R}$ given by $S_x(\bar{c}, \bar{a}) = \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{i, \bar{a}}(x))$ is Lipschitz continuous by Lemma 2.5.

We introduce the address function $\pi : X \rightarrow \mathcal{I}$ given by

$$\pi(x) = e \in \mathcal{I} \Leftrightarrow \tau_e T(x) = x,$$

which is well defined for T because, in each point, there exists a left inverse function.

To study the iteration of points in Λ we compute, for each $^1 \bar{b} = (b_{-1}, b_{-2}, \dots) \in \mathcal{C}^{\mathbb{N}}$ the image $G(x, S_x(\bar{c}, \bar{a}), \bar{b}) = (T(x), S_{T(x)}(b_{-1} * \bar{c}, e_{-1} * \bar{a}), \sigma \bar{b})$ where $e_{-1} = \pi(x) \in \mathcal{I}$ (the unique element in \mathcal{I} , such that, $\tau_{e_{-1}} T(x) = x$), $b_{-1} * \bar{c} = (b_{-1}, c_0, c_1, \dots)$ and $e_{-1} * \bar{a} = (e_{-1}, a_0, a_1, \dots)$. For an arbitrary n we get $G^n(x, S_x(\bar{c}, \bar{a}), \bar{b})$ as

$$(T^n(x), S_{T^n(x)}(b_{-n} \cdots b_{-1} * \bar{c}, e_{-n} \cdots e_{-1} * \bar{a}), \sigma^n \bar{b})$$

where $e_{-i} = \pi(T^{i-1}(x))$, $i = 1, \dots, n$.

This shows that G has a simple behavior in Λ . Therefore is natural to consider a conjugation between G and the symbolic dynamics. To do that, we define the map $\theta_b : X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \rightarrow X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}}$ by $\theta_b(x, \bar{a}, \bar{c}) = (T(x), \pi(x) * \bar{a}, b * \bar{c})$, for any $b \in \mathcal{C}$. We also define the map

$$\theta(x, \bar{a}, \bar{c}, \bar{b}) = (\theta_b(x, \bar{c}, \bar{a}), \sigma(\bar{b})) = (T(x), \pi(x) * \bar{a}, b_{-1} * \bar{c}, \sigma(\bar{b}))$$

and the map $\Psi : X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \rightarrow X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}$ by $\Psi(x, \bar{a}, \bar{c}, \bar{b}) = (x, S_x(\bar{c}, \bar{a}), \bar{b})$.

Lemma 2.11. *Consider the maps G, Ψ, θ above defined. Then, the following properties are true:*

- a) $S_{T(x)}(b * \bar{c}, \pi(x) * \bar{a}) = A_b(x) + \lambda S_x(\bar{c}, \bar{a})$, for any $b \in \mathcal{C}$.

¹We use a little bit different indexation for b starting from $\bar{b} = (b_{-1}, b_{-2}, \dots)$ instead $\bar{b} = (b_0, b_1, \dots)$. This is made in order to avoid confusion on the concatenation operation $b_{-1} * \bar{c} = (b_{-1}, c_0, c_1, \dots)$.

b) $G \circ \Psi = \Psi \circ \theta$.

Proof. (a) We already proved that.

(b) To see that, we compute both sides of the equation

$$\begin{aligned} G \circ \Psi(x, \bar{a}, \bar{c}, \bar{b}) &= G(x, S_x(\bar{c}, \bar{a}), \bar{b}) = (T(x), S_{T(x)}(b_{-1} * \bar{c}, \pi(x) * \bar{a}), \sigma \bar{b}) = \\ &= \Psi(T(x), \pi(x) * \bar{a}, b_{-1} * \bar{c}, \sigma \bar{b}) = \Psi \circ \theta(x, \bar{a}, \bar{c}, \bar{b}). \end{aligned}$$

■

Theorem 2.12. Consider G, Ψ, θ as in Lemma 2.11. Recalling that $\mathcal{C} = \{0, \dots, m-1\}$ and $\mathcal{I} = \{0, 1\}$, let ν be the uniform Bernoulli measure on $\mathcal{C}^{\mathbb{N}}$ given by $\nu(j) = \frac{1}{m}$, η be the uniform Bernoulli measure on $\mathcal{I}^{\mathbb{N}}$ given by $\eta(i) = \frac{1}{2}$ and ℓ be the normalized Lebesgue measure on X . The probability measure μ in $X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}$ given by

$$\int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} g(x, y, \bar{b}) d\mu(x, y, \bar{b}) = \int_{X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}}} g(\Psi(x, \bar{a}, \bar{c}, \bar{b})) d\ell(x) d\eta(\bar{a}) d\nu(\bar{c}) d\nu(\bar{b})$$

that is, $\mu = \Psi(\ell \times \eta \times \nu^2)$ is ergodic with respect to G . In particular, there exists a set $\Omega \subseteq X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}$ with $\mu(\Omega) = 1$, such that, for all $(x, y, \bar{b}) \in \Omega$,

$$\frac{1}{N} \sum_{j=0}^{N-1} g(G^j(x, y, \bar{b})) \rightarrow \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} g(x, y, \bar{b}) d\mu(x, y, \bar{b}),$$

for all $g \in L^1(\mu)$. This measure is called the **random SRB** measure for the IFS $R = (X \times \mathbb{R}, G_c(x, y))_{c \in \mathcal{C}}$.

Proof. The proof is a consequence of the Birkhoff theorem if we write our map in a clever way. Consider the map $\theta'(x, \bar{a}) = (T(x), \pi(x) * \bar{a})$ introduced by [Tsu01]. He proves that $\ell \times \eta$ is ergodic for θ' . We also define $\theta''(\bar{c}, \bar{b}) = (b_{-1} * \bar{c}, \sigma(\bar{b})) = ((b_{-1}, c_0, c_1, \dots), (b_{-2}, b_{-3}, \dots))$ that is the two sided-shift in $\mathcal{C}^{\mathbb{Z}}$. It is a known fact that ν^2 is the uniform Bernoulli measure which is ergodic with respect to the bilateral shift in \mathbb{Z} . As $\theta = \theta' \times \theta''$ is a factor map, the product $\ell \times \eta \times \nu^2$ is ergodic for θ . From Lemma 2.11 we can transfer the ergodicity and the invariance of θ with respect to $\ell \times \eta \times \nu^2$ to G with respect to μ . Indeed,

$$\begin{aligned} \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} g \circ G(x, y, \bar{b}) d\mu(x, y, \bar{b}) &= \int_{X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}}} g \circ G(\Psi(x, \bar{a}, \bar{c}, \bar{b})) d\ell(x) d\eta(\bar{a}) d\nu(\bar{c}) d\nu(\bar{b}) = \\ &= \int_{X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}}} g \circ \Psi(\theta(x, \bar{c}, \bar{a}, \bar{b})) d\ell(x) d\eta(\bar{a}) d\nu(\bar{c}) d\nu(\bar{b}) = \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} g(x, y, \bar{b}) d\mu(x, y, \bar{b}). \end{aligned}$$

Thus, we can apply the ergodic Birkhoff theorem and the result follows. ■

Remark 2.13. We notice that the measure μ given by Theorem 2.12, and consequently the set Ω depends on λ because $\mu = \Psi(\ell \times \eta \times \nu^2)$ and $\Psi(x, \bar{a}, \bar{c}, \bar{b}) = (x, S_x(\bar{c}, \bar{a}), \bar{b})$. When $\lambda \rightarrow 1$ the discounted sum $S_x(\bar{c}, \bar{a})$ does not converges.

Corollary 2.14. *Let μ be the random SRB measure given by Theorem 2.12. The following statements are true:*

a)
$$\int_X v_{\lambda}^{-}(x) d\ell(x) \leq \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} y d\mu(x, y, \bar{b}) \leq \int_X v_{\lambda}^{+}(x) d\ell(x).$$

b)
$$\Pi_2(x, y, \bar{b}) = y \in L^1(\mu).$$

c) *There exists a set $\Omega \subseteq X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}$ ($\mu(\Omega) = 1$), such that, for all $(x, \bar{b} = (b_{-1}, b_{-2}, \dots)) \in \Pi_{\{1,3\}}\Omega$*

$$\frac{1}{N} \sum_{j=1}^{N-1} A_{b_{-j}}(T^{j-1}(x)) \rightarrow \frac{(1-\lambda)}{\lambda} \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} y d\mu(x, y, \bar{b}).$$

d) *The limit in (c) can be improved by $\forall \varepsilon > 0, \exists \lambda_0, \forall \lambda > \lambda_0$ exists a set $\Omega_{\lambda} \subseteq X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}$ ($\mu(\Omega_{\lambda}) = 1$), such that, for all $(x, \bar{b} = (b_{-1}, b_{-2}, \dots)) \in \Pi_{\{1,3\}}\Omega_{\lambda}, \exists N_0$, such that,*

$$\frac{1}{N} \sum_{j=1}^{N-1} A_{b_{-j}}(T^{j-1}(x)) \leq \bar{u} + \varepsilon,$$

for all $N \geq N_0$, where $\bar{u} = \sup_{\mu \in \mathcal{H}} \int A_c(\tau_a(x)) d\mu(x, c, a)$.

Proof.

(a)-(b) We claim that $g \in L^1(\mu)$ where $g(x, y, \bar{b}) = \Pi_2(x, y, \bar{b}) = y$. To see this we recall that

$$\begin{aligned} \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} \Pi_2(x, y, \bar{b}) d\mu(x, y, \bar{b}) &= \int_{X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}}} \Pi_2(\Psi(x, \bar{a}, \bar{c}, \bar{b})) d\ell(x) d\eta(\bar{a}) d\nu(\bar{c}) d\nu(\bar{b}) = \\ &= \int_{X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}}} S_x(\bar{c}, \bar{a}) d\ell(x) d\eta(\bar{a}) d\nu(\bar{c}) d\nu(\bar{b}) \in \left[\int_X v_{\lambda}^{-}(x) d\ell(x), \int_X v_{\lambda}^{+}(x) d\ell(x) \right], \end{aligned}$$

because $v_{\lambda}^{-}(x) \leq S_x(\bar{c}, \bar{a}) \leq v_{\lambda}^{+}(x)$ from Proposition 2.8.

(c) Using Theorem 2.12 we get

$$\frac{1}{N} \sum_{n=0}^{N-1} \Pi_2(G^n(x, y, \bar{b})) \rightarrow \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} \Pi_2(x, y, \bar{b}) d\mu(x, y, \bar{b}),$$

where

$$\Pi_2 \circ G^n(x, y, \bar{b}) = \Pi_2(T^n(x), y^n, \sigma^n \bar{b}) = y^n.$$

We notice that $y^0 = y$ and for $n \geq 1$ we have the formula

$$y^n = \lambda^n y + \sum_{j=1}^n \lambda^{n-j} A_{b_{-j}}(T^{j-1}(x)),$$

so

$$\begin{aligned} \frac{1}{N} \sum_{n=0}^{N-1} \Pi_2(G^n(x, y, \bar{b})) &= \frac{y}{N} + \frac{1}{N} \sum_{n=1}^{N-1} y^n = \frac{y}{N} + \frac{1}{N} \sum_{n=1}^{N-1} \left[\lambda^n y + \sum_{j=1}^n \lambda^{n-j} A_{b_{-j}}(T^{j-1}(x)) \right] = \\ &= y \frac{1}{N} \sum_{n=0}^{N-1} \lambda^n + \frac{1}{N} \sum_{n=1}^{N-1} \left[\sum_{j=1}^n \lambda^{n-j} A_{b_{-j}}(T^{j-1}(x)) \right]. \end{aligned}$$

As $y \frac{1}{N} \sum_{n=0}^{N-1} \lambda^n \rightarrow 0$ for any y , we obtain

$$\frac{1}{N} \sum_{n=1}^{N-1} \left[\sum_{j=1}^n \lambda^{n-j} A_{b_{-j}}(T^{j-1}(x)) \right] \rightarrow \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} y d\mu(x, y, \bar{b}).$$

Rewriting the above sum we obtain

$$\begin{aligned} \frac{1}{N} \sum_{n=1}^{N-1} \left[\sum_{j=1}^n \lambda^{n-j} A_{b_{-j}}(T^{j-1}(x)) \right] &= \frac{1}{N} \sum_{j=1}^{N-1} \left[\sum_{i=1}^{N-1-j} \lambda^i \right] A_{b_{-j}}(T^{j-1}(x)) = \\ &= \frac{1}{N} \sum_{j=1}^{N-1} \left[\frac{\lambda^{N-j} - \lambda}{\lambda - 1} \right] A_{b_{-j}}(T^{j-1}(x)) = \\ &= \frac{1}{N(\lambda - 1)} \sum_{j=1}^{N-1} \lambda^{N-j} A_{b_{-j}}(T^{j-1}(x)) - \frac{\lambda}{N(\lambda - 1)} \sum_{j=1}^{N-1} A_{b_{-j}}(T^{j-1}(x)). \end{aligned}$$

We notice that $|A_{b_{-j}}(T^{j-1}(x))| \leq \|A_c\|_{\infty}$ so

$$\lim_{N \rightarrow \infty} \frac{1}{N(\lambda - 1)} \sum_{j=1}^{N-1} \lambda^{N-j} A_{b_{-j}}(T^{j-1}(x)) = 0.$$

Using this, we conclude that

$$\frac{1}{N} \sum_{j=1}^{N-1} A_{b_{-j}}(T^{j-1}(x)) \rightarrow \frac{(1 - \lambda)}{\lambda} \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} y d\mu(x, y, \bar{b}).$$

(d) From (c) we obtain a set $\Omega_{\lambda} \subseteq X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}$ ($\mu_{\lambda}(\Omega_{\lambda}) = 1$), such that, for all $(x, \bar{b} = (b_{-1}, b_{-2}, \dots)) \in \Pi_{\{1,3\}}\Omega$

$$\frac{1}{N} \sum_{j=1}^{N-1} A_{b_{-j}}(T^{j-1}(x)) \rightarrow \frac{(1 - \lambda)}{\lambda} \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} y d\mu(x, y, \bar{b}).$$

On the other hand, we know that $(1 - \lambda) \max v_{\lambda}^+ \rightarrow \bar{u}$, where \bar{u} is the maximum of the integrals of $(x, c, a) \rightarrow A_c(\tau_a(x))$, for all the holonomic measures over $X \times \mathcal{C} \times \mathcal{I}$. Therefore, it is also true

that, $\frac{(1-\lambda)}{\lambda} \max v_\lambda^+ \xrightarrow{\lambda \rightarrow 1} \bar{u}$. So, for any $\varepsilon > 0$ there exists $\lambda_0 \in (0, 1)$, such that, for all $\lambda > \lambda_0$ we have $\frac{(1-\lambda)}{\lambda} \max v_\lambda^+ \leq \bar{u} + \varepsilon$.

We notice that

$$\begin{aligned} \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} y d\mu(x, y, \bar{b}) &= \int_{X \times \mathcal{I}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}} \times \mathcal{C}^{\mathbb{N}}} S_x(\bar{c}, \bar{a}) d\ell(x) d\eta(\bar{a}) d\nu(\bar{c}) d\nu(\bar{b}) \leq \\ &\leq \int_X v_\lambda^+(x) d\ell(x) \leq \max v_\lambda^+. \end{aligned}$$

Multiplying the above equation by $\frac{(1-\lambda)}{\lambda}$ and taking limits we get $N_0 \in \mathbb{N}$, such that, $\frac{1}{N} \sum_{j=1}^{N-1} A_{b_{-j}}(T^{j-1}(x)) \leq \bar{u} + \varepsilon$, for all $N > N_0$. ■

2.5 The associated ergodic optimization problem

In ergodic optimization the central problem is to find the maximum of the integral of a potential function $A : X \rightarrow \mathbb{R}$ with respect to any probability measure μ which is invariant under a dynamic $T : X \rightarrow X$, that is,

$$m_T(A) = \sup_{T(\mu)=\mu} \int_X A(x) d\mu(x).$$

Obviously, this problem has very different solutions depending strongly on A and on T . Moreover, the Birkhoff theorem tells us that invariant measures can be understood using empirical averages, that is,

$$\frac{1}{N} \sum_{i=0}^{N-1} A(T^i(x)) \rightarrow \hat{A}(x)$$

a.e. and $\int_X \hat{A}(x) d\mu(x) = \int_X A(x) d\mu(x)$. In particular, $\hat{A}(x) = \int_X A(x) d\mu(x)$ a.e., when μ is ergodic.

In physical applications or optimal controlling problems these empirical averages represents evaluations of an observable information along of the dynamical trajectory $x_0 = x, x_1 = T(x), x_2 = T^2(x), \dots$. So, it is very natural to suppose that the value A can be influenced by some noise producing a range of possible variations $A_c : X \rightarrow \mathbb{R}$ for $c \in \mathcal{C}$. In such case, the empirical averages will be controlled by a sequence $(c_0, c_1, \dots) \in \mathcal{C}^{\mathbb{N}}$:

$$\frac{1}{N} \sum_{i=0}^{N-1} A_{c_i}(T^i(x)).$$

From Theorem 2.12 we can give an immediate solution for this problem for $X = \mathbf{S}^1$. Is easy to see that if $g(x, y, \bar{b}) = A_{b_{-1}}(x) \in L^1(\mu)$, then there exists a set $\Omega \subseteq X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}$, with $\mu(\Omega) = 1$, such that, for all $(x, y, \bar{b}) \in \Omega$,

$$\begin{aligned} \frac{1}{N} \sum_{j=1}^{N-1} A_{b_{-j}}(T^{j-1}(x)) &\rightarrow \int_{X \times \mathbb{R} \times \mathcal{C}^{\mathbb{N}}} A_{b_{-1}}(x) d\mu(x, y, \bar{b}) = \\ &= \int_{\mathcal{C}^{\mathbb{N}}} \left(\int_X A_{b_{-1}}(x) d\ell(x) \right) d\nu(\bar{b}) = \int_X \frac{1}{m} \sum_{c=0}^{m-1} A_c(x) d\ell(x). \end{aligned}$$

3 IFS Ergodic optimization for a finite family of potentials

In this section we formulate the IFS ergodic optimization problem for a finite family of potentials. Our results naturally generalizes the theory of IFS ergodic optimization developed in [LO09].

As usual, the endomorphism dynamics $x_0 = x, x_1 = T(x), x_2 = T^2(x), \dots$ can be replaced by an IFS dynamics $(X, \tau_i)_{i \in \mathcal{I}}$, that is, $x_0 = x, x_1 = \tau_{a_0}(x_0), x_2 = \tau_{a_1}(x_1), \dots, x_{i+1} = \tau_{a_i}(x_i), \dots$. In this way the empirical averages

$$\frac{1}{N} \sum_{i=0}^{N-1} A_{c_i}(\tau_{a_i}(x_i)),$$

will be controlled by a sequence $(c_0, a_0, c_1, a_1, \dots) \in (\mathcal{C} \times \mathcal{I})^{\mathbb{N}}$.

We want to investigate the variational problem associated to these averages because it is closely related to the behavior of our skew product IFS when the discount parameter λ is close to 1. We call this “**the IFS ergodic optimization for a finite family of potentials**”, $A_c : X \rightarrow \mathbb{R}$, for $c \in \mathcal{C}$. To do that, we need to define what is the appropriated notion of invariant measure and a variational formulation.

Given a finite set of potentials $A_c : X \rightarrow \mathbb{R}$ for $c \in \mathcal{C}$ and an IFS $R = (X, \tau_a)_{a \in \mathcal{I}}$ we want to find the value

$$m(R) = \sup_{\mu \in \mathcal{H}} \int A_c(\tau_a x) d\mu(x, c, a),$$

over the set of holonomic probabilities

$$\mathcal{H} = \left\{ \mu \in \text{Prob}(X \times \mathcal{C} \times \mathcal{I}) \mid \int d_x g(a) d\mu(x, c, a) = 0, \forall g \in C^0(X, \mathbb{R}) \right\},$$

and the optimal measures $\mu_\infty \in \mathcal{H}$, such that, $m(R) = \int A_c(\tau_a x) d\mu_\infty(x, c, a)$.

In the following we are going to show that \mathcal{H} is not empty.

3.1 Empirical holonomic measures

Since not all holonomic measure is the projection of a invariant one for the map $(x, \bar{c}, \bar{a}) \rightarrow (\tau_a x, \sigma \bar{c}, \sigma \bar{a})$, we can not use the Birkhoff theorem to characterize the holonomic measures. However, we can get some similar results introducing the set of empirical holonomic measures.

We consider, for each $n \in \mathbb{N}$, $\bar{c} \in \mathcal{C}^{\mathbb{N}}$ and $\bar{a} \in \mathcal{I}^{\mathbb{N}}$ the empirical measure $\mu_{\bar{c}, \bar{a}, x_0}^n$ given by

$$\int g(x, c, a) d\mu_{\bar{c}, \bar{a}, x_0}^n = \frac{1}{n} \sum_{i=0}^{n-1} g(x_i, c_i, a_i)$$

where x_i is the iteration by the IFS R of some point x_0 , controlled by \bar{a} .

We define

$$\mathcal{H}_0 \equiv \left\{ \mu \mid \begin{array}{l} \mu \text{ is a cluster point, in the weak-}^* \text{ topology,} \\ \text{of the sequence } (\mu_{\bar{c}, \bar{a}, x_0}^n)_{n \in \mathbb{N}} \end{array} \right\}$$

the set of all the empirical holonomic measures.

There exists another ways to obtain holonomic measures. The map $\hat{\sigma}(x, \bar{c}, \bar{a}) = (\tau_a x, \sigma \bar{c}, \sigma \bar{a})$ is continuous and $X \times \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}}$ is compact, thus there exists invariant measures. Moreover, if $M(\hat{\sigma})$ is the set of invariant measures for $\hat{\sigma}$, then $\text{ex}(M(\hat{\sigma}))$, the set of ergodic measures with respect to $\hat{\sigma}$, is not empty. It is easy to see that, for any $\xi \in M(\hat{\sigma})$, the push-forward $\Pi^\sharp(\xi) \in \text{Prob}(X \times \mathcal{C} \times \mathcal{I})$ given by $\int g(x, c, a) d\Pi^\sharp(\xi) = \int g(\Pi(x, \bar{c}, \bar{a})) d\xi$, where $\Pi(x, \bar{c}, \bar{a}) = (x, c_0, a_0)$, is holonomic (by integrating $h(x, \bar{c}, \bar{a}) = d_x w(a_0)$ and using the invariance of $\hat{\sigma}$).

Additionally, if $\xi \in M(\hat{\sigma})$ ergodic, then we get, from the Birkhoff theorem, that there exists a set $\Omega \subset X \times \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}}$, with $\xi(\Omega) = 1$ such that, for all $(x_0, \bar{c}, \bar{a}) \in \Omega$

$$\frac{1}{n} \sum_{i=0}^{n-1} g(\Pi(\hat{\sigma}^i(x_0, \bar{c}, \bar{a}))) \rightarrow \int g(\Pi(\hat{\sigma}(x, \bar{c}, \bar{a}))) d\xi = \int g(x, c, a) d\Pi^\sharp(\xi)$$

for all $g \in C^0(X \times \mathcal{C} \times \mathcal{I}, \mathbb{R})$. Since $\frac{1}{n} \sum_{i=0}^{n-1} g(\Pi(\hat{\sigma}^i(x_0, \bar{c}, \bar{a}))) = \int g(x, c, a) d\mu_{\bar{c}, \bar{a}, x_0}^n$ we conclude that $\Pi^\sharp(\xi) \in \mathcal{H}_0$, because it is a cluster point. Therefore, $\Pi^\sharp(\text{ex}(M(\hat{\sigma}))) \subseteq \mathcal{H}_0$. As Π^\sharp is a linear operator, we get $\Pi^\sharp(M(\hat{\sigma})) \subseteq \mathcal{H}_0$.

The next theorem shows that we can solve the IFS ergodic optimization problem in \mathcal{H}_0 but we do not know if there exists other solutions in $\mathcal{H}/\mathcal{H}_0$.

Theorem 3.1. *The following properties are true:*

- a) $\mathcal{H}_0 \subset \mathcal{H}$.
- b) $\int A_c(\tau_a(x)) d\mu \leq \bar{u}$ for all $\mu \in \mathcal{H}_0$.
- c) If \bar{c} and \bar{a} are optimal in x_0 and $\mu_{\bar{c}, \bar{a}, x_0}^n \rightarrow \mu_\infty$ (up to subsequence $n_k \rightarrow \infty$) then $\int A_c(\tau_a(x)) d\mu_\infty = \bar{u}$.
- d) $\sup_{\mu \in \mathcal{H}_0} \int A_c(\tau_x) d\mu(x, c, a) = \bar{u}$.
- e) $\bar{u} = m(R)$.

Proof. (a) We can easily see that $\mu_{\bar{c}, \bar{a}, x_0}^n$ is a probability, but not necessarily a holonomic measure. But, if $\mu_{\bar{c}, \bar{a}, x_0}^n \rightarrow \mu$, then $\mu \in \mathcal{H}$.

(b) Integrating $g(x, c, a) = A_c(\tau_a(x))$ with respect to $\mu_{\bar{c}, \bar{a}, x_0}^n$, we have

$$\int A_c(\tau_a(x)) d\mu_{\bar{c}, \bar{a}, x_0}^n = \frac{1}{n} \sum_{i=0}^{n-1} A_{c_i}(\tau_{a_i}(x_i)) = \frac{1}{n} \sum_{i=0}^{n-1} A_{c_i}(x_{i+1}),$$

and using the sub-action equation we get

$$\begin{aligned} \bar{u} &\geq d_{x_0} b(a_0) + A_{c_0}(\tau_{a_0}(x_0)) = d_{x_0} b(a_0) + A_{c_0}(x_1), \\ \bar{u} &\geq d_{x_1} b(a_1) + A_{c_1}(\tau_{a_1}(x_1)) = d_{x_1} b(a_1) + A_{c_1}(x_2), \text{ etc.} \end{aligned}$$

Adding these sequences of inequalities we obtain $n\bar{u} \geq b(x_n) - b(x_0) + \sum_{i=0}^{n-1} A_{c_i}(x_{i+1})$, with equality, if and only if, \bar{c} and \bar{a} are optimal in x_0 . Rewriting the above inequality we obtain

$$\bar{u} \geq \frac{b(x_n) - b(x_0)}{n} + \frac{1}{n} \sum_{i=0}^{n-1} A_{c_i}(x_{i+1}) = \frac{b(x_n) - b(x_0)}{n} + \int A_c(\tau_a(x)) d\mu_{\bar{c}, \bar{a}, x_0}^n.$$

As b is continuous and X is compact we can take the limit and conclude that $\int A_c(\tau_a(x))d\mu \leq \bar{u}$.
(c) If \bar{c} and \bar{a} are optimal in x_0 and $\mu_\infty = \lim \mu_{\bar{c}, \bar{a}, x_0}^n$, then $\int A_c(\tau_a(x))d\mu_\infty = \bar{u}$.

(d) From (b) and (c) we obtain that $\sup_{\mu \in \mathcal{H}_0} \int A_c(\tau_x)d\mu(x, c, a) = \bar{u}$.

(e) From the Bellman equation $\bar{u} \geq d_x b(a) + A_c(\tau_a(x))$, we obtain that, for all $\mu \in \mathcal{H}$, $\int A_c(\tau_x)d\mu(x, c, a) \leq \bar{u}$. From (a), $\mathcal{H}_0 \subset \mathcal{H}$, thus $m(R) = \bar{u}$. \blacksquare

3.2 Empirical discounted holonomic measures

By analogy with [Gom08] we define the set of discounted holonomic measures with discount, $0 < \lambda < 1$, and trace $\nu \in \text{Prob}(X)$, as being the set

$$\mathcal{H}^\lambda(\nu) = \left\{ \mu \in \text{Prob}(X \times \mathcal{C} \times \mathcal{I}) \mid \int d_x^\lambda w(a) d\mu(x, c, a) = \right. \\ \left. -(1 - \lambda) \int w(x) d\nu(x), \forall w \in C^0(X, \mathbb{R}) \right\},$$

where $d_x^\lambda w(a) = \lambda w(\tau_a x) - w(x)$ is the discounted discrete differential of a continuous function w . By an abuse of notation, we can write $d_x^\lambda w(a) = d_x w(a)$, then $\mathcal{H}^1(\nu) = \mathcal{H}$, for any ν . As before, we can formulate the discounted IFS ergodic optimization problem for a finite family of potentials as follows:

Given a finite set of potentials $A_c : X \rightarrow \mathbb{R}$, for $c \in \mathcal{C}$, a discount $0 < \lambda < 1$, a trace $\nu \in \text{Prob}(X)$ and an IFS $R = (X, \tau_a)_{a \in \mathcal{I}}$, we want to find the value

$$m_\lambda(R) = \sup_{\mu \in \mathcal{H}^\lambda(\nu)} \int A_c(\tau_a x) d\mu(x, c, a),$$

over the set of discounted holonomic probabilities $\mathcal{H}^\lambda(\nu)$ and the optimal measures $\mu_\infty \in \mathcal{H}^\lambda(\nu)$, such that, $m_\lambda(R) = \int A_c(\tau_a x) d\mu_\infty(x, c, a)$.

Theorem 3.2. *The following properties are true:*

a) For any trace measure $\nu \in \text{Prob}(X)$, we get $\gamma \in \mathcal{H}^\lambda(\nu)$, where $\gamma : C^0(X, \mathbb{R}) \rightarrow \mathbb{R}$ is given by $\gamma(g) = (1 - \lambda) \int_X \sum_{i=0}^{\infty} \lambda^i g(x_i, c_i, a_i) d\nu(x)$, for a fixed pair $(\bar{c}, \bar{a}) \in \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}}$ and $x_0 = x \in X$. In particular the set $\mathcal{H}^\lambda(\nu)$ is not empty.

b) For any trace $\nu \in \text{Prob}(X)$, and γ given in (a), we have $m_\lambda(R) = (1 - \lambda) \int v_\lambda(x) d\nu(x)$ where $v_\lambda(x)$ is the solution of the Bellman equation $0 = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) + d_x^\lambda v_\lambda(a)$.

c) For any $z \in X$ the value of the problem for $\mathcal{H}^\lambda(\delta_z)$ is $m_\lambda(R) = (1 - \lambda)v_\lambda(z)$.

d) We have the formula

$$v_\lambda(z) = \inf_{w \in C^0(X, \mathbb{R})} \left[w(z) + \frac{1}{1 - \lambda} \sup_{x \in X} h_w(x) \right],$$

where $h_w(x) = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} \{d_x^\lambda w(a) + A_c(\tau_a x)\}$.

Proof. (a) To see that $\mathcal{H}^\lambda(\nu)$ is not empty, we consider the functional $\gamma : C^0(X, \mathbb{R}) \rightarrow \mathbb{R}$ given by $\gamma(g) = (1-\lambda) \int_X \sum_{i=0}^{\infty} \lambda^i g(x_i, c_i, a_i) d\nu(x)$, for a fixed pair $(\bar{c}, \bar{a}) \in \mathcal{C}^\mathbb{N} \times \mathcal{I}^\mathbb{N}$ and $x_0 = x \in X$. This linear functional is well defined and continuous, because $x \rightarrow \sum_{i=0}^{\infty} \lambda^i g(x_i, c_i, a_i)$ is bounded. Moreover, γ is positive and $\gamma(1) = 1$. Therefore, γ is a probability. Computing $\sum_{i=0}^{n-1} \lambda^i d_{x_i}^\lambda w(a_i) = \lambda^n w(x_n) - w(x)$

and taking the limit when $n \rightarrow \infty$ we can see that $\int d_x^\lambda w(a) d\gamma(x, c, a) = (1-\lambda) \int_X w(x) d\nu(x)$, thus $\gamma \in \mathcal{H}^\lambda(\nu)$.

(b) From the Bellman equation $0 = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) + d_x^\lambda v_\lambda(a)$ we get the inequality $A_c(\tau_a(x)) \leq -d_x^\lambda v_\lambda(a)$. Integrating the above inequality with respect to any $\mu \in \mathcal{H}^\lambda(\nu)$, we get $m_\lambda(R) \leq (1-\lambda) \int v_\lambda(x) d\nu(x)$. To obtain the equality we choose, for each $x \in X$, an optimal pair $(\bar{c}, \bar{a}) \in \mathcal{C}^\mathbb{N} \times \mathcal{I}^\mathbb{N}$, that is, $A_{\bar{c}_i}(\tau_{\bar{a}_i}(x_i)) = -d_{x_i}^\lambda v_\lambda(a_i)$ for all i , for each $x_0 = x$, defining a multifunction $x \rightarrow (\bar{c}, \bar{a})$. This u.s.c. multifunction (see the Maximum Theorem and Theorem 3, Chap. VI, §3, in [Ber97] for selective properties) has a continuous selection $x \rightarrow (\bar{c}(x), \bar{a}(x))$, making the correspondence $x \rightarrow \sum_{i=0}^{\infty} \lambda^i g(x_i, c_i, a_i)$ continuous. Therefore $m_\lambda(R) = (1-\lambda) \int v_\lambda(x) d\nu(x)$.

(c) It is a consequence of (b), when $\nu = \delta_z$.

(d) Using a duality argument similar to [Gom08] we prove, in Section 4 Theorem 4.2, that the above problem always has solution

$$\inf_{w \in C^0(X, \mathbb{R})} \left[(1-\lambda) \int w(x) d\nu(x) + \sup_{x \in X, c \in \mathcal{C}, a \in \mathcal{I}} \{d_x^\lambda w(a) + A_c(\tau_a x)\} \right] = \max_{\mu \in \mathcal{H}^\lambda(\nu)} \int A_c(\tau_a x) d\mu(x, c, a),$$

for any trace measure ν . Using this formula for $\nu = \delta_z \in \text{Prob}(X)$, for an arbitrary $z \in X$, we get a variational formula for v_λ

$$v_\lambda(z) = \inf_{w \in C^0(X, \mathbb{R})} \left[w(z) + \frac{1}{1-\lambda} \sup_{x \in X} h_w(x) \right],$$

where $h_w(x) = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} \{d_x^\lambda w(a) + A_c(\tau_a x)\}$. ■

Inspired by Theorem 3.2, we are going to consider the particular case, $\nu = \delta_{x_0} \in \text{Prob}(X)$, where $x_0 = \text{argmax}_{x \in X} v_\lambda(x)$.

Definition 3.3. We consider, for each $\bar{c} \in \mathcal{C}^\mathbb{N}$ and $\bar{a} \in \mathcal{I}^\mathbb{N}$, the **empirical discounted probability measure** $\mu_{x_0, \bar{c}, \bar{a}}^\lambda$ given by

$$\int g(x, c, a) d\mu_{x_0, \bar{c}, \bar{a}}^\lambda = (1-\lambda) \sum_{i=0}^{\infty} \lambda^i g(x_i, c_i, a_i)$$

where, x_i is the i -iterate by the IFS R of some point $x_0 \in X$.

These measures are quite different from the empirical ones because each one is not an average in n , or subsequences, but they are defined only for $\lambda < 1$. Moreover, if $\mu = \mu_{x_0, \bar{c}, \bar{a}}^\lambda$, then

$$\int A_c(\tau_a x) d\mu(x, c, a) = (1 - \lambda) \sum_{i=0}^{\infty} \lambda^i A_{c_i}(\tau_{a_i} x_i) = (1 - \lambda) S_{x_0}(\bar{c}, \bar{a}),$$

showing the connection between this maximization problem and the superior boundary of the set Λ .

Theorem 3.4. *We consider, for each $\bar{c} \in \mathcal{C}^{\mathbb{N}}$ and $\bar{a} \in \mathcal{I}^{\mathbb{N}}$, the empirical discounted probability measure $\mu_{x_0, \bar{c}, \bar{a}}^\lambda$. Then, the following statements are true:*

- a) $\mu_{x_0, \bar{c}, \bar{a}}^\lambda \in \mathcal{H}^\lambda(\delta_{x_0})$.
- b) Consider $x_0 = \operatorname{argmax}_{x \in X} v_\lambda(x)$ and $\nu = \delta_{x_0} \in \operatorname{Prob}(X)$. If the pair $(\bar{c}, \bar{a}) \in \mathcal{C}^{\mathbb{N}} \times \mathcal{I}^{\mathbb{N}}$ is optimal then, $m_\lambda(R) = (1 - \lambda) \max_{x \in X} v_\lambda(x)$ and $\mu^\lambda = \mu_{x_0, \bar{c}, \bar{a}}^\lambda \in \mathcal{H}^\lambda(\delta_{x_0})$ solves the discounted IFS ergodic optimization problem for the finite family of potentials.

Proof. (a) Note that

$$\begin{aligned} \int d_x^\lambda w(a) d\mu_{x_0, \bar{c}, \bar{a}}^\lambda &= (1 - \lambda) \sum_{i=0}^{+\infty} \lambda^i d_{x_i}^\lambda w(a_i) = (1 - \lambda) \lim_{n \rightarrow +\infty} (\lambda^n w(x_n) - w(x_0)) = \\ &= -(1 - \lambda) w(x_0) = -(1 - \lambda) \int w(x) d\delta_{x_0}(x). \end{aligned}$$

Thus, $\mu_{x_0, \bar{c}, \bar{a}}^\lambda \in \mathcal{H}^\lambda(\delta_{x_0})$.

(b) We recall that the integral $\int A_c(\tau_a x) d\mu_{x_0, \bar{c}, \bar{a}}^\lambda$ is the limit

$$\int A_c(\tau_a x) d\mu_{x_0, \bar{c}, \bar{a}}^\lambda = (1 - \lambda) \lim_{n \rightarrow +\infty} \sum_{i=0}^{n-1} \lambda^i A_{c_i}(\tau_{a_i} x_i).$$

We can use the Bellman equation $v_\lambda(x) = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) + \lambda v_\lambda(\tau_a(x))$ to construct a special discounted measure. We observe that the Bellman equation could be rewritten as

$$0 = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) + \lambda v_\lambda(\tau_a(x)) - v_\lambda(x) = \sup_{c, a \in \mathcal{C} \times \mathcal{I}} A_c(\tau_a(x)) + d_x^\lambda v_\lambda(a).$$

In particular, $A_c(\tau_a(x)) \leq -d_x^\lambda v_\lambda(a)$, $\forall (c, a) \in \mathcal{C} \times \mathcal{I}$, and the equality is attained for some pair $(c_0, a_0) \in \mathcal{C} \times \mathcal{I}$ in each point $x \in X$.

From the above inequality we obtain that for any measure μ the following inequality holds

$$\int A_c(\tau_a(x)) d\mu(x, c, a) \leq \int -d_x^\lambda v_\lambda(a) d\mu(x, c, a).$$

If we additionally suppose that $\mu \in \mathcal{H}^\lambda(\nu)$, then

$$\int A_c(\tau_a(x))d\mu(x, c, a) \leq -(1 - \lambda) \int -v_\lambda(x)d\nu(x).$$

Thus, $m_\lambda(R) \leq (1 - \lambda) \int v_\lambda(x)d\nu(x) = (1 - \lambda)v_\lambda(x_0)$ in the case where $\nu = \delta_{x_0}$.

We take the measure $\mu^\lambda = \mu_{x_0, \bar{c}, \bar{a}}^\lambda$ by choosing $x_0 = \operatorname{argmax}_{x \in X} v_\lambda(x)$ and \bar{c} and \bar{a} in, such way, that $0 = A_{c_i}(\tau_{a_i}(x_i)) + d_{x_i}^\lambda v_\lambda(a_i)$, or $A_{c_i}(\tau_{a_i}(x_i)) = -d_{x_i}^\lambda v_\lambda(a_i)$, for all $i \in \mathbb{N}$. Thus,

$$\int A_c(\tau_a x) d\mu^\lambda = (1 - \lambda) \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} \lambda^i (-d_{x_i}^\lambda v_\lambda(a_i)) = (1 - \lambda) \max_{x \in X} v_\lambda(x).$$

It means that $m_\lambda(R) = (1 - \lambda) \max_{x \in X} v_\lambda(x)$ and then μ^λ solves the discounted ergodic optimization problem for the finite family of potentials for $\nu = \delta_{x_0} \in \operatorname{Prob}(X)$. \blacksquare

Remark 3.5. From [CO17] we know that $(1 - \lambda) \max_{x \in X} v_\lambda(x) \rightarrow \bar{u}$, thus if $\mu^\lambda \rightarrow \mu$, through some subsequence $\lambda_j \rightarrow 1$, then $\mu \in \mathcal{H}$ and $\int A_c(\tau_a x) d\mu = \lim_{\lambda_j \rightarrow 1} \int A_c(\tau_a x) d\mu^{\lambda_j} = (1 - \lambda_j) \max_{x \in X} v_{\lambda_j}(x) = \bar{u}$. So, we can obtain maximizing holonomic measures as limit of discounted holonomic measures with trace δ_{x_0} , arising from the solutions of Bellman's equation.

4 Appendix: Duality for Discounted Holonomic Measures

For the sake of completeness, we state the duality result used in the Theorem 3.2. See [Vil03] for a proof of the Fenchel-Rockafellar duality theorem. See [Gom02] and [Gom08] for applications in optimization and variational problems.

Theorem 4.1 (Fenchel-Rockafellar duality). *Suppose that E is a normed vector space, Γ and Φ are two convex functions defined on E taking values in $\mathbb{R} \cup \{+\infty\}$. Denote Γ^* and Φ^* , respectively, the Legendre-Fenchel transform of Γ and Φ . Suppose there exists $\psi_0 \in E$, such that, $\Gamma(\psi_0) < +\infty$, $\Phi(\psi_0) < +\infty$, and that Γ is continuous on ψ_0 . Then,*

$$\inf_{\psi \in E} [\Gamma(\psi) + \Phi(\psi)] = \sup_{\pi \in E^*} [-\Gamma^*(-\pi) - \Phi^*(\pi)] \quad (1)$$

Moreover, the supremum in (1) is attained in at least one element in $\pi \in E^*$.

Given a finite set of potentials $A_c : X \rightarrow \mathbb{R}$ for $c \in \mathcal{C}$, a discount $0 < \lambda < 1$, a trace $\nu \in \operatorname{Prob}(X)$ and an IFS $R = (X, \tau_a)_{a \in \mathcal{I}}$ we want to find the dual of

$$m_\lambda(R) = \sup_{\mu \in \mathcal{H}^\lambda(\nu)} \int A_c(\tau_a x) d\mu(x, c, a),$$

over the set of discounted holonomic probabilities $\mathcal{H}^\lambda(\nu)$ satisfying $\mu \in \mathcal{H}^\lambda(\nu)$

$$\int d_x^\lambda w(a) d\mu(x, c, a) = -(1 - \lambda) \int w(x) d\nu(x), \quad (2)$$

for all $w \in C^0(X, \mathbb{R})$. Here, $d_x^\lambda w(a) = \lambda w(\tau_a x) - w(x)$ be the discounted discrete differential of a continuous function w .

Theorem 4.2. *Under the above conditions, there exists at least one μ satisfying the equation (2), such that,*

$$\inf_{w \in C^0(X, \mathbb{R})} \left[(1 - \lambda) \int w(x) d\nu(x) + \sup_{x \in X, c \in \mathcal{C}, a \in \mathcal{I}} \{d_x^\lambda w(a) + A_c(\tau_a x)\} \right] = \max_{\mu \in \mathcal{H}^\lambda(\nu)} \int A_c(\tau_a x) d\mu(x, c, a). \quad (3)$$

Proof. We define for $\psi \in C(X \times \mathcal{C} \times \mathcal{I})$ the maps

$$\Gamma(\psi) = \sup_{X \times \mathcal{C} \times \mathcal{I}} \{ \psi(x, c, a) + A_c(\tau_a x) \},$$

and

$$\Phi(\psi) = \begin{cases} (1 - \lambda) \int w(x) d\nu(x), & \text{if } \psi(x, c, a) = d_x^\lambda w(a) \text{ for some } w(x) \\ +\infty, & \text{otherwise} \end{cases}.$$

Φ is well defined because if π_0 is any measure with trace ν and if $\psi = d_x^\lambda w_1(a) = d_x^\lambda w_2(a)$, then, $\int d_x^\lambda w_1(a) d\pi_0 = \int d_x^\lambda w_2(a) d\pi_0$, therefore

$$(1 - \lambda) \int w_1(x) d\nu(x) = (1 - \lambda) \int w_2(x) d\nu(x).$$

The functions Γ and Φ are trivially convex. To fulfill the hypothesis of Theorem 4.1 we need to find one function ψ , such that, $\Gamma(\psi), \Phi(\psi) < \infty$ and Γ is continuous in ψ . Then, take any ψ in the form $\psi = d_x^\lambda w(a)$, for some $w(x)$, and use that Γ is continuous for the supremum norm.

Now we study $\Gamma^*(\pi)$. First suppose that π is not a positive functional. Then, there exists $u \leq 0$, such that, $\pi(u) > 0$. We write $\psi_t(x, c, a) = tu(x, c, a) - A_c(\tau_a x)$. Then,

$$\begin{aligned} \Gamma^*(\pi) &= \sup_{\psi} \left\{ \int \psi d\pi - \Gamma(\psi) \right\} \geq \limsup_{t \rightarrow \infty} \int \psi_t(x, y) d\pi - \Gamma(\psi_t) \\ &= \limsup_{t \rightarrow \infty} \int \lambda u - A_c(\tau_a x) d\pi - \Gamma(tu - A_c(\tau_a x)) = \limsup_{t \rightarrow \infty} \int -A_c(\tau_a x) d\pi + t \int u d\pi - t \sup u \\ &\geq \limsup_{t \rightarrow \infty} \int -A_c(\tau_a x) d\pi + t \int u d\pi = +\infty. \end{aligned}$$

Suppose that π is a positive functional. If $\int 1 d\pi \neq 1$, then for any $t \in \mathbb{R}$:

$$\begin{aligned} \Gamma^*(\pi) &= \sup_{\psi} \left\{ \int \psi d\pi - \Gamma(\psi) \right\} \geq \left\{ \int t - A_c(\tau_a x) d\pi - \Gamma(t - A_c(\tau_a x)) \right\} \\ &= \int -A_c(\tau_a x) d\pi + \int t d\pi - t = \int -A_c(\tau_a x) d\pi + t \left(\int 1 d\pi - 1 \right). \end{aligned}$$

If $\int 1 d\pi > 1$, we get $\Gamma^*(\pi) \geq \limsup_{t \rightarrow +\infty} \int -A_c(\tau_a x) d\pi + t(\int 1 d\pi - 1) = +\infty$. If $\int 1 d\pi < 1$, we get $\Gamma^*(\pi) \geq \limsup_{t \rightarrow -\infty} \int -A_c(\tau_a x) d\pi + t(\int 1 d\pi - 1) = +\infty$. We conclude that if $\int 1 d\pi \neq 1$, then, $\Gamma^*(\pi) = +\infty$.

Now, we suppose that π is a probability. Then:

$$\Gamma^*(\pi) = \sup_{\psi} \left\{ \int \psi d\pi - \Gamma(\psi) \right\} =$$

$$\begin{aligned}
&= \sup_{\psi} \left\{ \int -A_c(\tau_a x) d\pi + \int \psi(x, y) + A_c(\tau_a x) d\pi - \Gamma(\psi) \right\} = \\
&= \sup_{\psi} \left\{ \int -A_c(\tau_a x) d\pi + \int (\psi + A_c(\tau_a x)) - \Gamma(\psi) d\pi \right\} \leq \\
&\leq \sup_{\psi} \left\{ \int -A_c(\tau_a x) d\pi + 0 \right\} = \int -A_c(\tau_a x) d\pi.
\end{aligned}$$

On the other hand,

$$\Gamma^*(\pi) = \sup_{\psi} \left\{ \int \psi d\pi - \Gamma(\psi) \right\} \geq \left\{ \int -A_c(\tau_a x) d\pi - \Gamma(-A_c(\tau_a x)) \right\} = \int -A_c(\tau_a x) d\pi$$

Thus, we conclude that

$$\Gamma^*(\pi) = \begin{cases} -\int A_c(\tau_a x) d\pi(x, c, a) & \text{if } \pi \text{ is a probability} \\ +\infty & \text{otherwise} \end{cases}.$$

We recall that $\Phi(\psi) = (1 - \lambda) \int w(x) d\nu(x)$ if $\psi(x, c, a) = d_x^\lambda w(a)$, for some $w(x)$ and $+\infty$, otherwise. As the Legendre transform is

$$\Phi^*(-\pi) = \sup_{\psi} [\langle -\pi, \psi \rangle - \Phi(\psi)],$$

we must consider the supremum just for $\psi(x, c, a) = d_x^\lambda w(a)$, thus,

$$= \sup_w \left[-\int d_x^\lambda w(a) d\pi - ((1 - \lambda) \int w(x) d\nu(x)) \right] = 0,$$

so

$$\Phi^*(-\pi) = \begin{cases} 0, & \text{if } \pi \text{ is holonomic} \\ +\infty, & \text{otherwise} \end{cases}.$$

Then,

$$\inf_{\psi \in E} [\Gamma(\psi) + \Phi(\psi)] = \inf_{w(x)} \left[(1 - \lambda) \int w(x) d\nu(x) + \sup_{(x, c, a)} \{d_x^\lambda w(a) + A_c(\tau_a x)\} \right]$$

and

$$\sup_{\pi \in E^*} [-\Gamma^*(-\pi) - \Phi^*(\pi)] \stackrel{\pi \leftrightarrow -\pi}{=} \sup_{\pi \in E^*} [-\Gamma^*(\pi) - \Phi^*(-\pi)] = \sup_{\pi \in \mathcal{H}^\lambda(\nu)} \int A_c(\tau_a x) d\pi.$$

From equation (1) we conclude the proof. ■

References

- [Ber97] Claude Berge. *Topological spaces*. Dover Publications, Inc., Mineola, NY, 1997. Including a treatment of multi-valued functions, vector spaces and convexity, Translated from the French original by E. M. Patterson, Reprint of the 1963 translation.

- [BKRLU06] Rodrigo Bamón, Jan Kiwi, Juan Rivera-Letelier, and Richard Urzúa. On the topology of solenoidal attractors of the cylinder. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 23(2):209–236, 2006.
- [BV11] Michael F. Barnsley and Andrew Vince. The chaos game on a general iterated function system. *Ergodic Theory Dynam. Systems*, 31(4):1073–1079, 2011.
- [CO17] L. Cioletti and E. R. Oliveira. Applications of variable discounting dynamic programming to iterated function systems and related problems. *Preprint*, 2017.
- [DaGR17] L. J. Díaz, K. Gelfert, and M. Rams. Topological and ergodic aspects of partially hyperbolic diffeomorphisms and nonhyperbolic step skew products. *Tr. Mat. Inst. Steklova*, 297(Poryadok i Khaos v Dinamicheskikh Sistemakh):113–132, 2017.
- [GL08] E. Garibaldi and A. O. Lopes. On the Aubry-Mather theory for symbolic dynamics. *Ergodic Theory Dynam. Systems*, 28(3):791–815, 2008.
- [GLT09] E. Garibaldi, A. O. Lopes, and Ph. Thieullen. On calibrated and separating sub-actions. *Bull. Braz. Math. Soc. (N.S.)*, 40(4):577–602, 2009.
- [Gom02] Diogo Aguiar Gomes. A stochastic analogue of Aubry-Mather theory. *Nonlinearity*, 15(3):581–603, 2002.
- [Gom05] Diogo A. Gomes. Viscosity solution methods and the discrete Aubry-Mather problem. *Discrete Contin. Dyn. Syst.*, 13(1):103–116, 2005.
- [Gom08] Diogo Aguiar Gomes. Generalized Mather problem and selection principles for viscosity solutions and Mather measures. *Adv. Calc. Var.*, 1(3):291–307, 2008.
- [JMN14] Anna Jaśkiewicz, Janusz Matkowski, and Andrzej S. Nowak. On variable discounting in dynamic programming: applications to resource extraction and other economic models. *Ann. Oper. Res.*, 220:263–278, 2014.
- [LO09] Artur O. Lopes and Elismar R. Oliveira. Entropy and variational principles for holonomic probabilities of IFS. *Discrete Contin. Dyn. Syst.*, 23(3):937–955, 2009.
- [LO17] Artur O. Lopes and Elismar R. Oliveira. On the thin boundary of the fat attractor. In *Modeling, Dynamics, Optimization and Bioeconomics III*. Springer International Publishing, 2017.
- [Ram03] Michał Rams. Absolute continuity of the SBR measure for non-linear fat baker maps. *Nonlinearity*, 16(5):1649–1655, 2003.
- [Tsu01] Masato Tsujii. Fat solenoidal attractors. *Nonlinearity*, 14(5):1011–1027, 2001.
- [Vil03] Cédric Villani. *Topics in optimal transportation*, volume 58 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2003.