

On Darboux integrable semi-discrete systems of exponential type

Kostyantyn Zheltukhin¹ and Ergun Bilen²

¹ Department of Mathematics, Faculty of Science,
Middle East Technical University 06531 Ankara, Turkey.
e-mail: zheltukh@metu.edu.tr

² Department of Mathematics, Faculty of Science,
Middle East Technical University 06531 Ankara, Turkey.
e-mail: ergun.bilen@metu.edu.tr

November 14, 2021

Abstract

In the present paper we consider a discretization of hyperbolic systems of exponential type. We proved that, in the case of 2×2 systems, the resulting semi-discrete system is Darboux integrable only if it corresponds to a Cartan matrix of a semi-simple Lie algebra.

2000 Mathematics Subject Classification: 37K10, 17B80, 39A99.

Keywords: Hyperbolic semi-discrete equations; Darboux integrability; Characteristic ring, Cartan Matrices.

1 Introduction

In a recent paper [1] Habibullin et al proposed a method for discretization of hyperbolic type exponential systems

$$r_{xy}^i = e^{\sum a_{ij} r^j}. \quad (1)$$

It was conjectured that the method preserves integrability of such systems. The integrability of system (1) and its discretization is understood as Darboux integrability [2]. It was shown in [3] that system (1) is Darboux integrable if and only if A is a Cartan matrix of a semi-simple Lie algebra. To check for Darboux integrability we use the notion of characteristic rings introduced in [3], [4]. Later the notion of characteristic ring was applied to classify Darboux integrable hyperbolic equations in [5]-[10], see also a review paper [11]. For discrete equations the Darboux integrability was considered in [12], [14]. The notion of characteristic rings for discrete systems and its application to classification of such systems was developed in [13]-[21]

We note that the discrete analogs of system (1) find applications in many areas of physics, see a review paper [22].

Let us give necessary definitions. Consider an $N \times N$ matrix $A = (a_{ij})$ and a corresponding exponential system. Following [1] we decompose A into a sum $A = A_+ + A_-$, where $A_+ = (a_{ij}^+)$ is an upper triangular matrix with all diagonal entries equal to 1 and $A_- = (a_{ij}^-)$ is a lower triangular matrix. The corresponding differential-difference exponential type system is

$$q_{1x}^i - q_x^i = e^{\sum a_{ij}^+ q_1^j + \sum a_{ij}^- q^j}. \quad (2)$$

In this equation $q^i(n, x)$, $i = 1, \dots, N$, are unknown functions depending on discrete variable n and continuous variable x , and $q_1^i = q^i(n+1, x)$ denotes a shift in n and $q_x^i = \frac{d}{dx} q^i(n, x)$ denotes derivative with respect to x .

Conjecture 1. (see [1]) *A system (2) is Darboux integrable if and only if A is a Cartan matrix of a semi-simple Lie algebra.*

As was shown in [1] if $N = 2$ and A is a Cartan matrix of a semi-simple Lie algebra then the corresponding system (2) is Darboux integrable. In the present paper we prove that for $N = 2$ Darboux integrability of the system (2) implies that A is a Cartan matrix of a semi-simple Lie algebra. Our proof is based on the notion of the characteristic x -ring. We recall that a hyperbolic system is Darboux integrable if and only if its characteristic x -

and n -rings are finite dimensional (see [14]). It turns out that it is enough to consider only the characteristic x -ring. The characteristic x -ring is finite dimensional if and only if A is a Cartan matrix of a semi-simple Lie algebra.

Theorem 2. *When $N = 2$, the system (2) is Darboux integrable if and only if the matrix $A = (a_{ij})_{2 \times 2}$ is the Cartan matrix.*

The paper is organized as follows in Section 2 we describe the characteristic x -ring corresponding to the system (2) with general matrix A , with $N = 2$, and in Section 3 we determine for which matrices A the characteristic x -ring is finite dimensional and we prove the above theorem.

2 Characteristic x -ring

Let $A = (a_{ij})$, $i, j = 1, 2$, be an arbitrary matrix. Consider a system

$$\begin{aligned} u_{1x} - u_x &= e^{(a_{11}-1)u+u_1+a_{12}v_1} \\ v_{1x} - v_x &= e^{a_{21}u+(a_{22}-1)v+v_1}. \end{aligned} \quad (3)$$

We assume that $a_{12}, a_{21} \neq 0$, otherwise the system reduces to a pair of ordinary differential equations. In what follow we use notation $u_k = u(n+k, x)$, $v_k = v(n+k, x)$ and $u_{kx} = \frac{d}{dx}u_k$, $v_{kx} = \frac{d}{dx}v_k$.

The characteristic x -ring for system (3), denoted \mathcal{X} , is generated by the vector field (see [14])

$$Z = \frac{\partial}{\partial x} + \sum_{k=-\infty}^{\infty} (u_{kx} \frac{\partial}{\partial u_k} + v_{kx} \frac{\partial}{\partial v_k}),$$

which corresponds to the total derivative with respect to x , and vector fields

$$Y_1 = \frac{\partial}{\partial u_x}, \quad Y_2 = \frac{\partial}{\partial v_x}.$$

It follows from (3) that

$$\begin{aligned} u_{kx} &= u_x + \sum_{i=1}^k e^{(a_{11}-1)u_{i-1}+u_i+a_{12}v_i}, & u_{-kx} &= u_x - \sum_{i=1}^k e^{(a_{11}-1)u_{-i}+u_{-i+1}+a_{12}v_{-i+1}} \\ v_{kx} &= v_x + \sum_{i=1}^k e^{(a_{22}-1)v_{i-1}+v_i+a_{21}u_{i-1}}, & v_{-kx} &= v_x - \sum_{i=1}^k e^{(a_{22}-1)v_{-i}+v_{-i+1}+a_{21}u_{-i}}. \end{aligned}$$

For convenience we define

$$\begin{aligned} M_i &= e^{(a_{11}-1)u_{i-1}+u_i+a_{12}v_i}, & M_{-i} &= e^{(a_{11}-1)u_{-i}+u_{-i+1}+a_{12}v_{-i+1}} \\ N_i &= e^{(a_{22}-1)v_{i-1}+v_i+a_{21}u_{i-1}}, & N_{-i} &= e^{(a_{22}-1)v_{-i}+v_{-i+1}+a_{21}u_{-i}}, \end{aligned}$$

$i = 1, 2, \dots$. So the vector field Z takes form

$$\begin{aligned} Z &= \frac{\partial}{\partial x} + u_x \frac{\partial}{\partial u} + \sum_{k=1}^{\infty} \left(u_x + \sum_{i=1}^k M_i \right) \frac{\partial}{\partial u_k} + \sum_{k=1}^{\infty} \left(u_x - \sum_{i=1}^k M_{-i} \right) \frac{\partial}{\partial u_{-k}} \\ &\quad + v_x \frac{\partial}{\partial v} + \sum_{k=1}^{\infty} \left(v_x + \sum_{i=1}^k N_i \right) \frac{\partial}{\partial v_k} + \sum_{k=1}^{\infty} \left(v_x - \sum_{i=1}^k N_{-i} \right) \frac{\partial}{\partial v_{-k}}. \end{aligned} \quad (4)$$

Taking commutator of Z with Y_1 and Y_2 we obtain vector fields

$$A = [Y_1, Z] = \sum_{j=-\infty}^{\infty} \frac{\partial}{\partial u_j}, \quad B = [Y_2, Z] = \sum_{j=-\infty}^{\infty} \frac{\partial}{\partial v_j}$$

The commutators of Z with A and B give us vector fields

$$\begin{aligned} [A, Z] &= a_{11} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k M_i \right) \frac{\partial}{\partial u_k} - a_{11} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k M_{-i} \right) \frac{\partial}{\partial u_{-k}} \\ &\quad + a_{21} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k N_i \right) \frac{\partial}{\partial v_k} - a_{21} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k N_{-i} \right) \frac{\partial}{\partial v_{-k}} \end{aligned}$$

and

$$\begin{aligned} [B, Z] &= a_{12} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k M_i \right) \frac{\partial}{\partial u_k} - a_{12} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k M_{-i} \right) \frac{\partial}{\partial u_{-k}} \\ &\quad + a_{22} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k N_i \right) \frac{\partial}{\partial v_k} - a_{22} \sum_{k=1}^{\infty} \left(\sum_{i=1}^k N_{-i} \right) \frac{\partial}{\partial v_{-k}}. \end{aligned}$$

It is convenient to replace vector fields $[A, Z]$ and $[B, Z]$ by their linear combinations

$$P_1 = \frac{1}{a_{22}a_{11} - a_{21}a_{12}} (a_{22}[A, Z] - a_{21}[B, Z])$$

and

$$P_2 = \frac{1}{a_{22}a_{11} - a_{21}a_{12}}(-a_{12}[A, Z] + a_{11}[B, Z]).$$

So, vector fields P_1 and P_2 are

$$P_1 = \sum_{k=1}^{\infty} \left(\sum_{i=1}^k M_i \right) \frac{\partial}{\partial u_k} - \sum_{k=1}^{\infty} \left(\sum_{i=1}^k M_{-i} \right) \frac{\partial}{\partial u_{-k}}$$

and

$$P_2 = \sum_{k=1}^{\infty} \left(\sum_{i=1}^k N_i \right) \frac{\partial}{\partial v_k} - \sum_{k=1}^{\infty} \left(\sum_{i=1}^k N_{-i} \right) \frac{\partial}{\partial v_{-k}}.$$

Since $Z = \frac{\partial}{\partial x} + u_x A + v_x B + P_1 + P_2$, we can consider vector fields $\frac{\partial}{\partial x}$, Y_1 , Y_2 , A , B , P_1 and P_2 as generators of the characteristic x -ring \mathcal{X} .

The vector fields $\frac{\partial}{\partial x}$, Y_1 , Y_2 , A and B are pairwise commuting and vector fields $\frac{\partial}{\partial x}$, Y_1 , Y_2 commute with vector fields P_1 , P_2 . The commutators of A , B with P_1 , P_2 are

$$[A, P_1] = a_{11}P_1, \quad [B, P_1] = a_{12}P_1, \quad [A, P_2] = a_{21}P_2, \quad [B, P_2] = a_{22}P_2.$$

Thus the characteristic x -ring \mathcal{X} is finite dimensional if and only if the ring generated by vector fields P_1 , P_2 , denoted \mathcal{A} , is finite dimensional.

If we define a sequence of linear subspaces

$$\mathbb{X}_0 = \text{Lin}\{P_1, P_2\},$$

and

$$\mathbb{X}_n = \text{Lin}\{[P_i, V] : V \in \mathbb{X}_{n-1}, i = 1, 2\}, \quad n = 1, 2, \dots$$

then it is easy to see that

$$\mathcal{A} = \mathbb{X}_0 \oplus \mathbb{X}_1 \oplus \mathbb{X}_2 + \dots \quad (5)$$

We define sequences of vector fields T_n , R_n , $n = 1, 2, \dots$, by

$$T_n = [P_1, T_{n-1}], \quad n = 2, 3, \dots \quad \text{and} \quad T_1 = [P_1, P_2],$$

$$R_n = [P_2, R_{n-1}], \quad n = 2, 3, \dots \quad \text{and} \quad R_1 = T_1$$

3 Proof of Theorem 2

From the decomposition (5) we see that \mathcal{A} is finite dimensional if and only if $\mathbb{X}_n = 0$ for some n . The following lemma (see [14]) is very useful to derive conditions arising from $\mathbb{X}_n = 0$.

Lemma 3. *Suppose that the vector field*

$$K = \sum_{k=1}^{\infty} \left(\alpha_k \frac{\partial}{\partial u_k} + \alpha_{-k} \frac{\partial}{\partial u_{-k}} \right) + \sum_{k=1}^{\infty} \left(\beta_k \frac{\partial}{\partial v_k} + \beta_{-k} \frac{\partial}{\partial v_{-k}} \right)$$

satisfies the equality $DKD^{-1} = hK$, where h is a function depending on shifts and derivatives of variables u and v , then $K = 0$.

To apply this lemma we evaluate how the transformation $D(*)D^{-1}$ acts on vector fields. For vector fields A, B, P_1 and P_2 we have

$$DAD^{-1} = A, \quad DBD^{-1} = B \quad (6)$$

and

$$DP_1D^{-1} = P_1 - M_1A, \quad DP_2D^{-1} = P_2 - N_1B \quad (7)$$

Using (6) and (7) we calculate action of the transformation $D(*)D^{-1}$ on the vector fields T_n and R_n .

First we consider vector fields $T_n, n = 1, 2, \dots$. We find

$$\begin{aligned} DT_1D^{-1} &= D[P_1, P_2]D^{-1} = [DP_1D^{-1}, DP_2D^{-1}] = [P_1 - M_1A, P_2 - N_1B] \\ &= T_1 + a_{12}N_1P_1 - a_{21}M_1P_2 + a_{21}M_1N_1B. \end{aligned} \quad (8)$$

In the same way as for T_1 we find

$$\begin{aligned} DT_2D^{-1} &= T_2 - (2a_{21} + a_{11})M_1T_1 - a_{12}(a_{11} + 2a_{21})P_1 + a_{21}(a_{11} + a_{21} - 1)M_1^2P_2 + \\ &\quad a_{21}(1 - a_{11} - a_{21})M_1^2N_1B \end{aligned} \quad (9)$$

For other vector fields T_n we have following lemma.

Lemma 4. *For the vector fields $T_n, n=3,4,\dots$, we have*

$$DT_nD^{-1} = T_n + \alpha_n M_1 T_{n-1} + \beta_n M_1^2 T_{n-2} + \dots, \quad (10)$$

where

$$\alpha_n = -\frac{n(n-1)}{2}a_{11} - na_{21}, \quad (11)$$

$$\begin{aligned} \beta_n = \frac{1}{24}n \left(-8a_{11} + 12na_{11} - 4n^2a_{11} - 2a_{11}^2 + 9na_{11}^2 - 10n^2a_{11}^2 + 3n^3a_{11}^2 \right. \\ \left. + 12a_{21} - 12na_{21} + 12a_{11}a_{21} - 24na_{11}a_{21} + 12n^2a_{11}a_{21} - 12a_{21}^2 + 12na_{21}^2 \right). \end{aligned} \quad (12)$$

Proof. For $n = 3$ we have

$$\begin{aligned} DT_3D^{-1} &= [DP_1D^{-1}, DT_2D^{-1}] = \\ &[P_1 - M_1A, T_2 - (2a_{21} + a_{11})T_1 - a_{12}(a_{11} + 2a_{21})P_1 + a_{21}(a_{11} + a_{21} - 1)P_2 + \dots] \\ &= T_3 + \alpha_3T_2 + \beta_3T_1 + \dots \end{aligned} \quad (13)$$

where $\alpha_3 = -3(a_{11} + a_{21})$ and $\beta_3 = (a_{11}(2a_{11} + 6a_{21} - 1) + a_{21}(3a_{21} - 3))$. Thus, α_3 satisfies (11), β_3 satisfies (12).

For $n \geq 4$, we can show, using induction, that the coefficients α_n and β_n satisfy recursion relations

$$\alpha_n = \alpha_{n-1} - (na_{11} + a_{21}), \quad (14)$$

$$\beta_n = \beta_{n-1} + (1 - na_{11} - a_{21})\alpha_{n-1} \quad (15)$$

Solving the above recursion relations we obtain that α_n satisfies (11) and β_n satisfies (12). \square

Now we consider vector fields R_n , $n = 1, 2, \dots$. We have

$$DR_1D^{-1} = DT_1D^{-1} = R_1 + a_{12}N_1P_1 - a_{21}M_1P_2 + a_{21}M_1N_1B. \quad (16)$$

In the same way as in (8) we find

$$DR_2D^{-1} = R_2 - (2a_{12} + a_{22})N_1R_1 + a_{12}(1 - a_{22} - a_{12})N_1^2P_2 \quad (17)$$

and

$$\begin{aligned} DR_3D^{-1} &= R_3 - (3a_{12} + 3a_{22})N_1R_2 + (a_{12} + 2a_{22} - 1)(2a_{12} + a_{22})N_1^2R_1 \\ &+ a_{12}(2 - 3a_{22})(1 - a_{12} - a_{22})N_1^3P_2 + a_{12}(1 - a_{12} - a_{22})N_1^4B \end{aligned} \quad (18)$$

For other vector fields R_n we have the following lemma.

Lemma 5. For the vector fields R_n , $n=4,5 \dots$, we have

$$DR_n D^{-1} = R_n + \tilde{\alpha}_n N_1 R_{n-1} + \tilde{\beta}_n N_1^2 R_{n-2} + \tilde{\gamma}_n N_1^3 R_{n-3} + \dots, \quad (19)$$

where

$$\tilde{\alpha}_n = -\frac{n(n-1)}{2}a_{22} - na_{12}, \quad (20)$$

$$\begin{aligned} \tilde{\beta}_n = \frac{1}{24}(-2+n) & (3n^3 a_{22}^2 - 4n^2 a_{22}(1 + a_{22} - 3a_{12}) + 12a_{12}(-1 + a_{22} + a_{12}) \\ & + n(4a_{22} + a_{22}^2 + 12(-1 + a_{12})a_{12})) \end{aligned} \quad (21)$$

$$\begin{aligned} \tilde{\gamma}_n = -\frac{1}{48}(-3+n)(-2+n) & (n^4 a_{22}^3 - 2n^3 a_{22}^2(2 + a_{22} - 3a_{12}) \\ & + 8a_{12}(4 - 5a_{22} + a_{22}^2 - 6a_{12} + 3a_{22}a_{12} + 2a_{12}^2) \\ & + n^2 a_{22}(4 + 6a_{22} + a_{22}^2 - 20a_{12} - 2a_{22}a_{12} + 12a_{12}^2) \\ & + 2n(a_{22}^2(-1 + 6a_{12}) + 4a_{12}(2 - 3a_{12} + a_{12}^2) + a_{22}(-2 - 6a_{12} + 6a_{12}^2))). \end{aligned} \quad (22)$$

Proof. By direct calculations for $n = 4$ we have

$$DR_4 D^{-1} = R_4 + \tilde{\alpha}_4 R_3 + \tilde{\beta}_4 R_2 + \tilde{\gamma}_4 R_1 + \dots \quad (23)$$

where $\tilde{\alpha}_4 = -(6a_{22} + 4a_{12})$, $\tilde{\beta}_4 = (11a_{22}^2 + 5(-1 + a_{12})a_{12} + a_{22}(-4 + 17a_{12}))$ and $\tilde{\gamma}_4 = -(-1 + 2a_{22} + a_{12})(-2 + 3a_{22} + a_{12})(a_{22} + 2a_{12})$. It is easy to check that $\tilde{\alpha}_4$, $\tilde{\beta}_4$ and $\tilde{\gamma}_4$ satisfy equalities (20), (21) and (22). For $n \geq 5$ we can show, using induction, that the coefficients $\tilde{\alpha}_n$, $\tilde{\beta}_n$ and $\tilde{\gamma}_n$ satisfy recursion relations

$$\tilde{\alpha}_{n+1} = \tilde{\alpha}_n - (na_{22} + a_{12}), \quad (24)$$

$$\tilde{\beta}_{n+1} = \tilde{\beta}_n + (1 - na_{22} - a_{12})\tilde{\alpha}_n, \quad (25)$$

$$\tilde{\gamma}_{n+1} = \tilde{\gamma}_n + (2 - na_{22} - a_{12})\tilde{\beta}_n. \quad (26)$$

Solving these recursion relations we obtain equalities (20), (21) and (22). \square

The previous two lemmas allow us to specify matrix $A = (a_{ij})$ in the case of finite characteristic x -ring.

Lemma 6. *Let \mathcal{A} be a finite dimensional ring then for some integer $n \geq 2$ we have $T_n = 0$ and $T_{n-1} \neq 0$. In that case*

$$a_{11} = 2, \quad a_{12} = -1, \quad (27)$$

$$a_{21} = n - 2, \quad a_{22} = 2. \quad (28)$$

Proof. First we show that $a_{11} = 2$ and $a_{21} = n - 2$. If $n = 2$ we have $T_2 = 0$ and $DT_2D^{-1} = 0$. So coefficients of T_1 and P_2 in (9) must be zero. Thus

$$2a_{11} + a_{21} = 0, \quad (29)$$

$$a_{11} + a_{21} - 1 = 0. \quad (30)$$

Solving these equalities for a_{11} and a_{12} we get $a_{11} = 2$ and $a_{21} = n - 2$. In the same way for $n > 2$, we have $T_n = 0$ and $DT_nD^{-1} = 0$. Hence coefficients α_n and β_n in the equality (10) must be zero. Using equalities (11) and (12) we have

$$-\frac{n(n-1)}{2}a_{11} - na_{21} = 0,$$

$$\begin{aligned} \beta_n = \frac{1}{24}n \left(-8a_{11} + 12na_{11} - 4n^2a_{11} - 2a_{11}^2 + 9na_{11}^2 - 10n^2a_{11}^2 + 3n^3a_{11}^2 \right. \\ \left. + 12a_{21} - 12na_{21} + 12a_{11}a_{21} - 24na_{11}a_{21} + 12n^2a_{11}a_{21} - 12a_{21}^2 + 12na_{21}^2 \right) = 0. \end{aligned} \quad (31)$$

Solving these equalities for a_{11} and a_{21} we get $a_{11} = 2$ and $a_{21} = n - 2$.

Now let us show that $a_{22} = 2$ and $a_{12} = -1$. Since \mathcal{A} is finite dimensional there exists $m \geq 2$ such that $R_m = 0$. It turns out that $m = 2$.

Indeed, assume $m > 2$. If $m = 3$ then we have $R_3 = 0$. We have $DR_3D^{-1} = 0$. So coefficients of R_1 , R_2 and P_2 in (18) must be zero. Equating the coefficients to zero we get overdetermined system

$$\begin{aligned} 3a_{12} + 3a_{22} &= 0, \\ (a_{12} + 2a_{22} - 1)(2a_{12} + a_{22}) &= 0, \\ (2 - 3a_{22})(1 - a_{12} - a_{22}) &= 0 \end{aligned}$$

that has no solution. Thus it is not possible to have $m = 3$.

In the same way if $m > 3$, we have $R_m = 0$ and $DR_mD^{-1} = 0$. Hence coefficients $\tilde{\alpha}_m$, $\tilde{\beta}_m$ and $\tilde{\gamma}_m$ in the equality (19) must be zero. Using equalities

(20), (21) and (22) for $\tilde{\alpha}_m$, $\tilde{\beta}_m$ and $\tilde{\gamma}_m$ respectively we get an overdetermined system

$$\tilde{\alpha}_m = 0, \quad \tilde{\beta}_m = 0, \quad \tilde{\gamma}_m = 0 \quad (32)$$

that has no solution. Thus it is not possible to have $m > 3$.

If $m = 2$ then we have $R_2 = 0$ and $DR_2D^{-1} = 0$. So coefficients of R_1 and P_2 in (17) must be zero. Thus

$$2a_{12} + a_{22} = 0, \quad (33)$$

$$1 - a_{22} - a_{12} = 0. \quad (34)$$

Solving these equalities for a_{11} and a_{12} we get $a_{11} = 2$ and $a_{21} = n - 2$. \square

In the proof of previous lemma we see that in case of finite dimensional characteristic x -ring all vector fields $R_n = 0$, $n \geq 2$. The similar statement is true for vector fields T_n , $n \geq 4$. To show that we have the following lemma.

Lemma 7. *Let \mathcal{A} be a finite dimensional ring. Then $T_n = 0$, $n \geq 4$.*

Proof. Let us introduce sequences

$$W_n = [P_2, W_{n-1}], \quad n = 4, 5 \dots \quad \text{with} \quad W_3 = T_3,$$

and

$$V_n = [P_2, V_{n-1}], \quad n = 5, 6 \dots \quad \text{with} \quad V_4 = T_4.$$

First we show that $W_n = 0$, for $n \geq 5$. We have

$$[P_2, T_2] = [P_2, [P_1, T_1]] = -([T_1, [P_2, P_1]] + [P_1, [T_1, P_2]]) = [T_1, T_1] + [P_1, R_2] = 0,$$

since $R_2 = 0$ in the case of the finite dimensional ring \mathcal{A} . Taking commutator of P_1 with $[P_2, T_2]$ we get

$$[P_1, [P_2, T_2]] = -([T_2, [P_1, P_2]] + [P_2, [T_2, P_1]]) = 0,$$

which gives $W_4 = -[T_1, T_2]$. So, for W_5 we have

$$W_5 = [P_2, -[T_1, T_2]] = [T_1, [T_2, P_2]] + [T_2, [P_2, T_1]] = -[T_1, [P_2, T_2]] + [T_2, R_2] = 0.$$

This shows that $W_5 = 0$ and $W_n = 0$ for $n > 5$.

Now we show that the assumption $T_4 \neq 0$ leads to a contradiction. Assume $T_4 \neq 0$. Since \mathcal{A} is finite dimensional we have $V_n = 0$ for some $n \geq 5$. Assume $V_5 = 0$. Then considering

$$DV_5D^{-1} = V_5 - (4a_{12} + a_{22})N_1T_4 + \dots,$$

we have $(4a_{12} + a_{22}) = 0$ and by Lemma 6 we have $a_{12} = -1$ and $a_{22} = 2$, which is not possible. Assume $V_n = 0$, for $n > 5$. Then evaluating DV_nD^{-1} , using induction, we get

$$DV_nD^{-1} = V_n + \frac{1}{2}[(42 - 8n)a_{22} + (7n - n^2 - 22)a_{12}]N_1V_{n-1} + \dots$$

We have $[(42 - 8n)a_{22} + (7n - n^2 - 22)a_{12}] = 0$ and by Lemma 6 we have $a_{12} = -1$ and $a_{22} = 2$, which is not possible. \square

Proof of Theorem 2. Since \mathcal{A} is finite dimensional, using Lemma 6 and Lemma 7 we find that

$$\begin{aligned} a_{11} &= 2, & a_{12} &= -1, \\ a_{21} &= -c, & a_{22} &= 2, \end{aligned} \tag{35}$$

where $c = 1, 2, 3$. For each value of c , A is the Cartan matrix and the corresponding x - and n - integrals are described in [1]. Hence that completes the proof of the main theorem.

References

- [1] Habibullin I., Zheltukhin K. and Yangubaeva M. *Cartan matrices and integrable lattice Toda field equations* J. Phys. A **44**, 465202 (2011)
- [2] Darboux G. *Leconsur la theory generale des surface et les application-geometriques du calculus infinitesimal* vol. **2** (Paris: Gautier Villas) (1915)
- [3] Shabat A. B., Yamilov R. I. *Exponential systems of type I and Cartan matrices* (Russian) *Preprint* BBAS USSR Ufa (1981)
- [4] Leznov A.N., Smirnov V.G. and Shabat A.B. *Internal symmetry group and integrability conditions for two-dimensional dynamical systems* (Russian) *Teoret. Mat. Fiz.* **51**, 10-21 (1982)

- [5] Sokolov V.V., Zhiber A.V. *On the Darboux integrable hyperbolic equations* Phys. Lett. A **208**, 303-308 (1995)
- [6] Zhiber A.V., Sokolov V.V. and Startsev S.Ya. *On nonlinear Darboux-integrable hyperbolic equations*, (Russian) Dokl. Akad. Nauk **343**, 746-748 (1995)
- [7] Zhiber A. V., Sokolov V. V. *Exactly integrable hyperbolic equations of Liouville type*, Russian Math. Surveys **56**, 61-101 (2001)
- [8] Zhiber A.V., Murtazina R.D. *On the characteristic Lie algebras for the equations $u_{xy} = f(u, u_x)$* , J. Math. Sci. (N. Y.) **151**, 3112-3122 (2008)
- [9] Murtazina R. D. *Nonlinear hyperbolic equations with characteristic ring of dimension 3*, (Russian) Ufa Math. J. **3**, 113-118 (2011)
- [10] Kostrigina O.S., Zhiber A.V. *Darboux-integrable two-component nonlinear hyperbolic systems of equations*, J. Math. Phys. **52**, 033503 (2011)
- [11] Zhiber A.B., Murtazina R.D., Habibullin I.T. and Shabat A.B. *Characteristic Lie rings and integrable models in mathematical physics*, Ufa Math. J. **4**, 17-85 (2012)
- [12] Adler V. E., Startsev S.Ya. *On discrete analogues of the Liouville equation*, Theoret. and Math. Phys. **121**, 1484-1495 (1999)
- [13] Habibullin I.T. *Characteristic algebras of fully discrete hyperbolic type equations*, SIGMA Symmetry Integrability Geom.: Methods Appl. **1**, 9 (2005)
- [14] Habibullin I.T., Pekcan A. *Characteristic Lie algebra and the classification of semi-discrete models*, Theoret. and Math. Phys. **151**, 781-790 (2007)
- [15] Habibullin I.T. *Characteristic algebras of discrete equations. In: Difference equations, special functions and orthogonal polynomials*, Hackensack NJ World Sci. Publ., 249-257 (2007)
- [16] Habibullin I.T. and Gudkova E.V. *Classification of integrable discrete Klein-Gordon models*, Physica Scripta bf 81, 045003 (2011)

- [17] Habibullin I.T., Zheltukhina N. and Pekcan A. *On some algebraic properties of semi-discrete hyperbolic type equations*, Turkish J. Math. **32**, 277-292 (2008)
- [18] Habibullin I.T., Zheltukhina N. and Pekcan A. *On the classification of Darboux integrable chains*, J. Math. Phys. **49**, 102702 (2008)
- [19] Habibullin I.T., Zheltukhina N. and Pekcan A. *Complete list of Darboux integrable chains of the form $t_{1x} = t_x + d(t, t_1)$* , J. Math. Phys. **50**, 102710 (2009)
- [20] Habibullin I.T., Zheltukhina N. and Sakieva A. *On Darboux-integrable semi-discrete chain*, J. Phys. A **43**, 434017 (2010)
- [21] Habibullin I.T. and Zheltukhina N. *Discretization of Liouville type nonautonomous equations*, J. Nonlinear Math. Phys. **23**, 620-642 (2016)
- [22] Kuniba A. Nakanishi T. and Suzuki J. *T-systems and Y-systems in integrable systems*, J. Phys., A **44** no. 10, 103001 (2011).