# Discovery of 36 eclipsing EL CVn binaries found by the Palomar Transient Factory

J. van Roestel<sup>1\*</sup>, T. Kupfer<sup>2</sup>, R. Ruiz-Carmona<sup>1</sup>, P.J. Groot<sup>1</sup>, T.A. Prince<sup>2</sup>, K. Burdge<sup>2</sup>, R. Laher<sup>3</sup>, D.L. Shupe<sup>4</sup>, E. Bellm<sup>5</sup>

- Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O.Box 9010, 6500 GL, Nijmegen, NL
- <sup>2</sup>Cahill Center for Astronomy and Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA
- <sup>3</sup>Spitzer Science Center, California Institute of Technology, Pasadena, CA 91125, USA
- <sup>4</sup>Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, USA
- <sup>5</sup>Department of Astronomy, University of Washington, Box 351580, Seattle, WA 98195, USA

Accepted XXX. Received YYY; in original form ZZZ

#### ABSTRACT

We report the discovery and analysis of 36 new eclipsing EL CVn-type binaries, consisting of a core helium-composition pre-white dwarf and an early-type main-sequence companion, more than doubling the known population of these systems. We have used supervised machine learning methods to search 0.8 million lightcurves from the Palomar Transient Factory, combined with SDSS, Pan-STARRS and 2MASS colours. The new systems range in orbital periods from 0.46 to 3.8 d and in apparent brightness from ~14 to 16 mag in the PTF R or  $g^\prime$  filters. For twelve of the systems, we obtained radial velocity curves with the Intermediate Dispersion Spectrograph at the Isaac Newton Telescope. We modelled the lightcurves, radial velocity curves and spectral energy distributions to determine the system parameters. The radii  $(0.3–0.7\,R_\odot)$  and effective temperatures  $(8000–17000\,\mathrm{K})$  of the pre-He-WDs are consistent with stellar evolution models, but the masses  $(0.12–0.28\,\mathrm{M}_\odot)$  show more variance than models have predicted. This study shows that using machine learning techniques on large synoptic survey data is a powerful way to discover substantial samples of binary systems in short-lived evolutionary stages.

 $\mathbf{Key}$  words: binaries: close – binaries: eclipsing – white dwarfs – stars: individual:  $\mathrm{EL}\ \mathrm{CVn}$ 

#### 1 INTRODUCTION

EL CVn binaries are eclipsing binaries containing a low mass ( $\sim 0.15-0.33\,\mathrm{M}_{\odot}$ ) pre-helium white dwarf (pre-He-WD) and an A/F-type main sequence star. The prototype system, EL CVn, is part of a sample of 17 EL CVn systems (Maxted et al. 2014a) discovered by SWASP (Pollacco et al. 2006) with magnitudes in the range of 9<V<13. All lightcurves show "boxy", shallow eclipses ( $\lesssim 0.1\,\mathrm{mag}$  depth) with periods between  $\sim 0.5\mathrm{d}$  and  $\sim 3\,\mathrm{d}$ , and in most cases ellipsoidal variation due to the deformation of the A/F-star. The low radial velocity amplitudes ( $\sim 15-30\,\mathrm{km\,s}^{-1}$ ) of the primaries confirm the low mass nature of the pre-He-WDs.

A total of 10 EL CVn systems were found in the Kepler survey: KOI-74 (van Kerkwijk et al. 2010; Bloemen et al. 2012); KOI-81 (van Kerkwijk et al. 2010; Matson et al. 2015); KOI-1375 (Carter et al. 2011); KOI-1224 (Breton et al. 2012); KIC-9164561, KIC-10727668 (Rappaport et al. 2015);

KIC-4169521, KOI-3818, KIC-2851474, and KIC-9285587 (Faigler et al. 2015). All these systems were studied in great detail, and by modelling the Kepler lightcurves in combination with radial velocity curves, all system parameters have been determined. Four of these systems contain small pre-He-WDs (<  $0.05\,R_\odot$ ) and as a consequence their lightcurves feature shallow eclipses only detectable from space. The fact that 10 EL CVn-like systems are found in the Kepler field suggests that there should be many more in our Galaxy, in line with an estimate of the local space density from stellar evolution and population synthesis models,  $4-10\times10^{-6}\,\mathrm{pc}^{-3}$  (Chen et al. 2017).

Besides the samples found by Kepler and SWASP, there were serendipitous discoveries of binaries related to EL CVn systems. The star V209 in  $\omega$  Cen is likely an EL CVn binary (Kaluzny et al. 2007), but the primary does not seem to be a typical main sequence star: its mass is  $0.95 \rm M_{\odot}$  but it has a temperature of 9370 K. OGLE–BLG–RRLYR–02792 is an eclipsing binary which contains a large pre-He-WD which seems to be pulsating like an RR-Lyrae star (Pietrzyński

<sup>\*</sup> j.vanroestel@astro.ru.nl

et al. 2012). A possible non-eclipsing variant of an EL CVn binary is the star Regulus ( $\alpha$  Leo). Gies et al. (2008) and Rappaport et al. (2009) found that Regulus A is a singlelined spectroscopic binary with a period of 40 d, consisting of an A-type primary and a companion with a mass of  $>0.3\,\mathrm{M}_{\odot}$ , at the upper end of the pre-He-WDs mass range.

EL CVn binaries share many characteristics with a new type of binary: "R CMa"-type binaries are Algol binaries with a bloated, low mass, donor (e.g. Budding & Butland 2011; Lee et al. 2016). They are very similar to EL CVn systems, except that they are semi-detached, and therefore considered the progenitors of EL CVn systems. Two "detached R CMa" systems have been identified using Kepler photometry and are now considered to be newly born EL CVn binaries (KIC-10661783; Lehmann et al. 2013, and KIC-8262223; Guo et al. 2017).

EL CVn systems are part of a larger family of binaries where one component of the binary is an extremely low mass white dwarf (ELMWD). The majority of ELMWDcontaining binaries without a main-sequence companion have white dwarf or neutron star companions instead (e.g. Marsh et al. 1995; van Kerkwijk et al. 2005). In these systems, the ELMWD dominates the luminosity, making them identifiable with a single spectrum. The ELM survey (Brown et al. 2010) uses this approach and has been successful in finding many ELMWDs in binary white dwarf systems.

In this paper, we present system parameters for 36 new EL CVn systems, all eclipsing, discovered using the Palomar Transient Factory (PTF). In Section 2 we describe the identification of the systems using supervised machine learning classifiers. In Section 3 we discuss the spectroscopic followup of 12 of the new systems. In Section 4 we discuss the analysis of the lightcurves, spectra and spectral energy distributions, and we present the results in Section 5. In Section 6 we compare our results with theoretical prediction and compare our sample with already known EL CVn binaries. We end with a summary and conclusion in Section 7.

#### TARGET SELECTION

#### The Palomar Transient Factory

The Palomar Transient Facility (PTF) used the 1.2 m Oschin Telescope at Palomar Observatory with a mosaic camera consisting of 11 CCDs. The CCDs have 4Kx2K pixels and the camera has a pixel scale of 1.02"/pixel, giving it a total field of view of 7.26 square degrees. PTF uses an automated image processing pipeline which does bias and flatfield corrections, source finding and photometry. All data is automatically processed, see Rau et al. (2009); Law et al. (2009) for further information.

#### 2.2Data

For all objects detected by PTF, lightcurves are automatically generated (see Laher et al. 2014) and lightcurve statistics are calculated. These statistics include, among others, the mean, root-mean-square, percentiles,  $\chi^2$ -statistic, see Masci & Bellm (2016) for a full list. These lightcurve statistics are based on the lightcurve features used in Richards et al. (2011, 2012), that are useful to distinguish different

Table 1. The number of objects after our initial selection with PTF lightcurves (>40 epochs,  $\chi^2_{\rm reduced}$  > 10, <16 mag). The percentage for which additional colour information is available is shown in the table below.

Filter	# objects	$_{ugriz}^{\mathrm{SDSS}}$	NOMAD JHK	$\begin{array}{c} \text{Pan-STARRS} \\ grizy \end{array}$
R	532 477	43.65%	97.58%	98.92%
g'	257 918	55.45%	98.69%	99.26%
$R \cap g'$	36 943	64.39%	96.48%	98.66%

types of variable stars. Important to note is that we do not use features related to any periodicity in the lightcurve. This has a practical reason; it is very difficult to automatically obtain a reliable period for all the PTF lightcurves because they are sparsely sampled and span many years.

For this study, we used all available lightcurve data that was obtained between the start of PTF in December 2008 and March 2016. We treat the data for the R and g' filter as two separate datasets in the subsequent analysis. These datasets are very substantial (R:  $\sim$ 250 million, g':  $\sim$ 50 million objects). We make an initial cut and select only objects which are variable by requiring that  $\chi^2_{\rm reduced} > 10$ , that lightcurves have more than 40 epochs, and that objects are brighter than 16 mag in either PTF R or PTF g'. This still leaves more than  $\sim 10^5$  candidates (see Table 1 and Fig. 1).

We match the objects in these datasets to the latest SDSS catalogue (ugriz bands, DR13, SDSS Collaboration et al. 2016), the NOMAD catalogue (JHK bands, Zacharias et al. 2004), and the Pan-STARRS catalogue (grizy bands, Chambers et al. 2016). An overview of the total number of objects and the colour availability are given in Table 1.

#### 2.3 Machine Learning Classification

To cut back on the number of candidates for an initial visual lightcurve inspection, we use supervised machine learning classifiers to make a pre-selection. The idea is that instead of finding EL CVn binaries by using fixed, pre-defined, usersupplied selection criteria, a sample of known EL CVn binaries and not-EL CVn objects (a 'training set') is provided and a machine learning code ('classifier') decides what the best way is to separate the two groups given the characteristics (called 'features', e.g. g - r colour or the lightcurve's root-mean-square value). There are many different types of classifiers, and the behaviour of each classifier can be adjusted by changing so-called hyperparameters. Setting the correct hyperparameters is required to avoid over- or underfitting of the data. For an introduction to machine learning in astronomy, see Ivezić et al. (2014), and for a practical guide to machine learning (with Python) see Andreas C. Müller (2016).

Because supervised machine learning classifiers can process huge amounts of data very quickly, they have become a popular tool to handle with the large amount of lightcurves produced by survey telescopes. Many different techniques have been tried for lightcurve classification (e.g. Debosscher 2009; Palaversa et al. 2013; Angeloni et al. 2014; Peters et al. 2015; Mackenzie et al. 2016; Armstrong et al. 2016; Sesar et al. 2017). In recent years, the Random Forest method (Breiman 2001) has become very popular as it typically per-




Figure 1. All objects in the PTF sample after our initial cut (>40 epochs,  $\chi^2_{\text{reduced}}$ > 10, <16 mag), red for PTF R, green for PTF g'. The EL CVn binaries we discovered in the data are shown as white dots. The black lines show Galactic latitudes of -15°, 0°, and 15°.

forms the best and is also easy to interpret (e.g. Richards et al. 2011; Masci et al. 2014).

To find EL CVn binaries we have experimented with three different supervised machine learning classifiers based on combining decision trees: the standard 'Random Forest', an 'Extra-Trees' classifier (Geurts et al. 2006), both implemented in the PYTHON package SKLEARN (Pedregosa et al. 2011), and the more sophisticated 'Gradient boosted decision tree' classifier, implemented in XGBOOST (Chen & Guestrin 2016). All three classifiers combine many randomised decision trees, which are a sequence of binary decisions.

Here we briefly discuss the differences between the methods. Both Random Forest and Extra-Trees combine the prediction of many independent, randomised decision trees. The larger the number of trees the better but at the cost of increased computation time. For both methods, each tree is built using only a subset of all features (rule-of-thumb is the square root of the total number of features). Random Forest uses the best possible split of the data given the available features and uses that to separate the different classes. Extra-Trees differs from Random Forest as it does not use the best split, but a random split. This extra randomization step has the consequence that decision boundaries are more smooth compared to Random Forest. Both methods are relatively simple; they have only a few hyperparameters and are relatively robust against overfitting. XGBOOST also uses many randomised decision trees. But instead of combining many independent trees, new trees are created to optimally complement the existing trees. This is done by giving samples which were wrongly classified by the previous trees a larger weight when building the next tree. The next tree is, therefore, more likely to classify these objects correctly. The disadvantage of this method is that it is more sensitive to overfitting compared to Random Forest. The XGBoost implementation has many hyperparameters which can be set to counteract this, but it can be difficult to determine the best values for these parameters. The advantages of all three methods are that they are insensitive to uninformative features, do not require scaling of the data, and are easy to interpret: they automatically determine the importance of features.

#### 2.4 EL CVn identification

Because supervised machine learning algorithms require a training sample we first need to identify EL CVn binaries in our data. There are no known EL CVn binaries in the PTF magnitude range, so we need to find new ones the oldfashioned way. We do this by selecting a sample of A-type main sequence stars using SDSS colours (0.8  $< u - g_{SDSS} <$ 1.5 and  $-0.5 < g - r_{\rm SDSS} < 0.2$ ) and require that Stetson-K > 0.6 (one of the lightcurve statistics, see also Stetson 1996). To limit the sample size and have increased post-facto confidence in the selected objects, we further require that the light curve is significantly variable ( $\chi^2>40$ ) and with more than 150 epochs in R and 100 in g'. EL CVn binaries are characterized by their  $\lesssim 0.1\,\mathrm{mag}$ , flat-bottomed primary eclipse and slightly shallower secondary eclipse. We, therefore, do a period search using both Analysis-of-Variance and Boxed-Least-Square methods (AoV, Schwarzenberg-Czerny 1989; Devor 2005; BLS, Kovács et al. 2002, VARTOOLS implementation Hartman & Bakos 2016) on each of the lightcurves and inspect each folded lightcurve for these criteria. In case of doubt the candidate was included in the lightcurve modelling (see Section 4.1). If the lightcurve fitting showed a

'V'-shaped, non-total, eclipse, we rejected it from the sample, as these systems could also be regular MS-MS binary. In other words, we require our systems to be totally eclipsing.

Using this method, we found 6 EL CVn binaries, which we then used as a training set for a Random Forest classifier, combined with a sample of 4000 randomly chosen objects (which we confirmed are not EL CVn binaries). Since the training set is so small, we do not attempt any parameter optimisation, but used the default hyperparameters (500 trees, the square-root of the total number of features as the number of features per tree, and no limits on the tree depth). We applied the classifier to the data (the PTF R and PTF g' lightcurve statistics combined with SDSS colours) and inspected the ~100 best candidates identified by the classifiers. We added newly found EL CVn candidates to the training sample and repeated the procedure an additional two times. This resulted in the discovery of an additional 11 systems, bringing the total to 17.

Because we required that SDSS colours were available, we only inspected roughly half of the data so far (see Table 1). We, therefore, replaced the SDSS colours with the BVRHJK colours from the NOMAD catalogue (Pan-STARRS colours were not yet available at this time). We again checked the best 100 candidates in an iterative way, adding the new EL CVn systems to the training sample. The combined SDSS and NOMAD process resulted in a total of 30 EL CVn binaries.

With this sample, we trained the three different classifiers (Random Forest, Extra-Trees and XGBoost) and determined the best hyperparameters. We use the PTF variability features combined with the Pan-STARRS colours. The goal of our classifier is not to classify all samples correctly (high precision), but instead to rank the candidates according to 'EL CVn' likeness. We, therefore, do not optimise the precision of our classifier, but instead we use the 'area-under-curve' for the 'receiver operating characteristic' (roc-auc). We do this using stratified K-fold cross-validation to calculate the roc-auc score. For more details on classifier metrics (like roc-auc) and model optimisation, see Ivezić et al. (2014); Andreas C. Müller (2016).

For both Random Forest and Extra-Trees we find similar optimal hyperparameters. Using more than 600 trees does not improve performance significantly. The number of features per tree influences the roc-auc score marginally, but there is a clear preference for only using two features per tree. We checked different hyperparameters that limit the depth or complexity of the tree, but we find that the roc-auc score only decreases when the tree depth or complexity is limited using any of the hyperparameters.

For XGBoost there are more hyperparameters to tune. We start by optimising the most important three: the number of estimators, the learning rate and the tree depth while setting the other parameters to typical values. After finding the optimal combination of these hyperparameters we continue to optimise the minimum child weight, sub-sampling fraction and the column sub-sample fraction.

After training all classifiers we selected the top 1000 candidates (in both R and g' datasets) from the three classifiers and visually inspected their lightcurves. We found an additional 6 EL CVn binaries, bringing the final number to 36 systems, listed in Table 2.

**Table 2.** Overview of the EL CVn binaries we discovered in the PTF data. In the rest of the paper we will use the PTF name. The "PTF R"-column lists the median magnitude of the light curve in R-band.

PTF name	IAU name (PTF1 J)	P (d)	PTF R
PTFS1600y	J004040.23+412521.61	1.184	13.7
PTFS1600ad	J004300.75+381537.26	1.084	14.7
PTFS1700do	J005424.06+411126.98	3.051	15.7
PTFS1600aa	J005659.72+130920.66	0.693	15.9
PTFS1601p	J011909.91+435907.11	1.222	15.3
PTFS1501bh	J012814.72+040551.90	0.620	13.9
PTFS1601q	J013336.92+470600.18	1.252	16.2
PTFS1601cl	J014839.10+382314.56	0.892	13.6
PTFS1402de	J021913.15 + 215921.98	0.619	15.0
PTFS1607aa	J071207.01 + 211654.98	0.846	15.0
PTFS1607v	J075310.42+835154.79	0.720	15.3
PTFS1607t	J075642.49 + 162143.99	0.876	14.2
PTFS1607ab	J075950.03+154319.09	0.773	14.0
PTFS1608ab	J080425.26 + 070845.24	0.610	14.6
PTFS1612al	J121254.27+363341.76	0.637	15.7
PTFS1512bf	J124154.58 + 001333.06	0.607	14.2
PTFS1613s	J133220.59 + 352847.28	1.142	14.3
PTFS1613u	J133929.37 + 455055.64	0.564	15.3
PTFS1615ag	J150041.84 - 191417.23	0.681	14.3
PTFS1615v	J150327.61 + 460322.78	0.559	15.9
PTFS1515ay	J150336.10 + 195842.16	0.464	14.8
PTFS1615w	J152726.81 + 120453.54	1.441	14.9
PTFS1615ao	J152758.90 + 190751.63	0.895	15.0
PTFS1615u	J153005.01 + 202157.06	0.778	15.8
PTFS1616cr	J162342.13 + 231456.58	0.565	14.0
PTFS1617n	J173257.98 + 403600.93	2.337	15.3
PTFS1617m	J175433.50 + 230041.83	3.773	14.7
PTFS1619l	J191826.08 + 485302.94	1.160	13.7
PTFS1521ct	J213318.98 + 254126.30	1.172	15.8
PTFS1621ax	J213534.11 + 233313.86	1.018	15.0
PTFS1521cm	J214858.33 + 030417.50	0.685	15.1
PTFS1622by	J220719.56 + 085415.66	0.749	15.8
PTFS1522cc	J225539.41 + 342137.72	0.572	14.7
PTFS1622aa	J225652.53 + 390822.70	0.766	15.6
PTFS1622bt	J225755.64 + 310133.67	0.688	15.1
PTFS1723aj	J231010.08+331249.78	1.109	14.8

A quick comparison between the classifiers shows that both the Extra-Trees classifier and the XGBOOST classifier perform equally well while the Random Forest performs a bit worse. This is confirmed by the ranking of the last six discovered EL CVn binaries that were all further down the list for the Random Forest method. Although Extra-Trees and XGBOOST performance is comparable, tuning the XGBOOST classifier took significantly more time and effort. Due to the combination of yield versus investment, we deem the Extra-Trees classifier the best (in this particular case).

PTF observed the Kepler field and has thus observed the EL CVn binaries found by Kepler. None of these were recovered by our search and we investigated the reason why. First of all, most Kepler systems feature eclipses much shallower than PTF can detect. The Kepler EL CVn systems with deep enough eclipses to be detected by PTF were not recovered because either the star was saturated in the PTF data, or the object was not observed at a sufficient number of epochs.




Figure 2. The left panel shows the weighted root-mean-squared of the PTF lightcurve versus the normalised 95th percentile of the lightcurve (percentile 95 minus median, divided by 90 percentile range, see Table A1), the right panel shows 2MASS J-PTF R versus Pan-STARRS g-r colour-colour space. The red dots show all EL CVn binaries, the black contours show all samples in the PTF R dataset, with the black contours containing 25, 50, and 75 per cent of the data, samples outsides the contours are indicated with black points. The background colour indicates the 'EL CVn'-score by the Extra-Trees classifier with grey lines at every 0.1 score interval. The score is calculated assuming the median values of the EL CVns for all parameters, except the parameters on the x- and y-axis.

#### 3 SPECTROSCOPY

For 19 of our EL CVn systems, we obtained phase-resolved spectroscopy with the Isaac Newton Telescope (INT). We used the Intermediate Dispersion Spectrograph (IDS) with the R632V grating (0.90 Å pixel $^{-1}$ , 3800–5800 Å) for 8 bright nights and the R900V grating for 9 bright nights on 3 separate runs (0.63 Å pixel $^{-1}$ , 4000–5500 Å). Conditions were good with seeing of  $\lesssim 1~\rm arcsec$ , except for the last four nights. During these nights the seeing was 2–5 arcsec and two nights were mostly clouded. An overview of the spectroscopic runs, the set-up and the weather quality is given in Table A2.

Since the orbital period and phase for all systems is determined very precisely from the photometry (see Section 5), we timed the observations such that we observed the systems around orbital phases 0.25 and 0.75. The signal—to—noise per pixel of each spectrum ranges between 40 and 80, sufficient to detect the weaker metal lines in the A/F-star's spectrum. Spectra were taken in pairs and before or after each stellar spectrum a calibration lamp spectrum (CuAr) was obtained to make sure the wavelength calibration was stable.

The data were reduced using IRAF. We used L.A.Cosmic (van Dokkum 2001) to remove cosmic rays and performed the standard bias and flat calibrations. For the wavelength calibration, we used ~40 arc lines, which resulted in a typical root-mean-square uncertainty on the wavelength solution of  $\lesssim\!\!0.1$  pixels (4-6 km s $^{-1}$ ).

#### 4 METHODS AND ANALYSIS

### 4.1 Lightcurve

By modelling the lightcurves we put strong constraints on the system parameters. To construct a model lightcurve given a set of binary star parameters, we use LCURVE (by T.R. Marsh and collaborators, see Copperwheat et al. 2010, see also Copperwheat et al. 2011; Parsons et al. 2011). The LCURVE code uses grids of points to model the two stars.

The shape of the stars in the binary is set by a Roche potential. We assume that the orbit is circular and that the rotation periods of the stars are synchronised to the orbital period. We discuss the validity of the latter assumption in Section 6.1. We calculate the lightcurves assuming the effective wavelength of the PTF filters; 4641 Å for the g' filter and 6516 Å for the R filter. In this section (and the rest of the paper), we refer to the A/F-type main sequence as the primary (subscript '1') and the pre-He-WD as the secondary (subscript '2').

The free parameters of the model are: the orbital period (P) and mid-eclipse time  $(t_0)$ , both in BMJD<sub>TDB</sub> (the barycentric Julian date in the terrestrial dynamic time frame, minus 2400000.5), the effective temperatures of both stars  $(T_{1,2})$ , the scaled radii of both stars  $(r_{1,2} = R_{1,2}/a, \text{ where }$ a is the binary separation), the inclination angle (i), the mass ratio  $(q = M_2/M_1)$ , an albedo (absorption) for both stars, a linear limb darkening coefficient  $(x_{1,2})$ , and a gravity darkening coefficient  $(y_{1,2})$  in the relation  $I \propto g^y$  (where g is the local surface gravity, von Zeipel 1924). Not all these parameters are well constrained by the data and therefore we fix or set an allowed range for some parameters. We constrain the temperature of the primary star  $(T_1)$  to 6500–10000 K; the temperature range of A/F-type main-sequence stars. This is needed because with only a lightcurve the temperature ratio is well constrained, but the absolute values of the temperatures of each star are not. We will not use the resulting temperatures of the lightcurve fit, but instead determine the effective temperatures of both stars by modelling the spectral energy distribution (see Section 4.2). We fix the limb darkening coefficient of star 2  $(x_2)$  to 0.5, since the effect on the lightcurve is minimal. We allow the limb darkening coefficient of the A/F-star  $(x_1)$  to vary between 0.08 to 1.05, the lowest and highest values for stars in the allowed temperature range (Gianninas et al. 2013).

To determine the uncertainty on the parameters we use the Markov Chain Monte Carlo (MCMC) method as implemented by EMCEE (Foreman-Mackey et al. 2013). The standard method to determine the uncertainties on the parameters is by using the least-square  $(\chi^2)$  statistic. However, this assumes that the uncertainty estimates of the data are correct and Gaussian distributed. This is not the case for the PTF lightcurves (as in many observational datasets). Ignoring this problem leads to an underestimate of the uncertainties in the derived parameters, and can in some cases also change the solution. To solve this problem we add additional white noise<sup>1</sup> to our model (see Section 8 in Hogg et al. 2010 and for a simple example see Foreman-Mackey 2013). This means that our model has the white noise amplitude as an extra parameter, which we can simply optimize over, exactly the same as for the lightcurve parameters.

This method requires the following modification to the standard least-square function:

$$\tilde{\chi}^2 = \sum_n \frac{(y_n - m_n(p))^2}{\sigma_n^2 + f^2 m_n(p)^2} + \log(\sigma_n^2 + f^2 m_n(p)^2) \tag{1}$$

where  $y_n$  is the data,  $m_n$  the lightcurve model as a function of the lightcurve parameters p,  $\sigma_n$  the uncertainties, and f a factor which adds an extra noise source. Note that the first term of the equation is almost the same as in a regular least-square  $(\chi^2)$  regression, except for the additional noise term  $[f^2m_n(p)^2]$ . The first term can be minimised by letting f go to infinity, instead of minimising the difference between data and model  $[y_n - m_n(p)]$ . Therefore, the second term is needed to penalise models with a large value of f. Using this equation, the optimal amount of white noise is added to account for any difference between the data and model. To obtain the best model, we simply minimise  $\tilde{\chi}^2$  over the lightcurve parameters p and the parameter f, just like regular least-square regression.

For each of the systems, we first find the approximate solution using a simple simplex minimiser of the modified least-square function. We then use EMCEE to find the best set of parameters of all the available lightcurves for that system. For each filter we use different values for  $x_1$ ,  $y_1$ , and 'absorb', while the rest of the parameters are filter independent. We use 256 parallel MCMC chains (called 'walkers') and use at least 2000 generations or more if needed. Any further calculations are done using the last 2560 positions of the walkers.

#### 4.2 Effective temperature

To determine the temperatures of both components we fit the spectral energy distribution of the target with model spectra, similar as in Maxted et al. (2011). We use data from GALEX (far UV & near UV, Bianchi et al. 2014), SDSS DR13 (ugriz, SDSS Collaboration et al. 2016), Pan-STARRS (grizy, Chambers et al. 2016), 2MASS (HJK, Skrutskie et al. 2006), and WISE (W1 & W2, Wright et al. 2010) for each target (where available). We used as model spectra the

BaSeL3.1 spectral library (Westera et al. 2002). To calculate the flux per band, we convolve the model spectra with each band's response curve.

The overall spectrum is the sum of two model spectra of a given temperature and metallicity, created using bilinear interpolation from the BaSeL library. With only an SED, it is not possible to measure the metallicity of the stars reliably. However, since metallicity and temperature are correlated, we treat the metallicity of both stars as free parameters and marginalise over them in the final result. For the surface gravity, we assume  $\log g=4$  for the A/F-star and  $\log g=5$ for the pre-He-WD. We set the relative contribution to the total light by the ratio between  $r_1$  and  $r_2$  obtained from the lightcurve. At first, we also used the temperature ratio obtained from the lightcurve, but we learnt that this gave inconsistent predictions for the eclipse depths. This is likely due to the use of blackbody spectra by LCURVE. Instead, we directly use the eclipse depth of the primary eclipse instead of the temperature ratio. The final variable is the extinction, set by E(B-V). To calculate the reddening following from the extinction we used the reddening law by Cardelli et al. (1989) with  $R_V = 3.1$  (as implemented by PYSYNPHOT).

To determine the temperatures of both stars we minimized the function:

$$\tilde{\chi}^2 = \sum_n \frac{(y_n - m_n(p))^2}{\sigma_n^2 + f^2 m_n(p)^2} + \log(\sigma_n^2 + f^2 m_n(p)^2)$$
+ prior( $r_1/r_2$ , E<sub>R-V</sub>) (2)

with y the data, m the model, and f an additional noise factor. We used a value for E(B-V) according to Schlafly & Finkbeiner (2011), with an uncertainty of 0.034 (as in Maxted et al. 2011). For some added flexibility in our model, we added an extra term of uncertainty to the magnitudes (f), similar to the way it was applied in Equation 1. We again use EMCEE to determine the best values and uncertainties, similar as in Section 4.1.

#### 4.3 Radial velocity

To obtain the radial velocity curve of the primary star, we cannot use the Balmer absorption lines in the spectrum because these are present both in the A/F-star and the pre-He-WD. Using these would not yield reliable results. Instead, we use the metal lines present in the spectra of the A/Ftype stars. We cross-correlate the spectra with a template; a high-resolution spectrum of the A5 star HD145689 (Bagnulo et al. 2003). We first interpolate the target spectrum to the (much higher) sampling of HD145689. We then remove the continuum with a low-order polynomial and determine the radial velocity shift using cross-correlation. To estimate the uncertainty on the radial velocity shift, we add random Gaussian noise to the target spectra according to the uncertainty per pixel and measure the radial velocity shift. We repeat this process 11 times and use the standard deviation of the results as the uncertainty. We use the metal lines between 4150–4301 Å, 4411–4791 Å, and 4941–5400 Å to get three separate measurements of the radial velocity shift. The radial velocity measurements are corrected to the heliocentric velocity frame with the RVCORRECT task in IRAF.

To determine the radial velocity amplitude, we fit a sinusoidal curve with a fixed value for the period and phase to

<sup>&</sup>lt;sup>1</sup> If the noise cannot be treated as white noise, but the noise is correlated (red noise), Gaussian process regression can be used. See for a simple example Foreman-Mackey (2014) and an example of this method used to model flickering in a cataclysmic variable by McAllister et al. (2017).

the measurements. This leaves only the amplitude and systemic velocity as free parameters. We again use a modified least-squares objective function which can also take into account underestimated uncertainties (similar to Equation 1):

$$\tilde{\chi}^2 = \sum_n \frac{(y_n - m_n(p))^2}{\sigma_n^2 + f^2} + \log(\sigma_n^2 + f^2)$$
 (3)

with  $y_n$  the radial velocity measurements,  $\sigma$  the statistical uncertainty on the cross-correlation results and f the extra white noise. Fitting the data shows (Table 3) that f ranges between 7 and  $12 \,\mathrm{km}\,\mathrm{s}^{-1}$ , a factor of 2 higher than the statistical noise  $\sigma$ . This extra noise is partially due to the uncertainties in the wavelength calibration (4–6 km s<sup>-1</sup>) but does not account for all residual variance. This means that we either underestimate the uncertainties in the cross-correlation procedure (for example by the normalisation of the spectra) or that we underestimate the uncertainties in the calibration process of the spectra. This could be due to instabilities of the optical elements in the IDS/INT combination, which typically changes for each observing run. Since we combine data from four different observing runs, this could result in minor differences in the setup. A potential method to verify this is to check the wavelength of sky emission lines, but these are not available in the spectral range we use.

#### 4.4 Galactic kinematics

For the 12 systems for which we have obtained a radial velocity measurement, we calculate their Galactic location and velocity. We determine the distance to the systems by using the K-band magnitude and absolute radius, combined with the K-band surface brightness calibration by Kervella & Fouqué (2008). The proper motions of the systems are taken from the UCAC5 catalogue (Zacharias et al. 2017) or the Initial Gaia Source List (Smart & Nicastro 2013). Combined with right ascension and declination, we calculate velocity in the direction of the Galactic Centre  $(V_{\rho})$  and the Galactic rotation direction  $(V_{\phi})$ , the Galactic orbital eccentricity (e), and the angular momentum in the Galactic z direction  $(J_z)$ . The Galactic radial velocity  $V_Q$  is negative towards the Galactic centre, while stars that are revolving on retrograde orbits around the Galactic Centre have negative  $V_{\phi}$ . Stars on retrograde orbits have positive  $J_{7}$ . Thin disk stars generally have very low eccentricities e. Population membership can be derived from the position in the  $V_0$ -  $V_{\phi}$  diagram and the  $J_z$  - e diagram (Pauli et al. 2006).

#### 4.5 Masses and radii

To fully solve for the elements of the binary system, we need to combine the information from the lightcurve fit with an additional piece of information to set the scale of the system. This is typically done by measuring the radial velocity amplitude of both stars. We only have the radial velocity amplitude of one of the stars in the binary. In principle, we can combine this with the mass-ratio q, but the uncertainties on the mass-ratio derived from the lightcurve fitting are high, and the uncertainty on the masses scale with a high power of q (for low q:  $M_1 \propto K_1^2 q^{-3}$ ,  $M_2 \propto K_1^3 q^{-2}$ ), and are therefore not constraining.

To circumvent this problem, we make use of the assumption that the primary star is a main-sequence star. Using only the lightcurve parameters, we can calculate the average density of the main-sequence component:

$$\langle \rho \rangle = \frac{3\pi}{GP^2 r_1^3 (1+q)} \tag{4}$$

To propagate the uncertainties correctly, we calculate the stellar density for each point in the MCMC chain and assign a random temperature according to our measurement of the SED. With the average density and temperature of the main-sequence star, we can use stellar models to determine its mass. We use the Yale-Potsdam stellar models (Spada et al. 2017) and follow the same procedure as in Breton et al. (2012) to make a continues mapping of the mass in  $T - \langle \rho \rangle$  space. We convolve each track with a Gaussian probability function with a standard deviation of 200K in temperature and 0.1 dex in density. For each point in the temperature-density grid, we assign the mass with the highest probability. We use this mapping to calculate the primary mass for the posterior distribution of the lightcurve fits (see Section 4.1). As can be seen in Fig. 4, most but not all measurements agree with the models. Two systems, PTFS1612al and PTFS1615u, have slightly higher densities than would be the case for a solar metallicity composition for zero-age main sequence models. For these two systems, we extrapolate the models to determine the mass.

With the mass of the primary  $(M_1)$  combined with q, i and P, we calculate the semi-major axis (a) using Kepler's law

$$a^{3} = GM_{1}(1+q) \left(\frac{2\pi}{P}\right)^{2} \tag{5}$$

Note that in both equations the mass ratio is present in the form of 1+q, and since the mass ratio is small  $(q \sim 0.1)$ , the high uncertainty on q only mildly affects the accuracy on a and  $M_1$ . However, the uncertainty on the pre-He-WDs mass  $(M_2 = qM_1)$  is proportional to the uncertainty on q, which means that the uncertainty on  $M_2$  is too high to be constraining.

This can be solved by including the measured radial velocity  $(K_1)$  in our calculation, which is available for 12 systems. We use an iterative approach to find the optimal solution as in Rappaport et al. (2015), again for each sample from the lightcurve fit posterior distribution. This involves calculating  $M_1$ , q and  $\langle \rho \rangle$  until the solution converges, which it does after 2 iterations.

#### 5 RESULTS

For the 36 EL CVn binaries in the PTF data, we fit the lightcurves with a binary star model, see Fig. A1 and Table A4 in the Appendix. The best model fits to the lightcurves all show a flat-bottomed primary eclipse and a round-bottom secondary eclipse. The orbital period of the binary, the radii of both stars, and the orbital inclination are typically well constrained, but the uncertainty on the massratios of the systems is typically  $\gtrsim 10\,\mathrm{per}$  cent. The extra noise term in the fit for the lightcurves is typically  $\lesssim 1\,\mathrm{per}$  cent. This is consistent with the expected uncertainty in the absolute photometric calibration which is not part of the

Table 3. Radial velocity amplitude of the A/F-star, the systemic velocity, the residual variance of the fit, the derived distance, the measured proper motions and the associated stellar population for the 12 EL CVn systems with radial velocity curves. The proper motion is taken from either the UCAC5 catalogue (a, Zacharias et al. 2017) or or the Initial Gaia Source List (b, Smart & Nicastro

ID	$(\text{km s}^{-1})$	$(\text{km s}^{-1})$	$(\mathrm{km}\mathrm{s}^{-1})$	Distance (pc)	$\mu_{\alpha}\cos(\delta)$ (mas yr <sup>-1</sup> )	$\mu_{\delta}$ (mas yr <sup>-1</sup> )	Pop.
PTFS1600y	$22.8 \pm 0.9$	$-88.7 \pm 0.9$	6.9	$2340 \pm 70$	$-7.3 \pm 2.0^{a}$	$-4.2 \pm 1.3^{a}$	Thin
PTFS1600ad	$29.7 \pm 1.4$	$-23.3 \pm 1.2$	7.2	$3770 \pm 180$	$1.1 \pm 1.5^a$	$-1.8 \pm 1.5^{a}$	Thin
PTFS1601p	$18.4 \pm 2.1$	$-45.9 \pm 1.8$	12.4	$4960 \pm 500$	$1.4 \pm 4.6^{a}$	$-7.6 \pm 4.2^{a}$	Thin/Thick
PTFS1501bh	$24.0 \pm 1.5$	$16.6 \pm 1.2$	8.2	$1280 \pm 70$	$10.1 \pm 1.5^a$	$-4.9 \pm 1.5^{a}$	Thin
PTFS1601cl	$35.2 \pm 2.1$	$-14.4 \pm 1.2$	9.1	$2890 \pm 90$	$2.0 \pm 1.3^{a}$	$2.2 \pm 1.3^{a}$	Thin
PTFS1607t	$26.7 \pm 1.7$	$31.0 \pm 1.2$	5.5	$2160 \pm 70$	$-2.1 \pm 0.3^a$	$1.6 \pm 1.6^{a}$	Thin
PTFS1607ab	$32.7 \pm 1.3$	$-37.6 \pm 1.1$	6.3	$1810 \pm 70$	$-3.1 \pm 2.6^{b}$	$-8.2 \pm 3.1^{b}$	Thin
PTFS1512bf	$31.4 \pm 1.9$	$70.3 \pm 1.5$	11.8	$1820 \pm 50$	$-19.2 \pm 6.4^{a}$	$4.7 \pm 5.7^{a}$	Thick
PTFS1617n	$17.7 \pm 1.7$	$-198.7 \pm 1.9$	7.2	$5700 \pm 380$	$-1.8 \pm 2.6^a$	$-2.9 \pm 2.7^{a}$	Thick/Halo
PTFS1617m	$13.1 \pm 2.4$	$-40.4 \pm 2.1$	10.9	$4060 \pm 180$	$-0.9 \pm 1.5^a$	$-9.2 \pm 1.7^{a}$	Thin/Thick
PTFS1619l	$22.7 \pm 1.3$	$-12.5 \pm 1.0$	5.8	$2040 \pm 140$	$-1.9 \pm 1.6^a$	$-2.0 \pm 1.6^{a}$	Thin
PTFS1521cm	$34.7 \pm 2.0$	$-6.5 \pm 1.9$	7.9	$2870 \pm 110$	$9.6 \pm 2.8^a$	$-7.8\pm2.7^a$	Thick




Figure 3.  $V_{\phi}$ – $V_{\rho}$  (left) and e– $J_z$  diagrams (right). The solid and dotted ellipses render the  $3\sigma$  thin and thick disk contours in the  $V_{\phi}$ – $V_{\rho}$ diagram, while the solid box in the e– $J_z$  marks the thick disk region as specified by Pauli et al. (2006).

error bars of the lightcurves. The orbital periods of the binaries range from 0.46 d to 3.8 d, with inclinations between 74-90 degrees. The radii of the primary stars divided by the semi-major axis  $(r_1)$  are typically 0.2–0.5, and the primary stars fill about 0.4-0.9 of their Roche lobe. The average density derived from the lightcurve is typically between 10-70 per cent of Solar density, consistent with A/F-type main sequence stars. The mass-ratio as determined from the lightcurves are typically between 0.08-0.2, but there are outliers to larger ratios. However, the uncertainties on the outliers are high. The mass-ratio determined from the amplitude of the inter-eclipse variability, which is in some cases not significant (e.g. 1700do) and explains the high uncertainty on the value for the mass-ratio is some cases. From the lightcurve we determined the temperature ratio of the two stars, assuming blackbody spectral energy distributions is typically 0.5–0.95.

The results of the SED fitting are shown in Table 4 and Fig. A2. The temperatures of the A/F-stars in the EL CVn systems range between 6600-10000 K, consistent with temperatures for A/F-type main sequence stars (Ftype:  $6000-7350\,\mathrm{K},\,\mathrm{A}$ -type:  $7350-10000\,\mathrm{K},\,\mathrm{Pecaut}\,\,\&\,\,\mathrm{Mama}$ -

jek 2013). The temperature of the pre-He-WDs range from 7900 to 17000 K. In all systems, the pre-He-WDs are hotter than the A/F-star companion. The uncertainty on the A/Fstar's temperature is typically 100-200 K. The temperature of the pre-He-WD is less well constrained (100–1400 K), because it depends on the availability of UV data and on how accurately the eclipse depth can be measured from the lightcurves. The RMS scatter between the data and model is typically  $\lesssim 5$  per cent, with a few outliers to 10 per cent (see Table A3). This residual scatter can be due to calibration differences between telescopes, but also because the observations are taken at a random phase. An observation taken in-eclipse results in a ~10 per cent lower flux as out of eclipse.

For 12 of the EL CVn systems we obtain usable radial velocity curves and measure the radial velocity amplitude, see Fig. A3. The remaining 7 systems were not enough measurements were obtained or they were observed at unfavourable orbital phases, precluding an accurate radial velocity measurement. However, all radial velocity amplitudes are low, in the range of 20–40 km s<sup>-1</sup>. This confirms that the secondary stars in these binaries are indeed low mass stars.

**Table 4.** System parameters of all EL CVn systems and the uncertainty (standard deviation) on the parameters. Systems for which a radial velocity measurement is used to calculate the parameters is indicated with the 'RV' superscript. This mainly affects the reliability and systematics on the mass and surface gravity of the pre-He-WD  $(M_2 \text{ and } \log g_2)$ .

Name	$P\left(\mathrm{d}\right)$	i (°)	$M_1  ({ m M}_{\odot})$	$M_2  ({ m M}_{\odot})$	$R_1 (R_{\odot})$	$R_2 (R_{\odot})$	$T_1(K)$	$T_2(K)$	$\log g_1$	$\log g_2$
1600y <sup>RV</sup>	1.1838920	84.5	1.62	0.17	2.41	0.46	6930	8900	3.88	4.33
	0.0000008	2.7	0.04	0.01	0.07	0.02	100	110	0.02	0.03
$1600 \mathrm{ad}^{\mathrm{RV}}$	1.0840448	86.5	1.76	0.23	1.83	0.35	8050	10490	4.16	4.72
	0.0000010	2.2	0.04	0.01	0.05	0.02	120	200	0.02	0.04
1700do	3.0507582	87.4	2.40	0.81	2.34	0.33	9890	17100	4.08	5.31
	0.0000278	1.8	0.06	0.25	0.13	0.03	90	1400	0.04	0.15
1600aa	0.6934558	78.7	1.67	0.50	1.67	0.55	7880	9300	4.21	4.67
	0.0000006	0.9	0.05	0.09	0.04	0.02	190	400	0.02	0.09
$1601\mathrm{p}^\mathrm{RV}$	1.2215885	83.8	1.82	0.14	1.65	0.34	8600	11700	4.26	4.54
	0.0000051	3.2	0.06	0.02	0.14	0.04	160	500	0.06	0.10
$1501\mathrm{bh^{RV}}$	0.6204144	78.4	1.30	0.12	1.23	0.20	6870	11100	4.38	4.91
	0.0000005	1.9	0.04	0.01	0.07	0.01	110	400	0.04	0.06
1601q	1.2515058	80.5	1.85	0.30	1.93	0.46	8300	10700	4.13	4.58
	0.0000051	2.9	0.08	0.18	0.15	0.04	200	700	0.05	0.40
$1601 {\rm cl}^{\rm RV}$	0.8917354	82.9	2.02	0.28	2.44	0.52	8290	10100	3.97	4.45
	0.0000005	2.9	0.06	0.01	0.07	0.02	200	300	0.02	0.03
1402de	0.6189694	87.0	1.61	0.36	1.56	0.45	7860	9300	4.27	4.69
	0.0000011	2.5	0.04	0.13	0.07	0.02	150	300	0.03	0.23
1607aa	0.8463120	84.6	1.85	0.30	1.81	0.38	8470	10300	4.19	4.76
	0.0000016	3.4	0.05	0.07	0.08	0.02	160	300	0.03	0.13
1607v	0.7198356	82.6	1.58	0.20	1.83	0.16	7260	10900	4.11	5.32
	0.0000020	5.8	0.06	0.05	0.16	0.03	120	500	0.06	0.24
$1607 t^{\rm RV}$	0.8759507	76.6	1.40	0.16	1.87	0.38	6600	8600	4.04	4.48
	0.0000004	1.0	0.05	0.01	0.05	0.01	140	200	0.02	0.03
$1607 \mathrm{ab^{RV}}$	0.7730986	83.8	1.40	0.19	1.45	0.32	6980	8810	4.26	4.71
	0.0000002	2.3	0.03	0.01	0.05	0.01	100	80	0.03	0.04
1608ab	0.6101718	86.8	1.50	0.11	1.39	0.52	7400	7900	4.32	4.04
	0.0000014	1.9	0.06	0.10	0.04	0.02	200	400	0.02	0.24
1612al	0.6369260	86.8	1.38	0.16	1.16	0.37	7280	10300	4.45	4.50
	0.0000006	2.0	0.08	0.09	0.05	0.02	110	300	0.03	0.22
$1512\mathrm{bf}^{\mathrm{RV}}$	0.6074343	87.2	1.39	0.17	1.53	0.32	6910	9740	4.21	4.65
	0.0000002	1.9	0.02	0.01	0.02	0.01	90	180	0.01	0.04
1613s	1.1420695	76.2	1.83	0.17	2.72	0.25	7350	13700	3.83	4.88
	0.0000024	6.0	0.08	0.05	0.30	0.05	140	800	0.08	0.28
1613u	0.5644902	81.6	1.52	0.27	1.65	0.37	7340	9690	4.19	4.73
	0.0000003	2.7	0.02	0.04	0.04	0.01	70	160	0.02	0.08
1615ag	0.6806897	85.7	1.52	0.27	1.63	0.32	7370	10200	4.20	4.87
	0.0000046	3.3	0.05	0.07	0.05	0.02	200	400	0.02	0.13
1615v	0.5594054	73.7	1.39	0.13	1.50	0.33	6920	9400	4.23	4.54
	0.0000003	1.4	0.03	0.03	0.04	0.01	120	300	0.02	0.17
1515ay	0.4642873	89.0	1.33	0.15	1.30	0.46	6800	7930	4.33	4.27
	0.0000001	1.1	0.03	0.04	0.02	0.01	100	150	0.01	0.10
1615w	1.4407151	77.7	1.61	0.24	2.59	0.40	6690	10300	3.82	4.63
	0.0000024	2.7	0.05	0.05	0.14	0.03	110	200	0.04	0.14
1615ao	0.8954515	77.6	1.64	0.41	1.82	0.64	7580	8700	4.13	4.43
	0.0000007	0.8	0.05	0.12	0.05	0.03	170	160	0.02	0.15
1615u	0.7777349	82.4	1.50	0.24	1.27	0.16	7400	12200	4.40	5.41
	0.0000011	4.2	0.09	0.10	0.14	0.02	200	600	0.08	0.27
1616cr	0.5649690	82.5	1.40	0.07	1.36	0.46	7060	8000	4.32	3.93
	0.0000002	0.9	0.03	0.02	0.02	0.01	120	170	0.01	0.07

Table 4 - continued

Name	$P\left(\mathrm{d}\right)$	i (°)	$M_1  ({\rm M}_{\odot})$	$M_2  ({ m M}_{\odot})$	$R_1 (R_{\odot})$	$R_2 (R_{\odot})$	$T_1(K)$	$T_2(K)$	$\log g_1$	$\log g_2$
1617n <sup>RV</sup>	2.3367776	87.3	1.80	0.18	2.41	0.38	7500	11600	3.93	4.55
	0.0000052	2.1	0.04	0.02	0.12	0.03	110	400	0.04	0.07
$1617\mathrm{m}^\mathrm{RV}$	3.7728999	87.8	1.68	0.14	2.57	0.69	6990	9320	3.84	3.89
	0.0000083	1.5	0.06	0.03	0.08	0.03	190	190	0.02	0.10
1619l <sup>RV</sup>	1.1599993	83.2	1.56	0.17	2.13	0.34	6870	9200	3.97	4.60
	0.0000017	4.1	0.05	0.01	0.14	0.04	120	150	0.05	0.09
1521ct	1.1724964	83.0	1.82	0.36	1.72	0.56	8520	9800	4.23	4.50
	0.0000014	1.4	0.06	0.29	0.11	0.04	190	300	0.04	0.28
1621ax	1.0181522	84.0	1.69	0.30	2.14	0.17	7350	11800	4.00	5.48
	0.0000045	4.6	0.06	0.07	0.14	0.03	170	700	0.04	0.22
$1521\mathrm{cm}^{\mathrm{RV}}$	0.6854774	80.0	1.49	0.21	1.49	0.43	7290	9240	4.27	4.49
	0.0000002	1.0	0.02	0.01	0.03	0.01	70	90	0.02	0.03
1622by	0.7486683	85.8	1.69	0.31	1.84	0.33	7700	11100	4.13	4.88
	0.0000016	3.4	0.07	0.07	0.07	0.02	300	1400	0.03	0.13
1522cc	0.5717853	81.2	1.40	0.26	1.62	0.27	6860	9570	4.17	4.99
	0.0000003	3.1	0.04	0.04	0.05	0.01	120	190	0.02	0.08
1622aa	0.7661291	84.7	1.60	0.16	1.74	0.26	7500	10900	4.16	4.85
	0.0000038	4.0	0.08	0.05	0.10	0.03	300	1300	0.04	0.19
1622bt	0.6884160	79.2	1.65	0.29	1.74	0.29	7700	12200	4.18	4.97
	0.0000004	2.0	0.06	0.04	0.05	0.01	200	1000	0.02	0.08
1723aj	1.1088064	85.6	1.57	0.18	2.51	0.23	6640	11000	3.84	4.98
	0.0000009	3.3	0.05	0.03	0.07	0.02	130	400	0.02	0.11




Figure 4. The temperature versus the average density of the main-sequence stars of the binary system, indicated with a black or white dot. The black lines show main-sequence evolution tracks (solar metallicity) from Spada et al. (2017) between 1.0 and  $2.5\,\mathrm{M}_\odot$  with  $0.1\,\mathrm{M}_\odot$  intervals. The colour map shows the mass of the star according to the interpolation method by Breton et al. (2012). The dashed lines are isochrones of 0, 0.1, 0.2, 0.5, 1, and 2 Gyr since the start of the main sequence.

Using all information available, we determine the stellar parameters of the stars in the EL CVn systems, summarised in Table 4. The masses of the A/F-type star range between 1.3 and  $2.4\,M_{\odot}$ . The radii of these stars  $(1.15{-}2.7\,R_{\odot})$  are consistent with these stars being regular main-sequence stars.

The radii of the pre-He-WDs range between 0.17–0.65  $R_{\odot}$ . To calculate the mass of the pre-He-WD  $(M_2)$ , we include the measured radial velocity amplitude if available,

which 'replaces' the uncertain mass-ratio measurement from the lightcurve. For most of the systems we do not have a radial velocity amplitude measurement, so we do depend on the mass-ratio to determine the mass of the pre-He-WDs, which range between 0.12 and  $0.5\,\rm M_{\odot}$ . As discussed in Section 4.1 and 4.5, the mass determination of the secondary using only the mass-ratio is very uncertain because of the high uncertainty on the mass-ratio. If we limit ourselves only to systems for which we have a radial velocity amplitude, the mass range is  $0.12\text{--}0.28\,\rm M_{\odot}$ , significantly smaller.

For the sample for which we have radial velocity curves, we determine the motion in the Galactic plane and derive their population membership as described in Sec. 4.4 and shown in Table 3. Fig. 3 shows that more than half of the systems are part of the thin disk population. A few are part of the thick disk, and PTFS1617n could also be a halo object.

#### 6 DISCUSSION

#### 6.1 Co-rotation

In the lightcurve modelling (Section 4.1), we assume that both stars are synchronised with the orbit. Previous studies of EL CVn binaries have made the same assumption, but all authors acknowledge that it might not be correct, since mass-accretion can spin up the A/F-star significantly (see Section 6.2). van Kerkwijk et al. (2010) extensively discuss how all parameter estimates are affected by incorrectly assuming co-rotation. Since the precision of PTF lightcurves is far lower than the precision of the Kepler lightcurves, the only significant effect this assumption has in our analysis is on the estimate of the mass-ratio. If a star is rotating faster than the orbital period, the mass-ratio (q) is overestimated.

We quantify this by simulating a typical EL CVn lightcurve with a primary star which is rotating 2 and 4 times faster than the orbital period, while keeping all other parameters the same. Fitting these lightcurves with the model assuming co-rotation, results in values for q of 0.02 and 0.10 higher the initial value of q=0.17. All other lightcurve parameters do not change significantly. We therefore conclude that for mildly faster-than-synchronous rotating primary stars ( $P_{\rm rot}/P_{\rm orb}>0.5$ ), the effect on the mass-ratio is similar or smaller than the statistical uncertainty on the mass-ratio. If the primary star is rotating faster, the mass-ratio is overestimated.

This overestimate propagates through to the rest of our parameter estimates; the average density of the primary is overestimated, and therefore the mass of the primary underestimated (Fig. 4), and the semi-major axis is overestimated. However, the effect is small since these parameters only weakly depend on the mass-ratio (see Equations 4 and 5). For a large part of our sample, we do not have any radial velocity amplitudes, and for these systems we rely on the mass-ratio to calculate the mass of the pre-He-WD. As mentioned in Section 4.5, the mass of the pre-He-WD depends on the mass-ratio to the third power. This, combined with a high statistical and systematic uncertainty on the mass-ratio, makes the calculations of  $M_2$  (without a radial velocity amplitude) unreliable.

To check if the A/F-stars are rotating faster than synchronous, we compare the rotation periods to the orbital period for stars in known EL CVn systems. The orbital period of the main sequence star has been determined for five Kepler EL CVn binaries by measuring the projected rotational velocity  $(v \sin i)$ : 1.79(60) d, 0.79(14) d, 5.0(2.4) d, and 1.71(62) d; (Faigler et al. 2015) and (0.93 d, Lehmann et al. 2013). In addition, the rotational period has tentatively been identified from a frequency analysis for KOI-81 (0.48 d Matson et al. 2015), KOI-74 (0.59 d Bloemen et al. 2012), KOI-1224 (3.49 d Breton et al. 2012), and KIC-8262223 (0.62 d Guo et al. 2017). All rotational periods are of the same order as the orbital period. A detailed comparison of the rotational and orbital period shows that most stars rotate faster than synchronous, but in one case the rotation period is longer than the orbital period. For three cases (all from Faigler et al. 2015) the rotation period is consistent with the orbital period of the binary, but since the uncertainties on the rotational periods are large, it is difficult to say if they are synchronised. The data therefore indicates that at least some (or maybe most) of the A/F-stars are not synchronised with the orbital period.

There is however an important difference between the PTF sample and the sample of EL CVn systems with known rotation periods of the primary (all found by Kepler). The relative size of the A/F-star  $(r_1)$  is a factor of  $\sim 3$  larger in the PTF sample, which strongly affects the synchronisation timescale of the star ( $\propto r_1^{-8.5}$ , Zahn 1977). We used the equation by Zahn (1977) and tabulated values for E<sub>2</sub> from Claret (2004) to calculate the synchronisation timescale for each of the EL CVn systems in our sample. This shows that the synchronisation timescale of the A/F-type star is less than 10 Myr in 20 systems, and less than 100 Myr for 32 systems. For these 32 systems, the synchronisation time is shorter than the time since mass transfer (0–260 Myr, see Fig. 5). The remaining 4 systems (with the smallest values for  $r_1$ )

have synchronisation timescales that are significantly longer than the estimated age. Based on this theoretical prediction, we can assume that that most of the A/F-stars are rotating synchronously. Whether this is actually the case requires an independent measurement of the rotation period.

#### 6.2 Binary evolution and stellar parameters

In the canonical formation channel of EL CVn binaries (e.g. Chen et al. 2017), two main-sequence stars of similar mass are born at a short orbital period of a few days. The more massive star evolves faster and increases in radius. Before it ascends the red giant branch (RGB), it fills its Roche lobe and starts stable mass-transfer to the lower mass secondary star. This process continues until almost the complete outer envelope is transferred (identified as "R CMa"-type binaries, e.g. Lee et al. 2016). The remnant of the initially more massive star has become a pre-white dwarf with a helium core and a thick hydrogen envelope ( $\approx 0.01 - 0.04 \,\mathrm{M}_{\odot}$ , see Istrate et al. 2016a; Chen et al. 2017). The accretor has become a rejuvenated main-sequence star of spectral type A or F which dominates the luminosity of the system. If present in a specific stellar population as e.g. found in clusters, such a system would be identified as a 'blue straggler'. If the orbital inclination is such that it shows eclipses, we identify it as an EL CVn binary.

The structure and evolution of pre-He-WDs have been extensively studied, as they also occur at more advanced binary evolutionary stages with either a white dwarf (Marsh et al. 1995) or a neutron star (van Kerkwijk et al. 2005) as companions. Modelling of the formation process of binaries with a low mass pre-He-WD (Althaus et al. 2013; Istrate et al. 2014a,b, 2016a; Chen et al. 2017) shows that there are strong correlations between the binary orbital period, and the mass, temperature, radius and age of the pre-He-WD. First, higher mass pre-He-WDs are formed at longer orbital periods. This is a direct result of the mass accretion process. This relation was already found in pre-He-WD – neutron star binaries and has been parametrised by Lin et al. (2011). Binary evolution studies by Istrate et al. (2014b) and Chen et al. (2017) also predict this P-M relation for EL CVn binaries. The latter shows that the relation between orbital period and mass is very robust, but at the low-mass end of the relation  $(0.16-0.20M_{\odot})$  there is some spread.

Pre-He white dwarfs (of a given mass) are also predicted to follow a particular evolutionary track, corresponding to a particular combination of radius and temperature as a function of age. The temperature and radius are directly related to the envelope mass and core mass of the white dwarf. Directly after the mass accretion process ends, the pre-He-WD is large ( $\gtrsim 0.5\,R_{\odot}$ ) and has a low ( $\lesssim 8000\,\mathrm{K}$ ) surface temperature. While the hydrogen envelope is slowly being consumed by shell burning, the pre-He-WD shrinks and increases in temperature while maintaining an approximately constant luminosity (this phase is often referred to as the constant luminosity phase). When almost the entire envelope has been consumed, the pre-He-WD starts to cool down while the radius keeps decreasing (the cooling track). At the beginning of the cooling track, multiple hydrogen shell flashes (H-flash, e.g. Driebe et al. 1998) can occur in the more massive pre-He-WDs. These flashes briefly increase the temperature and radius of the star, after which the white




Figure 5. Temperature versus radius of the pre-He-WDs, with the colours indicating different orbital periods. The coloured lines indicate evolution tracks by Althaus et al. (2013) for different masses. The dots indicate pre-He-WDs from this study, triangles are other low mass pre-He-WDs in EL CVn systems; upward pointing triangles indicate Kepler discoveries, downward pointing triangles indicate SWASP discoveries (for references, see Sec. 1). During the evolution of pre-He-WDs, multiple hydrogen shell flashes can occur, indicated as grey lines. The tracks before the first H-flash and after the last H-flash are shown as coloured lines. Isochrones are shown as dashed lines, counted from the end of mass-transfer in the binary. The solid black line indicates the approximate detection limits, estimated by assuming a  $T_1 = 7000 \,\mathrm{K}$ ,  $R_1 = 1.5 \,\mathrm{R}_{\odot}$  primary star. The bottom boundary is set by an eclipse depth of 0.03 mag in R-band, the top boundary is set by the requirement of a flat-bottom eclipse ( $R_1 > 2R_2$ ), and the left limit is set by the requirement that the flat-bottom eclipse is deeper than the secondary eclipse ( $T_1 < T_2$ ).

dwarf settles back on the cooling track. The exact mass boundary at which this starts to occur is uncertain. Models by Althaus et al. (2013) show shell flashes for masses above  ${\sim}0.18 M_{\odot},$  while Istrate et al. (2014b) put this boundary at  ${\sim}0.21 M_{\odot}.$ 

Fig. 5 shows the temperature versus radius of the pre-He-WD, with the colour of the points indicating the orbital period of the system. This shows that the temperatures and radii we find are consistent with predictions for pre-He-WDs in the constant luminosity phase, and before the occurrence of any H-flash. While some of the measurements are also consistent with pre-He-WDs undergoing an H-flash (grev lines), the short time spent in this phase makes this extremely unlikely. For the PTF sample, the orbital period (indicated by the colours) follows the same trend as the models, with long period systems containing larger and hotter pre-He-WDs. To test if the data match the models, we interpolate the models in orbital period, which allows us to test directly how well the radius and temperature match the model for a given orbital period. The fraction of measurements within 1, 3, and 5 standard deviations is 25, 75 and 86 per cent. Given the fact that we interpolate, and the uncertainties on radius and temperature could contain some systematic uncertainties, we conclude that most of the measurements agree with the models. This comparison to the models also allows us to infer the time since the end of mass transfer, which ranges from 0 to  $260\,\mathrm{Myr}$  with an average of  $110\,\mathrm{Myr}$ .

Fig. 6 shows the orbital period of the binary versus the mass of the pre-He-WD. The left panel shows the measured values, while the right panel shows the expected masses using the models inferred from the measured temperature and radius (Fig. 5). The measured values indicate that all pre-He-WD are low mass systems, but the PTF sample scatters around the model predictions. The right panel shows that if the radius and temperature measurements and models are used to derive the mass, the results fall within 10 per cent of the prediction of the mass-period relation for pre-He-WDs.

There are two possible explanations for this discrepancy. Either we have underestimated the uncertainties on the mass measurements, or there is some additional intrinsic spread on the predicted mass versus period, radius, and temperature not properly modelled. There are a number of possible systematic uncertainties that could affect the mass determination. First of all, we have assumed that the A/F-star is a regular main sequence star with a solar metallicity to estimate its mass (see Fig. 4). In Fig. 6 we have indicated how the mass estimate changes if we assume a different metallicity. If the real metallicity is lower than assumed, the mass is




Figure 6. Pre-He-WD mass versus the orbital period of the binary. The left panel shows the measured mass (using the radial velocity amplitude) of the PTF sample, the right panel shows the mass derived from the radius and temperature models (see Fig. 5). Large points indicate PTF systems with a radial velocity measurement, small points indicate systems without a radial velocity measurement (not shown in the right panel because of the high uncertainties). Grey triangles indicate mass-period measurements of other EL CVn systems; upward pointing triangles indicate Kepler discoveries, downward pointing symbols indicate SWASP discoveries (for references, see Sec. 1). The black line shows the period-mass relation by Lin et al. (2011), and the shaded area indicates a 10 per cent uncertainty on this relation.

overestimated. This could explain a part of the inconsistency with the theory, but extreme metallicities would be needed to explain the largest outliers. Since thick disk systems generally have a lower metallicity, the masses for these systems could be overestimated. However, the thick disk systems do not show any particular trend, indicating that this assumption may not be the dominant cause of the inconsistency with the model predictions.

Another possibility is that we have underestimated our measurement uncertainties. The mass of the pre-He-WD is mostly determined by the radial velocity measurement. As shown in Fig. A3 in the Appendix, we need to add an additional uncertainty to the formal uncertainties in order to explain all variance in the radial velocity measurements. For PTFS1601cl (one of the outliers), where we did not obtain radial velocity measurements at the quadratures, small systematic offsets between measurements can have a large impact on the radial velocity amplitude. We did check the radial velocity amplitude measurements of PTFS1512bf by obtaining a few high-resolution spectra with the Echellette Spectrograph and Imager (ESI) on the Keck telescope. The resulting radial velocity amplitude measured from these spectra is consistent with the result from the IDS spectra, which leads to the conclusion that uncertainties due to an unstable detector are most likely very small.

An alternative explanation is that there is some intrinsic variance between mass and period, radius and temperature. For example, Istrate et al. (2016b) shows that assumptions about rotation, diffusion and metallicity give different results when modelling the mass, radius and temperature of pre-He-WDs. The magnitude of the effect is estimated to be low, about 10 per cent. This would be enough to explain the variance in the right panel but cannot explain the outliers on the left panel.

To solve this ambiguity, a measurement of both the main-sequence and pre-He-WD radial velocity is needed. This allows the mass of both stars to be calculated by only using Kepler's law (combined with the period and inclination measurement from the lightcurve). This has been done for SWASP J0247-25 (Maxted et al. 2013), KOI-81 (Matson et al. 2015), KIC-10661783, (Lehmann et al. 2013), and KIC-8262223 (Guo et al. 2017). For SWASP J0247-25, KIC-10661783, and KIC-10661783 the mass of the pre-He-WD agrees well with the P-M relation, but for KOI-81, the mass is significantly lower  $(0.10 \rm M_{\odot})$  than the P-M relation predicts. This hints that there is more scatter in the P-M relation than models estimated.

#### 6.3 Galactic population and space density

Using stellar evolution and population synthesis codes, Chen et al. (2017) predict a space density of  $4-10 \times 10^{-6} \,\mathrm{pc}^{-3}$  for EL CVn binaries (including non-eclipsing ones) with orbital periods less than 2.2 d. In addition, they predicted that EL CVn binaries should mainly be found in young stellar populations, and therefore be more abundant in the thin disk. We use the Galaxy model based on SDSS data by Jurić et al. (2008) to estimate how many EL CVn binaries we would expect to see given this space density and in what ratios between thin disk, thick disk and halo. We populate our model Galaxy with stars with absolute magnitudes according to a normal distribution with a mean and standard deviation of  $R = 2.46 \pm 0.54$  mag, values determined from our sample of 36 systems. We simulate the PTF coverage by using (overly) simple requirements: |b| > 15,  $\delta > 0$ , 13.5 < R < 16 (see Fig. 1). We ignore the Galactic Plane because these fields tend to be observed only ~50 times. The minimum number of epochs in our uncovered sample is 58, indicating that at least 58 observations are needed to identify an EL CVn binary. Using the 58 epoch limit, we derive an effective coverage of 32.8 per cent for the remaining area. We also correct for the requirement that the systems must be eclipsing. This decreases the number of observable EL CVn systems by a factor of 0.307; determined from our sample using radii and inclination. Even if the binaries are eclipsing, if the pre-He-WD is too small (and thus old), we would not find it in the PTF data. To correct for this, we assume a lifetime of EL CVn binaries of 1 Gyr (the main sequence lifetime of a  $2 \,\mathrm{M}_{\odot}$ star) and compare this to the typical age of PTF EL CVn binaries (0-260 Myr). We therefore assume that PTF can only detect 26 percent of all EL CVn binaries.

According to the model and the assumed selection criteria, 26 per cent of the PTF sample should be from the thin disk. If we assume our model is correct, there is only a 1.8 per cent chance  $(\sum_{n=7}^{12} \binom{12}{n} 0.26^n [1-0.26]^{12-n})$  to find  $\geq 7$  thin disk systems out of a total of 12 EL CVn systems. If any of the ambiguous cases are from the thin disk, this probability drops well below 1 per cent. This indicates that our model is unlikely to be correct, and confirms that EL CVn systems are more abundant in the thin disk compared to the average stellar population, as was already suggested by Chen et al. (2017).

Using the model and estimated PTF detection efficiency, we also predict that we should have found  $\sim 300-750$ EL CVn systems, a factor of  $\sim 10-25$  higher than we actually recovered. This could simply be because we are overpredicting the contributions of the thick disk and halo. However, even if we assume a factor of 4 higher contribution from the thin disk (to bring the model in line with with the ratio of thin to thick disk systems), the model still predicts at least a factor of 5 - 12 more EL CVn systems than we found. Another uncertain estimate which could explain the discrepancy is the assumed efficiency of PTF in finding EL CVn binaries. The PTF observing cadence and coverage is highly inhomogeneous, and the assumptions we have used are very simple approximations. Assuming that we can find all EL CVn systems observed more than 58 times and are vounger than 200 Myr is overly optimistic, and could explain the discrepancy of a factor of 5 (or more).




Figure 7. (top) The temperature of the A/F-star versus the orbital period of the EL CVn system. (bottom) The distribution of orbital periods of the PTF sample (red) and the SWASP sample (blue). In the top left, the result of the KS-test and AD-test are shown (see text). The top panel shows that there is a strong correlation between orbital period and temperature, which is a result of the binary evolution process (see text). The histograms show that the PTF data is more biased to the short period systems compared to the SWASP sample. It also shows a possible gap at periods of 1 day, caused by a detection bias against these systems. For comparison, we also plotted the detection probability of an eclipsing population with well-sampled lightcurves ( $\propto P^{-2/3}$ , dotted line), and for lightcurves with a limited amount of epochs ( $\propto P^{-4/3}$ , dashed line).

The inhomogeneity of the PTF dataset makes it difficult to do a proper study of the Galactic distribution and space density of EL CVn systems. We do find tentative results that EL CVn systems occur more often in the thin disk, as was predicted by Chen et al. 2017. We also find that the space density is at the lower bound, or even below the prediction of  $4{\text -}10 \times 10^{-6}\,{\rm pc^{-3}}$ . To properly measure the properties of the population of EL CVn systems, a larger sample of EL CVn binaries is needed, preferably from a more uniform variability survey.

#### 6.4 Comparison with the SWASP sample

To better understand the biases of our survey, we compare the PTF sample to the sample found by SWASP (Maxted et al. 2014a). While we both have used the lightcurve characteristics to identify EL CVn binaries, there are some intrinsic differences between the surveys, and therefore different biases in finding EL CVn systems. The most obvious difference between the surveys is the magnitude range; SWASP probes magnitudes between 9–13 mag, while our sample is fainter, between 13.5 and 16 mag. A second major difference is the cadence and the number of epochs in a lightcurve; PTF lightcurves have an irregular cadence and a low num-

ber of epochs (~100) compared to the regular cadence and better-sampled SWASP lightcurves (~4000–13000 epochs).

There are indeed differences between the two samples. First , the distance range for the SWASP sample is 100–1200 pc, while the PTF sample reaches 1200–5000 pc. This is expected given the different magnitude range of the two surveys. We therefore also expect to find relatively more thick disk and halo systems compared to thin disk systems in the PTF sample. However, using the one-sided Fisher's exact test (Fisher 1934), we find no significant difference between the relative number of thin disk systems. This is consistent with our finding that EL CVn systems are more numerous in the thin disk compared to the average stellar population and explains why at larger distances it is still the most dominant population.

The two samples are also different with regards to orbital period and temperature of the A/F-star (which dominates the luminosity), see Fig 7. We performed a Kolmogorov-Smirnov (KS) test and Anderson-Darling (AD) test (e.g. Sec. 3.1 in Feigelson & Babu 2012) to compare the distribution of orbital periods. Both tests show that it is unlikely that the samples are drawn from the same distribution ( $p_{KS} = 0.013$  and  $p_{AD} = 0.005$ ). The histogram in Fig. 7 shows that we find more short orbital period systems and the top panel in Fig. 7 shows that at short orbital periods, the temperature of the primary star is low. This correlation can be understood because the mass of A/F-star is correlated with the orbital period. High mass main-sequence stars (2 M<sub>O</sub>) begin their main-sequence lifetime at a temperature of 9500 K, and cool down to 7500 K towards the end of their main sequence lifetime (see Fig. 4). Main sequence stars of  $1.3\,M_{\odot}$  start at a temperature of  $7500\,\mathrm{K}$  and only cools by 500 K during their time on the main sequence. Therefore, PTFs sensitivity for lower luminosity (lower temperature) EL CVn systems (partially) explains why the PTF sample contains more short period systems.

A second explanation is that, because of the sparse sampling, it is harder to find long period systems with PTF compared to SWASP. Both surveys use eclipses to find EL CVn binaries, and therefore are biased towards short period systems (Prob<sub>ecl</sub>  $\propto R_1 P^{-2/3}$ ). In addition, short period systems spend a larger fraction of their orbit in eclipse ( $\tau_{\rm ecl} \propto R_1 P^{-2/3}$ ). This does not bias the SWASP search since the lightcurves are well sampled. With PTF however, a lack of observations during the eclipse can hinder the identification of a system as an EL CVn binary.

The difference between the two samples shows that selection effects make it difficult to determine the intrinsic properties of EL CVn binaries. To do so requires an integrated approach: stellar evolution and population synthesis models should be used to simulate a sample of EL CVn binaries, which are then 'observed' by simulating the variability survey which was used to find the real sample. Such a calculation is difficult given the in-homogeneity of the PTF sample, and beyond the scope of this work.

#### 6.5 Pulsations

Pre-He-WDs are predicted to exhibit both p- and g-mode pulsations (e.g. Córsico & Althaus 2014, 2016; Córsico et al. 2016; Istrate et al. 2016b). Pulsations have been found in two of the SWASP EL CVn binaries: WASP 0247-25 (Maxted

et al. 2013; Istrate et al. 2017) and WASP 1628+10 (Maxted et al. 2014b). The pulsation periods are 5–10 minutes and the amplitudes  $\sim 1-2$  per cent of the pre-He-WD luminosity. Models of pre-He-WDs predict that in a large area in T-log g space, pre-He-WDs should feature pulsations (Fig. 10 in Córsico et al. 2016; Istrate et al. 2016b). Many of the pre-He-WDs in the PTF sample lie in this region, making them useful to test the general predictions for pulsation theory. In addition, because stellar parameters can be measured very precisely, a pulsating pre-He-WD in an eclipsing binary is extremely useful to test evolutionary and seismic models in great detail (e.g. Istrate et al. 2017).

Unfortunately, the very sparse sampling of the PTF lightcurves makes it very difficult to identify such pulsations. We did attempt to find pulsations by using a Lomb-Scargle algorithm (Lomb 1976; Scargle 1982, implementation by VanderPlas & Ivezić 2015) on the residuals of the lightcurves. We found periodic behaviour in the residuals at predicted periods of  $\sim\!10$  minutes for a number of the systems, but because of the sparse sampling and low amplitude, it is difficult to determine if these are real or not. High cadence follow-up photometry is needed to establish the reality of these pulsations.

#### 7 SUMMARY AND CONCLUSION

In this paper, we report the discovery and analysis of 36 new EL CVn systems extracted from Palomar Transient Factory data. With this sample, we more than double the number of known EL CVn systems. To find the EL CVn systems we used machine learning classifiers to make a pre-selection of candidates. This has proven to be an efficient method to minimise the number of lightcurves that have to be visually inspected.

The radii  $(0.16-0.7\,\mathrm{R}_{\odot})$  and temperatures (8000-17000 K) of the pre-He-WDs in the sample indicate they are all young systems in the "constant luminosity" phase (0-250 Myr) of their evolution. The masses of the pre-He-WDs are all low ( $<0.3M_{\odot}$ ), but our measurements show a large spread around the predicted mass-period relation, which remains unexplained. If we use the measured radii and temperatures combined with models, we do find masses consistent with the mass-period relation. This discrepancy is either due to systematic or underestimated uncertainties in our measurements, or there is more variance in the masses than the stellar evolution models predict. This problem can be resolved by obtaining more accurate radial velocity measurements (ideally for both stars in the binary to obtain an independent mass ratio measurement), and by more extensively testing the effect on the mass-period relation of e.g. different metallicities and rotation rates.

Although a detailed population study is difficult with the PTF dataset, we find that EL CVn binaries occur more often in the thin disk than an average Galactic stellar population. In addition, we find that the space density is most likely lower than the predicted value of  $4{\text -}10\times10^{-6}\,{\rm pc}^{-3}.$  To properly determine the properties of the EL CVn population, a more systematic search combined with stellar and Galactic modelling is required.

This new sample of young pre-He-WDs will be useful to put many theoretical models to the test, including stellar structure models for low mass white dwarfs, pulsation models, and binary evolution models. In addition, the methods we have used to identify EL CVn systems can easily be adapted to find other rare types of variable stars and these (and similar machine learning methods) will be vital to fully utilise (future) variability surveys like ZTF (Bellm 2014), NGTS (Wheatley et al. 2017), GOTO (Steeghs & Galloway 2017), BlackGEM (Bloemen et al. 2015), TESS (Ricker et al. 2015), PLATO (Rauer et al. 2014), and LSST (Ivezić et al. 2008).

#### ACKNOWLEDGEMENTS

We thank the referee, Simon Jeffery, for thoroughly reading the manuscript and providing us with useful comments and suggestions.

We thank Adam Miller, Fabian Gieseke and Tom Heskes for many useful discussions about machine learning classification. We thank Tom Marsh for the use of LCURVE. We thank Luc Hendriks for suggesting to use XGBOOST.

JvR acknowledges support from the Netherlands Research School of Astronomy (NOVA) and Foundation for Fundamental Research on Matter (FOM), and also the California Institute of Technology were a large part of this work was conducted. JvR and PJG thank the University of Cape Town for their hospitality: this work was finalised while visiting UCT, supported by the NWO-NRF Bilateral Agreement in Astronomy. This research was supported in part by the National Science Foundation under Grant No. NSF PHY-1125915 and PJG thanks the Kavli Institute for Theoretical Physics at the University of California, Santa Barbara for a productive stay.

The Intermediate Palomar Transient Factory project is a scientific collaboration among the California Institute of Technology, Los Alamos National Laboratory, the University of Wisconsin, Milwaukee, the Oskar Klein Center, the Weizmann Institute of Science, the TANGO Program of the University System of Taiwan, and the Kavli Institute for the Physics and Mathematics of the Universe.

Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS-IV acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS web site is www.sdss.org.

The Pan-STARRS1 Surveys (PS1) and the PS1 public science archive have been made possible through contributions by the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max-Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, the Queen's University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under Grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation Grant No.

AST-1238877, the University of Maryland, Eotvos Lorand University (ELTE), the Los Alamos National Laboratory, and the Gordon and Betty Moore Foundation.

This publication made use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

Based on observations made with the NASA Galaxy Evolution Explorer. GALEX is operated for NASA by the California Institute of Technology under NASA contract NAS5-98034.

This publication made use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

This research has made use of the VizieR catalogue access tool, CDS, Strasbourg, France. The original description of the VizieR service was published in A&AS 143, 23.

This research has made use of the SVO Filter Profile Service (http://svo2.cab.inta-csic.es/theory/fps/) supported from the Spanish MINECO through grant AyA2014-55216.

This research made use of Scikit-learn (Pedregosa et al. 2011).

This research made use of Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013).

IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under cooperative agreement with the National Science Foundation (Tody 1993).

#### REFERENCES

Althaus L. G., Miller Bertolami M. M., Córsico A. H., 2013, A&A, 557, A19

Andreas C. Müller S. G., 2016, Introduction to Machine Learning with Python. O'Reilly

Angeloni R., et al., 2014, A&A, 567, A100

Armstrong D. J., et al., 2016, MNRAS, 456, 2260

Astropy Collaboration et al., 2013, A&A, 558, A33

Bagnulo S., Jehin E., Ledoux C., Cabanac R., Melo C., Gilmozzi R., ESO Paranal Science Operations Team 2003, The Messenger, 114, 10

Bellm E., 2014, in Wozniak P. R., Graham M. J., Mahabal A. A., Seaman R., eds, The Third Hot-wiring the Transient Universe Workshop. pp 27–33 (arXiv:1410.8185)

Bianchi L., Conti A., Shiao B., 2014, VizieR Online Data Catalog, 2335

Bloemen S., et al., 2012, MNRAS, 422, 2600

Bloemen S., Groot P., Nelemans G., Klein-Wolt M., 2015, in Rucinski S. M., Torres G., Zejda M., eds, Astronomical Society of the Pacific Conference Series Vol. 496, Living Together: Planets, Host Stars and Binaries. p. 254

Breiman L., 2001, Machine Learning, 45, 5

Breton R. P., Rappaport S. A., van Kerkwijk M. H., Carter J. A., 2012, ApJ, 748, 115 Brown W. R., Kilic M., Allende Prieto C., Kenyon S. J., 2010, ApJ, 723, 1072

Budding E., Butland R., 2011, MNRAS, 418, 1764

Cardelli J. A., Clayton G. C., Mathis J. S., 1989, ApJ, 345, 245

Carter J. A., Rappaport S., Fabrycky D., 2011, ApJ, 728, 139

Chambers K. C., et al., 2016, preprint, (arXiv:1612.05560)

Chen T., Guestrin C., 2016, preprint, (arXiv:1603.02754)

Chen X., Maxted P. F. L., Li J., Han Z., 2017, MNRAS, 467, 1874 Claret A., 2004, A&A, 424, 919

Copperwheat C. M., Marsh T. R., Dhillon V. S., Littlefair S. P., Hickman R., Gänsicke B. T., Southworth J., 2010, MNRAS, 402, 1824

Copperwheat C. M., et al., 2011, MNRAS, 410, 1113

Córsico A. H., Althaus L. G., 2014, A&A, 569, A106

Córsico A. H., Althaus L. G., 2016, A&A, 585, A1

Córsico A. H., Althaus L. G., Serenelli A. M., Kepler S. O., Jeffery C. S., Corti M. A., 2016, A&A, 588, A74

Debosscher J., 2009, PhD thesis, Institute of Astronomy, Katholieke Universiteit Leuven, Belgium

Devor J., 2005, ApJ, 628, 411

Driebe T., Schoenberner D., Bloecker T., Herwig F., 1998, A&A, 339, 123

Faigler S., Kull I., Mazeh T., Kiefer F., Latham D. W., Bloemen S., 2015, ApJ, 815, 26

Feigelson E. D., Babu G. J., 2012, Modern Statistical Methods for Astronomy

Fisher R., 1934, Statistical Methods for Research Workers, 5 edn Foreman-Mackey D., 2013, Example: Fitting a Model to http://dan.iel.fm/emcee/current/user/line/ #maximum-likelihood-estimation

Foreman-Mackey D., 2014, Tutorial: model fitting with correlated noise, http://dfm.io/george/current/user/model/

Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, PASP, 125, 306

Geurts P., Ernst D., Wehenkel L., 2006, Mach. Learn., 63, 3 Gianninas A., Strickland B. D., Kilic M., Bergeron P., 2013, ApJ,

766, 3 Gies D. R., et al., 2008, ApJ, 682, L117

Guo Z., Gies D. R., Matson R. A., García Hernández A., Han Z., Chen X., 2017, ApJ, 837, 114

Hartman J. D., Bakos G. Á., 2016, Astronomy and Computing, 17, 1

Hogg D. W., Bovy J., Lang D., 2010, preprint, (arXiv:1008.4686)

Istrate A. G., Tauris T. M., Langer N., Antoniadis J., 2014a, A&A, 571, L3

Istrate A. G., Tauris T. M., Langer N., 2014b, A&A, 571, A45 Istrate A. G., Marchant P., Tauris T. M., Langer N., Stancliffe

Istrate A. G., Fontaine G., Gianninas A., Grassitelli L., Marchant P., Tauris T. M., Langer N., 2016b, A&A, 595, L12

Istrate A. G., Fontaine G., Heuser C., 2017, ApJ, 847, 130

R. J., Grassitelli L., 2016a, A&A, 595, A35

Ivezić v., et al., 2008

Ivezić Ž., Connolly A., Vanderplas J., Gray A., 2014, Statistics, Data Mining and Machine Learning in Astronomy. Princeton University Press

Jurić M., et al., 2008, ApJ, 673, 864

 Kaluzny J., Rucinski S. M., Thompson I. B., Pych W., Krzeminski W., 2007, AJ, 133, 2457

Kervella P., Fouqué P., 2008, A&A, 491, 855

Kovács G., Zucker S., Mazeh T., 2002, A&A, 391, 369

Laher R. R., et al., 2014, PASP, 126, 674

Law N. M., et al., 2009, PASP, 121, 1395

Lee J. W., Kim S.-L., Hong K., Koo J.-R., Lee C.-U., Youn J.-H., 2016, AJ, 151, 25

Lehmann H., Southworth J., Tkachenko A., Pavlovski K., 2013, A&A, 557, A79

Lin J., Rappaport S., Podsiadlowski P., Nelson L., Paxton B., Todorov P., 2011, ApJ, 732, 70

Lomb N. R., 1976, Ap&SS, 39, 447

Mackenzie C., Pichara K., Protopapas P., 2016, ApJ, 820, 138

Marsh T. R., Dhillon V. S., Duck S. R., 1995, MNRAS, 275, 828 Masci F., Bellm E., 2016, PTF light curve statistics, http: //www.ptf.caltech.edu/system/media\_files/binaries/30/ original/Objects SourcesTable cols v3.html

Masci F. J., Hoffman D. I., Grillmair C. J., Cutri R. M., 2014, A.J. 148, 21

Matson R. A., Gies D. R., Guo Z., Quinn S. N., Buchhave L. A., Latham D. W., Howell S. B., Rowe J. F., 2015, ApJ, 806, 155

Maxted P. F. L., et al., 2011, MNRAS, 418, 1156

Maxted P. F. L., et al., 2013, Nature, 498, 463

Maxted P. F. L., et al., 2014a, MNRAS, 437, 1681

Maxted P. F. L., Serenelli A. M., Marsh T. R., Catalán S., Mahtani D. P., Dhillon V. S., 2014b, MNRAS, 444, 208

McAllister M. J., et al., 2017, MNRAS, 464, 1353

Palaversa L., et al., 2013, AJ, 146, 101

Parsons S. G., Marsh T. R., Gänsicke B. T., Drake A. J., Koester D., 2011, ApJ, 735, L30

Pauli E.-M., Napiwotzki R., Heber U., Altmann M., Odenkirchen M., 2006, A&A, 447, 173

Pecaut M. J., Mamajek E. E., 2013, ApJS, 208, 9

Pedregosa F., et al., 2011, Journal of Machine Learning Research, 12, 2825

Peters C. M., et al., 2015, ApJ, 811, 95

Pietrzyński G., et al., 2012, Nature, 484, 75

Pollacco D. L., et al., 2006, PASP, 118, 1407

Rappaport S., Podsiadlowski P., Horev I., 2009, ApJ, 698, 666 Rappaport S., Nelson L., Levine A., Sanchis-Ojeda R., Gandolfi D., Nowak G., Palle E., Prsa A., 2015, ApJ, 803, 82

Rau A., et al., 2009, PASP, 121, 1334

Rauer H., et al., 2014, Experimental Astronomy, 38, 249

Richards J. W., et al., 2011, ApJ, 733, 10

Richards J. W., et al., 2012, ApJ, 744, 192

Ricker G. R., et al., 2015, Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003

SDSS Collaboration et al., 2016, preprint, (arXiv:1608.02013) Scargle J. D., 1982, ApJ, 263, 835

Schlafly E. F., Finkbeiner D. P., 2011, ApJ, 737, 103

Schlegel D. J., Finkbeiner D. P., Davis M., 1998, ApJ, 500, 525 Schwarzenberg-Czerny A., 1989, MNRAS, 241, 153

Sesar B., et al., 2017, AJ, 153, 204

Skrutskie M. F., et al., 2006, AJ, 131, 1163

Smart R. L., Nicastro L., 2013, VizieR Online Data Catalog, 1324 Spada F., Demarque P., Kim Y.-C., Boyajian T. S., Brewer J. M., 2017, ApJ, 838, 161

Steeghs D., Galloway D., 2017, Gravitational wave Optical Transient Observatory, https://www2.warwick.ac.uk/fac/ sci/physics/research/astro/research/goto/

Stetson P. B., 1996, PASP, 108, 851

Tody D., 1993, in Hanisch R. J., Brissenden R. J. V., Barnes J., eds. Astronomical Society of the Pacific Conference Series Vol. 52, Astronomical Data Analysis Software and Systems II. p. 173

VanderPlas J. T., Ivezić Ž., 2015, ApJ, 812, 18

Westera P., Lejeune T., Buser R., Cuisinier F., Bruzual G., 2002, A&A, 381, 524

Wheatley P. J., et al., 2017, preprint, (arXiv:1710.11100)

Wright E. L., et al., 2010, AJ, 140, 1868

Zacharias N., Monet D. G., Levine S. E., Urban S. E., Gaume R., Wycoff G. L., 2004, in American Astronomical Society Meeting Abstracts. p. 1418

Zacharias N., Finch C., Frouard J., 2017, VizieR Online Data Catalog, 1340

Zahn J.-P., 1977, A&A, 57, 383

van Dokkum P. G., 2001, PASP, 113, 1420

#### 18 J. van Roestel

van Kerkwijk M. H., Bassa C. G., Jacoby B. A., Jonker P. G., 2005, in Rasio F. A., Stairs I. H., eds, Astronomical Society of the Pacific Conference Series Vol. 328, Binary Radio Pulsars. p. 357 (arXiv:astro-ph/0405283)

van Kerkwijk M. H., Rappaport S. A., Breton R. P., Justham S., Podsiadlowski P., Han Z., 2010, ApJ, 715, 51 von Zeipel H., 1924, MNRAS, 84, 665

## APPENDIX A: ADDITIONAL TABLES AND FIGURES

 ${\bf Table}\ {\bf A1.}\ {\bf List}$  of a features used by the machine learning classifiers.

#### PTF variability

wRMS (mag)	weighted root-mean-square of the PTF
	lightcurve
skew (mag)	skewness of the PTF lightcurve
medAbsDev (mag)	median absolute deviation of the PTF
	lightcurve
StetsJ	Stetson-J statistic of the PTF lightcurve
StetsK	Stetson-K statistic of the PTF lightcurve
Neumann	the Von Neumann ratio statistic of the
	PTF lightcurve
MBR.	median buffer range: the fraction of points
	more than 20% of the lightcurve ampli-
	tude from the weighted mean magnitude,
	divided by total number of epochs.
$AMBS\{1,2,3\}$	The fraction of lightcurve points that are
111111111111111111111111111111111111111	# standard deviation above the mean
	magnitude.
BMBS{1,2,3}	The fraction of lightcurve points that are
DMD5(1,2,5)	# standard deviation below the mean
	magnitude.
	S
$prange\{\#\} (mag)$	range containing {90,80,65,50,35,20} per
( // )	cent of the data.
$percentile\{\#\}$	#th percentile minus the me-
	dian of the PTF lightcurve, di-
	vided by prange90 , with # in
	$\{5,10,17.5,25,32.5,40,60,67.5,75,82.5,90,95\}.$

#### PAN-STARRS colours

PSr (mag)	Pan-STARRS $r$ - median of the lightcurve
PSgr (mag)	Pan-STARRS $g-r$
PSri (mag)	Pan-STARRS $r-i$
PSiz (mag)	Pan-STARRS $i-z$
PSzy (mag)	Pan-STARRS $z-y$
2MASS colours	
J (mag)	2MASS $J$ - median of the lightcurve
JH (mag)	2MASS $J - H$
HK (mag)	2MASS $H - K$

Table A2. Overview of the nights at the INT with the IDS.

Date	Grating	CCD	seeing (")	weather
2016-09-07	R632V	RED+2	0.6	excellent
2016-09-08	R632V	RED+2	0.7	excellent
2016-09-09	R632V	RED+2	0.6	excellent
2016-09-10	R632V	RED+2	0.7	good
2016-09-11	R632V	RED+2	0.6 - 1.0	good
2016-09-12	R632V	RED+2	0.7 - 1.0	good
2016-09-13	R632V	RED+2	0.8 – 1.2	good
2016-09-14	R632V	RED+2	1.0	good
2016-12-14	R900V	RED+2	0.8-1.4	good
2016-12-15	R900V	RED+2	1.4	ok
2017-01-09	R900V	RED+2	0.8	good
2017-01-10	R900V	RED+2	1.2 – 2.6	ok-bad
2017-03-10	R900V	EEV10	1.5-3.0	bad
2017-03-11	R900V	EEV10	_	clouds
2017-03-12	R900V	EEV10	2-4	bad
2017-03-13	R900V	EEV10	1.5	bad

**Table A3.** The temperatures of the A/F-star  $(T_1)$  and pre-Helium white dwarf  $(T_2)$  determined from the spectral energy distribution of the binary stars, see Fig. A2. The E(B-V) values are taken from Schlegel et al. (1998); Schlafly & Finkbeiner (2011), with an uncertainty of 0.034 (as in Maxted et al. 2011). The 'RMS' column indicates the additional uncertainty added to account for all variance, which is achieved by the parameter f in Equation 2.

ID	$T_1$ (K)	$T_2$ (K)	E(B-V)	RMS
1600y	$6930 \pm 100$	$8900 \pm 110$	0.047	0.05
1600ad	$8050 \pm 120$	$10490 \pm 200$	0.024	0.04
1700 do	$9890 \pm 110$	$17100 \pm 1400$	0.015	0.03
1600aa	$7890 \pm 190$	$9300 \pm 400$	0.107	0.02
1601p	$8600 \pm 160$	$11700 \pm 500$	0.030	0.03
1501bh	$6870 \pm 110$	$11100 \pm 400$	0.035	0.03
1601q	$8300 \pm 230$	$10700 \pm 700$	0.081	0.03
1601cl	$8280 \pm 200$	$10100 \pm 300$	0.030	0.04
1402 de	$7870 \pm 150$	$9300 \pm 300$	0.100	0.03
1607aa	$8470 \pm 160$	$10300 \pm 300$	0.087	0.04
1607v	$7260 \pm 130$	$10900 \pm 500$	0.090	0.03
1607t	$6600 \pm 140$	$8600 \pm 200$	0.009	0.10
1607ab	$6980 \pm 100$	$8810 \pm 80$	0.005	0.09
1608ab	$7360 \pm 240$	$7900 \pm 400$	0.037	0.05
1612al	$7280 \pm 110$	$10300 \pm 300$	0.039	0.01
1512bf	$6920 \pm 90$	$9740 \pm 180$	0.022	0.03
1613s	$7350 \pm 140$	$13700 \pm 800$	0.051	0.04
1613u	$7340 \pm 70$	$9690 \pm 160$	0.006	0.04
1615ag	$7380 \pm 200$	$10200 \pm 400$	0.093	0.05
1615v	$6920 \pm 120$	$9400 \pm 300$	0.030	0.03
1515ay	$6800 \pm 100$	$7930 \pm 150$	0.029	0.04
1615w	$6690 \pm 110$	$10300 \pm 200$	0.046	0.04
1615ao	$7580 \pm 170$	$8700 \pm 160$	0.070	0.03
1615u	$7400 \pm 200$	$12200 \pm 600$	0.069	0.06
1616cr	$7060 \pm 120$	$8000 \pm 170$	0.095	0.10
1617n	$7500 \pm 110$	$11600 \pm 400$	0.022	0.03
$1617 \mathrm{m}$	$6990 \pm 190$	$9320 \pm 190$	0.071	0.04
1619l	$6870 \pm 120$	$9200 \pm 150$	0.031	0.04
1521ct	$8520 \pm 180$	$9800 \pm 300$	0.090	0.02
1621ax	$7340 \pm 170$	$11800 \pm 700$	0.108	0.04
$1521 \mathrm{cm}$	$7290 \pm 80$	$9240 \pm 90$	0.051	0.02
1622by	$7730 \pm 260$	$11100 \pm 1400$	0.079	0.07
1522cc	$6860 \pm 120$	$9570 \pm 190$	0.042	0.06
1622aa	$7490 \pm 290$	$10900 \pm 1300$	0.160	0.11
1622bt	$7750 \pm 220$	$12200 \pm 1000$	0.061	0.07
1723aj	$6640 \pm 130$	$11000\pm400$	0.061	0.04

**Table A4.** The parameters of the lightcurve models shown in Fig. A1. This table shows the median and root-mean-square of final 5120 points in our MCMC chains. Note that these distributions are not normally distributed and parameters can be strongly correlated.

ID Band	$P(\mathbf{d})$	$t_0  (\mathrm{BMJD_{tdb}})$	i (°)	q	$\langle r_1 \rangle$	$\langle r_2 \rangle$	$T_2/T_1$	$\operatorname{absorb}_R$	absorb <sub>g'</sub>	$\log(f_R)$	$\log(f_{g'})$	fill	$\rho(\rho_{\odot})$
1600y R+g	1.1838920 0.0000008	55570.2084 0.0006	84.5 2.7	0.12 0.02	0.421 0.010	0.081 0.003	0.73 0.03	0.8 0.2	0.8 0.2	-2.1 0.0	-2.5 0.1	0.748 0.019	0.114 0.007
1600ad R+g	1.0840448 0.0000010	56247.4677 0.0006	86.5 2.2	0.11 0.03	0.328 0.008	0.063 0.003	0.80 0.03	1.3 0.5	1.0 0.5	-2.3 0.0	-2.4 0.1	0.575 0.025	0.292 0.019
1700do R	3.0507595 0.0000281	55556.8044 0.0014	87.4 1.8	0.33 0.10	0.178 0.010	0.025 0.002	0.77 0.04	2.8 1.2		-2.3 0.0		0.380 0.020	0.187 0.024
1600aa R+g	0.6934558 0.0000006	56892.6368 0.0003	78.7 0.8	0.29 0.05	0.391 0.007	0.128 0.004	0.94 0.01	3.6 0.8	3.2 0.8	-2.5 0.2	-2.6 0.4	0.801 0.026	0.362 0.021
1601p R+g	1.2215885 0.0000051	57152.5232 0.0010	84.0 3.1	0.21 0.09	0.274 0.019	0.056 0.005	0.84 0.03	1.4 1.0	1.1 1.1	-2.2 0.0	-2.3 0.3	0.534 0.036	0.358 0.059
1501bh R+g	0.6204144 0.0000005	55097.3927 0.0008	78.4 1.9	0.14 0.04	0.356 0.017	0.058 0.004	0.64 0.04	0.5 0.3	0.4 0.2	-2.1 0.1	$-2.1 \\ 0.0$	0.646 0.023	0.674 0.085
1601q R+g	1.2515054 0.0000051	57190.1373 0.0012	80.6 2.7	0.12 0.07	0.308 0.021	0.074 0.007	0.91 0.03	1.6 1.1	2.2 1.2	-2.3 0.1	-2.5 0.3	0.535 0.049	0.266 0.051
1601cl R+g	0.8917354 0.0000005	56063.3087 0.0004	82.9 2.9	0.11 0.02	0.475 0.011	0.101 0.004	0.95 0.02	2.3 0.8	1.9 0.7	-2.1 0.0	-2.3 0.0	0.830 0.021	0.142 0.009
$\begin{array}{c} 1402 \mathrm{de} \\ \mathrm{R} \end{array}$	0.6189694 0.0000011	55768.8152 0.0008	87.0 2.4	0.17 0.07	0.405 0.015	0.119 0.006	0.91 0.03	2.5 1.1		-2.9 0.3		0.757 0.052	0.454 0.049
1607aa R+g	0.8463124 0.0000017	56246.6579 0.0007	84.4 3.6	0.14 0.03	0.375 0.017	0.079 0.005	0.92 0.03	2.2 1.0	3.7 0.9	-2.6 0.3	-2.6 0.2	0.678 0.024	0.314 0.035
1607v R+g	0.7198355 0.0000020	55769.1206 0.0014	82.7 5.7	0.13 0.04	0.447 0.037	0.039 0.007	0.66 0.08	2.0 1.0	2.2 1.0	-2.4 0.1	-2.3 0.0	0.808 0.038	0.256 0.048
1607t R+g	0.8759507 0.0000004	56158.7102 0.0004	76.6 1.0	0.09 0.02	0.417 0.009	0.085 0.003	0.74 0.03	0.7 0.2	0.5 0.2	-2.3 $0.0$	-2.3 0.0	0.712 0.020	0.220 0.014
$^{1607ab}_{R+g}$	0.7730986 0.0000002	55151.7862 0.0004	83.8 2.3	0.18 0.03	0.351 0.010	0.077 0.003	0.77 0.03	1.9 0.4	1.5 0.4	-2.3 0.1	$-2.2 \\ 0.0$	0.665 0.020	0.439 0.035
1608ab R	0.6101718 0.0000014	57034.9178 0.0003	86.8 1.9	$0.07 \\ 0.07$	0.390 0.006	0.146 0.004	$0.90 \\ 0.02$	0.4 0.3		-3.0 0.3		0.656 0.058	0.551 0.044
1612al R	0.6369260 0.0000006	55782.6928 0.0007	86.8 2.0	0.12 0.06	0.322 0.009	0.103 0.004	0.69 0.03	0.1 0.1		-2.3 0.1		0.574 0.043	0.880 0.084
1512bf R+g	0.6074343 0.0000002	56100.9311 0.0002	87.2 1.9	0.10 0.02	0.438 0.005	0.091 0.003	0.66 0.03	0.1 0.1	0.1 0.1	-2.1 0.0	-2.2 0.0	0.762 0.017	0.391 0.012
1613s R+g	1.1420695 0.0000024	56511.1762 0.0014	76.2 6.0	0.10 0.03	0.470 0.050	0.044 0.008	0.60 0.05	1.2 0.5	1.2 0.5	-2.2 0.1	$-2.2 \\ 0.0$	0.812 0.050	0.091 0.023
1613u R+g	0.5644902 0.0000003	56787.5255 0.0003	81.6 2.7	0.18 0.03	0.472 0.012	0.107 0.004	0.77 0.03	1.4 0.4	1.6 0.4	-2.4 0.1	-2.5 0.2	0.889 0.023	0.340 0.024
1615ag R+g	0.6806898 0.0000046	55380.8645 0.0010	85.8 3.3	0.17 0.04	0.410 0.013	0.079 0.005	0.76 0.05	2.2 1.0	1.9 0.8	-2.2 0.3	-2.4 0.1	0.772 0.026	0.358 0.029
1615v R+g	0.5594054 0.0000003	54962.6621 0.0007	73.7 1.4	0.10 0.02	0.457 0.012	0.099 0.004	0.71 0.03	0.2 0.1	0.1 0.1	-2.2 0.1	-2.9 0.3	0.784 0.029	0.410 0.031
1515ay R+g	0.4642873 0.0000001	56138.1751 0.0002	89.0 1.1	0.11 0.03	0.452 0.003	0.161 0.002	0.83 0.02	0.0 0.0	0.1 0.1	-2.2 0.1	$-1.9 \\ 0.0$	0.794 0.027	0.606 0.018
1615w R+g	1.4407151 0.0000024	56530.5582 0.0010	77.7 2.7	0.15 0.04	0.393 0.021	0.060 0.004	0.60 0.04	0.7 0.3	0.4 0.2	-2.9 0.3	-2.5 0.1	0.723 0.026	0.092 0.015
1615ao R+g	0.8954515 0.0000007	56308.1400 0.0005	77.6 0.8	0.24 0.06	0.368 0.007	0.129 0.005	0.94 0.02	2.1 1.1	1.9 1.0	-2.4 0.3	-2.5 0.2	0.729 0.038	0.273 0.020
1615u R+g	0.7777349 0.0000011	56185.6219 0.0011	82.4 4.2	0.16 0.07	0.297 0.034	0.038 0.006	0.59 0.05	1.4 0.7	1.8 0.8	$-2.1 \\ 0.1$	$-2.2 \\ 0.0$	0.554 0.041	0.726 0.191
$_{\rm R+g}^{1616\rm cr}$	0.5649690 0.0000002	55972.1677 0.0002	82.5 0.9	0.05 0.01	0.416 0.005	0.141 0.002	0.84 0.02	0.4 0.2	0.5 0.1	-2.3 0.2	-3.0 0.2	0.657 0.021	0.558 0.025

 ${\bf Table}~{\bf A4}-{\it continued}$ 

ID Band	$P\left(\mathbf{d}\right)$	$t_0 \left( \mathrm{BMJD}_{\mathrm{tdb}} \right)$	i (°)	q	$\langle r_1 \rangle$	$\langle r_2 \rangle$	$T_2/T_1$	$\operatorname{absorb}_R$	absorb <sub>g'</sub>	$\log(f_R)$	$\log(f_{g'})$	fill	$\rho(\rho_{\odot})$
1617n	2.3367776	55591.4345	87.3	0.21	0.258	0.040	0.63	2.6	1.5	-2.1	-2.4	0.501	0.117
R+g	0.0000052	0.0027	2.1	0.07	0.012	0.003	0.04	0.8	0.5	0.1	0.1	0.025	0.013
$_{\rm R+g}^{1617\rm m}$	3.7728999 0.0000083	56584.2146 0.0010	87.8 1.5	0.24 0.15	0.206 0.005	$0.056 \\ 0.002$	0.68 0.03	0.6 0.5	0.7 0.4	-2.1 0.1	$-2.2 \\ 0.1$	0.410 0.046	0.085 0.011
1619l R+g	1.1599993 0.0000017	56560.2644 0.0016	82.7 4.2	0.14 0.04	$0.385 \\ 0.022$	0.061 0.006	0.67 0.04	1.2 0.4	3.6 0.8	-2.2 0.1	-2.3 0.3	0.704 0.029	0.153 0.021
1521ct	1.1724964	56907.7345	83.2	0.17	0.282	0.093	0.92	1.2	2.2	-2.3	-2.9	0.528	0.373
R+g	0.0000013	0.0005	1.4	0.09	0.009	0.004	0.02	0.8	1.1	0.1	0.3	0.046	0.044
1621ax	1.0181525	56741.7492	83.1	0.17	0.401	0.029	0.57	2.9	2.2	-2.2 0.0	-2.2	0.758	0.171
R+g	0.0000044	0.0017	4.8	0.04	0.028	0.006	0.07	0.9	1.1		0.1	0.031	0.027
1521cm	0.6854774	56068.9363	80.0	0.22	0.381	0.110	0.78	0.7	0.8	-2.9	-2.6	0.740	0.428
R+g	0.0000002	0.0002	1.0	0.04	0.008	0.003	0.02	0.3	0.2	0.3	0.2	0.025	0.027
1622by	0.7486683	55718.0157	85.6	0.16	0.424	0.076	0.81	3.5	3.1	-2.4	-2.9	0.791	0.270
R+g	0.0000015	0.0010	3.6	0.03	0.017	0.005	0.04	0.9	0.8	0.3	0.3	0.024	0.025
1522cc R+g	0.5717853 0.0000003	56641.9749 0.0003	81.2 3.1	0.18 0.03	0.472 0.013	0.078 0.004	0.69 0.04	1.4 0.3	1.2 0.3	-2.4 0.1	-2.6 0.2	0.894 0.018	0.331 0.023
1622aa	0.7661291	57136.1682	84.7	0.10	0.410	0.060	0.75	0.6	1.4	-2.4 0.2	-2.9	0.716	0.302
R+g	0.0000038	0.0014	4.0	0.03	0.022	0.006	0.05	0.5	0.7		0.3	0.026	0.038
1622bt	0.6884160	56746.2432	79.2	0.17	0.425	0.071	0.70	1.7	1.4	-2.3	-2.5	0.795	0.314
R+g	0.0000004	0.0003	2.0	0.02	0.012	0.003	0.03	0.4	0.3	0.0	0.2	0.017	0.023
1723aj R+g	1.1088064 0.0000009	56733.1351 0.0004	85.6 3.3	0.11 0.02	0.460 0.011	0.042 0.003	0.52 0.05	0.5 0.2	0.5 0.2	-2.4 0.0	-3.1 0.2	0.818 0.022	0.100 0.006




Figure A1. The PTF lightcurves in R (left) and g' (right) with the best model over-plotted (see Table A4).




Figure A1-continued




Figure A1-continued







Figure A1 - continued




 ${\bf Figure}~{\bf A1}-{\it continued}$ 




MNRAS 000, 1–?? (2017) Figure A2. The spectral energy distribution and the best-fitting model spectra (see Table A3). The grey lines show the SED of the A/F-star and pre-He-WD. The black line shows the sum of both components. The A/F-star dominates the SED over the whole wavelength range, except in the far-UV in some of the cases.




 $\mathbf{Figure}\ \mathbf{A2}-\mathit{continued}$ 



 $\mathbf{Figure}\ \mathbf{A2}-\mathit{continued}$ 



 $\mathbf{Figure}\ \mathbf{A2}-\mathit{continued}$ 






Figure A3. The radial velocity measurements with the INT and the best fitting model. Black errorbars show the estimated uncertainty from the cross-correlation procedure, while the red errorbars show the uncertainties required to account for all residual variance. The shaded grey contours show the 1, 2 and 3 standard deviation intervals of model, obtained using the larger uncertainties.




Figure A3 - continued